

Evolution of Maud Rise Polynya during the last 250 years – a multiproxy ice core reconstruction from coastal Dronning Maud Land, Antarctica

Rahul Dey^{1,2}, Chavarukonam M. Laluraj¹, Kenichi Matsuoka³, Ashish Paiguinkar¹, Bhikaji L Redkar¹ and Meloth Thamban¹

¹National Centre for Polar and Ocean Research (NCPOR), Ministry of Earth Sciences, Vasco da Gama, Goa 403804, India

²Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen, Copenhagen N 2200, Denmark

³Norwegian Polar Institute, Framsentret, Postboks 6606, Langnes, 9296 Tromsø, Norway

Correspondence to: Rahul Dey (rdey1801@gmail.com)

Abstract

Open ocean polynyas drive deep ocean convection, influencing regional carbon and heat budgets, which in turn influence the ocean circulation and overall climate of Antarctica. The Maud Rise Polynya (MRP), also known as the Weddell Polynya, is one such polynya that forms in the Southern Ocean during early spring or winter months. The extensively studied MRP opening, which occurred during 2016–2017 and 1974–1976–76, triggered intense convection, ventilating heat from the deep ocean and modifying water mass properties. However, polynya evolution ~~beyond~~before the satellite era remains poorly understood. Here, we develop a polynya index using multiple proxy records from an ice core in coastal Dronning Maud Land, East Antarctica. Our approach—integrating records of snow accumulation, $\delta^{18}\text{O}$, deuterium excess, ~~NassNa~~ flux, and Na/SO₄ ratios—enhances polynya reconstruction, thereby overcoming the limitations of single-proxy methods. The index replicates the 1974–1976–76 polynya and extends the record to 1774, revealing three major events comparable to the 1974–1976–76 great polynya event, in the past 250 years, totalling ~~47~~twenty five polynya years-(likelihood > 0.6). We identified distinct clusters of polynya activity, possibly corresponding to a specific combination of atmospheric circulation patterns and oceanographic preconditioning for MRP development. This study offers a long-term perspective on MRP variability, providing insights into its drivers and climate-related impacts.

1. Introduction

Open ocean polynyas represent one of the most dynamic and influential features in the Southern Ocean system, serving as critical regulators of heat and gas exchange between the ocean and atmosphere. These persistent areas of open water within the sea ice cover act as windows through which complex oceanographic and atmospheric processes interact, influencing global ocean circulation, carbon cycling, and climate patterns (Bennetts et al., 2024; The SO-CHIC consortium, 2023; Zheng et al., 2021). The significance of polynyas extends beyond their immediate vicinity, influencing deep-water formation, marine ecosystem dynamics, and ~~global climate processes~~the intensification of sea-to-air heat and moisture fluxes during the winter months. In the Southern Ocean, polynyas manifest in two distinct forms: (1) coastal polynyas, which form along the Antarctic margin through mechanical forcing by katabatic winds, and (2) open-ocean polynyas, which develop hundreds of kilometres offshore through complex thermodynamic processes. While coastal polynyas occur regularly and have been extensively studied (Arrigo and van Dijken, 2003; Årthun et al., 2013; Jacobs et al., 1979; Visbeck et al., 1996; Xu et al., 2023), open-ocean polynyas represent rare but profoundly influential events that can significantly impact global ocean circulation patterns. The Maud Rise Polynya (MRP), also known as Weddell Polynya, occurring over the Maud Rise seamount in the eastern Weddell Sea (Fig. 1), is the most enigmatic open-ocean polynya in the Southern Ocean (Holland, 2001). First documented through satellite observations in the 1970s, the MRP reached its largest extent during the winter and early spring months from 1974 to 1976, exceeding 300,000 km² at its peak (Carsey, 1980; Zwally et al., 1985). This event drove intense ocean convection, reaching depths of over 3000 meters, and fundamentally altered water mass properties throughout the region, influencing global ocean circulation patterns (Gordon and Comiso, 1988).

Following this dramatic episode, the Maud Rise region has exhibited intermittent polynya activity, typically manifesting as halos of reduced sea ice concentration (Lindsay et al., 2004; McHedlishvili et al., 2022). However, the phenomenon garnered renewed attention when a large polynya exceeding 40,000 km² reappeared during the two winters of 2016 and 2017 (Jena et al., 2019). Early polynya studies focused primarily on documenting polynya occurrence and extent through satellite imagery (Carsey, 1980; Comiso and Gordon, 1987; Zwally et al., 1985), while more recent works have attempted to unravel the complex mechanisms controlling polynya formation and maintenance (de Lavergne et al., 2014; Wilson

et al., 2019). Satellite observations have revealed that open-ocean polynya formation often follows a characteristic pattern, beginning with the appearance of small-scale openings that can rapidly expand under favourable conditions, such as wind-driven upwelling (de Lavergne et al., 2014; Jena et al., 2019), baroclinic instabilities (Akitomo, 2006; Campbell et al., 2019), and deep warm water intrusions (Gülk et al., 2024; Heuzé et al., 2021). However, the precise triggers initiating this process remain incompletely understood, because the relative importance of these processes likely varies temporally and may be interconnected (Campbell et al., 2019; McPhee, 2003; Narayanan et al., 2024), and also because large polynyas openings are rare and satellite-based observations range only for the past few decades, limiting the observational opportunity. Similarly, model-based assessments are limited in their ability to understand polynya formation in the weakly stratified Southern Ocean (Heuzé et al., 2013; Sallée et al., 2013). This modelling challenge hampers our ability to assess both the role of polynyas in natural climate variability and their potential responses to future climate change.

To obtain longer polynya records, Goosse et al. (2021) examined surface mass balance (SMB) records reconstructed from six ice cores and two automatic weather stations. In these records, they ~~found that~~observed increased snowfall during the 2016-17 MRP opening and assumed that ~~this had also~~similar anomalies occurred ~~in~~during the past MRP opening periods. Their study constructed a polynya index using two data assimilation approaches and one statistical method to reconstruct historical climate states based on SMB reconstruction from ice cores. Their study found multiple polynya openings since 1250 CE, but observed that large polynya openings are rare. However, they rely only on snow accumulation records, which can be influenced by many factors other than polynya occurrences. The polynya signal in their ice-core SMB records is also weak compared to natural atmospheric variability, making it difficult to distinguish polynya-induced anomalies from SMB caused by other climatic fluctuations. The short instrumental record from automatic weather stations prevents a robust calibration of polynya reconstructions, increasing the risk of false positives (unrelated high snowfall events misinterpreted as polynya activity) and false negatives (missed polynya events). Additionally, the ice-core and instrumental records they used were collected from a large region, including the Weddell Sea sector and the ice sheet inland, where the surface elevation exceeds 2000 m a.s.l., which can hardly reflect changes in the vapour sources in the MRP regions directly (Fig. 1).

To overcome the limitations of the previous study, we develop a novel, multi-proxy approach to reconstruct past MRP activity using a high-resolution ice core, IND-36/9, from Djupranen Ice Rise in coastal Dronning Maud Land. While polynya opening results in increased heat and moisture exchange, the polynya years, characterised by a decrease in sea ice cover, also result in increased sea salt production. These signatures can be carried to the coastal region of Antarctica and deposited in the low-elevation ice shelves and ice rises. The back-trajectory analysis (Section 2.5) clearly indicates the direct atmospheric link between MRP and the IND-36/9 core site (Fig. 1). Our ice core site is situated within ~1000 km of the typical MRP location, close enough to potentially capture atmospheric signatures transported from polynya formation near the Maud Rise. By combining records of snow accumulation, water isotopes, and major ion concentrations, we aim to provide a more robust reconstruction of polynya activity. This approach leverages the strengths of selecting an ~~idea~~ideal core site and employing multiple proxies to overcome the limitations of single-proxy reconstructions, thereby capturing the complex signatures of polynya events that extend beyond the last few decades, when satellite and instrumental records are available.

2. Study Area and Methods

2.1. Study area

The DML coast is characterised by distinct topographic features like ice rises having associated local ice flow, climate regime and SMB variability (Drews et al., 2015; Goel et al., 2017; Lenaerts et al., 2014; Matsuoka et al., 2015; Pratap et al., 2022; Rignot et al., 2019). As part of the Indo-Norwegian project MADICE, an ice core was drilled on the summit of the Djupranen Ice Rise (70.18° S, 9.18° E; elevation 321 m a.s.l.), at the western margin of the Nivlisen Ice Shelf in coastal Dronning Maud Land (DML), East Antarctica (Fig. 1). Our site survey found that this ice rise summit has been at the current position at least in the past few millennia (Pratap et al., 2022). This location also offers several advantages for polynya reconstruction: its high SMB allows annual layer counting (Dey et al., 2023) and its coastal proximity may provide sensitivity to maritime signals (Ejaz et al., 2021; Wauthy et al., 2024), while its elevation causes insignificant surface melt and local noise while maintaining regional signal strengths (Dey, 2023). Its position is also within primary atmospheric transport pathways from the Maud Rise region (Fig. 1).

2.2. Ice core: from the field to the laboratory

The drill site of the Djupranen ice rise was located at its summit, based on analysis of satellite altimetry data and satellite image analysis, followed by an ice-penetrating radar survey (Pratap et al., 2022). The radar survey also ~~revealed~~shows a flat bed topography below the summit (Goel et al., 2026) and the presence of Raymond arches, indicative of a relatively stable summit position in the past (Goel et al., 2020). The ice core was drilled using an electromechanical ice core drilling system (Model D2, GeoTec, Japan). Over nine days, a 122 m ice core (IND 36/B9; hereafter IND36/9) was retrieved. The drilled ice core sections were sealed in high-density polyethylene core bags, packed in expanded polypropylene boxes, and stored in a reefer container at -20°C until their transport to the National Centre for Polar and Ocean Research (NCPOR), Goa, India, where samples were stored in the in-house Ice Core Laboratory maintained at -20°C . The ice cores were processed in the -15°C core processing facility at NCPOR. The cores were initially cut into 3.5 cm thick, 10 cm wide slabs for line scanning and then subsequently sub-sampled at 5 cm resolution for stable isotope and chemical analysis. The samples for chemical analysis were cut into cuboids; the three dimensions of the samples were measured using a calliper and weighed using a weighing balance. The density for the samples was calculated as mass divided by the volume of each sample. The error in measuring the sample dimensions was ± 0.5 mm, while the weighing balance's uncertainty was ± 0.1 g for measurements up to 100 g. As a result, the density measurements have a propagated uncertainty of 5%.

Major inorganic anions (Cl^{-} , SO_4^{2-} , MSA^{-} and NO_3^{-}) and cations (Na^{+} , NH_4^{+} , K^{+} , Mg^{2+} , and Ca^{2+}) were measured in the samples using an ICS 5000+ ion chromatograph (Thermo Dionex) equipped with a conductivity detector. Anions were separated on an AS11 (2 mm) column with potassium hydroxide as eluent and an AG11 (2 mm) guard column with AERS 500, 2 mm suppressor. Cations were separated on a CS17 (0.4 mm) capillary column with methane sulphonic acid as the eluent and the CG17 (0.4 mm) capillary guard column with a CCES 300 capillary suppressor. A 10 mg/L stock solution of Na^{+} , NH_4^{+} , K^{+} , Ca^{2+} , and Mg^{2+} was mixed and then diluted with MiliQ ultrapure water to prepare standards for cation exchange chromatography. The anion standards were prepared from 10 ppm stock solutions of MSA^{-} , Cl^{-} , SO_4^{2-} , and NO_3^{-} . The dilutions were conducted volumetrically and were freshly prepared within a few days of each run, ranging from 5 ppb to 1 ppm. Eight standards were

selected from this range for calibration. Before analysis, the ice core samples were melted in a Class 100 clean room. The analytical precision for all ions was better than 10%.

The ice core samples were analysed for oxygen and hydrogen isotopic ratios at NCPOR using a Triple Isotope Water Analyser (TIWA-45EP from Los Gatos Research, USA), which works on the principle of off-axis integrated cavity output spectroscopy (OA-ICOS). The melted ice core samples were introduced into the TIWA-45EP without sample conversion through a PAL HTC-xt auto-injector (CTC Analytics) equipped with a heated ($\approx$ approximately 85 °C) injector block (LGR) (Berman et al., 2013). Using a Hamilton 1.2 μ L, zero-dead-volume syringe, samples were injected into the injector block and evaporated for direct isotope analysis. Measurements were completed at a speed of $\sim$ 90 s per individual injection. To eliminate sample-to-sample memory, a total of nine injections were made, with the first three being discarded for analysis. The last six injections were averaged to produce a single, high-throughput (HT) sample measurement. One commercially available working standard from LGR1C and two in-house laboratory standards (CDML1 and HL1) with known isotopic composition, spanning the entire range of our sample measurements (-46.19‰ to -19.49‰ for $\delta^{18}\text{O}$ and -362.85‰ to -154.0‰ for δD) were analysed routinely as reference waters after every five ice core samples to check the instrument performance. Laboratory standards are calibrated on the VSMOW/SLAP scale. The external precision obtained using our laboratory standards (CDML1 and HL1) for $\delta^{18}\text{O}$ was $\pm 0.046\text{‰}$ and $\pm 0.068\text{‰}$, respectively, and for δD was $\pm 0.32\text{‰}$ and $\pm 0.23\text{‰}$ (1σ standard deviation) for 30 samples. Replicate analyses performed based on ten samples yield repeatability of $\pm 0.76\text{‰}$ for δD and $\pm 0.09\text{‰}$ for $\delta^{18}\text{O}$. All the raw instrumental OA-ICOS data were processed in the LGR post-analysis software. Any measured injection with water number density outside the manufacturer's suggested range of $2\text{--}4.5 \times 10^{16}$ H_2O molecules/ cm^3 was discarded. Injections with incomplete evaporation were detected by examining the standard deviation of the measured water number density (σ_{nmeas}) as reported by the instrument (Berman et al., 2013). Processed raw data directly gives $\delta^{18}\text{O}$ and δD , which are further used to calculate deuterium excess [$\text{d-excess} = \delta\text{D}_{\text{ice}} - 8 \times \delta^{18}\text{O}_{\text{ice}}$].

2.3. Chronology development

The chronology of the IND36/9 core is based on a multiproxy approach involving annual layer determination from the stratigraphy of $\delta^{18}\text{O}$, major ions, and pixel intensity data from the line scanner, following Dey et al. (2023). The core has experienced limited surface

melt, with annual melt proportion varying between 0 and 4.4%, with a median melt proportion of 0.25% (Dey et al., 2023). Since diffusion in the firn attenuates high-frequency water-isotope information in ice cores (The Firn Symposium team, 2024), even in high accumulation sites of coastal Antarctica (Mahalinganathan et al., 2022), we diffusion-corrected our water isotope records following Jones et al. (2023) as detailed in the supplementary materials (Fig. S1). The shift in seasonal peaks in our corrected record is less than 5 cm, which falls within the sampling interval of 5 cm and is therefore insignificant in affecting the accuracy of the chronology.

A five-point smoothing was applied to the pixel intensity data to simplify annual layer counting and reduce noise in the record. We also used age tie points of known volcanic eruptions identified from non-sea-salt sulphate (nssSO_4) peaks, as well as the tritium bomb peak of 1962. Volcanic indicators (nssSO_4) have been used to identify specific, dated volcanic eruptions, allowing us to reduce the uncertainties resulting from the relative dating procedure. However, unambiguous eruption identifications are challenging in ice cores from coastal regions, where the nssSO_4 background signals are commonly highly variable due to the proximity of the ocean and ocean-related MSA products (Philippe et al., 2016).

A preliminary chronology was obtained from the annual counts, which was then refined using the nssSO_4 peaks (Fig. 2) from the well-established volcanic eruptions of Pinatubo (1991), El Chichon (1982), Agung (1963), Cerro Azul (1932), Santa Maria (1902), Krakatoa (1882), Cosiguina/Babuyan (1834), Tambora (1815) and unknown volcanic eruption (1809). Similar to Dey et al. (2023), we refined the chronology between the volcanic and Tritium tie points using the StratiCounter algorithm (Winstrup et al., 2012). The manual counts provide a basic framework from which StratiCounter develops and refines its statistical characterisation of annual layers, enabling adaptation to varying layer properties with depth (Winstrup et al., 2012). To minimise reliance on initial manual inputs, we conducted multiple iterations using refined layer templates derived from previous algorithm outputs. The chronology of the IND36/9 ice core provides a robust temporal framework that extends back to 1774 CE at a depth of 122m (Fig. 3).

2.4. Satellite-based polynya metrics

We followed Heuzé et al. (2021) to define the "polynya-prone" region of the Weddell Sea (6°W to 12°E, 68°S to 60°S, approximately 600 km x 900 km), focusing on the winter and early spring months (June 1 to October 31). This region was selected based on historical observations of polynya formation and its significance in Antarctic bottom water production (Campbell et al., 2019; Heuzé et al., 2021). Polynya activity was quantified using two independent metrics derived from satellite data. These metrics complement each other to provide a comprehensive assessment of polynya dynamics.

~~The first metric, Polynya Days, represents the annual count of days (between June 1 and October 31) with sea ice concentration below 60% in our study zone. This threshold was determined through sensitivity analysis of satellite imagery and validation against in situ observations from previous studies (Heuzé et al., 2021). Sea ice concentration data were derived from passive microwave measurements using the NASA Team algorithm (Comiso and Nishio, 2008), which has demonstrated robust performance in discriminating between open water and sea ice in polar regions.~~ To quantify the polynya metrics, we use sea ice concentration data derived from passive microwave measurements using the NASA Team algorithm (Comiso and Nishio, 2008), processed at a spatial resolution of 25 x 25 km per pixel. The first metric, Polynya Days, represents the annual count of days (between June 1 and October 31), with a minimum sea ice concentration in the polynya-prone region below 60%. The second metric, Cumulative Polynya Area, measures the total extent of all polynya occurrences within a year (June 1 to October 31). This ~~metric~~ was calculated by summing the daily open water areas within the polynya ~~areas~~prone region, defined as ~~contiguous regions~~the cumulative sum of pixel area with sea ice concentration below the 60% threshold. ~~The combination of these metrics allows us to characterise both the temporal persistence and spatial extent of polynyas, providing insights into their formation mechanisms and potential impact on regional oceanographic processes.~~ The combination of these metrics allows us to characterise both the temporal persistence and spatial extent of polynyas, providing insights into their formation mechanisms and potential impact on regional oceanographic processes.

2.5. Air mass trajectory modelling

The HYbrid Single-Particle Lagrangian Integrated Trajectory (HySPLIT) model,

developed by the NOAA Air Resources Laboratory (ARL), provides a means of generating back trajectories to identify the source of moisture uptake for precipitation (Markle et al., 2012). We used HYSPLIT version 5.3 for back trajectory computations, initialized with conditions from the 2.5 by 2.5-degree resolution NCEP/National Center for Atmospheric Research (NCAR) reanalysis data for the period 1948 to 2016. Forward trajectories were run for 240 hours (10 days), initialised every hour, starting over the polynya-prone region of the Weddell Sea. Groups of forward trajectories were computed with initial altitudes of 100, 200, 300, 400, and 500 meters above ground level, and in all cases, the vertical velocity from the meteorological data was used as the input for vertical motion.

We calculated trajectory density from multiple trajectories using HYSPLIT (Miller et al., 2002), because individual trajectories are highly sensitive to meteorological uncertainties, turbulence, and small initial condition variations, often leading to misleading or unreliable airflow pathways and potential position errors of up to 20% of the distance travelled. It also aids in ~~discriminating~~differentiating between local circulation patterns and regional-scale flow features (Dorling and Davies, 1995) and has been previously applied to interpret polar ice core paleoclimate records (Dixon et al., 2012; Ejaz et al., 2021; Neff and Bertler, 2015). We calculated air mass transport densities for each year 1979 – 2016 from the HySPLIT output. The trajectory ~~endpoints~~densities in each equal-area (1 by 1 degree) ~~oceanic~~ pixel were summed and divided by the total number of air mass trajectories.

3. Results

3.1. Identification of polynya years in satellite data

Satellite imageries ~~are~~is crucial for polynya studies as they provide continuous, high-resolution observations of sea ice dynamics, allowing researchers to monitor polynya formation, extent, and variability over time. Satellite observations spanning four decades reveal distinct periods of polynya activity in the Weddell Sea region since 1979 (Fig. 4). The most notable and well-documented event occurred during 1974–~~1976~~ – 76, when the polynya reached an exceptional size of 300,000 km² (Carsey, 1980). This event, often referred to as the Great Weddell Polynya, represented a significant perturbation to the regional ocean-atmosphere system and has served as a benchmark for subsequent polynya observations. Following this major event, several smaller polynyas and polynya-like features ("halos") have

been observed during the late 1980s to early 1990s and early 2000s (Heuzé et al., 2021). The two independent metrics used for polynya detection—Polynya Days and Cumulative Area—reveal distinct but complementary temporal patterns. This dual-metric approach enables a more comprehensive understanding of polynya dynamics than either metric alone could provide. While the magnitudes of these metrics do not exhibit direct correlation ($r = 0.42$, $p < 0.01$), both metrics successfully identify known major polynya/halo events between 1979 and 2016, and provide unique insights into their temporal and spatial characteristics. The Polynya Days metric effectively captures persistent small-scale features, while the Cumulative Area metric better represents brief but large openings. While the record of polynya openings from satellite imagery provides crucial insights, their influence on coastal Antarctic ice cores could depend majorly on atmospheric transport.

3.2. Air mass transport patterns

We use forward air trajectory analysis to examine the transport pathways and determine whether polynya-derived signals can reach and be recorded in the ice cores. This approach allows us to trace the path of air masses from the polynya-prone area to potential ice core sites, providing insight into which locations are most likely to capture polynya signals. Forward trajectory analysis reveals consistent and well-defined transport pathways from the Maud Rise region to coastal Dronning Maud Land (Fig. 5). Frequency analysis ~~indicates~~shows that our study site (1 x 1 degree box surrounding the ice core site) receives approximately 2% of all trajectories originating from the MRP region, a statistically significant proportion ($p < 0.001$) that indicates reliable capture of polynya-related atmospheric signals. This percentage remains relatively stable across different seasons and years, suggesting a robust atmospheric connection between the source and deposition regions. The trajectory density analysis indicates that our ice core location falls within a primary atmospheric transport corridor originating from the MRP region. This corridor exhibits enhanced stability during the winter months when polynya formation typically occurs. However, trajectory calculations are susceptible to significant spatial error of 15 – 30% of distance travelled (Draxler, 2008). Therefore, a combination of forward and backward trajectories would provide a better overview of the travel pathways. Our back-trajectory analysis (Fig. 6) further confirms the findings from the forward trajectory patterns, showing trajectories originating from the polynya-prone region influencing our ice core site. We observe that air masses reaching our site also originate from more distant open-

water areas (even during the winter months) in the Southern Ocean. However, these distal sources remain relatively consistent across the study period and are located significantly further from the IND-36/9 site compared to the polynya-prone region. Consequently, these background marine inputs are less likely to drive the synchronized, high-magnitude anomalies observed across all proxies, particularly the sharp increases in ssNa flux, that characterize the polynya years. Trajectory analysis of the transport pathways, therefore, confirms that winds can carry polynya-derived signals from the open water to our study site and preserve them in the ice core records, regardless of polynya presence.

3.3. Development of the polynya index

3.3.1. Ice core record and proxy selection

The ice core proxy dataset from 1774 to 2016 exhibits distinct variability across all measured properties. We choose ~~five~~^{four} major proxies for developing our polynya index: sea-salt sodium (ssNa), deuterium excess (d-excess), $\delta^{18}\text{O}$, and snow accumulation, ~~and the Na/SO₄ ratio~~ (Fig. 7a). Sodium concentration ranges from a low of 19.401 to 218.646 ppb, with a mean value of 89.879 ppb. Some periods show sustained higher values, such as the late 1700s and early 1800s, while others, including the late 19th and 20th centuries, display more variable levels. D-excess varies between -1.838 to 10.758‰, with a mean of 4.313‰, showing notable peaks in the late 19th century and sharp declines in the early 20th century. $\delta^{18}\text{O}$ values fluctuate between -21.646 to -15.656‰, with a mean of -17.858‰, displaying alternating periods of enrichment and depletion, including a gradual decline over recent decades. Annual snow accumulation ranges from 0.12 to 0.83 m w.e., with a mean of 0.39 m w.e., exhibiting distinct fluctuations, including lower values in the early 20th century and a more variable pattern in recent decades. ~~The Na/SO₄ ratio ranges from 0.02 to 1.31, with a mean of 0.42, exhibiting periodic fluctuations throughout the dataset.~~

~~Our polynya index incorporates the five selected parameters, each chosen based on established physical mechanisms that link polynya formation to specific signatures in ice core records.~~ The opening of a polynya results in increased heat and moisture exchange between the warm, open water and the cold air. Therefore, during polynya years (decrease in the sea ice cover), there is also an increase in local precipitation in the Weddell Sea region (Moore et al., 2002) and resultant snow accumulation further inland (Goosse et al., 2021). The water isotope

ratios of precipitation are often related to temperature at the precipitation site (Ejaz et al., 2022; Naik et al., 2010) and the distance of transport (Goursaud et al., 2017; Klein et al., 2019), and the pattern of annual mean water isotopes of precipitation associated with polynya formation would relatively be similar to that for increased temperature from the heat exchange due to polynya opening and shorter transport distance, especially over a coastal site. Extensive While sea ice typically acts as a physical barrier to air-sea exchange, a polynya creates a localized window for enhanced aerosol production via multiple distinct mechanisms. First, the open-water surface provides a direct source where high-speed cyclonic winds and storm activity trigger bubble-bursting processes, injecting sea-salt particles into the marine boundary layer. Second, extensive open water polynyas can also act as a large factory for sea ice production, comparable to the production of the largest coastal polynyas (Zhou et al., 2023). ~~Previous studies have demonstrated that the surface of fresh sea ice, including frost flowers, is the major~~ (Zhou et al., 2023). Previous studies have shown that the surface of fresh sea ice, including frost flowers, is an important source of sea salt to the Antarctic, and the production of frost flowers is controlled by the amount of new sea ice production (Wolff et al., 2003). ~~Therefore, the formation of large open-ocean polynyas would lead to an increased supply of sea-salt sodium to the ice core site.~~ (Wolff et al., 2003). Furthermore, blowing snow across these newly formed, brine-wetted ice surfaces facilitates the sublimation of salty snow particles, which is also the major source of sea salt aerosol to the inland sites (Frey et al., 2020). Because these processes occur proximately to the drilling site and are situated directly along the atmospheric trajectories identified in Figure 6, they result in high-magnitude pulses of ssNa that are chemically distinct from the far, year-round open-ocean baseline. We however use ssNa flux instead of concentration to mitigate the potentially confounding effects of varying annual snow accumulation rates. By using flux, we ensure that a dilution effect during high-snowfall years doesn't artificially lower our ssNa signal, and conversely, that a low-snowfall year doesn't create a false peak.

~~Additionally, Sulphate (SO₄) fractionation during freezing of seawater (ice) is another well-known process to characterise sea ice coverage (Richardson, 1976). It is driven by the crystallisation of Na₂SO₄·10H₂O (mirabilite) occurring at temperatures below -8.2°C. Since the salt crystals are associated with the ice lattice, the remaining brine becomes increasingly depleted in SO₄ (and to a lesser extent in Na) as it progresses below this critical temperature. This process, therefore, increases the Na/SO₄ ratio of the sea salt source relative to bulk~~

seawater by preferentially removing sulphate ions (Wagenbach et al., 1998). Since this process only occurs at temperatures below -8.2°C , temperatures not encountered at the summer ocean surface, but frequently encountered at the winter sea ice surface (Levine et al., 2014). The anomalously high Na/SO_4 ratio can be used as an indicator of sea salt sourced from the production of fresh sea ice at the polynyas.

3.3.2. Development of the integrated polynya index

Our ice core record shows an anomalous increase in all proxy records during the prominent polynyas of 1964 and 1974–1976 (Fig. 7a). However, in the case of the transient polynyas and halos in more recent years, the relative changes are weaker but still identifiable. This could be due in part to the inherent complexity of air–ocean interactions and transportation processes, as well as the uncertainty associated with the measurements and chronology of the ice core. Therefore, to develop a more robust proxy, rather than using the proxies individually, we derive a multi-proxy index to quantify the likelihood of polynya occurrence more accurately.

The polynya index is defined using anomalous changes in snow accumulation, $\delta^{18}\text{O}$, deuterium excess, Na_{flux} , and Na/SO_4 ratio as possible indicators of polynya activity. We used Monte Carlo simulations to account for uncertainties associated with these individual proxies and ultimately to constrain better the influence of these uncertainties on the multi-proxy polynya index. This is because individual measurement uncertainties are independent of chronological uncertainties. We run ten thousand simulations to minimise statistical bias and define background data as the sum of a 30-year running median (RM) and two times of median absolute deviations (MAD). To reconstruct past MRP activity, we developed a multi-proxy index using a cumulative-anomaly approach, calculated as the means of rolling z-scores. This allows us to isolate anomalies that are synchronized across proxies from long-term climate trends. We use four independent proxies: sea-salt sodium (ssNa) flux, $\delta^{18}\text{O}$, deuterium excess (d-excess), and annual snow accumulation rates for our polynya index calculation. Prior to index calculation, all proxies are scaled to 0 – 1 to remove bias toward proxies with greater variability. For each proxy, we first calculate a 30-year rolling z-score to identify annual departures from the local background state. This accounts for non-stationary signals in the 250-year record, such as the gradual depletion of water isotopes or recent decreases in snow accumulation rates. As explained in section 3.3.1, MRP openings are linked to positive

anomalies in ssNa flux, $\delta^{18}\text{O}$, d-excess, and annual snow accumulation rates; therefore, we used only positive z-scores for the calculation. Negative z-scores were set to zero to prevent them from cancelling out physically significant peaks in other proxies. To test the dependence of the polynya index on snow accumulation, we calculate two versions of the index: one using all four selected proxies and another excluding annual snow accumulation rates. The polynya index is a mean of all individual rolling z-scores, which is then normalized so that the final values are such that the largest known polynya opening of 1974 – 76 is set to 1. A polynya year (a year when a polynya activity is recorded) is defined as a year in which the polynya index exceeds the likelihood threshold (0.6, 0.8, or 1). The polynya index is, therefore, an indicator of polynya occurrence from ice core observations and not a measure of the absolute extent or duration of the polynya events.

3.4 Past Maud Rise Polynya activity

Our ~250-year reconstruction reveals complex and significant temporal variability in polynya occurrence, shedding new light on the long-term behaviour of the Maud Rise Polynya (MRP). We identified ~~forty seven distinct twenty-five potential~~ polynya years from 1774 to 2016 (likelihood >0.6; Fig. 7b); ~~however, the period from 1920 to 1950 was the longest interval without significant polynya activity in our record. The 1830s were the most active decade, with seven polynya years recorded between 1830 and 1840. This high frequency of three events suggests that atmospheric and oceanic conditions during this period were highly conducive to polynya formation. Similar clusters occurred in the 1880s, with five polynya years, and in the 1970s, with four polynya years, including the well documented 1974–1976 sequence (Fig. 7b). These clusters suggest potential links to large-scale climate oscillations or recurring oceanographic conditions that favour polynya development. The reconstruction also identifies two past events (1833–1884 and 1911) that produced index values comparable to or (1788, 1834 and 1901) exceeding those the threshold of 1.0, indicating that these were events of possibly similar magnitude as the well documented 1974–1976 – 76 polynya opening.~~ The high index values for these events likely indicate that they were large-scale and persistent openings; however, exact quantification of the polynya extent is beyond the scope of this study. During the late 18th to mid-19th century, the polynya exhibited a high-frequency variability, frequently dropping to near-zero levels as seen in 1782 and the early 1800, while also producing moderate activity clusters above the 0.60 threshold in the 1810s and late 1820s. The late 19th century

transitioned into a period of more consistent activity, with 1887 being a notable high-activity year exceeding 0.80. The early 20th century was marked by a prominent polynya in 1901 followed by significant bursts of activity exceeding the 0.60 threshold in 1911, 1924, and 1926. Mid-century activity was relatively suppressed, though a significant spike above 0.90 occurred in 1951. The lesser known 1964 polynya opening is also recorded in our polynya index (1965 in our polynya index); however, the likelihood is relative lower (>0.6) than the 1974 – 76 event. Following the 1974 – 76 event, the post-1980 era has generally shifted toward lower average values, though moderate spikes exceeding the 0.60 threshold occurred in 1982, 2003, 2006, and 2010, corresponding to know halo years.

4. Discussion

4.1. Performance of the new polynya index

We compared our polynya index with those from Goosse et al. (2021) and found that the major polynyas identified in our study over the common time period are mostly identified in their reconstruction. A direct annual correlation is however statistically limited by the independent age-scale uncertainties, typically $\pm 1-2$ years, inherent in all ice core records. These chronological offsets cause high-magnitude events to appear out of phase between two indices/studies, which artificially degrades linear correlation coefficients despite clear physical synchronization in the raw signals. To provide a more robust and physically meaningful assessment, we used a binary presence-absence matrix assuming a ± 2 years lead/lag window at threshold levels of 0.6, 0.7, 0.8, 0.9 and 1. Therefore, we look for polynya events at different threshold levels in our record and then look for polynya events with a threshold of 0.6 from the polynya index from Goosse et al. (2021) and found that all major polynyas identified in their study over the common time period are also identified in this study. within a ± 2 years window. For simplicity, we focus on only one index from each study, simple average of the standardized time series (Stat) from Goosse et al. (2021) and polynya index including accumulation from this study. We find that nine polynya years (threshold = 0.6) in our index that are also corroborated by the Goosse et al. (2021) record (Figure 9a). Similarly, seven polynya years are corroborated at a threshold of 0.7 and three at 0.8. At the threshold of 0.9 and 1, only the polynya event of 1974 is corroborated, which is expected as both studies use this event as an assumed truth and normalize all event to it. Similarly, when we repeat the same process for polynyas identified in Goosse et al. (2021) (Figure 9b), we find that 17 polynya years are

corroborated at 0.6 threshold, 12 polynya years at 0.7, 9 polynya years at 0.8, eight at 0.9 and seven polynya years with a threshold of 1 are also identified in our polynya index.

This provides a degree of validation for both approaches, suggesting that they are capturing similar large-scale polynya events. However, our index shows enhanced sensitivity to polynya formations, identifying the polynya occurrence of 1964, even though with a ~~low~~ lower likelihood. This low likelihood of the 1964 polynya is possibly due to the lower persistence as compared to the well-documented events during 1974 – ~~1976~~ 76 (Meier et al., 2013). Furthermore, our index shows better performance in identifying transient polynyas and halos, which were missed in the reconstruction by Goosse et al. (2021). These shorter-lived or less intense polynya events, while not as dramatic as major openings, play a crucial role in regional oceanography and climate dynamics. Their detection provides a more comprehensive picture of polynya activity over time.

~~Visual inspection shows that some years show synchronised peaks across the indices from the two studies, with leads/lags associated with the timescale of the ice cores (Fig. 8). However, some peaks are present in our polynya index which are absent/muted in Goosse et al. (2021). This discrepancy between the two records is possibly due to the selection of ice core sites used for calculating the polynya index. Beyond chronological offsets, discrepancies between the two indices likely reflect differences in regional sensitivity and site selection.~~ Our analysis of air mass transport frequency from the MRP regions shows a sink in the coastal regions of Dronning Maud Land, with a significant proportion of the trajectories reaching our study area. In contrast, four out of the six ice core sites used in Goosse et al. (2021) fall outside the primary sink of these trajectories (Fig. 5). This difference in site selection could be a major reason for the observed discrepancy between the two polynya indices. Ice cores from sites that rarely receive air masses from the polynya region may not be reliably used to reconstruct polynya events or may do so with reduced sensitivity. Whereas our ice core site is well located to capture the occurrence of the MRP opening, as it frequently receives air masses originating from the polynya region. ~~However, they~~ However, Goosse et al. (2021) still manage to detect the large, multi-year polynyas as they lead to a widespread positive anomaly in precipitation over the coastal and continental region. Consequently, while our index captures both high-magnitude regional events (e.g., 1974) and more localized openings (e.g., 1964), inland records may only register the most extreme regional anomalies. This suggests that the lack of a perfect

year-to-year correlation is not a limitation of the proxies themselves, but rather a reflection of the better spatial representativeness of the IND-36/9 site for capturing the nuances of the Maud Rise polynya system. Another possible reason for this observed difference in sensitivity could be the robustness of our multiproxy approach in identifying polynyas, as it is less susceptible to noise or artefacts in any single proxy. It is crucial to note that the observed differences in peak occurrences do not necessarily invalidate either dataset but rather highlight the complexity in reconstructing past polynya activity. These discrepancies underscore the challenges inherent in paleoclimate reconstruction and the potential complementarity of different methodologies in building a comprehensive understanding of past polynya dynamics.

4.2. Historical polynya activity and its link to maritime climate

Our polynya reconstruction aligns with and extends satellite era observations of MRP formation mechanisms. Recent studies have shed light on the complex interplay of oceanic and atmospheric factors controlling MRP formation and persistence (Campbell et al., 2019; Jena et al., 2019). However, extending these insights to longer timescales remains challenging. Our multi-century reconstruction offers a unique opportunity to explore the long-term drivers of MRP variability. Autonomous profiling float observations during 2016 and 2017 revealed that the MRP were initiated and modulated by the passage of severe storms, and the intense heat loss drove deep overturning within them (Campbell et al., 2019). Wind-driven upwelling of record strength weakened haline stratification in the upper ocean, thus favouring destabilisation. A recent study, however, highlighted the role of salt transport through northward Ekman transport as an additional mechanism of the polynya formation and these processes were driven by intensified eastward surface stresses during 2015 – 2018 (Narayanan et al., 2024).

~~To investigate these factors during the past polynya openings, we used ocean reanalysis outputs from the Simple Ocean Data Assimilation (SODA) version 2.2.4, which spans the period from 1871 to 2010 (Carton and Giese, 2008; Giese and Ray, 2011). It is forced using data assimilation of in-situ temperature and salinity profiles and in-situ and satellite SSTs. This long-term dataset provides monthly fields of ocean temperature, salinity, currents, and sea surface height at a horizontal resolution of $\sim 0.5^\circ$, making it particularly suited for studying large-scale oceanic features and their variability over decadal time scales. Our analysis reveals that periods with high likelihood of polynya occurrences are consistently characterised by~~

increased eastward zonal wind stress due to westerly wind, enhanced northward water flow, and elevated sea surface salinity (Fig. 9). This consistency lends credence to current theories of MRP formation mechanisms and suggests that similar processes operate on multi-decadal to centennial timescales. However, we also observed multiple periods of increased meridional velocity, zonal wind stress, and sea surface salinity that did not invariably lead to a high likelihood of polynya occurrence. This non-linearity underscores the complexity of polynya formation processes and suggests the existence of critical thresholds in wind stress, ocean circulation, or stratification that must be exceeded for polynya formation to occur. The temporal sequencing and duration of favourable conditions also play crucial roles in polynya initiation and maintenance. It could also mean that there are still one or more unexplored processes that control the opening of major polynyas in the Weddell Sea. Our multiproxy approach, combining snow accumulation, water isotopes, and major ion concentrations, provides a more robust reconstruction of polynya activity compared to single proxy studies. This approach allows us to capture diverse signatures of polynya events and mitigate the limitations of individual proxies.

Although beyond the scope of this study, Therefore, it is crucial to understand the maritime climate variability during the polynya and non-polynya years. Our reconstruction shows alternation between polynya-active and quiescent periods, which may provide valuable context for assessing how future MRP behaviour may evolve under anthropogenic climate change. The temporal clustering of MRPs in our record possibly indicates that once favourable preconditioning develops, the region remains susceptible to repeated polynya openings over several years or decades. As global temperatures rise, projected changes in Southern Ocean wind patterns, sea-ice extent, and surface freshening may shift the thresholds required for deep convection, thereby altering both the frequency and persistence of MRPs. Understanding how atmospheric and oceanic conditions differ between the polynya and non-polynya periods identified in this study will be critical for refining projections. Integrating these insights with high-resolution coupled ocean-atmosphere models could help identify the physical thresholds and feedbacksfeedback governing MRP formation. Such integration is crucial for predicting how rare but climatically significant features such as the Maud Rise Polynya may influence Southern Ocean overturning, carbon exchange, and global climate in a warming world.

5. Conclusions and Outlook

We provide new insights into the temporal variability of Maud Rise Polynya, one of the largest and well-known open ocean ~~polynyas~~polynyas, by integrating multiple proxies using an ice core from coastal Dronning Maud Land, representing the past nearly 250 years. We identified multiple polynya openings during the 1830s, 1880s, and 1970s. These clusters of polynya openings possibly correspond to a specific combination of atmospheric circulation patterns and oceanographic preconditioning, suggesting a more deterministic framework for MRP development. Over the 250-year record, we identify large events during 1833 – 34 and 1911, which were comparable in magnitude to the documented 1974–~~1976~~–76 polynya, indicating that such openings are not isolated anomalies but represent recurring features of the Southern Ocean system. These events are characterised by proxy signatures consistent with modern polynya conditions, supporting the persistence of underlying formation mechanisms throughout the observational period. Conversely, the interval from 1920 to 1950, marked by a notable absence of polynya activity, coincides with documented changes in Southern Ocean circulation and atmospheric forcing. This prolonged quiescent phase provides valuable baseline information for assessing natural variability in MRP activity on multi-decadal timescales.

Our multiproxy approach significantly enhances the sensitivity of the reconstruction, particularly in detecting smaller-magnitude and shorter-duration events that may have been overlooked in previous studies. As a result, the reconstructed history reveals a more dynamic and variable pattern of MRP activity than previously recognised, with implications for understanding its role in ocean ventilation, carbon cycling, and regional climate variability. Our study reveals that synthesising paleoclimate records, modern observations, and modelling approaches can provide a robust foundation for advancing the understanding of the MRP as a critical and recurrent component of the Southern Ocean climate system.

Acknowledgements

The ice core was drilled as part of the India–Norway MADICE project, with support from the Ministry of Earth Sciences, Government of India, and the Research Council of Norway. Ice core drilling and field expeditions have been made successful with the support of the NCPOR logistics leaders, the Maitri logistics team, and all field participants. K. Mahalinganathan and

Tariq Ejaz are acknowledged for their contributions to ice core processing and analysis. We acknowledge the use of imagery from the NASA Worldview application (<https://worldview.earthdata.nasa.gov>), part of the NASA Earth Science Data and Information System (ESDIS). [This is NCPOR contribution no. xxxx](#)

Financial support

This research has been supported by the Ministry of Earth Sciences, India (grant no. MoES/Indo-Nor/PS-3/2015) and the Research Council of Norway (grant no. 248780) for the joint India–Norway project, ‘Mass balance, dynamics, and climate of the central Dronning Maud Land coast, East Antarctica’ (MADICE).

Author contributions

RD and MT defined the study objectives. RD led the ice core processing and analysis with support from AP, BLR and CML. RD led the data analysis and interpretation with inputs from KM and CML. RD prepared the manuscript with feedback from all co-authors. MT and KM were the project leaders

Data Availability

All ice core data and polynya indices are being prepared for submission to the NCPOR Polar Data Centre (<https://data.ncpor.res.in>). For final archival, open data formats will be used. A DOI will be generated and included in the final version of the paper. Data access can be granted to reviewers promptly upon request.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.





Figure 1 | Study area and wind-trajectory patterns. (a) Maud Rise Polynya during its largest recent opening on 25 September 2017, detected with Moderate Resolution Imaging Spectroradiometer (MODIS) on NASA's [TerraAqua](https://worldview.earthdata.nasa.gov/) satellite (from NASA Worldview, <https://worldview.earthdata.nasa.gov/>, last access: ~~15 September 2024~~. 01 April 2026). The polynya is visible as the dark, open-water feature amidst the surrounding white sea ice (b) MODIS Satellite imagery taken on the same day in a non-polynya year, 25 September 2022.

611 The MODIS corrected reflectance imagery provides a true colour representation where the
612 open ocean appears dark gray or black, while sea ice and clouds appear white or light gray.
613 Images are overlaid with the ~~endpoint~~trajectory frequency of back-trajectories ending at the
614 Djupranen ice core site (green star) from June to October ~~of the years~~in 2017 and 2022,
615 respectively (see methods). The Indian Maitri Station is marked by a yellow square. The
616 colourmap is on a logarithmic scale. The inset shows the location of the main map. Grounding
617 line and calving front are marked ~~(Matsuoka et al., 2015).~~(Matsuoka et al., 2015).






Figure 2 | Tie points for ice core chronology. Volcanic events are identified from the non-sea-salt sulphate flux records. Only major volcanic events used as age tie points for reconstructing the chronology are marked.



624

625 **Figure 3 | Age-depth scale for the Djupranen ice core.** Tie points used for chronology (Fig.
626 2) are shown with dashed lines. Major time markers are shown using the black dashed lines.
627 Minimal deviation from the intersection points of the vertical and horizontal lines indicates the
628 robustness of the chronology.






Figure 4 | Satellite polynya metrics. Two different annual metrics of satellite-derived polynya

632 occurrence. Since the polynya is highly dynamic temporally and spatially, the metrics do not
633 show a one-to-one resemblance. The red vertical bar shows the 1974 – 76 polynya event, which
634 reached a peak extent of approximately 300,000 km² (Carsey 1980), more than ten times of the
635 maximum cumulative polynya area observed between 1979 – 2016.



637

638 **Figure 5 | Airmass transport pathways originating from MRP: EndpointTrajectory**
 639 frequency of forward trajectories originating over the polynya-prone region (Heuzé et al.,
 640 2021) of the Weddell Sea, showing the prevalent transport pathway during the winter to early
 641 spring months for the period 1948 – 2016. A significant proportion of the trajectories end over
 642 our ice core site (red diamond). ~~Location of ice cores used by Goosse et al., (2021) are shown~~
 643 ~~with green circles. The gray region shows area with no trajectory endpoints~~Location of ice
 644 cores used by Goosse et al. (2021) are shown with green circles. The black trapezium shows
 645 the polynya prone regions. The gray region shows area with no trajectories.






Figure 6 | Airmass transport pathways ending at the ice core site: ~~Endpoint~~Trajectory frequency of back trajectories originating from the ice core site during a well-known polynya year (2017; left) and a non-polynya year (2022; right) is shown. The ~~endpoint~~trajectory frequency pattern during both years is very similar, indicating a consistent transport pathway during most of the years and suggesting that our ice core records activity over the polynya-prone region of the Weddell Sea (black trapezium) in all years. The gray region shows areas with no ~~trajectory~~endpoint~~trajectories~~. The dashed lines show the NSIDC median sea ice extent for 1980 – 2010, with green indicating September (maximum) and red representing February (minimum) sea ice extent.








Figure 7 | (a) Ice core proxies and polynya identification. Annual variability of selected ice core proxy variability from 1960 – 2016. Known events from the satellite data are marked with dark gray patches for polynya years and light gray patches for halo years (Heuzé et al., 2021). proxies and polynya index from 1960 – 2016. The red curve is the moving median over a 30-year window. The two known polynya events of 1964 and 1974 – 76 are marked with gray patches. **(b) Ice core derived polynya indices:** Ice core derived polynya indices in the past 250 years using the anomalies in the annual record of five ice-core proxies: Na, deuterium

666 excess, $\delta^{18}\text{O}$, and snow accumulation ~~and Na/SO₄ ratio. In order to. To~~ test the dependence of
667 the polynya index on snow accumulation, we calculate two indices, one using snow
668 accumulation (blue curve) and another without (red curve). The two indices behave similarly
669 over the entire time period; however, the index without snow accumulation shows a higher
670 range of variability. ~~Three different thresholds, 0.6, 0.8, and 1, are set for detecting polynyas.~~



671




Figure 8 | Comparison of the polynya indices. Polynya indices reconstructed in this study (upper panel) are shown with the polynya indices from Goosse et al. (2021) (lower panel). The polynya indices from Goosse et al. (2021) are based on six surface mass balance records using data assimilation with two control simulations performed with the SPEAR (Seamless system for Prediction and EArth system Research) global climate model, SPEAR_AM2 (DAAM2; Blue) and SPEAR_LO (DALO; green), and a simple average of the standardized time series (Stat, red).

The known polynya events of 1964 (1965 in our record and absent in Goosse et al. (2021)) and 1974 – 76 (1974 in our record and 1975 in Goosse et al. (2021)) are labelled.




Figure 9 | Factors influencing polynya formations (a) Mean meridional velocity, (b) zonal wind stress, and (c) sea surface salinity record over the polynya-prone region of the Weddell Sea. (d) Polynya index reconstructed in this study. Grey bars are the major polynya events prior to 1976.




Figure 9 | Comparison of synchronous polynya events. Binary presence-absence analysis comparing the reconstructed polynya index from this study and the index from Goose et al. (2021) across five threshold levels (0.6, 0.7, 0.8, 0.9, and 1.0) and assuming a lead/lag of ± 2 years. (a) Temporal distribution of polynya events identified at different threshold in this study and corroborated by the Goose et al. (2021) record (threshold ≥ 0.6). (b) Distribution of polynya events from Goose et al. (2021) at different threshold corroborated by this study (threshold ≥ 0.6).

696 References

- 697 Akitomo, K.: Thermobaric deep convection, baroclinic instability, and their roles in vertical heat
698 transport around Maud Rise in the Weddell Sea, *Journal of Geophysical Research: Oceans*, 111,
699 2006.
- 700 Arrigo, K. R. and van Dijken, G. L.: Phytoplankton dynamics within 37 Antarctic coastal polynya
701 systems, *Journal of Geophysical Research: Oceans*, 108, 2003.
- 702 Årthun, M., Holland, P. R., Nicholls, K. W., and Feltham, D. L.: Eddy Driven Exchange between the
703 Open Ocean and a Sub-Ice Shelf Cavity, *Journal of Physical Oceanography*, 43, 2372–2387,
704 2013.
- 705 Bennetts, L. G., Shakespeare, C. J., Vreugdenhil, C. A., Foppert, A., Gayen, B., Meyer, A., Morrison,
706 A. K., Padman, L., Phillips, H. E., Stevens, C. L., Toffoli, A., Constantinou, N. C., Cusack, J.
707 M., Cyriac, A., Doddridge, E. W., England, M. H., Evans, D. G., Heil, P., Hogg, A. M., Holmes,
708 R. M., Huneke, W. G. C., Jones, N. L., Keating, S. R., Kiss, A. E., Kraitzman, N., Malyarenko,
709 A., McConnochie, C. D., Meucci, A., Montiel, F., Neme, J., Nikurashin, M., Patel, R. S., Peng,
710 J. P., Rayson, M., Rosevear, M. G., Sohail, T., Spence, P., and Stanley, G. J.: Closing the Loops
711 on Southern Ocean Dynamics: From the Circumpolar Current to Ice Shelves and From Bottom
712 Mixing to Surface Waves, *Reviews of Geophysics*, 62, e2022RG000781, 2024.
- 713 Berman, E. S. F., Levin, N. E., Landais, A., Li, S., and Owano, T.: Measurement of $\delta^{18}\text{O}$, $\delta^{17}\text{O}$, and
714 ^{47}O excess in Water by Off-Axis Integrated Cavity Output Spectroscopy and Isotope Ratio Mass
715 Spectrometry, *Analytical Chemistry*, 85, 10392–10398, 2013.
- 716 Campbell, E. C., Wilson, E. A., Moore, G. W. K., Riser, S. C., Brayton, C. E., Mazloff, M. R., and
717 Talley, L. D.: Antarctic offshore polynyas linked to Southern Hemisphere climate anomalies,
718 *Nature*, 570, 319–325, 2019.
- 719 Carsey, F. D.: Microwave Observation of the Weddell Polynya, *Monthly Weather Review*, 108, 2032–
720 2044, 1980.
- 721 Carton, J. A. and Giese, B. S.: A Reanalysis of Ocean Climate Using Simple Ocean Data Assimilation
722 (SODA), *Monthly Weather Review*, 136, 2999–3017, 2008.
- 723 Comiso, J. C. and Gordon, A. L.: Recurring polynyas over the Cosmonaut Sea and the Maud Rise,
724 *Journal of Geophysical Research: Oceans*, 92, 2819–2833, 1987.
- 725 Comiso, J. C. and Nishio, F.: Trends in the sea ice cover using enhanced and compatible AMSR-E,
726 SSM/I, and SMMR data, *Journal of Geophysical Research: Oceans*, 113, 2008.
- 727 de Lavergne, C., Palter, J. B., Galbraith, E. D., Bernardello, R., and Marinov, I.: Cessation of deep
728 convection in the open Southern Ocean under anthropogenic climate change, *Nature Climate*
729 *Change*, 4, 278–282, 2014.
- 730 Dey, R.: Reconstruction of Antarctic climate variability using high-resolution ice core stratigraphy,
731 2023. Goa University, 2023.
- 732 Dey, R., Thamban, M., Laluraj, C. M., Mahalinganathan, K., Redkar, B. L., Kumar, S., and Matsuoka,
733 K.: Application of visual stratigraphy from line-scan images to constrain chronology and melt
734 features of a firn core from coastal Antarctica, *Journal of Glaciology*, 69, 179–190, 2023.

- Dixon, D. A., Mayewski, P. A., Goodwin, I. D., Marshall, G. J., Freeman, R., Maasch, K. A., and Sneed, S. B.: An ice core proxy for northerly air mass incursions into West Antarctica, *International Journal of Climatology*, 32, 1455–1465, 2012.
- Dorling, S. R. and Davies, T. D.: Extending cluster analysis—synoptic meteorology links to characterise chemical climates at six northwest European monitoring stations, *Atmospheric Environment*, 29, 145–167, 1995.
- Draxler, R.: NOAA Air resources laboratory FAQ How do i estimate the absolute (in km) and relative (%) errors when using the HYSPLIT trajectory model. Sept, 2008.
- Drews, R., Matsuoka, K., Martín, C., Callens, D., Bergeot, N., and Pattyn, F.: Evolution of Derwael Ice Rise in Dronning Maud Land, Antarctica, over the last millennia, *Journal of Geophysical Research: Earth Surface*, 120, 564–579, 2015.
- Ejaz, T., Rahaman, W., Laluraj, C. M., Mahalinganathan, K., and Thamban, M.: Rapid Warming Over East Antarctica Since the 1940s Caused by Increasing Influence of El Niño Southern Oscillation and Southern Annular Mode, *Frontiers in Earth Science*, Volume 10—2022, 2022.
- Ejaz, T., Rahaman, W., Laluraj, C. M., Mahalinganathan, K., and Thamban, M.: Sea Ice Variability and Trends in the Western Indian Ocean Sector of Antarctica During the Past Two Centuries and Its Response to Climatic Modes, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033943, 2021.
- Giese, B. S. and Ray, S.: El Niño variability in simple ocean data assimilation (SODA), 1871–2008, *Journal of Geophysical Research: Oceans*, 116, 2011.
- Goel, V., Brown, J., and Matsuoka, K.: Glaciological settings and recent mass balance of Blåskimen Island in Dronning Maud Land, Antarctica, *The Cryosphere*, 11, 2883–2896, 2017.
- Goel, V., Matsuoka, K., Berger, C. D., Lee, I., Dall, J., and Forsberg, R.: Characteristics of ice rises and ice rumples in Dronning Maud Land and Enderby Land, Antarctica, *Journal of Glaciology*, 66, 1064–1078, 2020.
- Goosse, H., Dalaiden, Q., Cavitte, M. G. P., and Zhang, L.: Can we reconstruct the formation of large open-ocean polynyas in the Southern Ocean using ice core records?, *Clim. Past*, 17, 111–131, 2021.
- Gordon, A. L. and Comiso, J. C.: Polynyas in the Southern Ocean, *Scientific American*, 258, 90–97, 1988.
- Goursaud, S., Masson-Delmotte, V., Favier, V., Preunkert, S., Fily, M., Gallée, H., Jourdain, B., Legrand, M., Magand, O., Minster, B., and Werner, M.: A 60 year ice core record of regional climate from Adélie Land, coastal Antarctica, *The Cryosphere*, 11, 343–362, 2017.
- Gülk, B., Roquet, F., Naveira-Garabato, A. C., Bourdallé-Badie, R., Madec, G., and Giordani, H.: Impacts of Vertical Convective Mixing Schemes and Freshwater Forcing on the 2016–2017 Maud Rise Polynya Openings in a Regional Ocean Simulation, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS004106, 2024.
- Heuzé, C., Heywood, K. J., Stevens, D. P., and Ridley, J. K.: Southern Ocean bottom water characteristics in CMIP5 models, *Geophysical Research Letters*, 40, 1409–1414, 2013.
- Heuzé, C., Zhou, L., Mohrmann, M., and Lemos, A.: Spaceborne infrared imagery for early detection

- of Weddell Polynya opening, *The Cryosphere*, 15, 3401–3421, 2021.
- Holland, D. M.: Explaining the Weddell Polynya—a Large Ocean Eddy Shed at Maud Rise, *Science*, 292, 1697–1700, 2001.
- Jacobs, S. S., Gordon, A. L., and Ardai, J. L.: Circulation and Melting Beneath the Ross Ice Shelf, *Science*, 203, 439–443, 1979.
- Jena, B., Ravichandran, M., and Turner, J.: Recent Reoccurrence of Large Open Ocean Polynya on the Maud Rise Seamount, *Geophysical Research Letters*, 46, 4320–4329, 2019.
- Jones, T. R., Cuffey, K. M., Roberts, W. H. G., Markle, B. R., Steig, E. J., Stevens, C. M., Valdes, P. J., Fudge, T. J., Sigl, M., Hughes, A. G., Morris, V., Vaughn, B. H., Garland, J., Vinther, B. M., Rozmiarek, K. S., Brashear, C. A., and White, J. W. C.: Seasonal temperatures in West Antarctica during the Holocene, *Nature*, 613, 292–297, 2023.
- Klein, F., Abram, N. J., Curran, M. A. J., Goosse, H., Goursaud, S., Masson-Delmotte, V., Moy, A., Neukom, R., Orsi, A., Sjolte, J., Steiger, N., Stenni, B., and Werner, M.: Assessing the robustness of Antarctic temperature reconstructions over the past 2 millennia using pseudoproxy and data assimilation experiments, *Clim. Past*, 15, 661–684, 2019.
- Lenaerts, J. T. M., Brown, J., Van Den Broeke, M. R., Matsuoka, K., Drews, R., Callens, D., Philippe, M., Gorodetskaya, I. V., Van Meijgaard, E., Reijmer, C. H., Pattyn, F., and Van Lipzig, N. P. M.: High variability of climate and surface mass balance induced by Antarctic ice rises, *Journal of Glaciology*, 60, 1101–1110, 2014.
- Levine, J. G., Yang, X., Jones, A. E., and Wolff, E. W.: Sea salt as an ice core proxy for past sea ice extent: A process-based model study, *Journal of Geophysical Research: Atmospheres*, 119, 5737–5756, 2014.
- Lindsay, R. W., Holland, D. M., and Woodgate, R. A.: Halo of low ice concentration observed over the Maud Rise seamount, *Geophysical Research Letters*, 31, 2004.
- Markle, B. R., Bertler, N. A. N., Sinclair, K. E., and Sneed, S. B.: Synoptic variability in the Ross Sea region, Antarctica, as seen from back-trajectory modeling and ice core analysis, *Journal of Geophysical Research: Atmospheres*, 117, 2012.
- Matsuoka, K., Hindmarsh, R. C. A., Moholdt, G., Bentley, M. J., Pritchard, H. D., Brown, J., Conway, H., Drews, R., Durand, G., Goldberg, D., Hattermann, T., Kingslake, J., Lenaerts, J. T. M., Martin, C., Mulvaney, R., Nicholls, K. W., Pattyn, F., Ross, N., Scambos, T., and Whitehouse, P. L.: Antarctic ice rises and rumples: Their properties and significance for ice sheet dynamics and evolution, *Earth Science Reviews*, 150, 724–745, 2015.
- McHedlishvili, A., Spreen, G., Melsheimer, C., and Huntemann, M.: Weddell Sea polynya analysis using SMOS–SMAP apparent sea ice thickness retrieval, *The Cryosphere*, 16, 471–487, 2022.
- McPhee, M. G.: Is thermobaricity a major factor in Southern Ocean ventilation?, *Antarctic Science*, 15, 153–160, 2003.
- Meier, W. N., Gallaher, D., and Campbell, G. G.: New estimates of Arctic and Antarctic sea ice extent during September 1964 from recovered Nimbus I satellite imagery, *The Cryosphere*, 7, 699–705, 2013.
- Moore, G. W. K., Alverson, K., and Renfrew, I. A.: A Reconstruction of the Air–Sea Interaction

- Associated with the Weddell Polynya, *Journal of Physical Oceanography*, 32, 1685–1698, 2002.
- Naik, S. S., Thamban, M., Laluraj, C. M., Redkar, B. L., and Chaturvedi, A.: A century of climate variability in central Dronning Maud Land, East Antarctica, and its relation to Southern Annular Mode and El Niño–Southern Oscillation, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- Narayanan, A., Roquet, F., Gille, S. T., Güllk, B., Mazloff, M. R., Silvano, A., and Naveira Garabato, A. C.: Ekman driven salt transport as a key mechanism for open ocean polynya formation at Maud Rise, *Science Advances*, 10, eadj0777, 2024.
- Neff, P. D. and Bertler, N. A. N.: Trajectory modeling of modern dust transport to the Southern Ocean and Antarctica, *Journal of Geophysical Research: Atmospheres*, 120, 9303–9322, 2015.
- Philippe, M., Tison, J. L., Fjøsne, K., Hubbard, B., Kjær, H. A., Lenaerts, J. T. M., Drews, R., Sheldon, S. G., De Bondt, K., Claeys, P., and Pattyn, F.: Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica, *The Cryosphere*, 10, 2501–2516, 2016.
- Pratap, B., Dey, R., Matsuoka, K., Moholdt, G., Lindbäck, K., Goel, V., Laluraj, C. M., and Thamban, M.: Three decade spatial patterns in surface mass balance of the Nivlisen Ice Shelf, central Dronning Maud Land, East Antarctica, *Journal of Glaciology*, 68, 174–186, 2022.
- Richardson, C.: Phase Relationships in Sea Ice as a Function of Temperature, *Journal of Glaciology*, 17, 507–519, 1976.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, 2019.
- Sallée, J. B., Shuckburgh, E., Bruneau, N., Meijers, A. J. S., Bracegirdle, T. J., Wang, Z., and Roy, T.: Assessment of Southern Ocean water mass circulation and characteristics in CMIP5 models: Historical bias and forcing response, *Journal of Geophysical Research: Oceans*, 118, 1830–1844, 2013.
- The Firn Symposium team: Firn on ice sheets, *Nature Reviews Earth & Environment*, 5, 79–99, 2024.
- The SO-CHIC consortium: Southern ocean carbon and heat impact on climate, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 381, 20220056, 2023.
- Visbeck, M., Marshall, J., and Jones, H.: Dynamics of Isolated Convective Regions in the Ocean, *Journal of Physical Oceanography*, 26, 1721–1734, 1996.
- Wagenbach, D., Dueroz, F., Mulvaney, R., Keek, L., Minikin, A., Legrand, M., Hall, J. S., and Wolff, E. W.: Sea salt aerosol in coastal Antarctic regions, *Journal of Geophysical Research: Atmospheres*, 103, 10961–10974, 1998.
- Wauthy, S., Tison, J. L., Inoue, M., El Amri, S., Sun, S., Fripiat, F., Claeys, P., and Pattyn, F.: Spatial and temporal variability of environmental proxies from the top 120 m of two ice cores in Dronning Maud Land (East Antarctica), *Earth Syst. Sci. Data*, 16, 35–58, 2024.
- Wilson, E. A., Riser, S. C., Campbell, E. C., and Wong, A. P. S.: Winter Upper Ocean Stability and Ice–Ocean Feedbacks in the Sea Ice Covered Southern Ocean, *Journal of Physical*

- Oceanography, 49, 1099–1117, 2019.
- Winstrup, M., Svensson, A. M., Rasmussen, S. O., Winther, O., Steig, E. J., and Axelrod, A. E.: An automated approach for annual layer counting in ice cores, *Clim. Past*, 8, 1881–1895, 2012.
- Wolff, E. W., Rankin, A. M., and Röthlisberger, R.: An ice core indicator of Antarctic sea ice production?, *Geophysical Research Letters*, 30, 2003.
- Xu, Y., Zhang, W., Maksym, T., Ji, R., and Li, Y.: Stratification Breakdown in Antarctic Coastal Polynyas. Part I: Influence of Physical Factors on the Destratification Time Scale, *Journal of Physical Oceanography*, 53, 2047–2067, 2023.
- Zheng, W. E. I., Zhaoru, Z., Timo, V., Xiaoqiao, W., and Yuanjie, C.: An overview of Antarctic polynyas: sea ice production, forcing mechanisms, temporal variability and water mass formation, *Advances in Polar Science*, 32, 295–311, 2021.
- Zhou, L., Heuzé, C., and Mohrmann, M.: Sea Ice Production in the 2016 and 2017 Maud Rise Polynyas, *Journal of Geophysical Research: Oceans*, 128, e2022JC019148, 2023.
- Zwally, H. J., Comiso, J. C., and Gordon, A. L.: Antarctic Offshore Leads and Polynyas and Oceanographic Effects. In: *Oceanology of the Antarctic Continental Shelf*, 1985.
- Akitomo, K.: Thermobaric deep convection, baroclinic instability, and their roles in vertical heat transport around Maud Rise in the Weddell Sea, *Journal of Geophysical Research: Oceans*, 111, 2006. <https://doi.org/10.1029/2005JC003284>
- Arrigo, K. R. and van Dijken, G. L.: Phytoplankton dynamics within 37 Antarctic coastal polynya systems, *Journal of Geophysical Research: Oceans*, 108, 2003. <https://doi.org/10.1029/2002JC001739>
- Årthun, M., Holland, P. R., Nicholls, K. W., and Feltham, D. L.: Eddy-Driven Exchange between the Open Ocean and a Sub-Ice Shelf Cavity, *Journal of Physical Oceanography*, 43, 2372–2387, 2013. <https://doi.org/10.1175/JPO-D-13-0137.1>
- Bennetts, L. G., Shakespeare, C. J., Vreugdenhil, C. A., Foppert, A., Gayen, B., Meyer, A., Morrison, A. K., Padman, L., Phillips, H. E., Stevens, C. L., Toffoli, A., Constantinou, N. C., Cusack, J. M., Cyriac, A., Doddridge, E. W., England, M. H., Evans, D. G., Heil, P., Hogg, A. M., Holmes, R. M., Huneke, W. G. C., Jones, N. L., Keating, S. R., Kiss, A. E., Kraitzman, N., Malyarenko, A., McConnochie, C. D., Meucci, A., Montiel, F., Neme, J., Nikurashin, M., Patel, R. S., Peng, J.-P., Rayson, M., Rosevear, M. G., Sohail, T., Spence, P., and Stanley, G. J.: Closing the Loops on Southern Ocean Dynamics: From the Circumpolar Current to Ice Shelves and From Bottom Mixing to Surface Waves, *Reviews of Geophysics*, 62, e2022RG000781, 2024. <https://doi.org/10.1029/2022RG000781>
- Berman, E. S. F., Levin, N. E., Landais, A., Li, S., and Owano, T.: Measurement of $\delta^{18}\text{O}$, $\delta^{17}\text{O}$, and ^{17}O -excess in Water by Off-Axis Integrated Cavity Output Spectroscopy and Isotope Ratio Mass Spectrometry, *Analytical Chemistry*, 85, 10392–10398, 2013. <https://doi.org/10.1021/ac402366t>

Campbell, E. C., Wilson, E. A., Moore, G. W. K., Riser, S. C., Brayton, C. E., Mazloff, M. R., and Talley, L. D.: Antarctic offshore polynyas linked to Southern Hemisphere climate anomalies, *Nature*, 570, 319-325, 2019. <https://doi.org/10.1038/s41586-019-1294-0>

Carsey, F. D.: Microwave Observation of the Weddell Polynya, *Monthly Weather Review*, 108, 2032-2044, 1980. [https://doi.org/10.1175/1520-0493\(1980\)108<2032:MOOTWP>2.0.CO;2](https://doi.org/10.1175/1520-0493(1980)108<2032:MOOTWP>2.0.CO;2)

Comiso, J. C. and Gordon, A. L.: Recurring polynyas over the Cosmonaut Sea and the Maud Rise, *Journal of Geophysical Research: Oceans*, 92, 2819-2833, 1987. <https://doi.org/10.1029/JC092iC03p02819>

de Lavergne, C., Palter, J. B., Galbraith, E. D., Bernardello, R., and Marinov, I.: Cessation of deep convection in the open Southern Ocean under anthropogenic climate change, *Nature Climate Change*, 4, 278-282, 2014. <https://doi.org/10.1038/nclimate2132>

Dey, R.: Reconstruction of Antarctic climate variability using high resolution ice core stratigraphy, 2023. Goa University, 2023. <https://doi.org/10.5281/zenodo.12705334>

Dey, R., Thamban, M., Laluraj, C. M., Mahalinganathan, K., Redkar, B. L., Kumar, S., and Matsuoka, K.: Application of visual stratigraphy from line-scan images to constrain chronology and melt features of a firn core from coastal Antarctica, *Journal of Glaciology*, 69, 179-190, 2023. <https://doi.org/10.1017/jog.2022.59>

Dixon, D. A., Mayewski, P. A., Goodwin, I. D., Marshall, G. J., Freeman, R., Maasch, K. A., and Sneed, S. B.: An ice-core proxy for northerly air mass incursions into West Antarctica, *International Journal of Climatology*, 32, 1455-1465, 2012. <https://doi.org/10.1002/joc.2371>

Dorling, S. R. and Davies, T. D.: Extending cluster analysis—synoptic meteorology links to characterise chemical climates at six northwest European monitoring stations, *Atmospheric Environment*, 29, 145-167, 1995. [https://doi.org/10.1016/1352-2310\(94\)00251-F](https://doi.org/10.1016/1352-2310(94)00251-F)

Draxler, R.: NOAA-Air resources laboratory-FAQ-How do i estimate the absolute (in km) and relative (%) errors when using the HYSPLIT trajectory model. Sept, 2008.

Drews, R., Matsuoka, K., Martín, C., Callens, D., Bergeot, N., and Pattyn, F.: Evolution of Derwael Ice Rise in Dronning Maud Land, Antarctica, over the last millennia, *Journal of Geophysical Research: Earth Surface*, 120, 564-579, 2015. <https://doi.org/10.1002/2014jf003246>

Ejaz, T., Rahaman, W., Laluraj, C. M., Mahalinganathan, K., and Thamban, M.: Rapid Warming Over East Antarctica Since the 1940s Caused by Increasing Influence of El Niño

- Southern Oscillation and Southern Annular Mode, *Frontiers in Earth Science*, Volume 10 - 2022, 2022. <https://doi.org/10.3389/feart.2022.799613>
- Ejaz, T., Rahaman, W., Laluraj, C. M., Mahalinganathan, K., and Thamban, M.: Sea Ice Variability and Trends in the Western Indian Ocean Sector of Antarctica During the Past Two Centuries and Its Response to Climatic Modes, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033943, 2021. <https://doi.org/10.1029/2020JD033943>
- Frey, M. M., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Yang, X., Jones, A. E., Nerentorp Mastromonaco, M. G., Jones, D. H., and Wolff, E. W.: First direct observation of sea salt aerosol production from blowing snow above sea ice, *Atmos. Chem. Phys.*, 20, 2549–2578, 2020. <https://doi.org/10.5194/acp-20-2549-2020>
- Goel, V., Brown, J., and Matsuoka, K.: Glaciological settings and recent mass balance of Blåskimen Island in Dronning Maud Land, Antarctica, *The Cryosphere*, 11, 2883-2896, 2017. <https://doi.org/10.5194/tc-11-2883-2017>
- Goel, V., Matsuoka, K., Berger, C. D., Lee, I., Dall, J., and Forsberg, R.: Characteristics of ice rises and ice rumpled in Dronning Maud Land and Enderby Land, Antarctica, *Journal of Glaciology*, 66, 1064-1078, 2020. <https://doi.org/10.1017/jog.2020.77>
- Goel, V., Martín, C., Matsuoka, K., Pratap, B., Moholdt, G., Dey, R., Laluraj, C. M., and Thamban, M.: A new coastal ice-core site identified in Dronning Maud Land, Antarctica, for high-resolution climate reconstructions to the Last Glacial Maximum, *The Cryosphere*, 20, 1363–1378, 2026. <https://doi.org/10.5194/tc-20-1363-2026>
- Goosse, H., Dalaiden, Q., Cavitte, M. G. P., and Zhang, L.: Can we reconstruct the formation of large open-ocean polynyas in the Southern Ocean using ice core records?, *Clim. Past*, 17, 111-131, 2021. <https://doi.org/10.5194/cp-17-111-2021>
- Gordon, A. L. and Comiso, J. C.: Polynyas in the Southern Ocean, *Scientific American*, 258, 90-97, 1988.
- Goursaud, S., Masson-Delmotte, V., Favier, V., Preunkert, S., Fily, M., Gallée, H., Jourdain, B., Legrand, M., Magand, O., Minster, B., and Werner, M.: A 60-year ice-core record of regional climate from Adélie Land, coastal Antarctica, *The Cryosphere*, 11, 343-362, 2017. <https://doi.org/10.5194/tc-11-343-2017>
- Gülk, B., Roquet, F., Naveira Garabato, A. C., Bourdallé-Badie, R., Madec, G., and Giordani, H.: Impacts of Vertical Convective Mixing Schemes and Freshwater Forcing on the 2016–2017 Maud Rise Polynya Openings in a Regional Ocean Simulation, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS004106, 2024. <https://doi.org/10.1029/2023MS004106>
- Heuzé, C., Heywood, K. J., Stevens, D. P., and Ridley, J. K.: Southern Ocean bottom water

- characteristics in CMIP5 models, *Geophysical Research Letters*, 40, 1409-1414, 2013.
<https://doi.org/10.1002/grl.50287>
- Heuzé, C., Zhou, L., Mohrmann, M., and Lemos, A.: Spaceborne infrared imagery for early detection of Weddell Polynya opening, *The Cryosphere*, 15, 3401-3421, 2021.
<https://doi.org/10.5194/tc-15-3401-2021>
- Holland, D. M.: Explaining the Weddell Polynya--a Large Ocean Eddy Shed at Maud Rise, *Science*, 292, 1697-1700, 2001. <https://doi.org/10.1126/science.1059322>
- Jacobs, S. S., Gordon, A. L., and Ardai, J. L.: Circulation and Melting Beneath the Ross Ice Shelf, *Science*, 203, 439-443, 1979. <https://doi.org/10.1126/science.203.4379.439>
- Jena, B., Ravichandran, M., and Turner, J.: Recent Reoccurrence of Large Open-Ocean Polynya on the Maud Rise Seamount, *Geophysical Research Letters*, 46, 4320-4329, 2019.
<https://doi.org/10.1029/2018GL081482>
- Jones, T. R., Cuffey, K. M., Roberts, W. H. G., Markle, B. R., Steig, E. J., Stevens, C. M., Valdes, P. J., Fudge, T. J., Sigl, M., Hughes, A. G., Morris, V., Vaughn, B. H., Garland, J., Vinther, B. M., Rozmiarek, K. S., Brashear, C. A., and White, J. W. C.: Seasonal temperatures in West Antarctica during the Holocene, *Nature*, 613, 292-297, 2023.
<https://doi.org/10.1038/s41586-022-05411-8>
- Klein, F., Abram, N. J., Curran, M. A. J., Goosse, H., Goursaud, S., Masson-Delmotte, V., Moy, A., Neukom, R., Orsi, A., Sjolte, J., Steiger, N., Stenni, B., and Werner, M.: Assessing the robustness of Antarctic temperature reconstructions over the past 2 millennia using pseudoproxy and data assimilation experiments, *Clim. Past*, 15, 661-684, 2019. <https://doi.org/10.5194/cp-15-661-2019>
- Lenaerts, J. T. M., Brown, J., Van Den Broeke, M. R., Matsuoka, K., Drews, R., Callens, D., Philippe, M., Gorodetskaya, I. V., Van Meijgaard, E., Reijmer, C. H., Pattyn, F., and Van Lipzig, N. P. M.: High variability of climate and surface mass balance induced by Antarctic ice rises, *Journal of Glaciology*, 60, 1101-1110, 2014.
<https://doi.org/10.3189/2014JoG14J040>
- Lindsay, R. W., Holland, D. M., and Woodgate, R. A.: Halo of low ice concentration observed over the Maud Rise seamount, *Geophysical Research Letters*, 31, 2004.
<https://doi.org/10.1029/2004GL019831>
- Markle, B. R., Bertler, N. A. N., Sinclair, K. E., and Sneed, S. B.: Synoptic variability in the Ross Sea region, Antarctica, as seen from back-trajectory modeling and ice core analysis, *Journal of Geophysical Research: Atmospheres*, 117, 2012.
<https://doi.org/10.1029/2011JD016437>
- Matsuoka, K., Hindmarsh, R. C. A., Moholdt, G., Bentley, M. J., Pritchard, H. D., Brown, J.,

- Conway, H., Drews, R., Durand, G., Goldberg, D., Hattermann, T., Kingslake, J., Lenaerts, J. T. M., Martín, C., Mulvaney, R., Nicholls, K. W., Pattyn, F., Ross, N., Scambos, T., and Whitehouse, P. L.: Antarctic ice rises and rumples: Their properties and significance for ice-sheet dynamics and evolution, *Earth-Science Reviews*, 150, 724-745, 2015. <https://doi.org/10.1016/j.earscirev.2015.09.004>
- McHedlishvili, A., Spreen, G., Melsheimer, C., and Huntemann, M.: Weddell Sea polynya analysis using SMOS–SMAP apparent sea ice thickness retrieval, *The Cryosphere*, 16, 471-487, 2022. <https://doi.org/10.5194/tc-16-471-2022>
- McPhee, M. G.: Is thermobaricity a major factor in Southern Ocean ventilation?, *Antarctic Science*, 15, 153-160, 2003. <https://doi.org/10.1017/S0954102003001159>
- Meier, W. N., Gallaher, D., and Campbell, G. G.: New estimates of Arctic and Antarctic sea ice extent during September 1964 from recovered Nimbus I satellite imagery, *The Cryosphere*, 7, 699-705, 2013. <https://doi.org/10.5194/tc-7-699-2013>
- Moore, G. W. K., Alverson, K., and Renfrew, I. A.: A Reconstruction of the Air–Sea Interaction Associated with the Weddell Polynya, *Journal of Physical Oceanography*, 32, 1685-1698, 2002. [https://doi.org/10.1175/1520-0485\(2002\)032<1685:AROTAS>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<1685:AROTAS>2.0.CO;2)
- Naik, S. S., Thamban, M., Laluraj, C. M., Redkar, B. L., and Chaturvedi, A.: A century of climate variability in central Dronning Maud Land, East Antarctica, and its relation to Southern Annular Mode and El Niño–Southern Oscillation, *Journal of Geophysical Research: Atmospheres*, 115, 2010. <https://doi.org/10.1029/2009JD013268>
- Narayanan, A., Roquet, F., Gille, S. T., Gülk, B., Mazloff, M. R., Silvano, A., and Naveira Garabato, A. C.: Ekman-driven salt transport as a key mechanism for open-ocean polynya formation at Maud Rise, *Science Advances*, 10, eadj0777, 2024. <https://doi.org/10.1126/sciadv.adj0777>
- Neff, P. D. and Bertler, N. A. N.: Trajectory modeling of modern dust transport to the Southern Ocean and Antarctica, *Journal of Geophysical Research: Atmospheres*, 120, 9303-9322, 2015. <https://doi.org/10.1002/2015jd023304>
- Philippe, M., Tison, J. L., Fjøsne, K., Hubbard, B., Kjær, H. A., Lenaerts, J. T. M., Drews, R., Sheldon, S. G., De Bondt, K., Claeys, P., and Pattyn, F.: Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica, *The Cryosphere*, 10, 2501-2516, 2016. <https://doi.org/10.5194/tc-10-2501-2016>
- Pratap, B., Dey, R., Matsuoka, K., Moholdt, G., Lindbäck, K., Goel, V., Laluraj, C. M., and Thamban, M.: Three-decade spatial patterns in surface mass balance of the Nivlisen Ice Shelf, central Dronning Maud Land, East Antarctica, *Journal of Glaciology*, 68, 174-186,

2022. <https://doi.org/10.1017/jog.2021.93>
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, 2019. <https://doi.org/10.1073/pnas.1812883116>
- Sallée, J.-B., Shuckburgh, E., Bruneau, N., Meijers, A. J. S., Bracegirdle, T. J., Wang, Z., and Roy, T.: Assessment of Southern Ocean water mass circulation and characteristics in CMIP5 models: Historical bias and forcing response, *Journal of Geophysical Research: Oceans*, 118, 1830–1844, 2013. <https://doi.org/10.1002/jgrc.20135>
- The Firn Symposium team: Firn on ice sheets, *Nature Reviews Earth & Environment*, 5, 79–99, 2024. <https://doi.org/10.1038/s43017-023-00507-9>
- The SO-CHIC consortium: Southern ocean carbon and heat impact on climate, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 381, 20220056, 2023. <https://doi.org/10.1098/rsta.2022.0056>
- Visbeck, M., Marshall, J., and Jones, H.: Dynamics of Isolated Convective Regions in the Ocean, *Journal of Physical Oceanography*, 26, 1721–1734, 1996. [https://doi.org/10.1175/1520-0485\(1996\)026<1721:DOICRI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1996)026<1721:DOICRI>2.0.CO;2)
- Wauthy, S., Tison, J. L., Inoue, M., El Amri, S., Sun, S., Fripiat, F., Claeys, P., and Pattyn, F.: Spatial and temporal variability of environmental proxies from the top 120 m of two ice cores in Dronning Maud Land (East Antarctica), *Earth Syst. Sci. Data*, 16, 35–58, 2024. <https://doi.org/10.5194/essd-16-35-2024>
- Wilson, E. A., Riser, S. C., Campbell, E. C., and Wong, A. P. S.: Winter Upper-Ocean Stability and Ice–Ocean Feedbacks in the Sea Ice–Covered Southern Ocean, *Journal of Physical Oceanography*, 49, 1099–1117, 2019. <https://doi.org/10.1175/JPO-D-18-0184.1>
- Winstrup, M., Svensson, A. M., Rasmussen, S. O., Winther, O., Steig, E. J., and Axelrod, A. E.: An automated approach for annual layer counting in ice cores, *Clim. Past*, 8, 1881–1895, 2012. <https://doi.org/10.5194/cp-8-1881-2012>
- Xu, Y., Zhang, W., Maksym, T., Ji, R., and Li, Y.: Stratification Breakdown in Antarctic Coastal Polynyas. Part I: Influence of Physical Factors on the Destratification Time Scale, *Journal of Physical Oceanography*, 53, 2047–2067, 2023. <https://doi.org/10.1175/JPO-D-22-0218.1>
- Zheng, W. E. I., Zhaoru, Z., Timo, V., Xiaoqiao, W., and Yuanjie, C.: An overview of Antarctic polynyas: sea ice production, forcing mechanisms, temporal variability and water mass formation, *Advances in Polar Science*, 32, 295–311, 2021. <https://doi.org/10.13679/j.advps.2021.0026>

1066 Zwally, H. J., Comiso, J. C., and Gordon, A. L.: Antarctic Offshore Leads and Polynyas and
1067 Oceanographic Effects. In: Oceanology of the Antarctic Continental Shelf, 1985.
1068 <https://doi.org/10.1029/AR043p0203>