

Evolution of Maud Rise Polynya during the last 250 years – a multiproxy ice core reconstruction from coastal Dronning Maud Land, Antarctica

Rahul Dey^{1,2}, Chavarukonam M. Laluraj¹, Kenichi Matsuoka³, Ashish Paiguinkar¹, Bhikaji L Redkar¹ and Meloth Thamban¹

¹National Centre for Polar and Ocean Research (NCPOR), Ministry of Earth Sciences, Vasco da Gama, Goa 403804, India

²Physics of Ice, Climate and Earth, Niels Bohr Institute, University of Copenhagen, Copenhagen N 2200, Denmark

³Norwegian Polar Institute, Framsentret, Postboks 6606, Langnes, 9296 Tromsø, Norway

Correspondence to: Rahul Dey (rdey1801@gmail.com)

Abstract

Open ocean polynyas drive deep ocean convection, influencing regional carbon and heat budgets, which in turn influence the ocean circulation and overall climate of Antarctica. The Maud Rise Polynya (MRP), also known as the Weddell Polynya, is one such polynya that forms in the Southern Ocean during early spring or winter months. The extensively studied MRP opening, which occurred during 2016 – 2017 and 1974 – 76, triggered intense convection, ventilating heat from the deep ocean and modifying water mass properties. However, polynya evolution before the satellite era remains poorly understood. Here, we develop a polynya index using multiple proxy records from an ice core in coastal Dronning Maud Land, East Antarctica. Our approach—integrating records of snow accumulation, $\delta^{18}\text{O}$, deuterium excess, ssNa flux, and Na/SO₄ ratios—enhances polynya reconstruction, thereby overcoming the limitations of single-proxy methods. The index replicates the 1974 – 76 polynya and extends the record to 1774, revealing three major events comparable to the 1974 – 76 great polynya event, in the past 250 years, totalling twenty five polynya years (likelihood > 0.6). We identified distinct clusters of polynya activity, possibly corresponding to a specific combination of atmospheric circulation patterns and oceanographic preconditioning for MRP development. This study offers a long-term perspective on MRP variability, providing insights into its drivers and climate-related impacts.

1. Introduction

Open ocean polynyas represent one of the most dynamic and influential features in the Southern Ocean system, serving as critical regulators of heat and gas exchange between the ocean and atmosphere. These persistent areas of open water within the sea ice cover act as windows through which complex oceanographic and atmospheric processes interact, influencing global ocean circulation, carbon cycling, and climate patterns (Bennetts et al., 2024; The SO-CHIC consortium, 2023; Zheng et al., 2021). The significance of polynyas extends beyond their immediate vicinity, influencing deep-water formation, marine ecosystem dynamics, and the intensification of sea-to-air heat and moisture fluxes during the winter months. In the Southern Ocean, polynyas manifest in two distinct forms: (1) coastal polynyas, which form along the Antarctic margin through mechanical forcing by katabatic winds, and (2) open-ocean polynyas, which develop hundreds of kilometres offshore through complex thermodynamic processes. While coastal polynyas occur regularly and have been extensively studied (Arrigo and van Dijken, 2003; Årthun et al., 2013; Jacobs et al., 1979; Visbeck et al., 1996; Xu et al., 2023), open-ocean polynyas represent rare but profoundly influential events that can significantly impact global ocean circulation patterns. The Maud Rise Polynya (MRP), also known as Weddell Polynya, occurring over the Maud Rise seamount in the eastern Weddell Sea (Fig. 1), is the most enigmatic open-ocean polynya in the Southern Ocean (Holland, 2001). First documented through satellite observations in the 1970s, the MRP reached its largest extent during the winter and early spring months from 1974 to 1976, exceeding 300,000 km² at its peak (Carsey, 1980; Zwally et al., 1985). This event drove intense ocean convection, reaching depths of over 3000 meters, and fundamentally altered water mass properties throughout the region, influencing global ocean circulation patterns (Gordon and Comiso, 1988).

Following this dramatic episode, the Maud Rise region has exhibited intermittent polynya activity, typically manifesting as halos of reduced sea ice concentration (Lindsay et al., 2004; McHedlishvili et al., 2022). However, the phenomenon garnered renewed attention when a large polynya exceeding 40,000 km² reappeared during the two winters of 2016 and 2017 (Jena et al., 2019). Early polynya studies focused primarily on documenting polynya occurrence and extent through satellite imagery (Carsey, 1980; Comiso and Gordon, 1987; Zwally et al., 1985), while more recent works have attempted to unravel the complex mechanisms controlling polynya formation and maintenance (de Lavergne et al., 2014; Wilson

et al., 2019). Satellite observations have revealed that open-ocean polynya formation often follows a characteristic pattern, beginning with the appearance of small-scale openings that can rapidly expand under favourable conditions, such as wind-driven upwelling (de Lavergne et al., 2014; Jena et al., 2019), baroclinic instabilities (Akitomo, 2006; Campbell et al., 2019), and deep warm water intrusions (Gülk et al., 2024; Heuzé et al., 2021). However, the precise triggers initiating this process remain incompletely understood, because the relative importance of these processes likely varies temporally and may be interconnected (Campbell et al., 2019; McPhee, 2003; Narayanan et al., 2024), and also because large polynyas openings are rare and satellite-based observations range only for the past few decades, limiting the observational opportunity. Similarly, model-based assessments are limited in their ability to understand polynya formation in the weakly stratified Southern Ocean (Heuzé et al., 2013; Sallée et al., 2013). This modelling challenge hampers our ability to assess both the role of polynyas in natural climate variability and their potential responses to future climate change.

To obtain longer polynya records, Goosse et al. (2021) examined surface mass balance (SMB) records reconstructed from six ice cores and two automatic weather stations. In these records, they observed increased snowfall during the 2016-17 MRP opening and assumed that similar anomalies occurred during the past MRP opening periods. Their study constructed a polynya index using two data assimilation approaches and one statistical method to reconstruct historical climate states based on SMB reconstruction from ice cores. Their study found multiple polynya openings since 1250 CE but observed that large polynya openings are rare. However, they rely only on snow accumulation records, which can be influenced by many factors other than polynya occurrences. The polynya signal in their ice-core SMB records is also weak compared to natural atmospheric variability, making it difficult to distinguish polynya-induced anomalies from SMB caused by other climatic fluctuations. The short instrumental record from automatic weather stations prevents a robust calibration of polynya reconstructions, increasing the risk of false positives (unrelated high snowfall events misinterpreted as polynya activity) and false negatives (missed polynya events). Additionally, the ice-core and instrumental records they used were collected from a large region, including the Weddell Sea sector and the ice sheet inland, where the surface elevation exceeds 2000 m a.s.l., which can hardly reflect changes in the vapour sources in the MRP regions directly (Fig. 1).

To overcome the limitations of the previous study, we develop a novel, multi-proxy approach to reconstruct past MRP activity using a high-resolution ice core, IND-36/9, from Djupranen Ice Rise in coastal Dronning Maud Land. While polynya opening results in increased heat and moisture exchange, the polynya years, characterised by a decrease in sea ice cover, also result in increased sea salt production. These signatures can be carried to the coastal region of Antarctica and deposited in the low-elevation ice shelves and ice rises. The back-trajectory analysis (Section 2.5) clearly indicates the direct atmospheric link between MRP and the IND-36/9 core site (Fig. 1). Our ice core site is situated within ~1000 km of the typical MRP location, close enough to potentially capture atmospheric signatures transported from polynya formation near the Maud Rise. By combining records of snow accumulation, water isotopes, and major ion concentrations, we aim to provide a more robust reconstruction of polynya activity. This approach leverages the strengths of selecting an ideal core site and employing multiple proxies to overcome the limitations of single-proxy reconstructions, thereby capturing the complex signatures of polynya events that extend beyond the last few decades, when satellite and instrumental records are available.

2. Study Area and Methods

2.1. Study area

The DML coast is characterised by distinct topographic features like ice rises having associated local ice flow, climate regime and SMB variability (Drews et al., 2015; Goel et al., 2017; Lenaerts et al., 2014; Matsuoka et al., 2015; Pratap et al., 2022; Rignot et al., 2019). As part of the Indo-Norwegian project MADICE, an ice core was drilled on the summit of the Djupranen Ice Rise (70.18° S, 9.18° E; elevation 321 m a.s.l.), at the western margin of the Nivlisen Ice Shelf in coastal Dronning Maud Land (DML), East Antarctica (Fig. 1). Our site survey found that this ice rise summit has been at the current position at least in the past few millennia (Pratap et al., 2022). This location also offers several advantages for polynya reconstruction: its high SMB allows annual layer counting (Dey et al., 2023) and its coastal proximity may provide sensitivity to maritime signals (Ejaz et al., 2021; Wauthy et al., 2024), while its elevation causes insignificant surface melt and local noise while maintaining regional signal strengths (Dey, 2023). Its position is also within primary atmospheric transport pathways from the Maud Rise region (Fig. 1).

2.2. Ice core: from the field to the laboratory

The drill site of the Djupranen ice rise was located at its summit, based on analysis of satellite altimetry data and satellite image analysis, followed by an ice-penetrating radar survey (Pratap et al., 2022). The radar survey also shows a flat bed topography below the summit (Goel et al., 2026) and the presence of Raymond arches, indicative of a relatively stable summit position in the past (Goel et al., 2020). The ice core was drilled using an electromechanical ice core drilling system (Model D2, GeoTec, Japan). Over nine days, a 122 m ice core (IND 36/B9; hereafter IND36/9) was retrieved. The drilled ice core sections were sealed in high-density polyethylene core bags, packed in expanded polypropylene boxes, and stored in a reefer container at -20°C until their transport to the National Centre for Polar and Ocean Research (NCPOR), Goa, India, where samples were stored in the in-house Ice Core Laboratory maintained at -20°C . The ice cores were processed in the -15°C core processing facility at NCPOR. The cores were initially cut into 3.5 cm thick, 10 cm wide slabs for line scanning and then subsequently sub-sampled at 5 cm resolution for stable isotope and chemical analysis. The samples for chemical analysis were cut into cuboids; the three dimensions of the samples were measured using a calliper and weighed using a weighing balance. The density for the samples was calculated as mass divided by the volume of each sample. The error in measuring the sample dimensions was ± 0.5 mm, while the weighing balance's uncertainty was ± 0.1 g for measurements up to 100 g. As a result, the density measurements have a propagated uncertainty of 5%.

Major inorganic anions (Cl^- , SO_4^{2-} , MSA^- and NO_3^-) and cations (Na^+ , NH_4^+ , K^+ , Mg^{2+} , and Ca^{2+}) were measured in the samples using an ICS 5000+ ion chromatograph (Thermo Dionex) equipped with a conductivity detector. Anions were separated on an AS11 (2 mm) column with potassium hydroxide as eluent and an AG11 (2 mm) guard column with AERS 500, 2 mm suppressor. Cations were separated on a CS17 (0.4 mm) capillary column with methane sulphonic acid as the eluent and the CG17 (0.4 mm) capillary guard column with a CCES 300 capillary suppressor. A 10 mg/L stock solution of Na^+ , NH_4^+ , K^+ , Ca^{2+} , and Mg^{2+} was mixed and then diluted with MiliQ ultrapure water to prepare standards for cation exchange chromatography. The anion standards were prepared from 10 ppm stock solutions of MSA^- , Cl^- , SO_4^{2-} , and NO_3^- . The dilutions were conducted volumetrically and were freshly prepared within a few days of each run, ranging from 5 ppb to 1 ppm. Eight standards were

selected from this range for calibration. Before analysis, the ice core samples were melted in a Class 100 clean room. The analytical precision for all ions was better than 10%.

The ice core samples were analysed for oxygen and hydrogen isotopic ratios at NCPOR using a Triple Isotope Water Analyser (TIWA-45EP from Los Gatos Research, USA), which works on the principle of off-axis integrated cavity output spectroscopy (OA-ICOS). The melted ice core samples were introduced into the TIWA-45EP without sample conversion through a PAL HTC-xt auto-injector (CTC Analytics) equipped with a heated ($\approx$ approximately 85 °C) injector block (LGR) (Berman et al., 2013). Using a Hamilton 1.2 μ L, zero-dead-volume syringe, samples were injected into the injector block and evaporated for direct isotope analysis. Measurements were completed at a speed of $\sim$ 90 s per individual injection. To eliminate sample-to-sample memory, a total of nine injections were made, with the first three being discarded for analysis. The last six injections were averaged to produce a single, high-throughput (HT) sample measurement. One commercially available working standard from LGR1C and two in-house laboratory standards (CDML1 and HL1) with known isotopic composition, spanning the entire range of our sample measurements (-46.19‰ to -19.49‰ for $\delta^{18}\text{O}$ and -362.85‰ to -154.0‰ for δD) were analysed routinely as reference waters after every five ice core samples to check the instrument performance. Laboratory standards are calibrated on the VSMOW/SLAP scale. The external precision obtained using our laboratory standards (CDML1 and HL1) for $\delta^{18}\text{O}$ was $\pm 0.046\text{‰}$ and $\pm 0.068\text{‰}$, respectively, and for δD was $\pm 0.32\text{‰}$ and $\pm 0.23\text{‰}$ (1σ standard deviation) for 30 samples. Replicate analyses performed based on ten samples yield repeatability of $\pm 0.76\text{‰}$ for δD and $\pm 0.09\text{‰}$ for $\delta^{18}\text{O}$. All the raw instrumental OA-ICOS data were processed in the LGR post-analysis software. Any measured injection with water number density outside the manufacturer's suggested range of $2\text{--}4.5 \times 10^{16}$ H_2O molecules/ cm^3 was discarded. Injections with incomplete evaporation were detected by examining the standard deviation of the measured water number density (σ_{nmeas}) as reported by the instrument (Berman et al., 2013). Processed raw data directly gives $\delta^{18}\text{O}$ and δD , which are further used to calculate deuterium excess [$\text{d-excess} = \delta\text{D}_{\text{ice}} - 8 \times \delta^{18}\text{O}_{\text{ice}}$].

2.3. Chronology development

The chronology of the IND36/9 core is based on a multiproxy approach involving annual layer determination from the stratigraphy of $\delta^{18}\text{O}$, major ions, and pixel intensity data from the line scanner, following Dey et al. (2023). The core has experienced limited surface

melt, with annual melt proportion varying between 0 and 4.4%, with a median melt proportion of 0.25% (Dey et al., 2023). Since diffusion in the firn attenuates high-frequency water-isotope information in ice cores (The Firn Symposium team, 2024), even in high accumulation sites of coastal Antarctica (Mahalinganathan et al., 2022), we diffusion-corrected our water isotope records following Jones et al. (2023) as detailed in the supplementary materials (Fig. S1). The shift in seasonal peaks in our corrected record is less than 5 cm, which falls within the sampling interval of 5 cm and is therefore insignificant in affecting the accuracy of the chronology.

A five-point smoothing was applied to the pixel intensity data to simplify annual layer counting and reduce noise in the record. We also used age tie points of known volcanic eruptions identified from non-sea-salt sulphate (nssSO₄) peaks, as well as the tritium bomb peak of 1962. Volcanic indicators (nssSO₄) have been used to identify specific, dated volcanic eruptions, allowing us to reduce the uncertainties resulting from the relative dating procedure. However, unambiguous eruption identifications are challenging in ice cores from coastal regions, where the nssSO₄ background signals are commonly highly variable due to the proximity of the ocean and ocean-related MSA products (Philippe et al., 2016).

A preliminary chronology was obtained from the annual counts, which was then refined using the nssSO₄ peaks (Fig. 2) from the well-established volcanic eruptions of Pinatubo (1991), El Chichon (1982), Agung (1963), Cerro Azul (1932), Santa Maria (1902), Krakatoa (1882), Cosiguina/Babuyan (1834), Tambora (1815) and unknown volcanic eruption (1809). Similar to Dey et al. (2023), we refined the chronology between the volcanic and Tritium tie points using the StratiCounter algorithm (Winstrup et al., 2012). The manual counts provide a basic framework from which StratiCounter develops and refines its statistical characterisation of annual layers, enabling adaptation to varying layer properties with depth (Winstrup et al., 2012). To minimise reliance on initial manual inputs, we conducted multiple iterations using refined layer templates derived from previous algorithm outputs. The chronology of the IND36/9 ice core provides a robust temporal framework that extends back to 1774 CE at a depth of 122m (Fig. 3).

2.4. Satellite-based polynya metrics

We followed Heuzé et al. (2021) to define the "polynya-prone" region of the Weddell Sea (6°W to 12°E, 68°S to 60°S, approximately 600 km x 900 km), focusing on the winter and early spring months (June 1 to October 31). This region was selected based on historical observations of polynya formation and its significance in Antarctic bottom water production (Campbell et al., 2019; Heuzé et al., 2021). Polynya activity was quantified using two independent metrics derived from satellite data. These metrics complement each other to provide a comprehensive assessment of polynya dynamics.

To quantify the polynya metrics, we use sea ice concentration data derived from passive microwave measurements using the NASA Team algorithm (Comiso and Nishio, 2008), processed at a spatial resolution of 25 x 25 km per pixel. The first metric, Polynya Days, represents the annual count of days (between June 1 and October 31), with a minimum sea ice concentration in the polynya-prone region below 60%. The second metric, Cumulative Polynya Area, measures the total extent of all polynya occurrences within a year (June 1 to October 31). This was calculated by summing the daily open water areas within the polynya prone region, defined as the cumulative sum of pixel area with sea ice concentration below the 60% threshold. The combination of these metrics allows us to characterise both the temporal persistence and spatial extent of polynyas, providing insights into their formation mechanisms and potential impact on regional oceanographic processes. The combination of these metrics allows us to characterise both the temporal persistence and spatial extent of polynyas, providing insights into their formation mechanisms and potential impact on regional oceanographic processes.

2.5. Air mass trajectory modelling

The HYbrid Single-Particle Lagrangian Integrated Trajectory (HySPLIT) model, developed by the NOAA Air Resources Laboratory (ARL), provides a means of generating back trajectories to identify the source of moisture uptake for precipitation (Markle et al., 2012). We used HYSPLIT version 5.3 for back trajectory computations, initialized with conditions from the 2.5 by 2.5-degree resolution NCEP/National Center for Atmospheric Research (NCAR) reanalysis data for the period 1948 to 2016. Forward trajectories were run for 240 hours (10 days), initialised every hour, starting over the polynya-prone region of the Weddell Sea. Groups of forward trajectories were computed with initial altitudes of 100, 200,

300, 400, and 500 meters above ground level, and in all cases, the vertical velocity from the meteorological data was used as the input for vertical motion.

We calculated trajectory density from multiple trajectories using HYSPLIT (Miller et al., 2002), because individual trajectories are highly sensitive to meteorological uncertainties, turbulence, and small initial condition variations, often leading to misleading or unreliable airflow pathways and potential position errors of up to 20% of the distance travelled. It also aids in differentiating between local circulation patterns and regional-scale flow features (Dorling and Davies, 1995) and has been previously applied to interpret polar ice core paleoclimate records (Dixon et al., 2012; Ejaz et al., 2021; Neff and Bertler, 2015). We calculated air mass transport densities for each year 1979 – 2016 from the HySPLIT output. The trajectory densities in each equal-area (1 by 1 degree) pixel were summed and divided by the total number of air mass trajectories.

3. Results

3.1. Identification of polynya years in satellite data

Satellite imagery is crucial for polynya studies as they provide continuous, high-resolution observations of sea ice dynamics, allowing researchers to monitor polynya formation, extent, and variability over time. Satellite observations spanning four decades reveal distinct periods of polynya activity in the Weddell Sea region since 1979 (Fig. 4). The most notable and well-documented event occurred during 1974 – 76, when the polynya reached an exceptional size of 300,000 km² (Carsey, 1980). This event, often referred to as the Great Weddell Polynya, represented a significant perturbation to the regional ocean-atmosphere system and has served as a benchmark for subsequent polynya observations. Following this major event, several smaller polynyas and polynya-like features ("halos") have been observed during the late 1980s to early 1990s and early 2000s (Heuzé et al., 2021). The two independent metrics used for polynya detection—Polynya Days and Cumulative Area—reveal distinct but complementary temporal patterns. This dual-metric approach enables a more comprehensive understanding of polynya dynamics than either metric alone could provide. While the magnitudes of these metrics do not exhibit direct correlation ($r = 0.42$, $p < 0.01$), both metrics successfully identify known major polynya/halo events between 1979 and 2016, and provide unique insights into their temporal and spatial characteristics. The Polynya Days metric

effectively captures persistent small-scale features, while the Cumulative Area metric better represents brief but large openings. While the record of polynya openings from satellite imagery provides crucial insights, their influence on coastal Antarctic ice cores could depend majorly on atmospheric transport.

3.2. Air mass transport patterns

We use forward air trajectory analysis to examine the transport pathways and determine whether polynya-derived signals can reach and be recorded in the ice cores. This approach allows us to trace the path of air masses from the polynya-prone area to potential ice core sites, providing insight into which locations are most likely to capture polynya signals. Forward trajectory analysis reveals consistent and well-defined transport pathways from the Maud Rise region to coastal Dronning Maud Land (Fig. 5). Frequency analysis shows that our study site (1 x 1 degree box surrounding the ice core site) receives approximately 2% of all trajectories originating from the MRP region, a statistically significant proportion ($p < 0.001$) that indicates reliable capture of polynya-related atmospheric signals. This percentage remains relatively stable across different seasons and years, suggesting a robust atmospheric connection between the source and deposition regions. The trajectory density analysis indicates that our ice core location falls within a primary atmospheric transport corridor originating from the MRP region. This corridor exhibits enhanced stability during the winter months when polynya formation typically occurs. However, trajectory calculations are susceptible to significant spatial error of 15 – 30% of distance travelled (Draxler, 2008). Therefore, a combination of forward and backward trajectories would provide a better overview of the travel pathways. Our backward trajectory analysis (Fig. 6) further confirms the findings from the forward trajectory patterns, showing trajectories originating from the polynya-prone region influencing our ice core site. We observe that air masses reaching our site also originate from more distant open-water areas (even during the winter months) in the Southern Ocean. However, these distal sources remain relatively consistent across the study period and are located significantly further from the IND-36/9 site compared to the polynya-prone region. Consequently, these background marine inputs are less likely to drive the synchronized, high-magnitude anomalies observed across all proxies, particularly the sharp increases in ssNa flux, that characterize the polynya years. Trajectory analysis of the transport pathways, therefore, confirms that winds can carry polynya-derived

signals from the open water to our study site and preserve them in the ice core records, regardless of polynya presence.

3.3. Development of the polynya index

3.3.1. Ice core record and proxy selection

The ice core proxy dataset from 1774 to 2016 exhibits distinct variability across all measured properties. We choose four major proxies for developing our polynya index: sea-salt sodium (ssNa), deuterium excess (d-excess), $\delta^{18}\text{O}$, and snow accumulation (Fig. 7a). Sodium concentration ranges from a low of 19.1 to 218.6 ppb, with a mean value of 89.9 ppb. Some periods show sustained higher values, such as the late 1700s and early 1800s, while others, including the late 19th and 20th centuries, display more variable levels. D-excess varies between -1.8 to 10.8‰, with a mean of 4.3‰, showing notable peaks in the late 19th century and sharp declines in the early 20th century. $\delta^{18}\text{O}$ values fluctuate between -21.6 to -15.6‰, with a mean of -17.8‰, displaying alternating periods of enrichment and depletion, including a gradual decline over recent decades. Annual snow accumulation ranges from 0.12 to 0.83 m w.e., with a mean of 0.39 m w.e., exhibiting distinct fluctuations, including lower values in the early 20th century and a more variable pattern in recent decades.

The opening of a polynya results in increased heat and moisture exchange between the warm, open water and the cold air. Therefore, during polynya years (decrease in the sea ice cover), there is also an increase in local precipitation in the Weddell Sea region (Moore et al., 2002) and resultant snow accumulation further inland (Goosse et al., 2021). The water isotope ratios of precipitation are often related to temperature at the precipitation site (Ejaz et al., 2022; Naik et al., 2010) and the distance of transport (Goursaud et al., 2017; Klein et al., 2019), and the pattern of annual mean water isotopes of precipitation associated with polynya formation would relatively be similar to that for increased temperature from the heat exchange due to polynya opening and shorter transport distance, especially over a coastal site. While sea ice typically acts as a physical barrier to air-sea exchange, a polynya creates a localized window for enhanced aerosol production via multiple distinct mechanisms. First, the open-water surface provides a direct source where high-speed cyclonic winds and storm activity trigger bubble-bursting processes, injecting sea-salt particles into the marine boundary layer. Second, extensive open water polynyas can also act as a large factory for sea ice production, comparable

to the production of the largest coastal polynyas (Zhou et al., 2023). Previous studies have shown that the surface of fresh sea ice, including frost flowers, is an important source of sea salt to the Antarctic, and the production of frost flowers is controlled by the amount of new sea ice production (Wolff et al., 2003). Furthermore, blowing snow across these newly formed, brine-wetted ice surfaces facilitates the sublimation of salty snow particles, which is also the major source of sea salt aerosol to the inland sites (Frey et al., 2020). Because these processes occur proximately to the drilling site and are situated directly along the atmospheric trajectories identified in Figure 6, they result in high-magnitude pulses of ssNa that are chemically distinct from the far, year-round open-ocean baseline. We however use ssNa flux instead of concentration to mitigate the potentially confounding effects of varying annual snow accumulation rates. By using flux, we ensure that a dilution effect during high-snowfall years doesn't artificially lower our ssNa signal, and conversely, that a low-snowfall year doesn't create a false peak.

3.3.2. Development of the integrated polynya index

To reconstruct past MRP activity, we developed a multi-proxy index using a cumulative-anomaly approach, calculated as the means of rolling z-scores. This allows us to isolate anomalies that are synchronized across proxies from long-term climate trends. We use four independent proxies: sea-salt sodium (ssNa) flux, $\delta^{18}\text{O}$, deuterium excess (d-excess), and annual snow accumulation rates for our polynya index calculation. Prior to index calculation, all proxies are scaled to 0 – 1 to remove bias toward proxies with greater variability. For each proxy, we first calculate a 30-year rolling z-score to identify annual departures from the local background state. This accounts for non-stationary signals in the 250-year record, such as the gradual depletion of water isotopes or recent decreases in snow accumulation rates. As explained in section 3.3.1, MRP openings are linked to positive anomalies in ssNa flux, $\delta^{18}\text{O}$, d-excess, and annual snow accumulation rates; therefore, we used only positive z-scores for the calculation. Negative z-scores were set to zero to prevent them from cancelling out physically significant peaks in other proxies. To test the dependence of the polynya index on snow accumulation, we calculate two versions of the index: one using all four selected proxies and another excluding annual snow accumulation rates. The polynya index is a mean of all individual rolling z-scores, which is then normalized so that the final values are such that the largest known polynya opening of 1974 – 76 is set to 1. A polynya year (a year when a polynya

activity is recorded) is defined as a year in which the polynya index exceeds the likelihood threshold (0.6, 0.8, or 1). The polynya index is, therefore, an indicator of polynya occurrence from ice core observations and not a measure of the absolute extent or duration of the polynya events.

3.4 Past Maud Rise Polynya activity

Our ~250-year reconstruction reveals complex and significant temporal variability in polynya occurrence, shedding new light on the long-term behaviour of the Maud Rise Polynya (MRP). We identified twenty-five potential polynya years from 1774 to 2016 (likelihood >0.6; Fig. 7b); with three events (1788, 1834 and 1901) exceeding the threshold of 1.0, indicating that these were events of possibly similar magnitude as the 1974 – 76 polynya opening. The high index values for these events likely indicate that they were large-scale and persistent openings; however, exact quantification of the polynya extent is beyond the scope of this study. During the late 18th to mid-19th century, the polynya exhibited a high-frequency variability, frequently dropping to near-zero levels as seen in 1782 and the early 1800, while also producing moderate activity clusters above the 0.60 threshold in the 1810s and late 1820s. The late 19th century transitioned into a period of more consistent activity, with 1887 being a notable high-activity year exceeding 0.80. The early 20th century was marked by a prominent polynya in 1901 followed by significant bursts of activity exceeding the 0.60 threshold in 1911, 1924, and 1926. Mid-century activity was relatively suppressed, though a significant spike above 0.90 occurred in 1951. The lesser known 1964 polynya opening is also recorded in our polynya index (1965 in our polynya index); however, the likelihood is relative lower (>0.6) than the 1974 – 76 event. Following the 1974 – 76 event, the post-1980 era has generally shifted toward lower average values, though moderate spikes exceeding the 0.60 threshold occurred in 1982, 2003, 2006, and 2010, corresponding to known halo years.

4. Discussion

We compared our polynya index with those from Goosse et al. (2021) and found that the major polynyas identified in our study over the common time period are mostly identified in their reconstruction. A direct annual correlation is however statistically limited by the independent age-scale uncertainties, typically ± 1 –2 years, inherent in all ice core records. These chronological offsets cause high-magnitude events to appear out of phase between two

indices/studies, which artificially degrades linear correlation coefficients despite clear physical synchronization in the raw signals. To provide a more robust and physically meaningful assessment, we used a binary presence-absence matrix assuming a ± 2 years lead/lag window at threshold levels of 0.6, 0.7, 0.8, 0.9 and 1. Therefore, we look for polynya events at different threshold levels in our record and then look for polynya events with a threshold of 0.6 from the polynya index from Goosse et al. (2021) within a ± 2 years window. For simplicity, we focus on only one index from each study, simple average of the standardized time series (Stat) from Goosse et al. (2021) and polynya index including accumulation from this study. We find that nine polynya years (threshold = 0.6) in our index that are also corroborated by the Goosse et al. (2021) record (Figure 9a). Similarly, seven polynya years are corroborated at a threshold of 0.7 and three at 0.8. At the threshold of 0.9 and 1, only the polynya event of 1974 is corroborated, which is expected as both studies use this event as an assumed truth and normalize all event to it. Similarly, when we repeat the same process for polynyas identified in Goosse et al. (2021) (Figure 9b), we find that 17 polynya years are corroborated at 0.6 threshold, 12 polynya years at 0.7, 9 polynya years at 0.8, eight at 0.9 and seven polynya years with a threshold of 1 are also identified in our polynya index.

This provides a degree of validation for both approaches, suggesting that they are capturing similar large-scale polynya events. However, our index shows enhanced sensitivity to polynya formations, identifying the polynya occurrence of 1964, even though with a lower likelihood. This low likelihood of the 1964 polynya is possibly due to the lower persistence as compared to the well-documented events during 1974 – 76 (Meier et al., 2013). Furthermore, our index shows better performance in identifying transient polynyas and halos, which were missed in the reconstruction by Goosse et al. (2021). These shorter-lived or less intense polynya events, while not as dramatic as major openings, play a crucial role in regional oceanography and climate dynamics. Their detection provides a more comprehensive picture of polynya activity over time. Beyond chronological offsets, discrepancies between the two indices likely reflect differences in regional sensitivity and site selection. Our analysis of air mass transport frequency from the MRP regions shows a sink in the coastal regions of Dronning Maud Land, with a significant proportion of the trajectories reaching our study area. In contrast, four out of the six ice core sites used in Goosse et al. (2021) fall outside the primary sink of these trajectories (Fig. 5). This difference in site selection could be a major reason for the observed discrepancy between the two polynya indices. Ice cores from sites that rarely receive air masses

from the polynya region may not be reliably used to reconstruct polynya events or may do so with reduced sensitivity. Whereas our ice core site is well located to capture the occurrence of the MRP opening, as it frequently receives air masses originating from the polynya region. However, Goosse et al. (2021) still manage to detect the large, multi-year polynyas as they lead to a widespread positive anomaly in precipitation over the coastal and continental region. Consequently, while our index captures both high-magnitude regional events (e.g., 1974) and more localized openings (e.g., 1964), inland records may only register the most extreme regional anomalies. This suggests that the lack of a perfect year-to-year correlation is not a limitation of the proxies themselves, but rather a reflection of the better spatial representativeness of the IND-36/9 site for capturing the nuances of the Maud Rise polynya system. Another possible reason for this observed difference in sensitivity could be the robustness of our multiproxy approach in identifying polynyas, as it is less susceptible to noise or artefacts in any single proxy. It is crucial to note that the observed differences in peak occurrences do not necessarily invalidate either dataset but rather highlight the complexity in reconstructing past polynya activity. These discrepancies underscore the challenges inherent in paleoclimate reconstruction and the potential complementarity of different methodologies in building a comprehensive understanding of past polynya dynamics.

Our polynya reconstruction aligns with and extends satellite era observations of MRP formation mechanisms. Recent studies have shed light on the complex interplay of oceanic and atmospheric factors controlling MRP formation and persistence (Campbell et al., 2019; Jena et al., 2019). However, extending these insights to longer timescales remains challenging. Our multi-century reconstruction offers a unique opportunity to explore the long-term drivers of MRP variability. Autonomous profiling float observations during 2016 and 2017 revealed that the MRP were initiated and modulated by the passage of severe storms, and the intense heat loss drove deep overturning within them (Campbell et al., 2019). Wind-driven upwelling of record strength weakened haline stratification in the upper ocean, thus favouring destabilisation. A recent study, however, highlighted the role of salt transport through northward Ekman transport as an additional mechanism of the polynya formation and these processes were driven by intensified eastward surface stresses during 2015 – 2018 (Narayanan et al., 2024). Therefore, it is crucial to understand the maritime climate variability during the polynya and non-polynya years. Our reconstruction shows alternation between polynya-active and quiescent periods, which may provide valuable context for assessing how future MRP

behaviour may evolve under anthropogenic climate change. The temporal clustering of MRPs in our record possibly indicates that once favourable preconditioning develops, the region remains susceptible to repeated polynya openings over several years or decades. As global temperatures rise, projected changes in Southern Ocean wind patterns, sea-ice extent, and surface freshening may shift the thresholds required for deep convection, thereby altering both the frequency and persistence of MRPs. Understanding how atmospheric and oceanic conditions differ between the polynya and non-polynya periods identified in this study will be critical for refining projections. Integrating these insights with high-resolution coupled ocean–atmosphere models could help identify the physical thresholds and feedback governing MRP formation. Such integration is crucial for predicting how rare but climatically significant features such as the Maud Rise Polynya may influence Southern Ocean overturning, carbon exchange, and global climate in a warming world.

5. Conclusions and Outlook

We provide new insights into the temporal variability of Maud Rise Polynya, one of the largest and well-known open ocean polynyas, by integrating multiple proxies using an ice core from coastal Dronning Maud Land, representing the past nearly 250 years. We identified multiple polynya openings during the 1830s, 1880s, and 1970s. These clusters of polynya openings possibly correspond to a specific combination of atmospheric circulation patterns and oceanographic preconditioning, suggesting a more deterministic framework for MRP development. Over the 250-year record, we identify large events during 1833 – 34 and 1911, which were comparable in magnitude to the documented 1974 – 76 polynya, indicating that such openings are not isolated anomalies but represent recurring features of the Southern Ocean system. These events are characterised by proxy signatures consistent with modern polynya conditions, supporting the persistence of underlying formation mechanisms throughout the observational period. Conversely, the interval from 1920 to 1950, marked by a notable absence of polynya activity, coincides with documented changes in Southern Ocean circulation and atmospheric forcing. This prolonged quiescent phase provides valuable baseline information for assessing natural variability in MRP activity on multi-decadal timescales.

Our multiproxy approach significantly enhances the sensitivity of the reconstruction, particularly in detecting smaller-magnitude and shorter-duration events that may have been overlooked in previous studies. As a result, the reconstructed history reveals a more dynamic

and variable pattern of MRP activity than previously recognised, with implications for understanding its role in ocean ventilation, carbon cycling, and regional climate variability. Our study reveals that synthesising paleoclimate records, modern observations, and modelling approaches can provide a robust foundation for advancing the understanding of the MRP as a critical and recurrent component of the Southern Ocean climate system.

Acknowledgements

The ice core was drilled as part of the India–Norway MADICE project, with support from the Ministry of Earth Sciences, Government of India, and the Research Council of Norway. Ice core drilling and field expeditions have been made successful with the support of the NCPOR logistics leaders, the Maitri logistics team, and all field participants. K. Mahalinganathan and Tariq Ejaz are acknowledged for their contributions to ice core processing and analysis. We acknowledge the use of imagery from the NASA Worldview application (<https://worldview.earthdata.nasa.gov>), part of the NASA Earth Science Data and Information System (ESDIS).

Financial support

This research has been supported by the Ministry of Earth Sciences, India (grant no. MoES/Indo-Nor/PS-3/2015) and the Research Council of Norway (grant no. 248780) for the joint India–Norway project, ‘Mass balance, dynamics, and climate of the central Dronning Maud Land coast, East Antarctica’ (MADICE).

Author contributions

RD and MT defined the study objectives. RD led the ice core processing and analysis with support from AP, BLR and CML. RD led the data analysis and interpretation with inputs from KM and CML. RD prepared the manuscript with feedback from all co-authors. MT and KM were the project leaders

Data Availability

All ice core data and polynya indices are being prepared for submission to the NCPOR Polar Data Centre (<https://data.ncpor.res.in>). For final archival, open data formats will be used. A DOI will be generated and included in the final version of the paper. Data access can be granted

517 to reviewers promptly upon request.

518 **Declaration of competing interest**

519 The authors declare that they have no known competing financial interests or personal
520 relationships that could have appeared to influence the work reported in this paper.




Figure 1 | Study area and wind-trajectory patterns. (a) Maud Rise Polynya during its largest recent opening on 25 September 2017, detected with Moderate Resolution Imaging Spectroradiometer (MODIS) on NASA's Aqua satellite (from NASA Worldview, <https://worldview.earthdata.nasa.gov/>, last access: 01 April 2026). The polynya is visible as the dark, open-water feature amidst the surrounding white sea ice (b) MODIS Satellite imagery taken on the same day in a non-polynya year, 25 September 2022. The MODIS corrected reflectance imagery provides a true colour representation where the open ocean appears dark gray or black, while sea ice and clouds appear white or light gray. Images are overlaid with the trajectory frequency of back-trajectories ending at the Djupranen ice core site (green star) from June to October in 2017 and 2022, respectively (see methods). The Indian Maitri Station is marked by a yellow square. The colourmap is on a logarithmic scale. The inset shows the location of the main map. Grounding line and calving front are marked (Matsuoka et al., 2015).





Figure 2 | Tie points for ice core chronology. Volcanic events are identified from the non-sea-salt sulphate flux records. Only major volcanic events used as age tie points for reconstructing the chronology are marked.



539

540 **Figure 3 | Age-depth scale for the Djupranen ice core.** Tie points used for chronology (Fig.
 541 2) are shown with dashed lines. Major time markers are shown using the black dashed lines.
 542 Minimal deviation from the intersection points of the vertical and horizontal lines indicates the
 543 robustness of the chronology.





Figure 4 | Satellite polynya metrics. Two different annual metrics of satellite-derived polynya occurrence. Since the polynya is highly dynamic temporally and spatially, the metrics do not show a one-to-one resemblance. The red vertical bar shows the 1974 – 76 polynya event, which reached a peak extent of approximately 300,000 km² (Carsey 1980), more than ten times of the maximum cumulative polynya area observed between 1979 – 2016.

550



551

552 **Figure 5 | Airmass transport pathways originating from MRP:** Trajectory frequency of
 553 forward trajectories originating over the polynya-prone region (Heuzé et al., 2021) of the
 554 Weddell Sea, showing the prevalent transport pathway during the winter to early spring months
 555 for the period 1948 – 2016. A significant proportion of the trajectories end over our ice core
 556 site (red diamond). Location of ice cores used by Goose et al. (2021) are shown with green
 557 circles. The black trapezium shows the polynya prone regions. The gray region shows area
 558 with no trajectories.




Figure 6 | Airmass transport pathways ending at the ice core site: Trajectory frequency of back trajectories originating from the ice core site during a well-known polynya year (2017; left) and a non-polynya year (2022; right) is shown. The trajectory frequency pattern during both years is very similar, indicating a consistent transport pathway during most of the years and suggesting that our ice core records activity over the polynya-prone region of the Weddell Sea (black trapezium) in all years. The gray region shows areas with no trajectories. The dashed lines show the NSIDC median sea ice extent for 1980 – 2010, with green indicating September (maximum) and red representing February (minimum) sea ice extent.





Figure 7 | (a) Ice core proxies and polynya identification. Annual variability of selected ice core proxies and polynya index from 1960 – 2016. The red curve is the moving median over a 30-year window. The two known polynya events of 1964 and 1974 – 76 are marked with gray patches. **(b) Ice core derived polynya indices:** Ice core derived polynya indices in the past 250 years using the anomalies in the annual record of five ice-core proxies: Na, deuterium excess, $\delta^{18}\text{O}$, and snow accumulation. To test the dependence of the polynya index on snow accumulation, we calculate two indices, one using snow accumulation (blue curve) and another

576 without (red curve). The two indices behave similarly over the entire time period; however, the
577 index without snow accumulation shows a higher range of variability.




Figure 8 | Comparison of the polynya indices. Polynya indices reconstructed in this study (upper panel) are shown with the polynya indices from Goosse et al. (2021) (lower panel). The polynya indices from Goosse et al. (2021) are based on six surface mass balance records using data assimilation with two control simulations performed with the SPEAR (Seamless system for Prediction and Earth system Research) global climate model, SPEAR_AM2 (DAAM2; Blue) and SPEAR_LO (DALO; green), and a simple average of the standardized time series (Stat, red). The known polynya events of 1964 (1965 in our record and absent in Goosse et al. (2021) and 1974 – 76 (1974 in our record and 1975 in Goosse et al. (2021)) are labelled.




Figure 9 | Comparison of synchronous polynya events. Binary presence-absence analysis comparing the reconstructed polynya index from this study and the index from Goosse et al. (2021) across five threshold levels (0.6, 0.7, 0.8, 0.9, and 1.0) and assuming a lead/lag of ± 2 years. (a) Temporal distribution of polynya events identified at different threshold in this study and corroborated by the Goosse et al. (2021) record (threshold ≥ 0.6). (b) Distribution of polynya events from Goosse et al. (2021) at different threshold corroborated by this study (threshold ≥ 0.6).

596 **References**

- 597 Akitomo, K.: Thermobaric deep convection, baroclinic instability, and their roles in vertical
 598 heat transport around Maud Rise in the Weddell Sea, *Journal of Geophysical Research: Oceans*, 111, 2006. <https://doi.org/10.1029/2005JC003284>
 599
- 600 Arrigo, K. R. and van Dijken, G. L.: Phytoplankton dynamics within 37 Antarctic coastal
 601 polynya systems, *Journal of Geophysical Research: Oceans*, 108, 2003.
 602 <https://doi.org/10.1029/2002JC001739>
- 603 Årthun, M., Holland, P. R., Nicholls, K. W., and Feltham, D. L.: Eddy-Driven Exchange
 604 between the Open Ocean and a Sub-Ice Shelf Cavity, *Journal of Physical Oceanography*,
 605 43, 2372-2387, 2013. <https://doi.org/10.1175/JPO-D-13-0137.1>
- 606 Bennetts, L. G., Shakespeare, C. J., Vreugdenhil, C. A., Foppert, A., Gayen, B., Meyer, A.,
 607 Morrison, A. K., Padman, L., Phillips, H. E., Stevens, C. L., Toffoli, A., Constantinou, N.
 608 C., Cusack, J. M., Cyriac, A., Doddridge, E. W., England, M. H., Evans, D. G., Heil, P.,
 609 Hogg, A. M., Holmes, R. M., Huneke, W. G. C., Jones, N. L., Keating, S. R., Kiss, A. E.,
 610 Kraitzman, N., Malyarenko, A., McConnochie, C. D., Meucci, A., Montiel, F., Neme, J.,
 611 Nikurashin, M., Patel, R. S., Peng, J.-P., Rayson, M., Rosevear, M. G., Sohail, T., Spence,
 612 P., and Stanley, G. J.: Closing the Loops on Southern Ocean Dynamics: From the
 613 Circumpolar Current to Ice Shelves and From Bottom Mixing to Surface Waves, *Reviews*
 614 *of Geophysics*, 62, e2022RG000781, 2024. <https://doi.org/10.1029/2022RG000781>
- 615 Berman, E. S. F., Levin, N. E., Landais, A., Li, S., and Owano, T.: Measurement of $\delta^{18}\text{O}$,
 616 $\delta^{17}\text{O}$, and ^{17}O -excess in Water by Off-Axis Integrated Cavity Output Spectroscopy and
 617 Isotope Ratio Mass Spectrometry, *Analytical Chemistry*, 85, 10392-10398, 2013.
 618 <https://doi.org/10.1021/ac402366t>.
- 619 Campbell, E. C., Wilson, E. A., Moore, G. W. K., Riser, S. C., Brayton, C. E., Mazloff, M.
 620 R., and Talley, L. D.: Antarctic offshore polynyas linked to Southern Hemisphere climate
 621 anomalies, *Nature*, 570, 319-325, 2019. <https://doi.org/10.1038/s41586-019-1294-0>
- 622 Carsey, F. D.: Microwave Observation of the Weddell Polynya, *Monthly Weather Review*,
 623 108, 2032-2044, 1980. [https://doi.org/10.1175/1520-0493\(1980\)108<2032:MOOTWP>2.0.CO;2](https://doi.org/10.1175/1520-0493(1980)108<2032:MOOTWP>2.0.CO;2)
 624
- 625 Comiso, J. C. and Gordon, A. L.: Recurring polynyas over the Cosmonaut Sea and the Maud
 626 Rise, *Journal of Geophysical Research: Oceans*, 92, 2819-2833, 1987.
 627 <https://doi.org/10.1029/JC092iC03p02819>
- 628 de Lavergne, C., Palter, J. B., Galbraith, E. D., Bernardello, R., and Marinov, I.: Cessation of
 629 deep convection in the open Southern Ocean under anthropogenic climate change, *Nature*

Climate Change, 4, 278-282, 2014. <https://doi.org/10.1038/nclimate2132>

Dey, R.: Reconstruction of Antarctic climate variability using high resolution ice core stratigraphy, 2023. Goa University, 2023. <https://doi.org/10.5281/zenodo.12705334>

Dey, R., Thamban, M., Laluraj, C. M., Mahalinganathan, K., Redkar, B. L., Kumar, S., and Matsuoka, K.: Application of visual stratigraphy from line-scan images to constrain chronology and melt features of a firn core from coastal Antarctica, *Journal of Glaciology*, 69, 179-190, 2023. <https://doi.org/10.1017/jog.2022.59>

Dixon, D. A., Mayewski, P. A., Goodwin, I. D., Marshall, G. J., Freeman, R., Maasch, K. A., and Sneed, S. B.: An ice-core proxy for northerly air mass incursions into West Antarctica, *International Journal of Climatology*, 32, 1455-1465, 2012. <https://doi.org/10.1002/joc.2371>

Dorling, S. R. and Davies, T. D.: Extending cluster analysis—synoptic meteorology links to characterise chemical climates at six northwest European monitoring stations, *Atmospheric Environment*, 29, 145-167, 1995. [https://doi.org/10.1016/1352-2310\(94\)00251-F](https://doi.org/10.1016/1352-2310(94)00251-F)

Draxler, R.: NOAA-Air resources laboratory-FAQ-How do i estimate the absolute (in km) and relative (%) errors when using the HYSPLIT trajectory model. Sept, 2008.

Drews, R., Matsuoka, K., Martín, C., Callens, D., Bergeot, N., and Pattyn, F.: Evolution of Derwael Ice Rise in Dronning Maud Land, Antarctica, over the last millennia, *Journal of Geophysical Research: Earth Surface*, 120, 564-579, 2015. <https://doi.org/10.1002/2014jf003246>

Ejaz, T., Rahaman, W., Laluraj, C. M., Mahalinganathan, K., and Thamban, M.: Rapid Warming Over East Antarctica Since the 1940s Caused by Increasing Influence of El Niño Southern Oscillation and Southern Annular Mode, *Frontiers in Earth Science*, Volume 10 - 2022, 2022. <https://doi.org/10.3389/feart.2022.799613>

Ejaz, T., Rahaman, W., Laluraj, C. M., Mahalinganathan, K., and Thamban, M.: Sea Ice Variability and Trends in the Western Indian Ocean Sector of Antarctica During the Past Two Centuries and Its Response to Climatic Modes, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033943, 2021. <https://doi.org/10.1029/2020JD033943>

Frey, M. M., Norris, S. J., Brooks, I. M., Anderson, P. S., Nishimura, K., Yang, X., Jones, A. E., Nerentorp Mastromonaco, M. G., Jones, D. H., and Wolff, E. W.: First direct observation of sea salt aerosol production from blowing snow above sea ice, *Atmos. Chem. Phys.*, 20, 2549–2578, 2020. <https://doi.org/10.5194/acp-20-2549-2020>

Goel, V., Brown, J., and Matsuoka, K.: Glaciological settings and recent mass balance of Blåskimen Island in Dronning Maud Land, Antarctica, *The Cryosphere*, 11, 2883-2896,

2017. <https://doi.org/10.5194/tc-11-2883-2017>
- Goel, V., Matsuoka, K., Berger, C. D., Lee, I., Dall, J., and Forsberg, R.: Characteristics of ice rises and ice rumpled in Dronning Maud Land and Enderby Land, Antarctica, *Journal of Glaciology*, 66, 1064-1078, 2020. <https://doi.org/10.1017/jog.2020.77>
- Goel, V., Martín, C., Matsuoka, K., Pratap, B., Moholdt, G., Dey, R., Laluraj, C. M., and Thamban, M.: A new coastal ice-core site identified in Dronning Maud Land, Antarctica, for high-resolution climate reconstructions to the Last Glacial Maximum, *The Cryosphere*, 20, 1363–1378, 2026. <https://doi.org/10.5194/tc-20-1363-2026>
- Goosse, H., Dalaiden, Q., Cavitte, M. G. P., and Zhang, L.: Can we reconstruct the formation of large open-ocean polynyas in the Southern Ocean using ice core records?, *Clim. Past*, 17, 111-131, 2021. <https://doi.org/10.5194/cp-17-111-2021>
- Gordon, A. L. and Comiso, J. C.: Polynyas in the Southern Ocean, *Scientific American*, 258, 90-97, 1988.
- Goursaud, S., Masson-Delmotte, V., Favier, V., Preunkert, S., Fily, M., Gallée, H., Jourdain, B., Legrand, M., Magand, O., Minster, B., and Werner, M.: A 60-year ice-core record of regional climate from Adélie Land, coastal Antarctica, *The Cryosphere*, 11, 343-362, 2017. <https://doi.org/10.5194/tc-11-343-2017>
- Gülk, B., Roquet, F., Naveira Garabato, A. C., Bourdallé-Badie, R., Madec, G., and Giordani, H.: Impacts of Vertical Convective Mixing Schemes and Freshwater Forcing on the 2016–2017 Maud Rise Polynya Openings in a Regional Ocean Simulation, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS004106, 2024. <https://doi.org/10.1029/2023MS004106>
- Heuzé, C., Heywood, K. J., Stevens, D. P., and Ridley, J. K.: Southern Ocean bottom water characteristics in CMIP5 models, *Geophysical Research Letters*, 40, 1409-1414, 2013. <https://doi.org/10.1002/grl.50287>
- Heuzé, C., Zhou, L., Mohrmann, M., and Lemos, A.: Spaceborne infrared imagery for early detection of Weddell Polynya opening, *The Cryosphere*, 15, 3401-3421, 2021. <https://doi.org/10.5194/tc-15-3401-2021>
- Holland, D. M.: Explaining the Weddell Polynya--a Large Ocean Eddy Shed at Maud Rise, *Science*, 292, 1697-1700, 2001. <https://doi.org/10.1126/science.1059322>
- Jacobs, S. S., Gordon, A. L., and Ardai, J. L.: Circulation and Melting Beneath the Ross Ice Shelf, *Science*, 203, 439-443, 1979. <https://doi.org/10.1126/science.203.4379.439>
- Jena, B., Ravichandran, M., and Turner, J.: Recent Reoccurrence of Large Open-Ocean Polynya on the Maud Rise Seamount, *Geophysical Research Letters*, 46, 4320-4329, 2019.

699 <https://doi.org/10.1029/2018GL081482>

700 Jones, T. R., Cuffey, K. M., Roberts, W. H. G., Markle, B. R., Steig, E. J., Stevens, C. M.,
701 Valdes, P. J., Fudge, T. J., Sigl, M., Hughes, A. G., Morris, V., Vaughn, B. H., Garland,
702 J., Vinther, B. M., Rozmiarek, K. S., Brashear, C. A., and White, J. W. C.: Seasonal
703 temperatures in West Antarctica during the Holocene, *Nature*, 613, 292-297, 2023.
704 <https://doi.org/10.1038/s41586-022-05411-8>

705 Klein, F., Abram, N. J., Curran, M. A. J., Goosse, H., Goursaud, S., Masson-Delmotte, V.,
706 Moy, A., Neukom, R., Orsi, A., Sjolte, J., Steiger, N., Stenni, B., and Werner, M.:
707 Assessing the robustness of Antarctic temperature reconstructions over the past
708 2 millennia using pseudoproxy and data assimilation experiments, *Clim. Past*, 15, 661-
709 684, 2019. <https://doi.org/10.5194/cp-15-661-2019>

710 Lenaerts, J. T. M., Brown, J., Van Den Broeke, M. R., Matsuoka, K., Drews, R., Callens, D.,
711 Philippe, M., Gorodetskaya, I. V., Van Meijgaard, E., Reijmer, C. H., Pattyn, F., and Van
712 Lipzig, N. P. M.: High variability of climate and surface mass balance induced by
713 Antarctic ice rises, *Journal of Glaciology*, 60, 1101-1110, 2014.
714 <https://doi.org/10.3189/2014JoG14J040>

715 Lindsay, R. W., Holland, D. M., and Woodgate, R. A.: Halo of low ice concentration observed
716 over the Maud Rise seamount, *Geophysical Research Letters*, 31, 2004.
717 <https://doi.org/10.1029/2004GL019831>

718 Markle, B. R., Bertler, N. A. N., Sinclair, K. E., and Sneed, S. B.: Synoptic variability in the
719 Ross Sea region, Antarctica, as seen from back-trajectory modeling and ice core analysis,
720 *Journal of Geophysical Research: Atmospheres*, 117, 2012.
721 <https://doi.org/10.1029/2011JD016437>

722 Matsuoka, K., Hindmarsh, R. C. A., Moholdt, G., Bentley, M. J., Pritchard, H. D., Brown, J.,
723 Conway, H., Drews, R., Durand, G., Goldberg, D., Hattermann, T., Kingslake, J., Lenaerts,
724 J. T. M., Martín, C., Mulvaney, R., Nicholls, K. W., Pattyn, F., Ross, N., Scambos, T., and
725 Whitehouse, P. L.: Antarctic ice rises and rumples: Their properties and significance for
726 ice-sheet dynamics and evolution, *Earth-Science Reviews*, 150, 724-745, 2015.
727 <https://doi.org/10.1016/j.earscirev.2015.09.004>

728 McHedlishvili, A., Spreen, G., Melsheimer, C., and Huntemann, M.: Weddell Sea polynya
729 analysis using SMOS–SMAP apparent sea ice thickness retrieval, *The Cryosphere*, 16,
730 471-487, 2022. <https://doi.org/10.5194/tc-16-471-2022>

731 McPhee, M. G.: Is thermobaricity a major factor in Southern Ocean ventilation?, *Antarctic*
732 *Science*, 15, 153-160, 2003. <https://doi.org/10.1017/S0954102003001159>

733 Meier, W. N., Gallaher, D., and Campbell, G. G.: New estimates of Arctic and Antarctic sea

ice extent during September 1964 from recovered Nimbus I satellite imagery, *The Cryosphere*, 7, 699-705, 2013. <https://doi.org/10.5194/tc-7-699-2013>

Moore, G. W. K., Alverson, K., and Renfrew, I. A.: A Reconstruction of the Air–Sea Interaction Associated with the Weddell Polynya, *Journal of Physical Oceanography*, 32, 1685-1698, 2002. [https://doi.org/10.1175/1520-0485\(2002\)032<1685:AROTAS>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<1685:AROTAS>2.0.CO;2)

Naik, S. S., Thamban, M., Laluraj, C. M., Redkar, B. L., and Chaturvedi, A.: A century of climate variability in central Dronning Maud Land, East Antarctica, and its relation to Southern Annular Mode and El Niño–Southern Oscillation, *Journal of Geophysical Research: Atmospheres*, 115, 2010. <https://doi.org/10.1029/2009JD013268>

Narayanan, A., Roquet, F., Gille, S. T., Gülk, B., Mazloff, M. R., Silvano, A., and Naveira Garabato, A. C.: Ekman-driven salt transport as a key mechanism for open-ocean polynya formation at Maud Rise, *Science Advances*, 10, eadj0777, 2024. <https://doi.org/10.1126/sciadv.adj0777>

Neff, P. D. and Bertler, N. A. N.: Trajectory modeling of modern dust transport to the Southern Ocean and Antarctica, *Journal of Geophysical Research: Atmospheres*, 120, 9303-9322, 2015. <https://doi.org/10.1002/2015jd023304>

Philippe, M., Tison, J. L., Fjøsne, K., Hubbard, B., Kjær, H. A., Lenaerts, J. T. M., Drews, R., Sheldon, S. G., De Bondt, K., Claeys, P., and Pattyn, F.: Ice core evidence for a 20th century increase in surface mass balance in coastal Dronning Maud Land, East Antarctica, *The Cryosphere*, 10, 2501-2516, 2016. <https://doi.org/10.5194/tc-10-2501-2016>

Pratap, B., Dey, R., Matsuoka, K., Moholdt, G., Lindbäck, K., Goel, V., Laluraj, C. M., and Thamban, M.: Three-decade spatial patterns in surface mass balance of the Nivlisen Ice Shelf, central Dronning Maud Land, East Antarctica, *Journal of Glaciology*, 68, 174-186, 2022. <https://doi.org/10.1017/jog.2021.93>

Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095-1103, 2019. <https://doi.org/10.1073/pnas.1812883116>

Sallée, J.-B., Shuckburgh, E., Bruneau, N., Meijers, A. J. S., Bracegirdle, T. J., Wang, Z., and Roy, T.: Assessment of Southern Ocean water mass circulation and characteristics in CMIP5 models: Historical bias and forcing response, *Journal of Geophysical Research: Oceans*, 118, 1830-1844, 2013. <https://doi.org/10.1002/jgrc.20135>

The Firn Symposium team: Firn on ice sheets, *Nature Reviews Earth & Environment*, 5, 79-99, 2024. <https://doi.org/10.1038/s43017-023-00507-9>

- 769 The SO-CHIC consortium: Southern ocean carbon and heat impact on climate, *Philosophical*
770 *Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*,
771 381, 20220056, 2023. <https://doi.org/10.1098/rsta.2022.0056>
- 772 Visbeck, M., Marshall, J., and Jones, H.: Dynamics of Isolated Convective Regions in the
773 Ocean, *Journal of Physical Oceanography*, 26, 1721-1734, 1996.
774 [https://doi.org/10.1175/1520-0485\(1996\)026<1721:DOICRI>2.0.CO;2](https://doi.org/10.1175/1520-0485(1996)026<1721:DOICRI>2.0.CO;2)
- 775 Wauthy, S., Tison, J. L., Inoue, M., El Amri, S., Sun, S., Fripiat, F., Claeys, P., and Pattyn, F.:
776 Spatial and temporal variability of environmental proxies from the top 120 m of two ice
777 cores in Dronning Maud Land (East Antarctica), *Earth Syst. Sci. Data*, 16, 35-58, 2024.
778 <https://doi.org/10.5194/essd-16-35-2024>
- 779 Wilson, E. A., Riser, S. C., Campbell, E. C., and Wong, A. P. S.: Winter Upper-Ocean Stability
780 and Ice–Ocean Feedbacks in the Sea Ice–Covered Southern Ocean, *Journal of Physical*
781 *Oceanography*, 49, 1099-1117, 2019. <https://doi.org/10.1175/JPO-D-18-0184.1>
- 782 Winstrup, M., Svensson, A. M., Rasmussen, S. O., Winther, O., Steig, E. J., and Axelrod, A.
783 E.: An automated approach for annual layer counting in ice cores, *Clim. Past*, 8, 1881-
784 1895, 2012. <https://doi.org/10.5194/cp-8-1881-2012>
- 785 Xu, Y., Zhang, W., Maksym, T., Ji, R., and Li, Y.: Stratification Breakdown in Antarctic
786 Coastal Polynyas. Part I: Influence of Physical Factors on the Destratification Time Scale,
787 *Journal of Physical Oceanography*, 53, 2047-2067, 2023. [https://doi.org/10.1175/JPO-D-](https://doi.org/10.1175/JPO-D-22-0218.1)
788 22-0218.1
- 789 Zheng, W. E. I., Zhaoru, Z., Timo, V., Xiaoqiao, W., and Yuanjie, C.: An overview of Antarctic
790 polynyas: sea ice production, forcing mechanisms, temporal variability and water mass
791 formation, *Advances in Polar Science*, 32, 295-311, 2021.
792 <https://doi.org/10.13679/j.advps.2021.0026>
- 793 Zwally, H. J., Comiso, J. C., and Gordon, A. L.: Antarctic Offshore Leads and Polynyas and
794 Oceanographic Effects. In: *Oceanology of the Antarctic Continental Shelf*, 1985.
795 <https://doi.org/10.1029/AR043p0203>