



Improved constraints on ammonia emissions and deposition from co-assimilating NH₃ and NO₂ satellite observations over the Netherlands

Tyler Wizenberg¹, Enrico Dammers^{1,2}, Arjo Segers¹, Mark W. Shephard⁴, Pierre Coheur⁵, Lieven Clarisse⁵, Martin Van Damme^{5,6}, Henk Eskes⁷, Roy Wichink Kruit⁸, Shelley van der Graaf⁸, and Martijn Schaap^{1,3}

¹Air Quality & Emissions Research, TNO, Utrecht, the Netherlands

²Institute of Environmental Sciences, Universiteit Leiden, Leiden, the Netherlands

³Freie Universiteit Berlin (FUB), Berlin, Germany

⁴Environment & Climate Change Canada (ECCC), Toronto, Canada

⁵Université libre de Bruxelles (ULB), BLU-ULB research Center, Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing (SQUARES), Brussels, Belgium

⁶Royal Belgian Institute for Space Aeronomy (BIRA-IASB), Brussels, Belgium

⁷Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

⁸National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

Correspondence: Tyler Wizenberg (tyler.wizenberg@tno.nl)

Abstract. Ammonia (NH₃) and nitrogen dioxide (NO₂) are key components of reactive nitrogen, strongly affecting air quality and ecosystem health. However, long-term constraints on ammonia emissions and deposition remain uncertain due to sparse in situ measurements and limitations of individual satellite products. We jointly assimilate five years (2018–2022) of NH₃ and NO₂ satellite observations over the Netherlands to improve constraints on reactive nitrogen concentrations, emissions, and deposition. NH₃ retrievals from the Infrared Atmospheric Sounding Interferometer (IASI) and the Cross-Track Infrared Sounder (CrIS) are combined with NO₂ observations from the TROPOspheric Monitoring Instrument (TROPOMI) within the LOTOS-EUROS chemical transport model using a Local Ensemble Transform Kalman Filter. The co-assimilation produces coherent year-to-year adjustments in modeled NH₃ concentration, emission, and deposition fields. Validation against measurements from the Dutch National Air Quality Monitoring Network (LML) shows reduced biases, clearer diurnal cycles, and improved correlations. Sensitivity experiments demonstrate that including TROPOMI NO₂ alongside IASI and CrIS NH₃ yields the lowest NH₃ surface bias versus LML, highlighting the added value of coupling chemically related satellite observations. Comparisons with monthly Measurements of Ammonia in Nature (MAN) observations showed improved correlations but persistent spatial biases due to representativeness differences, while MAN sensors co-located with LML stations exhibited consistent improvements. These results demonstrate that co-assimilating complementary satellite observations can substantially improve constraints on ammonia emissions and deposition, with direct relevance for air-quality assessment and nitrogen policy applications.



1 Introduction

Ammonia (NH_3) and nitrogen dioxide (NO_2) are key reactive nitrogen species that play central roles in atmospheric chemistry and air quality. Through dry and wet deposition, these gases contribute substantially to the transfer of reactive nitrogen to the Earth's surface, where they influence both terrestrial and aquatic ecosystems. Although reactive nitrogen is essential for plant growth and ecosystem functioning, excessive deposition can cause environmental degradation, including soil acidification, eutrophication, and biodiversity loss (Galloway et al., 2008; Sutton et al., 2011).

NH_3 is the most abundant alkaline gas in the atmosphere, primarily emitted from agricultural activities such as livestock farming and fertilizer application (Erisman et al., 2011; Galloway et al., 2008). In agriculture, NH_3 is released through the decomposition of animal waste and the volatilization from nitrogen-rich fertilizers applied to soils (Erisman et al., 2008, 2011; Sutton et al., 2011; Paulot et al., 2014). This gas plays a vital role in the formation of secondary inorganic aerosols (SIAs) by reacting with acidic compounds such as sulfuric and nitric acids to form ammonium sulfate and ammonium nitrate (Erisman et al., 2011). SIAs contribute to fine particulate matter ($\text{PM}_{2.5}$), which affects atmospheric visibility and poses significant health risks, including respiratory and cardiovascular diseases (Pope and Dockery, 2006; Pope et al., 2009).

NO_2 in the atmosphere originates primarily from nitric oxide (NO), which is emitted during fossil fuel combustion in vehicles, power plants, and industrial processes and rapidly oxidized in air. Together, NO and NO_2 form the NO_x family, a key precursor to tropospheric ozone and nitrate aerosols (Crutzen, 1979; Ojeda-Castillo et al., 2025; Zara et al., 2021). NO_2 participates in photochemical reactions leading to the formation of ozone and particulate nitrates, contributing to air pollution and smog formation (Zha, 2011). In addition, NO_2 indirectly influences the atmospheric lifetime of NH_3 by modulating the availability of nitric acid (HNO_3): increased oxidation of NO_2 consumes HNO_3 , leaving less available to form ammonium nitrate from NH_3 . Elevated levels of NO_2 and its reaction products are associated with respiratory problems such as asthma and decreased lung function, and contribute to environmental degradation through acid deposition and nutrient imbalances in ecosystems (Eum et al., 2022; Jonson et al., 2017; Liu et al., 2021).

Understanding the spatial and temporal distributions of NH_3 and NO_2 is essential for accurate atmospheric modeling and the development of effective emissions control strategies and regulations. However, the reactive nature and short atmospheric lifetimes of NH_3 and NO_2 pose significant challenges to monitoring and modeling of these species. Emissions inventories derived from bottom-up approaches often have large uncertainties associated with them, particularly in the case of NH_3 where these uncertainties can be greater than 100% for some emissions sectors (Kuenen et al., 2022). Ground-based measurement networks provide high-accuracy observations but differ greatly in spatial density; NO_2 is routinely monitored at hundreds of sites across Europe, whereas NH_3 is measured at far fewer locations, resulting in much sparser spatial coverage and limited temporal resolution. Moreover, these networks generally measure surface concentrations and do not provide full atmospheric column information. Space-based observations from instruments such as IASI on the MetOp series, CrIS on the Suomi-NPP and NOAA-20/21 platforms, and TROPOMI on Sentinel-5 Precursor offer broader spatial coverage, but with limited temporal sampling due to discrete overpass times. Satellite measurements of this type can be subject to errors resulting from, for example, instrumental noise, retrieval biases, and residual cloud cover.



Inverse modeling methods present a powerful approach to overcome these limitations by integrating observational data into chemical transport models to yield optimized estimates of atmospheric constituents. By assimilating large numbers of NH_3 and NO_2 measurements, it is possible to improve the accuracy of simulated concentrations, diurnal cycles, and associated emission and deposition fields. Assimilating NO_2 observations in particular helps to better constrain the NO_x budget, which in turn reduces uncertainties in the formation and loss of nitric acid (HNO_3) and thus indirectly constrains the NH_3 fields through their chemical coupling. Through such an approach, air quality predictions can be improved and policy-relevant insights can be gained for mitigating the impacts of reactive nitrogen on ecosystems and human health.

In this paper, we perform a co-assimilation of measurements of NH_3 from IASI, CrIS and NO_2 from TROPOMI in the LOTOS-EUROS local ensemble transform Kalman filter (LETKF) over a model domain encompassing the Netherlands. We evaluate resulting optimized emissions and deposition fields, with a focus on NH_3 , and we compare the results against independent observations from ground-based measurements networks.

2 Methodology

2.1 The LOTOS-EUROS model

In this study, we utilized the LOTOS-EUROS (LONg Term Ozone Simulation-EUROpean Operational Smog) v2.2.009 chemical transport model to simulate atmospheric concentrations over the study region (Manders et al., 2017). LOTOS-EUROS is a three-dimensional Eulerian model designed for regional air quality assessments and operational forecasting in Europe. It effectively simulates the dispersion, chemical transformation, and deposition of atmospheric pollutants, including gases and aerosols. LOTOS-EUROS is part of the Copernicus Atmospheric Monitoring Service (CAMS) European air quality ensemble (Colette et al., 2025). This service provides forecasts for the main air pollutants using an ensemble of state-of-the-science CTMs. Within CAMS, LOTOS-EUROS is regularly validated against in-situ observations and TROPOMI satellite data, as well as evaluation against the other ensemble members (Peuch et al., 2022; Colette et al., 2025). LOTOS-EUROS also has participated in numerous model inter-comparisons, typically showing a strong performance (Bessagnet et al., 2016; Colette et al., 2017; Vivanco et al., 2017).

The model incorporates detailed representations of atmospheric processes such as advection, diffusion, bi-directional fluxes, and chemical reactions. It uses the Carbon Bond Meism IV (CBM-IV) for gas-phase chemistry, which includes a comprehensive set of reactions relevant to ozone formation and other photochemical oxidants. Aerosol dynamics are modeled using size-resolved modules that account for primary emissions, secondary formation, and processes like coagulation and deposition.

Model outputs, including concentrations of key pollutants such as NH_3 , NO_2 , ozone (O_3), and particulate matter (PM_{10} and $\text{PM}_{2.5}$), are regularly validated against observational data from the Dutch LML air quality monitoring network, measurements from the German environmental agency (the Umweltbundesamt; UBA) and the EBAS network throughout the European Union. The NH_3 , NO_2 and particulate matter components from the model are also frequently validated against the EMEP model and other models within the CAMS model ensemble (Schaap et al., 2008; Manders et al., 2017; Tsyro et al., 2022).



2.1.1 The Local ensemble transform Kalman filter

The Ensemble Kalman filter (Evensen, 2003) is a sequential data assimilation method that combines observations (in this case, from satellite instruments) with a CTM to optimize the concentration and emission fields of the underlying model. The Local Ensemble Transform Kalman Filter (LETKF) integrates model simulations with observations to optimize the state estimation of the system. The LOTOS-EUROS LETKF v3.0.7 was used in this study, and it has previously been applied to studies of particulate matter (Lopez-Restrepo et al., 2020), NO₂ (Timmermans et al., 2019), and NH₃ (Van Der Graaf et al., 2022), and is based on the formulation described in Hunt et al. (2007) and follows the implementation described in Shin et al. (2016). In comparison to the standard Ensemble Kalman Filter, the LETKF is more computationally efficient and performs the analysis on a per grid-cell basis using only nearby observations, which are determined using a specified spatial correlation length (discussed in further detail in Section 2.1.2. The temporal variability in the emissions are specified in the assimilation by the temporal correlation length τ (set to 3 days for NH₃ based on Van Der Graaf et al. (2022), and 1 day for NO₂) with the temporal correlation coefficient α that is defined as (Lopez-Restrepo et al., 2020; Van Der Graaf et al., 2022):

$$\alpha_k = e^{-|t_k - t_{k-1}|/\tau} \quad (1)$$

where t_k and t_{k-1} are time-points k and $k-1$. This formulation ensures that, in the absence of new observations for an extended period, the influence of past updates diminishes and the system progressively returns toward the a priori emission state.

The state vector \mathbf{x} of the LETKF includes three-dimensional trace-gas concentrations and two-dimensional perturbation factors (β) for the emissions. These are represented as an ensemble of N members, which capture the uncertainties in both the model and the observations. In this study, $N = 12$ was used as this was found by Van Der Graaf et al. (2022) to be a sufficient trade-off between an appropriate statistical representation of the model uncertainty in the system and overall computational efficiency. For the initialization of the ensemble, the perturbation factors are sampled from a normal distribution with a mean of zero, and a standard deviation of one. The values of the mean and standard deviation for the ensemble can be adjusted if desired.

The LETKF operates in two sequential steps, the forecast and the analysis. In the forecast step, the state vector ensemble is propagated forward in time using the model dynamics (Van Der Graaf et al., 2022):

$$\mathbf{X}^f = \{\mathbf{x}_1^f, \mathbf{x}_2^f, \dots, \mathbf{x}_N^f\},$$

where \mathbf{X}^f is the forecast ensemble and \mathbf{x}_i^f is the i -th forecast ensemble member. Each analyzed ensemble member from time t_{k-1} is propagated forward to t_k by the following:

$$\mathbf{x}_i^f(k) = M_{k-1}(\mathbf{x}_i^a(k-1)), \quad (2)$$

where the model operator M_{k-1} describes the forward model simulation, and also includes the application of the emissions perturbation factors, and $\mathbf{x}_i^a(k-1)$ denotes the i -th analyzed ensemble member from time t_{k-1} . Once the ensemble has been



propagated forwards in time, the ensemble mean $\bar{\mathbf{x}}$ and forecast error covariance \mathbf{P} are then calculated as

$$\bar{\mathbf{x}} = \frac{1}{N} \sum_{i=1}^N \mathbf{x}_i^f, \quad (3)$$

115

$$\mathbf{P} = \frac{1}{N-1} \sum_{i=1}^N (\mathbf{x}_i^f - \bar{\mathbf{x}})(\mathbf{x}_i^f - \bar{\mathbf{x}})^T, \quad (4)$$

When new observations \mathbf{y}^{obs} are available, the analysis ensemble \mathbf{X}^a is obtained by updating each ensemble member according to (Van Der Graaf et al., 2022):

$$\mathbf{x}_i^a = \mathbf{x}_i^f + \mathbf{P}^a \mathbf{H}^T \mathbf{R}^{-1} (\mathbf{y}^{\text{obs}} - \mathbf{h}(\mathbf{x}_i^f)), \quad (5)$$

120 where $\mathbf{h}(\mathbf{x}_i^f)$ denotes the model-simulated equivalent of the satellite retrieval, \mathbf{H} is the linearized observation operator, and \mathbf{R} is the observation error covariance matrix.

The simulated satellite observations are computed following the averaging-kernel formalism of Rodgers (2000) and Rodgers and Connor (2003). First, the model state is interpolated to the retrieval grid using the gridding operator \mathbf{G} :

$$\mathbf{x}_{r,i} = \mathbf{G} \mathbf{x}_i^f. \quad (6)$$

125 To ensure that the comparison with the satellite product is made at the same effective vertical resolution as the retrieval, the averaging kernel \mathbf{A} is applied:

$$\mathbf{h}(\mathbf{x}_i^f) = \mathbf{x}_a + \mathbf{A}(\mathbf{x}_{r,i} - \mathbf{x}_a), \quad (7)$$

where \mathbf{x}_a is the a priori profile used in the retrieval. In the linear limit, the observation operator becomes:

$$\mathbf{H} = \mathbf{A} \mathbf{G}. \quad (8)$$

130 The observations are assumed to satisfy

$$\mathbf{y}^{\text{obs}} = \mathbf{h}(\mathbf{x}^{\text{true}}) + \mathbf{v}, \quad \mathbf{v} \sim \mathcal{N}(0, \mathbf{R}), \quad (9)$$

with \mathbf{R} taken from the retrieval error covariance matrices of the satellite data products and representing both measurement and representativeness errors (Van Der Graaf et al., 2022).

The analysis error covariance used in Eq. 5 is computed from

$$135 \mathbf{P}^a = [\mathbf{P} \mathbf{H}^T \mathbf{R}^{-1} \mathbf{H} + \mathbf{I}]^{-1} \mathbf{P}. \quad (10)$$

Once the update is complete, the analyzed ensemble \mathbf{X}^a becomes the initial condition for the next forecast step.



2.1.2 Spatial localization

To ensure computational efficiency and avoid spurious correlations, the LETKF applies spatial and temporal localization in a per grid cell approach following the method described by Shin et al. (2016). The temporal localization was described in the previous section, and is applied using Equation 1. For the spatial localization, the simulated observations are first computed for all ensemble members, and then for the given grid cell to be analyzed, all observations (both simulated and real) within 3.5 ρ distance are gathered, with ρ being the specified correlation length. The analysis is then performed using the observations collected, with the weight of the additional observations being limited by a Gaussian decay function: (Shin et al., 2016):

$$w(\Delta d) = \exp\left(-\frac{\Delta d^2}{2\rho^2}\right),$$

where Δd is the distance between the observation and the model grid point. In this study, the spatial correlation lengths are chosen based on the mean footprint sizes of the corresponding satellite instruments. This choice reflects the fact that each retrieval represents an average over its footprint and is treated as an independent observation. Using a localization length much larger than the footprint would allow a single observation to influence the analysis over spatial scales that are not resolved by the measurement, potentially producing spurious long-range increments and unrealistically smooth updates. To ensure spatial consistency between the retrieval resolution and its influence in the LETKF, we use $\rho = 15$ km for CrIS and IASI NH₃ and a smaller $\rho = 5$ km for TROPOMI NO₂, reflecting the higher spatial resolution of the latter.

2.1.3 Model configuration

In this section, a short summary of the most important model inputs and configuration parameters are provided. For a more detailed description of the LOTOS-EUROS model we direct the reader to Manders et al. (2017). The LETKF v3.0.7 is coupled to version 2.2.009 of the LOTOS-EUROS model. An initial long simulation covering the period of 2018–2022 was performed on a domain that covers the majority of Europe (15°W–35°E; 35–70°N) with a resolution of approximately 25 × 25 km², and the output from this run was utilized as boundary conditions for the assimilation run which was performed on a domain covering most of North-western Europe (2–16°E; 47–56°N) with a resolution of 7 × 7 km². The model is driven by meteorological fields obtained from the ECMWF short-term forecast model at a 3-hourly temporal resolution which is then interpolated to an hourly frequency within the model. The simulations were conducted using 12 vertical levels, extending from the ground to about 10 km above the earth's surface, matching the vertical layer structure of the ECMWF meteorology dataset.

The base emission dataset used in the model simulations combines the European-scale CAMS-REG-v5.1 inventory (Kuenen et al., 2022) with higher-resolution national inventories for the Netherlands (Emissieregistratie; ER) and Germany (Gridding Emission Tool for ArcGIS; GRETA). This combined CAMS GrETa-ER emissions dataset was developed within the National Kennisprogramma Stikstof (NKS) funded by the Dutch Ministry of Agriculture, Fisheries, Food Security and Nature (LVVN). For years after 2019, the 2019 emission totals are used as a baseline but are adjusted dynamically according to meteorological conditions. As a result, year-to-year variations in the base emissions remain relatively small and primarily reflect meteorological influences rather than structural changes in activity or policy. Temporally, the emissions are distributed using hourly time factors



specific to aggregated source categories. For agricultural NH_3 emissions, a meteorologically dependent parameterization is applied that accounts for weather-driven shifts in fertilizer application timing, following the approach of Ge et al. (2020). Emissions are also vertically distributed according to sector-specific release heights, which is particularly relevant for industrial and power-generation sources where average stack heights determine the initial plume elevation.

The model output for all major nitrogen species consists of simulated concentrations (surface, and all vertical layers) and wet and dry deposition, as well as concentrations matched at the footprints of all satellite products (i.e. IASI, CrIS, TROPOMI). To match the model with the satellite footprints, and for the ingestion of these observations in the LETKF, the CAMS Satellite Operator (CSO) is used. CSO (<https://ci.tno.nl/gitlab/cams/cso>) is an open-access tool developed at TNO and implemented to facilitate fast intercomparisons between modelled and satellite concentrations. The tool consists of two entities: a pre-processor to download, select, and convert satellite observations into a common format, accompanied with a post-processing tools to aggregate and visualize the data; and a source code that can be used within regional air quality modelling and assimilation systems such as LOTOS-EUROS. The CSO module is able to read the files created by the pre-processor, simulate satellite observations using model variables, and apply observational operators where applicable.

2.2 Satellite datasets

2.2.1 IASI NH_3

The IASI instruments onboard the MetOp-A, -B, and -C satellites are in Sun-synchronous orbits, passing locations twice daily with Equator crossing times at 09:30 and 21:30 local time, and with a time difference of approximately 45 minutes between them (Clerbaux et al., 2009). The data products of IASI-A, -B, and -C span the periods October 2007–October 2021 (IASI-A), March 2013 onward (IASI-B), and September 2019 onward (IASI-C), with the latter two instruments still operational. Each of the three IASI instruments has an observational swath width exceeding 2000 km, with a pixel footprint of approximately 12 km in diameter at nadir, increasing to around $\sim 20 \times 40 \text{ km}^2$ at the swath edges. In this study, we utilize the most recent IASI product, the Artificial Neural Network for IASI (ANNI)v4 (Clarisse et al., 2023), which is an updated version of the earlier IASI-NNv2 and LUT products (Van Damme et al., 2014, 2017).

Similar to its predecessors, the IASI-ANNIv4 retrieval involves two steps. First, the Hyperspectral Range Index (HRI) is calculated to characterize the NH_3 signal strength in each spectrum. The second step utilizes a neural network trained on a large dataset of modeled data, which links the HRI to the NH_3 total column. The primary improvement in this version is the addition of a column averaging kernel to facilitate comparisons with in-situ and model data, enabling the effects of the a-priori profile shape to be considered. Earlier ANNI NH_3 products were evaluated in previous studies by Dammers et al. (2016); Guo et al. (2021); Kutzner et al. (2021); Herrera et al. (2022), including against ground-based FTIR observations from the Network for the Detection of Atmospheric Composition ge (NDACC). The validation of IASI NH_3 against NDACC measurements by Dammers et al. (2016) showed on average favorable correlations ($R = 0.80$), with a mean low bias on the order of 35%.

In the current study, we apply the recommended data quality filters, namely, with the pre-filter and post-filter set to 1, and only observations with a cloud_fraction $< 25\%$ are used.



2.2.2 CrIS NH₃

The CrIS-1 instrument on the Suomi-NPP and CrIS-2 on NOAA-20 were launched in October 2011 and November 2017, respectively, with CrIS-Fast Physical Retrieval (CFPR) NH₃ data products starting in May 2012 and March 2019. Both instruments are in sun-synchronous orbits, providing global coverage twice daily at around 13:30 and 1:30 local solar time, with overpasses within 45 minutes of each other. They offer observations with circular pixel footprints of approximately 14 km at nadir over a 2200 km wide swath. This study utilizes the CFPR NH₃ product version 1.6.4 (Shephard and Cady-Pereira, 2015; Shephard et al., 2020). The CFPR method involves a physical retrieval based on the Rodgers (2000) optimal estimation method combined with a fast optimal spectral sampling forward model (Moncet et al., 2008), minimizing the residual between measured and simulated spectra. The excellent signal-to-noise ratio (~ 1600) of the CrIS instrument in the ammonia spectral region enables detection sensitivities of approximately 0.5 ppbv near the surface, or 3.5×10^{15} molec cm⁻² for total columns, under typical atmospheric conditions ($\sim 50\%$ detection rate). Under highly favorable infrared remote sensing conditions (e.g., strong thermal contrast) the detection limit can improve to about 0.2 ppbv, corresponding to a $\sim 10\%$ detection rate (Shephard et al., 2025). Since the last major validation study by Dammers et al. (2017), the product has undergone several iterations, including the addition of a cloud flag based on VIIRS data, non-detects, and a quality flag (White et al., 2023). The validation study by Dammers et al. (2017) reported a good correlation between FTIR and satellite observations ($R \approx 0.8$) with a slight high bias (slope = 1.02). For higher column concentrations, CrIS observations showed a small positive difference with the ground-based FTIR measurements around 25 – 50%, while for lower concentrations, the bias increased to 2.5×10^{15} molecules cm⁻² with a standard deviation of around 50-100%. This study only includes observations with a quality_flag of ≥ 3 and with a cloud_flag equal to 0 (clear-sky observations). Only daytime observations are used in the analysis. It should be noted that the CrIS-1 instrument suffered a failure of its mid-wave IR (MWIR) band from 26 March to 24 June 2019, leading to a data gap in this period (Iturbide-Sanchez et al., 2022).

2.2.3 TROPOMI NO₂

The TROPOMI instrument, aboard the Sentinel-5 Precursor (S-5P) polar-orbiting satellite, is a nadir-viewing spectrometer designed for atmospheric observations. It crosses the equator at approximately 13:30 local time. The instrument measures radiation across the ultraviolet, visible, and infrared spectral ranges, enabling the monitoring of atmospheric trace gases and aerosols (Veefkind et al., 2012). Retrieval of NO₂ columns follows a three-step process. First, the NO₂ slant column density is calculated from the L1b spectra recorded by TROPOMI using a DOAS fitting algorithm. This slant column is then separated into stratospheric and tropospheric components through data assimilation using the TM5-MP model, which operates at a horizontal resolution of $1^\circ \times 1^\circ$ (Williams et al., 2017). Finally, slant column densities are converted to vertical column densities (VCD) by applying total and altitude-dependent air mass factors (AMFs). These AMFs are influenced by several factors, including NO₂ vertical profiles obtained from TM5-MP, the satellite's viewing geometry, surface albedo, surface pressure, and characteristics of clouds and aerosols. Further details on the retrieval process can be found in van Geffen et al. (2022) and in the algorithm theoretical baseline document (van Geffen et al., 2024).



235 Routine validation of TROPOMI NO₂ measurements against ground-based MAX-DOAS observations from 29 stations has revealed a mean bias of -28%, increasing to -40% in regions with heavy pollution (Lambert and et al., 2024). This bias largely stems from the TM5-MP vertical profiles, which inadequately resolve high-concentration hot-spots and show deviations in the profile shape, particularly near the surface (Chan et al., 2020; Verhoelst et al., 2021). To address these discrepancies, the a-priori vertical profile can be updated using one derived from a higher-resolution air quality model, which has been shown to partially mitigate biases (Griffin et al., 2019; Zhao et al., 2020; Judd et al., 2020; Douros et al., 2023). This correction process employs TROPOMI averaging kernels and is described in detail in the TROPOMI NO₂ Product User Manual (Eskes et al., 2024).

In this study, we used the reprocessed TROPOMI NO₂ version 2.4.0 dataset. To ensure data reliability, observations with a quality assurance value below 0.75 were excluded. This threshold effectively eliminates pixels with cloud radiance fractions exceeding 0.5, thereby reducing the impact of uncertain retrievals (van Geffen et al., 2022).

2.3 Ground-based in-situ measurements

2.3.1 The LML network

The Dutch National Air Quality Monitoring Network, known as the "Landelijk Meetnet Luchtkwaliteit" (LML) (Elskamp, 1989; Elzakker and Buijsman, 1999), is a comprehensive ground-based measurement network designed to monitor air quality across the Netherlands. Operated by Rijksinstituut voor Volksgezondheid en Milieu (RIVM), the LML network consists of a large number of monitoring stations distributed across urban, suburban, and rural areas. These stations continuously collect data on various air pollutants, including PM₁₀, PM_{2.5}, NO₂, NH₃, O₃, sulfur dioxide (SO₂), carbon monoxide (CO), and volatile organic compounds (VOCs).

The network provides real-time data, which is crucial for assessing the air quality in different regions and understanding the impact of pollution on public health and the environment. LML stations employ state-of-the-art sensors and analytical techniques to ensure high data accuracy and consistency, enabling authorities to monitor trends, detect exceedances of air quality standards, and develop policy interventions when necessary. NH₃ measurements are made using the miniDOAS, an active instrument that utilizes the differential optical absorption spectroscopy (DOAS) measurement technique. These instruments have an NH₃ detection limit of roughly 0.25 μg m⁻³ and an estimated precision of 0.1 μg m⁻³ for hourly averaged observations (Berkhout et al., 2017) however, no measurement uncertainties are provided in the dataset.

Data from the LML are publicly accessible (Luchtmeetnet.nl), allowing citizens, researchers, and policymakers to track air quality levels in near real-time. This transparency helps raise awareness about air pollution issues and supports efforts toward improving air quality across the country. The LML network is also integrated with broader European air quality initiatives, contributing to the wider understanding of transboundary pollution and climate change mitigation efforts. The LML NH₃ measurements were previously applied to study long-term trends in the Netherlands by van Zanten et al. (2017).

Ten LML sites were selected in total for the comparisons with the model simulations, and the coordinates, details, and species measured for each of these sites are provided in Table 1. Six of the chosen LML sites provide hourly NH₃ surface con-



Site Name	Coordinates	Species Measured	Type
Biest Houtakker-Biestsestraat	51.15° N, 5.15° E	NH ₄ ⁺ (wet)	Rural/Agricultural
De Bilt-Wilheminalaan	52.10° N, 5.17° E	NH ₄ ⁺ (wet)	Urban
De Zilk-Vogelaarsdreef	52.30° N, 4.51° E	NH ₃ , NH ₄ ⁺ (wet)	Coastal
Philippine-Stelleweg	51.29° N, 3.7° E	NH ₄ ⁺ (wet)	Rural/Agricultural
Speuld-Garderenseweg	52.27° N, 3.7° E	NH ₄ ⁺ (wet)	Rural/forested area
Valthermond-Noorderdiep	52.88° N, 5.72° E	NH ₃ , NH ₄ ⁺ (wet)	Rural/Agricultural
Vredepeel-Vredeweg	51.54° N, 5.85° E	NH ₃ , NH ₄ ⁺ (wet)	Rural/Agricultural
Wekerom-Riemterdijk	52.11° N, 5.71° E	NH ₃	Rural/Agricultural
Wieringerwerf-Medeblikkerweg	52.80° N, 5.05° E	NH ₃ , NH ₄ ⁺ (wet)	Coastal/Agricultural
Zegveld-Oude Meije	52.14° N, 4.84° E	NH ₃	Rural/Agricultural

Table 1. Locations and details of LML sites used for comparisons with the LOTOS-EUROS LETKF simulations.

centration measurements; De Zilk-Vogelaarsdreef, Valthermond-Noorderdiep, Vredepeel-Vredeweg, Wekerom-Riemterdijk, Wieringerwerf-Medeblikkerweg, and Zegveld-Oude Meije. These sites were selected because they provided hourly NH₃ measurements over a long period (>1 year). In addition to the NH₃ measurements, wet deposition measurements of dissolved ammonium (NH₄⁺) concentrations in precipitation are made periodically (i.e., at irregular intervals) at Biest Houtakker-Biestsestraat, De Bilt-Wilheminalaan, Philippine-Stelleweg, Speuld-Garderenseweg, Valthermond-Noorderdiep, Vredepeel-Vredeweg, and Wieringerwerf-Medeblikkerweg. Wet deposition is also measured at the De Zilk-Vogelaarsdreef site, which serves as a European Monitoring and Evaluation Programme (EMEP) location where dissolved ammonium is monitored on a daily basis. The observed NH₄⁺ measurements were paired with the model output and converted to monthly mean fluxes using the corresponding measured and modeled precipitation amounts. The modeled and observed precipitation agreed well on average, though some transient mismatches occurred. To minimize their influence on the wet deposition comparisons, a strict filter was applied: cases where the measured and modeled precipitation exceeded a mean absolute deviation of 1σ were excluded, as were measurements with very low precipitation (<0.1 mm). The locations of the sites within the Netherlands are shown on a map in Figure 1.

The NH₃ measurement sites occasionally experience local-scale enhancement events (i.e., nearby manure dumping events), which are not spatially representative of the model resolution. For 2020 onward, official flags are provided in the LML dataset to enable the filtering of such events, but for the years prior to this no such information is available. As a result, additional filtering was manually applied, and any hourly NH₃ observations with concentrations exceeding 100 μg m⁻³ are removed and excluded from the comparisons.




Figure 1. Map of the locations of the selected LML sites in the Netherlands used for comparisons with the model simulations. Underlying basemap data sourced from © OpenStreetMap contributors (2017).

2.3.2 The MAN network

The "Measuring Ammonia in Nature" (MAN) network was established in 2005 to monitor atmospheric ammonia concentrations in nature reserve areas across the Netherlands, with a particular focus on nitrogen-sensitive Natura2000 areas (Lolkema et al., 2015). The network provides essential data for assessing national ammonia concentration trends, validating air quality models, and analyzing regional variability.

The network employs commercial Gradko passive ammonia samplers, which are cost-effective, easy to deploy, and well-suited for large-scale monitoring (Lolkema et al., 2015; Noordijk et al., 2020). These samplers are calibrated monthly using active ammonia sampling devices within the LML network (Lolkema et al., 2015). Local volunteers, often conservation wardens, handle the monthly exchange of samplers, ensuring consistent data collection across diverse habitats.

Currently, the MAN network includes over 300 sampling sites across the Netherlands. The ammonia concentration data gathered by the network facilitates the identification of spatial concentration patterns and regional anomalies throughout the country. Note that in comparison to LML, the MAN samplers are typically placed in nature areas and away from source regions.



The MAN network's data plays a critical role in assessing the effectiveness of environmental policies aimed at reducing nitrogen emissions and in analyzing long-term trends in nitrogen deposition. The network can detect annual trends as low as 300 3% over extended time series, making it a valuable tool for air quality management and biodiversity conservation in nitrogen-sensitive areas (Lolkema et al., 2015).

To reduce meteorological influences on the passive samplers, the MAN network is calibrated against high-performance reference instruments for NH_3 at six locations in the Netherlands (see Berkhout et al. (2017)). The calibration process is detailed in Noordijk et al. (2020). The uncertainty in the MAN measurements consists of two components: the measurement 305 uncertainty and the calibration uncertainty. The random uncertainty is approximately $0.9 \mu\text{g m}^{-3}$ for a single monthly value and $0.32 \mu\text{g m}^{-3}$ for annual averages. The systematic uncertainty is estimated at 28% for monthly values and 10% for annual averages (Noordijk et al., 2020).

For comparison with the LETKF simulations, all available MAN data from the period 2018–2022 was used, comprising a total of 309 standard sites, and 6 additional MAN calibration sensors located at LML measurement sites. While not all sites 310 provide uninterrupted time series over the full period, the large number of sites and monthly measurements ensures statistically robust results.

3 Results and Discussion

3.1 Optimized emission fields

We first examine the NH_3 emission fields pre- and post-assimilation to evaluate the impact of ingesting satellite observations 315 on the model simulation. The base and optimized NH_3 yearly total emissions for each individual year and for the mean over 2018–2022 are shown in Figure 2. The relative difference plots in Figure 2(c) reveal a consistent spatial pattern: increases across much of the south and east of the Netherlands and decreases in the north and the adjacent regions of Germany. The largest mean increase occurred in 2020 (+14.3%), the smallest in 2021 (+3.0%), with a period-average change of +8.0%. It should be noted that 2020 was an exceptional year, with unusually warm and sunny summer months, which in turn led to 320 increased volatilization of NH_3 and higher emissions. This is likely one of the key factors contributing to the higher emissions change post-assimilation in that particular year.

The persistent increases in the south-eastern Netherlands mirror the patterns reported by Ge et al. (2020), who found that incorporating detailed agricultural activity data into the MACC inventory led to higher emissions in this region due to improved spatial allocation of sources, more accurate representation of manure management and application timing, and region-specific 325 regulatory constraints. They also showed that emissions here are disproportionately influenced by intensive pig and poultry farming, whereas dairy cattle dominate in much of the rest of the country. In contrast, the decreases we find in the north are consistent with their observation that refined allocation can reduce emissions where earlier inventories overestimated activity, particularly for dairy cattle. These parallels suggest that the positive adjustments in our assimilation likely reflect structural biases in the base inventory, both in the spatial distribution and the livestock-sector partitioning of emissions, rather than being 330 solely an artifact of the assimilation.




Figure 2. (a) The total NH_3 emissions by year and the mean emissions 2018–2022 period from the base CAM5 GRETA-ER inventory, (b) the same but for the optimized emissions from the LETKF analysis, and (c) the mean relative differences between the base and the optimized emission fields.




Figure 3. A time-series of monthly total NH_3 emissions calculated over the model domain shown in Figure 2 for the 2018–2022 period from the base simulation and the optimized LETKF analysis. The relative difference (optimized - base)/base in % is shown by the dashed line.

Meteorological effects can also have a significant impact on NH_3 emissions, with temperature, precipitation, and wind speed strongly influencing volatilization rates and atmospheric transport (Søgaard et al., 2002; Gyldenkerne et al., 2005; Ge et al.,



2023). Warm, dry conditions can substantially enhance emissions above climatological norms, while precipitation events may suppress them or trigger post-event peaks. Such variability may contribute to the year-to-year differences in our assimilation adjustments, and could help explain the particularly large increase in 2020, when the Netherlands experienced a very sunny spring and an unusually warm, dry summer.

In addition to the annual emission changes, it is also informative to examine the temporal evolution of emissions at finer timescales. Figure 3 presents the time series of monthly NH_3 emission totals, aggregated over the same region shown in Figure 2, providing further insight into how the assimilation influences variability across individual months. A broadly consistent pattern is found across all years, with a decrease in the springtime emission peak in the optimized simulation relative to the base simulation, and a corresponding increase in the emissions during the summer months. Particularly large decreases in the springtime peaks are seen for 2018–2020, while a minimal decrease occurs in 2021 and a slight increase is observed in 2022 between the optimized and base simulations, respectively. As seen in Figure 2, 2020 shows the most significant emissions changes, with particularly large increases in the emissions (on the order of +70%) in the LETKF-optimized simulation between April and September. In contrast, 2021 shows the smallest emissions changes, with a smaller than average increase in the emissions in the optimized simulation in May and June, but mostly insignificant changes throughout the remainder of the year.

Similar NH_3 seasonal cycles to that of the optimized simulation have been presented in recent modeling and observational studies of the region. The updated temporal emission profiles presented by Ge et al. (2020), and which are implemented in the model simulations used in this study, show shifts in the early-year emission peaks to later in the springtime, and increases in the overall summertime emissions. A long-term time-series (2008–2020) of IASI NH_3 total column measurements over the Netherlands in Van Damme et al. (2022) displays a springtime peak in April and a slightly smaller but somewhat comparable secondary peak in July to August. A similar pattern of reduced spring peaks and enhanced summertime emissions was also found by Ding et al. (2024) after assimilating CrIS NH_3 observations into the DECSO system. In addition, a recent two-year field campaign in key agricultural regions of the Netherlands by Lô et al. (2025) observed strong peaks in surface NH_3 concentrations during the summer months, driven by increased volatilization under warmer conditions. Despite the inclusion of the Ge et al. (2020) temporal profiles, the model simulations in this study still underestimate summertime emissions relative to these independent datasets, suggesting that the seasonal emission distribution in LOTOS-EUROS may require a further upward adjustment during summer. The broadly consistent pattern of change across most years in Figure 3 supports the need to redistribute a portion of emissions from early spring toward the summer months to better align the model with satellite and in-situ observations.

3.2 Impact on NH_x deposition fields

Maps of the total NH_x deposition for the base and optimized simulations are shown in Figures 4(a) and (b). The relative difference plots in Figure 4(c) displays a similar general spatial pattern as for emissions, whereby the most significant increases in the modeled deposition occur in the south and east of the Netherlands, while smaller or negative differences are found in the north of the country. The largest changes in the total deposition are seen in 2020 and 2021 with differences of +10.4% and +9.6%, respectively. Very small relative differences are seen in 2019 and 2021 of +3.8% and +2.7%, respectively. The




Figure 4. (a) The modeled total NH_x deposition (dry + wet) by year and the mean for the 2018–2022 period from the base un-optimized simulation, (b) the same but for the optimized deposition from the LETKF analysis, and (c) the mean relative differences between the base and the optimized deposition fields.

mean relative change in the NH_x deposition over the full 2018–2022 period is +6.3%. It should be noted that a large relative difference is seen over the area of the IJsselmeer and along the Dutch coastline that is caused by the dry re-emission of NH_x , which is derived from standard maps in the LOTOS-EUROS model. However, the absolute differences (in terms of kg m^{-2}) in these areas are negligible. The reduced sensitivity of deposition to the assimilation is expected, since part of the emitted NH_3 is exported offshore.

Although the assimilation exhibits consistent patterns and adjustments to the emissions, and to some extent the deposition, across all years of the simulation, validation against independent ground-based measurements is necessary to fully assess the accuracy of the assimilated fields and whether an improvement over the original state is achieved. This will be the focus of Section 3.5.

3.3 Spatial distribution of NH_3 Concentrations

In this section, we evaluate the effects of the assimilation on the spatial distribution of NH_3 concentrations. Figure 5 shows the spatial distribution of NH_3 total column concentrations over the model domain from 2018 to 2022, comparing the baseline and LETKF optimized simulations. Panels (a) on the top row present the NH_3 total columns from the base model, panels (b) on the middle row show the updated concentration fields after integrating satellite observations, and panels (c) on the bottom row display the relative differences, highlighting the assimilation’s impact on simulated NH_3 concentrations.



Across all years, both the baseline and optimized NH_3 total column fields exhibit similar large-scale spatial patterns, with elevated NH_3 levels concentrated in regions known for intensive agricultural activity, such as northwestern Germany, and the south-eastern Netherlands. However, clear differences emerge between the prior and assimilated estimates, with the assimilation generally increasing NH_3 concentrations across most regions, as shown by the predominantly positive relative differences in panels (c). The largest change in the total columns between the base and optimized simulations is seen in 2020, where a mean increase of 29.3% is found, while the years with the smallest differences were 2019 and 2021 where an increase in the columns across the domain of roughly 3.5% in both years was found. The mean differences over the 2018–2022 period indicate a systematic increase in mean NH_3 column concentrations within the model domain of approximately 10.4%, with the largest increases seen over the south eastern Netherlands, suggesting that the base model likely underestimates emissions or overestimates deposition processes there. In some areas, namely in the northern and southern western parts of the domain in 2018 and 2019, slight reductions in the NH_3 total columns are seen after assimilation, which is likely due to an overestimation in the a priori emissions, and this broadly mirrors the pattern of the emissions changes from Figure 2. For 2020 and 2022, the mean differences in NH_3 total columns are positive across the domain, whereas the corresponding emission difference maps show localized decreases, particularly in the northeastern part of the model domain. This apparent mismatch likely reflects the influence of atmospheric transport, as NH_3 emitted in one region can be advected and deposited elsewhere, so column enhancements do not necessarily coincide spatially with the emission sources. The mean of the differences for all years reveals a relatively consistent year-to-year pattern of enhancement, reinforcing the robustness of these corrections over multiple years. The systematic increases in NH_3 after assimilation suggest that satellite-derived observations are providing important constraints to correct for underestimation in the base model and emissions, particularly in regions where agricultural sources dominate. These adjustments likely reflect the assimilation compensating for structural model biases such as underestimated volatilization, overly rapid deposition, spatial misallocation in emission inventories, or the fact that the current temporal NH_3 agricultural emission profiles do not take meteorological conditions sufficiently into account (Ge et al., 2023). As discussed in Section 2.2, satellite retrievals can carry systematic biases arising from factors such as a priori assumptions, poor observational conditions (e.g., low thermal contrast), or cloud screening, which may influence the resulting analysis fields. By assimilating several satellite datasets, the impact of biases in individual datasets can be reduced to some extent, and this will be explored in further detail in Section 3.5.2. Independent evaluation against ground-based observations therefore remains important to determine whether the observed increases reflect genuine atmospheric conditions.

3.4 DOFS and Observation Density

To evaluate the impact of the number of available satellite observations on the assimilation, we can estimate the number of independent pieces of information provided by the satellite measurements through the Degrees of Freedom for Signal (DOFS) in the model domain. Here, the DOFS are calculated on the model grid using the method described in Chen et al. (2023):

$$DOFS = \mathbf{I} - \hat{\mathbf{S}}' \mathbf{S}_a^{-1} \quad (11)$$




Figure 5. (a) The mean NH_3 total column concentrations by year and for the full 2018–2022 period for the LOTOS-EUROS baseline simulation, (b) the same but for the LETKF optimized simulation, and (c) the mean relative differences between optimized and baseline simulations.

where \mathbf{I} is the identity matrix, $\hat{\mathbf{S}}'$ is the posterior error covariance matrix, and \mathbf{S}_a is the a priori error covariance matrix, both
 415 of which are directly output per time-step from the LETKF.

Figure 6(a) presents the mean spatial distribution of the DOFS per each year of the simulation, as well as the mean over
 the full 2018–2022 period. Regions with high observation coverage, particularly over areas with higher NH_3 concentrations
 (and thus, more successful retrievals with lower uncertainty), exhibit elevated DOFS values, indicating a stronger observational
 influence. Conversely, regions with sparse observations, such as areas with lower retrieval sensitivity, show lower DOFS values,
 420 implying a stronger dependence on the model prior. A consistent pattern of higher DOFS in north-western German border
 region, and throughout much of the Netherlands, in particular along the western half of the country is seen for all years.
 Additionally, a higher DOFS on average is seen for the year of 2020, where the largest overlap of the CrIS and IASI instruments
 occurred, which is reflected in Figure 6(b). The CrIS-2 NH_3 data product begins in March 2019 and CrIS-1 observations end
 mid-2021 as a result of the instrument being decommissioned, while the IASI-C NH_3 data product starts in September 2019
 425 and the IASI-A product ended in October 2021 due to the MetOp-A platform reaching the end of its operational lifetime.

To assess the relationship between DOFS and observation density, a time-series of the total number of observations in the
 model domain is shown in Figure 6(b). A positive relationship is observed, particularly for the year of 2020, demonstrating that
 a higher density of available satellite measurements generally provides more independent pieces of information to constrain
 the state in the assimilation. This impact is also seen in Figures 2 and 5, where the largest adjustments in emissions and total



430 column concentrations relative to the original simulation were observed in 2020. It is a possibility that in the years with more
 limited observations the change in the emissions post-analysis is underestimated due to the lower amount of information being
 provided by the satellites. However, as discussed earlier, 2020 was an exceptional year with very warm summer months and
 particularly high NH_3 emissions, which likely contribute to the higher DOFS there. The relationship is not strictly linear, as
 the DOFS are also influenced by the prior and posterior states and error covariance structure as well as the relative uncertainty
 435 of the satellite observations. This means that the assimilation can be limited both by the density of available retrievals and by
 their accuracy: high observational coverage with large uncertainties provides limited benefit, while accurate retrievals at low
 coverage can still meaningfully constrain the system, but are less able to capture transient emission events or resolve daily and
 diurnal cycles. In practice, both factors act together, and the results emphasize the importance of maintaining high-density,




Figure 6. (a) The mean NH_3 Degrees of Freedom for Signal (DOFS) in the model domain, and (b) a time-series of the number of NH_3 satellite observations per month within the domain. The total number of observations per month is shown by the solid black line.



accurate satellite observations—ideally from multiple instruments—to more effectively constrain NH₃ emissions and capture
440 inter-annual variability.

3.5 Comparisons with ground-based observations

3.5.1 LML hourly surface concentrations

To accurately assess the impact of the assimilation of the satellite observations, and to evaluate whether the optimized con-
centration fields represent more realistic estimates of the true spatio-temporal distribution of NH₃, it is important to compare
445 the results against an independent dataset. The surface concentrations from the base and optimized model runs were compared
against ground-based surface observations from six sites in the LML network that were described in Section 2.3.1. These sites
were selected because they provide hourly NH₃ surface concentration measurements and have significant time-series (i.e., >1
year) of data. As the primary focus of this study is NH₃, detailed discussion of the corresponding NO₂ assimilation results is
beyond the scope of the present study. These results will be presented separately in a forthcoming paper to allow for a more
450 comprehensive treatment.

A correlation plot of the monthly temporal means is shown in Figure 7, and a corresponding plot of the spatial means (i.e.,
all sites averaged for a given month) is provided in Figure 8. All linear regressions were performed using an ordinary least
squares fitting approach.

Not all LML sites provided data for all months, leading to a total of 346 data points (compared to the expected 360) in Figure
455 7. An improvement in the mean bias from $-2.0 \mu\text{g m}^{-3}$ to $-1.0 \mu\text{g m}^{-3}$ relative to the LML measurements can be seen between
the baseline and optimized simulations in Figure 7, respectively. The correlation in the temporal means also improved slightly
from $R = 0.84$ to $R = 0.85$, and the slope of the regression also improved from 0.79 to 0.91. A larger improvement in the
agreement of the spatial means is observed in Figure 8, with an increase in the Pearson correlation coefficient from $R = 0.77$
to $R = 0.84$ and an improvement in the slope of the regression from 0.79 to 0.90 from the base simulation to the optimized run,
460 respectively. This indicates that the spatial pattern of the NH₃ surface concentrations in the Netherlands is being represented
more accurately in the LETKF optimized simulation than in the original simulation without assimilation.

A Taylor diagram illustrating the measurement-model comparisons before and after assimilation at each individual LML site
is shown in Figure 9. In general, the correlations between the measured and modeled surface NH₃ concentrations improve after
assimilation at all LML sites except Wekerom-Riemterdijk, and the largest improvements are seen at Zegveld Oude-Meije and
465 Valthermond-Noorderdiep. The correlation shows a small improvement at Vredepeel-Vredeweg and this was accompanied by
a significant reduction in the bias from $-3.10 \mu\text{g m}^{-3}$ to $0.20 \mu\text{g m}^{-3}$, but a large increase in the relative standard deviation
is also seen that is driven by an apparent larger number of short-term enhancement events being captured in the modeled
time-series after assimilation (shown in Appendix Figure A1). The mean over all sites displays a small improvement in the
correlation, but also a small increase in the standard deviation (also shown earlier in Figure 7).

470 Since the LML data are provided at an hourly frequency, the impact of assimilation on the diurnal cycles of NH₃ in the model
can also be investigated at each site. The mean diurnal cycles from the observations, the baseline model run, and the optimized




Figure 7. Scatter plot of monthly temporal means of (left) LOTOS-EUROS base NH₃, and (right) LOTOS-EUROS LETKF optimized NH₃ versus LML observed NH₃ surface concentrations for the period of January 2018 to December 2022.




Figure 8. Scatter plot of monthly spatial means of (left) LOTOS-EUROS base NH₃, and (right) LOTOS-EUROS LETKF optimized NH₃ versus LML observed NH₃ surface concentrations for the period of January 2018 to December 2022. Each data-point represents the mean calculated across all LML sites for a given month, and are colored corresponding to the month while the marker style indicates the year.



run calculated over the 2018–2022 period are shown in Figure 10. In most cases, the mean differences in the diurnal cycles showed improvement relative to the observations in the analysis run in comparison to the base simulation even though only morning and afternoon satellite overpasses were used. From Figure 10 it can be seen that in the base simulation, the diurnal cycles are in many cases largely underestimated relative to the surface observations, however after the assimilation of the satellite observations a much closer agreement between the measurements and the model is found. At Wekerom-Riemterdijk and Wieringerwerf-Medemblikkerweg the mean differences increased from $-2.03 \mu\text{g m}^{-3}$ to $-2.70 \mu\text{g m}^{-3}$ and $-2.83 \mu\text{g m}^{-3}$ to $-2.92 \mu\text{g m}^{-3}$, respectively. At all other LML sites a decrease in the mean difference of the diurnal cycles was observed in the analysis relative to the base run. The correlations in the mean diurnal cycles remain largely the same, with the exception of Wieringerwerf-Medemblikkerweg where a moderate improvement from $R = 0.72$ to $R = 0.77$ is found. However, the diurnal cycle at Wieringerwerf-Medemblikkerweg appears to be significantly underestimated in both the base model and the optimized simulation. These underestimations are likely related to local-scale NH_3 enhancements from the nearby farmlands in the Wiereingermeer polder being poorly captured by the model, coupled with complex local meteorological conditions due to the proximity of the site to the IJsselmeer and the North Sea which is roughly 15–20km away.

Although the assimilation of satellite observations improves the representation of NH_3 diurnal cycles relative to measurements at several LML sites, particularly with respect to systematic biases, it does not fully reproduce the observed variability. A fundamental limitation of Kalman filter–based approaches is that, although emissions are adjusted, there can be a temporal lag




Figure 9. Taylor diagram displaying the Pearson correlation coefficients and the relative standard deviations of the comparisons between the model and each LML site for the 2018–2022 period before (black) and after (red) assimilation.



associated with the transport of concentrations from source regions to the measurement locations, which is strongly modulated by local meteorological conditions. Furthermore, when the underlying diurnal cycle in the model is misrepresented, assimilation at only two main satellite overpass times may be insufficient to correct these deficiencies. An additional challenge is that NH_3 emissions originate from diverse sources such as livestock housing, manure application, and industry, each with distinct diurnal patterns, whereas the Kalman filter adjusts emissions uniformly without differentiating among source types. Overall, at sites located in regions of intensive agricultural activity such as Valthermond, Vredepeel, and Zegveld where midday enhancements are more likely to be captured by satellite observations, the influence of the assimilation on the diurnal cycles is more pronounced.

While these comparisons demonstrate clear improvements after assimilation, several further limitations should be acknowledged. First, the six LML sites provide only sparse coverage of the Netherlands and may not fully represent the strong spatial heterogeneity of NH_3 concentrations across the entire country, particularly in regions with intensive agricultural emissions. Second, the model–observation comparison involves a scale mismatch: the LML instruments measure point concentrations, while the LOTOS-EUROS model output represents grid-cell averages, which can potentially introduce representativity errors. In addition, the ground-based measurements themselves are not perfect and subject to calibration uncertainties and potential interferences. Lastly, because the statistics are based on monthly averages, the analysis does not explicitly resolve short-term (daily to synoptic) variability. Consequently, part of the observed agreement may reflect the influence of meteorological variability in addition to the direct effects of the assimilation. Taken together, the LML comparisons suggest that the assimilation generally improves the agreement of the model with independent ground-based observations, particularly in terms of spatial patterns. At the same time, the analysis also underlines the limitations of both the observational dataset and the comparison methodology, which should be taken into account when interpreting the results.

3.5.2 Impact of satellite selection on LML surface comparisons

To investigate the impact of the co-assimilation and the choice of satellites on the final optimized model state, assimilation runs were repeated for the year of 2020 using subsets of the satellite data products. These runs were: IASI only, CrIS only, and CrIS and IASI (without TROPOMI NO_2), and the comparisons against the hourly LML NH_3 observations were then repeated individually.

The temporal means for each of the four assimilation runs versus the LML observations are shown in Figure 11. From Figure 11, it can be seen that relative to the other assimilation runs, the assimilation performed using IASI, CrIS and TROPOMI leads to the lowest overall bias ($0.1 \mu\text{g m}^{-3}$) relative to the LML observations and a relatively strong correlation of $R = 0.88$. The simulation performed with IASI and CrIS without the assimilation of TROPOMI NO_2 yields a higher correlation of $R = 0.90$, but displays a higher bias of $0.7 \mu\text{g m}^{-3}$, a higher standard deviation, and a poorer regression slope. The simulations using only CrIS and only IASI display relatively higher mean biases of $1.4 \mu\text{g m}^{-3}$ and $-1.1 \mu\text{g m}^{-3}$, respectively. For comparison purposes, the temporal mean comparison for the base (unoptimized) simulation for 2020 is shown in the Appendix in panel (a) of Figure A2. Each of the simulations performed with assimilation show an improvement over the unoptimized base model run, which displays a correlation of $R = 0.82$ and a mean bias of $-1.9 \mu\text{g m}^{-3}$ relative to the LML observations.




Figure 10. Mean diurnal cycle of NH_3 for the period of 2018-2022 at each LML site from the observations (dark grey), the base model run (red), and the assimilation run (blue). On each panel, the difference between the mean diurnal cycles (Δ_{mean}) and the Pearson correlation coefficient (R) are provided for the base run (BG) in red and the assimilation run (AS) in blue.




Figure 11. Scatter plot of monthly temporal means of assimilation runs using (a) IASI and CrIS NH₃ and TROPOMI NO₂ observations, (b) IASI and CrIS NH₃ observations, (c) CrIS NH₃ observations only, and (d) IASI NH₃ observations only versus LML observed NH₃ surface concentrations for the year of 2020.

A similar plot of the spatial means for each of the assimilation runs versus the LML surface observations is provided in Figure 12. It can be seen that again from Figure 12(a), the assimilation performed using IASI, CrIS, and TROPOMI leads to the lowest overall bias ($0.1 \mu\text{g m}^{-3}$) of the four simulations and a relatively strong correlation of $R = 0.90$. In comparison, the assimilation run using IASI and CrIS observations but not TROPOMI has a slightly higher correlation ($R = 0.93$), but a higher bias of $0.7 \mu\text{g m}^{-3}$ and a poorer slope of regression (1.163 versus 0.977 from the IASI, CrIS and TROPOMI assimilation). The assimilation runs performed using CrIS only and IASI only (shown in panels (c) and (d) of Figure 12, respectively) have the same or poorer correlations than the IASI, CrIS and TROPOMI run, and relatively larger biases of $1.4 \mu\text{g m}^{-3}$ and $-1.1 \mu\text{g m}^{-3}$ for the CrIS-only and IASI-only runs, respectively. As was the case for the temporal mean comparisons, each of the optimized

525




Figure 12. Scatter plot of monthly spatial means of assimilation runs using (a) IASI and CrIS NH₃ and TROPOMI NO₂ observations, (b) IASI and CrIS NH₃ observations, (c) CrIS NH₃ observations only, and (d) IASI NH₃ observations only versus LML observed NH₃ surface concentrations for the year of 2020.

530 runs provides a significant improvement in comparison with the observations versus the unoptimized simulation (panel (b) of Figure A2), which had a weaker correlation of $R = 0.66$ and a mean bias of $-1.9 \mu\text{g m}^{-3}$ for 2020.

These comparisons highlight that the choice of satellite datasets used for assimilation is impactful, and that using just IASI or CrIS for the optimization may lead to greater biases than using a combined assimilation approach. Although assimilating a combination of CrIS and IASI observations leads to an improvement over just a single subset of the satellite data, the inclusion of TROPOMI NO₂ in the optimization aids in reducing the overall biases while still maintaining a strong correlation, particularly in comparison to the un-optimized LOTOS-EUROS simulation. The added value of assimilating NO₂ is that it helps to constrain the availability of HNO₃ through the NO₂–OH oxidation pathway, which in turn directly regulates the

535



partitioning of NH_3 into ammonium nitrate, an important sink of NH_3 (Finlayson-Pitts and Pitts, 2000). In this way, the NO_2 observations provide an indirect but important constraint on the fate of NH_3 , thereby improving the consistency of the coupled reactive nitrogen system in the model.

3.5.3 LML NH_x wet deposition comparisons

To evaluate the impact of the assimilation on the deposition of NH_x within the Netherlands, comparisons were performed against dissolved NH_4^+ concentrations measured in precipitation. These measurements are made at irregular intervals at several LML sites, and for the comparisons, these observations were paired with the modeled deposition interpolated to a daily frequency and converted to wet deposition fluxes (in $\text{kg N ha}^{-1} \text{ yr}^{-1}$) using the measured precipitation amounts.

Figure 13 shows the comparison of the measured and modeled temporal means of NH_4^+ wet deposition flux before and after assimilation. The baseline simulation displays a moderate correlation with the observations of $R = 0.51$ which improves to $R = 0.56$ after assimilation, while the mean bias improved slightly from $-1.0 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ to $-0.8 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ before and after assimilation, respectively. The scatter plot also shows a particularly large overestimation in the modeled deposition flux for several LML sites in the spring months of 2018 and 2020 which are reduced after assimilation, while there is a general underestimation during much of the remainder of the year. A similar pattern of overestimation and underestimation was found in the wet deposition comparisons in the earlier LOTOS-EUROS LETKF assimilation study by Van Der Graaf et al. (2022), which they attribute to a potential overestimation of the springtime NH_3 emissions and an underestimation in the later part of the year. A scatterplot of the spatial means is shown in Figure 14. A notable improvement is found in the correlation coefficients of the spatial means which went from $R = 0.58$ in the base model run to $R = 0.68$ in the assimilation run, as well as a small decrease in the mean bias from $-1.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ before assimilation to $-0.9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ after assimilation. The standard deviation of the differences also decreased from $3.0 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ to $2.4 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ before and after assimilation, respectively, indicating a lower variance in the model measurement differences.

In general, the wet deposition comparisons against the LML observations presented here indicate an improved spatial and temporal representation of the nitrogen wet deposition fluxes in the model after the assimilation of the satellite observations, further highlighting the value of satellite datasets for constraining not only emissions but also the resulting deposition.

3.5.4 MAN monthly NH_3 comparisons

The baseline and optimized simulations were also evaluated against monthly MAN NH_3 surface observations from a total of 315 sites (309 standard sites, and 6 MAN instruments located at LML sites) across the Netherlands for 2018–2022. Figure 15 shows the mean observed and modeled NH_3 concentrations, along with their differences. The baseline simulation exhibits a clear spatial bias, underestimation along the western coast and overestimation in the east, while the optimized run amplifies this pattern, consistent with the emission and concentration increases in the southern and eastern Netherlands. No comparable bias pattern was observed for the hourly LML sites or the CrIS and IASI total column comparisons.

Figure 16 compares monthly spatial means across all MAN sites, and for comparison purposes, the LML and MAN statistics are also summarized together in Table 2. The assimilation substantially improves spatial correlation ($R = 0.79$ to 0.89), but the




Figure 13. Scatter plot of monthly temporal means of (left) base LOTOS-EUROS NH₄⁺ wet deposition flux, and (right) LOTOS-EUROS LETKF optimized NH₄⁺ wet deposition flux versus LML surface observations for the period of January 2018 to December 2022.




Figure 14. Scatter plot of monthly spatial means of (left) base LOTOS-EUROS NH₄⁺ wet deposition flux, and (right) LOTOS-EUROS LETKF optimized NH₄⁺ wet deposition flux versus LML surface observations for the period of January 2018 to December 2022. Each data-point represents the mean calculated across all LML sites for a given month.




Figure 15. Spatial maps of the mean NH_3 surface concentrations from (a) MAN observations, (b) the base simulation, and (c) the optimized simulation calculated over the period of 2018–2022. The absolute differences (in $\mu\text{g m}^{-3}$) are shown in the second row for (d) the optimized run versus the base run, (e) the base run versus the observations, and (f) the optimized run versus the observations, and the relative differences (in %) are shown for the same comparisons in panels (g) to (i).




Figure 16. Scatter plot of monthly spatial means of (left) base LOTOS-EUROS NH₃, and (right) LOTOS-EUROS LETKF optimized NH₃ versus MAN observed NH₃ surface concentrations for the period of January 2018 to December 2022. Each data-point represents the mean calculated across all MAN sites for a given month, and are colored corresponding to the month while the marker style indicates the year.

mean bias increases from $+1.2 \mu\text{g m}^{-3}$ to $+2.4 \mu\text{g m}^{-3}$, and the slope of regression grows from 1.39 to 1.65. This suggests that the assimilation enhances large-scale spatial and seasonal variability, but also introduces a systematic offset. The LML surface NH₃ comparisons showed the bias and slope improving post-assimilation from $-2.0 \mu\text{g m}^{-3}$ to $-1.0 \mu\text{g m}^{-3}$, and 0.79 to 0.90, respectively. Supporting analyses using individual-site MAN monthly means and time series of national averages (appendix 575 Figs. B1 and B2) show similar behavior, with the optimized run better reproducing inter-annual variability but maintaining a high bias, particularly during summer months of 2020 and 2022, the years with the strongest emission adjustments (see Figure 2).

Additionally, to support a clearer comparison with the LML results and to assess the spatial representativity of the MAN network, we evaluated observations from the six MAN sensors located at LML sites against both the base and optimized 580 LETKF simulations. These six sensors are not part of the standard MAN dataset but are used exclusively for the monthly calibration of the remaining 309 MAN sites, as described in Noordijk et al. (2020). A spatial-mean scatterplot and monthly mean time-series are provided in the appendix as Figs. B3 and B4, respectively, and the associated statistics are summarized in Table 2.

The comparison of the MAN calibration sensors at the LML sites with the model simulations reveals patterns consistent with 585 the LML analysis. Prior to assimilation, the MAN-model mean bias is $-2.4 \mu\text{g m}^{-3}$, which is reduced to $-1.0 \mu\text{g m}^{-3}$ after assimilating the satellite observations. The spatial correlation also improves from ($R = 0.66$) to ($R = 0.76$), and the temporal agreement in the time-series (Fig. B4) is also improved, especially during the summer months. These values closely match the



LML results, particularly the post-assimilation bias of $-1.0 \mu\text{g m}^{-3}$. In contrast, the full MAN network exhibits positive biases of $+1.2 \mu\text{g m}^{-3}$ and $+2.4 \mu\text{g m}^{-3}$ against the base and optimized simulations, respectively, highlighting the different behavior of the co-located MAN sensors compared to the broader network.

Table 2. Summary of model performance statistics for NH_3 surface concentration comparisons against LML and MAN observations for the 2018–2022 period. Reported values correspond to spatial mean comparisons.

Network	Comparison Type	R	Slope	Mean Bias ($\mu\text{g m}^{-3}$)
LML	NH_3 surface (Base)	0.77	0.79	-2.0
LML	NH_3 surface (Optimized)	0.84	0.90	-1.0
MAN	NH_3 surface (Base)	0.79	1.39	+1.2
MAN	NH_3 surface (Optimized)	0.89	1.65	+2.4
MAN @ LML	NH_3 surface (Base)	0.66	0.67	-2.4
MAN @ LML	NH_3 surface (Optimized)	0.76	0.83	-1.0

Part of this contrast likely stems from measurement uncertainties in the passive Gradko samplers, but representativeness effects are expected to dominate. The LML stations and the satellite footprints sample air masses that are more representative of the broader agricultural landscape, whereas most MAN sites are located within Natura2000 areas, frequently in or near forested environments where canopy shielding and strong local heterogeneity depress NH_3 concentrations relative to the surrounding terrain. These sheltered conditions are highly localized and not captured by the coarse spatial scales of the regional model ($7 \times 7 \text{ km}^2$) or the 12–14 km nadir footprint of the satellite retrievals. As a result, the MAN measurements at these locations tend to reflect sub-grid processes rather than the landscape-average conditions that the model and satellite products are designed to represent. Remaining differences may also relate to how canopy deposition or near-surface mixing is expressed at these fine spatial scales, rather than broad model biases.

Overall, the MAN comparisons reaffirm that the LETKF enhances the spatial coherence of the NH_3 fields and performs consistently when evaluated against regionally representative measurements such as the LML sites. The discrepancies observed at many MAN locations primarily reflect representativeness limitations inherent in comparing coarse-scale model and satellite products with low-temporal-resolution passive samplers situated in highly heterogeneous environments. Reducing these mismatches will likely require finer-scale process representation and more explicit treatment of land-cover-dependent effects in future assimilation studies.

4 Conclusions

This study demonstrates that assimilating NH_3 and NO_2 satellite observations into the LOTOS-EUROS model using the LETKF framework substantially improves the representation of reactive nitrogen dynamics over the Netherlands. By co-assimilating IASI, CrIS, and TROPOMI retrievals over 2018–2022, the system produced optimized emissions, deposition, and concentration fields. The optimized emission fields showed consistent spatial structures across years, with persistent increases



in the southern and eastern Netherlands, and exhibited notable temporal shifts such as a reduced springtime emission peak and enhanced summertime emissions. These patterns are broadly consistent with recent observational and modeling studies evaluating NH_3 emissions over the region (Ge et al., 2020; Van Der Graaf et al., 2022; Ding et al., 2024).

Validation against independent LML surface observations at six sites showed clear post-assimilation improvements, with
615 mean biases reduced from -2.0 to $-1.0 \mu\text{g m}^{-3}$ and spatial correlations strengthened from $R = 0.77$ to $R = 0.84$. Assimilation
also improved the representation of diurnal cycles, reducing systematic underestimation and bringing simulated amplitudes
closer to observations, although some residual site-specific variability remained. Comparisons with dissolved NH_4^+ in precip-
itation likewise indicated enhanced performance, including an increase in the spatial correlation from $R = 0.58$ to $R = 0.68$.
Sensitivity tests for 2020 confirmed that the multi-satellite configuration performed best, with joint assimilation of IASI and
620 CrIS NH_3 and TROPOMI NO_2 producing the lowest biases and most robust improvements relative to LML.

Comparisons with monthly MAN observations showed improved spatial correlations after assimilation ($R = 0.79$ to $R =$
 0.89), but also revealed a systematic positive bias that increased from $+1.2$ to $+2.4 \mu\text{g m}^{-3}$. These biases primarily reflect rep-
resentativeness differences: many MAN sites are located within Natura2000 areas, often in forested or sheltered environments
where canopy effects and fine-scale heterogeneity suppress local NH_3 levels relative to the surrounding agricultural landscape.
625 Such localized variability is not resolved at the model grid scale or by satellite footprints, leading to systematic overestimation
when landscape-scale fields are evaluated against sheltered MAN locations.

In contrast, the MAN calibration sensors co-located at six LML stations which are situated in more open, source-proximal
environments, showed behavior consistent with the LML observations themselves. At these sites, the pre-assimilation mean
bias of $-2.4 \mu\text{g m}^{-3}$ improved to $-1.0 \mu\text{g m}^{-3}$, and the spatial correlation increased from $R = 0.66$ to $R = 0.76$. These re-
630 sults confirm that, when evaluated at regionally representative sites, the LETKF adjustments produce coherent and physically
consistent improvements. Together, the LML and MAN comparisons highlight both the strengthened large-scale spatial and
temporal structure of NH_3 after assimilation and the ongoing challenges of reconciling coarse-resolution model and satellite
fields with passive samplers located in highly heterogeneous environments. Further progress may be enabled by adopting a
label-based Kalman filtering approach. The labeling functionality introduced in LOTOS-EUROS v2.3 could be extended to the
635 LETKF, allowing sector-specific emission optimization and supporting finer-scale improvements.

The results also underscore the importance of satellite observation density, particularly for NH_3 . The year 2020, character-
ized by the highest availability of CrIS and IASI retrievals, showed the largest emission and concentration adjustments. The
recent launch of MTG-IRS on the MTG-S1 platform (1 July 2025) presents a major opportunity: its half-hourly $4 \times 4 \text{ km}$ NH_3
retrievals will provide unprecedented spatial and temporal coverage over Europe, enabling improved constraints on diurnal
640 variability, better characterization of emission events, and more rigorous evaluation of emission inventories. A dedicated NH_3
assimilation study using MTG-IRS is recommended once data become available.

In addition to these advancements, several refinements warrant further investigation. These include improving the represen-
tativeness between model and surface observations, enhancing the treatment of diurnal variability across source sectors, and
conducting a full system simulation experiment (SSE) to rigorously evaluate LETKF performance.



645 Overall, these findings highlight the essential role of satellite constraints in advancing chemical transport modeling and
nitrogen budget estimation. Co-assimilating complementary satellite instruments provides a pathway toward more accurate and
internally consistent representations of reactive nitrogen, strengthening our ability to constrain emissions, evaluate inventories,
and assess deposition at regional to national scales. The demonstrated improvements offer a strong foundation for exploiting
emerging missions such as MTG-IRS, enabling continued progress toward capturing fine-scale processes and further improving
650 assimilation performance in future studies.

Code availability. The use of the LOTOS-EUROS open-source version is regulated via registration. The open-source version of the model
can be obtained from <https://airqualitymodeling.tno.nl/lotos-euros/open-source-version/>. Access to the LOTOS-EUROS LETKF can be pro-
vided upon formal request to the authors. The CAMS Satellite Operator (CSO) was used in this work and is an open-access tool developed
at TNO and implemented to facilitate fast intercomparisons between modelled and satellite concentrations. CSO can be downloaded from:
655 (<https://ci.tno.nl/gitlab/cams/cso>).

Data availability. The CrIS NH₃ v1.6.4 data from SNPP and NOAA-20 created by Environment and Climate Change Canada are currently
publicly available upon request (mark.shephard@canada.ca). The IASI-NH₃ v4 ANNI datasets (Clarisse et al., 2023) are available from the
AERIS data infrastructure (<https://iasi.aeris-data.fr/nh3/>, last access: 26 August 2025). The TROPOMI NO₂ version 2.4 data (van Geffen
et al., 2022) are available on the Copernicus website (<https://dataspace.copernicus.eu/>, last access: 26 August 2025). The NH₃ concentration
660 and NH₄⁺ wet deposition data from the LML network are available on the RIVM website (<https://data.rivm.nl/data/luchtmeetnet/>; last access:
26 August 2025) (LML, 2025). The monthly NH₃ surface data from the MAN network are available at <https://man.rivm.nl> (last access: 26
August 2025) (MAN, 2025). Map data copyrighted OpenStreetMap contributors and available from <https://www.openstreetmap.org/>.

Author contributions. T. Wizenberg prepared the manuscript with contributions from all authors. T. Wizenberg, E. Dammers, and M. Schaap
designed the experiment and provided scientific guidance during the project. A. Segers developed the LOTOS-EUROS code and provided
665 assistance with performing the assimilation runs. M. Shephard, P. Coheur, M. Van Damme, L. Clarisse, and H. Eskes developed the satellite
retrievals and provided the data. R. Wichink Kruit and S. van der Graaf provided the LML and MAN data and provided feedback on the
analysis of these datasets. T. Wizenberg performed the formal analysis and presentation of the results.

Competing interests. The authors declare that they have no competing interests.

Acknowledgements. This study was funded by the Dutch Ministry of Agriculture, Fisheries, Food Security and Nature (LVVN), within the
670 framework of the National Nitrogen Knowledge Programme (NKS), project NKS-SAGEN, on satellite observations and ensemble modeling.

<https://doi.org/10.5194/egusphere-2025-5926>

Preprint. Discussion started: 11 January 2026

© Author(s) 2026. CC BY 4.0 License.



Lieven Clarisse is a senior research associate supported by the Belgian F.R.S.-FNRS. Generative AI tools were used in the drafting and editing process of this manuscript.



Appendix A: Additional figures for LML network comparisons

A1 Vredepeel-Vredeweg hourly NH_3 timeseries




Figure A1. Timeseries of hourly NH_3 surface concentration values at the Vredepeel-Vredeweg LML site from (black) the observations, (red) the background LOTOS-EUROS simulation, and (blue) the analysis run with assimilation.



675 A2 LML comparisons for 2020 with base model simulation




Figure A2. Scatter plot of (a) monthly temporal means of LOTOS-EUROS simulated NH₃ versus LML observed NH₃ surface concentrations, and (b) monthly spatial means of LOTOS-EUROS simulated NH₃ versus LML observed NH₃ surface concentrations for the period of January to December 2020.



Appendix B: Additional figures for MAN network comparisons

B1 Monthly mean scatter plot pre- and post-assimilation




Figure B1. Scatter plot comparing monthly means of observed MAN NH₃ mass concentrations with (left) base LOTOS-EUROS surface NH₃ mass concentration, and (right) LOTOS-EUROS LETKF optimized NH₃ mass concentration. Each data-point represents a mean at a single MAN site.

B2 Monthly mean timeseries




Figure B2. Time-series of monthly mean NH_3 surface mass concentrations and differences calculated across all MAN sites during 2018–2022. The shaded regions indicate the standard deviations of the monthly means.



B3 MAN @ LML sites: spatial correlations




Figure B3. Scatter plot of monthly spatial means of (left) base LOTOS-EUROS NH₃, and (right) LOTOS-EUROS LETKF optimized NH₃ versus MAN observed NH₃ surface concentrations at the locations of LML sites for the period of January 2018 to December 2022. Each data-point represents the mean calculated across the 6 MAN calibration sensors located at the LML sites for a given month, and are colored corresponding to the month while the marker style indicates the year.



680 B4 MAN @ LML sites: monthly mean time-series




Figure B4. Time-series of monthly mean NH_3 surface mass concentrations and differences calculated across the 6 MAN calibration sensors located at the LML sites during 2018–2022. The shaded regions indicate the standard deviations of the monthly means.



References

- Improved estimate of the policy-relevant background ozone in the United States using the GEOS-Chem global model with $1/2^\circ \times 2/3^\circ$ horizontal resolution over North America, *Atmospheric Environment*, 45, 6769–6776, <https://doi.org/10.1016/j.atmosenv.2011.07.054>, 2011.
- 685 LML: Dutch Air Quality Monitoring, <https://data.rivm.nl/data/luchtmeetnet/>, 2025.
- MAN: Dutch Measuring Ammonia in Nature, <https://man.rivm.nl>, 2025.
- Berkhout, A. J., Swart, D. P., Volten, H., Gast, L. F., Haaima, M., Verboom, H., Stefess, G., Hafkenscheid, T., and Hoogerbrugge, R.: Replacing the AMOR with the miniDOAS in the ammonia monitoring network in the Netherlands, *Atmospheric Measurement Techniques*, 10, 4099–4120, <https://doi.org/10.5194/amt-10-4099-2017>, 2017.
- 690 Bessagnet, B., Pirovano, G., Mircea, M., Cuvelier, C., Aulinger, A., Calori, G., Ciarelli, G., Manders, A., Stern, R., Tsyro, S., García Vivanco, M., Thunis, P., Pay, M.-T., Colette, A., Couvidat, F., Meleux, F., Rouil, L., Ung, A., Aksoyoglu, S., Baldasano, J. M., Bieser, J., Briganti, G., Cappelletti, A., D’Isidoro, M., Finardi, S., Kranenburg, R., Silibello, C., Carnevale, C., Aas, W., Dupont, J.-C., Fagerli, H., Gonzalez, L., Menut, L., Prévôt, A. S. H., Roberts, P., and White, L.: Presentation of the EURODELTA III intercomparison exercise – evaluation of the chemistry transport models’ performance on criteria pollutants and joint analysis with meteorology, *Atmospheric Chemistry and*
- 695 *Physics*, 16, 12 667–12 701, <https://doi.org/10.5194/acp-16-12667-2016>, 2016.
- Chan, K. L., Wiegner, M., van Geffen, J., De Smedt, I., Alberti, C., Cheng, Z., Ye, S., and Wenig, M.: MAX-DOAS measurements of tropospheric NO_2 and HCHO in Munich and the comparison to OMI and TROPOMI satellite observations, *Atmospheric Measurement Techniques*, 13, 4499–4520, <https://doi.org/10.5194/amt-13-4499-2020>, 2020.
- Chen, Z., Jacob, D. J., Gautam, R., Omara, M., Stavins, R. N., Stowe, R. C., Nesser, H., Sulprizio, M. P., Lorente, A., Varon, D. J., Lu,
- 700 X., Shen, L., Qu, Z., Pendergrass, D. C., and Hancock, S.: Satellite quantification of methane emissions and oil-gas methane intensities from individual countries in the Middle East and North Africa: implications for climate action, *Atmospheric Chemistry and Physics*, 23, 5945–5967, <https://doi.org/10.5194/acp-23-5945-2023>, 2023.
- Clarisse, L., Franco, B., Van Damme, M., Di Gioacchino, T., Hadji-Lazaro, J., Whitburn, S., Noppen, L., Hurtmans, D., Clerbaux, C., and Coheur, P.: The IASI NH_3 version 4 product: averaging kernels and improved consistency, *Atmospheric Measurement Techniques*, 16,
- 705 5009–5028, <https://doi.org/10.5194/amt-16-5009-2023>, 2023.
- Clerbaux, C., Boynard, A., Clarisse, L., George, M., Hadji-Lazaro, J., Herbin, H., Hurtmans, D., Pommier, M., Razavi, A., Turquety, S., Wespes, C., and Coheur, P.-F.: Monitoring of atmospheric composition using the thermal infrared IASI/MetOp sounder, *Atmospheric Chemistry and Physics*, 9, 6041–6054, <https://doi.org/10.5194/acp-9-6041-2009>, 2009.
- Colette, A., Andersson, C., Manders, A., Mar, K., Mircea, M., Pay, M.-T., Raffort, V., Tsyro, S., Cuvelier, C., Adani, M., Bessagnet, B.,
- 710 Bergström, R., Briganti, G., Butler, T., Cappelletti, A., Couvidat, F., D’Isidoro, M., Doumbia, T., Fagerli, H., Granier, C., Heyes, C., Klimont, Z., Ojha, N., Otero, N., Schaap, M., Sindelarova, K., Stegehuis, A. I., Roustan, Y., Vautard, R., Van Meijgaard, E., Vivanco, M. G., and Wind, P.: EURODELTA-Trends, a multi-model experiment of air quality hindcast in Europe over 1990–2010, *Geoscientific Model Development*, 10, 3255–3276, <https://doi.org/10.5194/gmd-10-3255-2017>, 2017.
- Colette, A., Collin, G., Besson, F., Blot, E., Guidard, V., Meleux, F., Royer, A., Petiot, V., Miller, C., Fermond, O., Jeant, A., Adani, M.,
- 715 Arteta, J., Benedictow, A., Bergström, R., Bowdalo, D., Brandt, J., Briganti, G., Carvalho, A. C., Christensen, J. H., Couvidat, F., D’Elia, I., D’Isidoro, M., van der Gon, H. D., Descombes, G., Tomaso, E. D., Douros, J., Escribano, J., Eskes, H., Fagerli, H., Fatahi, Y., Flemming, J., Friese, E., Frohn, L., Gauss, M., Geels, C., Guarnieri, G., Guevara, M., Guion, A., Guth, J., Hänninen, R., Hansen, K., Im, U., Janssen,



- R., Jeoffrion, M., Joly, M., Jones, L., Jorba, O., Kadantsev, E., Kahnert, M., Kaminski, J. W., Kouznetsov, R., Kranenburg, R., Kuenen, J., Lange, A. C., Langner, J., Lannuque, V., Macchia, F., Manders, A., Mircea, M., Nyiri, A., Olid, M., García-Pando, C. P., Palamarchuk, Y., Piersanti, A., Raux, B., Razingger, M., Robertson, L., Segers, A., Schaap, M., Siljamo, P., Simpson, D., Sofiev, M., Stangel, A., Struzewska, J., Tena, C., Timmermans, R., Tsikerdekis, T., Tsyro, S., Tyuryakov, S., Ung, A., Uppstu, A., Valdebenito, A., van Velthoven, P., Vitali, L., Ye, Z., Peuch, V.-H., and Rouil, L.: Copernicus Atmosphere Monitoring Service – Regional Air Quality Production System v1.0, *Geoscientific Model Development*, 18, 6835–6883, <https://doi.org/10.5194/gmd-18-6835-2025>, 2025.
- Crutzen, P. J.: The Role of NO and NO₂ in the Chemistry of the Troposphere and Stratosphere, *Annual Review of Earth and Planetary Sciences*, 7, 443–472, <https://doi.org/doi.org/10.1146/annurev.ea.07.050179.002303>, 1979.
- Dammers, E., Palm, M., Van Damme, M., Vigouroux, C., Smale, D., Conway, S., Toon, G. C., Jones, N., Nussbaumer, E., Warneke, T., Petri, C., Clarisse, L., Clerbaux, C., Hermans, C., Lutsch, E., Strong, K., Hannigan, J. W., Nakajima, H., Morino, I., Herrera, B., Stremme, W., Grutter, M., Schaap, M., Wichink Kruit, R. J., Notholt, J., Coheur, P.-F., and Erisman, J. W.: An evaluation of IASI-NH₃ with ground-based Fourier transform infrared spectroscopy measurements, *Atmospheric Chemistry and Physics*, 16, 10 351–10 368, <https://doi.org/10.5194/acp-16-10351-2016>, 2016.
- Dammers, E., Shephard, M. W., Palm, M., Cady-Pereira, K., Capps, S., Lutsch, E., Strong, K., Hannigan, J. W., Ortega, I., Toon, G. C., Stremme, W., Grutter, M., Jones, N., Smale, D., Siemons, J., Hrpcek, K., Tremblay, D., Schaap, M., Notholt, J., and Erisman, J. W.: Validation of the CrIS fast physical NH₃ retrieval with ground-based FTIR, *Atmospheric Measurement Techniques*, 10, 2645–2667, <https://doi.org/10.5194/amt-10-2645-2017>, 2017.
- Ding, J., A, R. V. D., Eskes, H., Dammers, E., Shephard, M., Kruit, R. W., Guevara, M., and Tarrason, L.: Ammonia emission estimates using CrIS satellite observations over Europe, *Atmospheric Chemistry and Physics*, 24, 10 583–10 599, <https://doi.org/10.5194/acp-24-10583-2024>, 2024.
- Douros, J., Eskes, H., van Geffen, J., Boersma, K. F., Compennolle, S., Pinardi, G., Blechschmidt, A.-M., Peuch, V.-H., Colette, A., and Veefkind, P.: Comparing Sentinel-5P TROPOMI NO₂ column observations with the CAMS regional air quality ensemble, *Geoscientific Model Development*, 16, 509–534, <https://doi.org/10.5194/gmd-16-509-2023>, 2023.
- Elskamp, H.: National Air Quality Monitoring Network. Technical description, Report 228702017, <http://hdl.handle.net/10029/259431>, 1989.
- Elzakker, B. and Buijsman, E.: Meetactiviteiten in 1999 in het Landelijk Meetnet Luchtkwaliteit, Report 723101032, <https://www.rivm.nl/bibliotheek/rapporten/729999002.pdf>, 1999.
- Erisman, J. W., Sutton, M. A., Galloway, J., Zbigniew, K., and Winiwarter, W.: How a century of ammonia synthesis changed the world, *Nature Geoscience*, 1, 636–639, <https://doi.org/doi.org/10.1038/ngeo325>, 2008.
- Erisman, J. W., van Grinsven, H., Grizzetti, B., Bouraoui, F., Powlson, D., Sutton, M. A., Bleeker, A., and Reis, S.: The European nitrogen problem in a global perspective, pp. 9–31, Cambridge University Press, <https://doi.org/10.1017/cbo9780511976988.005>, 2011.
- Eskes, H., van Geffen, J., Boersma, K., Eichmann, K.-U., Apituley, A., Pedergnana, M., Sneep, M., Veefkind, J. P., and Loyola, D.: Sentinel-5 precursor/TROPOMI Level 2 Product User Manual Nitrogen dioxide, Tech. Rep. SSP-KNMI-L2-0021-MA, Koninklijk Nederlands Meteorologisch Instituut (KNMI), issue 4.3.0, processor version 2.7.1, CI-7570-PUM, date 4 April 2024, <https://sentiwiki.copernicus.eu/web/s5p-products#S5P-Products-L2>, 2024.
- Eum, K. D., Honda, T. J., Wang, B., Kazemiparkouhi, F., Manjourides, J., Pun, V. C., Pavlu, V., and Suh, H.: Long-term nitrogen dioxide exposure and cause-specific mortality in the U.S. Medicare population, *Environmental Research*, 207, <https://doi.org/10.1016/j.envres.2021.112154>, 2022.



- Evensen, G.: The Ensemble Kalman Filter: theoretical formulation and practical implementation, *Ocean Dynamics*, 53, 343–367, <https://doi.org/10.1007/s10236-003-0036-9>, 2003.
- Finlayson-Pitts, B. J. and Pitts, J. N.: Chemistry of the upper and lower atmosphere: theory, experiments, and applications, Academic Press, San Diego, 2000.
- 760 Galloway, J. N., Townsend, A. R., Erisman, J. W., Bekunda, M., Cai, Z., Freney, J. R., Martinelli, L. A., Seitzinger, S. P., and Sutton, M. A.: Transformation of the Nitrogen Cycle: Recent Trends, Questions, and Potential Solutions, *Science*, 320, 889–892, <https://doi.org/10.1126/science.1136674>, 2008.
- Ge, X., Schaap, M., Kranenburg, R., Segers, A., Reinds, G. J., Kros, H., and Vries, W. D.: Modeling atmospheric ammonia using agricultural emissions with improved spatial variability and temporal dynamics, *Atmospheric Chemistry and Physics*, 20, 16055–16087,
765 <https://doi.org/10.5194/acp-20-16055-2020>, 2020.
- Ge, X., Schaap, M., Dammers, E., Shephard, M., and de Vries, W.: Impact of interannual weather variation on ammonia emissions and concentrations in Germany, *Agricultural and Forest Meteorology*, 334, <https://doi.org/10.1016/j.agrformet.2023.109432>, 2023.
- Griffin, D., Zhao, X., McLinden, C. A., Boersma, F., Bourassa, A., Dammers, E., Degenstein, D., Eskes, H., Fehr, L., Fioletov, V., Hayden, K., Kharol, S. K., Li, S.-M., Makar, P., Martin, R. V., Mihele, C., Mittermeier, R. L., Krotkov, N., Sneep, M., Lamsal, L. N., Linden, M. t.,
770 Geffen, J. v., Veefkind, P., and Wolde, M.: High-Resolution Mapping of Nitrogen Dioxide With TROPOMI: First Results and Validation Over the Canadian Oil Sands, *Geophysical Research Letters*, 46, 1049–1060, <https://doi.org/https://doi.org/10.1029/2018GL081095>, 2019.
- Guo, X., Wang, R., Pan, D., Zondlo, M. A., Clarisse, L., Damme, M. V., Whitburn, S., Coheur, P. F., Clerbaux, C., Franco, B., Golston, L. M., Wendt, L., Sun, K., Tao, L., Miller, D., Mikoviny, T., Müller, M., Wisthaler, A., Tevlin, A. G., Murphy, J. G., Nowak, J. B., Roscioli, J. R., Volkamer, R., Kille, N., Neuman, J. A., Eilerman, S. J., Crawford, J. H., Yacovitch, T. I., Barrick, J. D., and Scarino, A. J.: Validation of
775 IASI Satellite Ammonia Observations at the Pixel Scale Using In Situ Vertical Profiles, *Journal of Geophysical Research: Atmospheres*, 126, <https://doi.org/10.1029/2020JD033475>, 2021.
- Gyldenkerne, S., Skjøth, C. A., Hertel, O., and Ellermann, T.: A dynamical ammonia emission parameterization for use in air pollution models, *Journal of Geophysical Research Atmospheres*, 110, 1–14, <https://doi.org/10.1029/2004JD005459>, 2005.
- Herrera, B., Bezanilla, A., Blumenstock, T., Dammers, E., Hase, F., Clarisse, L., Magaldi, A., Rivera, C., Stremme, W., Strong, K., Viatte,
780 C., Van Damme, M., and Grutter, M.: Measurement report: Evolution and distribution of NH₃ over Mexico City from ground-based and satellite infrared spectroscopic measurements, *Atmospheric Chemistry and Physics*, 22, 14119–14132, <https://doi.org/10.5194/acp-22-14119-2022>, 2022.
- Hunt, B. R., Kostelich, E. J., and Szunyogh, I.: Efficient data assimilation for spatiotemporal chaos: A local ensemble transform Kalman filter, *Physica D: Nonlinear Phenomena*, 230, 112–126, <https://doi.org/10.1016/j.physd.2006.11.008>, 2007.
- 785 Iturbide-Sanchez, F., Strow, L., Tobin, D., Chen, Y., Tremblay, D., Knuteson, R. O., Johnson, D. G., Buttes, C., Suwinski, L., Thomas, B. P., Rivera, A. R., Lynch, E., Zhang, K., Wang, Z., Porter, W. D., Jin, X., Predina, J. P., Eresmaa, R. I., Collard, A., Ruston, B., Jung, J. A., Barnet, C. D., Beierle, P. J., Yan, B., Mooney, D., and Revercomb, H.: Recalibration and Assessment of the SNPP CrIS Instrument: A Successful History of Restoration After Midwave Infrared Band Anomaly, *IEEE Transactions on Geoscience and Remote Sensing*, 60, <https://doi.org/10.1109/TGRS.2021.3112400>, 2022.
- 790 Jonson, J. E., Borken-Kleefeld, J., Simpson, D., Nyíri, A., Posch, M., and Heyes, C.: Impact of excess NO_x emissions from diesel cars on air quality, public health and eutrophication in Europe, *Environmental Research Letters*, 12, <https://doi.org/10.1088/1748-9326/aa8850>, 2017.



- Judd, L. M., Al-Saadi, J. A., Szykman, J. J., Valin, L. C., Janz, S. J., Kowalewski, M. G., Eskes, H. J., Veeffkind, J. P., Cede, A., Mueller, M., Gebetsberger, M., Swap, R., Pierce, R. B., Nowlan, C. R., Abad, G. G., Nehrir, A., and Williams, D.: Evaluating Sentinel-5P TROPOMI tropospheric NO₂ column densities with airborne and Pandora spectrometers near New York City and Long Island Sound, *Atmospheric Measurement Techniques*, 13, 6113–6140, <https://doi.org/10.5194/amt-13-6113-2020>, 2020.
- Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J. P., Super, I., and Gon, H. D. V. D.: CAMS-REG-v4: a state-of-the-art high-resolution European emission inventory for air quality modelling, *Earth System Science Data*, 14, 491–515, <https://doi.org/10.5194/essd-14-491-2022>, 2022.
- Kutzner, R. D., Cuesta, J., Chelin, P., Petit, J.-E., Ray, M., Landsheere, X., Tournadre, B., Dupont, J.-C., Rosso, A., Hase, F., Orphal, J., and Beekmann, M.: Diurnal evolution of total column and surface atmospheric ammonia in the megacity of Paris, France, during an intense springtime pollution episode, *Atmospheric Chemistry and Physics*, 21, 12 091–12 111, <https://doi.org/10.5194/acp-21-12091-2021>, 2021.
- Lambert, J.-C. and et al.: Quarterly Validation Report of the Copernicus Sentinel-5 Precursor Operational Data Products 24: April 2018 – August 2024, S5P MPC Routine Operations Consolidated Validation Report series, Issue 24, Version 24.00.00, 212 pp., 16 September 2024, 2024.
- Liu, S., Jørgensen, J. T., Ljungman, P., Pershagen, G., Bellander, T., Leander, K., Magnusson, P. K., Rizzuto, D., Hvidtfeldt, U. A., Raaschou-Nielsen, O., Wolf, K., Hoffmann, B., Brunekreef, B., Strak, M., Chen, J., Mehta, A., Atkinson, R. W., Bauwelinck, M., Varraso, R., Boutron-Ruault, M.-C., Brandt, J., Cesaroni, G., Forastiere, F., Fecht, D., Gulliver, J., Hertel, O., de Hoogh, K., Janssen, N. A., Katsouyanni, K., Ketzel, M., Klompmaker, J. O., Nagel, G., Oftedal, B., Peters, A., Tjønneland, A., Rodopoulou, S. P., Samoli, E., Kristoffersen, D. T., Sigsgaard, T., Stafoggia, M., Vienneau, D., Weinmayr, G., Hoek, G., and Andersen, Z. J.: Long-term exposure to low-level air pollution and incidence of asthma: the ELAPSE project, *European Respiratory Journal*, 57, 2003 099, <https://doi.org/10.1183/13993003.030992020>, 2021.
- Lolkema, D. E., Noordijk, H., Stolk, A. P., Hoogerbrugge, R., Zanten, M. C. V., and Pul, W. A. V.: The Measuring Ammonia in Nature (MAN) network in the Netherlands, *Biogeosciences*, 12, 5133–5142, <https://doi.org/10.5194/bg-12-5133-2015>, 2015.
- Lopez-Restrepo, S., Yarce, A., Pinel, N., Quintero, O. L., Segers, A., and Heemink, A. W.: Forecasting PM₁₀ and PM_{2.5} in the Aburrá Valley (Medellín, Colombia) via EnKF based data assimilation, *Atmospheric Environment*, 232, <https://doi.org/10.1016/j.atmosenv.2020.117507>, 2020.
- Lô, S., Dohmen, W., Heederik, D., Weijers, E., van Schothorst, I., Bleeker, A., Vermeulen, R., and Hoek, G.: Spatial and temporal variability of atmospheric ammonia using a dense network in an area with livestock, residential and natural environments intertwined, *Atmospheric Environment*, 360, <https://doi.org/10.1016/j.atmosenv.2025.121394>, 2025.
- Manders, A. M. M., Builtjes, P. J. H., Curier, L., Denier Van Der Gon, H. A. C., Hendriks, C., Jonkers, S., Kranenburg, R., Kuenen, J. J. P., Segers, A. J., Timmermans, R. M. A., Visschedijk, A. J. H., Wichink Kruit, R. J., Van Pul, W. A. J., Sauter, F. J., Van Der Swaluw, E., Swart, D. P. J., Douros, J., Eskes, H., Van Meijgaard, E., Van Ulft, B., Van Velthoven, P., Banzhaf, S., Mues, A. C., Stern, R., Fu, G., Lu, S., Heemink, A., Van Velzen, N., and Schaap, M.: Curriculum vitae of the LOTOS–EUROS (v2.0) chemistry transport model, *Geoscientific Model Development*, 10, 4145–4173, <https://doi.org/10.5194/gmd-10-4145-2017>, 2017.
- Moncet, J.-L., Uymin, G., Lipton, A. E., and Snell, H. E.: Infrared Radiance Modeling by Optimal Spectral Sampling, *Journal of the Atmospheric Sciences*, 65, 3917–3934, <https://doi.org/10.1175/2008JAS2711.1>, 2008.
- Noordijk, H., Braam, M., Rutledge-Jonker, S., Hoogerbrugge, R., Stolk, A. P., and van Pul, W. A.: Performance of the MAN ammonia monitoring network in the Netherlands, *Atmospheric Environment*, 228, <https://doi.org/10.1016/j.atmosenv.2020.117400>, 2020.



- 830 Ojeda-Castillo, V., Murillo-Tovar, M. A., Hernández-Mena, L., Saldarriaga-Noreña, H., Vargas-Amado, M. E., Herrera-López, E. J., and Díaz, J.: Tropospheric NO₂: Anthropogenic Influence, Global Trends, Satellite Data, and Machine Learning Application, *Remote Sensing*, 17, <https://doi.org/10.3390/rs17010049>, 2025.
- OpenStreetMap contributors: Planet dump retrieved from <https://planet.osm.org>, <https://www.openstreetmap.org>, 2017.
- Paulot, F., Jacob, D. J., Pinder, R. W., Bash, J. O., Travis, K., and Henze, D. K.: Ammonia emissions in the United States, European Union, and China derived by high-resolution inversion of ammonium wet deposition data: Interpretation with a new agricultural emissions inventory (MASAGE_NH₃), *Journal of Geophysical Research*, 119, 4343 – 4364, <https://doi.org/10.1002/2013JD021130>, 2014.
- Peuch, V.-H., Engelen, R., Rixen, M., Dee, D., Flemming, J., Suttie, M., Ades, M., Agustí-Panareda, A., Ananasso, C., Andersson, E.,
835 Armstrong, D., Barré, J., Bousserez, N., Dominguez, J. J., Garrigues, S., Inness, A., Jones, L., Kipling, Z., Letertre-Danczak, J., Parrington, M., Razinger, M., Ribas, R., Vermoote, S., Yang, X., Simmons, A., Garcés De Marcilla, J., and Thépaut, J.-N.: The Copernicus Atmosphere Monitoring Service: From Research to Operations, *Bulletin of the American Meteorological Society*, 103, E2650–E2668, <https://doi.org/10.1175/BAMS-D-21-0314.1>, 2022.
- Pope, C. A. and Dockery, D. W.: Health effects of fine particulate air pollution: Lines that connect, *Journal of the Air and Waste Management Association*, 56, 709–742, <https://doi.org/10.1080/10473289.2006.10464485>, 2006.
840 Association, 56, 709–742, <https://doi.org/10.1080/10473289.2006.10464485>, 2006.
- Pope, C. A., Ezzati, M., and Dockery, D. W.: Fine-Particulate Air Pollution and Life Expectancy in the United States, *New England Journal of Medicine*, 360, 376–386, <https://doi.org/10.1056/nejmsa0805646>, 2009.
- Rodgers, C. D.: Inverse Methods for Atmospheric Sounding: Theory and Practice, vol. 2 of *Series on Atmospheric, Oceanic and Planetary Physics*, WORLD SCIENTIFIC, <https://doi.org/10.1142/3171>, 2000.
- 845 Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002jd002299>, 2003.
- Schaap, M., Timmermans, R. M., Roemer, M., Boersen, G., Bultjes, P. J., Sauter, F. J., Velders, G. J., Beck, J. P., and Sauter, P.: The LOTOS-EUROS model: description, validation and latest developments The LOTOS-EUROS model, Tech. rep., 2008.
- Shephard, M. W. and Cady-Pereira, K. E.: Cross-track Infrared Sounder (CRIS) satellite observations of tropospheric ammonia, 8, 1323–1336,
850 <https://doi.org/10.5194/amt-8-1323-2015>, 2015.
- Shephard, M. W., Dammers, E., Cady-Pereira, K. E., Kharol, S. K., Thompson, J., Gainariu-Matz, Y., Zhang, J., McLinden, C. A., Kovachik, A., Moran, M., Bittman, S., Sioris, C. E., Griffin, D., Alvarado, M. J., Lonsdale, C., Savic-Jovicic, V., and Zheng, Q.: Ammonia measurements from space with the Cross-track Infrared Sounder: characteristics and applications, *Atmospheric Chemistry and Physics*, 20, 2277–2302, <https://doi.org/10.5194/acp-20-2277-2020>, 2020.
- 855 Shephard, M. W., Kharol, S. K., Dammers, E., Sioris, C. E., Bell, A., Jansen, R., Caron, J., Snel, R., Palombo, E., Cady-Pereira, K. E., McLinden, C. A., Lutsch, E., and Knuteson, R. O.: Infrared Satellite Detection Limits for Monitoring Atmospheric Ammonia, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 18, 10272–10291, <https://doi.org/10.1109/JSTARS.2025.3557240>, 2025.
- Shin, S., Kang, J. S., and Jo, Y.: The Local Ensemble Transform Kalman Filter (LETKF) with a Global NWP Model on the Cubed Sphere, *Pure and Applied Geophysics*, 173, 2555–2570, <https://doi.org/10.1007/s00024-016-1269-0>, 2016.
- 860 Sutton, M. A., Oenema, O., Erisman, J. W., Leip, A., van Grinsven, H., and Winiwarter, W.: Too much of a good thing, *Nature*, 472, 159–161, <https://doi.org/10.1038/472159a>, 2011.
- Søgaard, H. T., Sommer, S. G., Hutchings, N. J., Huijsmans, J. F. M., Bussink, D. W., and Nicholson, F.: Ammonia volatilization from field-applied animal slurry-the ALFAM model, Tech. rep., www.alfam.dk, 2002.



- Timmermans, R., Segers, A., Curier, L., Abida, R., Attié, J. L., Amraoui, L. E., Eskes, H., Haan, J. D., Kujanpää, J., Lahoz, W., Nijhuis, A. O.,
865 Quesada-Ruiz, S., Ricaud, P., Veefkind, P., and Schaap, M.: Impact of synthetic space-borne NO₂ observations from the Sentinel-4 and
Sentinel-5P missions on tropospheric NO₂ analyses, *Atmospheric Chemistry and Physics*, 19, 12 811–12 833, <https://doi.org/10.5194/acp-19-12811-2019>, 2019.
- Tsyro, S., Aas, W., Colette, A., Andersson, C., Bessagnet, B., Ciarelli, G., Couvidat, F., Cuvelier, K., Manders, A., Mar, K., Mircea, M., Otero,
N., Pay, M. T., Raffort, V., Roustan, Y., Theobald, M. R., Vivanco, M. G., Fagerli, H., Wind, P., Briganti, G., Cappelletti, A., D’Isidoro, M.,
870 and Adani, M.: Eurodelta multi-model simulated and observed particulate matter trends in Europe in the period of 1990–2010, *Atmospheric
Chemistry and Physics*, 22, 7207–7257, <https://doi.org/10.5194/acp-22-7207-2022>, 2022.
- Van Damme, M., Clarisse, L., Heald, C. L., Hurtmans, D., Ngadi, Y., Clerbaux, C., Dolman, A. J., Erisman, J. W., and Coheur, P. F.: Global
distributions, time series and error characterization of atmospheric ammonia (NH₃) from IASI satellite observations, *Atmospheric Chemistry
and Physics*, 14, 2905–2922, <https://doi.org/10.5194/acp-14-2905-2014>, 2014.
- 875 Van Damme, M., Whitburn, S., Clarisse, L., Clerbaux, C., Hurtmans, D., and Coheur, P.-F.: Version 2 of the IASI NH₃ neural network retrieval
algorithm: near-real-time and reanalysed datasets, *Atmospheric Measurement Techniques*, 10, 4905–4914, <https://doi.org/10.5194/amt-10-4905-2017>, 2017.
- Van Damme, M., Clarisse, L., Stavroukou, T., Wichink Kruit, R., Sellekaerts, L., Viatte, C., Clerbaux, C., and Coheur, P. F.: On the weekly cycle
of atmospheric ammonia over European agricultural hotspots, *Scientific Reports*, 12, <https://doi.org/10.1038/s41598-022-15836-w>, 2022.
- 880 Van Der Graaf, S., Dammers, E., Segers, A., Kranenburg, R., Schaap, M., Shephard, M. W., and Erisman, J. W.: Data assimilation of CrIS
NH₃ satellite observations for improving spatiotemporal NH₃ distributions in LOTOS-EUROS, *Atmospheric Chemistry and Physics*, 22,
951–972, <https://doi.org/10.5194/acp-22-951-2022>, 2022.
- van Geffen, J. H. G. M., Eskes, H. J., Boersma, K. F., Maasackers, J. D., and Veefkind, J. P.: Sentinel-5P TROPOMI ATBD
for the Total and Tropospheric NO₂ Data Products (Version 2.8.0, Report S5P-KNMI-L2-0005-RP), Technical documentation,
885 KNMI, https://sentiwiki.copernicus.eu/_attachments/1673595/S5P-KNMI-L2-0005-RP%20-%20Sentinel-5P%20TROPOMI%20ATBD%20NO2%20data%20products%202024%20-%202.8.0.pdf, released 18 Nov 2024, 2024.
- van Geffen, J., Eskes, H., Compernelle, S., Pinardi, G., Verhoelst, T., Lambert, J.-C., Sneep, M., ter Linden, M., Ludewig, A., Boersma, K. F.,
and Veefkind, J. P.: Sentinel-5P TROPOMI NO₂ retrieval: impact of version v2.2 improvements and comparisons with OMI and ground-
based data, *Atmospheric Measurement Techniques*, 15, 2037–2060, <https://doi.org/10.5194/amt-15-2037-2022>, 2022.
- 890 van Zanten, M. C., Kruit, R. J. W., Hoogerbrugge, R., der Swaluw, E. V., and van Pul, W. A.: Trends in ammonia measurements in the
Netherlands over the period 1993–2014, *Atmospheric Environment*, 148, 352–360, <https://doi.org/10.1016/j.atmosenv.2016.11.007>, 2017.
- Veefkind, J., Aben, I., McMullan, K., Förster, H., de Vries, J., Otter, G., Claas, J., Eskes, H., de Haan, J., Kleipool, Q., van Weele, M., Hasekamp,
O., Hoogeveen, R., Landgraf, J., Snel, R., Tol, P., Ingmann, P., Voors, R., Kruizinga, B., Vink, R., Visser, H., and Levelt, P.: TROPOMI on the
ESA Sentinel-5 Precursor: A GMES mission for global observations of the atmospheric composition for climate, air quality and ozone layer
895 applications, *Remote Sensing of Environment*, 120, 70–83, <https://doi.org/https://doi.org/10.1016/j.rse.2011.09.027>, the Sentinel Missions -
New Opportunities for Science, 2012.
- Verhoelst, T., Compernelle, S., Pinardi, G., Lambert, J.-C., Eskes, H. J., Eichmann, K.-U., Fjæraa, A. M., Granville, J., Niemeijer, S., Cede,
A., Tiefengraber, M., Hendrick, F., Pazmiño, A., Bais, A., Bazureau, A., Boersma, K. F., Bogner, K., Dehn, A., Donner, S., Elokhov, A.,
Gebetsberger, M., Goutail, F., Grutter de la Mora, M., Gruzdev, A., Gratsea, M., Hansen, G. H., Irie, H., Jepsen, N., Kanaya, Y., Karagkiozidis,
900 D., Kivi, R., Kreher, K., Levelt, P. F., Liu, C., Müller, M., Navarro Comas, M., PETERS, A. J. M., Pommereau, J.-P., Portafaix, T., Prados-Roman,
C., Puentedura, O., Querel, R., Remmers, J., Richter, A., Rimmer, J., Rivera Cárdenas, C., Saavedra de Miguel, L., Sinyakov, V. P., Stremme,



- W., Strong, K., Van Roozendaal, M., Veefkind, J. P., Wagner, T., Wittrock, F., Yela González, M., and Zehner, C.: Ground-based validation of the Copernicus Sentinel-5P TROPOMI NO₂ measurements with the NDACC ZSL-DOAS, MAX-DOAS and Pandora global networks, *Atmospheric Measurement Techniques*, 14, 481–510, <https://doi.org/10.5194/amt-14-481-2021>, 2021.
- 905 Vivanco, M., Bessagnet, B., Cuvelier, C., Theobald, M., Tsyro, S., Pirovano, G., Aulinger, A., Bieser, J., Calori, G., Ciarelli, G., Manders, A., Mircea, M., Aksoyoglu, S., Briganti, G., Cappelletti, A., Colette, A., Couvidat, F., D’Isidoro, M., Kranenburg, R., Meleux, F., Menut, L., Pay, M., Rouil, L., Silibello, C., Thunis, P., and Ung, A.: Joint analysis of deposition fluxes and atmospheric concentrations of inorganic nitrogen and sulphur compounds predicted by six chemistry transport models in the frame of the EURODELTAIII project, *Atmospheric Environment*, 151, 152–175, <https://doi.org/10.1016/j.atmosenv.2016.11.042>, 2017.
- 910 White, E., Shephard, M. W., Cady-Pereira, K. E., Kharol, S. K., Ford, S., Dammers, E., Chow, E., Thiessen, N., Tobin, D., Quinn, G., O’Brien, J., and Bash, J.: Accounting for Non-Detects: Application to Satellite Ammonia Observations, *Remote Sensing*, 15, 2610, <https://doi.org/10.3390/rs15102610>, 2023.
- Williams, J. E., Boersma, K. F., Le Sager, P., and Verstraeten, W. W.: The high-resolution version of TM5-MP for optimized satellite retrievals: description and validation, *Geoscientific Model Development*, 10, 721–750, <https://doi.org/10.5194/gmd-10-721-2017>, 2017.
- 915 Zara, M., Boersma, K. F., Eskes, H., van der Gon, H. D., de Arellano, J. V.-G., Krol, M., van der Swaluw, E., Schuch, W., and Velders, G. J.: Reductions in nitrogen oxides over the Netherlands between 2005 and 2018 observed from space and on the ground: Decreasing emissions and increasing O₃ indicate changing NO_x chemistry, *Atmospheric Environment: X*, 9, <https://doi.org/10.1016/j.aeaoa.2021.100104>, 2021.
- Zhao, X., Griffin, D., Fioletov, V., McLinden, C., Cede, A., Tiefengraber, M., Müller, M., Bogner, K., Strong, K., Boersma, F., Eskes, H., Davies, J., Ogyu, A., and Lee, S. C.: Assessment of the quality of TROPOMI high-spatial-resolution NO₂ data products in the Greater Toronto Area, 920 *Atmospheric Measurement Techniques*, 13, 2131–2159, <https://doi.org/10.5194/amt-13-2131-2020>, 2020.