

# Investigating the Role of Ammonia in Enhancing Secondary Organic Aerosol Formation from the Co-photooxidation of Anthropogenic and Biogenic VOCs

Yongxin Yan<sup>1</sup>, Junling Li<sup>1\*</sup>, Yufei Song<sup>1</sup>, Yushi Gong<sup>1</sup>, Shudan Wei<sup>1</sup>, Zhaolin Wang<sup>1</sup>, Haijie Zhang<sup>1</sup>,  
5 Yanqin Ren<sup>1</sup>, Maofa Ge<sup>2</sup>, and Hong Li<sup>1\*</sup>

<sup>1</sup> State Key Laboratory of Environmental Criteria and Risk Assessment, Chinese Research Academy of Environmental Sciences, Beijing, 100012, China

<sup>2</sup> Institute of Chemistry, Chinese Academy of Sciences, Beijing, 100190, China

*Correspondence to:* Hong Li (lihong@craes.org.cn), Junling Li (lijl@craes.org.cn)

10 **Abstract.** Ammonia (NH<sub>3</sub>) plays a crucial role in the complex physicochemical processes occurring in the atmosphere, but the mechanisms governing secondary organic aerosol (SOA) formation from NH<sub>3</sub>-involved interactions between mixed anthropogenic and biogenic organic compounds remain poorly understood, thereby limiting the predictive capacity for air quality and climate. Previous studies have demonstrated that NH<sub>3</sub> can alter the oxidation pathways of single aromatic hydrocarbon precursor, promoting particle formation and growth. However, its role in mixed organic precursor systems has  
15 not been systematically explored. This study aims to elucidate these mechanisms through photooxidation experiments conducted in a large outdoor photochemical smog chamber, investigating mixtures of n-heptylcyclohexane (anthropogenic) and  $\alpha$ -pinene (biogenic) under varying NH<sub>3</sub> conditions. The results indicated that NH<sub>3</sub> obviously accelerated VOC degradation and significantly contributed to SOA enhancement through facilitating nucleation and participating in particle-phase reactions in the mixed system. The presence of NH<sub>3</sub> could not only promote the generation of intermediates such as  
20 aldehydes and ketones, but also lead to an increase in SOA mass and number concentration, particularly nitrogen-containing light-absorbing substances like imidazoles. This research can provide a scientific basis for systematically assessing how NH<sub>3</sub> affects the co-oxidation of ambient anthropogenic and biogenic gases, and deepen the understanding of its role in SOA generation, particularly light-absorbing aerosols, in the AVOC-BVOC mixed system.

## 1 Introduction

25 The atmospheric photochemical oxidation of anthropogenic and biogenic volatile organic compounds (AVOCs, BVOCs) can lead to the formation of secondary organic aerosols (SOA), which have been widely considered important for atmospheric chemistry, global climate, and human health (Ziemann and Atkinson, 2012; Shrivastava et al., 2017; Liu et al., 2022). Recent laboratory experiments have demonstrated that the contribution of specific VOC species to SOA varies under different conditions, including oxidant species and concentrations, seed particles, inorganic gas, temperature, relative humidity, and  
30 radiation intensity (Zhang et al., 2019; Palm et al., 2017; Zhang et al., 2023; Li et al., 2021a). Nevertheless, the evidence that

current model predictions of SOA are generally lower than field observations suggests that significant nonlinear effects occur in the formation of SOA in the mixed VOCs systems (Hallquist et al., 2009; Xu et al., 2021). Hence, knowledge gaps still persist regarding the complex oxidation mechanisms of multiple coexisting gas- and particle-phase substances in the atmosphere, along with the properties, composition, and formation mechanisms of the resulting SOA, which poses 35 challenges for precisely evaluating the global SOA budget, origins, and climatic implications.

It is well known that ammonia ( $\text{NH}_3$ ) is a common primary alkaline pollutant in the atmosphere, primarily emitted from agricultural sources. Moreover, on-road mobile sources of  $\text{NH}_3$  are becoming increasingly important with the rise in urban automobile ownership (Sun et al., 2017), significantly contributing to the pollution levels of fine particulate matter ( $\text{PM}_{2.5}$ ) (Wen et al., 2023). Recent studies have revealed that  $\text{NH}_3$  can influence the formation and chemical composition of single 40 VOC-derived SOA (Du et al., 2023).  $\text{NH}_3$  can promote new particle formation (NPF) and particulate growth during the photo-oxidation of aromatic hydrocarbons (e.g., toluene and xylene) (Liu et al., 2021a). Additionally,  $\text{NH}_3$  directly contributes to SOA formation through acid-base reactions with organic compounds during the photo-oxidation of biogenic olefins (e.g.,  $\alpha$ -pinene) (Hao et al., 2020). Tunnel measurements show that  $\text{NH}_3$  emitted by traffic rose from 21~225 ppb at the entrance to 40~2776 ppb at the exit (Liu et al., 2014). Therefore, investigating the role of high concentration  $\text{NH}_3$  in the 45 photochemical oxidation of mixed VOCs would benefit to deepen our understanding of the complex atmospheric oxidation mechanisms.

In recent years, the relative contribution of SOA to  $\text{PM}_{2.5}$  mass concentration has significantly increased (Ming et al., 2017). Long chain alkanes have been found to not only significantly affect the SOA yield, but also play an important role in atmospheric oxidation (Donahue et al., 2009; Tang et al., 2019). Long chain alkanes in the atmosphere come from a wide 50 range of sources, especially in the actual urban atmosphere, including volatile chemicals such as coatings and exhaust emissions from motor vehicles (Fang et al., 2021; Wang et al., 2021a; Qi et al., 2019; Deng et al., 2017b). Research has shown that the amount of SOA generated by long-chain alkanes is comparable to that of polycyclic aromatic hydrocarbons and monocyclic aromatic hydrocarbons, and even more significant in some cases (Luo et al., 2019; Agrawal et al., 2008; Siegl et al., 1999; Yuan et al., 2013; Deng et al., 2017a). The photochemical reactivity and SOA generation potential of long- 55 chain alkanes are greatly influenced by their chemical structural characteristics. Cycloalkanes have the highest activity and SOA generation potential, followed by branched alkanes, and finally straight chain alkanes (Hu et al., 2022). As a mixed hydrocarbon compound with both cyclic and branched alkyl structures, n-heptylcyclohexane can serve as an important reference for studying anthropogenic SOA derived from long-chain alkanes. On a global scale, the BVOC emissions account for over 90% of the total VOCs worldwide (Shao et al., 2022; Ahlberg et al., 2017), making them the most important source 60 of atmospheric organic compounds and a significant contributor to global SOA production (Chen and Jang, 2012). Under real atmospheric conditions, when BVOCs are mixed with anthropogenic emissions, their oxidation pathways and potential for generating SOA can be altered (Donahue et al., 2006; Emanuelsson et al., 2013; John H. Offenberg et al., 2007; Voliotis et al., 2021; Li et al., 2022b). Among them,  $\alpha$ -pinene has the characteristics of high emissions, high reactivity, and high SOA contribution, making it a very important BVOCs species in the atmospheric environment (Wang et al., 2021b). At present,

65 our understanding of the mixed oxidation mechanism of biogenic olefins and long-chain alkanes, as well as their derived SOA components and properties, is still limited. Moreover, there is less research on the impact of NH<sub>3</sub> on AVOC-BVOC mixed systems, and the mechanism of SOA formation in these systems is not yet clear.

Herein, a large outdoor photochemical smog chamber simulation was conducted to explore the SOA formation from the mixed system of n-heptylcyclohexane (representing AVOCs) and  $\alpha$ -pinene (representing BVOCs) under the presence of NH<sub>3</sub>.  
70 Then an in-depth analysis of the SOA composition was conducted, further clarifying the potential chemical mechanisms. This study aims to comprehensively reveal the important enhancing role of NH<sub>3</sub> on the co-photooxidation of anthropogenic-biogenic mixed system, and furtherly improve the understanding of the mechanisms and influencing factors of SOA in the system.

## 2 Materials and Methods

### 75 2.1 Experimental conditions

The experiments were conducted in a large outdoor photochemical smog chamber located on one rooftop of the building at the Chinese Research Academy of Environmental Sciences (CRAES). It was a 56 m<sup>3</sup> Teflon chamber covered with a well-controlled enclosure, inside where three fans were fixed to ensure uniform mixing (Li et al., 2021b). Before each experiment, the chamber was purged with zero air at a flow rate of 100 L min<sup>-1</sup> for at least 24 hours to ensure cleanliness. Herein, all the  
80 experiments were performed from October to the ensuing January, and the initial conditions and experimental results could be found in Table S1. As significant organic precursors of SOA, the liquid-phase n-heptylcyclohexane, and/or  $\alpha$ -pinene with known volumes were injected into a U-shaped tube and then blown into the chamber with zero air after heating. In all experiments, the initial concentration of n-heptylcyclohexane (TCI Development Co., Ltd, Shanghai) was in the range of 49.0~119.8 ppb, while  $\alpha$ -pinene (Beijing Inno Chem Science & Technology Co., Ltd) was in the range of 43.7~86.7 ppb.  
85 With HONO as the precursor for OH radicals and NO, it was prepared by the dropwise addition of 0.2 mL 5 wt % NaNO<sub>2</sub> into 0.4 mL 30 wt % H<sub>2</sub>SO<sub>4</sub>, formed NO, NO<sub>2</sub>, and HONO from which were blown into the chamber by zero air likewise. The measured initial NO<sub>x</sub> concentration in the chamber ranged from 89.9 to 157 ppb. NH<sub>3</sub> was introduced from a 500 ppm standard gas cylinder, the concentration of which was calculated based on the chamber volume and added volume of NH<sub>3</sub>. The initial NH<sub>3</sub> concentration was in the range of 74.85~748.5 ppb. When the target species that introduced into the chamber  
90 were thoroughly mixed, the enclosure was opened, thus the photochemical reaction began. The entire photochemical reaction process of each experiment lasted about 7 h (usually from 10 a.m. to 17 p.m.), with a maximum temperature (T) of 14.4~40.7 °C and a maximum relative humidity (RH) of less than 15% inside the chamber. The sunlight was utilized as the natural light source at JNO<sub>2</sub> of 0.9~4.4 ×10<sup>-3</sup> s<sup>-1</sup>. Sampling commenced immediately upon opening of the enclosure. The first filter collected samples continuously for 3.5 hours (i.e., the first half of the experiment, designated as "former") and was  
95 then replaced with a fresh one for an additional 3.5 hours of continuous collection (i.e., the second half of the experiment, designated as "latter"). Each set of experiments was conducted at least 2 times to verify reproducibility and ensure robust

results. Under background conditions (i.e., blank measurements), the chamber air contained  $\text{SO}_2 \leq 0.5$  ppb,  $\text{NO}_x \leq 2$  ppb, and  $\text{O}_3 \leq 0.5$  ppb, while released or formed VOCs and particles were negligible ( $< 0.01 \mu\text{g m}^{-3}$ ).

## 2.2 Instrumentation

100 The collected precursor samples (n-heptylcyclohexane and  $\alpha$ -pinene) in the Tenax TA tube underwent quantitative analysis using thermal desorption gas chromatography (TD-GC, GC, 8890, Agilent, USA; TD, UNITY-xr, Germany) to obtain precursor concentrations and consumption rates within the chamber. The mass concentration, number concentration, surface mean diameter, and volume concentration of formed particles were monitored with a scanning mobility particle sizer (SMPS, model 3080, model 3081, and model 3772, TSI Inc., USA). Additionally, the qualitative analysis of organic chemical  
105 compounds, which were subjected to ultrasonic extraction from quartz films using 10 mL of methanol (20 minutes), was performed using an electrospray ionization quadrupole time-of-flight mass spectrometer (ESI-Q-ToF-MS, Bruker, Germany). Furthermore, an FTIR spectrometer (Bruker, Germany) equipped with a RTDLATGs detector and a UV-Vis light spectrometer (Hitachi, U-3900) were applied to the qualitative detection of potential functional groups in the organic components. The qualitative and quantitative analysis of eluted aldehyde-ketones from collected samples were carried out  
110 using an HPLC-UV/MS (Shimadzu, Japan). The T and RH inside the chamber were monitored using the temperature and humidity sensors (Beijing Star Sensor Technology Co., Ltd.), while the irradiance was detected with a  $\text{JNO}_2$  filter radiometer (Metcon, Germany). The consumption and generation of gaseous  $\text{SO}_2$ ,  $\text{NO}_x$  and  $\text{O}_3$  inside the chamber were monitored in real-time by online  $\text{SO}_2$  (EC 9850, Ecotech, Australia),  $\text{NO}_x$  (EC 9841, Ecotech, Australia) and  $\text{O}_3$  analyzers (EC 9830, Ecotech, Australia).

## 115 2.3 Calculation

In this work, the OH exposure during the photo-oxidation process of precursor compounds were quantified by utilizing TD-GC, as illustrated by Eq. (S1) (Mao et al., 2009):

$$OH \text{ exposure} = \frac{\ln\left[\frac{[tracer]_0}{[tracer]_t}\right]}{k}, \quad (\text{S1})$$

where k denoted the rate constant for the reaction of the tracer and the OH radical.

120 Here, the tracer referred to a precursor compound capable of reacting with OH radicals. Given the rapid photo-oxidation rate of  $\alpha$ -pinene, n-heptylcyclohexane was appointed as the trace. In experiments without n-heptylcyclohexane,  $\alpha$ -pinene served as the tracer. At the standard temperature (298 K), the reaction rate constant for OH radical with n-heptylcyclohexane was  $1.91 \times 10^{-11} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$ , and with  $\alpha$ -pinene was  $5.23 \times 10^{-11} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$  (Atkinson and Arey, 2003). Furthermore, the physical significance of OH exposure was the total OH concentration from the onset to the detection period;  
125 hence, the cumulative time was employed in the calculation of OH exposure.

The SOA yield (Y) was defined as the fraction of reactive organic gases (ROG,  $\mu\text{g m}^{-3}$ ) converted into aerosols, and it could be calculated by Eq. (S2):

$$Y = \frac{\Delta Mo}{\Delta ROG}, \quad (S2)$$

where  $\Delta Mo$  ( $\mu\text{g m}^{-3}$ ) was the maximum mass concentration of SOA, obtained by deriving the data from the Scanning  
 130 Mobility Particle Sizer (SMPS);  $\Delta ROG$  is the amount of precursor substances involved in the reaction.

The absorption of SOA generated in the experiment across the 300 to 700 nm was measured using a UV-Vis absorption spectrophotometer. The specific absorption coefficient, denoted as  $Abs_\lambda$ , was calculated based on Eq. (S3) (Liu et al., 2021b):

$$Abs_\lambda = (A_\lambda - A_{700}) \cdot \frac{V_1}{V_a \cdot L} \cdot \ln 10, \quad (S3)$$

where  $A_\lambda$  was the light absorption coefficient of SOA at the  $\lambda$  wavelength,  $A_{700}$  was the light absorption intensity background  
 135 value,  $V_1$  was the volume of methanol with dissolved aerosols,  $V_a$  was the volume of the sampled air, and  $L$  (1 cm) was the optical path length.

Based on the UV-Vis absorption spectra, the mass absorption efficiency ( $MAE_\lambda$ ;  $\text{m}^2 \text{g}^{-1}$ ) was characterized the ultraviolet absorption intensity of SOA, which was calculated according to Eq. (S4) (Liu et al., 2021b):

$$MAE_\lambda = \frac{Abs_\lambda}{M}, \quad (S4)$$

140 where  $Abs_\lambda$  ( $\text{m}^{-1}$ ) was the light absorption coefficient of SOA at the  $\lambda$  wavelength, and  $M$  ( $\mu\text{g m}^{-3}$ ) was the mass concentration of methanol-soluble organic matter.


Mass defect was defined as the difference between exact mass and nominal mass of a certain compound (Roach et al., 2011).

### 3 Results and discussion

#### 3.1 General results

145 The experiments were carried out according to the scheme of n-heptylcyclohexane + HONO (AL), n-heptylcyclohexane + HONO +  $\text{NH}_3$  (AL-N),  $\alpha$ -pinene + HONO (AP),  $\alpha$ -pinene + HONO +  $\text{NH}_3$  (AP-N), n-heptylcyclohexane +  $\alpha$ -pinene + HONO (MIX) and n-heptylcyclohexane +  $\alpha$ -pinene + HONO +  $\text{NH}_3$  (MIX-N), and the initial conditions and results were summarized in Table S1. Since consistent results from replicate experiments effectively ruled out interference from random  
 150 uncertainties, herein selected one representative dataset for presentation. In the background experiment, the mass concentration of inorganic particles formed from the photochemical reaction of  $\text{NH}_3$  with  $\text{NO}_x$  was below  $10 \mu\text{g m}^{-3}$ , and the number concentration was  $< 2000 \# \text{cm}^{-3}$  (Fig. S1(a)). This is approximately 1~2 orders of magnitude lower than the total particle concentration in the organic reaction system. Therefore, the contribution of inorganic particle formation in the organic precursor system is considered negligible. By the conclusion of the reaction, the maximum particle diameters at  
 155 varying  $\text{NH}_3$  concentration converged to statistically similar values (Fig. S1(b)). As showed in Fig.1(a), the maximum of OH exposure was in the range of  $(5.6\sim 7.9) \times 10^{10} \text{ molecules cm}^{-3} \text{ s}$  corresponding to 10.4~14.6 simulated hours, assuming a diurnal average OH concentration of  $1.5 \times 10^6 \text{ molecules cm}^{-3}$  (Mao et al., 2009; Li et al., 2024). The OH exposure continuously increased over time, leading to an enhanced atmospheric oxidation capacity. The reaction commenced with a

decline in the concentration of reactants, and the consumption rate of which seemed to be accelerated by the addition of  $\text{NH}_3$  (Fig. 1(b)). Upon illumination, the SOA mass concentration rose rapidly while  $\text{NO}_x$  levels declined. Compared with the  $\text{NH}_3$ -free system, the  $\text{NH}_3$ -containing system produced significantly higher SOA mass concentrations; moreover, the mixed organic system yielded higher SOA mass concentrations than the single one.



165 **Figure 1: (a) The OH exposure (b) Precursor concentration (c) SOA mass concentration and (d)  $\text{NO}_x$  concentration in the MIX-N (n-heptylcyclohexane +  $\alpha$ -pinene + HONO +  $\text{NH}_3$ ) experiment, MIX (n-heptylcyclohexane +  $\alpha$ -pinene + HONO) experiment, AL-N (n-heptylcyclohexane + HONO +  $\text{NH}_3$ ), AL (n-heptylcyclohexane + HONO) experiment, AP-N ( $\alpha$ -pinene + HONO +  $\text{NH}_3$ ) experiment and AP ( $\alpha$ -pinene + HONO) experiment. In Fig.1(b), the triangle represented n-heptylcyclohexane, and the circle referred to  $\alpha$ -pinene. In Fig(b) and (c), the solid and hollow symbols denoted  $\text{NH}_3$ -free and  $\text{NH}_3$ -containing experimental systems, respectively.**

170 In terms of particle size, the average SOA diameter in the MIX-N experiment was distinctly larger than that in the AL-N experiment but almost identical to that in the AP-N experiment, implying that the disparity stemmed primarily from precursor type rather than  $\text{NH}_3$ . Previous work likewise indicated that the average SOA diameter in mixed systems was governed by the highly reactive BVOC precursors (Cui et al., 2024). As showed in Fig.2, the average diameter of particles from photochemical oxidation ranged from 300 to 450 nm in the MIX-N and MIX experiment, ranged from 200 to 300 nm in the AL-N and AL experiment, and ranged from 250 to 450 nm in the AP-N and AP experiment. Obviously, the absence of

175 any appreciable shift in average diameter between  $\text{NH}_3$ -present and  $\text{NH}_3$ -free conditions further corroborated that  $\text{NH}_3$   
 exerted no significant enhancement on particle size. However, Li et al. (2018) reported that  $\text{NH}_3$  could increase SOA size-  
 growth potential; this discrepancy was attributable to their lower reactive organic compounds-to-primary  $\text{NO}_x$  concentration  
 ratio ( $\text{VOC}/\text{NO}_x$ ) ( $1.6\sim 4.9$  ppbC  $\text{ppb}^{-1}$ ). Under such high- $\text{NO}_x$  conditions, the photo-oxidation of organics preferentially  
 drove post-nucleation growth, an effect that was largely muted in our experiments with higher  $\text{VOC}/\text{NO}_x$  ( $3.1\sim 16.2$  ppbC  
 180  $\text{ppb}^{-1}$ ) (Li et al., 2018; Li et al., 2022c).





Figure 2: The number concentration distributions of particles at different diameters in the (a) MIX-N (n-heptylcyclohexane +  $\alpha$ -pinene + HONO +  $\text{NH}_3$ ) experiment, (b) MIX (n-heptylcyclohexane +  $\alpha$ -pinene + HONO) experiment, (c) AL-N (n-

185 heptylcyclohexane + HONO + NH<sub>3</sub>), (d) AL (n-heptylcyclohexane + HONO) experiment, (e) AP-N ( $\alpha$ -pinene + HONO + NH<sub>3</sub>)  
experiment and (f) AP ( $\alpha$ -pinene + HONO) experiment.

In the process of each experiment, particles were generated rapidly within the first hour of the reaction (Fig. 2). When the maximum number concentration reached approximately  $10^4 \text{ \# cm}^{-3}$ , strong particle growth appeared as a clear banana shape. Compared to the NH<sub>3</sub>-free experiments, the maximum SOA yield of the mixed organic system rose from 14.3 % to 40 % when NH<sub>3</sub> was present, indicating that the introduction of NH<sub>3</sub> led to a marked escalation in SOA production (Fig. 3(a)).  
190 Although temperature could significantly affect the partitioning process of organic compounds and lower temperatures could sensibly facilitated the formation of particles (Li et al., 2022d), the particle mass concentration exhibited a pronounced enhancement in the NH<sub>3</sub>-present systems under identical organic precursor as well as comparable temperature and VOC/NO<sub>x</sub> ratio, suggesting that NH<sub>3</sub> played a facilitative role in the particle formation aligned with other research (Xu et al., 2021). It was probably because NH<sub>3</sub> could react with certain compounds found in SOA, contributing to convert itself into  
195 nitrogen-containing organic compounds (NOC) that remained in particles (Laskin et al., 2015), such as imines and (O'brien et al., 2013). On the other hand, both nuclei coagulation and condensation caused by acid–base reactions between NH<sub>3</sub> and organic acids (such as pinic acid and pinonic acid) were responsible for the SOA growth (Xu et al., 2021).




200 **Figure 3.** The (a) SOA yield and (b) relationship between SOA formation and NH<sub>3</sub>, VOC/NO<sub>x</sub> ratio, as well as temperature (T) in the MIX-N (n-heptylcyclohexane +  $\alpha$ -pinene + HONO + NH<sub>3</sub>) experiment, MIX (n-heptylcyclohexane +  $\alpha$ -pinene + HONO)  
experiment, AL-N (n-heptylcyclohexane + HONO + NH<sub>3</sub>), AL (n-heptylcyclohexane + HONO) experiment, AP-N ( $\alpha$ -pinene + HONO + NH<sub>3</sub>) experiment and AP ( $\alpha$ -pinene + HONO) experiment. In Fig. 3(b), the temperature and SOA mass concentration  
205 here referred to the maximum value during the reaction process. The red circles referred to MIX-N experiments, and the red filled circles referred to MIX experiments. The blue circles referred to AL-N experiments, and the blue filled circles referred to AL experiments. The yellow circles referred to AP-N experiments, and the yellow filled circles referred to AP experiments. The size of the circle represented the level of temperature. The larger the circle, the higher the experimental temperature.

### 3.2 Light absorption of SOA

Identifying the chemical class of SOA was essential to elucidating how NH<sub>3</sub> influences the photo-oxidation of binary organic mixtures; to this end, we derived functional group information of the SOA by resolving the absorption spectrum. The  
210 infrared absorption spectrum featured peaks that correspond to the functional groups of organic matter in the SOA. Various

characteristic functional groups of the main products could be differentiated by identifying the positions of these peaks (Fig. 4). The peak observed at  $2923\text{ cm}^{-1}$  and  $2855\text{ cm}^{-1}$  originated from the O–H stretching vibration. The characteristic peak at  $1645\text{ cm}^{-1}$  corresponded to the C=O stretching vibrations of the carbonyl group in either aldehyde or amide. The peak around  $1054\text{ cm}^{-1}$  was considered as the O–H bending vibration or C–O stretching vibration, while the distinctive peak at 215  $1382\text{ cm}^{-1}$  represented the —NO<sub>2</sub> stretching vibration of the nitro group (Babar et al., 2017; Huang et al., 2018). These peaks were consistently found in both MIX-N, AL-N and AP-N experiments, indicating that the predominant products from the NH<sub>3</sub>-containing systems were compounds with carbonyl functionalities, along with certain NOC (Liu et al., 2015). Especially in systems containing NH<sub>3</sub>, the enhanced peak at  $1382\text{ cm}^{-1}$  suggested that the introduction of NH<sub>3</sub> led to an increased light absorption of SOA, resulting from the enhancement of NOC formation (Wang et al., 2022).

220 In the UV absorption profiles, the peak at 300 nm was found when NH<sub>3</sub> participating in the reaction, which suggested the formation of carbonyl groups (Martinez et al., 1992). However, the MAE in each system were relatively small, and the light absorption within the UV-visible spectrum were weak. This could be attributed to the absence of conjugated double bonds in the SOA derived from n-heptylcyclohexane and  $\alpha$ -pinene (Lambe et al., 2012). Notably, the latter phase of each experiment showed a more pronounced light absorption than the former phase. The MAE at 300 nm during the former half of the 225 experiments followed the sequence MIX-N > AL-N > AP-N, whereas followed MIX-N  $\approx$  AP-N > AL-N in the latter half. These observations implied that more light-absorbing substances generated as the reaction progressed. Furthermore, the relative magnitude of SOA mass concentrations can be inferred based on the MAE values. The SOA generated from the n-heptylcyclohexane and  $\alpha$ -pinene mixture contained a higher concentration of light-absorbing substances per unit mass.




230 Figure 4. Infrared absorption spectrogram of generated particles in the (a) MIX-N (n-heptylcyclohexane +  $\alpha$ -pinene + HONO +  $\text{NH}_3$ ) experiment, (b) AL-N (n-heptylcyclohexane + HONO +  $\text{NH}_3$ ), (c) AP-N ( $\alpha$ -pinene + HONO +  $\text{NH}_3$ ) experiment and ultraviolet absorption spectrogram of generated particles in the (d) MIX-N (n-heptylcyclohexane +  $\alpha$ -pinene + HONO +  $\text{NH}_3$ ) experiment, (e) AL-N (n-heptylcyclohexane + HONO +  $\text{NH}_3$ ), (f) AP-N ( $\alpha$ -pinene + HONO +  $\text{NH}_3$ ) experiment. “Former” referred to the former half of the reaction process, and “Latter” referred to the latter half of the reaction process.

### 235 3.3 Chemical compositions of SOA

#### 3.3.1 Molecular composition


For a deeper examination on the characteristics of organic compounds within the particulate phase, mass spectrometry was utilized to analyse the particulate matter under both positive and negative ionization modes, as shown in Fig. 5. Under positive mode, a notable formation of trimer was observed in both  $\text{NH}_3$ -containing and  $\text{NH}_3$ -free mixed system, with  $m/z$

240 spanning ranges of 0~300 Da, 300~450 Da, and 500~550 Da. However, the signal intensity of SOA polymeric products slightly decreased in the NH<sub>3</sub>-containing mixed system, indicating that participation of NH<sub>3</sub> in photochemical reactions may have competitively interrupted the cross-reactions between n-heptylcyclohexane and  $\alpha$ -pinene and their intermediates, thereby suppressing the formation of high-molecular-weight organics to some extent.



245 **Figure 5.** Mass spectra of generated SOA under positive (+) / negative (-) mode in the (a) MIX-N (n-heptylcyclohexane +  $\alpha$ -pinene + HONO + NH<sub>3</sub>) experiment, (b) MIX (n-heptylcyclohexane +  $\alpha$ -pinene + HONO) experiment, (c) AL-N (n-heptylcyclohexane + HONO + NH<sub>3</sub>), (d) AL (n-heptylcyclohexane + HONO) experiment, (e) AP-N ( $\alpha$ -pinene + HONO + NH<sub>3</sub>) experiment and (f) AP ( $\alpha$ -pinene + HONO) experiment.

As Venn diagram displayed in the Fig. 6, the positive mode revealed a higher detection of compounds with mass spectrometry signal relative intensities exceeding 0.05 compared to the negative mode, with a limited overlap in compound detection between the two modes. This indicated a prevalence of compounds with protonatable functional groups (such as —NH—, —O—, —CH<sub>2</sub>, C=O, etc.) in the reaction systems involving n-heptylcyclohexane and  $\alpha$ -pinene, while compounds with alcoholic and carboxylic acid functionalities were relatively less abundant (Glasius, 1999, J. Chromatogr. A; Steckel and Schlosser, 2019, Molecules). In positive mode, 414 compounds were totally detected, with 234 compounds consistently present across both former and latter half of the reaction, representing over 56% of the total compounds identified (Table S2). In negative mode, 182 compounds were totally identified, with 74% of these compounds being consistently detected throughout the reaction phases. The presence of unique compounds in each stage, observed in both ionization modes, was likely due to the cross-reactions of intermediates thus the generation of new substances (Shao et al., 2022).




260 **Figure 6. Number of common discrete compounds and unique compounds in single and mixed experiments detected under positive (+) and negative (-) mode, the signal relative intensity of which were greater than 0.05. A relative intensity of 1 was assigned to the most intense peak that was non-impurity in each sample group.**

In the presence of NH<sub>3</sub> within the reaction system, the species common to both the MIX experiments and the AP experiments were 1.8 to 12 times more abundant than those common to the MIX experiments and the AL experiments. This finding was similar to the 16-fold increase reported by Shao et al. (2022) in their experiments with mixed  $\alpha$ -pinene and o-cresol, underscoring the prominence of  $\alpha$ -pinene oxidation in mixed systems of AVOC and BVOC. Conversely, in the absence of NH<sub>3</sub>, a higher number of shared compounds between the AL and MIX experiments were noted in positive mode, whereas a greater overlap was observed between the AP and MIX experiments in negative mode, indicating that the

270 participation of  $\text{NH}_3$  could promote the formation of a greater number of compounds bearing readily protonatable functional groups.

### 3.3.2 Elementary composition

An in-depth analysis of the SOA chemical composition across the systems was performed by categorizing the detected molecular formulas according to their elemental makeup, primarily into CHO and CHON groups (with C, H, O, N corresponding to the constituent atoms in the molecules), and further divided into six classes based on carbon count. The observed peak area for each compound was normalized against the total peak area of the sample, as illustrated in Fig. 7. Among molecules containing solely C, H, and O,  $\text{C}_3\sim\text{C}_5$  exhibited lower signal intensities. The MIX-N experiment predominantly comprised  $\text{C}_6\sim\text{C}_{10}$ , whereas the MIX experiment was enriched with  $\text{C}_{11}\sim\text{C}_{15}$  molecules. This served as compelling evidence that the presence of  $\text{NH}_3$  strongly promoted the termination of cross-reactions or self-oxidation of organic precursors and their intermediates, thereby suppressing the formation of high-molecular-weight compounds. For molecules that include C, H, O, and N, the majority in each system were composed of compounds with 9 to 10 carbon atoms.



**Figure 7. The normalized signal intensity distribution of different compound categories (CHO and CHON) for various single and mixed precursor systems in positive and negative mode.**

$C_6$ ~ $C_{10}$  compounds included those produced through both functionalization (addition oxygenated function group) and fragmentation (cleavage of C—C bond) pathways during  $\alpha$ -pinene oxidation (Eddingsaas et al, 2012). The main routes of the  $\alpha$ -pinene degradation were the OH addition channel and an H-abstraction channel (Capouet et al., 2004), via which particle-phase dimerization and oligomerization reactions finally occurred (e.g. alcohol + carbonyl to form hemiacetals and acetals, hydroperoxide + carbonyl to form peroxyhemiacetals and peroxyacetals, carboxylic acid + alcohol to form esters, and aldehyde self-reactions to form aldols) (Ziemann and Atkinson, 2012; Gao et al., 2004a, b), resulting in formation of large molecules ( $>C_{10}$ ) and potentially accounting for the  $C_{16}$ ~ $C_{20}$  abundance. Additionally, the gas-phase autoxidation of  $\alpha$ -pinene might contribute to the formation of CHO products with 16~20 carbon atoms (Berndt, 2021; Ehn et al., 2014).


In general, the heptyl-chain of n-heptylcyclohexane could afford it many sites of reaction without the potential of disrupting the stability of the  $C_6$  ring, and continued oxidation of branch chain would tend to eventually cause fragmentation (Capouet et al., 2004). This could account for the predominant presence of  $C_3$ ~ $C_5$  in AL experiments, which was likely associated with the oxidation of the branched structure of n- heptylcyclohexane. Obviously, the presence of  $NH_3$  led to a significant enhancement in the abundance of  $C_3$ ~ $C_5$  molecules, as  $NH_3$  could promote the partitioning of these small molecules from the gas phase into the particulate phase (Li et al., 2022a).

In the MIX experiments, products containing only CHO elements exhibited a higher abundance of  $C_{11}$  to  $C_{15}$  compounds, which could be attributed to the cross-products of n-heptylcyclohexane and  $\alpha$ -pinene oxidation and their intermediates, leading to the formation of high carbon number compounds. However, upon the addition of  $NH_3$ , it promoted the transformation of high molecular weight products into smaller ones, resulting in a decrease in the abundance of  $C_{11}$  to  $C_{15}$ . This was consistent with the previous conclusion.

### 3.3.3 Compounds composition

Furthermore, we identify SOA compounds by dividing regions based on different O/C, H/C, and OSC values. (Fig. 8). The nitrogen-number scaling was intentionally applied to visually distinguish how molecular species distributions vary across different nitrogen -number ranges, thereby potentially offering preliminary clues for subsequent analysis of  $NH_3$ -involved reaction pathways in mixed organic oxidation systems. Given that the reaction precursors did not contain aromatic compounds, naturally no aromatic hydrocarbons were formed in region A1. Clearly under the positive mode, the products of n-heptylcyclohexane and  $\alpha$ -pinene were mainly found in region A2 and B1, suggesting that the products from both single and mixed systems were primarily composed of aliphatic compounds and oxygenated organic molecules (OOMs) with more reducing functional groups. Scaling by nitrogen number allows for a more intuitive view of the NOC species distribution. Since the SOA collected in this experiment covered entire reaction period, it might contain some intermediates with unstable structures. Furthermore, the oxidation conditions in our photochemical experiments were significantly more complex than those in the ozonolysis of single  $\alpha$ -pinene. Consequently, the average H:C ratio was likely to be slightly higher than

315 reference values (Putman et al., 2012). In the presence of  $\text{NH}_3$ , there were fewer compounds in B2 than in the experiments without  $\text{NH}_3$  (Fig. S2) presenting fewer OOMs with more oxidizing functional groups (alcohols, ethers, peroxides, carbonyls) in the products. This could be attributed to the suppression effect of OOMs formation by  $\text{NH}_3$ . As for the negative mode, the products of n-heptylcyclohexane and  $\alpha$ -pinene were major detected as aliphatic compounds (Fig. S3 and S4). Furthermore, the mass defect plot revealed that products bearing a higher oxygen content were clearly detected in the absence of  $\text{NH}_3$ , whereas the maximum oxygen content of the products was markedly lower when  $\text{NH}_3$  was present, implying that  $\text{NH}_3$  could terminate the multistep oxidation of oligomeric intermediates during the reaction (Fig. S5 and S6). This also suggested that the presence of  $\text{NH}_3$  inhibited the formation of OOMs. In summary, the photo-oxidation reaction of the n-heptylcyclohexane and  $\alpha$ -pinene single / mixed systems yielded aliphatic compounds and OOMs, and the participation of  $\text{NH}_3$  in the reaction could suppress the formation of OOMs.



325

330 **Figure 8.** Van Krevelen diagram of compounds detected single / mixed experiments of n-heptylcyclohexane and  $\alpha$ -pinene under positive mode. The lower left region (A1) with low H/C and low O/C region ( $\text{O/C} \leq 0.5$ ,  $\text{H/C} \leq 1.0$ ) indicated the low oxygen aromatic hydrocarbons. The region A2 with high H/C and low O/C region ( $\text{O/C} \leq 0.5$ ,  $\text{H/C} \geq 1.5$ ) represented aliphatic compounds (Mazzoleni et al., 2010; Mazzoleni et al., 2012). OOMs contained a great fraction of reducing functional groups in the B1 region ( $\text{O/C} \geq 0.6$ ,  $\text{H/C} > 1.2$ ,  $\text{OSc} < 0$ ), oxidizing functional groups in the B2 region ( $\text{O/C} \geq 0.6$ ,  $\text{H/C} > 1.2$ ,  $\text{OSc} > 0$ ). The moderate oxidation level OOMs were in B3 region ( $\text{OSc} \geq 0$ ,  $\text{H/C} \leq 1.2$ ) (Tu et al., 2016)


### 3.4 Generation of OVOCs Intermediates

To gain a more comprehensive understanding of the mechanisms of SOA formation, we conducted an analysis of oxygenated VOCs (OVOCs) intermediates, key carbonyl compounds identified in the reaction products (Fig. 9). In the MIX experiment, hexanal peaked at  $1.1 \mu\text{g m}^{-3}$  during the former half of the reaction, and was subsequently consumed by fragmentation or further oxidation. The highest concentration in the latter half of the reaction was heptanal ( $2.3 \mu\text{g m}^{-3}$ ), which could be attributed to the fact that the heptane branch was first broken and further oxidation was required to break into smaller branches, but sufficient oxidation environment cannot be provided by largely consumed oxidants in the later stage of the reaction. In this system, the total amount of aldehydes and ketones was relatively small, presumably because the aldehydes/ketones reacted with intermediates or products and were transferred to the particle phase. When  $\text{NH}_3$  was present, the mass concentrations of all carbonyls rose sharply, indicating that  $\text{NH}_3$  either initiated cross reactions and side-reactions within the mixture or reacted with specific intermediates, driving more oxidation/fragmentation of the intermediates into aldehydes and ketones.

In the AL experiment, every carbonyl except crotonaldehyde and heptanal was more abundant in the latter half of the reaction. This might be due to the unstable structure of these two compounds, which caused their rapid oxidation to other substances. Moreover, relative to the  $\text{NH}_3$ -free case, the  $\text{NH}_3$ -present system produced substantially more aldehydes and ketones with  $\leq 4$  carbon atoms, demonstrating that  $\text{NH}_3$  enhanced carbonyl compounds formation in the system, especially that with small molecule.

In the AP experiment, compared with the  $\text{NH}_3$ -free system, every carbonyl except acetaldehyde increased when  $\text{NH}_3$  was present. The decrease of acetaldehyde presumably because  $\text{NH}_3$  reacted directly with it to form nitrogen-containing products that partitioned into the particle phase, lowering its gas-phase concentration.

In summary,  $\text{NH}_3$  redirected the oxidation pathway through competitive reactions, yielding a larger fraction of lower molecular weight products and small carbonyls.



355 **Figure 9. Mass concentration of aldehydes and ketones generated by photo oxidation in the (a) MIX-N (n-heptylcyclohexane +  $\alpha$ -pinene + HONO + NH<sub>3</sub>) experiment, (b) MIX (n-heptylcyclohexane +  $\alpha$ -pinene + HONO) experiment, (c) AL-N (n-heptylcyclohexane + HONO + NH<sub>3</sub>), (d) AL (n-heptylcyclohexane + HONO) experiment, (e) AP-N ( $\alpha$ -pinene + HONO + NH<sub>3</sub>) experiment and (f) AP ( $\alpha$ -pinene + HONO) experiment.**

### 3.5 Possible photochemical oxidation paths

360 Under high-NO<sub>x</sub> conditions, the photo-oxidation reaction of n-heptylcyclohexane and  $\alpha$ -pinene mixture began with H-abstraction by OH radical and thus reaction with O<sub>2</sub> to form an alkyl peroxy radical (RO<sub>2</sub>). When NH<sub>3</sub> co-existed with n-heptylcyclohexane,  $\alpha$ -pinene and NO<sub>x</sub>, it markedly enhanced SOA production from the photochemical oxidation of the binary mixture. Cross-reactions between n-heptylcyclohexane,  $\alpha$ -pinene and their organic intermediates promoted nucleation and growth of high-molecular-weight oligomers; in particular, peroxy radicals derived from n-heptylcyclohexane added


365 across the double bond of  $\alpha$ -pinene, yielding mixed oxidation products such as alkylated terpene oxides. Concurrently, NH<sub>3</sub> neutralised organic acids to form low-volatility ammonium salts, strongly depressing their saturation vapour pressures. NH<sub>3</sub> also reacted with OH radicals or organic radicals to generate nitrogen-containing radicals that further evolved into N-

containing molecules, again lowering volatility. Moreover,  $\text{NH}_3$  participated in Maillard chemistry with carbonyls, producing N-heterocycles (e.g. imidazoles, Fig. 10) that subsequently fragmented into smaller N-heterocycles and organic acids.

370 Existing observational evidence had indicated that the presence of  $\text{NH}_3$  could substantially enhance the formation of imidazole compounds (Liu et al., 2023). These low-volatility products underwent further multigenerational oxidation, giving highly oxygenated and nitrated species. Through these intertwined pathways  $\text{NH}_3$  substantially reduced the overall volatility of the photo-oxidation products from the n-heptylcyclohexane and  $\alpha$ -pinene mixture, efficiently enhancing SOA formation.

By reacting with organic peroxy radicals formed during oxidation,  $\text{NH}_3$  produced low volatility compounds that partitioned

375 directly into the particle phase. This partitioning interrupted self-/cross-reactions of  $\text{RO}_2\cdot$  and suppressed the formation of large OOMs. Meanwhile, multi-step oxidation initiated by  $\text{NH}_3$  reacting with carbonyls additionally generated aldehydic and ketonic intermediates, further enriching the gas-phase carbonyl compounds.



380 **Figure 10. Proposed possible reactions of the mixture experiment in the presence of  $\text{NH}_3$  ( $\text{R}_1$  was alkyl groups,  $\text{P}_{\text{N}1}$  was organic nitrate esters,  $\text{P}_{\text{N}2}$  was organic ammonium salt,  $\text{P}_{\text{N}3}$  was imidazole compounds,  $\text{P}_{\text{N}4}$  was multi-generational compounds of imidazole cleavage).**

## 4 Conclusions

In the present work, a large-scale outdoor photochemical smog chamber was utilized to study the effect of inorganic gases (NH<sub>3</sub>) on the photochemical process of mixed AVOC (n-heptylcyclohexane) and BVOC ( $\alpha$ -pinene). After the reaction begins, NH<sub>3</sub> promoted nucleation and participated in particle-phase reactions, leading to the rapid formation and growth of SOA particles within one hour, consequently accelerating precursor consumption. In contrast to previous findings, the influence of NH<sub>3</sub> on the surface mean particle size of precursor-derived SOA was relatively small, primarily because a higher VOC/NO<sub>x</sub> ratio could weaken the enhancement of new particle growth under high-NO<sub>x</sub> conditions. The primary products from experiments with mixed n-heptylcyclohexane and  $\alpha$ -pinene were gas-phase carbonyl compounds, as well as particle-phase aliphatic compounds, NOCs including imidazoles, and oxygenated organic molecules (OOMs). Reactions between NH<sub>3</sub> and carbonyls enhanced the production of aldehydes and ketones, while its acid-base reactions with organic acids, its radical reactions, and its Maillard reactions with carbonyls promoted SOA formation in the binary system. Furthermore, reactions between NH<sub>3</sub> and organic radicals terminated the cross-reactions and autoxidation of RO<sub>2</sub>, thereby suppressing OOMs formation.

This study explored the influence of NH<sub>3</sub> on photochemical oxidation reactions and products in mixed systems of cyclic branched alkanes and biogenic alkenes under high-NO<sub>x</sub> conditions, improving upon previous studies of single organic systems by better simulating real atmospheric conditions. The findings can deepen the understanding of SOA formation in regions impacted by complex emissions and further clarify the mechanisms through which NH<sub>3</sub> affects mixed oxidation reactions. This study reveals potential pathways for NH<sub>3</sub> participation in mixed oxidation reactions. Future research should conduct in-depth analyses and quantification of product characteristics to elucidate definite pathways, thereby achieving a comprehensive understanding of chem-physical processes in the real atmosphere.

In recent years, global NH<sub>3</sub> emissions have shown an overall upward trend, with agricultural emissions remaining dominant. South and East Asia (e.g., India, Bangladesh, and China) are regions with the highest global NH<sub>3</sub> concentrations, while significant concentrations have also been observed in biomass burning areas of North Africa, Central Africa, and South America (Khan et al., 2020). Because NH<sub>3</sub> influences aerosol formation and evolution in multiple ways, investigating its atmospheric behavior and accurately identifying its chemical evolution mechanisms in complex atmospheric environments can help to mitigate aerosol impacts on climate and improve the accuracy of regional air quality forecasting.

### Data availability

The data used in this study are available upon request from the corresponding author.

## 410 Acknowledgements

This work was financially supported by the National Natural Science Foundation of China (42405104), National Key Research and Development Program of China (No. 2024YFC3712902), the National Natural Science Foundation of China (42130606, 42175133, 22306179). The authors thank Mr. Chunshan Liu of Beijing Convenient Environmental Tech Co. Ltd (<http://www.bjkwnt.com/>) for their help and support in smog chamber setup.

## 415 Author contributions

J.L. conceived the study. Y.Y. performed the experiments and conducted the data analyses. Y.Y. and J.L. interpreted and discussed the data results. Y.Y. wrote the manuscript. J.L. and H.L. revised the manuscript. All authors contributed to the final paper.

## Competing interests

420 The authors declare no competing interests.

## References

- Agrawal, H., Sawant, A. A., Jansen, K., Miller, J. W., and Cocker, D. R.: Characterization of chemical and particulate emissions from aircraft engines, *Atmospheric Environment*, 42, 4380-4392, 10.1016/j.atmosenv.2008.01.069, 2008.
- 425 Ahlberg, E., Falk, J., Eriksson, A., Holst, T., Brune, W. H., Kristensson, A., Roldin, P., and Svenningsson, B.: Secondary organic aerosol from VOC mixtures in an oxidation flow reactor, *Atmospheric Environment*, 161, 210-220, 10.1016/j.atmosenv.2017.05.005, 2017.
- Atkinson, R. and Arey, J.: Atmospheric Degradation of Volatile Organic Compounds, *Chemical Reviews*, 103, 4605–4638, 2003.
- 430 Babar, Z. B., Park, J.-H., and Lim, H.-J.: Influence of NH<sub>3</sub> on secondary organic aerosols from the ozonolysis and photooxidation of  $\alpha$ -pinene in a flow reactor, *Atmospheric Environment*, 164, 71-84, 10.1016/j.atmosenv.2017.05.034, 2017.
- Capouet, M., Peeters, J., Nozière, B., and Müller, J. F.: Alpha-pinene oxidation by OH: simulations of laboratory experiments, *Atmospheric Chemistry and Physics*, 4, 2285-2311, 10.5194/acp-4-2285-2004, 2004.
- Chen, T. and Jang, M.: Secondary organic aerosol formation from photooxidation of a mixture of dimethyl sulfide and isoprene, *Atmospheric Environment*, 46, 271-278, 10.1016/j.atmosenv.2011.09.082, 2012.
- 435 Cui, Y., Chen, K., Zhang, H., Lin, Y. H., and Bahreini, R.: Chemical Composition and Optical Properties of Secondary Organic Aerosol from Photooxidation of Volatile Organic Compound Mixtures, *ACS EST Air*, 1, 247-258, 10.1021/acsestair.3c00041, 2024.
- Deng, W., Liu, T. Y., Zhang, Y. L., Situ, S. P., Hu, Q. H., He, Q. F., Zhang, Z., Lü, S. J., Bi, X. H., Wang, X. M., Boreave, A., George, C., Ding, X., and Wang, X. M.: Secondary organic aerosol formation from photo-oxidation of toluene with NO<sub>x</sub> and SO<sub>2</sub>: Chamber simulation with purified air versus urban ambient air as matrix, *Atmospheric Environment*, 150, 67-76, 10.1016/j.atmosenv.2016.11.047, 2017a.

- Deng, W., Hu, Q., Liu, T., Wang, X., Zhang, Y., Song, W., Sun, Y., Bi, X., Yu, J., Yang, W., Huang, X., Zhang, Z., Huang, Z., He, Q., Mellouki, A., and George, C.: Primary particulate emissions and secondary organic aerosol (SOA) formation from idling diesel vehicle exhaust in China, *Sci Total Environ*, 593-594, 462-469, 10.1016/j.scitotenv.2017.03.088, 2017b.
- 445 Donahue, N. M., Robinson, A. L., and Pandis, S. N.: Atmospheric organic particulate matter: From smoke to secondary organic aerosol, *Atmospheric Environment*, 43, 94-106, 10.1016/j.atmosenv.2008.09.055, 2009.
- Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environmental Science & Technology*, 40, 2635-2643, 10.1021/es052297c, 2006.
- 450 Du, L., Xu, L., Li, K., George, C., and Ge, M.: NH<sub>3</sub> Weakens the Enhancing Effect of SO<sub>2</sub> on Biogenic Secondary Organic Aerosol Formation, *Environmental Science & Technology Letters*, 10, 145-151, 10.1021/acs.estlett.2c00959, 2023.
- Emanuelsson, E. U., Hallquist, M., Kristensen, K., Glasius, M., Bohn, B., Fuchs, H., Kammer, B., Kiendler-Scharr, A., Nehr, S., Rubach, F., Tillmann, R., Wahner, A., Wu, H. C., and Mentel, T. F.: Formation of anthropogenic secondary organic aerosol (SOA) and its influence on biogenic SOA properties, *Atmospheric Chemistry and Physics*, 13, 2837-2855, 10.5194/acp-13-2837-2013, 2013.
- 455 Fang, H., Huang, X., Zhang, Y., Pei, C., Huang, Z., Wang, Y., Chen, Y., Yan, J., Zeng, J., Xiao, S., Luo, S., Li, S., Wang, J., Zhu, M., Fu, X., Wu, Z., Zhang, R., Song, W., Zhang, G., Hu, W., Tang, M., Ding, X., Bi, X., and Wang, X.: Measurement report: Emissions of intermediate-volatility organic compounds from vehicles under real-world driving conditions in an urban tunnel, *Atmospheric Chemistry and Physics*, 21, 10005-10013, 10.5194/acp-21-10005-2021, 2021.
- 460 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmospheric Chemistry and Physics*, 9, 5155-5236, 10.5194/acp-9-5155-2009, 2009.
- 465 Hao, L. Q., Kari, E., Leskinen, A., Worsnop, D. R., and Virtanen, A.: Direct contribution of ammonia to  $\alpha$ -pinene secondary organic aerosol formation, *Atmospheric Chemistry and Physics*, 20, 14393-14405, 10.5194/acp-20-14393-2020, 2020.
- Hu, W., Zhou, H., Chen, W., Ye, Y., Pan, T., Wang, Y., Song, W., Zhang, H., Deng, W., Zhu, M., Wang, C., Wu, C., Ye, C., Wang, Z., Yuan, B., Huang, S., Shao, M., Peng, Z., Day, D. A., Campuzano-Jost, P., Lambe, A. T., Worsnop, D. R., Jimenez, J. L., and Wang, X.: Oxidation Flow Reactor Results in a Chinese Megacity Emphasize the Important Contribution of S/IVOCs to Ambient SOA Formation, *Environ Sci Technol*, 56, 6880-6893, 10.1021/acs.est.1c03155, 2022.
- 470 Huang, M., Xu, J., Cai, S., Liu, X., Hu, C., Gu, X., Zhao, W., Fang, L., and Zhang, W.: Chemical analysis of particulate products of aged 1,3,5-trimethylbenzene secondary organic aerosol in the presence of ammonia, *Atmospheric Pollution Research*, 9, 146-155, 10.1016/j.apr.2017.08.003, 2018.
- John H. Offenberg, Charles W. Lweis, Michael Lewandowski, Mohammed Jaoui, Tadeusz E. Kleindienst, and Edney, E. O.: Contributions of toluene and  $\alpha$ -Pinene to SOA formed in an irradiated toluene/ $\alpha$ -pinene/NO<sub>x</sub>/ air mixture: Comparison of results using 14C content and SOA organic tracer methods, *Environmental Science & Technology* 41, 3972-3976, 2007.
- 475 Khan, M. A. H., Lowe, D., Derwent, R. G., Foulds, A., Chhantyal-Pun, R., McFiggans, G., Orr-Ewing, A. J., Percival, C. J., and Shallcross, D. E.: Global and regional model simulations of atmospheric ammonia, *Atmospheric Research*, 234, 10.1016/j.atmosres.2019.104702, 2020.
- 480 Lambe, A. T., Onasch, T. B., Croasdale, D. R., Wright, J. P., Martin, A. T., Franklin, J. P., Massoli, P., Kroll, J. H., Canagaratna, M. R., Brune, W. H., Worsnop, D. R., and Davidovits, P.: Transitions from Functionalization to Fragmentation Reactions of Laboratory Secondary Organic Aerosol (SOA) Generated from the OH Oxidation of Alkane Precursors, *Environ. Sci. Technol.*, 46, 5430-5437, 10.1021/es300274t, 2012.
- Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chemical Reviews*, 115, 4335-4382, 10.1021/cr5006167, 2015.
- 485 Li, J., Xu, W., You, B., and Sun, Y.: Dynamic variations of ammonia in various life spaces and seasons and the influences of human activities, *Building and Environment*, 212, 10.1016/j.buildenv.2022.108820, 2022a.
- Li, J., Li, K., Li, H., Wang, X., Wang, W., Wang, K., and Ge, M.: Long-chain alkanes in the atmosphere: A review, *J Environ Sci (China)*, 114, 37-52, 10.1016/j.jes.2021.07.021, 2022b.
- 490 Li, J., Li, K., Li, H., Wang, X., Wang, W., Wang, K., and Ge, M.: Long-chain alkanes in the atmosphere: A review, *Journal of Environmental Sciences*, 114, 37-52, 10.1016/j.jes.2021.07.021, 2022c.

- Li, J., Li, H., Li, K., Chen, Y., Zhang, H., Zhang, X., Wu, Z., Liu, Y., Wang, X., Wang, W., and Ge, M.: Enhanced secondary organic aerosol formation from the photo-oxidation of mixed anthropogenic volatile organic compounds, *Atmospheric Chemistry and Physics*, 21, 7773-7789, 10.5194/acp-21-7773-2021, 2021a.
- 495 Li, J., Li, H., Wang, X., Wang, W., Ge, M., Zhang, H., Zhang, X., Li, K., Chen, Y., Wu, Z., Chai, F., Meng, F., Mu, Y., Mellouki, A., Bi, F., Zhang, Y., Wu, L., and Liu, Y.: A large-scale outdoor atmospheric simulation smog chamber for studying atmospheric photochemical processes: Characterization and preliminary application, *Journal of Environmental Sciences*, 102, 185-197, 10.1016/j.jes.2020.09.015, 2021b.
- 500 Li, J., Li, K., Zhang, H., Zhang, X., Ji, Y., Chu, W., Kong, Y., Chu, Y., Ren, Y., Zhang, Y., Zhang, H., Gao, R., Wu, Z., Bi, F., Chen, X., Wang, X., Wang, W., Li, H., and Ge, M.: Effects of OH radical and SO<sub>2</sub> concentrations on photochemical reactions of mixed anthropogenic organic gases, *Atmospheric Chemistry and Physics*, 22, 10489-10504, 10.5194/acp-22-10489-2022, 2022d.
- 505 Li, K., Chen, L., White, S. J., Yu, H., Wu, X., Gao, X., Azzi, M., and Cen, K.: Smog chamber study of the role of NH<sub>3</sub> in new particle formation from photo-oxidation of aromatic hydrocarbons, *Sci Total Environ*, 619-620, 927-937, 10.1016/j.scitotenv.2017.11.180, 2018.
- Li, K., Zhang, J., Bell, D. M., Wang, T., Lamkaddam, H., Cui, T., Qi, L., Surdu, M., Wang, D., Du, L., El Haddad, I., Slowik, J. G., and Prevot, A. S. H.: Uncovering the dominant contribution of intermediate volatility compounds in secondary organic aerosol formation from biomass-burning emissions, *Natl Sci Rev*, 11, nwae014, 10.1093/nsr/nwae014, 2024.
- 510 Liu, F., Xu, T., Ng, N. L., and Lu, H.: Linking Cell Health and Reactive Oxygen Species from Secondary Organic Aerosols Exposure, *Environ. Sci. Technol.*, 57, 1039-1048, 10.1021/acs.est.2c05171, 2022.
- Liu, S., Huang, D., Wang, Y., Zhang, S., Liu, X., Wu, C., Du, W., and Wang, G.: Synergetic effects of NH<sub>3</sub> and NO<sub>x</sub> on the production and optical absorption of secondary organic aerosol formation from toluene photooxidation, *Atmospheric Chemistry and Physics*, 21, 17759-17773, 10.5194/acp-21-17759-2021, 2021a.
- 515 Liu, S., Wang, Y., Wang, G., Zhang, S., Li, D., Du, L., Wu, C., Du, W., and Ge, S.: Enhancing effect of NO<sub>2</sub> on the formation of light-absorbing secondary organic aerosols from toluene photooxidation, *Science of the Total Environment*, 794, 148714, 10.1016/j.scitotenv.2021.148714, 2021b.
- Liu, T., Wang, X., Wang, B., Ding, X., Deng, W., Lu, S., and Zhang, Y.: Emission factor of ammonia (NH<sub>3</sub>) from on-road vehicles in China: tunnel tests in urban Guangzhou, *Environmental Research Letters*, 9, 10.1088/1748-9326/9/6/064027, 2014.
- 520 Liu, X. D., Wang, H. Y., Wang, F. L., Lv, S. J., Wu, C., Zhao, Y., Zhang, S., Liu, S. J., Xu, X. B., Lei, Y. L., and Wang, G. H.: Secondary Formation of Atmospheric Brown Carbon in China Haze: Implication for an Enhancing Role of Ammonia, *Environmental Science & Technology*, 57, 11163-11172, 10.1021/acs.est.3c03948, 2023.
- Liu, Y., Liggio, J., Staebler, R., and Li, S. M.: Reactive uptake of ammonia to secondary organic aerosols: kinetics of organonitrogen formation, *Atmospheric Chemistry and Physics*, 15, 13569-13584, 10.5194/acp-15-13569-2015, 2015.
- 525 Luo, H., Jia, L., Wan, Q., An, T. C., and Wang, Y. J.: Role of liquid water in the formation of O<sub>3</sub> and SOA particles from 1,2,3-trimethylbenzene, *Atmospheric Environment*, 217, 10.1016/j.atmosenv.2019.116955, 2019.
- Mao, J., Ren, X., Brune, W. H., Olson, J. R., Crawford, J. H., Fried, A., Huey, L. G., Cohen, R. C., Heikes, B., Singh, H. B., Blake, D. R., Sachse, G. W., Diskin, G. S., Hall, S. R., and Shetter, R. E.: Airborne measurement of OH reactivity during INTEX-B, *Atmos. Chem. Phys.*, 9, 163-173, 10.5194/acp-9-163-2009, 2009.
- 530 Martinez, R. D., Buitrago, A. A., Howell, N. W., Hearn, C. H., and Joens, J. A.: THE NEAR UV ABSORPTION-SPECTRA OF SEVERAL ALIPHATIC-ALDEHYDES AND KETONES AT 300-K, *Atmospheric Environment Part a-General Topics*, 26, 785-792, 10.1016/0960-1686(92)90238-g, 1992.
- Ming, L., Jin, L., Li, J., Fu, P., Yang, W., Liu, D., Zhang, G., Wang, Z., and Li, X.: PM<sub>2.5</sub> in the Yangtze River Delta, China: Chemical compositions, seasonal variations, and regional pollution events, *Environmental Pollution*, 223, 200-212, 10.1016/j.envpol.2017.01.013, 2017.
- 535 O'Brien, R. E., Laskin, A., Laskin, J., Liu, S., Weber, R., Russell, L. M., and Goldstein, A. H.: Molecular characterization of organic aerosol using nanospray desorption/electrospray ionization mass spectrometry: CalNex 2010 field study, *Atmospheric Environment*, 68, 265-272, 10.1016/j.atmosenv.2012.11.056, 2013.
- Palm, B. B., Campuzano-Jost, P., Day, D. A., Ortega, A. M., Fry, J. L., Brown, S. S., Zarzana, K. J., Dube, W., Wagner, N. L., Draper, D. C., Kaser, L., Jud, W., Karl, T., Hansel, A., Gutiérrez-Montes, C., and Jimenez, J. L.: Secondary organic

- aerosol formation from in situ OH, O<sub>3</sub> and NO<sub>3</sub> oxidation of ambient forest air in an oxidation flow reactor, *Atmospheric Chemistry and Physics*, 17, 5331-5354, 10.5194/acp-17-5331-2017, 2017.
- Putman, A. L., Offenberg, J. H., Fisseha, R., Kundu, S., Rahn, T. A., and Mazzoleni, L. R.: Ultrahigh-resolution FT-ICR mass spectrometry characterization of  $\alpha$ -pinene ozonolysis SOA, *Atmospheric Environment*, 46, 164-172, 10.1016/j.atmosenv.2011.10.003, 2012.
- 545 Qi, L., Liu, H., Shen, X., Fu, M., Huang, F., Man, H., Deng, F., Shaikh, A. A., Wang, X., Dong, R., Song, C., and He, K.: Intermediate-Volatility Organic Compound Emissions from Nonroad Construction Machinery under Different Operation Modes, *Environ Sci Technol*, 53, 13832-13840, 10.1021/acs.est.9b01316, 2019.
- Roach, P. J., Laskin, J., and Laskin, A.: Higher-Order Mass Defect Analysis for Mass Spectra of Complex Organic Mixtures, *Anal. Chem.*, 83, 4924-4929, 10.1021/ac200654j, 2011.
- 550 Shao, Y., Voliotis, A., Du, M., Wang, Y., Pereira, K., Hamilton, J., Alfarrá, M. R., and McFiggans, G.: Chemical composition of secondary organic aerosol particles formed from mixtures of anthropogenic and biogenic precursors, *Atmospheric Chemistry and Physics*, 22, 9799-9826, 10.5194/acp-22-9799-2022, 2022.
- Shrivastava, M., Cappa, C. D., Fan, J., Goldstein, A. H., Guenther, A. B., Jimenez, J. L., Kuang, C., Laskin, A., Martin, S. T., 555 Ng, N. L., Petaja, T., Pierce, J. R., Rasch, P. J., Roldin, P., Seinfeld, J. H., Shilling, J., Smith, J. N., Thornton, J. A., Volkamer, R., Wang, J., Worsnop, D. R., Zaveri, R. A., Zelenyuk, A., and Zhang, Q.: Recent advances in understanding secondary organic aerosol: Implications for global climate forcing, *Reviews of Geophysics*, 55, 509-559, 10.1002/2016rg000540, 2017.
- Siegl, W. O., Hammerle, R. H., Herrmann, H. M., Wenclawiak, B. W., and Luers-Jongen, B.: Organic emissions profile for a light-duty diesel vehicle, *Atmospheric Environment*, 33, 797-805, 10.1016/s1352-2310(98)00209-x, 1999.
- 560 Sun, K., Tao, L., Miller, D. J., Pan, D., Golston, L. M., Zondlo, M. A., Griffin, R. J., Wallace, H. W., Leong, Y. J., Yang, M. M., Zhang, Y., Mauzerall, D. L., and Zhu, T.: Vehicle Emissions as an Important Urban Ammonia Source in the United States and China, *Environ. Sci. Technol.*, 51, 2472-2481, 10.1021/acs.est.6b02805, 2017.
- Tang, R. Z., Wang, H., Liu, Y., and Guo, S.: Constituents of Atmospheric Semi-Volatile and Intermediate Volatility Organic 565 Compounds and Their Contribution to Organic Aerosol, *Progress in Chemistry*, 31, 180-190, 10.7536/pc180431, 2019.
- Voliotis, A., Wang, Y., Shao, Y., Du, M., Bannan, T. J., Percival, C. J., Pandis, S. N., Alfarrá, M. R., and McFiggans, G.: Exploring the composition and volatility of secondary organic aerosols in mixed anthropogenic and biogenic precursor systems, *Atmospheric Chemistry and Physics*, 21, 14251-14273, 10.5194/acp-21-14251-2021, 2021.
- Wang, H., Guo, S., Yu, Y., Shen, R., Zhu, W., Tang, R., Tan, R., Liu, K., Song, K., Zhang, W., Zhang, Z., Shuai, S., Xu, H., 570 Zheng, J., Chen, S., Li, S., Zeng, L., and Wu, Z.: Secondary aerosol formation from a Chinese gasoline vehicle: Impacts of fuel (E10, gasoline) and driving conditions (idling, cruising), *Sci Total Environ*, 795, 148809, 10.1016/j.scitotenv.2021.148809, 2021a.
- WANG, K., WANG, W., LI, J., LIU, X., YANG, W., LI, J., and GE, M.: Research progress of intermediate volatility organic compounds, *Environmental Chemistry and Ecotoxicology*, 40, 10.7524/j.issn.0254-6108.2021032906, 2021b.
- 575 Wang, Y., Liu, S., and Wang, G.: Effects of NH<sub>3</sub> on the formation of SOA derived from xylene photochemical oxidation, *Journal of Atmospheric and Environmental Optics*, 17, 185-194, 2022.
- Wen, Y., Zhang, S., Wu, Y., and Hao, J.: Vehicular ammonia emissions: an underappreciated emission source in densely populated areas, *Atmospheric Chemistry and Physics*, 23, 3819-3828, 10.5194/acp-23-3819-2023, 2023.
- Xu, L., Du, L., Tsona, N. T., and Ge, M.: Anthropogenic Effects on Biogenic Secondary Organic Aerosol Formation, *Advances in Atmospheric Sciences*, 38, 1053-1084, 10.1007/s00376-020-0284-3, 2021.
- 580 Yuan, B., Hu, W. W., Shao, M., Wang, M., Chen, W. T., Lu, S. H., Zeng, L. M., and Hu, M.: VOC emissions, evolutions and contributions to SOA formation at a receptor site in eastern China, *Atmospheric Chemistry and Physics*, 13, 8815-8832, 10.5194/acp-13-8815-2013, 2013.
- Zhang, P., Chen, T., Liu, J., Liu, C., Ma, J., Ma, Q., Chu, B., and He, H.: Impacts of SO<sub>2</sub>, Relative Humidity, and Seed Acidity on Secondary Organic Aerosol Formation in the Ozonolysis of Butyl Vinyl Ether, *Environ. Sci. Technol.*, 53, 8845-8853, 10.1021/acs.est.9b02702, 2019.
- 585 Zhang, S., Du, L., Yang, Z., Tchinda, N. T., Li, J., and Li, K.: Contrasting impacts of humidity on the ozonolysis of monoterpenes: insights into the multi-generation chemical mechanism, *Atmospheric Chemistry and Physics*, 23, 10809-10822, 10.5194/acp-23-10809-2023, 2023.

590 Ziemann, P. J. and Atkinson, R.: Kinetics, products, and mechanisms of secondary organic aerosol formation, *Chemical Society Reviews*, 41, 10.1039/c2cs35122f, 2012.