

Impacts of Secondary Ice Production on the Microphysics and Dynamics of Deep Convective Clouds in Different Environments

Deepak Waman ¹, Julian Meusel ¹, Behrooz Keshtgar ¹, Gabriella Wallentin ¹, Christian Barthlott ¹, Sachin Patade ², Sonali Shete ², Thara Prabhakaran ², Romain Fievet ³, Declan Finney ⁴, Alan Blyth ⁴, and Corinna Hoose ¹

¹Institute of Meteorology and Climate Research Troposphere Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

²Indian Institute of Tropical Meteorology, Ministry of Earth Sciences, Pune, India

³Max Planck Institute for Meteorology, Hamburg, Germany

⁴National Centre for Atmospheric Science, University of Leeds, Leeds, United Kingdom

Correspondence: Deepak Waman (deepak.waman@kit.edu)

Abstract.

This study numerically investigates the impact of secondary ice production (SIP) on cloud microphysical and diabatic properties in continental and marine deep convective clouds (DCCs). Four cases are simulated using the Icosahedral Nonhydrostatic (ICON) model with a 2-moment cloud microphysics scheme at 2 1.6 km horizontal grid spacing. ICON forms secondary ice via rime splintering, fragmentation during raindrop freezing (RDF), ice-ice collision, and sublimational breakupsublimation. A more detailed RDF scheme (Phillips et al., 2018, PHIL18) is implemented and compared to the existing simpler scheme. Both schemes predict similar overall properties in the simulated DCCs, suggesting that a simpler scheme can represent RDF in numerical models. (Sullivan et al., 2018, SULL18).

In the simulated DCCs, SIP processes accurately reproduce observed ice number concentrations, with PHIL18 yielding closer agreement with observations than SULL18. SIP enhances ice numbers by 10^{-10} to 10^{-3} , decreasing (increasing) supercooled-liquid (ice) mass by 10-30%, leading to sustained upper-level glaciationand prolonged convective activity. Including SIP increases surface precipitation by 4% in marine DCCs, with no significant change in continental DCCs. SIP enhance enhances longwave absorption in the mixed-phase region and increased increases (20 % in continental and 40 % in marine DCCs) cloud radiative heating. SIP intensifies latent heating by up to 20 %, reaching 20-40 K d⁻¹ in continental and 80 K d⁻¹ in marine DCCs, from increased depositional growth of ice particles. This enhanced diabatic heating increases buoyancy in the stratiform regions, leading to a 10 % rise in mean vertical velocity, strengthening convectionstratiform ascent. In contrast, in convective regions, SIP increases condensate loading, which suppresses buoyancy, leading to similarly weaker convective ascent. These findings highlight the pivotal role of SIP in shaping the microphysical structure and dynamical behavior of deep convection, highlighting the need for its accurate representation in numerical models.

20 1 Introduction

Clouds play a crucial role in Earth’s energy balance by influencing radiative fluxes and latent heating (Lohmann et al., 2016). The ~~release of latent heat~~ **latent heat released** during cloud formation modifies atmospheric circulation and vertical energy distribution. Clouds influence both incoming shortwave (SW) and outgoing longwave (LW) radiation (Trenberth et al., 2009), affecting atmospheric heating rates at different heights. This interaction of clouds with radiation is quantified as ‘cloud radiative heating’ (CRH), which is the difference between atmospheric radiative heating rates in cloudy and clear-sky conditions, explaining how clouds locally heat or cool the atmosphere. This vertical redistribution of radiative energy depends on cloud properties (Sullivan et al., 2023; Keshtgar et al., 2024).

Deep convective clouds (DCCs) can produce heavy precipitation via convective cores and stratiform regions, while their associated long-lived and extensive, non-precipitating cirrus anvils play a crucial role in the radiation budget (Feng et al., 2011). DCCs range from small individual cells (≤ 1 km) to large ($\sim$ few hundred km) mesoscale convective systems (MCS) and can form in the tropics, subtropics, and midlatitudes (Wilcox et al., 2023). An MCS evolves through three phases: early growth, mature, and dissipation (Houze, 1982). The early growth phase is associated with individual cells that transport the condensate to higher levels and produce intense surface precipitation. In its mature stage, these cells organize into convective cores and a broad stratiform region that produces light precipitation. In the dissipation phase, both of these regions remain surrounded by persistent cirrus anvil clouds. In the tropics, MCSs significantly influence latent heating and radiative forcing (Roca et al., 2017; Wilcox et al., 2023), thereby modifying the thermodynamic structure of the atmosphere (Bouniol et al., 2016).

On a global scale, ~~the majority~~ **a significant fraction** of surface precipitation is ~~from the ice-crystal (‘cold rain’) process~~ (Lau and Wu, 2003; Field and Heymsfield, 2015) **thought to originate from the ‘ice-crystal’ process, yet the dominant pathway varies significantly by region and convective conditions** (Lau and Wu, 2003; Field and Heymsfield, 2015; Gupta et al., 2023). The ice-crystal process involves the growth of ice-crystals through ~~vapour~~ **vapor** deposition, in which water ~~vapour~~ **vapor** directly deposits onto ice-crystals or through riming, ~~in which~~ **which involves freezing of** supercooled cloud droplets and drizzle freeze upon collision with ice-crystals. These processes form larger snowflakes or heavily rimed ice particles (e.g., graupel/hail) that may melt and form cold rain.

The processes governing ice formation in clouds are also crucial for understanding their radiative properties. Ice formation in clouds can occur through 1) the homogeneous nucleation of aqueous aerosol particles (APs) and supercooled cloud droplets, which typically occur above the -38°C level. In contrast, at mixed-phase levels (between 0 and -38°C), mostly insoluble APs, known as ice nucleating particles (INPs), initiate ice through 2) heterogeneous ice nucleation (DeMott et al., 2010; Kanji et al., 2017). However, in the mixed-phase region, it is often reported that the measured ice particle number concentration (INC) is about $10^2 - 10^4$ times higher than the available active INP number concentration (Sotiropoulou et al., 2020; Lawson et al., 2015; Waman et al., 2022). This discrepancy can be explained by the multiplication processes that involve secondary ice production (SIP) from pre-existing ice (Takahashi et al., 1995; Field et al., 2017; Korolev and Leisner, 2020).

Several SIP processes have been proposed so far (Korolev and Leisner, 2020). For example,

1. The Hallett-Mossop rime-splintering process (HM, Hallett and Mossop, 1974)
- 55 2. Fragmentation of freezing raindrops or drizzle (RDF) (Phillips et al., 2018; Kleinheins et al., 2021) (Johnson and Hallett, 1968; Takahashi and Yamashita, 1977; Phillips et al., 2018; Kleinheins et al., 2021)
3. Fragmentation upon ice-ice collision (IIC) (Vardiman, 1978; Takahashi et al., 1995) (Vardiman, 1978; Takahashi et al., 1995)
4. Sublimational fragmentation of dendritic snow and graupel (SBF) (Oraltay and Hallett, 1989; Bacon et al., 1998; Deshmukh et al., 2022)

60 SIP through rime-splintering happens between -3 and -8°C , and involves ejection of ice splinters when supercooled droplets freeze upon collision with ice particles (Hallett and Mossop, 1974). The HM process has been recognized as important for rapid ice enhancement in the early growth of DCCs (Patade et al., 2016; Jackson et al., 2018; Lasher-Trapp et al., 2021; Waman et al., 2022; Han et al., 2024) Previous field observations provide direct evidence for the presence of the HM process across diverse cloud environments. For example, aircraft measurements in convective (Bower et al., 1996) and marine boundary layer (Huang et al., 2021; Järvinen et al., 2022) clouds showed that the measured INCs exceed INPs by several orders of magnitude within the HM regime. Radar measurements of tropical marine cumuli by Lasher-Trapp et al. (2016) also observed INCs consistent with active rime-splintering, supported by radar evidence of graupel presence within the HM regime. Previous modelling studies found that the HM process contributes significantly to the overall INC (e.g., Patade et al., 2016; Lasher-Trapp et al., 2021), especially during the early growth stage of DCCs (Waman et al., 2022; Han et al., 2024). Although a recent laboratory experiment by 70 Seidel et al. (2024) mimicking convective conditions found no clear fragmentation via splinter production through the HM process, the occasional observations of splinters in this experiment indicate the mechanism may remain active in natural clouds with more complex dynamics and a broader range of droplet sizes. its relevance in natural clouds remains uncertain.

Furthermore, IIC describes SIP from a collision between two ice particles (Grzegorzczuk et al., 2023; Gautam et al., 2024) (Phillips et al., 2017a; Grzegorzczuk et al., 2023; Gautam et al., 2024). Previous modelling studies (Zhao and Liu, 2021; Han et al., 2024; Yang et al., 2024; Patade et al., 2025) 75 Phillips et al., 2017b; Zhao and Liu, 2021; Han et al., 2024; Yang et al., 2024; Patade et al., 2025) suggest that IIC can account for over 60 % of secondary ice in the simulated DCCs, especially in their mature stage (Waman et al., 2022). Two parameterizations are available to represent SIP via IIC in numerical models: 1) a temperature-dependent scheme by Takahashi et al. (1995), and 2) a more extensive parameterization, based on the principle of energy conservation by Phillips et al. (2017b). The rate of IIC depends on the properties of the colliding particles, such as size, shape, density, roughness, 80 temperature, and fall velocity.

SIP via RDF occurs during the freezing of drizzle or raindrops, as evident from laboratory studies by Mason and Maybank (1960) and Keinert et al. (2020). In RDF, the internal pressure from trapped liquid causes the ice shell to break, ejecting secondary ice splinters and liquid splashes that may later freeze. Lawson et al. (2015), using aircraft observations and a modelling approach, shows that in the studied marine DCCs, parcel model simulations of tropical marine convection with warm cloud base (about 85 17°C), inferred that RDF dominates SIP between -4 and -15°C , and may facilitate rapid cloud glaciation. Furthermore, recent modelling studies of DCCs (Huang et al., 2021; Waman et al., 2022; Patade et al., 2025) reported that RDF dominates overall SIP in their early stages. contributes notably to overall SIP (Huang et al., 2021; Patade et al., 2025), particularly as cloud tops reach the -10 to

–20°C during early growth stages. Also, Waman et al. (2022) found that while the HM process dominates in clouds with warmer tops ($> -10^{\circ}\text{C}$), RDF may become increasingly important as the cloud deepens.

90 The importance of SBF in deep convective clouds, which occurs DCCs, which occur in unsaturated cloudy air (Korolev and Leisner, 2020), remains uncertain. While laboratory experiments (Oraltay and Hallett, 1989; Bacon et al., 1998; Dong et al., 1994) Previous laboratory experiments (Oraltay and Hallett, 1989; Dong et al., 1994; Bacon et al., 1998) demonstrated fragmentation of sublimating dendrites and rimed particles under relative humidity with respect to ice (RH_i) 50–90%, field observations (Korolev and Isaac, 2004) and modeling studies (Waman et al., 2022; Yang et al., 2024) found minimal contribution of SBF to total ice formation in clouds. . Based on these studies, Deshmukh et al. (2022) developed a theoretical parameterization for SBF during sublimation of dendritic snow and graupel based on the laboratory findings, though its applicability . However, field observations by Korolev and Isaac (2004) suggested that SBF contributes only marginally to SIP, consistent with recent modelling studies by Waman et al. (2022) and Yang et al. (2024) employing the SBF scheme of Deshmukh et al. (2022). Waman et al. (2022) further showed that while SBF forms little ice in convective updraft regions, it can account for the majority of SIP in downdraft and stratiform regions following IIC. Therefore, the applicability of the currently available SBF scheme (Deshmukh et al., 2022) to different natural DCCs remains to be validatedevaluated.

SIP may also occur from INP activation in transient supersaturation around a freezing drop, where the release of latent heat warms the surface of the drop, causing vapor diffusion and local supersaturation (Korolev and Leisner, 2020). Also, rapid temperature changes during riming can cause thermal shocks in ice crystals, leading to fracturing and splintering. However, these processes remain poorly understood and lack parameterizations in current numerical models.

105 SIP can alter diabatic heating in DCCs through increased growth of ice splinters via deposition and riming, which may strengthen updrafts and modify dynamics (Qu et al., 2022; Grzegorzczak et al., 2025). Recent modelling studies show that SIP can influence cloud dynamics, precipitation production, and radiative properties. Zhao and Liu (2021) reported that on a global scale, SIP increases the net cloud radiative forcing by 1.1 W m^{-2} annually via microphysical changes. Moreover, in tropical convection, Qu et al. (2022) reported that SIP produces the observed ice mass and number, enhances buoyancy, and prolongs cloud lifetime. In contrast, an idealized simulation by Grzegorzczak et al. (2023) found that SIP may lower the cloud top height without notably altering ice mass and buoyancy. These findings suggest that SIP plays a key role in the microphysics, dynamics, and radiative properties of clouds under different atmospheric conditions.

Hence, SIP can may induce positive feedback in DCCs, where increased growth of extra ice particles enhances diabatic heating and updrafts, intensifying convection. This affects can affect their development, organization, precipitation efficiency, and radiative properties. Yet, its impact on DCCs remains insufficiently well quantified across diverse environmental regimes. Previous studies have largely focused on idealized or isolated cases, leaving gaps in our understanding of how SIP modulates cloud dynamics and radiative effects in both continental and marine DCCs. Therefore, this study aims to investigate the possible impact of SIP on the dynamics, convective intensity, and radiative properties of DCCs, using simulations of three continental and one marine DCCs with the Icosahedral Nonhydrostatic (ICON, Zängl et al., 2015) model in limited area mode (LAM). Section 2 describes the observed DCCs and numerical model setup. Results are presented in Sec. 3, followed by conclusions in Sec. 4. The acronyms and symbols used are defined in Tables A1 and A2 respectively.

2 Field Details of field Campaigns and Numerical Model

The present study focuses on four different campaigns (Table 1) involving DCCs from various locations and periods. These are the three continental cases, including 1) the fourth phase Cloud Aerosol Interaction and Precipitation Enhancement Experiment (CAIPEEX-IV) over Solapur, India, 2) Deep Convective Microphysics Experiment (DCMEX) over Magdalena Mountains, New Mexico, 3) Midlatitude Continental Convective Clouds Experiment (MC3E) over Oklahoma, USA, and one marine deep convection observed during 4) Organized Convection and EarthCARE Studies over the Tropical Atlantic (ORCESTR). The selection of these cases is based on the intensity of convection, availability of high-quality observations from aircraft, satellite, and ground-based instruments (discussed below), spanning both stratiform and deep convective regions (especially in CAIPEEX, MC3E, and ORCESTR). As this study aims to assess the impacts of SIP on the microphysical and dynamical properties of DCCs in contrasting environments, the ORCESTR case (3 September 2024) is selected as the only marine deep convective case, providing a thermodynamic contrast to the three continental cases (CAIPEEX, DCMEX, MC3E).

These cases are described in the following section.

Table 1. Summary of the selected convective cases (continental cases are denoted by * and marine by ‡), the associated cloud base and top temperatures, convective available potential energy (CAPE) from ICON control runs (Figs. 1–4b), and the instruments used during the campaigns, mounted on aircraft (marked with *)and , satellite (denoted by †), and Research Vessel (RV) meteor (denoted by +). The particle sampling range used in the present study for the optical probes (Cloud-liquid and ice probes are denoted with ⊗ and ⊕ respectively) mounted on the aircraft is also shown.

Case/Region/Time	CAPE (J kg ^{−1})	Cloud base temperature (°C)	Cloud Min- imum cloud top tempera- ture (°C)	Aircraft*/ Satellite†	Cloud-liquid [⊗] Ice particle [⊕] probes
CAIPEEX* (Peninsular India, 30 September 2019, 06–20 UTC)	1210	20	−60	Beechcraft B200*	CDP [⊗] (0.002–0.05 mm) CIP [⊕] (0.4–0.93 mm) , PIP [⊕] (0.4–6 mm)
DCMEX* (New Mexico, USA 2 August 2022, 18–23 UTC)	669	5	−60	BAe-146*	CDP [⊗] (0.002–0.05 mm) 2DS [⊕] (0.1–1.28 mm) , CIP [⊕] (0.1CIP15 [⊕] (0.015– 0.93 mm) CIP100 [⊕] (0.1–6.2 mm) CPI [⊕] (0.01–2.3 mm) Nevrozov ^{⊕⊗} (0.1–2 g m ^{−3})
MC3E* (Oklahoma, USA, 11 May 15 UTC –12 May 2011 03 UTC)	1224	17	−60	Citation*	CDP [⊗] (0.002–0.05 mm) CIP [⊕] (0.2–1.5 mm) , HVPS-3 [⊕] (0.2–19.2 mm)
ORCESTRAT‡ (Tropical Atlantic, 3 September 00 UTC –4 September 2024 00 UTC)	1006	23	−60	EarthCARE† BOWTIE-RV Meteor+	CPR ^{⊗⊕} (EarthCARE) Cloud-radar retrieval ^{⊗⊕}

2.1.1 CAIPEEX

During CAIPEEX-IV, the Beechcraft B200 aircraft (Fig. 1a) measured aerosol and cloud properties, along with meteorological parameters, between cloud base and -15°C . The B200 aircraft performed multiple passes near stratus layers with embedded convection (Prabhakaran et al., 2023; Patade et al., 2025). Details of the optical probes mounted on the B200 during CAIPEEX are summarized in Table 1. The cloud droplet probe (CDP) sampled cloud droplet properties, such as cloud droplet number concentration (CDNC) and liquid water content (LWC). The cloud imaging probe (CIP) and precipitation imaging probe (PIP) measured drizzle and ice properties, including INC, whereas ice water content (IWC) is derived using the mass-size relationship from PIP's size distribution (Patade et al., 2025). The CIP was equipped with anti-shatter tips (Korolev et al., 2011), whereas the PIP data were processed following Patade et al. (2025). To minimize biases from artificial shattering (collision with probe tips), only ice particles larger than $400\text{ }\mu\text{m}$ (NI_{400}) from PIP and CIP are used to validate the simulated INC. In CAIPEEX-IV, a dual-polarization, C-band radar ($75\text{--}76^{\circ}\text{E}$, $17\text{--}18^{\circ}\text{N}$) observed the convective system (Prabhakaran et al., 2023). Concentrations of INP for immersion freezing mode were measured during CAIPEEX-IV using a Spectrometer for Ice Nuclei (SPIN). For validation of simulated active INPs, only SPIN measurements between -14 and -20°C are considered, as observed water subsaturation at lower temperatures would lead to droplet evaporation (Shete et al., 2026).

To minimize biases from artificial shattering (collision with probe tips), only ice particles larger than $400\text{ }\mu\text{m}$ (NI_{400}) from PIP and CIP are used for model validation.

The case examined here was observed on 30 September 2019 during CAIPEEX-IV over Solapur, primarily involving DCCs. The pre-convective air and dewpoint temperatures from the control run (Sec. 2.2, Table 3) are shown in Fig. 61b. For the CAIPEEX case, the lifting condensation level (LCL) is at 20°C with tops reaching up to -60°C (Fig. 1b), with convective available potential energy (CAPE) of 1210 J kg^{-1} (Table 1).

Figure 1c shows the particle habits at different cloudy levels observed by the CIP during the CAIPEEX campaign. It is seen that, in this post-monsoon convection, the warm levels ($> 0^{\circ}\text{C}$) were dominated by drizzles and raindrops of diameters between 0.1 and 0.9 mm. Also, between 0 and -10°C , the coexistence of both liquid and ice particles was observed, indicating the presence of well-developed, mixed-phase conditions. At -4°C , ice particles were chiefly needles/columns, plates, and heavily rimed graupels, suggesting possible formation of secondary ice SIP via rime-splintering. Furthermore, colder levels were highly populated with relatively large ($> 0.7\text{ mm}$ in diameter) rimed particles, highlighting the presence of both warm and cold rain processes in such clouds.




Figure 1. (a) Flight track of the B200 aircraft, (b) simulated (red line) and B200 observed (black line) vertical profile of the air temperature, simulated dew point temperature (blue line), and LCL (orange circle) at 09 UTC on 30 September 2019. In (b), the simulated temperature profiles are from the control run (Table 3) of the CAIPEEX case. Also (c) hydrometeor images by the CIP probe observed at different levels during the CAIPEEX convection.

2.1.2 DCMEX

The DCMEX was conducted during July-August 2022 over the Magdalena Mountains, New Mexico, USA, by the UK National Environment Research Council to understand the aerosols, microphysics, and dynamics involved in the development of DCCs (Finney et al., 2024). The DCMEX aimed to enhance the understanding of microphysical processes and their representation in climate models, to address uncertainties associated with cloud-radiative effects.

The DCMEX case analyzed here was observed on 02 August 2022. On this day, the measurements of aerosols and cloud microphysics were made by the UK's BAe146 aircraft between 1526 and 2000 UTC (Fig. 2a). INP spectra for immersion freezing during DCMEX were measured from aerosol samples collected on Teflon and polycarbonate filters aboard the BAe146 aircraft (Daily et al., 2024). The INC was measured by the optical probes mounted on the BAe146 aircraft (Finney et al., 2024), such as the 2-dimensional stereo probe (2DS), and CIP (ice particles larger than $100\text{ }\mu\text{m}$ (NI_{100})), cloud particle imager (CPI) probe, CIP15, and CIP100, equipped with anti-shatter tips and corrected for any possible artificial shattering (Field et al., 2006) (Table 1). From CPI, CIP15, and CIP100, only particles larger than $100\text{ }\mu\text{m}$ (NI_{100}) are included to evaluate the simulated INC. Furthermore, the CDP measured CDNC, whereas the Nevzorov hot wire probe was used to measure their LWC and IWC (Finney et al., 2024). Also, the ground-based C-band, dual-polarimetric Shared Mobile Atmospheric Research and Teaching radar, unit 1 (Finney et al., 2024) was deployed at Socorro Airport (34.022° N , 106.898° W), which provided continuous monitoring of cloud evolution before, during, and after the aircraft observations.

For the selected DCMEX case, the simulated LCL is at 5°C , and the maximum cloud tops are reaching up to -60°C , whereas the CAPE is 670 J kg^{-1} (Fig. 2b).

Figure 2c presents the particle habits observed by the CIP 2DS probe at different subzero levels for DCMEX clouds. At levels near cloud base, between 1 and -2°C , drizzles were observed. At levels colder than -2°C , the presence of irregular, large ice particles was prominent, along with small, pristine. The warmer levels (-5 and -6°C) are dominated by large, irregularly shaped particles consistent with rimed ice or frozen drops, mixed with smaller fragments and a few column-like structures. At the colder level -23°C , particles are notably smaller, with quasi-spherical shapes likely corresponding to frozen droplets and irregular ice fragments, and a general absence of large pristine habits such as plates or dendrites.




Figure 2. As in Fig. 1, but in (a) the flight path is for the BAE146 aircraft. In (b), observed air temperature (black line) is from the University of Wyoming archive [35.038° N, 106.623° W] at 00 UTC on 02 August 2022. The particle habits in (c) are from 2DS probe mounted on the BAE146 aircraft.

2.1.3 MC3E

The MC3E campaign (Jensen et al., 2016) was conducted from 22 April to 6 June 2011 over the Southern Great Plains, Oklahoma, USA, to study the complex physical processes that drive midlatitude DCCs and associated precipitation. It was jointly
190 organized by the U.S. Department of Energy Atmospheric Radiation Measurements (ARM) and the National Aeronautics and Space Administration's (NASA) Global Precipitation Measurement (GPM) mission.

During MC3E, the University of North Dakota Citation aircraft (Table 1) measured the cloud microphysical properties at levels between cloud base (17°C) and -30°C (Fig. 3a). Mounted on the Citation aircraft, the high volume precipitation spectrometer version 3 (HVPS-3) and CIP measured the IWC and INC, whereas the CDP measured the LWC and CDNC. During
195 MC3E, the HVPS-3 was equipped with anti-shattering tips (Korolev et al., 2011). The possible bias due to artificial shattering of ice particles is minimized by considering ice concentrations of particles larger than $200\text{ }\mu\text{m}$ (NI_{200}). Additionally, the simulated radar reflectivity is validated against the reflectivity measured by the Ka-band ARM zenith radar (KAZR, Widener et al., 2012). Since INP measurements were not available during MC3E, to validate simulated INPs, the present study uses INP concentrations measured by a continuous flow diffusion chamber (CFDC) over the study region for May 2014 (DeMott
200 et al., 2015).

Here, an MCS observed on 11 May 2011 during the MC3E campaign, characterized by relatively high CAPE (1224 J kg^{-1}), is analyzed. For this MCS, the LCL was at 17°C and the tops were extending up to -60°C (Fig. 3b, Table 1).

For the observed MCS during MC3E, Fig. 3c shows the hydrometeor habits from the CIP measurements at different levels. The levels near cloud base (17°C) were dominated by liquid particles larger than 0.2 mm , indicating the presence of smaller
205 drizzle drops or raindrops. At -5°C , the co-existence of large aggregates, with rare pristine ice crystals, and supercooled liquid was observed, favoring mixed-phase cloud processes such as riming. The presence of larger aggregates at these levels chiefly suggests stratiform cloud conditions. Also, at colder mixed-phase levels, between -16 and -22°C , hydrometeors appeared to be more irregular, with dendritic and aggregated particles observed, suggesting vigorous ice crystal growth via aggregation at these levels.




Figure 3. As in Fig. 1, but in (a) the flight path is for the Citation aircraft. In (b), the observed air temperature (black line) is from the University of Wyoming archive [35.18° N, 97.44° W] at 12 UTC on 11 May 2011.

The ORCESTRA campaign (Stevens et al., 2026), conducted over the tropical Atlantic (Fig. 4a) from 10 August to 30 September 2024, was led by the Max Planck Institute for Meteorology (MPI-M). Its main objective was to improve the understanding of the mesoscale organization of tropical convection, including its interactions with tropical waves and air–sea processes, and its impacts on climate, the Earth’s radiative budget, and tropical cyclogenesis. During ORCESTRA, aircraft and ground-based
 215 observations were carried out across the tropical Atlantic Ocean, including regions near Barbados and the Cape Verde Islands (Fig. 4b). These observations are also used to calibrate measurements from the European Space Agency and Japan Aerospace Exploration Agency (JAXA) Cloud Aerosol Radiation Explorer satellite (EarthCARE, Wehr et al., 2023).

The present study analyzes an MCS observed on 3 September 2024, during ORCESTRA (Fig. 4a). The ICON simulated air temperature profile is compared with the corresponding observations for 00 UTC on 3 September 2024 (Fig. 4b). For these
 220 clouds, the base and top temperatures were at about 23°C, and −60°C respectively, with CAPE of about 1100 1000 J kg^{−1}.

The EarthCARE (Table 1) is a sun-synchronous satellite, launched on 28 May 2024, and is equipped with the cloud profiling radar (CPR, Wehr et al., 2023)(CPR, Wehr et al., 2023; Sato et al., 2025). The main aim of EarthCARE is to study the vertical structure of clouds and aerosols and the radiation budget at the top of the atmosphere to enhance relationships between clouds, aerosols, and radiation, improving
 our understanding of how aerosols and clouds influence the Earth’s climate and to improve enhancing the accuracy of
 225 climate models through these observations.

The present study uses cloud properties observed by the EarthCARE CPR (Fig. 4a) to evaluate and validate the corresponding ICON simulation. Cloud properties from the EarthCARE JAXA CPR Level-2A cloud profile product (CPR_CLP_2A, baseline Bb, processor version 1.1) are used to evaluate the ICON simulation for the ORCESTRA case (Fig. 4a). The CPR was jointly developed by JAXA and the National Institute of Information and Communications Technology, Japan, and operates at a high frequency of 94 GHz
 230 (Wehr et al., 2023). It is designed to measure the cloud properties, moisture content, the vertical structure, and motions of the hydrometeors. The simulated radar reflectivity is compared with the EarthCARE CPR-observed reflectivity, averaged over 1 km along-track. Radar reflectivity observations from the High Altitude and Long Range (HALO) research aircraft, equipped with the MIRA cloud radar (Ka-band, 35 GHz, nadir-pointing, Ewald et al., 2019; Ewald, 2025), are also used for comparison. The HALO aircraft was operational at 13.9 km, and its flight track was closely collocated with the
 235 EarthCARE CPR overpass during 16:05–16:16 UTC on 3 September 2024. Additionally, the simulated surface precipitation is validated against the coincident observations from the GPM satellite measurements, which uses the dual-frequency radar and multi-channel radiometer. Launched in February 2014, the GPM is led by NASA and JAXA and designed to provide global observations of rain and snow.

Also, vertical profiles of LWC and IWC are obtained from Cloudnet retrievals based on cloud radar measurements
 240 aboard the Research Vessel Meteor during the Beobachtung von Ozean und Wolken - Das Trans ITCZ Experiment (BOWTIE) campaign (Foth, 2026; Stevens et al., 2026), providing additional observational constraints. Since in-situ INP measurements were unavailable during ORCESTRA, INP concentrations for immersion freezing are obtained from a

previous study by (Price et al., 2018), who measured INPs in the marine boundary layer close to the study region (Cape Verde). An EarthCARE overpass coinciding with active deep convection over the tropical Atlantic on the selected day enabled the use of CPR_CLP_2A microphysical retrievals.




Figure 4. (a) Geographical location of the entire ORCESTRa campaign (dashed black box). The yellow line in (a) shows the track of the research vessel RV Meteor during the BOWTIE campaign, the cyan line indicates the HALO aircraft track, and the red line shows the EarthCARE overpass over the selected ORCESTRa region (solid black box), and ‘x’ shows the Radiosonde location on 3 September 2024. Also shown are the vertical profiles of (b) the simulated (red line) and observed (black line, Wyoming Sonde archive [8.47° N, 29.206° W]) air and simulated dew point temperature (blue line) at 00 UTC on 03 September 2024. In (b), the orange circle denotes LCL, and the simulated temperature profiles are from the control (Table 3) run of the ORCESTRa case.

2.2 Numerical Model

The present study uses the ICON (Zängl et al., 2015) model to simulate the convective cases (Sec . 2.1). ICON is developed by the German weather service (Deutscher Wetterdienst), in collaboration with the MPI-M, German Climate Computing Center (Deutsches Klimarechenzentrum, DKRZ), Center for Climate Systems Modeling, and Karlsruhe Institute of Technology.

250 The non-hydrostatic dynamical core of ICON solves fully compressible, non-hydrostatic primitive equations (Gassmann and Herzog, 2008) on an unstructured triangular grid and with a C-type staggering based on successive refinement of a spherical icosahedron (Wan et al., 2013), to avoid singularity issues at the poles.

Here, the ICON NWP physics package is used, with a 2-moment bulk microphysics scheme (Seifert and Beheng, 2006) which follows the generalized gamma distribution to describe the particle size distributions (PSD) of hydrometeors. The

255 diameter-mass and fall velocity-mass relationships are parameterized with power laws (Han et al., 2024, Table 1 therein). In the 2-moment microphysics scheme, ICON predicts the mass and number mixing ratios of cloud droplets and ice ('crystal'), snow, graupel, hail, and rain.

ICON uses the ecRad radiation scheme (Hogan and Bozzo, 2018), where SW and LW radiative transfer are governed by the effective radii of cloud liquid and ice-crystal. In this study, these effective radii are derived from both their mass mixing ratios

260 and number concentrations, enabling a more consistent coupling between radiation and microphysics.

In ICON, the CCN activation scheme by Segal and Khain (2006) initiates cloud droplets based on aerosol properties (CCN concentration and log-normal size distribution width) and vertical velocity at the cloud base. This scheme uses regression equations and lookup tables derived from a 2000-bin microphysics parcel model, separating clouds into four regimes: maritime, continental, intermediate, and polluted continental. Furthermore, at subzero levels warmer than -38°C , ICON forms primary

265 (heterogeneous) ice by activating INP through immersion freezing and deposition nucleation (Hande et al., 2015), depending on the temperature and saturation ratio with respect to ice (S_i). [based on assumptions on aerosol composition and loading.](#) In all simulated DCCs, the immersion freezing mode of Hande et al. (2015) is modified using a temperature-dependent quadratic scaling factor constrained by observed INP concentrations, as discussed in Sec. 3. Also, above the -38°C level, ICON initiates homogeneous ice through freezing of water droplets (Jeffery and Austin, 1997; Cotton and Field, 2002)

270 and aqueous aerosols. Homogeneous freezing of raindrops is not considered, as raindrops freeze rapidly at warmer levels before reaching the threshold for homogeneous nucleation (Seifert and Beheng, 2006). The heterogeneous raindrop freezing is parameterized following Bigg (1953).

2.2.1 Secondary ice processes implemented in ICON

ICON initiates secondary ice through four ice multiplication processes. These are

275 1. Hallett-Mossop (HM) process of rime-splintering

In ICON, the rate of splinter formation during riming of supercooled cloud droplets larger than $24\text{ }\mu\text{m}$, through the HM

process, is represented as,

$$\frac{\partial N_{sec}}{\partial t} = \aleph_{hm} \times w_{hm}(T) \times q_{rime} \quad (1)$$

Here, N_{sec} is the number of secondary ice fragments from the HM process, $\aleph_{hm}$ is a constant (3.5×10^8 fragments per kg, Table A2), q_{rime} is the rimed ice mixing ratio, and $w_{hm}(T)$ is a temperature-dependent function given by,

$$w_{hm}(T) = \begin{cases} 0 & T < 265 \text{ K} \\ \left(\frac{T-265}{3} \times \frac{T-270}{-2} \right) & 265 \text{ K} < T < 270 \text{ K} \\ 0 & T > 270 \text{ K} \end{cases} \quad (2)$$

2. Fragmentation in raindrop freezing

In the ICON model, SIP is also initiated via RDF, following the parameterization by Sullivan et al. (2018, hereafter SUL18). The SUL18 Sullivan et al. (2018, hereafter SULL18). The SULL18 is a simplified scheme that forms secondary ice during the immersion freezing of raindrops, dependent on the temperature and shattering probability of a freezing raindrop. More details are from Sullivan et al. (2018) SULL18 and Han et al. (2024).

To examine how the existing parameterizations of RDF impact ice formation, dynamics, and radiative properties in the simulated clouds, a new, more detailed parameterization, proposed by Phillips et al. (2018, hereafter PHIL18), is newly implemented in ICON and compared with the predictions of SUL18SULL18.

The RDF proposed by PHIL18 forms secondary ice through two different modes. The first mode involves SIP in collisions between supercooled drops (0.05 – 5 mm diameter (D)) and a less massive ice particle or in immersion freezing of a raindrop. The number of secondary ice fragments (N_{sec}) from mode 1 is given by,

$$N_{sec} = \Xi_{RDF}(D) \times \Omega(T) \left[\frac{\zeta \eta^2}{(T - T_0)^2 + \eta^2} + \beta_{RDF} \times T \right] \forall m > m_i, D < 1.6 \text{ mm} \quad (3)$$

Here, T is temperature in $^{\circ}\text{C}$, and the quantities such as Ξ_{RDF} , β_{RDF} , ζ , and η are dependent on D , and Ω is a function of T (Phillips et al., 2018, Sec. 4 and Appendix B therein). (PHIL18, Sec. 4 and Appendix B therein).

In mode 1, both small and big fragments can be formed, denoted as ‘S’ and ‘B’ respectively, during the freezing of a raindrop. The number of big fragments, added to the graupel category, is given as ,

$$N_B = \min \left\{ \Xi_{RDF}(D) \Omega(T) \left[\frac{\zeta_B \eta_B^2}{(T - T_{B,0})^2 + \eta_B^2} \right], N_{sec} \right\} \quad (4)$$

And, the number of small fragments, identified as ice-crystals, can be derived as,

$$N_S = N_{sec} - N_B \quad (5)$$

In mode 2, secondary ice splinters are initiated via the emission and eventual freezing of splashes formed during the collision between a larger ice particle and a raindrop. The number of fragments (N_{sec}) from the second mode is given by,

$$N_{sec} = 3\Phi(T) \times [1 - f(T)] \times \max(DE - DE_{crit}, 0) \forall m < m_i \text{ and } D > 0.15 \text{ mm}. \quad (6)$$

305 Here, Φ is the probability of any drop fragment in a splash containing ice, which is set to 0.3 following James et al. (2021). And, DE is dimensionless energy defined by,

$$DE = \frac{K_0}{\gamma_{\text{liq}} \pi D^2} \quad (7)$$

and, the collision kinetic energy (CKE, K_0) is,

$$K_0 = \frac{1}{2} \frac{mm_i}{(m + m_i)} (v - v_i)^2 \quad (8)$$

310 where m and m_i are the masses of liquid and ice particles, respectively. Table A2 describes the other symbols, and additional details are from PHIL18.

3. Fragmentation in ice-ice collision

Fragment formation during collisions between ice particles is included as a SIP process in ICON following Phillips et al. (2017b). This formulation links SIP in IIC to the CKE available during collision. In ICON, this scheme is applied within
315 the bulk microphysics framework, by discretizing each colliding ice particle into emulated bins that follow a gamma size distribution (Han et al., 2024). The rime fraction is assumed to be constant and set to 0.4. The fragments from the collision between all possible pairs of bins are added together to obtain the total number of fragments from ice-ice collisions. For a collision between two ice particles, the expected number of secondary ice fragments (N_{sec}) is given as,

$$N_{\text{sec}} = \alpha A(M) \left[1 - \exp \left(- \left(\frac{CK_0}{\alpha A(M)} \right)^\gamma \right) \right] \quad (9)$$

320 Here, K_0 is the CKE (Eq. 8) of the colliding ice particles, α is the surface area of the smaller particle, γ is the shape parameter, and C is the asperity-fragility coefficient. For either ice-crystal or snow, $\gamma = 0.5 - 0.25\Psi$, where Ψ is the rime fraction given by Gautam et al., 2024 (Table 2 therein) given by,

$$\Psi = \begin{cases} 0 & \forall D < D_3 \\ \Psi_1 & \forall D > D_4 \\ \Psi_1 \left(\frac{D - D_3}{D_4 - D_3} \right) & \forall D_3 < D < D_4 \end{cases} \quad (10)$$

Here, D is the mean size of ice particle, $D_3 = 0.3 \text{ mm}$, and $D_4 = 2 \text{ mm}$ are empirical constants for particle size, Ψ_1 is the maximum rime fraction, which is 0.18 for dendritic and 0.3 for non-dendritic crystal. The other symbols are described in Table A2. In ICON, fragments are formed in collisions between ice particles of different or the same habits, except for the crystal-crystal collision. Further details of the physical assumptions and parameter choices are described in Phillips et al. (2017b, Sec. 4 and Table 1 therein) and Han et al. (2024).

4. Fragmentation in sublimation of graupel and dendritic snow

330

In ICON, an empirical formulation proposed by Deshmukh et al. (2022) is used to represent fragmentation during sublimation of dendritic snow and graupel. According to this formulation, the rate of fragment production during sublimation of dendritic snow is,

$$\frac{\partial N_{sec}}{\partial t} = \beta_{SBF} \times d^\gamma \times v' \times (100 - RH_i) \times f_v \times \Xi_{SBF} \quad \forall 261K > T > 256K \quad (11)$$

335 and, during sublimation of graupel,

$$\frac{\partial N_{sec}}{\partial t} = \left(\frac{\rho_{D0}}{\rho_r} \right) \times \beta_{SBF} \times d^\gamma \times v^* \times (100 - RH_i) \times f_v \times \Xi_{SBF} \quad \forall T \quad (12)$$

Where d is the [crystal diameter](#). [diameter of a sublimating particle](#). Following Waman et al. (2022), only 20% of secondary ice crystals from SBF are assumed to survive the subsaturated regions in which they form. Further details on the symbols are given in Table A2. Additional details are from Deshmukh et al. (2022).

340 The fragments formed in these four SIP processes are added to the cloud-ice category, allowing for a possible cascade effect. Also, the contribution to the total ice from homogeneous and heterogeneous nucleation, and the four SIP processes, is estimated using their process rates.

2.2.2 Experimental Design

In this study, each case (Sec. 2.1) is simulated in the LAM of ICON, with four two-way nested domains (Fig. 5), with the
 345 horizontal grid spacing halved from 13 km in the outermost (2500 km × 2500 km) domain to 1.6 km in the innermost (500 km × 500 km) domain. With a 21 km model top, the vertical grid spacing is gradually increased from the surface to the model top using 65 vertical levels. The model integration time step is 10 s. For the simulated DCCs, the initial and lateral boundary conditions (LBCs) are derived from the ECMWF atmospheric reanalysis version 5 (ERA5, Hersbach et al., 2020) data and updated every three hours at the lateral boundaries for the outer domain. These nested domains are coupled online, and the
 350 LBCs for the inner domain are derived from the corresponding outer domain. [In addition, all All](#) domains are simulated with the 2-moment microphysics scheme (Seifert and Beheng, 2006). For the inner two domains (3.2 and 1.6 km), shallow convection (Sakradzija et al., 2015) is parameterized, and deep convection is assumed to be explicitly resolved.




Figure 5. Outlines of the geographical areas used as domains for the simulated events of convection during (a) CAIPEEX, (b) DCMEX, (c) MC3E, and (d) ORCESTRA. The dashed red line and thick cyan lines in (d) show the EarthCARE overpass and HALO aircraft track over the study location between 1606 and 1610 UTC on 3 September 2024.

Due to their similar aerosol conditions, the simulated CAIPEEX, DCMEX, and MC3E cases are classified as ‘continental’, while ORCESTRA is categorized as ‘marine’ convection. Table 2 summarizes the details of the simulation time for each convective case. For the continental DCCs, the spin-up time is 3–6 h. In contrast, no spin-up time is considered for the ORCESTRA simulation, as ICON realistically reproduces the observed convection that was already present at initialization (Fig. 9f).

Table 2. Summary of simulation and spin-up time of the simulated DCCs.

Case	Spin-up time(h)	Simulation analysis start/end time
CAIPEEX	6 0-6 UTC, 30 Sept. 2019	06-20 6-20 UTC 30 September Sept. 2019
DCMEX	6 0-6 UTC, 30 Aug. 2022	16 6 UTC 2 August to 00 Aug. to 0 UTC 3 August Aug. 2022
MC3E	3 12-15 UTC, 11 May 2011	12 15 UTC 11 May to 12 UTC 12 May 2011
ORCESTRA	0	00 0 UTC 3 September to 00 Sept. to 0 UTC 4 September Sept. 2024

Here, the The control run (Table 3) is performed with four SIP processes (Sec. 2.2.1), with RDF represented using the SUL18 PHIL18 scheme. Also, to assess the role of SIP on cloud and radiative properties, a sensitivity run (‘No SIP’ run) is performed

in which SIP is turned off and compared with the control run. Furthermore, To examine the influence of the choice of RDF scheme on the simulated cloud properties, another sensitivity run (PHIL18 SULL18 run) is performed in which the SUL18 PHIL18 scheme is replaced with the PHIL18 SULL18 scheme.

The results discussed in the following sections are from the innermost (1.6 km) domain.

Table 3. Details of the simulations performed.

Case	Simulation analysis start/end time
Control	Includes all four SIP processes with RDF represented with the SUL18 PHIL18 scheme
No SIP	Excludes all four SIP processes
PHIL18 SULL18	Includes all four SIP processes with RDF representedwith the PHIL18 by the SULL18 scheme

3 Results from the control simulations

3.1 Model Validation

This section compares the micro- and macrophysical properties of the simulated DCCs (Sec. 2.1), with aircraft (for CAIPEEX, DCMEX, MC3E) and satellite (for ORCESTRA) observations that are temporally and spatially coincident . (supplementary Fig. S1).

3.1.1 CAIPEEX

For the simulated convection in CAIPEEX CAIPEEX DCCs, Fig. 6shows a a compares the original and scaled Hande et al. (2015) active INP concentration from immersion freezing parameterizations against the measurements from SPIN. The original scheme (gray dashed line) underestimates the observed INP concentrations by about an order of magnitude over the temperature range -14 to -20°C , while the scaled formulation (solid black line) shows improved agreement with the observations. The scaled scheme is extended beyond the observational range of the SPIN instrument (-14 to -20°C) using the derived quadratic scale factor, as indicated by the dashed black line in Fig. 6a. Figure 6b compares the cumulative frequency distribution (CDF) of simulated vertical velocity against the corresponding observations from the B200 aircraft. The simulation reasonably reproduces the observed updraft distribution, including the fractions of moderate ($w > 1 \text{ m s}^{-1}$) and strong ($w > 5 \text{ m s}^{-1}$) updrafts. However, the model underestimates the occurrence and magnitude of observed downdrafts, with maximum downdrafts reaching about 10 m s^{-1} in the observations but only 1 m s^{-1} in the control simulation. This weaker simulated downdrafts may partly be due to the limited ability of kilometer-scale simulations to resolve the turbulent structures associated with deep convection (Wang et al., 2022).

Also, Fig. 6(c-f) shows a comparison of the control simulation (Table 3) with the coincident aircraft observations, conditionally averaged over the cloudy updrafts ($w > 1 \text{ m s}^{-1}$). The simulated properties of cloud-liquid (LWC and CDNC, LWC (Fig. 6(a, b)) are c) is seen to be in good agreement with observations from the CDP, with errors less than $\pm 30\%$. Furthermore, overall, whereas the simulated CDNC is about 80% lower than the observed values (Fig. 6d). Furthermore, the simulated total INC agrees well with the coincident observations of NI_{400} from CIP and PIP (Fig. 6f), with their means differing less than $\pm 40\%$ at all sampling levels. This agreement between simulated and observed NI_{400} is consistent across different cloudy convective updraft regimes (supplementary Fig. S4). The same agreement is reported between the simulated and observed IWC (Fig. 6e).

As shown in the contoured frequency by altitude diagrams (CFADs), the simulated radar reflectivity (Fig. 6h) shows the general vertical structure of CAIPEEX clouds as observed by the C-band radar (Fig. 6g). The observed reflectivity shows a distinct bright band at about 5 km with a maximum reflectivity exceeding 25 dBZ. Although the simulation underestimates the high reflectivity values below 6 km, it adequately reproduces the overall shape and vertical distribution of reflectivity, with both showing cloud tops at similar altitudes (16 km). The median profiles show that both simulated and observed reflectivities decrease with height above the melting level (5 km), the simulated convection exhibits slightly weaker reflectivity, chiefly at higher ($> 6 \text{ km}$) altitudes, suggesting possible differences in ice-phase processes.

Also, the simulated reflectivity distribution above the freezing level in Fig. 6h is broader than observed. This can be partly attributed to C-band signal attenuation in convective cores and mixed-phase regions (Tian and Srivastava, 1997), and limitations of the bulk microphysics scheme in representing the ice particle size distribution at these levels (Ori et al., 2020).




Figure 6. Conditionally averaged predictions (minimum a) Observed INP number concentrations from Shete et al. (2026) compared against the original (dotted linesdashed gray line), mean Hande et al. (2015) immersion freezing scheme and its scaled version (solid linesblack line) and maximum fitted to match the observations. All simulations of CAIPEEX are performed with this scaled version of Hande et al. (2015) INP activation scheme. (dashed linesb) Cumulative frequency distributions comparing simulated (black line) vertical velocity with collocated B200 aircraft observations (blue line). Conditionally averaged mean profiles (black lines) from the control simulation of the CAIPEEX case, over cloudy convective updrafts ($w > 1 \text{ m s}^{-1}$) for of the (ac) liquid water content (LWC), and (bd) cloud droplet number concentration (CDNC), compared against the coincident CDP observations, and (ce) ice water content (IWC) and compared with the PIP while in (df), total ice particle number concentration (INC) NI_{400} is compared with against the corresponding observations from PIP probe (for IWC), and CIP and PIP (for NI_{400} INC) probes mounted on the B200 aircraft during CAIPEEX. In (d), only particles larger than 0.4 mm are shown from both CIP and PIP probes. Also, the simulated radar reflectivity from the control run (fh) is compared with the corresponding observations (eg) from the C-band dual polarization ground-based radar operational during CAIPEEX, averaged over 75–76°E and 17–18°N in both the simulation and the observations. Error bars and shaded regions in (c–f) denote the standard deviations of the observations and simulations, respectively. Sample sizes are shown next to the observational data points.

3.1.2 DCMEX

400 For the simulated DCMEX convection, the scaled Hande et al. (2015) immersion freezing scheme shows improved agreement with the observed (2.1.2) INP concentrations (Fig. 7a). The CDF of vertical velocity (Fig 7b) shows that the BAe146 aircraft chiefly sampled the stratiform regions of the DCMEX convection, whereas ICON simulates relatively stronger convection while still reproducing the observed fraction of strong updrafts ($w > 5 \text{ m s}^{-1}$).

Figure 7(c-f) compares the results from the control simulation of DCMEX DCCs on 2 August 2022 with BAe146 aircraft
405 observations, focusing on the cloudy updraft regions ($w > 1 \text{ m s}^{-1}$). The model simulates the magnitude of observed cloud-liquid (mean droplet sizes, LWC, CDNC) properties at levels between the cloud base (5°C) and -10°C (Fig. 7 (a, b, c, d)). However, at colder levels ($< -10^\circ\text{C}$), limited sampling size (< 10) of the CDP results in larger model-observation discrepancies ($> 50\%$).

Also, at such cold levels, the simulated IWC is about four orders an order of magnitude higher than observed by the Nevzorov probe (Fig. 7e), whereas the medians means differ by less than 50% at warmer subzero levels ($> -10^\circ\text{C}$). The same is true for the
410 INC At these levels, simulated and observed NI_{100} are in the same orders of magnitude. At colder levels ($< -10^\circ\text{C}$) in the convective updrafts, the simulated NI_{100} is about 1-2 orders of magnitude higher than the observations from different optical probes (CIP, 2DS) (Fig. 7d). Notably, the observed INC differs by more than an order of magnitude at all subzero levels. This discrepancy may partly result from f). This discrepancy persists in regions of higher convective updrafts (supplementary Fig. S4), and may reflect both ICON overestimating INC at these colder levels, and significant spatial variability in observed INC due to localized evolution
415 of individual convective turrets, especially below -15°C level, as well as significant uncertainties in the in-situ measurements, making it challenging to evaluate model performance against the observed ice properties.

In different stratiform regions (supplementary Fig. S2, S4), ICON generally reproduces the measured microphysical properties within observational uncertainty. However, discrepancies among the observed NI_{100} and INP indicate considerable uncertainty in the observed ice concentrations, particularly at levels above -15°C , which complicates accurate
420 evaluation of simulated INCs (supplementary Fig. S2d).

The CFAD (Fig. 7 (e, fg, h)) shows that the distribution of the simulated radar reflectivity agrees well with the observations from the dual-polarimetric, C-band Doppler radar, with both showing a similar cloud-top altitude of about 16 km. The maximum radar reflectivity is up to 40 dBZ, both in the simulation and observation.




Figure 7. As in Fig. 6 but for DCMEX case observed on 2 August 2022. During DCMEX, the LWC, IWC, and CDNC were measured by the Nevzorov and CDP probes, respectively mounted on the BAe146 aircraft. **The** In (a) INPs measured by Teflon and polycarbonate filters aboard the BAe-146 FAAM aircraft (Daily et al., 2024). In (f), the black line indicates mean INC of for particles larger than $100\ \mu m$ from the control simulation, and particles exceeding $0.1\ mm$ was measured by CIP from the CPI (green circles), CIP15 (blue squares), and CIP100 (brown asterisks) probes are shown, whereas 2DS (yellow diamonds) includes all particles. The radar observations are from the C-band, dual polarization, Shared Mobile Atmospheric Research and Teaching ground-based Doppler radar, unit 1, operational during DCMEX.

3.1.3 MC3E

425 For the simulated MCS, observed on 11 May 2011 during MC3E, the scaled Hande et al. (2015) scheme for immersion freezing mode agrees with the observed (Sec. 2.1.3) INP concentrations (Fig. 8a). Also, as evident from Fig.8b, the Citation aircraft primarily sampled the stratiform regions of the observed MCS, with ICON showing similar fraction (20-27%) of moderate updrafts ($w > 1 \text{ m s}^{-1}$) but relatively higher fraction of stronger updrafts ($w > 5 \text{ m s}^{-1}$).

Figure 8(c-f) presents a comparison between the simulated cloud-liquid and ice properties and those measured by the Citation aircraft, in the cloudy updraft regions ($w > 1 \text{ m s}^{-1}$). The simulated and observed properties of cloud-liquid (LWC and CDNC) are in the same orders of magnitude, with the simulation consistently exhibiting slightly higher values than those measured (Fig. 8 (a, bc, d)). Also, the simulated total INC is IWC and NI_{200} are compared with the NI_{200} corresponding observations from HPVS-v3 and CIP (Fig. 8 (c, de, f)). The simulated INC IWC and NI_{200} at observational levels reaches , with NI_{200} reaching 10 L^{-1} , comparable in magnitude to HPVS-v3 and CIP measurements. , across different convective cloudy updraft conditions (supplementary Fig. S4(j-l)).

In the control simulation of MC3E, the simulated reflectivity agrees well with the observation from the Ka-band Zenith radar, as evident in Fig. 8(e, rg, h). The model adequately predicts the overall frequency distribution and the vertical extent of the observed reflectivity. In addition, the maximum reflectivity is located below the freezing level (4.5 km), both in observation (25 dBZ) and simulation ($> 40 \text{ dBZ}$), and the reflectivity values decrease with altitude. However, the higher simulated reflectivity relative to KAZR observations can be partly attributed to Ka-band signal attenuation in heavy precipitation (Chandra et al., 2015) and the single-column sampling limitation of the vertically-pointing KAZR, whose observations are restricted to clouds passing through its narrow vertical sampling volume.




Figure 8. As in Fig. 6 but for MC3E case observed on 11 May 2011. In (a) INP observations (DeMott et al., 2015; Waman et al., 2022) are from the same month but different year (2014). During MC3E, the LWC and CDNC were measured by the CDP probes, respectively mounted on the Citation aircraft. The INC of particles larger than $0.2\ \mu m$ was measured by In (f), the black line represents mean NI_{200} from the control simulation and from both the CIP and HVPS-v3 probes. The radar observations are from the Ka-band zenith radar ($97.485^{\circ}\ W, 36.605^{\circ}\ N$) operational during MC3E.

3.1.4 ORCESTRA

For the ORCESTRA case, the immersion freezing parameterization of Hande et al. (2015) is compared and scaled against marine INP observations of Price et al. (2018) (Sec. 2.1.4, Fig. 9a). The CDF of vertical velocity (Fig. 9b) indicates that ICON underestimates the frequency of both moderate ($w > 1 \text{ m s}^{-1}$) and stronger ($w > 5 \text{ m s}^{-1}$) updrafts, with simulated moderate updraft fraction of only 0.3% compared to observations (23%, Fig. 9b). This discrepancy can be attributed to both weaker simulated convection in this marine case and the limitations of CPR Doppler retrieval of vertical motion in precipitating clouds (Kim et al., 2025). The observed microphysical and radar properties are derived from the CPR onboard the EarthCARE satellite, the airborne HALO-MIRA cloud radar, and shipborne measurements from the RV Meteor during the BOWTIE campaign (Sec. 2.1, Fig. 9 (a-c, f-c-e, g-h)). Also, the simulated total precipitation is compared with the measurements from the GPM. The simulated surface precipitation is validated against GPM measurements (Fig. 9d-f).

From these profiles, it is evident that ICON predicts the observed vertical distribution of the LWC (Fig. 9a-c) and total IWC (Fig. 9b-d) with reasonable accuracy. Also, the simulated total INC agrees well with the CPR observations at most altitudes (Fig. 9c-e). However, between -20 and -40°C levels, ICON predicts the total INC that is up to two orders of magnitude higher than those from the CPR. This discrepancy is likely because ICON overpredicts due to ICON overpredicting INC at these levels as well as the limited sensitivity of the CPR to large concentrations of small ice particles, which, despite being numerous, contribute little to radar reflectivity. This results in an underestimation by CPR of the total INC in cloudy regions dominated by small ice-crystals.

Furthermore, Fig. 9d-f shows a good agreement between the temporal evolution of the domain-averaged surface precipitation between the simulation and the GPM observation. ICON adequately captures the observed peak and decrease in precipitation with time, with generally higher precipitation than the corresponding observation. This is attributed to the fact that at such a higher spatial resolution (1.6 km) than GPM (10 km), ICON captures intense convective cores that may produce larger and more localized precipitation maxima (supplementary Fig. S1)S5(d-e)).

Also, for this For this marine convection, Fig. 9(e, f-g-i) shows a CFAD comparison of the radar reflectivity between radar reflectivity from the control simulation and the CPR observations, (Fig. 9i), EarthCARE-CPR (W-band, Fig. 9g), and the airborne HALO-MIRA cloud radar (Ka-band, Fig. 9h) between 16:00 and 05–16:15 16 UTC. The CPR CFAD shows a broad distribution of reflectivity, with a mean value simulated reflectivity distribution agrees well with HALO-MIRA, with both exhibiting pronounced convective features and median values of about 10 dBZ at most vertical levels. The median of the CPR reflectivity remains consistently below 10 dBZ at all levels, mainly due to its sensitivity to a wide range of hydrometeors, including stratiform precipitation, as well as its ability to detect localized and small-scale convection.

In contrast, the CFAD from the control simulation shows a relatively broader and higher reflectivity distribution, mainly below 8 km, with a median exceeding 10 dBZ. This is chiefly because at higher spatial resolution (1.6 km), ICON can capture strong and localized convective cores that may experience signal attenuation at the 94 GHz frequency of CPR, particularly in regions of In contrast, EarthCARE-CPR shows a larger occurrence of weak echoes and relatively lower median reflectivity. Although a direct quantitative comparison is not feasible due to differences in radar frequency and attenuation characteristics, the CPR CFAD is consistently shifted toward weaker reflectivity than those of both simulation and HALO-

MIRA. This may be attributed to stronger attenuation of the W-band signal in heavy precipitation (Sasikumar et al., 2025). The simulation, therefore, may exhibit pronounced convective features, whereas CPR measurements may show stratiform features. Despite these differences, the overall vertical structure of the simulated reflectivity compares adequately with the CPR observations, with both showing a gradual decrease in reflectivity with altitude and maximum frequency between the lower and mid-troposphere. Notably, ICON shows the observed stratiform and embedded This is particularly evident below 4 km, where HALO-MIRA detects substantially higher echoes (exceeding 20 dBZ) associated with precipitation, while the EarthCARE-CPR signal is highly attenuated at these levels. As a result, while the simulation and HALO-MIRA exhibit convective features of the ORCESTRA storm, as evident by regions of lower reflectivity (0-10 dBZ) at mid-levels (5-10 km)selected ORCESTRA convection, the EarthCARE-CPR emphasizes the broader stratiform structure.




Figure 9. (a) Observed INP number concentrations from Price et al. (2018) compared against the original (dashed gray line) Hande et al. (2015) immersion freezing scheme and its scaled version (solid black line) fitted to match the observations. (b) CDF comparing the simulated (black line) vertical velocity against the collocated observations from EarthCARE-CPR (blue line). Comparison of the simulated mean properties (mean- solid black lines, min- dotted lines, and max- dashed lines), such as (a-c) LWC, (b-d) total IWC, with coincident observations from EarthCARE CPR (c-yellow squares) and Cloudnet radar retrievals from BOWTIE campaign (blue circles). In (e), the total INC is compared with the coincident observations from EarthCARE CPR (yellow squares). Also, (d-f) the simulated domain-averaged surface precipitation rate (solid black line) is compared with the corresponding observations from the GPM (solid blue line). Furthermore, the contoured frequency by altitude diagram (CFAD) of radar reflectivity of the (e-g) EarthCARE-CPR is compared with the (h) HALO-MIRA Ka-band radar, and (i) control simulation of ORCESTRa for the period between 16:05:04 and 16:16:16 UTC. In (c-e), observational sample sizes in each temperature (for EarthCARE-CPR, marked in yellow) and altitude (RV-Meteor, marked in blue) bins are shown next to the observational data points. The simulated microphysical properties in (a-c-b-f, g-i) and radar reflectivity are plotted corresponding to the region of EarthCARE, HALO-MIRA, and RV Meteor pass, as shown in Fig. 4a.

In summary, ICON adequately reproduces the observed microphysical properties and distribution of radar reflectivity for the simulated DCCs including different SIP processes in the simulated DCCs improves the agreement between simulated and observed properties. Also, in their mixed-phase regions (0 and -38°C), the combination of various SIP processes (Sec. 2.2.1) allows ICON to simulate the observed INC, which cannot be explained by heterogeneous ice nucleation alone in these DCCs (supplementary Fig. S2 Figs. S3, S4).

3.2 Microphysical analyses of the control runs

In Fig. 10, the process rates of INCs from the primary (heterogeneous) ice and from various SIP processes are analyzed from the control runs of the simulated DCCs.

In the mixed-phase region of these DCCs, SIP in HM and RDF dominates ice formation at levels warmer than -10°C , whereas at colder levels IIC dominates the overall ice formation (Fig. 10), whereas primary . Primary ice contributes least to the total ice, especially at levels warmer than -20°C . Also, in all simulated clouds, it is seen that each SIP process exhibits a peak at different altitudes/levels. For example, IIC peaks at much colder levels (-30°C -25°C), forming about 99% of the total secondary ice there. Furthermore, RDF peaks at relatively warmer temperatures (-20°C) and forms about 1, at -3°C (in CAIPEEX, DCMEX, and ORCESTRA) and around -5°C (in MC3E), forming over 80% of the total secondary ice . at these warmer subzero levels. It is further reported that the contribution of RDF to total SIP varies with the scheme used.

The HM process is found to peak at much warmer levels (between -8 and -10°C -5 and -8°C), contributing over 3070% to the total secondary ice at these levels. Moreover, SBF produces ice splinters at a quasi-steady rate at all subzero levels warmer than -38°C , forming less than 1% of the total secondary ice. Hence, depending on their contribution to the total secondary ice, IIC can be ranked as the first, the HM process as the second, RDF as the third, and SBF as the fourth most important SIP process, as also reported by Waman et al. (2022).




Figure 10. Profiles of the number process rates of various SIP processes, such as fragmentation in raindrop freezing (RDF, dashed black brown lines), Hallett-Mossop rime-splintering (HM, solid gray blue lines), ice-ice collision (IIC, solid black lines), and sublimation (SBF, dashed gray yellow lines), averaged over convective ($w > 1 \text{ m s}^{-1}$) cloudy grid points for the control simulations of the DCCs, such as (a) CAIPEEX, (b) DCMEX, (c) MC3E, and (d) ORCESTR. Also shown is the process rate for heterogeneously nucleated ice (Primary ice, dotted gray green lines). Process rates represent time-integrated values over the simulation period, while vertical velocity corresponds to instantaneous model output.

3.3 Impact of SIP on microphysics, precipitation and dynamics of DCCs

505 To assess the role of ice multiplication (Sec. 2.2.1) in modulating the microphysical, dynamical, and radiative properties of the simulated DCCs, a sensitivity test was performed. This involved modifying the control run to create a perturbation simulation ('No SIP' run), in which all SIP processes are excluded, which is then compared with the control run.

3.3.1 Impact of SIP on microphysical properties

For the simulated CAIPEEX convection, Fig. 11 illustrates the time-height evolution of the simulated microphysical properties
510 (LWC, total IWC, and INC) in the control and No SIP simulations. Additionally, corresponding changes due to SIP (control minus No SIP) are also shown in the same figures. In the mixed-phase region of the simulated CAIPEEX these DCCs, supercooled liquid is about 40-40% higher without SIP than with it, chiefly due to decreased depositional growth of ice crystals there (Fig. 11 (a-c)). In contrast, in the mixed-phase levels in this monsoon convection, the exclusion of SIP predicts excluding SIP causes a 20% and 70% decrease in the total IWC and INC, respectively, than those with SIP (Fig. 11 (d-i)).




Figure 11. Domain averaged time-height profiles of the (a) LWC, (d) total IWC, and (g) total INC from the control simulation of CAIPEEX convection. The same information is shown from the No SIP run in (b, d, h). Furthermore, the change (control minus No SIP) in LWC, total IWC, and total INC is shown in (c), (f), and (i), respectively. Also, in these plots, the freezing and -38°C levels are denoted by the horizontal black and gray lines. In (i), the color bar scale is plotted as $\text{sign}(\Delta\text{Total INC}) \times \log(1 + |\Delta\text{Total INC}|)$, where Δ denotes change.

515 In the **lower** mixed-phase region ($> -10^{\circ}\text{C}$) of the simulated DCMEX clouds, **during convection (18-23 UTC)**, the absence of SIP causes about a 20% increase (**decrease**) in the supercooled liquid **mass**, **whereas it increases by up to 10% at higher levels** (Fig. 12(a-c)). Without SIP, the total ice mass reduces by about 30% in the mixed-phase region (**total ice**) mass (Fig. 12(a-f-f)). Furthermore, at these levels, **when present**, SIP produces as high as 10^3 L^{-1} **ice number concentration INCs** over a broad vertical extent (4.2 4 to 16 km) and a prolonged period (18-23 UTC, Fig. 12 (g-h)). In contrast, without SIP, the maximum INC is 50 L^{-1} **at**

520 **homogeneous levels**, **primarily at levels of homogeneous freezing**, with a more confined vertical extent and a relatively shorter duration of enhancement.




Figure 12. As in Fig. 11, but for DCMEX convection.

In an MCS during MC3E, the exclusion of SIP substantially alters the simulated microphysical structure. The absence of SIP in the mixed-phase regions of this MCS, excluding SIP causes over 20% decrease in MC3E predicts over 40% increase in the supercooled liquid mass for extended vertical levels, and period (11 May 12 UTC to 12 May 04 UTC) than with SIP (Fig. 13 (a-c)), whereas the total ice mass decreases by the same factor, especially in the mixed-phase region (Fig. 13 (d-f)). Additionally, in these DCCs, INC extends vertically from near the freezing level to 18 16 km, with about 3 orders of magnitude less INC in the mixed-phase region without SIP than with it (Fig. 13 (g-i)). Also, the exclusion of SIP from MC3E exhibits reduced vertical spread as well as temporal extent of these clouds. Hence, in these clouds, sustained glaciation facilitated by SIP forms a broad and long-lived anvil structure.




Figure 13. As in Fig. 11, but for MC3E convection.

In these the marine DCCs of ORCESTRA, the absence of SIP predicts excluding SIP shows over 20% more supercooled liquid above 530 5 km, while a 10% increase is seen below it (it decreases with a similar fraction below it Fig. 14(a-c)).

In contrast, excluding SIP predicts a significant . Also, the absence of SIP in this marine convection exhibits a substantial change in the distribution and magnitude of the total IWC (Fig. 14(d-f)) and INC (Fig. 14(g-i)). In the absence of SIP, at the mixed-phase levels, the region, without SIP, the total IWC is about 50% lower, mainly due to the reduced depositional growth of ice particles , than in the control run. Also At these levels, excluding SIP reduces suppresses INC by a factor of 10^2 - 10^3 . Moreover, the The presence 535 of SIP sustains the overall ice mass over time and altitude in the control run, which would otherwise rapidly diminish after the convective peak (3-9 UTC). This is because, when present, SIP allows for can be attributed to continuous multiplication of ice via various processes SIP processes (Sec. 2.2) throughout the life cycle of this marine convection, which sustains overall IWC and INC through depositional growth.




Figure 14. As in Fig. 11, but for ORCESTRA convection.

In summary, SIP significantly **impacts** **affects** the microphysical evolution of the simulated DCCs by promoting upper-level glaciation, which extends cloud lifetime and enhances both temporal longevity and spatial structure. In all simulated DCCs, excluding SIP leads to a 20% reduction in ice mass, a corresponding increase in liquid mass, and a more than tenfold decrease in ice particle numbers. This effect is most pronounced in MC3E, with up to 50% decrease (increase) in ice (liquid) mass without SIP than with it. These microphysical changes are hypothesized to influence the macrophysical and dynamical properties of these DCCs, as discussed in the following sections.

3.4 Impact of SIP on surface precipitation

In this section, the impact of SIP on the accumulated surface precipitation is discussed for the simulated DCCs.

Figure 15 illustrates domain-averaged surface accumulated precipitation in the control runs (black bars) of the simulated DCCs. It is reported that the amount of precipitation differs greatly among these clouds, with ORCESTRA representing intense tropical marine convection and producing the highest (42 43.6 mm) precipitation. In addition, MC3E forms about 6 6.8 mm surface precipitation, while CAIPEEX and DCMEX show more suppressed convective (< 4 mm) conditions.

Furthermore, in these DCCs, the impact of excluding SIP on the simulated surface precipitation is shown in Fig. 15 (gray bars). In short-lived, weakly precipitating (< 1 mm h⁻¹) DCCs of CAIPEEX (Patade et al., 2025) and DCMEX (Huihui et al., 2025), excluding SIP causes a negligible change in the surface precipitation, due to relatively weaker SIP activity. In contrast, in long-lived MCSs of MC3E and ORCESTRA, the exclusion of SIP reduces the surface precipitation by 1-5 1-4%, mainly due to reduced ice-crystal growth via deposition and riming(Fig.S3 in the supplement). .




Figure 15. Total accumulated surface precipitation from the control (black bars) and No SIP (gray bars) simulations of CAIPEEX, DCMEX, MC3E, and ORCESTRA clouds.

3.5 Impact of SIP on cloud dynamics

This section discusses dynamical properties, such as latent heating, CRH, mean vertical velocity, and cloud top altitude, from the control simulations of the DCCs. In addition, the sensitivity of these dynamical properties to SIP in the simulated DCCs is analyzed.

560 Figure 16 (left panel) illustrates the evolution of latent heating in the control runs of the simulated storm. The contour lines in Fig. 16 represent key microphysical processes contributing to total latent heating: condensation or evaporation (solid black line), vapor deposition (dashed black line), riming of cloud-liquid and rain (solid gray line), and evaporation and melting of precipitating hydrometeors (dash-dotted black line). This temporal evolution of latent heating reflects the life cycle of simulated DCCs, from initiation and growth to dissipation (Sec. 1), characterized by strong latent heating and cooling, respectively.

565 The simulated latent heating reveals strong variability in intensity and vertical extent across the simulated clouds. As convection intensifies in the control runs of CAIPEEX (between 7 and 12 UTC, Fig. 16a) and DCMEX (between 17 and 20 UTC, Fig. 16d), a pronounced heating core develops between 1-12 km, peaking at 30 K d^{-1} (in CAIPEEX) and 20 K d^{-1} (in DCMEX). In these DCCs, this heating is primarily driven by the heat release in droplet activation (1-2 km, not shown here), condensation (2-12 km), and depositional growth of ice-crystals (9-12 km), and further supported by riming at mid-levels.
570 After 12 UTC in CAIPEEX and 20 UTC in DCMEX, the heating weakens and contracts, and regions of cooling, driven by evaporation and melting, become more prominent, reaching up to -10 K d^{-1} below the freezing level.

Similarly, in its control run (Fig. 16g), MC3E convection exhibits vertically extensive and persistent latent heating, peaking at 40 K d^{-1} above 7 km between 18-23 UTC on 11 May 2011. During its developing phase (15-17 UTC), heating extends from the surface to 10 km, chiefly from nucleation and condensation (up to 8 K d^{-1}). In its mature phase (17-23 UTC), the heating
575 further strengthens to 50 K d^{-1} at 7 km, driven by condensation in the lower troposphere, and riming and deposition aloft (Fig. 16g). Notably, the anvil region ($> 10 \text{ km}$) shows sustained heating from the depositional growth of ice-crystals. After 23 UTC, as the system transitions to a stratiform regime, upper tropospheric heating weakens and spreads horizontally due to reduced deposition, while cooling due to evaporation and melting intensifies below the freezing level, indicating downdrafts and weakening convection.

580 In contrast, among the simulated DCCs, marine convection of ORCESTRA exhibits the strongest ($> 50 \text{ K d}^{-1}$, Fig. 16j) and most sustained diabatic heating. This is chiefly because, in such marine clouds, higher moisture facilitates condensation, which causes up to 80 K d^{-1} heat release at 3 km, between 3-8 UTC. Above the 6 km level, depositional growth of ice crystals dominates overall diabatic heating, with a sustained peak of 80 K d^{-1} between 4-8 UTC. In the mixed-phase region of ORCESTRA, riming (8 K d^{-1}) contributes up to 10% to total latent heating.

585 In the marine convection of ORCESTRA, abundant moisture below cloud base limits cooling due to evaporation and melting, as a shallow subsaturated layer allows precipitation to fall into already humid air. Consequently, evaporation and melting produce weaker cooling than in drier continental environments (Fig. 16 (a-i)). This likely weakens downdrafts and favors

ascending convective turrets, enhancing vertical transport of heat and moisture. Overall, the presence of extra moisture in these marine DCCs boosts latent heating and enables a more vertically extensive, organized convective system.

590 The level of peak latent heating also differs between continental and marine DCCs. In the investigated continental cases, mean diabatic heating peaks at colder temperatures ($< -38^{\circ}\text{C}$) compared to the marine case (ORCESTR), where it peaks at significantly warmer temperatures ($< -20^{\circ}\text{C}$) (supplementary Fig. s3S6). This difference is primarily due to contrasting microphysical processes driven by aerosol conditions. Continental DCCs, characterized by higher aerosol loading, form numerous small droplets that remain supercooled at colder ($< -38^{\circ}\text{C}$) levels, where they freeze homogeneously and subsequently grow
595 by vapor deposition. Conversely, marine DCCs with lower aerosol concentrations and higher moisture availability produce fewer, larger droplets that promote glaciation via enhanced collision-coalescence and subsequent raindrop/drizzle freezing. The growth of these frozen hydrometeors through riming and vapor deposition causes latent heating to peak at relatively warmer subzero levels (-20°C).

Figure 16 (b, e, h, k) shows diabatic heating profiles from No SIP runs of the simulated DCCs. In the mixed-phase regions of
600 continental DCCs, the exclusion of SIP reduces latent heating, mainly during the early growth phase, by over 10% in CAIPEEX and DCMEX, and up to 50% in MC3E. Notably, in these regions, latent heating from deposition is weaker than the cooling from sublimation, both with and without SIP (supplementary Figs. s2S6(a-f)), especially during the mature and decay phases (Figs. 16 (a-i)). However, without SIP, sublimation is more pronounced due to fewer ice particles, which reduces the surface area available for vapor deposition, than with SIP. Additionally, at these levels, the absence of SIP increases diabatic heating
605 from condensation by 10% compared to with SIP, as more moisture remains available to condense into supercooled liquid (Figs. 16 (b, e, h)). Also, in continental DCCs, excluding SIP has little effect on latent heat release during riming.

Below freezing levels in these continental DCCs, excluding SIP leads to a small (5%) increase in overall latent heating, mainly due to enhanced evaporation and melting of precipitation compared to with SIP. In contrast, at -40°C in these continental clouds, excluding SIP increases total latent heating by up to 10%, mainly due to enhanced vapor deposition (Fig. 16,
610 Fig. s3S6(a-f) in the supplement). This is mainly because, without SIP, homogeneously formed ice crystals tend to grow larger and more efficiently via vapor deposition, releasing more latent heat than with SIP.

In contrast, in the developing and mature phases (0-9 UTC) of ORCESTR marine DCCs (Fig. 16 (j, k)), excluding SIP results in up to 15% lower latent heating (Fig. 16l) compared to the control run, chiefly in the mixed-phase region. This decrease is due to a 70% reduction in depositional growth of ice particles, as fewer INC form without SIP (Figs. 16l, supplementary
615 Fig. s3S6(g, h)). However, at these levels, latent heating from condensation doubles when SIP is excluded, as more moisture is available instead of being consumed by ice-crystal growth.

In ORCESTR, excluding SIP increases supercooled liquid availability above 7 km (Fig. 14a), enhancing ice growth via riming and resulting in about 20% more diabatic heating. Also, below the freezing level in this marine convection, the exclusion of SIP causes little change ($\pm 1\%$) in total latent heating, as evaporation of precipitating hydrometeors is minimal in the
620 saturated environment.




Figure 16. Domain averaged time-height profiles of the total latent heating in the (a) control, and (b) No SIP simulations of the CAIPEEX clouds. Also, the change in latent heating from the exclusion of SIP (Control minus No SIP) for the CAIPEEX case is shown in (c). The contour lines in (a, b, c) show the contribution to the latent heating from major microphysical processes, such as condensation/evaporation (positive/negative values shown in solid black lines), deposition/sublimation (positive/negative values shown in dashed black lines), riming (dashed-dot black lines), and evaporation and melting of precipitation (solid gray lines). In (c), only major processes, such as Condensation/Evaporation, Deposition/Sublimation, and Evaporation and melting of precipitation, are shown. The other processes, such as rain freezing, homogeneous and heterogeneous nucleation, are not shown as they contribute little ($< 2\%$) to the overall latent heating. The same information is shown for the simulated (d, e, f) DCMEX, (g, h, i) MC3E, and (j, k, l) ORCESTRA clouds in their control and No SIP runs.

Figure 17 illustrates the vertical profiles of mean CRH from the control runs (solid black line) of DCCs. The maximum CRH reaches up to 3 K d^{-1} at -40°C in the continental DCCs (CAIPEEX, DCMEX, MC3E), and at -20°C in the marine case (ORCESTR), coinciding with the respective peaks in latent heating (supplementary Fig. s3S6) due to the contrasting microphysical processes described above.

- 625 The impact of SIP on CRH is assessed in Fig. 17 by comparing the control simulation with the No SIP run (solid gray line). It is reported that, in dashed lines. In the mixed-phase region (0 to -38°C), excluding SIP reduces CRH by up to 2010% in continental DCCs and 4050% in marine DCCs. This reduction is primarily attributed to lower INC (Figs. 11-14) at these levels in the absence of SIP, which weakens CRH through reduced (to depositional growth of ice particles (by up to 20%)) depositional growth of ice particles, thereby lowering the diabatic heating associated with the ice phase.
- 630 SIP exclusion exhibits the most pronounced impact on LW CRH (Fig. 17), which decreases by up to 15% (in continental DCCs) and 50% (in marine DCCs) due to reduced absorption of outgoing LW radiation. In continental DCCs, excluding SIP causes minimal change ($< 5\%$) in SW CRH. In contrast, excluding SIP from ORCESTR marine DCCs exhibits enhanced SW radiative warming (from strong cooling to weak warming), primarily due to increased incoming SW radiation through the cloud column. This is because of a significant reduction in INC (by a factor of 10^3 , supplementary Fig. s2S3(d)), which causes
- 635 optically thinner mixed-phase clouds and thereby allows, thereby allowing more SW radiation to reach and absorb at lower levels.




Figure 17. Simulated domain-averaged net cloud radiative heating (CRH, K d⁻¹, black lines) from the control (solid black lines) and No SIP (solid gray dashed lines) simulations of (a) CAIPEEX, (b) DCMEX, (c) MC3E, and (d) ORCESTRA clouds. Also shown are the SW (dashed blue lines) and LW (dotted yellow lines) components of CRH.

To quantify the impact of SIP on vertical motions, buoyancy is computed as $B = g(\theta_v - \theta_{v,mean})/\theta_{v,mean}$, where $\theta_v = \theta(1 + 0.61q_v - qt)$ is the virtual potential temperature, $\theta_{v,mean}$ is its horizontal domain mean, q_v , q_t are the vapor and total condensate mixing ratios, and g is the gravitational acceleration. Excluding SIP in the simulated DCCs causes noticeable changes in the 90th-percentile vertical velocities exhibits notable changes in buoyancy (Fig. 18), thereby influencing the overall, left panel), which in turn influences both stratiform ($0 < w < 1 \text{ m s}^{-1}$) and convective ($w > 1 \text{ m s}^{-1}$) ascent Fig. 18 (middle and right panels), thereby modulating the convective strength. In the mixed-phase regions of continental DCCs At levels colder than -10°C in all simulated cases, excluding SIP weakens vertical motion by about 5-10%, while in the marine convection of ORCESTRAT, the reduction exceeds 20% systematically reduces the 90th percentile stratiform ascent (Fig. 18, middle panel), with reductions of 5% in CAIPEEX and DCMEX, and up to 30% in MC3E and ORCESTRAT. This is explicable in terms of reduced diabatic heating (Figs attributed to reduced latent heating in the absence of SIP (Fig. 16, 17), which lowers parcel buoyancy in the 'No SIP' simulations. Noticeably, below the freezing level in MC3E the net positive buoyancy, suppressing stratiform ascent.

In contrast, in strong convective regions ($w > 1 \text{ m s}^{-1}$) (Fig. 18, right panel), excluding SIP reduces glaciation and total condensate loading in the mixed-phase regions (0 to -20°C). Although this reduces latent heating (Fig. 16), the associated decrease in suspended condensate lowers the downward drag sufficiently to produce a net increase in buoyancy without SIP (Fig. 18 a), SIP exclusion causes a little (3%) increase in vertical velocity, due to (left panel)), thereby enhancing convective ascent by over 20% less diabatic cooling there %.

In DCCs of ORCESTRAT, excluding SIP produces a vertically varying response in convective ascent (Fig. 18l). At warmer subzero levels ($> -20^\circ\text{C}$), where latent heating reduction dominates, convective ascent reduces by 20%. In contrast, at colder levels, a 20% increase in convective ascent is attributed to elevated buoyancy due to reduced condensate loading (Fig.S3 (b, f) in the supplement 18j).

Thus, above the freezing level, all simulated DCCs show weaker updrafts when SIP is excluded, primarily because reduced diabatic heating through diminished microphysical processes Unlike the other cases, excluding SIP in CAIPEEX reduces convective ascent by up to 10% in their mixed-phase region (Fig. 16). Similar changes appear in the maximum updraft and downdraft magnitudes 18c), driven by reduced buoyancy in convective regions (Fig.S4 in the supplement 18a). This suggests that in this relatively less intense CAIPEEX convection, the latent heating reduction from excluding SIP dominates over the condensate loading effect, resulting in weaker convective updrafts.




Figure 18. Domain-averaged profiles of the **vertical velocity** buoyancy (left panel), stratiform ($0 < w < 1 \text{ m s}^{-1}$, middle panel) and convective ($w > 1 \text{ m s}^{-1}$, right panel) ascents during convective periods from the control (black lines) and No SIP (gray lines) of the DCCs during (a-a-c) CAIPEEX, (b-d-f) DCMEX, (c-g-i) MC3E, and (d-j-l) ORCESTRA. In the left panel, the thick and thin lines denote buoyancy in convective and stratiform ascents, respectively.

The impact of SIP on the domain-mean cloud top height (CTH) in the simulated DCCs for all simulated cases is shown in Fig. 19. The CTH at each grid cell is defined as the highest level where total water content exceeds 0.001 g m^{-3} . Overall, excluding SIP decreases the CTH by about 1 km in continental DCCs, with a slight increase in CAIPEEX DCCs in their mature stage. This decrease is 2-3 km in ORCESTRA marine convection throughout the simulations. In contrast, when SIP is excluded from the marine ORCESTRA case, a pronounced reduction in CTH is seen. During the active convective stage, both control and No SIP simulations reach a similar peak CTH of 15 km, but from the mature stage onward, excluding SIP leads to a progressive reduction in CTH, reaching about 4 km lower CTH than the control by the end of the simulation. This is due to reduced latent heat release consistent with decreased latent heating without SIP, which weakens buoyancy and limits the vertical development of these DCCs when SIP is absent. This effect is more pronounced in ORCESTRA convection stratiform ascent and cloud depth in the marine environment, where cloud growth is more sensitive to latent heating and ice-crystal processes, as noted in Figs. 19 (j-l). Without SIP, the suppression of ice-crystal processes weakens convective intensity, leading to shallower cloud tops. These findings are in contrast to a modeling study by Grzegorzczuk et al. (2025), who reported stronger latent heating and a 1.5 km higher cloud top higher CTH when SIP was excluded in simulated DCCs.

In summary, in the simulated DCCs, the exclusion of SIP significantly reduces diabatic (latent and radiative) heating, which weakens the strength of convection through decreased vertical motion. This reduction is primarily driven by suppressed ice-crystal processes, such as depositional growth and riming, which lowers latent heat release and CRH. In such DCCs, SIP results in enhanced absorption of outgoing LW radiation, increasing net CRH by over 20%. With SIP, the increase in latent heat release increases the CTH by 5-30%, both in continental and marine DCCs, the present study finds a negligible impact on CTH in continental DCCs but a larger reduction (up to 4 km) in the marine ORCESTRA case when SIP is excluded.




Figure 19. Time series of the cloud top height in the control (black lines) and No SIP (gray lines) simulations of (a) CAIPEEX, (b) DCMEX, (c) MC3E, and (d) ORCESTRA DCCs. The same information is shown for (e, f) DCMEX, (g, h) MC3E, and (i, j) ORCESTRA clouds. The cloud top height at each grid cell is defined as the highest level where total water content exceeds $10^{-3-2} \text{ g m}^{-3}$ following Grzegorzcyk et al. (2025).

3.6 Comparison of SIP parameterizations for raindrop shattering

680 In the control runs (Table 3), SIP during fragmentation in RDF is represented using the SUL18 PHIL18 parameterization. To assess the impact of the choice of SIP parameterization of RDF fragmentation on the simulated cloud properties, these convective cases are simulated with the PHIL18 SULL18 scheme (Sec. 2.2.1, PHIL18 SULL18 run) and compared with their corresponding control (SUL18) PHIL18 runs.

Figure 20 shows the domain-averaged total INC simulated in the control (black blue line), PHIL18 (black SULL18 (orange line), and No SIP (grey gray line) runs. Between 0 and The process rate analysis (supplementary Fig. S8) reveals that RDF from PHIL18 and SULL18 contributes approximately 80% and 1% to the total secondary ice, respectively, at levels warmer than -10°C levels in . At these levels, SULL18 forms approximately 1.5 times lower overall INCs across all simulated DCCs (Fig. 20(a,d,g,j)), both SUL18 and a,c,e,g), with mean INCs in convective updrafts about 10 times lower than those formed by PHIL18 schemes predict about 1.5 times higher mean INC than SUL18.

690 (Supplementary Fig. S3). PHIL18 also exhibits closer agreement with the observed INCs (Fig. S3). This difference is because PHIL18 forms secondary ice splinters during both immersion freezing of raindrops and collisions between raindrops and ice particles. In comparison, SUL18 SULL18 only considers SIP during immersion freezing of raindrops.

Moreover, in ln the mixed-phase region of these DCCs, the additional INC in PHIL18 enhances relatively lower INC in SULL18 dampens depositional growth (not shown) and results in 2-10% more 2-5% less diabatic heating than the control run PHIL18 (Fig. 20 (h, k)). This enhanced decreased diabatic heating with PHIL18 boosts SULL18 weakens buoyancy, which leads to an increase a decrease in vertical motion compared to SUL18 (not shown here). PHIL18 supplementary Fig. S8.

In summary, the choice of the RDF SIP scheme has a moderate impact on the simulated properties in these DCCs . more pronounced impact on INC at warmer subzero temperatures, whereas its influence on the simulated dynamical properties of these DCCs remains moderate. This highlights the sensitivity of simulated SIP through RDF to the choice of scheme. This may remain true for parameterizations of other SIP processes.

700




Figure 20. Profiles of the means of the domain-averaged total ice number concentrations INCs from the control (which include Su18 PHIL18 RDF, solid black blue lines), Phil18 SULL18 RDF (solid gray yellow lines), and No SIP (dashed gray lines) runs in the (a) CAIPEEX, (c) DCMEX, (e) MC3E, and (g) ORCESTRA clouds. Also, the domain-averaged (b, d, f, h) latent heating rate in the control ('Su18 PHIL18 RDF'), Phil18 SULL18 RDF, and No SIP runs are shown for these the simulated clouds.

4 Summary and conclusions

This study assesses the impact of SIP on the microphysics and dynamics of DCCs by simulating four contrasting cloud cases with the ICON model. These include three aircraft-observed continental cases, observed in (i) CAIPEEX over India on 30 September 2019, (ii) DCMEX over New Mexico on 2 August 2022, (iii) MC3E over Oklahoma on 11 May 2011, and (iv) marine convection on 3 September 2024 during ORCESTRA over the tropical Atlantic region, observed by the EarthCARE satellite and research vessel RV Meteor. For all simulated DCCs, the simulated microphysical properties (LWC, CDNC, IWC, INC) and radar reflectivity are adequately validated against coincident observations. Additionally, a new SIP treatment for RDF, A SIP treatment of RDF proposed by PHIL18, is newly implemented in ICON and compared to the existing scheme of SUL18/SULL18.

In the simulated DCCs, SIP from IIC accounts for over 90% of secondary ice, dominating overall INC in their mixed-phase regions (Fig. 10). In addition, the HM process and RDF processes contribute to over 30-70% in the lower mixed-phase region, while RDF and SBF contribute less than 2% to total secondary ice. Two factors may explain this minimal contribution of SBF to total INC in these DCCs: parameterization uncertainties in the Deshmukh et al. (2022) formulation, or the physical limitation that most fragments from SBF evaporate sublime before reaching supersaturated regions where fragment growth can occur (Korolev and Leisner, 2020).

The findings of the present study are;

1. In the simulated continental DCCs of CAIPEEX, DCMEX, and MC3E, the impact of SIP on microphysical properties varies in the range of simulated clouds. In their mixed-phase regions, without SIP, the supercooled liquid mass increases by 30-60-20-40% due to enhanced condensation of vapor. In contrast, in a relatively saturated environment of marine DCCs of ORCESTRA, excluding SIP causes little (10%) increase in forms about 20% more supercooled liquid at subzero levels. Also, in the mixed-phase region of these DCCs, the absence of SIP causes a decrease in the IWC (INC) by 10-30% ($10\text{-}10^3\text{ L}^{-1}$) compared to with SIP.
2. Furthermore, the sensitivity of the surface precipitation to SIP is simulated to depend closely on the convective intensity and longevity. In continental DCCs of CAIPEEX, DCMEX and MC3E, prohibiting SIP predicts no significant change in the accumulated surface precipitation, whereas it decreases by about 4% in more vigorous, long-lived continental MC3E and marine ORCESTRA convection marine convection of ORCESTRA.
3. In the simulated DCCs, excluding SIP reduces total latent heating by 10–20%, consistent with Qu et al. (2022) and Grzegorzczuk et al. (2025), with the magnitude depending on convection intensity and duration. The reduction arises primarily from weakened ice-crystal growth by vapor deposition, leading to suppressed depositional heating and enhanced sublimational cooling in the mixed-phase region. These effects are modest (20%) in short-lived, weakly precipitating systems (CAIPEEX, DCMEX) but can reach up to 50% in sustained, moderately to strongly precipitating DCCs (MC3E, ORCESTRA). At the mixed-phase levels, enhanced vapor condensation partly offsets the cooling, indicating that the absence of SIP inhibits ice-phase processes and promotes the warm-rain pathway.

4. Excluding SIP from the simulated DCCs causes about a 15-40% decrease in the net CRH, mainly through LW CRH (20-50% decrease without SIP). This can be attributed to reduced absorption of outgoing LW radiation due to fewer ice particle mass and number in the absence of SIP.
5. In these DCCs, reduced CRH (by 15-40%) and latent heat release (by 10-20%) without SIP decreases buoyancy and leads to shallower cloud tops (up to 1-3 km). This study also found that excluding SIP weakens (strengthens) stratiform (convective) ascent across all simulated DCCs. A 5-20% increase in stratiform ascent without SIP is driven by reduced latent heating and buoyancy in those regions. In convective regions, however, the response is more complex. Although latent heating and CRH are reduced without SIP, the accompanying decrease in condensate loading sufficiently reduces downward drag, resulting in stronger convective ascent than with SIP. This contrasts with Grzegorzczuk et al. (2025), who reported shallower cloud top in the presence of SIP due to enhanced latent heating in an idealized simulation. This reduced buoyancy without SIP also lowers the mean vertical velocity by about 10%, indicating lower convective vigor than with SIP. Collectively, these findings highlight that excluding SIP suppresses convective intensity and leads to significantly shallower cloud development.
6. In continental DCCs, prohibiting SIP found a modest ($< 5\%$) increase in CTH. In contrast, the marine ORCESTRAs case exhibits a substantial reduction (by up to 3 km) in CTH during the mature stage when SIP is disabled. This remains consistent with reduced diabatic heating, which weakens stratiform ascent and limits cloud deepening. The lower cloud tops also likely contribute to the reduced LW CRH, as warmer cloud tops emit more longwave radiation to space. However, without SIP, the larger reduction in CTH in this marine convection suggests that the response may not be attributed to the absence of SIP alone, and may also reflect case-specific dynamical influences.
7. In all simulated DCCs, both SUL18 and PHIL18 schemes produce similar dynamical properties despite PHIL18 schemes of SIP through RDF predict similar INC and dynamical properties. Therefore, a simpler scheme by SUL18 for SIP in RDF can be used in numerical models to simulate the observed IWC and INC and associated radiative and dynamical properties. generating higher INCs at temperatures warmer than -10°C . PHIL18 also shows closer agreement with collocated INC observations when combined with other SIP processes.
8. This study further confirms the negligible impact of SIP in SBF on overall INC in a range of simulated DCCs, consistent with the findings of Waman et al. (2022) and Yang et al. (2024).

To conclude, this study reveals that SIP impact varies greatly among DCCs in different environmental conditions and convective strengths. By constraining SIP effects across four DCCs under different environments (from continental to marine), this study demonstrates that the magnitude of SIP influence on microphysics, dynamics, and diabatic properties systematically scales with convective evolution, intensity, and longevity, ranging from negligible effects in weak, short-lived systems to pronounced impacts in vigorous, long-lived DCCs. The exclusion of SIP leads to a substantial decrease in both temporal and vertical extent of convection in the MC3E and ORCESTRAs clouds, chiefly due to reduced glaciation in their mixed-phase regions, as also reported by Qu et al. (2022). This study also found that, in the simulated DCCs, without SIP, the reduction in latent heating and CRH decreases

buoyancy and mean vertical velocity at subzero levels, resulting in shallower cloud tops in sustained convection. This comprehensive assessment across diverse convective regimes establishes that SIP fundamentally controls the intensity, longevity, and radiative properties of sustained deep convection.

770 This study further confirms the critical role of various SIP processes in numerical models to accurately simulate the observed INC in natural DCCs (Fig. S2). It should be noted that uncertainties remain in existing SIP schemes (Seidel et al., 2024; Korolev and Leisner, 2020), which can significantly limit their accuracy and efficiency. These uncertainties are primarily from factors such as variability in cloud conditions, irregular particle habits in natural clouds, the nonlinear nature of microphysical processes, and various model assumptions and simplifications. Consequently, these challenges may hinder the realistic representation of these processes in numerical models, which underscores the need for additional observational and experimental studies to increase their fidelity.

Table A1: List of acronyms

Acronym
2DS
AP
BOWTIE
CAIPEEX
CCN
CDP
CIP CDNC
PIP CIP
2DS CKE
HVPS-3 CPI
CPR
NASA CRH
JAXA DCCs
DCMEX
DE
DKRZ
EarthCARE
MPI-M Max Planck Institute for Meteorology DKRZ Deutsches Klimarechenzentrum NWP Numerical Weather Prediction DE Dimensionless Energy CKE Collision Kinetic Energy LBC Lateral
ERA5
LWC HALO
CDNC HM
HVPS-3
ICON
IIC
INC
INP
IWC
INC JAXA
CAIPEEX LBC

Continued on next page

Acronym
DCMEX LW
LWC
MC3E
MPI-M
NASA
NWP
ORCESTR
ICON PHIL18
SIP PIP
HM Hallett-Mossop RDF
IIC Fragmentation in Ice-Ice Collision SBF
SUL18 SIP
PHIL18 SPIN
LW SULL18
SW
CRH Cloud Radiative Heating DCCs Deep Convective Clouds AP Aerosol Particle INP Ice Nucleating Particle CCN Cloud Condensation Nuclei

Table A2. List of symbols and parameters

Symbol	Description	Unit
D	Diameter of freezing drizzle/raindrop	mm
RH_i	Relative Humidity over ice	%
S_i	Saturation ratio with respect to ice	–
N_{sec}	Number of secondary ice fragments	–
T	Temperature	K (except where marked as °C)
N_{hm}	3.5×10^8	fragments per kg
q_{rime}	Rimed ice mass mixing ratio	kg kg^{-1}
$\Xi_{RDF}, \zeta, \beta_{RDF}, \eta$	D dependent polynomials from PHIL18	–
Ω	T dependent polynomial from PHIL18	–
N_S, N_B	Number of small and big fragments in RDF mode 1 by PHIL18	–
T_0	Value of T at maximum of Lorentzian for N_{sec}	°C
$T_{B,0}$	Value of T at maximum of Lorentzian of N_B	°C
ζ_B, η_B	D dependent polynomials for N_B	–
DE	Dimensionless Energy	–
DE_{crit}	Critical DE	–
K_0	Collision Kinetic Energy	kg m^{-2}
m, m_i	Mean masses of raindrop and colliding ice particle	kg
v, v_i	Mean velocities of raindrop and colliding ice particle	m s^{-1}
c_w	Specific heat capacity of liquid water	$4200 \text{ J K}^{-1} \text{ kg}^{-1}$
L_f	Specific latent heat of freezing	$3.3 \times 10^5 \text{ J kg}^{-1}$
$f(T)$	Frozen fraction at end of (stage 1) freezing; $f(T) = -c_w T / L_f$	–
C	Asperity-fragility coefficient	J^{-1}
α	Surface area (equivalent spherical) of smaller particle	m^2
$A(M)$	Number density of breakable asperities in region of contact	m^{-2}
n_{branch}	Number density of branches on smaller particles	m^{-2}
b_1	Fraction of surface of smaller particle in region of impact	–
γ	Empirical parameters (Deshmukh et al., 2022)	–
β_{SBF}	Empirical parameters (Deshmukh et al., 2022)	$\text{m}^{-0.5} \text{ s}^{-1}$
d	Crystal diameter	m
v', v^*	Onset transition factors for dendritic snow and graupel (Deshmukh et al., 2022)	–
Ξ_{SBF}	Emission factor (Deshmukh et al., 2022)	–
ρ_r	Bulk density of rimed particle	kg m^{-3}
ρ_{D0}	Density of a rimed particle by Dong et al. (1994)	kg m^{-3}
f_v	Ventilation coefficient for vapor diffusion	–

Code availability. The present study used the October 2024 release of the ICON model, which is available at: <https://gitlab.dkrz.de/icon/icon-model.git>. The raw simulation outputs will be archived on the high-performance storage system at the DKRZ. The post-processing codes can be accessed: https://github.com/Deepak-Waman/ACP_SIP_DYNAMICS_2025.git. The post-processed data will be made available on KIT-Open.

780 *Data availability.* The aircraft/radar data for the CAIPEEX case is available on request. The EarthCARE data for ORCESTRa is available at <https://gportal.jaxa.jp/gpr/user/regist1?lang=en>. The DCMEX data can be accessed through <https://data.ceda.ac.uk/badc/dcmex/data>, and MC3E at <https://www.earthdata.nasa.gov/data/projects/mc3e/data-access-tools>. The Radiosonde data for the simulated cases is obtained from <https://weather.uwyo.edu/upperair/sounding.shtml>.

Author contributions. DW and CH conceptualized the study. DW, JM, and GW implemented the SIP processes in ICON. BK and CB were
785 involved in the ICON setup and experiment design, and BK further offered guidance for the analysis of the simulations. TP made the observational data for CAIPEEX available, while DF and AB provided data for DCMEX. SS and SP helped in processing CAIPEEX aircraft and radar data. RF offered guidance on the EarthCARE data. DW conducted the simulations and analysis, and wrote the manuscript with input from all co-authors. CH supervised the project and secured funding.

Competing interests. The authors declare that they have no conflict of interests.

790 *Acknowledgements.* This work was funded through the project WarmWorld (Grant 01LK2202C) of the German Federal Ministry of Research, Technology and Space. This work used resources of the Deutsches Klimarechenzentrum (DKRZ) granted by its Scientific Steering Committee (WLA) under project ID bk1341. CH and GW acknowledge funding through the Horizon Europe Cluster 5 project CleanCloud (Grant no. 101137639). The authors thank Sagar Pokale, David Dufton, and Aaron Bansemer for their advice on the use of ground-based, flight, and satellite data. Bathymetric/topographic data were obtained from the General Bathymetric Chart of the Oceans (GEBCO, 2025) grid (doi: 10.5285/37c52e96-24ea-67ce-e063-7086abc05f29). The authors thank the two anonymous reviewers and the editor Prof. Ivy Tan for their constructive and insightful comments, which have helped improve the quality and clarity of the manuscript. During the writing process, artificial intelligence large language models (ChatGPT, Grammarly, Claude) were used to detect and correct grammatical mistakes.

References

- 800 Bacon, N., Swanson, B., Baker, M., and Davis, E.: Breakup of levitated frost particles, *Atmospheric Research*, 103, 13 763–13 775, <https://doi.org/10.1029/98JD01162>, 1998.
- Bigg, E. K.: The formation of atmospheric ice crystals by the freezing of droplets, *Quarterly Journal of the Royal Meteorological Society*, 79, 510–519, <https://doi.org/10.1002/qj.49707934207>, 1953.
- Bouniol, D., Roca, R., Fiolleau, T., and Poan, D. E.: Macrophysical, microphysical, and radiative properties of tropical mesoscale convective
805 systems over their life cycle, *Journal of Climate*, 29, 3353–3371, <https://doi.org/10.1175/JCLI-D-15-0551.1>, 2016.
- Bower, K. N., Moss, S., Johnson, D., Choulaton, T., Latham, J., Brown, P., Blyth, A., and Cardwell, J.: A parametrization of the ice water content observed in frontal and convective clouds, *Quarterly Journal of the Royal Meteorological Society*, 122, 1815–1844, 1996.
- Chandra, A., Zhang, C., Kollias, P., Matrosov, S., and Szyrmer, W.: Automated rain rate estimates using the Ka-band ARM zenith radar (KAZR), *Atmospheric Measurement Techniques*, 8, 3685–3699, 2015.
- 810 Cotton, R. J. and Field, P. R.: Ice nucleation characteristics of an isolated wave cloud, *Quarterly Journal of the Royal Meteorological Society*, 128, 2417—2437, <https://doi.org/10.1256/qj.01.150>, 2002.
- Daily, M. I., Robinson, J., Finney, D., Raif, E., McQuaid, J. B., Murray, B. J., and Blyth, A.: Dataset for Airborne observations of ice-nucleating particles during the 2022 DCMEX campaign, New Mexico, 2024.
- DeMott, P., Prenni, A., Liu, X., Kreidenweis, S., Petters, M., Twohy, C., Richardson, M., Eidhammer, T., and Rogers, D.: Predicting global
815 atmospheric ice nuclei distributions and their impacts on climate, *Proceedings of the National Academy of Sciences*, 107, 11 217–11 222, <https://doi.org/10.1073/pnas.0910818107>, 2010.
- DeMott, P., Suski, K., Hill, T., and Levin, E.: Southern Great Plains ice nuclei characterization experiment final campaign summary, Tech. rep., DOE Office of Science Atmospheric Radiation Measurement (ARM) Program . . . , 2015.
- Deshmukh, A., Phillips, V. T. J., Bansemer, A., Patade, S., and Waman, D.: New empirical formulation for the sublimational breakup of
820 graupel and dendritic snow, *Journal of Atmospheric Sciences*, 79, 317–336, <https://doi.org/10.1175/JAS-D-20-0275.1>, 2022.
- Dong, Y. Y., Oraltay, R. G., and Hallett, J.: Ice particle generation during evaporation, *Atmospheric Research*, 32, 45–53, [https://doi.org/10.1016/0169-8095\(94\)90050-7](https://doi.org/10.1016/0169-8095(94)90050-7), 1994.
- Ewald, F.: MIRA Cloud Radar Moments from the HALO Microwave Package (Level 2), oRCESTRA campaign dataset. Available at: <ipfs://bafybeifxtmq5mpn7vwiiwl4vlpoil7rgm2tnhmkeyqsyudleqegxzvwl3a>, 2025.
- 825 Ewald, F., Groß, S., Hagen, M., Hirsch, L., Delanoë, J., and Bauer-Pfundstein, M.: Calibration of a 35 GHz airborne cloud radar: lessons learned and intercomparisons with 94 GHz cloud radars, *Atmospheric Measurement Techniques*, 12, 1815–1839, 2019.
- Feng, Z., Dong, X., Xi, B., Schumacher, C., Minnis, P., and Khaiyer, M.: Top-of-atmosphere radiation budget of convective core/stratiform rain and anvil clouds from deep convective systems, *Journal of Geophysical Research*, 116, <https://doi.org/10.1029/2011JD016451>, 2011.
- Field, P. R. and Heymsfield, A. J.: Importance of snow to global precipitation, *Geophysical Research Letters*, 42, 9512–9520,
830 <https://doi.org/10.1002/2015GL065497>, 2015.
- Field, P. R., Heymsfield, A. J., and Bansemer, A.: Shattering and particle interarrival times measured by optical array probes in ice clouds, *Journal of Atmospheric and Oceanic Technology*, 23, 1357–1371, <https://doi.org/10.1175/JTECH1922.1>, 2006.
- Field, P. R., Lawson, R. P., and et. al.: Secondary ice production: Current state of the science and recommendations for the future, *Meteorological Monographs*, 58, 7.1–7.20, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0014.1>, 2017.

- 835 Finney, D. L., Blyth, A., and et. al.: Deep Convective Microphysics Experiment (DCMEX) coordinated aircraft and ground observations: microphysics, aerosol, and dynamics during cumulonimbus development, *Earth System Science Data*, 16, 2141–2163, <https://doi.org/10.5194/essd-16-2141-2024>, 2024.
- Foth, A.: Cloud radar and Cloudnet on RV Meteor during BOWTIE, <https://doi.org/10.82246/bafybeidtfyua2vmmrpnhl6nj57x2ynowaceghdapyetc6lnrpji>, 2026.
- 840 Gassmann, A. and Herzog, H. J.: Towards a consistent numerical compressible non-hydrostatic model using generalized Hamiltonian tools, *Quarterly Journal of the Royal Meteorological Society*, 134, 1597–1613, <https://doi.org/10.1002/qj.297>, 2008.
- Gautam, M., Waman, D., Patade, S., Deshmukh, A., Phillips, V., Jackowicz-Korczynski, M., Paul, F. P., Smith, P., and Bansemer, A.: Fragmentation in Collisions of Snow with Graupel/Hail: New Formulation from Field Observations, *Journal of the Atmospheric Sciences*, 81, 2149–2164, 2024.
- 845 Grzegorzczak, P., Yadav, S., Zanger, F., Theis, A., Mitra, S., Borrmann, S., and Szakal, M.: Fragmentation of ice particles: laboratory experiments on graupel–graupel and graupel–snowflake collisions, *Atmospheric Chemistry and Physics*, 23, 13 505—13 521, <https://doi.org/10.5194/acp-23-13505-2023>, 2023.
- Grzegorzczak, P., Wobrock, W., Canzi, A., Niquet, L., Tridon, F., and Planche, C.: Investigating secondary ice production in a deep convective cloud with a 3D bin microphysics model: Part II-Effects on the cloud formation and development, *Atmospheric Research*, 314, 107 797, 850 2025.
- Gupta, A. K., Deshmukh, A., Waman, D., Patade, S., Jadav, A., Phillips, V. T. J., Bansemer, A., Martins, J. A., and Goncalves, F. L. T.: The microphysics of the warm-rain and ice crystal processes of precipitation in simulated continental convective storms, *Communication Earth Environment*, 4, <https://doi.org/10.1038/s43247-023-00884-5>, 2023.
- Hallett, J. and Mossop, S.: Production of secondary ice particles during the riming process, *Nature*, 249, 26–28, 855 <https://doi.org/10.1038/249026a0>, 1974.
- Han, C., Hoose, C., and Durlich, V.: Secondary ice production in simulated deep convective clouds: a sensitivity study, *Journal of Atmospheric Sciences*, 81, 5, <https://doi.org/10.1175/JAS-D-23-0156.1>, 2024.
- Hande, L. B., Hoose, C., and Tegen, I.: Seasonal variability of Saharan desert dust and ice nucleating particles over Europe, *Atmospheric Chemistry and Physics*, 15, 4389—4397, <https://doi.org/10.5194/acp-15-4389-2015>, 2015.
- 860 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *Quarterly journal of the royal meteorological society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hogan, R. and Bozzo, A.: A Flexible and Efficient Radiation Scheme for the ECMWF Model, *Proceedings of the National Academy of Sciences*, 10, 1990–2008, <https://doi.org/10.1029/2018MS001364>, 2018.
- Houze, R. A. J.: Cloud clusters and large-scale vertical motions in the tropics, *Journal of the Meteorological Society of Japan*, 60, 396–410, 865 https://doi.org/10.2151/jmsj1965.60.1_396, 1982.
- Huang, Y., Wu, W., McFarquhar, G. M., Wang, X., Morrison, H., Ryzhkov, A., Hu, Y., Wolde, M., Nguyen, C., Schwarzenboeck, A., and Milbrandt, J.: Microphysical processes producing high ice water contents (HIWCs) in tropical convective clouds during the HAIC-HIWC field campaign: evaluation of simulations using bulk microphysical schemes, *Atmospheric Chemistry and Physics*, 21, 6919–6944, <https://doi.org/10.5194/acp-21-6919-2021>, 2021.
- 870 Huihui, W., Marsden, N., and co authors: The role of aerosols and meteorological conditions in shaping cloud droplet development in New Mexico summer deep-convective systems, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2025-2600>, 2025.

- Jackson, R., French, R., Leon, D. C., Plummer, D. M., Lasher-Trapp, S., Blyth, A., and Korolev, A.: Observations of the microphysical evolution of convective clouds in the southwest of the United Kingdom, *Atmospheric Chemistry and Physics*, 18, 15 329—15 344, <https://doi.org/10.5194/acp-18-15329-2018>, 2018.
- 875 James, R. L., Phillips, V. T., and Connolly, P. J.: Secondary ice production during the break-up of freezing water drops on impact with ice particles, *Atmospheric Chemistry and Physics*, 21, 18 519—18 530, <https://doi.org/10.5194/acp-21-18519-2021>, 2021.
- Järvinen, E., McCluskey, C. S., Waitz, F., Schnaiter, M., Bansemer, A., Bardeen, C. G., Gettelman, A., Heymsfield, A., Stith, J. L., Wu, W., et al.: Evidence for secondary ice production in Southern Ocean maritime boundary layer clouds, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD036 411, 2022.
- 880 Jeffery, C. A. and Austin, P. H.: Homogeneous nucleation of supercooled water: Results from a new equation of state, *Quarterly Journal of the Royal Meteorological Society*, 102, 25 269–25 279, <https://doi.org/10.1029/97JD02243>, 1997.
- Jensen, M. P., Peterson, A., and et. al.: The Midlatitude Continental Convective Clouds Experiment (MC3E), *Bulletin of the Meteorological Society*, 97, 1667–1686, <https://doi.org/10.1175/BAMS-D-14-00228.1>, 2016.
- Johnson, D. and Hallett, J.: Freezing and shattering of supercooled water drops, *Quarterly Journal of the Royal Meteorological Society*, 94, 885 468–482, 1968.
- Kanji, Z., Ladino, L., Wex, H., Boose, Y., Burkert-Kohn, M., Cziczo, D., and Krämer, M.: Overview of ice nucleating particles, *Proceedings of the National Academy of Sciences*, 58, 1.1–1.33, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0006.1>, 2017.
- Keinert, A., Spannagel, D., Leisner, T., and Kiselev, A.: Secondary ice production upon freezing of freely falling drizzle droplets, *Journal of the Atmospheric Sciences*, 77, 2959–2967, <https://doi.org/10.1175/JAS-D-20-0081.1>, 2020.
- 890 Keshtgar, B., Voigt, A., Mayer, B., and Hoose, C.: Uncertainties in cloud-radiative heating within an idealized extratropical cyclone, *Atmospheric Chemistry and Physics*, 24, 2451–2469, <https://doi.org/10.5194/acp-24-4751-2024>, 2024.
- Kim, J., Kollias, P., Puigdomènech Treserras, B., Battaglia, A., and Tan, I.: Evaluation of the EarthCARE Cloud Profiling Radar (CPR) Doppler velocity measurements using surface-based observations, *Atmospheric Chemistry and Physics*, 25, 15 389–15 402, 2025.
- Kleinheins, J., Kiselev, A., Keinert, A., Kind, M., and Leisner, T.: Thermal Imaging of Freezing Drizzle Droplets: Pressure Release Events 895 as a Source of Secondary Ice Particles, *Proceedings of the National Academy of Sciences*, 78, 1703—1713, <https://doi.org/10.1175/JAS-D-20-0323.1>, 2021.
- Korolev, A. and Isaac, G. A.: Observations of sublimating ice particles in clouds, in: *Proceedings of the 14th International Conference on clouds and precipitation*, pp. 808–811, 2004.
- Korolev, A. and Leisner, T.: Review of experimental studies of secondary ice production, *Atmospheric Chemistry and Physics*, 20, 11 767– 900 11 797, <https://doi.org/10.5194/acp-20-11767-2020>, 2020.
- Korolev, A. V., Emery, E. F., Strapp, W., Cober, S. G., Issac, G. A., Wasley, E., and Marcotte, D.: Small Ice Particles in Tropospheric Clouds: Fact or Artifact?, *Bulletin of the American Meteorological Society*, 92, 967–973, <https://doi.org/10.1175/2010BAMS3141.1>, 2011.
- Lasher-Trapp, S., Leon, D. C., DeMott, P. J., Villanueva-Birriel, C. M., Johnson, A. V., Moser, D. H., Tully, C. S., and Wu, W.: A multisensor investigation of rime splintering in tropical maritime cumuli, *Journal of the Atmospheric Sciences*, 73, 2547–2564, 2016.
- 905 Lasher-Trapp, S., Scott, E. L., Järvinen, M., Schnaiter, F., Waitz, P. J., DeMott, P. J., McCluskey, C. S., and J., H. T. C.: Observations and Modeling of Rime Splintering in Southern Ocean Cumuli, *Journal of Geophysical Research*, 126, <https://doi.org/10.1029/2021JD035479>, 2021.
- Lau, K. M. and Wu, H. T.: Warm rain processes over tropical oceans and climate implications, *Geophysical Research Letter*, 30, 7.3–7.5, <https://doi.org/10.1029/2003GL018567>, 2003.

- 910 Lawson, P., Woods, S., and Morisson, H.: The Microphysics of Ice and Precipitation Development in Tropical Cumulus Clouds, *Journal of Atmospheric Sciences*, 72, 2429–2445, <https://doi.org/10.1175/JAS-D-14-0274.1>, 2015.
- Lohmann, U., Lüönd, F., and Mahrt, F.: *An introduction to clouds: From the microscale to climate*, Cambridge University Press, 2016.
- Mason, B. J. and Maybank, J.: The fragmentation and electrification of freezing water drops, *Quarterly Journal of the Royal Meteorological Society*, 86, 1703—1713, <https://doi.org/10.1002/qj.49708636806>, 1960.
- 915 Oraltay, R. G. and Hallett, J.: Evaporation and melting of ice crystals: A laboratory study, *Atmospheric Research*, 24, 169–189, [https://doi.org/10.1016/0169-8095\(89\)90044-6](https://doi.org/10.1016/0169-8095(89)90044-6), 1989.
- Ori, D., Schemann, V., Karrer, M., Dias Neto, J., von Terzi, L., Seifert, A., and Kneifel, S.: Evaluation of ice particle growth in ICON using statistics of multi-frequency Doppler cloud radar observations, *Quarterly Journal of the Royal Meteorological Society*, 146, 3830–3849, 2020.
- 920 Patade, S., Shete, S., Malap, N., Kulkarni, G., and Prabha, T. V.: Observational and simulated cloud microphysical features of rain formation in the mixed phase clouds observed during CAIPEEX, *Atmospheric Research*, 169, 32—45, <https://doi.org/10.1016/j.atmosres.2015.09.018>, 2016.
- Patade, S., Kulkarni, G., Patade, S. Waman, D., Sotiropoulou, G., Samanta, S., and Malap, N.: Importance of secondary ice production in mixed-phase monsoon clouds over the Indian subcontinent, *Atmospheric Research*, 315, 107 890, 925 <https://doi.org/10.1016/j.atmosres.2024.107890>, 2025.
- Phillips, V. T. J., Yano, J. J., Formenton, M., Ilotoviz, E., Kanawade, V., Kudzotsa, I., Sun, J., Bansemer, A., Detwiler, A. G., Khain, A., and Tessendorf, S.: Ice multiplication by breakup in ice–ice collisions. Part II: Numerical simulations, *Journal of Atmospheric Sciences*, 74, 2789–2811, <https://doi.org/10.1175/JAS-D-16-0223.1>, 2017a.
- Phillips, V. T. J., Yano, J. J., and Khain, A.: Ice multiplication by breakup in ice–ice collisions. Part I: Theoretical formulation, *Journal of* 930 *Atmospheric Sciences*, 74, 1705–1719, <https://doi.org/10.1175/JAS-D-16-0224.1>, 2017b.
- Phillips, V. T. J., Patade, S., Gutierrez, J., and Bansemer, A.: Secondary ice production by fragmentation of freezing drops: Formulation and theory Possible High Ice Particle Production during Graupel–Graupel Collisions, *Journal of Atmospheric Sciences*, 75, 3031–3070, <https://doi.org/10.1175/JAS-D-17-0190.1>, 2018.
- Prabhakaran, T., Murugavel, P., Konwar, M., and coauthors: CAIPEEX: Indian Cloud Seeding Scientific Experiment, *Bulletin of the American Meteorological Society*, 104, E2095—E2120, <https://doi.org/10.1175/BAMS-D-21-0291.1>, 2023.
- 935 Price, H., Baustian, K., McQuaid, J., Blyth, A., Bower, K., Choularton, T., Cotton, R., Cui, Z., Field, P., Gallagher, M., et al.: Atmospheric ice-nucleating particles in the dusty tropical Atlantic, *Journal of Geophysical Research: Atmospheres*, 123, 2175–2193, 2018.
- Qu, Z., Korolev, A., Milbrandt, J., Heckman, I., Huang, Y., McFarquhar, G., Morrison, H., Wold, M., and Nguyen, C.: The impacts of secondary ice production on microphysics and dynamics in tropical convection, *Atmospheric Chemistry and Physics*, 22, 12 287–12 310, 940 <https://doi.org/10.5194/acp-22-12287-2022>, 2022.
- Roca, R., Fiolleau, T., and Bouniol, D.: A simple model of the life cycle of mesoscale convective systems cloud shield in the tropics, *Journal of Climate*, 30, 4283–4298, <https://doi.org/10.1175/JCLI-D-16-0556.1>, 2017.
- Sakradzija, M., Seifert, A., and Heus, T.: Fluctuations in a quasi-stationary shallow cumulus cloud ensemble, *Proceedings of the National Academy of Sciences*, 22, 65–85, <https://doi.org/10.5194/npg-22-65-2015>, 2015.
- 945 Sasikumar, S., Battaglia, A., Puigdomènech, B., and Kollias, P.: The Estimation of Path Integrated Attenuation for the EarthCARE Cloud Profiling Radar, *EGUsphere* [preprint], pp. 2025–3573, <https://doi.org/10.5194/egusphere-2025-3573>, 2025.

- Sato, K., Okamoto, H., Nishizawa, T., Jin, Y., Nakajima, T. Y., Wang, M., Satoh, M., Roh, W., Ishimoto, H., and Kudo, R.: JAXA Level 2 cloud and precipitation microphysics retrievals based on EarthCARE radar, lidar, and imager: the CPR_CLP, AC_CLP, and ACM_CLP products, *Atmospheric Measurement Techniques*, 18, 1325–1338, 2025.
- 950 Segal, Y. and Khain, A.: Dependence of droplet concentration on aerosol conditions in different cloud types: Application to droplet concentration parameterization of aerosol conditions, *Journal of Geophysical Research*, 111, 1–11, <https://doi.org/10.1029/2005JD006561>, 2006.
- Seidel, J., Kiselev, A., Keinert, A., Stratmann, F., Leisner, T., and Hartmann, S.: Secondary ice production – no evidence of efficient rim-splintering mechanism, *Atmospheric Chemistry and Physics*, 24, 5247–5263, <https://doi.org/10.5194/acp-24-5247-2024>, 2024.
- 955 Seifert, A. and Beheng, K. D.: A two-moment cloud microphysics parameterization for mixed-phase clouds. Part 1: Model description, *Meteor. Atmos. Phys.*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- Shete, S., Patade, S., Malap, N., Bankar, S., Jaykumar, S., Baumgardner, D., Axisa, D., Murugavel, P., and Prabhakaran, T.: Observations and simulations of immersion freezing INPs and their impact on cloud properties over the rain shadow region of India, *Atmospheric Research*, p. 109056, 2026.
- 960 Sotiropoulou, G., S., S., Savre, J., Lloyd, G., Lachlan-Cope, T., Ekman, A. M., and Nenes, A.: The impact of secondary ice production on Arctic stratocumulus, *Atmospheric Chemistry and Physics*, 20, 1301–1316, <https://doi.org/10.5194/acp-20-1301-2020>, 2020.
- Stevens, B., Bony, S., Gross, S., Klocke, D., Windmiller, J. M., Wing, A. A., Von Bismarck, J., Brito, E., David, R. O., Delanoë, J., et al.: Orcestra: Organized convection and earthcare studies over the tropical Atlantic, *Tellus*, 79, 2026.
- Sullivan, S., Keshtgar, B., and coauthors.: How does cloud-radiative heating over the North Atlantic change with grid spacing, convective parameterization, and microphysics scheme in ICON version 2.1.00?, *Geoscientific Model Development*, 16, 3535–3551, <https://doi.org/10.5194/gmd-16-3535-2023>, 2023.
- 965 Sullivan, S. C., Barthlott, C., Crosier, J., Zhukov, I., Nenes, A., and Hoose, C.: The effect of secondary ice production parameterization on the simulation of a cold frontal rainband., *Atmospheric Chemistry and Physics*, 18, 16 461–16 480, <https://doi.org/10.5194/acp-18-16461-2018>, 2018.
- 970 Takahashi, C. and Yamashita, A.: Production of ice splinters by the freezing of water drops in free fall, *Journal of the Meteorological Society of Japan. Ser. II*, 55, 139–141, 1977.
- Takahashi, T., Nagao, Y., and Kushiya, Y.: Possible High Ice Particle Production during Graupel–Graupel Collisions, *Journal of Atmospheric Sciences*, 52, 4523–4527, [https://doi.org/10.1175/1520-0469\(1995\)052<4523:PHIPPD>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<4523:PHIPPD>2.0.CO;2), 1995.
- Tian, L. and Srivastava, R.: Measurement of attenuation at C band in a convective storm by a dual-radar method, *Journal of Atmospheric and*
- 975 *Oceanic Technology*, 14, 184–196, 1997.
- Trenberth, K. E., Fasullo, J. T., and Kiehl, J.: Earth’s global energy budget, *Bulletin of the American meteorological society*, 90, 311–324, <https://doi.org/10.1175/2008BAMS2634.1>, 2009.
- Vardiman, L.: The generation of secondary ice particles in clouds by crystal–crystal collision, *Journal of Atmospheric Sciences*, 35, 2168–2180, [https://doi.org/10.1175/1520-0469\(1978\)035<2168:TGOSIP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2168:TGOSIP>2.0.CO;2), 1978.
- 980 Waman, D., Patade, S., Jadav, A., Deshmukh, A., Gupta, A., Phillips, V. T. J., Bansemer, A., and DeMott, P. J.: Dependencies of Four Mechanisms of Secondary Ice Production on Cloud-Top Temperature in a Continental Convective Storm, *Journal of the Atmospheric Sciences*, 79, 3375–3404, <https://doi.org/10.1175/JAS-D-21-0278.1>, 2022.
- Wan, H., Giorgetta, M. A., and et. al.: The ICON-1.2 hydrostatic atmospheric dynamical core on triangular grids – Part 1: Formulation and performance of the baseline version, *Geoscientific Model Development*, 6, 735–763, <https://doi.org/10.5194/gmd-6-735-2013>, 2013.

- 985 Wang, D., Prein, A. F., Giangrande, S. E., Ramos-Valle, A., Ge, M., and Jensen, M. P.: Convective updraft and downdraft characteristics of continental mesoscale convective systems in the model gray zone, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036746, 2022.
- Wehr, T., Kubota, T., and coauthors: The EarthCARE mission – science and system overview, *Atmospheric Measurement Techniques*, 16, 3581–3608, <https://doi.org/10.5194/amt-16-3581-2023>, 2023.
- 990 Widener, K., Bharadwaj, N., and Johnson, K.: Ka-band ARM Zenith Radar (KAZR) instrument handbook, Tech. rep., DOE Office of Science Atmospheric Radiation Measurement (ARM) Program . . . , <https://doi.org/10.2172/1035855>, 2012.
- Wilcox, E. M., Yuan, T., and Song, H.: Deep convective cloud system size and structure across the global tropics and subtropics, *Atmospheric Measurement Techniques*, 16, 5387–5401, <https://doi.org/10.5194/amt-16-5387-2023>, 2023.
- Yang, J., Huang, S., Yang, T., Zhang, Q., Deng, Y., and Liu, Y.: Impact of ice multiplication on the cloud electrification of a cold-season
995 thunderstorm: a numerical case study, *Atmospheric Chemistry and Physics*, 24, 5989–6010, <https://doi.org/10.5194/acp-24-5989-2024>, 2024.
- Zhao, X. and Liu, X.: Global importance of secondary ice production, *Geophysical Research Letters*, 48, e2021GL092581, <https://doi.org/10.1029/2021GL092581>, 2021.
- Zängl, G., Reinert, D., Ripodas, P., , and Baldauf, M.: The ICON (ICOsahedral Nonhydrostatic) modelling framework of DWD and
1000 MPI-M: Description of the nonhydrostatic dynamical core, *Quarterly Journal of the Royal Meteorological Society*, 141, 563–579, <https://doi.org/10.1002/qj.2378>, 2015.