

Implementing methane dynamics into the LPJmL6 model

Sibyll Schaphoff¹, David Höttner^{1,2}, Christoph Müller¹, Dieter Gerten^{1,3}, Sebastian Ostberg¹, and Werner von Bloh¹

¹Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, Potsdam, Germany

²Freie Universität Berlin (FU Berlin)

³Geography Department, Humboldt-Universität zu Berlin, 10099 Berlin

Correspondence: Sibyll Schaphoff (Sibyll.Schaphoff@pik-potsdam.de)

Abstract. We extend the Dynamic Global Vegetation Model LPJmL to version 6.0 by explicitly representing methane (CH₄) dynamics within the coupled carbon–nitrogen–water system. The implementation (i) prognoses water-table depth and wetland extent using a CTI–TOPMODEL framework, (ii) solves sub-daily, vertically explicit mass balances for CH₄ and O₂ including diffusion, ebullition, and plant-mediated transport, (iii) represents methanogenesis and methanotrophy with temperature- and moisture-dependent kinetics, and (iv) integrates land-use and rice management effects alongside inundation-tolerant plant functional types. This architecture enables consistent simulation of CH₄, carbon dioxide (CO₂) and nitrous oxide (N₂) emissions from natural wetlands and managed systems together with the soil CH₄ sink. Extensive benchmarking against global datasets shows that LPJmL6 reproduces the magnitude and regional–temporal variability of CH₄ flux pathways while maintaining strong skill in the simulated terrestrial carbon, nitrogen, and water budgets. The model thus provides a coherent, process-based framework to quantify CH₄ within the coupled carbon–nitrogen–water system, elucidate interactions with vegetation and soils, and assess how land-use, wetland conservation and restoration, and rice management options affect methane and overall greenhouse-gas budgets in support of climate-mitigation strategies.

1 Introduction

Wetlands, including ponds, lakes, rivers, and marshes are responsible for approximately 30% of global methane (CH₄) emissions (Saunois et al., 2020). These unique ecosystems are characterized by water-saturated conditions that create anoxic environments promoting methanogenic processes. However, wetland dynamics, such as changes in its water balance and vegetation, can significantly impact CH₄ emissions (Yang et al., 2022).

The variability in wetland CH₄ emissions is governed by several environmental factors, including water table depth, temperature, vegetation type, nutrient availability, and air pressure (Whalen and Reeburgh, 1991; Bartlett and Harriss, 1992; Moore et al., 2011; Turetsky et al., 2014; Bloom et al., 2017b). These factors regulate the balance between oxic and anoxic decomposition, which determine CH₄ production and release under anoxic conditions and carbon dioxide production and release under oxic conditions. Water table depth, controlled by precipitation, evapotranspiration, and soil conditions, is a key driver of anoxic conditions necessary for methanogenesis. A high

25 water table limits O_2 diffusion, promoting CH_4 production, while a low water table fosters oxic decomposition and
 CH_4 oxidation (Turetsky et al., 2014; Whalen and Reeburgh, 1991). Temperature also influences CH_4 emissions by
affecting microbial activity, with soil temperature driving methanogenesis rates and water temperature influencing
gas solubility and diffusion (Moore et al., 2011). Vegetation impacts CH_4 fluxes by transporting CH_4 via aerenchyma
30 tissues and contributing organic substrates that fuel microbial processes (Bartlett and Harriss, 1992). Nutrient
availability further shapes microbial activity, influencing the balance between CH_4 production and oxidation (Bloom
et al., 2017a). Atmospheric pressure and vapor pressure deficit (VPD) indirectly affect emissions by influencing gas
diffusion and soil moisture dynamics (Moore et al., 2011).

These interrelated factors underscore the complexity of CH_4 emissions from wetlands and their role in global
greenhouse gas budgets. While the emission sources are diverse (including fossil fuel and waste (Saunio et al., 2020)),
35 this paper focuses on the specific issue of CH_4 emissions from land use (excluding direct livestock emissions, which are
governed by feed quantity, quality and livestock stocking densities (Heinke et al., 2023)) and from natural peatlands.
Understanding and quantifying these emissions is vital for developing effective climate mitigation strategies and
improving insights into the carbon cycle and its role in global climate systems.

Process-based representations of wetland CH_4 emissions in land surface and Earth system models have developed
40 substantially over the past two decades. Early model formulations established the core mechanistic structure still
used in many current approaches, resolving CH_4 production under anoxic conditions, oxidation in oxic soil layers,
and the main transport pathways to the atmosphere, including diffusion, ebullition, and plant-mediated transport
(Walter and Heimann, 2000). These concepts were subsequently embedded in dynamic vegetation and land surface
models, allowing wetland CH_4 emissions to respond interactively to soil temperature, water table dynamics, sub-
45 strate availability, vegetation composition, and permafrost or peatland processes. Examples include LPJ-WHyMe,
which introduced a methane module into a dynamic global vegetation model with explicit peatland hydrology and
permafrost processes (Wania et al., 2010); ORCHIDEE-based wetland methane formulations used to investigate
climate- CH_4 feedbacks (Ringeval et al., 2011); CLM4Me, which coupled CH_4 production, oxidation, and transport
to land surface biogeochemistry and hydrology (Riley et al., 2011); and later peatland- or wetland-specific devel-
50 opments such as ORCHIDEE-PEAT/ORCHIDEE-PCH4 and WETMETH (Salmon et al., 2022; Nzotungicimpaye
et al., 2021). Model intercomparison projects such as WETCHIMP further demonstrated that differences in simulated
wetland extent, inundation dynamics, soil thermal and hydrological states, and process parameterizations remain
major sources of uncertainty in global wetland CH_4 estimates (Melton et al., 2013; Wania et al., 2013). These studies
show that a realistic representation of wetland CH_4 emissions requires tight coupling between hydrology, soil redox
55 conditions, carbon substrate supply, vegetation transport pathways, and soil-atmosphere gas exchange.

We have implemented the main processes controlling CH_4 dynamics in an updated version of the Dynamic Global
Vegetation Model, LPJmL (Lund-Potsdam-Jena managed land; Schaphoff et al., 2018; von Bloh et al., 2018). This
enhanced model simulates the intricate dynamics of natural and land-use-related CH_4 emissions within the terrestrial
biosphere, integrating wetland and permafrost dynamics as part of the coupled global carbon and water cycles.

60 In this work, we extend the LPJmL model from version 5.10 (von Bloh et al., 2018; Lutz et al., 2019; Wirth et al., 2024) to version 6.0 by incorporating processes that represent CH_4 dynamics. Specifically, we introduce a dynamic representation of the water table to determine the surface area influenced by the unconfined aquifer. This, combined with simulated O_2 and CH_4 concentrations in soil layers, enables the estimation of wetland distribution and anaerobic soil conditions. Consequently, the model now simulates CH_4 production, soil-layer CH_4 concentrations, CH_4 oxidation, and fluxes to the atmosphere through diffusion, ebullition, and plant-mediated transport, fully integrated with the hydrological and biogeochemical processes of LPJmL. Land-use-related CH_4 emissions considered in the present study include rice systems. Livestock-related CH_4 emissions are represented in a previous LPJmL5.0 development (Heinke et al., 2023), but are not evaluated or further described here. Sub-grid open-water CH_4 emissions from lakes, ponds, and river networks are not explicitly represented in the current implementation. Following the documentation of this new implementation, we provide a global evaluation of key processes related to wetland and CH_4 dynamics, along with other relevant processes, to assess the overall performance of LPJmL6.

Following the documentation of this new implementation, we provide a global evaluation of key processes related to wetland and CH_4 dynamics, along with other relevant processes, to assess the overall performance of LPJmL6.

2 Module description

75 This module description focuses on the newly developed part of the model. A general description of the LPJmL model can be found in Schaphoff et al. (2018) and in subsequent descriptions and evaluations of further model extensions, i.e. von Bloh et al. (2018); Lutz et al. (2019); Wirth et al. (2024). Additional changes are documented in the Supplement and together constitute LPJmL5.10.2. Wetlands are the primary natural sources of CH_4 production, while agricultural land is a major source of human-caused CH_4 emissions. In LPJmL6, we consider both natural and anthropogenic sources of CH_4 emissions. Given the large computational cost of the simulation of CH_4 soil processes, this can be turned off for simulations with LPJmL6 that are not aiming at studying CH_4 dynamics (see the Supplement).

Each grid cell is divided into several sub-regions, called stands, which represent different land-cover and land-use types. We distinguish a natural wetland stand, natural upland stand, several agricultural stands, pasture stands and setaside stands. Managed land-cover types are further distinguished into rainfed and irrigated stands. The introduction of the natural wetland stand makes the conversion of different land-use types more complex in LPJmL6 compared to earlier versions (see section 2.5).

2.1 Wetland representation

We compute the water-saturated area in each 0.5° grid cell annually from the compound topographic index (CTI), which describes the propensity of a location to accumulate water (Beven and Kirkby, 1979) and is defined as

$$\text{CTI}_i = \ln \left(\frac{\alpha_i}{\tan(\beta_i)} \right), \quad (1)$$

where α_i is the upslope contributing area and β_i the local surface slope at point i . In our implementation, CTI and slope statistics at 0.5° resolution are taken from the topographic datasets described in the section 4.1. Within each grid cell, CTI values are used to represent subgrid-scale variations in wetness and to derive the saturated fraction following a TOPMODEL-type approach (Beven and Kirkby, 1979; Sivapalan et al., 1987; Kleinen et al., 2012).

To relate CTI_i to water-table depth we follow the TOPMODEL framework Beven and Kirkby (1979):

$$z_{\Delta_i} = \bar{z}_{\Delta} + \frac{1}{f} \cdot (\text{CTI}_i - \overline{\text{CTI}}), \quad (2)$$

where z_{Δ_i} is the local stand specific water table depth at point i (measured relative to the soil surface, with $z_{\Delta_i} = 0$ at the surface), $\bar{z}_{\Delta}$ is the annual mean water-table depth simulated for the 0.5° grid cell, $\overline{\text{CTI}}$ is the grid cell mean CTI, and f (m^{-1}) is a soil-dependent parameter describing the exponential decline of transmissivity with depth (see Table S1 in the Supplement). Areas are considered saturated when the water table reaches the soil surface, i.e. when $z_{\Delta_i} = 0$. The corresponding threshold CTI value for saturation CTI_{sat} follows from Eq. 2:

$$0 = \bar{z}_{\Delta} + \frac{1}{f} \cdot (\text{CTI}_{\text{sat}} - \overline{\text{CTI}}) \quad (3)$$

Rearranging Eq. 3 yields:

$$\text{CTI}_{\text{sat}} = \overline{\text{CTI}} - f \cdot \bar{z}_{\Delta}. \quad (4)$$

Within each grid cell, the gamma distribution is parameterized from the stored moments of the sub-grid CTI field, namely its mean, standard deviation, and skewness (see Sect. 4.1). For a given simulated grid-cell mean water-table depth, the corresponding saturation threshold CTI_{sat} is computed from Eq. 4, following the TOPMODEL concept that local saturation is related to the sub-grid distribution of topographic wetness (Beven and Kirkby, 1979; Sivapalan et al., 1987; Kleinen et al., 2012). The saturated fraction of the grid cell is then determined as the probability that $\text{CTI}_i \geq \text{CTI}_{\text{sat}}$, i.e. as the upper tail of the prescribed gamma distribution computed from its cumulative distribution function. This probability represents the sub-grid fraction where local topographic conditions are sufficiently wet for the local water table to reach the soil surface and thus defines the inundated area of the grid cell. The resulting fraction is therefore not prescribed directly by CTI, but emerges from the combination of the prescribed CTI distribution and the dynamically simulated grid-cell mean water-table depth, and defines the simulated wetland fraction.

The calculation is repeated annually using the annual mean of $\bar{z}_{\Delta}$. Thus, temporal changes in simulated water-table depth translate into changes in wetland extent, while the CTI distribution provides the fixed sub-grid topographic constraint on where saturation is most likely to occur. The computation of $\bar{z}_{\Delta}$ is described in section 2.2.

2.2 Soil water balance

120 Soil water dynamics are represented by a multi-layer model with layer thicknesses of 200, 300, 500, 1000, and 1000 mm. Vertical soil moisture transport in this model is governed by infiltration and percolation. Water is lost from the soil column through transpiration, soil evaporation, surface and lateral runoff, and sub-surface drainage, which in LPJmL6 is influenced by interactions with groundwater (see below).

2.2.1 Infiltration

125 The processes of vertical water movement in the soil column have remained unchanged since their initial description by Schaphoff et al. (2018) and Lutz et al. (2019). However, infiltration of daily rainfall and applied irrigation amounts (P ; in mm) is now influenced by topography and is modified by accounting for a contribution to surface runoff (β_{HAG}), which depends on the mean slope gradient of the stand (β):

$$infil = P \cdot (1 - \beta_{\text{HAG}} + \alpha \cdot \beta_{\text{HAG}}), \quad (5)$$

130 where α is the factor representing the dependence on soil water conditions as described in Schaphoff et al. (2018):

$$\alpha = \left(1 - \frac{W_{\text{liq}(1)} + W_{\text{ice}(1)}}{(\theta_{\text{sat}(1)} - w_{\text{pwp}(1)}) \cdot z_{(1)}} \right)^{\frac{1}{\Phi_{\text{inf}}}}, \quad (6)$$

based on the first layer's relative volumetric soil water content at saturation ($\theta_{\text{sat}(1)}$), the relative soil water content at the wilting point ($w_{\text{pwp}(1)}$), and its actual soil water content ($W_{\text{liq}(1)}$ in mm) and the actual ice content ($W_{\text{ice}(1)}$ in mm) of the first layer. $z_{(1)}$ is the layer thickness (in mm). The factor α is scaled with Φ_{inf} , a dimensionless
 135 infiltration parameter that controls the sensitivity of infiltration to the relative soil water content of the upper soil layer and is adjusted according to the actual above-ground litter layer cover fraction as described in (Lutz et al., 2019). To improve numerical stability during vertical soil-water routing, precipitation was internally partitioned into increments of at most 4 mm, which were sequentially routed through the soil column within the daily time step. The newly added Haggard function relates the slope gradient (β) to the fraction of non-infiltrating water contributing
 140 to surface runoff (Haggard et al., 2005):

$$\beta_{\text{HAG}} = \frac{0.06 \cdot \ln(\beta + 0.1) + 0.22}{0.43}. \quad (7)$$

Infiltrated water is routed through the soil column depending on the hydraulic conductivity of the soil layer and the soil moisture of the layer below (Schaphoff et al., 2018; Lutz et al., 2019) down to the layer of the water table. The percolation of the last unsaturated layer recharges the ground water, and thus the model is able to simulate
 145 the temporal variation of the water stored in the unconfined aquifer (W_a). Lateral runoff is determined by the

groundwater–soil water interactions which is adopted from the CLM4.5 model (Riley et al., 2011; Niu and Yang, 2006; Niu et al., 2005).

2.2.2 Drainage

Subsurface drainage (i.e. groundwater discharge) (q_{drain}) is distinguished between completely and partially frozen
150 and completely unfrozen conditions of the layers below the groundwater table. In the unfrozen state:

$$q_{\text{drain}} = q_{\text{drain,max}} \cdot \exp(-f_{\text{drain}} \cdot z_{\Delta}), \quad (8)$$

where $f_{\text{drain}} = 2.0 \text{ m}^{-1}$ is the decay factor controlling the exponential reduction of subsurface drainage with increasing mean water-table depth ($\bar{z}_{\Delta}$), following the default CLM 2.0 parameterization described by Niu et al. (2005). The local maximum subsurface drainage rate ($q_{\text{drain,max}}$ at $z_{\Delta} = 0$) depends on the stand-specific topographic slope β :

$$155 \quad q_{\text{drain,max}} = \text{max}_{\text{drain}} \cdot \tan^2(\beta). \quad (9)$$

The reference parameter $\text{max}_{\text{drain}}$ is set to 50 mm day^{-1} following Christen et al. (2006). For agricultural land, $\text{max}_{\text{drain}}$ is increased to 100 mm day^{-1} and interpreted as an effective upper limit for managed drainage rather than as a site-specific drainage parameter. This reflects the assumption of enhanced water removal on agricultural land and prevents unrealistically persistent waterlogging under managed conditions. The subsurface drainage rate for the
160 at least partially frozen state is:

$$q_{\text{perched}} = \Theta_{\text{ice}} \cdot \text{max}_{\text{perch}} \cdot k_s \cdot (z_{\text{frost}} - z_{\Delta}). \quad (10)$$

Lateral drainage above the frozen soil column depends on the saturated hydraulic conductivity (k_s), and the maximum rate $\text{max}_{\text{perch}}$, equivalent to $\text{max}_{\text{drain}}$ for sub-soil dynamics, is assumed to be 50 mm day^{-1} , but is strongly limited by Θ_{ice} that gives the ice impedance factor (Swenson et al., 2012a). z_{frost} is the depth at which the soil is
165 frozen. Drainage only occur when $z_{\text{frost}} > z_{\Delta}$. Θ_{ice} is calculated from the ice content of the layer that is partly frozen. It describes the increased tortuosity of the water flow when parts of the pore space are filled with ice:

$$\Theta_{\text{ice}} = 10^{-\Omega \cdot \text{icefrac}} \quad (11)$$

and

$$\text{icefrac} = \int_0^{\text{soildepth}} F_{\text{ice}}(z) / \text{soildepth} \, dz, \quad (12)$$

170 where $F_{\text{ice}} = \frac{\theta_{\text{ice}}}{\theta_{\text{sat}}}$ is the frozen part of the water filled pore space relating θ_{sat} , the relative water content at saturation (m^3 water pro m^3 soil), to the ice content relative to the soil porosity. If the soil is saturated and all water is frozen F_{ice} is at unity. Ω is a parameter set to 6 following Swenson et al. (2012b), to meet the impedance of frozen ground.

2.2.3 Water table depth calculation

This section describes how the water table depth (and therefore the groundwater level) changes depending on
 175 the exchange of water between the saturated zone (below the water table) and the unsaturated zone (above it). The equilibrium matric potential and corresponding water content follow the hydrostatic formulation of Zeng and Decker (2009), as implemented in the Community Land Model-CLM4.5 (Oleson et al., 2013) , and used for dynamic groundwater–soil coupling and recharge calculation following Swenson and Lawrence (2014). The recharge rate q_{recharge} of the water table depth can be determined by applying Darcy’s law to z_{Δ} :

$$180 \quad q_{\text{recharge}} = -k_{\text{aq}} \cdot \frac{\Psi_{\Delta} - \Psi_{(\text{lwt})}}{z_{\Delta} - z_{(\text{lwt})}}, \quad (13)$$

where Ψ_{Δ} and $\Psi_{(\text{lwt})}$ are the soil matric potential at z_{Δ} and of the layer above z_{Δ} , where lwt is the index of this layer. k_{aq} is the hydraulic conductivity of the layer which contains the water table and is calculated as follows:

$$k_{\text{aq}} = \Theta_{\text{ice},(\text{lwt}+1)} \cdot k_s \cdot \left(\frac{\theta_{(\text{lwt}+1)}}{\theta_{\text{sat},(\text{lwt}+1)}} \right)^{2 \cdot B_{(\text{lwt}+1)} + 3}. \quad (14)$$

$\Psi_{(\text{lwt})}$ is not simply a constant, it depends on how the soil would behave under hydrostatic equilibrium at the
 185 current water–table depth (z_{Δ}). $\Psi_{(\text{lwt})}$ represents the equilibrium potential at the interface between saturated and unsaturated zones and is described by the saturated soil matric potential of the different soil types (Ψ_{sat}), the relation of the volumetric water content above the water table ($\theta_{(\text{lwt})}$ in m^3 water pro m^3 soil) and θ_{sat} :

$$\Psi_{(\text{lwt})} = \Psi_{\text{sat},(\text{lwt})} \cdot \left(\frac{\theta_{(\text{lwt})}}{\theta_{\text{sat},(\text{lwt})}} \right)^{-B_{(\text{lwt})}}. \quad (15)$$

The Clapp – Hornberger exponent ($B_{(\text{lwt})}$) is an empirical parameter that controls the steepness of the soil water
 190 retention curve (Clapp and Hornberger, 1978), describing how rapidly soil suction increases as the soil dries. When the soil is saturated ($\theta = \theta_{\text{sat}}$), the matric potential is close to zero and the soil water is under little or no tension. As the soil dries, the matric potential becomes increasingly negative, reflecting stronger capillary and adsorptive forces that hold water within smaller pores. Following the approach from Lawrence and Slater (2008), both $B_{(\text{lwt})}$ and $\Psi_{\text{sat},(\text{lwt})}$ are adjusted based on the soil’s fraction of organic material using a weighted combination approach:

$$195 \quad B_{(\text{lwt})} = (1 - f_{\text{sc},(\text{lwt})}) \cdot B_{\text{min}} + f_{\text{sc},(\text{lwt})} \cdot B_{\text{sc}} \quad (16)$$

and

$$\Psi_{\text{sat},(\text{lwt})} = (1 - f_{\text{sc},(\text{lwt})}) \cdot \Psi_{\text{min}} + f_{\text{sc},(\text{lwt})} \cdot \Psi_{\text{sc}}, \quad (17)$$

where $f_{\text{sc},(\text{lwt})}$ is the fraction of organic material in this layer and B_{min} is the Clapp – Hornberger exponent for mineral soils (TableS1) and $B_{\text{sc}} = 2.7$ for organic soils respectively Ψ_{min} the saturated matric potential for mineral
 200 soils and $\Psi_{\text{sc}} = 10.3$ for organic soils (parameters for mineral soils are given in TableS1).

Within the soil bucket, the equilibrium state between the water table and the layers above is reached when the water entering the soil layer equals the water leaving it. The adjustment of the water table reflects the balance of water fluxes between the saturated zone and the unsaturated zone.

q_{recharge} determines from the hydraulic properties (Eq. 13) if the water table rises or falls. z_{Δ} is initially set to the
 205 maximum soil depth and gradually reaches equilibrium during the spin-up simulation. This adjustment is based on changes in saturated zone storage. From the balance of the recharge rate (q_{recharge}) and drainage (q_{drain} or q_{perched} for partly frozen soils) the water table can be adjusted using the following equation:

$$\Delta z_{\Delta} = \frac{q_{\text{recharge}} - q_{\text{drain}}}{S_y}, \quad (18)$$

where S_y is the specific yield ranging between 0 and 1. It depends on the water table location and saturated soil
 210 properties:

$$S_y = \theta_{\text{sat},(\text{lwt})} \left(1 - \left(1 + \frac{z_{\Delta}}{\Psi_{\text{sat},(\text{lwt})}} \right)^{\frac{-1}{B_{(\text{lwt})}}} \right). \quad (19)$$

The stand-specific aquifer (W_a in mm) represents a vertically and spatially constrained water pool that is associated with individual stands within a grid cell. Its volume is limited to a maximum of 5000 l and is updated by the balance of q_{recharge} and q_{drain} if the water table is below the calculated soil column W_a :

$$215 \quad W_a = q_{\text{recharge}} - q_{\text{drain}}. \quad (20)$$

W_a is stand specific and becomes hydraulically active and declines only when the water table falls below the maximum soil depth or the maximum storage is not reached, allowing the compartment to store additional water or release it through drainage. For the ground water dynamics an additional pool is introduced as described in section 2.2.4

220 2.2.4 Groundwater Pool Representation and Baseflow Recession Approach

A spatially explicit groundwater storage compartment was implemented for each grid cell to account for subsurface water dynamics. The groundwater pool is recharged by two components: (i) the percolation flux, defined as the

downward drainage from the bottom soil layer, and (ii) the lateral runoff generated in the lowest saturated soil layer, representing subsurface flow contributing to groundwater recharge.

225 To simulate the temporal evolution of groundwater discharge, we apply a baseflow-recession approach (Beck et al., 2013). The dimensionless baseflow recession coefficient ($k_{\text{bf}} \in [0, 1]$) characterises the rate at which baseflow declines (Fig. A1). The specific formulation used here was developed in the ERMITAGE project (ERMITAGE project report (2013)) and is defined separately for each grid cell to account for spatial heterogeneity in hydrogeological properties such as aquifer transmissivity, soil texture and topographic gradient. High values of k_{bf} denote fast-draining soils
 230 or shallow aquifers, leading to rapid groundwater release, whereas low values represent deep or low-conductivity aquifers with slow recession and more sustained baseflow contributions. The release of groundwater to streamflow (baseflow) is described by a first-order linear recession function:

$$Q_{\text{bf}}(t) = k_{\text{bf}} \cdot GW(t), \quad (21)$$

where $Q_{\text{bf}}(t)$ denotes the baseflow at time t , $GW(t)$ is the current groundwater storage.

235 The residual groundwater storage after baseflow release is retained in the groundwater compartment and contributes to subsequent discharge events, thereby introducing a memory effect in the hydrological system. This parameterization ensures both spatial differentiation and temporal delay in the subsurface runoff component, improving the realism of long-term streamflow simulations.

In cells with irrigated agriculture the groundwater storage pool also serves as an additional source for irrigation
 240 that is accessed only when surface water availability is insufficient to meet irrigation demand.

2.2.5 Stand-specific slope calculation

As the slope has an enormous importance for the infiltration and the runoff formation, but there is no information about the exact location of different stands within the grid cell, we distinguish only the mean slope for the wetland (β_{wet}) and for the upland area (β_{up}), respectively. To derive stand-specific slopes for wetlands and uplands, we
 245 therefore make a simple statistical assumption about the subgrid distribution of slope. Within each grid cell, the subgrid slope m is treated as a random variable which is assumed to follow a negative exponential probability distribution with cumulative distribution function (CDF)

$$F(m) = 1 - \exp(-\lambda m). \quad (22)$$

The rate parameter λ is chosen such that the mean of this distribution equals the DEM-derived grid-cell mean slope
 250 (β_{mean})

$$\mathbb{E}[m] = \frac{1}{\lambda} = \beta_{\text{mean}}. \quad (23)$$

where $\mathbb{E}[m]$ denotes the expectation (statistical mean) of m . After computing the wetland area fraction (A_{wet}) of the grid cell (see section 2.1), we determine a threshold slope $m_{\text{w,max}}$ such that $F(m_{\text{w,max}}) = A_{\text{wet}}$, assuming that the fraction of the cell with $m \leq m_{\text{w,max}}$ is equal to the wetland area fraction. Using the inverse CDF of the exponential distribution this yields

$$m_{\text{w,max}} = -\frac{1}{\lambda} \ln(1 - A_{\text{wet}}), \quad (24)$$

which is then constrained to the range $[m_{\text{min}}, m_{\text{max}}]$ of the grid cell. Slopes between m_{min} and $m_{\text{w,max}}$ are assigned to the wetland stand, and slopes between $m_{\text{up,min}}$ and m_{max} to the upland (natural or agricultural) stand. The mean slope of the wetland part is obtained from the truncated exponential distribution over $[m_{\text{min}}, m_{\text{w,max}}]$ as

$$\beta_{\text{wet}} = \frac{\int_{m_{\text{min}}}^{m_{\text{w,max}}} m \lambda e^{-\lambda m} dm}{\int_{m_{\text{min}}}^{m_{\text{w,max}}} \lambda e^{-\lambda m} dm}, \quad (25)$$

and is additionally constrained not to exceed β_{mean} . In the code these integrals are evaluated analytically using the antiderivative $-\exp(-\lambda m)(m + 1/\lambda)$. Analogously, the upland mean slope β_{up} is computed as the conditional mean of the same exponential distribution over $[m_{\text{w,max}}, m_{\text{max}}]$ (see Fig. 1). The DEM-derived minimum and maximum slopes of the cell are denoted m_{min} and m_{max} . All non-rice agricultural stands are assigned the upland mean slope, whereas rice paddies are assumed to be levelled fields and therefore receive a slope of exactly zero.

2.3 Methane cycle

CH_4 dynamics within the terrestrial biosphere arise from both natural biogeochemical cycles and anthropogenic influences. Natural CH_4 emissions are predominantly driven by anaerobic microbial methanogenesis in O_2 -deprived environments such as wetlands, peatlands, and rice paddies, where the decomposition of organic matter under anoxic conditions facilitates CH_4 production. Anthropogenic processes—such as those associated with intensive livestock farming, rice cultivation, fossil fuel extraction, and land-use modifications—contribute additional CH_4 through altered substrate availability and ecosystem disruption. Within LPJmL 6.0, soil CH_4 dynamics are resolved at sub-daily time steps that simulate the CH_4 pool by accounting for concurrent production and oxidation (see Fig. 2). The CH_4 flux to the atmosphere is computed as the sum of ebullition, molecular diffusion, and plant-mediated transport. The magnitude of each pathway responds to soil moisture (saturation and air-filled porosity), substrate availability (labile carbon), and the soil redox state (oxygen availability), which together regulate both production and oxidation rates.




Figure 1. Determination of the slope values of wetland and upland stands depending on the fraction of the wetland area within a grid cell




Figure 2. Newly added or modified soil processes in the LPJmL6 model. CO_2 diffusion is assumed to be instantaneous as in earlier versions of LPJmL (Schaphoff et al., 2018) and there is no interaction of CO_2 with the soil during the diffusion process, whereas the diffusion processes of oxygen and CH_4 are explicitly modeled because the time and position in the soil column is important for the CH_4 dynamics.

2.3.1 Balance of Methane and Oxygen Concentrations in Soils

The evolution of the CH_4 concentration in the soil is characterized by the diffusive transport of CH_4 through the soil column (first part of the equation) and the production of CH_4 (methanogenesis (R_{prod})) and losses through CH_4 oxidation above the water saturated layer if enough oxygen is available (R_{oxic}), the de-gassing through forming of bubbles in water filled pores (ebullition (Q_{ebull})), and the plant mediated transport ($Q_{\text{plant,CH}_4}$) by vascular plants through their aerenchyma:

$$\begin{aligned}
\epsilon_{\text{CH}_4} \frac{\partial C_{\text{CH}_4}(z, t)}{\partial t} &= \frac{\partial}{\partial z} \left[D_{\text{CH}_4}(z) \frac{\partial C_{\text{CH}_4}(z, t)}{\partial z} \right] \\
&\quad - Q_{\text{ebull}}^{\text{out}}(z, t) + Q_{\text{ebull}}^{\text{in}}(z, t) - Q_{\text{plant,CH}_4}(z, t) \\
&\quad + R_{\text{prod}}(z, t) - R_{\text{oxic}}(z, t).
\end{aligned} \tag{26}$$

285 The O_2 concentration is calculated analogously to the CH_4 concentration equation including the O_2 consuming processes of oxic decomposition of soil organic matter (R_{CO_2}), the plant mediated transport ($Q_{\text{plant,O}_2}$), nitrification of NH_4^+ to NO_3^- ($R_{\text{NO}_3^-}$), and aerobic root respiration (R_{root}):

$$\begin{aligned}
\epsilon_{\text{O}_2}(z) \frac{\partial C_{\text{O}_2}(z, t)}{\partial t} &= \frac{\partial}{\partial z} \left[D_{\text{O}_2}(z) \frac{\partial C_{\text{O}_2}(z, t)}{\partial z} \right] - Q_{\text{plant,O}_2}(z, t) \\
&\quad - \frac{W_{\text{O}_2}}{W_{\text{C}}} R_{\text{CO}_2}(z, t) - 2 \frac{W_{\text{O}_2}}{W_{\text{CH}_4}} R_{\text{oxic}}(z, t) \\
&\quad - 2 \frac{W_{\text{O}_2}}{W_{\text{N}}} R_{\text{NO}_3^-}(z, t) - \frac{W_{\text{O}_2}}{W_{\text{C}}} R_{\text{root}}(z, t).
\end{aligned} \tag{27}$$

where ϵ_{O_2} resp. ϵ_{CH_4} are the O_2 and CH_4 porosities, that convert concentration per soil volume to the gas concentration per air volume and are calculated as

$$\epsilon_{\text{O}_2} = v + \phi \cdot \theta_{\text{sat}} \cdot B_{\text{O}_2}, \quad \epsilon_{\text{CH}_4} = v + \phi \cdot \theta_{\text{sat}} \cdot B_{\text{CH}_4}, \tag{28}$$

where ϕ is the volumetric soil water content expressed per unit pore volume (mm^3 of water / mm^3 of soil porous volume; here θ_{sat} is used as an equivalent measure of the pore-volume fraction), and v is the soil air content (m^3 air per m^3 soil). W_{C} , W_{O_2} , and W_{CH_4} are the molar masses of Carbon (12 g mol^{-1}), O_2 (32 g mol^{-1}), and CH_4 (16 g mol^{-1}), respectively. The Bunsen coefficients of O_2 (B_{O_2}) and CH_4 (B_{CH_4}) describe the solubility of these gases in water as a function of temperature. Due to the decreasing capacity of water to dissolve gases with rising temperature, this dependence is commonly expressed using exponential decay functions. Following Yamamoto et al. (1976) and Wiesenburg and Guinasso Jr (1979), we apply the following empirical relationships:

$$B_{\text{O}_2} = 0.0647 \cdot e^{-0.0257 \cdot T_{\text{soil}}}, \quad B_{\text{CH}_4} = 0.0523 \cdot e^{-0.0236 \cdot T_{\text{soil}}}. \tag{29}$$

300 The net CH_4 flux to the atmosphere, F_{atm} , represents the total upward CH_4 transfer from the soil to the atmosphere. It is considered positive when the soil acts as a net CH_4 source and negative when it acts as a net sink. This flux is composed of three contributing terms

$$F_{\text{atm}}(t) = \left(-D_{\text{CH}_4}(z) \frac{\partial C_{\text{CH}_4}(z, t)}{\partial z} \right)_{z=0} + Q_{\text{ebull}}^{\text{out}}(0, t) + F_{\text{plant}}(t), \tag{30}$$

where the diffusive flux at the soil-atmosphere interface, $-D_{\text{CH}_4}(z) \frac{\partial C_{\text{CH}_4}(z,t)}{\partial z}$, and the ebullition flux, $Q_{\text{ebull}}^{\text{out}}(z=0,t)$,
 305 escape directly to the atmosphere. Note that ebullition contributes directly to the atmospheric flux only from the
 surface layer ($z=0$), as ebullition from deeper layers is transferred to the layer above instead of escaping directly
 (see section 2.3.5). The third term, $F_{\text{plant}}(t)$, represents the flux of CH_4 transported through plant tissues (see
 section 2.3.6), particularly aerenchyma structures, which provide a low-resistance conduit for CH_4 transfer from
 anaerobic soil layers directly to the atmosphere, bypassing diffusion and ebullition pathways.

310 2.3.2 Soil organic matter decomposition into CO_2 and CH_4

Oxic decomposition of organic carbon follows the principles described by Schaphoff et al. (2018). The anoxic decom-
 position model is adapted from Khvorostyanov et al. (2008). Aerobic conditions promote the generation of CO_2 in all
 soil carbon pools and anaerobic conditions lead to CH_4 production. The soil carbon pools consist of an intermediate
 and a slow pool with turnover rates for the oxic decomposition of 0.03 and 0.001 respectively at 10°C and a fast so-
 315 called active pool with a turnover rate of 0.3 (Schaphoff et al., 2018). For realistic decomposition dynamics, Schaphoff
 et al. (2013) introduced a vertical soil distribution (Jobbagy and Jackson, 2000) which improved the simulation of
 the carbon, which can be released by permafrost thawing (see also Schaphoff et al. (2013)). Decomposition follows
 first order kinetics:

$$\frac{\partial C(z,t)}{\partial t} = R_{\text{CO}_2}(z,t) - R_{\text{prod}}(z,t) \cdot \frac{W_{\text{C}}}{W_{\text{CH}_4}}, \quad (31)$$

320 where oxic (R_{CO_2}) and anoxic decomposition (R_{prod}) is reciprocally exclusive.

Soil respiration – oxic decomposition

Oxic decomposition is computed as

$$R_{\text{CO}_2}(z,t) = \min \left[k_1(z,t) \cdot C(z,t), \frac{\partial C_{\text{max}}(z,t)}{\partial t} \right] \quad (32)$$

where C is the carbon pool size at time t and soil depth z , k_1 is the daily oxic decomposition rate, and ∂C_{max} is the
 325 mass of carbon corresponding to O_2 which is available for oxic decomposition:

$$\frac{\partial C_{\text{max}}(z,t)}{\partial t} = \text{O}_2(z,t) \cdot \frac{W_{\text{C}}}{W_{\text{O}_2}} \quad (33)$$

$\text{O}_2(z)$ is the mass of molecular soil O_2 at depth z . The oxic decomposition rate is expressed as:

$$k_1(z,t) = k \cdot g(T_{\text{soil}}(z,t)) \cdot f(\phi(z)). \quad (34)$$

330 k_1 incorporates the rate constant (k) that represents the reciprocal of the mean residence time τ for the litter components, as described by Schaphoff et al. (2018) for the litter components. Additionally, k_1 is influenced by the temperature dependency function $g(T_{\text{soil}}(z, t))$, and the soil moisture function $f(\phi_z)$ as outlined in Schaphoff et al. (2018) related to the non-frozen water. The temperature dependence function is based on an Arrhenius relationship as described in Schaphoff et al. (2018)

$$g(T_{\text{soil}}(z, t)) = \exp \left[308.56 \cdot \left(\frac{1}{56.02} - \frac{1}{T_{\text{soil}}(z, t) + 46.02} \right) \right], \quad (35)$$

335 where T_{soil} is the soil temperature in degree Celsius. The soil moisture function has its maximum around field capacity of the soil and is given by

$$f(\phi(z)) = 0.0402 - 5.005 \cdot \phi(z)^3 + 4.269 \cdot \phi(z)^2 + 0.719 \cdot \phi(z).$$

Methanogenesis – anoxic decomposition

The anoxic decomposition (R_{prod}) is calculated as follows:

$$340 \quad R_{\text{prod}}(z, t) = k_2(z, t) \cdot C(z, t). \quad (36)$$

The methanogenesis rate (k_2) is expressed as

$$k_2(z, t) = k \cdot r_k \cdot \exp \left(\frac{-C_{\text{O}_2}(z, t)}{O_2^*} \right) \cdot g(T_{\text{soil}}(z, t)). \quad (37)$$

345 $C_{\text{O}_2}(z, t)$ is the O_2 concentration at depth z , while O_2^* represents a threshold O_2 concentration of 10 g m^{-3} . As demonstrated by Duval and Goodwin (2000) methanogenesis is significantly suppressed by higher O_2 concentrations. r_k is an empirical parameter reflecting the reduced methanogenesis rate under anoxic conditions, which is considerably slower than oxic decomposition and is set to 0.1 for the fast soil carbon pool and 0.5 for the fast decomposing litter pool. Components of the carbon pool, such as lignin, degrade much more slowly in anoxic environments due to their recalcitrant structure and resistance to microbial breakdown, further limiting methanogenesis. The decomposition rate under anoxic conditions is 4–10 times lower than under oxic conditions, with reported rates ranging from 0.02 to $1.44 \text{ mg CH}_4 \text{ g}^{-1}$ over 500 days (Lee et al., 2012). This corresponds to a decomposition rate between 1.46×10^{-5} and 0.00105. Methanogenesis rate k_2 has the same temperature dependence $g(T_{\text{soil}}(z, t))$ and rate constant (k) as the oxic decomposition.

2.3.3 Methane oxidation (R_{oxic})

The representation of the CH_4 oxidation (R_{oxic}) follows the approach by Walter and Heimann (2000). R_{oxic} occurs
 355 above the saturated soil layer, where enough O_2 is available.

$$R_{\text{oxic}}(z, t) = \min \left[\frac{V_{\text{max}} \cdot W_{\text{CH}_4} \cdot C_{\text{CH}_4}(z, t)}{K_m \cdot W_{\text{CH}_4} + C_{\text{CH}_4}(z, t)} \cdot g(T_{\text{soil}}(z, t)), C_{\text{O}_2}(z, t) \cdot \frac{W_{\text{CH}_4}}{2W_{\text{O}_2}} \cdot \text{O}_{2, \text{frac}} \right], \quad (38)$$

where K_m and V_{max} are the Michaelis-Menten coefficients. These are defined in the model for $K_m = 5 \mu\text{M}$ and $V_{\text{max}} = 20 \mu\text{M h}^{-1}$ (Walter and Heimann, 2000). C_{CH_4} is the CH_4 concentration at time t and depth z in g m^{-3} . $g(T_{\text{soil}})$ is the
 360 temperature response function given in Eq. 35. V_{max} and K_m need to be multiplied by the molecular mass of CH_4 . O_2 consumption is limited by $\text{O}_{2, \text{frac}} = 0.95$, accounting for O_2 use for oxidation of reduced compounds, where 5% is assumed (Conrad, 1996; Xu et al., 2010; Saunio et al., 2020). O_2 consumption from aerobic microbial respiration, plant root respiration, nitrification and CH_4 oxidation is taken into account by reducing the O_2 content of the respective layer.

365 For the temperature dependency of CH_4 oxidation a Q_{10} relationship is used

$$g(T_{\text{soil}}(z, t)) = Q_{10}^{\frac{T_{\text{soil}}(z, t) - T_{\text{soil, mean}}(z)}{10}} \quad (39)$$

where $T_{\text{soil, mean}}$ is the annual mean soil temperature and the Q_{10} is set to 1.8.

2.3.4 Diffusion of Methane from soil (F_{diff})

CH_4 diffusion, which describes the movement of gas from higher to lower concentrations between layers and the air,
 370 is governed by Fick's law:

$$F_{\text{diff}}(z, t) = D_{\text{CH}_4}(z) \cdot \frac{\partial C_{\text{CH}_4}}{\partial x}(z, t), \quad (40)$$

where $D_{\text{CH}_4}(z)$ is the diffusivity at depth z and $C_{\text{CH}_4}(z, t)$ is the CH_4 concentration at time t and depth z . D_{CH_4} is calculated depending on soil water conditions:

$$D_{\text{CH}_4} = (D_{\text{airCH}_4} \cdot v + D_{\text{waterCH}_4} \cdot w \cdot \theta_{\text{sat}} \cdot B_{\text{CH}_4}) \cdot \eta \quad (41)$$

375 where D_{airCH_4} is the diffusivity of air ($1.82\text{e-}5 \text{ m}^2\text{s}^{-1}$) and D_{waterCH_4} the diffusivity of water ($1.6\text{e-}9 \text{ m}^2\text{s}^{-1}$ Khvorostyanov et al. (2008)). B_{CH_4} is the Bunsen coefficient for CH_4 (see Eq. 29) and $\eta = 2/3$ denotes the soil tortuosity. O_2 diffusion is calculated analogous to Eqs. 40 and 41. Initial conditions of all soil layers are:

$$C_{\text{CH}_4} = 1.207 \text{ g m}^{-3},$$

$$C_{\text{O}_2} = 266 \text{ g m}^{-3}$$

380 The lower boundary for both CH_4 and O_2 conditions are:

$$\frac{\partial C_{\text{CH}_4}(z = \text{soildepth})}{\partial z} = \frac{\partial C_{\text{O}_2}(z = \text{soildepth})}{\partial z} = 0 \quad (42)$$

and the upper boundary conditions are:

$$C(z = 0) = \frac{p_s}{R \cdot T_{\text{air}}} \cdot \text{AC}_{\text{gas}} \cdot W, \quad (43)$$

where p_s is the atmospheric surface pressure (101000 Pa), T_{air} is the air temperature (in K), AC_{gas} is the atmospheric
 385 content (0.209 for O_2 , transient over time for CH_4), W is the molar mass of O_2 and CH_4 , respectively and R is the
 universal gas constant ($8.314 \text{ J mol}^{-1} \text{ K}^{-1}$).

2.3.5 Methane ebullition (Q_{ebull})

CH_4 ebullition is the process by which CH_4 gas produced in anaerobic soil environments escapes into the atmosphere
 through gas bubbles. This occurs when the concentration of dissolved CH_4 in soil porewater exceeds its solubility
 390 threshold, causing bubble nucleation and release. The process is modulated by factors such as CH_4 concentration,
 soil moisture, and vegetation cover, which influence both the formation and migration of gas bubbles. The change in
 CH_4 concentration in a soil layer over time due to ebullition is governed by a balance between CH_4 loss via ebullition
 and CH_4 flux received from the layer below.

Ebullition is active only when the local CH_4 concentration exceeds a critical threshold required for bubble formation
 395 (Walter and Heimann, 2000), and the outgoing ebullition flux is then calculated as

$$Q_{\text{ebull}}^{\text{out}}(z, t) = k_e \cdot \max((C_{\text{CH}_4}(z, t) - C_{\text{thresh}}), 0), \quad (44)$$

where k_e is the ebullition efficiency constant controlling the rate of CH_4 release, which typically ranges between 0.1
 to 1 per day, and C_{thresh} is the threshold CH_4 concentration for bubble formation. In this study we set $k_e = 1$. The
 functional form of the threshold concentration is given by

$$400 \quad C_{\text{thresh}} = C_{\text{min}} \cdot \left(1 + 1 - \sum_{\text{pft}=1}^{n_{\text{PFT}}} (\alpha_{\text{pft}}^e \cdot \text{FPC}_{\text{pft}}) \right), \quad (45)$$

Experimental studies have shown that the presence of vegetation can substantially suppress CH₄ ebullition due to competition with plant-mediated transport and structural alterations to sediment or soil (Deshmukh et al., 2020). Therefore, to represent the vegetation-dependent reduction in ebullition flux, we introduce plant functional type (PFT)-specific suppression (α^e) factors in the model (see Tables ?? and ?? for parameter values). These account for the
 405 observed differences in ebullition across vegetated and non-vegetated systems, as supported by field measurements. FPC is the fractional plant cover of the PFT, which influences bubble formation by altering gas exchange dynamics. $C_{\min}$ is the minimum CH₄ concentration required for bubble formation under standard conditions and has been observed to range between 500 and 1000 $\cdot 10^{-6}$ M (Khvorostyanov et al., 2008) which corresponds to 0.008 to 0.016 g m⁻³. We use 0.012 g m⁻³ in LPJmL6. Eq. 44 ensures that ebullition only occurs when the gas pressure exceeds
 410 the critical pressure threshold.

The outgoing ebullition flux is transferred upwards and is added to the incoming ebullition flux $Q_{ebull}^{in}(z, t)$ in a layer above in case of $z < 0$, or to the atmosphere in case of $z = 0$. At the shallowest soil depth, there is no overlying material to impede the upward migration of gas bubbles. Therefore, the ebullition flux originating from this depth contributes directly to the CH₄ flux emitted to the atmosphere.

415 At deeper soil depths, the fate of CH₄ released via ebullition depends on the moisture content of the overlying soil column. If the overlying layer is not saturated ($\theta < 0.9$), the bubbles are trapped and remain in the overlying layer, preventing direct transfer. However, if the overlying layer is water-saturated (soil moisture > 0.9), bubbles migrate further upward and are transferred to the next layer above.

This recursive upward propagation continues through the saturated soil profile until an unsaturated soil layer
 420 is found or the surface is reached. This mechanism determines whether CH₄ from deeper soil zones contributes to atmospheric emissions or remains retained within the subsurface environment.

2.3.6 Plant-Mediated Transport (Q_{plant})

Certain vascular plants possess aerenchyma, a specialized tissue forming interconnected air-filled channels within their stems, leaves, and roots. This tissue facilitates the movement of O₂ and CH₄ between the plant and its
 425 environment, playing a crucial role in gas exchange, particularly in waterlogged or anaerobic soils. O₂ is transported from the atmosphere through the stems and leaves into the roots, supporting root respiration. Conversely, CH₄ produced in anaerobic soils can diffuse into the aerenchyma through the roots and be transported upward through the plant, being released into the atmosphere, bypassing the oxidizing zones in the soil. We hypothesize that this gas exchange via aerenchyma is more prominent in grasses compared to trees, as aerenchyma is predominantly observed
 430 in herbaceous plants and is largely absent in woody species (Takahashi et al., 2014; Colmer, 2003). To quantify these processes, we adopt the approach outlined by Wania et al. (2010) which shows that gas diffusion through aerenchyma (F_{plant}) depends on both the cross-sectional area of plant tillers available for gas transport (A_{tiller} , in m²) and the volumetric soil liquid water content (θ_{liq} , in m³):

$$F_{\text{plant}}(t) = \int_{\text{rootingdepth}}^{\text{soil surface}} Q_{\text{plant}}(z, t) \cdot dz, \quad (46)$$

435 where

$$Q_{\text{plant}}(z, t) = (C_{\text{gas}}(z, t) - C_{\text{gas,air}}) \cdot \exp\left(\frac{-k_{\text{gas}} \cdot A_{\text{tiller}}}{\theta_{\text{liq}}(z, t)}\right). \quad (47)$$

This equation is applied to both O₂ and CH₄ in the model, reflecting the bidirectional transport of these gases. A_{tiller} is defined as:

$$A_{\text{tiller}} = \max(0.001, \pi \cdot r_{\text{tiller}}^2 \cdot N_{\text{tiller}} \cdot \phi_{\text{tiller}}), \quad (48)$$

440 where r_{tiller} is the radius of an individual plant tiller defined to be $r_{\text{tiller}} = 0.0345$ m (Nascimento et al., 2021). The variable ϕ_{tiller} is the porosity of the tiller structure, expressed as a dimensionless fraction ($\phi_{\text{tiller}} = 0.7$). The use of the max function ensures that A_{tiller} maintains a minimum value of 0.001 m^2 , preventing numerical instability when the calculated tiller area becomes very small. N_{tiller} represents the tiller fraction, which depends on plant biomass and root distribution within the soil layer:

$$445 \quad N_{\text{tiller}} = C_{\text{leaf}} \cdot (\gamma / W_{\text{tiller}}) \cdot F_{\text{root}(l)}, \quad (49)$$

where C_{leaf} is the leaf carbon per square meter for PFTs that are able to form aerenchyma and γ is the phenology status of the PFT. $W_{\text{tiller}} = 0.22$ g (Wania et al., 2010) is the weight of the tillers and $F_{\text{root}(l)}$ is the fraction of the root in the respective layer.

The gas exchange coefficient, (k_{gas} in m d^{-1}), represents the piston velocity between the water and the atmosphere. 450 Following Wania et al. (2010), (k_{gas}) is obtained by scaling a reference transfer velocity for a Schmidt number (Sc) of 600, ($k_{600} = 2.07 \text{ m d}^{-1}$) with the actual Schmidt number of the respective gas:

$$k_{\text{gas}} = k_{600} \cdot \left(\frac{Sc_{\text{gas}}}{600}\right)^{n_p}, \quad (50)$$

where n_p is the Schmidt-number exponent, here set to -0.5 . Sc can be calculated specifically for O₂ and CH₄:

$$Sc_{\text{CH}_4} = 1898 - 110.1 \cdot T_{\text{air}} + 2.834 \cdot T_{\text{air}}^2 - 0.02791 \cdot T_{\text{air}}^3 \quad (51)$$

455 and

$$Sc_{\text{O}_2} = 1800.6 - 120.1 \cdot T_{\text{air}} + 3.7818 \cdot T_{\text{air}}^2 - 0.047608 \cdot T_{\text{air}}^3 \quad (52)$$

In summary, CH₄ and O₂ fluxes are based on concentration gradients, gas transfer velocities, and tiller geometry, while accounting for soil moisture, porosity, and air content. These fluxes reflect the dynamic movement of gases between the soil and atmosphere, influenced by soil properties, plant structure, and environmental conditions.

460 2.3.7 Oxygen consumption by nitrification and root respiration

Dissolved oxygen in each soil layer is consumed by nitrification of ammonium and aerobic root respiration. Nitrification is represented by the overall reaction



i.e. one mole of N nitrified requires two moles of O₂. For the nitrification rate $R_{\text{NO}_3^-}(z, t)$ (g N m⁻³ s⁻¹), the
465 associated O₂ consumption is

$$S_{\text{O}_2}^{\text{nitr}}(z, t) = 2 \frac{W_{\text{O}_2}}{W_{\text{N}}} R_{\text{NO}_3^-}(z, t), \quad (53)$$

where W_{N} is the molar mass of N (14 g mol⁻¹). In the numerical implementation $R_{\text{NO}_3^-}(z, t)$ corresponds to the depth-dependent analogue of the nitrification flux $F_{\text{NO}_3^-}$ limited by O₂, NH₄⁺ availability, soil moisture, temperature and pH (see von Bloh et al. (2018)).

470 Aerobic root respiration represents the metabolic consumption of O₂ by living roots to sustain maintenance and growth. The model provides a depth-dependent root respiration rate $R_{\text{root}}(z, t)$ (g C m⁻³ s⁻¹), which is distributed over the soil profile according to the vertical root density. Because the O₂ balance is formulated in units of O₂ mass, this carbon-based respiration flux is converted into an equivalent O₂ sink $S_{\text{O}_2}^{\text{root}}$ using a constant mass-conversion factor, yielding

$$475 \quad S_{\text{O}_2}^{\text{root}}(z, t) = \frac{W_{\text{O}_2}}{W_{\text{C}}} R_{\text{root}}(z, t). \quad (54)$$

2.3.8 Numerical scheme

Numerical solutions to the CH₄ and O₂ diffusion equations are obtained using the finite difference method for the heat diffusion problem described by Alexiades and Solomon (1993, Sec. 4.1). For time stepping the forward Euler, Crank-Nicolson and backward Euler methods were tested. The backward Euler scheme was the only method that
480 provided sufficient stability and was thus chosen for the solution of the diffusion problems. All other elements of the methane and oxygen balance, including the reaction terms, are solved using the forward Euler method.

Experiments with the time step showed that the default daily time step of LPJmL is not sufficient to obtain accurate and stable numerical solutions for the coupled methane and oxygen balance. A baseline sub-daily time step

of 4 hours was therefore adopted. Additionally, an adaptive sub-stepping scheme was implemented that rejects this
 485 baseline time step and reduces it to six 40-minutes increments whenever the baseline time step causes a relative
 change of CH₄ or O₂ concentration larger than 20% in any soil layer.

The numerical errors resulting from the temporal discretization were estimated by performing a global run of
 LPJmL6 using a smaller baseline time step of 80 minutes while leaving the adaptive sub-stepping scheme untouched.
 The global sum and temporal mean of CH₄ and O₂ concentrations in soils changed by less than 1% compared to
 490 the default 4-hour baseline time step. Global methane emissions decreased by 5.2% when switching to the increased
 temporal resolution.

2.4 Inundation feedback on vegetation

Inundation harms trees, grasses and crops that are not tolerant of water-saturated soils. These damages can cause
 (potentially lethal) productivity losses. The most important factor determining plant survival is the duration of
 495 inundation (Parolin and Wittmann, 2010). To account for the effect of inundation on vegetation we have introduced
 a feedback of inundation on gross primary production (GPP) and added three new PFTs which are tolerant to
 inundation: a tropical broadleaved evergreen flood-tolerant tree, a C3 graminoid herbaceous plant and sphagnum
 moss. The last two were already implemented in the LPJWhy model by Wania et al. (2009). The tropical tree is
 parameterized similarly to the default tropical broadleaved evergreen tree in previous LPJmL versions but with a
 500 parameterization tolerant of inundations.

Inundation areas are often not flooded constantly. To account for inundation stress we have introduced two new
 parameters as suggested by Wania et al. (2009). The inundation stress height (ist_m) defines at which water table
 depth a PFT experiences inundation stress and the inundation duration (idt_d) defines how long a PFT can endure
 inundation. The inundation-tolerant PFTs have a higher inundation stress height and endure longer inundation
 505 periods (TableS3). The stress factor (S_{inun}) is applied to GPP whenever the water table exceeds ist_m ($GPP =$
 $GPP \cdot (1 - S_{inun})$):

$$S_{inun} = \min \left(1, \max \left(0, \left[\frac{1}{1 + \exp(-k \cdot (N_{inun} - idt_d))} - 0.5 \right] \times 2 \right) \right), \quad (55)$$

where N_{inun} counts the number of days with inundation stress. TableS3 gives the parameter values of idt_d for the
 different PFTs, representing their tolerable inundation duration. Values from Wania et al. (2009) were adjusted, i.a.
 510 because PFTs TrBrEv and PoHe are not present in the implementation of Wania et al. (2009). The parameters for
 crop functional types (CFTs) in TableS4 take into account that the different CFTs suffer differently from inundation
 (Kaur et al., 2020). The parameter k (0.2 in day⁻¹) controls how rapidly stress increases once the inundation duration
 exceeds the threshold, with larger values producing a steeper rise. When the inundation threshold (ist_m) is no longer
 exceeded, the duration counter decreases in accelerated daily steps (five per day), representing a faster recovery of
 515 the plants relative to the length of the preceding inundation period.

Additionally, inundation stress increases tree mortality, contributing an additional mortality component ($\text{mort}_{\text{inun}}$) that increases the overall tree mortality rate. This contribution is capped at a maximum increase (inun_{max}) of 10% relative to the potential mortality rate (mort_{max}):

$$\text{mort}_{\text{inun}} = \text{mort}_{\text{max}} \cdot (1 - \text{inun}_{\text{max}}^{S_{\text{inun}}}), \quad (56)$$

520 where mort_{max} represents the PFT-specific baseline mortality rate as described in Schaphoff et al. (2018).

2.5 Land-use change and rice cultivation

To account for the actual land cover, the model had to be extended for land-use change on wetlands. Previous versions of LPJmL distinguish a rainfed and an irrigated stand on which rainfed and irrigated crop cultivation as well as the off season is simulated, respectively. Here we assume that rice cultivation takes place on wetland areas if natural wetlands are present in the respective grid cell. Therefore, a new wetland setaside stand is introduced from which cropland is sourced for rice cultivation upon sowing/transplanting. If not enough upland (dry) natural land is available in the cell to satisfy land use demand, wetland area can also be converted to managed grasslands and, as a last resort, to cropland for crops other than rice. Soil water on rice fields is filled up above field capacity through irrigation to simulate permanent inundation. There is no explicit simulation of drainage systems for non-rice cultivation on wetland areas. A land use overview scheme is given in the Supplement Fig. S1.

3 Modelling protocol

To demonstrate the performance of LPJmL6 to model the newly implemented process dynamics, it was initialized with a spin-up period of 4000 years, using repeated climatic data from 1901 to 1930 to simulate pre-industrial conditions and achieve dynamic equilibrium in vegetation structure and composition as well as in carbon and nitrogen pools, representing a hypothetical stable pre-industrial state. To account for historical land-use dynamics, an additional 420 years of spin-up simulation were performed between the initial spin-up and the transient historical simulation, ensuring accurate representation of land-use history and its effects on biogeochemical pools and cycles. Following this, a transient simulation was conducted, covering the period 1901 to 2019.

4 Data used

540 4.1 Input data

LPJmL6 was here driven by GSWP3-W5E5 climate data (Lange et al., 2023) at a daily time step. This dataset combines W5E5 v2.0 (Cucchi et al., 2020; Lange et al., 2021) for the period 1979–2019 with GSWP3 v1.09 (Kim, 2017) for the period 1901–1978. GSWP3 data have been bias-adjusted towards W5E5 to reduce discontinuities

at the 1978/1979 transition (Mengel et al., 2021). Annual atmospheric CO₂ concentrations are taken from the
545 TRENDY project (Friedlingstein et al., 2023). Annual atmospheric CH₄ concentrations are taken from the NOAA
Global Monitoring Laboratory (NOAA Global Monitoring Laboratory, 2025). Land-use data are generated with the
LandInG toolbox (Ostberg et al., 2023). LandInG integrates and harmonises a number of gridded source datasets such
as cropland and pasture extent from HYDE v3.2.1 (Klein Goldewijk et al., 2017a), multiple-cropping suitability from
GAEZ v3 (IIASA/FAO, 2012), and crop-specific harvested areas for the year 2000 (Monfreda et al., 2008), with census
550 data at national scale from FAOSTAT (FAO, 2020b), AQUASTAT (FAO, 2020a) and MIRCA2000 (Portmann et al.,
2010a, b) to derive gridded time series of crop-specific growing patterns (with a distinction of rain-fed and irrigated
areas) and pasture areas. As fallow land is currently not supported as a separate land-use category by LPJmL,
fallow land from LandInG is assigned to the "others" crop category in LPJmL. LandInG also integrates crop-specific
fertilizer application patterns for the year 2000 (Mueller et al., 2012; Mueller, 2012) with national-scale fertilizer
555 application trends (Hurt et al., 2020) to derive gridded time series of crop-specific nitrogen fertilizer application
rates. Manure application rates are based on Zhang et al. (2017); Zhang et al. (2017). Crop-specific growing periods
are based on the GGCM phase 3 crop calendar (Jägermeyr et al., 2021). Grazing management is simulated on
prescribed pasture areas (see above), with livestock densities based on Heinke (2025). Soil texture and soil acidity
(pH) inputs are based on HWSD v1.21 (FAO/IIASA/ISRIC/ISSCAS/JRC, 2012), which has been aggregated from
560 the source resolution by selecting parameters representative of the most prevalent soil type in each LPJmL grid cell
(Ostberg et al., 2023). River network topology is based on STN-30 (Vörösmarty et al., 2000). Prescribed lake surface
areas are based on GLWD (Lehner and Döll, 2004). Dam and reservoir characteristics are based on GRanD (Lehner
et al., 2011; Biemans et al., 2011). Water use for households, industry and livestock in each grid cell is taken from
Flörke et al. (2013) (prioritized over irrigation; accounting for 201 km³ in the year 2000). We use the ISLSCP II
565 HYDRO1k elevation-derived products at 0.5°spatial resolution (Verdin et al., 2011), which provide, for each grid
cell, statistics of the compound topographic index (CTI) including its mean, standard deviation and skewness; these
moments are used to parameterise the gamma distribution of CTI in our wetland scheme (section 2.1). The ISLSCP
II product is derived from the original 1-km HYDRO1k database (U.S. Geological Survey, 2007b). The slope values
are taken directly from the USGS fine-resolution 30-arc-second GTOPO30 slope product (U.S. Geological Survey,
570 2007a), from which we derive the mean, maximum and minimum slope of each 0.5°cell for the slope-dependent
wetland parameterisation.

4.2 Validation data

A combination of geographically detailed data is used to evaluate the LPJmL6 model’s ability to simulate wetland
extent. Kaplan (2007) provides a global wetland map at 0.5 degree resolution, combining regional and global invento-
575 ries, geomorphic and hydrological data, and satellite observations. It broadly defines wetlands, including permanently
and seasonally inundated areas, and waterlogged soils. The Global Lakes and Wetlands Database (GLWD) by Lehner
and Döll (2004) offers high-resolution (1 km) data on lakes, reservoirs, and wetlands, classifying water bodies by

size, hydrology, and ecological features. While we use the GLWD lake area as an input to the model (section 4.1) we use the GLWD wetland extent for validating the simulated global dynamics of wetland extent. Gumbrecht et al. (2017) present the SWAMP dataset, a high-resolution (231 m) map of tropical and subtropical wetlands, integrating geomorphic classification, hydrological modeling, and satellite imagery. Zhang et al. (2021) developed WAD2M, a dataset on monthly wetland extent for 2000–2018 at 25 km resolution (using the mean annual extent), focused on permanently vegetated wetlands by excluding permanent water bodies, coastal wetlands, and rice paddies.

Estimates of CH₄ emissions from rice cultivation were derived from a combination of empirical field data, synthesis studies, and global-scale inverse modeling frameworks, as follows. The resulting CH₄ emission values represent cumulative seasonal emissions (in gCH₄ m⁻²) from flooded rice systems, calculated by multiplying observed or reported mean daily CH₄ emission rates (gCH₄ m⁻² day⁻¹) with region-specific growing season lengths. This approach allows for spatially explicit interregional comparison of seasonal CH₄ release, facilitating benchmarking of biogeochemical models and identifying emission hotspots. A comprehensive synthesis by Wang et al. (2018) compiled over 600 site-years of CH₄ flux data collected from 1990 to 2014, encompassing diverse rice-growing environments and management regimes. Their dataset provided country-level mean emission rates and associated 95 % confidence intervals, reflecting spatial heterogeneity across sites. These values were further stratified by continent and climate zone to support global upscaling. Complementing this synthesis, Zhu and Li (2024) incorporated more recent CH₄ flux observations from East Asia, including multi-cropping systems and long-term experimental sites. Their dataset included some of the highest recorded emissions, particularly in southern China, and expanded the representativeness of regional variability. In South Asia, Gupta et al. (2021) reported empirical data on daily CH₄ fluxes from irrigated rice paddies in northern India under varying water and residue management practices, highlighting the mitigation potential of alternate wetting and drying regimes. Earlier field-based estimates provided by Wassmann et al. (2000) focused on major rice ecosystems in Southeast Asia, using controlled experiments and standardized chamber measurements under the IRRI-GTZ collaboration. Their regional emission ranges from the 1990s remain influential for emission factor development and baseline scenario assessment.

While these sources generally provide spatial uncertainty bounds, they do not fully capture interannual variability, which may be significant under shifting climate and agronomic conditions. To contextualize and constrain bottom-up estimates, we integrated global atmospheric inversion data from the CarbonTracker-CH₄ framework (NOAA Global Monitoring Laboratory, NOAA Global Monitoring Laboratory (n.d.); Bruhwiler et al. (2014)). This system assimilates atmospheric CH₄ concentrations from NOAA’s global flask and in situ measurement network (NOAA Global Monitoring Laboratory, 2025) to infer regional and global emission patterns. Wetland and rice cultivation emissions were treated as distinct sectors, enabling partial attribution of seasonal CH₄ enhancements to rice-growing regions in Asia. Additional global source data and attribution uncertainty ranges were taken from the Global Methane Budget reports by Saunois et al. (2020, 2025), which combine bottom-up inventories, top-down atmospheric inversions, and process-based models. To analyze how CH₄ emissions vary with latitude across different datasets, the LPJmL6 model is compared with the WetCHARTs v1.0 dataset provided by Bloom et al. (2017a). WetCHARTs (Wetland

and Climate CH₄ Intercomparison of Models) offers a global dataset of wetland CH₄ emissions at a 0.5° resolution, along with associated uncertainties. This dataset integrates multiple emission models, remote sensing data, and atmospheric observations to provide a comprehensive and harmonized view of CH₄ emissions from wetlands.

Monthly river discharge observations were used to evaluate monthly simulated river discharge at the basin scale. The observational compilation comprises 285 gauging stations with reported upstream drainage areas exceeding 10,000 km². The dataset was assembled by combining (i) pan-Arctic discharge records from R-ArcticNET (v4.0) (Lammers et al., 2001, 2016) and (ii) global monthly discharge records from the RivDIS database (v1.1) distributed by the ORNL DAAC (NASA Earthdata) (Vörösmarty et al., 1998), where the latter was restricted to stations belonging to the dataset’s major basins.

We used eddy-covariance flux tower data to benchmark simulated fluxes of net ecosystem exchange (NEE), evapotranspiration (ET), and methane (CH₄), providing direct and continuous measurements of ecosystem-atmosphere exchanges of carbon dioxide, CH₄, and water vapor at the ecosystem scale. In this study, tower-derived NEE, CH₄, and ET were compared against the corresponding model outputs. While the evaluation allows for a direct assessment of the model’s ability to reproduce observed flux dynamics, it is subject to several limitations. These include differences in spatial scale between the grid-based simulations and the footprint of the flux towers, potential mismatches in vegetation composition due to the model’s dynamic PFT representation, and the use of large-scale meteorological forcing instead of site-specific weather observations. For CH₄ in particular, this scale mismatch can be especially relevant because tower footprints often represent small, locally heterogeneous wetland areas that may not be representative of the grid-cell mean wetland fraction and hydrological conditions simulated by the model.

To assess vegetation structure, we contrasted LPJmL6 PFT distributions with the ESA CCI-derived product of Harper et al. (2023), a 29-year (1992–2020) 300 m dataset that harmonizes land cover into fractional PFTs, enabling scale-aware, spatially explicit comparisons with model output (Harper et al., 2023).

For system-level context, we drew on the Global Carbon Budget (GCP, Friedlingstein et al., 2023), which synthesizes atmospheric CO₂ growth, ocean uptake, and land-sink estimates together with fossil and land-use emissions and reports an annual budget imbalance as an integrated constraint. We also applied the International Land Model Benchmarking framework (ILAMB), detailed at <https://www.ilamb.org/>, to place our results in a standardized multi-metric context (bias, RMSE, seasonal phase/amplitude, spatial distribution) and to summarize performance with an overall score (see Collier et al., 2018 and ILAMB documentation).

5 Model evaluation

5.1 ILAMB comparison

As part of our evaluation process, we conducted an extensive benchmarking analysis using the ILAMB framework, [full report is available at Zenodo: <https://doi.org/10.5281/zenodo.17877397>]. The ILAMB project provides a standardized system for assessing the performance of land surface models against a variety of observational datasets.

Table 1. Wetland area (in 10^6 km²) according to different data products, compared to simulations with LPJmL6 (with land-use change) and LPJmL6-PNV (potential natural vegetation only).

	GLWD	SWAMP	Kaplan (2007)	Zhang et al. (2021)	LPJmL6	LPJmL6-PNV
Global	11.4	6.0	5.7	3.6	5.6	5.6
>60°N	2.3		1.1	0.1	0.2	0.2
>30°N - 60°N	4.8	0.7*	1.9	1.2	1.2	1.5
<30°N	4.3	5.3	2.7	2.3	4.2	3.9

* area only partly represented

This framework is widely recognized for its ability to objectively evaluate model fidelity across multiple environmental and ecological variables, including carbon, water, and energy fluxes. For our analysis, we applied the ILAMB methodology to the LPJmL6 model in comparison to LPJmL5.10 shown for separate simulations with actual land use and for potential natural vegetation, producing comprehensive benchmarking outputs to compare its performance against observational data. This approach ensures a rigorous and transparent evaluation, highlighting both strengths and areas for improvement in LPJmL6’s ability to simulate terrestrial ecosystem processes. The results of this benchmarking exercise serve as a valuable reference point for future model development and intercomparison efforts, contributing to the broader goal of improving Land Surface Model performance.

5.2 Wetland area and distribution

Wetland distribution simulated by LPJmL6 depend on the CTI of the cell and the simulated water table depth as described in section 2.1. In the comparison of simulated and observed wetland area we do not exclude grid cells with rice cultivation. Because rice is typically established on wet or formerly wet soils, an expansion of rice area can therefore appear as an increase in wetland area in our evaluation against simulations for natural vegetation (LPJmL6-PNV), even though these sites often represent wetlands that have been converted to cropland. The simulated wetland area is overall comparable to other independent estimates (Table 1), but systematic discrepancies between datasets are evident. SWAMP includes only the wetland area south of 39°N, which explains the absence of high-latitude wetlands in this dataset and its focus on tropical regions, particularly the Amazon (Gumbricht et al., 2017). In contrast, GLWD indicates by far the greatest extent north of 60°N, reflecting its strong emphasis on boreal and permafrost wetlands (Lehner and Döll, 2004). LPJmL6 produces a much lower extent in these regions, closer to the estimates by (Zhang et al., 2021), yet this underestimation of boreal wetlands compared to GLWD indicates that LPJmL6 does not translate permafrost-induced flooding into wetlands. Conversely, LPJmL6 is able to reproduce the strong tropical maximum in the Amazon region that dominates the SWAMP dataset.

The latitudinal distribution (Fig. 3) highlights these contrasting signals: GLWD exhibits a pronounced peak in the high northern latitudes, SWAMP primarily represents tropical wetlands, and LPJmL6 shows a more balanced

670 distribution with clear tropical hotspots and additional mid-latitude wetlands in Eurasia. These differences also have
 direct implications for global methane budgets, as both tropical floodplains and high-latitude permafrost wetlands
 are recognized as key but highly uncertain contributors to natural CH₄ emissions (Saunois et al., 2020; Hugelius
 et al., 2020). The strong disagreement in global totals across models and observational datasets (ranging from 3.6 to
 11.4×10^6 km²) underlines the need for improved observational constraints and harmonized definitions of wetlands.
 675 To complement the latitudinal evaluation, we provide additional spatial and topographic diagnostics in the Supple-
 ment. These include maps of wetland fraction and absolute differences between LPJmL6 and four observation-based
 or inventory-based wetland products (Fig. S2) as well as binned comparisons of the wetland-fraction anomaly against
 mean grid-cell slope and CTI (Figs. S3–S4). These diagnostics indicate where the largest spatial disagreements occur
 and whether they are systematically associated with the topographic controls used in the wetland-fraction parame-
 680 terization.




Figure 3. Latitudinal distribution of zonal wetland area from different data products and from LPJmL6 (with land-use change) and LPJmL6-PNV (potential natural vegetation). Wetland area is integrated along longitude for each half degree latitude band (km²).

5.3 Methane budget

In this section, we confront the land CH_4 budget simulated by LPJmL6 with independent constraints on the global CH_4 budget. The global budget is determined by the balance between all surface sources and atmospheric sinks of CH_4 and is constrained by the observed atmospheric CH_4 burden, its temporal growth and estimates of the atmospheric CH_4 lifetime (Stevenson et al., 2020; Saunois et al., 2020; He et al., 2021; Saunois et al., 2025). These constraints define a range of total CH_4 emissions and, by subtraction of quantified source categories, a residual range attributable to net land sources.

LPJmL6 represents a subset of the global CH_4 cycle, namely emissions from natural wetlands, rice cultivation, wildfires and other agricultural soils, as well as aerobic soil CH_4 uptake. We refer to the sum of these components (emissions minus soil uptake) as the LPJmL6 land CH_4 budget. To relate this modelled land budget to the global CH_4 budget, we combine LPJmL6 fluxes with independent estimates of other anthropogenic and natural sources that are not simulated by LPJmL6 (enteric fermentation and manure management, landfills and waste, fossil fuel-related emissions, biomass and biofuel burning, and additional natural sources). This allows a consistent comparison of LPJmL6 with bottom-up (BU) and top-down (TD) global estimates and with the lifetime-based constraints on total emissions.

5.3.1 Methane emissions

The global constraints on CH_4 sources and sinks outlined above allow a more detailed comparison of LPJmL6-simulated emissions with other global estimates. LPJmL6 simulates CH_4 emissions from wetlands, rice fields, and wildfires, and can distinguish between emissions from agricultural land and natural land. Furthermore, LPJmL6 explicitly represents soil CH_4 uptake via diffusion from the atmosphere into the soil. To derive the possible range of CH_4 emissions compatible with observations, source categories that are not explicitly calculated by LPJmL6 have to be added. These include anthropogenic emissions from enteric fermentation and manure management, and from landfills and waste (about 170–205 $\text{TgCH}_4 \text{ a}^{-1}$), as well as fossil fuel-related emissions (about 145 $\text{TgCH}_4 \text{ a}^{-1}$) (NOAA Global Monitoring Laboratory, 2025) and biomass combustion (about 28 $\text{TgCH}_4 \text{ a}^{-1}$). Together, these account for 462–481 $\text{TgCH}_4 \text{ a}^{-1}$ (Saunois et al., 2025, 2020; Li et al., 2021; Stanley et al., 2016). Additional natural sources not represented in LPJmL6, such as emissions from freshwater systems, geological sources, termites and the ocean, account for a further 64–259 $\text{TgCH}_4 \text{ a}^{-1}$, with particularly large uncertainty for freshwater emissions. For the illustrative period 2010–2019, the atmospheric CH_4 sink is estimated to be between 523 and 649 $\text{TgCH}_4 \text{ a}^{-1}$.

Depending on the estimated CH_4 lifetime in the atmosphere given by Stevenson et al. (2020), between 31 and 242 $\text{TgCH}_4 \text{ a}^{-1}$ remain attributable to land net sources, for which LPJmL6 simulates about 142.1 $\text{TgCH}_4 \text{ a}^{-1}$ (wetlands, fire, rice cultivation, and other agricultural soil emissions and the explicitly simulated soil sink; mean over the years 2000 to 2019; see Table 3). If, however, we enforce global budget consistency by combining the lower (upper) estimates of the atmospheric sink with the lower (upper) estimates of CH_4 sources not simulated by LPJmL6, the




Figure 4. Global methane budget calculated from the atmospheric methane concentration and estimated lifetime of methane and the atmospheric growth resulting in the estimated overall methane sink that gives the range of the assumed methane emissions. Methane sources are anthropogenic (enteric fermentation& manure, landfills & waste, biomass & biofuel burning) and natural (incl. freshwater, geological, wildfires, termites, oceanic) (Saunio et al., 2025) and the range of the atmospheric methane lifetime is given by Stevenson et al. (2020). The two blue lines (labelled lifetime = 8 yr and lifetime = 10 yr) and the blue shaded area between them represent the atmospheric constraint, indicating the range of total global methane emissions, i.e. the region in which the sum of all sources and the soil sink should lie for a closed global methane budget. Bottom-up (BU) and top-down (TD) estimates, given as decadal means for 2000–2009 and 2010–2019, are converted to continuous annual values by linear interpolation between these two means; the resulting BU and TD time series are added to the simulated LPJmL6 methane sources and define the range shown by the orange and dark red lines. LPJmL6 simulated net methane sources include emissions from wetlands, rice cultivation, and wildfires and the soil sink.

residual range attributable to land net sources narrows to 116–157 TgCH₄ a^{−1}, which still encompasses the LPJmL6
715 estimate (see Fig. 4).

When all emissions simulated by LPJmL6 are combined with estimates for the remaining anthropogenic and natural sources, the resulting overall CH₄ budget matches well with the observationally constrained overall emissions Fig. 4. We construct two such combined estimates: one using bottom-up (BU) values for the non-LPJmL sources and one using top-down (TD) values. Adding the same LPJmL6 land emissions to these two source data sets shifts
720 both curves upwards by the modelled land flux; in our set-up this makes the BU-based combination systematically

higher and the TD-based combination lower. Consequently, the BU+LPJmL6 series forms the upper envelope and the TD+LPJmL6 series the lower envelope of the total-budget range shown by the red and orange lines in Fig. 4, respectively.

Table 2 summarizes CH₄ emissions from wetlands as estimated by the Global Methane Budget project (Kirschke et al., 2013; Saunio et al., 2016, 2020, 2025). For the period 2000–2009, BU estimates range between 147 and 175 TgCH₄ a^{−1}, whereas TD estimates range from 180 to 217 TgCH₄ a^{−1}. The discrepancy between these methodologies is evident across all decades examined. Notably, the differences are largest in the 1980s and 1990s, exceeding 50 TgCH₄ a^{−1}. More recent estimates show reduced discrepancies, with TD estimates decreasing significantly while BU estimates remain relatively stable. LPJmL6 aligns closely with BU estimates, reflecting its process-based methodology and explicit integration of wetland hydrology.

Table 2. Global methane emissions in TgCH₄ a^{−1} from natural wetlands across four decades, presenting values separately for bottom-up (BU) and top-down (TD) methods. The variation reflects the methodological differences and uncertainties in emission processes.

References	1980-1989		1990-1999		2000-2009		2003-2012		2010-2019	
	BU	TD	BU	TD	BU	TD	BU	TD	BU	TD
Kirschke et al. (2013)	167 ⁺⁶⁴ _{−52}	225 ⁺⁴¹ _{−42}	150 ⁺¹⁰ _{−6}	206 ⁺⁵⁹ _{−37}	175 ⁺³³ _{−33}	217 ⁺⁶⁷ _{−40}	167 ⁺³⁵ _{−40}	185 ⁺¹¹² _{−32}	159 ⁺⁴⁴ _{−40}	165 ⁺²⁰ _{−49}
Saunio et al. (2016)					166 ⁺³⁸ _{−41}	183 ⁺³⁹ _{−32}				
Saunio et al. (2020)					147 ⁺³² _{−45}	180 ⁺¹⁶ _{−27}				
Saunio et al. (2025)					153 ⁺³⁶ _{−37}	158 ⁺¹⁴ _{−13}				
This study*	126		134		155		164		180	

* here we report data from the simulation with actual vegetation, but exclude emissions from rice fields for consistency with the reference data

The site-level comparison, shown in the Supplement (Figs. S5–S21), indicates that LPJmL6 reproduces the observed seasonal and cumulative CH₄ emissions at some sites, for example RU–Ch2, US–Tw1, and US–Myb, but performs less well at others. This behaviour is expected because eddy-covariance CH₄ measurements often represent relatively small and heterogeneous source areas, whereas LPJmL6 simulates grid-cell mean fluxes. Individual tower footprints may therefore be dominated by local wetland patches, drainage features, vegetation composition, management, or water-table dynamics that are not representative of the corresponding 0.5° grid cell. Consequently, mismatches at individual sites do not necessarily indicate a failure of the large-scale model process representation, but may reflect a scale mismatch between local observations and grid-cell averages.

The cumulative comparison is often more robust than the month-to-month comparison because it integrates over short-term variability and gaps in the observations. However, the FLUXNET-CH₄ records are frequently incomplete, and missing periods can affect both monthly sums and cumulative totals, especially when data gaps occur during

peak emission periods. Therefore, the site-level evaluation should mainly be interpreted as a diagnostic of seasonal timing and order-of-magnitude agreement, rather than as a strict validation of local flux magnitudes at every site.

Figure 5 compares the spatial and latitudinal distribution of wetland CH₄ emissions simulated by LPJmL6 with the WetCHARTs ensemble (Bloom et al., 2017b). Both data sets show a pronounced tropical maximum in the latitudinal profile, reflecting the dominant contribution of tropical wetlands to the global wetland CH₄ emissions. In terms of zonally summed emissions, LPJmL6 agrees reasonably well with the WetCHARTs ensemble median across broad latitude bands and remains mostly within the WetCHARTs min–max range, particularly outside the tropical peak region.

However, the spatial comparison indicates that similar latitudinal totals can arise from different regional emission patterns. The anomaly map shows regional offsets between LPJmL6 and the WetCHARTs median, including areas where LPJmL6 underestimates or overestimates emissions relative to WetCHARTs despite comparable zonal sums. This suggests that agreement in the latitudinal profile should not be interpreted as full spatial agreement. Part of this mismatch is expected because the comparison is made only against the WetCHARTs ensemble median, while the ensemble spread reflects substantial uncertainty in wetland extent and CH₄ flux estimates. In addition, LPJmL6 is run on a coarse spatial grid and does not explicitly represent bank infiltration or river–floodplain exchange processes, which limits its ability to resolve narrow floodplains, riparian wetlands, and seasonal inundation features that can contribute substantially to local CH₄ emissions.

5.3.2 Methane emissions from the different flux components

Quantifying the exact global contribution of each pathway is challenging because of spatial and temporal variability in wetland extent, hydrology and vegetation, as well as differences in wetland types and environmental conditions. Table 3 therefore summarises the LPJmL6 global methane budget separated into atmospheric flux pathways (diffusion, plant-mediated transport, ebullition and fire emissions) and source categories (natural wetlands, rice fields, grasslands and other cropland), together with soil oxidation and the resulting net methane sink. This pathway- and source-resolved budget provides a process-level decomposition of the simulated CH₄ emissions and complements the spatial evaluation against WetCHARTs. It allows the relative importance of the main atmospheric transport pathways to be assessed separately from the contributions of different land-use and wetland source categories. In this way, the table provides an additional diagnostic for interpreting the model behaviour beyond the total wetland CH₄ flux and the latitudinal emission patterns.

The atmospheric flux partitioning shows that diffusion remains by far the dominant CH₄ emission pathway, with global emissions increasing from 117.6 to 157.5 Tg CH₄ a^{−1} between the 1980s and 2010–2019. Plant-mediated transport roughly doubles from 19.5 to 40.6 Tg CH₄ a^{−1} over the same period, whereas ebullition remains comparatively small (about 2 Tg CH₄ a^{−1}) and exhibits no clear trend, in line with the minor contribution of ebullition reported by Bieniada et al. (2021). Fire emissions are simulated at around 10 Tg CH₄ a^{−1}, consistent with independent estimates




Figure 5. Spatial and latitudinal comparison of wetland CH_4 emissions simulated by LPJmL6 with the WetCHARTs ensemble (Bloom et al., 2017b). Maps show (a) LPJmL6 wetland CH_4 emissions, (b) the WetCHARTs ensemble median, and (c) the anomaly between LPJmL6 and WetCHARTs, calculated as LPJmL6 minus the WetCHARTs median. Map values are shown as annual flux densities in $\text{g CH}_4 \text{ m}^{-2} \text{ yr}^{-1}$. The latitudinal profile shows area-weighted zonal sums in $\text{Tg CH}_4 (0.5^\circ)^{-1} \text{ yr}^{-1}$. The WetCHARTs min-max range is shown as a shaded band, and the ensemble median is shown as a solid line.

Table 3. LPJmL6 estimates of global methane balance differentiated for different components in $\text{TgCH}_4 \text{ a}^{-1}$.

	1980-1989	1990-1999	2000-2009	2010-2019
diffusion	117.6	125.1	141.9	157.5
plant mediated transport	19.5	21.3	28.2	40.6
ebullition	1.8	2.3	2.2	2.7
fire emissions	9.9	10.3	9.8	9.8
total emissions	148.8	159.0	182.1	210.5
net methane sink	-47.0	-51.6	-53.3	55.1
emission on the remaining natural wetlands	117.1	123.6	141.9	166.0
emission from rice fields	13.3	14.9	17.5	20.9
emission from grasslands	6.7	7.5	8.6	9.4
emission from the remaining cropland	2.0	2.7	4.2	4.5
oxidation through the soil	137.0	145.5	148.3	147.6

775 of global wildfire CH_4 emissions of approximately $10 \text{ Tg CH}_4 \text{ a}^{-1}$ and contributing about 5% of the total natural source (Saunois et al., 2025).

The soil methane sink in LPJmL6 can be compared with independent constraints. Murguia-Flores et al. (2018) estimated a global soil CH_4 sink of $33.5 \text{ Tg CH}_4 \text{ a}^{-1}$ for 1990–2009, whereas LPJmL6 simulates a stronger net uptake of about $52 \text{ Tg CH}_4 \text{ a}^{-1}$ over the same period (Table 3). A key difference is that LPJmL6 does not explicitly represent
780 the atmospheric CH_4 sink associated with methanotrophic activity in well-aerated soils; instead, CH_4 oxidation is treated as part of the soil CH_4 balance. The gross soil oxidation flux in LPJmL6 is close to $150 \text{ Tg CH}_4 \text{ a}^{-1}$, reflecting the substantial fraction of CH_4 that is oxidized while diffusing through the unsaturated zone. Despite these structural differences, the simulated net soil sink lies within, albeit towards the upper end of, the range of $9\text{--}51 \text{ Tg CH}_4 \text{ a}^{-1}$ reported by atmospheric inversions and bottom-up estimates (Saunois et al., 2025).

785 Because LPJmL6 explicitly represents both natural and managed ecosystems, including land use and land-use change, it can attribute global CH_4 emissions to distinct source categories. In our simulations, emissions from natural wetlands increase from 117 to $166 \text{ Tg CH}_4 \text{ a}^{-1}$ between the 1980s and 2010–2019, while emissions from rice fields rise from 13.3 to $20.9 \text{ Tg CH}_4 \text{ a}^{-1}$. This corresponds to an increase of about 57% in rice-field emissions compared to only about 13% growth in harvested rice area. Emissions from other agricultural land (excluding rice and grasslands)
790 grow from 2.0 to $4.5 \text{ Tg CH}_4 \text{ a}^{-1}$, and grassland emissions from 6.7 to $9.4 \text{ Tg CH}_4 \text{ a}^{-1}$. For comparison, EPA (2012) estimated that other agricultural CH_4 sources contribute roughly $20 \text{ Tg CH}_4 \text{ a}^{-1}$ for 2005–2020, which is higher than the LPJmL6 estimate but broadly consistent when considering that the EPA category also includes savanna and residue burning. Overall, all simulated source categories exhibit increasing CH_4 emissions over recent decades, likely reflecting the combined effects of climate change and evolving land management; however, the relative contributions
795 of these drivers warrant further investigation.

Another important aspect of LPJmL6 is the explicit representation of both natural and managed ecosystems, including land-use and land-use change dynamics. This structure allows us to attribute simulated changes in global CH_4 emissions to distinct source categories and regions across the terrestrial surface and to link emerging trends to underlying drivers. Such source-resolved information is essential for understanding, and ultimately mitigating,
800 the role of wetlands and agricultural systems in global greenhouse gas budgets. The LPJmL6 model allows us to separately quantify global CH_4 emissions from natural wetlands, rice fields, grasslands and other agricultural land. In the simulations, emissions from natural wetlands increase from $117 \text{ Tg CH}_4 \text{ a}^{-1}$ in the 1980s to $166 \text{ Tg CH}_4 \text{ a}^{-1}$ in 2010–2019. Emissions from rice fields rise from 13.3 to $20.9 \text{ Tg CH}_4 \text{ a}^{-1}$ over the same period, corresponding to an increase of about 57%, whereas harvested rice area increases by only about 13%. Emissions from other
805 agricultural land (excluding rice and grasslands) grow from 2.0 to $4.5 \text{ Tg CH}_4 \text{ a}^{-1}$, and grassland emissions from 6.7 to $9.4 \text{ Tg CH}_4 \text{ a}^{-1}$. Over the same period, simulated global grassland area increases by only about 5%, implying that the roughly 40% rise in grassland CH_4 emissions is driven primarily by changing environmental conditions and management rather than by areal expansion alone. For comparison, EPA (2012) estimated that other agricultural CH_4 sources contribute roughly $20 \text{ Tg CH}_4 \text{ a}^{-1}$ for 2005–2020, which is higher than the LPJmL6 estimate but

810 broadly consistent when considering that the EPA category also includes savanna and residue burning. Overall,
all simulated source categories exhibit increasing CH₄ emissions over recent decades, consistent with the combined
effects of changing climate conditions and evolving agricultural practices; however, the relative contributions of these
drivers require further investigation.

5.3.3 Emission from rice cultivation, grasslands and other agricultural land

815 Table 4 summarizes literature estimates of global CH₄ emissions from rice cultivation, aggregated over the growing
seasons, together with the corresponding LPJmL6 simulations. Most literature values are multi-year means for the
period indicated in the table (rather than single-year snapshots). They are based on a range of methodological
approaches, including emission-factor inventories, statistical upscaling from field measurements, and FAOSTAT-
based accounting, and span roughly 18.3 to 38.8 TgCH₄ a⁻¹. LPJmL6 simulates an increase from a global mean of
820 13.0 TgCH₄ a⁻¹ in the 1980s to 20.4 TgCH₄ a⁻¹ during 2010–2019, which places our results toward the lower end
of the published range (Table 4). A likely explanation is that the current representation of rice production systems
assumes one cropping season per grid cell, whereas multi-cropping (double or even triple rice) is widespread in parts
of South and Southeast Asia and eastern China (Waha et al., 2025, 2020). In such regions, annual emissions exceed
the values simulated by LPJmL6 because emissions from multiple growing seasons within a calendar year are not
825 explicitly represented. In annual sums of LPJmL6 simulated emissions, we combine planted rice area with those
from wetland-setaside emissions on these fields, which captures major biogeochemical controls but may still under-
represent management effects such as straw incorporation, organic amendments, and deliberate drainage. In the
LPJmL6 simulations, the unproductive (non-growing) period contributes about 4 TgCH₄ a⁻¹. While the emission
estimates obtained with other methodologies are strongly influenced by how the contributing days are defined, the
830 very wide range of 8–78 Tg CH₄ a⁻¹ simulated by by the process-based model CH4MOD (Hu et al., 2024) is mainly
driven by differing assumptions about management.

Table 4. Global CH₄ emissions from rice cultivation during the growing season

Source	Global Emissions (TgCH ₄ a ⁻¹)	years	Methodology
Yan et al. (2009)	18.3–38.8	2000	IPCC Tier 2, based on global EF database
Hu et al. (2024)	8–78	2008–2017	Process-Based Model (CH4MOD) under different management
Pazhanivelan et al. (2024)	19.9	2017–2018	Tier 2/Remote Sensing
FAO (2020c)	24.4	2010–2019	Modeled emissions based on FAOSTAT data
EPA (2012)	18.4	2015	IPCC-based bottom-up modeling
Wang et al. (2023a)	27 ± 6	2008–2017	IPCC Inventory Guidelines (Tier 2/3)
Saunois et al. (2025)	25–37	2010–2019	Integrated from national inventories
This study	20.9	2010–2019	simulated by LPJmL6

Table 5. Country-level cumulative CH₄ emissions from rice fields during the growing season simulated by LPJmL6 (1990–2014 mean, gCH₄ m⁻² a⁻¹). For each country, the table reports the mean, minimum, and maximum emissions within the national rice area.

Country	Mean	Min	Max
Bangladesh	13.62	3.46	39.24
Brazil	75.84	4.21	222.90
China	17.62	0.77	52.91
India	11.18	0.94	34.08
Indonesia	33.58	0.35	238.21
Italy	19.52	0.06	104.86
Japan	7.83	0.01	48.53
Philippines	3.45	0.58	9.42
South Korea	10.15	0.09	29.94
Spain	10.99	0.22	38.08
Uruguay	110.20	3.10	328.15
Vietnam	21.18	1.04	46.90

Across rice-growing regions, methane emissions from rice fields exhibit strong spatial and temporal variability. LPJmL6 simulates country-mean cumulative growing-season emissions that range from a few to more than 100 gCH₄ m⁻² a⁻¹, with local grid-cell values reaching up to 328 gCH₄ m⁻² a⁻¹ in the most strongly emitting areas (Table 5). The observed values compiled in Table 6 span a comparable order of magnitude, with maximum values extending from about 13 up to 347.5 gCH₄ m⁻² a⁻¹, indicating pronounced sub-national heterogeneity within the sampled regions. Independent field studies further highlight the very strong variability of rice CH₄ emissions. Wang et al. (2022) report substantial year-to-year differences in growing-season emissions from Chinese rice fields, with high emissions of 36.8 gCH₄ m⁻² a⁻¹ in 2018 and low emissions of 5.3 gCH₄ m⁻² a⁻¹ in 2019, while Qian et al. (2023) document even higher plot-scale values approaching 580 gCH₄ m⁻² a⁻¹ under specific conditions, illustrating the potential for extreme fluxes in certain regions or management settings. It is important to note that LPJmL6 always provides spatially complete coverage of each national rice area, whereas most literature values in Table 6 are derived from individual sites or sub-regions and therefore cannot be interpreted as true national means, even in the comparatively extensive dataset of Wang et al. (2018). This mismatch in spatial representativeness, together with differences in local management practices, environmental conditions and measurement protocols, explains part of the discrepancies between simulated and observed ranges, while the overall overlap in magnitude supports the ability of LPJmL6 to capture first-order patterns of rice CH₄ emissions.

Table 6: Observed cumulative CH₄ emissions from rice fields during the rice-growing season (gCH₄ m⁻² a⁻¹), compiled from field studies and grouped by region. Reported minimum and maximum values denote the lowest and highest growing-season emissions observed across the years covered in each study.

Region	Country	Mean	Min	Max	Reference
East Asia	China	–	0.3	127.4	Wang et al. (2018)
East Asia	China	19.9	1.0	97.0	Zhu and Li (2024)
East Asia	Japan	–	0.6	86.2	Wang et al. (2018)
East Asia	South Korea	–	9.7	214.5	Wang et al. (2018)
South Asia	India	–	0.8	347.5	Wang et al. (2018)
South Asia	India	–	3.0	15.0	Gupta et al. (2021)
South Asia	India	7.7	0.4	19.0	Bhatia (2013)
South Asia	Bangladesh	–	11.6	13.2	Wang et al. (2018)
Southeast Asia	Philippines	–	0.3	95.6	Wang et al. (2018)
Southeast Asia	Philippines	–	2.00	55.0	Wassmann et al. (2000)
Southeast Asia	Vietnam	–	3.1	35.3	Wang et al. (2018)
Southeast Asia	Indonesia	–	2.6	72.2	Wang et al. (2018)
South America	Brazil	–	4.6	51.7	Wang et al. (2018)
South America	Uruguay	–	9.33	24.9	Wang et al. (2018)
Europe	Spain	–	6.9	97.2	Wang et al. (2018)
Europe	Italy	–	0.8	81.9	Wang et al. (2018)

Globally, grasslands exhibit simulated CH₄ emissions that are half those from rice paddies, as indicated in Table 3. This important source of CH₄ emissions arises from two primary factors. First, the vast spatial extent of grasslands worldwide provides a substantial area for CH₄ production and emission, despite the generally lower CH₄ fluxes per unit area compared to rice paddies. Second, many grassland regions, particularly those converted from historical peatlands, are characterized by drained organic soils. The drainage of these former peatlands for agricultural use not only alters the hydrological regime but also influences the soil carbon dynamics, creating conditions conducive to CH₄ emissions (van den Pol-van Dasselaar, 1998). Residual croplands constitute a small but non-zero CH₄ source that is highly sensitive to field-scale hydrology. Even in nominally aerated systems, shallow or perched water tables, episodic waterlogging after irrigation or storms, and poorly drained depressions can create anoxic microsites where methanogenesis occurs. The resulting emissions are intermittent and heterogeneous, governed primarily by water-table depth and inundation frequency. Although flux magnitudes are typically far lower than those from wetlands, such emissions can be regionally relevant in poorly drained landscapes and in wet seasons, and they carry high spatial and interannual variability.

Table 7. Net biome production-Terrestrial land sink including land use change emissions estimated by Global Carbon Budget 2023 and simulated by LPJmL6

	GCB 2023		LPJmL6
	DGVM (net)	DGVM-LU-BK	
1990s	0.7 ±0.5	0.9 ±0.9	1.1
2000s	1.1 ±0.4	1.4 ±1.0	1.3
2010s	1.6 ±0.6	2.0 ±1.0	2.0

5.4 Carbon pools and fluxes

The implementation of the anoxic decomposition also influences the organic carbon pools and the CO₂ fluxes into the atmosphere. Therefore the LPJmL6 model needs to be benchmarked for the other important carbon cycle variables as well.

5.4.1 Net biome production

The release of soil carbon is decelerated due to the representation of wetlands in the LPJmL6 models. The anoxic decomposition is about 10 times slower and thus carbon is sequestered in soils for a much longer time. Thus the model can well reproduce the magnitude of the global carbon sink – including land-use change (LUC) emissions – within the estimate range of the Global Carbon Budget (GCB) 2023 (Friedlingstein et al., 2023) from Dynamic Global Vegetation Models (DGVM) and from the bookkeeping approach (DGVM-LU-BK) (see Table 7 and Fig. 6). Table 7 shows that LPJmL6 reproduces decadal terrestrial sink magnitudes within the GCB 2023 DGVM and DGVM-LU-BK uncertainty ranges and exhibits the expected strengthening from the 1990s to the 2010s; across the two GCB methods, residual differences can be traced to land-use-change treatment—bookkeeping versus implicit DGVM dynamics—and no systematic decadal bias is evident; this also holds for LPJmL6, indicating robust, method-invariant trends in the net land carbon sink. In Fig. 6, the simulated terrestrial sink follows the independently constrained apparent sink and the atmospheric CO₂ growth rate with comparable interannual variability, indicating that LPJmL6 captures the major climate-driven fluctuations and the multi-decadal increase in sink strength without long-term divergence from independent estimates.

5.4.2 Net ecosystem exchange

We used eddy-covariance flux tower data to benchmark simulated NEE (Fig. 7; full data description in section 4.2). As noted there, comparisons are constrained by scale mismatches between grid cells and tower footprints, possible




Figure 6. Terrestrial sink estimated by LPJmL6 and Global Carbon Budget 2023 (Friedlingstein et al., 2023).

vegetation (PFT) mismatches, and the use of large-scale meteorological forcing rather than site-specific observations, which limits the ability to capture site-specific variability in carbon fluxes.

Nevertheless, the comparison demonstrates that correlations between simulated and observed fluxes are generally high across many sites, indicating that the model captures the overall temporal dynamics of NEE reasonably well. However, the centered root mean square difference (CRMSD) and the normalized standard deviation reveal larger discrepancies, suggesting that while the timing of seasonal and interannual variability is reproduced, the magnitude of fluxes is often either over- or underestimated.

5.4.3 GPP

The ILAMB comparison demonstrates that the model successfully reproduces gross primary production (GPP) within the same order of magnitude as reported by Jung et al. (2011) (global mean $119 \pm 6 \text{ PgC a}^{-1}$ for 1982–2008 vs. 126 PgC a^{-1} simulated by LPJmL6). This agreement is also in line with later FLUXCOM syntheses that place global GPP roughly in the $106\text{--}130 \text{ PgC a}^{-1}$ range, depending on predictors and machine-learning methods used (Jung et al., 2020).




Figure 7. Taylor diagram showing the comparison of simulated net ecosystem exchange (NEE) against eddy-covariance flux tower measurements. Observational data are derived from the FLUXNET network, which provides standardized eddy-covariance measurements of ecosystem-atmosphere carbon and water fluxes (Pastorello et al., 2020)

895 5.4.4 Aboveground biomass (AGB)

Across the ILAMB benchmark, reference biomass products differ in scope (forest AGB vs. all vegetation; above- vs. below-ground components), spatial/temporal coverage, and retrieval/inversion methodology, yielding a wide spread in implied global stocks. LPJmL6 estimates global woody biomass to be 422 PgC (mean 2000–2019). Among reference datasets, products that resolve high biomass in humid tropical and high-latitude forests report totals comparable

900 to LPJmL6, for example, GEOCARBON ~ 408 PgC (GEO, 2014); the Xu–Saatchi synthesis 381 ± 2 PgC for 2000–2019 (Xu et al., 2021). By contrast, ESA CCI Biomass provides forest AGB of ~ 300 PgC (above-ground forest only) (GEO, 2014; Santoro et al., 2021), and NBCD2000 is regional (conterminous U.S., ~ 2000 baseline), serving primarily for spatial pattern evaluation rather than global sums (Kelldorfer et al., 2012). These definitional and methodological contrasts account for the spread in ILAMB intercomparisons and explain why LPJmL6 aligns more
 905 closely with datasets emphasizing woody stocks across humid tropical and boreal forests, whereas ESA CCI implies lower global totals.

From the reference data sets used in ILAMB, only the Xu–Saatchi (Xu et al., 2021) product (381 PgC) aligns closely with the LPJmL6 simulation result of 422 PgC, whereas most other datasets imply substantially higher global biomass totals ranging from 300 to 408 PgC.

910 5.5 Water fluxes

We assess the large-scale terrestrial water cycle to ensure that the CH_4 implementation operates within a physically consistent land-surface framework. The analysis combines global budgets with spatial diagnostics (renewable water resources and groundwater recharge) and skill metrics for multi-decadal river discharge and evapotranspiration (ET) simulations.

915 5.5.1 Global water fluxes

Table 8 synthesizes ranges of data-driven and model-based estimates for the principal terrestrial water fluxes – evapotranspiration (ET), transpiration (T), total runoff from land, groundwater recharge, and irrigation withdrawals. ET ($62 \times 10^3 \text{ km}^3 \text{ a}^{-1}$) and T ($39 \times 10^3 \text{ km}^3 \text{ a}^{-1}$) fall within published model ranges and are near the lower bounds of data-driven syntheses, indicating a conservative global partitioning of ET. Total runoff and river discharge (47
 920 and $40 \times 10^3 \text{ km}^3 \text{ a}^{-1}$) lie near the centers of reported ranges, consistent with independent constraints. Groundwater recharge ($31 \times 10^3 \text{ km}^3 \text{ a}^{-1}$) exceeds data-driven and modeled estimates; this difference reflects varying definitions (diffuse vs. focused recharge and return flows), considered land areas, and averaging periods, also examined spatially in section 5.5.2, with discharge and ET evaluations provided in sections 5.5.3 and 5.5.4. Nevertheless, the LPJmL6 estimate is near the upper end of published values and coincides with greater groundwater storage in the simulations
 925 than reported Müller Schmied et al. (2021) (Fig. 8).

5.5.2 Groundwater recharge and renewable storage

Fig. 8 compares long-term mean groundwater storage (GWS) and groundwater recharge for 1981–2010 from WaterGAP 2.2 (Müller Schmied et al., 2021) and LPJmL6. Recharge is highest in humid tropics and monsoon regions (Amazon, Congo, SE Asia, eastern India), and is moderate in temperate wet zones, and near zero across arid belts
 930 (Sahara, Arabian Peninsula, central Australia); orographic bands (Andes, Himalaya) are evident. Storage broadly

Table 8. Simulated global terrestrial water fluxes, and ranges of data-driven and modeled estimates (in $10^3 \text{ km}^3 \text{ a}^{-1}$). Evapotranspiration and transpiration represent long-term land averages (1982–2011); runoff and discharge are multi-decadal climatologies (1982–2011 land means); infiltration denotes diffuse groundwater recharge; irrigation withdrawals (around 2010) include both surface and groundwater abstraction.

Flux	Data-driven ($10^3 \text{ km}^3 \text{ a}^{-1}$)	Modelled ($10^3 \text{ km}^3 \text{ a}^{-1}$)	LPJmL6 ($10^3 \text{ km}^3 \text{ a}^{-1}$)
Evapotranspiration (ET) ^{1,2}	64–69	50–70	62
Transpiration (T) ^{3,4}	55 ± 12	39–52	39
Total runoff (land) ^{5,6,7}	41–46	40–50	47
Groundwater recharge ^{9,10}	12–13	~11–30	31*
Irrigation withdrawals (blue water) ^{11–14}	2.4–2.9	2.3–3.3	2.2

References: ¹ Pan et al. (2020); ² Wild et al. (2015); ³ Jasechko et al. (2013); ⁴ Good et al. (2015); ⁵ Schneider et al. (2017); ⁶ Godoy et al. (2021); ⁷ Döll et al. (2016); ⁸ Mohan et al. (2018); ⁹ Döll and Fiedler (2008); ¹⁰ FAO (2020a); ¹¹ Wada and Bierkens (2014a); ¹² Wada and Bierkens (2014b); ¹³ Huang et al. (2018); ¹⁴ FAO (2014);

*including return flows from irrigation

tracks persistently wet lowlands and large sedimentary basins, with minima in deserts and high-latitude regions. LPJmL6 tends to display sharper spatial gradients and stronger orographic signals, while WaterGap is smoother and often more conservative in semi-arid interiors.

5.5.3 Discharge for large river basins

935 The comparison of discharge data shows that LPJmL6 is able to reproduce the seasonal hydrographs (see Fig. 9 and in the Supplement Fig. S28, especially in the southern latitudes. There are small improvements compared to LPJmL5.10. The far northern discharge levels are also reproduced quite well, with the largest discrepancy in the mid-latitudes. In some regions with low R^2 , errors are dominated by timing mismatch: the model typically leads the observed cycle (peaks occur earlier), while amplitudes are reasonably captured (see the Supplement Fig. S28).

940 5.5.4 Evapotranspiration fluxes

Simulated ET was compared against eddy-covariance flux tower measurements. While towers provide high-frequency and highly localized ET estimates, the coarse spatial resolution of the simulations introduces a scale mismatch similar to that encountered for NEE.

Model-data agreement of ET is even higher than for NEE, in terms of both correlation coefficients (the model reproduces the temporal variability of ET with greater accuracy) and the CRMSD and the normalized standard deviation (the magnitude of ET fluxes is represented more consistently across sites). This likely reflects the fact that ET is primarily controlled by atmospheric demand and canopy conductance, processes that are more directly




Figure 8. Comparison of WaterGAP 2.2 (Müller Schmied et al., 2021) and LPJmL6 maps of (a,b) renewable groundwater storage (GWS; multi-year mean, 1981–2010) and (c,d) groundwater recharge (mean annual, 1981–2010). Colors show discrete classes for storage in mm; for recharge the units are mm a⁻¹.

constrained by meteorological forcing and vegetation structure in the model, whereas NEE additionally depends on complex carbon turnover processes and ecosystem-specific dynamics.

950 5.6 Vegetation distribution

A comparison of LPJmL6-simulated actual land cover (natural vegetation plus managed cropland and grassland) with ESA CCI LC4 land-cover product (Harper et al., 2023) reveals a generally good match in spatial patterns for PFTs (Fig. 11). However, in the temperate and southern boreal zone, ESA CCI LC4 data predominantly show broadleaved summergreen trees, while LPJmL6 simulates a substantially lower abundance of this PFT. Part of this




Figure 9. Comparison of simulated discharge with 286 gauges provided by ArcticNET and UNH/GRDC. Stations with basin area $> 10000 \text{ km}^2$ are taken into account. Gauges are ordered from north to south by color (see legend).




Figure 10. Taylor diagram showing the comparison of simulated evapotranspiration fluxes against eddy-covariance flux tower measurements. Observational data are derived from the FLUXNET network, which provides standardized eddy-covariance measurements of ecosystem-atmosphere carbon and water fluxes (Pastorello et al., 2020)

divergence likely stems from the LC-to-PFT reclassification applied to ESA CCI LC4, which can inflate broadleaved summergreen fractions in mosaic and open-forest landscapes and has been shown to introduce spatially structured errors, including overestimation near the taiga-tundra ecotone (Wang et al., 2023b). In the inner tropics, LPJmL6 dominantly simulates broadleaved evergreen trees in line with the observed data, yet it also produces a minor fraction of broadleaved deciduous cover, consistent with the common deciduous-evergreen coexistence in tropical forests (Muehleisen et al., 2024), reflecting the influence of rainfall variability. Mora et al. (2014) suggest that remote




Figure 11. Vegetation distribution of LPJmL6 compared to ESACCI-LC-L4 data (Harper et al., 2023)

sensing data, such as the ESA CCI LC4 dataset, may overestimate the extent of broadleaved evergreen trees in these regions due to challenges in distinguishing between evergreen and deciduous species under dense canopies and during certain seasons. Factors contributing to this misclassification include mixed pixel composition, seasonal variations in leaf phenology, and limitations in sensor resolution. For instance, the complexity of tropical forest structures and the similarity in spectral signatures between different tree species can lead to inaccuracies in classification (Zhong et al., 2024). Finally, while the model effectively simulates the presence of needleleaved deciduous trees in appropriate areas, aligning well with ESA observations, LPJmL6 tends to over-represent their overall abundance in the high-latitude boreal forests of Eurasia and North America.

Uncertainties also arise from the representation of land use, both in observational products and in the prescribed
970 land-use forcing used to drive LPJmL6. Reconstructions and scenarios of cropland and pasture area differ substan-
tially among datasets, particularly in regions with smallholder or mosaic agriculture, leading to divergent estimates
of the area available for natural and managed PFTs (e.g. Prestele et al., 2016; Hurtt et al., 2020; Klein Goldewijk
et al., 2017b). These discrepancies propagate into both the LC-to-PFT reclassification of ESA CCI LC4 and the
land-use trajectories imposed on LPJmL6, so part of the mismatch between simulated and observed PFT patterns
975 likely reflects uncertainties in land-use data rather than Vegetation dynamics alone.

6 Conclusions

We present LPJmL version 6, including a newly developed process-consistent CH_4 module that links hydrological,
vegetation dynamical and biogeochemical processes to atmospheric CH_4 exchange. A new wetland scheme combines
a dynamic water table with a CTI-based inundation fraction that is recalculated from the grid-cell CTI distribution,
980 enabling the simulation of spatially and temporally varying anoxic areas within each cell.

Evaluation demonstrates (i) agreement in global wetland area with several inventories, while highlighting a per-
sistent, first-order observational divergence (3.6–11.4 million km^2) that translates into different emission potentials;
LPJmL6 captures the tropical signal and part of the mid-latitude peak but remains conservative in boreal zones
relative to GLWD. (ii) global CH_4 emissions that are consistent with atmospheric measurement constraints when
985 combined with non-modeled sectors that bring the modeled net flux into the 506–630 $\text{Tg CH}_4 \text{ a}^{-1}$ corridor for
2008–2017. (iii) mechanistic attribution showing diffusion as a leading pathway and ebullition as small, with fires
contributing 5% of natural sources.

LPJmL6 successfully simulates the interactions between environmental factors such as temperature, soil moisture,
and vegetation, which govern CH_4 production, oxidation, and transport. Its ability to capture regional and temporal
990 emission trends aligns well with observational datasets and other global estimates, particularly in high-emission re-
gions like tropical wetlands and rice cultivation areas. At the same time, the model highlights remaining uncertainties
in CH_4 emissions from agricultural land and high-latitude regions, emphasizing the need for improved constraints
on wetland dynamics, managed water fluxes, and small-scale CH_4 sources. Uncertainties are dominated by wetland
extent in high latitudes and by agricultural sources, including rice-area dynamics and management, which LPJmL6
995 now resolves explicitly but for which observational constraints remain limited.

The wetland evaluation should also be interpreted in light of the prescribed CTI representation used by the
TOPMODEL formulation. In LPJmL6, CTI does not directly prescribe wetland extent, but provides a fixed sub-
grid topographic template that translates the simulated grid-cell mean water-table depth into a saturated fraction.
Regional biases in simulated wetland extent may therefore arise either from errors in the simulated water balance and
1000 water-table depth, or from limitations of the prescribed CTI distribution and its assumed gamma-shaped sub-grid
variability. These limitations are particularly relevant in regions where wetland occurrence is not controlled by local

topographic convergence alone. In very flat lowlands, small uncertainties in elevation or slope can strongly affect the inferred CTI distribution, whereas in steep terrain narrow riparian or valley-bottom wetlands may remain unresolved at the model resolution. In floodplains, coastal regions, and managed systems, wetland extent may additionally
1005 depend on river inundation, groundwater convergence, backwater effects, drainage, or water management, which are only partly represented by a static CTI-based approach. Consequently, agreement with observed large-scale latitudinal wetland patterns does not imply that the spatial wetland distribution is equally reliable in all regions.

A related uncertainty concerns the agricultural water-input threshold used to represent enhanced drainage on managed land. This threshold is a pragmatic model assumption rather than a parameter derived from site-specific
1010 drainage data. It is interpreted as an effective upper limit intended to account for enhanced drainage and water management on agricultural land. Future applications could refine this representation using spatially explicit information on drainage infrastructure, irrigation, and soil hydraulic properties.

Despite these uncertainties, LPJmL6 aligns with bottom-up wetland emission estimates across decades, while recent top-down estimates have converged downward, reducing the historical bottom-up–top-down gap and supporting
1015 the process-based approach implemented here. A systematic sensitivity analysis of the newly introduced and modified parameters is an important next step. Such an analysis would help quantify the robustness of the simulated wetland extent, agricultural CH₄ emissions, and total land–atmosphere CH₄ exchange to individual parameter choices and their interactions. Future model development should therefore focus on identifying the most influential parameters and constraining them with independent hydrological, biogeochemical, land-management, and CH₄ flux observations
1020 where available.

The implementation of CH₄ dynamics in LPJmL6 represents a significant advancement in the modeling of terrestrial land surface dynamics and, in particular, CH₄ emissions. By integrating CH₄ processes within the broader framework of the global carbon, nitrogen, and water cycles, the model allows for detailed assessments of emissions from different land-use groups, here reported separately for natural wetlands, rice paddies, grasslands, and the remaining cropland. The dynamic representation of wetlands and permafrost processes, combined with hydrological
1025 modeling, enhances the understanding of CH₄ flux variability across diverse ecosystems.

CH₄ emissions from land use and wetlands are a significant contributor to climate change. Forested wetlands, wetland dynamics, permafrost, and various anthropogenic and natural sources contribute all to the global CH₄ budget. By understanding the complex dynamics of CH₄ emissions and their environmental regulation, effective
1030 strategies for mitigating climate change can be developed. Continued research and monitoring of CH₄ emissions are essential for improving our understanding of this potent greenhouse gas and its impact on the planet.

LPJmL6 is a robust tool for understanding the complexities of CH₄ dynamics under current and future climate conditions. These advances allow for more accurate projections of CH₄ emissions, critical for formulating effective climate-mitigation policies. Continued refinement of CH₄-related processes, such as improved parameterization of
1035 wetland extents and anthropogenic emission factors, will further increase the realism of LPJmL6-derived surface fluxes. When these fluxes are used in, or coupled to, atmospheric transport and chemistry modules within Earth

system modelling frameworks, they can help to better constrain the global CH₄ budget and attribution of observed concentration trends, thereby supporting more robust climate-impact assessments. In addition to enhancing CH₄ representation, the improvements made to the soil water status significantly advance the model’s ability to capture land surface dynamics. This enhancement is particularly impactful for simulating water stress in natural vegetation and crops, leading to more accurate representation of plant-water interactions. Notably, this advancement represents a co-benefit of the CH₄-focused implementation, offering improvements beyond CH₄ simulations by strengthening the model’s capacity to simulate hydrological processes and their influence on terrestrial ecosystems.

Code and data availability. The LPJmL6 model version further developed and used in this study is archived on Zenodo: <https://zenodo.org/records/17911633> (Schaphoff et al., 2025b) and is also available from GitHub (<https://github.com/PIK-LPJmL/LPJmL>). The evaluation and analysis scripts, including all ILAMB validation scripts and configuration files required to reproduce the diagnostics, are likewise archived on Zenodo: <https://doi.org/10.5281/zenodo.17877397> (Schaphoff et al., 2025a). The LPJmL6 model output used for the evaluation in this paper is provided in NetCDF format via Zenodo: <https://doi.org/10.5281/zenodo.17877397> (Schaphoff et al., 2025a). Together, these repositories contain all model code, scripts, and simulation data necessary to reproduce the results presented here, apart from the external observational and benchmark data sets, which are available from the original providers cited in the main text.

All observational, remote-sensing, and synthesis products used for model evaluation are third-party datasets and are not redistributed with this manuscript; users must obtain them directly from the original providers under their respective licenses/usage policies. Wetland extent evaluation used: the Global Lakes and Wetlands Database (GLWD; <https://www.worldwildlife.org/pages/global-lakes-and-wetlands-database>, accessed 2018-08-15; the Kaplan wetland map (<https://doi.org/10.5281/zenodo.18274983>); the SWAMP “Tropical and Subtropical Wetlands Distribution” dataset (<https://doi.org/10.17528/CIFOR/DATA.00058>, accessed 2024-11-24); and WAD2M wetland dynamics (<https://zenodo.org/records/5553187>, accessed 2024-09-03). Latitudinal CH₄ emission benchmarking used WetCHARTs v1.0 (<https://doi.org/10.3334/ORNLDAAAC/1502>, accessed 2017-06-01). Atmospheric constraints used CarbonTracker-CH₄ (<https://gml.noaa.gov/ccgg/carbontracker-ch4/>, accessed 2024-02-28) and NOAA CH₄ trend products (https://gml.noaa.gov/ccgg/trends_ch4/, accessed 2024-02-28). Eddy-covariance NEE and ET were taken from FLUXNET2015 (<https://fluxnet.org/data/fluxnet2015-dataset/>, accessed 2017-04-17). Vegetation structure evaluation used ESA CCI Land Cover “Global Plant Functional Types (PFT) Dataset” (<http://maps.elie.ucl.ac.be/CCI/viewer/download.php>, accessed 2022-08-19). System-level synthesis constraints used the Global Carbon Budget data (<https://doi.org/10.18160/GCP-2023>, accessed 2024-03-15) and the Global Methane Budget data portal (<https://www.globalcarbonproject.org/methanebudget/> and data page <https://www.globalcarbonproject.org/methanebudget/24/data.htm>, accessed 2024-03-18). R-ArcticNET v4.0 station attributes and monthly discharge tables are available from the project download at <https://www.r-arcticnet.sr.unh.edu/v4.0/AllData/index.html>, accessed 2012-05-15 (R-ArcticNET Project). RivDIS v1.1 can be accessed via <https://doi.org/10.3334/ORNLDAAAC/199>, accessed 2015-03-14 (Vörösmarty et al., 1998).

Multi-metric benchmarking used ILAMB software and its reference dataset collection (<https://www.ilamb.org/>; datasets: <https://www.ilamb.org/datasets.html>).




Figure A1. Base-flow recession coefficient determining the outflow rate from groundwater stores as a fraction of the store (ERMITAGE project report (2013),).

Author contributions. SiS conceived and developed the model, performed the analyses, and led the writing of the manuscript. DH and WvB co-developed the model and provided expert guidance on the mathematical methods. All co-authors offered substantial feedback on successive drafts and contributed to the design and interpretation of the process evaluation.

Competing interests. The authors declare that they have no conflict of interest. CM is member of the editorial board of GMD.

1075 Acknowledgements. SO was supported by the Conservation International Foundation, grant number CI-114129 and from the Future of Life Institute through the CODEC project.

References

- State of Food and Agriculture. Biofuels: prospects, risks and opportunities, FAO, <http://www.fao.org/docrep/011/i0100e/i0100e00.htm>.
- 1080 GEOCARBON Global Forest Aboveground Biomass (Mg/ha), UNEP/GRID Geneva GeoNetwork, <https://datacore-gn.unepgrid.ch/geonetwork/srv/api/records/5e695176-266d-4697-bc06-c2d9196845b4>, global forest AGB map produced in the EU FP7 GEOCARBON project by combining Avitabile et al. (2016; pan-tropics) and Santoro et al. (2015; northern hemisphere) products, 2014.
- Alexiades, V. and Solomon, A.: Mathematical Modeling Of Melting And Freezing Processes, Hemisphere, 1993.
- 1085 Bartlett, K. B. and Harriss, R. C.: Review and assessment of methane emissions from wetlands, *Chemosphere*, 26, 261–320, [https://doi.org/10.1016/0045-6535\(93\)90427-7](https://doi.org/10.1016/0045-6535(93)90427-7), 1992.
- Beck, H. E., van Dijk, A. I. J. M., Miralles, D. G., de Jeu, R. A. M., (Sampurno) Bruijnzeel, L. A., McVicar, T. R., and Schellekens, J.: Global patterns in base flow index and recession based on streamflow observations from 3394 catchments, *Water Resources Research*, 49, 7843–7863, <https://doi.org/10.1002/2013WR013918>, 2013.
- 1090 Beven, K. and Kirkby, M.: A physically-based variable contributing area model of basin hydrology., *Hydrological Sciences Bulletin*, 24(1), 43–69, 1979.
- Bhatia, A. e. a.: Methane and nitrous oxide emissions from Indian rice paddies, and their global warming potential, *Current Science*, 105, 661–666, 2013.
- Biemans, H., Haddeland, I., Kabat, P., Ludwig, F., Hutjes, R. W. a., Heinke, J., von Bloh, W., and Gerten, D.: Impact of
1095 reservoirs on river discharge and irrigation water supply during the 20th century, *Water Resources Research*, 47, W03509, <https://doi.org/10.1029/2009WR008929>, 2011.
- Bieniada, A., Green, S. M., Peacock, M., et al.: Steady and ebullitive methane fluxes from active, restored and unrestored fens in a hemiboreal region, *Ecological Engineering*, 164, 106209, <https://doi.org/10.1016/j.ecoleng.2021.106209>, quantified CH₄ ebullition vs diffusion; vascular cover (reed, sedge) strongly affects ebullition fraction., 2021.
- 1100 Bloom, A., Bowman, K., Lee, M., Turner, A., Schroeder, R., Worden, J., Weidner, R., McDonald, K., and Jacob, D.: CMS: Global 0.5-deg Wetland Methane Emissions and Uncertainty (WetCHARTs v1.0), <https://doi.org/10.3334/ORNLDAAAC/1502>, 2017a.
- Bloom, A. A., Bowman, K. W., Lee, M., Turner, A. J., Schroeder, R., Worden, J. R., Weidner, R., McDonald, K. C., and Jacob, D. J.: A global wetland methane emissions and uncertainty dataset for atmospheric chemical transport models
1105 (WetCHARTs version 1.0), *Geoscientific Model Development*, 10, 2141–2156, <https://doi.org/10.5194/gmd-10-2141-2017>, 2017b.
- Bruhwyler, L. M. P., Basu, S., Bergamaschi, P., Bousquet, P., Dlugokencky, E., Houweling, S., Janardanan, R., Miller, S., Pandey, S., Prigent, C., et al.: CarbonTracker-CH₄: An assimilation system for estimating emissions of atmospheric methane, *Atmospheric Chemistry and Physics*, 14, 8269–8293, <https://doi.org/10.5194/acp-14-8269-2014>, 2014.
- 1110 Christen, E. W., Chung, S.-O., and Quayle, W.: Simulating the fate of molinate in rice paddies using the RICEWQ model, *Agricultural Water Management*, 85, 38–46, <https://doi.org/10.1016/j.agwat.2006.03.008>, 2006.
- Clapp, R. B. and Hornberger, G. M.: Empirical equations for some soil hydraulic properties, *Water Resour. Res.*, 14, 601–604, <https://doi.org/10.1029/wr014i004p00601>, 1978.

- Collier, N., Hoffman, F. M., Lawrence, D. M., Keppel-Aleks, G., Koven, C. D., Riley, W. J., Randerson, J. T., Swenson, S. C., Thornton, P. E., Wieder, W. R., Xu, C., Yang, X., and Zeng, X.: The International Land Model Benchmarking (ILAMB) System: Design, Theory, and Implementation, *Journal of Advances in Modeling Earth Systems*, 10, 2731–2750, <https://doi.org/10.1029/2018MS001354>, 2018.
- Colmer, T. D.: Long-distance transport of gases in plants: a perspective on internal aeration and radial oxygen loss from roots, *Plant, Cell & Environment*, 26, 17–36, <https://doi.org/10.1046/j.1365-3040.2003.00846.x>, 2003.
- Conrad, R.: Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O, and NO), *Microbiological Reviews*, 60, 609–640, <https://doi.org/10.1128/mr.60.4.609-640.1996>, 1996.
- Cucchi, M., Weedon, G. P., Amici, A., Bellouin, N., Lange, S., Müller Schmied, H., Hersbach, H., and Buontempo, C.: WFDE5: bias-adjusted ERA5 reanalysis data for impact studies, *Earth System Science Data*, 12, 2097–2120, <https://doi.org/10.5194/essd-12-2097-2020>, 2020.
- D. Gerten, S. Schaphoff, M. Bonsch, A. Popp, P. Holden, C. Wallace, K. Hiscock, T. Osborn; ed. N. Edwards, B. Pizzileo: Enhancing Robustness and Model Integration for The Assessment of Global Environmental Change, <https://arquivo.pt/wayback/20160421111120/http://ermitage.cs.man.ac.uk/?q=node/52>, grant Agreement No. 265170; archived at Arquivo.pt.
- Duval, B. and Goodwin, S.: Methane production and release from two New England peatlands, *International Microbiology*, 3, 89–96, 2000.
- Döll, P. and Fiedler, K.: Global-scale modeling of groundwater recharge, *Hydrology and Earth System Sciences*, 12, 863–885, <https://doi.org/10.5194/hess-12-863-2008>, 2008.
- Döll, P., Douville, H., Güntner, A., Schmied, H. M., and Wada, Y.: Modelling freshwater resources at the global scale: Challenges in integrating hydrological components, *Surveys in Geophysics*, 37, 195–221, <https://doi.org/10.1007/s10712-015-9343-1>, 2016.
- EPA: Global Anthropogenic Non-CO₂ Greenhouse Gas Emissions: 1990–2030, Report EPA 430-R-12-006, U.S. Environmental Protection Agency (EPA), Washington, D.C., <https://www.epa.gov/global-mitigation-non-co2-greenhouse-gases/global-non-co2-ghg-emissions-1990-2030>, accessed 2025-12-02, 2012.
- FAO: AQUASTAT database - Food and Agriculture Organization of the United Nations (FAO), <http://www.fao.org/nr/water/aquastat/data/query/index.html?lang=en>, 2014.
- FAO: AQUASTAT database, Food and Agriculture Organization of the United Nations, Rome, <http://www.fao.org/nr/water/aquastat/data/query/index.html?lang=en>, accessed 2020-04-17, 2020a.
- FAO: FAOSTAT database, Food and Agriculture Organization of the United Nations, Rome, <http://www.fao.org/faostat/en/>, accessed 2020-01-27, 2020b.
- FAO: FAOSTAT Emissions Data, <https://www.fao.org/faostat/en/#data/GT>, 2020c.
- FAO/IIASA/ISRIC/ISSCAS/JRC: Harmonized World Soil Database (version 1.21), FAO, Rome, Italy and IIASA, Laxenburg, Austria, <https://iiasa.ac.at/web/home/research/researchPrograms/water/HWSD.html>, accessed 2021-03-11, 2012.
- Flörke, M., Kynast, E., Bärlund, I., Eisner, S., Wimmer, F., and Alcamo, J.: Domestic and industrial water uses of the past 60 years as a mirror of socio-economic development: A global simulation study, *Global Environmental Change*, 23, 144–156, <https://doi.org/10.1016/j.gloenvcha.2012.10.018>, 2013.

- Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, O., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pocock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth System Science Data*, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, 2023.
- Godoy, A. R., Castanheira, B. G., Paiva, R. C. D., and Collischonn, W.: Global river discharge reanalysis based on multiple observation datasets, *Water Resources Research*, 57, e2020WR028427, <https://doi.org/10.1029/2020WR028427>, 2021.
- Good, S. P., Noone, D., and Bowen, G.: Hydrologic connectivity constrains partitioning of global terrestrial water fluxes, *Science*, 349, 175–177, <https://doi.org/10.1126/science.aaa5931>, 2015.
- Gumbrecht, T., Román-Cuesta, R., Verchot, L., Herold, M., Wittmann, F., Householder, E., Herold, N., and Murdiyarso, D.: Tropical and Subtropical Wetland Distribution, <https://doi.org/10.17528/CIFOR/DATA.00058>, sWAMP, 2017.
- Gupta, K., Kumar, R., Baruah, K. K., Hazarika, S., Karmakar, S., and Bordoloi, N.: Greenhouse gas emission from rice fields: a review from Indian context, *Environmental Science and Pollution Research*, 28, 30 551–30 572, <https://doi.org/10.1007/s11356-021-13935-1>, 2021.
- Haggard, B., Moore Jr, P., and Brye, K.: Effect of slope on runoff from a small variable-slope box, *Journal of Environmental hydrology*, 13, 25–45, 2005.
- Harper, K. L., Lamarche, C., Hartley, A., Peylin, P., Ottlé, C., Bastrikov, V., San Martín, R., Bohnenstengel, S. I., Kirches, G., Boettcher, M., Shevchuk, R., Brockmann, C., and Defourny, P.: A 29-year time series of annual 300 m resolution plant-functional-type maps for climate models, *Earth System Science Data*, 15, 1465–1499, <https://doi.org/10.5194/essd-15-1465-2023>, 2023.
- He, J., Naik, V., and Horowitz, L. W.: Hydroxyl Radical (OH) Response to Meteorological Forcing and Implication for the Methane Budget, *Geophysical Research Letters*, 48, e2021GL094140, <https://doi.org/10.1029/2021GL094140>, e2021GL094140 2021GL094140, 2021.
- Heinke, J.: A livestock density input for LPJmL5.5, <https://doi.org/10.5281/zenodo.14946695>, 2025.
- Heinke, J., Rolinski, S., and Müller, C.: Modelling the role of livestock grazing in C and N cycling in grasslands with LPJmL5.0-grazing, *Geoscientific Model Development*, 16, 2455–2475, <https://doi.org/10.5194/gmd-16-2455-2023>, 2023.

- Hu, Q., Li, J., Xie, H., Huang, Y., Canadell, J. G., Yuan, W., Wang, J., Zhang, W., Yu, L., Li, S., Lu, X., Li, T., and Qin, Z.: Global methane emissions from rice paddies: CH4MOD model development and application, *iScience*, 27, 111 237, <https://doi.org/https://doi.org/10.1016/j.isci.2024.111237>, 2024.
- 1190 Huang, Z., Hejazi, M., Li, X., Tang, Q., Vernon, C., Leng, G., Liu, Y., Döll, P., Eisner, S., Gerten, D., Hanasaki, N., and Wada, Y.: Reconstruction of global gridded monthly sectoral water withdrawals for 1971–2010 and analysis of their spatiotemporal patterns, *Hydrology and Earth System Sciences*, 22, 2117–2133, <https://doi.org/10.5194/hess-22-2117-2018>, 2018.
- Hugelius, G., Loisel, J., Chadburn, S., Jackson, R. B., Jones, M., MacDonald, G. M., Marushchak, M., Olefeldt, D., Packalen, M., Siewert, M. B., Treat, C., and Turetsky, M.: Large stocks of peatland carbon and nitrogen are vulnerable to permafrost thaw, *Proceedings of the National Academy of Sciences*, 117, 20 438–20 446, <https://doi.org/10.1073/pnas.1916387117>, 2020.
- 1195 Hurtt, G. C., Chini, L., Sahajpal, R., Frolking, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori, S., Klein Goldewijk, K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J. O., Kennedy, J., Krisztin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X.: Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6, *Geosci. Model Dev.*, 13, 5425–5464, <https://doi.org/10.5194/gmd-13-5425-2020>, data available at <https://luh.umd.edu/>, accessed 2021-07-08, 2020.
- 1200 IIASA/FAO: Global Agro-ecological Zones (GAEZv3.0), IIASA, Laxenburg, Austria and FAO, Rome, Italy, <http://www.gaez.iiasa.ac.at/>, accessed 2020-07-17, 2012.
- Jasechko, S., Sharp, Z. D., Gibson, J. J., Birks, S. J., Yi, Y., and Fawcett, P. J.: Terrestrial water fluxes dominated by transpiration, *Nature*, 496, 347–350, <https://doi.org/10.1038/nature11983>, 2013.
- 1205 Jobbagy, E. G. and Jackson, R. B.: The Vertical Distribution of Soil Organic Carbon and Its Relation to Climate and Vegetation, *Ecological Applications*, 10, 423–436, <http://www.jstor.org/stable/2641104>, articleType: research-article / Full publication date: Apr., 2000 / Copyright © 2000 Ecological Society of America, 2000.
- Jung, M., Reichstein, M., Margolis, H. A., Cescatti, A., Richardson, A. D., Arain, M. A., Arneth, A., Bernhofer, C., Bonal, D., Chen, J., Gianelle, D., Gobron, N., Kiely, G., Kutsch, W., Lasslop, G., Law, B. E., Lindroth, A., Merbold, L., Montagnani, L., Moors, E. J., Papale, D., Sottocornola, M., Vaccari, F., and Williams, C.: Global patterns of land-atmosphere fluxes of carbon dioxide, latent heat, and sensible heat derived from eddy covariance, satellite, and meteorological observations, *Journal of Geophysical Research: Biogeosciences*, 116, n/a–n/a, <https://doi.org/10.1029/2010JG001566>, 2011.
- 1210 Jung, M., Schwalm, C., Migliavacca, M., Walther, S., Camps-Valls, G., Koirala, S., Anthoni, P., Besnard, S., Bodesheim, P., Carvalhais, N., Chevallier, F., Gans, F., Goll, D. S., Haverd, V., Köhler, P., Ichii, K., Jain, A. K., Liu, J., Lombardozzi, D., Nabel, J. E. M. S., Nelson, J. A., O’Sullivan, M., Pallandt, M., Papale, D., Peters, W., Pongratz, J., Rödenbeck, C., Sitch, S., Tramontana, G., Walker, A., Weber, U., and Reichstein, M.: Scaling carbon fluxes from eddy covariance sites to globe: synthesis and evaluation of the FLUXCOM approach, *Biogeosciences*, 17, 1343–1365, <https://doi.org/10.5194/bg-17-1343-2020>, 2020.
- 1215 Jägermeyr, J., Müller, C., Minoli, S., Ray, D., and Siebert, S.: GGCMI Phase 3 crop calendar, <https://doi.org/10.5281/zenodo.5062513>, 2021.
- 1220 Kaplan, J.: A composite map of global wetland distribution at 0.5 degree resolution, <https://doi.org/10.5281/zenodo.18274983>, 2007.

- Kaur, G., Singh, G., Motavalli, P. P., Nelson, K. A., Orlowski, J. M., and Golden, B. R.: Impacts and management strategies for crop production in waterlogged or flooded soils: A review, *Agronomy Journal*, 112, 1475–1501, 2020.
- Kellndorfer, J., Walker, W., Kirsch, K., Fiske, G., Bishop, J., LaPoint, E., Hoppus, M., and Westfall, J.: NACP Aboveground Biomass and Carbon Baseline Data (NBCD 2000), U.S.A., 2000, <https://doi.org/10.3334/ORNLDAAAC/1081>, 2012.
- Khvorostyanov, D. V., Krinner, G., Ciais, P., M. Heimann, and Zimov, S. A.: Vulnerability of permafrost carbon to global warming. Part I: model description and role of heat generated by organic matter decomposition, *Tellus B*, 60, 250–264, <http://dx.doi.org/10.1111/j.1600-0889.2007.00333.x>, 2008.
- Kim, H.: Global Soil Wetness Project Phase 3 Atmospheric Boundary Conditions (Experiment 1) [Data set], <https://doi.org/10.20783/DIAS.501>, 2017.
- Kirschke, S., Bousquet, P., Ciais, P., Saunio, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron-Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J.-F., Langenfelds, R. L., Le Quere, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nature Geosci*, 6, 813–823, <http://dx.doi.org/10.1038/ngeo1955>, 2013.
- Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the Holocene - HYDE 3.2, *Earth Syst. Sci. Data*, 9, 927–953, <https://doi.org/10.5194/essd-9-927-2017>, data available at <https://www.pbl.nl/en/image/links/hyde>, accessed 2017-10-25, 2017a.
- Klein Goldewijk, K., Beusen, A., Doelman, J., and Stehfest, E.: Anthropogenic land use estimates for the Holocene – HYDE 3.2, *Earth System Science Data*, 9, 927–953, <https://doi.org/10.5194/essd-9-927-2017>, 2017b.
- Kleinen, T., Brovkin, V., and Schuldt, R. J.: A dynamic model of wetland extent and peat accumulation: results for the Holocene, *Biogeosciences*, 9, 235–248, <https://doi.org/10.5194/bg-9-235-2012>, 00017, 2012.
- Lammers, R. B., Shiklomanov, A. I., Vörösmarty, C. J., Fekete, B. M., and Peterson, B. J.: Assessment of contemporary Arctic river runoff based on observational discharge records, *Journal of Geophysical Research: Atmospheres*, 106, 3321–3334, <https://doi.org/10.1029/2000JD900444>, 2001.
- Lammers, R. B., Shiklomanov, A. I., Vörösmarty, C. J., Fekete, B. M., and Peterson, B. J.: R-ArcticNet, A Regional Hydrographic Data Network for the Pan-Arctic Region (ISO-image of CD-ROM), <https://doi.org/10.1594/PANGAEA.859422>, accessed 2026-01-14, 2016.
- Lange, S., Menz, C., Gleixner, S., Cucchi, M., Weedon, G. P., Amici, A., Bellouin, N., Schmied, H. M., Hersbach, H., Buontempo, C., and Cagnazzo, C.: WFDE5 over land merged with ERA5 over the ocean (W5E5 v2.0), <https://doi.org/10.48364/ISIMIP.342217>, 2021.
- Lange, S., Mengel, M., Treu, S., and Büchner, M.: ISIMIP3a atmospheric climate input data, <https://doi.org/10.48364/ISIMIP.982724.2>, 2023.
- Lawrence, D. and Slater, A.: Incorporating organic soil into a global climate model, *Climate Dynamics*, 30, 145–160, <http://dx.doi.org/10.1007/s00382-007-0278-1>, 10.1007/s00382-007-0278-1, 2008.
- Lee, H., Schuur, E. A., Inglett, K. S., Lavoie, M., and Chanton, J. P.: The rate of permafrost carbon release under aerobic and anaerobic conditions and its potential effects on climate, *Global Change Biology*, 18, 515–527, 2012.

- Lehner, B. and Döll, P.: Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296, 1–22, <https://doi.org/10.1016/j.jhydrol.2004.03.028>, data available at <https://www.worldwildlife.org/pages/global-lakes-and-wetlands-database>, accessed 2018-08-15, 2004.
- 1265 Lehner, B., Liermann, C. R., Revenga, C., Vörösmarty, C., Fekete, B., Crouzet, P., Döll, P., Endejan, M., Frenken, K., and Magome, J.: High-resolution mapping of the world's reservoirs and dams for sustainable river-flow management, *Frontiers in Ecology and the Environment*, 9, 494–502, 2011.
- Li, Y., Shang, J., Zhang, C., Zhang, W., Niu, L., Wang, L., and Zhang, H.: The role of freshwater eutrophication in greenhouse gas emissions: A review, *Science of The Total Environment*, 768, 144 582, <https://doi.org/10.1016/j.scitotenv.2020.144582>,
1270 2021.
- Lutz, F., Herzfeld, T., Heinke, J., Rolinski, S., Schaphoff, S., Von Bloh, W., Stoorvogel, J. J., and Müller, C.: Simulating the effect of tillage practices with the global ecosystem model LPJmL (version 5.0-tillage), *Geoscientific Model Development*, 12, 2419–2440, 2019.
- Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen,
1275 G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753–788, <https://doi.org/10.5194/bg-10-753-2013>, 2013.
- Mengel, M., Treu, S., Lange, S., and Frieler, K.: ATTRICI v1.1 – counterfactual climate for impact attribution, *Geoscientific Model Development*, 14, 5269–5284, <https://doi.org/10.5194/gmd-14-5269-2021>, 2021.
1280
- Mohan, C., Western, A. W., Wei, Y., and Saft, M.: Predicting groundwater recharge for varying land cover and climate conditions – a global meta-study, *Hydrology and Earth System Sciences*, 22, 2689–2703, <https://doi.org/10.5194/hess-22-2689-2018>, 2018.
- Monfreda, C., Ramankutty, N., and Foley, J. a.: Farming the planet: 2. Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000, *Global Biogeochem. Cycles*, 22, GB1022, <https://doi.org/10.1029/2007GB002947>, data available from <http://www.earthstat.org/>, accessed 2018-10-30, 2008.
1285
- Moore, T. R., Roulet, N. T., and Waddington, J. M.: Uncertainty in predicting the effect of climatic change on the carbon cycling of Canadian peatlands, *Climatic Change*, 40, 229–245, <https://doi.org/10.1023/A:1005439401383>, 2011.
- Mora, B., Tsendbazar, N.-E., Herold, M., and Arino, O.: Global land cover mapping: Current status and future trends, *Land use and land cover mapping in Europe: practices & trends*, pp. 11–30, 2014.
1290
- Muehleisen, A. J., Schwartz, N. B., Stump, S. M., and Staver, A. C.: Rainfall variability and deciduous-evergreen coexistence in tropical forests, *Theoretical Ecology*, 17, 255–268, <https://doi.org/10.1007/s12080-024-00588-2>, 2024.
- Mueller, N.: Crop-specific global fertilizer application rates from "Closing yield gaps through nutrient and water management", Version 1, Zenodo, <https://doi.org/10.5281/zenodo.5260732>, accessed 2022-03-08, 2012.
- 1295 Mueller, N. D., Gerber, J. S., Johnston, M., Ray, D. K., Ramankutty, N., and Foley, J. A.: Closing yield gaps through nutrient and water management, *Nature*, 490, 254–257, <https://doi.org/10.1038/nature11420>, 2012.
- Müller Schmied, H., Cáceres, D., Eisner, S., Flörke, M., Herbert, C., Niemann, C., Peiris, T. A., Popat, E., Portmann, F. T., Reinecke, R., Schumacher, M., Shadkam, S., Telteu, C.-E., Trautmann, T., and Döll, P.: The global water resources

- and use model WaterGAP v2.2d: model description and evaluation, *Geoscientific Model Development*, 14, 1037–1079, <https://doi.org/10.5194/gmd-14-1037-2021>, 2021.
- Murguia-Flores, F., Arndt, S., Ganesan, A. L., Murray-Tortarolo, G., and Hornibrook, E. R. C.: Soil Methanotrophy Model (MeMo v1.0): a process-based model to quantify global uptake of atmospheric methane by soil, *Geoscientific Model Development*, 11, 2009–2032, <https://doi.org/10.5194/gmd-11-2009-2018>, 2018.
- Nascimento, H. L. B., Pedreira, B. C., Sollenberger, L. E., Pereira, D. H., Magalhães, C. A. S., and Chizzotti, F. H. M.: Herbage accumulation, canopy structure and tiller morphology of marandu palisadegrass growing in open pasture and in silvopasture, *Agroforestry Systems*, 95, 339–352, <https://doi.org/10.1007/s10457-020-00590-7>, 2021.
- Niu, G.-Y. and Yang, Z.-L.: Effects of frozen soil on snowmelt runoff and soil water storage at a continental scale, *Journal of Hydrometeorology*, 7, 937–952, 2006.
- Niu, G.-Y., Yang, Z.-L., Dickinson, R. E., and Gulden, L. E.: A simple TOPMODEL-based runoff parameterization (SIMTOP) for use in global climate models, *Journal of Geophysical Research: Atmospheres*, 110, 2005.
- NOAA Global Monitoring Laboratory: Trends in Atmospheric Methane (CH₄), https://gml.noaa.gov/ccgg/trends_ch4/, accessed: 2025-07-10, 2025.
- NOAA Global Monitoring Laboratory: CarbonTracker-CH₄ Flux Histograms – Global Region, <https://gml.noaa.gov/ccgg/carbontracker-ch4/fluxhistograms.php?region=GLOB>, accessed: 2025-07-10, n.d.
- Nzotungicimpaye, C.-M., Zickfeld, K., MacDougall, A. H., Melton, J. R., Treat, C. C., Eby, M., and Lesack, L. F. W.: WET-METH 1.0: a new wetland methane model for implementation in Earth system models, *Geoscientific Model Development*, 14, 6215–6240, <https://doi.org/10.5194/gmd-14-6215-2021>, 2021.
- Oleson, K. W., Lawrence, D. M., Bonan, G. B., Drewniak, B., Huang, M., Koven, C. D., Levis, S., Li, F., Riley, W. J., Subin, Z. M., Swenson, S. C., Thornton, P. E., Bozbiyik, A., Fisher, R., Kluzek, E., Lamarque, J.-F., Lawrence, P. J., Muszala, S., Ricciuto, D. M., Sacks, W. J., Sun, Y., Tang, J., and Yang, Z.-L.: Technical Description of version 4.5 of the Community Land Model (CLM4.5), NCAR Technical Note NCAR/TN-503+STR, National Center for Atmospheric Research (NCAR), Boulder, Colorado, USA, <https://doi.org/10.5065/D6RR1W7M>, nCAR Earth System Laboratory, Climate and Global Dynamics Division, 2013.
- Ostberg, S., Müller, C., Heinke, J., and Schaphoff, S.: LandInG 1.0: a toolbox to derive input datasets for terrestrial ecosystem modelling at variable resolutions from heterogeneous sources, *Geoscientific Model Development*, 16, 3375–3406, <https://doi.org/10.5194/gmd-16-3375-2023>, 2023.
- Pan, S., Pan, N., Tian, H., Friedlingstein, P., Sitch, S., Shi, H., Arora, V. K., Haverd, V., Jain, A. K., Kato, E., Lienert, S., Lombardozi, D., Nabel, J. E. M. S., Ottlé, C., Poulter, B., Zaehle, S., and Running, S. W.: Evaluation of global terrestrial evapotranspiration using state-of-the-art approaches in remote sensing, machine learning and land surface modeling, *Hydrology and Earth System Sciences*, 24, 1485–1509, <https://doi.org/10.5194/hess-24-1485-2020>, 2020.
- Parolin, P. and Wittmann, F.: Struggle in the flood: tree responses to flooding stress in four tropical floodplain systems, *AoB PLANTS*, 2010, plq003, <https://doi.org/10.1093/aobpla/plq003>, 2010.
- Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., Poindexter, C., Chen, J., Elbashandy, A., Humphrey, M., Isaac, P., Polidori, D., Ribeca, A., van Ingen, C., Zhang, L., Amiro, B., Ammann, C., Arain, M. A., ArdÃ¶, J., Arkebauer, T., Arndt, S. K., Arriga, N., Aubinet, M., Aurela, M., Baldocchi, D., Barr, A., Beamesderfer, E., Marchesini, L. B., Bergeron, O., Beringer, J., Bernhofer, C., Berveiller, D., Billesbach, D., Black, T. A., Blanken, P. D., Bohrer, G.,

- Boike, J., Bolstad, P. V., Bonal, D., Bonnefond, J.-M., Bowling, D. R., Bracho, R., Brodeur, J., Brämmer, C., Buchmann, N., Burban, B., Burns, S. P., Buysse, P., Cale, P., Cavagna, M., Cellier, P., Chen, S., Chini, I., Christensen, T. R., Cleverly, J., Collalti, A., Consalvo, C., Cook, B. D., Cook, D., Coursolle, C., Cremonese, E., Curtis, P. S., Da Andrea, E., da Rocha, H., Dai, X., Davis, K. J., De Cinti, B., de Grandcourt, A., De Ligne, A., De Oliveira, R. C., Delpierre, N., Desai, A. R., Di Bella, C. M., di Tommasi, P., Dolman, H., Domingo, F., Dong, G., Dore, S., Duce, P., Dufrêne, E., Dunn, A., Duarek, J., Eamus, D., Eichelmann, U., ElKhidir, H. A. M., Eugster, W., Ewenz, C. M., Ewers, B., Famulari, D., Fares, S., Feigenwinter, I., Feitz, A., Fensholt, R., Filippa, G., Fischer, M., Frank, J., Galvagno, M., Gharun, M., Gianelle, D., Gielen, B., Gioli, B., Gitelson, A., Goded, I., Goeckede, M., Goldstein, A. H., Gough, C. M., Goulden, M. L., Graf, A., Griebel, A., Gruening, C., Grunwald, T., Hammerle, A., Han, S., Han, X., Hansen, B. U., Hanson, C., Hatakka, J., He, Y., Hehn, M., Heinesch, B., Hinko-Najera, N., Hörtnagl, L., Hutley, L., Ibrom, A., Ikawa, H., Jackowicz-Korczynski, M., Janoušek, D., Jans, W., Jassal, R., Jiang, S., Kato, T., Khomik, M., Klatt, J., Knohl, A., Knox, S., Kobayashi, H., Koerber, G., Kolle, O., Kosugi, Y., Kotani, A., Kowalski, A., Kruijt, B., Kurbatova, J., Kutsch, W. L., Kwon, H., Launiainen, S., Laurila, T., Law, B., Leuning, R., Li, Y., Liddell, M., Limousin, J.-M., Lion, M., Liska, A. J., Lohila, A., López-Ballesteros, A., López-Blanco, E., Loubet, B., Loustau, D., Lucas-Moffat, A., Lüscher, J., Ma, S., Macfarlane, C., Magliulo, V., Maier, R., Mammarella, I., Manca, G., Marcolla, B., Margolis, H. A., Marras, S., Massman, W., Mastepanov, M., Matamala, R., Matthes, J. H., Mazzenga, F., McCaughey, H., McHugh, I., McMillan, A. M. S., Merbold, L., Meyer, W., Meyers, T., Miller, S. D., Minerbi, S., Moderow, U., Monson, R. K., Montagnani, L., Moore, C. E., Moors, E., Moreaux, V., Moureaux, C., Munger, J. W., Nakai, T., Neiryneck, J., Nesic, Z., Nicolini, G., Noormets, A., Northwood, M., Nosetto, M., Nouvellon, Y., Novick, K., Oechel, W., Olesen, J. E., Ourcival, J.-M., Papuga, S. A., Parmentier, F.-J., Paul-Limoges, E., Pavelka, M., Peichl, M., Pendall, E., Phillips, R. P., Pilegaard, K., Pirk, N., Posse, G., Powell, T., Prasse, H., Prober, S. M., Rambal, S., Rannik, A., Raz-Yaseef, N., Reed, D., de Dios, V. R., Restrepo-Coupe, N., Reverter, B. R., Roland, M., Sabbatini, S., Sachs, T., Saleska, S. R., Sanchez-Cañete, E. P., Sanchez-Mejia, Z. M., Schmid, H. P., Schmidt, M., Schneider, K., Schrader, F., Schroder, I., Scott, R. L., Sedláček, P., Serrano-Ortiz, P., Shao, C., Shi, P., Shironya, I., Siebicke, L., Åsigt, L., Silberstein, R., Sirca, C., Spano, D., Steinbrecher, R., Stevens, R. M., Sturtevant, C., Suyker, A., Tagesson, T., Takanashi, S., Tang, Y., Tapper, N., Thom, J., Tiedemann, F., Tomassucci, M., Tuovinen, J.-P., Urbanski, S., Valentini, R., van der Molen, M., van Gorsel, E., van Huissteden, K., Varlagin, A., Verfaillie, J., Vesala, T., Vincke, C., Vitale, D., Vygodskaya, N., Walker, J. P., Walter-Shea, E., Wang, H., Weber, R., Westermann, S., Wille, C., Wofsy, S., Wohlfahrt, G., Wolf, S., Woodgate, W., Li, Y., Zampedri, R., Zhang, J., Zhou, G., Zona, D., Agarwal, D., Biraud, S., Torn, M., and Papale, D.: The FLUXNET2015 dataset and the ONEFlux processing pipeline for eddy covariance data, *Scientific Data*, 7, 225, <https://doi.org/10.1038/s41597-020-0534-3>, 2020.
- Pazhanivelan, S., Sudarmanian, N. S., Geethalakshmi, V., Deiveegan, M., Ragunath, K., Sivamurugan, A. P., and Shanmugapriya, P.: Assessing Methane Emissions from Rice Fields in Large Irrigation Projects Using Satellite-Derived Land Surface Temperature and Agronomic Flooding: A Spatial Analysis, *Agriculture*, 14, <https://doi.org/10.3390/agriculture14030496>, 2024.
- Portmann, F. T., Siebert, S., and Döll, P.: MIRCA2000-Global monthly irrigated and rainfed crop areas around the year 2000: A new high-resolution data set for agricultural and hydrological modeling, *Global Biogeochem. Cycles*, 24, GB1011, <https://doi.org/10.1029/2008GB003435>, 2010a.

- Portmann, F. T., Siebert, S., and Döll, P.: MIRCA2000, Version 1.1 [Data set], Zenodo,
1375 <https://doi.org/10.5281/zenodo.7422506>, accessed 2018-03-19, 2010b.
- Prestele, R., Alexander, P., Rounsevell, M. D. A., Arneth, A., Calvin, K., Doelman, J., Eitelberg, D. A., Engström, K.,
Fujimori, S., Hasegawa, T., Havlik, P., Humpenöder, F., Jain, A. K., Krisztin, T., Kyle, P., Meiyappan, P., Popp, A.,
Sands, R. D., Schaldach, R., Schüngel, J., Stehfest, E., Tabeau, A., Van Meijl, H., Van Vliet, J., and Verburg, P. H.:
Hotspots of uncertainty in land-use and land-cover change projections: a global-scale model comparison, *Global Change*
1380 *Biology*, 22, 3967–3983, <https://doi.org/https://doi.org/10.1111/gcb.13337>, 2016.
- Qian, H., Zhu, X., Huang, S., Linquist, B., Kuzyakov, Y., Wassmann, R., Minamikawa, K., Martinez-Eixarch, M., Yan, X.,
Zhou, F., Sander, B. O., Zhang, W., Shang, Z., Zou, J., Zheng, X., Li, G., Liu, Z., Wang, S., Ding, Y., van Groenigen,
K. J., and Jiang, Y.: Greenhouse gas emissions and mitigation in rice agriculture, *Nature Reviews Earth & Environment*,
4, 716–732, <https://doi.org/10.1038/s43017-023-00482-1>, 2023.
- 1385 R-ArcticNET Project: R-ArcticNET v4.0 Available Data (Site Attributes and Discharge Tables), <https://www.r-arcticnet.sr.unh.edu/v4.0/AllData/index.html>, accessed 2026-01-14.
- Riley, W., Subin, Z., Lawrence, D., Swenson, S., Torn, M., Meng, L., Mahowald, N., and Hess, P.: Barriers to predicting
changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in
CESM, *Biogeosciences*, 8, 1925–1953, 2011.
- 1390 Ringeval, B., Friedlingstein, P., Koven, C., Ciais, P., de Noblet-Ducoudré, N., Decharme, B., and Cadule, P.: Climate-
CH₄ feedback from wetlands and its interaction with the climate-CO₂ feedback, *Biogeosciences*, 8, 2137–2157,
<https://doi.org/10.5194/bg-8-2137-2011>, 2011.
- Salmon, E., Jégou, F., Guenet, B., Jourdain, L., Qiu, C., Bastrikov, V., Ducharne, A., Ciais, P., Zhu, D., Peng, S., Krinner,
G., Peylin, P., Bousquet, P., Janssens, I. A., Kleinen, T., Koven, C. D., Luyssaert, S., McGuire, A. D., Melton, J. R.,
1395 Ringeval, B., Saunois, M., Schneider von Deimling, T., Serça, D., Spahni, R., Treat, C. C., Wania, R., Zhang, Z., and
Ziemblińska, K.: Assessing methane emissions for northern peatlands in ORCHIDEE-PEAT revision 7020, *Geoscientific*
Model Development, 15, 2813–2838, <https://doi.org/10.5194/gmd-15-2813-2022>, 2022.
- Santoro, M., Cartus, O., Carvalhais, N., Rozendaal, D. M. A., Avitabile, V., Araza, A., de Bruin, S., Herold, M., Quegan, S.,
Rodríguez-Veiga, P., Balzter, H., Carreiras, J., Schepaschenko, D., Korets, M., Shimada, M., Itoh, T., Moreno Martínez,
1400 Á., Cavlović, J., Cazzolla Gatti, R., and others: The global forest above-ground biomass pool for 2010 estimated from high-
resolution satellite observations, *Earth System Science Data*, 13, 3927–3950, <https://doi.org/10.5194/essd-13-3927-2021>,
2021.
- Saunois, M., Bousquet, P., Poulter, B., Peregon, A., Ciais, P., Canadell, J. G., Dlugokencky, E. J., Etiope, G., Bastviken,
D., Houweling, S., Janssens-Maenhout, G., Tubiello, F. N., Castaldi, S., Jackson, R. B., Alexe, M., Arora, V. K., Beerling,
1405 D. J., Bergamaschi, P., Blake, D. R., Brailsford, G., Brovkin, V., Bruhwiler, L., Crevoisier, C., Crill, P., Covey, K., Curry,
C., Frankenberg, C., Gedney, N., Höglund-Isaksson, L., Ishizawa, M., Ito, A., Joos, F., Kim, H.-S., Kleinen, T., Krummel,
P., Lamarque, J.-F., Langenfelds, R., Locatelli, R., Machida, T., Maksyutov, S., McDonald, K. C., Marshall, J., Melton,
J. R., Morino, I., Naik, V., O’Doherty, S., Parmentier, F.-J. W., Patra, P. K., Peng, C., Peng, S., Peters, G. P., Pison, I.,
Prigent, C., Prinn, R., Ramonet, M., Riley, W. J., Saito, M., Santini, M., Schroeder, R., Simpson, I. J., Spahni, R., Steele,
1410 P., Takizawa, A., Thornton, B. F., Tian, H., Tohjima, Y., Viovy, N., Voulgarakis, A., van Weele, M., van der Werf, G. R.,
Weiss, R., Wiedinmyer, C., Wilton, D. J., Wiltshire, A., Worthy, D., Wunch, D., Xu, X., Yoshida, Y., Zhang, B., Zhang, Z.,

- and Zhu, Q.: The global methane budget 2000–2012, *Earth System Science Data*, 8, 697–751, <https://doi.org/10.5194/essd-8-697-2016>, 2016.
- Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J.,
1415 Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler,
L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G.,
Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout,
G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T.,
Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F.,
1420 Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet,
M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton,
B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der
Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B.,
Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth System Science Data*, 12, 1561–1623,
1425 <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P. A., Regnier, P., Canadell, J. G., Jackson, R. B., Patra, P. K.,
Bousquet, P., Ciais, P., Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, D. J., Belikov, D. A., Blake,
D. R., Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgerson,
M. A., Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P. B.,
1430 Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R., Mühle, J., Müller, J., Murguia-Flores, F., Niwa, Y., Noce,
S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers,
A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., van der Werf, G. R.,
Worthy, D. E. J., Xi, Y., Yoshida, Y., Zhang, W., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget
2000–2020, *Earth System Science Data*, 17, 1873–1958, <https://doi.org/10.5194/essd-17-1873-2025>, 2025.
- 1435 Schaphoff, S., Heyder, U., Ostberg, S., Gerten, D., Heinke, J., and Lucht, W.: Contribution of permafrost soils to the global
carbon budget, *Environmental Research Letters*, 8, 014026, <http://stacks.iop.org/1748-9326/8/i=1/a=014026>, 2013.
- Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr, J., Knauer,
J., Langerwisch, F., Lucht, W., Müller, C., Rolinski, S., and Waha, K.: LPJmL4 – a dynamic global vegetation model with
managed land – Part 1: Model description, *Geoscientific Model Development*, 11, 1343–1375, [https://doi.org/10.5194/gmd-](https://doi.org/10.5194/gmd-11-1343-2018)
1440 11-1343-2018, 2018.
- Schaphoff, S., Hötten, D., Müller, C., Gerten, D., Ostberg, S., and von Bloh, W.: Code and data for: "Implementing methane
dynamics into the LPJmL6 model", <https://doi.org/10.5281/zenodo.17877397>, 2025a.
- Schaphoff, S., von Bloh, W., Rammig, A., Thonicke, K., Biemans, H., Forkel, M., Gerten, D., Heinke, J., Jägermeyr,
J., Langerwisch, F., Lucht, W., Rolinski, S., Waha, K., Ostberg, S., Wirth, S. B., Fader, M., Drüke, M., Jans, Y.,
1445 Lutz, F., Herzfeld, T., Minoli, S., Porwollik, V., Stehfest, E., de Waal, L., Beringer (Erbrecht), T., Rost (Jachner), S.,
Gumpenberger, M., Heyder, U., Werner, C., Braun, J., Breier, J., Stenzel, F., Mathesius, S., Hemmen, M., Billing, M.,
Oberhagemann, L., Sakschewski, B., and Müller, C.: LPJmL: central open-source github repository of LPJmL at PIK,
<https://doi.org/10.5281/zenodo.17911633>, 2025b.

- Schneider, U., Becker, A., Finger, P., Meyer-Christoffer, A., Ziese, M., and Rudolf, B.: GPCC Full Data Monthly Product Version 2018 at 1.0°: Monthly Land-Surface Precipitation from Rain-Gauges built on GTS-based and Historical Data, Global Precipitation Climatology Centre (GPCC), https://doi.org/10.5676/DWD_GPCC/FD_M_V2018_100, 2017.
- Sivapalan, M., Beven, K., and Wood, E. F.: On hydrologic similarity: 2. A scaled model of storm runoff production, *Water Resources Research*, 23, 2266–2278, <https://doi.org/10.1029/WR023i012p02266>, 1987.
- Stanley, E. H., Casson, N. J., Christel, S. T., Crawford, J. T., Loken, L. C., and Oliver, S. K.: The ecology of methane in streams and rivers: patterns, controls, and global significance, *Ecological Monographs*, 86, 146–171, <https://doi.org/10.1890/15-1027>, 2016.
- Stevenson, D. S., Zhao, A., Naik, V., O'Connor, F. M., Tilmes, S., Zeng, G., Murray, L. T., Collins, W. J., Griffiths, P. T., Shim, S., Horowitz, L. W., Sentman, L. T., and Emmons, L.: Trends in global tropospheric hydroxyl radical and methane lifetime since 1850 from AerChemMIP, *Atmospheric Chemistry and Physics*, 20, 12 905–12 920, <https://doi.org/10.5194/acp-20-12905-2020>, 2020.
- Swenson, S. C. and Lawrence, D. M.: Assessing a groundwater and runoff parameterization using GRACE-based groundwater storage estimates, *Journal of Advances in Modeling Earth Systems*, 6, 779–799, <https://doi.org/10.1002/2013MS000282>, 2014.
- Swenson, S. C., Lawrence, D. M., and Lee, H.: Improved simulation of the terrestrial hydrological cycle in permafrost regions by the Community Land Model, *Journal of Advances in Modeling Earth Systems*, 4, <https://doi.org/https://doi.org/10.1029/2012MS000165>, 2012a.
- Swenson, S. C., Lawrence, D. M., and Lee, H.: Improved simulation of the terrestrial hydrological cycle in permafrost regions by the Community Land Model, *Journal of Advances in Modeling Earth Systems*, 4, n/a–n/a, <https://doi.org/10.1029/2012MS000165>, 2012b.
- Takahashi, H., Yamauchi, T., Colmer, T. D., and Nakazono, M.: Aerenchyma Formation in Plants, in: *Low-Oxygen Stress in Plants: Oxygen Sensing and Adaptive Responses to Hypoxia*, pp. 247–265, Springer Vienna, Vienna, https://doi.org/10.1007/978-3-7091-1254-0_13, 2014.
- Turetsky, M. R., Kotowska, A., Bubier, J., Dise, N. B., Crill, P., Hornibrook, E. R. C., Minkinen, K., Moore, T. R., Myers-Smith, I. H., Nykänen, H., Olefeldt, D., Rinne, J., Saarnio, S., Shurpali, N., Tuittila, E.-S., Waddington, J. M., White, J. R., Wickland, K. P., and Wilmking, M.: A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands, *Global Change Biology*, 20, 2183–2197, <https://doi.org/10.1111/gcb.12580>, 2014.
- U.S. Geological Survey: USGS EROS Archive - Digital Elevation - Global 30 Arc-Second Elevation (GTOPO30), https://www.usgs.gov/centers/eros/science/usgs-eros-archive-digital-elevation-global-30-arc-second-elevation-gtopo30?qt-science_center_objects=0#qt-science_center_objects, accessed: 2018, 2007a.
- U.S. Geological Survey: USGS EROS Archive - Digital Elevation - HYDRO1k, <https://www.usgs.gov/centers/eros/science/usgs-eros-archive-digital-elevation-hydro1k#overview>, accessed: 2010, 2007b.
- van den Pol-van Dasselaar, A.: Methane emissions from grasslands, internal phd, wu, Netherlands, <https://doi.org/10.18174/201574>, wU thesis 2483 Proefschrift Wageningen, 1998.
- Verdin, K. L., Hall, F. G., Collatz, G. J., Meeson, B. W., Los, S. O., Brown de Colstoun, E., and Landis, D. R.: ISLSCP II HYDRO1k Elevation-derived Products, ISLSCP Initiative II Collection, Oak Ridge National Laboratory Distributed Active Archive Center (ORNL DAAC), Oak Ridge, Tennessee, USA, doi:10.3334/ORNLDAAC/1007, 2011.

- von Bloh, W., Schaphoff, S., Müller, C., Rolinski, S., Waha, K., and Zaehle, S.: Implementing the nitrogen cycle into the dynamic global vegetation, hydrology, and crop growth model LPJmL (version 5.0), Geoscientific Model Development, 11, 2789–2812, <https://doi.org/10.5194/gmd-11-2789-2018>, 2018.
- 1490 Vörösmarty, C. J., Fekete, B. M., and Tucker, B. A.: Global River Discharge, 1807–1991, Version 1.1 (RivDIS), <https://doi.org/10.3334/ORNLDAAAC/199>, 1998.
- Vörösmarty, C. J., Fekete, B. M., Meybeck, M., and Lammers, R. B.: Global system of rivers: Its role in organizing continental land mass and defining land-to-ocean linkages, Global Biogeochem. Cycles, 14, 599–621, <https://doi.org/10.1029/1999GB900092>, 2000.
- 1495 Wada, Y. and Bierkens, M. F. P.: Sustainability of global water use: past reconstruction and future projections, Environmental Research Letters, 9, 104003, <https://doi.org/10.1088/1748-9326/9/10/104003>, 2014a.
- Wada, Y. and Bierkens, M. F. P.: Integrated assessment of global water resources using PCR-GLOBWB: Current state and future projections, Hydrology and Earth System Sciences, 18, 2853–2873, <https://doi.org/10.5194/hess-18-2853-2014>, 2014b.
- 1500 Waha, K., Dietrich, J. P., Portmann, F. T., Siebert, S., Thornton, P. K., Bondeau, A., and Herrero, M.: Multiple cropping systems of the world and the potential for increasing cropping intensity, Global Environ. Change, 64, 102131, <https://doi.org/10.1016/j.gloenvcha.2020.102131>, 2020.
- Waha, K., Folberth, C., Biemans, H., Boere, E., Bondeau, A., Hartley, A. J., Hoogenboom, G., Jägermeyr, J., Liu, Y., Mathison, C., Müller, C., Nkwasa, A., Olin, S., Ruane, A. C., De Vos, K., White, J. W., Williams, K., and Yu, Q.: Land use
1505 modelling needs to better account for multiple cropping to inform pathways for sustainable agriculture, Commun. Earth Environ., 6, 1–13, <https://doi.org/10.1038/s43247-025-02724-0>, 2025.
- Walter, B. P. and Heimann, M.: A process-based, climate-sensitive model to derive methane emissions from natural wetlands: Application to five wetland sites, sensitivity to model parameters, and climate, Global Biogeochem. Cycles, 14, 745–765, <http://dx.doi.org/10.1029/1999GB001204>, 2000.
- 1510 Wang, J., Akiyama, H., Yagi, K., and Yan, X.: Controlling variables and emission factors of methane from global rice fields, Atmospheric Chemistry and Physics, 18, 10419–10431, <https://doi.org/10.5194/acp-18-10419-2018>, 2018.
- Wang, J., Ciais, P., Smith, P., Yan, X., Kuzyakov, Y., Liu, S., Li, T., and Zou, J.: The role of rice cultivation in changes in atmospheric methane concentration and the Global Methane Pledge, Global Change Biology, 29, 2776–2789, <https://doi.org/https://doi.org/10.1111/gcb.16631>, 2023a.
- 1515 Wang, L., Arora, V. K., Bartlett, P., Chan, E., and Curasi, S. R.: Mapping of ESA’s Climate Change Initiative land cover data to plant functional types for use in the CLASSIC land model, Biogeosciences, 20, 2265–2282, <https://doi.org/10.5194/bg-20-2265-2023>, 2023b.
- Wang, Y., Hu, Z., Liu, C., Wu, Z., and Chen, S.: Methane emissions in japonica rice paddy fields under different elevated CO₂ concentrations, Nutrient Cycling in Agroecosystems, 122, 173–189, <https://doi.org/10.1007/s10705-022-10197-2>, 2022.
- 1520 Wania, R., Ross, I., and Prentice, I. C.: Integrating peatlands and permafrost into a dynamic global vegetation model: 2. Evaluation and sensitivity of vegetation and carbon cycle processes, Global Biogeochem. Cycles, 23, GB3015–, <http://dx.doi.org/10.1029/2008GB003413>, 2009.
- Wania, R., Ross, I., and Prentice, I.: Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHyMe v1. 3, Geoscientific Model Development Discussions, 3, 1–59, 2010.

- 1525 Wania, R., Melton, J., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R. E., Bohn, T., Avis, C. A., Chen, G., Eliseev, A. V., Hopcroft, P. O., Riley, W. J., Subin, Z. M., Tian, H., van Bodegom, P. M., Kleinen, T., Yu, Z. C., Singarayer, J. S., Zürcher, S., Lettenmaier, D., Beerling, D. J., Denisov, S. N., Prigent, C., Papa, F., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: methodology of a model inter-comparison project (WETCHIMP), *Geoscientific Model Development*, 6, 617–641, <https://doi.org/10.5194/gmd-6-617-2013>, 2013.
- 1530 Wassmann, R., Neue, H.-U., Lantin, R. S., Buendia, L. V., and Rennenberg, H.: Characterization of Methane Emissions from Rice Fields in Asia. I. Comparison among Field Sites in Five Countries, *Nutrient Cycling in Agroecosystems*, 58, 1–12, <https://doi.org/10.1023/A:1009848813994>, 2000.
- Whalen, S. C. and Reeburgh, W. S.: Methane emissions from boreal wetlands, *Journal of Geophysical Research: Atmospheres*, 96, 16 971–16 982, <https://doi.org/10.1029/91JD00048>, 1991.
- 1535 Wiesenburg, D. A. and Guinasso Jr, N. L.: Equilibrium solubilities of methane, carbon monoxide, and hydrogen in water and sea water, *Journal of chemical and engineering data*, 24, 356–360, 1979.
- Wild, M., Folini, D., Hakuba, M. Z., Schär, C., Seneviratne, S. I., Kato, S., Rutan, D., Ammann, C., Wood, E. F., and König-Langlo, G.: The energy balance over land and oceans: An assessment based on direct observations and CMIP5 climate models, *Climate Dynamics*, 44, 3393–3429, <https://doi.org/10.1007/s00382-014-2430-z>, 2015.
- 1540 Wirth, S. B., Braun, J., Heinke, J., Ostberg, S., Rolinski, S., Schaphoff, S., Stenzel, F., von Bloh, W., Taube, F., and Müller, C.: Biological nitrogen fixation of natural and agricultural vegetation simulated with LPJmL 5.7.9, *Geoscientific Model Development*, 17, 7889–7914, <https://doi.org/10.5194/gmd-17-7889-2024>, 2024.
- Xu, L., Saatchi, S. S., Yang, Y., Yu, Y., Pongratz, J., Bloom, A. A., Bowman, K., Worden, J., Liu, J., Yin, Y., Domke, G., McRoberts, R. E., Woodall, C., Nabuurs, G.-J., de Miguel, S., Keller, M., Harris, N., Maxwell, S., and Schimel, D.: Dataset for “Changes in Global Terrestrial Live Biomass over the 21st Century”, <https://doi.org/10.5281/zenodo.4161694>, data set, 2021.
- 1545 Xu, R., Dai, J., Luo, W., Yin, X., Li, Y., Tai, X., Han, L., Chen, Y., Lin, L., and Li, G.: A photothermal model of leaf area index for greenhouse crops, *Agricultural and Forest Meteorology*, 150, 541–552, <https://doi.org/10.1016/j.agrformet.2010.01.019>, 2010.
- 1550 Yamamoto, S., Alcauskas, J. B., and Crozier, T. E.: Solubility of methane in distilled water and seawater, *Journal of Chemical and Engineering Data*, 21, 78–80, 1976.
- Yan, X., Akiyama, H., Yagi, K., and Akimoto, H.: Global estimations of the inventory and mitigation potential of methane emissions from rice cultivation conducted using the 2006 IPCC Guidelines, *Global Biogeochemical Cycles*, 23, GB2002, <https://doi.org/10.1029/2008GB003299>, 2009.
- 1555 Yang, T., He, Q., Jiang, J., Sheng, L., Jiang, H., and He, C.: Impact of Water Table on Methane Emission Dynamics in Terrestrial Wetlands and Implications on Strategies for Wetland Management and Restoration, *Wetlands*, 42, 120, <https://doi.org/10.1007/s13157-022-01634-7>, 2022.
- Zeng, X. and Decker, M.: Improving the numerical solution of soil moisture-based Richards equation for land models with a deep or shallow water table, *Journal of Hydrometeorology*, 10, 308–319, <https://doi.org/10.1175/2008JHM1011.1>, 2009.
- 1560 Zhang, B., Tian, H., Lu, C., Dangal, S. R. S., Yang, J., and Pan, S.: Manure nitrogen production and application in cropland and rangeland during 1860–2014: A 5-minute gridded global data set for Earth system modeling, *Earth Syst. Sci. Data Discuss.*, pp. 1–35, <https://doi.org/10.5194/essd-2017-11>, 2017.

- Zhang, B., Tian, H., Lu, C., Dangal, S. R. S., Yang, J., and Pan, S.: Manure nitrogen production and application in cropland and rangeland during 1860 - 2014: A 5-minute gridded global data set for Earth system modeling, [Data set], PANGAEA, 1565 <https://doi.org/10.1594/PANGAEA.871980>, accessed 2022-05-04, 2017.
- Zhang, Z., Fluet-Chouinard, E., Jensen, K., McDonald, K., Hugelius, G., Gumbrecht, T., Carroll, M., Prigent, C., Bartsch, A., and Poulter, B.: Development of the global dataset of Wetland Area and Dynamics for Methane Modeling (WAD2M), Earth System Science Data, 13, 2001–2023, <https://doi.org/10.5194/essd-13-2001-2021>, 2021.
- Zhong, L., Dai, Z., Fang, P., Cao, Y., and Wang, L.: A Review: Tree Species Classification Based on Remote Sensing Data 1570 and Classic Deep Learning-Based Methods, Forests, 15, <https://doi.org/10.3390/f15050852>, 2024.
- Zhu, Y. and Li, H.: Methane emissions from rice paddies in the Yangtze River Delta region of China: synthesis of new estimates, International Journal of Environmental Science and Technology, <https://doi.org/10.1007/s13762-024-06050-4>, 2024.