

Supplementary information to :

Southern Ocean biological pump over the last glacial cycle from new diatom transfer functions

Mathieu Rembauville¹, Sylvain Pichat^{1,2}

¹Univ. Lyon, ENS de Lyon, Université Lyon 1, CNRS, UMR 5276 LGL-TPE, 69364, Lyon, France

²Climate Geochemistry Department, Max Planck Institute for Chemistry, Mainz, Germany

Correspondance to : Mathieu Rembauville (mathieu.rembauville@ens-lyon.fr)

Diatom species names updated following Algebase recommendations (www.algeabase.org)

Actinocyclus ehrenbergii (Ralfs) was updated to *Actinocyclus octonarius* (Ehrenberg).

Actinocyclus ingens (Rattray) was updated to *Actinocyclus rattrayi* (Williams).

Coscinodiscus tabularis (Grunow) was updated to *Azpeitia tabularis* (Grunow).

Corethron criophilum (Castracane) was updated to *Corethron pennatum* (Grunow).

Fragilariopsis sublinearis (Van Heurck) was updated to *Fragilariopsis obliquecostata* (Van Heurck).

Hemidiscus cuneiformis (Wallich) was updated to *Actinocyclus cuneiformis* (Wallich).

Thalassiosira trifulta (Fryxell) was updated to *Shionodiscus trifultus* (Fryxell).

Tropidoneis antarctica (Grunow) was updated to *Membraneis challengerii* (Grunow).

Full list of diatom species/taxa groups reported in the sediment trap and sediment core datasets

Actinocyclus actinochilus (Ehrenberg)

Actinocyclus cuneiformis (Wallich)

Actinocyclus curvatulus (Janisch)

Actinocyclus exiguus (Fryxell & Semina)

Actinocyclus octonarius (Ehrenberg)

Actinocyclus rattrayi (Williams)

Actinocyclus spp.

Alveus marinus (Kaczmarska & Fryxell)

Amphora copulata (Schoeman & R.E.M. Archibald)

Asteromphalus heptactis (Brébisson)

Asteromphalus hookeri (Ehrenberg)

Asteromphalus hyalinus (Karsten)

Asteromphalus parvulus (Karsten)

Asteromphalus spp.

Azpeitia tabularis (Grunow)
Banquisia belgicae (Van Heurck)
Chaetoceros aequatorialis var. *antarcticus* (Manguin)
Chaetoceros atlanticus (Cleve)
Chaetoceros bulbosus (Ehrenberg)
Chaetoceros criophilus (Castracane)
Chaetoceros decipiens (Cleve)
Chaetoceros diadema (Ehrenberg)
Chaetoceros dichchaeta (Ehrenberg)
Chaetoceros peruvianus (Brightwell)
Chaetoceros radicans (Schütt)
Chaetoceros subgenus *Hyalochaete*
Chaetoceros subgenus *Hyalochaete* resting spore
Chaetoceros subgenus *Phaeoceros*
Cocconeis costata (Gregory)
Cocconeis dallmannii (Al-Handal, Riaux-Gobin, Romero & Wulff)
Cocconeis fasciolata (Ehrenberg)
Cocconeis californica (Grunow)
Cocconeis californica var. *keruelensis* (Heiden)
Cocconeis melchioroides (Al-Handal, Riaux-Gobin, Romero & Wulff)
Cocconeis spp.
Corethron inerme (Karsten)
Corethron pennatum (Grunow)
Coscinodiscus asteromphalus (Ehrenberg)
Coscinodiscus cruvatulus (Grunow)
Coscinodiscus marginatus (Ehrenberg)
Coscinodiscus spp.
Cyclotella spp.
Dactyliosolen antarcticus (Castracane)
Denticulopsis spp.
Diploneis bombus (Ehrenberg)
Eucampia antarctica (Castracane)
Eucampia antarctica resting spore
Eupyxidicula turris (Greville)
Fallacia marnieri (Manguin)
Fragilariopsis bicapitata (unreferenced)
Fragilariopsis curta (Van Heurck)
Fragilariopsis cylindrus (Grunow)

Fragilariopsis doliolus (Wallich)
Fragilariopsis kerguelensis (O'Meara)
Fragilariopsis obliquecostata (Van Heurck)
Fragilariopsis panduriformis (unreferenced)
Fragilariopsis pseudonana (Hasle)
Fragilariopsis rhombica (O'Meara)
Fragilariopsis ritsheri (Hustedt)
Fragilariopsis separanda/rhombica
Fragilariopsis separanda (Hustedt)
Fragilariopsis vanheurckii (Peragallo)
Fragilariopsis spp.
Gomphomenopsis littoralis (unreferenced)
Guinardia cylindrus (Cleve)
Guinardia cylindrus (Cleve)
Gyrosigma spp.
Haslea trompii (Cleve)
Leptocylindrus spp.
Licmophora gracilis (Ehrenberg)
Manguinea fusiformis (Manguin)
Melosira adeliae (Manguin)
Membraneis challengeri (Grunow)
Navicula directa (Smith)
Navicula spp.
Nitzschia bicapitata (Cleve)
Nitzschia braarudii (Hasle)
Nitzschia interruptestriata (Simonsen)
Nitzschia kolaczekii (Grunow)
Nitzschia sicala (Castracane)
Nitzschia taeniiformis (Simonsen)
Nitzschia spp.
Odontella weissflogii (Grunow)
Odontella weissflogii resting spore
Paralia sulcata (Ehrenberg)
Paralia spp.
Plagiotropis gaussii (Heiden)
Pleurosigma directum (Grunow)
Pleurosigma spp.
Porosira glacialis (Grunow)

Porosira pseudodenticulata (Hustedt)
Proboscia alata (Brightwell)
Proboscia inermis (Castracane)
Proboscia truncata (Karsten)
Proboscia spp.
Psammodictyon panduriforme (Gregory)
Pseudogomphonema kamtschaticum (Grunow)
Pseudo-nitzschia heimii (Manguin)
Pseudo-nitzschia lineola (Cleve)
Pseudo-nitzschia heimii/lineola
Pseudo-nitzschia prolongatoides (Hasle)
Pseudo-nitzschia turgiduloides (Hasle)
Rhizosolenia antennata (Ehrenberg)
Rhizosolenia bergonii (Peragallo)
Rhizosolenia chunii (Karsten)
Rhizosolenia costata (Gersonde)
Rhizosolenia crassa (Schimper ex Karsten)
Rhizosolenia curvata (Zacharias)
Rhizosolenia hebetata (Bailey)
Rhizosolenia polydactyla (Castracane)
Rhizosolenia sima (Castracane)
Rhizosolenia simplex (Karsten)
Rhizosolenia styliformis (Brightwell)
Rhizosolenia spp.
Roperia tessellata (Roper)
Rouxia spp.
Shionodiscus gracilis (Karsten)
Shionodiscus gracilis var. *expectus* (VanLandingham)
Shionodiscus frenguelliopsis (Fryxell & Johansen)
Shionodiscus oestrupii (Ostenfeld)
Shionodiscus trifultus (Fryxell)
Stellarima microtrias (Ehrenberg)
Stellarima stellaris (Roper)
Synedropsis laevis (Heiden)
Synedropsis recta (Hasle, Medlin & Syversten)
Synedropsis spp.
Thalassionema nitzschioides (Grunow)
Thalassionema nitzschioides var. *capitulata* (Schrader)

Thalassionema nitzschioides var. *lanceolata* (Grunow)
Thalassionema nitzschioides var. *parva* (Heiden & Kolbe)
Thalassiosira antarctica (Karsten)
Thalassiosira decipiens (Grunow)
Thalassiosira eccentrica (Ehrenberg)
Thalassiosira ferelineata (Hasle & Fryxell)
Thalassiosira gracilis (Karsten)
Thalassiosira gracilis var. *expecta* (Van Landingham)
Thalassiosira gravida (Cleve)
Thalassiosira lentiginosa (Janisch)
Thalassiosira leptopus (Grunow)
Thalassiosira lineata (Jousé)
Thalassiosira maculata (Fryxell & Johansen)
Thalassiosira oestrupii (Ostenfeld)
Thalassiosira oliverana (O'Meara)
Thalassiosira scotia (Fryxell & Hoban)
Thalassiosira symmetrica (Fryxell & Hasle)
Thalassiosira tumida (Janisch)
Thalassiosira spp.
Thalassiosira resting spore
Thalassiothrix antarctica (Schimper)
Trachyneis aspera (Ehrenberg)
Trichotoxon reinboldii (Van Heurck)
Unidentified centric < 20 µm
Unidentified centric > 20 µm

Supplementary figures




Figure S1. Scree plot representing eigenvalues associated with the first 10 components of the factorial analysis. Dotted line denotes the eigenvalue of 1, a threshold used to select the informative factors used in the multiple linear regressions (MLR).




Figure S2. Variance explained by the first 10 components of the partial least square regression (PLSR). Line and dots represent variance distribution according to a null broken stick model that evenly distributes a decreasing variance among the components (see materials and methods for explanations). PLSR components explaining more variance than the null model were considered informative and kept for the transfer function calibration.




Figure S3. Parameters selection for the gradient boosting regression (GBR). a) Effect of tree depth on the root mean square error (RMSE) of the prediction. b) Effect of the number of estimators on the RMSE. Each line is the mean from the bootstrapping technique (10 000 permutations). RMSE were standardized (mean subtracted, divided by standard deviation) to allow a direct comparison of the two variables.