

Climate changes in Anatolia (Türkiye) across during the late Eocene and the Eocene-Oligocene Transition: successive warming and cooling, aridification, and implications for the westward dispersal of Asian terrestrial mammals

Paul Botté¹, Alexis Licht¹, Anne-Lise Jourdan¹, Leny Montheil¹, François Demory¹, Mustafa Kaya², Faruk Ocakoğlu³, Mehmet Serkan Akkiraz⁴, Deniz İbilioglu⁴, Pauline Coster⁵, Grégoire Métais⁶, Benjamin Raynaud⁶ & K. Christopher Beard^{7,8}

¹ Aix Marseille University, CNRS, IRD, INRAE, CEREGE, Aix-en-Provence, France

² Geological Institute, RWTH Aachen University, Aachen, Germany

³ Department of Geological Engineering, Eskişehir Osmangazi University, Eskişehir, Türkiye

⁴ Department of Geological Engineering, Kütahya Dumlupınar University, Kütahya, Türkiye

⁵ Réserve naturelle nationale géologique du Luberon, Unesco Global Geopark, Parc naturel régional du Luberon, Apt, France

⁶ Centre de recherche en paléontologie - Paris (CR2P), Muséum national d'Histoire naturelle, Sorbonne Université, CNRS, Paris, France

⁷ Biodiversity Institute, University of Kansas, Lawrence, Kansas, USA

⁸ Department of Ecology and Evolutionary Biology, University of Kansas, Lawrence, Kansas, USA

Correspondence to: Paul Botté (botte@cerege.fr)

Abstract. The Eocene–Oligocene Transition (EOT dated at ~34 Ma) represents one of the most significant climatic shifts of the Cenozoic, marking the transition from the last warmhouse state to a coolhouse state. This global cooling had major consequences for terrestrial ecosystems and was synchronous with the dispersal of numerous Asian mammalian clades towards western Europe. [However, the terrestrial expression of the EOT exhibits strong regional heterogeneity questioning its role in establishing dispersal corridors associated with floral and faunal turnovers. However, the terrestrial expression of the EOT exhibits strong regional heterogeneity. Consequently, its role in establishing dispersal corridors associated with the Grande Coupure floral and faunal turnover remains unclear.](#)

Here, we describe, date, and document the paleoenvironments of a continental sedimentary section from Balkanatolia, a biogeographic province [corresponding to the present-day Balkans \(NE Mediterranean region\) and Anatolia \(Türkiye\).](#) This [region](#) that most likely functioned as a critical stepping stone for the dispersal of Asian mammals toward western Europe. Our sedimentary record represents a fluvio-lacustrine [system exposed over a ~250m section located in Büyükteflek \(Çiçekdağı/Kırşehir area, Türkiye\).](#) system dated by magnetostratigraphy to the Priabonian and the lower Rupelian, including the Oi-1 glaciation (~33.65 Ma). Clumped isotopic analyses on pedogenic carbonates across our record show evidence for a Late Eocene Warming starting during the middle Priabonian (ca. 37 Ma), followed by a marked cooling event at the Early Eocene–Oligocene Glacial Maximum (EOGM). Stable isotopic data and sedimentary facies further indicate that this complete interval is associated with a long-term aridification trend, starting during the Late Eocene warming and culminating

at the EOT. Our results provide the first quantitative record of late Eocene warming on land, and our temperature estimates for the earliest Oligocene cooling are consistent with other Eurasian clumped-isotope records. These temperature shifts and associated aridification steps may have acted as contributing drivers of the late Eocene decline of Balkanatolian endemic taxa and likely facilitated the westward expansion of Asia-derived mammals ultimately resulting in the colonization of western Europe.

1 Introduction

The Eocene-Oligocene Transition (EOT) is one of the main climatic events of the Cenozoic. It marks the transition between the last Warmhouse state of the Eocene to a Coolhouse state in the Oligocene (Westerhold et al., 2020). The EOT is commonly interpreted as a phase of accelerated climatic and biotic change, linked to an abrupt global drop in temperatures (Coxall and Pearson, 2007). This drop is primarily explained by a decline in atmospheric CO₂ concentrations and changes in oceanic gateways, which, in combination with orbital forcing, acted as triggers for the first major glaciation of Antarctica (DeConto and Pollard, 2003). Within the deep sea record, the EOT is defined by a homogeneous and marked positive excursion of $\delta^{18}\text{O}$ from ~ 34.4 to 33.6 Ma (Katz et al., 2008; Hutchinson et al., 2021). The EOT is followed by the Early Oligocene Glacial maximum (EOGM), dated from 33.65 to ~ 33.16 Ma, corresponding to the peak of glacial expansion of the Oligocene associated with the major growth of the Antarctic ice sheet (Miller et al., 1991; Zachos et al., 1996; Hutchinson et al., 2021).

While the EOT and the EOGM are well documented in marine records, their impact on land remains poorly understood. This lack of understanding is mainly due to the heterogeneous nature of terrestrial responses: studies conducted so far show contrasting results regarding changes, or their absence, in seasonality, temperatures, and their timing of these changes, if they occur, or the absence of these changes (e.g., Pound and Saltzmann, 2017). Pollen compilations have shown a heterogeneous response of vegetation, especially in Eurasia where ecosystems show no, little, or a gradual shift to flora adapted to colder and more arid conditions (Pound and Saltzmann, 2017). Stable and clumped isotope analysis of continental carbonates (such as pedogenic and lacustrine carbonates) provides complementary insights into surface temperature and hydroclimate evolution, further highlighting and also underlines this heterogeneity. In North and South America, stable and/or clumped isotopic carbonate records from pedogenic carbonates indicate a cooling of temperatures and changes in seasonality, although with varying intensity and chronology (Kohn et al., 2015; Fan et al., 2018; Antoine et al., 2021; Meijer et al., 2025). In Eurasia, the response to the EOT is much more heterogeneous. In Western Europe, geochemical proxies for weathering intensity in paleosols and stable isotopes in pedogenic carbonates in the Ebro Basin (Spain) and on the Isle of Wight (UK) indicate no significant changes in the paleoenvironmental or climatic parameters such as the Mean Annual Temperature (MAT) or in seasonality in the Ebro Basin (Spain) and on the Isle of Wight (UK) (Sheldon et al., 2009; Sheldon et al., 2012). Other Alternative investigations on lacustrine carbonates, however, suggest a progressive shift toward more arid conditions, as observed in the Vistrenque Basin (France; Semmani et al., 2024), with marked aridity peaks predating the EOT, such as in the Alès Basin (France; Lettéron et al., 2017). In addition, clumped isotope analyses of gastropod shells

from the Hampshire Basin document a decrease in MAT by ~4-6°C across the EOT (Hren et al., 2013). Further east, in the central Asian Tajik Basin (Tajikistan), Wang et al. (2020) found evidence for increasing aridification beginning prior to the EOT in the Tajik Basin (Tajikistan), based on carbon isotopes in pedogenic carbonates. Studies conducted in the Qaidam Basin (north-central Tibet) provide further evidence of dramatic climatic changes across the late Eocene and the early Oligocene. Kent-Corson et al. (2009) documented a decrease in oxygen isotope values prior to the EOT, interpreted as reflecting a decline in temperatures. By contrast, Sun et al. (2020) reported a positive shift in oxygen isotopes from carbonate mudstones spanning the late Eocene to the middle Oligocene, suggesting a long-term trend toward enhanced aridity. In the nearby Xining Basin (northeastern Tibet), Page et al. (2019) documented a temperature decrease of ~20°C occurring both prior to the EOT and during the EOGM. This decline is interpreted as resulting from both a shift in the seasonality of carbonate formation and a decrease in surface temperatures, estimated at ~9°C. In the Lanzhou Basin (northeasternmost Tibet), Li et al. (2016) reported a shift toward more positive oxygen isotope values in pedogenic carbonates after 33 Ma, suggesting the onset of aridification following the EOT.

This heterogeneous Eurasian response to global climate change at the EOT and EOGM has been traditionally linked to regional changes in atmospheric circulation impacting monsoonal rainfall in eastern Asia (Page et al., 2019), and to Paratethys sea retreat impacting specific areas of central Asia and the Middle East (Toumoulin et al., 2022). It has also been linked to the growing role of topography, impacting atmospheric moisture sources and pathways dampening or enhancing global temperature and rainfall, especially in Europe around the nascent Alps where moisture sources significantly shifted. These topographic changes may also have influenced rainfall sources across northern and southwestern Europe between 35 and 31 Ma (Kocsis et al., 2014). As suggested by Kocsis et al. (2014), these topographic changes may have influenced rainfall water sources across northern and southwestern Europe between 35 and 31 Ma (Kocsis et al., 2014; Eronen et al., 2015), or to Paratethys sea retreat impacting specific areas of central Asia and the Middle East (Toumoulin et al., 2022).

The variable variability and regionality of the climatic and vegetation responses in Eurasia contrasts with the dramatic faunal changes that are witnessed at the continental scale. In Mongolia, a major faunal turnover, known as the Mongolian Remodelling, is interpreted as a response to the climatic shift toward more arid conditions (Meng & McKenna, 1998). But the most important faunal turnover spanning the late Priabonian and the earliest Oligocene is known as the “Grande Coupure” and corresponds to the extinction of 55% of European placental mammals (Weppe et al., 2023), coinciding with the arrival of Asia-derived mammals. Recent diversification modeling highlights the role of climate stress on western European taxa as a main driver for their decline, rather than competition with the arriving taxa (Weppe et al., 2023). Moreover, climate modeling suggests that the two proposed dispersal routes for dispersal (Fig. 1), via high latitudes in north-central Asia (Becker, 2009; Tissier et al., 2018; Mennecart et al., 2021) and via mid latitudes through Balkanatolia (Grandi and Bona, 2017; Licht et al., 2022a) were both significantly impacted by pCO₂ changes and changes by orbital and pCO₂ changes-orbital forcing during the latest Eocene. These factors affected biome stability along each route, limiting dispersal opportunities influenced the biome stability of each routes, dampening dispersal opportunities in one or the other (Tardif et al., 2021). Collectively, these changes Climate change through the EOT might have favored the dispersal of Asia-

a mis en forme : Indice

103 derived taxa towards western Europe during the EOT, though the ecosystem and climatic evolution is yet poorly documented
 104 in these two areas, where no terrestrial EOT record currently exists.
 105 This study provides the first magnetostratigraphically dated continental section from Balkanatolia that records a hydroclimatic
 106 response to the EOT, investigated. Our analyses combine through sedimentological facies description with stable and clumped
 107 isotope analyses of pedogenic carbonates. We compare our hydroclimatic records with other Eurasian records to better
 108 characterize the regional climatic evolution around the EOT, and assess how it could have impacted the faunal dispersal
 109 occurring in the region.
 110



111
 112 **Figure 1: Paleogeographic map of Balkanatolia and the Neotethysian realm at 34-33.5 Ma, modified from Montheil et al. (2025a).**
 113 **Arrows indicate the proposed dispersal routes of the “Grande Coupure” event. The studied location is shown with a red star.**
 114

115 2 Geological context

116 2.1 Balkanatolia

117 Balkanatolia is a semi-continuous strip of land extending from the Alpine region to the Lesser Caucasus (Licht et al., 2022a)
 118 (Figure 2). It comprised various terranes of Laurasian (Tisza, Dacia, Rhodope, Strandja Massif and the Pontides) and
 119 Gondwanan (Alcapa, Greater Adria, Anatolide-Tauride, South Armenia) affinities that progressively accreted throughout the
 120 Cretaceous and the early Paleogene due to the northeastward motion of Afro-Arabia and Greater Adria and consequent closure
 121 of the Anatolian Neotethys and the Sava Ocean (Van Hinsbergen et al., 2020; Mueller et al., 2022; see Figure 2B).
 122 During the Paleocene and Eocene, Balkanatolia consisted of large islands isolated from- Afro-Arabia, Western Europe and
 123 Eastern Asia by the Paratethys to the north and eastwest, the Neotethys to the south, and the Dobrujan, Valaisan and Carpathian
 124 seaways to the northwesteast (Barrier et al., 2018; Palcu and Krijgsman, 2022; Montheil et al., 2025a). This geographic
 125 isolation led to faunal endemism that prevailed until the Late Paleogene (Métais et al., 2018).

126 Tectonic activity and sea-level fluctuations created a dynamic landscape, often depicted as a mosaic of islands during the
 127 Paleocene and Early Eocene (Popov et al., 2004). These fluctuations were caused by alternating phases of extension, thrusting,
 128 and shortening, resulting in submergence and emergence of the terranes (Van Hinsbergen et al., 2020). Between the Lutetian
 129 and Bartonian, the retreat of shallow seaways separating Balkanatolian islands enhanced land connectivity (Licht et al., 2022a;
 130 Montheil et al., 2025a). As a result of renewed crustal shortening and sea level fall, terrestrial continuity with Laurasia was
 131 progressively re-established from the late Middle Eocene to Early Oligocene: first in the east via Anatolia and the Cimmerian
 132 terranes (Barrier et al., 2018), then in the west through continental sediment infill of the Luda Kamchiya Trough, connecting
 133 to the Moesian Platform (Doglioni et al., 1996). The first arrival of Asia-derived taxa in Balkanatolia is dated to the Bartonian
 134 (40-38 Ma) and associated with the decline of Balkanatolian endemic taxa (Licht et al., 2022a). This first wave of dispersal
 135 from Asia did not reach western Europe and Balkanatolia acted as a cul de sac (Métais et al., 2023). [The study of rodent](#)
 136 [assemblages of Balkanatolia shows that the Asian newcomers remained largely confined to Balkanatolia until the early](#)
 137 [Oligocene \(Maridet et al. 2025\). Only a few Asia-derived mammals colonized western Europe by passing through Balkanatolia](#)
 138 [during the late Priabonian \(mammalian biohorizon MP18-MP20; ca. 37-33.9 Ma\). These mammals include the anthracotheriid](#)
 139 [artiodactyl *Elomeryx*, possibly the gelocid artiodactyl *Phaneromeryx*, and the amphicyonid carnivoran *Cynodictis* \(Métais et](#)
 140 [al., 2023\). Faunal exchanges through Balkanatolia appear to have declined during the EOT and the EOGM in favor of the](#)
 141 [higher latitude routes, and resumed later during the Rupelian \(See Figure 1; Mennecart et al., 2021; Métais et al., 2023, Maridet](#)
 142 [et al. 2025\). The first dispersal event of Asia-derived mammals to western Europe and passing through Balkanatolia is dated](#)
 143 [to the late Priabonian \(mammalian biohorizon MP18-MP20; eg. 37-33.9 Ma\) and involved anthracotheriidae \(artiodactyla\),](#)
 144 [gelocidae \(perissodactyla\) and amphicyonidae \(Carnivora; Métais et al., 2023\). Faunal exchanges through Balkanatolia appear](#)
 145 [to have reduced during the EOT and the EOGM in favor of the higher latitude routes, and resumed later during the Rupelian](#)
 146 [\(See Figure 1; Mennecart et al., 2021; Métais et al., 2023\).](#)

148 2.2 The Çankiri Basin

149 Anatolia, as we know it today, is divided into the Pontides and the Anatolide-Taurides terranes, located in ~~its the~~ northern and
 150 southern parts, respectively. The separation of these two terranes is marked by the Izmir-Ankara-Erzincan Suture (IAES). In
 151 central Anatolia, the Kırşehir Massif (also known as the Central Anatolian Crystalline Complex, CACC) is sandwiched
 152 between the Pontide and Anatolide-Taurides terranes. The Kırşehir Massif has been interpreted either as a former volcanic arc
 153 located between the Pontides and the Anatolide-Taurides (~~Lefebvre et al., 2013~~) or as the subducted and exhumed northern tip
 154 of the Taurides (~~Lefebvre et al., 2013~~; Van Hinsbergen et al., 2016). The exhumation of the CACC was followed by renewed
 155 contraction related to a Paleogene collision of the massif with the Pontides (Gülyüz et al., 2013).

156 The Çankırı Basin, at the northern edge of the CACC, is a sedimentary basin that can be subdivided in ~~to~~ three main structural
 157 and stratigraphic units cropping out along its southern border, each reflecting distinct tectonic settings and depositional
 158 histories (Akgün et al., 2002; Kaymakcı et al., 2003) (Fig. 2C). The (1) *Çiçekdağı Belt*, represents the basin basement and
 159 belongs to the crystalline core of the CACC. This unit corresponds to Cretaceous volcanic arc units (~~Lefebvre et al., 2013~~;

a mis en forme : Police :Italique

a mis en forme : Police :Italique

a mis en forme : Police :Italique

160 Akgün et al., 2002) or is considered part of the Central Taurides ([Lefebvre et al., 2013](#); van Hinsbergen et al., 2020), which
161 was accreted onto the Pontides during ~~its the~~ collision with the Taurides. (~~Licht et al., 2017~~). It is overlain by the (2) *Çankırı*
162 *Basin Fill*, a Paleogene sequence filled with shallow marine to continental deposits (Akgün et al., 2002). This unit records
163 alternating episodes of marine and lacustrine transgression and regression [events](#) throughout the Paleogene. The lower part
164 consists of a Paleocene-Eocene regressive sequence, transitioning from flysch to molasse, and later overlain by nummulitic
165 limestones. From the Middle Eocene to the Oligocene, the sea progressively retreated, giving way to predominantly more
166 continental red clastics, intercalated with and eventually overlain by Oligocene evaporites (Kaymakçı et al., 2003). The third
167 unit, referred to as the (3) *cover series*, overlies the Cretaceous–Paleogene deposits and began to accumulate during the Early
168 Miocene with fluvio-lacustrine sediments alternating with evaporitic layers, continuing up to the Plio-Quaternary (Kaymakçı
169 et al., 2003; Akgün et al., 2002).

170 During most of the Eocene, the basin was bounded by the sea: the northern part of Çankırı corresponded to a deep-sea
171 environment, which was tectonically active due to its proximity to the IAES zone, while the southern part corresponded to a
172 partly shallow marine to continental deltaic environment, which was relatively tectonically quiescent ([Erdogan et al.,](#)
173 [1996reference here](#)). During the late Eocene (Priabonian), marine deposition had ceased, and the basin had lost its connection
174 to the sea, evolving into a large, closed [fluvio-lacustrine](#) system (Campbell et al., 2025).

175 Our study site is located in the Çiçekdağı area (Fig. 2C). In the area, the *Çiçekdağı Belt* crops out along the Çiçekdağı anticline,
176 which separates the Çiçekdağı Syncline and the Yerköy Syncline, located to the south and north of the Çiçekdağı area,
177 respectively, where Paleogene deposits of the *Çankırı Basin fill* are exposed (Gülyüz et al., 2013; Licht et al., 2022a).

178 The sedimentary sequence begins with the Yoncalı Formation, which contains conglomerates with volcanic clasts and lignites
179 (Lichte et al., 2022). It is overlain by the Late Lutetian–Early Priabonian marine limestones of the Kocaçay Formation, which
180 are rich in nummulites (Akgün et al., 2002; Gülyüz et al., 2013; Licht et al., 2022a). Above these beds, lie the continental
181 deposits of the Incik Formation, where fossils of ~~Asian mammals from the~~ Brontotheriidae and Hyracodontidae

(*Perissodactyls*) of Asian affinity families have recently been discovered (Licht et al., 2022a). These are then overlain by lacustrine deposits including marls, gypsum beds, and tuffs, known as the Sekili Member of the Incik Formation (Fig. 2C).




Figure 2: (A) Current structural overview of the Balkanatolian terranes. Terrane abbreviations: ST (Standja Massif), An-TD (Anatolide-Tauride), D (Dacia), SA (South Armenia), [LK\(Luda Kamchiya Trough\)](#), [MP \(Moesian Platform\)](#), Pd (Pontides), R (Rhodope), ACP (Alcapa), T (Tisza), GA (Greater Adria). (B) Middle Eocene structural overview of Balkanatolia (Based on Licht et al. (2022a) and updated by Montheil et al. (2025ba)). The star indicates the studied locality of Büyüktelek. (C) Simplified geological map of the Çiçekdağı Basin, with the synthetic stratigraphic log and the different basin units. [Maps in \(A\) and \(B\) are modified from Montheil et al., 2025b; geological map in \(C\) is modified from Gülyüz et al., \(2013\)](#) ~~Maps (A) and (B) maps modified from Montheil et al., 2025b; (C) geological map modified from Gülyüz et al., (2013)~~

2.3 The Büyükteflek section

195 Our study section is located within the Yerköy Syncline, near the village of Büyükt~~e~~eflek (See Fig. 2C; See Supplementary
196 Figures Fig. S1), and spans the Incik Formation and the Sekili Member. In this area, the Incik Formation is represented by red
197 beds, sandstones, conglomerates, and pedogenic carbonates. In the upper part of the section, the base of the Sekili Member is
198 marked by thick conglomeratic units and sandstone bodies containing reworked carbonate clasts and invertebrate fossils
199 derived from the upper Kocaçay Formation, where caliche horizons are commonly developed (Licht et al., 2022a). This section
200 has been previously studied by Gülyüz et al. (2013) and Licht et al. (2022a). The latter work provided improved constraints
201 on the basal part of the section, assigning a Priabonian age based on marine microfossils (ostracod and planktonic foraminifera)
202 assemblage within the uppermost beds of the Kocaçay Formation. The upper part of the section, covering the Sekili Member,
203 was dated using U–Pb analyses of interbedded tuff layers, yielding ages of 32.1 ± 1.1 Ma and 32.6 ± 1.1 Ma and was correlated
204 with the reversed polarity chron C12r (early Rupelian) based on magnetostratigraphic data. The lower part of the section,
205 extending from the Incik Formation to the base of the Sekili Member, was insufficiently sampled, preventing a precise
206 assignment by magnetostratigraphy within the Priabonian and the early Rupelian. Given the age constraints for the lower
207 (Priabonian) and the upper (Rupelian) parts, the section should cover the EOT.

209 3 Methods

210 3.1 Sedimentary logging

211 We measured and logged a 1975 m section spanning the Incik Formation and the Sekili Member at ~50 cm resolution, located
212 at the Büyütefle locality of Licht et al. (2022a) (See Supplementary Figures Fig. S1 for the overall picture of the Büyükteflek
213 section). Our ~~work consists of a~~ detailed stratigraphic log starting~~ed~~ at the level 14850m of ~~the one of~~ Licht et al. (2022a),
214 ending at the level 345m. ~~The aim was to improve the resolution of the log in order to . The sedimentary facies of these sections~~
215 ~~were characterized and classified the sedimentary facies of this section. These facies were classified~~ into 6 distinct lithofacies,
216 adapted ~~according to on~~ the classification of Miall (2013). ~~Photos Pictures~~ of the sedimentary facies are ~~provided displayed~~ in
217 Fig. 3, microphotographs and complementary ~~photos pictures~~ are available in the Supplementary figures Fig. S2.

218 3.2 X-Ray diffraction (XRD)

219 245 carbonate samples were collected all along the section and consist of nodules or blocks from continuous massive,
220 carbonated layers. All the samples were reduced to powder with a Dremel, and filtered with a 160 µm sieve. The mineralogy
221 of the carbonate samples was characterized using an X'Pert Pro MPD X-ray diffractometer (Malvern Panalytical), equipped
222 with a cobalt X-ray tube (wavelength $\lambda = 1.79$ Å), operating at 40 kV and 40 mA. Detailed procedures~~s~~ can be found in Couturier
223 et al. (2025). Sample powder was deposited onto low-background Silicium (Si) plates. The powders were homogenized using
224 drops of ethanol and allowed to dry. During analysis, samples were rotated at 15 rpm to improve measurement quality.
225 XRD scans were conducted over a 2θ range from 34° to 38°, with a step size of 0.026°. Each scan lasted 10 minutes, allowing
226 for the identification of carbonate mineral phases based on their primary diffraction peaks ~~(See XRD patterns in~~
227 ~~SupplementarySupplementary data 13).~~

a mis en forme : Police :10 pt

a mis en forme : Police :10 pt

a mis en forme : Police :10 pt

3.3 Magnetostratigraphy

For the magnetostratigraphic study, 110 rock samples were collected using a portable electric drill (core diameter of 1 inch [i.e. 2.54 cm](#)) and oriented with a magnetic compass. Samples were retrieved from red clastic layers ranging from clay to silt grain size and from pedogenic carbonates (from nodules to well-developed caliches) developed on silty sandstones or red clastics. We also collected [nine](#) oriented blocks in clay rocks ~~when they that when rocks~~ were too soft to be drilled.

Natural Remanent magnetization (NRM) and either its stepwise alternating field (AF) demagnetization with field increments of 5 mT from 0 to 70 mT or its stepwise thermal (TH) demagnetization with temperature increments of 20°C, 30°C, 40°C or 50°C from room temperature (~20°C) up to 660°C were measured with the Superconducting Rock Magnetometer SRM760R (2G enterprises) of CEREGE (Aix-Marseille University). Remanent signal from some clay samples was resistant to alternating field (AF) demagnetization and consequently required thermal demagnetization.

89 samples were demagnetized using ~~alternating field (AF)~~ AF demagnetization, and [16](#) samples were subjected to thermal demagnetization. ~~Among these, of which~~ 14 samples were analyzed using both methods. Data were plotted in orthogonal vector diagrams (Zijderveld, 1967) and stereographic projections and interpreted using the PuffinPlot software (Lurcock and Wilson, 2012). ~~The Characteristic Remanent Magnetization (ChRM) ChRM~~ directions were then calculated using the principal component analysis of Kirschvink (1980). ~~Samples exhibiting low magnetic intensity or lacked a coherent directional trend, likely reflecting local noise rather than a primary signal, were excluded from the dataset. Reversal tests and calculations of reversal angles were performed using both custom Python scripts and the Paleomagnetism.org platform. The dataset of this study has been supplemented with the paleomagnetic data from Licht et al. (2022a); (60 samples). The sample levels were adjusted to match our stratigraphic log using field-related descriptions.~~

~~Samples exhibiting low magnetic intensity or lacking a coherent directional trend, which is likely reflects reflecting local noise rather than a primary signal, were excluded from the dataset. The Component Remanent Magnetization (ChRM) for each sample was isolated using progressive thermal or alternating field (AF) demagnetization, performed in at least four steps covering the ranges 120–660°C or 5–70 mT (See demagnetization steps, e.g. samples BT 82, BT 48 2 and BT28 2 Fig. 5A,E,F; See Supplementary table 1 and Supplementary data 2 & 3), respectively, to ensure stable endpoint determination. Reliability was assessed based on the following criteria: the stability of demagnetization trajectories (ensuring the absence of erratic behaviour or secondary components) and, a maximum Maximum Angular Deviation (MAD) of 15° (Fisher, 1953; McFadden & McElhinny, 1990). Polarity consistency was further validated through reversal tests, reversal angle calculations, and Fisher direction statistics, performed using custom Python scripts and the Paleomagnetism.org platform and reversal tests and reversal angle calculations performed via custom Python scripts and the Paleomagnetism.org platform, to validate polarity consistency (Koyman et al., 2016; Fisher, 1953; Arason et al., 2010; Deenen et al., 2011; Helsop et al., 2023; McFadden et al., 1990; King, 1955; Tauxe, 2010; Tauxe et al., 2010; Tauxe & Watson, 1994; Tauxe et al., 2008; Zijderveld, 1967). These tests were applied to both our data and those the one from Licht et al. (2022a), and 28 samples were subsequently classified as outliers. Given the intrinsic uncertainties in paleomagnetic measurements, which can exceed ±30° in Virtual Geomagnetic~~

a mis en forme : Police :Italique

263 Pole (VGP) latitude for equatorial sites due to shallowing and low inclination resolution (Tauxe et al., 2010), VGPs with
264 latitudes between -30° and 30° were classified as uncertain, and no polarity interpretation was attempted for these cases.

266 3.4 Stable and clumped Isotopes

267 234 carbonate sample powders identified as pure calcite from XRD, and one composed of approximately equal proportions of
268 calcite and dolomite, were selected for stable isotope analysis. The samples were analysed on a Thermo 253+ mass
269 spectrometer attached to a Thermo Kiel IV device, run under the Qtegra™ software and was in Dual Inlet mode at CEREGE.
270 Sample powder corresponding to around 80-120 µg of pure carbonate was loaded into glass vials and placed on a carousel
271 along with international standards (IAEA-603, NBS18) and laboratory standards (BDH and VIA-1, which are made of pure
272 calcite) in the Kiel oven held at 70°C. After each vial is in turn put under vacuum, the powder reacts with 2 to 3 drops of
273 105% phosphoric acid at 70°C to produce CO₂ for analysis. The released CO₂ is then trapped and cleaned via a series of cryo-
274 traps, and finally sent to the Thermo 253+ mass spectrometer for measurements of the oxygen and carbon isotopes values.
275 Isotopic values are reported relative to IAEA-603 international standards in the V-PDB (Vienna Pee Dee Belemnite) reference
276 frame materials.

277 14 of these samples were selected for clumped isotope analysis based on their carbonate content and their distribution along
278 our stratigraphic log. They and were treated overnight with 3-8% NaOCl, rinsed and centrifuged 4 times and dried with a
279 freeze-drier, to remove nitrate contamination (Fiebig et al., 2024). The samples were then analysed at CEREGE on the same
280 previously described system. Details about the clumped isotope procedure at CEREGE are given in Supplementary Text S1.
281 Briefly, primary standards (ETH-1 to -4), secondary standards and unknown sample powders were loaded into the glass vials
282 and placed on the carousel of the Kiel oven still held at 70°C. The procedure is similar to the one described above for stable
283 isotope measurement with the exception of an additional step of cryotrapping involving a trap filled with a Porapak molecular
284 sieve, held at -20°C by a Peltier system. The resulting purified CO₂ gas is transferred to the 253+, where extra Faraday cups
285 will allow the measurement of clumped isotopes (m/z 47, 48, 49) on top of carbon and oxygen isotopes (m/z 44, 45, 46) and
286 pressure baseline corrections (m/z 47.5 and 48.5).

287 Baseline-corrected δ data are screened with an in-house Matlab script for statistical outliers and are then translated into δ¹⁸O,
288 δ¹³C and Δ₄₇ projected to the InterCarb-Carbon Dioxide Equilibrium Scale (I-CDES90; Bernasconi et al., 2021) using
289 D47crunch (Daëron, 2021), pooling over data from all sessions of the dataset. In total, we run 10 to 20 replicates per unknown
290 sample spread over 28 sessions, with an inter-session repeatability of 30 ppm in Δ₄₇ for all replicates (unknown and standards).
291 Paleotemperatures are calculated using the unified Δ₄₇-T calibration of Anderson et al. (2021). Uncertainties around
292 temperature estimates are provided with and without propagating intersession-related uncertainty (difference of +/-1 to 2 °C
293 at 2s).

294 Note that to test the reproducibility of our results and the impact of the NaOCl pre-treatment, three samples were divided into
295 two aliquots: one with pre-treatment, and one without. Two out of three samples show no statistical difference with or
296 without pre-treatment (difference < 2s); For a third sample, pre-treated and non-treated aliquots lie within 3s of each other. We

a mis en forme : Indice

a mis en forme : Exposant

a mis en forme : Exposant

a mis en forme : Indice

a mis en forme : Indice

a mis en forme : Indice

297 found no statistical difference between pre-treated and non-treated samples. While some samples show statistically significant
298 differences between pre-treated and non-treated values, there is no systematic offset across the dataset (See Supplementary
299 table 2)

300 Data presented in the manuscript are only from pre-treated samples. Raw data for unknown samples and standards are provided
301 together with interpreted data, to allow possible future reprocessing of Δ_{47} values and paleo-temperature estimates, in
302 Supplementary table 24.

303
304 **4 Results**
305 **4.1 Sedimentology and XRD analysis of the Büyükteflekk section**

306 The lower part of the studied section (from ~100 to 263 m) is characterized by three distinct sedimentary facies (Table 1). The
307 first facies (Fmp; see Fig. 3B,C,D) consists of clayey to silty red beds, generally massive in structure, and occasionally
308 exhibiting discontinuous planar laminations. These deposits may contain root traces and reduced mottling due to reducing
309 conditions. Above level 240 m, they display well more developed pedogenic features, including small carbonate nodules to
310 well-developed caliche horizons. X-ray diffraction (XRD) analysis indicates that all carbonate samples from these beds are
311 composed exclusively of calcite. The second facies (St; see Fig. 3G,H) is composed of channelized, trough cross-bedded
312 sandstones, with grain-size ranging from very fine to medium sand. Some beds contain thin layers of gravel or pebbles. The
313 fossiliferous horizon described by Licht et al. (2022a), containing Brontotheriidae (*Embolotherium aff. andrewsi*) and
314 Hyracodontidae (*Prohyracodon* sp.), occurs at approximately the 215-meter level within these facies. The third facies (Sm;
315 see Fig. 3D) consists of structureless greenish medium to coarse sandstones, containing isolated clasts ranging in size from
316 gravel to approximately 2 cm. The three facies Fmp-St-Sm are grouped under Facies Association FA1 (Table 2).

317 The level at 263 meters is marked by a field-identifiable surface, corresponding to a channelized erosional surface developed
318 above a well-developed caliche. This hiatus was interpreted by Licht et al. (2022a) as the boundary between the Incik
319 Formation and the Sekili Member (Hiatus 1 on Fig. 4). From 263 m to 292 m, the facies consist of channelized sandstones (St)
320 and pedogenized clay to silty red beds (Fmp). This interval marks the appearance of three new facies. The first one (Smp; see
321 Fig. 3A,E,F) consists of reddish to pinkish, massive, very fine- to fine-grained sands. These beds are strongly pedogenized
322 and feature carbonate nodules, well-developed caliche horizons or diffuse carbonate. The second facies (Gmm; see Fig. 3G),
323 which is coarser, is composed of massive, matrix-supported conglomerates with a sandy matrix, containing clasts ranging from
324 1 mm to 5–10 cm and exhibiting weak normal grading. Some of these conglomerates include reworked *Nummulites*, probably
325 derived from underlying marine deposits. The last facies (Gmc; see Fig. 3B,E,G) consists of clast-supported conglomerates
326 with a sandy matrix, containing imbricated clasts ranging from 1 mm to 5–20 cm, and may also contain reworked *Nummulites*.
327 These three new facies, together with the channelized sandstone Smp-St-Gmm-Gmc are grouped under Facies Association
328 FA2 (Table 2). X-ray diffraction analysis indicates that the nodules found at these depths contain only calcite.

329 At 292 meters, a second unconformity is observed, marked by the development of a 2–4 m thick caliche (Hiatus 2; see Fig. 4).
330 This paleosol is the thickest of the section and forms a topographic feature that can be tracked over hundreds of meters (Fig.

a mis en forme : Indice

a mis en forme : Police :Italique

a mis en forme : Police :Italique

331 3A). Five facies are present from this unconformity up to the top of the logged section in this study (level 355 m). These
332 include the same channelized sandstones (St), along with pedogenized red beds assigned to the Fmp facies. The Smp facies is
333 also identified, which is particularly abundant in this part of the section. X-ray diffraction analysis shows that carbonate in
334 these beds may contain either calcite (from 292 m to 325 m) or dolomite (from 302 m to 355 m). In addition, the two
335 conglomerate facies, Gmm and Gmc, are present within this part of the section. This interval of the log includes both facies
336 associations FA1 and FA2.
337
338

Facies	Grain Size	Bed Thickness	Description	Pedogenic Features	Interpretation	References for the interpretation
Fmp	Clay to silt	10 cm to 1 m	Massive red beds, occasionally displaying discontinuous planar laminations.	Root traces, reduced mottles, carbonate nodules to well-developed caliche horizons.	Overbank deposits in floodplain or delta plain environments.	Bhattacharya (2010); Miall (2010); Leeder (2016)
St	Very fine to medium sand	10 to 50 cm	Trough cross-bedded sandstones organized in cosets. Some layers contain thin beds of gravels or pebbles.	None	Sand bars in river channels, delta distributary channels or at delta mouth bars	Bhattacharya (2010); Miall (2010)
Sm	Medium sand to gravel	10 to 50 cm	Structureless greenish sandstones containing gravel-sized clasts up to 2 cm. sometimes carbonated.	None	Sediment gravity-flow deposits.	Miall (2010)
Smp	Very fine to fine sand	5 to 20 cm (sets)	Reddish to pinkish massive sandstones, which may either be strongly pedogenized with well-developed caliche horizons predominantly composed of calcite) or exhibit diffuse carbonate cementation	Pedogenic carbonate nodules to well-developed caliche horizons.	Sand bars in river channels, delta distributary channels, delta mouth bars, or distal fan gravity flow deposits that have undergone later emersion and pedogenesis	Leeder (2016); Miall (2010)

			(predominantly composed of dolomite).			
Gmm	Conglomerate (matrix-supported)	2 to 5 m (to confirm)	Massive, matrix-supported conglomerates with a sandy matrix, occasionally displaying cross-bedding. Clasts range from 1 mm to 5–10 cm and exhibit weak normal grading. These deposits occasionally contain reworked Nummulites derived from underlying marine beds	Occasional well-developed caliches	High-strength cohesive debris-flow deposits.	Miall (2010)
Gmc	Conglomerate (clast-supported)	2 to 5 m (to confirm)	Clast-supported conglomerates with a sandy matrix, containing imbricated clasts ranging from 1 mm to 5–20 cm. May also contain reworked <i>Nummulites</i> .	None	Pseudoplastic debris flows (inertial bedload, turbulent flow).	Miall (2010)

Table 1: Sedimentary facies of the Büyükteflek section

Facies Association Name	Facies	Interpretation	Sources/References
FA1	Fmp-St-Sm	Floodplain/alluvial plain	Miall (2010); Leeder(2016);
FA2	Smp-St-Gmm-Gmc	Lacustrine fan-delta	Miall (2010); Renaut and Gierlowski-Kordesch (2010)


Table 2 : Facies associations of the Büyükteflek section
Figure 3 : Examples of sedimentary facies described in Table 1. (A) Well-developed caliche (facies Smp); Hiatus 2, marked by red dotted lines, corresponds to the stratigraphic level at 292 m; hammer for scale. (B) Well-developed caliche (facies Fmp), eroded at the top by channelized conglomerates (facies Gmc); this surface corresponds to Hiatus 1 at 263 m; hammer for scale. (C) Massive clayish red beds (facies Fmp); hammer for scale. (D) Massive greenish sandstones containing gravel-sized clasts up to 2 cm (facies Sm), overlying red beds (facies Fmp); knife for scale. (E) Clast-supported conglomerates overlying caliche developed in sandstones (facies Smp); hammer for scale. (F) Massive sandstone with diffuse carbonate development (facies Smp); knife for scale. (G)

a mis en forme : Gauche, Interligne : simple, Bordure : Haut: (Pas de bordure), Bas: (Pas de bordure), Gauche: (Pas de bordure), Droite: (Pas de bordure), Entre : (Pas de bordure)
a mis en forme : Police :10 pt, Couleur de police : Noir



Alternation of matrix-supported (facies Gmm) clast-supported conglomerates (facies Gmc) and the channelized sandstones (facies St); Leny Montheil (180 cm) for scale. (H) Trough cross-bedded sandstones (facies St); hammer for scale.

Figure 4: Stratigraphic log of the Büyükteflek section (See Supplementary Text S2 for the exact location of hiatuses and tuffs) (S).



Colour coding illustrates the different facies and facies associations. The VGP latitude diagram shows the paleomagnetic data and the corresponding magnetozones (R1, N1, N2, R2, N3, N4).

4.2 Magnetostratigraphy of the Büyüktefle section

The magnetostratigraphic data presented here complement the work of Licht et al. (2022a) on the same section. Magnetic intensity measurements/Natural Remanent Magnetization (NRM) measurements revealed that clay samples generally exhibited weaker intensities, ranging from 6.9×10^{-4} to 1.5×10^{-1} A/m, while coarser samples, from silt to fine sand, displayed higher intensities, ranging from 1.5×10^{-2} to 8.9×10^{-1} A/m. Some samples showed a low-coercivity secondary evidence of a viscous component in both fine and coarse grain sizes (e.g., BT82, Fig. 5A). For the silt to fine sand fraction, demagnetization typically shows a stable decay toward the origin (e.g., BT23 Fig. 5B). Similarly, several clay samples also display a consistent decay toward the origin, though often accompanied by a higher degree of noise (e.g., BT71, BT80, Fig. 5C,D). Thermal demagnetization of selected clay samples was carried out between 120°C and 500°C (e.g., BT48_2, BT28_2, Fig. 5E, F; See Supplementary table 1 & 6 and Supplementary data 1 & 2).

These thermal demagnetization (performed from 120 to 660°C) results suggest the presence of both magnetite and hematite, with sometimes the unblocking temperature ranges supporting their coexistence. Zijderveld diagrams for these samples typically show a single-component decay, indicating the presence of a stable Characteristic Remanent Magnetization (ChRM) (see representative example in Fig. 5).

More samples recorded a primary-normal geomagnetic polarity than a reverse geomagnetic polarity, consistent with the investigated time interval, which is characterized by a predominance of long-lasting normal polarity chrons. This result is in good agreement with the investigated time interval characterized by a predominance of long-lasting normal polarity chrons. To complete the geomagnetic polarity record, we added the magnetostratigraphic dataset from Licht et al. (2022a), which covers the upper part of the section and includes a higher proportion of reverse magnetized samples attributed to Chron C12r.

Reversal tests and calculations of reversal angles were performed using both custom Python scripts and the Paleomagnetism.org platform. The reversal test yielded positive results.

Regarding ChRM directional statistics (See Supplementary Figures Fig. S5), the inclination derived from geographic coordinates (53.2°) is consistent with the site's latitude of the site, whereas it drops to 26.5° after tectonic correction, which is inconsistent with the expected value for the site's location. Notably, the α_{95} value of inclination data shows minimal change (from 3.2° to 4.7°) between when switching from geographic and tectonically corrected coordinates, indicating not significant tectonic influence, suggesting. The inclination discrepancy, most likely attributable to the shallowing effect commonly observed in laminated lake sediments (e.g. Philippe et al., 2023), thus reflects a systematic bias rather than increased data scatter. This suggests that the inclination discrepancy is not driven by increased data scatter but rather reflects a systematic bias. The discrepancy between the expected and calculated inclinations is most likely due to a shallowing effect often observed in laminated lake sediments (e.g. Philippe et al., 2023).

a mis en forme : Justifié


388 The observed angle between the mean normal and reversed (corrected) directions is $\gamma = 21.0^\circ$, whereas the critical angle at the
389 95% confidence level is $\gamma_c = 13^\circ$, resulting in a formal failure of the reversal test. The datasets are strongly unbalanced ($N=85$
390 vs. $N = 21$), and the reversed population shows greater dispersion ($k = 7.51$; $\alpha_{95} = 12.4^\circ$) than the normal population ($k =$
391 17.49 ; $\alpha_{95} = 3.8^\circ$; see Supplementary Figures Fig. S5). This statistical failure likely reflects the smaller sample size and higher
392 scatter of the reversed dataset rather than a true departure from geomagnetic antipodality, and the angular difference between
393 the means remains moderate. Given that this study is based on continental sedimentary rocks, such angular differences between
394 mean directions remain within acceptable limits (Tauxe et al., 2010). Fisher statistics indicate a mean inclination of 28.7° for
395 the normal samples and 34.7° for the reversed (corrected) samples (See Supplementary Figures Fig. S5). These relatively low
396 and comparable values further support the presence of inclination shallowing. Consequently, the tilting observed in the section
397 indicates that the magnetization was acquired prior to tilting rather than post-tilting. ~~While normal polarity samples show~~
398 ~~consistent and well-clustered directions, reverse polarity samples are more scattered, likely due to the limited number of data~~
399 ~~points. Nonetheless, the antipodal relationship between the normal and reversed directions supports the interpretation of a~~
400 ~~primary ChRM (See Supplementary Figures Fig. S5).~~
401 ~~We compared geographic coordinates with tectonically corrected coordinates. When switching between these two reference~~
402 ~~frames, the~~ Regarding the statistics of ChRM directions and focusing on inclination only (See Supplementary Figures Fig. S5)
403 ~~(The α_{95} value shows no significant change (from 3.24 to 4.733.42 to 4.39) when switching between geographic coordinates~~
404 ~~and tectonically corrected coordinates. The inclination derived from geographic coordinates (53.277°) is consistent with the~~
405 ~~site's latitude of the site. However, after tectonic correction, the inclination drops to 26.5457° after tectonic correction, which~~
406 ~~is inconsistent with the expected value for the site's location. This discrepancy is most likely due to a shallowing effect.~~
407 ~~To further investigate, we focused on inclination only statistics. Then, no significant changes in α_{95} were observed (from 3.24~~
408 ~~to 4.73), while the inclination shifted from 53.2° (geographic) to 26.54° (tectonic corrected). These results alone do not allow~~
409 ~~us to discriminate between primary and potential secondary magnetization. The primary character of the magnetization is~~
410 ~~further supported by the presence of several magnetic reversals along the section.~~ Furthermore, samples containing both
411 magnetite and hematite exhibit a single-component magnetization, suggesting that hematite recorded the ambient magnetic
412 field during or shortly after deposition, rather than during later mineralogical alteration, ~~and thus supporting a primary nature~~
413 ~~for the magnetization.~~ ~~This supports the interpretation of the ChRM as a relatively primary signal.~~
414 ~~Five-Four~~ distinct magnetic chrons have been identified within the Sekili Member interval of the studied section. These chrons
415 are defined based on the presence of at least ~~three-five~~ consecutive, reliable samples exhibiting consistent magnetic polarity.
416 In contrast, the lower part of the section, corresponding to the Incik Formation, displays less well-defined magnetic chrons,
417 ~~possibly reflecting either likely due to~~ lower sampling resolution ~~and/or enhanced diagenetic~~ ~~enhanced~~ ~~increased diagenetic~~
418 overprinting. However, the dominance of normal polarity intervals in this portion of the section led us to interpret it as
419 representing a normal chron.

a mis en forme : Police :10 pt

a mis en forme : Justifié, Espace Avant : 0 pt, Après : 0 pt, Interligne : 1.5 ligne

a mis en forme : Police : (Par défaut) Arial, 11.5 pt, Couleur de police : Violet

422
423



424

425 **Figure 5: Representative intensity plots and Zijderveld diagrams obtained after Alternating Field (AF) and Thermal**
426 **demagnetization.**
427

428 **4.3 Stable and Clumped isotopes**

429 The stable isotope composition of calcite samples ranges from -6.5‰ to $-4.95 \pm 0.1\text{‰}$ for $\delta^{18}\text{O}$ (on average: -5.7‰ V-PDB) and
430 from -7.8‰ to -5.2‰ for $\delta^{13}\text{C}$ (‰ V-PDB). Calcite $\delta^{18}\text{O}$ values do not exhibit any clear stratigraphic trend throughout the
431 section. In contrast, $\delta^{13}\text{C}$ values for calcite are slightly more variable. Below Hiatus 1, $\delta^{13}\text{C}$ values average $-6.8 \pm 0.3\text{‰}$.
432 Between Hiatus 1 and Hiatus 2, these values increase slightly, with a mean of $-6.4 \pm 0.6\text{‰}$. Above Hiatus 2, $\delta^{13}\text{C}$ values
433 display a decreasing trend, ranging from -5.99‰ to -7.8‰ with an average value of $-6.99\text{‰} \pm 0.8\text{‰}$. Among the samples
434 above Hiatus 2, one contains both calcite and dolomite in approximately equal proportions. Given its comparable clumped
435 isotopic values to those of other samples, it was retained in the results and discussion (See Supplementary Table 3). Clumped
436 isotope-derived temperatures (Δ_{47}) from pedogenic calcite carbonates fall within the range of Earth's surface temperatures
437 ($<40^\circ\text{C}$). The Δ_{47} results can be divided into three distinct groups. The first group (all samples below hiatus 1, but the very last
438 sample; 4 samples in total) shows relatively stable values, with a mean temperature of $24.4^\circ\text{C} \pm 0.7$ (2s). A significant increase
439 in surface temperature is observed in the second group, corresponding to the last sample below hiatus 1 and all the samples
440 situated between Hiatus 1 and Hiatus 2 (5 samples in total). This interval yields a higher and relatively consistent mean
441 temperature of $34^\circ\text{C} \pm 3.5$ (2s), representing a $\sim 9^\circ\text{C}$ increase compared to the interval below Hiatus 1. Then the third group
442 shows a decreasing trend in temperatures. The third group (5 samples), located above Hiatus 2, displays decreasing
443 temperatures of $\sim 7\text{--}8^\circ\text{C}$. This group yields a mean Δ_{47} -derived temperature of $27.4^\circ\text{C} \pm 3.6$ (2s), with respectively the highest
444 to the lowest values from $31.9 \pm 5^\circ\text{C}$ to $21.6^\circ\text{C} \pm 4.2^\circ\text{C}$. Clumped isotopic results from the sample with a mixture of dolomite
445 and calcite (See supplementary table 24 sample BTCARB16) sample yield similar Δ_{47} values than for pure calcite samples of
446 the same interval, suggesting that the CO_2 extracted during the acid reaction at 70°C is prominently calcitic (e.g. Li et al.,
447 2023). The exclusion of the sample from our dataset does not alter the average temperature of the interval (e.g. 27.5 ± 4.1 2s
448 without the sample instead of 27.4 ± 3.6 2s-with it). Student's t -tests were performed between the first and second groups,
449 and between the second and third groups. The three groups show statistically significant differences. The comparison between
450 Group 1 and Group 2 yielded a p -value of 0.005 and an effect size (Cohen's d) of $|d| = 3.33$, indicating a strong statistical
451 difference between these two groups. Similarly, the comparison between Group 2 and Group 3 produced a t -test p -value of
452 0.030 and an effect size of $|d| = 1.668$, also suggesting a substantial difference between the (See Supplementary table 4).
453 Based on clumped and stable isotope data, $\delta^{18}\text{O}_{\text{water}}$ values of soil water ($\delta^{18}\text{O}_{\text{water}}$) values (V-SMOW) have been reconstructed
454 according to Kim and O'Neil (1997) recalculation (See Fig. 6). These values can be divided into three groups, consistent with
455 the grouping defined from clumped isotope temperatures. Group 1 displays a mean $\delta^{18}\text{O}_{\text{water}}$ value of -3.6‰ (V-SMOW). A
456 marked increase is observed at the final sample just before Hiatus 1, with Group 2 showing a mean value of -1.9‰ . This is
457 followed by a slight decrease in Group 3, which exhibits a mean $\delta^{18}\text{O}_{\text{water}}$ value of -2.7‰ (See Supplementary table 45).
458
459

a mis en forme : Indice

a mis en forme : Indice

a mis en forme : Indice

a mis en forme : Indice

482 magnetozone N34, therefore postdate this age. This interval contains mostly normal polarity samples, with only a few isolated
483 reversed ones, suggesting that magnetic reversals may not have been preserved due to erosion, non-deposition, or limited
484 sampling resolution. Several correlation scenarios were considered using these constraints (See Supplementary Figures Fig.
485 S2, Fig. S3 and Supplementary Table 5). Before correlating our magnetozones, we have first considered all the deposits below
486 the second hiatus the section below R1 as a single, continuous normal magnetozone N1bis (thus excluding the reverse samples
487 defining R2). If correlated to chron C13n (as N1), assuming it corresponds to chron C13n. (This approach yielded a minimum
488 sedimentation rate of 58.6 cm/ka for N1bis, which is unrealistically high compared to the upper part of the section and thus
489 geologically implausible. Consequently, these rates do not support an Oligocene age for the underlying section and ensure that
490 the section below hiatus 2 is partly PriabonianEocene. , and We then retained the interpretation that yields the most stable
491 sediment accumulation rates through the section (See Supplementary Table 52). Just below Hiatus 2, Below hiatus 2, the
492 reverse normal polarity magnetozone RN2 has to be correlated to one of the five reverse chron between C13r and the base of
493 the Priabonian (C16r to C13r). The correlation for R2 that yields the most consistent accumulation rates when compared with
494 the rates of the higher part of the section (4.5 cm/kyr for R1) is is correlated with Chron C16n.1r (35.77–35.72 Ma;
495 accumulation rate: 5.4 cm/kyr for R2), and, the reversed magnetozone R2 with Chron C16n.1r (35.77–35.72 Ma), and the
496 underlying normal magnetozone N3 with Chron C16n.2n (36.35–35.77 Ma). We acknowledge that other hypotheses of
497 correlation are possible (See Supplementary Figures Fig. S2 and S3 for a second and third hypothesis), but all other correlations
498 yield much lower accumulation rates (e.g. < 1.8 cm/kyr for R2). The basal portion magnetozone N3 below the reverse
499 magnetozone R2 of the section could not be confidently correlated with a specific chron due to the scarcity of reversed-polarity
500 samples, suggesting that one or more magnetic reversals may have gone unrecorded or unsampled in this lower interval (See
501 Fig. 7). Consequently, The N34 interval and the marine beds are is should fall within the interval between correlated with the
502 mid-part of Chron C17n.2n base of Priabonian and, as the marine beds are Priabonian in age, to the base of Chron C16n.1r
503 Chron C17n.1n (e.g. within the ~37.71–35.7736–57 Ma window).
504 This magnetostratigraphic framework allows for temporal subdivision of the Büyüktefleken section. The basal unit, below the
505 first unconformity, corresponds to the early to middle Priabonian. The layer at 215 m, which yielded fossils of Brontotheriidae
506 (*Embolotherium* aff. *andrewsi*) and Hyracodontidae (*Prohyracodon* sp.), belong to magnetozone N3 is correlated with the
507 interval ranging from the mid-part of Chron C17n.2n base of the Priabonian to the base of the Chron C16n.1n. (This
508 magnetostratigraphic position constrains the age of the fossil-bearing horizon to the approximately 37.71 to 35.6.757 Ma
509 window (the Priabonian part of C17n). The interval between the two unconformities is constrained by the correlation of
510 magnetozone R2 to chron C16n.1r; it likely spans spans a time frame around ~36 Ma and corresponds to Chrons C16n.2n,
511 C16n.1r, and parts of the early-middle Priabonian (see Fig. 7) C16n.1n. Based on the average accumulation rate across the
512 section, this interval is expected to last around 675 kyr. The upper part of the section, above the second unconformity, is
513 attributed to chrons C13n and C12r. Chron C13n coincides with the EarlyEocene Oligocene Glacial Maximum (EOGM;
514 Hutchinson et al., 2021), while the uppermost part corresponds to the later, warmer times of the
515 RupelianOligocene.

a mis en forme : Police :(Par défaut) Times New Roman, 10 pt, Couleur de police : Automatique

a mis en forme : Police :(Par défaut) Times New Roman, 10 pt, Couleur de police : Automatique

a mis en forme : Police :(Par défaut) Times New Roman, 10 pt, Couleur de police : Automatique

a mis en forme : Police :(Par défaut) Times New Roman, 10 pt, Couleur de police : Automatique

a mis en forme : Police :(Par défaut) Times New Roman, 10 pt, Couleur de police : Automatique

a mis en forme : Police :(Par défaut) Times New Roman, 10 pt, Couleur de police : Automatique

a mis en forme : Police :(Par défaut) Arial, 11.5 pt, Couleur de police : Violet




Figure 7: Stratigraphic log of the Büyükteflek section showing the magnetostratigraphic correlation with associated accumulation rate (R1;R2) and minimum accumulation rates (N1;N2;N3;N4), calibrated with the Paleogene Geomagnetic Polarity Time Scale (Gradstein et al., 2020).

a mis en forme : Police :Gras, Couleur de police : Noir

5.2 Evolution of the depositional system and origin of the unconformities

Previous work by Gülyüz et al. (2013); suggested that episodes of folding, thrusting, and uplift along the Çiçekdağı anticline, located directly to the south of our section, began between ~38 and 35 Ma. They also identified a reversal of the direction of paleocurrents from predominantly southward to northward flow direction within the Sekili Member, suggesting a local reorganization of paleodrainages. It is thus very likely that the two unconformities recorded in our section are at least partly related to on-going tectonic uplift and deformation of the southern margin of the basin. Our magnetostratigraphic correlation does not allow us to determine indicates the exact duration of that the first unconformity (Hiatus 1), which spans part of the early-mid Priabonian, C16n.2n, the entire C16r, and a portion of the C17n.1n, most likely beginning in the upper part of C17n.1n. Otherwise, the accumulation rates for magnetozone N4 would be unrealistically high compared to those of the surrounding intervals. This hiatus corresponds to a minimum time interval extending from just before the onset of C16r (~36.6 Ma) to 36.4 Ma. The second unconformity (Hiatus 2) spans part of the early C13n, the whole of C13r, C15n, C15r, and the upper part of the C16n.1n. This corresponds to a minimum interval ranging from 35.6 to 33.8 Ma, including the latest Priabonian and most of the EOT (34.44 to 33.65 Myr, sensu Hutchinson et al., 2021).

The first unconformity is associated with a marked shift in prominent sedimentary facies and is associated with a change of depositional system. Deposits below Hiatus 1 only display one facies association (FA1), which is interpreted as reflecting fine-grained floodplain or alluvial plain deposits (Table 1.; Miall 2010; Leeder 2016). The deposits above Hiatus 1 are dominated by coarser, more energetic, and more pedogenized/carbonated facies grouped in Facies Association 2 (FA2), which is interpreted as a streamflow-dominated alluvial fan/delta with subaqueous portions extending into the lake. This more subaqueous or distal part of the fan-delta is supported by the presence of the facies Smp (See Fig. 4 and Tables 1, 2; Miall 2010; Renault and Gierlowski-Kordesch 2010).

According to Gülyüz et al. (2013), the local reversal in paleocurrents around that time corresponds to a shift from southward to northward and is interpreted as marking the uplift of the Cicekdagi anticline directly south of our section. This uplift is corroborated by the reworking of nummulitic limestones, starting with Hiatus 1. We thus suggest that the change of sedimentary facies relates to a change in the fluvial network and the regional hydrosystem. This reorganization is consistent with the uplift of the Çiçekdağı anticline and the associated re-routing of regional drainage systems (Gülyüz et al., 2013; Licht et al., 2022a). During the early parts of the Priabonian, the area was the locus of a southward-oriented, medium-distance drainage network exiting to the southern edge of the Cankiri basin, where the youngest marine deposits are found (Licht et al., 2022a). Hiatus 1 marks the uplift of the southern margin of the basin, the disconnection of this river network and the setup of a local, proximal-sourced northward-oriented drainage exiting to a newly formed lacustrine system occupying the Cankiri Basin.

Pedogenesis and carbonate cementation are progressively observed into coarser sediments through the section. At the base, pedogenic features of the Fmp facies (Table 1) are developed within clayey deposits in a floodplain environment. In the upper

part of the succession, beginning above Hiatus 1, pedogenesis and carbonate cementation are instead mainly associated with coarser gravity-flow deposits, represented by sandstones of the Smp facies (Table 1). These sandstones accumulated in the distal portions of a fan-delta system, near the lake margin or even within its subaqueous domain. Owing to this transitional position, such deposits were highly sensitive to lake-level fluctuations. The occurrence of well-developed caliches or carbonate-cemented horizons within these sandstones likely indicates phases of lake retreat, during which pedogenesis could develop on exposed surfaces.

The second unconformity is not associated with any major change of sedimentary facies but the appearance of dolomite. From the Hiatus 2 onward, almost all of our carbonate-rich sandy facies (Smp) consist predominantly of dolomite. When not related to diagenetic processes, dolomite can precipitate in hypersaline environments such as lagoons or saline lakes (García-Ruiz et al., 2023) depending on local salinity and pH fluctuations (Kim et al., 2023). We thus interpret the appearance of dolomite as reflecting an increase of lake salinity and a shift toward higher evaporation relative to precipitation, where water loss concentrates dissolved salts faster than they are diluted by freshwater inputs (Guo et al., 2023). This interpretation is further supported by the occurrence of Oligocene evaporites in other parts of the Çankırı Basin (Kaymakçı et al., 2003; Karadenizli, 2011). Together, these observations provide multiple independent sedimentological indicators of lake-level fluctuations under increasingly arid conditions.

The synchronicity between the second unconformity and the latest Priabonian and the EOT, a period of documented increase of aridity in many places in Eurasia (Abels et al., 2011; Page et al., 2019; Lettéron et al., 2022; Li et al., 2016; Sun et al., 2020; Semmani et al., 2024), also supports this interpretation. The sedimentary hiatus at the second unconformity was thus likely the combination of a favorable tectonic regime and a period of climate-controlled lake-level drop. Interestingly, the peak of lake retreat starts during the latest Priabonian and covers the EOT, and thus predates the EOGM and the global low of eustatic level (Miller et al., 2008); lake levels increase again during the EOGM. The lake level drop at the second unconformity seems thus uncorrelated with glacio-eustatic levels, and rather related to an earlier, late Priabonian and EOT aridity crisis.

5.3 A first record of the Eocene-Oligocene Transition in Anatolia

~~Our Δ_{47} -derived temperatures indicate a drop of $\sim 7^\circ\text{C}$ in soil carbonate growth temperature across the second unconformity, spanning the late Priabonian and the EOT, and persisting through the onset of the EOGM. Potential diagenetic processes, including burial recrystallization or hydrothermal alteration, could affect isotopic values in pedogenic carbonates. However, such processes are typically associated with the development of sparry calcite textures and the resetting of clumped isotope signatures at elevated temperatures (Henkes et al., 2014; Stolper and Eiler, 2015), which are not observed in this study. The preservation of micritic fabrics and surface-like clumped isotope temperatures –pedogenic morphologies therefore supports minimal post-depositional alteration.~~

~~Our Δ_{47} -derived temperatures indicate a drop of $\sim 7^\circ\text{C}$ in soil carbonate growth temperature across the second unconformity, spanning the late Priabonian and the EOT, and persisting through the onset of the EOGM. This temperature drop is similar to~~

a mis en forme : Indice

what is observed in several other clumped isotope records of the EOT world-wide. A comparable cooling of $\sim 7^\circ\text{C}$ in surface temperatures across the Eocene-Oligocene Transition was documented by Fan et al. (20187) in eastern Wyoming, inferred by clumped isotopes on pedogenic carbonates. In Western Europe, Δ_{47} measurements on freshwater gastropods from the Hampshire Basin (England) reveal a $\sim 4\text{--}6^\circ\text{C}$ decrease (Hren et al., 2013). In northeast Tibet, analyses from the Xining Basin indicate an apparent drop of $\sim 20^\circ\text{C}$, which has been interpreted as resulting from both a shift in the carbonate growth season and an actual temperature decline, leading to an estimated $\sim 9^\circ\text{C}$ decrease in surface temperatures (Page et al., 2019; See Supplementary figures-Table 6 for additional continental temperatures across the EOT).

Studies have shown that modern soil carbonates typically form during periods of soil dewatering during warm and dry seasons, and are commonly found in mid-latitude seasonal climate (Methner et al., 2016). This leads to a Δ_{47} bias toward Warm Month Mean Temperatures (WMMT) rather than MAT (Methner et al., 2016; Hough et al., 2014; Passey et al., 2010; Peters et al., 2013; Quade et al., 2013). In some instances, particularly under sub-humid climates with alternating wet and dry seasons, carbonate formation may instead reflect values closer to the mean annual temperature (Breecker et al., 2009; Peters et al., 2013). Under humid tropical monsoonal regimes, the bias can even shift toward recording winter temperatures, more closely reflecting Cold Month Mean Temperatures (CMT; Licht et al., 2022b). We compared our reconstructed surface temperatures with Eocene temperature estimates from pollen assemblages in Anatolia. There are no paleotemperature estimates for the Bartonian and Priabonian of central Anatolia, but Raynaud et al. (20265) report paleotemperatures for the Lutetian (48 to 41 Ma), stage that is globally warmer than the later stages of the Eocene (Westerhold et al., 2020). Lutetian Mean Annual Temperatures (MAT) range from 17.2 to 21.9°C , while Warm Month Mean Temperatures (WMMT) range from 24.7 to 28.1°C .— Our Δ_{47} -derived temperatures thus likely reflect carbonate growth during the warm season (See Fig. 8). We thus interpret the drop of $\sim 7^\circ\text{C}$ in our record as a drop in summer temperatures.

Stable isotope data from pedogenic carbonates reveal no clear change in $\delta^{18}\text{O}$ values across the second unconformity, and a slight decrease of approximately 0.7‰ is observed in $\delta^{13}\text{C}$ values. Reconstructed $\delta^{18}\text{O}_{\text{water}}$ values exhibit marked variability both before and after the second unconformity, ranging from -1‰ to -4‰ . These values contrast with previous reconstructions for the same region and time interval, which yielded consistently lower values. For example, Licht et al. (2017) reported $\delta^{18}\text{O}_{\text{water}}$ values between -5‰ and -7‰ from late Lutetian lacustrine carbonates, while Robert et al. (20265) obtained values of -5.3‰ to -8‰ from late Lutetian Balkanatolian mammal tooth enamel. The latter are also consistent with modern $\delta^{18}\text{O}$ values of meteoric waters in coastal Mediterranean Anatolia (Schemmel et al., 2013), after applying a $\sim 1\text{‰}$ correction for differences in seawater isotopic composition during the Eocene (Tindall et al., 2010), and with Oligocene meteoric water estimates (Lüdecke et al., 2013). The discrepancy between our reconstructed $\delta^{18}\text{O}_{\text{water}}$ values and those reported in earlier studies may reflect a proxy-related bias. Reconstructions are expected to capture primary meteoric water signatures, yet the $\delta^{18}\text{O}_{\text{water}}$ is influenced both by meteoric input and by evaporation. As emphasized by Kelson et al. (2023), evaporation in arid environments can enrich soil water $\delta^{18}\text{O}_{\text{water}}$ by $1\text{--}5\text{‰}$, even at depths > 40 cm. Such an effect could account for the offset between our estimates and previous reconstructions, and can explain the great variability in reconstructed soil water $\delta^{18}\text{O}_{\text{water}}$ values.

a mis en forme : Indice

a mis en forme : Indice

a mis en forme : Indice

620 The slight decrease in $\delta^{18}\text{O}_{\text{water}}$ values recorded across the unconformity, from mean values of -1.9‰ to -2.7‰ is thus difficult
621 to interpret considering the significant imprint of evaporative effects in our record. We note however that a $\sim 7^\circ\text{C}$ air
622 temperature drop would likely decrease rainwater isotopic composition by ca. -2‰ at these latitudes, and could thus partly
623 explain this decline (Bowen, 2008).
624 Our record thus indicates that the Eocene-Oligocene Transition and the EOGM in central Anatolia are associated with summer
625 temperature cooling of $\sim 7^\circ\text{C}$ and an increase of aridity marked first by a temporary lake retreat during the EOT, followed by
626 a lake rewatering during the EOGM, though marked with a persistent increase in lake salinity compared to the late Priabonian
627 lake system, as evidenced by the appearance of dolomite beds.

629 **5.4 A Late Eocene Warming?**

630 The portion of our section surrounding the first hiatus exhibits more pronounced shifts in isotopic data. Clumped isotope-
631 derived surface temperature estimates reveal a $\sim 9^\circ\text{C}$ increase beginning with the last paleosol below Hiatus 1, and warm
632 temperatures persist within all paleosols between hiatus 1 and 2. Previous studies (e.g., Bohaty and Zachos, 2003) have
633 documented a ‘Late Eocene Warming’ around $\sim 37\text{--}36\text{ Ma}$ seen in benthic foraminiferal records from the Southern Hemisphere.
634 Their reconstructions suggest a $\sim 3\text{--}5^\circ\text{C}$ temperature increase under ice-free conditions. The more recent benthic foraminiferal
635 compilation of Westerhold et al. (2020) dates this warming episode to ~ 37.5 to $\sim 36.5\text{ Ma}$. Tremblin et al. (2016) suggest a
636 significant warming of approximately 4°C in equatorial and North Atlantic regions starting during this time window and
637 lasting until at least 35 Ma , and possibly up to the EOT. The Recent pCO₂ reconstruction of cenCO2PIP (2023) suggests an
638 increase of pCO₂ during the latest Eocene before the EOT, though the amplitude and chronology of this increase are still
639 poorly constrained. By contrast, Tremblin et al. (2016) suggests that this warming episode could result from a reorganization
640 of ocean circulation in the North Atlantic. Numerous paleoceanographic records in the North Atlantic suggest an onset of a
641 regional subtropical gyre or an early version of the Atlantic meridional overturning circulation (AMOC) around 36 Ma , which
642 could lead to regional warming (Coxall et al., 2018; Liu et al., 2018).

643 Other records of the late Eocene warming on land are sparse. A recent study made in the Paris Basin records a decrease of
644 $\delta^{18}\text{O}$ values on bulk carbonate, also interpreted as a record of the late Eocene Warming during the C16r chron (Le Callonnec
645 et al., 2025). Clumped isotopic data from Tibetan soil carbonates highlight a warming trend across the late Priabonian,
646 culminating between 35 and 34 Ma (Page et al., 2019), and starting after a significant aridification step near the top of chron
647 C17n.1n (Abels et al., 2011), coherent with the beginning of the warming trend observed here.

648 In our record, this warming is accompanied by a mean $\delta^{18}\text{O}_{\text{water}}$ increase of 1.7‰ , along with enhanced variability. Given the
649 proximity of the study area to both the Neotethys and Paratethys domains, major changes in moisture source would require
650 substantial paleogeographic reorganization, and can be thus disregarded as a cause for the $\delta^{18}\text{O}_{\text{water}}$ change (Kaiseri-Özer et al.,
651 2013). The observed increase in $\delta^{18}\text{O}_{\text{water}}$ values is more consistent with progressive evaporative enrichment under increasingly
652 arid conditions, as frequently seen in soil carbonates (Kelson et al., 2023). The pronounced variability in $\delta^{18}\text{O}_{\text{water}}$ values is
653 also consistent with increased soil evaporation. While the The combined increase in mean $\delta^{18}\text{O}_{\text{water}}$ rises in parallel with

a mis en forme : Police :7 pt, Indice

a mis en forme : Indice

a mis en forme : Indice

654 temperature and, the pronounced variability during the late Eocene rather suggests increased seasonality and aridity, likely
 655 reflecting an increase of evaporative effects in soil carbonates (Kelson et al., 2023). In this context, the observed increase in
 656 $\delta^{18}\text{O}_{\text{water}}$ values is more consistent with progressive evaporative enrichment under increasingly arid conditions than with a shift
 657 in moisture source. This change is associated with a moderate rise in $\delta^{13}\text{C}$ values of approximately 0.7‰. This shift could be
 658 explained by a slight increase in plant water stress, also pointing to a trend toward increased aridity (Cerling and Quade, 1993;
 659 Kohn, 2010).
 660 The synchronicity of this apparent increase in temperature and aridity with the first unconformity suggests that it might have
 661 partly controlled the chronology of the hiatus. Though the uplift of the basin's southern margin is likely the primary cause for
 662 the discordance, the unconformity might have been enhanced by a base-level drop related to an increase of aridity. Overall,
 663 our record highlights an important climatic transition during the early-middle Priabonian within chron C17n.1n and preceding
 664 the EOT, marking the onset of an irreversible shift toward more arid conditions, coupled with elevated surface temperatures.
 665 This event can be hypothetically attributed to a “late Eocene Warming” impacting Eurasia, though its chronology and origin
 666 remain to be more precisely constrained. The subsequent cooling during the EOT brought back clumped isotope temperatures
 667 to their earlier, early Priabonian values, but accentuated a long-term trajectory toward increasing aridity.
 668
 669




Figure 8: Clumped isotope (Δ_{47}) paleotemperatures from pedogenic carbonates in the Büyüktefleken section (Çiçekdağı Basin, Turkey), spanning the Early–Middle Priabonian (~37.8–36.6 Ma) to the Early Rupelian (~33.4 Ma). The orange line shows the moving average; error bars = 1 standard error (SE). Temperature estimates for the Lutetian (48–41 Ma), derived from paleobotanical data reported by Raynaud et al. (20265). (WMT: Warm Month Temperatures; MAT = Mean Annual Temperatures; CMT = Cold Month Temperatures).

a mis en forme : Indice

5.5 Regional climatic response to cooling and implications for Asian terrestrial fauna dispersal

As many previous Eurasian records, our terrestrial section highlights the existence of a profound aridity crisis during the Priabonian, marked by two aridification steps during the middle (Hiatus 1) and latest (Hiatus 2) Priabonian. We show that the first aridification step is associated with a significant warming event, that we attribute to the Late Eocene Warming.

This first step likely initiated between prior to the base of the Chron C16n.1r (35.774 Ma) and 36.5 Ma, approximately corresponding to chron C17n.1n.

While quantitative paleontological data for Balkanatolia are sparse (fewer than ten Priabonian sites; Licht et al., 2022), regional trends suggest a significant impact of these climatic shifts on mammalian assemblages. In Western Europe, a comparable aridification phase is observed: Lettéron et al. (2022) describe an increase in aridity within the ASCI (Alès, Saint-Chaptes, and Issirac; France) lake system (Alès, Saint-Chaptes, and Issirac; France) between 37.1 and 35.2 Ma around the MP17, accompanied by a major turnover in floral assemblages based on palynological data (Alabouvette et al., 1983; Lettéron et al., 2018). In this region, the Grande Coupure (MP20) also coincides with a decline of 77% in endemic artiodactyl diversity (Weppe et al., 2023), which may have been linked to aridity and cooling. Other records from Central Asia indicate a similar increase in aridity during this interval. Wang et al. (2020) document enhanced aridity in the Tajik Basin, inferred from pedogenic carbonates showing reduced plant-derived soil respiration, correlated with the retreat of the Paratethys Sea during chron C17n.1n. Studies from the Xining Basin likewise record an early phase of aridification occurring during chron C17 (Page et al., 2019; Yang et al., 2022).

The link between this mid Priabonian aridity crisis and the late Eocene warming event is yet difficult to draw. Warming in the North Atlantic related to a proto-AMOC or a regional gyre, the only current explanation for the late Eocene warming, should lead to more regional evaporation and more precipitation on land (Elsworth et al., 2017). It thus cannot explain alone an Eurasian-wide aridity crisis. A tectonically driven Paratethys sea retreat episode in central Asia, as suggested by Wang et al. (2020), could partly explain a regional increase in aridity concomitant with the warmer temperatures of the late Eocene Warming, though its far-flung effects up to Balkanatolia and Europe are difficult to predict.

Regardless of its origin, the warming phase observed in our record covers mammalian biohorizons MP18 and MP19 and the first arrival of selected Asia-derived taxa in western Europe such as anthracotheriids, amphicyonids, and possibly gelocids, all of which likely dispersed through Balkanatolia (Métais et al., 2023). This synchronicity suggests that the late Eocene warming could have favored the westward dispersal of several Asia-derived taxa, which were already present on Balkanatolia since the

Bartonian (Licht et al., 2022a), and remained endemic to this biogeographic province during the early Oligocene (Van der Weerd et al. 2023; Maridet et al., 2025). The environmental stress associated with the warming event and the aridity crisis could have exacerbated the decline of endemic Balkanatolian taxa and further favored the penetration of Asia-derived taxa (Raynaud et al., 2025). The quasi-complete disappearance of Balkanatolian endemic taxa during the Priabonian (Licht et al., 2022a) further supports this interpretation, albeit Priabonian Balkanatolian faunas remain poorly documented (fewer than ten Priabonian sites; Licht et al., 2022a). We note however that the MP18 and MP19 interval corresponds to the peak of endemic mammalian diversity for large clades in western Europe like artiodactyls (Weppe et al., 2023). The late Eocene warming seems thus to have contrasted impacts: in Balkanatolia, it may have stressed endemic taxa and facilitated Asia-derived dispersal, whereas in western Europe, endemic diversity peaked before collapsing at the Grande-Coupure. The second aridity step is synchronous with the latest Priabonian and EOT and starts in our record at ~35.2 Ma. It is associated with a ~7°C cooling in carbonate growth temperatures leading to the EOGM, which resembles other terrestrial clumped isotope records covering this time interval in Eurasia (Page et al., 2019; Hren et al., 2013), and is compatible in amplitude with the rare European paleobotanical records that show some sensitivity to the EOT (Teodoridis and Kvaček, 2015; Suc et al., 2025). It is yet unclear if cooling started with the beginning of the second aridity step associated with the lake retreat at ~35.2 Ma, or, more likely, with the end of the event through the end of the EOT and early EOGM, just before the lake rewating.

Regardless of its origin, the warming phase observed in our record covers mammalian biohorizons MP18 and MP19 and the first arrival of selected Asia-derived taxa in western Europe such as Brontotheriidae, Amynodontidae or Anthracotheriids, gelocids or amphicyonidsae, which likely went through Balkanatolia (Metais et al., 2023). This synchronicity suggests that the late Eocene warming could have favored the westward dispersal of several Asia-derived taxa, which were already present on Balkanatolia since the Bartonian (Licht et al., 2022a). The environmental stress associated with the warming event and the aridity crisis could have enhanced the decline of endemic Balkanatolian taxa and further favored the penetration of Asia-derived taxa in the region (Raynaud et al., 20265). The quasi-complete disappearance of Balkanatolian endemic taxa during the Priabonian (Licht et al., 2022a) The last occurrence of Balkanatolian endemic taxa (a derived embrithopod) such as Palaeomasia embrithopoda is recorded around the EOT (Sanders et al., 2014) and further supports this interpretation, albeit Priabonian Balkanatolian fauna are poorly less documented. Quantitative paleontological data for Balkanatolia are sparse (fewer than ten Priabonian sites; Licht et al., 2022). We note however that MP18 and MP19 times correspond to the peak of endemic mammalian diversity for numerous large clades in western Europe like artiodactyls (Weppe et al., 2023). The late Eocene warming seems thus to have contrasting impacts: in Balkanatolia, it may have stressed endemic taxa and facilitated Asia-derived dispersal, whereas in western Europe, endemic diversity peaked before collapsing at the Grande-Coupure, a different impact in western Europe and Balkanatolia.

The second aridity step is synchronous with the latest Priabonian and EOT and starts in our record at ~35.2 Ma. It is associated with a ~7°C cooling in carbonate growth temperatures leading to the EOGM, which is similar with all other terrestrial clumped isotope records covering the time interval in Eurasia (Page et al., 2019; Hren et al., 2013), and is compatible in amplitude with the rare European paleobotanical records that show some sensitivity to the EOT (Teodoridis and Kvaček, 2015; Sue et al.,

2025). It is yet unclear if cooling started with the beginning of the second aridity step associated with the lake retreat at ~35.2 Ma, or, more likely, with the end of the event through the end of the EOT and early EOGM, just before the lake rewet-
Interestingly, the second aridity step predates the EOGM by at least 1.5 Myr, suggesting that its origin is not directly related to the Antarctic glaciation. Rather, we suggest that it reflects the hydroclimatic response to an earlier cooling event in the North Atlantic. Indeed, cooling in the North Atlantic seems to have preceded the EOT and to have been heterogeneously distributed both spatially and temporally, starting at least 1 Myr before the Eocene-Oligocene boundary (Sliwinska et al., 2023). This heterogeneity is so far explained by a complex response of the North Atlantic gyres to decreasing pCO_2 , which changes the location, strength and structure of these gyres (Sliwinska et al., 2023). Regardless of its origin, the second aridity step covers mammalian biohorizons MP20 and the onset of biodiversity decline for western European endemic artiodactyls, which culminates in the earliest Oligocene with a decline of 77% in artiodactyl diversity (Weppe et al., 2023). We thus suggest that the environmental stress associated with the second aridity step could have enhanced the decline of western European taxa. While it is unclear how this environmental stress impacted Balkanatolian fauna due to the sparse paleontological record of the Priabonian and Rupelian in the area, the last known occurrence of an endemic Balkanatolian taxon (the embrithopod *Axainamasia sandersi*, Métais et al. 2024) is recorded around the EOT (Sanders et al., 2014) and suggests that local ecological or paleogeographic factors allowed some endemic Balkanatolian species to persist well beyond the colonization of Balkanatolia by Asian predators and competitors. The aridification step and the consecutive EOGM correspond to periods without faunal exchanges between Balkanatolia and western Europe (Mennecart et al., 2021; Métais et al., 2023). This timing is counter-intuitive because the eustatic drop at the EOGM fully connected western Balkanatolia to western Europe (Licht et al., 2022a; Montheil et al., 2025b). This suggests that this climatic event resulted in the set-up of a persistent environmental barrier in central and southern Europe, blocking the westward expansion of Asia-derived mammals. This paleoenvironmental barrier may have been enhanced by orographic effects on regional precipitation, caused by the nascent Alps (Kocsis et al., 2014).
Interestingly, the second aridity step predates the EOGM by at least 1.5 Myr, suggesting that its origin is not directly related to the Antarctic glaciation. We suggest that they reflect the hydroclimatic response to an earlier cooling event in the North Atlantic. Indeed, cooling in the North Atlantic seems to have preceded the EOT and to have been heterogeneously spatially and temporally distributed, starting at least 1 Myr before the Eocene-Oligocene boundary (Sliwinska et al., 2023). This heterogeneity is so far explained by a complex response of the north Atlantic gyres to decreasing CO_2 , which changes gyres' location, strength and structure (Sliwinska et al., 2023). Regardless of its origin, the second aridity step covers mammalian biohorizons MP20 and the onset of biodiversity decline for western European endemic artiodactyls, which culminates in the earliest Oligocene with a decline of 77% in artiodactyl diversity (Weppe et al., 2023). We thus suggest that the environmental stress associated with the second aridity step could have enhanced the decline of western European taxa. While it is unclear how this environmental stress impacted Balkanatolian fauna due to the sparse paleontological record of the Priabonian and Rupelian in the area, the very last occurrence of endemic balkanatolian taxa such as *Palaeoamasia* (a derived embrithopod) is recorded around the EOT (Sanders et al., 2014) and suggests that the local fauna was also impacted. It is unclear how this

a mis en forme : Indice

a mis en forme : Police :Italique

environmental stress impacted Balkanatolian fauna as the paleontological record of the Priabonian and Rupelian of the area is less complete. However, reminding we note that the last occurrence of Balkanatolian endemic taxa, the Palaeoamasia embriothopoda, is dated around the EOT (Sanders et al., 2014), suggesting that the local fauna was also impacted. The aridification step and the consecutive EOGM correspond to periods without faunal exchanges between Balkanatolia and western Europe (Mennecart et al., 2021; Metais et al., 2023). This synchronicity is counter-intuitive because the eustatic drop at the EOGM fully connected western Balkanatolia to western Europe (Licht et al., 2022a; Montheil et al., 2025b). This suggests that this climatic event resulted in the set-up of a persistent environmental barrier in central and southern Europe, blocking the westward penetration of Asia-derived mammals. This paleoenvironmental barrier might have been enhanced by orographic effects on regional precipitation, caused by the nascent Alps (Kocsis et al., 2014).

6 Conclusion

Our sedimentological and stable isotope records document two major phases of aridification during the late Eocene. The first, occurring around 37–36.5 Ma prior to the base of chron C16n.1r (35.774 Ma), is associated with a marked increase of ~9°C in summer surface temperatures recorded by clumped isotope data, representing the first evidence of the Late Eocene Warming in western Eurasia. The second phase took place during the latest Priabonian, extending through the EOT and into the EOGM, and is characterized by a subsequent decline of ~7°C in summer surface temperatures during the EOGM.

These climatic transitions likely reflect large-scale reorganizations in atmospheric and oceanic circulation in the North Atlantic preceding the EOGM. Such environmental changes may have contributed to the decline of endemic fauna in Balkanatolia and western Europe during the late Eocene. However, the precise chronology and extent of these events require further refinement through integration with additional continental records to strengthen this interpretation.

Data availability

The data and supplementary figures are available as a supplementary material available online.

Acknowledgements

This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 101043268) to AL. Our geological and paleontological field work has also been supported by grants from the National Science Foundation (EAR 1543684 and 2141115) and the David B. Jones Foundation. We thank Corinne Sonzogni, Anne Alexandre, Mathieu Daeron, Jens Fiebig, Pascal Parmentier, Patricia Rieu et Sylvie de Freitas for their help setting up the Clumped isotope line at CEREGE. We also thank Julien Longerey for helping with sample preparation and imaging at CEREGE, Daniel Borschneck for his help with XRD analysis, Elodie Gazquez for her administrative support and Daniel Robert for his assistance with sample analyses.

Financial support
This work has been supported by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No. 101043268) to AL. Our geological and paleontological field work has also been supported by grants from the National Science Foundation (EAR 1543684 and 2141115) and the David B. Jones Foundation.

Disclaimer
Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher

Author contributions
PB participated in fieldwork; sample preparation (palaeomagnetism, pedogenic carbonates); data analyses (palaeomagnetism, stable and clumped isotopes); data interpretation; and discussion and writing of the manuscript. AL participated in fieldwork; clumped isotope data processing; data interpretation; discussion and writing; and supervised the project. ALJ participated in stable and clumped isotope analyses; clumped isotope data processing; and discussion and writing. LM participated in fieldwork; data interpretation; discussion and writing. FD participated in data interpretation; discussion and writing. MK, FO, MSA, GM, BR, and KCB participated in fieldwork; discussion and writing. DI and PC participated in discussion and writing.

Competing interests
The authors declare that they have no conflict of interest.

826
827
828
829
830
831
832
833
834
835
836
837

838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871

References

Abels, H. A., Dupont-Nivet, G., Xiao, G., Bosboom, R., & Krijgsman, W. (2011). Step-wise change of Asian interior climate preceding the Eocene–Oligocene Transition (EOT). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 299(3-4), 399-412. <https://doi.org/10.1016/j.palaeo.2010.11.028>

Akgün, F., Akay, E., & Erdoğan, B. (s. d.). Tertiary Terrestrial to Shallow Marine Deposition in Central Anatolia : A Palynological Approach.

[Alabouvette, B., Chedhomme, J., Frédet, J.M. and Lartaud, V. \(1983\) Inventaire des ressources nationales de charbon : lignites du fossé d'Alès-Barjac. Bur. Rech. Géol. Min. Mém., 83, Orléans, 63 pp.](#)

Anderson, N. T., Kelson, J. R., Kele, S., Daëron, M., Bonifacie, M., Horita, J., ... & Bergmann, K. D. (2021). A unified clumped isotope thermometer calibration (0.5–1,100 C) using carbonate-based standardization. *Geophysical Research Letters*, 48(7), e2020GL092069.

Antoine, P.-O., Yans, J., Castillo, A. A., Stutz, N., Abello, M. A., Adnet, S., Custódio, M. A., Benites-Palomino, A., Billet, G., Boivin, M., Herrera, F., Jaramillo, C., Martínez, C., Moreno, F., Navarrete, R. E., Negri, F. R., Parra, F., Pujos, F., Rage, J.-C., ... Marivaux, L. (2021). Biotic community and landscape changes around the Eocene–Oligocene transition at Shapaja, Peruvian Amazonia: Regional or global drivers? *Global and Planetary Change*, 202, 103512. <https://doi.org/10.1016/j.gloplacha.2021.103512>

[Arason, Þ. Levi, S. Maximum likelihood solution for inclination-only data in paleomagnetism, *Geophysical Journal International*, Volume 182, Issue 2, 753–771. \(2010\)](#)

Barrier, E., Vrielynck, B., Brouillet, J., Brunet, M., 2018. Paleotectonic reconstruction of the central tethyan realm. Tectono-sedimentary-palinspastic maps from late Permian to Pliocene. CCGM/CGMW, Paris. Atlas of 20.

Bernasconi, S. M., Daëron, M., Bergmann, K. D., Bonifacie, M., Meckler, A. N., Affek, H. P., ... & Ziegler, M. (2021). InterCarb: A community effort to improve interlaboratory standardization of the carbonate clumped isotope thermometer using carbonate standards. *Geochemistry, Geophysics, Geosystems*, 22(5), e2020GC009588.

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

Bhattacharya JP. 2010. *Deltas. Facies models*, 4, 233–264.

Bohaty, S. M., & Zachos, J. C. (2003). Significant Southern Ocean warming event in the late middle Eocene. *Geology*, 31(11), 1017. <https://doi.org/10.1130/G19800-1> <https://doi.org/10.1130/G19800.1>

Böhme, M., Aiglstorfer, M., Antoine, P. O., Appel, E., Havlik, P., Métais, G., ... & Prieto, J. (2013). Na Duong (northern Vietnam)-an exceptional window into Eocene ecosystems from Southeast Asia. *Zitteliana*, 53, 121-167.

Bowen, G. J. (2008). Spatial analysis of the intra-annual variation of precipitation isotope ratios and its climatological corollaries. *Journal of Geophysical Research: Atmospheres*, 113(D5), 2007JD009295. <https://doi.org/10.1029/2007JD009295>

Breecker, D. O., Sharp, Z. D., & McFadden, L. D. (2009). Seasonal bias in the formation and stable isotopic composition of pedogenic carbonate in modern soils from central New Mexico, USA. *Geological Society of America Bulletin*, 121(3-4), 630-640. <https://doi.org/10.1130/b26413.1>

Becker, D. 2009. Earliest record of rhinocerotoids (Mammalia: Perissodactyla) from Switzerland: systematics and biostratigraphy. *Swiss Journal of Geosciences*, 102: 489–504.

Campbell, C. F., Taylor, M. H., Mueller, M. A., Licht, A., Ocakoğlu, F., & Beard, K. C. (2026). The Late Paleogene Çankırı Basin of Türkiye : The geologic record of lithospheric foundering beneath the Central Anatolian Plateau. *Geological Society of America Bulletin*, 138, 2071-2086. <https://doi.org/10.1130/B38453.1>

Campbell, C. F., Taylor, M. H., Mueller, M. A., Licht, A., & Ocakoğlu, F. (s. d.) (2025). The Late Paleogene Çankırı Basin of Türkiye : The geologic record of lithospheric foundering beneath the Central Anatolian Plateau. *Geological Society of America Bulletin*, 136.

Cerling, T. E., & Quade, J. (1993). Stable carbon and oxygen isotopes in soil carbonates. In P. K. Swart, K. C. Lohmann, J. McKenzie, & S. Savin (Eds.), *Climate change in continental isotopic records* (Vol. 78, pp. 217–231). American Geophysical Union

Couturier, J., Levard, C., Collin, B., Chaurand, P., Vidal, V., Mathon, O., Duvivier, A., Angeletti, B., Borschneck, D., Rose, J., Pellet-Rostaing, S., & Arrachart, G. (2025). Dissolution of rare earth elements : Exploring the ability of deep eutectic solvents and organic acid solutions, the case of lactic acid. *Separation and Purification Technology*, 366, 132740. <https://doi.org/10.1016/j.seppur.2025.132740>

Coxall, H. K., Huck, C. E., Huber, M., Lear, C. H., Legarda-Lisarrri, A., O'Regan, M., Sliwinska, K. K., Van De Flierdt, T., De Boer, A. M., Zachos, J. C., & Backman, J. (2018). Export of nutrient rich Northern Component Water preceded early Oligocene Antarctic glaciation. *Nature Geoscience*, 11(3), 190-196. <https://doi.org/10.1038/s41561-018-0069-9>

Daëron, M. (2021). Full propagation of analytical uncertainties in $\Delta 47$ measurements. *Geochemistry, Geophysics, Geosystems*, 22(5), e2020GC009592.

Deenen, M.H.L., Langereis, C.G., van Hinsbergen, D.J.J., and Biggin, A.J., *Geomagnetic secular variation and the statistics of paleomagnetic directions*, *Geophysical Journal International* 186, p. 509-520 (2011)

Doglioni, C., Busatta, C., Bolis, G., Marianini, L., Zanella, M., 1996. Structural evolution of the eastern Balkans (Bulgaria). *Mar. Pet. Geol.* 13 (2), 225–251.

- a mis en forme : Surlignage
- a mis en forme : Police :Italique
- a mis en forme : Police :Italique
- a mis en forme : Police :(Par défaut) Times New Roman
- a mis en forme : Police :(Par défaut) Times New Roman
- a mis en forme : Police :(Par défaut) Times New Roman
- a mis en forme : Police :(Par défaut) Times New Roman
- a mis en forme : Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204))
- a mis en forme : Police :Non Italique, Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204))
- a mis en forme : Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204))
- a mis en forme : Police :Italique
- a mis en forme : Police :Italique
- a mis en forme : Police :(Par défaut) Times New Roman
- a mis en forme : Police :(Par défaut) Times New Roman

906 Elsworth, G., Galbraith, E., Halverson, G., & Yang, S. (2017). Enhanced weathering and CO2 drawdown caused by latest
907 Eocene strengthening of the Atlantic meridional overturning circulation. *Nature Geoscience*, 10(3), 213-216.
908 <https://doi.org/10.1038/ngeo2888>

909 [Erdogan, B., Akay, E., & Ugur, M. S. \(1996\). Geology of the Yozgat region and evolution of the collisional Çankiri](#)
910 [Basin. *International Geology Review*, 38\(9\), 788-806.](#)

911 [Eronen, J. T., Janis, C. M., Chamberlain, C. P., & Mulch, A. \(2015\). Mountain uplift explains differences in Palaeogene](#)
912 [patterns of mammalian evolution and extinction between North America and Europe. *Proceedings of the Royal Society B:*](#)
913 [*Biological Sciences*, 282\(1809\), 20150136. <https://doi.org/10.1098/rspb.2015.0136>](#)

914 Fan, M., Ayyash, S. A., Tripathi, A., Passey, B. H., & Griffith, E. M. (2018). Terrestrial cooling and changes in hydroclimate
915 in the continental interior of the United States across the Eocene-Oligocene boundary. *GSA Bulletin*, 130(7-8), 1073-1084.
916 <https://doi.org/10.1130/b31732.1>

917 Fiebig, J., Bernecker, M., Meijer, N., Methner, K., Staudigel, P. T., Davies, A. J., ... & Petersen, S. V. (2024). Carbonate
918 clumped isotope values compromised by nitrate-derived NO2 interferent. *Chemical Geology*, 670, 122382.

919 [Fisher, R. A. \(1953\). Dispersion on a sphere. *Proceedings of the royal society of London. Series A. Mathematical and physical*](#)
920 [*sciences*, 217\(1130\), 295-305.](#)

921 García-Ruiz, J. M. (2023). A fluctuating solution to the dolomite problem. *Science*, 382(6673), 883-884.
922 <https://doi.org/10.1126/science.adl1734>

923 [Gradstein, F. M., Ogg, J. G., Schmitz, M. D., & Ogg, G. M. \(Eds.\). \(2020\). *Geologic time scale 2020*. Elsevier.](#)

924 [Gülyüz, E., Kaymakci, N., Meijers, M. J. M., Van Hinsbergen, D. J. J., Lefebvre, C., Vissers, R. L. M., Hendriks, B. W. H.,](#)
925 [& Peynircioğlu, A. A. \(2013\). Late Eocene evolution of the Çiçekdağı Basin \(central Turkey\): Syn-sedimentary compression](#)
926 [during microcontinent-continent collision in central Anatolia. *Tectonophysics*, 602, 286-299.](#)
927 <https://doi.org/10.1016/j.tecto.2012.07.003>

928 [Grandi, F. and Bona, F. 2017. *Prometherium dalmatinum* from the late Eocene of Grancona \(Vicenza, NE Italy\). The oldest](#)
929 [terrestrial mammal of the Italian peninsula. *Comptes Rendus Palevol* 16 \(7\): 738-745.](#)

930 Guo, P., Wen, H., Li, C., He, H., & Sánchez-Román, M. (2023). Lacustrine dolomite in deep time : What really matters in
931 early dolomite formation and accumulation? *Earth-Science Reviews*, 246, 104575.
932 <https://doi.org/10.1016/j.earscirev.2023.104575>

933 Gürer, D., Van Hinsbergen, D. J. J., Matenco, L., Corfu, F., & Cascella, A. (2016). Kinematics of a former oceanic plate of
934 the Neotethys revealed by deformation in the Ulukışla basin (Turkey). *Tectonics*, 35(10), 2385-2416.
935 <https://doi.org/10.1002/2016TC004206>

936 [Heslop, D., et al., A bootstrap common mean direction test. *Journal of Geophysical Research: Solid Earth*, 128\(8\),](#)
937 [p.e2023JB026983 \(2023\).](#)

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : Italique

a mis en forme : Police : Italique

a mis en forme : Police : (Par défaut) Times New Roman

a mis en forme : Police : 10 pt

a mis en forme : Interligne : 1.5 ligne, Bordure : Haut: (Pas de bordure), Bas: (Pas de bordure), Gauche: (Pas de bordure), Droite: (Pas de bordure), Entre : (Pas de bordure), Motif : Transparente

a mis en forme : Police : 10 pt, Couleur de police : Automatique

Hough, B. G., Fan, M., & Passey, B. H. (2014). Calibration of the clumped isotope geothermometer in soil carbonate in Wyoming and Nebraska, USA : Implications for paleoelevation and paleoclimate reconstruction. *Earth and Planetary Science Letters*, 391, 110-120. <https://doi.org/10.1016/j.epsl.2014.01.008>

Hren, M. T., Sheldon, N. D., Grimes, S. T., Collinson, M. E., Hooker, J. J., Bugler, M., & Lohmann, K. C. (2013). Terrestrial cooling in Northern Europe during the Eocene–Oligocene transition. *Proceedings of the National Academy of Sciences*, 110(19), 7562-7567. <https://doi.org/10.1073/pnas.1210930110>

Hutchinson, D. K., Coxall, H. K., Lunt, D. J., Steinthorsdottir, M., de Boer, A. M., Baatsen, M., von der Heydt, A., Huber, M., Kennedy-Asser, A. T., Kunzmann, L., Ladant, J.-B., Lear, C. H., Moraweck, K., Pearson, P. N., Piga, E., Pound, M. J., Salzmann, U., Scher, H. D., Sijp, W. P., ... Zhang, Z. (2021). The Eocene–Oligocene transition : A review of marine and terrestrial proxy data, models and model–data comparisons. *Climate of the Past*, 17(1), 269-315. <https://doi.org/10.5194/cp-17-269-2021>

Judd, E. J., Tierney, J. E., Lunt, D. J., Montañez, I. P., Huber, B. T., Wing, S. L., & Valdes, P. J. (2024). A 485-million-year history of Earth's surface temperature. *Science*, 385(6715), eadk3705. <https://doi.org/10.1126/science.adk3705>

Karadenizli, L. (2011). Oligocene to Pliocene palaeogeographic evolution of the Çankırı-Çorum Basin, central Anatolia, Turkey. *Sedimentary Geology*, 237(1-2), 1-29.

Katz, M. E., Miller, K. G., Wright, J. D., Wade, B. S., Browning, J. V., Cramer, B. S., & Rosenthal, Y. (2008). Stepwise transition from the Eocene greenhouse to the Oligocene icehouse. *Nature Geoscience*, 1(5), 329-334. <https://doi.org/10.1038/ngeo179>

Kaymakci, N., White, S. H., & Vandijk, P. M. (2003). Kinematic and structural development of the Çankiri Basin (Central Anatolia, Turkey) : A paleostress inversion study. *Tectonophysics*, 364(1-2), 85-113. [https://doi.org/10.1016/S0040-1951\(03\)00043-X](https://doi.org/10.1016/S0040-1951(03)00043-X)

Kayseri-Özer, M. S. (2013). Spatial distribution of climatic conditions from the Middle Eocene to Late Miocene based on palynoflora in Central, Eastern and Western Anatolia. *Geodinamica Acta*, 26(1-2), 122-157.

Kelson, J. R., Huth, T. E., Passey, B. H., Levin, N. E., Petersen, S. V., Ballato, P., Beverly, E. J., Breecker, D. O., Hoke, G. D., Hudson, A. M., Ji, H., Licht, A., Oerter, E. J., & Quade, J. (2023). Triple oxygen isotope compositions of globally distributed soil carbonates record widespread evaporation of soil waters. *Geochimica et Cosmochimica Acta*, 355, 138-160. <https://doi.org/10.1016/j.gca.2023.06.034>

Kent-Corson, M. L., Ritts, B. D., Zhuang, G., Bovet, P. M., Graham, S. A., & Page Chamberlain, C. (2009). Stable isotopic constraints on the tectonic, topographic, and climatic evolution of the northern margin of the Tibetan Plateau. *Earth and Planetary Science Letters*, 282(1-4), 158-166. <https://doi.org/10.1016/j.epsl.2009.03.011>

Kim, J., Kimura, Y., Puchala, B., Yamazaki, T., Becker, U., & Sun, W. (2023). Dissolution enables dolomite crystal growth near ambient conditions. *Science*, 382(6673), 915-920. <https://doi.org/10.1126/science.adi3690>

R. F. King The Remanent Magnetism of Artificially Deposited Sediments *Mon. Nat. Roy. astr. Soc., Geophys. Suppl.*, 7, 115–134. (1955).

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :Italique

a mis en forme : Police :Italique

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Couleur de police : Couleur personnalisée(RVB(34;34;34)), Surlignage

a mis en forme : Police :Italique, Couleur de police : Couleur personnalisée(RVB(34;34;34)), Surlignage

a mis en forme : Couleur de police : Couleur personnalisée(RVB(34;34;34)), Surlignage

a mis en forme : Police :Italique, Couleur de police : Couleur personnalisée(RVB(34;34;34)), Surlignage

a mis en forme : Couleur de police : Couleur personnalisée(RVB(34;34;34)), Surlignage

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

972 Kocsis, L., Ozsvárt, P., Becker, D., Ziegler, R., Scherler, L., & Codrea, V. (2014). Orogeny forced terrestrial climate variation
973 during the late Eocene–early Oligocene in Europe. *Geology*, 42(8), 727–730. <https://doi.org/10.1130/G35673.1>
974 Kohn, M. J. (2010). Carbon isotope compositions of terrestrial C3 plants as indicators of (paleo) ecology and (paleo) climate.
975 *Proceedings of the National Academy of Sciences*, 107(46), 19691–19695.
976 Kohn, M. J., Strömberg, C. A. E., Madden, R. H., Dunn, R. E., Evans, S., Palacios, A., & Carlini, A. A. (2015). Quasi-static
977 Eocene–Oligocene climate in Patagonia promotes slow faunal evolution and mid-Cenozoic global cooling. *Palaeogeography*,
978 *Palaeoclimatology*, *Palaeoecology*, 435, 24–37.
979 <https://doi.org/10.1016/j.palaeo.2015.05.028><https://doi.org/10.1016/j.palaeo.2015.05.028>
980 <https://doi.org/10.1134/S0869593824700357><https://doi.org/10.1134/S0869593824700357>
981 <https://doi.org/10.1134/S0869593824700357>
982 <https://doi.org/10.1134/S0869593824700357>
983 Laurence Le Callonnec, Châteauneuf, J.-J., Renard, M., Zuddas, P., Emmanuel, L., & Riveline, J. (2025). New Data on the
984 Recording of Global Climatic Events at the Eocene–Oligocene Transition in the Sediments of the Paris Basin (Cormeilles-en-
985 Paris Section). *Stratigraphy and Geological Correlation*, 33(2), 257–277.
986 <https://doi.org/10.1134/S0869593824700357><https://doi.org/10.1134/S0869593824700357>
987 <https://doi.org/10.1134/S0869593824700357>
988 Lefebvre, C., Meijers, M. J. M., Kaymakci, N., Peynircioğlu, A., Langereis, C. G., & Van Hinsbergen, D. J. J. (2013).
989 Reconstructing the geometry of central Anatolia during the late Cretaceous : Large-scale Cenozoic rotations and deformation
990 between the Pontides and Taurides. *Earth and Planetary Science Letters*, 366, 83–98.
991 <https://doi.org/10.1016/j.epsl.2013.01.003>
992 Lettéron, A., Fournier, F., Hamon, Y., Villier, L., Margerel, J.-P., Bouche, A., Feist, M., & Joseph, P. (2017). Multi-proxy
993 paleoenvironmental reconstruction of saline lake carbonates : Paleoclimatic and paleogeographic implications (Priabonian-
994 Rupelian, Issirac Basin, SE France). *Sedimentary Geology*, 358, 97–120. <https://doi.org/10.1016/j.sedgeo.2017.07.006>
995 Lettéron, A., Hamon, Y., Fournier, F., Demory, F., Séranne, M., & Joseph, P. (2022). Stratigraphic architecture of a saline
996 lake system : From lake depocentre (Alès Basin) to margins (Saint-Chartes and Issirac basins), Eocene–Oligocene transition,
997 south-east France. *Sedimentology*, 69(2), 651–695. <https://doi.org/10.1111/sed.12920>
998 Lettéron, A., Hamon, Y., Fournier, F., Séranne, M., Pellenard, P., & Joseph, P. (2018). Reconstruction of a saline, lacustrine
999 carbonate system (Priabonian, St-Chartes Basin, SE France) : Depositional models, paleogeographic and paleoclimatic
1000 implications. *Sedimentary Geology*, 367, 20–47. <https://doi.org/10.1016/j.sedgeo.2017.12.023>
1001 Li, B., Sun, D., Wang, X., Zhang, Y., Hu, W., Wang, F., Li, Z., Ma, Z., & Liang, B. (2016). $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ records from a
1002 Cenozoic sedimentary sequence in the Lanzhou Basin, Northwestern China : Implications for palaeoenvironmental and
1003 palaeoecological changes. *Journal of Asian Earth Sciences*, 125, 22–36. <https://doi.org/10.1016/j.jseas.2016.05.010>

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

1004 Li, Q., Zhang, Y., Dong, L., & Guo, Z. (2018). Oligocene syn depositional lacustrine dolomite : A study from the southern
1005 Junggar Basin, NW China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 503, 69-80.
1006 <https://doi.org/10.1016/j.palaeo.2018.04.004>

1007 Li, S., & Guan, P. (2023). Advances in carbon and oxygen isotopes and clumped isotope of lacustrine carbonates. *Acta*
1008 *Scientiarum Naturalium Universitatis Pekinensis*, 59(6), 1052-1068.

1009 Licht, A., Coster, P., Ocakoğlu, F., Campbell, C., Métais, G., Mulch, A., Taylor, M., Kappelman, J., & Beard, K. C. (2017).
1010 Tectono-stratigraphy of the Orhaniye Basin, Turkey : Implications for collision chronology and Paleogene biogeography of
1011 central Anatolia. *Journal of Asian Earth Sciences*, 143, 45-58. <https://doi.org/10.1016/j.jseas.2017.03.033>

1012 Licht, A., Kelson, J., Bergel, S., Schauer, A., Petersen, S. V., Capirala, A., Huntington, K. W., Dupont-Nivet, G., Win, Z., &
1013 Aung, D. W. (2022). Dynamics of Pedogenic Carbonate Growth in the Tropical Domain of Myanmar. *Geochemistry,*
1014 *Geophysics, Geosystems*, 23(7), e2021GC009929. <https://doi.org/10.1029/2021GC009929>

1015 Licht, A., Métais, G., Coster, P., İbilioğlu, D., Ocakoğlu, F., Westerweel, J., Mueller, M., Campbell, C., Mattingly, S., Wood,
1016 M. C., & Beard, K. C. (2022a). Balkanatolia : The insular mammalian biogeographic province that partly paved the way to the
1017 Grande Coupure. *Earth-Science Reviews*, 226, 103929. <https://doi.org/10.1016/j.earscirev.2022.103929>

1018 Licht, A., Kelson, J., Bergel, S., Schauer, A., Petersen, S. V., Capirala, A., Huntington, K. W., Dupont-Nivet, G., Win, Z., &
1019 Aung, D. W. (2022b). Dynamics of Pedogenic Carbonate Growth in the Tropical Domain of Myanmar. *Geochemistry,*
1020 *Geophysics, Geosystems*, 23(7), e2021GC009929. <https://doi.org/10.1029/2021GC009929>

1021

1022 Liu, Z., Huang, C., Algeo, T. J., Liu, H., Hao, Y., Du, X., Lu, Y., Chen, P., Guo, L., & Peng, L. (2018). High-resolution
1023 astrochronological record for the Paleocene-Oligocene (66–23 Ma) from the rapidly subsiding Bohai Bay Basin, northeastern
1024 China. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 510, 78-92. <https://doi.org/10.1016/j.palaeo.2017.10.030>

1025 Lurcock, P. C. and G. S. Wilson (2012). PuffinPlot: A versatile, user-friendly program for paleomagnetic analysis. *Geochemistry,*
1026 *Geophysics, Geosystems*, 13, Q06Z45. doi:10.1029/2012GC004098.

1027 Lüdecke, T. (2013). Stable isotope-based reconstruction of Oligo-Miocene paleoenvironment and paleohydrology of Central
1028 Anatolian lake basins (Turkey). *TURKISH JOURNAL OF EARTH SCIENCES*. <https://doi.org/10.3906/yer-1207-11>

1029 C, M.W. Classification of the reversal test in palaeomagnetism *Geophysical Journal International* 103 (3), pp. 725-729 (1990).

1030 Maridet, O., Codrea, V. A., Fărcaș, C., Solomon, A. A., Venczel, M., & Tissier, J. (2025). The record of cricetid rodents across
1031 the Eocene–Oligocene transition in Transylvania, Romania: implications for the “Grande Coupure” at European scale. *Acta*
1032 *Palaeontologica Polonica*, 70, 291-327.

1033 Meijer, N., Methner, K., Seymour, N. M., Bernecker, M., Fiebig, J., Voigt, S., Chamberlain, C. P., & Mulch, A. (2025).
1034 Oligocene cooling in the North American cordillera (SW Montana, USA) revealed by dual clumped isotope thermometry.
1035 *Earth and Planetary Science Letters*, 671, 119618. <https://doi.org/10.1016/j.epsl.2025.119618>

1036 Meng, J., & McKenna, M. C. (1998). Faunal turnovers of Palaeogene mammals from the Mongolian Plateau. *Nature*,
1037 394(6691), 364-367. <https://doi.org/10.1038/28603>

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman, 9 pt

a mis en forme : Police :(Par défaut) Times New Roman, 9 pt, Italique

a mis en forme : Police :(Par défaut) Times New Roman, 9 pt

a mis en forme : Police :9 pt

a mis en forme : Police :Italique

a mis en forme : Police :Italique

a mis en forme : Police :(Par défaut) Times New Roman

1038 Menecart, B., Aiglstorfer, M., Li, Y., Li, C., & Wang, S. (2021). Ruminants reveal Eocene Asiatic palaeobiogeographical
1039 provinces as the origin of diachronous mammalian Oligocene dispersals into Europe. *Scientific Reports*, 11(1), 17710.
1040 <https://doi.org/10.1038/s41598-021-96221-x>

1041 Métais, G., Coster, P., Licht, A., Ocakoğlu, F., & Beard, K. C. (2023). Additions to the late Eocene Süngülü mammal fauna in
1042 Easternmost Anatolia and the Eocene-Oligocene transition at the periphery of Balkanatolia. *Comptes Rendus Palevol*, 35.
1043 <https://doi.org/10.5852/cr-palevol2023v22a35>

1044 Métais, G., Coster, P. M., Kappelman, J. R., Licht, A., Ocakoğlu, F., Taylor, M. H., & Beard, K. C. (2018a). Eocene
1045 metatherians from Anatolia illuminate the assembly of an island fauna during Deep Time. *PLOS ONE*, 13(11), e0206181.
1046 <https://doi.org/10.1371/journal.pone.0206181>

1047 Métais, G., Coster, P. M., Kappelman, J. R., Licht, A., Ocakoğlu, F., Taylor, M. H., & Beard, K. C. (2018b). Eocene
1048 metatherians from Anatolia illuminate the assembly of an island fauna during Deep Time. *PLOS ONE*, 13(11), e0206181.
1049 <https://doi.org/10.1371/journal.pone.0206181>

1050 Métais, G., Coster, P., Kaya, M., Licht, A., Miller, K., Ocakoğlu, F., Rust, K., & Beard, K. C. (2024). Rapid colonization and
1051 diversification of a large-bodied mammalian herbivore clade in an insular context: New embrithopods from the Eocene of
1052 Balkanatolia. *Journal of Mammalian Evolution*, 31, 15. <https://doi.org/10.1007/s10914-024-09711-w>

1053 Methner, K., Mulch, A., Fiebig, J., Wacker, U., Gerdes, A., Graham, S. A., & Chamberlain, C. P. (2016). Rapid Middle Eocene
1054 temperature change in western North America. *Earth and Planetary Science Letters*, 450, 132-139.
1055 <https://doi.org/10.1016/j.epsl.2016.05.053>

1056 Miall AD. 2010. Alluvial deposits. Facies models 4: 105–138.

1057 Miall AD. 2013. *The geology of fluvial deposits: sedimentary facies, basin analysis, and petroleum geology*: Springer, 582 p.

1058 Miller, K. G., Wright, J. D., and Fairbanks, R. G.: Unlocking the Ice House: Oligocene-Miocene oxygen isotopes, eustasy, and
1059 margin erosion, *J. Geophys. Res.-Sol. Ea.*, 96, 6829–6848, <https://doi.org/10.1029/90JB02015>, 1991.

1060 Miller, K. G., Browning, J. V., Aubry, M.-P., Wade, B. S., Katz, M. E., Kulpecz, A. A., & Wright, J. D. (2008). Eocene-
1061 Oligocene global climate and sea-level changes : St. Stephens Quarry, Alabama. *Geological Society of America Bulletin*,
1062 120(1-2), 34-53. <https://doi.org/10.1130/B26105.1><https://doi.org/10.1130/B26105.1>

1063 Montheil, L., Licht, A., Beard, K. C., Métais, G., Coster, P., Vaes, B., Donnadieu, Y., Pineau, E., Husson, L., & Dupont-Nivet,
1064 G. (2025ab). Across ancient oceans: Eocene dispersal routes of Asian terrestrial mammals to Europe, Afro-Arabia and South
1065 America. *Earth-Science Reviews*. <https://doi.org/10.1016/j.earscirev.2025.105352>

1066 Montheil, L., Licht, A., İbilioğlu, D., Botté, P., Ocakoğlu, F., Demory, F., Ruffet, G., Guihou, A., Kaya, M., Raynaud, B.,
1067 Akkiraz, M. S., Deschamps, P., Métais, G., Coster, P., & Beard, K. C. (2025ba). Updating the timeline of faunal endemism in
1068 Balkanatolia, the biogeographic province connecting Europe, Asia and Africa. *Journal of Asian Earth Sciences*, 290, 106661.
1069 <https://doi.org/10.1016/j.jseaes.2025.106661>

a mis en forme : Police :Italique

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :Italique

a mis en forme : Police :(Par défaut) Times New Roman

Montheil, L., Licht, A., Beard, K. C., Métais, G., Coster, P., Vaes, B., Donnadiou, Y., Pineau, E., Husson, L., & Dupont-Nivet, G. (2025ab). Across ancient oceans: Eocene dispersal routes of Asian terrestrial mammals to Europe, Afro-Arabia and South America. *Earth Science Reviews*. <https://doi.org/10.1016/j.earscirev.2025.105352>

Mueller, M. A., Licht, A., Campbell, C., Ocakoğlu, F., Akşit, G. G., Métais, G., Coster, P. M. C., Beard, K. C., & Taylor, M. H. (2022). Sedimentary Provenance From the Evolving Forearc-to-Foreland Central Sakarya Basin, Western Anatolia Reveals Multi-Phase Intercontinental Collision. *Geochemistry, Geophysics, Geosystems*, 23(3), e2021GC010232. <https://doi.org/10.1029/2021GC010232>

Page, M., Licht, A., Dupont-Nivet, G., Meijer, N., Barbolini, N., Hoon, C., Schauer, A., Huntington, K., Bajnai, D., Fiebig, J., Mulch, A., & Guo, Z. (2019). Synchronous cooling and decline in monsoonal rainfall in northeastern Tibet during the fall into the Oligocene icehouse. *Geology*, 47(3), 203-206. <https://doi.org/10.1130/G45480.1>

Palcu, D. V., & Krijgsman, W. (2023). The dire straits of Paratethys : Gateways to the anoxic giant of Eurasia. *Geological Society, London, Special Publications*, 523(1), 111-139. <https://doi.org/10.1144/SP523-2021-73>

Passey, B. H., Levin, N. E., Cerling, T. E., Brown, F. H., & Eiler, J. M. (2010). High-temperature environments of human evolution in East Africa based on bond ordering in paleosol carbonates. *Proceedings of the National Academy of Sciences*, 107(25), 11245-11249. <https://doi.org/10.1073/pnas.1001824107>

Peters, N. A., Huntington, K. W., & Hoke, G. D. (2013). Hot or not? Impact of seasonally variable soil carbonate formation on paleotemperature and O-isotope records from clumped isotope thermometry. *Earth and Planetary Science Letters*, 361, 208-218. <https://doi.org/10.1016/j.epsl.2012.10.024>

Philippe, É. G., St-Onge, G., Valet, J. P., Godbout, P. M., Egli, R., Francus, P., & Roy, M. (2023). Influence of Seasonal Post-Depositional Processes on the Remanent Magnetization in Varved Sediments From Glacial Lake Ojibway (Canada). *Geochemistry, Geophysics, Geosystems*, 24(3), e2022GC010707.

Popov, S.V., R`ogl, F., Rozanov, A.Y., Steininger, F.F., Shcherba, I.G., Kovac, M., 2004. Lithological-Paleogeographic Maps of Paratethys-10 Maps Late Eocene to Pliocene.

Pound, M. J., & Salzmann, U. (2017). Heterogeneity in global vegetation and terrestrial climate change during the late Eocene to early Oligocene transition. *Scientific Reports*, 7(1), 43386. <https://doi.org/10.1038/srep43386>

Quade, J., Eiler, J., Daëron, M., & Achyuthan, H. (2013). The clumped isotope geothermometer in soil and paleosol carbonate. *Geochimica et Cosmochimica Acta*, 105, 92-107. <https://doi.org/10.1016/j.gca.2012.11.031>

Raynaud, B., Akkiraz, M.S., Boura, A., Hoon, C., Gibson, M., Giobbini, A., Botté, P., Montheil, L., Kaya, M., Ocakoğlu, F., İbilioglu, D., Métais, G., Beard, K.C., Coster, P., Licht, A., 2026. Lutetian swamp-freshwater palynoflora from Bultu-Zile (central Anatolia, Türkiye): Implications for Eocene ecosystems of Balkanatolia. *Journal of Asian Earth Sciences* 300, 106980. <https://doi.org/10.1016/j.jseaes.2026.106980>

Raynaud, B., Akkiraz, M. S., Boura, A., Hoon, C., Gibson, M., Giobbini, A., Botté, P., Montheil, L., Kaya, M., Ocakoğlu, F., İbilioglu, D., Métais, G., Beard, K. C., Coster, P., & Licht, A. Lutetian swamp-freshwater palynoflora from Bultu-Zile (central Anatolia, Türkiye): implications for Eocene ecosystems of Balkanatolia. *Journal of Asian Earth Sciences (in Review)*.

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204)), Non Surlignage

a mis en forme : Police :Non Italique, Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204)), Non Surlignage

a mis en forme : Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204)), Non Surlignage

a mis en forme : Police :Non Italique, Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204)), Non Surlignage

a mis en forme : Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204)), Non Surlignage

a mis en forme : Soulignement , Couleur de police : Couleur personnalisée(RVB(17;85;204))

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Non souligné, Couleur de police : Automatique

Renaut RW, Gierlowski-Kordes EH, 2010. Lakes. Facies models 4: 541–575.

Robert, D., M.A. Birmingham, A. Licht, A. Olcott, A.-L. Jourdan, A. Guihou, Marina Suarez, R. Amiot, A. Vinçon-Laugier, P. Botté, F. Ocakoğlu, L. Montheil, M. Kaya, B. Raynaud, M.S. Akkiraz, P. Deschamps, G. Métais, P. Coster, & K.C. Beard (2026). Isotopic Constraints on the Ecology and Dispersal of Lutetian Embrithopods from Balkanatolia. *Palaaios*.

Sanders, W. J., Nemec, W., Aldinucci, M., Janbu, N. E., & Ghinassi, M. (2014). Latest evidence of *Palaeoamasia* (Mammalia, Embrithopoda) in Turkish Anatolia. *Journal of Vertebrate Paleontology*, 34(5), 1155–1164. <https://doi.org/10.1080/02724634.2014.850430>

Schemmel, F., Mikes, T., Rojay, B., & Mulch, A. (2013). The impact of topography on isotopes in precipitation across the Central Anatolian Plateau (Turkey). *American Journal of Science*, 313(2), 61–80. <https://doi.org/10.2475/02.2013.01>

Semmani, N., Fournier, F., Suc, J.-P., Fauquette, S., Séranne, M., Léonide, P., Marié, L., & Borgomano, J. (2024). Continental depositional record of climate and tectonic evolution around the Eocene-Oligocene transition in southeast France : Perspectives from the Vistrenque Basin (Camargue). *BSGF - Earth Sciences Bulletin*, 195, 8. <https://doi.org/10.1051/bsgf/2024005>

Sheldon, N. D. (2009). Nonmarine records of climatic change across the Eocene-Oligocene transition. In C. Koeberl & A. Montanari, *The Late Eocene Earth—Hothouse, Icehouse, and Impacts*. Geological Society of America. [https://doi.org/10.1130/2009.2452\(15\)](https://doi.org/10.1130/2009.2452(15))

Sheldon, N. D., Costa, E., Cabrera, L., & Garcés, M. (2012). Continental Climatic and Weathering Response to the Eocene-Oligocene Transition. *The Journal of Geology*, 120(2), 227–236. <https://doi.org/10.1086/663984>

Śliwińska, K. K., Coxall, H. K., Hutchinson, D. K., Liebrand, D., Schouten, S., & De Boer, A. M. (2023). Sea surface temperature evolution of the North Atlantic Ocean across the Eocene–Oligocene transition. *Climate of the Past*, 19(1), 123–140. <https://doi.org/10.5194/cp-19-123-2023>

[Speijer, R. P., Pälke, H., Hollis, C. J., Hooker, J. J., & Ogg, J. G. \(2020\). The paleogene period. In *Geologic time scale 2020* \(pp. 1087-1140\). Elsevier.](#)

[Stolper, D. A., & Eiler, J. M. \(2015\). The kinetics of solid-state isotope-exchange reactions for clumped isotopes: A study of inorganic calcites and apatites from natural and experimental samples. *American Journal of Science*, 315\(5\), 363-411.](#)

Suc, J.-P., Fauquette, S., Popescu, S.-M., Melinte-Dobrinescu, M. C., Huet, B., Sorrel, P., Semmani, N., Fournier, F., Bessedik, M., Zheng, Z., Truc, G., Ferry, S., Rubino, J.-L., Girard, V., & Séranne, M. (2025). Late Eocene to Early Miocene environments, vegetation and climate in Southeastern France documented by palynology. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 659, 112625. <https://doi.org/10.1016/j.palaeo.2024.112625>

Sun, Y., Liu, J., Liang, Y., Ji, J., Liu, W., Aitchison, J. C., Sun, J., Lu, J., Song, B., Xu, Y., Zhang, K., & Liu, Z. (2020). Cenozoic moisture fluctuations on the northeastern Tibetan Plateau and association with global climatic conditions. *Journal of Asian Earth Sciences*, 200, 104490. <https://doi.org/10.1016/j.jseas.2020.104490>

Tardif, D., Toumoulin, A., Fluteau, F., Donnadieu, Y., Le Hir, G., Barbolini, N., Licht, A., Ladant, J.-B., Sepulchre, P., Viovy, N., Hoorn, C., & Dupont-Nivet, G. (2021). Orbital variations as a major driver of climate and biome distribution during the greenhouse to icehouse transition. *Science Advances*, 7(43), eabh2819. <https://doi.org/10.1126/sciadv.abh2819>

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman, Italique

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman, Italique

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Couleur de police : Couleur personnalisée(RVB(34;34;34)), Surlignage

1138 [Tauxe, L., with contributions from Butler, R.F., Van der Voo, R., and Banerjee, S.K., Essentials of Paleomagnetism, University](#)
1139 [of California Press, ISBN: 9780520260313 \(2010\)](#)
1140 [Tauxe, L. Essentials of Paleomagnetism, 2010. University of California press,](#)
1141 [Tauxe, L., Watson, G.S. The fold test: an eigen analysis approach Earth and Planetary Science Letters 122 \(3-4\), pp. 331-341](#)
1142 [\(1994\).](#)
1143 [Tauxe, L., Kodama, K., & Kent, D. V. Testing corrections for paleomagnetic inclination error in sedimentary rocks: a](#)
1144 [comparative approach. Phys. Earth Planet. Int., 169, 152–165. \(2008\)](#)
1145 Teodoridis, V., & Kvaček, Z. (2015). Palaeoenvironmental evaluation of Cainozoic plant assemblages from the Bohemian
1146 Massif (Czech Republic) and adjacent Germany. *Bulletin of Geosciences*, 695-720. <https://doi.org/10.3140/bull.geosci.1553>
1147 The Cenozoic CO Proxy Integration Project (CenCOPIP) Consortium*†, Hönlisch, B., Royer, D. L., Breecker, D. O., Polissar,
1148 P. J., Bowen, G. J., Henahan, M. J., Cui, Y., Steinthorsdottir, M., McElwain, J. C., Kohn, M. J., Pearson, A., Phelps, S. R.,
1149 Uno, K. T., Ridgwell, A., Anagnostou, E., Auermann, J., Badger, M. P. S., Barclay, R. S., ... Zhang, L. (2023). Toward a
1150 Cenozoic history of atmospheric CO₂. *Science*, 382(6675), eadi5177. <https://doi.org/10.1126/science.adi5177>
1151 Tindall, J., Flecker, R., Valdes, P., Schmidt, D. N., Markwick, P., & Harris, J. (2010). Modelling the oxygen isotope
1152 distribution of ancient seawater using a coupled ocean–atmosphere GCM: Implications for reconstructing early Eocene
1153 climate. *Earth and Planetary Science Letters*, 292(3-4), 265-273.
1154 <https://doi.org/10.1016/j.epsl.2009.12.049><https://doi.org/10.1016/j.epsl.2009.12.049>
1155 Tissier, J., Becker, D., Codrea, V.A., Costeur, L., Fărcaș, C., Solomon, A., Venczel, M., and Maridet, O. 2018. New data on
1156 *Amynodontidae* (Mammalia, Perissodactyla) from Eastern Europe: Phylogenetic and palaeobiogeographic implications around
1157 *the Eocene–Oligocene transition. PLOS ONE 13: e0193774.*
1158 Toumoulin, A., Tardif, D., Donnadiou, Y., Licht, A., Ladant, J.-B., Kunzmann, L., & Dupont-Nivet, G. (2022). Evolution of
1159 continental temperature seasonality from the Eocene greenhouse to the Oligocene icehouse –a model–data comparison.
1160 *Climate of the Past*, 18(2), 341-362. <https://doi.org/10.5194/cp-18-341-2022>
1161 Tremblin, M., Hermoso, M., & Minoletti, F. (2016). Equatorial heat accumulation as a long-term trigger of permanent
1162 Antarctic ice sheets during the Cenozoic. *Proceedings of the National Academy of Sciences*, 113(42), 11782-11787.
1163 <https://doi.org/10.1073/pnas.1608100113><https://doi.org/10.1073/pnas.1608100113>
1164 [Van de Weerd, A. A., de Bruijn, H., & Wessels, W. \(2023\). A small assemblage of early Oligocene rodents and insectivores](#)
1165 [from the Sivas basin, Turkey. Palaeobiodiversity and Palaeoenvironments, 103\(3\), 609-632.](#)
1166 Van Hinsbergen, D. J. J., Kaymakci, N., Spakman, W., & Torsvik, T. H. (2010). Reconciling the geological history of western
1167 Turkey with plate circuits and mantle tomography. *Earth and Planetary Science Letters*, 297(3-4), 674-686.
1168 <https://doi.org/10.1016/j.epsl.2010.07.024>
1169 Van Hinsbergen, D. J. J., Maffione, M., Plunder, A., Kaymakci, N., Ganerod, M., Hendriks, B. W. H., Corfu, F., Gürer, D.,
1170 De Gelder, G. I. N. O., Peters, K., McPhee, P. J., Brouwer, F. M., Advokaat, E. L., & Vissers, R. L. M. (2016). Tectonic

- a mis en forme : Police :10 pt
- a mis en forme : Espace Après : 0 pt, Interligne : 1.5 ligne, Bordure : Haut: (Pas de bordure), Bas: (Pas de bordure), Gauche: (Pas de bordure), Droite: (Pas de bordure), Entre : (Pas de bordure), Motif : Transparente
- a mis en forme : Police :10 pt, Couleur de police : Automatique
- a mis en forme : Police :10 pt
- a mis en forme : Police :10 pt, Couleur de police : Automatique
- a mis en forme : Police :10 pt
- a mis en forme : Police :10 pt, Couleur de police : Automatique
- a mis en forme : Police : (Par défaut) Times New Roman
- a mis en forme : Police : (Par défaut) Times New Roman
- a mis en forme : Police : (Par défaut) Times New Roman
- a mis en forme : Police : (Par défaut) Times New Roman
- a mis en forme : Police :Italique
- a mis en forme : Police : (Par défaut) Times New Roman
- a mis en forme : Police : (Par défaut) Times New Roman
- a mis en forme : Surlignage
- a mis en forme : Police :Italique
- a mis en forme : Police :Italique
- a mis en forme : Police : (Par défaut) Times New Roman
- a mis en forme : Police : (Par défaut) Times New Roman

1171 evolution and paleogeography of the Kırşehir Block and the Central Anatolian Ophiolites, Turkey : TECTONIC EVOLUTION
1172 OF CENTRAL ANATOLIA. *Tectonics*, 35(4), 983-1014. <https://doi.org/10.1002/2015TC004018>
1173 Van Hinsbergen, D. J. J., Torsvik, T. H., Schmid, S. M., Mañenco, L. C., Maffione, M., Vissers, R. L. M., Gürer, D., &
1174 Spakman, W. (2020). Orogenic architecture of the Mediterranean region and kinematic reconstruction of its tectonic evolution
1175 since the Triassic. *Gondwana Research*, 81, 79-229. <https://doi.org/10.1016/j.gr.2019.07.009>
1176 Wang, X., Carrapa, B., Sun, Y., Dettman, D. L., Chapman, J. B., Caves Rugenstein, J. K., Clementz, M. T., DeCelles, P. G.,
1177 Wang, M., Chen, J., Quade, J., Wang, F., Li, Z., Oimuhammadzoda, I., Gadoev, M., Lohmann, G., Zhang, X., & Chen, F.
1178 (2020). The role of the westerlies and orography in Asian hydroclimate since the late Oligocene. *Geology*, 48(7), 728-732.
1179 <https://doi.org/10.1130/G47400.1>
1180 Weppe, R., Condamine, F. L., Guinot, G., Maugoust, J., & Orliac, M. J. (2023). Drivers of the artiodactyl turnover in insular
1181 western Europe at the Eocene–Oligocene Transition. *Proceedings of the National Academy of Sciences*, 120(52),
1182 e2309945120. <https://doi.org/10.1073/pnas.2309945120>
1183 Westerhold, T., Marwan, N., Drury, A. J., Liebrand, D., Agnini, C., Anagnostou, E., Barnet, J. S. K., Bohaty, S. M., De
1184 Vleeschouwer, D., Florindo, F., Frederichs, T., Hodell, D. A., Holbourn, A. E., Kroon, D., Lauretano, V., Littler, K., Lourens,
1185 L. J., Lyle, M., Pälike, H., ... Zachos, J. C. (2020). An astronomically dated record of Earth's climate and its predictability
1186 over the last 66 million years. *Science*, 369(6509), 1383-1387. <https://doi.org/10.1126/science.aba6853>
1187 Yang, Y., Fang, X., Han, W., Wang, Y., & Galy, A. (2022). Terrestrial carbonate oxygen isotopes constraints on the interplay
1188 between westerlies and monsoonal rains modulating the Cenozoic climate on the northeastern Tibetan Plateau.
1189 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 608, 111289. <https://doi.org/10.1016/j.palaeo.2022.111289>
1190 Zachos, J. C., Quinn, T. M., & Salamy, K. A. (1996). High-resolution (10⁴ years) deep-sea foraminiferal stable isotope records
1191 of the Eocene-Oligocene climate transition. *Paleoceanography*, 11(3), 251-266. <https://doi.org/10.1029/96PA00571>
1192 [Zijderveld, J.D.A. \(1967\): A.C. demagnetization of rocks: analysis of results. In D.W. Collinson, K.M. Crees and S.K. Runcorn](#)
1193 [\(Eds.\) Methods in Paleomagnetism \(pp. 254-286\). Elsevier, Amsterdam,](#)

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :(Par défaut) Times New Roman

a mis en forme : Police :10 pt

a mis en forme : Espace Après : 0 pt, Interligne : 1.5
ligne, Bordure : Haut: (Pas de bordure), Bas: (Pas de
bordure), Gauche: (Pas de bordure), Droite: (Pas de
bordure), Entre : (Pas de bordure), Motif : Transparente

a mis en forme : Police :10 pt, Couleur de police :
Automatique