# Retreating glaciers and snow cover are amplifying summer droughts in the upper Adige River Basin (Italy)

Susen Shrestha[1,2], Stefano Terzi[1*], Davide Zoccatelli[4], Mattia Zaramella[2], Marco Borga[2], Andrea Galletti[1], Mattia Callegari[5], Roberto Dinale[6], Massimiliano Pittore[1], Giacomo Bertoldi[3]

[1]Eurac Research, Center for Climate Change and Transformation (CCT), Bolzano, Italy
[2]Department of Land, Environment, Agriculture and Forestry, University of Padova, Italy
[3]Eurac Research, Institute for Alpine Environment, Bolzano, Italy
[4]Luxembourg Institute of Science and Technology (LIST), Luxembourg
[5]Eurac Research, Institute for Earth Observation, Bolzano, Italy
[6]Ufficio Dighe e Idrologia, Provincia Autonoma di Bolzano, Italy

*Correspondence to:* Stefano Terzi (stefano.terzi@eurac.edu)

**Abstract.**

Snow and glacier meltwater play a critical role in sustaining summer streamflow in mountains and downstream regions. Yet, understanding glaciers' contributions to buffer river streamflow during droughts remains limited and pose major barriers to improve present and future water management within the context of climate change.

This study evaluates the contribution of glacier melt to summer flows and its mitigation effect of hydrological droughts in the upper Adige River basin, (Italy). We developed and implemented a new dynamic glacier module into the ICHYMOD-TOPMELT hydrological model, annually updating glacier area and improving the quantification of meltwater contributions under progressive glacier retreat (from 111 km2 in 1997 to 79 km2 in 2017).

The hydrological model exhibited robust performances (KGE=0.89 for 1997-2022 for the whole upper Adige River basin at the Bronzolo closing station; 0.88 in summers) capturing observed glacier area, mass balance, and seasonal melt trends. Results show that glacier melt in the upper Adige River basin contributed to an average of 4.5% to total summer streamflow during 1997–2022, with significant spatial variability and reaching 30% in glacierized subbasins. During the severe droughts of 2003, 2005 and 2022, glacier melt contributions ranged

between 4 and 12% at the upper Adige closing section. In 2003, high temperatures and limited Snow Water Equivalent (SWE) led to glacier melt accounting for 11.67% of summer flows. By contrast, the significant runoff deficit in 2005 was driven by colder temperatures, ~~-~~which ~~in 2005~~ reduced glacier melt contributions to 4.85%, combined ~~-compounding~~ with low SWE conditions ~~and leading to a significative runoff deficit~~. In 2022, the combination of low precipitation, low snow cover and high temperature drove glacier melt. ~~Differently~~ In contrast to ~~than~~ the 2003 drought, reduced glacier areas led to lower absolute contributions (8.17%).

Our findings reveal that despite the increased melt rates in recent warm years, retreating glacier areas have reduced their absolute buffering effect. Glacier retreat is weakening the~~ir~~ contribution of glaciers to summer flows, increasing the upper Adige River basin's dependence on precipitation and snowmelt, the latter of which ~~which~~ is also showing a decreasing trend. For these reasons, accounting for dynamic glacier and snow changes is essential to improve future drought projections and inform adaptive water management in glacier-fed basins within the context of climate and anthropogenic changes.

**Keywords:** Glacier melt, Adige River basin, droughts, hydrological modelling, climate change~~water scarcity~~**.**

## 1 Introduction

Recent droughts in Europe caused widespread socio-ecological impacts in the Italian Alps and downstream areas (Avanzi et al., 2026, 2024; Biella et al., 2025; Montanari et al., 2023). ~~The unprecedented 2022 droughts in Europe was one of the worst event ever recorded for large parts of Europe caused widespreadwith severe socio-ecological impacts in the Italian Alps and related downstream areas~~ T~~.~~

~~The combination of below-average snowfall in winter, limited rainfall and exceptionally high temperature during spring and summer led to widespread drought impacts across multiple sectors (Avanzi et al., 2025, 2024). In those areas where glaciers are still present, t~~he increase in~~in~~ meltwater from snow and glaciers temporarily and partially offset the lack of water from below average snowfall in winter, limited rainfall and exceptionally high temperatures during spring and summer ~~some of the summer drought impacts~~ (Leone et al., 2026).

However, rising temperature~~s increase~~ and ~~variation~~ changes ~~of~~ in precipitation patterns related to climate change are progressively shrinking ~~glaciers~~ and pushing glaciers ~~them~~ to a point of no return (Hock et al., 2022; Pepin et al., 2022; Van Tricht et al., 2026). From 2000 to 2023, 39% of the European glacier~~s~~ mass was already lost (The GlaMBIE Team, 2025) and according to the IPCC projections most of the glaciers in the European Alps are expected to disappear by 2100 (Hock et al., 2022). Of particular concerns are the potential consequences of Alpine

glacier~~s~~ recession~~s~~ on water provisioning to downstream areas, both on the long term and during drought events, when water is most needed (Ultee et al., 2022).

Yet, understanding ~~the~~ glaciers' contributions to buffer~~ing~~ river streamflow during droughts remains limited and poses a major barrier~~s~~ to improv~~inge~~ present and future water management. On the one hand, this limitation is mainly due to difficulties ~~to~~ in dynamically captur~~inge~~ glacier evolution~~s~~ within hydrological models, ~~finding~~ requiring a trade-off ~~among~~ between the limited data availability on glaciers, high model~~s~~ parametrization, and computational costs (Seibert et al., 2018). ~~.~~

In particular, models often rely on either highly paramet~~e~~rized energy-balance equations (Frans et al., 2016; Naz et al., 2014), fully distributed (Maussion et al., 2019) or data-intensive statistical approaches (Shen et al., 2023), which can lead to high computational cost~~ly~~ and/or ~~difficult~~ limited interpretability o~~fn~~ glacier dynamics and their contributions within hydrological models. Indeed, existing applications have increasingly integrated glacier~~s~~ evolution schemes into hydrological models to simulate mass balance, including equilibrium-line altitude shifts (Linsbauer et al., 2013), volume-area scaling (Luo et al., 2013; Radić and Hock, 2010; Stahl et al., 2008) and the Δh parameterization (Huss et al., 2010, 2008).

On the other hand, existing applications integrating glacier~~s~~ changes into semi-distributed hydrological models (Li et al., 2015; Seibert et al., 2018) often ~~do not~~neither consider dynamic glacier evolution ~~and do not~~nor focus on glacier contributions during droughts. Therefore, there is ~~the~~ a need for a simplified, yet reliable glacier dynamic representation in semi-distributed hydrological models to improve the understanding of the hydro climatic conditions (temperature, precipitation and snow cover anomalies) affecting glacier melt buffering contributions to summer droughts.

The critical 2022 drought situation for the Italian Alps calls for improving our understanding o~~fn~~ how much, where and when glaciers contributed to mitigat~~inge~~ drought impacts. The upper Adige River Basin in the Italian Alps represents a case in point (Figure 1). Even though it has been considered an area with plenty of ~~available~~ water ~~coming~~ from accumulated snow and glacier melt sustaining summer streamflow and supporting extensive~~the high~~ water uses ~~from~~ across multiple sectors (Lemus-Canovas et al., 2025; Zebisch et al., 2025), drought events in 2015, 2017 and~~,~~ 2022 challenged water management and have been raising concerns ~~on~~ about the high sensitivity of the basin to droughts in the context of ever reducing contributions from snow and glacier melt (Chiogna et al., 2015; Colombo et al., 2023; Stephan et al., 2021).

For these reasons, our research aims to improve long-term predictions of glacier meltwater contributions to streamflow during droughts in the upper Adige River Basin. Moreover, we propose a modelling approach that can

be applied to other glacierized catchments to evaluate the effect of glacier melt in mitigating hydrological droughts under ~~retreating glaciers~~ conditions of glacier retreat.

To do so, we integrated a novel dynamic glacier module within the existing ICHYMOD-TOPMELT semi-distributed hydrological model (Norbiato et al., 2009, 2008) that annually updates glacier extent using a volume-area scaling relationship to better characterize in space and time glacier melt contributions to streamflow conditions in the upper part of the Adige River Basin (sections 2.2, 3.1 and 3.2). We focused on major drought years including the unprecedented 2022, investigating the role of multiple hydro-climatic conditions (temperature, precipitation and snow cover anomalies) in contributing to droughts (sections 2.5). By doing so, we estimated glacier contributions during critical conditions hence highlighting potential ~~cases of~~ vulnerabilities associated with ~~to~~ future glacier retreat~~s~~ (section 3.3). Finally, we investigated a counterfactual scenario considering the 1997 glacier extent~~sion~~ during the 2022 drought to estimate ~~their~~ its potential buffering effects during extreme conditions (section 3.3.3).

## 2 Material and methods

### 2.1 Study area and input data

The Adige River flows from the eastern Italian Alps to the Adriatic Sea for 410 km, making it the second longest river in Italy with an overall drainage area of 12200 km$^2$ (Lemus-Canovas et al., 2025; Navarro-Ortega et al., 2015). The upper part of the Adige River Basin (6924 km$^2$, Figure 1) is characterized by different climatic conditions, with a semi-continental climate in low elevation areas with mild winters and warm summers, and an alpine climate in higher elevations with cold winters and summers (Bertoldi et al., 2023; Tscholl et al., 2025). At higher altitudes, significant water resources accumulate during the winter season in the form of snowfall, which are mobilised during spring. This situation determines the alpine nivo-glacial hydrological regime, characterised by two peaks, one in spring due to snow and glacier melt and the other in autumn due to cyclonic storms (Larsen et al., 2021), which provide abundant water for the development of various and intensive anthropogenic activities. The large availability of water shapes these areas, ranging from reservoirs, hydropower plants and managed pastures at high elevation to irrigated apple orchards and vineyards intensively cultivated at valley bottoms.

Hydrological alterations are mainly induced by various artificial dams with the three largest forming the Lake Resia ~~lake~~ (116 Mm$^3$), Lake Vernago ~~lake~~ (44 Mm$^3$) and Lake Zoccolo ~~lake~~ (34 Mm$^3$) storing water which is mainly used for hydropower production. Most of the lakes receive water from upstream glaciers, the extent~~sion~~ of which was mapped to the decreasing values of 111, 98 and 79 km$^2$ in~~during~~ 1997, 2005 and 2017 (Galos et al.,

(2022), further ~~data on~~ annual mass balance data for the whole Adige River Basin are available from Dussaillant et al., (2025)). The study area includes a dense network of hydro-meteorological stations, comprising 88 rain gauges (~~equivalent~~ approximately to 1 per 72 km2), 124 temperature gauges (approximately 1 per 55 km2) and 50 streamflow gauges (1 per 23 km of river), with hourly time series of precipitation, temperature and streamflow, most of them available from the ~~Civil Protection Agency of the province of Bolzano~~Civil Protection Agency of the province of Bolzano (https://meteo.provincia.bz.it/download-dati.asp).





**Figure 1 - Map of the upper part of the Adige River Basin (Italy) with glacier extension (Galos et al., 2022) precipitation and river gauge stations, the 16 sub-basins used in the hydrological modelling ~~as well as~~and the digital elevation model (NASA Shuttle Radar Topography Mission (SRTM) Global. Distributed by OpenTopography, 2013) as a background layer.**

## 2.2 Hydrological model

The simulation of water availabilitye in the upper part of the Adige River Basin starts from the combination of the hydrological model ICHYMOD (Norbiato et al., 2008, Norbiato et al., 2009), with the TOPMELT snow module (Zaramella et al., 2019). The resulting ICHYMOD-TOPMELT is a computationally efficient, semi-

distributed conceptual hydrological model that simulates hydro-meteorological state variables (i.e., precipitation, temperature, Snow Water Equivalent (SWE), glacier melt and runoff) at hourly resolution while representing spatial variability in snow and ice melt within subbasins.

The model has been operationally employed by the Province of Bolzano ~~e~~Civil Pprotection Oeffice for flood forecasting, and previous extensive model validation in the same region has been performed for runoff production (Dalla Torre et al., 2024; Stergiadi et al., 2020), snow and ice melt modelling (Di Marco et al., 2021).

Runoff generation, soil moisture, and evapotranspiration processes are lumped at the sub-catchment scale using the Probability Distributed soil Model (PDM, ~~—~~ (Moore, (2007, 1985)~~Moore, 2007, Moore, 1985~~). Catchment outflows include direct runoff, baseflow, and evapotranspiration (ET), which is estimated using the Hargreaves method and adjusted for soil moisture availability.

In order to properly model snow and ice accumulation and melt processes in complex terrain, ~~we divided~~ the subbasins have been subdivided into~~by~~ elevation and radiation classes, which are updated monthly using clear-sky potential solar radiation. Elevation drives ~~a~~ gradients in precipitation and temperature, using regionally calibrated lapse rates. ~~We calculated~~ Ssnow and glacier processes are calculated independently ~~on~~ for each combination of elevation and radiation using a modified temperature-index model based on the snowpack formulation of Cazorzi and Dalla Fontana, (1996).

Snow and ice melt are estimated from both radiative and thermal energy exchanges, with separate formulations for daytime, nighttime, and rainy conditions (Zaramella et al., 2019). Snow and ice differ in their reflectivity: snow albedo decreases over time as the surface ages and warms, following Brock et al., (2000), whereas ice maintains a ~~higher~~ lower, fixed albedo that enhances radiation absorption and accelerates melt. In areas covered by glaciers, glacier ice melt begins when snow water equivalent in a radiation–elevation band falls below a threshold value of 5 mm as defined by (Zaramella et al., 2019). Moreover, we updated the snow-to-ice transformation for each elevation and energy band following a 1:1 volumetric conversion of available SWE to ice depth. Finally, SWE and glacier melt values, based on the elevation and radiation classes for each subbasin, were aggregated to obtain basin-scale ~~have an overall~~ values.

### 2.2.1 The new g~~G~~lacier dynamics module

We developed the simulation of glaciers area dynamics over time and integrated it as a new module into the semi-distributed hydrological model ICHYMOD (Norbiato et al., 2009, 2008). At the foundation of the glacier module is the relationship between glacier volume and area, which was applied at the subbasin level, with the annual hydrological balance used to dynamically update ~~the~~ glacier volume.

We used the glacier outline ~~of~~ from 1997 and the Digital Elevation Model (DEM) from 1994 to create the initial ice thickness map. This ~~is~~ was achieved ~~done~~ by applying the flow law described by (Glen, 1955) and Farinotti et al., (2009), relying only on glacier surface topography. We divided each glacier into $i$ contour lines, where the specific ice flux $\overline{q}_i$ was~~is~~ estimated based on their distance from the equilibrium-line altitude (ELA), following the approach of Huss and Farinotti, (2012). The ice thickness $h_i$ is then calculated as:

$$h_i = \sqrt[n+2]{\frac{\overline{q}_i}{2A}\frac{(n+2)}{(C\rho g \sin\overline{\alpha})^n}}$$

Where $A$ is a flow rate factor calibrated over the whole region ($2.5*10^{-8}$), $C$ is a correction factor for basal sliding, $\rho$ is the ice density, $g$ is the gravitational acceleration, $\overline{\alpha}$ the mean slope in a direction perpendicular to the contour line and $n$ is 3 as standard value adopted in glacier modelling following (Farinotti et al., 2009; Huss and Farinotti, 2012). The ice thickness is then smoothed across contour lines, corrected for local surface slope, distributed using a triangular profile along the contour line, and interpolated across the original grid resolution. The glacier area is reported for each ~~unit of~~ elevation unit and time-invariant radiation classes, assumed representative for the melt period, that form the basis ~~are at the base~~ of ICHYMOD-TOPMELT. We validated the results for the~~on~~ initial glacier depth by comparing them with the 1997 South Tyrol glacier inventory, which estimates a total glacier volume of 3326 million $m^3$ (Galos et al., 2015; Knoll and Kerschner, 2009).

The initial ice thickness is reduced gradually using the $\Delta h$ methodology developed by Huss et al., (2010):

$$\Delta h = (h_r - 0.3)^2 + 0.6(h_r - 0.3) + 0.09$$

Where $\Delta h$ is the normalized variation of ice thickness and $h_r$ is the normalized elevation range as in Huss et al., (2010). The parameters used are those recommended for small glaciers, as most of the glaciers have an area below 5 $km^2$. Using this method, we gradually reduced glacier~~s'~~ volume while updating glacier~~their~~ 3D geometry, creating a relationship between glacier volume i~~o~~n each basin and area within ~~on~~ each class of time-invariant elevation and radiation. At the end of each hydrological year, we computed the glacier~~al~~ balance for each basin, which is the net balance between the glacier melt and the snow accumulation~~ed~~ o~~ver~~ the glacier area, assuming that any snow remaining on the glacier at the end of the year directly transforms into ice. We updated the previous-year glacier volume with the glacier~~al~~ balance. For each ~~unit of~~ elevation and energy class, we calculated the new glacier area by linear interpolation between the available steps in the volume-area relationship. Finally, we calculated the glacier volume and area for the whole study basin by summing up all individual subbasins.

### 2.3 Simulation setup

Following the semi-distributed approach of the ICHYMOD-TOPMELT model, we performed hydrological simulations considering 615 hydrological response units (hereafter referred to as subbasins) within the upper Adige River Basin. For each subbasin, we computed the vertical water mass balance before routing the resulting flows along the river network. From a temporal point of view, we considered the hydrological year from October to September of the following year. In order to effectively start our assessment in October 1997 (year of the first available glacier area observation), we raun the model at 1-hour time steps from 1996 until 2022, using the first year as a spin-up period and hence discarding it from subsequent analyses.

The parametrization of the hydrological kernel of ICHYMOD-TOPMELT was obtained by maximizing NSE at 16 stream gauges over the period 2003–2019. As the simulations of glacier dynamics wereas highly sensitive to the glacier albedo parameter, we calibrated it through a sensitivity analysis (by testing values the rangingе from 0.4 to 0.8, consistent with those reported in studies conducted in similar areas, e.g., (Hartl et al., (2025); Krampe et al., (2022); Richter et al., (2026)). We then selected the albedo value that provided the, assessing what value ensured the closest match to the available glacier area observations over the entire simulation span (1997-2022). Finally, an additional simulation considered the same glacier area, as it was recorded in 1997, for the whole simulation period from 1997 until to 2022. By doing so, we aimed to better understand how the drought severity would have been different due to changes in glacier area that occurred over in the last 26 years.

### 2.4 Predictions performance metrics

We considered the Kling-Gupta Efficiency (KGE), as defined by Gupta et al., (2009), tofor evaluateing the hydrological model performance over the entire simulation period (1997-2022). To quantify the benefits of shifting from static to dynamic glacier modelling, we considered percentage changes in KGE as well as in the mean bias ratio (MBR) of modelled runoff, for both for the entirewhole study period and for the selected summer months of 2003, 2005 and 2022. We evaluated the glacier module performance of the glacier module against available observations of glacier area, volume and annual mass balance. In addition, we validated the modelled cumulative mass balance against the annual glacier mass balance dataset of Dussaillant et al., (2025) for the entire study area, covering the period from 1997 onwards.

### 2.5 Hydro-climatic drivers of droughts

For the analysis and characterization of past hydrological droughts occurrence and evolution, we considered a set of hydro-climatic drivers describing the physical conditions that led to droughts. The drivers were implemented and analysed both spatially on the whole upper Adige River Basin and temporally over multiple years and seasons, with a focus on the summer season. In addition, the analysis focused on the 2003, 2005 and 2022 ~~most severe~~ drought years, ~~-~~which were selected as the most severe based on the lowest summer Streamflow Standardized Index (SSI, Modarres, 2007; Telesca et al., 2012), computed for ~~with reference to~~to the July-August-September months (JAS) at the Bronzolo closing section (further information is provided in the S~~s~~upplementary M~~m~~aterial). The hydro-climatic drivers analysed for past major drought events were precipitation, temperature, SWE at the end of June (assumed to be a proxy for summer snow coverage and therefore for the ~~the snow budget resulting from winter-spring prior to the melting season~~ amount of ice melt contributing to summer runoff (Penna et al., 2017)), glacier melt and runoff (at the basin outlet). In particular, for each of the ~~outlet of the~~ 615 ~~subbasins~~ outlets, we computed the area-weighted averages of all the hydro-climatic variables as:

$$Variables_i^{weighted} = \frac{\sum_{k=1}^{i}(Variables_k \times Area_k)}{\sum_{k=1}^{i} Area_k}$$

Where $i$ is the given subbasin number while $k$ considers all the subbasins nested within subbasin $i$. By doing so, we aggregated ~~and made all~~the variables and made them comparable according to the runoff routing pathway. Further, we analysed multiple hydro-climatic drivers (total precipitation, temperature anomalies, and SWE at the end of June) and their z-score values for all ~~the~~ modelled years from 1996 to 2023. In particular, the z-score metric (Kreyszig et al., 2011), evaluates how many standard deviations a data point is from the mean of its own dataset, effectively capturing whether~~if their~~ values are above or below the climatological mean for each month (positive and negative z-scores, respectively). The z~~Z~~-score provides a standardized interannual comparison of precipitation, temperature, SWE, runoff and glacier melt patterns and captures drought patterns in relation to glacier melt conditions during summer months (i.e., July, August and September).

Finally, we provide spatially distributed representations of summer climatologies as well as focused~~s~~ analyses of~~on~~ the specific droughts in 2003, 2005 and 2022. We reported precipitation, SWE, runoff and glacier melt contributions as percentage differences from the climatological mean, and temperature as an absolute difference~~s~~.


## 3 Results

### 3.1 Glacier module performance

The dynamic glacier module was initialized with the 1997 glacier area observation, iteratively updated and run until 2022 (Figure 2). The calibration of the glacier albedo parameter resulted in a value of 0.65 as it achieved an overall $R^2$ of 0.84 between the simulated glacier mass balance and the median value in Dussaillant et al., (2025) glacier mass balance observations (bottom panel in Figure 2), while ensuring a close fit to the 2005 and 2017 glacier area observations (top panel in Figure 2). ~~was calibrated to best fit the available glacier area observations for 2005 and 2017. Since a~~highlighting that albedo is a key calibration parameter in the model~~, we performed a~~ The ~~sensitiviy~~sensitivity ~~analnsiys~~analysis~~-~~ on the ~~considering a range between 0.4 and 0.8 on the basis of the glacier albedo values present in the (REF). W~~The ~~resulting value of 0.65 achieved an overall $R^2$ equal to 0.84 between the simulated glacier mass balance and the median value in Dussaillant et al., (2025) glacier mass balance observations (bottom panel in Figure 2), while ensuring a close fit to the 2005 and 2017 glacier area observations (top panel in Figure 2). Figure 2 also shows the uncertainty band for the 0.4-0.8 albedo values highlighting how albedo is a key calibration parameter of the model. The sensitivity analysis show how changing the~~ albedo ~~from 0.4 to 0.8~~ range 0.4-0.8 (grey band in Figure 2) shows potential ~~could imply a~~ glacier area reduction going from 28% to 49% and mass loss ranging from 15 to 27 m w.e. ~~durng~~at the end of the 26-year study period.

~~We selected the 0.65 led to glacier area and mass balance closest to the observed data: three available glacier measurements and the Dussaillant dataset estimate of glacier mass (upper and lower panel in ).~~

The simulated glacier values showed a generalized reduction in~~of~~ both glacier area (-48.93 $km^2$) and mass (-1.99 $km^3$) over the 26 years of simulation, corresponding to a mean areal reduction rate of 1.95 $km^2$/year and mean mass reduction of 0.079 $km^3$/year, with only a few years showing ~~except a few cases of~~ positive glacier mass accumulation in the basin, such as ~~in~~ 2000/2001.


Glacier Area (km²)
Albedo Range (0.40–0.80)
Albedo: 0.65
Observation
R² (Dussaillant median & albedo 0.65) = 0.838
Cum. Mass Balance (m w.e.)
Albedo Range (0.40–0.80)
Albedo: 0.65
Dussaillant Range
Dussaillant Median
Year

260





**Figure 2 - Time series of glacier area (top panel) and mass balance (bottom panel) from 1997 to 2022 with modelled values (red line) and related uncertainty band (grey area), different albedo values from 0.4 to 0.8 (light blue area), glacier area observations (Galos et al., (2022), black dots) and annual glacier mass changes from Dussaillant et al., (2025, black line).**

### 3.2 Integrated hydrological model results

The hydrological model simulations considered both ~~a~~ dynamic and static glacier module~~s~~. We compared their results to evaluate ~~the~~ glaciers contributions at the subbasins level, over the simulation period and especially during specific drought events. When integrating the dynamic glacier module, the hydrological model showed KGE values ranging from 0.40 to 0.86 for all the glaciated basins, with an average value of 0.70 (Figure 3).





**Figure 3 – The Kling-Gupta Efficiency (KGE) performance metric (left y-axis) for the streamflow simulation for the whole period (1998-2022), integrating a dynamic glacier module (light blue dots) and a static glacier module (blue dots) according to increasing values of glacier area over the each subbasin area on the x-axis. The bar plot represents KGE variation (%, right y-axis) between the static and dynamic ~~two~~ glacier modelling ~~setups~~modules, with specific reference to the summer droughts of 2003, 2005 and 2022 and darker red colour the greater the model improvements.**

Over the entire simulation period, the differences between the integration of the dynamic or static glacier modules are limited. For three subbasins, the overall KGE performance has slightly increased, with values below +0.01, while for others it slightly decreased~~s~~, by less than -0.006. The latter subbasins with declining KGE values for the dynamic module are those showing a negative runoff bias for simulations with the static glacier module (see ~~more information in~~ Tables S3~~2~~s and S4~~3~~s in~~of~~ the Ssupplementary M~~m~~aterial). While the dynamic glacier module improved the representation of reduced glacier area conditions, it also implies a decrease in~~of~~ the glacier melt contribution, thereby~~and hence~~ increasing the already negative runoff bias, ultimately leading to a decrease of the KGE.

These results become even more pronounced when focusing on the specific summer months (JAS) of drought events in 2003, 2005 and 2022. During that time, streamflow reached its lowest observed levels over the simulated period, hence making the glacial meltwater contribution proportionally more relevant. During the three considered

droughts, the simulations with the dynamic glacier module showed improved KGE performance for all glacierized subbasins, except for two, as depicted in the bar plot of Figure 3. In particular, the largest increases were observed in subbasin n.11 (Passirio, +21.7%), n.5 (Rio Riva a Caminata, +16.5%) and n.13 (Aurino a S.Giorgio, +9.8%), while a slight KGE decrease was recorded for the subbasins n.8 (Aurino a Cadipietra, -0.87%) and n.2 (Rio Riva a Seghe, -1.14%). This latter subbasin shows the lowest percentage~~%~~ KGE decrease, despite having the highest proportion of glacier coverage. In particular, the KGE value slightly decreases from 0.66 to 0.65 due to a decrease in~~of 0.78 to 0.77~~ MBR from 0.78 to 0.77 when ~~in case of~~ shifting from the static to the dynamic glacier module. ~~The findings confirm the previously identified tendency: subbasins with a positive runoff bias in the static glacier module improve their performance when including the dynamic module and vice-versa (information on KGE and MBR values for each subbasin in Table 2s and 3s of the supplementary material).~~

### ~~3.4~~3.3 Evaluating the hydro-climatic drivers of droughts

#### ~~3.4.1~~3.3.1 Temporal patterns of hydro-climatic variables in the upper Adige

Results from the simulations were aggregated at ~~for~~ the ~~whole~~ basin scale to support the temporal analysis of hydro-climatic variables driving droughts. ~~–~~The purpose was~~is~~ to understand the hydroclimatic conditions that lead to the maximum glacier compensation effect during~~on~~ droughts. Figure 4 (panel a) shows a heatmap of interannual~~year to year~~ z-score values over summer seasons (JAS) from 1998 to 2022 for all hydro-climatic variables (with the exception of SWE, for which the z-score was evaluated based on its value at the end of June) together with~~and~~ their related mutual correlation values (panel b). In the heatmap, no single hydro-climatic variable can fully capture drought conditions, as high SWE can compensate for below-average summer precipitation ~~during summers~~ (e.g., years 2001 and 2014), while winters with below~~-~~-average snow can be compensated for by a rainy summer (e.g., years 2012 and 2017). Nevertheless, an overview on the correlation analysis between the hydro-climatic variables provides further insights.





**Figure 4 - Heatmap of the z-score values (green negative, purple positive) for the 5 hydro-climatic drivers of droughts for the whole upper Adige River Basin from 1998 to 2022 calculated for the summers months JJA and at end of June for SWE (panel a). Mann–Kendall correlations (from −1 to 1) between variables and significance: ∗, ∗∗ and ∗∗∗ refer to significant p-values ≤ 0.05, ≤ 0.01 and ≤ 0.001, respectively (panel b). Scatterplot of z-scores for SWE and glacier melt with colours for the temperature z-scores (panel c).**

In particular, runoff shows a ~~strong~~moderate positive correlation with precipitation (r=0.47, p<0.001) with decreasing correlation values with SWE (r=0.32, p<0.05), while it appears uncorrelated with glacier melt (p>0.05, Figure 4, panel b). On the one hand, SWE and glacier melt show a moderate~~strong~~ and highly significant negative correlation (r=-0.55, p<0.001), recalling the blanketing effect of abundant snow cover, which ~~preserving~~ protect

glacier from melting. On the other hand, glacier melt is driven by summer temperature (r=0.42, p<0.01) as it is highlighted by the conditions in 2003 and 2022 conditions, when high temperature combined with low SWE led to ~~higher than~~above-average glacier melt during severe hydrological droughts.

We further examined the apparent ~~strong~~ relationship between SWE, temperature, and glacier melt visualizing each summer in a scatterplot (Figure 4, panel c). High glaciers melt rates typically occur~~s~~ under~~with~~ above-average temperatures and below-average SWE at the end of June, confirming the behaviour suggested by the correlation analysis.


### ~~3.4.2~~3.3.2 Focus on the major past hydrological droughts

In the selected years of 2003, 2005 and 2022, different temporal and spatial dynamics led to runoff deficit and droughts across ~~in~~ the study area (Figure 4 and Figure 5). The year 2003 was characterized by exceptionally hot and dry spring and summer conditions triggering a hydrological drought. In particular, *Val d'Ultimo* showed a reduction of ~~-~~60% compared to ~~the~~ its average summer (JAS) runoff of 75 mm and *Val Sarentino* showed a ~~-~~58% reduction relative to ~~of~~ its runoff average of 68 mm (Figure 5, further information on subbasins locations and names are available in Figure S1 and Table S2 of the Supplementary Material)~~ of runoff~~. Other areas in the north-east also showed ~~-50%~~ runoff reductions of around 50% relative to ~~from~~ an average runoff of 99 mm, despite increased glacier melt contributions. The runoff decline was primarily driven by strong precipitation deficits that~~,~~ ~~which~~ affected the entire study area. The most severe deficit was observed in *Val d'Ultimo*, with a summer precipitation reduction of ~~-~~77% below ~~for~~ the average ~~summer precipitation~~value of 106 mm. Temperature anomalies were above ~~the~~ average, with a mean increase~~s~~ of ~~+~~0.89 °C across the upper Adige River Basin ~~region~~. Snow Water Equivalent (SWE) at the end of June showed negative anomalies throughout the region, with 14 subbasins recording values ~~-~~100% below their climatological mean~~compared to average summer values~~. The 14~~se~~ subbasins that showed the lowest decrease in SWE during summer 2003 were also those that showed the ~~also registered the~~ lowest ~~average~~ climatological SWE values over ~~in~~ the 26-year simulation period. Due to the hot ~~2003~~ summer conditions in 2003, glacier melt strongly contributed across the region. Glacierized subbasins in the southwest showed values ~~+~~101% higher melt compared to the summer average values of 0.7–1.2 mm, while northeastern glaciers showed even stronger increases reaching ~~glacier melt of +~~180% above the average value of 0.7 mm.

In 2005, the runoff deficits across the region were the result of different factors compared to the 2003 drought. The most pronounced runoff reductions occurred in the already dry western areas, with runoff deficits of ~~-~~55%

relative to the ~~of its~~ average runoff of 107 mm ~~of runoff~~ in *Val di Solda* and -54% relative to the average runoff of 81 mm in the lower ~~part of the~~ *Val di Solda*.

In contrast, the eastern part of the region, where both precipitation and glacier melt anomalies were mild or even positive, experienced ~~smaller~~ more limited runoff deficits of around -20%. These patterns were influenced by a less severe precipitation deficit than ~~compared to~~in 2003. The western part of the region consistently experienced dry conditions, with precipitation deficits down to -40%, while the southern part recorded positive anomalies of up to ~~+~~22% and the eastern part up to ~~+~~18%.


Preci
Temp
SWE
Glaciermelt
Runoff
mm
°C
%
Valle dei Molini
Val di Solda
Val Martello
Val d'Ultimo
Val di Mezzo
Val Casies
Val Sarentino
Val di Vizze
Val d'Ega
JAS
2003
2005
2022





**Figure 5 – Spatial representation of climatological mean (first column on the left) and anomalies (three columns on the right) computed for the average summer values from 1997-2023 within the upper Adige River Basin of the hydro-climatic variables. Only SWE consider values at the end of June. The anomalies refer to the selected drought years of 2003, 2005 and 2022.**


Temperature anomalies were negative across the entire area, with a widespread cooling of ~~-~~0.8~~/~~-1.4 °C below average. This colder-than-average summer had a notable impact on other hydrological components. Despite the cold summer temperatures ~~in summer~~, SWE at the end of June was similar to that of 2003, with deficits ranging from -100% to -60% across the region, particularly in headwater subbasins. Differently than in 2003, glacier melt in 2005 showed lower values, largely driven by the cooler temperatures. However, some eastern subbasins still exhibited glacier melt anomalies up to ~~+~~31%, contributing to the relatively smaller runoff deficits in that part of the region.

During the severe 2022 drought, precipitation anomalies, especially in the eastern part, compounded with higher-than-average temperature (+1 °C) and one of the most severe SWE deficits, especially for the western part. Runoff deficits reached~~showed~~ -60% relative to ~~for~~ the average summer runoff of 81 mm in *Val Casies*, ~~-~~48% relative to

~~for~~ 111 mm in *Val di Vizze*, ~~-~~47% relative to ~~for~~ 48 mm in *Val d'Ega* and ~~-~~46% relative to the ~~for~~ average summer runoff of 75 mm in *Val d'Ultimo*. Precipitation deficits were mostly located in the centre-eastern part. *Val Casies* and *Val di Mezzo* recorded the lowest values with deficits of ~~-~~45% and ~~-~~42% respectively, relative to~~for~~ summer averages of 121 mm and 113 mm. Positive temperature anomalies were widespread across the region, with average values of +1 °C, contributing to accelerated snow- and ice-~~-~~melt processes. SWE at the end of June showed one of the most severe reductions, particularly in the western part. Overall, 75% of the subbasins showed SWE deficits below ~~-~~90%, except for the north- and central~~mid~~-eastern areas. Glacier melt was only partially affected by the warmer conditions and reduced~~tion~~ ~~of~~ snow cover~~age~~. Increased glacier melt was~~ere~~ observed~~showed~~ in some areas, such as in *Val Ridanna*, which recorded ~~+~~156% increase relative to~~of~~ its summer average of 7.1 mm. However, in other subbasins, glacier melt was negative, including reduction of ~~-~~100% in *Val di Anterselva*, ~~-~~62% in *Val di Fleres*, and ~~-~~41% in *Val Aurina*.

### ~~3.4.3~~3.3.3 What if glaciers did not retreat?

To further quantify the effects of glacier retreat on droughts in the upper Adige, we compared runoff and glacier melt conditions from 1997 to 2023 considering a counterfactual scenario of constant glacier area and volume as it was in 1997. ~~The r~~Results ~~reveal~~ show progressively increasing differences both in terms of glacier melt and runoff during the selected drought events~~,~~. This condition highlight~~sing~~ the growing influence of glacier recession and the buffering capacity lost that would have been available during drought years. During the 2003 drought, ~~In particular,~~ average summer glacier melt from the dynamic~~al~~ ~~glacier module~~simulation was 7.81% lower than under 1997 glacier conditions, indicating a relatively limited loss of buffering capacity. The difference increases to 15% in 2005 and reached 44.7% in 2022, demonstrating the growing impact of glacier retreat on meltwater availability during dry summers ~~shows increasingly differences compared to the results of the 1997 constant glacier extent model: 7.81, 15.03 and 44.69% for the 3 selected drought years of 2003, 2005 and 2022, respectively, as detailed in (~~**Figure *6***~~)Table 1~~.




395

**Figure 6 – Simulating the lost buffering capacity (light blue shaded area) in terms of glaciermelt, runoff and glaciermelt contribution to runoff at the Bronzolo gauge station considering the 1997 glacier areas (red lines/bar) and yearly updating the glacier area (blue lines/bar).**

400 **Table 1 – Values of glacier melt and runoff for the three selected drought years of 2003, 2005 and 2022 at the Bronzolo gauge station and the percentage decrease of glacier melt contribution for the dynamic glacier module compared to the static module with 1997 levels.**

| Years | Average summer flow (mm) | | | | | | Glacier melt % of runoff | | |
|---|---|---|---|---|---|---|---|---|---|
| | Glacier melt | | | Runoff | | | | | |
| | Dynamic module | 1997 glaciers | Δ [%] | Dynamic module | 1997 glaciers | Δ [%] | Dynamic module | 1997 glaciers | Δ [%] |
| 2003 | 5.80 | 6.29 | **-7.81** | 49.68 | 50.17 | **-0.99** | 11.67 | 12.54 | **-6.89** |
| 2005 | 2.52 | 2.96 | **-15.03** | 51.86 | 52.32 | **-0.87** | 4.85 | 5.66 | **-14.29** |
| 2022 | 4.43 | 8.00 | **-44.69** | 54.19 | 57.79 | **-6.23** | 8.17 | 13.85 | **-41.01** |

The loss of glacier buffering capacity translated into reductions in summer runoff, although the magnitude of the effect Average summer runoff proved to be less sensitive to glacier retreatwas smaller than that observed for

405 glacier melt., with notable differences among the three simulated drought years. In 2003, runoff was showed a change of -0.99% lower in between the dynamic simulationand static simulations., while iIn 2005, both scenarios

produced~~vided~~ similar runoff values mainly due to lower contributions from glaciers during the cold 2005 drought conditions~~ characterizing the 2005 drought~~. In 2022, runoff was ~~decreased by~~ 6.23% lower than in the counterfactual scenario~~between the dynamic and static glacier simulation~~.

The percentage contribution of glacier -melt ~~over~~ to ~~the~~ total runoff most clearly shows the erosion of glacier buffering capacity~~declined significantly over time between the dynamic and 1997 glacier modules~~. In relation to the 1997-glacier counterfactual, the contribution of glacier melt to total runoff was reduced by ~~In 2003, the difference was~~ 6.89% during the 2003 drought, ~~increasing to~~ 14.29% in 2005 and ~~reaching~~ 41.01% in 2022. This result indicates that by 2022, more than two-fifth of the potential glacier buffering capacity that would have been available under the 1997 glacier extension had already been lost. While the reduction in~~of~~ glacier -melt is mainly attributable~~ed~~ to ~~the~~ shrinking glacier area, the ~~increasing differences between the dynamic and 1997-glacier scenario indicate that the~~ longer the simulation period and the wider the gap between the two simulations from 2003 to 2022 dynamic and 1997-glacier scenario, highlighting ~~greater the glacier retreat~~ the reducing capacity of the upper Adige river basin to sustain streamflow during dry and warm summers. ~~and hence the larger is the error in terms of glaciers mitigating summer drought introduced by considering a static glacier area.~~

## 4 Discussions

This study highlights the evolving role of glaciers in buffering hydrological responses to drought in the upper Adige River Basin. During~~In the past~~ major past droughts, glacier melt contributed~~ion~~ ~~ranged~~ between 4% and 13% of summer runoff~~, with variations according to the underlying temperature, precipitation, and SWE conditions~~. However, this buffering capacity is diminishing due to ongoing glacier retreat. Over the 26 years of simulation, glaciers in the upper Adige River Basin ~~have been~~ shru~~i~~nk~~ing~~ at an average rate of 1.95 km² per year (Figure 2), significantly reducing their ability to sustain summer flows during dry periods~~, as shown for the 2022 drought, where we estimated a reduction of 44% of the glacier melt contribution to runoff compared to the 1997 glacier extent.~~

At the single summer drought event, glacier melt contributions is shaped by the interaction among cryospheric and climatic conditions (Figure 4 and Figure 5). In particular, the amount of snow cover at the end of June and the high temperature conditions during July-August-September months play a critical role in determining summer glacier melt within the upper Adige River Basin (Figure 4). Reduced snow cover shortens the duration of snow melt contribution while also exposing glacier ice to high temperatures earlier in the summer and hence increasing glacier melt rates. This result aligns with previous studies on factors driving glacier compensation effects in other glacierized areas in the world (Jenicek et al., 2016; Van Tiel et al., 2021).

The comparison of the 2003, 2005, and 2022 major drought events also provides evidence that the buffering role of glaciers is changing over time. Although 2022 experienced temperature conditions comparable to those of 2003, the compensatory effect provided by glaciers declined from 12% of summer runoff in 2003 to 8% in 2022. While the inherent uncertainty embedded in multiple factors such as glacier extent, elevation range, aspect, glacier melt data and model uncertainties, the implemented glacier dynamic module consistently indicates a lower glacier melt contribution in summer 2022 than in 2003. This result points to the reduction in glacierized area as an important factor limiting meltwater availability during recent droughts.

The exceptional importance of glacier melt during the 2022 drought has also been identified in other glacierized catchments in northern Italy, although the magnitude of the estimated anomaly differs among regions. In the upper Adige River basin, results indicate that 2022 summer glacier melt contribution was 280% above the climatological mean, whereas (Leone et al., (2026) found an anomaly of about 350% for other glacierized catchments in Valle d'Aosta and Valtellina in northern Italy. Despite these quantitative differences, both studies indicate that enhanced glacier melt substantially sustained summer streamflow during the 2022 drought.

The declining glacier melt contribution to summer streamflow (Figure 5 and Figure 6) between the two similar droughts of 2003 and 2022, suggests that the glacier peak water in the upper Adige River basin may already have been reached or is being approached. Similar patterns have been reported across the Alps, where glacier melt remains an important component of summer streamflow during dry years, even though its absolute contribution is already declining because of glacier retreats (Huss and Hock, 2018; Sommer et al., 2020; van Tiel et al., 2026).

~~We showed that the amount of snow cover at the end of June and the high temperature conditions during July-August-September months play a critical role in determining summer glacier melt within the upper Adige River Basin (Figure 4). A reduced snow cover leads to a shorter duration of snow melt contribution while also earlier exposing glacier ice to high temperatures and hence increasing glacier melt rates. This result aligns with previous studies on factors driving glacier compensation effects in other glacierized areas in the world (Jenicek et al., 2016; Van Tiel et al., 2021).~~

~~The comparison of the 2003, 2005, and 2022 droughts illustrates how the combination of varying cryospheric and climatic conditions shape runoff deficits. The 2003 drought, characterized by extreme precipitation deficits and high temperatures, resulted in the most severe runoff reductions in the upper Adige River basin (up to -60% in *Val d'Ultimo* and *Val Sarentino*). Despite the lowest runoff values, glacier melt provided a compensatory effect contributing to 12% of runoff and, hence mitigating the hydrological deficit. Differently, the 2005 drought, despite cooler temperatures and a lower precipitation deficit, was exacerbated by a significative reduction in snow and glacier melt. Such conditions highlight the importance of meltwater contributions to sustain streamflow during~~

~~droughts. The 2022 drought, although similarly warm as 2003, showed lower glacier melt contributions (8%) than 2003. Our results show some differences from similar studies in other regions in the north of Italy. Results indicate that 2022 summer glacier melt contribution was 280% above the climatological mean, whereas Leone et al., (2025) found an anomaly of about 350% for other glacierized catchments (Valle d´Aosta and Valtellina) in northern Italy, indicating for all regions the importance of glacier melt during the 2022 drought. Despite the inherent uncertainty embedded in multiple factors such as glacier extent, elevation range, orientation as well as glacier melt data and model uncertainties, the implemented glacier dynamic module highlights the importance of areal glacier reduction affecting glacier melt contribution in 2022 compared to 2003.~~For these reasons, the continued loss of this natural water storage could have important implications for water-demanding sectors in the future, particularly hydropower generation and irrigated agriculture, whose demands can remain high during periods of limited summer availability (Clason et al., 2022; Schaefli et al., 2019). Moreover, sharp declines during drought periods could strengthen upstream-downstream dependencies and increase the potential for competition among water uses. These pressures are likely to become more relevant under future climate and socio-economic changes. Future water management plans should therefore consider the progressive loss of cryospheric buffering and combine short-term improvements in water allocation with longer-term adaptation and transformation strategies addressing water demand, sectoral priorities and basin-wide coordination.

### ~~4.2~~4.1 Limitations

While the integration of the glacier module into the ICHYMOD-TOPMELT hydrological model has provided valuable insights into glacier melt contributions and drought responses in the upper Adige River Basin, some limitations calling for future improvements must be acknowledged.

The glacier module is based on the empirical relationship between glacier volume and area at the catchment scale. Although it performs well in simulating overall glacier melt contributions, it cannot capture the full complexity of glacier dynamics, such as internal flows, at the individual glacier level. Moreover, the module assumes that available SWE over glacier surfaces at the end of each hydrological year, directly contributes to glacier mass balance. This approach shows limitations to accurately simulate specific years with net glacier mass gain, as observed in certain periods. Additionally, while the model reproduces glacier area evolution reliably, its representation of mass balance is constrained by uncertainties in the Dussaillant dataset used for calibration (Figure 2). Nonetheless, the cumulative mass balance estimates from the model fall within the uncertainty range of Dussaillant-derived values, supporting its value for basin-scale assessments.

The use of a ~~The choice of in the glacier model only a~~ fixed ~~albeod~~albedo value in the glacier model constitutes a model simplification~~is a strong sempificatio~~ compared to ~~n with respect~~ state-of the art glaciers models (Maussion et al., 2019)~~(Maussoion et at. 2029)~~. Although glacier albedo ~~this parameter~~ may vary ~~in~~ spatially~~ce~~ and ~~time~~temporally, we tested~~dhose to explore~~ a range of spatio-temporally fixed parameters in order to ~~since this would~~ (i) substantially reduce the computational effort and (ii) avoid a clear risk of equifinality, given the ~~very~~ limited ~~set of~~ validation data available relative to the large number~~information at our disposal, compared to an almost endless set~~ of potential spatio~~ce~~-temporal~~ime~~ ~~combinations of~~ albedo configurations~~values~~.~~–~~ ~~Indeed, s~~Several other parameters equally influence glacier melt dynamics, namely combined melt factor for combined radiation-temperature driven melt, rain melt factor and night melt factor, all handling ice melt in different conditions. However, these parameters are coupled for snow and ice melt by model design, and therefore simultaneously ~~effectively~~ affecting~~controlling~~ the ~~behavior~~behaviour of both storages~~ at once~~. We consequently ~~Therefore, the calibration effort~~ focused the calibration on glacier albedo, which is the only parameter in the model framework that allows glacier melt to be adjusted independently of snowmelt~~the sole decoupled parameter in our framework~~.

T~~We are aware that our~~he selected calibration value of 0.65 is ~~slightly~~ higher than usual values typically associated with bare glacier ice and is ~~used for ice and more similar~~closer to those reported ~~values used~~ for firn. ~~(snow one year old).~~ Nevertheless,~~However,~~ it lies ~~the selected value falls~~ within the range reported by studies ~~defined~~conducted in comparable Alpine environments ~~in other studies in similar areas~~ (e.g., ~~Richter et a~~(Hartl et al., 2025; Krampe et al., 2022; Richter et al., 2026)~~l., 2026)~~ and remains below~~lower than~~ the fresh snow albedo value of 0.9 used in the model.

The performance of the integrated hydrological model shows a decreasing trend in Kling-Gupta Efficiency (KGE) values for subbasins with glacier coverage exceeding 1%. These subbasins are typically small, high-elevation catchments where modelling challenges are amplified due to limited data availability, meteorological forcings uncertainty and the complexity of snow- and ice-related processes. In this regard, only eight glacierized subbasins had complete runoff observations for the summer periods of 2003, 2005, and 2022, restricting the robustness of model validation and of the analyses focused on these years, as detailed in the S~~s~~upplementary M~~m~~aterial.

Despite these limitations, including a dynamic glacier module improved streamflow representation during low-flow summer events. The improved model enhanced the glacier melt quantification during droughts and provided a valuable tool for understanding the conditions that lead to high or low glacier melt contributions to summer

drought, that can be applied to other mountain glacierized catchments. The tool can be used for adapting water resources management in alpine regions within climate change context, where glacier retreat is altering seasonal water availability.

## 5 Conclusions

This study ~~Our analysis offers~~provides a modelling approach to quantify the role of glaciers and snow cover in buffering summer droughts, implementing a novel dynamic glacier module into the ICHYMOD-TOPMELT hydrological model. The application in the upper Adige River Basin shows an almost monotone trend of glacier retreat for consecutive 26 years. Focusing on the 2003, 2005 and 2022 most severe droughts, we ~~Our analysis~~ show~~s~~ how the combination of multiple hydroclimatic variables (i.e., summer precipitation, temperature and especially snow water equivalent at the beginning of ~~the~~ summers) drives the amount of glacier ~~ice~~melt that mitigat~~cing to~~ hydrological droughts.

~~Simulations in the upper Adige River basin in the Italian Alps for the 2003, 2005 and 2022 most severe droughts showed how~~ Gglacier melt contributions ranged between 4 and 12% at the upper Adige closing section while the ~~. The ongoing decrease of~~ glacier extension decreases (from 111 km2 in 1997 to 79 km2 in 2017) is leading to a significant~~ive~~ decline~~rease~~ in glacier melt contribution to the overall runoff, especially during acute hot and dry events. This condition suggests that the glacier peak water in the upper Adige River basin may already have been reached or is being approached. Moreover, our counterfactual analysis showed that, if glaciers had retained their 1997 extent, their contribution to total runoff during the ~~fir~~ 2022 drought could have been 41.01% higher than the contribution estimated under the actual 2022 glacier extent. For these reasons, the continued glacier loss and its implications on water management need to be considered in future plans, ~~This implies increasing vulnerabilities for water management in the area under drought conditions and~~ particularly with increasing ~~downstream~~ demand across multiple sectors and for downstream users. Given the increasing emissions and global temperature and consequent glaciers disappearance (Van Tricht et al., 2026), our results call for proactive adaptation and transformation strategies within water demand planning accounting for the decreasing buffering glaciers role to sustain summer flows and mitigate drought conditions.

## Data and code availability

Data and code are available by request to the Authors. ~~We are in the process of releasing all the model code, input and output simulation data in an open access public repository.~~

**Financial support**

The publication of this research was partially funded by the NEXOGENESIS project (grant agreement number 101003881), financed by the European Commission under the H2020 programme. Further support was provided by the European Space Agency (ESA) under the EO4MULTIHA project (2023–2025), contract number 4000141754/23/I-DT, by the project PE00000005_RETURN: "Multi-Risk sciEnce for resilient communities undeR a changing Climate" PNRR, under the Missione 4, Componente 2, Investimento 1.3, call n. 341 del 15/03/2022 CUP: D53C22002510002 declaration, by the PRIN 2022 project finanziamenti PNRR – Missione 4 "Istruzione e Ricerca" Componente C2 Investimento 1.1 "Fondo per il programma Nazionale di Ricerca e Progetti di Rilevante Interesse Nazionale" (PRIN) – CUP C53D23001990006, the PNRR project "Interconnected Nord-Est Innovation Ecosystem (iNEST)" Spoke 1 – RT1: Safety and quality of life in mountain environments CUP C43C22000340006 – financed by the European Union – Next Generation EU and by the project Nextwater_ST, funded by the Autonomous Province of Bolzano under Decree No. 23075/2024, CUP D53C24005810003.

**Conflict of interest/Competing interests**

The authors declare no conflict of interest.

**Author contribution**

SS, ST and GB conceived the study, and SS performed the analysis with the significant~~ive~~ support from ST, GB, AG, MZ, DZ. SS and ST wrote the first manuscript draft with support from AG, DZ and GB. MB, RD, MP contributed with insights, discussed the argument and review the manuscript. ST secured funding. SS and ST equally contributed to the study.

**References**

Avanzi, F., Munerol, F., Milelli, M., Gabellani, S., Massari, C., Girotto, M., Cremonese, E., Galvagno, M., Bruno, G., Morra di Cella, U., Rossi, L., Altamura, M., Ferraris, L., 2024. Winter snow deficit was a harbinger of summer 2022 socio-hydrologic drought in the Po Basin, Italy. Commun. Earth Environ. 5, 1–12. https://doi.org/10.1038/s43247-024-01222-z

Avanzi, F., Terzi, S., Castelli, M., Munerol, F., Andreaggi, M., Galvagno, M., Galletti, A., Maurer, T., Massari, C., Carlson, G., Girotto, M., Bertoldi, G., Cremonese, E., Gabellani, S., di Cella, U.M., Altamura, M., Rossi, L., Ferraris, L., 2026. Impacts of Mediterranean snow droughts on mountain socio-ecohydrology. Hydrol. Earth Syst. Sci. 30, 5769–5790. https://doi.org/10.5194/hess-30-5769-2026

Bertoldi, G., Bozzoli, M., Crespi, A., Matiu, M., Giovannini, L., Zardi, D., Majone, B., 2023. Diverging snowfall trends across months and elevation in the northeastern Italian Alps. Int. J. Climatol. 43, 2794–2819. https://doi.org/10.1002/joc.8002

Biella, R., Shyrokaya, A., Ionita, M., Vignola, R., Sutanto, S.J., Todorovic, A., Teutschbein, C., Cid, D., Llasat, M.C., Alencar, P., Matanó, A., Ridolfi, E., Moccia, B., Pechlivanidis, I., van Loon, A., Wendt, D.E., Stenfors, E., Russo, F., Vidal, J.-P., Barker, L., de Brito, M.M., Lam, M., Bláhová, M., Trambauer, P., Hamed, R., McGrane, S.J., Ceola, S., Bakke, S.J., Krakovska, S., Nagavciuc, V., Tootoonchi, F., Di Baldassarre, G., Hauswirth, S., Maskey, S., Zubkovych, S., Wens, M., Tallaksen, L.M., 2025. The 2022 drought needs to be a turning point for European drought risk management. Nat. Hazards Earth Syst. Sci. 25, 4475–4501. https://doi.org/10.5194/nhess-25-4475-2025

Brock, B.W., Willis, I.C., Sharp, M.J., 2000. Measurement and parameterization of albedo variations at Haut Glacier d'Arolla, Switzerland. J. Glaciol. 46, 675–688. https://doi.org/10.3189/172756500781832675

Cazorzi, F., Dalla Fontana, G., 1996. Snowmelt modelling by combining air temperature and a distributed radiation index. J. Hydrol. 181, 169–187. https://doi.org/10.1016/0022-1694(95)02913-3

Chiogna, G., Majone, B., Paoli, K.C., Diamantini, E., Stella, E., Mallucci, S., Lencioni, V., Zandonai, F., Bellin, A., 2015. A review of hydrological and chemical stressors in the Adige catchment and its ecological status. Sci. Total Environ. 540, 429—443. https://doi.org/10.1016/j.scitotenv.2015.06.149

Clason, C., Rangecroft, S., Owens, P.N., Łokas, E., Baccolo, G., Selmes, N., Beard, D., Kitch, J., Dextre, R.M., Morera, S., Blake, W., 2022. Contribution of glaciers to water, energy and food security in mountain regions: current perspectives and future priorities. Ann. Glaciol. 63, 73–78. https://doi.org/10.1017/aog.2023.14

Colombo, N., Guyennon, N., Valt, M., Salerno, F., Godone, D., Cianfarra, P., Freppaz, M., Maugeri, M., Manara, V., Acquaotta, F., Petrangeli, A.B., Romano, E., 2023. Unprecedented snow-drought conditions in the Italian Alps during the early 2020s. Environ. Res. Lett. 18, 074014. https://doi.org/10.1088/1748-9326/acdb88

Dalla Torre, D., Di Marco, N., Menapace, A., Avesani, D., Righetti, M., Majone, B., 2024. Suitability of ERA5-Land reanalysis dataset for hydrological modelling in the Alpine region. J. Hydrol. Reg. Stud. 52, 101718. https://doi.org/10.1016/j.ejrh.2024.101718

Di Marco, N., Avesani, D., Righetti, M., Zaramella, M., Majone, B., Borga, M., 2021. Reducing hydrological modelling uncertainty by using MODIS snow cover data and a topography-based distribution function snowmelt model. J. Hydrol. 599, 126020. https://doi.org/10.1016/j.jhydrol.2021.126020

Dussaillant, I., Hugonnet, R., Huss, M., Berthier, E., Bannwart, J., Paul, F., Zemp, M., 2025. Annual mass change of the world's glaciers from 1976 to 2024 by temporal downscaling of satellite data with in situ observations. Earth Syst. Sci. Data 17, 1977–2006. https://doi.org/10.5194/essd-17-1977-2025

Farinotti, D., Huss, M., Bauder, A., Funk, M., Truffer, M., 2009. A method to estimate the ice volume and ice-thickness distribution of alpine glaciers. J. Glaciol. 55, 422–430. https://doi.org/10.3189/002214309788816759

Frans, C., Istanbulluoglu, E., Lettenmaier, D.P., Clarke, G., Bohn, T.J., Stumbaugh, M., 2016. Implications of decadal to century scale glacio-hydrological change for water resources of the Hood River basin, OR, USA. Hydrol. Process. 30, 4314–4329. https://doi.org/10.1002/hyp.10872

Galos, S.P., Klug, C., Dinale, R., 2022. 20 Years of Glacier Change: The Homogenized Glacier Inventories for South Tyrol 1997-2005-2017. Geogr. Fis. E Din. Quat. 45, 171–183. https://doi.org/10.4461/gfdq.2022.45.6

Galos, S.P., Klug, R., Prinz, R., Rieg, L., Dinale, R., Sailer, R., Kaser, G., 2015. Recent glacier changes and related contribution potential to river discharge in the Vinschau/Val Venosta, Italian Alps. Geogr. Fis. E Din. Quat. 143–154. https://doi.org/10.4461/GFDQ.2015.38.13

Glen, J.W., 1955. The creep of polycrystalline ice. Proc. R. Soc. Lond. Math. Phys. Sci. 228, 519–538. https://doi.org/10.1098/rspa.1955.0066

Gupta, H.V., Kling, H., Yilmaz, K.K., Martinez, G.F., 2009. Decomposition of the mean squared error and NSE performance criteria: Implications for improving hydrological modelling. J. Hydrol. 377, 80–91. https://doi.org/10.1016/j.jhydrol.2009.08.003

Hartl, L., Covi, F., Stocker-Waldhuber, M., Baldo, A., Fugazza, D., Di Mauro, B., Naegeli, K., 2025. Loss of accumulation zone exposes dark ice and drives increased ablation at Weißseespitze, Austria. The Cryosphere 19, 3329–3353. https://doi.org/10.5194/tc-19-3329-2025

Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., Jackson, M., Kääb, A., Kang, S., Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B., Steltzer, H., 2022. High Mountain Areas. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]., 1st ed. Cambridge University Press. https://doi.org/10.1017/9781009157964

Huss, M., Farinotti, D., 2012. Distributed ice thickness and volume of all glaciers around the globe. J. Geophys. Res. Earth Surf. 117. https://doi.org/10.1029/2012JF002523

Huss, M., Farinotti, D., Bauder, A., Funk, M., 2008. Modelling runoff from highly glacierized alpine drainage basins in a changing climate. Hydrol. Process. 22, 3888–3902. https://doi.org/10.1002/hyp.7055

Huss, M., Hock, R., 2018. Global-scale hydrological response to future glacier mass loss. Nat. Clim. Change 8, 135–140. https://doi.org/10.1038/s41558-017-0049-x

Huss, M., Jouvet, G., Farinotti, D., Bauder, A., 2010. Future high-mountain hydrology: A new parameterization of glacier retreat. Hydrol. Earth Syst. Sci. 14, 815—829. https://doi.org/10.5194/hess-14-815-2010

Jenicek, M., Seibert, J., Zappa, M., Staudinger, M., Jonas, T., 2016. Importance of maximum snow accumulation for summer low flows in humid catchments. Hydrol. Earth Syst. Sci. 20, 859–874. https://doi.org/10.5194/hess-20-859-2016

Knoll, C., Kerschner, H., 2009. A glacier inventory for South Tyrol, Italy, based on airborne laser-scanner data. Ann. Glaciol. 50, 46–52. https://doi.org/10.3189/172756410790595903

Krampe, D., Arndt, A., Schneider, C., 2022. Energy and glacier mass balance of Fürkeleferner, Italy: past, present, and future. Front. Earth Sci. 10. https://doi.org/10.3389/feart.2022.814027

Kreyszig, E., Kreyszig, H., Norminton, E.J., 2011. Advanced Engineering Mathematics, Tenth. ed. Wiley, Hoboken, NJ. ISBN 978-0-470-45836-5

Larsen, S., Majone, B., Zulian, P., Stella, E., Bellin, A., Bruno, M.C., Zolezzi, G., 2021. Combining Hydrologic Simulations and Stream-network Models to Reveal Flow-ecology Relationships in a Large Alpine Catchment. Water Resour. Res. 57. https://doi.org/10.1029/2020wr028496

Lemus-Canovas, M., Crespi, A., Maines, E., Terzi, S., Pittore, M., 2025. More intense heatwaves under drier conditions: a compound event analysis in the Adige River basin (Eastern Italian Alps). Hydrol. Earth Syst. Sci. 29, 6781–6809. https://doi.org/10.5194/hess-29-6781-2025

Leone, M., Avanzi, F., Morra di Cella, U., Gabellani, S., Cremonese, E., Isabellon, M., Pogliotti, P., Scotti, R., Monti, A., Ferraris, L., Colombo, R., 2026. The 2022–2023 snow drought in the Italian Alps doubled glacier contribution to summer streamflow. Hydrol. Earth Syst. Sci. 30, 4649–4666. https://doi.org/10.5194/hess-30-4649-2026

Li, H., Beldring, S., Xu, C.-Y., Huss, M., Melvold, K., Jain, S.K., 2015. Integrating a glacier retreat model into a hydrological model – Case studies of three glacierised catchments in Norway and Himalayan region. J. Hydrol. 527, 656–667. https://doi.org/10.1016/j.jhydrol.2015.05.017

Linsbauer, A., Paul, F., Machguth, H., Haeberli, W., 2013. Comparing three different methods to model scenarios of future glacier change in the Swiss Alps. Ann. Glaciol. 54, 241–253. https://doi.org/10.3189/2013AoG63A400

Luo, Y., Arnold, J., Liu, S., Wang, X., Chen, X., 2013. Inclusion of glacier processes for distributed hydrological modeling at basin scale with application to a watershed in Tianshan Mountains, northwest China. J. Hydrol. 477, 72–85. https://doi.org/10.1016/j.jhydrol.2012.11.005

Maussion, F., Butenko, A., Champollion, N., Dusch, M., Eis, J., Fourteau, K., Gregor, P., Jarosch, A.H., Landmann, J., Oesterle, F., Recinos, B., Rothenpieler, T., Vlug, A., Wild, C.T., Marzeion, B., 2019. The Open Global Glacier Model (OGGM) v1.1. Geosci. Model Dev. 12, 909–931. https://doi.org/10.5194/gmd-12-909-2019

Modarres, R., 2007. Streamflow drought time series forecasting. Stoch. Environ. Res. Risk Assess. 21, 223–233. https://doi.org/10.1007/s00477-006-0058-1

Montanari, A., Nguyen, H., Rubinetti, S., Ceola, S., Galelli, S., Rubino, A., Zanchettin, D., 2023. Why the 2022 Po River drought is the worst in the past two centuries. Sci. Adv. 9, eadg8304. https://doi.org/10.1126/sciadv.adg8304

Moore, R.J., 2007. The PDM rainfall-runoff model. Hydrol. Earth Syst. Sci. 11, 483–499. https://doi.org/10.5194/hess-11-483-2007

Moore, R.J., 1985. The probability-distributed principle and runoff production at point and basin scales. Hydrol. Sci. J. 30, 273–297. https://doi.org/10.1080/02626668509490989

NASA Shuttle Radar Topography Mission (SRTM) Global. Distributed by OpenTopography, 2013. Shuttle Radar Topography Mission (SRTM). https://doi.org/10.5069/G9445JDF

Navarro-Ortega, A., Acuña, V., Bellin, A., Burek, P., Cassiani, G., Choukr-Allah, R., Dolédec, S., Elosegi, A., Ferrari, F., Ginebreda, A., Grathwohl, P., Jones, C., Rault, P.K., Kok, K., Koundouri, P., Ludwig, R.P., Merz, R., Milacic, R., Muñoz, I., Nikulin, G., Paniconi, C., Paunović, M., Petrovic, M., Sabater, L., Sabater, S., Skoulikidis, N.Th., Slob, A., Teutsch, G., Voulvoulis, N., Barceló, D., 2015. Managing the effects of multiple stressors on aquatic ecosystems under water scarcity. The GLOBAQUA project. Sci. Total Environ., Towards a better understanding of the links between stressors, hazard assessment and ecosystem services under water scarcity 503–504, 3–9. https://doi.org/10.1016/j.scitotenv.2014.06.081

Naz, B.S., Frans, C.D., Clarke, G.K.C., Burns, P., Lettenmaier, D.P., 2014. Modeling the effect of glacier recession on streamflow response using a coupled glacio-hydrological model. Hydrol. Earth Syst. Sci. 18, 787–802. https://doi.org/10.5194/hess-18-787-2014

Norbiato, D., Borga, M., Esposti, S.D., Gaume, E., Anquetin, S., 2008. Flash flood warning based on rainfall thresholds and soil moisture conditions: An assessment for gauged and ungauged basins. J. Hydrol. 362, 274–290. https://doi.org/10.1016/j.jhydrol.2008.08.023

Norbiato, D., Borga, M., Merz, R., Blöschl, G., Carton, A., 2009. Controls on event runoff coefficients in the eastern Italian Alps. J. Hydrol. 375, 312–325. https://doi.org/10.1016/j.jhydrol.2009.06.044

Penna, D., Engel, M., Bertoldi, G., Comiti, F., 2017. Towards a tracer-based conceptualization of meltwater dynamics and streamflow response in a glacierized catchment. Hydrol. Earth Syst. Sci. 21, 23–41. https://doi.org/10.5194/hess-21-23-2017

Pepin, N.C., Arnone, E., Gobiet, A., Haslinger, K., Kotlarski, S., Notarnicola, C., Palazzi, E., Seibert, P., Serafin, S., Schöner, W., Terzago, S., Thornton, J.M., Vuille, M., Adler, C., 2022. Climate Changes and Their Elevational Patterns in the Mountains of the World. Rev. Geophys. 60, e2020RG000730. https://doi.org/10.1029/2020RG000730

Radić, V., Hock, R., 2010. Regional and global volumes of glaciers derived from statistical upscaling of glacier inventory data. J. Geophys. Res. Earth Surf. 115. https://doi.org/10.1029/2009JF001373

Richter, J.N., Arndt, A., Ban, N., Gampierakis, N., Maussion, F., Prinz, R., Scheiter, M., Umlauf, N., Nicholson, L., 2026. Towards the Bayesian calibration of a glacier surface energy balance model for unmonitored glaciers. EGUsphere 1–47. https://doi.org/10.5194/egusphere-2025-6249

Schaefli, B., Manso, P., Fischer, M., Huss, M., Farinotti, D., 2019. The role of glacier retreat for Swiss hydropower production. Renew. Energy 132, 615–627. https://doi.org/10.1016/j.renene.2018.07.104

Seibert, J., Vis, M.J.P., Kohn, I., Weiler, M., Stahl, K., 2018. Technical note: Representing glacier geometry changes in a semi-distributed hydrological model. Hydrol. Earth Syst. Sci. 22, 2211–2224. https://doi.org/10.5194/hess-22-2211-2018

Shen, C., Appling, A.P., Gentine, P., Bandai, T., Gupta, H., Tartakovsky, A., Baity-Jesi, M., Fenicia, F., Kifer, D., Li, L., Liu, X., Ren, W., Zheng, Y., Harman, C.J., Clark, M., Farthing, M., Feng, D., Kumar, P., Aboelyazeed, D., Rahmani, F., Song, Y., Beck, H.E., Bindas, T., Dwivedi, D., Fang, K., Höge, M., Rackauckas, C., Mohanty, B., Roy, T., Xu, C., Lawson, K., 2023. Differentiable modelling to unify machine learning and physical models for geosciences. Nat. Rev. Earth Environ. 4, 552–567. https://doi.org/10.1038/s43017-023-00450-9

Sommer, C., Malz, P., Seehaus, T.C., Lippl, S., Zemp, M., Braun, M.H., 2020. Rapid glacier retreat and downwasting throughout the European Alps in the early 21st century. Nat. Commun. 11, 3209. https://doi.org/10.1038/s41467-020-16818-0

Stahl, K., Moore, R.D., Shea, J.M., Hutchinson, D., Cannon, A.J., 2008. Coupled modelling of glacier and streamflow response to future climate scenarios. Water Resour. Res. 44. https://doi.org/10.1029/2007WR005956

Stephan, R., Erfurt, M., Terzi, S., Žun, M., Kristan, B., Haslinger, K., Stahl, K., 2021. An Alpine Drought Impact Inventory to explore past droughts in a mountain region. Nat. Hazards Earth Syst. Sci. Discuss. 2021, 1–25. https://doi.org/10.5194/nhess-2021-24

Stergiadi, M., Di Marco, N., Avesani, D., Righetti, M., Borga, M., 2020. Impact of Geology on Seasonal Hydrological Predictability in Alpine Regions by a Sensitivity Analysis Framework. Water 12, 2255. https://doi.org/10.3390/w12082255

Telesca, L., Lovallo, M., Lopez-Moreno, I., Vicente-Serrano, S., 2012. Investigation of scaling properties in monthly streamflow and Standardized Streamflow Index (SSI) time series in the Ebro basin (Spain). Phys. Stat. Mech. Its Appl. 391, 1662–1678. https://doi.org/10.1016/j.physa.2011.10.023

The GlaMBIE Team, 2025. Community estimate of global glacier mass changes from 2000 to 2023. Nature 639, 382–388. https://doi.org/10.1038/s41586-024-08545-z

Tscholl, S., Marsoner, T., Bertoldi, G., Bottarin, R., Egarter Vigl, L., 2025. A High-Resolution Climatic Water Balance for Eco-Hydrological Inference in the Upper Adige Catchment (Italy). Geosci. Data J. 12, e70007. https://doi.org/10.1002/gdj3.70007

Ultee, L., Coats, S., Mackay, J., 2022. Glacial runoff buffers droughts through the 21st century. Earth Syst. Dyn. 13, 935–959. https://doi.org/10.5194/esd-13-935-2022

van Tiel, M., Huss, M., Zappa, M., Jonas, T., Farinotti, D., 2026. Swiss glacier mass loss during the 2022 drought: persistent streamflow contributions amid declining melt water volumes. Hydrol. Earth Syst. Sci. 30, 23–43. https://doi.org/10.5194/hess-30-23-2026

Van Tiel, M., Van Loon, A.F., Seibert, J., Stahl, K., 2021. Hydrological response to warm and dry weather: do glaciers compensate? Hydrol. Earth Syst. Sci. 25, 3245–3265. https://doi.org/10.5194/hess-25-3245-2021

Van Tricht, L., Zekollari, H., Huss, M., Rounce, D.R., Schuster, L., Aguayo, R., Schmitt, P., Maussion, F., Tober, B., Farinotti, D., 2026. Peak glacier extinction in the mid-twenty-first century. Nat. Clim. Change 16, 143–147. https://doi.org/10.1038/s41558-025-02513-9

765 Zaramella, M., Borga, M., Zoccatelli, D., Carturan, L., 2019. TOPMELT 1.0: a topography-based distribution function approach to snowmelt simulation for hydrological modelling at basin scale. Geosci. Model Dev. 12, 5251—5265. https://doi.org/10.5194/gmd-12-5251-2019

Zebisch, M., Pedoth, L., Pörnbacher, M., Renner, K., Fritsch, U., Alberton, M., Bertoldi, G., Bisignano, J., Corradini, P., Crespi, A., Crippa, C., Ducati, V., Eisendle, F., Hilpold, A., Hoffmann, C., Horvath, K., 770 Kircher, V., Lun, G., Maino, F., Misconel, S., Obojes, N., Omizzolo, A., Pellegrini, C., Perkmann, U., Prina, M.G., Roveri, G., Sparber, W., Strapazzon, G., Thaler, F., 2025. Rischi climatici e adattamento - Verso un Alto Adige resiliente al clima 188. https://doi.org/10.57749/Q8Y8-TV70