

**Assessing phytoplankton community composition using in-situ  
Evaluating a multispectral miniaturised fluorometer with three  
excitation fluorescence and potential channels for application  
to predicting phytoplankton community structure indices from BGC-  
Argo profiling floats float observations**

Flavien Petit<sup>1,2</sup>, Julia Uitz<sup>1</sup>, Louison Dufour<sup>3,4</sup>, Collin Roesler<sup>5</sup>, Frédéric Partensky<sup>3</sup>, Laurence Garczarek<sup>3</sup>, Priscillia Gourvil<sup>6</sup>, Céline Dimier<sup>7</sup>, Melek Golbol<sup>7,8</sup>, Vincenzo Vellucci<sup>7,9</sup>, David Antoine<sup>1,10</sup>, Christophe Penker<sup>h1,11</sup>, Vincent Taillandier<sup>1</sup>, Hervé Claustre<sup>1</sup>

<sup>1</sup> Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, LOV, F-06230 Villefranche-sur-Mer, France

10 <sup>2</sup>National Oceanography Centre, Southampton, UK

<sup>3</sup>Sorbonne Université, CNRS, UMR7144 Adaptation and Diversity in the Marine Environment (AD2M), ECOMAP team, Station Biologique de Roscoff (SBR), 29680 Roscoff, France.

<sup>4</sup>Centre Algatech, Institute of Microbiology of the Czech Academy of Sciences, Novohradská 237, Třeboň 1437901, Czech Republic

15 <sup>5</sup>Department of Earth and Oceanographic Science, Bowdoin College, Brunswick, ME, USA

<sup>6</sup>Sorbonne Université, CNRS FR2424, Roscoff Culture Collection (RCC), Station Biologique de Roscoff (SBR), 29680 Roscoff, France.

<sup>7</sup>Sorbonne Université, CNRS, Institut de la Mer de Villefranche, IMEV, F-06230 Villefranche-sur-Mer, France

20 <sup>8</sup> Sorbonne Université, MNHN, CNRS, IRD, Laboratoire d'Océanographie et du Climat : Expérimentations et Approches Numériques, LOCEAN, F-75005 Paris, France

<sup>9</sup> Sorbonne Université, CNRS, OSU Stations Marines, STAMAR, 4 Place Jussieu, F-75252 Paris, France

<sup>10</sup> Remote Sensing and Satellite Research Group, School of Earth and Planetary Sciences, Curtin University, Perth, WA 6845, Australia

<sup>11</sup> INSU Division Technique (DT-INSU), UAR 855, CNRS, Plouzané, France

25 *Correspondence to:* Flavien Petit ([flavien.petit@noc.ac.uk](mailto:flavien.petit@noc.ac.uk))([flavien.petit@noc.ac.uk](mailto:flavien.petit@noc.ac.uk))

## Abstract.

30 Phytoplankton community composition is a key determinant of ocean biogeochemical cycles, yet its observation from autonomous platforms remains challenging. In this study, we assessed the potential of *in situ* multispectral excitation fluorescence (MXF) to ~~discriminate~~predict phytoplankton ~~assemblages~~ community structure indices in the Northwestern Mediterranean Sea, ~~with~~ With a view toward applications on Biogeochemical-Argo (BGC-Argo) profiling floats, we evaluated a miniaturised, three-excitation-channel fluorometer. Laboratory measurements on ten phytoplankton strains confirmed that MXF ratios at 440, 470, and 532 nm provide taxon-specific signatures, especially for picocyanobacteria and green algae. Field  
35 observations of phytoplankton pigments were clustered into four ecologically distinct phytoplankton communities ~~along~~across the seasonal cycle, which defined the targeted phytoplankton community structure indices. A machine learning model was then trained to classify these clusters using MXF and additional bio-optical indices. Results show that existing BGC-Argo configurations (single-wavelength fluorescence, particulate backscattering, and beam attenuation coefficients) reliably distinguish broad community structures, such as pico- versus microphytoplankton dominance, but resolving finer pigment-  
40 based differences requires the ~~added~~additional spectral information ~~of~~provided by MXF. The different excitation channels contributed unequally: 440 and 470 nm provided robust pigment sensitivity across communities, while 532 nm was particularly informative for detecting phycoerythrin-~~and chlorophyll b~~-rich taxa. Overall, combining MXF with bio-optical proxies improved classification performance by integrating pigment-specific and size-structure information, demonstrating the potential of MXF to enhance autonomous monitoring of phytoplankton community dynamics and their role in ocean  
45 biogeochemical cycles.

## 1 Introduction

Phytoplankton play a key role in global biogeochemical cycles, particularly in the carbon cycle. They fix dissolved inorganic carbon through photosynthesis and transfer a portion of it to higher trophic levels, initiating the biological carbon pump. This mechanism is pivotal in regulating the ocean's carbon storage. However, primary production, i.e., the rate at which  
50 phytoplankton produce organic carbon, varies significantly across different time and space scales. This variation is attributed to environmental changes that induce changes in phytoplankton community structure and biomass (Rousseau and Gregg, 2014). Therefore, in the current context of climate change, monitoring phytoplankton dynamics on a global scale is crucial. The emergence of new observation platforms, such as BioGeoChemical-Argo (BGC-Argo) profiling floats equipped with miniaturized bio-optical sensors, offers the possibility to collect continuous vertical profiles of optical measurements that serve  
55 as proxies of biogeochemical variables (Biogeochemical-Argo Planning Group, 2016; Claustre et al. 2020). Fluorescence is a widely used proxy of chlorophyll-*a* (Chl<sub>a</sub>) concentration, a ubiquitous pigment in phytoplankton organisms, which is in turn used as an indicator of phytoplankton biomass. Equipped with (single channel) fluorometers, autonomous platforms thus allow

the observation of phytoplankton biomass variability across a wide range of spatial and temporal scales (e.g. Boss et al., 2008; Barbieux et al., 2019; Cornec et al., 2021; Bock et al., 2022).

60 Information on phytoplankton biomass only ~~is~~, however, is insufficient to understand the links between phytoplankton and the carbon cycle. Indeed, the composition of phytoplankton communities is known to be a critical determinant of the carbon cycle since several key processes largely vary between phytoplankton size classes or phylogenetic groups, such as CO<sub>2</sub> fixation through photosynthesis (Cermeño et al., 2005; Uitz et al., 2008), trophic interactions (Cushing, 1989; Finkel, 2007), elemental cycling (Morel 2008, Litchman et al., 2015), or carbon transfer to the deep ocean (Buesseler et al., 1998; Guidi et al., 2009; Henson et al., 2012; Bonnet et al., 2023).

~~However~~Yet, the composition of phytoplankton communities cannot be measured directly from the sensors currently implemented on BGC-Argo floats. Only a few methods have been proposed so far to overcome this challenge and go beyond the mere estimation of Chla biomass from bio-optical measurements of BGC-Argo floats. Specifically, Sauzède et al. (2015) developed a neural network algorithm using the vertical shape of the *in-situ* fluorescence profile as input to retrieve the relative contribution to the Chla of the three phytoplankton size classes (pico-, nano- and microphytoplankton). Cetinić et al. (2015) proposed a simple community index based on the ratio of the fluorescence signal to the particulate backscattering coefficient. Similarly, Terrats et al. (2020) used this ratio to detect coccolithophore blooms. Finally, Rembauville et al. (2017) developed a regional approach to estimate the stock of particulate organic carbon (POC) of bacteria and three phytoplankton size classes. This approach has been developed for applications to BGC-Argo floats that measure not only the fluorescence and particulate backscattering coefficient, but also the beam attenuation coefficient. More recently, Brewin et al. (2022) proposed an analytical framework to partition vertical chlorophyll profiles measured by BGC-Argo floats into contributions from two phytoplankton communities associated with the mixed layer and the deep chlorophyll maximum, combining chlorophyll and particulate backscattering observations. This model was later extended to three communities in an oxygen minimum zone context (Cox et al., 2023). Yet, while those methods provide useful information about phytoplankton community composition, they mostly rely on regional empirical relationships between phytoplankton community composition and bio-optical indices.

Multispectral excitation fluorescence (MXF) is an alternative approach to retrieve information on the relative pigment composition of the phytoplankton assemblage, from which major taxa can be discriminated. MXF consists in measuring *in-situ* fluorescence signals in response to excitation at different wavelengths, corresponding to the absorption peaks of different accessory pigments used as biomarkers of specific taxa in the phytoplankton community (e.g. Yentsch and Phinney, 1985; Bricaud et al., 2004; Brewin et al., 2014). A combination of three wavebands centred around 440, 470 and 532 nm was previously investigated in freshwater environments (Proctor and Roesler, 2010) and in the Arabian Sea (Thibodeau et al., 2014), providing promising results and paving the way for use in open ocean waters. Multi-channel instruments such as the FluoroProbe (bbe Moldaenke, 5 excitation wavelengths) have also been widely used for *in-situ* taxonomic discrimination in freshwater and coastal environments (Beutler et al., 2002; Catherine et al., 2012). More recently, multi-wavelength excitation fluorometer applications have extended to open ocean discrimination of phytoplankton assemblages, demonstrating their capacity to resolve major taxonomic groups including diatoms, dinoflagellates, as well as phycoerythrin-containing

(*Synechococcus*, *Trichodesmium*, etc.) or lacking (*Prochlorococcus*) cyanobacteria. prochlorophytes (Xu et al., 2022; Kodama and Taniuchi, 2022). Concurrently, laboratory-based studies have demonstrated that portable, low-cost multi-excitation devices combined with machine learning classifiers can achieve high-accuracy taxonomic discrimination across a wide range of phyla (Magalhães et al., 2025; Zhang et al., 2025), further highlighting the potential of this approach for scalable, cost-effective phytoplankton monitoring. The present study aims at extending the approach of Proctor and Roesler (2010) and assessing the potential of *in-situ* three-excitation-channel MXF as a proxy of phytoplankton taxonomic composition in view of future applications to BGC-Argo floats. Higher-resolution multispectral fluorometers, such as the FluoroProbe and MultiExciter (9 channels), offer enhanced taxonomic discrimination (e.g., Garrido et al., 2019, Kodama and Taniuchi, 2022; Xu et al., 2022). However, their significant weight, large physical footprint, and high power consumption limit their use primarily to shipboard or short-term logging applications. In contrast, the SeaBird ECO sensor used in the present study is miniaturized and optimized for the stringent power and payload constraints of the global BGC-Argo fleet (Bittig et al., 2019). Our study therefore evaluates whether a reduced 3-channel spectral subset sensor, ready for integration on profiling floats, can provide sufficient biological resolution for global-scale autonomous observations.

Although standard BGC-Argo floats are all equipped with a single-channel fluorometer (with excitation at 470 nm) and a backscattering sensor (e.g., Bittig et al., 2019), part of the BGC-Argo fleet is also instrumented with additional sensors that may provide useful information on phytoplankton composition indicators, such as a beam transmissometer (Rembauville et al. 2017) or a dual-channel fluorometer with excitation at 440 and 470 nm (see ) around 440 and 470 nm. Of note, the nominal wavelength of the blue excitation channel is sometimes referred to as 435 nm in instrument metadata (see <https://vocab.nerc.ac.uk/collection/R27/current/>) but both values correspond to the same blue excitation band used in these fluorometers.

In this context, the present study aims to evaluate the potential of bio-optical measurements from the sensors currently implemented on BGC-Argo profiling floats, either alone or in combination with MXF, for retrieving information on phytoplankton community composition. Hence, we designed a predictive machine learning model to assess the ability of MXF combined with other bio-optical observations to infer taxonomic information.

For this purpose, we combined laboratory experiments and fieldwork conducted in the Northwestern (NW) Mediterranean Sea. The NW Mediterranean Sea provides a good case study because its pronounced seasonal phytoplankton biomass cycle (e.g., D'Ortenzio et al., 2005; Lavigne et al., 2015) and ecological succession (Vidussi et al. 2001; Marty et al., 2002; Mayot et al., 2017) are comparable to those observed in the temperate regions of the open ocean.

For the laboratory work, we selected ten phytoplankton strains representative of the various taxa observed along the seasonal succession of the NW Mediterranean Sea. We measured the MXF response of each strain under controlled conditions, and measured the variability among taxa and strains. This allowed us to characterize the MXF sensor and validate the analysis of the field data.

During our fieldwork, we collected concomitant pigment concentrations and bio-optical parameters using a sensor package comprising a MXF sensor, a single-channel fluorometer, a backscatterometer and a transmissometer over an annual

cycle in the NW Mediterranean Sea. This *in-situ* dataset was then used to develop and test a phytoplankton community composition discrimination model. Ultimately, we provide recommendations for the use of MXF, alone or in combination with other bio-optical indicators, to infer the taxonomic composition of phytoplankton communities from BGC-Argo profiling float observations.

## 130 2. Material & methods

### 2.1. Laboratory work

#### 2.1.1. Phytoplankton strains and culture conditions

135 For laboratory experiments, we selected ten phytoplankton strains provided by the Roscoff Culture Collection (RCC; <https://roscoff-culture-collection.org/>). These strains ~~are~~ were selected as being representative of the taxonomic diversity of the main eukaryotic and prokaryotic phytoplankton organisms encountered in open-ocean waters, and particularly at the BOUSSOLE site in the NW Mediterranean Sea, based on previous pigment-based studies in the region (Marty et al., 2002; Mayot et al., 2017). The selected strains include three diatom species, one pelagophyte, one dinoflagellate and five  
140 photosynthetic prokaryotes (three *Synechococcus* and two *Prochlorococcus* strains; Table 1).

All strains were grown at a constant temperature of 21°C, under 50  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$  continuous white light provided by a white-blue-green LED system (Alpheus, France), and in either K+Si (Keller et al., 1987) or PCR-S11 culture medium (Rippka et al., 2000) for eukaryotes and prokaryotes, respectively. As fluorescence is significantly influenced by the physiology of phytoplankton cells, we used cultures in stable physiological status as assessed by a high PSII quantum yield  
145 ( $F_v/F_M$ ) using a Phyto-PAM-II fluorometer (Walz, Effeltrich, Germany). The  $F_v/F_M$  parameter was calculated as  $(F_M - F_0)/F_M$  (Pittera et al., 2014), where  $F_0$  is the dark-adapted basal fluorescence,  $F_M$  is the maximal fluorescence associated with the closing of photosynthetic reaction centres, and  $F_v$  is the variable fluorescence.  $F_M$  was measured after exposure to saturating light pulses and addition of 100  $\mu\text{M}$  of the photosystem II inhibitor 3'-(3,4-dichlorophenyl)-1',1'-dimethylurea (DCMU; Parkhill et al., 2001). The  $F_v/F_M$  parameter was measured concomitantly to cell counts made using a  
150 Guava EasyCyte flow cytometer (Luminex Corporation, USA) all over the growth of each phytoplankton culture (Marie et al., 2001). The MXF protocol (see Section 2.1.2) was applied to each culture in the middle to late exponential growth phase, just before the drop of the  $F_v/F_M$  index. The MXF protocol was repeated three times on distinct replicate culture vessels for each strain (biological triplicates).

155 Table 1: Name, taxonomy, pigment composition as detected by ~~HPLC~~High Performance Liquid Chromatography (HPLC) pigment analysis, and size class of the ten phytoplankton strains used for the laboratory experiments. Pico ~~stands~~ for picophytoplankton (0.2-2 µm), Nano for nanophytoplankton (2-20 µm) and Micro for microphytoplankton (20-200 µm); HL stands for high-light adapted; LL stands for low-light adapted. The pigments measured are Chlorophyll a (Chla), Fucoxanthin (Fuco), Diadinoxanthin (Diad), Diatoxanthin (Diat), 19'-HF (19'-Hexanoyloxyfucoxanthin), 19'-BF (19'-  
160 Butanoyloxyfucoxanthin), Peridinin (Peri), Zeaxanthin (Zea), Divinyl-chlorophyll a and b (DV-Chla, DV-Chlb), Chlorophyll c1 and c2 or c3 (Chlc1 + c2, Chlc3).

Species name	Class	RCC strain number	Other names	Main pigments	Size class
<i>Conticribra (Thalassiosira) weissflogii</i>	Mediophyceae (diatom)	RCC76	CCMP1336	Fuco, Diad, Chlc1 + c2, Diat	Micro
<i>Chaetoceros diadema</i>	Mediophyceae (diatom)	RCC1717	RA080513-06	Fuco, Diad, Chlc1 + c2	Micro
<i>Pelagomonas calceolata</i>	Pelagophyceae	RCC100	CCMP1214	<del>Fuco</del> , Chlc3, Chlc1 + c2, 19'-BF, <del>Fuco</del> , Diad, Diat	Nano
<i>Scrippsiella</i> sp.	Dinophyceae	RCC3006	VFAC24-3	<del>Peri</del> , Chlc1 + c2, <del>Peri</del> , Diad	Nano
<i>Minidiscus</i> sp.	Mediophyceae (diatom)	RCC4213	MACUMBA-SC18	<del>Fuco</del> , Chlc1 + c2, <del>Fuco</del> , Diad, Diat	Nano
<i>Prochlorococcus marinus</i> (LL)	Cyanophyceae	RCC156	SS120-04/95	Zea, DV- <del>Chlb</del> Chla, DV- <del>Chla</del> Chlb	Pico
<i>Prochlorococcus marinus</i> (HL)	Cyanophyceae	-	PCC 9511	Zea, DV-Chlb, DV-Chla	Pico
<i>Synechococcus</i> sp.	Cyanophyceae	RCC2319	MINOS11	Zea	Pico
<i>Synechococcus</i> sp.	Cyanophyceae	RCC2374	A15-62	Zea	Pico
<i>Synechococcus</i> sp.	Cyanophyceae	RCC2379	BOUM118	Zea	Pico

### 2.1.2. Multispectral fluorescence measurements

165 All MXF measurements were performed using an ECO 3X1M fluorometer (Sea-Bird electronics, USA), with three excitation  
wavebands centred ~~on~~ onto 440, 470, and 532 nm, and emission onto 695 nm, with a 10-nm bandwidth. This three-channel  
design has been chosen for its immediate capacity to be integrated on a BGC-Argo float, or equivalent autonomy asset, due to  
its miniaturisation and low power consumption, the ECO series sensor already being part of the BGC-Argo payload.  
To determine the fluorescence to Chla slope factor (here expressed in fluorescence per Chla unit), the MXF measurements  
170 were collected for each culture over a 5-point dilution series ranging from 0.1 to ~~10 mg Chla m<sup>-3</sup>~~ 10 mg Chla m<sup>-3</sup>. Each culture  
was dark acclimated for 2 h before dilution and MXF measurements. The MXF measurements were  
performed immediately after dilution to avoid any dilution-induced physiological stress. The ECO 3X1M sensor outputs were  
recorded with the TeraTerm® software. Each culture was diluted in a 1L glass beaker that was then placed under constant  
slow stirring. The multispectral fluorometer was placed at the centre of the beaker, and the optical window was immersed 5  
175 mm below the surface. Blank measurements were performed with culture media and were then subtracted from the culture  
measurements to remove any possible fluorescence signal from colour-dissolved organic matter. Blank values were within a  
few counts of the dark reading, indicating that the measurements were not subject to optical interferences from the beaker edge  
or benchtop scattering. For each culture triplicate, on each dilution, we measured the fluorescence response during three series  
of one minute of continuous acquisition at 1 Hz, each separated by two minutes of darkness. The signal did not decrease  
180 significantly during acquisition, indicating that there was no quenching during the protocol application. The ECO 3X1M was  
used with factory calibration coefficients. Temperature effects on the sensor were considered negligible as only the optical  
window was immersed in the culture medium, and never for more than one minute continuously. The stability of the signal  
over each one-minute acquisition period confirmed the absence of any thermal drift.

### 2.1.3. Laboratory data processing

185 For each of the ten selected phytoplankton strains grown in culture, the MXF measurements were processed as follows. First,  
for each dilution series, a blank value was recorded by measuring the average response of the culture medium alone and was  
then subtracted from the raw sensor output acquired for the culture as described above. The MXF measurements collected over  
the three consecutive acquisition periods were averaged to obtain a single fluorescence value, expressed in digital counts (DC).  
Finally, the dilution series was used to define a Chla-specific calibration value, expressed in units of DC (mg Chla m<sup>-3</sup>)<sup>-1</sup>, for  
190 each of the three excitation wavelengths and each of the ten selected strains. This calibration value represents the coefficient  
of a linear regression between the fluorescence response at a given wavelength expressed in DC and the Chla concentration in  
mg m<sup>-3</sup> for the entire dilution range and for each replicate of a given phytoplankton strain. For the sake of simplicity, the raw  
fluorescence signal, in DC, at an excitation wavelength  $\lambda$ , will be noted as  $F_\lambda$  (i.e.,  $F_{440}$ ,  $F_{470}$ , and  $F_{532}$  for the excitation  
wavelengths 440 nm, 470 nm, and 532 nm, respectively) and the Chla-specific calibration value will be noted as  $F^*_\lambda$  (i.e.,  
195  $F^*_{440}$ ,  $F^*_{470}$ , and  $F^*_{532}$ ).

## 2.2. Field measurements of bio-optical and biogeochemical variables

### 2.2.1. Sampling strategies

200 Concomitant phytoplankton pigment determinations and bio-optical measurements were performed at sea every month, from  
December 2020 to October 2021 at the BOUSSOLE station (Buoy for the acquisition of long-term optical time series), a long-  
term monitoring site located at 7°54'E, 43°22'N in the Ligurian (NW Mediterranean) Sea (Antoine et al., 2008). On each  
monthly cruise (GOLBOL Melek, VELLUCCI Vincenzo, ANTOINE David (2000)  
BOUSSOLE, <https://doi.org/10.18142/1>), a CTD-rosette equipped with an optical sensor package  
205 was used to perform casts from the surface down to 400 m depth. The optical package included an MXF sensor (the same ECO  
3X1M as used for the laboratory experiments), an ECO FLBB sensor, and a C-Rover beam transmissometer (both Sea-Bird  
Scientific). The ECO FLBB measures the Chla fluorescence at one excitation (470 nm) and one emission (695 nm)  
wavelengths, as well as the particulate backscattering coefficient at 700 nm ( $b_{bp}$ , see section 2.2.2). ~~We note that among~~ It is  
worth noting that among the three excitation channels of the ECO 3X1M (440, 470, and 532 nm), the first two (440 and 470  
210 nm) are shared with the two types of dual-channel fluorometers (Sea-Bird Scientific ECO FLBBFL<sub>τ</sub> or RBR Tridente)  
~~now currently~~ implemented on some BGC-Argo floats. This correspondence allows us to also test the potential of dual-  
channel fluorometers for inferring phytoplankton composition indicators in our analysis. The C-Rover transmissometer  
measures the light beam transmitted between the emitter and receptor (at 650 nm and over an optical path length of 25 cm),  
allowing the calculation of the attenuation coefficient. From this, the particulate beam attenuation coefficient ( $c_p$ ) is calculated  
215 by removing the attenuation of pure seawater. Both sensors have already been mounted on several BGC-Argo floats for  
different biogeochemical applications (e.g., Mignot et al., 2014; Rembauville et al., 2017; Barbieux et al., 2022).  
Concomitantly, seawater was sampled from the Niskin bottles attached to the CTD-rosette at ten discrete depths for pigment  
identification and quantification by ~~High-Performance-Liquid-Chromatography (HPLC)~~ HPLC.

### 2.2.2 In-situ data processing

220 The factory-determined dark value of the ECO FLBB sensor was validated in the laboratory using black tape to cover the  
optical window, then subtracted from the raw DC following BGC-Argo data management recommendations (Schmechtig et  
al., 2018a). The optical backscattering coefficient was measured during the CTD-rosette upcast, which was used for seawater  
sampling. The angular scattering coefficient ( $\beta$ ) was recorded every second at a central angle of 124° and a wavelength of 700  
nm. To obtain the particulate angular scattering coefficient ( $\beta_p$ ), the contribution of pure ~~seawater~~ water, dependent on  
225 temperature and salinity (Zhang et al., 2009), was subtracted from  $\beta$ . The  $\beta_p$  coefficient was then converted into  $b_{bp}$  following  
standard conversion guidelines and applying a  $\chi$  factor of 1.076 (Schmechtig et al., 2018b).

The ECO 3X1M sensor was mounted on the CTD-rosette frame, providing simultaneous MXF measurements. The same black tape procedure was used to subtract dark values. The raw fluorescence values, expressed as counts, were directly used as the fluorescence signal.

230 The particulate attenuation coefficient was corrected for sensor drift and calculated from total beam transmittance as in Barnes & Antoine (2014).

The outliers in fluorescence,  $c_p$  and  $b_{bp}$  datasets were detected and removed using a threshold of 1.5 simple moving average ( $\Delta$ depth = 3 m). Each profile was then smoothed using a simple moving average ( $\Delta$ depth = 3 m).

### 235 2.3. Determination of phytoplankton pigments

For both laboratory and field samples, Chla and accessory pigments were identified and quantified by HPLC analysis. Briefly, seawater from discrete field samples (~2L) or cultures (0.1 to 0.5L, depending on culture biomass concentration) was filtered onto glass fibre filters (GF/F Whatman 25 mm), stored in liquid nitrogen during cruises and then transferred at  $-80^\circ\text{C}$  in the laboratory until further analysis at the SAPIGH HPLC analytical facility of the Institut de la Mer de Villefranche (IMEV; <https://lov.imev-mer.fr/web/facilities/sapigh/>). Phytoplankton pigments were extracted by sonication in 100% 3 mL methanol (100%) at  $-20^\circ\text{C}$  for 2 hours clarified by vacuum filtration (GF/F Whatman 25 mm), and finally separated and quantified analyzed within 24 hours by HPLC using an Agilent Technologies 1200 Series system. More details about the HPLC analytical protocol can be found in Ras et al. (2008). The total chlorophyll *a* concentration, [Chla], is defined as the sum of chlorophyll *a*, divinyl-chlorophyll *a*, and chlorophyllid *a* concentrations.

245 For *in-situ* samples, we specifically investigated the distribution of seven diagnostic pigments (DP): peridinin (Peri), 19'-butanoyloxyfucoxanthin (19-BF), fucoxanthin (Fuco), 19'-hexanoyloxyfucoxanthin (19-HF), alloxanthin (Allo), zeaxanthin (Zea), divinyl-chlorophyll *b* (DV-Chla, DV-Chlb), and chlorophyll *b* (Chlb), with total chlorophyll *b* (TChlb) defined as the sum of DV-Chlb and Chlb, as well as divinyl-chlorophyll *a* (DV-Chla). These pigments are defined as biomarkers of major phytoplankton taxa and were further grouped into three phytoplankton size classes, i.e. micro- ( $>20\ \mu\text{m}$ ), 250 nano- (2-20  $\mu\text{m}$ ) and picophytoplankton ( $<2\ \mu\text{m}$ ), according to the approach of Claustre (1994) and Vidussi et al. (2001). Following the equations given in Uitz et al. (2006), the DP-based method allowed the estimation of the relative contribution to the [Chla] of the three size classes. Because it relies on biomarker pigment concentrations, this approach yields an average, synthetic estimate of both the taxonomic and size composition of the phytoplankton communities. Although it has limits because some phytoplankton taxa may occasionally span over several size classes and some DP may be found in several taxa (e.g. Vidussi et al., 2001; Chase et al., 2020), this approach has been shown to provide reliable, quantitative provides a widely 255 used information for analyses about the phytoplankton community structure at large spatial and temporal scales (e.g., Vidussi et al., 2001; Bricaud et al., 2004; Uitz et al., 2006; Brewin et al., 2014). Moreover, the BOUSSOLE site and the NW Mediterranean Sea have been the subject of extensive prior characterization of phytoplankton seasonal dynamics (e.g., Marty

et al., 2002; Organelli et al., 2013; Latasa et al., 2022), providing a well-documented ecological context against which the  
260 outputs of this method can be interpreted.

## 2.4. Statistical analyses

To explore the potential of deriving phytoplankton community composition from MXF and other bio-optical measurements, we employed a ~~multi~~two-step analytical approach.

265 First, we derived phytoplankton community indices, defined as clusters with similar pigment composition. Phytoplankton pigment data were analyzed using Correspondence Analysis (CA) to identify similarities in pigment composition among samples. The first three dimensions of the CA were then subjected to a Hierarchical Ascending Classification (HAC) to define distinct clusters, corresponding to typical phytoplankton assemblages observed in the NW Mediterranean at the BOUSSOLE site (following Kramer and Siegel, 2019; Uitz et al., 2023). ~~These~~

270 Then, these clusters were ~~subsequently~~ used as categorical targets to evaluate the ability of MXF and bio-optical descriptors to infer phytoplankton composition through a Histogram Gradient Boosting (HGB) classification model.

This clustering-based strategy offers ~~two~~three key advantages: (1) it reduces the number of target variables, effectively adapting to the reduced number of degrees of freedom imposed by the limited number of excitation channels; (2) it transforms the prediction task from a regression to a classification problem, ~~eliminating~~ reduces the direct influence of biomass effects ~~magnitude~~ and allowing ~~allows~~ the model to focus solely on taxonomic composition indices; and (23) it ensures that the model captures the most significant source of taxonomic variability in the dataset, thereby reducing the influence of minor variance components and mitigating the risk of overfitting.

### 2.4.1. Correspondence analysis

We used correspondence analysis (CA) to visualize the main similarities among samples based on their relative pigment  
280 concentrations. This method generates linear combinations of relative pigment contributions, creating a new multidimensional space where sample projections reflect their resemblance in pigment composition.

In this transformed space, samples with similar pigment signatures are located close to each other, while those with distinct compositions are further apart. The first dimensions of the analysis capture the most significant variance in the dataset, effectively summarizing the dominant patterns in pigment distribution. By reducing data complexity, this approach provides a  
285 clearer interpretation of how different phytoplankton communities are structured based on their pigment signatures.

We applied this statistical method for two key objectives. First, we used it to compare the pigment composition of cultured phytoplankton strains with that of field samples, assessing whether the selected strains accurately represent the seasonal phytoplankton succession at the sampled location in the NW Mediterranean Sea. This CA was performed using the seven DP concentrations. Field samples were projected in the transformed CA projection space, and the distance between field and

290 laboratory samples will be discussed. Second, we conducted a CA exclusively on field samples, using the same set of pigments, and extracted the first three components as inputs for our clustering method described hereafter.

#### 2.4.2. Clustering of phytoplankton pigment data

The pigment data from NW Mediterranean field samples, consisting of the set of pigment concentrations detailed in the previous section, were clustered to identify major phytoplankton assemblages along the seasonal cycle using the CA (cf. 295 Section 2.4.1). The first three dimensions of the CA were used to quantify the resemblance in pigment composition across samples. A Hierarchical Ascending Classification (HAC) was then applied to these three dimensions, grouping samples based on their relative pigment composition rather than absolute pigment concentrations. The resulting cluster dendrogram was cut at a height of 20, minimising the intra-cluster variance, and yielding three initial clusters.

Given the distinct pigment composition of prokaryotic picophytoplankton versus micro-/nanophytoplankton communities, we 300 repeated the clustering after excluding picophytoplankton-dominated samples to refine the classification within the micro-/nanophytoplankton group. This second clustering step divided the micro-/nanophytoplankton samples into two additional clusters. In total, four distinct phytoplankton communities were identified.

#### 2.4.3. Classification of phytoplankton groups based on MXF and additional bio-optical proxies

Here we evaluated the possibility of using MXF measurements alone, or in combination with other bio-optical proxies 305 measured by BGC-Argo floats to retrieve information on phytoplankton community composition. For this purpose, measurements of  $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $b_{bp}$  and  $c_p$  were used as inputs of a model aiming at predicting the four different clusters identified with the method described in the previous section (cf. Section 2.5.2). We tested its performance using different sets of inputs, corresponding to either already deployed or feasible BGC-Argo sensor combinations, and varying levels of prediction task complexity (i.e., different number of clusters to predict).

310 The classification of *in-situ* phytoplankton communities (i.e., the prediction of a categorical target variable) based on MXF and additional bio-optical measurements was performed using a Histogram Gradient Boosting (HGB) algorithm. This type of machine learning model is particularly well-suited for tabular data, where each sample (row) is characterized by a consistent set of features (columns), and the dataset contains a relatively low number of observations (Chen and Guestrin, 2016; Shwartz-Ziv and Armon, 2022).

315 The model's performance depends not only on the total number of observations but also on the distribution of observations across target classes (here, phytoplankton community clusters). An imbalance in class representation can bias the model toward the dominant class, leading to an artificial overestimation of its performance for that group while reducing accuracy for underrepresented classes. To mitigate this imbalance, we applied the Synthetic Minority Oversampling Technique (SMOTE) (Chawla et al., 2002), which generates synthetic samples for minority classes to improve classification fairness and overall 320 model performance. In the end, each cluster was represented by 32 samples.

Because the dataset consists of a time series of a single year of phytoplankton community succession, the phytoplankton biomass was strongly correlated with the community composition. Since the five different measured variables (i.e.,  $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $b_{bp}$ ,  $c_p$ ) are significantly correlated with phytoplankton biomass, we used biomass-specific ratios to avoid overfitting and highlight intrinsic optical properties. First,  $F_{440}$  and  $F_{532}$  were normalized to  $F_{470}$ , which is the channel typically used to estimate Chla concentration from single-channel fluorometers. Second, each of the three fluorescence signals ( $F_{440}$ ,  $F_{470}$  and  $F_{532}$ ) was normalized to  $b_{bp}$  and  $c_p$ , used as particulate biomass proxies. The refractive index, representative of the composition of the particulate pool and estimated as a function of  $b_{bp}/c_p$ , was also used (Twardowski et al., 2001; Boss et al., 2004).

The hyperparameters of the model, i.e. the parameters influencing the learning process, were defined using a cross-validation grid search. In brief, the model has a learning rate of 0.05, to ensure stable convergence and 1400 estimators, and to capture non-linear relationships without overfitting, constrained by a maximum depth of 8 to maintain model interpretability and generalizability. The influence of the different descriptors was inspected through the mean impurity index, which reflects the importance of each descriptor in the succession of the decision trees. The model was validated with 20 cross-validations, using a stratified shuffle split method with a test size of 20%, which allows one to obtain the same proportion of the four clusters in each learning and testing dataset with random sampling. The classification results can be categorized into four different categories: True Positive (TP) corresponding to the accurate prediction of the presence of a class of a given phytoplankton assemblage, True Negative (TN) to the accurate prediction of the absence of a class, False Positive (FP) to the wrong prediction of the presence of a class, and False Negative (FN) to the wrong prediction of the absence of a class. The performance of the classification method was assessed through two different parameters, precision and recall, defined as follows:

$$\text{Precision} = \text{TP} / (\text{TP} + \text{FP}) \quad (1)$$

$$\text{Recall} = \text{TP} / (\text{TP} + \text{FN}) \quad (2)$$

The precision can be interpreted as the fraction of positive predictions of the model that were accurate, while the recall can be interpreted as the fraction of positive samples that have been correctly predicted by the model. A precision or recall value of 1 indicates perfect classification, meaning no false positives (for precision) or false negatives (for recall). Conversely, a low precision value suggests a high number of false positives, whereas a low recall value indicates that many actual positive cases were missed by the model.

The performance of the HGB classification model was tested for six different combinations of optical properties (e.g.  $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $c_p$ , and  $b_{bp}$ ), each consistent with potential future applications to BGC-Argo profiling floats (Table 2). Ultimately, we tested the performance of the model in predicting different numbers of clusters, ranging from 4 down to 2, each reflecting a different level of predictive complexity.

Table 2: List of sensor configurations tested in this study for retrieving phytoplankton community composition. For each configuration labelled A to F, we provide the measured variables along with examples of sensors that are available for integration, or already integrated, on BGC-Argo profiling floats.

Sensor configuration	F440	F470	F532	b <sub>bp</sub>	c <sub>p</sub>	Combination of sensors
A	X	X	X	X	X	MXF sensor Backscatterometer Transmissometer
B	X	X		X	X	Dual channel fluorometer Backscatterometer Transmissometer
C	X	X	X		X	MXF sensor Transmissometer
D	X	X		X		Dual channel fluorometer Backscatterometer
E	X	X	X			MXF sensor
F		X		X	X	Single channel fluorometer Backscatterometer Transmissometer

### 3. Results and discussion

355

#### 3.1. MXF signal in laboratory-controlled conditions

After quantifying the calibrated fluorescence for each of the ten phytoplankton strains and for each excitation wavelength, we considered two Chl<sub>a</sub>-specific fluorescence ratios  $F^*_{532} / F^*_{470}$  and  $F^*_{440} / F^*_{470}$ , as described in Proctor and Roesler (2010). Both ratios varied by a factor of approximately 2 when all taxa were considered (Fig. 1). The three *Synechococcus* strains consistently showed high  $F^*_{532} / F^*_{470}$  ratios (1.37 +/- 0.1) and low  $F^*_{440} / F^*_{470}$  ratios (0.91 +/- 0.06). By contrast, *Prochlorococcus* strains exhibited intermediate  $F^*_{532} / F^*_{440}$  and high  $F^*_{440} / F^*_{470}$  ratios (1.27 +/- 0.08 and 1.22 +/- 0.07 respectively). The diatom strains showed low to intermediate values (1 +/- 0.05 for both ratios). The dinoflagellates strain had similar average ratios as the diatoms, but replicates were a bit more variable. Finally,

365 the pelagophyceae strains had the lowest  $F^*_{532} / F^*_{470}$  and  $F^*_{440} / F^*_{470}$  ratios (0.71 +/- 0.02 and 0.95 +/- 0.12, respectively) of all taxa.


The higher  $F^*_{532} / F^*_{470}$  values observed for the *Synechococcus* taxon may be explained by their higher fluorescence at 532 nm, induced by the presence of phycoerythrin. Indeed, this phycobiliprotein is systematically found in open ocean *Synechococcus* and binds two chromophores, phycourobilin ( $\lambda_{\max}$  ~495 nm) and phycoerythrobilin ( $\lambda_{\max}$  ~545 nm), the latter thus being the most excited at 532 nm (Six et al., 2007; Grébert et al. 2018). It should be noted, however, that the phycoerythrobilin-to-phycourobilin ratio can vary in response to ambient light quality in some open-ocean *Synechococcus* strains through chromatic acclimation (Palenik, 2001; Humily et al. 2013), a capacity which may modulate the F532 response of individual strains and introduce variability in the signal beyond what is captured by our laboratory measurements under fixed white light conditions. Consequently, the phycourobilin-rich strain RCC2379 expectedly exhibited a lower average  $F^*_{532} / F^*_{470}$  ratio than the two other *Synechococcus* strains both being chromatic acclimators which, in white light, exhibit a low phycourobilin to phycoerythrobilin ratio (Palenik, 2001; Six et al. 2004; Humily et al. 2013). ~~Although the~~ The fairly high  $F^*_{532} / F^*_{470}$  ratio observed in *Prochlorococcus* strains are harder to explain, given ~~the~~ their very low ~~amounts of~~ phycoerythrin ~~present in these strains~~ content (Steglich et al. 2003, 2005). In contrast, the higher  $F^*_{440} / F^*_{470}$  ratio of the HL-adapted PCC 9511 compared to the LL-adapted RCC156 is consistent with ~~may be explained by~~ the much higher DV-Chla ( $\lambda_{\max}$  ~450 nm) to DV-Chlb ( $\lambda_{\max}$  ~475 nm) ratio of the former strain (Moore et al. 1995). The differences in fluorescence responses among diatoms, dinoflagellates, and pelagophyceae are also likely related to their distinct content in accessory chlorophylls, ~~yet these differences are somewhat subtle and harder to interpret in terms of~~ and carotenoids. In particular fucoxanthin in diatoms and pelagophytes, and peridinin in dinoflagellates, absorb light in the 440-532 nm range and transfer energy to Chla via fluorescence properties (resonance energy transfer, potentially contributing to the fluorescence signal measured at these excitation wavelengths (e.g., Bidigare et al., 1989; Ras et al., 2002; Bricaud et al., 2004); Menehgin et al., 2018). However, the efficiency of this energy transfer varies across taxa and physiological states, making the contributions of these carotenoids more challenging to interpret quantitatively than those of chlorophylls.

Our results are in line with numerous previous studies that demonstrated that pigment composition influences both light absorption and fluorescence emission spectra in phytoplankton, leading to taxon-specific fluorescence signatures (Yentsch and Menzel, 1963; Johnsen and Sakshaug, 2007; Hu et al., 2010; MacIntyre et al., 2010; Proctor and Roesler, 2010). More specifically, some laboratory studies using monospecific cultures demonstrated that fluorescence spectra vary significantly across taxa when multiple excitation and emission wavelengths are used (Yentsch and Menzel, 1963; Johnsen and Sakshaug, 2007; Poryvkina et al., 1994). More recent work has expanded on these findings by incorporating mixed communities, showing potential for determination of natural assemblages of phytoplankton (Hu et al., 2010; Escoffier et al., 2015). While Hu et al. (2010) included both monospecific and mixed cultures, as well as coastal marine samples, most of these studies were conducted under controlled laboratory conditions rather than in open-ocean environments.

Only a few studies have investigated fluorescence responses in natural, mixed phytoplankton communities (Seppälä and Balode, 1998; Hu et al, 2010; Proctor and Roesler, 2010; Thibodeau et al. 2014). Some of these studies benefited from

conditions that enhanced pigment-specific fluorescence signals, such as reversed filtration to increase pigment concentration (Seppälä and Balode, 1998), high resolution of fluorescence excitation/emission spectra (Seppälä and Balode, 1998; Hu et al, 2010) or naturally high chlorophyll concentrations in bloom conditions (Proctor and Roesler, 2010). These conditions contrast sharply with those of the open ocean, where phytoplankton communities are more diverse, pigment concentrations are lower, and taxonomic differentiation based on MXF alone is more challenging.

Despite this complexity, our MXF measurements on mono-specific cultures are promising and coherent with Proctor and Roesler (2010), who further demonstrated that intra-taxon variance in specific fluorescence ratios was lower than inter-taxon variance. Both studies their study and the present one suggest that a multispectral MXF sensor with three excitation channels (440, 470, and 532 nm) can provide sufficient sensitivity to distinguish taxa in controlled conditions even though natural communities consist of mixed assemblages of taxa with complex pigment signatures, leading to less contrasted fluorescence response than for laboratory cultures. Therefore, in the following section, we investigate whether it is possible to resolve taxonomic composition from *in-situ* phytoplankton fluorescence signals by analysing a year-long dataset from the NW Mediterranean Sea. Additionally, we assess whether combining MXF measurements with other bio-optical proxies could improve classification performance, in anticipation of future deployment of multispectral fluorometers on BGC-Argo profiling floats.



415


Figure 1: Scatterplot of F\*440/F\*470 vs. F\*532/F\*470 ratios for each phytoplankton strain grown in culture. The colour code indicates the taxon to which each strain belongs; the symbols indicate the strains, using the Roscoff Culture Collection code, when available (see Table 1).

420

### 3.2. Phytoplankton communities in the Northwestern Mediterranean Sea

Surface-Surface chlorophyll concentration at the BOUSSOLE site can reach values up to 5 mg Chla m<sup>-3</sup> during the spring bloom and drop below 0.1 mg Chla m<sup>-3</sup> in summer, along with seasonal changes of the phytoplankton community composition (Marty et al., 2002; Antoine et al. 2020). In winter, pigment indicators of prymnesiophyte, namely 19'-HF and 19'-BF, are observed alongside fucoxanthin, a marker of diatoms, and alloxanthin, a marker of cryptophytes (Marty et al., 2002; Mayot et al., 2017). During the spring phytoplankton bloom, fucoxanthin concentrations increase (Marty et al., 2002), which is likely related to an increase in diatoms. In summer, the mixed layer is phosphate-limited and phytoplankton communities are representative of stratified oligotrophic regions, with a prevalence of pigments specific to pico-phytoplankton, diviny-chlorophyll-a and zeaxanthin (Marty et al., 2008). This phytoplankton diversity is comparable to what is observed in temperate open-ocean regions (Vidussi et al., 2001; Marty et al., 2002; Lavigne et al., 2015; Mayot et al., 2017).

The composition of the phytoplankton communities in the field samples was analysed using a pigment-based clustering approach (see Section 2.4.2). The clustering allowed grouping samples with similar pigment composition and led to the discrimination of four distinct phytoplankton assemblages (clusters) over the year (Fig. 2a-d). All clusters are dominated by nanophytoplankton but vary significantly in the partitioning between micro- and picophytoplankton (Fig. 2e-h). The first cluster corresponds to winter communities as well as deep autumn communities, with a large proportion of picophytoplankton and a significant contribution of Chlb, a pigment typical of green microalgae, mostly flagellates (Bustillos-Guzmán et al., 1995). The second cluster coincides with the bloom community with a shared contribution of micro- and nanophytoplankton. This assemblage is characterized by a high fucoxanthin contribution, typically associated with diatoms. The third cluster is associated with summer communities located at and below the level of the deep chlorophyll maximum (DCM) and also exhibits a mixed composition of micro- and nanophytoplankton. Finally, the fourth cluster is characteristic of picophytoplankton communities in surface waters from summer to autumn, typically dominated by picocyanobacteria (Barlow et al., 1997).



445

450

Figure 2: Distribution of the phytoplankton communities as determined from the cluster analysis applied to the field pigment data: (a-d) Vertical distribution of the four pigment-determined clusters indicative of the main phytoplankton communities encountered over an annual cycle in the northwestern Mediterranean Sea (BOUSSOLE site). The size of the dots indicates the Chla concentration, used as a proxy of the phytoplankton biomass. (e-h) Tree map of the relative pigment concentration of each cluster (areas delimited by the grey lines), with the size class corresponding to the pigment taxa affiliation (colored).

455

The results from the culture experiments demonstrated that phytoplankton taxa can be distinguished by exhibit distinct MXF at the taxonomic level of the Class for eukaryotic phytoplankton (e.g., Diatoms, Dinoflagellates, Pelagophyceae) and Genus for picocyanobacteria (*Synechococcus* and *Prochlorococcus*). We then fluorescence ratios under controlled conditions (Fig. 1). So, we compared the relative accessory pigment composition of the *in-situ* clusters to that of the five laboratory characterized taxa. This analysis seeks to determine whether the field samples fall in the range of variability of accessory pigment composition of the culture samples.

460

Similar to the field samples, a correspondence analysis (see Section 2.4.1) was applied to the pigment composition of the ten phytoplankton strains grown in the laboratory. This method allows a visualization of the different strains in a space where the distance between two samples reflects their relative pigment composition similarity (Fig. 3). We observe three distinct groups corresponding to the different taxa represented by the ten selected strains. One is composed of Diatoms diatoms and Pelagophyceae, while the two others correspond to *Synechococcus* and *Prochlorococcus*, respectively. This highlights a

strong contrast in pigment composition between *Synechococcus* which has Zea, *Prochlorococcus* which has DV-Chla, DV-Chlb and Zea, and the nano- and microphytoplankton taxa that share many different pigments (Jeffrey et al., 1997; Veldhuis et al., 2005).

The culturelaboratory results provide a reference from which the projection space is computed, on which the field samples are projected (Fig. 3). The field data are evenly spread in the centre of the plan, indicating that the variability in the pigment composition of the field samples is similar to that observed in the laboratory cultures. Moreover, the four field-based clusters are fairly well distinguished in the CA. ~~The only exception being, except for~~ a few samples of cluster 1 with high CA2 values, due to the presence of 19'-HF, a pigment ~~that is not found in~~ absent from our culturedreference strains. However, this pigment has a very similar absorption signature to other carotenoids like 19'-BF or peridinin. We thus expect that the MXF sensor is sensitive enough to discriminate between different phytoplankton groups (clusters) characterized by distinct pigment composition in field samples.

In addition, we note that the *in-situ* samples have lower eigenvalues (i.e., absolute values on CA axes) than the laboratory samples indicating that the pigment variability is less contrasted in the field than in laboratory samples. This is not surprising because of the complexity of the pigment composition in natural samples associated with mixed phytoplankton assemblages, instead of single taxa in monospecific cultures. This characteristic of open-ocean samples could somewhat hamper the possibilities of inferring information on phytoplankton community composition from MXF measurements, a hypothesis that is tested in the next section.






Figure 3: Correspondence analysis of the pigment concentrations of the strains grown in culture. The pigment concentrations measured in the northwestern Mediterranean (BOUSSOLE site) seawater samples are [projected as supplementary observations](#) and represented using the same colour code as in Figure 2 ~~and projected as supplementary observations~~.

### 3.3. Discrimination of phytoplankton taxa from *in-situ* MXF and additional bio-optical variables

The predictive model (see Section 2.4.3) was tested using as inputs measurements from the MXF sensor alone ( $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ), or in combination with a backscatterometer ( $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $b_{bp}$ ) and a transmissometer ( $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $b_{bp}$ ,  $c_p$ ). Additionally, the model was also tested using measurements from a dual excitation channel fluorometer and backscatterometer ( $F_{440}$ ,  $F_{470}$ ,  $b_{bp}$ ), or combined with a transmissometer ( $F_{440}$ ,  $F_{470}$ ,  $b_{bp}$ ,  $c_p$ ) (Fig. 4 and Table A1). These two configurations are of particular interest as many BGC-Argo floats have been ~~and will continue to be~~, deployed with this specific set of sensors.

Considering the most comprehensive configuration, A (i.e.,  $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $b_{bp}$ ,  $c_p$ ), the precision and recall scores of the HGB model are homogeneous among all four clusters with values between 60% and 85% (+/- 15%) (Fig. 4 and Table A1). These results based on annual sampling in the NW Mediterranean are quite robust, showing sufficient precision to demonstrate that MXF, when used in combination with a backscatterometer and a transmissometer, effectively allow four phytoplankton taxonomic groups to be distinguished. If we remove either the MXF sensor or the transmissometer, leading to measurements of  $F_{440}$ ,  $F_{470}$ ,  $b_{bp}$  and  $c_p$ , or  $F_{440}$ ,  $F_{470}$ ,  $F_{532}$  and  $c_p$  (i.e., configurations B and C, respectively), we observe similar precision and recall scores, and a marked variability between the clusters (Fig. 4 and Table A1). However, all four clusters displayed a mean precision of 67.25% (+/- 6.9%) and 69% (+/- 12%) for configurations B and C, respectively. When considering configuration D, which corresponds to a sensor load with a dual-channel fluorometer and a backscatterometer (i.e.,  $F_{440}$ ,  $F_{470}$  and  $b_{bp}$ ), the scores are significantly lower with a mean precision of 62% (+/- 8.7%). The use of the MXF sensor only (configuration E, i.e.  $F_{440}$ ,  $F_{470}$  and  $F_{532}$ ) led to variable performances depending on the cluster. Thus, removing the transmissometer ( $c_p$ ) or the 532-nm excitation fluorescence channel ( $F_{532}$ ) seemingly induces a significant decrease in the general accuracy and recall scores of the model (Fig. 4). Finally, the configuration with only single-channel fluorescence measurements and two bio-optical indices (configuration F, i.e.  $F_{470}$ ,  $b_{bp}$ ,  $c_p$ ) led to lower performance for every cluster, except for the picophytoplankton dominated one (i.e., cluster 4; see Fig. 2). This suggests that a configuration without MXF performs better when the phytoplankton communities have contrasted size structures.






515

Figure 4. Precision and recall scores for phytoplankton community classification using different sensor package configurations. (A) Precision and (B) recall scores for predicting four phytoplankton clusters using various sensor combinations, either currently equipped on BGC-Argo profiling floats or suitable for future installations: full configuration (F440, F470, F532, b<sub>bp</sub>, c<sub>p</sub>); dual channel fluorescence with backscatterometer and transmissometer (F440, F470, b<sub>bp</sub>, c<sub>p</sub>); MXF with c<sub>p</sub> (F440, F470, F532, c<sub>p</sub>); dual channel fluorometer with backscatterometer (F440, F470, b<sub>bp</sub>); MXF only (F440, F470, F532) and single channel fluorescence with backscatterometer and transmissometer (F470, b<sub>bp</sub>, c<sub>p</sub>). Precision represents the fraction of positive predictions that were correct, while recall indicates the fraction of true positives correctly identified. Error bars reflect variability across cross-validation runs. Significance of paired t.test, [testing for](#)

525 significant difference within each cluster, is indicated with stars (ns: p.value > 0.05 ; \*: p.value <= 0.05 ; \*\*: p.value <= 0.01 ; \*\*\*: p.value <= 0.001 ; \*\*\*\*: p.value <= 0.0001).

Based on configuration A, which combines MXF and all bio-optical indicators, we can evaluate the predictive power of each descriptor. The mean importance of the descriptors in the discrimination of distinct phytoplankton clusters, as indicated by  
530 the impurity metrics (Fig. 5), highlights a significant role of all selected descriptors ( $b_{bp}/c_p$ ,  $F_{440}/F_{470}$ ,  $F_{440}/c_p$ ,  $F_{532}/F_{470}$ ,  $F_{532}/b_{bp}$ ,  $F_{470}/c_p$ ,  $F_{470}/b_{bp}$ ,  $F_{440}/b_{bp}$ ), and particularly of  $b_{bp}/c_p$ ,  $F_{440}/F_{470}$ ,  $F_{440}/c_p$  and  $F_{532}/F_{440}$  ratios. The high importance of the  $b_{bp}/c_p$  ratio was expected as it has been already described as an indicator of the particle size distribution (Dall’Olmo et al., 2009; Slade and Boss, 2015; Organelli et al., 2020). Interestingly, the second and fourth strongest predictive descriptors correspond to the fluorescence ratios from the MXF sensor ( $F_{440}/F_{470}$  and  $F_{532}/F_{470}$ ). This indicates the strong added value of the MXF descriptors  
535 in the predictive model, in line with our laboratory results. These results also support the findings of both Proctor and Roesler (2010) and Thibodeau et al. (2014) who showed that the  $F_{440}/F_{470}$  and  $F_{532}/F_{470}$  fluorescence ratios are related to the taxonomic composition of phytoplankton communities. Finally, we note that the commonly used phytoplankton community index,  $F_{470}/b_{bp}$  ratio (Cetinić et al., 2015; Lacour et al., 2019; Terrats et al., 2020), has a notably lower predictive robustness in our analysis- than all other ratios except  $F_{440}/b_{bp}$ . Therefore, the use of new sensor configurations, including a multiple excitation  
540 channel fluorometer and a beam attenuation transmissometer implemented on BGC-Argo floats, may have a stronger potential than previously described methods to detect seasonal succession of phytoplankton groups, with markedly different pigment composition or cell sizes (here picophytoplankton-dominated vs. microphytoplankton-dominated communities).



545 Figure 5: Importance of the different descriptors in the classification model, expressed as the mean decrease in impurity. The mean decrease in impurity reflects how much each descriptor contributes to improving the purity of the splits in the decision tree. A higher value indicates that the descriptor plays a more significant role in distinguishing (i.e., pigment-based clusters) in the model.

### 3.4. Model tuning to predict phytoplankton communities

550 In ~~Section 3.3, we considered four clusters found along the annual phytoplankton succession. We now evaluate the previous~~ section, the ~~same classification model and hyperparameters was evaluated~~ for predicting two or three clusters (instead the prediction of four), to assess whether reducing the level of phytoplankton clusters representing the main communities observed over the annual cycle. Here, we further explore the sensitivity of the model to the predictive complexity ~~improves the by~~ progressively reducing the number of clusters from four to three and two. This allows us to assess how model performance varies under ~~different sensor configurations simpler ecological scenari~~. First, we reduced the number of clusters from four to three by merging Clusters 2 and 3, the two clusters with the smallest distance, representing mixed nano- and microphytoplankton-dominated communities. The main distinction between these two clusters lies in their dominant pigments. Cluster 2 is indeed characterized by a higher relative contribution of alloxanthin, while Cluster 3 is dominated by 19'-HF. The three resulting clusters correspond to the surface summer picophytoplankton community associated with large concentrations of zeaxanthin (Cluster 4), the winter and deep summer picophytoplankton community with Chlb and DV-Chlb (Cluster 1), and the mixed micro- and nanophytoplankton community (Clusters 2 and 3 grouped together). Second, we reduced the initial number of four clusters to two, by merging Cluster 1 (dominated by Chlb-containing picophytoplankton) and Cluster 4 (dominated by zeaxanthin-containing picophytoplankton) to a picophytoplankton-dominated cluster, essentially grouping *Synechococcus*, *Prochlorococcus* and chlorophytes. The second cluster consists of the previously merged cluster (initially Clusters 2 and 3) composed of micro- and nanophytoplankton.

565 ~~Like the approach~~As presented in ~~Section~~section 3.3, we evaluated the classification model performances for different sensor configurations using the mean balanced recall metrics, which reflects the percentage of correctly classified samples. The same cross-validation method was applied to ensure consistency (Fig. 6). We first observed that a decrease in prediction complexity (specifically, a reduction in the number of clusters) does not consistently lead to an improvement in the classification model performance, and that the effect of such a reduction depends on the sensor configuration. When all sensors are included (configuration A), the model performs best in predicting 2 or 3 clusters than 4 clusters, achieving above 75% recall. Similarly, the model using as inputs two fluorescence wavelengths, as well as the  $b_{bp}$  and  $c_p$  coefficients (i.e., configuration B) achieves above 75% recall when predicting 2 or 3 clusters.

575 It is worth noting that the model using as inputs bio-optical measurements from sensors already implemented on some BGC-Argo floats (i.e, configuration F with single-channel fluorescence,  $b_{bp}$  and  $c_p$ ) demonstrates strong predictive capabilities for 2 or 3 clusters, achieving recall values of 72% and 74%, respectively. In the two-cluster prediction scenario, one


cluster predominantly comprises picophytoplankton, such as *Synechococcus* and *Prochlorococcus*, while the other includes a mix of nano- and microphytoplankton. Interestingly, previous studies have shown that the combination of  $F_{470}$ ,  $b_{bp}$ , and  $c_p$  effectively correlate with phytoplankton community size structure (Veldhuis et al., 2005; Brewin et al. 2011; Cetinić et al. 2015; Sauzède et al. 2015; Rembauville et al. 2017, Terrats et al. 2023) supporting our findings. In the three-cluster scenario, the model ~~has to~~ must distinguish between two different types of picophytoplankton communities rather than one. These communities exhibit markedly different photoacclimation profiles, with deep-water communities displaying a  $F_{470}/b_{bp}$  ratio significantly distinct from that of surface communities (Bellacicco et al. 2016; Graff et al. 2016). Thus, the model successfully discriminates phytoplankton communities according to their average size and photoacclimation status. However, when the nano- and microphytoplankton-dominated cluster is further divided into two distinct communities based on varying carotenoid composition, the model performance declines markedly, resulting in a recall score of only 40%. This low score, compared to the high predictive performance of the other models, that all include multiple fluorescence with at least two wavelengths (i.e., configurations A to E), highlights the importance of MXF for pigment-based remote classification of phytoplankton communities.

When using MXF data only (i.e., configuration E), the model recall performance drops to approximately 50%, whatever the number of predicted clusters. In comparison, models ~~conFig-d~~ with the MXF sensor and a transmissometer (i.e., configurations C) reach around 65% recall, regardless of the ~~amount~~ number of ~~cluster~~ clusters to be predicted. Adding the particulate backscattering coefficient ( $b_{bp}$ , configuration A) slightly increases the recall score to around 75% when predicting 2 or 3 clusters. This highlights the importance of incorporating additional bio-optical indices with MXF, such as  $b_{bp}$  and  $c_p$ , to enhance the phytoplankton group classification model. Furthermore, when the backscatterometer and transmissometer are present, the MXF 532-nm fluorescence channel does not improve the model performance (configuration A and B). This indicates that, overall, when bio-optical sensors measuring  $b_{bp}$  and  $c_p$  can be used as inputs, only two fluorescence wavelengths are needed to achieve an optimal classification of phytoplankton communities.

These findings demonstrate that meaningful phytoplankton taxonomic information can be retrieved from MXF signals. If the standard BGC-Argo configuration (i.e., configuration F) with one single-channel fluorescence and two optical indices ( $b_{bp}$  and  $c_p$ ) shows good prediction performances for predicting two and three clusters, adding one more fluorescence wavelength (i.e., configurations A to E) significantly improves the performance when predicting four clusters. Thus, our results demonstrate that MXF is a promising avenue for remote classification of phytoplankton community composition.

Our results also highlight that the predictive skill of individual descriptors, particularly the 532 nm fluorescence channel, depends on both the way clusters are defined and ~~on~~ the ecological context in which communities occur. When clusters retain a strong contribution of Chl $b$ -containing taxa (e.g., chlorophytes) or phycoerythrin-rich picocyanobacteria (*Synechococcus*), the  $F_{532}/F_{470}$  ratio provides valuable information on community composition. However, when such groups are merged into broader assemblages, the influence of  $F_{532}$  diminishes and the model relies more heavily on size- or biomass-sensitive proxies such as  $c_p$  and  $b_{bp}$ . Ecologically, this reflects the fact that pigment-based contrasts are strongest when communities differ in accessory pigment composition, while size-structure proxies dominate when communities are merged across pigment gradients. From a

broader perspective, this sensitivity also explains why the usefulness of  $F_{532}$  will vary geographically: in regions where *Synechococcus* or green flagellates are recurrent and occasionally abundant (e.g., coastal upwelling systems), including 532 nm excitation is expected to significantly improve classification performance the inclusion of a 532nm excitation channel may improve the discrimination of these specific groups by enhancing the excitation of accessory pigments that are less efficiently excited at shorter wavelengths, as discussed in previous studies (Morel, 1997; Saito et al., 2005). In contrast, in persistently oligotrophic waters such as the subtropical gyres, the added value of  $F_{532}$  is likely reduced.



More broadly, our results allow us to draw a general framework linking sensor configuration to achievable phytoplankton classification performance in the context of BGC-Argo observations. The level of community discrimination that can realistically be achieved depends on two interacting factors: the ecological contrast between the communities to be distinguished (e.g., differences in pigmentation, size, or photoacclimation status), and the spectral and optical information available from the sensor suite. When communities differ primarily in size structure, such as pico- versus microphytoplankton dominance, bio-optical proxies alone (i.e.,  $b_{bp}$  and  $c_p$ ) provide sufficient discriminatory power, while the MXF contributes only

630 limited additional value. In this regime, the standard BGC-Argo configuration (configuration F) already achieves recall scores above 70%. MXF becomes essential when communities share similar size structures but differ in pigment composition, for instance, when distinguishing between nano- and microphytoplankton assemblages characterized by contrasting carotenoid signatures, or between distinct picophytoplankton communities dominated by *Prochlorococcus* vs. *Synechococcus*. In these cases, adding even a single fluorescence channel (440 or 470 nm) to the standard configuration provides a statistically significant improvement in classification performance. The 532 nm channel provides additional value specifically in environments where phycoerythrin-rich taxa such as *Synechococcus* or Chlb-containing organisms are recurrent and abundant, but its contribution diminishes when such groups are absent or merged into broader assemblages. In terms of minimum sensor requirements, our results suggest that combining two fluorescence channels (440 and 470 nm) with  $b_{bp}$  and  $c_p$  represents an optimal and achievable configuration for BGC-Argo platforms. This configuration achieves over 75% recall in scenarios distinguishing (1) two phytoplankton clusters (i.e. picophytoplankton vs mixed nano- plus microphytoplankton assemblages) or (2) three clusters (i.e. a mixed nano- plus microphytoplankton, a *Synechococcus* like and a *Prochlorococcus* like assemblages). This configuration corresponds to dual-channel fluorometers already implemented on part of the BGC-Argo

635 fleet, suggesting that meaningful phytoplankton community information could be extracted from existing deployments without requiring new instrumentation. Adding a third fluorescence channel at 532 nm provides incremental benefit specifically for pigment-based discrimination at finer taxonomic resolution, and would be the recommended configuration for regions or seasons where *Synechococcus* or green flagellates are ecologically important.

640

645




Figure 6: Values of the mean weighted recall resulting from cross-validation with different numbers of clusters and different sensor combinations, either currently equipped on BGC-Argo profiling floats or feasible for future deployments: full configuration - A (F<sub>440</sub>, F<sub>470</sub>, F<sub>532</sub>, b<sub>bp</sub>, c<sub>p</sub>); dual-channel fluorescence with backscatterometer and transmissometer - configuration B (F<sub>440</sub>, F<sub>470</sub>, b<sub>bp</sub>, c<sub>p</sub>); MXF with cp - configuration C (F<sub>440</sub>, F<sub>470</sub>, F<sub>532</sub>, ~~ep~~b<sub>bp</sub>); dual-channel fluorometer with backscatterometer - configuration D (F<sub>440</sub>, F<sub>470</sub>, b<sub>bp</sub>); MXF only - configuration E (F<sub>440</sub>, F<sub>470</sub>, F<sub>532</sub>); and single-channel fluorescence with backscatterometer and transmissometer - configuration F (F<sub>470</sub>, b<sub>bp</sub>, c<sub>p</sub>). Significance of paired t-test, testing significant different within each sensor configuration, is indicated with stars (ns: p.value > 0.05; \*: p.value <= 0.05 ; \*\*: p.value <= 0.01 ; \*\*\*: p.value <= 0.001 ; \*\*\*\*: p.value <= 0.0001).

655

#### 4. Conclusion and perspectives

Monitoring phytoplankton community composition is essential for understanding marine biogeochemical cycles, particularly those related to oceanic carbon dynamics. However, this remains challenging in open-ocean environments due to the limitations of current autonomous sensor technologies and the complex bio-optical signature of mixed phytoplankton assemblages. ~~In this study, we developed and tested a machine learning (HGB) model to classify phytoplankton assemblages using *in situ* MXF and additional bio-optical measurements. The goal was to assess the potential of MXF for taxonomic discrimination of phytoplankton communities from autonomous platforms, particularly BGC-Argo profiling floats. The HGB model uses MXF signals, specifically fluorescence excited at 440, 470, and 532 nm, alongside particulate backscattering ( $b_{bp}$ ) and beam attenuation ( $c_p$ ) coefficients as input features. Taxonomic information was assessed through pigment-based clustering of field samples collected over a full annual cycle in the NW Mediterranean Sea.~~

~~Our primary objective was to evaluate the predictive skill of MXF measurements. Under laboratory conditions, MXF provided sufficient sensitivity to distinguish five key phytoplankton groups. When applied to field data, MXF alone had lower performance due to the presence of complex mixed phytoplankton assemblages, compared to monospecific cultures. However, combining MXF with additional bio-optical indices improved discrimination of phytoplankton communities.~~

~~This study specifically evaluated whether multispectral fluorescence (MXF), using three excitation channels (440, 470, and 532 nm), alone or in combination with other bio-optical measurements readily available on BGC-Argo platforms, can provide meaningful taxonomic discrimination of phytoplankton assemblages. We used a gradient boosting model, chosen for its high performance and good explainability, to assess the potential of MXF for autonomous, in-situ monitoring of phytoplankton community structure. We assessed the model's performance across different sensor configurations, reflecting realistic BGC-Argo float payloads, and across varying levels of classification complexity; (i.e., number of predicted pigment-based clusters). While existing sensor/lab conditions confirmed MXF sensitivity across five groups, field predictions required combining MXF with bio-optical indices to extract meaningful information from complex, mixed phytoplankton assemblages. Sensor configurations already implemented on BGC-Argo, such as single-wavelength fluorescence with  $b_{bp}$  and  $c_p$ , performed well for simple phytoplankton communities (two or three clusters); full separation of more complex community types (four clusters) required the richer spectral information provided by MXF. Notably, MXF enabled taxonomic discrimination within micro and nanophytoplankton dominated assemblages, even when using only two fluorescence channels, 440 and 470 nm. In contrast, bio-optical properties such as  $b_{bp}$  and  $c_p$ , which are robust proxies for particle size and biomass, were more effective at distinguishing between communities with pronounced size differences, such as pico versus microphytoplankton, i.e. strong size contrast between predicted assemblages). However, MXF becomes essential when communities share similar size distributions but differ in pigment composition, such as when distinguishing nano- from microphytoplankton assemblages or contrasting picophytoplankton communities. In these cases, combining two fluorescence channels (440 and 470 nm) with  $b_{bp}$  and  $c_p$  represents an optimal and realistic configuration, achieving over 75% recall across multiple community scenarios and corresponding to dual-channel fluorometers already deployed on part of the BGC-Argo fleet.~~

Overall, our results suggest that integrating MXF into BGC-Argo platforms could significantly enhance our ability  
690 to monitor shifts in phytoplankton communities, particularly among groups with overlapping size distributions. This  
advancement would provide valuable insights into ecosystem dynamics and the role of phytoplankton taxonomic composition  
across broad spatial and temporal scales, enabled by the autonomous sampling capabilities of BGC-Argo floats, in open-ocean  
biogeochemical cycles.

695 Future work should aim to generalize these findings across diverse oceanic regions and refine the predictive model to account  
for a broader variety of phytoplankton taxa. Building a standardized dataset linking phytoplankton community composition,  
from pico- to micro-size classes, to MXF and bio-optical measurements will be crucial for improving our understanding of  
phytoplankton dynamics through autonomous observations.

**Table A1-: Mean and standard deviation of precision and recall scores from the cross validation (repeated prediction of clusters from the machine learning model using randomly picked samples for training and testing). The standard deviation is indicated in parentheses.** Each line corresponds to a specific combination of sensors, either currently deployed on BGC-Argo profiling floats or suitable for future deployments: full configuration ( $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $b_{bp}$ ,  $c_p$ ); dual channel fluorescence with backscatterometer and transmissometer ( $F_{440}$ ,  $F_{470}$ ,  $b_{bp}$ ,  $c_p$ ); MXF with backscatterometer ( $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ,  $b_{bp}$ ); dual channel fluorometer with backscatterometer ( $F_{440}$ ,  $F_{470}$ ,  $b_{bp}$ ); MXF only ( $F_{440}$ ,  $F_{470}$ ,  $F_{532}$ ) and single channel fluorescence with backscatterometer and transmissometer ( $F_{470}$ ,  $b_{bp}$ ,  $c_p$ ).

Configuration		Cluster 1		Cluster 2		Cluster 3		Cluster 4	
		Precision	Recall	Precision	Recall	Precision	Recall	Precision	Recall
$F_{440} + F_{470} + F_{532} + b_{bp} + c_p$	A	0.68 (0.15)	0.68 (0.18)	0.67 (0.15)	0.67 (0.16)	0.86 (0.14)	0.86 (0.14)	0.58 (0.12)	0.58 (0.1)
$F_{440} + F_{470} + b_{bp} + b_{bp}$	B	0.58 (0.2)	0.59 (0.21)	0.68 (0.17)	0.67 (0.17)	0.75 (0.19)	0.74 (0.19)	0.68 (0.2)	0.66 (0.18)
$F_{440} + F_{470} + F_{532} + c_p$	C	0.62 (0.19)	0.63 (0.2)	0.71 (0.19)	0.72 (0.17)	0.86 (0.14)	0.87 (0.14)	0.57 (0.19)	0.57 (0.19)
$F_{440} + F_{470} + b_{bp}$	D	0.56 (0.18)	0.59 (0.21)	0.63 (0.19)	0.63 (0.2)	0.74 (0.22)	0.76 (0.15)	0.55 (0.18)	0.54 (0.2)
$F_{440} + F_{470} + F_{532}$	E	0.37 (0.2)	0.37 (0.19)	0.81 (0.21)	0.83 (0.21)	0.51 (0.14)	0.51 (0.14)	0.65 (0.2)	0.64 (0.2)
$F_{470} + b_{bp} + c_p$	F	0.36 (0.26)	0.37 (0.16)	0.32 (0.26)	0.33 (0.25)	0.68 (0.22)	0.69 (0.21)	0.47 (0.19)	0.47 (0.17)

**Code, data, or code and data availability**

All the code used to process raw data, perform statistical analysis and make ~~fgures~~figures shown in that paper are available on github <https://github.com/Flavi1P/mf><https://github.com/Flavi1P/mf>. Data are available upon request to the corresponding author.

715 **Supplement link**

The link to the supplement will be included by Copernicus, if applicable.

**Author contributions**

720 Flavién Petit (corresponding author) designed the study, performed data acquisition and analysis, drafted and wrote the manuscript. Julia Uitz, Hervé Claustre, Colin Roesler, Frédéric Partensky and Laurence Garczarek were involved in the design of the study, data analysis and manuscript writing. Louison Dufour, Priscillia Gourvil, Céline Dimier, Vincenzo Vellucci, Christophe Penkerc'h and David Antoine were involved in the data acquisition and analysis and manuscript writing. All authors commented on the final manuscript.

**Competing interests**

This work does not present any conflict of interest.

725 **Disclaimer**

Copernicus Publications adds a standard disclaimer: “Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.”

730 Please feel free to add disclaimer text at your choice, if applicable.

**Acknowledgements**

This paper represents a contribution to the following projects: OBOO funded by CNRS INSU via the LEFE-CYBER program; ~~EFFICACY~~ funded by the French “Agence Nationale de la Recherche” (ANR--19--~~ee02~~—CE02-0019);  
735 REFINE funded by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement N° 834177); Argo-2030 with support from the French government within the framework of the “Investissements d’Avenir” program (PIA EquipEx+) integrated within France 2030, and managed by ANR under the reference “ANR-21-ESRE-0019”; and BOUSSOLE funded by CNES and ESA (contract N° 4000119096/17/I-BG) with support of the French Oceanographic Fleet. FP was funded by a PhD grant from Sorbonne Université (Ecole Doctorale 129)  
740 and a research scientist position funded by the National Oceanography Centre.

### Financial support

Support funds and grant agreement numbers are listed as specified upon manuscript registration and reported to FundRef upon publication.

### Review statement

745 The review statement will be added by Copernicus Publications listing the handling editor as well as all contributing referees according to their status anonymous or identified.

### References

- Antoine, D., P. Guevel, J.-F. Desté, G. Bécu, F. Louis, A. J. Scott, and P. Bardey. 2008. The  
750 “BOUSSOLE” Buoy—A New Transparent-to-Swell Taut Mooring Dedicated to Marine Optics: Design, Tests, and Performance at Sea. *Journal of Atmospheric and Oceanic Technology* **25**: 968–989. doi:[10.1175/2007JTECHO563.1](https://doi.org/10.1175/2007JTECHO563.1)
- Barbieux, M. and others. 2019. Bio-optical characterization of subsurface chlorophyll maxima in the Mediterranean Sea from a Biogeochemical-Argo float database. *Biogeosciences* **16**: 1321–1342. doi:[10.5194/bg-16-1321-2019](https://doi.org/10.5194/bg-16-1321-2019)
- 755 Barbieux, M. and others. 2022. Biological production in two contrasted regions of the Mediterranean Sea during the oligotrophic period: an estimate based on the diel cycle of optical properties measured by BioGeoChemical-Argo profiling floats. *Biogeosciences* **19**: 1165–1194. doi:[10.5194/bg-19-1165-2022](https://doi.org/10.5194/bg-19-1165-2022)
- 760 Barlow, R. G., R. F. C. Mantoura, D. G. Cummings, and T. W. Fileman. 1997. Pigment chemotaxonomic distributions of phytoplankton during summer in the western Mediterranean. *Deep Sea Research Part II: Topical Studies in Oceanography* **44**: 833–850. doi:[10.1016/S0967-0645\(96\)00089-6](https://doi.org/10.1016/S0967-0645(96)00089-6)

- 765 Bellacicco, M., G. Volpe, S. Colella, J. Pitarch, and R. Santoleri. 2016. Influence of photoacclimation on the phytoplankton seasonal cycle in the Mediterranean Sea as seen by satellite. *Remote Sensing of Environment* **184**: 595–604. doi:[10.1016/j.rse.2016.08.004](https://doi.org/10.1016/j.rse.2016.08.004)
- Bidigare, R. R., J. H. Morrow, and D. A. Kiefer. 1989. Derivative analysis of spectral absorption by photosynthetic pigments in the western Sargasso Sea. *J Mar Res* **47**: 323–341. doi:[10.1357/002224089785076325](https://doi.org/10.1357/002224089785076325)
- 770 Biogeochemical-Argo Planning Group. 2016. The scientific rationale, design and implementation plan for a Biogeochemical-Argo float array, Ifremer.
- Bittig, H. C. and others. 2019. A BGC-Argo Guide: Planning, Deployment, Data Handling and Usage. *Front. Mar. Sci.* **6**. doi:[10.3389/fmars.2019.00502](https://doi.org/10.3389/fmars.2019.00502)
- Bock, N., M. Cornec, H. Claustre, and S. Duhamel. 2022. Biogeographical Classification of the Global Ocean From BGC-Argo Floats. *Global Biogeochemical Cycles* **36**: e2021GB007233. doi:[10.1029/2021GB007233](https://doi.org/10.1029/2021GB007233)
- 775 Bonnet, S. and others. 2023. Diazotrophs are overlooked contributors to carbon and nitrogen export to the deep ocean. *ISME J* **17**: 47–58. doi:[10.1038/s41396-022-01319-3](https://doi.org/10.1038/s41396-022-01319-3)
- Boss, E., W. S. Pegau, M. Lee, M. Twardowski, E. Shybanov, G. Korotaev, and F. Baratange. 2004. Particulate backscattering ratio at LEO 15 and its use to study particle composition and distribution. *Journal of Geophysical Research: Oceans* **109**. doi:[10.1029/2002JC001514](https://doi.org/10.1029/2002JC001514)
- 780 Boss, E., D. Swift, L. Taylor, P. Brickley, R. Zaneveld, S. Riser, M. J. Perry, and P. G. Strutton. 2008. Observations of pigment and particle distributions in the western North Atlantic from an autonomous float and ocean color satellite. *Limnology and Oceanography* **53**: 2112–2122. doi:[10.4319/lo.2008.53.5\\_part\\_2.2112](https://doi.org/10.4319/lo.2008.53.5_part_2.2112)
- 785 Brewin, R. J. W. and others. 2011. An intercomparison of bio-optical techniques for detecting dominant phytoplankton size class from satellite remote sensing. *Remote Sensing of Environment* **115**: 325–339. doi:[10.1016/j.rse.2010.09.004](https://doi.org/10.1016/j.rse.2010.09.004)
- Brewin, R. J. W., S. Sathyendranath, P. K. Lange, and G. Tilstone. 2014. Comparison of two methods to derive the size-structure of natural populations of phytoplankton. *Deep Sea Research Part I: Oceanographic Research Papers* **85**: 72–79. doi:[10.1016/j.dsr.2013.11.007](https://doi.org/10.1016/j.dsr.2013.11.007)
- 790 [Brewin, R.J.W., Dall’Olmo, G., Gittings, J., Sun, X., Lange, P.K., Raitsos, D.E., Bouman, H.A., Hoteit, I., Aiken, J., Sathyendranath, S., 2022. A Conceptual Approach to Partitioning a Vertical Profile of Phytoplankton Biomass Into Contributions From Two Communities. \*Journal of Geophysical Research: Oceans\* 127, e2021JC018195. <https://doi.org/10.1029/2021JC018195>](https://doi.org/10.1016/j.dsr.2013.11.007)
- 795 <https://doi.org/10.1029/2021JC018195>
- Bricaud, A., H. Claustre, J. Ras, and K. Oubelkheir. 2004. Natural variability of phytoplanktonic absorption in oceanic waters: Influence of the size structure of algal populations. *Journal of Geophysical Research: Oceans* **109**. doi:[10.1029/2004JC002419](https://doi.org/10.1029/2004JC002419)
- 800 Buesseler, K., L. Ball, J. Andrews, C. Benitez-Nelson, R. Belostock, F. Chai, and Y. Chao. 1998. Upper ocean export of particulate organic carbon in the Arabian Sea derived from thorium-

234. Deep Sea Research Part II: Topical Studies in Oceanography **45**: 2461–2487. doi:[10.1016/S0967-0645\(98\)80022-2](https://doi.org/10.1016/S0967-0645(98)80022-2)
- 805 Bustillos-Guzmán, J., H. Claustre, and C. Marty. 1995. Specific phytoplankton signatures and their relationship to hydrographic conditions in the coastal northwestern Mediterranean Sea. *Marine Ecology Progress Series* **124**: 247–258. doi:[10.3354/meps124247](https://doi.org/10.3354/meps124247)
- Cermeño, P., E. Marañón, J. Rodríguez, and E. Fernández. 2005. Large-sized phytoplankton sustain higher carbon-specific photosynthesis than smaller cells in a coastal eutrophic ecosystem. *Marine Ecology Progress Series* **297**: 51–60. doi:[10.3354/meps297051](https://doi.org/10.3354/meps297051)
- 810 Cetinić, I., M. J. Perry, E. D’Asaro, N. Briggs, N. Poulton, M. E. Sieracki, and C. M. Lee. 2015. A simple optical index shows spatial and temporal heterogeneity in phytoplankton community composition during the 2008 North Atlantic Bloom Experiment. *Biogeosciences* **12**: 2179–2194. doi:[10.5194/bg-12-2179-2015](https://doi.org/10.5194/bg-12-2179-2015)
- 815 Chase, A. P., S. J. Kramer, N. Haëntjens, E. S. Boss, L. Karp-Boss, M. Edmondson, and J. R. Graff. 2020. Evaluation of diagnostic pigments to estimate phytoplankton size classes. *Limnology and Oceanography: Methods* **18**: 570–584. doi:[10.1002/lom3.10385](https://doi.org/10.1002/lom3.10385)
- Chawla, N. V., K. W. Bowyer, L. O. Hall, and W. P. Kegelmeyer. 2002. SMOTE: Synthetic Minority Over-sampling Technique. *jair* **16**: 321–357. doi:[10.1613/jair.953](https://doi.org/10.1613/jair.953)
- 820 Chen, T., and C. Guestrin. 2016. XGBoost: A Scalable Tree Boosting System. *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*. Proceedings of the KDD ’16: The 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. ACM. 785–794.
- 825 Claustre, H. 1994. The trophic status of various oceanic provinces as revealed by phytoplankton pigment signatures. *Limnol. Oceanogr.* **39**: 1206–1210. doi:[10.4319/lo.1994.39.5.1206](https://doi.org/10.4319/lo.1994.39.5.1206)
- Claustre, H., K. S. Johnson, and Y. Takeshita. 2020. Observing the Global Ocean with Biogeochemical-Argo. *Annual Review of Marine Science* **12**: 23–48. doi:[10.1146/annurev-marine-010419-010956](https://doi.org/10.1146/annurev-marine-010419-010956)
- 830 Cornec, M. and others. 2021. Deep Chlorophyll Maxima in the Global Ocean: Occurrences, Drivers and Characteristics. *Global Biogeochemical Cycles* **35**: e2020GB006759. doi:[10.1029/2020GB006759](https://doi.org/10.1029/2020GB006759)
- 835 Cremella, B., Y. Huot, and S. Bonilla. 2018. Interpretation of total phytoplankton and cyanobacteria fluorescence from cross-calibrated fluorometers, including sensitivity to turbidity and colored dissolved organic matter. *Limnology & Ocean Methods* **16**: 881–894. doi:[10.1002/lom3.10290](https://doi.org/10.1002/lom3.10290)
- Cushing, D. H. 1989. A difference in structure between ecosystems in strongly stratified waters and in those that are only weakly stratified. *Journal of Plankton Research* **11**: 1–13. doi:[10.1093/plankt/11.1.1](https://doi.org/10.1093/plankt/11.1.1)
- Dall’Olmo, G., T. K. Westberry, M. J. Behrenfeld, E. Boss, and W. H. Slade. 2009. Direct contribution of phytoplankton-sized particles to optical backscattering in the open ocean. doi:[10.5194/bgd-6-291-2009](https://doi.org/10.5194/bgd-6-291-2009)

- 840 [Das, S.K., Frank, H.A., 2002. Pigment Compositions, Spectral Properties, and Energy Transfer Efficiencies between the Xanthophylls and Chlorophylls in the Major and Minor Pigment-Protein Complexes of Photosystem II. \*Biochemistry\* 41, 13087–13095. <https://doi.org/10.1021/bi0204802>](#)
- 845 D’Ortenzio, F., D. Iudicone, C. de Boyer Montegut, P. Testor, D. Antoine, S. Marullo, R. Santoleri, and G. Madec. 2005. Seasonal variability of the mixed layer depth in the Mediterranean Sea as derived from *in situ* profiles. *Geophysical Research Letters* **32**. doi:[10.1029/2005GL022463](https://doi.org/10.1029/2005GL022463)[10.1029/2005GL022463](https://doi.org/10.1029/2005GL022463)
- 850 El Hourany, R., M. Abboud-Abi Saab, G. Faour, C. Mejia, M. Crépon, and S. Thiria. 2019. Phytoplankton ~~Diversity~~[diversity](#) in the Mediterranean Sea ~~From Satellite Data Using Self-Organizing Maps~~[from satellitedata using self-organizing maps](#). *Journal of Geophysical Research: Oceans* **124**: 5827–5843. doi:[10.1029/2019JC015131](https://doi.org/10.1029/2019JC015131)[10.1029/2019JC015131](https://doi.org/10.1029/2019JC015131)
- 855 Escoffier, N., C. Bernard, S. Hamlaoui, A. Groleau, and A. Catherine. 2015. Quantifying phytoplankton communities using spectral fluorescence: the effects of species composition and physiological state. *Journal of Plankton Research* **37**: 233–247. doi:[10.1093/plankt/fbu085](https://doi.org/10.1093/plankt/fbu085)
- Finkel, Z. V. 2007. CHAPTER 15 - Does Phytoplankton Cell Size Matter? The Evolution of Modern Marine Food Webs, p. 333–350. *In* P.G. Falkowski and A.H. Knoll [eds.], *Evolution of Primary Producers in the Sea*. Academic Press.
- 860 Frank, H. A., and R. J. Cogdell. 2012. 8.6 Light Capture in Photosynthesis, p. 94–114. *In* E.H. Egelman [ed.], *Comprehensive Biophysics*. Elsevier.
- Graff, J. R., T. K. Westberry, A. J. Milligan, M. B. Brown, G. Dall’Omo, K. M. Reifel, and M. J. Behrenfeld. 2016. Photoacclimation of natural phytoplankton communities. *Marine Ecology Progress Series* **542**: 51–62. doi:[10.3354/meps11539](https://doi.org/10.3354/meps11539)
- 865 Grébert, T. and others. 2018. Light color acclimation is a key process in the global ocean distribution of *Synechococcus* cyanobacteria. *Proceedings of the National Academy of Sciences* **115**: E2010–E2019. doi:[10.1073/pnas.1717069115](https://doi.org/10.1073/pnas.1717069115)
- 870 Guidi, L., L. Stemann, G. A. Jackson, F. Ibanez, H. Claustre, L. Legendre, M. Picheral, and G. Gorskya. 2009. Effects of phytoplankton community on production, size, and export of large aggregates: A world-ocean analysis. *Limnology and Oceanography* **54**: 1951–1963. doi:[10.4319/lo.2009.54.6.1951](https://doi.org/10.4319/lo.2009.54.6.1951)
- Henson, S. A., R. Sanders, and E. Madsen. 2012. Global patterns in efficiency of particulate organic carbon export and transfer to the deep ocean. *Global Biogeochemical Cycles* **26**. doi:[10.1029/2011GB004099](https://doi.org/10.1029/2011GB004099)
- 875 Holtrop, T. and others. 2021. Vibrational modes of water predict spectral niches for photosynthesis in lakes and oceans. *Nat Ecol Evol* **5**: 55–66. doi:[10.1038/s41559-020-01330-x](https://doi.org/10.1038/s41559-020-01330-x)

Hu, X., R. Su, F. Zhang, X. Wang, H. Wang, and Z. Zheng. 2010. Multiple excitation wavelength fluorescence emission spectra technique for discrimination of phytoplankton. *J. Ocean Univ. China* **9**: 16–24. doi:[10.1007/s11802-010-0016-x](https://doi.org/10.1007/s11802-010-0016-x)

880 Humily, F., F. Partensky, C. Six, G. K. Farrant, M. Ratin, D. Marie, and L. Garczarek. 2013. A Gene ~~Island~~island with ~~Two Possible Configurations Is Involved~~two possible configurations is involved in ~~Chromatic Acclimation~~chromatic acclimation in ~~Marine~~marine *Synechococcus* F. Rodriguez-Valera [ed.]. *PLoS ONE* **8**: e84459. doi:[10.1371/journal.pone.0084459](https://doi.org/10.1371/journal.pone.0084459)

885 Jeffrey, S. W., R. F. C. Mantoura, S. W. Wright, International Council of Scientific Unions, and Unesco, eds. 1997. *Phytoplankton pigments in oceanography: guidelines to modern methods*, UNESCO Publishing.

Johnsen, G., and E. Sakshaug. 2007. Biooptical characteristics of PSII and PSI in 33 species (13 pigment groups) of marine phytoplankton, and the relevance for pulse-amplitude-modulated and fast-repetition-rate ~~fluorometry~~fluorometry. *Journal of Phycology* **43**: 1236–1251. doi:[10.1111/j.1529-8817.2007.00422.x](https://doi.org/10.1111/j.1529-8817.2007.00422.x)

Keller, M. D., R. C. Selvin, W. Claus, and R. R. L. Guillard. 1987. Media for the Culture of Oceanic ~~Ultraplankton~~Ultraplankton. *Journal of Phycology* **23**: 633–638. doi:[10.1111/j.1529-8817.1987.tb04217.x](https://doi.org/10.1111/j.1529-8817.1987.tb04217.x)

895 Kodama, T., Taniuchi, Y., Kasai, H., Yamaguchi, T., Nakae, M., Okumura, Y., 2022. Empirical estimation of marine phytoplankton assemblages in coastal and offshore areas using an in situ multi-wavelength excitation fluorometer. PLOS ONE 17, e0257258. <https://doi.org/10.1371/journal.pone.0257258>

900 Kramer, S. J., and D. A. Siegel. 2019. How Can Phytoplankton Pigments Be Best Used to Characterize ~~Surface Ocean Phytoplankton Groups~~surface ocean phytoplankton groups for Ocean ~~Color Remote Sensing Algorithms~~color remote sensing algorithms? *Journal of Geophysical Research: Oceans* **124**: 7557–7574. doi:[10.1029/2019JC015604](https://doi.org/10.1029/2019JC015604)

905 Lacour, L., N. Briggs, H. Claustre, M. Ardyna, and G. Dall'Olmo. 2019. The Intraseasonal Dynamics of the ~~Mixed Layer Pump~~mixed layer pump in the ~~Subpolar~~subpolar North Atlantic Ocean: A Biogeochemical-Argo Float Approach. *Global Biogeochemical Cycles* **33**: 266–281. doi:[10.1029/2018GB005997](https://doi.org/10.1029/2018GB005997)

910 Latasa, M., Scharek, R., Morán, X.A.G., Gutiérrez-Rodríguez, A., Emelianov, M., Salat, J., Vidal, M., Estrada, M., 2022. Dynamics of phytoplankton groups in three contrasting situations of the open NW Mediterranean Sea revealed by pigment, microscopy, and flow cytometry analyses. Progress in Oceanography 201, 102737. <https://doi.org/10.1016/j.pocean.2021.102737>

Lavigne, H., F. D'Ortenzio, M. Ribera D'Alcalà, H. Claustre, R. Sauzède, and M. Gacic. 2015. On the vertical distribution of the chlorophyll *a* concentration in the Mediterranean Sea: a

- 915 basin-scale and seasonal approach. *Biogeosciences* **12**: 5021–5039. doi:[10.5194/bg-12-5021-2015](https://doi.org/10.5194/bg-12-5021-2015)
- Litchman, E., P. de Tezanos Pinto, K. F. Edwards, C. A. Klausmeier, C. T. Kremer, and M. K. Thomas. 2015. Global biogeochemical impacts of phytoplankton: a trait-based perspective. *Journal of Ecology* **103**: 1384–1396. doi:[10.1111/1365-2745.12438](https://doi.org/10.1111/1365-2745.12438)
- 920 Litchman, E., P. de Tezanos Pinto, C. A. Klausmeier, M. K. Thomas, and K. Yoshiyama. 2010. Linking traits to species diversity and community structure in phytoplankton, p. 15–28. *In* L. Naselli-Flores and G. Rossetti [eds.], *Fifty years after the “Homage to Santa Rosalia”*: Old and new paradigms on biodiversity in aquatic ecosystems. Springer Netherlands.
- 925 MacIntyre, H. L., E. Lawrenz, and T. L. Richardson. 2010. Taxonomic ~~Discrimination~~discrimination of ~~Phytoplankton~~phytoplankton by ~~Spectral Fluorescence~~spectral fluorescence, p. 129–169. *In* D.J. Suggett, O. Prášil, and M.A. Borowitzka [eds.], *Chlorophyll a Fluorescence in Aquatic Sciences: Methods and Applications*. Springer Netherlands.
- 930 Marie, D., F. Partensky, D. Vaultot, and C. Brussaard. 2001. Enumeration of ~~Phytoplankton~~phytoplankton, ~~Bacteria~~bacteria, and ~~Viruses~~viruses in ~~Marine Samples~~marine samples. *Current Protocols in Cytometry* **10**: 11.11.1–11.11.15. doi:[10.1002/0471142956.cy1111s10](https://doi.org/10.1002/0471142956.cy1111s10)
- Marty, J.-C., J. Chiavérini, M.-D. Pizay, and B. Avril. 2002. Seasonal and interannual dynamics of nutrients and phytoplankton pigments in the western Mediterranean Sea at the DYFAMED time-series station (1991–1999). *Deep Sea Research Part II: Topical Studies in Oceanography* **49**: 1965–1985. doi:[10.1016/S0967-0645\(02\)00022-X](https://doi.org/10.1016/S0967-0645(02)00022-X)
- 935 Marty, J.-C., N. Garcia, and P. Raimbault. 2008. Phytoplankton dynamics and primary production under late summer conditions in the NW Mediterranean Sea. *Deep Sea Research Part I: Oceanographic Research Papers* **55**: 1131–1149. doi:[10.1016/j.dsr.2008.05.001](https://doi.org/10.1016/j.dsr.2008.05.001)
- 940 Mayot, N. and others. 2017. Influence of the Phytoplankton Community Structure on the Spring and Annual Primary Production in the Northwestern Mediterranean Sea: PHYTOPLANKTON DYNAMICS IN THE NWM. *Journal of Geophysical Research: Oceans* **122**: 9918–9936. doi:[10.1002/2016JC012668](https://doi.org/10.1002/2016JC012668)
- 945 Mena, C., P. Reglero, M. Hidalgo, E. Sintes, R. Santiago, M. Martín, G. Moyà, and R. Balbín. 2019. Phytoplankton Community Structure Is Driven by Stratification in the Oligotrophic Mediterranean Sea. *Frontiers in Microbiology* **10**.
- [Meneghin, E., Volpato, A., Cupellini, L., Bolzonello, L., Jurinovich, S., Mascoli, V., Carbonera, D., Mennucci, B., Collini, E., 2018. Coherence in carotenoid-to-chlorophyll energy transfer. Nat Commun 9, 3160. https://doi.org/10.1038/s41467-018-05596-5](https://doi.org/10.3389/fmicb.2018.02668)
- 950 Mignot, A., H. Claustre, J. Uitz, A. Poteau, F. D’Ortenzio, and X. Xing. 2014. Understanding the seasonal dynamics of phytoplankton biomass and the deep chlorophyll maximum in oligotrophic environments: A Bio-Argo float investigation. *Global Biogeochemical Cycles* **28**: 856–876. doi:[10.1002/2013GB004781](https://doi.org/10.1002/2013GB004781)

- 955 Moore, L., R. Goericke, and S. Chisholm. 1995. Comparative physiology of *Synechococcus* and *Prochlorococcus*: influence of light and temperature on growth, pigments, fluorescence and absorptive properties. *Mar. Ecol. Prog. Ser.* **116**: 259–275. doi:[10.3354/meps116259](https://doi.org/10.3354/meps116259)
- Morel, A. 1997. Consequences of a *Synechococcus* bloom upon the optical properties of oceanic (case 1) waters. *Limnology & Oceanography* **42**: 1746–1754. doi:[10.4319/lo.1997.42.8.1746](https://doi.org/10.4319/lo.1997.42.8.1746)
- 960 Morel, F. M. M. 2008. The co-evolution of phytoplankton and trace element cycles in the oceans. *Geobiology* **6**: 318–324. doi:[10.1111/j.1472-4669.2008.00144.x](https://doi.org/10.1111/j.1472-4669.2008.00144.x)
- [Organelli, E., Bricaud, A., Antoine, D., Uitz, J., 2013. Multivariate approach for the retrieval of phytoplankton size structure from measured light absorption spectra in the Mediterranean Sea \(BOUSSOLE site\). \*Appl. Opt.\*, \*\*AO 52\*\*, 2257–2273. <https://doi.org/10.1364/AO.52.002257>](https://doi.org/10.1364/AO.52.002257)
- 965 [Organelli, E., G. Dall’Olmo, R. J. W. Brewin, F. Nencioli, and G. A. Tarran. 2020. Drivers of spectral optical scattering by particles in the upper 500 m of the Atlantic Ocean. \*Opt. Express\*, \*\*OE 28\*\*: 34147–34166. doi:10.1364/OE.408439](https://doi.org/10.1364/OE.408439)
- 970 [Palenik, B., 2001. Chromatic Adaptation in Marine \*Synechococcus\* Strains. \*Applied and Environmental Microbiology\* \*\*67\*\*, 991–994. <https://doi.org/10.1128/AEM.67.2.991-994.2001>](https://doi.org/10.1128/AEM.67.2.991-994.2001)
- Parkhill, J.-P., G. Maillet, and J. J. Cullen. 2001. Fluorescence-Based Maximal Quantum Yield-based maximal quantum yield for P<sub>siII</sub> as a Diagnostic of Nutrient Stress. *Journal of Phycology* **37**: 517–529. doi:[10.1046/j.1529-8817.2001.037004517.x](https://doi.org/10.1046/j.1529-8817.2001.037004517.x)
- 975 [Pittera, J., F. Humily, M. Thorel, D. Grulois, L. Garczarek, and C. Six. 2014. Connecting thermal physiology and latitudinal niche partitioning in marine \*Synechococcus\*. \*ISME J\* \*\*8\*\*: 1221–1236. doi:10.1038/ismej.2013.228](https://doi.org/10.1038/ismej.2013.228)
- Poryvkina, L., S. Babichenko, S. Kaitala, H. Kuosa, and A. Shalapjonok. 1994. Spectral fluorescence signatures in the characterization of phytoplankton community composition. *Journal of Plankton Research* **16**: 1315–1327. doi:[10.1093/plankt/16.10.1315](https://doi.org/10.1093/plankt/16.10.1315)
- 980 Proctor, C. W., and C. S. Roesler. 2010. New insights on obtaining phytoplankton concentration and composition from in situ multispectral Chlorophyll fluorescence: In situ phytoplankton composition. *Limnol. Oceanogr. Limnology and Oceanography Methods* **8**: 695–708. doi:[10.4319/lom.2010.8.0695](https://doi.org/10.4319/lom.2010.8.0695)
- 985 [Ras, J., H. Claustre, and J. Uitz. 2008. Spatial variability of phytoplankton pigment distributions in the Subtropical South Pacific Ocean: comparison between in situ and predicted data. 17.](https://doi.org/10.1002/2017JC013067)
- Rembauville, M. and others. 2017. Plankton Assemblage Estimated with BGC-Argo Floats in the Southern Ocean: Implications for Seasonal Successions and Particle Export: PLANKTON ASSEMBLAGE BGC-ARGO. *Journal of Geophysical Research: Oceans* **122**: 8278–8292. doi:[10.1002/2017JC013067](https://doi.org/10.1002/2017JC013067)
- 990 Reynolds, C. S. 2006. *The Ecology of Phytoplankton*, Cambridge University Press.

- 995 Rippka, R. and others. 2000. *Prochlorococcus marinus* Chisholm et al. 1992 *subsp. pastoris subsp. nov.* strain PCC 9511, the first axenic chlorophyll  $a_2/b_2$ -containing cyanobacterium (Oxyphotobacteria). International Journal of Systematic and Evolutionary Microbiology **50**: 1833–1847. doi:[10.1099/00207713-50-5-1833](https://doi.org/10.1099/00207713-50-5-1833)
- 1000 Roesler, C. S., and E. Boss. 5 In Situ Measurement of the ~~Inherent Optical Properties~~  
~~inherent optical properties~~ (IOPs) and ~~Potential~~  
~~potential~~ for ~~Harmful Algal Bloom Detection~~  
~~harmful algal bloom detection~~ and ~~Coastal Ecosystem Observations~~  
~~coastal ecosystem observations~~.
- Rousseaux, C. S., and W. W. Gregg. 2014. Interannual Variation in Phytoplankton Primary Production at A Global Scale. Remote Sensing **6**: 1–19. doi:[10.3390/rs6010001](https://doi.org/10.3390/rs6010001)
- 1005 Saito, M. A., G. Rocap, and J. W. Moffett. 2005. Production of cobalt binding ligands in a *Synechococcus* feature at the Costa Rica upwelling dome. ~~Limnol. Oceanogr.~~ **50**: ~~279–290.~~ doi:[10.4319/lo.2005.50.1.0279](https://doi.org/10.4319/lo.2005.50.1.0279)  
~~Limnology and Oceanography~~ **50**: ~~279–290.~~ doi:[10.4319/lo.2005.50.1.0279](https://doi.org/10.4319/lo.2005.50.1.0279)
- 1010 Sauzède, R., H. Claustre, C. Jamet, J. Uitz, J. Ras, A. Mignot, and F. D’Ortenzio. 2015. Retrieving the vertical distribution of chlorophyll *a* concentration and phytoplankton community composition from *in situ* fluorescence profiles: A method based on a neural network with potential for global-scale applications. Journal of Geophysical Research: Oceans **120**: 451–470. doi:[10.1002/2014JC010355](https://doi.org/10.1002/2014JC010355)
- Schmechtig, C., H. Claustre, A. Poteau, and F. D’Ortenzio. 2018a. Bio-Argo quality control manual for the Chlorophyll-A concentration, Ifremer.
- 1015 Schmechtig, C., A. Poteau, H. Claustre, F. D’Ortenzio, G. Dall’Olmo, and E. Boss. 2018b. Processing Bio-Argo particle backscattering at the DAC level, Ifremer.
- Seppälä, J., and M. Balode. 1998. The use of spectral fluorescence methods to detect changes in the phytoplankton community, p. 207–217. *In* T. Tamminen and H. Kuosa [eds.], Eutrophication in Planktonic Ecosystems: Food Web Dynamics and Elemental Cycling: Proceedings of the Fourth International PELAG Symposium, held in Helsinki, Finland, 26–30 August 1996. Springer Netherlands.
- 1020 Shima, S., R. P. Ilagan, N. Gillespie, B. J. Sommer, R. G. Hiller, F. P. Sharples, H. A. Frank, and R. R. Birge. 2003. Two-~~Photon~~  
~~photon~~ and ~~Fluorescence Spectroscopy~~  
~~fluorescence spectroscopy~~ and the ~~Effect~~  
~~effect~~ of ~~Environment~~  
~~environment~~ on the ~~Photochemical Properties~~  
~~photochemical properties~~ of ~~Peridinin~~  
~~peridinin~~ in ~~Solutions~~  
~~solution~~ and in the ~~Peridinin-Chlorophyll-Protein~~  
~~peridinin-chlorophyll-protein~~ from *Amphidinium carterae*. J. Phys. Chem. A **107**: 8052–8066. doi:[10.1021/jp022648z](https://doi.org/10.1021/jp022648z)
- 1025 Shwartz-Ziv, R., and A. Armon. 2022. Tabular data: Deep learning is not all you need. Information Fusion **81**: 84–90. doi:[10.1016/j.inffus.2021.11.011](https://doi.org/10.1016/j.inffus.2021.11.011)
- 1030 Six, C., J. Thomas, B. Brahamsha, Y. Lemoine, and F. Partensky. 2004. Photophysiology of the marine cyanobacterium *Synechococcus* sp. WH8102, a new model

organism. *Aquat. Microb. Ecol.* **35**: 17–29. doi:[10.3354/ame035017](https://doi.org/10.3354/ame035017)  
*Aquatic Microbial Ecology* **35**: 17–29. doi:[10.3354/ame035017](https://doi.org/10.3354/ame035017)

1035 Six, C., J.-C. Thomas, L. Garczarek, M. Ostrowski, A. Dufresne, N. Blot, D. J. Scanlan, and F. Partensky. 2007. Diversity and evolution of phycobilisomes in marine ~~*Synechococcus* spp.~~*Synechococcus* spp.: a comparative genomics study. *Genome Biology* **8**: R259. doi:[10.1186/gb-2007-8-12-r259](https://doi.org/10.1186/gb-2007-8-12-r259)  
[10.1186/gb-2007-8-12-r259](https://doi.org/10.1186/gb-2007-8-12-r259)

Slade, W. H., and E. Boss. 2015. Spectral attenuation and backscattering as indicators of average particle size. *Appl. Opt., AO* **54**: 7264–7277. doi:[10.1364/AO.54.007264](https://doi.org/10.1364/AO.54.007264)

1040 ~~Steglich, C., Mullineaux, C.W., Teuchner, K., Hess, W.R., Lokstein, H., 2003. Photophysical properties of Prochlorococcus marinus SS120 divinyl chlorophylls and phycoerythrin in vitro and in vivo. FEBS Letters 553, 79–84. https://doi.org/10.1016/S0014-5793(03)00971-2~~

1045 ~~Steglich, C., Frankenberg-Dinkel, N., Penno, S., Hess, W.R., 2005. A green light-absorbing phycoerythrin is present in the high-light-adapted marine cyanobacterium Prochlorococcus sp. MED4. Environmental Microbiology 7, 1611–1618. https://doi.org/10.1111/j.1462-2920.2005.00855.x~~

Terrats, L. and others. 2023. BioGeoChemical-Argo Floats Reveal Stark Latitudinal Gradient in the Southern Ocean ~~Deep Carbon Flux Driven by Phytoplankton Community Composition.~~*deep carbon flux driven by phytoplankton community composition.* *Global Biogeochemical Cycles* **37**: e2022GB007624. doi:[10.1029/2022GB007624](https://doi.org/10.1029/2022GB007624)  
[10.1029/2022GB007624](https://doi.org/10.1029/2022GB007624)

1050 Terrats, L., H. Claustre, M. Cornec, A. Mangin, and G. Neukermans. 2020. Detection of Coccolithophore Blooms With BioGeoChemical-Argo Floats. *Geophysical Research Letters* **47**: e2020GL090559. doi:[10.1029/2020GL090559](https://doi.org/10.1029/2020GL090559)

1055 Thibodeau, P. S., C. S. Roesler, S. L. Drapeau, S. G. Prabhu Matondkar, J. I. Goes, and P. J. Werdell. 2014. Locating *Noctiluca miliaris* in the Arabian Sea: An optical proxy approach. *Limnology and Oceanography* **59**: 2042–2056. doi:[10.4319/lo.2014.59.6.2042](https://doi.org/10.4319/lo.2014.59.6.2042)

1060 Twardowski, M. S., E. Boss, J. B. Macdonald, W. S. Pegau, A. H. Barnard, and J. R. V. Zaneveld. 2001. A model for estimating bulk refractive index from the optical backscattering ratio and the implications for understanding particle composition in case I and case II waters. *Journal of Geophysical Research: Oceans* **106**: 14129–14142. doi:[10.1029/2000JC000404](https://doi.org/10.1029/2000JC000404)

Uitz, J., H. Claustre, A. Morel, and S. B. Hooker. 2006. Vertical distribution of phytoplankton communities in open ocean: An assessment based on surface chlorophyll. *Journal of Geophysical Research* **111**. doi:[10.1029/2005JC003207](https://doi.org/10.1029/2005JC003207)

1065 Uitz, J. U., Y. Huot, F. Bruyant, M. Babin, and H. Claustre. 2008. Relating phytoplankton photophysiological properties to community structure on large scales. *Limnology and Oceanography* **53**: 614–630. doi:[10.4319/lo.2008.53.2.0614](https://doi.org/10.4319/lo.2008.53.2.0614)

Veldhuis, M. J. W., K. R. Timmermans, P. Croot, and B. van der Wagt. 2005. Picophytoplankton; a comparative study of their biochemical composition and photosynthetic properties. *Journal of Sea Research* **53**: 7–24. doi:[10.1016/j.seares.2004.01.006](https://doi.org/10.1016/j.seares.2004.01.006)

- 1070 Vidussi, F., H. Claustre, B. B. Manca, A. Luchetta, and J.-C. Marty. 2001. Phytoplankton pigment distribution in relation to upper thermocline circulation in the eastern Mediterranean Sea during winter. *J. Geophys. Res.* **106**: 19939–19956. doi:[10.1029/1999JC000308](https://doi.org/10.1029/1999JC000308)
- 1075 [Xu, Q., Wang, S., Sukigara, C., Goes, J.I., Gomes, H. do R., Matsuno, T., Zhu, Y., Xu, Y., Luang-on, J., Watanabe, Y., Yoo, S., Ishizaka, J., 2022. High-Resolution Vertical Observations of Phytoplankton Groups Derived From an in-situ Fluorometer in the East China Sea and Tsushima Strait. \*Front. Mar. Sci.\* \*\*8\*\*. <https://doi.org/10.3389/fmars.2021.756180>](https://doi.org/10.3389/fmars.2021.756180)
- Yentsch, C. S., and D. W. Menzel. 1963. A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. *Deep Sea Research and Oceanographic Abstracts* **10**: 221–231. doi:[10.1016/0011-7471\(63\)90358-9](https://doi.org/10.1016/0011-7471(63)90358-9)
- 1080 Yentsch, C. S., and D. A. Phinney. 1985. Spectral fluorescence: an ataxonomic tool for studying the structure of phytoplankton populations\*. *Journal of Plankton Research* **7**: 617–632. doi:[10.1093/plankt/7.5.617](https://doi.org/10.1093/plankt/7.5.617)
- Zhang, X., L. Hu, and M.-X. He. 2009. Scattering by pure seawater: Effect of salinity. *Opt. Express* **17**: 5698. doi:[10.1364/OE.17.005698](https://doi.org/10.1364/OE.17.005698)
- 1085 [Zhang, Q., Huang, Y., An, S., 2025. Quantification of phytoplankton groups using in-situ multi-excitation chlorophyll fluorescence measurements and machine learning \(mf-ML\). \*Algal Research\* \*\*90\*\*, 104155. <https://doi.org/10.1016/j.algal.2025.104155>](https://doi.org/10.1016/j.algal.2025.104155)
- 1090 Zhang, Y., X. He, Y. Bai, T. Li, X. Jin, D. Wang, and F. Gong. 2025. Satellite estimation of the spectral power exponent of particulate backscattering coefficient in the global ocean. *Opt. Express, OE* **33**: 5411–5428. doi:[10.1364/OE.545222](https://doi.org/10.1364/OE.545222)