

Resolving distribution and controls of terrigenous and marine particulate organic matter across an energetic shelf


Yu-Shih Lin¹, Shu-Ying Chuang¹, Yuan-Pin Chang¹, Chieh-Wei Hsu², Hui-Ling Lin¹, James T. Liu¹, Wei-Jen Huang¹

5 ¹Department of Oceanography, National Sun Yat-sen University, Kaohsiung, 804201, Taiwan

²Cross College Elite Tech Program, National Kaohsiung University of Science and Technology, Kaohsiung, 811532, Taiwan

Correspondence to: Wei-Jen Huang (wjhuang29@mail.nsysu.edu.tw)

Abstract. We assess the sources, distribution and controls of particulate organic matter (POM) across the northeastern Taiwan Strait, where monsoonal forcing, water-mass mixing, riverine inputs and sediment resuspension modulate particle dynamics. By integrating lignin biomarkers, bulk geochemistry, and sedimentary constraints within a two-step quantification approach, we demonstrate the influence of river discharge, plume intrusions, and seafloor resuspension on the distribution of terrigenous POM. Terrigenous particulate organic carbon (POC_{terr}) represents a minor component in most water samples but becomes substantial in resuspension-dominated layers. Combining estimated POC_{terr} with modeled current velocities yields an export flux of $\sim 243 \pm 56 \text{ kt C yr}^{-1}$, consistent with the regional imbalance between riverine input and sedimentary burial. After correction for terrigenous influence, bulk POM parameters reflect nutrient supply, photoacclimation, and temperature-dependent variation in stable carbon isotopic ($\delta^{13}\text{C}$) composition. Comparisons with co-sampled surface sediments show that biomarker signals are preserved more faithfully than $\delta^{13}\text{C}$ of organic matter, which is strongly modulated by lateral transport. This study provides a practical framework for quantifying terrigenous and marine POM in continental-shelf settings and offers improved constraints for interpreting source-to-sink processes and sedimentary archives.



20

Graphical abstract.

1 Introduction


Continental shelves are widely recognized as critical interfaces in the global carbon cycle, although they represent only 5 % of the ocean area (Dunne et al., 2007). These environments receive substantial inputs of terrigenous and marine organic matter (OM) and are estimated to account for up to 85 % of global organic carbon (OC) burial (Burdige, 2005, 2007). To constrain the fate and long-term sequestration efficiency of OM in individual shelf systems, quantitative characterization of marine and terrigenous OM sources is essential. Extensive research has focused on characterizing the sources of OM in shelf sediments (e.g., Bao et al., 2018; Tao et al., 2023; Wei et al., 2020), yet the provenance of particulate organic matter (POM) suspended in the water column remains insufficiently resolved. This knowledge gap limits our understanding of modern source-to-sink processes and undermines confidence in paleoenvironmental reconstructions based on sedimentary OM.

Bulk geochemical parameters of OM, such as atomic C/N ratio, the mass ratio of OC to chlorophyll a (OC/Chl), and the stable carbon isotopic ratio of OC ($\delta^{13}\text{C}_{\text{OC}}$), have been widely used to constrain POM sources (e.g., Gao et al., 2014; Guo et al., 2015; Lee et al., 2023). However, their capacity to distinguish complex POM mixtures in shelf waters is limited. Shelf waters receive terrigenous POM inputs from rivers and from sediment resuspension, and these two sources may differ in bulk properties depending on the extent of OM partitioning during transport and resuspension (Lin et al., 2025a). In addition, marine plankton communities, particularly primary producers, can produce bulk POM signatures that overlap with those of terrigenous OM (e.g., Geider, 1987; Laws et al., 1995; Martiny et al., 2013). This ambiguity is illustrated by the frequent observation of ^{13}C -depleted POM in subsurface and bottom shelf waters (e.g., Wu et al., 2003; Huang et al., 2020; Lee et al., 2023). In contrast to open-ocean settings, where a terrigenous source for the low- $\delta^{13}\text{C}$ POM in the lower euphotic zone can generally be excluded (Close and Henderson, 2020), the proximity of shelf waters to land makes it difficult to determine whether such isotopic signals primarily reflect terrigenous inputs or in situ marine processes. To alleviate these complications, previous studies have focused on sampling layers where a single POM source is expected to dominate, such as the deep Chl maximum (DCM) or the benthic nepheloid layer (e.g., Liu et al., 2018a; Sun et al., 2024). Although informative for process-oriented investigations, these strategies do not provide an integrated view of POM mixing and transport throughout the water column.

Two complementary approaches may help address this limitation. The first involves the use of biomarkers that provide diagnostic information on specific OM sources. Lignin is an abundant and stable biopolymer found exclusively in the cell walls of vascular plants (Benner et al., 1987; Hedges et al., 1997). Because vascular plants are largely restricted to land, lignin serves as unambiguous tracers for terrigenous OM. Lignin has been widely applied to trace terrigenous inputs in marine sediments (e.g., Bianchi et al., 2018), but its use in POM studies remains limited. The second approach involves characterizing surface sediments collected concurrently with POM to constrain the contribution from resuspension. This strategy was implemented by Sun et al. (2024), who quantitatively evaluated how sediment reworking influences the distribution and degradation of terrigenous POM. Together, these constraints offer a way to better distinguish terrigenous

contributions to POM and to clarify how bulk geochemical parameters relate to the condition of marine plankton
55 communities.

The overarching goal of this study is to provide a full-water-column assessment of the sources, distribution, and controls of
POM in the northeastern Taiwan Strait, a shallow and energetic shelf system (Fig. 1). The properties and spatial distribution
of OM in surface sediments collected concurrently with the POM samples were presented in our earlier study (Lin et al.,
2025a). The present work builds on this foundation through three objectives. First, we quantify terrigenous particulate
60 organic carbon (POC_{terr}), defined as the sum of biospheric and petrogenic OC, using lignin and the stable carbon isotopic
composition of organic carbon ($\delta^{13}\text{C}_{\text{OC}}$) as key tracers within an integrated two-step quantitative approach. The inclusion of
lignin provides an additional constraint for identifying the sources of low- $\delta^{13}\text{C}$ POM, which is also observed in our study
area. Second, we examine the biogeochemical characteristics of the remaining marine POM and evaluate the associated
physiological controls. Finally, we establish source-to-sink coherence by comparing POM characteristics in the water
65 column, which reflect short-term transport processes, with the OM composition recorded in surface sediments. This
comparison provides insights into OM transport dynamics and supports more robust interpretations of OM proxies in shelf
sediment archives.



70 Figure 1: Study area and sampling sites. (a) Study area relative to China and Taiwan. The SEATS station (Liu et al., 2007) is also
 75 shown. ECS, East China Sea; LS, Luzon Strait; SCS, South China Sea; TS, Taiwan Strait; WPS, West Philippine Sea. (b) Seawater sampling sites from the cruise NOR3-0104. The extent of mud belts (mean grain size < 63 μm) was based on the
 isosurface map of sediment grain size (Lin et al., 2025a). (c) Bathymetry and summer flow field (redrawn from Jan et al., 2010).
 Also shown are seabed sediment sampling sites from cruise NOR3-0104 (Lin et al., 2025a), legacy stations from cruise OR3-632
 (Ocean Data Bank, 2025), and Transect T1 from Wong et al. (2015). Bathymetric features: Changyun Rise (CYR), Kuanyin
 Depression (KYD), Penghu Channel (PHC), Penghu Islands (PHI), and Taiwan Bank (TB). Currents or water masses: KS, Kuroshio; SCS(S)W, South China Sea (Surface) Water.

2 Material and Methods

2.1 Study area

80 The Taiwan Strait is a shallow and energetic conduit linking the South China Sea and East China Sea (Fig. 1). Bounded by
 the Chinese coast to the west and the island of Taiwan to the east, the strait averages ~60 m in depth, extends ~350 km in
 length, and spans ~180 km in width (Jan et al., 2002). Its hydrography is highly dynamic, shaped by the interplay of complex
 bathymetry, monsoonal forcing, and riverine inputs.

Because sampling was conducted in summer, the following description emphasizes the hydrographic conditions of this season. During summer, under the influence of the southwest monsoon, the Taiwan Strait exhibits a net northeastward transport (Jan et al., 2002). Oceanic waters from the northern South China Sea enter the strait either along the shelf off southeastern China or through the funnel-shaped Penghu Channel, which serves as the primary pathway for volume transport during this season ($\sim 1.2 \times 10^6 \text{ m}^3 \text{ s}^{-1}$; Jan et al., 2002). The inflow is dominated by the South China Sea Water (SCSW), reflecting the limited intrusion of the Kuroshio into the northern South China Sea through the Luzon Strait during summer (Jan et al., 2006, 2010). The upper water column, typically above 40 m, is characterized by the South China Sea Surface Water (SCSSW), a brackish water mass influenced by Pearl River discharge (Bai et al., 2015; Jan et al., 2006). In the northeastern Taiwan Strait, which is the focus of this study, additional freshwater inputs are supplied by small rivers from Taiwan. Tidal dynamics are dominated by the semidiurnal M2 constituent, which has a period of 12.42 hours and produces strong oscillatory currents particularly in the Penghu Channel and Kuanyin Depression. Tidal current amplitudes decreased from $\sim 0.8 \text{ m s}^{-1}$ at the northeast and southeast entrances to $\sim 0.2 \text{ m s}^{-1}$ in the central strait (Wang et al., 2003).

The modern sedimentary regime of the Taiwan Strait has been documented previously (Huh et al., 2011; Liu et al., 2018b). The region receives substantial sediment from both the distal Yangtze River and the proximal small mountainous rivers of Taiwan, supporting a mud-rich deposition system that extends for more than 1,000 km. In the northeastern Taiwan Strait, sedimentation is organized into two mud belts that are primarily sourced from Taiwan (Horng and Huh, 2011; Lin et al., 2025a). The Taiwan Along-Shore Mud Belt, referred to here as the nearshore mud belt, forms as fine sediments are deposited near river mouths and subsequently transported northward through resuspension-advection processes. This belt is enriched in fresh terrigenous OM, largely derived from vascular plants. Further offshore, the Cross-Shelf Mud Belt, referred to here as the offshore mud belt, spatially overlaps with a depocenter characterized by high sedimentation rates and contains terrigenous OM predominantly of petrogenic origin. These characteristics suggest that this mud belt is formed through hypopycnal or other gravity-driven transport processes triggered by typhoons and floods.

Primary production, determined across the Changyun Rise, exhibits pronounced seasonality, with summer productivity reaching $664 \pm 270 \text{ mg C m}^{-2} \text{ d}^{-1}$ (Tseng et al., 2020). The summer maximum corresponds to enhanced offshore Chl concentrations integrated over the euphotic zone. The elevated offshore production during summer has been attributed to nutrient supply transported from upwelling zones near the Taiwan Bank and the Penghu Islands.

110 2.2 Fieldwork

A total of 21 sites in the northeastern Taiwan Strait were surveyed aboard the RV *New Ocean Researcher 3* during cruise NOR3-0104 (13–21 June 2022) (Table A1 in Appendix A). These sites were arranged along three SW–NE transects (X, Y, and Z), spanning the nearshore and offshore mud belts (Fig. 1b). At each station, we deployed a conductivity-temperature-depth (CTD) rosette system (SBE911plus, Sea-Bird Scientific) equipped with 12-L Niskin bottles for simultaneous acquisition of hydrographic data and water samples. Dissolved oxygen, fluorescence, beam attenuation, and photosynthetically active radiation were monitored using onboard sensors (SBE43 and C-Star, Sea-Bird Scientific; Aqua

Tracka III and PAR sensor, Chelsea Instruments Ltd.). Surface water samples from the air-sea interface were collected using a clean bucket, and their temperature and salinity were immediately measured with a portable conductivity meter (Cond 3110, WTW).

120 Each water sample was divided into three or four aliquots for different purposes. For bulk and Chl analyses, 1–2 L of water was filtered through pre-combusted glass fiber filters (GF75, Advantec; diameter: 25 mm, pore size: 0.3 μm). For lignin analysis, 3–4 L of water was filtered using the same filter type but with a larger diameter (47 mm). All filters were stored in the dark at $-20\text{ }^{\circ}\text{C}$ until analysis. Additional aliquots for carbonate chemistry ($n = 28$; Table A1) were collected following Huang et al. (2012, 2020). These samples were transferred into 250-mL borosilicate glass bottles fitted with a drip-free
125 polypropylene pouring ring, poisoned with 60 μL of saturated HgCl_2 solution, sealed with a screw cap, and stored in the dark at room temperature.

2.3 Analytical procedures

Total suspended matter (TSM) concentrations were determined gravimetrically. Filters were first examined under a microscope to remove visible zooplankton and plastic debris. Decalcified filters were analyzed for particulate organic carbon
130 (POC), particulate organic nitrogen (PON), and $\delta^{13}\text{C}_{\text{OC}}$ using an elemental analyzer coupled to an isotope ratio mass spectrometer (Flash 2000 and Delta V Plus; Thermo Fisher Scientific), following procedures detailed in Lin et al. (2020a). The relative analytical error was 3 % for POC and PON, yielding a 4 % relative uncertainty for calculated atomic C/N or N/C ratios. The absolute error for $\delta^{13}\text{C}_{\text{OC}}$ was 0.4 ‰.

For Chl analysis, filter samples were extracted in cold acetone within one month after collection, and Chl concentrations
135 were quantified fluorometrically (10-AU; Turner Designs, Inc.) following Aminot and Rey (2001).

Lignin phenols, recognized biomarkers diagnostic of vascular plant inputs, were extracted using cupric oxide oxidation (Hedges and Ertel, 1982). Filters and reagents were placed into a polytetrafluoroethylene liner, to which 8 mL of argon-purged 2 N NaOH was added. After purging the headspace with argon, the liner was sealed in a stainless-steel vessel and heated at $170\text{ }^{\circ}\text{C}$ for 3 h. The resulting hydrolysate was acidified with HCl and spiked with the internal standard ethyl
140 vanillin, and the target compounds were extracted into ethyl acetate via liquid-liquid extraction. The extracts, condensed passively at $40\text{ }^{\circ}\text{C}$, were derivatized using a 1:1 mixture of *N,O*-bis(trimethylsilyl)trifluoroacetamide and pyridine to a final volume of 10 μL , and analyzed by gas chromatography-mass spectrometry (7890A and 5957C; Agilent Technologies, Inc.) using an HP-5MS capillary column (30 m \times 0.25 mm, film thickness 0.25 μm ; Technologies, Inc.). Method detection limits, calculated as 3.14 times the standard deviation of replicate low-concentration standards ($n = 6$) following U.S. EPA
145 guidelines, were about 15 ng/L for the sum of eight lignin monomers. Replicate analysis of a sediment standard indicated 5–20 % variability among individual phenols. We report the sum of eight lignin monomers per unit volume of water ($\Sigma 8$), OC normalized concentration ($\Lambda 8$), and the mass ratio of vanillic acid to vanillin ($(\text{Ad}/\text{Al})_{\text{v}}$) (Hedges and Ertel, 1982). A higher $(\text{Ad}/\text{Al})_{\text{v}}$ ratio indicates more extensive lignin degradation.

Dissolved inorganic carbon (DIC) was measured using a DIC analyzer (AS-C3, Apollo SciTech, LLC). Samples were acidified with 10 % phosphoric acid, and the released CO₂ was quantified via a CO₂ analyzer (LI-7000, LI-COR, Inc.). Analytical quality was ensured using certified reference materials from Andrew Dickson at the Scripps Institute of Oceanography, yielding precision and accuracy better than 0.1 % (Huang et al., 2012). pH measurements were performed spectrophotometrically (Varian Cary-50, Agilent Technologies, Inc.) in a 10-cm quartz cell at 25 ± 0.5 °C (pH₂₅), following Clayton and Byrne (1993). Measurement precision, assessed using unpurified *m*-cresol purple, was < 0.003.

155 3 Data

3.1 Data sources and previously published data

A subset of the hydrographic observations used in this study was previously reported in Lin et al. (2025a), which focused primarily on sediment geochemistry. In that study, Chl, surface-water salinity, bottom-water temperature, and bottom-water TSM were used only to provide environmental context. Lin et al. (2025b), archived in Zenodo, is the companion dataset for the present study and provides the underlying measurements, including POM geochemical data analyzed and discussed here for the first time.

3.2 Processing of hydrographic data

A total of 64 samples were analyzed to calibrate sensor measurements of beam attenuation (Atten) and fluorescence (FL). The resulting calibration equations are:

$$165 \text{ TSM} = 2.6427 \times \text{Atten} + 0.8053, r^2 = 0.90 \quad (1)$$

$$\text{Chl} = 0.7176 \times \text{FL} + 0.2523, r^2 = 0.74 \quad (2)$$

where TSM is in mg L⁻¹, Atten in m⁻¹, Chl in µg L⁻¹, and FL in µg L⁻¹.

The euphotic depth (*Z_e*) was defined as the depth of 1 % surface light penetration. Photosynthetically active radiation measurements were available only for 11 daytime stations (Table A1). For the remaining sites, *Z_e* was estimated from bottom water depth (*Z_{bot}*) using the linear relationship derived from these 11 stations:

$$170 \text{ } Z_e = 0.5402 \times Z_{\text{bot}} + 3.0745, r^2 = 0.83 \quad (3)$$

where both *Z_e* and *Z_{bot}* are in meters.

To provide regional hydrographic context for our observations, the following summer temperature-salinity (T-S) datasets were incorporated into the analysis. The first consists of typical SCSW and Kuroshio Water compiled by Jan et al. (2010). The second comprises observations from Transect T1, which extends from the continental shelf west of the Taiwan Bank to the open South China Sea (Fig. 1c; Wong et al., 2015).

3.3 Processing of geochemical data

3.3.1 Estimation of POC_{terr}

We developed a two-step approach to quantify POC_{terr} (details in Texts B1 and B2 of Appendix B). Figure 2 summarizes the workflow. This approach combines the complementary strengths of lignin biomarkers and $\delta^{13}\text{C}$ in diagnosing and quantifying terrigenous OM in the water column.

In the first step, lignin concentrations were used primarily to assess the relative abundance of terrigenous OM, particularly in low- $\delta^{13}\text{C}$ samples where source attribution based on isotopic composition alone may be ambiguous. Elevated $\Delta 8$ values were expected if low- $\delta^{13}\text{C}$ POM contained a greater terrigenous contribution. The spatial patterns of lignin and its correlation with environmental variables indicated three potential sources (see Sect. 4.3; Table A2): *i*) Pearl River TSM for offshore surface waters, *ii*) Taiwanese river TSM for nearshore waters, and *iii*) seabed sediments for benthic nepheloid layers.

Following source identification, POC_{terr} was estimated from lignin concentrations assuming characteristic lignin-to- POC_{terr} ratios ($\Delta 8_{\text{terr}}$) in the source material:

$$\text{POC}_{\text{terr}} = \Sigma 8_{\text{m}} / \Delta 8_{\text{terr,s}} \quad (4)$$

where the subscript m denotes measured POM data and s the inferred source. Source-specific $\Delta 8_{\text{terr,s}}$ values were taken from Zhang et al. (2014) and Lin et al. (2025a) (Table A3). Mixing equations were then applied to derive POC, N/C, and $\delta^{13}\text{C}_{\text{OC}}$ values corrected for terrigenous inputs (POC_{corr} , N/C_{corr} , $\delta^{13}\text{C}_{\text{OC,corr}}$).


However, lignin-based estimates of POC_{terr} are subject to uncertainty because $\Delta 8_{\text{terr,s}}$ can vary during particle transport (Wakeham et al., 2009). This limitation motivated the second step, in which a $\delta^{13}\text{C}$ -based mixing model was applied to samples with high TSM. Based on their spatial distribution, these samples can be reasonably attributed to resuspended seabed sediment (cf. Sect. 4.3), allowing more reliable assignment of endmember $\delta^{13}\text{C}$ values. To define high-TSM samples operationally, we ranked the dataset by TSM and evaluated the correlation between $\delta^{13}\text{C}_{\text{OC,corr}}$ (from Step 1) and temperature (cf. Sect. 5.2.3) for subsets with progressively increasing upper TSM limits. The correlation deteriorated when samples with samples with $\text{TSM} \geq 4 \text{ mg L}^{-1}$ were included (Fig. C1 in Appendix C). Therefore, this value was adopted as the threshold separating low- and high-TSM samples.

The $\delta^{13}\text{C}$ -based binary mixing model treats POC as a mixture of sedimentary and marine OC. Sedimentary endmembers were site-specific, whereas marine endmembers were derived from the $\delta^{13}\text{C}_{\text{OC}}$ -temperature relationship of low-TSM samples (cf. Sect. 5.2.3). POC_{terr} was calculated as:

$$\text{POC}_{\text{terr}} = \text{POC}_{\text{m}} \times f_{\text{sed}} \times f_{\text{terr,sed}} \quad (5)$$

where POC_{m} is measured POC, f_{sed} the sedimentary fraction of POC derived from the binary mixing model, and $f_{\text{terr,sed}}$ the terrigenous fraction of sedimentary OC (Table A4). In this step, $\delta^{13}\text{C}_{\text{OC,corr}}$ could not be obtained because $\delta^{13}\text{C}_{\text{OC}}$ was prescribed from the $\delta^{13}\text{C}_{\text{OC}}$ -temperature relationship. Therefore, only POC_{corr} and N/C_{corr} were computed for high-TSM samples. The estimated f_{sed} values were further used to calculate the source signature of $\Delta 8$ ($\Delta 8_{\text{cal}}$) required to account for the measured $\Delta 8$ ($\Delta 8_{\text{m}}$) in high-TSM samples:

Comparison of $\Lambda\delta_{\text{m}}$ and the corresponding $\Lambda\delta_{\text{terr, s}}$ used in Step 1 helps evaluate the limitations and uncertainties of lignin-based POC_{terr} quantification.



215 **Figure 2:** Schematic diagram summarizing the data processing procedure for estimating POC_{terr} . EM, endmember; T, temperature.

3.3.2 Estimation of POC_{terr} export flux

To compare with the riverine flux from Taiwan, the advective export flux of POC_{terr} was estimated by combining POC_{terr} concentrations with flow data from the Hybrid Coordinate Ocean Model (HYCOM). The applicability of HYCOM for reconstructing water flux through the strait has been validated by Huang et al. (2019). A latitudinal transect between 120.40° and 121.04° E at 25.04° N was defined as the northern boundary of the study area (Fig. 1b), and export flux was calculated as the rate of material crossing this transect.

POC_{terr} estimates from Step 1 were used for low-TSM samples and those from Step 2 for high-TSM samples. POC_{terr} in offshore surface waters, attributable to remote Pearl River inputs (see Sect. 5.1.1), was excluded. The bottommost 5 m of the water column was also omitted due to *i*) the lack of CTD data, and *ii*) the higher likelihood of particle re-deposition near the seafloor. Calculation details are provided in Text B3.

3.3.3 Calculation of dissolved CO₂ concentration

For samples with carbonate chemistry measurements, we used CO2SYS Version 3 (Pierrot et al., 2021) to compute dissolved CO₂ (CO₂(aq)) concentrations from measured DIC and pH₂₅ (Table A5). Due to the lack of concurrent nutrient data, we incorporated historical records from Sites A to F, which were visited during the cruise OR3-632 of RV *Ocean Researcher III* in June 2000 (Fig. 1c). These sites exhibit similar water mass characteristics to those in our study area (Fig. C2). These data were accessed from the Ocean Data Bank (2025). Salinity-based empirical relationships for silica and phosphate were derived from these records (Fig. C3) and then applied to provide the nutrient inputs required for CO2SYS calculations for NOR3-0104 samples.

3.4 Statistical analysis and visualization

Statistical analyses were conducted using Excel® add-ons XLSTAT (Lumivero). The non-parametric Mann-Whitney test was used for comparing two datasets, and the Kruskal-Wallis test followed by Dunn's test was used for comparisons among three datasets. Differences between slopes of two linear regressions were tested using Student's *t*-test following Andrade and Estévez-Pérez (2014). Statistical significance was set at $\alpha = 0.05$. Bathymetric maps and topographic profiles were generated in QGIS (version 3.34.11) using digital elevation models from the Ocean Data Bank (2025) referenced to WGS 84. Transects with isopleths were created using Ocean Data View (version 5.6.3; Schlitzer, 2025).

4 Results

4.1 Hydrographic data

Hydrographic data are available in Lin et al. (2025b), and their vertical distributions are shown in Fig. 3. Temperature ranged from 23.4 to 30.0 °C (Fig. 3a), highest along the Taiwanese coast and lowest in the bottom water at Site X5. Salinity showed the opposite pattern, increasing from 32.8 nearshore to 34.3 offshore (Fig. 3b). In the T-S diagram (Fig. C2), most samples clustered near the typical SCSW curve, while two subsets exhibited lower salinity. One subset, with higher potential density (σ_θ) values (21.7–22.5 kg m⁻³), comprised upper-water-column samples from Transects X and Y and overlapped with middle-shelf waters in Transect T1. This subset is therefore attributed to the SCSSW. The other subset, having lower σ_θ values (< 21.7 kg m⁻³), occupied nearly the entire water column in Transect Z. Given its proximity to the Taiwanese coast, this subset is hereafter referred to as Taiwan Coastal Water (TCW).

After calibration (Eq. (1)), sensor-derived TSM ranged from 0.7 to 36.4 mg L⁻¹ (Fig. 3c). High TSM occurred in bottom waters (> 40 m) of Sites X1–X5 and Y1–Y4, and throughout Transect Z. At weakly stratified sites (X1, Y1, and Z10), bottom-enriched TSM reached the surface.

255 Calibrated Chl (Eq. (2)) ranged from 0.2 to 2.5 µg L⁻¹ (Fig. 3d). Chl-rich waters (exceeding ~1 µg L⁻¹) occurred at 10–40 m depths in Transects X and Y and throughout Transect Z, particularly near river mouths (Z1, Z2, Z5, Z8, and Z9).





Figure 3: Hydrographic conditions along three transects in the northeastern Taiwan Strait during summer 2022. (a) Temperature, with black dots marking CTD measurement points. (b) Salinity and distribution of South China Sea Surface Water (SCSSW) and Taiwan Coastal Water (TCW). (c) TSM and σ_{θ} . (d) Chl concentrations and Z_e . Grey dots indicate depths for discrete water sampling.

4.2 POM data

POM data are available in Lin et al. (2025b) and displayed in Fig. 4. POC ranged from 63 to 578 $\mu\text{g L}^{-1}$ (Fig. 4a), higher in nearshore than offshore waters. Surface POC (0–5 m) exceeded subsurface values ($p = 0.006$). Elevated bottom-water POC at Sites X1–X3, Z9, and Z10 coincided with high TSM concentrations (Fig. 3c). $\delta^{13}\text{C}_{\text{OC}}$ values ranged from -25.1 to -21.0 ‰, with heavier values nearshore than offshore (Fig. 4b). Most high-POC samples, except turbid samples at Sites Z9 and Z10, had $\delta^{13}\text{C}_{\text{OC}}$ values above -23 ‰. Low- $\delta^{13}\text{C}$ (< -24 ‰) POM occurred in subsurface and bottom waters of Transects X and Y. C/N ratios (6.4–10.0) lacked clear spatial or vertical patterns (Fig. 4c). C/N ratios > 8 occurred in surface waters of Y1–Y3 and Z2–Z4, in bottom waters of Y6, and throughout Z10. $\Sigma 8$ ranged from below detection to 2,753 ng L^{-1} (Fig. 4d). Where measurable, $\Lambda 8$ ranged from 0.12 to 8.6 $\text{mg lignin g}^{-1} \text{OC}$. In offshore Transects X and Y, $\Sigma 8$ were elevated in surface and bottom waters but remained below detection in subsurface waters. $\Lambda 8$ showed no significant difference between samples with $\delta^{13}\text{C}_{\text{OC}}$ values below and above -24 ‰ ($p = 0.489$). Transect Z exhibited higher $\Sigma 8$ than the offshore transects ($p \leq 0.001$), particularly at high-TSM sites (Z1–Z2, Z5, Z8–Z10). Most $(\text{Ad/Al})_{\text{V}}$ values ranged from 0.1 to 2; four anomalously high values (2.5–5) likely resulted from incomplete oxygen sparging during sample pretreatment. After excluding these samples, Transect Z ($(\text{Ad/Al})_{\text{V}} = 0.54 \pm 0.28$) had lower degradation state than the other transects (mean = 1.10; $p \leq 0.004$).



280 **Figure 4: POM characteristics along the three transects in the northeastern Taiwan Strait during summer 2022. (a) POC concentrations. (b) $\delta^{13}\text{C}_{\text{OC}}$ values. (c) Atomic C/N ratios. (d) $\Sigma 8$ concentrations. Grey dots indicate depths for discrete water sampling. MDL, method detection limit.**


4.3 Estimated POC_{terr} concentrations

Applying Eq. (4) requires defining terrigenous POM sources (Tables A2 and A3). $\Sigma 8$ correlated strongly with TSM ($r = 0.79$, $p < 0.0001$) and moderately with salinity ($r = -0.42$, $p < 0.0001$) (Figs. 5a–b), indicating sediment resuspension as the dominant lignin source. Muddy sediments, richer in lignin than sand (Lin et al., 2025a; Fig. C4), likely supplied most lignin.

285 Enrichment of lignin also occurred in surface waters. Surface lignin in Transect Z likely originated from Taiwanese rivers, whereas that in Transects X and Y is attributed to long-distance transport from the Pearl River, consistent with summer circulation (Fig. 1c).

Lignin-based POC_{terr} values and corrected POM parameters (POC_{corr}, N/C_{corr}, and $\delta^{13}\text{C}_{\text{OC,corr}}$) after Step 1 are listed in Table A2. POC_{terr} concentrations did not exceed $63 \mu\text{g L}^{-1}$ and were below $10 \mu\text{g L}^{-1}$ for most samples. The terrigenous fraction of POC (f_{terr}), calculated as POC_{terr}/POC_m, averaged 0.06 ± 0.07 for all samples. Relative differences between measured and corrected values averaged 6.4 % for POC and $< 1 \%$ for C/N and $\delta^{13}\text{C}_{\text{OC}}$.

The $\delta^{13}\text{C}_{\text{OC}}$ -based mixing model of the second step yielded POC_{terr} = 47–332 $\mu\text{g L}^{-1}$ and $f_{\text{terr}} = 0.47$ –0.92 for high-TSM samples (Table A4). For comparison, lignin-based f_{terr} values were 0.03–0.69 for the same set of samples. $\Delta 8_{\text{cal}}$ averaged only $17 \pm 10 \%$ of $\Delta 8_{\text{terr,s}}$ (Table A4). Combined results (Steps 1 and 2) show $f_{\text{terr}} > 0.5$ in offshore bottom waters along 295 Transect X, at Site Y5, and in nearshore waters at Z9–Z10 (Fig. 5c).



300 **Figure 5: Characteristics of terrigenous POM. (a) Relationship between $\Sigma 8$ and TSM. (b) Relationship between $\Sigma 8$ and salinity. (c) Spatial distribution of f_{terr} along the three transects in the northeastern Taiwan Strait during summer 2022. Grey dots indicate depths for discrete water sampling.**

4.4 Estimated POC_{terr} export flux

POC_{terr} concentrations derived from the combined approach, i.e. lignin-based estimates for low-TSM samples and $\delta^{13}\text{C}$ -based estimates for high-TSM samples, were used to calculate the POC_{terr} export flux. Along 25.04° N, POC_{terr} concentrations were elevated in nearshore and offshore bottom waters (Fig. C5a). The annually averaged flow was
305 northeastward, but highest velocities occurred where POC_{terr} was low (Fig. C5b). Thus, POC_{terr} export was controlled mainly by concentration rather than current speed (Fig. C5c). The total export flux was $243 \pm 56 \text{ kt C y}^{-1}$.

4.5 Carbonate chemistry

Carbonate chemistry data are available in Lin et al. (2025b). DIC ranged from 1,807 to 2,007 $\mu\text{mol kg}^{-1}$, lowest at 0 m of Site X3 and highest at 72 m of Site X5. pH_{25} ranged from 7.96 (72 m of Site X5) to 8.13 (0 m of Site Z5). Computed $\text{CO}_2(\text{aq})$
310 concentrations ranged from 8.7 to 13.9 $\mu\text{mol kg}^{-1}$ (equivalent to 8.9 to 14.2 $\mu\text{mol L}^{-1}$), with minimum and maximum values occurring at Sites Z5 and X5, respectively (Table A6).

5 Discussion

5.1 Distribution and budget of terrigenous POM

5.1.1 Persistence of lignin and POC_{terr} at low abundance during advective transport

315 The fate of terrigenous OM in the ocean has long intrigued geochemists (Bianchi, 2011; Hedges et al., 1997; Talling et al., 2024). Globally, rivers deliver $\sim 200 \text{ Mt yr}^{-1}$ of POC to the ocean, $\sim 80 \%$ biospheric and $\sim 20 \%$ petrogenic (Galy et al., 2015), yet only 62–90 Mt yr^{-1} is ultimately buried (Talling et al., 2024). One explanation for this imbalance is repeated settling-resuspension cycles in estuarine and coastal waters, which weaken organo-mineral associations and enhance OM bioavailability (Bao et al., 2018; Burdige, 2007; Sun et al., 2024). What remains uncertain is whether the terrigenous OM
320 released from disrupted aggregates is quickly degraded or can persist long enough to disperse across continental shelves.

Because of the overlapping bulk signatures of terrigenous and marine POM (Geider, 1987; Hedges et al., 1997; Laws et al., 1995; Martiny et al., 2013), we used lignin as a tracer of POC_{terr}. Our $\Sigma 8$ concentrations ($\leq 1,789 \text{ ng L}^{-1}$; Fig. 4d) agree well with literature values: lower than estuaries (1–40 $\mu\text{g L}^{-1}$; Reeves and Preston, 1989), comparable to shelf waters (10–1000 ng L^{-1} ; calculated from Bianchi et al., 1997), and higher the Pacific Ocean ($< 10 \text{ ng L}^{-1}$; Hernes and Benner, 2006).

325 $\Sigma 8$ correlated more strongly with TSM than salinity (Fig. 5), indicating resuspended sediment as the primary source, consistent with estuarine patterns (Reeves and Preston, 1989). Lignin was also detected in low-TSM waters, occurring in nearshore TCW as a result of Taiwanese river discharge and along-shore hypopycnal transport (Lin et al., 2025a), and in offshore SCSSW, in line with the inferred cross-shelf contribution from the Pearl River plume (Bai et al., 2015; Jan et al., 2006). This interpretation is supported by higher $(\text{Ad/Al})_V$ ratios offshore, implying longer exposure of lignin to
330 photochemical or biological degradation in SCSSW than in TCW. The offshore-nearshore difference in $(\text{Ad/Al})_V$ ratios is

consistent with longer water transit times from the Pearl River mouth to the northeast Taiwan Strait (~15 days; Bai et al., 2015) than from the outlets of Taiwanese rivers to the northern end of our transects (~5 days; estimated from Wang et al., 2003). In both cases, however, the transport times remain short relative to the slow degradation kinetics of lignin (Benner et al., 1987). Together, these observations indicate that lignin, and thus a fraction of terrigenous POM, can persist during both
335 along-shore and cross-shelf transport.

By contrast, lignin concentrations were minimal to negligible in most offshore subsurface waters. Although seemingly counterintuitive given their proximity to land, this pattern can be explained by water-mass origin, limited lignin supply, and weak hydrodynamic forcing during sampling. Offshore subsurface and bottom waters are derived from SCSW (Fig. C2), which is likely lignin-poor due to a greater contribution from Pacific-origin waters (Nan et al., 2015; You et al., 2005). The
340 more frequent detection of lignin in bottom than subsurface waters suggests that resuspension was insufficient to transport sediment higher into the water column, consistent with the calm sea state during the cruise. Offshore subsurface waters may also receive lignin from overlying river plumes, but low river discharge prior to and during sampling confined plume influence to a narrow coastal band (Lin et al., 2025a), limiting lignin supply offshore. This lignin-poor subsurface layer would likely contract under rougher conditions, which are common in summer. Notably, these waters also correspond to
345 where low- $\delta^{13}\text{C}$ POM is most frequently observed (Fig. 4). Their low lignin concentrations indicate that this isotopic signature is unlikely to reflect terrigenous inputs.

5.1.2 Uncertainty in POC_{terr} estimation

Lignin-based estimates yielded low POC_{terr} concentrations, with f_{terr} averaging 0.14 ± 0.08 and 0.06 ± 0.09 for high- and low-TSM samples, respectively (Table A2). In contrast, f_{terr} values of high-TSM samples were revised to 0.68 ± 0.14 using the
350 $\delta^{13}\text{C}_{\text{OC}}$ -based mixing model (Table A4). The higher f_{terr} values from the $\delta^{13}\text{C}$ -based approach are considered more reliable for high-TSM samples, as these are primarily sourced from seabed sediments known to contain a high proportion of terrigenous OM (Lin et al., 2025a). To investigate the discrepancy between the two approaches, we back-calculated $\Lambda_{8\text{cal}}$ that would reconcile the measured POM Λ_8 with the mixing-model results (Eq. (6)). $\Lambda_{8\text{cal}}$ averaged only $17 \pm 10\%$ of $\Lambda_{8\text{terr,s}}$ (Table A4), implying either lignin degradation in the water column and/or a greater contribution from lignin-depleted source
355 material. Elevated $(\text{Ad}/\text{Al})_{\text{V}}$ ratios in offshore surface waters, relative to those of Pearl River POM ($(\text{Ad}/\text{Al})_{\text{V}} < 0.75$; Zhang et al., 2014), support lignin degradation. However, samples that do not exhibit marked changes in $(\text{Ad}/\text{Al})_{\text{V}}$ ratios, such as Transect Z POM compared with underlying sediments or riverine TSM ($p = 0.80$ to 0.81), indicate that additional processes are also involved.

One possibility is the mixed contribution of lignin-rich and lignin-poor source materials. In estimating POC_{terr} (Table A2),
360 we adopted $\Lambda_{8\text{terr,s}}$ values from either riverine TSM ($4.6\text{--}11.2$ mg lignin g^{-1} OC) or seabed sediments ($5.3\text{--}72.4$ mg lignin g^{-1} OC). If the actual source is a mixture of both, POC_{terr} would be overestimated in surface waters but underestimated in subsurface and bottom waters. Given the larger volume of the latter, our estimates are likely biased low overall. Another possibility is preferential resuspension of lignin-poor OM. This is supported by density-fractionation experiments (Wakeham

et al., 2009), which showed that although lignin concentrations (normalized to sediment dry weight) increased in low-density
 365 fractions, Λ_8 decreased because OC was enriched more strongly than lignin. The magnitude of Λ_8 reduction relative to bulk
 sediments was highly variable, precluding a robust correction.

Given these uncertainties, we did not attempt a detailed correction of the POC_{terr} estimates for low-TSM samples, but instead
 provide a first-order assessment of the potential bias. If the average $\Lambda_{8\text{cal}}/\Lambda_{8\text{terr,s}}$ ratio were applied to low-TSM samples, f_{terr}
 would increase by approximately sixfold (1/0.17). Even under this scenario, terrigenous OM would remain a secondary
 370 component for low-TSM samples. Therefore, our results support previous findings that terrigenous POM makes only a
 limited contribution to shelf waters above the benthic nepheloid layer (e.g., Ho et al., 2021; Liu et al., 2018a, 2022).

5.1.3 Closing the regional POC_{terr} budget

Combining POC_{terr} concentrations with simulated flow data (Fig. C5), we estimated an advective flux of $243 \pm 56 \text{ kt y}^{-1}$ from
 the northeastern Taiwan Strait, providing a first-order approximation for the magnitude of POC_{terr} export. This value is
 375 broadly consistent with the regional imbalance between riverine input and burial (Table 1) and may help close the POC_{terr}
 budget. It is higher than our previous estimate that accounted for benthic oxidation (Lin et al., 2025a), although the two
 values likely overlap within uncertainties.

Notably, 48 % of exported POC_{terr} exited via waters above the narrow ($< 10 \text{ km}$) nearshore mud belt, underscoring its dual
 role as both a temporary sink of fluvial inputs and a source fuelling long-distance transport. An additional 36 % was
 380 exported via the nepheloid layer above the offshore mud belt, whereas only 17 % was carried by clear shelf waters. However,
 as discussed above, POC_{terr} concentrations in low-TSM samples are likely underestimated by the lignin-based approach. If
 the sixfold correction derived from the average $\Lambda_{8\text{cal}}/\Lambda_{8\text{terr,s}}$ ratio is applied, the contribution of clear shelf waters would
 increase to 43 %. This suggests that low-TSM shelf waters, though having a minor fraction of terrigenous OM, may play a
 non-negligible role in its advective transport.

385


Table 1: POC_{terr} budget of the northeastern Taiwan Strait

Item	POC_{terr} (kt y^{-1})	% riverine input	Source
Measured flux			
Riverine input	362 ± 75	100	Lin et al. (2020b)
Burial	75 ± 4.6	21 ± 4	Lin et al. (2025a)
Calculated flux			
Riverine input – burial	287 ± 75	79 ± 21	Based on Lin et al. (2025a)
Advective export	189 ± 83	52 ± 25	Based on Lin et al. (2025a)
Advective export	243 ± 56	67 ± 21	This study

5.2 Controlling factors of marine POM properties

We first compared POM characteristics before and after correction for terrigenous input. Here, "overall characteristics" refer to slopes derived from linear regressions of variables. The measured and corrected overall atomic C/N ratios were 7.67 ± 0.16 and 6.56 ± 0.15 , respectively, and differed significantly ($p < 0.0001$). Excluding high-TSM samples yielded comparable values (6.87 ± 0.15 and 6.61 ± 0.16 ; $p = 0.08$; Fig. 6a). After correction, overall POC/Chl ratios decreased from 92.7 ± 18.4 to 79.6 ± 10.9 g C g⁻¹ Chl, and for low-TSM samples from 91.8 ± 9.8 to 83.0 ± 9.1 g C g⁻¹ Chl (Fig. 6b). These differences were not significant ($p = 0.15$ to 0.16). Neither did the correction alter the $\delta^{13}\text{C}_{\text{OC}}$ -temperature relationship for low-TSM samples.

As discussed in Sect. 5.1.2, the lignin-based approach likely underestimates POC_{terr} , which may partly contribute to the limited differences observed after correction. However, even when accounting for the potential magnitude of this underestimation, terrigenous POM remains a secondary component in low-TSM samples. The weak correlations between POC and $\Sigma 8$ ($r = 0.14$, $p = 0.27$) and between $\delta^{13}\text{C}_{\text{OC}}$ and $\Lambda 8$ ($r = -0.04$, $p = 0.76$) in these samples further indicates that additional upward revision of lignin-based POC_{terr} estimates is unlikely to substantially alter the slopes of linear regressions used to assess overall characteristics. In the following discussion, we therefore focus on the corrected low-TSM dataset, while noting that the conclusions are insensitive to whether the correction is applied.



405 **Figure 6: Characteristics of marine POM. (a) Relationship between POC and PON. (b) Relationship between POC and Chl. (c) Relationship between $\delta^{13}C_{oc}$ and temperature. (d) Relationship between $\delta^{13}C_{oc}$ and $CO_2(aq)$. (e) Relationship between $\delta^{13}C_{oc}$ and $POC/CO_2(aq)$ (both in $\mu mol L^{-1}$). Linear regressions are based on low-TSM samples only. * denotes data excluded from the regression. References: 1 = Liu et al., 2018a; 2 = Liu et al., 2022.**

5.2.1 C/N ratios reflect nutrient supply ratios

The overall C/N ratio of 6.61 ± 0.16 (Fig. 6a) agrees with the global median of 6.5 (Martiny et al., 2013) and canonical
 410 Redfield ratio of 6.63, with no offshore-nearshore difference ($p = 0.09$). Regionally, our values resemble those from summer POM in the upper 200 m at SEATS in the South China Sea (6.44 ± 0.28 ; $p = 0.48$; Liu et al., 2007; Fig. 1a), but exceed those of the DCM in the East China Sea (5.76 ± 0.14 ; $p < 0.0001$; Liu et al., 2018a). Further north, the Yellow Sea DCM values (6.31 ± 0.53 ; Liu et al., 2022) are indistinguishable from our results ($p = 0.12$). Although phytoplankton C/N ratios are influenced by nutrients, irradiance, temperature, growth rate, and community structure, nutrient supply remains the dominant
 415 control in the global surface ocean (Moreno and Martiny, 2018; Tanioka and Matsumoto, 2020). Marine primary production is commonly N-limited in the world ocean, but N stress is alleviated in large plume-affected regions (Huang et al., 2019). This mechanism was invoked to explain lower POM C/N ratios in the western North Atlantic (Martiny et al., 2013). Across

the Pan-China Sea (cf. Bianchi et al., 2018), C/N variation is consistent with this pattern. Near Redfield ratios prevail in the South China Sea, Taiwan Strait, and Yellow Sea, where nutrient supply is relatively balanced (Huang et al., 2019; Liu et al., 2022; Wong et al., 2015), whereas lower ratios occur in the East China Sea owing to nitrogen enrichment from the Yangtze River (Zhong et al., 2025).

5.2.2 POC/Chl ratios track photoacclimation

Cellular OC/Chl ratio is a physiological indicator of phytoplankton growth constraints (Arteaga et al., 2014; Gui and Sun, 2024). It responds to light, nutrients, and temperature, although temperature effects are minor at regional scales (Wang et al., 2009). When light intensity is strong, phytoplankton down-regulates Chl synthesis to prevent damage from excess light and reallocates resources towards carbon fixation, elevating the ratio. With increasing water depth, the ratio decreases as cells invest nitrogen in light harvesting (Brown et al., 2003; Li et al., 2010). Under high nitrogen availability, more nitrogen is allocated to light harvesting, lowering the ratio; under nitrogen limitation, nitrogen is diverted to enzymes needed for nutrient acquisition, increasing the ratio (Arteaga et al., 2014). The ratio also underpins conversions from satellite Chl to phytoplankton carbon biomass (Carr et al., 2006). Microscopy and flow cytometry, along with assumptions of cell geometry and empirical equations for cellular carbon content, are the standard approaches to estimate phytoplankton carbon biomass required to compute OC/Chl ratio (e.g., Li et al., 2010). Here, we explore whether overall POC/Chl ratios can approximate phytoplankton OC/Chl ratios in shelf waters by comparing our values with cell-based estimates and model predictions.

The regression between POC and Chl shows a non-zero intercept (Fig. 6b), indicating contributions from detritus or heterotrophs. The overall POC/Chl ratio of $83.0 \pm 9.1 \text{ g C g}^{-1} \text{ Chl}$ is close to the OC/Chl value of nutrient-depleted surface waters ($\sim 90 \text{ g C g}^{-1} \text{ Chl}$; Eppley, 1968) and lies with the modeled range of $50\text{--}100 \text{ g C g}^{-1} \text{ Chl}$ for the nitrogen-limited subtropical western North Pacific (Arteaga et al., 2014).

Two sample subsets showed elevated Chl concentrations and merit further examination: nearshore waters and offshore DCM layers (Fig. 3d). Nearshore waters, enriched by riverine nitrogen input in summer ($1\text{--}3 \mu\text{mol L}^{-1}$ nitrate plus nitrite; Huang, 2022), had a ratio of $75.0 \pm 12.5 \text{ g C g}^{-1} \text{ Chl}$ (Fig. C6a). Offshore DCM layers, which are nitrogen-depleted ($< 1 \mu\text{mol L}^{-1}$ nitrate; Tseng et al., 2020), showed a lower ratio of $43.0 \pm 29.1 \text{ g C g}^{-1} \text{ Chl}$, although the POC-Chl relationship was not significant (Fig. C6b). This contrast is further supported by comparisons of measured POC/Chl ratios, which are higher in nearshore waters than in offshore DCM layers ($p < 0.001$). Overall POC/Chl ratios as low as $28\text{--}38 \text{ g C g}^{-1} \text{ Chl}$ have been reported for DCM layers in the East China Sea and Yellow Sea (Liu et al., 2018a, 2022). The pattern observed in our samples indicates that photoacclimation exerts a stronger control on the POC/Chl ratio than nutrient stress, based on the physiological considerations described above. Alternative explanations include shifts in phytoplankton community composition, as suggested by Li et al. (2010), or covariation between bacterial biomass and Chl, as demonstrated by Brown et al. (2003). Given the highly dynamic nature of planktonic communities in shelf waters, we refrain from relate our POC/Chl data to reported variability in community composition in the Taiwan Strait (e.g., Tong et al., 2024; Zhong et al.,

450 2020). Concurrent measurements of POM properties and planktonic community structure are therefore required to robustly resolve the mechanisms underlying POC/Chl ratio.

5.2.3 Temperature control on $\delta^{13}\text{C}_{\text{OC}}$

$\delta^{13}\text{C}$ of marine phytoplankton reflects the balance between growth demand and DIC supply in seawater (Fry, 1996). Under diffusive CO_2 transport, isotopic fractionation between CO_2 and cellular carbon increases when growth slows or $\text{CO}_2(\text{aq})$ concentration rises (i.e., a lower $\mu/\text{CO}_2(\text{aq})$ ratio, where μ is growth rate), producing more negative $\delta^{13}\text{C}$ (Laws et al., 1995). Biological parameters influencing fractionation include growth rate (Laws et al., 1995), cell geometry (Popp et al., 1998) and taxonomic physiology (Burkhardt et al., 1999), whereas relevant environmental factors include $\text{CO}_2(\text{aq})$, nutrients (Riebesell et al., 2000), and temperature (Rau et al., 1989). When $\text{CO}_2(\text{aq})$ is low, many phytoplankton activate carbon concentrating mechanisms (CCMs; Giordano et al., 2005), which reduce fractionation (Fielding et al., 1998; Sharkey and Berry, 1985) and weaken the predictability of $\delta^{13}\text{C}$ based on $\mu/\text{CO}_2(\text{aq})$ ratios (Burkhardt et al., 1999; Rau et al., 2001). A critical $\text{CO}_2(\text{aq})$ concentration of $\sim 10 \mu\text{mol L}^{-1}$ to induce CCMs has been suggested as the threshold for CCM induction, although CCMs may operate at concentrations above this level (Burkhardt et al., 1999).

Because $\delta^{13}\text{C}$ of DIC varies little in the global euphotic zone (Kroopnick, 1985) and the temperature effect on the variation of bicarbonate- $\text{CO}_2(\text{aq})$ isotopic equilibrium is small (0.12‰ K^{-1} ; Mook et al., 1974), $\delta^{13}\text{C}_{\text{OC}}$ of POM largely reflects biological fractionation. Globally, $\delta^{13}\text{C}_{\text{OC}}$ decreases toward higher latitudes, a trend attributed to increased CO_2 solubility in colder waters (Rau et al., 1989). Recent work by Liu et al. (2022) showed that temperature is a better predictor of POM $\delta^{13}\text{C}_{\text{OC}}$ than either $\text{CO}_2(\text{aq})$ or $\text{POC}/\text{CO}_2(\text{aq})$ ratio, a proxy for the $\mu/\text{CO}_2(\text{aq})$ ratio, in both shelf and open ocean environments. This is possibly because temperature integrates multiple biological and environmental controls on isotopic fractionation, irrespective of carbon acquisition modes (Liu et al., 2022).

470 Our dataset corroborates this view. Across all samples, $\delta^{13}\text{C}_{\text{OC}}$ correlates strongly with temperature (Fig. 6c). For samples with carbonate chemistry data, the r^2 values follow the order temperature (0.72), $\text{POC}/\text{CO}_2(\text{aq})$ ratio (0.65), and $\text{CO}_2(\text{aq})$ (0.35) (Figs. 6c to 6e), confirming temperature as the strongest predictor. The weak $\text{CO}_2(\text{aq})$ relationship likely reflects widespread CCM activity in the warm, high-light conditions of our study area. Comparisons with the East China and Yellow Seas show no clear regional trend in slope (Fig. 6c), limiting broader extrapolation. Even so, for regional POM studies, this relationship provides a practical constraint on the isotopic composition of marine POM (Fig. 2) and improves estimates of terrigenous POC in the water column (Sect. 3.3.2).

5.3 Comparison of POM and sedimentary OM

Differences between water-column POM and underlying sedimentary OM have long interested researchers studying particle transport, carbon preservation, and paleoenvironmental reconstruction (Hayes et al., 1999; Wakeham and McNichol, 2014; Wang et al., 2017). Concomitant measurements of POM and surface sediment OM from the Taiwan Strait (Lin et al., 2025a) allow us to assess how short-term hydrogeochemical signals in POM compare with the sedimentary archive of this shelf

environment. We examined three categories of parameter pairs: *i*) water-column integrated $\Sigma 8$ versus sedimentary lignin, *ii*) water-column integrated Chl versus sedimentary photosynthetic pigments, and *iii*) POC-weighted water-column $\delta^{13}\text{C}_{\text{OC}}$ versus sedimentary $\delta^{13}\text{C}_{\text{OC}}$. Sediment data represent the upper 1 cm, reflecting recent deposition. For categories *i* and *ii*, all relevant variants of sedimentary parameters reported in Lin et al. (2025a), including dry-weight concentration, OC-normalized concentration, and burial flux, were included. Sedimentary pigments comprise Chl and pheopigments a, derivatives of Chl formed during biological degradation (Stephens et al., 1997; Sun et al., 1993).

Correlation coefficients for all parameter pairs are listed in Table A7, and the best-correlated pair in each category is shown in Fig. 7. Water-column integrated stocks showed the strongest correlations with OC-normalized sedimentary biomarker concentrations. For lignin (Fig. 7a), the correlation reflects the role of seabed sediment as a major lignin source to the water column (Sect. 5.1.1) and indicates that lignin-rich sedimentary OM likely contains easily resuspended wood microdetritus.

For photosynthetic pigments (Fig. 7b), Lin et al. (2025a) attributed the relationship to the biological pump. Notably, nearshore "hot spots" of photosynthetic activity, characterized by high POC, heavy $\delta^{13}\text{C}_{\text{OC}}$ and low $\text{CO}_2(\text{aq})$ (Fig. 4), did not dominate the correlation; instead, sedimentary patterns were primarily controlled by euphotic-zone thickness, which also governs depth-integrated primary production in the region (Tseng et al., 2020).




Figure 7: Comparison of POM and sedimentary OM. (a) Water-column integrated $\Sigma 8$ ($\Sigma 8_{\text{int}}$) versus OC-normalized lignin concentration (A8) of the sediments (Sed). (b) Water-column integrated Chl (Chl_{int}) versus sedimentary OC-normalized pheopigments a (Pheo) concentration. (c) POC-weighted water-column $\delta^{13}\text{C}_{\text{OC}}$ versus sedimentary $\delta^{13}\text{C}_{\text{OC}}$.

By contrast, $\delta^{13}\text{C}_{\text{OC}}$ in POM and sedimentary OM shows a puzzling negative correlation (Fig. 7c). Because $\delta^{13}\text{C}_{\text{OC}}$ integrates OM from sources with distinct isotopic signatures, the result points to the competing influences of lateral and vertical transport in shaping sedimentary $\delta^{13}\text{C}_{\text{OC}}$ on shallow shelves. The absence of a positive relationship between sedimentary and water-column $\delta^{13}\text{C}_{\text{OC}}$, together with the temperature dependence of marine POM $\delta^{13}\text{C}_{\text{OC}}$, raises a fundamental question: Is it appropriate to assign a single marine $\delta^{13}\text{C}_{\text{OC}}$ endmember in isotope mixing models? Previous work suggests that incorporation of marine POM into sediments largely occurs during episodic, high-POC pulses (Berger and Wefer, 1990) with minimal isotopic fractionation (Kukert and Riebesell, 1998). Yet in environments where blooms are rare, as is likely the

case in parts of the northeastern Taiwan Strait, the isotopic signatures of marine OM ultimately buried on the seafloor remain poorly constrained. Revisiting published datasets containing $\delta^{13}\text{C}_{\text{OC}}$ of co-sampled suspended, sinking and deposited particles, along with hydrographic context, may help address this issue.

6 Conclusions

This study provides an integrated, full-water-column assessment of terrigenous and marine POM in the northeastern Taiwan Strait. The results demonstrate how hydrodynamics, water-mass structure, resuspension, and primary production jointly regulate POM sources, distribution, and export on this energetic shelf. The main findings are:

1. Lignin tracers reveal contributions from both Taiwanese rivers and the Pearl River plume. POC_{terr} was estimated using a two-step approach that incorporated both lignin and $\delta^{13}\text{C}_{\text{OC}}$ data. Terrigenous fractions are low in low-TSM waters but rise sharply in high-TSM layers due to sediment resuspension.
2. Combining POC_{terr} with HYCOM velocities yields an export flux of $\sim 243 \pm 56 \text{ kt C yr}^{-1}$, comparable to the mismatch between riverine supply and sedimentary burial. This flux helps close the regional POC_{terr} budget.
3. After correction for terrigenous contributions, C/N ratios reflect balanced nutrient supply, POC/Chl ratios respond to photoacclimation, and $\delta^{13}\text{C}_{\text{OC}}$ is better explained by temperature than $\text{CO}_2(\text{aq})$ -related variables, implying widespread CCM activity.
4. Sedimentary lignin and pigment concentrations correspond to their water-column stocks, but the negative POM-sediment $\delta^{13}\text{C}_{\text{OC}}$ correlation shows that lateral transport overprints vertical delivery in determining sedimentary isotopic composition.

Overall, this work refines the quantification of terrigenous and marine POM, clarifies source-to-sink connectivity, and strengthens the modern framework needed to interpret sedimentary OM proxies in shelf environments.

Appendix A

530 **Table A1: List of sampling sites from the cruise NOR3-0104**

Site	Date of cast	Longitude (°E)	Latitude (°N)	Water depth (m)	Ze ^a (m)	Depth of sampling ^b (m)
X1	16 Jun 2022	120.173467	24.342333	51	31*	<u>0</u> , <u>5</u> , 25, <u>48</u>
X2	15 Jun 2022	120.133383	24.502500	63	37*	0, 5, 35, 58
X3	15 Jun 2022	120.170100	24.668667	69	43	<u>0</u> , <u>5</u> , 30, 50, <u>65</u>
X4	15 Jun 2022	120.334433	24.837383	79	48	0, 5, 40, 73
X5	15 Jun 2022	120.433533	25.005617	79	46*	<u>0</u> , <u>5</u> , 10, <u>40</u> , <u>72</u>
Y1	16 Jun 2022	120.279550	24.210783	49	22	<u>0</u> , 5, 25, <u>45</u>
Y2	16 Jun 2022	120.337267	24.391367	52	27	0, 5, 20, 45
Y3	16 Jun 2022	120.407750	24.501300	53	34	<u>0</u> , 5, 25, <u>48</u>
Y4	16 Jun 2022	120.456050	24.618267	58	34*	0, 5, 25, 53
Y5	17 Jun 2022	120.613133	24.826467	72	52	0, 5, 40, 67
Y6	17 Jun 2022	120.736800	25.031917	96	51	<u>0</u> , 5, 35, <u>90</u>
Z1	19 Jun 2022	120.192467	23.875233	18	13*	<u>0</u> , <u>5</u> , <u>12</u>
Z2	19 Jun 2022	120.256733	24.033033	32	20*	0, 5, 15, 27
Z3	19 Jun 2022	120.311200	24.147400	50	21	<u>0</u> , 5, 17, <u>45</u>
Z4	19 Jun 2022	120.428150	24.266383	27	21	0, 5, 15, 22
Z5	18 Jun 2022	120.584500	24.424800	25	17*	<u>0</u> , 5, 10, <u>22</u>
Z6	18 Jun 2022	120.618033	24.465317	30	23	0, 5, 15, 25
Z7	18 Jun 2022	120.674333	24.558533	26	17*	0, 5, 10, 20
Z8	18 Jun 2022	120.919300	24.878467	25	17	<u>0</u> , 5, 12, <u>20</u>
Z9	17 Jun 2022	120.991300	24.980033	13	10*	0, 5, 9
Z10	17 Jun 2022	121.056183	25.060767	22	12*	<u>0</u> , 5, 10, 17

^aZe is the euphotic zone depth, defined as the depth of 1 % of surface photosynthetically active radiation. * denotes values derived from the linear regression between measured Ze and bottom-water depth (Eq. (3)).

^bUnderlined depths indicate subsampling depths for carbonate chemistry analysis.

535 **Table A2: Inferred sources of terrigenous OM and results of the lignin-based estimation**

Site	Depth (m)	Inferred source of terrigenous OM	POC _{terr} (μg L ⁻¹)	<i>f</i> _{terr} ^a	POC _{corr} (μg L ⁻¹)	C/N _{corr}	δ ¹³ C _{OC,corr} (‰)
X1	0	Pearl River TSM	3	0.03	105	6.54	-22.87
X1	5	X1 surface sediment	1	0.01	86	6.77	-23.29
X1	25	X1 surface sediment	BDL ^b	NA ^b	NA	NA	NA
X1	48	X1 surface sediment	3	0.03	96	7.76	-24.22
X2	0	Pearl River TSM	10	0.04	218	6.72	-21.18
X2	5	X2 surface sediment	1	0.01	103	7.37	-24.04
X2	35	X2 surface sediment	BDL	NA	NA	NA	NA
X2	58	X2 surface sediment	13	0.13	97	7.31	-23.21
X3	0	Pearl River TSM	7	0.06	113	7.23	-23.14
X3	5	X3 surface sediment	BDL	NA	NA	NA	NA
X3	30	X3 surface sediment	BDL	NA	NA	NA	NA
X3	50	X3 surface sediment	2	0.02	98	7.81	-23.59
X3	65	X3 surface sediment	8	0.08	93	8.05	-24.15
X4	0	Pearl River TSM	10	0.10	91	7.58	-24.09
X4	5	X4 surface sediment	3	0.05	61	7.10	-24.64
X4	40	X4 surface sediment	6	0.10	57	7.48	-24.48
X4	73	X4 surface sediment	20	0.28	71	8.01	-23.51
X5	0	Pearl River TSM	22	0.17	125	6.83	-23.63
X5	5	X5 surface sediment	2	0.02	82	7.57	-25.17
X5	10	X5 surface sediment	2	0.02	92	7.59	-24.97

Site	Depth (m)	Inferred source of terrigenous OM	POC _{terr} ($\mu\text{g L}^{-1}$)	f_{terr}^a	POC _{corr} ($\mu\text{g L}^{-1}$)	C/N _{corr}	$\delta^{13}\text{C}_{\text{OC,corr}}$ (‰)
X5	40	X5 surface sediment	5	0.07	71	6.59	-24.46
X5	72	X5 surface sediment	12	0.15	78	7.41	-23.46
Y1	0	Pearl River TSM	4	0.05	88	7.26	-22.37
Y1	5	Y1 surface sediment	4	0.05	80	7.37	-22.79
Y1	25	Y1 surface sediment	9	0.10	88	8.47	-23.86
Y1	45	Y1 surface sediment	6	0.06	92	7.81	-23.07
Y2	0	Pearl River TSM	7	0.07	109	7.82	-22.36
Y2	5	Y2 surface sediment	3	0.03	86	8.05	-22.88
Y2	20	Y2 surface sediment	BDL	NA	88	7.44	-23.32
Y2	45	Y2 surface sediment	3	0.04	74	7.59	-24.14
Y3	0	Pearl River TSM	7	0.05	128	7.93	-23.10
Y3	5	Y3 surface sediment	2	0.02	85	8.51	-23.43
Y3	25	Y3 surface sediment	BDL	NA	96	7.59	-24.12
Y3	48	Y3 surface sediment	4	0.05	77	8.06	-24.30
Y4	0	Pearl River TSM	11	0.12	98	6.94	-23.77
Y4	5	Y4 surface sediment	BDL	NA	115	8.14	-23.75
Y4	25	Y4 surface sediment	4	0.05	87	7.75	-23.90
Y4	53	Y4 surface sediment	BDL	NA	76	6.94	-23.34
Y5	0	Pearl River TSM	3	0.02	114	7.11	-23.61
Y5	5	Y4 surface sediment	BDL	NA	65	7.05	-24.61
Y5	40	Y4 surface sediment	8	0.13	59	7.47	-24.21
Y5	67	Y4 surface sediment	31	0.63	49	11.14	-25.08
Y6	0	Pearl River TSM	5	0.05	97	7.27	-24.01
Y6	5	Y4 surface sediment	3	0.05	65	7.71	-24.70
Y6	35	Y4 surface sediment	BDL	NA	NA	NA	NA
Y6	90	Y4 surface sediment	8	0.08	95	9.88	-25.02
Z1	0	Taiwanese rivers TSM	40	0.20	200	6.80	-21.07
Z1	5	Taiwanese rivers TSM	16	0.09	181	6.87	-20.54
Z1	12	Taiwanese rivers TSM	32	0.23	140	6.42	-20.56
Z2	0	Taiwanese rivers TSM	63	0.32	194	9.04	-23.26
Z2	5	Taiwanese rivers TSM	13	0.09	133	7.01	-21.70
Z2	15	Taiwanese rivers TSM	8	0.07	108	7.05	-22.88
Z2	27	Taiwanese rivers TSM	12	0.11	108	7.29	-22.44
Z3	0	Taiwanese rivers TSM	58	0.33	173	7.55	-21.25
Z3	5	Z3 surface sediment	3	0.02	145	7.56	-23.57
Z3	17	Z3 surface sediment	4	0.04	103	7.44	-23.17
Z3	45	Z3 surface sediment	2	0.02	106	7.95	-23.42
Z4	0	Taiwanese rivers TSM	39	0.22	175	10.50	-24.20
Z4	5	Z4 surface sediment	1	0.01	132	6.46	-22.54
Z4	15	Z4 surface sediment	1	0.01	118	6.94	-23.03
Z4	22	Z4 surface sediment	2	0.02	123	6.62	-22.89
Z5	0	Taiwanese rivers TSM	24	0.08	288	7.06	-20.66
Z5	5	Z5 surface sediment	1	0.01	267	7.10	-20.93
Z5	10	Z5 surface sediment	3	0.01	247	7.47	-21.81
Z5	22	Z5 surface sediment	4	0.02	182	6.75	-21.47
Z6	0	Taiwanese rivers TSM	15	0.08	194	6.89	-21.53
Z6	5	Z6 surface sediment	2	0.02	143	7.69	-22.72
Z6	15	Z6 surface sediment	1	0.01	135	7.55	-22.93
Z6	25	Z6 surface sediment	3	0.02	107	6.73	-22.76
Z7	0	Taiwanese rivers TSM	22	0.10	229	6.70	-20.94
Z7	5	Z7 surface sediment	1	0.004	223	6.87	-21.09
Z7	10	Z7 surface sediment	1	0.01	152	7.13	-22.06
Z7	20	Z7 surface sediment	1	0.004	121	7.11	-22.40
Z8	0	Z8 surface sediment	17	0.07	252	6.29	-22.75

Site	Depth (m)	Inferred source of terrigenous OM	POC _{terr} ($\mu\text{g L}^{-1}$)	$f_{\text{terr}}^{\text{a}}$	POC _{corr} ($\mu\text{g L}^{-1}$)	C/N _{corr}	$\delta^{13}\text{C}_{\text{OC,corr}}$ (‰)
Z8	5	Z8 surface sediment	9	0.07	140	6.59	-22.24
Z8	12	Z8 surface sediment	11	0.10	112	7.06	-22.63
Z8	20	Z8 surface sediment	3	0.03	123	7.71	-23.50
Z9	0	Z9 surface sediment	14	0.06	226	7.58	-22.80
Z9	5	Z9 surface sediment	21	0.13	157	7.39	-23.72
Z9	9	Z9 surface sediment	10	0.05	200	7.41	-23.40
Z10	0	Z10 surface sediment	59	0.18	331	8.41	-23.92
Z10	5	Z10 surface sediment	12	0.12	102	8.29	-24.22
Z10	10	Z10 surface sediment	10	0.03	330	8.63	-24.40
Z10	17	Z10 surface sediment	40	0.07	538	8.20	-23.69
<i>Mean</i>				<i>0.06</i>			
<i>σ</i>				<i>0.07</i>			

^aFractional contribution of POC_{terr} to measured POC.

^bBDL, below detection limit of lignin; NA, not applicable.

Table A3: Geochemical properties of potential terrigenous OM sources. Unless otherwise noted, data are from Lin et al. (2025a)

Type	Site(s) or location	M_z^{a} (μm)	$\Lambda\delta_{\text{terr,s}}$ (mg lignin g^{-1} OC)	N/C _{terr,s}	$\delta^{13}\text{C}_{\text{terr,s}}$ (‰)
Pearl River TSM ^b	LHS1	NA	11.2	0.083	-26.3
Taiwanese rivers TSM ^c	JSB, DJB, CCB	18	4.6	0.122	-25.42
Surface sediment ^d	X1	23	26.1	0.166	-23.97
Surface sediment	X2	19	11.4	0.160	-23.16
Surface sediment	X3	19	11.5	0.164	-23.24
Surface sediment	X4	19	5.9	0.160	-23.18
Surface sediment	X5	22	11.2	0.158	-23.05
Surface sediment	Y1	41	14.5	0.180	-24.29
Surface sediment	Y2	20	12.2	0.176	-23.71
Surface sediment	Y4	30	5.3	0.190	-23.18
Surface sediment	Z3	44	12.2	0.181	-23.71
Surface sediment	Z4	16	35.9	0.136	-23.89
Surface sediment	Z5	13	30.5	0.117	-24.79
Surface sediment	Z6	13	33.2	0.127	-25.45
Surface sediment	Z8	28	44.3	0.116	-25.19
Surface sediment	Z9	24	72.4	0.131	-24.20
Surface sediment	Z10	18	46.8	0.118	-24.59

540 ^aMean grain size.

^bFor Pearl River TSM, $\Lambda\delta_{\text{terr,s}}$, N/C_{terr,s}, and $\delta^{13}\text{C}_{\text{terr,s}}$ are equal to the measured values (Zhang et al., 2014).

^cFor Taiwanese rivers TSM, $\Lambda\delta_{\text{terr,s}}$ was calculated using $\Lambda\delta = 0.0490 \times Q_w^{0.9847}$ ($r^2 = 0.94$, $n = 7$; derived from Lin et al., 2025a), where Q_w is the fluvial water discharge ($\sim 100 \text{ m}^3 \text{ s}^{-1}$ prior to and during the cruise NOR3-0104). N/C_{terr,s} and $\delta^{13}\text{C}_{\text{terr,s}}$ are taken from measured values.

545 ^dFor seabed sediments of the northeastern Taiwan Strait, these parameters were derived from the output of the mixing models presented in Lin et al. (2025a).

Table A4: Endmember values and results of the binary mixing model for high-TSM ($\geq 4 \text{ mg L}^{-1}$) samples

Site	Depth (m)	$\delta^{13}\text{C}_{\text{sed}}^{\text{a}}$ (‰)	$\delta^{13}\text{C}_{\text{mar}}^{\text{b}}$ (‰)	$f_{\text{sed}}^{\text{c}}$	$f_{\text{terr, sed}}^{\text{d}}$	POC_{terr} ($\mu\text{g L}^{-1}$)	f_{terr}	POC_{corr} ($\mu\text{g L}^{-1}$)	C/N _{corr}	$\Lambda_{8\text{cal}}^{\text{e}}$ ($\text{mg g}^{-1} \text{ OC}$)	$\Lambda_{8\text{cal}}/\Lambda_{8\text{terr, s}}^{\text{f}}$
X2	58	-22.91	-24.91	0.86	0.92	88	0.79	23	16.71	1.7	0.15
X3	65	-22.96	-25.09	0.51	0.92	47	0.47	54	10.55	1.9	0.17
X4	73	-22.74	-25.34	0.75	0.86	59	0.65	32	12.22	2.0	0.34
X5	72	-22.87	-25.65	0.82	0.94	69	0.77	21	14.07	1.9	0.17
Z1	0	-25.42	-21.55	0.11	NA ^g	NA	NA	NA	NA	NA	NA
Z1	5	-25.42	-22.21	< 0	NA	NA	NA	NA	NA	NA	NA
Z1	12	-25.42	-22.45	< 0	NA	NA	NA	NA	NA	NA	NA
Z2	0	-25.42	-21.81	0.57	1.00	146	0.57	111	9.81	2.0	0.43
Z8	0	-25.14	-21.86	0.35	0.99	94	0.65	175	5.62	8.0	0.18
Z9	0	-24.16	-21.74	0.51	0.99	122	0.51	119	7.51	8.4	0.12
Z9	5	-24.16	-22.63	0.77	0.99	135	0.76	43	6.75	11.3	0.16
Z9	9	-24.16	-22.63	0.56	0.99	116	0.55	94	7.16	6.3	0.09
Z10	0	-24.54	-21.97	0.81	0.99	313	0.80	77	8.16	8.8	0.19
Z10	5	-24.54	-22.74	0.84	0.99	96	0.84	18	7.48	5.8	0.12
Z10	10	-24.54	-22.74	0.93	0.99	313	0.92	27	10.47	1.4	0.03
Z10	17	-24.54	-22.76	0.58	0.99	332	0.57	246	7.89	5.6	0.12
Mean							0.68				0.17
σ							0.14				0.10

550 ^aSite-specific endmember $\delta^{13}\text{C}$ values of sedimentary OC, taken from measured values reported in Lin et al. (2024).

^bTemperature-dependent endmember $\delta^{13}\text{C}$ values of marine OC (cf. Sect. 5.2.3).

^cSedimentary fraction of POC.

^dTerrigenous fraction of sedimentary OC, taken from the mixing model outputs presented in Lin et al. (2025a).

^eCalculated source Λ_8 signatures (Eq. (B6)) that account for the measured Λ_8 of the high-TSM samples.

555 ^f $\Lambda_{8\text{terr, s}}$ values are taken from Table A3.

^gNA, not applicable.

Table A5: Data sources and parameter settings used for the CO2SYS computations

Variable or constant	Value or constant setting	Note
DIC	Measured values	
pH	Measured values	
Salinity	Measured values	Sensor data
Input temperature	25 °C	Temperature at which the measurements were performed
Output temperature	Measured values	Sensor data
Input pressure	0 dbar	Pressure at which the measurements were performed
Output pressure	Measured values	Sensor data
Silica	Predicted values	Based on the salinity-silica relationships established using data from Sites A to F of the cruise OR3-632 (Fig. C3a)
Phosphate	Predicted values	Based on the salinity-phosphate relationship established using data from Sites A to F of the cruise OR3-632 (Fig. C3b)
Ammonium	0 $\mu\text{mol kg}^{-1}$	Presumed value
Hydrogen sulfide	0 $\mu\text{mol kg}^{-1}$	Presumed value
Input pH scale	Seawater scale	
K1, K2 constants	Lueker et al., 2000	
KSO4 constant	Dickson, 1990	
KF constant	Perez and Fraga, 1987	
Boron	Uppström, 1974	

Table A6: CO₂(aq) concentrations computed using the CO2SYS program

Site	Depth (m)	CO ₂ (aq) (μmol kg ⁻¹)
X1	0	10.8
X1	5	11.3
X1	25	11.7
X1	48	11.5
X3	0	10.0
X3	5	11.3
X3	65	12.8
X5	0	11.4
X5	40	12.8
X5	72	13.9
Y1	0	10.9
Y1	45	11.2
Y3	0	11.0
Y3	48	11.4
Y6	0	11.2
Y6	90	12.1
Z1	0	10.1
Z1	5	10.9
Z1	12	11.0
Z3	0	10.4
Z3	45	9.2
Z5	0	8.7
Z5	22	9.6
Z8	0	11.0
Z8	20	11.0
Z10	0	11.3
Z10	17	11.5

Table A7: Correlation^a between water-column and sedimentary OM parameters

Sedimentary parameter	Water-column parameter, full-depth integrated		
	Σ8 (mg m ⁻²)	Chl (mg m ⁻²)	POC-weighted δ ¹³ C _{OC} (‰)
<i>Category i</i>			
Σ8 (μg g ⁻¹ dw)	0.42	- ^b	-
Λ8 ^c (mg g ⁻¹ OC)	0.65*	-	-
Burial flux of lignin (mg m ⁻² y ⁻¹)	-0.01	-	-
<i>Category ii</i>			
Chl (ng g ⁻¹ dw)	-	-0.32	-
Chl (μg g ⁻¹ OC)	-	-0.26	-
Burial flux of Chl (μg m ⁻² y ⁻¹)	-	0.16	-
Pheo ^c (μg g ⁻¹ dw)	-	0.27	-
Pheo (mg g ⁻¹ OC)	-	0.54*	-
Burial flux of Pheo (mg m ⁻² y ⁻¹)	-	0.40	-
<i>Category iii</i>			
δ ¹³ C _{OC} (‰)	-	-	-0.75**

^aValues in bold indicate strong correlations ($r \geq 0.7$ or ≤ -0.7). ** and * denote significance at the 0.01 and 0.05 levels, respectively.

^bNot applicable.

^cΛ8, OC-normalized concentration of eight lignin monomers; Pheo, pheopigments a.

Appendix B

570 Text B1. Lignin-based estimation of POC_{terr}

Lignin is a powerful biomarker for diagnosing the relative abundance of terrigenous OM. Its use for quantifying POC_{terr} , however, is not straightforward and relies on two assumptions: *i*) $\Delta\delta_{\text{terr}}$ remains constant during transport and sorting from source regions to shelf waters, and *ii*) particulate lignin experiences negligible alteration (e.g., dissolution or oxidation). Observations in our samples that may challenge these assumptions, and their implications for POC_{terr} quantification, are discussed in Sect. 5.1.2.

After deriving POC_{terr} with Eq. (4), we calculated POC, N/C, and $\delta^{13}\text{C}_{\text{OC}}$ corrected for terrigenous inputs using following mixing equations:

$$\text{POC}_{\text{corr}} = \text{POC}_{\text{m}} - \text{POC}_{\text{terr}} \quad (\text{B1})$$

$$\text{N/C}_{\text{corr}} = [\text{POC}_{\text{m}} \times (\text{N/C})_{\text{m}} - \text{POC}_{\text{terr}} \times (\text{N/C})_{\text{terr,s}}] / \text{POC}_{\text{corr}} \quad (\text{B2})$$

$$580 \quad \delta^{13}\text{C}_{\text{OC,corr}} = (\text{POC}_{\text{m}} \times \delta^{13}\text{C}_{\text{OC,m}} - \text{POC}_{\text{terr}} \times \delta^{13}\text{C}_{\text{terr,s}}) / \text{POC}_{\text{corr}} \quad (\text{B3})$$

where subscripts m and corr denote measured and corrected values for POM, respectively, and s refers to the inferred source material.

Text B2. Binary mixing model with floating endmembers for estimating POC_{terr} of high-TSM samples

We applied a $\delta^{13}\text{C}$ -based binary mixing model that treats POC as a mixture of sedimentary (sed) and marine (mar) sources:

$$585 \quad f_{\text{sed}} + f_{\text{mar}} = 1 \quad (\text{B4})$$

$$f_{\text{sed}} \times \delta^{13}\text{C}_{\text{sed}} + f_{\text{mar}} \times \delta^{13}\text{C}_{\text{mar}} = \delta^{13}\text{C}_{\text{OC,m}} \quad (\text{B5})$$

where f is the fractional contribution of each OC component to bulk POC. Endmember $\delta^{13}\text{C}$ values (Table A4) vary spatially for $\delta^{13}\text{C}_{\text{sed}}$ and with temperature for $\delta^{13}\text{C}_{\text{mar}}$. Some samples produced negative f_{sed} values (Table A4), likely reflecting uncertainties in the $\delta^{13}\text{C}_{\text{OC}}$ -temperature relationship. Samples with f_{sed} near or below zero were grouped with low-TSM samples for evaluation of marine POM. f_{sed} values above 0.3 were used to calculate POC_{terr} (Eq. (5)), which was further used to derive POC_{corr} , N/C_{corr} , and $\Delta\delta_{\text{cal}}$ (Eqs. (B1), (B2), and (6); Table A4).

Text B3. Estimating the advective POC_{terr} flux out of the northeastern Taiwan Strait

We defined a latitudinal transect at 25.04° N (120.40–121.04° E; Fig. 1b) as the northern boundary of the study area and quantified the export flux as the rate of material crossing this transect.

590 A previous study of nutrient fluxes through the Taiwan Strait (Huang et al., 2019) used polynomial regressions linking measured concentrations to observed temperature and salinity, which were then related to HYCOM outputs of temperature and salinity. This approach was not suitable for POC_{terr} because of sparse data coverage and weak regression performance ($r^2 < 0.3$; data not shown). Instead, we used POC_{terr} concentrations derived from summer samples to calculate the annual flux. Interpolation along the transect used measurements from Sites X5, Y6, Z9, and Z10. To restrict sediment resuspension to the

600 known offshore and nearshore mud belts, dummy stations were added at the belt margins (Fig. 1b) and assigned the POC_{terr} value of Site Y6. Because our sampling took place under calm sea states, resuspension was likely mild, making the flux estimate conservative. Although simplified, the approach is sufficient for comparing the export flux with riverine inputs from Taiwan.

Flow rates along the transect were obtained from HYCOM, which provides $1/12^\circ$ horizontal resolution and 41 vertical layers. 605 Daily east-west (u) and north-south (v) velocity data for 2022 were averaged to derive the annual mean velocity (\bar{u} , \bar{v}) used in flux estimation.


Interpolation of POC_{terr} and flow speed was carried out using the Data Interpolating Variational Analysis (DIVA) function in Ocean Data View (Fig. C5). Outputs were exported as $1 \text{ km} \times 1 \text{ m}$ grid cells using the 2-D estimation function. The export flux (F) of each grid cell was calculated as:

$$610 \quad F = POC_{terr} \times A \times \sqrt{\bar{u}^2 + \bar{v}^2} \quad (B7)$$


A is the area of the grid cell. Figure C5c shows the resulting cross section of POC_{terr} export flux. The total flux was obtained by summing across all grid cells. The uncertainty was propagated from known sources, including analytical errors in POC (3 %) and lignin (20 %), as well as uncertainty in the $\delta^{13}C_{OC}$ -temperature relationship (11 %).

615


Appendix C



620 **Figure C1: Correlation between $\delta^{13}\text{C}_{\text{OC,corr}}$ (from Step 1) and temperature (T) versus the upper-bound TSM in each data subset. The correlation coefficient declined sharply when TSM exceeded 4 mg L^{-1} , so this value (dashed gray line) was used to separate low- and high-TSM samples.**




625 **Figure C2: Depth-referenced T-S diagram for the NOR3-0104 sampling sites in the northeastern Taiwan Strait. Characteristic summer T-S properties of South China Sea Water (SCSW) and Kuroshio Water (KW) were digitized from Jan et al. (2010). For comparison, T-S properties from Sites A to F of cruise OR3-632 and from Transect T1 (green shade; Wong et al., 2015) are also shown. Light green indicates water properties characteristic of middle-shelf stations in the northern South China Sea (SCS), whereas dark green represents those of open SCS. Samples with salinity lower than typical SCSW values were divided into two groups, one referred to as the Taiwan Coastal Water (TCW) and the other associated with the characteristics of South China Sea Surface Water (SCSSW).**




630

Figure C3: Salinity-based empirical relationships used to predict silica and phosphate concentrations. (a) Salinity-silica relationships. (b) Salinity-phosphate relationship. Data from Sites A to F of the cruise OR3-632.



635

Figure C4: Spatial distribution of $\Delta 8_{\text{terr}}$ in seabed sediments of the northeastern Taiwan Strait. White dots mark marine sediment sampling sites, and riverine TSM data (stars) are also shown with colors matching the legend. The thick gray line denotes the boundary (mean grain size = $63 \mu\text{m}$; Fig. 1b) of mud belts. Data from Lin et al. (2025a).



640 **Figure C5: Data used for estimating the POC_{terr} export flux. (a) POC_{terr} concentration. Black and white dots are CTD samples and dummy (labeled with d above the frame) stations, respectively. (b) Annual mean flow speed, with black dots marking HYCOM grid points. (c) Calculated POC_{terr} export flux from the northeastern Taiwan Strait. The bottommost 5 m of the water column were excluded from the calculation.**




Figure C6. POC-Chl relationships for different sample subsets. (a) Nearshore samples. (b) Offshore samples from the DCM layers. The black line denotes the POC-Chl relationship of low-TSM samples, as shown in Fig. 6b.

Data availability

The hydrographic, POM, and carbonate chemistry data are available from Lin et al. (2025b).

650 **Author contributions**

Yu-Shih Lin: Writing – original draft, Funding acquisition, Formal analysis, Data curation, Conceptualization. Shu-Ying Chuang: Writing – review and editing, Investigation, Formal analysis, Data curation. Yuan-Pin Chang: Writing – review and editing, Resources, Methodology. Chieh-Wei Hsu: Writing – review and editing, Investigation. Hui-Ling Lin: Writing – review and editing, Resources, Methodology. James T. Liu: Writing – review and editing, Resources, Project administration.

655 Wei-Jen Huang: Writing – original draft, Funding acquisition, Formal analysis, Data curation, Conceptualization.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We thank the captain and crew of RV *New Ocean Researcher 3* for their competent work. We thank Jia-Shan Weng for the assistance of fieldwork, Yi-Hsuan Chen for logistic support, Mei-Huei Huang and Ai-Lin Lyu for the assistance of laboratory work, and Wei-Teh Li and I-Huan Lee for providing the HYCOM data. During the preparation of this work, the authors used ChatGPT 5.1 to improve readability and language. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

660

Financial support

665 This work was supported by the National Science and Technology Council [grant number NSTC 111-2611-M-110-022 and 111-2611-M-110-024 to YSL, and MOST 107-2611-M-110-007 and 108-2611-M-110-010 to WJH], Taiwan.

References

Aminot, A. and Rey, F.: Chlorophyll a: Determination by spectroscopic methods, ICES Techniques in Marine Environmental Sciences, 30, 17 pp., [https://doi.org/ 10.25607/OBP-278](https://doi.org/10.25607/OBP-278), 2001.

- 670 Andrade, J. M. and Estévez-Pérez, M. G.: Statistical comparison of the slopes of two regression lines: A tutorial, *Anal. Chim. Acta*, 838, 1–12, <https://doi.org/10.1016/j.aca.2014.04.057>, 2014.
- Arteage, L., Phalow, M., and Oschlies, A.: Global patterns of phytoplankton nutrient and light colimitation inferred from an optimality-based model, *Global Biogeochem. Cycles*, 28, 648–661, <https://doi.org/10.1002/2013GB004668>, 2014.
- Bai, Y., Huang, T. H., He, X., Wang, S. L., Hsin, Y. C., Wu, C. R., Zhai, W., Lui, H. K., and Chen, C. T. A.: Intrusion of the Pearl River plume into the main channel of the Taiwan Strait in summer, *J. Sea Res.*, 95, 1–15, <https://doi.org/10.1016/j.seares.2014.10.003>, 2015.
- 675 Bao, R., van der Voort, T., Zhao, M., Guo, X., Montluçon, D. B., McIntyre, C., and Eglinton, T.: Influence of hydrodynamic processes on the fate of sedimentary organic matter on continental margins, *Global Biogeochem. Cycles*, 32, 1420–1432, <https://doi.org/10.1029/2018GB005921>, 2018.
- 680 Benner, R., Fogel, M. L., Sprague, E. K., and Hodson, R. E.: Depletion of ^{13}C in lignin and its implications for stable carbon isotope studies, *Nature*, 329, 708–710, <https://doi.org/10.1038/329708a0>, 1987.
- Berger, W. H. and Wefer, G.: Export production: seasonality and intermittency, and paleoceanographic implications, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 89, 245–254, [https://doi.org/10.1016/0031-0182\(90\)90065-F](https://doi.org/10.1016/0031-0182(90)90065-F), 1990.
- Bianchi, T. S.: The role of terrestrially derived organic carbon in the coastal ocean: A changing paradigm and the priming effect, *Proc. Natl. Acad. Sci. U.S.A.*, 108, 19473–19481, <https://doi.org/10.1073/pnas.1017982108>, 2011.
- 685 Bianchi, T. S., Cui, X., Blair, N. E., Burdige, D. J., Eglinton, T. I., and Galy, V.: Centers of organic carbon burial and oxidation at the land-ocean interface, *Org. Geochem.*, 115, 138–155, <https://doi.org/10.1016/j.orggeochem.2017.09.008>, 2018.
- Bianchi, T. S., Lambert, C. D., Santschi, P. H., and Guo, L.: Sources and transport of land-derived particulate and dissolved organic matter in the Gulf of Mexico (Texas shelf/slope): The use of lignin-phenols and loliolides as biomarkers, *Org. Geochem.*, 27, 65–78, [https://doi.org/10.1016/S0146-6380\(97\)00040-5](https://doi.org/10.1016/S0146-6380(97)00040-5), 1997.
- 690 Brown, S. L., Landry, M. R., Neveux, J., and Dupouy, C.: Microbial community abundance and biomass along a 180° transect in the equatorial Pacific during an El Niño–Southern Oscillation cold phase, *J. Geophys. Res.*, 108, C12, 8319, <https://doi.org/10.1029/2001JC000817>, 2003.
- 695 Burdige, D. J.: Burial of terrestrial organic matter in marine sediments: A re-assessment, *Global Biogeochem. Cycles*, 19, GB4011, <https://doi.org/10.1029/2004GB002368>, 2005.
- Burdige, D. J.: Preservation of organic matter in marine sediments: Controls, Mechanisms, and an imbalance in sediment organic carbon budgets? *Chem. Rev.*, 107, 467–485, <https://doi.org/10.1021/cr050347q>, 2007.
- Burkhardt, S., Riebesell, U., and Zondervan, I.: Effects of growth rate, CO_2 concentration, and cell size on the stable carbon isotope fractionation in marine phytoplankton, *Geochim. Cosmochim. Acta*, 63, 3729–3741, [https://doi.org/10.1016/S0016-7037\(99\)00217-3](https://doi.org/10.1016/S0016-7037(99)00217-3), 1999.
- 700 Carr, M.-E., Friedrichs, M. A. M., Schmeltz, M., Aita, M. N., Antoine, D., Arrigo, K. R., Asanuma, I., Aumont, O., Barber, R., Behrenfeld, M., Bidigare, R., Buitenhuis, E. T., Campbell, J., Ciotti, A., Dierssen, H., Dowell, M., Dunne, J., Esaias,

- W., Gentili, B., Gregg, W., Groom, S., Hoepffner, N., Ishizaka, J., Kameda, T., Le Quéré, C., Lohrenz, S., Marra, J.,
705 Mélin, F., Moore, K., Morel, A., Reddy, T. E., Ryan, J., Scardi, M., Smyth, T., Turpie, K., Tilstone, G., Waters, K., and
Yamanaka, Y.: A comparison of global estimates of marine primary production from ocean color, *Deep-Sea Res. II*, 53,
741–770, <https://doi.org/10.1016/j.dsr2.2006.01.028>, 2006.
- Clayton, T. D. and Byrne, R. H.: Spectrophotometric seawater pH measurements: Total hydrogen ion concentration scale
calibration of *m*-cresol purple and at-sea results, *Deep-Sea Res. I*, 40, 2115–2129, <https://doi.org/10.1016/0967->
710 0637(93)90048-8, 1993.
- Close, H. G. and Henderson, L. C.: Open-ocean minima in $\delta^{13}\text{C}$ values of particulate organic carbon in the lower euphotic
zone, *Front. Mar. Sci.*, 7, 540165, <https://doi.org/10.3389/fmars.2020.540165>, 2020.
- Dickson, A. G.: Standard potential of the reaction $\text{AgCl(s)} + 12\text{H}_2\text{(g)} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of
the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K, *J. Chem. Thermodyn.*, 22, 113–127,
715 [https://doi.org/10.1016/0021-9614\(90\)90074-Z](https://doi.org/10.1016/0021-9614(90)90074-Z), 1990.
- Dunne, J. P., Sarmiento, J. L., and Gnanadesikan, A.: A synthesis of global particle export from the surface ocean and
cycling through the ocean interior and on the seafloor, *Global Biogeochem. Cycles*, 21, GB4006,
<https://doi.org/10.1029/2006GB002907>, 2007.
- Eppley, R. W.: An incubation method for estimating the carbon content of phytoplankton in natural samples, *Limnol.*
720 *Oceanog.*, 13, 574–582, <https://doi.org/10.4319/lo.1968.13.4.0574>, 1968.
- Fielding, A. S., Turpin, D. H., Guy, R. D., Calvert, S. E., Crawford, D. W., and Harrison, P. J.: Influence of the carbon
concentrating mechanism on carbon stable isotope discrimination by the marine diatom *Thalassiosira pseudonana*, *Can. J.*
Bot., 76, 1098–1103, <https://doi.org/10.1139/b98-069>, 1998.
- Fry, B.: $^{13}\text{C}/^{12}\text{C}$ fractionation by marine diatoms, *Mar. Ecol. Prog. Ser.*, 134, 283–294, <https://doi.org/10.3354/meps134283>,
725 1996.
- Galy, V., Peucker-Ehrenbrink, B., and Eglinton, T.: Global carbon export from the terrestrial biosphere controlled by erosion,
Nature, 521, 204–207, <https://doi.org/10.1038/nature14400>, 2015.
- Gao, L., Li, D., and Ishizaka, J.: Stable isotope ratios of carbon and nitrogen in suspended organic matter: Seasonal and
spatial dynamics along the Changjiang (Yangtze River) transport pathway, *J. Geophys. Res. Biogeosci.*, 119, 1717–1737,
730 <https://doi.org/10.1002/2013JG002487>, 2014.
- Geider, R. J.: Light and temperature dependence of the carbon to chlorophyll a ratio in microalgae and cyanobacteria:
Implications for physiology and growth of phytoplankton, *New Phytol.*, 106, 1–34, <https://doi.org/10.1111/j.1469->
8137.1987.tb04788.x, 1987.
- Giordano, M., Beardall, J., and Raven, J. A.: CO_2 concentrating mechanisms in algae: Mechanisms, environmental
735 modulation, and evolution, *Annu. Rev. Plant Biol.*, 56, 99–131,
<https://doi.org/10.1146/annurev.arplant.56.032604.144052>, 2005.

- Gui, J. and Sun, J.: Phytoplankton carbon to chlorophyll a model development: a review, *Front. Mar. Sci.*, 11, 1466072, <https://doi.org/10.3389/fmars.2024.1466072>, 2024.
- 740 Guo, W., Ye, F., Xu, S., and Jia, G.: Seasonal variation in sources and processing of particulate organic carbon in the Pearl River estuary, South China, *Estuar. Coast. Shelf Sci.*, 167, 540–548, <https://doi.org/10.1016/j.ecss.2015.11.004>, 2015.
- Hayes, J. M., Strauss, H., and Kaufman, A. J.: The abundance of ^{13}C in marine organic matter and isotopic fractionation in the global biogeochemical cycle of carbon during the past 800 Ma, *Chem. Geol.*, 161, 103–125, [https://doi.org/10.1016/S0009-2541\(99\)00083-2](https://doi.org/10.1016/S0009-2541(99)00083-2), 1999.
- 745 Hedges, J. I. and Ertel, J. R.: Characterization of lignin by gas capillary chromatography of cupric oxide oxidation products, *Anal. Chem.*, 54, 174–178, <https://doi.org/10.1021/ac00239a007>, 1982.
- Hedges, J. I., Keil, R. G., and Benner, R.: What happens to terrestrial organic matter in the ocean? *Org. Geochem.*, 27, 195–212, [https://doi.org/10.1016/S0146-6380\(97\)00066-1](https://doi.org/10.1016/S0146-6380(97)00066-1), 1997.
- Hernes, P. J. and Benner, R.: Terrigenous organic matter sources and reactivity in the North Atlantic Ocean and a comparison to the Arctic and Pacific Oceans, *Mar. Chem.*, 100, 66–79, <https://doi.org/10.1016/j.marchem.2005.11.003>, 750 2006.
- Ho, P. C., Okuda, N., Yeh, C. F., Wang, P. L., Gong, G. C., and Hsieh, C. H.: Carbon and nitrogen isoscape of particulate organic matter in the East China Sea, *Prog. Oceanogr.*, 197, 102667, <https://doi.org/10.1016/j.pocean.2021.102667>, 2021.
- Horng, C. S. and Huh, C. A.: Magnetic properties as tracers for source-to-sink dispersal of sediments: A case study in the Taiwan Strait, *Earth Planet. Sci. Lett.*, 309, 141–152, <https://doi.org/10.1016/j.epsl.2011.07.002>, 2011.
- 755 Huang, C., Chen, F., Zhang, S., Chen, C., Meng, Y., Zhu, Q., and Song, Z.: Carbon and nitrogen isotopic composition of particulate organic matter in the Pearl River Estuary and the adjacent shelf, *Estuar. Coast. Shelf Sci.*, 246, 107003, <https://doi.org/10.1016/j.ecss.2020.107003>, 2020.
- Huang, T. H., Chen, C. T. A., Lee, J., Wu, C. R., Wang, Y. L., Bai, Y., He, X., Wang, S. L., Kandasamy, S., Lou, J. Y., Tsuang, B. J., Chen, H. W., Tseng, R. S., and Yang, Y. J.: East China Sea increasingly gains limiting nutrient P from 760 South China Sea, *Sci. Rep.*, 9, 5648, <https://doi.org/10.1038/s41598-019-42020-4>, 2019.
- Huang, W. J., Kao, K. J., Lin, Y. S., Chen, C. T. A., and Liu, J. T.: Daily to weekly impacts of mixing and biological activity on carbonate dynamics in a large river-dominated shelf, *Estuar. Coast. Shelf Sci.*, 245, 106914, <https://doi.org/10.1016/j.ecss.2020.106914>, 2020.
- Huang, W. J., Wang, Y., and Cai, W. J.: Assessment of sample storage techniques for total alkalinity and dissolved inorganic carbon in seawater, *Limnol. Oceanogr.: Methods*, 10, 711–717, <https://doi.org/10.4319/lom.2012.10.711>, 2012.
- Huang, Y. N.: Variability of hydrochemical parameters and dissolved trace element (Cd, Cu, Fe, Mn, Ni, Pb, Zn) concentrations in nearshore coastal waters off Taiwan, M.Sc. thesis, National Sun Yat-sen University, Taiwan, 196 pp., 2022.

- Huh, C. A., Chen, W., Hsu, F. H., Su, C. C., Chiu, J. K., Lin, S., Liu, C. S., and Huang, B. J.: Modern (<100 years) sedimentation in the Taiwan Strait: Rates and source-to-sink pathways elucidated from radionuclides and particle size distribution, *Cont. Shelf Res.*, 31, 47–63, <https://doi.org/10.1016/j.csr.2010.11.002>, 2011.
- 770 Jan, S., Sheu, D. D., and Kuo, H. M.: Water mass and throughflow transport variability in the Taiwan Strait, *J. Geophys. Res.*, 111, C12012, <https://doi.org/10.1029/2006JC003656>, 2006.
- 775 Jan, S., Tseng, Y. H., and Dietrich, D. E.: Sources of water in the Taiwan Strait, *J. Oceanogr.*, 66, 211–221, <https://doi.org/10.1007/s10872-010-0019-7>, 2010.
- Jan, S., Wang, J., Chern, C. S., and Chao, S. Y.: Seasonal variation of the circulation in the Taiwan Strait, *J. Mar. Syst.*, 35, 249–268, [https://doi.org/10.1016/S0924-7963\(02\)00130-6](https://doi.org/10.1016/S0924-7963(02)00130-6), 2002.
- Kroopnick, P. M.: The distribution of ^{13}C of ΣCO_2 in the world oceans, *Deep Sea Res. Part A. Oceanogr. Res. Pap.*, 32, 57–84, [https://doi.org/10.1016/0198-0149\(85\)90017-2](https://doi.org/10.1016/0198-0149(85)90017-2), 1985.
- 780 Kukert, H. and Riebesell, U.: Phytoplankton carbon isotope fractionation during a diatom spring bloom in a Norwegian fjord, *Mar. Ecol. Prog. Ser.*, 173, 127–137, <https://doi.org/10.3354/meps173127>, 1998.
- Laws, E. A., Popp, B. N., Bidigare, R. R., Kennicutt, M. C., and Macko, S. A.: Dependence of phytoplankton carbon isotopic composition on growth rate and $[\text{CO}_2]_{\text{aq}}$: Theoretical considerations and experimental results, *Geochim. Cosmochim. Acta*, 59, 1131–1138, [https://doi.org/10.1016/0016-7037\(95\)00030-4](https://doi.org/10.1016/0016-7037(95)00030-4), 1995.
- 785 Lee, J., Liu, J. T., Lin, Y. S., Chen, C. T. A., and Wang, B. S.: The contrast in suspended particle dynamics at surface and near-bottom on the river-dominated northern South China Sea shelf in summer: implication on physics and biogeochemistry coupling, *Front. Mar. Sci.*, 10, 1156915, <https://doi.org/10.3389/fmars.2023.1156915>, 2023.
- Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K_1 and K_2 : validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium, *Mar. Chem.*, 70, 105–119, [https://doi.org/10.1016/S0304-4203\(00\)00022-0](https://doi.org/10.1016/S0304-4203(00)00022-0), 2000.
- 790 Li, Q. P., Franks, P. J. S., Landry, M. R., Goericke, R., and Taylor, A. G.: Modeling phytoplankton growth rates and chlorophyll to carbon ratios in California coastal and pelagic ecosystems, *J. Geophys. Res.*, 115, G04003, <https://doi.org/10.1029/2009JG001111>, 2010.
- Lin, B., Liu, Z., Eglington, T. I., Kandasamy, S., Blattmann, T. M., Haghypour, N., Huang, K. F., and You, C. F.: Island-wide variation in provenance of riverine sedimentary organic carbon: A case study from Taiwan, *Earth Planet. Sci. Lett.*, 539, 116238, <https://doi.org/10.1016/j.epsl.2020.116238>, 2020b.
- 795 Lin, Y. S., Chuang, S. Y., Chang, Y. P., Hsu, C. W., Lin, H. L., Liu, J. T., and Huang, W. J.: Hydrographic, particulate organic matter, and carbonate chemistry data of the northeastern Taiwan Strait, Zenodo [data set], <https://doi.org/10.5281/zenodo.18066734>, 2025b.
- 800 Lin, Y. S., Lee, J., Lin, L. H., Fu, K. H., Chen, C. T. A., Wang, Y. H., and Lee, I. H.: Biogeochemistry and dynamics of particulate organic matter in a shallow-water hydrothermal field (Kueishantao Islet, NE Taiwan), *Mar. Geol.*, 422, 106121, <https://doi.org/10.1016/j.margeo.2020.106121>, 2020a.

- Lin, Y. S., Wei, C. L., Chang, Y. P., Wang, J. T., Chuang, S. Y., Wang, L. C., Liu, J. T., Li, H. C., and Hsu, C. W.: Hydrographic and sediment geochemical data of the northeastern Taiwan Strait, Zenodo [data set],
805 <https://doi.org/10.5281/zenodo.13336522>, 2024.
- Lin, Y. S., Wei, C. L., Chang, Y. P., Wang, J. T., Chuang, S. Y., Wang, L. C., Liu, J. T., Li, H. C., and Hsu, C. W.: Linking source-to-sink processes, organic matter composition, and oxygen consumption on a shallow passive margin shelf sustained by small mountainous rivers, *Geochim. Cosmochim. Acta*, 408, 40–55,
<https://doi.org/10.1016/j.gca.2025.09.031>, 2025a.
- 810 Liu, J. T., Hsu, R. T., Yang, R. J., Wang, Y. P., Wu, H., Du, X., Li, A., Chien, S. C., Lee, J., Yang, S., Zhu, J., Su, C. C., Chang, Y., and Huh, C. A.: A comprehensive sediment dynamics study of a major mud belt system on the inner shelf along an energetic coast, *Sci. Rep.*, 8, 4229, <https://doi.org/10.1038/s41598-018-22696-w>, 2018b.
- Liu, K. K., Kao, S. J., Hu, H. C., Chou, W. C., Hung, G. W., and Tseng, C. M.: Carbon isotopic composition of suspended and sinking particulate organic matter in the northern South China Sea—From production to deposition, *Deep-Sea Res.*,
815 54, 1504–1527, <https://doi.org/10.1016/j.dsr2.2007.05.010>, 2007.
- Liu, Q., Kandasamy, S., Lin, B., Wang, H., and Chen, C. T. A.: Biogeochemical characteristics of suspended particulate matter in deep chlorophyll maximum layers in the southern East China Sea, *Biogeosciences*, 15, 2091–2109,
<https://doi.org/10.5194/bg-15-2091-2018>, 2018a.
- Liu, Q., Kandasamy, S., Zhai, W., Wang, H., Veeran, Y., Gao, A., and Chen, C. T. A.: Temperature is a better predictor of
820 stable carbon isotopic compositions in marine particulates than dissolved CO₂ concentration, *Commun. Earth Environ.*, 3, 303, <https://doi.org/10.1038/s43247-022-00627-y>, 2022.
- Martiny, A. C., Vrugt, J. A., Primeau, F. W., and Lomas, M. W.: Regional variation in the particulate organic carbon to nitrogen ratio in the surface ocean, *Global Biogeochem. Cycles*, 27, 723–731, <https://doi.org/10.1002/gbc.20061>, 2013.
- Mook, W. G., Bommerson, J. C., and Staverman, W. H.: Carbon isotope fractionation between dissolved bicarbonate and
825 gaseous carbon dioxide, *Earth Planet. Sci. Lett.*, 22, 169–176, [https://doi.org/10.1016/0012-821X\(74\)90078-8](https://doi.org/10.1016/0012-821X(74)90078-8), 1974.
- Moreno, A. R. and Martiny, A. C.: Ecological stoichiometry of ocean plankton, *Annu. Rev. Mar. Sci.*, 10, 43–69,
<https://doi.org/10.1146/annurev-marine-121916-063126>, 2018.
- Nan, F., Xue, H., and Yu, F.: Kuroshio intrusion into the South China Sea: A review, *Prog. Oceanogr.*, 137, 314–333,
<https://doi.org/10.1016/j.pocean.2014.05.012>, 2015.
- 830 Ocean Data Bank: Ocean data bank/data-policy: v1.1.0, Zenodo [data set], <https://doi.org/10.5281/zenodo.15515930>, 2025.
- Perez, F. and Fraga, F.: Association constant of fluoride and hydrogen ions in seawater, *Mar. Chem.*, 21, 161–168,
[https://doi.org/10.1016/0304-4203\(87\)90036-3](https://doi.org/10.1016/0304-4203(87)90036-3), 1987.
- Pierrot, D., Epitalon, J. M., Orr, J. C., Lewis, E., and Wallace, D. W. R.: MS Excel program developed for CO₂ system calculations—version 3.0, GitHub repository [code], https://github.com/dpierrot/co2sys_xl, 2021.

- 835 Popp, B. N., Laws, E. A., Bidigare, R. R., Dore, J. E., Hanson, K. L., and Wakeham, S. G.: Effect of phytoplankton cell geometry on carbon isotopic fractionation, *Geochim. Cosmochim. Acta*, 62, 69–77, [https://doi.org/10.1016/S0016-7037\(97\)00333-5](https://doi.org/10.1016/S0016-7037(97)00333-5), 1998.
- Rau, G. H., Chavez, F. P., and Friederich, G. E.: Plankton $^{13}\text{C}/^{12}\text{C}$ variations in Monterey Bay, California: evidence of non-diffusive inorganic carbon uptake by phytoplankton in an upwelling environment, *Deep-Sea Res. I*, 48, 79–94, [https://doi.org/10.1016/S0967-0637\(00\)00039-X](https://doi.org/10.1016/S0967-0637(00)00039-X), 2001.
- 840 Rau, G. H., Takahashi, T., and Des Marais, D. J.: Latitudinal variations in plankton $\delta^{13}\text{C}$: implications for CO_2 and productivity in past oceans, *Nature*, 341, 516–518, <https://doi.org/10.1038/341516a0>, 1989.
- Reeves, A. D. and Preston, M. R.: The composition and lignin in estuarine suspended particulates and the distribution of particulate lignin in estuaries as determined by capillary gas chromatography of cupric oxide oxidation products, *Estuar. Coast. Shelf Sci.*, 29, 583–5599, [https://doi.org/10.1016/0272-7714\(89\)90012-7](https://doi.org/10.1016/0272-7714(89)90012-7), 1989.
- 845 Riebesell, U., Burkhardt, S., Dauelsberg, A., and Kroon, B.: Carbon isotope fractionation by a marine diatom: dependence on the growth-rate-limiting resource, *Mar. Ecol. Prog. Ser.*, 193, 295–303, <https://doi.org/10.3354/meps193295>, 2000.
- Schlitzer, R.: Ocean Data View, <https://odv.awi.de>, last access 27 December 2025.
- Sharkey, T. D. and Berry, J. A.: Carbon isotope fractionation of algae as influenced by an inducible CO_2 concentrating mechanism, in: *Inorganic Carbon Uptake by Aquatic Photosynthetic Organisms*, edited by: Lucas, W. J. and Berry, J. A., American Society of Plant Physiologists, Rockville, Md., 389–407, 1985.
- Stephens, M. P., Kadko, D. C., Smith, C. R., and Latasa, M.: Chlorophyll-a and pheopigments as tracers of labile organic carbon at the central equatorial Pacific seafloor, *Geochim. Cosmochim. Acta*, 61, 4605–4619, [https://doi.org/10.1016/S0016-7037\(97\)00358-X](https://doi.org/10.1016/S0016-7037(97)00358-X), 1997.
- 855 Sun, M. Y., Lee, C., and Aller, R. C.: Anoxic and oxic degradation of ^{14}C -labeled chloropigments and a ^{14}C -labeled diatom in Long Island Sound sediments, *Limnol. Oceanogr.*, 38, 1438–1451, <https://doi.org/10.4319/lo.1993.38.7.1438>, 1993.
- Sun, X., Hu, L., Fan, D., Wang, H., Yang, Z., and Guo, Z.: Sediment resuspension accelerates the recycling of terrestrial organic carbon at a large river-coastal ocean interface, *Global Biogeochem. Cycles*, 38, e2024GB008213, <https://doi.org/10.1029/2024GB008213>, 2024.
- 860 Talling, P. J., Hage, S., Baker, M. L., Bianchi, T. S., Hilton, R. G., and Maier, K. L.: The global turbidity current pump and its implications for organic carbon cycling, *Annu. Rev. Mar. Sci.*, 16, 105–133, <https://doi.org/10.1146/annurev-marine-032223-103626>, 2024.
- Tanioka, T. and Matsumoto, K.: A meta-analysis on environmental drivers of marine phytoplankton C:N:P, *Biogeosciences*, 17, 2939–2954, <https://doi.org/10.5194/bg-17-2939-2020>, 2020.
- 865 Tao, S., Wang, A., Liu, J. T., Ye, X., Blattmann, T. M., Ran, C., Liu, Z., Wang, L., Yin, X., Zhang, H., Li, L., Ning, X., Hung, C. C., and Haghypour, N.: Characteristics of sedimentary organic carbon burial in the shallow conduit portion of source-to-sink sedimentary systems in marginal seas, *Geochim. Cosmochim. Acta*, 353, 92–111, <https://doi.org/10.1016/j.gca.2023.05.006>, 2023.

- 870 Tong, Z., Wang, C., Lin, L., Ma, L., and Huang, B.: Nutrient depletion and phytoplankton shifts driven by the Pearl River
plume in the Taiwan Strait, *Front. Mar. Sci.*, 11, 1485670, <https://doi.org/10.3389/fmars.2024.1485670>, 2024.
- Tseng, H. C., You, W. L., Huang, W., Chung, C. C., Tsai, A. Y., Chen, T. Y., Lan, K. W., and Gong, G. C.: Seasonal
variations of marine environment and primary production in the Taiwan Strait, *Front. Mar. Sci.*, 7, 38,
<https://doi.org/10.3389/fmars.2020.00038>, 2020.
- 875 Uppström, L. R.: The boron/chlorinity ratio of deep-sea water from the Pacific Ocean. *Deep Sea Res. Oceanogr. Abstr.*, 21,
161–162, [https://doi.org/10.1016/0011-7471\(74\)90074-6](https://doi.org/10.1016/0011-7471(74)90074-6), 1974.
- Wakeham, S. G., Canuel, E. A., Lerberg, E. J., Mason, P., Sampere, T. P., and Bianchi, T. S.: Partitioning of organic matter
in continental margin sediments among density fractions, *Mar. Chem.*, 115, 211–225,
<https://doi.org/10.1016/j.marchem.2009.08.005>, 2009.
- Wakeham, S. G. and McNichol, A. P.: Transfer of organic carbon through marine water columns to sediments—insights
880 from stable and radiocarbon isotopes of lipid biomarkers, *Biogeosciences*, 11, 6895–6914, <https://doi.org/10.5194/bg-11-6895-2014>, 2014.
- Wang, A., Ye, X., Liu, J. T., Xu, Y., Yin, X., and Xu, X.: Sources of settling particulate organic carbon during summer in
the northern Taiwan Strait, *Estuar. Coast. Shelf Sci.*, 198, 487–496, <https://doi.org/10.1016/j.ecss.2016.10.008>, 2017.
- Wang, X. J., Behrenfeld, M., Le Borgne, R., Murtugudde, R., and Boss, E.: Regulation of phytoplankton carbon to
885 chlorophyll ratio by light, nutrients and temperature in the Equatorial Pacific Ocean: a basin scale model, *Biogeosciences*,
6, 391–404, <https://doi.org/10.5194/bg-6-391-2009>, 2009.
- Wang, Y. H., Jan, S., and Wang, D. P.: Transports and tidal current estimates in the Taiwan Strait from shipboard ADCP
observations (1999–2001), *Estuar. Coast. Shelf Sci.*, 57, 193–199, [https://doi.org/10.1016/S0272-7714\(02\)00344-X](https://doi.org/10.1016/S0272-7714(02)00344-X), 2003.
- Wei, B., Mollenhauer, G., Hefter, J., Grotheer, H., and Jia, G.: Dispersal and aging of terrigenous organic matter in the Pearl
890 River Estuary and the northern South China Sea Shelf, *Geochim. Cosmochim. Acta*, 282, 324–339,
<https://doi.org/10.1016/j.gca.2020.04.032>, 2020.
- Wong, G. T. F., Pan, X., Li, K. Y., Shiah, F. K., Ho, T. Y., and Guo, X.: Hydrography and nutrient dynamics in the northern
South China Sea Shelf-sea (NoSoCS), *Deep-Sea Res. II*, 117, 23–40, <https://doi.org/10.1016/j.dsr2.2015.02.023>, 2015.
- Wu, Y., Zhang, J., Li, D. J., Wei, H., and Lu, R. X.: Isotope variability of particulate organic matter at the PN section in the
895 East China Sea, *Biogeochemistry*, 65, 31–49, <https://doi.org/10.1023/A:1026044324643>, 2003.
- You, Y., Chern, C. S., Yang, Y., Liu, C. T., Liu, K. K., and Pai, S. C.: The South China Sea, a cul-de-sac of North Pacific
Intermediate Water, *J. Oceanogr.*, 61, 509–527, <https://doi.org/10.1007/s10872-005-0059-6>, 2005.
- Zhang, Y., Kaiser, K., Li, L., Zhang, D., Ran, Y., and Benner, R.: Sources, distributions, and early diagenesis of sedimentary
organic matter in the Pearl River region on the South China Sea, *Mar. Chem.*, 158, 39–48,
900 <https://doi.org/10.1016/j.marchem.2013.11.003>, 2014.
- Zhong, X., Liu, J., Shi, M., Liu, X., Lv, Z., and Ran, X.: Nutrient dynamics in the East China Sea: Seasonal changes, budget,
and ecological impacts, *Prog. Oceanogr.*, 234, 103463, <https://doi.org/10.1016/j.pocean.2025.103463>, 2025.

Zhong, Y., Liu, X., Xiao, W., Laws, E. A., Chen, J., Wang, L., Liu, S., Zhang, F., and Huang, B.: Phytoplankton community patterns in the Taiwan Strait match the characteristics of their realized niches, *Prog. Oceanogr.*, 186, 102366, 905 <https://doi.org/10.1016/j.pocean.2020.102366>, 2020.