



Ocean salinity across space-time scales: From water cycle indicator to dynamical driver

Lisan Yu¹

5 ¹Physical Oceanographic Department, Woods Hole Oceanographic Institution, Woods Hole, MA 2543, USA.

Correspondence to: Dr. Lisan Yu (lyu@whoi.edu)

Abstract. Ocean salinity plays a complementary role in the climate system: it integrates changes in the global water cycle while also helping drive ocean circulation through its control on seawater density. Salinity has long been viewed as the ocean's "rain gauge," a largely passive recorder of surface evaporation, precipitation, and runoff. Yet salinity also shapes the currents and mixing that redistribute heat and freshwater, raising a central question: when does salinity mainly record climate forcing, and when does it actively influence climate dynamics? This review synthesizes two decades of satellite and in situ observations within a regime-dependent framework in which salinity's function is set by the competition among freshwater forcing, advection, and mixing timescales. At basin scales (>1000 km) over decades, salinity tracks water-cycle change through pattern amplification, with fresh regions freshening and salty regions becoming saltier. At regional to mesoscale 10 (~10–500 km) and seasonal-to-interannual timescales, salinity traces circulation pathways; subsurface anomalies often reflect subduction and ventilation histories from years earlier. At submesoscales (O(10 km)) and synoptic timescales (hours to days), salinity becomes dynamically active, sharpening density fronts, modulating stratification, and altering mixing in ways that feed back on its own transport and air–sea exchange. Understanding ocean climate response requires resolving regime boundaries where these balances shift. The critical observational gap is global sea-surface salinity at O(10 km), 20 where salinity transitions from passive tracer to active driver yet current satellite products cannot resolve this scale. Observations at regime boundaries would show how water-cycle intensification and ocean circulation changes interact, improving projections of climate change, ocean heat storage and distribution, and ecosystem dynamics at regional and global scales.

25 **1 Introduction**

Ocean salinity occupies a unique position at the nexus of the global water cycle and ocean circulation. As the integrated expression of surface freshwater fluxes (evaporation, precipitation, river discharge, and ice melt), salinity has long been viewed as the ocean's natural rain gauge, faithfully recording the changes in Earth's hydrological cycle (Dickson et al. 1988; Curry et al. 2003; Boyer et al. 2005; Durack & Wijffels 2010; Helm et al. 2010; Skliris et al. 2014; Vinogradova & Ponte, 30 2017; Cheng et al. 2020; Yu et al. 2020). Yet through its fundamental role in setting seawater density alongside temperature, salinity also actively shapes ocean stratification, convection and mixing, and governs large-scale circulation (Talley, 2008; Mignot & Frankignoul 2010; Schanze et al. 2010; Kolodziejczyk & Gaillard, 2013). This dual character of being both



passive recorder and active drive raises a fundamental question: under what conditions does salinity reflect climate forcing, and when does it shape climate dynamics? This review synthesizes how observational advances over the past two decades
35 have revealed that the answer depends critically on spatial and temporal scale, with salinity's role emerging from the competition among forcing, advection, and mixing processes.

The traditional "rain gauge" perspective treated salinity as a passive consequence of atmospheric forcing in which net precipitation regions freshened, net evaporation regions became more saline, and water cycle processes left their signature on surface and subsurface patterns (Warren, 1983; Schmitt, 2008). This framework enabled insights into global water balance
40 (Baumgartner and Reichel, 1975) and paleoclimate reconstruction (Rohling and Bigg, 1998). The apparent correspondence between ocean salinity and evaporation-minus-precipitation (E–P) motivated attempts to infer water cycle variations from salinity observations (Gordon and Giulivi, 2008; Yu, 2011; Terray et al. 2012; Durack et al. 2012; Skliris et al. 2016; Vinogradova & Ponte, 2017; Fournier et al. 2023). Implicit in this framework was a largely one-way causality in which atmospheric forcing determined salinity, and salinity served primarily as a marker of forcing variability, with limited
45 consideration of feedbacks through ocean circulation.

This passive view neglects a fundamental reality: salinity anomalies do not remain where they are forced. They are advected by ocean currents, modified by mixing, and, through their impact on density, feed back on the circulation that transports them. High-latitude freshening can inhibit deep convection, weaken the meridional overturning circulation, and alter basin-scale heat transport (Rahmstorf, 1995; Barreiro et al. 2008; Holliday et al. 2020), while tropical freshening strengthens near-
50 surface stratification, reduces vertical mixing, and modifies upper-ocean thermal structure (Lukas and Lindstrom, 1991; Maes et al., 2002; Mignot et al. 2012; Camara et al. 2015). In evaporative subtropics, persistent E–P excess combined with Ekman convergence maintains subtropical salinity maxima (Qu et al. 2013; Gordon et al., 2015; Bingham et al., 2019; Aubone et al. 2021); subduction of these waters ventilates mode waters and shallow overturning cells, exporting surface freshwater anomalies to the interior and equatorward (Qu et al. 2016; Yu et al. 2018; Zika et al., 2018). At submesoscales,
55 salinity becomes equally important as temperature in driving density gradients, shaping stratification and mediating vertical exchange near surface (Rudnick & Ferrari, 1999; Timmermans & Winsor 2013; Jaeger & Mahadevan 2018; Coadou-Chaventon et al. 2024; Yu 2026). Unlike temperature, salinity variations lack strong restoring feedbacks, as freshwater fluxes force salinity anomalies but do not rapidly relax them, making salinity variations both more persistent and more strongly shaped by advection and mixing than their temperature counterparts (Yu 2011; Vinogradova & Ponte 2013; Lyu et al. 2025). The competition among forcing, advection, and mixing timescales determines whether salinity behaves primarily
60 as a rain gauge recording atmospheric forcing, a passive tracer marking circulation pathways, or a dynamical driver actively modifying ocean stratification and currents.

Over the past two decades, observational advances have transformed our ability to characterize salinity's scale-dependent roles systematically. The Argo program, with thousands of autonomous profiling floats sampling temperature and salinity in



65 the upper 2000 meters, provides near-real-time monitoring of subsurface variability (Roemmich & Gilson., 2009; Riser et al., 2016), while historical ship-based compilations provide century-long context for sea surface salinity changes (Bingham et al. 2002; Boyer et al. 2005; Ishii et al., 2006; Good et al., 2013; Friedman et al., 2017). In parallel, satellite salinity missions, including ESA's Soil Moisture and Ocean Salinity (SMOS) (2009 - present) (Reul et al. 2012), NASA's Aquarius (2011-2015) (Lagerloef et al., 2013), and NASA's Soil Moisture Active Passive (SMAP) (2015-present) (Entekhabi et al., 70 2010), have provided continuous, near-global sea surface salinity (SSS) that resolves mesoscale features, tracks seasonal to interannual variability, and reveals the surface imprint of water cycle forcing (Vinogradova et al., 2019; Reul et al. 2020; Boutin et al. 2021). The emerging autonomous uncrewed surface vehicles (Saildrones, Wave Gliders, etc) in recent years have extended high-resolution salinity sampling to regions undersampled by Argo and ships, capturing submesoscale to mesoscale variability along extended trajectories (Patterson et al. 2025). Together, in situ and satellite salinity observations 75 capture the spatial structure linking evaporation-precipitation patterns to surface salinity (Fig. 1). The large-scale correspondence between E-P and SSS, evident in the subtropical salinity maxima and tropical fresh pools, confirms the fundamental water balance relationship while revealing important regional departures driven by circulation and mixing. These observations, together with satellite measurements of terrestrial water storage from GRACE and GRACE FO (Tapley et al., 2004) and river discharge from SWOT (Morrow et al., 2019), form a global water cycle observing framework 80 spanning the land-ocean-atmosphere continuum (Vinogradova et al. 2025).




Figure 1. Time-mean fields. (a) sea surface salinity (SSS) and (b) evaporation minus precipitation (E – P) averaged over 2012-2022. Adapted from Yu (2023).



These observations reveal salinity's scale-dependent dynamics and climate feedbacks. At basin scales, the amplification of mean patterns (fresh regions freshening, salty regions salinifying) confirms water cycle intensification (Durack and Wijffels, 2010; Helm et al., 2010; Durack et al., 2012; Skliris et al., 2016; Cheng et al., 2020), extending to semi-enclosed basins where evaporative loss drives coordinated salinity and bottom pressure changes (Lehmann et al. 2022; Liu et al., 2025).

90 Basin-scale synchronization between subtropical maxima and tropical minima demonstrates coupling through circulation rather than local forcing alone (Hasson et al. 2018; Yu, 2023), while subsurface extremes reveal ventilation pathways throughout the water column (Skliris et al., 2014; Zika et al., 2015). Beyond recording water cycle change, salinity drives critical climate feedbacks: upper-ocean stratification changes modulate heat uptake and surface warming (Zika et al., 2018; Liu et al., 2023; Vogt et al., 2025), near-surface salinity anomalies predict continental precipitation (Li et al., 2016; Rathore

95 et al. 2021), barrier layers intensify tropical cyclones (Balaguru et al., 2020), and high-latitude freshening weakens the Atlantic meridional overturning circulation (Caesar et al., 2018), with ecosystem and biogeochemical consequences across all scales (Fournier et al. 2023; Röthig et al. 2023).

Understanding these feedbacks matters for climate prediction and projection. Inadequate representation of salinity feedbacks could bias regional climate projections, while salinity's potential for nonlinear responses (Rahmstorf et al., 2005; Weijer et al., 2019) raises questions about model's capability to capture climate trajectories. Conversely, salinity observations may provide predictive information for seasonal-decadal forecasting (Qu et al. 2014; Zhu et al. 2014; Hackert et al. 2020). Quantifying salinity's scale-dependent roles has direct consequences for interpreting observed climate variations, constraining model biases, and projecting regional climate change.

100

This review synthesizes two decades of observations into a regime-dependent framework for ocean salinity behavior. Section 2 reviews the observational foundation from satellites, Argo, and historical archives that enables systematic characterization of salinity variability. Section 3 examines where and when salinity records water-cycle change: basin-scale and multi-decadal patterns where forcing dominates, mesoscale and seasonal-to-interannual redistribution where circulation shapes pathways, and submesoscale and synoptic dynamics where salinity actively shapes mixing and stratification. Section 4 develops a unifying framework showing how competition among forcing, advection, and mixing timescales determines these regime transitions, with implications for predictability and future projections. Section 5 identifies observational priorities at regime boundaries, particularly sustained global sea-surface salinity at O(10 km) resolution, to constrain how water-cycle intensification couples with circulation change under warming.

105

110

2 Observational foundations: resolving scale-dependent salinity dynamics


Observing salinity's scale-dependent dynamics requires capturing the spatial and temporal scales where forcing, advection, and mixing compete. Basin-scale patterns demand decadal stability and global coverage to detect water cycle trends. Mesoscale circulation pathways require spatial continuity to resolve how anomalies propagate and organize. Submesoscale

115



120 dynamics require high-frequency temporal sampling to capture processes modifying stratification and mixing. No single observing platform spans basin-to-submesoscale spatial resolution (1000 km to 1 km) while maintaining decadal-to-synoptic temporal coverage (decades to hours), necessitating integration across complementary systems designed for specific scale regimes.

125 Three observational advances since 2000 have enabled this integration (Figure 2). The Argo array provides quasi-synoptic three-dimensional sampling of the upper 2000 m with approximately 3° spacing and 10-day repeat cycles (Roemmich et al., 2009; Riser et al., 2016). Satellite L-band radiometry missions (SMOS, Aquarius, SMAP) deliver continuous surface salinity fields at approximately 100 km spatial and weekly temporal resolution (Boutin et al. 2018; Vinogradova et al., 2019; Reul et al. 2020). Autonomous surface vehicles and gliders resolve submesoscale variability through intensive targeted deployments achieving kilometer spatial and hourly temporal sampling (Patterson et al., 2025). GRACE/GRACE-FO and SWOT provide basin-integrated ocean mass variations and terrestrial water storage changes that constrain regional freshwater budgets (Tapley et al., 2004; Morrow et al., 2019; Vinogradova et al., 2025).



130 **Figure 2. Timeline of major ocean observing systems for water cycle research (2000–2025).** The Argo profiling float array (hatched: pilot phase 2000–2004; solid: near-global coverage 2004–present) provides continuous subsurface temperature and salinity measurements to 2000 m depth. Satellite missions SMOS (2009–present), Aquarius (2011–2015), and SMAP (2015–present) measure sea surface salinity from L-band radiometry at ~40–60 km resolution. Uncrewed surface vehicles (USVs; 2017–present) capture submesoscale (1–10 km) surface variability in temperature and salinity.
135 GRACE/GRACE-FO (2002–present) constrains basin-integrated ocean mass changes that reflect E-P-R (evaporation minus precipitation minus runoff) net fluxes. SWOT (2022–present) observes sea surface height at submesoscale (~15 km) resolution and monitors terrestrial surface water storage and river discharge. These complementary systems span spatial



scales from submesoscale to basin and enable integrated characterization of ocean salinity variability and water cycle fluxes. Mission images courtesy of NASA/ESA/NOAA; USVs image from Patterson et al. (2025).

140

These systems address distinct observational requirements across scale regimes. Argo vertical profiles distinguish atmospheric forcing from ocean circulation. Forcing-driven anomalies remain confined to the mixed layer, while circulation-driven anomalies extend along isopycnals or track subsurface density structure. Isopycnal analysis isolates water-mass changes from vertical displacement of the density field (Bindoff & McDougall, 1994), enabling attribution of surface salinity trends to water cycle changes rather than circulation shifts (Qu et al. 2016; Zika et al., 2018). Satellite L-band radiometers (SMOS, Aquarius, SMAP) resolve the horizontal spatial coherence that sparse in situ profiling cannot capture, revealing mesoscale fronts, eddy-driven stirring, and major river plumes as organized circulation features rather than isolated point anomalies (Grotsky et al., 2012; Lee et al. 2012; Reul et al., 2014; Yu 2015; Fournier et al. 2017; Melnichenko et al., 2017). Combined with altimetric geostrophic currents, satellite-derived winds, and E-P fields, satellite SSS enables regional freshwater budgets that partition surface salinity tendencies into local atmospheric forcing versus remote ocean advection (Vinogradova & Ponte, 2013; Dong et al. 2014; Yu 2023). Autonomous platforms capture submesoscale features at their native scales, including rainfall-generated fresh lenses, sharp frontal salinity gradients, and barrier layers evolving at kilometers and hours (Drushka et al., 2019). Glider profiling and SWOT sea surface height observations reveal whether submesoscale salinity features remain surface-trapped or couple vertically to interior dynamics (Morrow et al., 2019; du Plessis et al., 2022).

Observational coverage remains fundamentally asymmetric across scale regimes. Basin-scale salinity patterns and multi-decadal trends are well constrained by Argo subsurface sampling and satellite surface coverage combined, enabling robust detection of pattern amplification and separation of externally forced trends from internal climate variability. Mesoscale surface salinity structure benefits from satellite spatial continuity but lacks corresponding subsurface resolution; reconstructing three-dimensional circulation pathways and quantifying subduction rates requires vertical sampling substantially denser than Argo's nominal 3° spacing. Submesoscale salinity features evolving at kilometer spatial scales and synoptic temporal scales are systematically undersampled by all sustained observing systems, captured only during intensive field campaigns with autonomous platforms. This observational asymmetry reflects fundamental technological and resource constraints rather than design limitations: simultaneous achievement of global basin coverage, mesoscale spatial resolution, and submesoscale temporal resolution exceeds current observing system capacity. Section 3 examines salinity behavior across these observationally defined regimes.

3 Where salinity indicates water-cycle changes: timescale-dependent forcing control



Salinity acts as a water-cycle indicator when ocean processes integrate freshwater forcing faster than they redistribute it. The governing equation for mixed-layer salinity evolution is:

$$\frac{\partial S}{\partial t} = \underbrace{-\mathbf{u}_h \cdot (\nabla_h S)}_{\text{horizontal advection}} - \underbrace{w \frac{\partial S}{\partial z}}_{\text{vertical advection}} + \underbrace{\nabla_h \cdot (K_h \nabla_h S)}_{\text{horizontal mixing}} + \underbrace{\frac{\partial}{\partial z} \left(K_v \frac{\partial S}{\partial z} \right)}_{\text{vertical mixing}} + \underbrace{\frac{S_0}{h} (E - P - R)}_{\text{surface freshwater forcing (FWF)}}$$

(1)

where \mathbf{u}_h and w are horizontal and vertical velocities, K_h and K_v are mixing coefficients, and $S_0/h (E-P-R)$ denotes the effect of evaporation (E), precipitation (P), and runoff (R) on mixed-layer salinity. The terms on the right-hand-side represent horizontal advection, vertical advection, horizontal mixing, vertical mixing, and surface freshwater forcing (FWF), respectively. The rain gauge approximation emerges when advection and mixing terms become small relative to forcing, reducing Eq.(1) to $\partial S/\partial t \approx S_0/h (E-P-R)$. Whether this holds depends critically on timescale: at what temporal scales does forcing accumulate faster than circulation redistributes it? Figure 3 addresses this question at seasonal and interannual timescales by correlating observed $\partial S/\partial t$ with $E-P$ at each ocean location, showing forcing controls salinity tendency across approximately 30% of ocean area.

3.1 Seasonal-interannual forcing dominance

Figure 3 shows where forcing controls salinity variability versus where ocean circulation redistributes it. Forcing-dominated regions exhibit statistically significant positive correlations ($r > 0.50$ seasonal, $r > 0.14$ interannual; $p < 0.1$) between salinity tendency and local $E-P$. Circulation-dominated regions show weak, non-significant correlations ($p > 0.1$), indicating advection and mixing redistribute freshwater before forcing signatures accumulate. Significant negative correlations ($r < -0.50$ seasonal, $r < -0.14$ interannual; $p < 0.1$) occur where circulation processes oppose surface forcing.

Forcing-dominated regions (33% of ocean area) concentrate in subtropical gyre interiors (20-40°N/S), semi-enclosed seas, and monsoon regions. Subtropical gyres maintain strong correlations ($r = 0.6-0.7$) at seasonal timescales and moderate correlations ($r = 0.3-0.5$) at interannual timescales, where weak advection (1-3 cm/s) (Yu 2023) allows $E-P$ signals to accumulate. Semi-enclosed seas exhibit strongest forcing control ($r > 0.7$ seasonal) because geometric constraints limit advective escape. Mediterranean evaporation (~10 cm/mon; Fig.1b) creates 0.3-0.5 psu seasonal amplitude correlating at $r = 0.85$ with $E-P$ (Skiriris et al., 2018). Monsoon regions show timescale-dependent behavior: Bay of Bengal exhibits strong seasonal correlation ($r = 0.5-0.7$) but weak interannual correlation (not significant). Monsoon precipitation (~22 cm/mon during June-September; Rao & Sivakumar 2003) creates 2-4 psu freshening on 3-month timescales shorter than lateral spreading timescales (6-9 months), enabling local accumulation. At interannual timescales, ENSO-driven current anomalies introduce variance comparable to precipitation anomalies, degrading correlation (Delcroix & Hénin, 1991; Hasson et al. 2014).




Figure 3. Correlation between $E-P$ and $\delta S/\delta t$ at seasonal and interannual timescales (2012-2022). Spatial distribution of correlation coefficients for (a) seasonal variability (demeaned) and (b) interannual variability (detrended and deseasonalized). (c) Percentage of ocean area showing statistically significant positive correlations ($p < 0.1$, orange), significant negative correlations ($p < 0.1$, blue), and weak, non-significant correlations ($p > 0.1$, green). Significant positive correlations occupy 33% of ocean area at both timescales, while weak correlations dominate (55% seasonal, 66% interannual). Correlation thresholds for significance ($p < 0.1$) are 0.50 for seasonal and 0.14 for interannual variability. Adapted from Yu et al. (2021).

205

Circulation-dominated regions (55-66%) include western boundary currents, equatorial upwelling zones, and subpolar gyres. Western boundary currents exhibit weak correlation because strong advection redistributes freshwater faster than forcing accumulates (Hogg & Johns 1995). Correlation degrades from subtropical gyre interiors toward western boundaries where Gulf Stream and Kuroshio advection dominates. Equatorial upwelling zones exhibit significant negative correlation ($r = -0.4$ to -0.6 seasonal; 5-10% of ocean area) where upwelling brings high-salinity subsurface water to the surface during precipitation seasons, notably in eastern equatorial Pacific (Maes et al. 2014). Subpolar gyres and Southern Ocean show


210



weak correlation despite large precipitation because strong currents, deep winter mixing (200-800 m) (de Boyer Montégut et al. 2004), and energetic mesoscale eddies redistribute freshwater (Müller et al. 2019) before forcing signatures accumulate.

3.2 Weak salinity pattern amplification at decadal timescales

215 Decadal-scale trends reveal contrasting behavior between atmospheric forcing and ocean salinity response. Analysis of
1993-2010 shows the ocean water cycle (E-P) pattern amplified by approximately 5%, with spatial correlation $r \approx 0.5$
between climatological E-P and observed trends (Vinogradova & Ponte, 2017), matching Clausius-Clapeyron expectations
for atmospheric moisture response to warming (Held and Soden, 2006). Surface salinity patterns, however, amplified by less
than 1% globally, with near-zero spatial correlation between climatological patterns and observed trends (Fig. 4). Ocean
220 basins show scattered responses: some amplify climatological patterns while others reverse them. This weak salinity pattern
amplification, despite coherent atmospheric forcing, indicates that ocean processes operating on decadal timescales prevent
surface salinity from tracking atmospheric forcing coherently.



225 **Figure 4. Pattern amplification in ocean water cycle versus surface salinity over 1993-2010.** (a,b) Regions where climatological patterns amplified (orange: wet regions got wetter and dry regions got drier for freshwater flux; fresh regions got fresher and salty regions got saltier for salinity) versus weakened (gray). (c,d) Strength of pattern amplification: x-axis



shows anomalies (relative to global average) in zonal ocean basin averages of climatological mean freshwater flux (c) or surface salinity (d); y-axis shows corresponding 1993-2010 changes. Colors denote ocean basins. Black line shows linear regression; slope gives total pattern amplification. Ocean water cycle amplified by 5% globally (c), but surface salinity patterns amplified by less than 1% globally (d). Adapted from Vinogradova & Ponte (2017).

Regional analysis reveals the physical basis for weak global pattern amplification. Pacific basin-average salinity decreased over 1970-2002, exceeding internal variability estimates and showing detectable anthropogenic influence, while Atlantic basin-average trends remained within internal variability range (Terray et al., 2012). This basin-scale contrast reflects differences in how circulation and natural variability compete with forcing accumulation. North Atlantic salinity over 1993-2012 exhibits large multiyear variability driven by NAO modulation of the North Atlantic Current, with circulation-driven anomalies dominating over surface freshwater forcing (Stendardo et al., 2016). Strong western boundary currents and active decadal modes redistribute North Atlantic freshwater on timescales comparable to 10-20 year forcing accumulation. The Pacific shows clearer forced trends during 2005-2015 (basin-average 2.2×10^{-3} psu/year) following earlier freshening in 1994-2005 (Li et al., 2019), suggesting forced signals can emerge where circulation redistribution operates more slowly.

Salinity budgets quantify the circulation-forcing competition. Pacific upper 200m analysis for 2005-2015 shows surface freshwater flux and ocean advection produce opposite effects, each with magnitudes approximately twice the net salinity tendency (Li et al., 2019). This opposing relationship indicates active redistribution rather than passive forcing integration. Where precipitation dominates, advection exports excess freshwater; where evaporation dominates, advection imports freshwater from elsewhere. Horizontal advection and diffusion control subtropical redistribution while vertical diffusion and entrainment control tropical and high-latitude redistribution (Lyu et al., 2025), creating regionally varying pathways that prevent globally coherent response to coherent forcing. Despite these complications, salinity exhibits higher signal-to-noise ratios than atmospheric variables, as detection requires fewer than three ensemble members for SSS versus substantially more for precipitation or E-P (Terray et al., 2012), because salinity integrates high-variance atmospheric forcing over time while spreading anomalies spatially.

Natural decadal variability operates on timescales comparable to 10-30 year analysis periods. The Interdecadal Pacific Oscillation shift from positive to negative phase around 1998-1999 contributes approximately 30% of Pacific salinity variance, exceeding 60% in equatorial regions (Vinogradova & Ponte, 2017). NAO variability similarly drives North Atlantic salinity through circulation changes independent of local forcing (Stendardo et al., 2016). At decadal timescales, forcing accumulation, circulation redistribution, and natural variability all operate on comparable timescales, preventing any single process from dominating. The result is coherent atmospheric forcing (+5% pattern amplification) producing weak



260 salinity response (<1% pattern amplification), showing that 10-30 year periods are insufficient for forcing signals to
overwhelm circulation redistribution and natural variability. Basin differences arise where local conditions favor one
process. Pacific shows partial forcing dominance where redistribution is slower, Atlantic shows variability dominance where
energetic currents operate faster. Longer integration periods are required for forcing to accumulate decisively beyond
redistribution timescales, as examined in Section 3.3 for multi-decadal scales.

3.3 Multi-decadal pattern emergence

265 Over multi-decadal timescales (50+ years), the influence of freshwater forcing on ocean salinity becomes clearly detectable,
although these patterns remain obscure at shorter timescales (Figs.3-4). Spatial correlation between 1950-2000 salinity trends
and climatological SSS patterns reaches $r = 0.7-0.8$, indicating fresh regions systematically freshened while salty regions
salinified over 50+ year periods (Durack and Wijffels, 2010; Durack et al., 2012). This pattern correlation contrasts sharply
with near-zero values at decadal timescales (Fig.4b;d), showing that forcing accumulation over 50+ years produces spatially
270 coherent salinity response despite circulation redistribution. The 145-year record of ocean salinity measurements, obtained
from HMS Challenger and SMS Gazelle expeditions (1870s) through modern observations, demonstrates that pattern
amplification has been detectable since the early industrial era, but with marked acceleration. Rates increased from $\sim 0.166 \text{ g kg}^{-1} \text{ century}^{-1}$ (1870s-1950s) to $\sim 0.306 \text{ g kg}^{-1} \text{ century}^{-1}$ (1950s-2010s), a $54 \pm 10\%$ acceleration (Gould and Cunningham,
2021). Spatial correlation between regional salinity changes across these periods ($r = 0.64$) indicates persistent forcing-
275 driven patterns since the 1870s, though with non-linear intensification. Recent decades show further acceleration, with post-
1991 rates nearly doubling those of 1960-1990 (Cheng et al., 2020; Douville and Cheng, 2024).

Regional patterns show quantitatively consistent amplification across independent analyses at 4-8% per °C (Durack and
Wijffels, 2010; Helm et al., 2010; Skliris et al., 2016), in line with Clausius-Clapeyron predictions for atmospheric moisture
response to warming (Held and Soden, 2006). Tropical and subpolar fresh regions (climatological $P > E$) freshened 0.2-0.4
280 psu over 1950-2000, while subtropical salinity maxima (climatological $E > P$ regions) increased 0.1-0.3 psu (Curry et al.
2003; Durack and Wijffels, 2010). However, pattern amplification exhibits marked asymmetry: freshening in low-salinity
regions proves far more robust across ocean basins with consistent spatial patterns and strong emergence from natural
variability, whereas salinification in subtropical gyres displays greater sensitivity to region definition and weaker signal-to-
noise ratios (Cheng et al., 2020; Douville and Cheng, 2024). Semi-enclosed basins provide particularly clear evidence where
285 geometric constraints enhance forcing signals. For instance, Mediterranean salinification (0.10-0.15 psu) tracks 10-15%
evaporation increases within observational uncertainty (Skliris et al., 2018), while Baltic freshening (0.10-0.20 psu) is
consistent with increased precipitation and runoff (Meier et al., 2006; Lehmann et al., 2022).

Attribution studies identify the different physical processes driving the observed pattern amplification. Observed multi-
decadal patterns have less than 5% probability of arising from internal variability alone when major climate modes are



290 removed, with approximately two-thirds of variance attributable to external forcing (Pierce et al., 2012; Terray et al., 2012).
CMIP5 simulations show the emergence of anthropogenic influence. Historical runs incorporating all forcings reproduce
observed spatial patterns and amplification magnitude, whereas natural-forcing-only simulations show no significant trends
(Cheng et al., 2020). Mechanistic decomposition using ocean model experiments reveals three contributing processes (Zika
et al., 2018): direct surface freshwater flux changes from water cycle intensification, ice mass loss contributions (relatively
295 minor), and ocean warming effects on stratification. Ocean warming proves particularly important because warming-induced
stratification inhibits vertical mixing, effectively preserving and amplifying surface salinity patterns. Warming explains
approximately half of surface salinity pattern changes from 1957-2016, with water cycle intensification of $3.6 \pm 2.1\%$ per
degree Celsius accounting for the remainder. These changes exceed 2σ natural variability ($>95\%$ confidence), showing
human influence on ocean salinity patterns since 1960.

300 The physical mechanisms enabling multi-decadal pattern emergence operate through timescale separation and stratification
enhancement rather than circulation suppression. Surface $E-P$ anomalies create near-surface salinity changes that
subsequently subduct along isopycnals into low-latitude subsurface layers via subtropical gyre ventilation (Durack et al.,
2012), advect poleward and downward through intermediate water formation driving broad 300-2000m freshening (Helm et
al., 2010), and interact with ocean warming to enhance high-latitude stratification and reduce vertical mixing that would
305 otherwise erode surface salinity contrasts (Zika et al., 2018). At multi-decadal scales, circulation contributions remain
important (Jarugula et al. 2025). The Atlantic-Pacific salinity contrast increased $5.9 \pm 0.6\%$ over 1965-2020, yet direct E-P-R
forcing explains less than half this change, with the remainder from circulation adjustments including thermocline heaving,
cross-basin moisture transport, and gyre intensification (Singh et al., 2016; Friedman et al., 2017; Zika et al., 2021; Lu et al.,
2024). What distinguishes multi-decadal from shorter timescales is not that circulation stops, but that forcing accumulates
310 over 50+ years while basin-scale redistribution operates on 5-20 year timescales. Because forcing persists much longer than
redistribution and mixing processes, it can build up a coherent signal despite ongoing circulation. Forcing controls only 30%
of ocean area at seasonal-interannual scales (Fig.3) and produces less than 1% pattern amplification at decadal scales (Fig.4),
but generates strong pattern correlations ($r = 0.7-0.8$) and 54% acceleration post-1950s at multi-decadal scales. This
progression shows that forcing fingerprints emerge when forcing timescales far exceed redistribution timescales, consistent
315 with the rain gauge paradigm over sufficiently long periods and revealing ocean warming as a critical amplifying
mechanism.

4 When circulation controls salinity: timescale competition and regime transitions

Section 3 documented a striking timescale dependence in how salinity responds to freshwater forcing. Forcing controls
salinity tendency across $\sim 30\%$ of ocean area at seasonal-interannual timescales (Fig. 3), yet produces $<1\%$ pattern
320 amplification at decadal scales despite 5% atmospheric forcing amplification (Fig. 4). At multi-decadal scales, however,



forcing fingerprints emerge with strong spatial pattern correlations ($r = 0.7-0.8$). What physical mechanisms explain these regime transitions?

325 The question is not whether forcing or circulation "controls" salinity, since both always operate. Rather, we ask: Does the spatial distribution of salinity anomalies match the spatial distribution of forcing, or does it trace the pathways by which circulation redistributes remotely forced anomalies? The answer lies in competition among four characteristic timescales governing Eq. (1). Table 1 defines these timescales: the forcing timescale τ_f measures how fast freshwater fluxes change, the advection timescale $\tau_{adv} = L/U$ measures how fast currents redistribute anomalies, the horizontal mixing timescale $\tau_{hmix} = L^2/K_h$ measures how fast stirring erodes gradients, and the vertical mixing timescale $\tau_{vmix} = h^2/K_v$ measures how fast vertical processes modify stratification.

330 Three primary competitions emerge from these timescales. First, τ_f versus τ_{adv} determines whether forcing accumulates coherently (rain gauge) or anomalies transport far from formation regions (passive tracer). Second, τ_{vmix} versus τ_{adv} controls whether subsurface waters preserve formation-era memory or respond to contemporary forcing. Third, convergence of all four timescales at submesoscales creates a dynamical regime where salinity actively shapes density structure. The horizontal mixing timescale τ_{hmix} (decades to centuries for basin scales) is typically much longer than τ_f , τ_{adv} , and τ_{vmix} , so while it
 335 determines whether anomalies maintain coherence during transport, it rarely competes directly except at submesoscales where all timescales converge.

Table 1. Fundamental timescales governing salinity evolution


Symbol	Timescale	Definition	Typical range	Physical meaning
τ_f	Forcing	Duration of $E-P-R$ events or trend	Days (storms), weeks–months (monsoon/ENSO), decades (climate trends)	Persistence of freshwater input
τ_{adv}	Advection	L/U	Weeks (eddies) to years (gyres)	Time for currents to redistribute anomalies
τ_{hmix}	Horizontal mixing	L^2/K_h	Months (mesoscale) to centuries (basin)	Time for stirring to smooth lateral gradients
τ_{vmix}	Vertical Mixing	h^2/K_v	Days (convection) to centuries (stratified)	Time for vertical processes to modify vertical structure

4.1 Amplitude-phase separation and horizontal pathways

340 **4.1.1 Amplitudes agree, phases differ**



Figure 5 reveals a fundamental separation in how salinity responds to freshwater forcing. Seasonal amplitude patterns are strikingly coherent across atmospheric forcing $E-P$ (Fig. 5a), mixed-layer forcing $FWF = S_0(E-P)/h$ (Fig. 5c), and observed salinity tendency $\partial S/\partial t$ (Fig. 5e), with amplitudes exceeding 0.2 psu/month in tropical convergence zones and 0,1 psu/month in subtropical net evaporation maxima. This amplitude coherence indicates that forcing magnitude controls salinity variance.



345

Figure 5. Amplitude-phase asymmetry in seasonal salinity variability demonstrates forcing sets variance amplitude but circulation determines spatial-temporal distribution. Harmonic analysis of (a,b) atmospheric freshwater forcing $E-P$, (c,d) freshwater flux $FWF = S_0(E-P)/h$, and (e,f) observed salinity tendency $\partial S/\partial t$ from satellites (2010-2020). Left panels show seasonal amplitude; right panels show phase (month of annual maximum). Adapted from Yu et al. (2021).

350

However, phase maps tell a different story. Across most ocean regions, $\partial S/\partial t$ phase (Fig. 5f) leads $E-P$ phase (Fig. 5b) by 1-3 months. The eastern tropical Pacific exemplifies this behavior. Local $E-P$ reaches maximum in March-April when the



ITCZ migrates southward (Fig. 5b), yet $\partial S/\partial t$ peaks earlier in January-March. Similarly, $\partial S/\partial t$ in the subtropical North Pacific leads $E-P$ by approximately 2 months. This phase lead is stronger evidence for circulation control than a lag would be. A phase lag might indicate slow local response to forcing, but a phase lead requires non-local processes. Salinity cannot respond to local forcing before that forcing occurs. The phase lead indicates that salinity changes reflect ocean processes, either through advection of anomalies from upstream regions where forcing peaked earlier in the seasonal cycle, or upwelling of subsurface water with different seasonal history, or remote forcing effects transported by circulation. Forcing creates the variance, but circulation determines when and where that variance manifests.

The progressive phase shift from $E-P \rightarrow FWF \rightarrow \partial S/\partial t$ in Figure 5b,d,f quantifies how ocean processes successively alter salinity timing. The $E-P$ to FWF phase shift reflects mixed-layer depth modulation, where deeper winter mixed layers dilute the same freshwater flux over larger volumes. The FWF to $\partial S/\partial t$ phase shift isolates pure circulation effects operating through horizontal advection, vertical exchanges, and mixing. Subtropical gyre interiors where all three phases align represent forcing-dominated regimes. Upwelling zones, boundary currents, and monsoon regions showing large phase differences between FWF and $\partial S/\partial t$ represent circulation-dominated regimes where advection competes directly with forcing accumulation.

4.1.2 Multi-decadal versus seasonal regimes

The timescale framework from Table 1 explains why forcing control strengthens from seasonal to multi-decadal timescales. At seasonal-interannual timescales, τ_f (seasonal forcing persistence, ~ 3 -12 months; ENSO, ~ 2 -5 years) proves comparable to τ_{adv} for surface redistribution (mesoscale eddies, \sim weeks to months; gyre-scale transport, \sim months to years). When $\tau_f/\tau_{adv} \lesssim 1$, forcing creates variance but advection redistributes it before coherent spatial patterns emerge. Figure 3 shows this regime where temporal correlation between $\partial S/\partial t$ and $E-P$ remains weak (< 0.4) across 70% of ocean area. The forcing signal exists but advection scrambles it spatially and temporally, as shown by the phase differences in Figure 5.

At multi-decadal timescales, the ratio shifts. Forcing accumulates over $\tau_f \sim 50$ -100 years while basin-scale redistribution through gyre circulation, subtropical cells, and meridional overturning operates on $\tau_{adv} \sim 5$ -20 years (McCreary and Lu, 1994; Gu and Philander, 1997; Qu et al., 2013; Fine et al., 2017). This gives $\tau_f/\tau_{adv} \approx 5$ -10. When forcing persists much longer than redistribution timescales, circulation repeatedly samples the same forcing pattern. Subtropical gyre water recirculates within the high-evaporation subtropical band every 5–10 years, experiencing persistent $E > P$ forcing with each circulation cycle. Over 50–100 years, this repeated exposure to the same forcing regime accumulates coherent basin-scale salinity trends that spatially match the forcing pattern. The spatial correlation emerges not because circulation stops redistributing anomalies, but because circulation pathways remain within forcing regimes long enough for forcing to accumulate systematically. The spatial pattern correlation between multi-decadal salinity trends and climatological $E-P$ reaches $r = 0.7$ -0.8 (Durack and Wijffels, 2010; Skliris et al., 2016; Cheng et al., 2020). Fresh regions systematically freshen,



385 salty regions systematically salinify. Circulation still redistributes anomalies actively, but forcing accumulates persistently, imposing its spatial fingerprint.

Figure 4 captures the critical transition at decadal timescales (10-30 years), where $\tau_f/\tau_{adv} \approx 1-2$. Atmospheric forcing amplifies coherently (5% pattern intensification) but ocean salinity responds incoherently (<1% pattern amplification). This asymmetric response reflects the fact that forcing and redistribution operate on comparable timescales. Basin-scale adjustment processes including gyre spinup, thermocline heaving, and ENSO-driven reorganization all operate over 5-20 years (Anderson & Gill 1975; Lysne & Deser 2002; Cessi & Otheguy 2003; McPhaden et al. 2006), meaning τ_{adv} and τ_f are comparable at decadal scales and neither process dominates. The atmosphere intensifies its hydrological cycle, but ocean circulation redistributes anomalies as soon as forcing creates patterns.

The progression from 30% spatial control (seasonal, $\tau_f/\tau_{adv} \lesssim 1$) through <1% pattern amplification (decadal, $\tau_f/\tau_{adv} \approx 1-2$) to $r = 0.7-0.8$ pattern correlation (multi-decadal, $\tau_f/\tau_{adv} \approx 5-10$) quantifies how the ratio τ_f/τ_{adv} controls whether forcing or circulation dominates salinity evolution. The ocean-atmosphere system operates identically at all timescales, but forcing accumulation becomes progressively more effective at longer integration periods because circulation can only redistribute anomalies over finite timescales while forcing persists continuously.

4.1.3 River plumes as extreme cases

River plumes illustrate the full spectrum of τ_f/τ_{adv} regimes within a single system and clarify how the "amplitudes agree, phases differ" behavior in Section 4.1.1 emerges from the competition between local forcing and redistribution. River discharge varies on daily to weekly timescales while plume evolution spans from near-field accumulation (days) through mid-field redistribution (weeks) to far-field dispersal (months to years) (Grotsky et al. 2014; Horner-Devine et al., 2015; Reul et al., 2014; Fournier et al., 2023). Because τ_f is set by discharge variability while τ_{adv} increases systematically with distance from the mouth, a single plume spans $\tau_f/\tau_{adv} < 1$, ≈ 1 , and > 1 as an observable horizontal gradient. This spatial progression produces regime transitions analogous to the seasonal–multi-decadal progression in Section 4.1.2, but with τ_{adv} varying in space rather than τ_f varying in time. Table 2 compares these river plume regimes with their open-ocean analogs.

Table 2. River plume spatial progression as analog of open-ocean timescale regimes

Field	Distance from mouth	τ_f/τ_{adv}	River Plume Behavior	Open-Ocean Analog	Timescale Basis
Near-field	<200 km	$\approx 0.3-0.5$	Local accumulation; $\partial S/\partial t$ tracks discharge (zero lag)	Multi-decadal basin trends	$\tau_f \ll \tau_{adv}$ (near-field) vs $\tau_f \gg \tau_{adv}$ (multi-decadal): both are "rain gauge" at opposite extremes



Mid-field	200–1000 km	≈ 1	Amplitude-phase separation; discharge sets variance, currents control timing	Seasonal-interannual variability	$\tau_f \sim \tau_{adv}$: forcing and advection compete equally
Far-field	>1000 km	$\approx 10\text{--}50$	Passive tracer; 6–12 month lag, encodes pathway history	Mode water subduction; gyre circulation	$\tau_f \ll \tau_{adv}$ (far-field) or $\tau_{vmix} \gg \tau_{adv}$ (subsurface): anomalies decouple from local forcing

410

The near-field (<200 km from river mouth) shows a forcing-dominated regime with $\tau_f/\tau_{adv} \approx 0.3\text{--}0.5$, where daily discharge fluctuations ($\tau_f \sim$ days) occur while boundary-current export requires weeks ($\tau_{adv} \sim 2\text{--}3$ weeks). Here redistribution is too slow to compete with local accumulation, so salinity tracks discharge with minimal phase lag (Figure 5–style phase alignment). To leading order, a simple volume-averaged salt balance gives $\partial S/\partial t \approx -S_0 Q/(hA)$, where Q is the river freshwater volume flux into the control region ($\text{m}^3 \text{s}^{-1}$), A is the horizontal area over which that freshwater is distributed, and h is the effective plume thickness (so $V = hA$). Equivalently, defining an area-normalized runoff flux $R \equiv Q/A$ (m s^{-1}) yields $\partial S/\partial t \approx -S_0 R/h$ (Horner-Devine et al., 2015), making explicit that the tendency scales with freshwater input per unit mixed-layer volume. In this “forcing dominance by slow removal” limit, advection is present but subdominant in the tendency budget and does not set the phase, contrasting with the multi-decadal forcing dominance in Section 4.1.2, where forcing wins by persisting longer than redistribution.

The mid-field (200–1000 km) marks the transition where $\tau_f \sim \tau_{adv}$, and discharge variability and boundary current transport both operate on weekly timescales, producing amplitude–phase separation. Here, discharge still largely sets the magnitude of salinity variance, but timing increasingly reflects redistribution by boundary currents, eddies, and rings, so salinity can retain discharge-like amplitudes while exhibiting phase shifts and reduced coherence with contemporaneous forcing (Fournier et al., 2017). This mirrors Figure 5, where forcing imprints variance, but circulation determines when that variance appears at a given location.

The far-field (>1000 km) exhibits passive-tracer behavior with $\tau_f/\tau_{adv} \approx 10\text{--}50$, where weekly discharge fluctuations become fast compared to 6–12 month transit times. In this regime, present-day salinity no longer tracks contemporaneous discharge timing; it primarily encodes water-mass age, pathway, and the integrated history of earlier forcing that has been advected and mixed. The Amazon plume detected in the Caribbean ~ 2000 km from the river mouth exemplifies this through lagged discharge–salinity relationships and ring-mediated transport (Hellweger and Gordon, 2002; Salisbury et al., 2011).



435 Pathway coherence depends on $\tau_f/\tau_{\text{hmix}}$, where $\tau_{\text{hmix}} = L^2/K_h$ determines survival against lateral stirring. For the Amazon plume extending ~ 1500 km, $\tau_{\text{hmix}} \sim 70$ years far exceeds $\tau_{\text{adv}} \sim 6$ months, allowing coherent transport. Satellite SSS and multisensor plume mapping highlight this coherence while also showing that mesoscale stirring and ring interactions redistribute freshwater laterally without immediately erasing the large-scale signal (Reul et al., 2014; Fournier et al., 2017). Subtropical-to-tropical cells show similar persistence, where $\tau_{\text{adv}} \sim 3\text{--}5$ years $\ll \tau_{\text{hmix}} \sim$ centuries preserves formation signatures during equatorward transit (McCreary and Lu, 1994; Qu et al., 2013).

4.2 Surface-subsurface differences and vertical pathways

440 Section 4.1 examined how horizontal advection competes with forcing to create spatial and temporal patterns. The vertical dimension introduces a fundamentally different competition between vertical mixing timescale τ_{vmix} and horizontal advection timescale τ_{adv} . At the surface where τ_{vmix} is short (days for wind mixing, weeks for convection) (Kraus & Turner, 1967; Niiler 1975; Marshall & Schott, 1999), salinity responds quasi-instantaneously to $E\text{--}P\text{--}R$ forcing. Below the mixed layer where stratification suppresses vertical exchange, τ_{vmix} increases dramatically (decades to centuries for diffusive mixing through the permanent pycnocline) (Ledwell et al., 1993; Wunsch & Ferrari 2004), while τ_{adv} remains moderate (years for 445 subtropical cells, decades for gyre circulation) (Fernandez et al. 2015). This reversal in timescale ratios creates a vertical regime boundary where salinity transitions from recording contemporary forcing to preserving formation-era conditions.

4.2.1 Surface-subsurface differences in observed trends

450 Figure 6 reveals systematic vertical structure in multi-decadal salinity trends that cannot be explained by local forcing. In subtropical gyres ($20\text{--}40^\circ\text{N/S}$) across all ocean basins, maximum trends occur at subsurface mode-water depths (100-500 m), vertically offset from the surface where $E\text{--}P$ forcing acts (Durack and Wijffels, 2010; Cheng et al. 2020). The Pacific (Fig. 6a) shows subsurface trend maxima of 0.08–0.1 psu per 70 years centered at 200-300 m depth near 30°N and 30°S , while surface trends are slightly weaker and of opposite sign. The Atlantic (Fig. 6b) has even stronger subsurface intensification, with trends exceeding 0.16 psu per 70 years at 150-250 m depth in the South Atlantic subtropical gyre. The Indian Ocean (Fig. 6c) shows a comparable vertical structure despite different surface forcing patterns.

455 This surface-subsurface difference indicates that observed trends reflect subduction of surface-formed anomalies along isopycnal pathways rather than in situ response to local forcing. If trends resulted from direct forcing response throughout the water column, maximum trends would occur at the surface where $E\text{--}P$ forcing is strongest and decrease monotonically with depth as the forcing signal diffuses downward. Instead, subsurface maxima indicate that surface anomalies are actively transported to depth along sloping isopycnals faster than vertical mixing can homogenize them (McCreary and Lu 1994; Qu 460 et al. 2013; Fine et al. 2017). The subsurface ocean thus preserves a multi-decadal memory of surface forcing history, with



trend magnitude and depth distribution reflecting the integrated history of surface forcing variations and subduction pathways.




Figure 6. Subsurface salinity trends reveal circulation memory through surface-subsurface asymmetry. Zonally-averaged linear trends (1950–2019) in the upper 2000 m for (a) Pacific, (b) Atlantic, (c) Indian, and (d) global ocean. Black contours show climatological mean salinity (psu); colors show trends (psu per 70 years); stippling indicates trends not significant at 90% confidence. Y-axis scale changes at 500 m depth. In subtropical gyres (20–40°N/S), maximum trends occur at subsurface mode-water depths (100–500 m), vertically offset from surface forcing. Data from EN4 analysis (Good et al., 2013).

470

4.2.2 Timescale competition: τ_{vmix} versus τ_{adv}

The surface-subsurface differences arise from competing vertical mixing and horizontal advection timescales. At the surface, $\tau_{\text{vmix}} \sim$ days to weeks (set by wind-driven turbulence and convective overturning) proves much shorter than $\tau_{\text{adv}} \sim$ weeks to months (set by mesoscale eddies and boundary currents redistributing surface anomalies horizontally) (Niiler, 1975;



475 Marshall & Schott, 1999; Abernathy & Marshall, 2013). This gives $\tau_{\text{mix}}/\tau_{\text{adv}} \ll 1$, allowing forcing signals to mix vertically through the mixed layer before horizontal advection can redistribute them laterally.

Below the seasonal thermocline, the ratio reverses. Stratification suppresses vertical exchange, increasing τ_{mix} to decades or centuries for diffusive mixing across the permanent pycnocline (Ledwell et al., 1993; Wunsch & Ferrari, 2004). Meanwhile, τ_{adv} for horizontal transport along isopycnals remains moderate at years to decades for subtropical cells (McCreary and Lu, 480 1994; Johnson and McPhaden, 1999) and decades for gyre-scale circulation (Qu et al. 2013; Fernandez et al. 2015; Fine et al., 2017). This gives $\tau_{\text{mix}}/\tau_{\text{adv}} \gg 1$ in the permanent thermocline, meaning horizontal advection redistributes anomalies along density surfaces much faster than vertical mixing can erase them. Water parcels retain their formation-region salinity signatures for years to decades as they travel along isopycnal pathways, creating the subsurface "memory" visible in Figure 6.

485 The regime boundary occurs where $\tau_{\text{mix}} \sim \tau_{\text{adv}}$, typically near the base of the winter mixed layer (50-200 m depth depending on latitude and basin). Above this depth, vertical processes dominate and salinity reflects contemporary forcing (Kraus & Turner, 1967; Niiler 1975). Below this depth, horizontal advection along isopycnals dominates and salinity reflects formation-era conditions from years to decades earlier (McCreary and Lu, 1994; Johnson and McPhaden, 1999). The exact depth of this transition varies seasonally (deepening in winter when convection penetrates deeper) and geographically 490 (deeper in subtropical mode water formation regions where vigorous winter convection creates thick homogeneous layers) (Hanawa & Talley, 2001; de Boyer Montégut et al., 2004).

4.2.3 Subduction and subsurface memory

Subduction physically accomplishes the $\tau_{\text{mix}} \gg \tau_{\text{adv}}$ regime described in Section 4.2.2 through a seasonal cycle of ventilation and capping. Each winter, convection homogenizes surface waters, creating mixed layers whose salinity reflects integrated 495 winter forcing. When spring arrives and the ocean restratifies, a seasonal thermocline reforms above these winter-formed waters, sealing them from further surface contact. These capped layers then spread horizontally along constant density surfaces into the ocean interior (Hanawa and Talley 2001).

In the North Atlantic, high evaporation creates surface salinity maxima that subduct to form Subtropical Underwater (STUW), a subsurface salinity maximum at 100-200m depth 20-40°N. The strong salinification trends (0.16-0.2 psu per 70 500 years) in this layer is visible in Fig.6b, reflecting both intensifying surface evaporation and poleward shifts in ventilation regions (Yu et al. 2018; Liu et al. 2019). The South Pacific operates similarly through subtropical mode water formation near 30-40°S, producing the subsurface maximum at 200-300m in Figure 6a. In both cases, the interior accumulates a long-term forcing signal while the surface experiences continual seasonal cycling (Marshall et al. 1993; Qiu and Huang 1995).



The persistence timescale follows from weak vertical exchange across the permanent thermocline. With diapycnal diffusivity
505 $K_v \sim 10^{-5} \text{ m}^2 \text{ s}^{-1}$ and thermocline thickness $h \sim 200\text{-}300 \text{ m}$, vertical mixing operates on $\tau_{\text{vmix}} = h^2/K_v \sim 50\text{-}100$ years (Ledwell
et al. 1993). Horizontal transport covers basin scales in $\tau_{\text{adv}} \sim 5\text{-}10$ years. The ratio $\tau_{\text{vmix}}/\tau_{\text{adv}} \sim 10$ means water parcels travel
thousands of kilometers before vertical exchange significantly modifies their properties. The seasonal thermocline acts as a
barrier, with rapid surface mixing (days) above and slow diffusion below.

This mechanism breaks down where stratification weakens. In subpolar regions, vigorous winter convection erodes the
510 permanent thermocline, reducing the vertical barrier. Near the equator, strong upwelling and tropical instability waves
enhance mixing to $K_v \sim 10^{-4} \text{ m}^2 \text{ s}^{-1}$, reducing τ_{vmix} to 5-10 years. When $\tau_{\text{vmix}} \sim \tau_{\text{adv}}$, vertical exchange modifies properties
during transit, explaining the nearly vertical trend structure in tropical regions visible in Figure 6.

4.3 Submesoscale transition: when all timescales converge

Sections 4.1 and 4.2 described regimes in which controlling timescales are cleanly ordered, so salinity behaves primarily as a
515 recorder. When $\tau_f \gg \tau_{\text{adv}}$, salinity integrates persistent freshwater forcing and large-scale patterns resemble the forcing
patterns (rain gauge behavior). When $\tau_f \ll \tau_{\text{adv}}$ or $\tau_{\text{vmix}} \gg \tau_{\text{adv}}$, salinity is exported and preserved along pathways, mapping
circulation and formation histories (passive tracer behavior).

At submesoscales $O(1\text{-}10 \text{ km})$ this ordering breaks down because the relevant processes operate on comparable hours-to-
520 days timescales. Mixed-layer instabilities, frontogenesis, and surface forcing all evolve fast enough that no single process
sets salinity before the others act (Boccaletti et al. 2007; Thomas and Ferrari 2008; McWilliams 2016). Salinity cannot
simply integrate forcing because forcing varies on the same timescale as advection redistributes anomalies. It cannot
passively inherit upstream properties because the upstream field is itself rapidly rearranged. Instead, salinity often becomes
dynamically consequential by sharpening horizontal buoyancy gradients and thereby strengthening ageostrophic secondary
circulations. These circulations can drive vertical velocities $w \sim O(10\text{-}100 \text{ m day}^{-1})$, ventilating the upper pycnocline far
525 more efficiently than mesoscale stirring at $w \sim O(1 \text{ m day}^{-1})$ (Mahadevan and Tandon 2006; McWilliams 2016).

High-resolution Saildrone observations during a 2019 North Atlantic crossing reveal this thermohaline control shift (Figure
7). At mesoscales exceeding 100 km, density gradients in the Gulf Stream and open ocean are predominantly temperature-
controlled with Turner angle $Tu > \pi/4$ and density ratio $R\rho > 1$. The continental shelf, however, shows salinity already
contributing significantly at these larger scales due to freshwater inputs and coastal fronts. Near 10 km the balance flips in all
530 three regimes and salinity variance dominates the density gradient budget, diagnosed by Tu dropping below $\pi/4$ and $R\rho < 1$
(Yu 2026). The transition clusters near the first baroclinic deformation radius ($R_d = 28 \text{ km}$), consistent with submesoscale
dynamics emerging as the flow approaches the breakdown of purely geostrophic adjustment (McWilliams 2016).



The sharpness of this transition is regime-dependent. In the Gulf Stream where thermal wind balance strongly dominates, Tu drops abruptly from greater than $5\pi/12$ to less than $\pi/4$ over just 10-20 km. On the continental shelf where salinity already plays a larger role, the transition spreads over 50+ km. This regime dependence emphasizes that 10 km marks where local nondimensional ratios (Rossby number, frontogenetic sharpening versus damping, mixing versus advection) cross order unity, not a fixed geometric constant (Callies and Ferrari 2013; McWilliams 2016).




Figure 7. Thermohaline regime transition at submesoscales. (a) Sea surface density gradient magnitude and (b) Turner angle (Tu) from satellite observations with 2019 North Atlantic Saildrone tracks overlaid (black lines). Tu less than 45° (blue) indicates salinity-dominated density; Tu greater than 45° (orange) indicates temperature-dominated density. (c) Median Turner angle (left axis) and density ratio $R\rho$ (right axis) versus wavelength computed from Saildrone measurements for three regimes: Gulf Stream winter (blue), open ocean summer (red), and continental shelf fall (green). Dashed line marks deformation radius ($R_d = 28 \text{ km}$); gray shading indicates submesoscale transition zone (5-15 km); horizontal line marks $Tu =$



545 $\pi/4$ threshold. All regimes shift from temperature-dominated ($Tu > \pi/4$) at mesoscales to salinity-dominated ($Tu \lesssim \pi/4$) at submesoscales. Transition sharpness varies with background conditions. Adapted from Yu (2026).

4.3.1 Why 10 km? Geometric and dynamical convergence

Three physical constraints converge near 10 km at mid-latitudes. First, the first baroclinic Rossby deformation radius $R_d =$
550 NH/f (where N is buoyancy frequency, H is ocean depth, and f is Coriolis parameter) equals approximately 10-30 km, setting
the scale below which ageostrophic motions and vertical exchange intensify (McWilliams 2016). Below R_d , horizontal flow
cannot maintain geostrophic balance, forcing vertical circulation. Second, frontogenesis compresses buoyancy gradients into
narrow widths approximately U/f (~ 1 -10 km for typical upper-ocean velocities $U \sim 0.1$ -0.5 m s^{-1} and Coriolis parameter $f \sim$
555 $5 \times 10^{-5} \text{ s}^{-1}$), where strain overwhelms diffusive smoothing (Thomas and Ferrari 2008). Third, the advective timescale $L/U \sim$
10 hours to 1 day at 10 km becomes comparable to the diurnal and storm forcing timescales, which modulate stratification
on the same window (Mahadevan et al. 2010).

This geometric convergence produces timescale convergence. At 10 km submesoscales, $\tau_{\text{adv}} = L/U \sim 1$ -10 days, $\tau_{\text{hmix}} = L^2/K_h$
 \sim days to weeks (for submesoscale stirring with $K_h \sim 10$ -100 $\text{m}^2 \text{ s}^{-1}$), $\tau_{\text{vmix}} = h^2/K_v \sim$ hours to days (for active mixing with
560 surface layer thickness $h \sim 20$ -50 m and enhanced $K_v \sim 10^{-3}$ - $10^{-2} \text{ m}^2 \text{ s}^{-1}$), and $\tau_f \sim$ hours to days (storm passage, diurnal
heating). When $\tau_f \sim \tau_{\text{adv}} \sim \tau_{\text{hmix}} \sim \tau_{\text{vmix}}$, all processes compete equally. Salinity evolution becomes the joint outcome of
forcing, stirring, and vertical exchange operating simultaneously rather than a hierarchy where one process dominates
(Boccaletti et al. 2007; Thomas and Ferrari 2008). This is fundamentally different from the regime separation in Sections 4.1
and 4.2 where one timescale dominated, and salinity could be understood through that dominance.

4.3.2 Differential damping: why salinity fronts persist

565 Salinity achieves comparable importance to temperature in density gradients at 10 km submesoscales even in regions where
temperature strongly dominates at mesoscales. The mechanism lies in differential damping of fronts. Frontogenesis sharpens
both temperature and salinity gradients through strain and confluence, but only temperature fronts experience strong negative
feedback from surface heat fluxes. Solar heating preferentially warms colder patches because they have lower heat content,
creating differential heating that erodes horizontal temperature gradients over days to weeks (Taylor and Ferrari 2010;
570 Mahadevan et al. 2010). Strong diurnal cycling and episodic mixing at fronts enhance this damping. The result is radiative
restoring that continuously weakens temperature fronts even as frontogenesis sharpens them.

Freshwater forcing provides no comparable damping mechanism. Precipitation does not preferentially target fresh patches,
and evaporation depends on atmospheric state (wind speed, humidity, air temperature) rather than on salinity itself. Salinity
fronts therefore persist and sharpen through frontogenesis without the radiative damping that weakens temperature fronts



575 (McWilliams 2016; Jaeger and Mahadevan 2018). Over the days-to-weeks timescales of submesoscale evolution ($\tau_{adv} \sim 1-10$ days, $\tau_{mix} \sim$ days to weeks), this differential damping allows salinity variance to accumulate while temperature variance degrades.

At scales below R_d where ageostrophic dynamics become strong, this differential damping shifts the thermohaline balance from temperature-dominated ($Tu > \pi/4$, $R\rho > 1$) at mesoscales to equal contribution ($Tu \sim \pi/4$) at submesoscales. This transition is dynamically consequential. Temperature-driven density fronts experience two-way coupling with the atmosphere through surface heat flux damping, which limits front intensity and vertical circulation strength. Salinity-driven density fronts lack this damping and can sustain sharper gradients. When both contribute equally to density at submesoscales, the undamped salinity component allows stronger frontogenesis and more intense ageostrophic secondary circulations than would occur under pure temperature control. The resulting vertical velocities reach $O(10-100 \text{ m day}^{-1})$, an order of magnitude stronger than mesoscale vertical exchange, ventilating the upper pycnocline and enhancing nutrient supply, carbon export, and mixed layer restratification. This mechanism operates most effectively in regions with strong near-surface salinity gradients such as river plume boundaries, marginal ice zones, and precipitation fronts (Yu 2026).

At scales below R_d , this differential damping shifts density control from temperature-dominated ($Tu > \pi/4$, $R\rho > 1$) at mesoscales to equal T-S contribution ($Tu \sim \pi/4$) at submesoscales (Ruddick, 1983; Rudnick & Ferrari 1999). Equal contribution is not merely shared control but a dynamical transition: mesoscale fronts are typically thermally controlled and remain tightly coupled to the atmosphere because surface heat fluxes (including radiative and turbulent components) and mixed-layer turbulence relax SST contrasts on $O(\text{days})$ timescales, limiting sustained thermal front sharpening (Taylor & Ferrari, 2010; Hausmann et al., 2017). By contrast, when submesoscale density fronts include a comparable haline contribution, the salinity component lacks an equally fast, state-dependent restoring and can therefore support sharper buoyancy gradients and stronger ageostrophic secondary circulations than temperature alone could maintain (Thomas & Ferrari, 2008; McWilliams, 2016). These overturnings commonly generate vertical velocity $w \sim O(10-100 \text{ m day}^{-1})$, far larger than mesoscale vertical exchange, and can rapidly ventilate the upper pycnocline, especially where near-surface salinity gradients are strong, such as river plumes, marginal ice zones/meltwater fronts, and precipitation-driven freshwater filaments (Mahadevan & Tandon, 2006; Boccaletti et al., 2007; Horner-Devine et al., 2015; Kozlov et al. 2020).

600 4.4 Synthesis: timescale ratios as regime boundaries

Across Sections 4.1–4.3, the same message keeps resurfacing: salinity behavior does not depend on absolute spatial or temporal scales (10 km versus 100 km, or days versus years); what matters is how forcing, advection, and mixing timescales are fast or slow relative to each other. In practice, three comparisons decide how to read a salinity signal. First, does freshwater forcing change slowly enough for anomalies to accumulate where they are made (τ_f/τ_{adv})? Second, once salinity anomalies subducted below the mixed layer, are they erased by vertical exchange or preserved as water-mass properties



(τ_{vmix}/τ_{adv})? Third, are we in a regime where the process hierarchy collapses because everything happens on similar timescales? Table 3 summarizes these boundaries and what they look like in data.

Table 3. Salinity regimes from timescale ratios

Regime	Controlling Ratio	Physical mechanism	Key signature
Rain Gauge	$\tau_f/\tau_{adv} \geq 5$	Forcing persists; circulation redistributes within forcing regimes	Pattern follows freshwater forcing ($r \approx 0.7-0.8$)
Transition	$\tau_f/\tau_{adv} \approx 1-3$	Forcing and advection both matter	Similar amplitudes, shifted phase ($\sim 1-3$ months)
Passive tracer	$\tau_{vmix}/\tau_{adv} \geq 10$	vertical isolation preserves formation history	Mismatch with local forcing; subsurface trends $\geq 2\times$ surface
Dynamical	$\tau_f \sim \tau_{adv} \sim \tau_{vmix} \sim \tau_{hmix}$ (hours–days)	No hierarchy; salinity contributes to buoyancy/frontogenesis (differential damping)	Near-equal T–S density control ($Tu \approx \pi/4$) at ~ 10 km mid-latitudes

610

This framework also makes the "rain gauge vs passive tracer" behavior easy to interpret. In the rain-gauge limit, anomalies stay close to where they are forced because forcing persists longer than circulation can export them from their formation region; the salinity map therefore resembles the E–P pattern. In the passive-tracer limit, anomalies are displaced from their formation region, either because lateral transport is faster than changes in forcing, or because the interior is vertically insulated. Salinity thus marks pathways and water-mass origin rather than local surface fluxes.

615

The submesoscale case is not just another point on the same continuum; it is where the interpretation changes. When the timescales converge, salinity cannot be treated as a record of forcing or a record of upstream conditions because both are evolving on the same hours-to-days window. Through differential damping (Section 4.3.2), salinity contributes comparably to density and enables vertical velocities $O(10-100 \text{ m day}^{-1})$ that ventilate the upper pycnocline, an order of magnitude stronger than mesoscale processes.

620

Two regions of the regime diagram remain the least well constrained. The $\tau_f/\tau_{adv} \sim O(1)$ transition on decadal scales, where substantial forcing amplification produces muted pattern response, still needs a clean mechanistic attribution (gyre adjustment, coupled variability, and/or changes in ventilation geometry are all plausible). Additionally, although the dynamical submesoscale regime is increasingly well observed, it remains poorly represented in many climate models; how



625 salinity-driven submesoscale overturning feeds back onto basin-scale stratification and biogeochemistry is still an open question.




Figure 8. Salinity regime diagram. Four regimes from timescale competition (Table 3). Rain gauge (surface, cyan-blue): $\tau_f / \tau_{adv} \geq 5$, forcing persists, pattern tracks *E-P* at basin-decadal scales. Passive tracer (subsurface, yellow-green): $\tau_{vmix} / \tau_{adv} \geq 10$, vertical isolation preserves formation history in mode waters. Transition (orange-yellow): $\tau_f / \tau_{adv} \sim 1-3$, forcing and advection compete, producing amplitude coherence with phase offsets over seasonal-decadal timescales. Dynamical (red): all timescales converge, equal T-S density control ($Tu \approx \pi/4$) at ~ 10 km submesoscales. Gradient shading shows gradual boundaries. Rain gauge and passive tracer coexist at decadal-basin scales as surface and subsurface regimes respectively. Diagram is schematic; boundaries vary with local conditions.

635

5 Synthesis and Outlook

This review shows that salinity's role is set by competition among three processes: freshwater forcing, advection by ocean currents, and mixing across gradients. Sections 3–4 show that their balance shifts systematically with scale. At submeso–mesoscales, salinity is dynamically active because it controls density, regulates stratification, and feeds back on mixing. At

640



seasonal–interannual scales, local forcing explains salinity evolution over only ~30% of the ocean surface; across the remaining ~70%, advection and mixing reshape anomalies before local forcing can dominate. At decadal scales, forcing control remains weak as many regions sit in transition. At multi-decadal scales, the forced imprint emerges as long-term salinity trend patterns correlate strongly ($r \approx 0.7–0.8$) with the climatological mean, yielding pattern amplification in which
645 where fresh regions freshen and salty regions become saltier.

Observations over the past two decades enabled this framework. The next step is observing the scales where the balance flips. Regime boundaries are expressed through fronts, filaments, and freshwater lenses that set stratification and mixing yet are smoothed by today's SSS resolution at 40–50 km. Without resolving these features, we map large-scale patterns but miss the mechanisms creating persistence, export, and vertical penetration. The most consequential gap is therefore specific thus:
650 sustained, global SSS at O(10 km) resolution, sampled frequently enough to track feature evolution and distinguish when salinity acts as a proxy for surface freshwater forcing, a tracer of pathways, or a driver of dynamics.

5.1 What the scale-dependent framework clarifies

The timescale competition framework clarifies three aspects of how salinity relates to forcing, circulation, and predictability.

Why spatial mismatch is informative. Interpreting salinity as a water-cycle indicator often focuses on whether it maps onto
655 E–P–R. The seasonal–interannual result (forcing-dominated behavior over ~30% of area; Fig.3) has been interpreted as evidence that salinity is unreliable across most of the ocean. But that interpretation assumes the only purpose of salinity is to mirror forcing. Sections 3–4 show that mismatch provides information. Where salinity departs from local forcing, water parcels have been transported or mixed before forcing accumulates locally. Those departures reveal quantities not directly observable, such as transit times, source regions, and coherence of pathways. Examples include thermocline salinity
660 correlating with subtropical surface conditions from several years earlier, subsurface salinity maxima recording the last surface contact of a water mass, and river plume far-fields whose phase lags trace boundary-current routes. The "70% mismatch" is signal, not noise.

Why salinity prediction depends on ocean state representation, not just atmospheric forcing. Salinity is a conservative variable. Once formed, an anomaly is transformed only by mixing, not by continuous exchange with the atmosphere (unlike
665 temperature). That conservation creates memory. A model can predict rainfall anomalies correctly yet still miss salinity because the salinity at any location reflects both local forcing and the water arriving from upstream (Fig.5). If the model misrepresents ocean currents or mixing, it delivers the wrong water to that location even when atmospheric forcing is correct. Conversely, salinity can be predictable even when rainfall is not, because water arriving today may have been formed years earlier in a distant region. This matters for practical prediction, as seasonal forecast systems can show good
670 performance in precipitation and SST yet fail in salinity because their circulation state, pathway geometry, or mixing



parameterizations are biased. Salinity forecast accuracy is therefore a coupled test: it requires both accurate forcing and ocean transport and mixing representations.

Why different studies reach different conclusions about forcing control. Basin-scale studies emphasizing robust pattern amplification and regional studies emphasizing advection and mixing are not disagreeing about physics; they examine different regime boundaries. At $O(1000 \text{ km})$ and multi-decadal scales, forcing evolves slowly enough that circulation and mixing redistribute anomalies repeatedly and forced patterns emerge. At $O(100 \text{ km})$ and seasonal scales, water travels large distances while forcing is evolving; local E–P–R competes directly with supply from upstream. Both are correct within their domains. The key question is not whether salinity indicates forcing, but which role dominates: forcing signal, pathway tracer, or dynamical driver..

680 **5.2 Resolving Regime Boundaries with $O(10 \text{ km})$ Satellite SSS**

Current satellite SSS products have an effective resolution of roughly $40\text{--}50 \text{ km}$ (Vinogradova et al. 2019; Reul et al. 2020). That scale is sufficient to map basin-scale patterns and many forced signals, but it smooths the features that often set the balance between forcing, advection, and mixing. The result is a persistent observational gap at $O(10 \text{ km})$, the range in which salinity often transitions from a largely passive indicator of freshwater fluxes to an active dynamical driver that shapes density, stratification, and vertical exchange.

The importance of $O(10 \text{ km})$ is dynamical, not simply a matter of detail. At these scales, three controls converge (McWilliams 2016). First, horizontal density gradients become steep enough to drive ageostrophic secondary circulations and vertical motions of order $10\text{--}100 \text{ m day}^{-1}$, directly coupling lateral structure to vertical exchange. Second, salinity contrasts between adjacent water masses become large enough that salinity can rival temperature in setting density. A 0.5 psu difference can produce a density effect comparable to 2°C , so neglecting salinity in the density budget becomes dynamically consequential. Third, freshwater-driven stratification reaches an intermediate regime where it is strong enough to inhibit routine wind mixing yet weak enough that storms can episodically overcome it. Whether freshwater anomalies persist, mix downward, or are exported laterally is therefore decided on storm timescales by the pre-existing frontal and stratification structure.

These controls are strongly nonlinear, and this is why coarse resolution is limiting. At $40\text{--}50 \text{ km}$, fronts, filaments, and freshwater lenses are blended into smooth gradients that do not represent either the sharp structures where buoyancy suppresses mixing or the well-mixed interiors where winds dominate. The relevant physics depends on the sharpness and vertical structure of the gradients, not on their spatial average. Storm response depends on the initial state. A sharp halocline at $10\text{--}20 \text{ m}$ can resist deepening and instead sharpen under shear, whereas a more gradual stratification profile allows mixing to deepen the mixed layer and entrain freshwater from below. Freshwater over saltwater creates stratification inhibiting



mixing; less mixing allows more freshwater to accumulate, strengthening stratification further (Drushka et al. 2016). This positive feedback operates only when gradients are sharp. Doubling the gradient does not double the stratification effect but can shift the regime from mixing-dominated to stratification-dominated.

705 A sustained $O(10\text{ km})$ SSS observing capability (Colliander et al., 2024) would turn these ambiguities into testable mechanisms. By tracking individual fronts and lenses through forcing events, it would quantify whether salinity stratification typically resists storm mixing or is routinely eroded, whether frontal structures sharpen through instabilities or broaden through wind stirring, and where and when the salinity contribution to density exceeds that of temperature. This is also the scale at which we can determine how much salinity variance, and therefore buoyancy variance, is currently unresolved. If the missing fraction is small, today's products and coarse models may be adequate for many climate diagnostics. If it is
710 substantial in key regions or seasons, then parameterized submesoscale processes are not a detail but a leading source of uncertainty in regional stratification, mixing, and coupled feedbacks.

This capability matters most where freshwater forcing and gradients are strongest. River-influenced shelves and boundary currents, tropical convergence zones with patchy rainfall, western boundary current extensions with intense fronts, and marginal ice zones where freshwater and heat fluxes interact all experience large freshwater fluxes per unit area where
715 salinity anomalies are strongest and gradients are sharpest. In these settings, $O(10\text{ km})$ SSS would resolve the nearshore-to-offshore transition from forcing-dominated plume cores to circulation-dominated export pathways, enabling quantitative estimates of export fractions, pathway timescales, and seasonal regime switching that current observations often describe only qualitatively. Improving to $O(10\text{ km})$ is not simply higher resolution but direct access to the mechanisms that control persistence, export, and vertical penetration of freshwater anomalies.

720 **5.3 Vertical Exchange and High-Latitude Regime Shifts**

Horizontal resolution addresses only part of the regime-boundary problem. Two additional observational gaps determine whether salinity anomalies remain surface-confined or become climate-relevant interior signals. Both involve transitions where competing processes cross critical thresholds, and both remain poorly constrained because current observing systems capture mean states rather than the events that drive transitions.

725 The first gap concerns mechanisms of vertical penetration. Subsurface salinity maxima at 100–200 m depth record surface forcing from years earlier (Fig.6), demonstrating that anomalies can penetrate and persist. Yet penetration depends on a buoyancy competition. Wintertime cooling removes buoyancy and destabilizes the water column, while freshwater input adds buoyancy and stabilizes it. Where haline stratification is weak, as in North Atlantic subtropical mode waters, cooling deepens mixed layers through hundreds of meters and subducts surface anomalies. Where haline stratification is strong, as in
730 the North Pacific, cooling cannot erode the fresh cap and surface anomalies remain shallow or are exported laterally.

Adjacent regions with similar atmospheric forcing show markedly different penetration depths, indicating that pre-existing subsurface structure and event timing matter as much as seasonal-mean forcing (Yeager & Large 2007).

735 The critical unknown is whether penetration occurs through gradual seasonal deepening or through episodic storms whose mechanical energy briefly overwhelms stratification (Dohan & Davis, 2011; Whitt & Taylor, 2017). If penetration is episodic, then a few intense storms per winter disproportionately set subduction and subsurface pathway memory. If gradual, then cumulative seasonal forcing sets penetration depth. The distinction matters for projections because climate models parameterize vertical mixing differently for gradual versus episodic processes, and future warming may strengthen haline stratification faster than it weakens thermal stratification. Testing these mechanisms requires high-frequency time series capturing individual storm responses with co-located measurements of winds, buoyancy fluxes, and vertical structure across
740 the seasonal cycle.

The second gap concerns high-latitude regime transitions. Arctic and subpolar seas switch between long stratified periods when freshwater accumulates near the surface and brief convective episodes when mixing homogenizes hundreds of meters within days (Marshall & Schott 1999). This switching controls whether freshwater anomalies influence deep water formation and overturning or remain surface properties. Arctic freshwater content varies by 10,000 km³ over decades, potentially affecting Atlantic overturning, but attribution requires understanding whether freshwater remains near-surface or penetrates to depth (Proshutinsky et al. 2009). Current observations are biased toward ice-free summer periods when stratification is strongest, missing winter convective events that may determine whether surface anomalies ventilate or remain trapped. Year-round sampling through under-ice platforms, moorings, and autonomous vehicles would capture regime shifts and test whether high-latitude regions are approaching thresholds where convective mixing weakens or freshwater trapping
750 strengthens under continued warming.

These gaps highlight that resolving regime boundaries requires observing not just finer spatial scales but also the temporal variability that drives transitions between states. The processes controlling vertical penetration operate on storm timescales; the processes controlling high-latitude freshwater pathways operate on seasonal cycles that include winter conditions that remain systematically undersampled.

755 **5.4 Model Uncertainties at Regime Boundaries**

Climate model ensembles reproduce basin-scale pattern amplification yet diverge sharply in regional projections at the regime boundaries emphasized in this review. Tropical differences trace to how models represent shallow overturning and the efficiency of exporting subtropical anomalies into the equatorial band (Wang et al. 2023). Barrier layer changes depend on how models balance precipitation increases against wind-driven mixing (Pang et al. 2023; Vogt et al. 2025). High-latitude
760 outcomes depend on whether models trigger or suppress episodic convection and how they handle ice-ocean freshwater



exchange (Haine et al. 2023). These divergences concentrate where observations are currently sparse or averaged over the critical scales.

765 Reducing projection uncertainty requires constraining regime-boundary physics rather than perfecting basin-mean trends. O(10 km) SSS observations would constrain front-resolving buoyancy structure at the surface, while coordinated vertical observing would constrain penetration mechanisms and high-latitude switching. Together, these would enable model evaluation to move beyond climatology to process-based metrics including phase lags between forcing and salinity response, depth and strength of subsurface salinity maxima, persistence statistics of freshwater lenses, and the contribution of salinity to density at fronts.

770 One path forward involves state-aware parameterizations. Rather than applying uniform mixing rules tied mainly to grid spacing, models could diagnose when a grid cell is front-like with strong gradients, lens-like with strong near-surface stratification, or quiescent with weak gradients, and adjust subgrid physics accordingly (Fox-Kemper et al. 2008). This approach would encode the framework's core insight that different balances apply in different regimes.

5.5 Broader implications

775 The scale- and regime-dependent framework developed here extends beyond salinity. Salinity is conservative enough to preserve formation history as water parcels move, changing primarily through boundary freshwater fluxes and mixing rather than continuous air-sea exchange. This conservation makes salinity an effective tracer of pathways and memory, and it explains why the same framework applies naturally to other quasi-conservative tracers such as nutrients, dissolved gases, and isotopes (Talley 2008). Water-cycle monitoring benefits from complementary constraints on the same forcing. At large scales and long timescales, freshwater fluxes alter both ocean mass through ocean bottom pressure and sea level and salt concentration through salinity (Liu et al. 2025). Where advection is slow relative to forcing accumulation, these constraints reinforce each other and can be used to evaluate and refine E-P-R products. Where advection is fast, both mass and salinity become mixed signals of forcing and circulation, and their joint interpretation becomes regime aware rather than purely local. Observing strategies that combine gravimetry, altimetry, and high-resolution SSS offer a practical route to separating water-cycle change from redistribution by circulation, provided the key regime boundaries are resolved.

785 Coupled feedbacks operate through stratification. Freshwater input strengthens stratification and suppresses vertical exchange; reduced exchange can trap heat and biogeochemical properties near the surface, further modifying density structure and mixing. This feedback is inherently coupled because temperature and salinity jointly set density, density regulates mixing, and mixing sets the evolution of both fields (Zika et al. 2018; Liu et al. 2023). At basin scales the coupling can appear weak because different forcings dominate the mean budgets, but at fronts and freshwater lenses it becomes 790 decisive on day-to-week timescales, precisely where nonlinearities are strongest. O(10 km) salinity observations target the



scales where coupling is most active and where small errors in stratification and mixing can propagate upscale through biases in heat uptake, ventilation, and tracer transport.

Physical-biological connections are similarly conditioned by regime boundaries, particularly by the persistence of physical structures relative to ecosystem response times (*ref*). Barrier layers and freshwater lenses create stratified surface habitats that persist for days to weeks, long enough for phytoplankton to respond but potentially too brief for higher trophic levels. Subsurface pathways transport nutrients and other tracers on multi-year routes, creating predictability only if mixing does not erase source signatures. In both cases, ecosystem predictability relates to how coherent pathways are and how fast mixing acts relative to transport. Advancing these connections involves observations that resolve the transition regimes where pathway coherence breaks down, where stratification and mixing flip between states, and where biological variability can be linked mechanistically to the evolving physical environment.

Concluding perspective

Over the past two decades, observations have turned salinity from a descriptive field into a physically interpretable signal. The organizing principle is competition among forcing, advection, and mixing timescales: where forcing evolves slowly relative to circulation and mixing, salinity records water-cycle change; where these timescales converge, salinity either reveals circulation pathways or actively shapes stratification and mixing. This framework explains why different studies reach different conclusions about forcing control by showing they examine different physical regimes.

The path forward is observation-driven model improvement. Resolving regime boundaries through $O(10 \text{ km})$ SSS, event-resolving vertical time series, and year-round high-latitude sampling would distinguish among mechanisms and enable models to represent physics as state dependent rather than spatially uniform. The questions this unlocks extend beyond salinity itself: how water-cycle intensification couples with circulation change, where stratification feedbacks amplify or dampen climate responses, and how regime transitions cascade through biogeochemical cycles.

Salinity's value lies in its bridging role across climate subsystems. It connects freshwater forcing to circulation pathways and to stratification-controlled exchange, revealing how the ocean stores, transports, and transforms water-cycle anomalies and how those transformations feed back on heat uptake, interior renewal, and ecosystems. Understanding salinity across scales provides a direct route to understanding ocean climate response.



820 **Code and data availability:** No new data or code used in this study.

Competing interests: The author declares no conflict of interest.

Author contribution: L.Y. is the sole author and responsible for all aspects of this work.

Acknowledgements

L.Y. gratefully acknowledges funding support from NASA Ocean Surface Salinity Science Team (OSSST) program (Grant
825 80NSSC22K0996)

References

- Abernathy, R. P. and Marshall, J.: Global surface eddy diffusivities derived from satellite altimetry, *J. Geophys. Res. Oceans*, 118, 901–916, <https://doi.org/10.1002/jgrc.20066>, 2013.
- 830 Anderson, D. L. T. and Gill, A. E.: Spin-up of a stratified ocean, with applications to upwelling, *Deep Sea Research and Oceanographic Abstracts*, 22, 583–596, [https://doi.org/10.1016/0011-7471\(75\)90046-7](https://doi.org/10.1016/0011-7471(75)90046-7), 1975.
- Aubone, N., Palma, E. D., and Piola, A. R.: The surface salinity maximum of the South Atlantic, *Progr. Oceanogr.*, 191, 102499, <https://doi.org/10.1016/j.pocean.2020.102499>, 2021.
- Balaguru, K., Foltz, G. R., Leung, L. R., Kaplan, J., Xu, W., Reul, N., and Chapron, B.: Pronounced Impact of Salinity on
835 Rapidly Intensifying Tropical Cyclones, *Bull. Amer. Meteor. Soc.*, 101, E1497–E1511, <https://doi.org/10.1175/BAMS-D-19-0303.1>, 2020.
- Barreiro, M., Fedorov, A., Pacanowski, R., and Philander, S. G.: Abrupt climate changes: How freshening of the northern Atlantic affects the thermohaline and wind-driven oceanic circulations, *Annual Review of Earth and Planetary Sciences*, 36, 33–58, <https://doi.org/10.1146/annurev.earth.36.090507.143219>, 2008.
- 840 Baumgartner, A. and Reichel, E.: *The World Water Balance*, Elsevier, New York, 179 pp., 1975.



- Bindoff, N. L. and McDougall, T. J.: Diagnosing Climate Change and Ocean Ventilation Using Hydrographic Data, *J. Phys. Oceanogr.*, 24, 1137–1152, [https://doi.org/10.1175/1520-0485\(1994\)024<1137:DCCA0V>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<1137:DCCA0V>2.0.CO;2), 1994.
- Bingham, F. M., Howden, S. D., and Koblinsky, C. J.: Sea surface salinity measurements in the historical database, *Journal of Geophysical Research*, 107, 8019, <https://doi.org/10.1029/2000JC000767>, 2002.
- 845 Bingham, F. M., Busecke, J. J. M., and Gordon, A. L.: Variability of the South Pacific subtropical surface salinity maximum, *Journal of Geophysical Research: Oceans*, 124, 6050–6066, <https://doi.org/10.1029/2018JC014598>, 2019.
- Boccaletti, G., Ferrari, R., and Fox-Kemper, B.: Mixed layer instabilities and restratification, *J. Phys. Oceanogr.*, 37, 2228–2250, <https://doi.org/10.1175/JPO3101.1>, 2007.
- Boutin, J., Vergely, J. L., Marchand, S., D'Amico, F., Hasson, A., Kolodziejczyk, N., et al.: New SMOS Sea Surface Salinity with reduced systematic errors and improved variability. *Remote Sensing of Environment*, 214, 115–134. <https://doi.org/10.1016/j.rse.2018.05.022>, 2018.
- 850 Boutin, J., Reul, N., Koehler, J., Martin, A., Catany, R., Guimbard, S., et al.: Satellite-based sea surface salinity designed for ocean and climate studies. *Journal of Geophysical Research: Oceans*, 126, e2021JC017676. <https://doi.org/10.1029/2021JC017676>, 2021.
- 855 Boyer, T. P., Levitus, S., Antonov, J. I., Locarnini, R. A., and Garcia, H. E.: Linear trends in salinity for the world ocean, 1955–1998, *Geophysical Research Letters*, 32, L01604, <https://doi.org/10.1029/2004GL021791>, 2005.
- Caesar, L., Rahmstorf, S., Robinson, A., Feulner, G., and Saba, V.: Observed fingerprint of a weakening Atlantic Ocean overturning circulation, *Nature*, 556, 191–196, <https://doi.org/10.1038/s41586-018-0006-5>, 2018.
- 860 Callies, J., and Ferrari, R.: Interpreting energy and tracer spectra of upper-ocean turbulence in the submesoscale range (1–200 km). *J. Phys. Oceanogr.*, 43(11), 2456–2474., <https://doi.org/10.1175/JPO-D-13-063.1>, 2013
- Camara, I., Kolodziejczyk, N., Mignot, J., Lazar, A., and Gaye, A. T.: On the seasonal variations of salinity of the tropical Atlantic mixed layer, *J. Geophys. Res. Oceans*, 120, 4441–4462, <https://doi.org/10.1002/2015JC010865>, 2015.
- Cessi, P. and Otheguy, P.: Oceanic Teleconnections: Remote Response to Decadal Wind Forcing, *J. Phys. Oceanogr.*, 33, 1604–1617, [https://doi.org/10.1175/1520-0485\(2003\)033<1604:OTRRTD>2.0.CO;2](https://doi.org/10.1175/1520-0485(2003)033<1604:OTRRTD>2.0.CO;2), 2003.
- 865 Cheng, L., Trenberth, K. E., Gruber, N., et al.: Improved estimates of changes in upper ocean salinity and the hydrological cycle, *J. Climate*, 33, 10357–10381, <https://doi.org/10.1175/JCLI-D-20-0366.1>, 2020.



- Coadou-Chaventon, S., Speich, S., Zhang, D., Rocha, C. B., and Swart, S.: Oceanic fronts driven by the Amazon freshwater plume and their thermohaline compensation at the submesoscale, *J. Geophys. Res. Oceans*, 129, e2024JC021326, <https://doi.org/10.1029/2024JC021326>, 2024.
- 870 Colliander, A., Crow, W., Entekhabi, D., Fournier, S., Harper, J., Holmes, T., et al.: Science of 10-km Resolution L-band Radiometry: Workshop Report, Jet Propulsion Laboratory, Pasadena, California, USA, <https://doi.org/10.48577/jpl.LY2KYW>, 2024.
- Curry, R., Dickson, B., and Yashayaev, I.: A change in the freshwater balance of the Atlantic Ocean over the past four decades, *Nature*, 426, 826–829, <https://doi.org/10.1038/nature02206>, 2003.
- 875 de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., and Iudicone, D.: Mixed layer depth over the global ocean: An examination of profile data and a profile-based climatology, *J. Geophys. Res.*, 109, C12003, <https://doi.org/10.1029/2004JC002378>, 2004.
- Delcroix, T. and Hénin, C.: Seasonal and interannual variations of sea surface salinity in the tropical Pacific Ocean, *J. Geophys. Res.*, 96, 22135–22150, <https://doi.org/10.1029/91JC02124>, 1991.
- 880 Dickson, R. R., Meincke, J., Malmberg, S. A., and Lee, A. J.: The "great salinity anomaly" in the Northern North Atlantic 1968–1982, *Progress in Oceanography*, 20, 103–151, [https://doi.org/10.1016/0079-6611\(88\)90049-3](https://doi.org/10.1016/0079-6611(88)90049-3), 1988.
- Dohan, K. and Davis, R. E.: Mixing in the transition layer during two storm events, *Journal of Physical Oceanography*, 41, 42–66, <https://doi.org/10.1175/2010JPO4253.1>, 2011.
- Dong, S., Baringer, M. O., Goni, G. J., Meinen, C. S., and Garzoli, S. L.: Seasonal variations in the South Atlantic meridional overturning circulation from observations and numerical models, *Geophys. Res. Lett.*, 41, 4611–4618, <https://doi.org/10.1002/2014GL060428>, 2014.
- Douville, H., and Cheng, L.: Asymmetric sea surface salinity response to global warming: “Fresh gets fresher but salty hesitates”. *Geophysical Research Letters*, 51, e2023GL107944. <https://doi.org/10.1029/2023GL107944>, 2024.
- 890 Drushka, K., Asher, W. E., Ward, B., and Walesby, K.: Understanding the formation and evolution of rain-formed fresh lenses at the ocean surface, *Journal of Geophysical Research*, 121, 2673–2689, <https://doi.org/10.1002/2015JC011527>, 2016.
- Drushka, K., Asher, W. E., Jessup, A. T., Thompson, E. J., Iyer, S., and Clark, D.: Capturing fresh layers with the surface salinity profiler, *Oceanography*, 32, 76–85, <https://doi.org/10.5670/oceanog.2019.215>, 2019.



- du Plessis, M. D., Swart, S., Biddle, L. C., Giddy, I. S., Monteiro, P. M. S., Reason, C. J. C., et al.: The daily-resolved Southern Ocean mixed layer: Regional contrasts assessed using glider observations, *Journal of Geophysical Research: Oceans*, 127, e2021JC017760, <https://doi.org/10.1029/2021JC017760>, 2022.
- 895
- Durack, P. J. and Wijffels, S. E.: Fifty-year trends in global ocean salinities and their relationship to broad-scale warming, *J. Climate*, 23, 4342–4362, <https://doi.org/10.1175/2010JCLI3377.1>, 2010.
- Durack, P. J., Wijffels, S. E., and Matear, R. J.: Ocean salinities reveal strong global water cycle intensification during 1950 to 2000, *Science*, 336, 455–458, <https://doi.org/10.1126/science.1212222>, 2012.
- 900 Entekhabi, D., Njoku, E. G., O'Neill, P. E., Kellogg, K. H., Crow, W. T., Edelstein, W. N., et al.: The Soil Moisture Active Passive (SMAP) mission, *Proc. IEEE*, 98, 704–716, 2010.
- Fernandez, A., Lapen, T. J., Andreasen, R., Swart, P. K., White, C. D., and Rosenheim, B. E.: Ventilation time scales of the North Atlantic subtropical cell revealed by coral radiocarbon from the Cape Verde Islands, *Paleoceanography*, 30, 938–948, <https://doi.org/10.1002/2015PA002790>, 2015.
- 905 Fine, R. A., Peacock, S., Maltrud, M. E., and Bryan, F. O.: A new look at ocean ventilation time scales and their uncertainties, *J. Geophys. Res.*, 122, 3771–3798, <https://doi.org/10.1002/2016JC012529>, 2017.
- Fournier, S., Lee, T., and Gierach, M. M.: Seasonal and interannual variations of sea surface salinity associated with the Mississippi River plume observed by SMOS and Aquarius, *Remote Sens. Environ.*, 180, 431–439, <https://doi.org/10.1016/j.rse.2016.02.050>, 2016.
- 910 Fournier, S., Vialard, J., Lengaigne, M., Lee, T., Gierach, M. M., and Chaitanya, A. V. S.: Modulation of the Ganges-Brahmaputra river plume by the Indian Ocean dipole and eddies inferred from satellite observations. *Journal of Geophysical Research: Oceans*, 122, 9591–9604, <https://doi.org/10.1002/2017JC013333>, 2017.
- Fournier, S., Reager, J. T., Chandanpurkar, H. A., Pascolini-Campbell, M., and Jarugula, S.: The salinity of coastal waters as a bellwether for global water cycle changes, *Geophysical Research Letters*, 50, e2023GL106684, <https://doi.org/10.1029/2023GL106684>, 2023.
- 915
- Fox-Kemper, B., Ferrari, R., and Hallberg, R.: Parameterization of Mixed Layer Eddies. Part I: Theory and Diagnosis, *J. Phys. Oceanogr.*, 38, 1145–1165, <https://doi.org/10.1175/2007JPO3792.1>, 2008.



- Friedman, A. R., Reverdin, G., Khodri, M., and Gastineau, G.: A new record of Atlantic sea surface salinity from 1896 to 2013 reveals the signatures of climate variability and long-term trends, *Geophys. Res. Lett.*, 44, 1866–1876, 920 <https://doi.org/10.1002/2017GL072582>, 2017.
- Good, S. A., Martin, M. J., and Rayner, N. A.: EN4: Quality controlled ocean temperature and salinity profiles and monthly objective analyses with uncertainty estimates, *Journal of Geophysical Research: Oceans*, 118, 6704–6716, <https://doi.org/10.1002/2013JC009067>, 2013.
- Gordon, A. L. and Giulivi, C. F.: Sea surface salinity trends over fifty years within the subtropical North Atlantic, 925 *Oceanography*, 21, 20–29, <https://doi.org/10.5670/oceanog.2008.64>, 2008.
- Gordon, A. L., Giulivi, C. F., Busecke, J., and Bingham, F. M.: Differences among subtropical surface salinity patterns, *Oceanography*, 28, 32–39, <https://doi.org/10.5670/oceanog.2015.02>, 2015.
- Gould, W. J. and Cunningham, S. A.: Global-scale patterns of observed sea surface salinity intensification since the 1870s, *Commun. Earth Environ.*, 2, 76, <https://doi.org/10.1038/s43247-021-00161-3>, 2021.
- 930 Grodsky, S. A., N. Reul, G. Lagerloef, G. Reverdin, J. A. Carton, B. Chapron, Y. Quilfen, V. N. Kudryavtsev, and H.-Y. Kao: Haline hurricane wake in the Amazon/Orinoco plume: AQUARIUS/SACD and SMOS observations, *Geophys. Res. Lett.*, 39, L20603, doi:10.1029/2012GL053335, 2012.
- Grodsky, S. A., Reverdin, G., Carton, J. A., and Coles, V. J.: Year-to-year salinity changes in the Amazon plume: Contrasting 2011 and 2012 Aquarius/SACD and SMOS satellite data, *Remote Sensing of Environment*, 140, 14–22, 935 <https://doi.org/10.1016/j.rse.2013.08.033>, 2014.
- Gu, D. and Philander, S. G. H.: Interdecadal climate fluctuations that depend on exchanges between the tropics and extratropics, *Science*, 275, 805–807, <https://doi.org/10.1126/science.275.5301.805>, 1997.
- Hackert, E., Kovach, R. M., Molod, A., Vernieres, G., Borovikov, A., Marshak, J., and Chang, Y.: Satellite sea surface salinity observations impact on El Niño/Southern Oscillation predictions: Case studies from the NASA GEOS seasonal 940 forecast system, *Journal of Geophysical Research: Oceans*, 125, e2019JC015788, <https://doi.org/10.1029/2019JC015788>, 2020.
- Haine, T. W. N., Siddiqui, A. H., and Jiang, W.: Arctic freshwater impact on the Atlantic Meridional Overturning Circulation: status and prospects, *Philos Trans A Math Phys Eng Sci*, 381, 20220185, <https://doi.org/10.1098/rsta.2022.0185>, 935 2023.



- 945 Hanawa, K. and Talley, L. D.: Mode waters, in: Ocean circulation and climate: Observing and modelling the global ocean, international geophysics series, edited by: Siedler, G., Church, J., and Gould, J., Vol. 77, Academic Press, London, 373–386, [https://doi.org/10.1016/S0074-6142\(01\)80129-7](https://doi.org/10.1016/S0074-6142(01)80129-7), 2001.
- Hasson, A., Delcroix, T., Boutin, J., Dussin, R., and Ballabrera-Poy, J.: Analyzing the 2010–2011 La Niña signature in the tropical Pacific sea surface salinity using in situ data, SMOS observations, and a numerical simulation, *Journal of Geophysical Research: Oceans*, 119, 3855–3867, <https://doi.org/10.1002/2013JC009388>, 2014.
- 950 Geophysical Research: Oceans, 119, 3855–3867, <https://doi.org/10.1002/2013JC009388>, 2014.
- Hasson, A., Puy, M., Boutin, J., Guilyardi, E., and Morrow, R.: Northward pathway across the tropical North Pacific Ocean revealed by surface salinity: How do El Niño anomalies reach Hawaii?, *Journal of Geophysical Research: Oceans*, 123, 2697–2715, <https://doi.org/10.1002/2017JC013423>, 2018.
- Hausmann, U., Czaja, A., and Marshall, J.: Mechanisms controlling the SST air-sea heat flux feedback and its dependence on spatial scale, *Clim Dyn*, 48, 1297–1307, <https://doi.org/10.1007/s00382-016-3142-3>, 2017.
- 955 on spatial scale, *Clim Dyn*, 48, 1297–1307, <https://doi.org/10.1007/s00382-016-3142-3>, 2017.
- Held, I. and Soden, B.: Robust responses of the hydrological cycle to global warming, *Journal of Climate*, 19, 5686–5699, <https://doi.org/10.1175/JCLI3990.1>, 2006.
- Hellweger, F. L. and Gordon, A. L.: Tracing Amazon River water into the Caribbean Sea, *J. Mar. Res.*, 60, 537–549, https://elischolar.library.yale.edu/journal_of_marine_research/2443, 2002.
- 960 Helm, K. P., Bindoff, N. L., and Church, J. A.: Changes in the global hydrological-cycle inferred from ocean salinity, *Geophys. Res. Lett.*, 37, L18701, <https://doi.org/10.1029/2010GL044222>, 2010.
- Hogg, N. G. and Johns, W. E.: Western boundary currents, *Rev. Geophys.*, 33, 1311–1334, <https://doi.org/10.1029/95RG00491>, 1995.
- Holliday, N.P., Bersch, M., Berx, B. et al.: Ocean circulation causes the largest freshening event for 120 years in eastern subpolar North Atlantic, *Nat Commun*, 11, 585, <https://doi.org/10.1038/s41467-020-14474-y>, 2020.
- 965 subpolar North Atlantic, *Nat Commun*, 11, 585, <https://doi.org/10.1038/s41467-020-14474-y>, 2020.
- Horner-Devine, A. R., Hetland, R. D., and MacDonald, D. G.: Mixing and transport in coastal river plumes, *Annu. Rev. Fluid Mech.*, 47, 569–594, <https://doi.org/10.1146/annurev-fluid-010313-141408>, 2015.
- Ishii, M., Kimoto, M., Sakamoto, K., and Iwasaki, S.-I.: Steric sea level changes estimated from historical ocean subsurface temperature and salinity analyses, *Journal of Oceanography*, 62, 155–170, <https://doi.org/10.1007/s10872-006-0041-y>, 2006.



- 970 Jaeger, G. S. and Mahadevan, A.: Submesoscale-selective compensation of fronts in a salinity-stratified ocean, *Sci. Adv.*, 4, e1701504, <https://doi.org/10.1126/sciadv.1701504>, 2018.
- Jarugula, S., Lee, T., Wang, O., and Fournier, S.: Maritime continent water cycle as a key forcing for decadal variation of upper-ocean salinity in the southeast Indian Ocean. *Journal of Geophysical Research: Oceans*, 130, e2025JC022733. <https://doi.org/10.1029/2025JC022733>, 2025.
- 975 Johnson, G. C., and McPhaden, M. J.: Interior Pycnocline Flow from the Subtropical to the Equatorial Pacific Ocean. *J. Phys. Oceanogr.*, 29, 3073–3089, [https://doi.org/10.1175/1520-0485\(1999\)029<3073:IPFFTS>2.0.CO;2](https://doi.org/10.1175/1520-0485(1999)029<3073:IPFFTS>2.0.CO;2), 1999.
- Kolodziejczyk, N. and Gaillard, F.: Variability of the heat and salt budget in the subtropical southeastern Pacific mixed layer between 2004 and 2010: Spice injection mechanism, *J. Phys. Oceanogr.*, 43, 1880–1898, <https://doi.org/10.1175/JPO-D-13-04.1>, 2013.
- 980 Kozlov, I. E., Plotnikov, E. V., and Manucharyan, G. E.: Brief Communication: Mesoscale and submesoscale dynamics in the marginal ice zone from sequential synthetic aperture radar observations, *The Cryosphere*, 14, 2941–2947, <https://doi.org/10.5194/tc-14-2941-2020>, 2020.
- Kraus, E. B. and Turner, J. S.: A one-dimensional model of the seasonal thermocline: II. The general theory and its consequences, *Tellus*, 19, 98–106, <https://doi.org/10.3402/tellusa.v19i1.9753>, 1967.
- 985 Lagerloef, G., DeCharon, A., and Lindstrom, E.: Ocean salinity and the Aquarius/SAC-D Mission: a new frontier in ocean remote sensing, *J. Mar. Technol. Soc.*, 47, 26–30, 2013.
- Ledwell, J., Watson, A., and Law, C.: Evidence for slow mixing across the pycnocline from an open-ocean tracer-release experiment, *Nature*, 364, 701–703, <https://doi.org/10.1038/364701a0>, 1993.
- Lee, T., Lagerloef, G., Gierach, M. M., Kao, H.-Y., Yueh, S., and Dohan, K.: Aquarius reveals salinity structure of tropical instability waves, *Geophys. Res. Lett.*, 39, L12610, <https://doi.org/10.1029/2012GL052232>, 2012.
- Lehmann, A., Myrberg, K., Post, P., Chubarenko, I., Dailidienė, I., et al.: Salinity dynamics of the Baltic Sea, *Earth Syst. Dynam.*, 13, 373–392, <https://doi.org/10.5194/esd-13-373-2022>, 2022.
- Li, G., Cheng, L., Zhu, J., Trenberth, K. E., Mann, M. E., and Abraham, J. P.: Increasing ocean stratification over the past half-century, *Nat. Clim. Change*, 10, 1116–1123, <https://doi.org/10.1038/s41558-020-00918-2>, 2020.



- 995 Li, L., Schmitt, R. W., Ummerhofer, C. C., and Karnauskas, K. B.: North Atlantic salinity as a predictor of Sahel rainfall, *Science Advances*, 2, e1501588, <https://doi.org/10.1126/sciadv.1501588>, 2016.
- Liu, H., Yu, L., and Lin, X.: Recent decadal change in the North Atlantic Subtropical Underwater associated with the poleward expansion of the surface salinity maximum, *J. Geophys. Res. Oceans*, 124, 4433–4448, <https://doi.org/10.1029/2018JC014508>, 2019.
- 1000 Liu, M., Soden, B. J., Vecchi, G. A., and Wang, C.: The spread of ocean heat uptake efficiency traced to ocean salinity, *Geophysical Research Letters*, 50, e2022GL100171, <https://doi.org/10.1029/2022GL100171>, 2023.
- Liu, C., Liang, X., and Yu, L.: Salinity trends and mass balances in the Mediterranean Sea: revisit the role of air-sea freshwater fluxes and oceanic exchange, *Ocean Sci.*, 21, 2069–2083, <https://doi.org/10.5194/os-21-2069-2025>, 2025.
- Lu, Y., Li, Y., Lin, P. et al: North Atlantic–Pacific salinity contrast enhanced by wind and ocean warming, *Nat. Clim. Chang.*, 14, 723–731, <https://doi.org/10.1038/s41558-024-02033-y>, 2024.
- 1005 Lukas, R. and Lindstrom, E.: The mixed layer of the western equatorial Pacific Ocean, *J. Geophys. Res.*, 96, 3343–3357, 1991.
- Lysne, J. A. and Deser, C.: Wind-Driven Thermocline Variability in the Pacific: A Model–Data Comparison, *J. Climate*, 15, 829–845, [https://doi.org/10.1175/1520-0442\(2002\)015<0829:WDTVIT>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0829:WDTVIT>2.0.CO;2), 2002.
- 1010 Lyu, Y., Bindoff, N. L., Mohapatra, S., Rathore, S., and Phillips, H. E.: Global water cycle pattern amplification: Contributing factors and mechanisms, *Journal of Geophysical Research: Oceans*, 130, e2024JC022278, <https://doi.org/10.1029/2024JC022278>, 2025.
- Maes, C., Picaut, J., and Belamari, S.: Salinity barrier layer and onset of El Niño in a Pacific coupled model, *Geophys. Res. Lett.*, 29, 2206, <https://doi.org/10.1029/2002GL016029>, 2002.
- 1015 Maes, C., Reul, N., Behringer, D., and O’kane, T.: The salinity signature of the equatorial Pacific cold tongue as revealed by the satellite SMOS mission, *Geoscience Letters*, 1, <https://doi.org/10.1186/s40562-014-0017-5>, 2014.
- Mahadevan, A. and Tandon, A.: An analysis of mechanisms for submesoscale vertical motion at ocean fronts, *Ocean Modelling*, 14, 241–256, <https://doi.org/10.1016/j.ocemod.2006.05.003>, 2006.
- Mahadevan, A., Tandon, A., and Ferrari, R.: Rapid changes in the mixed layer stratification driven by submesoscale instabilities and winds, *J. Geophys. Res. Oceans*, 115, C03017, 2010.
- 1020



- Marshall, J. C., Williams, R. G., and Nurser, A. J. G.: Inferring the Subduction Rate and Period over the North Atlantic. *J. Phys. Oceanogr.*, 23, 1315–1329, [https://doi.org/10.1175/1520-0485\(1993\)023<1315:ITSRAP>2.0.CO;2](https://doi.org/10.1175/1520-0485(1993)023<1315:ITSRAP>2.0.CO;2), 1993.
- Marshall, J. and Schott, F.: Open-ocean convection: Observations, theory, and models, *Rev. Geophys.*, 37, 1–64, <https://doi.org/10.1029/98RG02739>, 1999.
- 1025 McCreary, J. P. and Lu, P.: Interaction between the subtropical and equatorial ocean circulations: The subtropical cell, *J. Phys. Oceanogr.*, 24, 466–497, [https://doi.org/10.1175/1520-0485\(1994\)024<0466:IBTSAE>2.0.CO;2](https://doi.org/10.1175/1520-0485(1994)024<0466:IBTSAE>2.0.CO;2), 1994.
- McPhaden, M. J., Zebiak, S. E., and Glantz, M. H.: ENSO as an Integrating Concept in Earth Science, *Science*, 314, 1740–1745, <https://doi.org/10.1126/science.1132588>, 2006.
- McWilliams, J. C.: Submesoscale currents in the ocean, *Proc. Math. Phys. Eng. Sci.*, 472, 20160117, 1030 <https://doi.org/10.1098/rspa.2016.0117>, 2016.
- Meier, H. E. M., Kjellström, E., and Graham, L. P.: Estimating uncertainties of projected Baltic Sea salinity in the late 21st century, *Geophys. Res. Lett.*, 33, L15705, <https://doi.org/10.1029/2006GL026488>, 2006.
- Melnichenko, O., Amores, A., Maximenko, N., Hacker, P., and Potemra, J.: Signature of mesoscale eddies in satellite sea surface salinity data, *J. Geophys. Res. Oceans*, 122, 1416–1424, <https://doi.org/10.1002/2016JC012420>, 2017.
- 1035 Mignot, J., Frankignoul, C.: Local and remote impacts of a tropical Atlantic salinity anomaly. *Clim Dyn* 35, 1133–1147, <https://doi.org/10.1007/s00382-009-0621-9>, 2010.
- Mignot, J., Lazar, A., and Lacarra, M.: On the formation of barrier layers and associated vertical temperature inversions: A focus on the northwestern tropical Atlantic, *J. Geophys. Res.*, 117, C02010, <https://doi.org/10.1029/2011JC007435>, 2012.
- Morrow, R., Fu, L.-L., Arduin, F., Benkiran, M., Chapron, B., Cosme, E., et al.: Global observations of fine-scale ocean surface topography with the Surface Water and Ocean Topography (SWOT) mission, *Front. Mar. Sci.*, 6, 232, 1040 <https://doi.org/10.3389/fmars.2019.00232>, 2019.
- Müller, V., Kieke, D., Myers, P. G., Pennelly, C., Steinfeldt, R., and Stendardo, I.: Heat and freshwater transport by mesoscale eddies in the southern subpolar North Atlantic, *Journal of Geophysical Research: Oceans*, 124, 5565–5585, <https://doi.org/10.1029/2018JC014697>, 2019.
- 1045 Niiler, P. P.: Deepening of the wind-mixed layer, *Journal of Marine Research*, 33, https://elischolar.library.yale.edu/journal_of_marine_research/1328, 1975.



- Pang, S., Wang, X., and Vialard, J.: How Well Do CMIP6 Models Simulate Salinity Barrier Layers in the North Indian Ocean?, *J. Climate*, 37, 289–308, <https://doi.org/10.1175/JCLI-D-23-0366.1>, 2023.
- 1050 Patterson, R. G., Cronin, M. F., Swart, S., Beja, J., Edholm, J. M., et al.: Uncrewed surface vehicles in the global ocean observing system: A new Frontier for observing and monitoring at the air-sea interface, *Frontiers in Marine Science*, 12, <https://doi.org/10.3389/fmars.2025.1523585>, 2025.
- Pierce, D. W., Gleckler, P. J., Barnett, T. P., Santer, B. D., and Durack, P. J.: The fingerprint of human-induced changes in the ocean's salinity and temperature fields, *Geophys. Res. Lett.*, 39, L21704, <https://doi.org/10.1029/2012GL053389>, 2012.
- 1055 Proshutinsky, A., Krishfield, R., Timmermans, M.-L., Toole, J., Carmack, E., McLaughlin, F., Williams, W. J., Zimmermann, S., Itoh, M., and Shimada, K.: Beaufort Gyre freshwater reservoir: State and variability from observations, *J. Geophys. Res.*, 114, C00A10, <https://doi.org/10.1029/2008JC005104>, 2009.
- Qiu, B., and Huang, R. X.: Ventilation of the North Atlantic and North Pacific: Subduction Versus Obduction. *J. Phys. Oceanogr.*, 25, 2374–2390, [https://doi.org/10.1175/1520-0485\(1995\)025<2374:VOTNAA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1995)025<2374:VOTNAA>2.0.CO;2), 1995.
- 1060 Qu, T., Gao, S., and Fukumori, I.: Formation of salinity maximum water and its contribution to the overturning circulation in the North Atlantic as revealed by a global general circulation model, *Journal of Geophysical Research: Oceans*, 118, 1982–1994, <https://doi.org/10.1002/jgrc.20152>, 2013.
- Qu, T., Song, Y. T., and Maes, C.: Sea surface salinity and barrier layer variability in the equatorial Pacific as seen from Aquarius and Argo, *J. Geophys. Res. Oceans*, 119, 15–29, <https://doi.org/10.1002/2013JC009375>, 2014.
- 1065 Qu, T., Zhang, L., and Schneider, N.: North Atlantic subtropical underwater and its year-to-year variability in annual subduction rate during the Argo period, *Journal of Physical Oceanography*, 46, 1901–1916, <https://doi.org/10.1175/JPO-D-15-0246.1>, 2016.
- Rahmstorf, S.: Bifurcations of the Atlantic thermohaline circulation in response to changes in the hydrological cycle, *Nature*, 378, 145–149, <https://doi.org/10.1038/378145a0>, 1995.
- 1070 Rahmstorf, S., Crucifix, M., Ganopolski, A., Goosse, H., Kamenkovich, I., et al.: Thermohaline circulation hysteresis: A model intercomparison, *Geophys. Res. Lett.*, 32, L23605, <https://doi.org/10.1029/2005GL023655>, 2005.
- Rao, R. R. and Sivakumar, R.: Seasonal variability of sea surface salinity and salt budget of the mixed layer of the north Indian Ocean, *J. Geophys. Res.*, 108, 3009, <https://doi.org/10.1029/2001JC000907>, 2003.



- Rathore, S., Bindoff, N. L., Ummenhofer, C. C., Phillips, H. E., Feng, M., and Mishra, M.: Improving Australian rainfall prediction using sea surface salinity, *Journal of Climate*, 34, 2473–2490, <https://doi.org/10.1175/JCLI-D-20-0625.1>, 2021.
- 1075 Reul, N., Tenerelli, J., Chapron, B., Vandemark, D., Quilfen, Y., and Kerr, Y.: SMOS satellite L-band radiometer: a new capability for ocean surface remote sensing in hurricanes. *J. Geophys. Res.* 117:C02006. doi: 10.1029/2011JC007474, 2012.
- Reul, N., Chapron, B., Lee, T., Donlon, C., Boutin, J., and Alory, G.: Sea surface salinity structure of the meandering Gulf Stream revealed by SMOS sensor, *Geophys. Res. Lett.*, 41, 3141–3148, <https://doi.org/10.1002/2014GL059215>, 2014.
- Reul, N., Grodsky, S. A., Arias, M., et al.: Sea surface salinity estimates from spaceborne L-band radiometers: An overview
1080 of the first decade of observation (2010–2019). *Remote Sens. Environ.*, 242, 111769, <https://doi.org/10.1016/j.rse.2020.111769>, 2020.
- Riser, S., Freeland, H., Roemmich, D., Wijffels, S., Troisi, A., et al.: Fifteen years of ocean observations with the global Argo array, *Nature Clim Change*, 6, 145–153, <https://doi.org/10.1038/nclimate2872>, 2016.
- Rohling, E. J., and Bigg, G. R.: Paleosalinity and $\delta^{18}\text{O}$: A critical assessment, *J. Geophys. Res.*, 103(C1), 1307–1318,
1085 doi:10.1029/97JC01047, 1998.
- Roemmich, D. and Gilson, J.: The 2004–2008 mean and annual cycle of temperature, salinity, and steric height in the global ocean from the Argo Program, *Prog. Oceanogr.*, 82, 81–100, <https://doi.org/10.1016/j.pocean.2009.03.004>, 2009.
- Röthig, T., Trevathan-Tackett, S. M., Voolstra, C. R., Ross, C., Chaffron, S., et al.: Human-induced salinity changes impact marine organisms and ecosystems, *Global Change Biology*, 29, 4731–4749, <https://doi.org/10.1111/gcb.16859>, 2023.
- 1090 Ruddick, B. R.: A Practical Indicator of the Stability of the Water Column to Double-Diffusive Activity, *Deep-Sea Research*, 30, 1105–1107, [https://doi.org/10.1016/0198-0149\(83\)90063-8](https://doi.org/10.1016/0198-0149(83)90063-8), 1983.
- Rudnick, D. L. and Ferrari, R.: Compensation of horizontal temperature and salinity gradients in the ocean mixed layer, *Science*, 283, 526–529, <https://doi.org/10.1126/science.283.5401.526>, 1999.
- Salisbury, J., Vandemark, D., Campbell, J., Hunt, C., Wisser, D., Reul, N., and Chapron, B.: Spatial and temporal coherence
1095 between Amazon River discharge, salinity, and light absorption by colored organic carbon in western tropical Atlantic surface waters, *J. Geophys. Res.*, 116, C00H02, <https://doi.org/10.1029/2011JC006989>, 2011.
- Schanze, J. J., Schmitt, R. W., and Yu, L. L.: The global oceanic freshwater cycle: A state-of-the-art quantification, *J. Mar. Res.*, 68, 569–595, https://elischolar.library.yale.edu/journal_of_marine_research/280, 2010.



- 1100 Schmitt, R.W.: Salinity and the global water cycle. *Oceanography* 21(1):12–19, <https://doi.org/10.5670/oceanog.2008.63>,
2008.
- Singh, H. K. A., Donohoe, A., Bitz, C. M., Nusbaumer, J., and Noone, D. C.: Greater aerial moisture transport distances with warming amplify interbasin salinity contrasts, *Geophys. Res. Lett.*, 43, 8677–8684, doi:10.1002/2016GL069796, 2016.
- Skliris, N., Marsh, R., Josey, S. A., Good, S. A., Liu, C., and Allan, R. P.: Salinity changes in the World Ocean since 1950 in relation to changing surface freshwater fluxes, *Clim. Dynam.*, 43, 709–736, <https://doi.org/10.1007/s00382-014-2131-7>,
1105 2014.
- Skliris, N., Zika, J. D., Nurser, G., Josey, S. A., and Marsh, R.: Global water cycle amplifying at less than the Clausius-Clapeyron rate, *Sci. Rep.*, 6, 38752, <https://doi.org/10.1038/srep38752>, 2016.
- Skliris, N., Zika, J.D., Herold, L. et al.: Mediterranean sea water budget long-term trend inferred from salinity observations. *Clim Dyn* 51, 2857–2876, <https://doi.org/10.1007/s00382-017-4053-7>, 2018.
- 1110 Stendardo, I., Rhein, M., and Hollmann, R.: A high resolution salinity time series 1993–2012 in the North Atlantic from Argo and Altimeter data, *J. Geophys. Res. Oceans*, 121, 2523–2551, <https://doi.org/10.1002/2015JC011439>, 2016.
- Talley, L.D.: Freshwater Transport Estimates and the Global Overturning Circulation: Shallow, Deep and through Flow Components. *Progress in Oceanography*, 78, 257-303. <https://doi.org/10.1016/j.pocean.2008.05.001>, 2008.
- 1115 Tapley, B. D., Bettadpur, S., Ries, J. C., Thompson, P. F., and Watkins, M. M.: GRACE measurements of mass variability in the Earth system, *Science*, 305, 503–505, <https://doi.org/10.1126/science.1099192>, 2004.
- Taylor, J. R. and Ferrari, R.: Buoyancy and Wind-Driven Convection at Mixed Layer Density Fronts, *J. Phys. Oceanogr.*, 40, 1222–1242, <https://doi.org/10.1175/2010JPO4365.1>, 2010.
- Terray, L., Corre, L., Cravatte, S., Delcroix, T., Reverdin, G., and Ribes, A.: Near-surface salinity as Nature's rain gauge to detect human influence on the tropical water cycle, *J. Climate*, 25, 958–977, <https://doi.org/10.1175/JCLI-D-10-05025.1>,
1120 2012.
- Thomas, L. and Ferrari, R.: Friction, Frontogenesis, and the Stratification of the Surface Mixed Layer, *J. Phys. Oceanogr.*, 38, 2501–2518, <https://doi.org/10.1175/2008JPO3797.1>, 2008.
- Timmermans, M.-L. and Winsor, P.: Scales of horizontal density structure in the Chukchi Sea surface layer, *Cont. Shelf Res.*, 52, 39–45, <https://doi.org/10.1016/j.csr.2012.10.015>, 2013.



- 1125 Vinogradova, N. T. and Ponte, R. M.: Clarifying the link between surface salinity and freshwater fluxes on monthly to interannual time scales, *J. Geophys. Res.*, 118, 3190–3201, <https://doi.org/10.1002/jgrc.20200>, 2013.
- Vinogradova, N. T. and Ponte, R. M.: In search of fingerprints of the recent intensification of the ocean water cycle, *J. Climate*, 30, 5513–5528, <https://doi.org/10.1175/JCLI-D-16-0626.1>, 2017.
- Vinogradova, N., Lee, T., Boutin, J., Drushka, K., Fournier, S., et al.: Satellite salinity observing system: recent discoveries and the way forward, *Front. Mar. Sci.*, 6, 243, <https://doi.org/10.3389/fmars.2019.00243>, 2019.
- 1130 Vinogradova, N. T., Pavelsky, T. M., Farrar, J. T., Hossain, F., and Fu, L.-L.: A new look at Earth's water and energy with SWOT, *Nat Water*, 3, 27–37, <https://doi.org/10.1038/s44221-024-00372-w>, 2025.
- Vogt, L., Sallée, J.-B., and de Lavergne, C.: Stratification and overturning circulation are intertwined controls on ocean heat uptake efficiency in climate models, *Ocean Sci.*, 21, 1081–1103, <https://doi.org/10.5194/os-21-1081-2025>, 2025.
- 1135 Wang, F., Xu, X., Zhang, F., and Ma, L.: Structure of the Atlantic meridional overturning circulation in three generations of climate models, *Earth and Space Science*, 10, e2023EA002887, <https://doi.org/10.1029/2023EA002887>, 2023.
- Warren, B. A.: Why is no deep water formed in the North Pacific?, *J. Mar. Res.*, 41, 327–347, https://elischolar.library.yale.edu/journal_of_marine_research/1685, 1983.
- Weijer, W., Cheng, W., Drijfhout, S. S., Federov, A. V., Hu, A., et al.: Stability of the Atlantic Meridional Overturning Circulation: A review and synthesis, *Journal of Geophysical Research: Oceans*, 124, 5336–5375, <https://doi.org/10.1029/2019JC015083>, 2019.
- 1140 Whitt, D. B. and Taylor, J. R.: Energetic Submesoscales Maintain Strong Mixed Layer Stratification during an Autumn Storm, *J. Phys. Oceanogr.*, 47, 2419–2427, <https://doi.org/10.1175/JPO-D-17-0130.1>, 2017.
- Wunsch, C. and Ferrari, R.: Vertical mixing, energy, and the general circulation of the oceans, *Annual Review of Fluid Mechanics*, 36, 281–314, <https://doi.org/10.1146/annurev.fluid.36.050802.122121>, 2004.
- 1145 Yeager, S. G. and Large, W. G.: Observational Evidence of Winter Spice Injection, *J. Phys. Oceanogr.*, 37, 2895–2919, <https://doi.org/10.1175/2007JPO3629.1>, 2007.
- Yu, L.: A global relationship between the ocean water cycle and near-surface salinity, *Journal of Geophysical Research – Oceans*, 116, C10025, <https://doi.org/10.1029/2010JC006937>, 2011.



- 1150 Yu, L.: Sea-surface salinity fronts and associated salinity-minimum zones in the tropical ocean, *J. Geophys. Res. Oceans*, 120, 4205–4225, <https://doi.org/10.1002/2015JC010790>, 2015.
- Yu, L., Bingham, F. M., Lee, T., Dinnat, E. P., Fournier, S., Melnichenko, O., Tang, W., and Yueh, S. H.: Revisiting the global patterns of seasonal cycle in sea surface salinity, *J. Geophys. Res. Oceans*, 126, e2020JC016789, <https://doi.org/10.1029/2020JC016789>, 2021.
- 1155 Yu, L., Jin, X., and Liu, H.: Poleward shift in ventilation of the North Atlantic subtropical underwater, *Geophys. Res. Lett.*, 45, 258–266, <https://doi.org/10.1002/2017GL075772>, 2018.
- Yu, L., Josey, S. A., Bingham, F. M., and Lee, T.: Intensification of the global water cycle and evidence from ocean salinity: a synthesis review, *Ann. N.Y. Acad. Sci.*, 1472, 76–94, <https://doi.org/10.1111/nyas.14354>, 2020.
- Yu, L.: Connecting subtropical salinity maxima to tropical salinity minima: Synchronization between ocean dynamics and the water cycle, *Progress in Oceanography*, 219, 103172, <https://doi.org/10.1016/j.pocean.2023.103172>, 2023.
- 1160 Yu, L.: Meso–Submesoscale T–S Compensation and Density Variability in the North Atlantic from Saildrone, *J. Phys. Oceanogr.*, 56, 115–134, <https://doi.org/10.1175/JPO-D-25-0037.1>, 2026.
- Zhu, J., Huang, B., Zhang, R. H., Hu, Z. Z., Kumar, A., Balmaseda, M. A., Marx, L., and Kinter III, J. L.: Salinity anomaly as a trigger for ENSO events, *Sci Rep*, 4, 6821, <https://doi.org/10.1038/srep06821>, 2014.
- 1165 Zika, J. D., Skliris, N., Nurser, A. J. G., Josey, S. A., Mudryk, L., Laliberté, F., and Marsh, R.: Maintenance and Broadening of the Ocean's Salinity Distribution by the Water Cycle, *J. Climate*, 28, 9550–9560, <https://doi.org/10.1175/JCLI-D-15-0273.1>, 2015.
- Zika, J. D., Skliris, N., Blaker, A. T., Marsh, R., Nurser, A. G., and Josey, S. A.: Improved estimates of water cycle change from ocean salinity: The key role of ocean warming, *Environmental Research Letters*, 13, 074036, <https://doi.org/10.1088/1748-9326/aace42>, 2018.
- 1170 Zika, J. D., Gregory, J. M., McDonagh, E. L., Marzocchi, A., and Clément, L.: Recent Water Mass Changes Reveal Mechanisms of Ocean Warming, *J. Climate*, 34, 3461–3479, <https://doi.org/10.1175/JCLI-D-20-0355.1>, 2021.