

1 **Enhanced methane monitoring: A globally harmonized daily**  
2 **0.1° XCH<sub>4</sub> through machine learning-based fusion of GOSAT,**  
3 **GOSAT-2, and TROPOMI**

4 Jebun Naher Keya<sup>1,\*</sup>, Yejin Kim<sup>1,\*</sup>, Hyunyoung Choi<sup>1</sup>, Jungho Im<sup>1,2,3,4,\*</sup>

5 <sup>1</sup>Department of Civil, Urban, Earth, & Environmental Engineering, Ulsan National Institute of Science and  
6 Technology, Ulsan, 44919, Republic of Korea

7 <sup>2</sup>Graduate School of Carbon Neutrality, Ulsan National Institute of Science and Technology, Ulsan, 44919,  
8 Republic of Korea

9 <sup>3</sup>Artificial Intelligence Graduate School, Ulsan National Institute of Science and Technology, Ulsan, 44919,  
10 Republic of Korea

11 <sup>4</sup>Department of Environmental Resources Engineering, State University of New York College of Environmental  
12 Science and Forestry, Syracuse, NY 13210, USA

13  
14 \*Corresponding Author: Jungho Im ([ersgis@unist.ac.kr](mailto:ersgis@unist.ac.kr))

15 \*These authors contributed equally to this work.

16  
17  
18  
19  
20

21 **Supplementary**

22

23 **Equations S1–S3 (Statistical Metrics for Model Performance Evaluation)**

24 Model performance was evaluated using four standard statistical metrics: coefficient of  
25 determination ( $R^2$ ; Eq.1), root mean square error (RMSE; Eq. 2), mean bias error (MBE; Eq.  
26 3) and mean absolute error (MAE; Eq.4).

27 
$$R^2 = 1 - \frac{\sum_i (y_i - \hat{y}_i)^2}{\sum_i (y_i - \bar{y})^2} \quad (1)$$

28 
$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (\hat{y}_i - y_i)^2} \quad (2)$$

29 
$$MAE = \frac{1}{n} \sum_{i=1}^n |\hat{y}_i - y_i| \quad (3)$$

30

31 In the equation,  $\hat{y}_i$  represents the predicted value,  $y_i$  is observed value,  $\bar{y}$  is the mean  
32 of observed values, and  $n$  is the total number of samples

33

34 **Table S1.** TCCON (GGG2020 version) sites used for evaluation of satellite XCH<sub>4</sub> and machine  
 35 learning based bias correction.

TCCON site	Latitude, Longitude (°)	Elevation (m)	Data reference
Bremen <sup>a, b</sup>	53.10, 8.85	30	Notholt et al., 2022
Burgos	18.53, 120.65	35	Morino et al., 2022
Darwin	12.46, 130.93	37	Deutscher et al., 2023
East Trout Lake	54.35, -104.99	501.8	Wunch et al., 2022
Edwards <sup>c</sup>	34.96, -117.88	888	Iraci et al., 2022
Garmisch <sup>c</sup>	47.48, 11.06	743	Sussmann et al., 2025
Hefei	31.90, 117.17	30	Liu et al., 2023
Karlsruhe	49.10, 8.44	119	Hase et al., 2024
Lamont	36.60, -97.49	320	Wennberg et al., 2025
Lauder	-45.04, 169.68	370	Pollard et al., 2022
Nicosia <sup>a</sup>	35.14, 33.38	185	Petri et al., 2024
Orléans	47.97, 2.11	130	Warneke et al., 2024
Paris	48.85, 2.36	60	Té et al., 2022
Park Falls	45.95, -90.27	442	Wennberg et al., 2022
Pasadena	34.14, -118.13	210	Wennberg et al., 2022
Rikubetsu <sup>a</sup>	43.46, 143.77	380	Morino et al., 2022
Saga	33.24, 130.29	7	Shiomi et al., 2022
Sodankylä	67.37, 26.63	188	Kivi et al., 2022
Tsukuba	36.05, 140.12	31	Morino et al., 2022
Xianghe <sup>c</sup>	39.75, 116.96	36	Zhou et al., 2022

36 <sup>a</sup> Sites excluded from the GOSAT evaluation under the GOSAT collocation criteria. <sup>b</sup> Sites  
 37 excluded from the GOSAT-2 evaluation under the GOSAT-2 collocation criteria. <sup>c</sup>  
 38 Independent test sites not used for training/validation of the machine learning based bias-  
 39 correction model.

40  
 41

42 **Table S2.** GOSAT XCH<sub>4</sub> retrieval parameters used to estimate  $\Delta(\text{GOSAT} - \text{TCCON})$ .

<b>GOSAT predictor variables</b>	<b>Units</b>
1. Bias corrected XCO <sub>2</sub> retrieved	ppb
1. Aerosol Optical Thickness (AOT1 & 2)	-
2. Cirrus Optical Thickness	-
3. Longitude	degree
4 Reference CO <sub>2</sub> Profile (total)	ppb
5. Reference CO <sub>2</sub> Profile 1	Mol cm <sup>-2</sup>
6. Satellite zenith	degree
7. Reference pressure – retrieved pressure ( $\Delta P$ s)	hPa
8. Temperature shift	K

43  
44  
45  
46

47 **Table S3.** GOSAT-2 XCH<sub>4</sub> retrieval parameters used to estimate  $\Delta(\text{GOSAT-2} - \text{TCCON})$ .

<b>GOSAT-2 predictor variables</b>	<b>Units</b>
1. Reference pressure – retrieved pressure ( $\Delta P_s$ )	hPa
2. XCO <sub>2</sub> retrieved	-
3. H <sub>2</sub> O Profile uncertainty	ppb
4. Longitude	deg
5. XCH <sub>4</sub> Degrees of Freedom	-
6. Pressure level	hPa
7. Solar Zenith Angle (SZA)	deg
8. Column averaging kernel vertical level 1	-
9. Fluorescent Slope	-
10. Aerosol Optical Thickness Type1 + Type2 (total)	-
11. Residual Chi-Squared : Band 3	-

48  
49  
50

51 **Table S4.** TROPOMI XCH<sub>4</sub> retrieval parameters used to estimate  $\Delta(\text{TROPOMI} - \text{TCCON})$  in  
 52 step 1 and  $\Delta(\text{TROPOMI} - \text{GOSAT-2})$  in step 2

TROPOMI predictor variables (abbreviation)	Units
1. XCH <sub>4</sub> precision	ppb
2. Fluorescence	mol s <sup>-1</sup> m <sup>-2</sup> nm <sup>-1</sup> sr <sup>-1</sup>
3–4. CO total column and precision	mol m <sup>-2</sup>
5–6. H <sub>2</sub> O total column and precision	mol m <sup>-2</sup>
7–8. Aerosol size and precision	-
9–10. Aerosol height and precision	m
11–12. Aerosol column and precision	m <sup>-2</sup>
13–14. NIR surface albedo and precision	-
15–16. SWIR surface albedo and precision	-
17. NIR aerosol optical thickness	-
18. SWIR aerosol optical thickness	-
19. NIR chi-square	-
20. SWIR chi-square	-
22. Solar zenith angle (SOZ)	degree
23. Solar azimuth angle (SOA)	degree
24. Satellite zenith angle (SAZ)	degree
25. Satellite azimuth angle (SAA)	degree
26. Surface altitude	m
27. Surface classification	-
28. U10 wind speed (U10)	m s <sup>-1</sup>
29. V10 wind speed (V10)	m s <sup>-1</sup>
30. NIR cirrus reflectance	-
31. SWIR cirrus reflectance	-
32. XCH <sub>4</sub> profile a priori	mol m <sup>-2</sup>

53  
 54

55 **Table S5.** GOSAT XCH<sub>4</sub> retrieval parameters used to estimate  $\Delta(\text{GOSAT} - \text{GOSAT-2})$  in Step2

1. Surface pressure	hPa
2. Reference surface pressure	hPa
3. Total column of dry air molecules	Mol cm <sup>-2</sup>
4. $\Delta P$ s : reference pressure – retrieved pressure	hPa
5. Altitude (height)	m
6. Retrieved XCO <sub>2</sub>	ppb
7. Bias corrected XCO <sub>2</sub>	ppb
8–19. Reference CO <sub>2</sub> profile at pressure levels P4–P15	Mol cm <sup>-2</sup>
20. Total reference CO <sub>2</sub> profile column	Mol cm <sup>-2</sup>
21. Vertically integrated CH <sub>4</sub> column	Mol cm <sup>-2</sup>
22. Aerosol Optical Thickness type-2	-
23. Air mass	-
24–25. Residual mean square	-
26. Column averaging kernel vertical level 6	-

56  
57  
58  
59

60 **Table S6.** Cross-validation results comparing Random Forest (RF) and eXtreme Gradient  
 61 Boosting (XGBoost) for TROPOMI, GOSAT, and GOSAT-2 bias correction against TCCON.  
 62 For each metric, the better-performing value between the two algorithms is shown in bold.

		<b>RF</b>			<b>XGBoost</b>		
	CV type	LOSOCV <sup>a</sup>	LOMOCV <sup>b</sup>	LOYOCV <sup>c</sup>	LOSOCV	LOMOCV	LOYOCV
TROPOMI	N	5,156					
	R <sup>2</sup>	0.78	0.82	0.81	0.79	0.84	0.82
	MAE (ppb)	10.48	9.34	9.39	10.20	8.83	9.17
	RMSE (ppb)	13.94	12.78	12.87	13.58	12.06	12.57
	N	702					
GOSAT	R <sup>2</sup>	0.87	0.88	0.86	0.86	0.88	0.86
	MAE (ppb)	8.20	7.66	8.42	8.43	7.72	8.53
	RMSE (ppb)	10.85	10.16	11.09	11.13	10.22	11.19
	N	1,751					
GOSAT-2	R <sup>2</sup>	0.88	0.91	0.89	0.88	0.90	0.89
	MAE (ppb)	7.68	7.07	7.45	7.89	7.19	7.67
	RMSE (ppb)	10.96	9.86	10.65	11.22	10.04	10.84

63 <sup>a</sup> LOSOCV: Leave-One-Site-Out Cross-Validation

64 <sup>b</sup> LOMOCV: Leave-One-Month-Out Cross-Validation

65 <sup>c</sup> LOYOCV: Leave-One-Year-Out Cross-Validation

66

67

68 **Table S7.** Cross-validation results comparing Random Forest (RF), eXtreme Gradient  
 69 Boosting (XGBoost) and Light Gradient Boosting Machine (LightGBM) for TROPOMI and  
 70 GOSAT bias correction against best-performing ML-based bias-corrected GOSAT-2. For each  
 71 metric, the better-performing value between the three algorithms is shown in bold.

		<b>RF</b>			<b>XGBoost</b>		
CV type		LOSOCV <sup>a</sup>	LOMOCV <sup>b</sup>	LOYOCV <sup>c</sup>	LOSOCV	LOMOCV	LOYOCV
TROPOMI	N	183,550					
	R <sup>2</sup>	0.86	0.87	0.87	0.91	0.92	0.92
	MAE (ppb)	8.50	8.13	8.18	6.78	6.32	6.59
	RMSE (ppb)	11.18	10.72	10.78	9.07	8.49	8.82
GOSAT	N	32,244					
	R <sup>2</sup>	0.91	0.92	0.91	0.92	0.92	0.92
	MAE (ppb)	7.22	7.00	9.94	7.06	6.86	7.11
	RMSE (ppb)	9.80	9.50	7.38	9.56	9.33	9.64

72 <sup>a</sup> LOSOCV: Leave-One-Site-Out Cross-Validation

73 <sup>b</sup> LOMOCV: Leave-One-Month-Out Cross-Validation

74 <sup>c</sup> LOYOCV: Leave-One-Year-Out Cross-Validation

75

76


77

78 **Table S9.** Seasonal hemispheric and high-emission zone mean XCH<sub>4</sub> concentrations in 2023.  
 79 Values represent area-weighted seasonal means and spatial standard deviations indicating  
 80 spatial variability within each region. Amplitude represents the seasonal range (maximum  
 81 minus minimum concentration) for each region per season.

<b>Region</b>	<b>MAM (ppb)</b>	<b>JJA (ppb)</b>	<b>SON (ppb)</b>	<b>DJF (ppb)</b>	<b>Amplitude (ppb)</b>
0–40N	1903.71 ± 10.04	1901.18 ± 6.27	1917.13 ± 10.69	1917.68 ± 9.38	16.50
0–80N	1883.22 ± 25.05	1889.30 ± 16.68	1903.46 ± 21.91	1915.36 ± 10.09	32.14
60S–0	1850.66 ± 31.90	1857.43 ± 24.52	1856.62 ± 25.22	1854.55 ± 29.29	6.77


82 Note- NH: Northern Hemisphere (0–80°N); High-emission zone (0–40°N); SH: Southern  
 83 Hemisphere (60°S–0°). MAM, JJA, SON, and DJF represent March–May, June–August,  
 84 September–November, and December–February, respectively.  
 85  
 86  
 87

88 **Figure S1.** Spatial cross-validation (SPCV) fold distribution for latitudinal generalization  
89 assessment. The bias correction models were evaluated using leave-one-band-out cross-  
90 validation (LOBOCV) across eleven latitude-based folds spanning 60°S–80°N.  
91



92  
93  
94


95 **Figure S2.** Comparison of operational TROPOMI products (TROPOMI standard and bias  
 96 correction) and machine-learning based bias correction result (ML Bias correction) with  
 97 TCCON XCH<sub>4</sub> (GGG2020 version). Shown are the mean ( $\mu$ ) and standard deviation ( $\sigma$ ) of the  
 98 TROPOMI–TCCON XCH<sub>4</sub> differences for colocated observations during 2020–2023. Station  
 99 locations are summarized in Table S1 and mapped in Fig. 1.



100


101

102 **Figure S3.** Comparison of operational GOSAT products (GOSAT standard and bias correction)  
 103 and machine-learning based bias correction result (ML Bias correction) with TCCON XCH<sub>4</sub>  
 104 (GGG2020 version). Shown are the mean ( $\mu$ ) and standard deviation ( $\sigma$ ) of the GOSAT–  
 105 TCCON XCH<sub>4</sub> differences for colocated observations during 2020–2023. Station locations are  
 106 summarized in [Table S1](#) and mapped in [Fig. 1](#).




107  
 108  
 109  
 110  
 111  
 112  
 113  
 114  
 115  
 116  
 117  
 118  
 119  
 120  
 121  
 122  
 123  
 124  
 125  
 126  
 127  
 128  
 129  
 130  
 131

132 **Figure S4.** Comparison of operational GOSAT-2 standard product and machine-learning based  
 133 bias correction result (ML Bias correction) with TCCON XCH<sub>4</sub> (GGG2020 version). Shown  
 134 are the mean ( $\mu$ ) and standard deviation ( $\sigma$ ) of the GOSAT-2–TCCON XCH<sub>4</sub> differences for  
 135 colocated observations during 2020–2023. Station locations are summarized in [Table S1](#) and  
 136 mapped in [Fig. 1](#).



137  
 138  
 139  
 140  
 141  
 142  
 143  
 144  
 145

146 **Figure S5.** Global coverage comparison between TROPOMI and fused XCH<sub>4</sub> products (2020–  
 147 2023). (a) Annual global land coverage showing TROPOMI official product (grey) and fused  
 148 product (green), with percentage improvement indicated for each year. (b) Monthly relative  
 149 coverage improvement of fused product over TROPOMI for each year from 2020–2023,  
 150 demonstrating temporal resilience during TROPOMI data gaps in July–August 2022 and  
 151 August 2023 caused by missing VIIRS cloud screening data.  
 152



153  
 154

155

156