

Supplementary materials for

Cloud vertical structure across China from a national Ka-band cloud radar network: Thermodynamic, dynamical, and land-surface controls

Hui Xu¹, Jianping Guo^{1,2*}, Jianbo Deng³, Rongfang Yang^{4,5}, Deli Meng⁶, Zhen Zhang¹, Ning Li¹, Yuping Sun¹, Shuairu Jiang¹, Tianmeng Chen¹, Juan Chen⁷, Liping Zeng⁸, Yongshui Zhou⁹, Bing Tong¹⁰

¹State Key Laboratory of Severe Weather Meteorological Science and Technology, Chinese Academy of Meteorological Sciences, Beijing, 100081, China

²CMA Field Scientific Experiment Base for Low-Altitude Economy Meteorological Support of Unmanned Aviation in Guangdong-Hong Kong-Macao Greater Bay Area, Shenzhen 518108, China

³Hunan Meteorological Research Institute, Changsha, 410118, China

⁴China Meteorological Administration Xiong'an Atmospheric Boundary Layer Key Laboratory, Xiong'an New Area, 071800, China

⁵Hebei Meteorological Technology and Equipment Center, Shijiazhuang 050022, China

⁶National Institute of Natural Hazards, Ministry of Emergency Management of China, Beijing 100085, China

⁷AVIC Leihua Electronic Technology Research Institute, Wuxi 214063, China

⁸Guizhou New Meteorological Technology Co., Ltd, Guiyang 550001, China

⁹Meteorological Observatory of Guizhou, Guiyang 550001, China

¹⁰State Key Laboratory of Regional and Urban Ecology, Research Center for Eco-environmental Sciences, Chinese Academy of Sciences, Beijing, 100085, China

Correspondence to: Dr./Prof. Jianping Guo (Email: jpguocams@gmail.com)

▲ CBH(first layer) ▼ CTH(first layer) ▲ CBH(second layer) ▼ CTH(second layer) ▲ CBH(third layer) ▼ CTH(third layer)




Figure S1. Examples of cloud vertical structure (CVS) detection for three selected cases: (a) high cloud, (b) middle cloud, and (b) low cloud.




Figure S2. Seasonal geographic distribution of cloud top height (CTH) over China in 2024, derived from the 80 MMCR stations of the same network referenced in Figure 1. The four panels correspond to (a) spring (March–April–May), (b) summer (June–July–August), (c) autumn (September–October–November), and (d) winter (December–January–February). Note that each panel presents a histogram illustrating the probability distribution of cloud occurrence frequency, with the corresponding mean value and standard deviation provided for each distribution.




Figure S3. Seasonal geographic distribution of cloud thickness (CT) over China in 2024, derived from the 80 MMCR stations of the same network referenced in Figure 1. The four panels correspond to (a) spring (March–April–May), (b) summer (June–July–August), (c) autumn (September–October–November), and (d) winter (December–January–February). Note that each panel presents a histogram illustrating the probability distribution of cloud occurrence frequency, with the corresponding mean value and standard deviation provided for each distribution.