Multi-sensor satellite analysis reveals latitudinal and morphometric controls on ice phenology across 31,000 lakes on the Alaska North Slope

Alexander L. Nguyen[1*], Cesar G. Lopez[1*], Jennifer Melara-Valle[1], Eliza Ross[1], Alexander S. Bradley[1], Maggie R. Limbeck[1], Claire C. Masteller[1], Roger Michaelides[1]

[1]Department of Earth, Environmental, and Planetary Sciences, Washington University in St. Louis, Saint Louis, MO, 63130, USA

* These authors contributed equally to this work

Correspondence to: Roger Michaelides (roger.michaelides@wustl.edu)

Abstract. Thermokarst lakes are critical components of Arctic carbon cycling, yet their ice phenology remains poorly characterized at regional scales. We present the first comprehensive analysis of ice-on and ice-off timing across 30,774 lakes on the Alaska North Slope using Sentinel-1 synthetic aperture radar (SAR) classified by a Random Forest (RF) model trained using labels derived from Sentinel-2 optical imagery and ERA5 temperature data for the period 2019–2023. Our RF classifier achieved 94% accuracy for ice state detection, enabling phenology retrieval for 97% of lake-years. Results revealed a mean ice-free period of 114 days (standard deviation = 25 days), with ice-off occurring at day-of-year 159 (June 8) and ice-on at day-of-year 273 (September 30). Spatial analysis demonstrated strong latitudinal control on ice phenology, with ice-free duration decreasing by 23 days per degree northward. Lake morphology (area, circularity, convexity, and shoreline development index) showed modest but significant effects on ice timing after controlling for latitude effects, with shoreline development index and convexity each contributing ~three days variation across typical lake ranges. Comparison of the RF model and simplistic accumulated degree-day (ADD) model-detected ice phenology yielded a convincing match, where the offsets in ice phenology between the models fell w within twice the per-detection observation gap for approximately 80% of the lakes. Furthermore, these offsets exhibited the same strong latitudinal control and negligible effects of lake morphology. These lake-specific phenology dates provide timing and duration constraints for future methane studies using high-resolution sensors and provide baseline phenology data essential for understanding how continued Arctic warming will affect thermokarst lake dynamics and associated carbon cycle feedbacks.

1 Introduction

The Arctic is warming at nearly two to four times the global average rate (Rantanen et al., 2022), with profound implications for thaw on permafrost landscapes. This Arctic Amplification (Serreze and Barry, 2011) impacts Arctic landscapes through processes such as the permafrost carbon feedback (Schurr et al., 2015), alteration of surface hydrology (Lafrenière and Lamoureux, 2019), landscape changes (Bryant et al., 2025) and ground subsidence (Streletskiy et al., 2025). Formed by ground

subsidence following permafrost thaw, thermokarst lakes cover more than 20% of the Alaskan Arctic Coastal Plain (Arp et al., 2011) and approximately 17% of Arctic lowland terrain globally (Muster et al., 2017). The majority of North Slope lakes are thermokarst in origin (Farquharson et al., 2016; Jones et al., 2022), with aeolian processes also playing an important role in lake origin and thermokarst lake evolution and orientation (Carson and Hussey, 1962). Additionally, glacial processes are another likely source of lake origin, specifically in the foothills of the Brooks Range (Detterman et al., 1958).

Thermokarst lakes represent dynamic components of high-latitude carbon cycling, where permafrost thaw surrounding these lakes mobilizes stored soil organic material (In 'T Zandt et al., 2020). These lakes emit substantial quantities of methane (CH_4) and carbon dioxide (CO_2) to the atmosphere (Walter Anthony et al., 2018), where current estimates for yearly CH_4 emissions from thermokarst lakes larger than 0.1 km^2 are 24.0 ± 8.4 Tg CH_4 yr^{-1} (Zhuang et al., 2023). Emissions are strongly modulated by lake area and depth (Heslop et al. 2020; Manasypov et al., 2023; Matthews et al., 2020; West et al. 2015) and the duration of ice melt and seasonal ice cover (Mu et al., 2025; Wik et al., 2016; Guo et al., 2020). However, the influence of lake morphometrics on ice condition and timing (i.e., duration of ice melt and seasonal ice cover) is relatively understudied.

Ice phenology - the timing of ice formation (ice-on) and breakup (ice-off) - controls the duration of open-water conditions during which gas exchange with the atmosphere occurs. Despite the importance of ice timing for understanding carbon fluxes, comprehensive regional-scale phenology data remain scarce at high spatiotemporal resolution (< 0.25 km/pixel/week) (Matthews et al., 2020). Previous studies have either focused on direct observation of ice phenology for individual lakes using *in situ* observations (Arp et al., 2015) or use single-sensor/single-modality satellite approaches for lake populations with a relatively limited number of lakes (< 1000; Latifovic and Pouliot, 2007; Surdu et al., 2014, 2016; Tom et al., 2020). The challenge of monitoring hundreds of thousands of thermokarst lakes across the remote Arctic terrain, combined with persistent cloud cover and polar night conditions, has hindered systematic phenology characterization. This technical challenge limits our ability to resolve the timing and duration of freeze-thaw transitions and detect climate-driven shifts in ice phenology, constraining efforts to quantify lake-atmosphere carbon exchange and accurately represent thermokarst lake processes in regional and global Earth system models.

Current direct methods for constraining lake ice phenology typically utilize simplistic thermal diffusion metrics such as the modified Stefan equation for accumulating degree days (ADD) (ºC · d; Arp et al., 2015; Stefan, 1890). ADD metrics are determined from air temperature records and do not account for lake-specific morphology (USACE, 2004). More complex models may incorporate lake depth, which captures ice thickness, but do not generally account for other morphologic parameters such as area, circularity, convexity, or shoreline properties, as these features are difficult to estimate without *in situ* measurements (Duguay et al., 2003). As a result, the degree to which thermokarst lake morphology may influence ice phenology relative to estimates based on ADD metrics, which do not account for morphology, remains largely unknown. Previous studies suggest that lake morphology, especially area and depth, play an important role in modulating ice phenology (Basu et al., 2024; Higgins et al., 2021; Smits et al., 2021). Specifically, the relationship between lake depth and ice thickness - which dictates whether lake ice is bedfast or floating - exerts a strong control on ice-off timing, occurring on average 17 days earlier for bedfast ice lakes than floating ice lakes in Northern Alaska (Arp et al., 2015). Recent advances in satellite remote

sensing offer new opportunities for large-scale ice monitoring (Bartsch et al. 2023). The Sentinel-1 synthetic aperture radar (SAR) operates independently of solar illumination and cloud cover, providing year-round observations at ~10-20 m spatial resolution. The distinct microwave radar backscatter signatures of ice versus open water enable automated ice detection for thermokarst lakes (Bartsch et al., 2017; Tom et al. 2020), though interpretation can be complicated by factors including ice thickness and structure, snow cover, bulk water content, and surface roughness that all impact radar scattering properties (Hall et al., 1994; Jeffries et al., 1994). However, Sentinel-2 optical imagery provides complementary information through spectral indices such as the Normalized Difference Snow Index (NDSI), which discriminates ice from water under sun-lit, clear-sky conditions (Riggs et al., 2016). NDSI can aid in SAR interpretation and fill in dates within the Sentinel-1 6 day repeat time when using both polar orbiting ascending and descending tracks. By combining these observations with temperature data from atmospheric reanalysis products, robust ice classification can be achieved across the full annual cycle, allowing for a regional-scale exploration of the morphologic controls on lake ice phenology.

In this contribution, we present a multi-sensor fusion approach to characterize ice phenology and its relationship to morphological parameters across 30,774 thermokarst lakes on the Alaska North Slope. Our objectives are to: (1) develop and validate an automated ice detection algorithm combining SAR, optical, and temperature data; (2) quantify spatiotemporal patterns in ice-on and ice-off timing across the study region relative to ADD-predicted timing; (3) examine relationships between lake morphometry and ice phenology. This analysis provides the most comprehensive characterization of regional-scale thermokarst lake ice phenology to date, establishing baseline conditions against which future changes can be assessed.

2 Methods

2.1 Study area and lake dataset

The study area encompasses the Arctic Coastal Plain (ACP) of the North Slope of Alaska (~69 °N - 71.36 °N), a ~200,000 km^2 region of continuous permafrost extending from the Brooks Range foothills to the Arctic Ocean (Fig. 1). This low-relief coastal plain is characterized by hundreds of thousands of 2 – 4 meter deep thermokarst lakes formed in ice-rich permafrost deposits (Hinkel et al. 2012). The region experiences a polar climate with mean annual temperatures of -10 to -12 °C, though summer temperatures regularly exceed 15 °C.

We obtained lake boundaries from the Alaska Lake and Pond Occurrence Dataset (ALPOD; Levenson et al., 2025), which provides high-resolution (10 m/pixel) lake polygons derived from a U-Net lake detection model (Mullen et al. 2023) applied to multi-temporal Sentinel-2 imagery across Alaska that totals in 801,895 lakes. The selected study region (69 °N - 71.36 °N, 163.85 °W - 141.02 °W) contains 128,197 ALPOD lakes. We selected all lakes ≥ 0.02 km² (2 ha) to ensure each lake had a large enough number of pixels (200 for 10 m/pixel imagery) for reliable Sentinel-1/2 detection. This selection yielded 31,108 total lakes for analysis, which encompasses the upper ~25^{th} percentile of lake area distribution prior to area filtering (Fig. S10). From this, we were able to retrieve phenology from 30,774 lakes (98.9%), where lake morphometrics could be extracted using lake polygons from the ALPOD database.

For the subpopulation of lakes used in this study, lake area spans five orders of magnitude from 0.02 to 859 km² (median = 0.065 km²), with the upper bound corresponding to Teshekpuk Lake. Study lakes had a range of circularity, defined as $C = 4\pi A/P^2$, where P is lake perimeter (m), from 0.01 – 0.78, with smaller values indicative of more oblong lakes. Lake shoreline development indices, ($SDI = P/2\sqrt{\pi A}$; Hutchinson, 1957), which quantifies shoreline complexity, range from 1.13 – 10.02, with larger values indicating more complex shorelines. Measures of lake convexity range from 0.11 - 0.998, where values of 1.0 indicate more convex shorelines and values closer to zero signify higher concavity or shoreline irregularity (Bozeman and Pilling, 2013). Calculating both SDI and convexity allows for a better understanding of both shoreline complexity and irregularity, where SDI is more impacted by fractal dimensionality from higher resolution data and increases with increasing area of lakes (Seekell et al. 2022). Lake circularity, ~~SDI, and~~ convexity, and SDI for the subpopulation of lakes used in this study were consistent with metrics from the full ALPOD database (Fig. S10).

Formatted: Pattern: Clear




Figure 1: Study area (69 °N - 71.36 °N, 163.85 °W - 141.02 °W) showing the 31,108 lakes with area ≥ 0.02 km² selected for analysis, of which 30,774 (blue) yielded ice phenology. Lakes for which phenology was not successfully detected are shown in bronze. Inset depicts ~~mapped location~~lake details within ~~Alaska~~the study region. Sources: ALPOD and the Natural Earth.

2.2 Satellite data acquisition and preprocessing

We acquired Sentinel-1 Ground Range Detected (GRD) imagery from Google Earth Engine (GEE) (Gorelick et al., 2017) for the period January 2019 through December 2023. We selected this time scale based on the availability of the Harmonized Sentinel-2 MSI: MultiSpectral Instrument, Level-2A (SR) collection from Google Earth Engine. 2019 was selected as the start year, as the 2017-2018 data does not have global coverage in this collection. Using 2023 as the final year for this data date range ensured that there were at least five years of data to retrieve lake ice phenology from, where we sought to define a nominal five-year time span over which to perform this analysis. The loss of Sentinel-1B on December 23rd, 2021, affected our analysis in increasing the repeat time (from 6 days to 12 days) between acquisitions over our study area, thereby effectively decreasing the density of measurements after data transmission failure of the satellite.

We used images from both ascending and descending track satellite orbits acquired in Interferometric Wide Swath mode, providing both vertical transmit/receive (VV) and vertical transmit/horizontal receive (VH) polarizations at 10 m resolution. Images were accessed from the GEE Sentinel-1 GRD collection, which provides data pre-processed using the standard Sentinel-1 Toolbox workflow including thermal noise removal, radiometric calibration, and orthorectification terrain correction (Mullissa et al., 2021). We additionally applied an incidence angle mask (25–50°) to reduce edge effects, and we subtracted VH from VV images to generate VV/VH images. We generated standard falsecolor RGB composites of backscatter (i.e., red: VV, green: VH, blue: VV/VH) (see supplemental figure ~~S12~~S1 for the entire workflow). To account for typical Arctic lake backscatter ranges, we scaled each band (in order) for open water to rough ice (-20 to –5 dB; VV), volumetric scattering cross-polarization range (-28 to –12 dB; VH), and the polarization ratio (8 to 18 dB; VV/VH). We scaled each band to target specific dB ranges that aid in differentiation between ice and water as well as potential ice type/thickness. By scaling the data, we target these specific dynamic ranges to maximize differentiation between the expected minimum and maximum expected backscatter values for the ice/water distinction for a given band. Additionally, scaling the data reduces the impact of extraneous outliers outside of the expected backscatter ranges for each band (Surdu et al. 2015; Murfitt and Duguay 2020).

We obtained Sentinel-2 Level-2A surface reflectance imagery for the same time period between 2019 – 2023, where our study period date range is limited by Sentinel-2 Level-2A products being unavailable for 2017-2018. We then filtered to acquisitions with <80% scene cloud cover and pixel-level cloud and cloud shadow masking using the s2cloudless algorithm with a 20% cloud probability threshold (Sentinel-Hub, 2025; Skakun et al., 2022). Sentinel-2 coverage varies spatially across the study area due to orbital geometry at ~70 °N, with reduced coverage between –153 °W and –157 °W longitude; consequently, S2 observations serve as high-confidence training labels where available rather than as the primary detection method. We calculated the Normalized Difference Snow Index (NDSI) as NDSI = (Green − SWIR) / (Green + SWIR) using bands B3 (560 nm) and B11 (1610 nm) (Hall et al. 1995), and we classified NDSI values > 0.4 as snow/ice following established thresholds (Riggs et al., 2016). The output of the NDSI is a binarized classification that is later used to calculate ice fraction, as explained in section 2.3.

We obtained ERA5 hourly 2 m air temperature data at 0.25° (~28 km/pixel) resolution from the Google Cloud Analysis-Ready, Cloud-Optimized (ARCO) ERA5 archive (Carver and Merose, 2023; Herbach et al. 2017; Hersbach et al., 2020) and aggregated to daily means for each lake centroid location. Prior work has demonstrated excellent agreement between ERA5 and in situ weather stations across the Arctic (Clelland et al. 2024; Ibebuchi et al. 2024; Tarek et al. 2020; Pernov et al. 2024). For example, Clelland et al. 2024 has shown agreeance of ERA5 with in-situ data in Siberia, Ibebuchi et al. 2024 find ERA5 outperforms other reanalysis products for comparison with in-situ data across the contiguous United States, Tarek et al. 2020 found ERA5 hydrological modeling was comparable to in-situ data across North America, and Pernov et al. 2024 found a 0.94 R^2 for temperature when comparing ERA5 to 17 weather stations across the Arctic. Given the agreement of prior work of validating ERA5 in the Arctic, we use ERA5 2 m air temperature data for this study over the ACP.

2.3 Ice detection algorithm


Our ice detection approach uses a Random Forest (RF) classifier (Breiman, 2001) to predict ice state from Sentinel-1 SAR backscatter features. This approach follows machine learning methods developed for lake ice classification from SAR imagery (Hoekstra et al., 2020; Tom et al., 2020), adapted for regional-scale application. Due to variations in liquid water content, and therefore dielectric permittivity and radar backscatter (Unterschultz et al, 2009), during phase changes, it is particularly difficult to classify SAR imagery for phenology during the shoulder season (Posch et al., 2024). To mitigate the effects of these factors on ice detection, we introduce a series of temperature-based thresholds, consistency checks, and consecutive occurrences. First, we derived training labels from two sources: ERA5 temperature reanalysis and Sentinel-2 optical imagery. Sustained temperature periods (10+ consecutive days below −20°C or above +10°C) provided high-confidence labels during winter months and polar night when optical data were unavailable, comprising approximately 62% of training labels. Sentinel-2 NDSI classifications provided the remaining labels during cloud-free conditions (30% cloud threshold), with temperature-based quality control to exclude potentially erroneous detections (e.g., S2 labels flagged as water when air temperature was below −5°C, or as ice when temperature exceeded +5°C, were excluded to filter algal bloom misclassifications). This dual-source approach maximizes temporal coverage across the full annual cycle while maintaining label reliability.

We trained the RF model on six Sentinel-1-derived features: VV and VH backscatter (dB) measured over the lake interior (extracted from removing a 10 m interior buffer applied to each lake feature perimeter), VV and VH backscatter over a 100 m marginal landscape surrounding each lake (extracted from adding a 100 m exterior buffer applied to each lake feature perimeter), and the VV/VH polarization ratio for both lake and landscape zones. The classifier was trained using mean ice fraction values (0 – 1) from both the interior and exterior lake buffers, applied to the Sentinel-2 NDSI-derived ice labels and the corresponding Sentinel-1 training data. A given lake-date is labeled as either ice or water using a threshold of ice fraction > 80% being ice, and an ice fraction < 20% being water. As a result, the approach does not explicitly capture spatial variations in ice growth or melt within a lake, except to the degree that they are reflected in the buffer-averaged values. High values of VV co-polarization record rough ice, whereas low values record smooth ice and water. VH cross-polarization is sensitive to

volumetric scattering from vegetation, and the ratio between co- and cross-polarizations aids in determination between types of ice. Furthermore, the landscape features encapsulated through the exterior buffer capture contextual information about surrounding terrain conditions that may aid in distinguishing ice from open water (e.g., freeze/thaw state of the landscape).

Training data comprised 1.09 million lake-date observations from 24,886 lakes. An additional 274,079 observations from 6,222 held-out (validation) lakes were reserved for testing. This lake-based splitting strategy, implemented using grouped cross-validation, ensures no lake appears in both training and test sets, preventing data leakage from temporal autocorrelation within individual lake time series. We implemented the RF model using scikit-learn (Pedregosa et al., 2011) with 100 trees, a maximum depth of 15, a minimum split of 20 samples, and a minimum of 10 samples for leaves.

~~The RF classifier achieved 94% overall accuracy when evaluated against Sentinel-2 NDSI labels from the held-out/validation lakes not used in training (n = 106,527 observations from 2,322 lakes). Performance was higher for the ice class (precision = 0.97 recall = 0.95) than the water class (precision = 0.86, recall = 0.93), reflecting both the class imbalance in training data (65% ice, 35% water owing to the shorter ice-free season) and the greater difficulty of detecting open water conditions (Fig. 2A). The cross-validated F1 score was 0.977 ± 0.001. Feature importance for the RF model was consistently higher for the land buffers surrounding the lakes for VH (green), VV (red), and VV/VH ratio (blue), respectively, compared to the lake interiors (VV, VH, and VV/VH ratio), which may indicate true importance or potential bias in the number of pixels used in the land vs lake buffers (Fig. 2B). Accuracy estimates reflect agreement with optical classifications; no independent validation against field observations was performed. During inference, ice state is determined solely by the RF classifier applied to Sentinel-1 features; temperature is not used as a classification input, preserving independence for subsequent analysis of temperature-phenology relationships.~~



~~Figure 2: A) Random forest classification confusion matrix, where numbers within each box represent number of samples falling within each category. The boxes with darker saturation indicate where the RF model predicted the correct label based on the true~~

~~label, where correct water and ice predictions are represented by medium blue and dark blue, respectively. The light blue/gray boxes indicate where the RF model predicted incorrectly. B) Random forest feature importance using Gini, where land and lake refer to the exterior and interior lake buffers, respectively. VV, VH, and VV/VH ratio refer to the polarimetric SAR band, and db represents decibels.~~

205

2.4 Phenology extraction

Using the RF model predictions, we output the probability that an individual lake-date is covered with ice, *P*(ice). We determined the ice-on and ice-off dates from this classified ice probability time series, *P*(ice), for each lake-year. To avoid detecting spurious mid-winter thaws or brief summer re-freezing events, we constrained transition detection using hysteresis
210 logic, where a lake-date with *P*(ice) > 0.7 was labeled as ice and *P*(ice) < 0.3 as water. Lake-dates that have *P*(ice) greater than 0.3 but less than 0.7 are determined ambiguous.

We then use the date prior to transition with the first date post-transition to interpolate the day in which the bimodally-distributed *P*(ice) crosses 0.5 (as determined by the symmetric distribution around the consistently global trough at 0.5). To prevent logical inconsistencies in the detected phenology (e.g. negative ice free and ice-covered periods), we only defined ice-
215 off if it started and is observed for a given year. This is implemented in the case that ice-off for an individual lake may have been detected as open throughout the prior winter season, where we remove that given lake-year's phenology for the current year's ice-off. Functionally, ice-off is the first date when the lake transitioned from frozen to open water state and remained classified as water for at least two consecutive observations and satisfies the imposed hysteresis logic. Similarly, we defined ice-on for a lake as long as it started and is observed for a given year. Otherwise, the lake will have frozen into the next year's
220 late winter/early spring outside of that given year. In such cases, as for ice-off, we remove and exclude that lake-year's relevant phenological event. Likewise, ice-on is the first date of the subsequent freeze-up when the lake transitioned to frozen state and remained classified as ice for at least two consecutive observations. With both ascending and descending Sentinel-1 orbits providing a median observation gap of 4 days at this latitude, this criterion typically requires sustained state persistence of approximately 8–16 days, filtering transient events while preserving sensitivity to true phenological transitions.
225 We then calculated the ice-free period as the number of days between ice-off and ice-on for each lake-year. We assigned null values to lakes lacking clear transitions due to data gaps, persistent cloud cover, or ambiguous classifications and excluded them from phenology statistics. We computed summary statistics including mean, median, standard deviation, and interquartile range across all valid lake-years, as well as stratified by geographic subregion and lake size class.

2.5 Phenology model comparison

230 To compare the SAR RF model with other methods for ice phenological timings, we used ERA5 daily average 2 m temperatures (Carver and Merose, 2023; Herbach et al. 2017; Hersbach et al., 2020) to calculate a continuous time series of both the accumulated degree day of thaw (ADDT) and accumulated degree day of freezing (ADDF) for ice thickness (Arp et al., 2015; Stefan, 1890).Accumulated Degree Days of Thaw (ADDT) and Accumulated Degree Days of Freeze (ADDF)

were calculated from daily mean air temperatures (T_i) on day i as cumulative sums of positive and negative temperatures, using equations 1 and 2 respectively:

$$ADDT = \sum_{i=1}^{n} \begin{cases} T_i, & T_i > 0 \\ 0, & T_i \leq 0 \end{cases} (1)$$

$$ADDF = \sum_{i=1}^{n} \begin{cases} -T_i, & T_i < 0 \\ 0, & T_i \geq 0 \end{cases} (2)$$

We calculated both of these metrics for each lake-year combination and used a seven-day switch window between freeze-thaw/thaw-freeze to ensure a sufficient duration for cumulative ice growth/melt before switching between ADDF and ADDT. We then converted from the accumulating degree days of freeze to ice thickness, h_{ice} (cm), using equation 3:

$$h_{ice} = \alpha\sqrt{\overline{ADDF}} (3)$$

Where α is the Stefan equation coefficient, with a chosen value of 2. During the seven-day switch window, ice thickness was estimated using equation 4:

$$h_{ice} = \alpha\left(\sqrt{\overline{ADDF}} - \sqrt{\overline{ADDT}}\right) (4)$$

During the seven-day switch, ice thickness values were used if positive, if $h_{ice} \leq 0$, ice thickness was set to zero to avoid negative ice thicknesses. For similar reasons, ice thickness was set to 0 for the ADDT period. We utilize this method because ADD metrics only require daily air temperatures and the empirically derived α coefficient. This approach is advantageous across our large study where many of the variables utilized in more complex thermodynamic models are simply unavailable or poorly constrained given the limited in-situ data that exists across the North Slope. This method also allows us to capture first order patterns while remaining computationally efficient, which is the primary objective of this study. Consequently, the model provides robust first-order estimates of spatial and temporal variability in ice thickness while avoiding the substantial uncertainties associated with poorly constrained thermodynamic parameters. For inter-model comparison, we used the SAR ice-off and ice-on dates from the transition-based detection described in Sect. 2.4, and defined the ADD thaw and freeze onsets as the start and end of the longest contiguous thawing interval of each year (the principal warm season). Because neither estimate relies on a fixed calendar window, the SAR–ADD comparison is independent of the SAR detection method. We then computed the offsets between the SAR and ADD models by subtracting the ADD from the SAR dates, where positive offsets indicate the SAR model predicting later than the ADD model and vice versa. Finally, we employ the Kruskal-Wallis H-test, Mann-Whitney U-test, and Cliff's Delta (δ) to statistically evaluate any potential controls of latitude and lake morphometry on discrepancies with the RF model and ADD model estimates of ice-off and ice-off dates

3 Results

3.1 Random-forest-derived ice phenology across the Arctic Coastal Plain

Formatted: Font: Times New Roman, 10 pt, Bold, Font color: Auto

The RF classifier achieved 94% overall accuracy when evaluated against Sentinel-2 NDSI labels from the held-out/validation lakes not used in training (n = 106,527 observations from 2,322 lakes). Performance was higher for the ice class (precision = 0.97 recall = 0.95) than the water class (precision = 0.86, recall = 0.93), reflecting both the class imbalance in training data (65% ice, 35% water owing to the shorter ice-free season) and the greater difficulty of detecting open water conditions (Fig. 2A). The cross-validated F1 score was 0.977 ± 0.001. Feature importance for the RF model was consistently higher for the land buffers surrounding the lakes for VH (green), VV (red), and VV/VH ratio (blue), respectively, compared to the lake interiors (VV, VH, and VV/VH ratio), which may indicate true importance or potential bias in the number of pixels used in the land vs lake buffers (Fig. 2B). Accuracy estimates reflect agreement with optical classifications; no independent validation against field observations was performed. During inference, ice state is determined solely by the RF classifier applied to Sentinel-1 features; temperature is not used as a classification input, preserving independence for subsequent analysis of temperature-phenology relationships.





Figure 2: A) Random forest classification confusion matrix, where numbers within each box represent number of samples falling within each category. The boxes with darker saturation indicate where the RF model predicted the correct label based on the true label, where correct water and ice predictions are represented by medium blue and dark blue, respectively. The light blue/gray boxes indicate where the RF model predicted incorrectly. B) Random forest feature importance using Gini, where land and lake refer to the exterior and interior lake buffers, respectively. VV, VH, and VV/VH ratio refer to the polarimetric SAR band, and db represents decibels.

Ice phenology was successfully retrieved for 30,774 lakes (98.9% of the study population) for at least one year, with 94.17% having complete five-year records. For the 1.1% of lakes that do not have detected phenology for at least one year, their overall

area probability density distribution and median exhibit no discernible difference between the total population beyond the large difference in the number of samples (Fig. ~~S11~~S12). However, the probability density of SDI and convexity for these lakes without phenology have a medium degree of separability (Cohen's *d* of 0.57 and 0.52, respectively) compared to the total population (Fig. ~~S11~~S12). Furthermore, the lakes without phenology have a higher median SDI value of 2.1 compared to the total lake population of 1.68 which indicates the lakes lacking phenology estimates are comparatively more complex. Across all lakes and years, mean ice-off occurred on day-of-year (DOY) 159 (June 8; standard deviation = 13 days), where the 2σ range is DOY 132 to DOY 185. Mean ice-on occurred on DOY 273 (September 30; standard deviation = 17 days), where the 2σ range is DOY 237 to DOY 308. The resulting ice-free period averaged 114 days (standard deviation = 25 days), with individual lake-years spanning 3 to 331 days (Fig. ~~S1~~S2). Interannual variability in ice timing was substantial (Figs. 3 and S3). The earliest regional mean ice-off occurred in 2019 (DOY 157), while the latest occurred in 2022 (DOY 164) (Fig. ~~S3~~S4). Likewise, the earliest regional mean ice-on occurred in 2021 (DOY 266), while the latest occurred in 2019 (DOY 278). Ice-free duration ranged from a mean of 105 days in 2022 to 122 days in 2019 (Fig. ~~S3~~S4). Across all lake-years, ice-off and ice-on timing were moderately negatively correlated ($r = -0.49$) across complete paired lake-dates ($n = 150{,}942$), largely reflecting latitudinal gradients in both variables (Fig. 5).


Ice Phenology Distributions (2019-2023)
Ice-off: 159.0 ± 12.9 days (median 160)
Ice-on: 272.8 ± 16.6 days (median 272)
n = 150,942 lake-years
Ice-off / Ice-on
2019
2020
2021
2022
2023
Number of Lake-Years
Day of Year
0
5000
10000
15000
20000
25000
30000
0
50
100
150
200
250
300
350




Figure 3: Distribution of SAR RF model-detected ice phenology metrics across all lake-years (2019–2023), where increasing saturation corresponds to more-recent years. The red histogram indicates ice-off date, and the blue histogram indicates ice-on date.

Comparison of the ice phenology events derived from the RF model with the ADD metric shows relatively strong correspondence (Fig. 4). There were an identical number of detections valid for comparison for both thaw and freeze. Likewise, the median gap between phenology derived from the RF model and ADD metric was the same at roughly 10 days, varying spatially and after the loss of Sentinel-1B. For ice-off dates, 82% of lakes align with ADD estimates within twice the per-detection observation gap ~~(median 10 days, varying spatially and after the loss of Sentinel-1B),~~ and for ice-on dates, 75% of lakes align within twice the observation gap (roughly 12 days, but varying spatially and after loss of Sentinel-1 B) (Fig. 4). The median gap between the SAR detected ice-on or ice–off date and corresponding ADD metric was 4 days for thaws and 6 days for freezes (Fig. ~~S4~~S5). For ice-off dates, the RF model was earlier than the ADD estimate for 37% of the lake-years, later than the ADD estimate for 60% of the lake-years and selected within six days for 40% of the lake-years, suggesting a modest tendency toward delayed thaw detection relative to cumulative degree-day predictions (Fig. ~~S6~~S7). For ice-on dates, the RF model was earlier than the ADD estimate for 77% of the lake-years, later than the ADD estimate for 21% of lake-years and selected within six days for 38% of lake-years, reflecting predominately earlier calls from the SAR model during lake

freeze-up compared to the ADD model (Fig. ~~S6~~S7). See supplemental figure S13 for a visual representation of the SAR RF model ice phenology performance of a randomly-selected lake.




Figure 4: Histograms of the offset between ice phenology timings from SAR detection and the ADDT (a) and ADDF (b) algorithm. Negative values indicate SAR detects dates before ADDT/F, while positive values indicate SAR detects dates after the ADD metric. The histograms are shaded by the per-detection observation gap (i.e., % within 1 observation gap, within 2 observation gaps), where increasing saturation indicates data within shorter periods. The dashed vertical line is at zero which indicates where there is zero offset between the SAR and ADD phenology.

For all lake-years, for ice-off dates, although statistically significant differences exist in lake morphometrics between cases where the RF model was earlier versus later than the ADD metric (H- and U-test $p << 0.05$), the effect sizes are very small ($\delta < 0.15$). This indicates that lake morphology does not play a significant role in explaining the observed offset between RF and ADD ice-off estimates. In contrast, latitude has a more significant effect size (H- and U-test $p << 0.05$; $\delta = -0.3105$), indicating that latitudinal effects likely explain some of the observed mismatch between RF and ADD ice-off timing. Lakes that the RF model identifies as ice-free earlier than the ADD estimate are generally found at lower latitudes while lakes where the RF model identifies the ice-free date later than the ADD estimate are generally located at higher latitudes. Similarly, for ice-on dates across all lake-years, morphology again exhibits statistically significant differences between groups, but with very weak effect sizes (H- and U-test $p << 0.05$; $\delta < 0.15$), while latitude has a stronger effect ($\delta = 0.557$).

Lakes identified by the RF model as ice-on earlier than the ADD estimate occur at higher latitudes, whereas those identified as ice-on later tend to lie farther south within the study region. Consistently, lakes classified as ice-on earlier by the RF model are substantially farther north than lakes classified as ice-off earlier ($\delta = -0.3164$), and lakes classified as ice-on later are located at higher latitudes than lakes classified as ice-off later ($\delta = 0.5517$).

For lake-years where the offset between the RF model ice-on and ice-off dates and the ADD estimates is greater than 2 revisit gaps, the results are very similar. Within ice-off and ice-on dates, and comparisons between the two dates, morphometrics variables feature statistically significant differences between the groups but weak effect sizes (H- and U-test $p << 0.05$; $\delta < 0.18$) for differences between the RF model and ADD estimate dates. Latitude, however, continues to be an important variable for offsets between the RF model and ADD estimates. Within ice-off dates, the latitudinal statistical significance and large effect size between groups (H- and U-test $p << 0.05$; $\delta = -0.5687$) indicates that the RF model selects ice-off dates earlier than the ADD estimates at lower latitudes while selecting ice-off dates later than ADD estimates at higher latitudes. For ice-on dates, the RF model detects ice later than the ADD estimates ($\delta = 0.6378$) at lower latitudes and detects ice earlier than the ADD estimates at higher latitudes. Lakes where the RF detects ice-off earlier than the ADD estimates occur at lower latitudes ($\delta = -0.5182$) while lakes where the RF model detects ice-on earlier than the ADD occur at higher latitudes. Lakes where the RF detects ice-off later than the ADD estimates occur at higher latitudes ($\delta = 0.6813$) and lakes where the RF model detects ice-on later than the ADD estimates occur at lower latitudes.


3.2 Spatial patterns and latitudinal gradients in lake phenology


Ice phenology exhibited strong spatial structure across the study region (Fig. 5). Latitude was a dominant control on ice timing, with ice-free period decreasing by roughly 30 days per degree northward across all lakes ($r = -0.50$, $p < 0.001$; Fig. 5c, Fig. 6a); after excluding lakes with residuals outside the 5th–95th percentile, this gradient is 23 days per degree ($r = -0.46$; see below). This gradient reflects the combined effects of decreasing summer temperatures and shorter warm seasons with increasing latitude. Latitude correlations were strongest for ice-off timing ($r = 0.52$) and ice-free duration ($r = -0.50$), and weaker for ice-on timing ($r = -0.36$) (Fig. ~~S9~~S10). Diagnostic tests indicated significant residual spatial autocorrelation

Formatted: Left

(Moran's I = 0.35–0.45) reflecting unmeasured spatially-varying factors such as elevation, microclimate, permafrost properties, surface/subsurface hydrology, and topographic shading (Fig. ~~S5a~~S6a). Spatial Error Models (SEM) were therefore fit using generalized moments estimation with K-nearest neighbour spatial weights (k=8). The spatial error parameter $\lambda \approx 0.59$ - 0.66 confirmed strong spatial autocorrelation from unmeasured regional factors, and filtered residuals showed negligible spatial autocorrelation (Moran's I ≈ -0.004 to -0.02), indicating adequate model specification (Fig. ~~S5b~~S6b). Further, the residuals feature a unimodal normal distribution centered around 0, with no additional peak present for lakes in the foothills of the Brooks Range (Fig. 5c). Nonetheless, we also exclude all lakes with residuals outside of the 5th-95th percentile in our analysis, which serves to remove residuals greater than 35 days primarily found in the foothills of the Brooks Range (Fig. ~~S5d~~S6d). Latitude remains a key control on ice timing even after this exclusion, with the ice-free period

decreasing by 23 ~~days per~~daysper degree northward ($r = -0.46$, $p < 0.001$; Fig. 5b).





375 **Figure 5: Spatial patterns in mean SAR RF model-detected ice phenology (2019–2023): (a) ice-off date (day of year), (b) ice-on date, and (c) ice-free duration. Each point represents one lake.**





Figure 6: Relationship between latitude and ice-free duration, as determined by the SAR RF model. The black line shows the linear regression best-fit~~.~~ on the raw data. White circles mark the median ice-free days within 0.25° latitude bins, and horizontal bars show the interquartile range (25th–75th percentile); bins with fewer than 30 observations are omitted. Shading shows the number of lake-year observations in each cell of a 40 × 40 grid spanning the range of ice-free days and latitude. a.) All lakes for which phenology was extracted and b.) all lakes with residuals in the 5th-95th percentile. Latitude is displayed on the y-axis for a map view of ice-free duration and is the independent variable in this regression.

3.3 Morphometric relationships

Morphometric variables showed negligible correlations with mean spring temperature ($|r| \leq 0.11$ for ~~SDI,~~ circularity, convexity, SDI, and log-transformed area), confirming that morphometry is not confounded with spatial climate gradients. To assess the strength of morphometric controls on ice phenology, we modelled ice-off date, ice-on date, and ice-free duration as functions of mean spring temperature (April–June) and morphometric parameters using ordinary least squares regression. Lake area shows a statistically significant, but negligible correlation with ice-off and ice-on dates and ice-free duration (Fig. 7A-C), indicating that although the relationship is detectable given the large sample size, it explains minimal variation in phenology. Lake circularity, convexity, and SDI exhibit very weak but statistically significant correlations with ice-free duration~~, compared to the negligible association of ice-free duration and convexity that does not exhibit statistical significance (Fig. 7F).~~. Overall, morphometric correlation (area, circularity, ~~SDI, and~~ convexity, and SDI) with ice phenology (ice-off date, ice-on date, and ice-free duration) is weak ($< \pm 0.17$) (Fig. 8). Morphometry coefficients attenuated by approximately 20–25% under spatial correction but remained highly significant ($p<0.001$), confirming that morphometric effects on ice phenology are robust to spatial confounding. Multiple regression with temperature as a covariate showed that morphometric parameters explained an additional 1.5–2% of variance in ice phenology beyond temperature alone. OLS residuals showed significant spatial autocorrelation, so ~~Spatial Error Models~~SEMs were used; all morphometric predictors remained significant after this correction, with shoreline development index (SDI) and convexity showing the largest effects.

Effect sizes using the SEM coefficients were modest relative to temperature and interannual variability. Lakes with more complex shorelines (higher SDI) experienced shorter ice-free seasons (−4.3 days across the interquartile range), as did lakes with ~~lower~~higher convexity (−3.5 days). We note that the signs for correlation between morphometric variables and phenology retrievals all stayed the same apart from convexity and ice-free period, where the correlation coefficient is positive (Fig. 7e) prior to using the SEM and negative after accounting for spatial autocorrelation. Lake area and circularity showed smaller effects (+2.7 and +2.4 days, respectively) (Fig. 9). ~~Increasing~~Using the entire data distribution, increasing shoreline complexity and decreasing lake convexity shortens the ice-free period; while increasing lake area and circularity ~~these~~ lengthens the ice-free period~~.~~ (Fig. 7).


A)
Ice-off Date vs Lake Size
Spearman r=0.073
p=8.99e-177
Ice-off DOY
Lake Area (km²)
B)
Ice-on Date vs Lake Size
Spearman r=0.101
p=0.00e+00
Ice-on DOY
Lake Area (km²)
C)
Ice-free Duration vs Lake Size
Spearman r=0.023
p=9.22e-19
Ice-free Period (days)
Lake Area (km²)
D)
Ice-free Duration vs Circularity
r=0.131
p=0.00e+00
Ice-free Period (days)
Circularity
E)
Ice-free Duration vs Shoreline Development Index
r=-0.143
p=0.00e+00
Ice-free Period (days)
Shoreline Development Index
F)
Ice-free Duration vs Convexity
r=0.079
p=6.41e-209
Ice-free Period (days)
Convexity





Figure 7: A) Ice-off dates vs lake area (km2) for all lake-years. B) Ice-on dates vs lake area for all lake-years. C) Ice-free duration vs lake area for all lake-years. D) Ice-free period (days) vs lake circularity for all lake-years. E) Ice-free period vs lake ~~SDI for all lake-years. F) Ice-free period vs lake~~ convexity for all lake-years. F) Ice-free period vs lake SDI for all lake-years. Ice phenology data here is determined by the SAR RF model. White circles denote the binned median metric and horizontal bars show the interquartile range (25th–75th percentile); bins with fewer than 30 observations are omitted.


Correlation Matrix: Morphometry vs Phenology
log_area
circularity
sdi
convexity
centroid_lat
ice_off_doy
ice_on_doy
ice_free_days
1.00
-0.07 1.00
0.07 -0.88 1.00
0.17 0.84 -0.86 1.00
0.06 -0.04 0.04 0.03 1.00
0.06 -0.09 0.10 -0.03 0.52 1.00
0.07 0.13 -0.14 0.10 -0.36 -0.49 1.00
0.01 0.13 -0.14 0.08 -0.50 -0.82 0.90 1.00
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
−0.75
−1.00

420




Figure 8: Correlation matrix between lake morphometric variables (log transformed area, circularity, SDI, convexity), latitude (centroid latitude), and SAR RF model-detected ice phenology (ice-off dates, ice-on dates, ice-free days).

Overall, morphometry plays a subordinate role to temperature in determining ice phenology at the regional scale, with temperature explaining 13-19% of variance compared to 1.5–2% for morphometric parameters. The relatively modest variance explained by temperature reflects the spatial autocorrelation structure in the data: much of the temperature signal is captured by latitude, which is implicitly included through the spatial error model. The spatial error parameter confirms that regional factors beyond point-level temperature such as coastal proximity, local topography, and regional weather patterns contribute substantially to ice timing variability.




Figure 9: Spatially corrected effect of morphometric variables on SAR RF model-detected ice-free duration, calculated across the interquartile range.

4 Discussion

4.1 Model Performance

While previous studies achieved high accuracy (95.8%) using sophisticated segmentation algorithms and extensive texture features on individual lakes (Hoekstra et al., 2020), such approaches are computationally prohibitive for analysis of tens of thousands of water bodies. Our simplified RF framework using six backscatter features achieves 94% accuracy while enabling consistent classification of ~30,000 lakes across multiple years, including during polar night and cloudy conditions when optical data are unavailable, demonstrating the feasibility of regional-scale ice phenology monitoring from satellite observations. Previous single-sensor SAR studies typically achieved accuracies on the order of ~80–85% (Surdu et al., 2014). The use of Sentinel-2 optical data to generate training labels for the RF classifier, combined with temperature-based quality control, enabled reliable classification even during the challenging transition seasons. The RF model feature importance (Fig. 2B) further shows that using additional contextual information, such as land buffers surrounding the lakes, increased overall RF model accuracy compared to solely relying on lake pixels for ice-state determination (Tom et al., 2020).

The separability of the probability density functions for lakes without ice phenology compared the total population for SDI and convexity, but not area (Fig. ~~S11~~S12), is likely reflective of complex shorelines negatively impacting our ice phenology detection. For example, our algorithm relies on interior buffers that may not perform as well when shorelines are increasingly complex. However, convexity and SDI are likely not the sole reason nor explanation for the phenology not being detected for these lakes, where there are other plausible mechanisms that could explain our algorithm failing for these lakes. The mean ice-free period of 114 days is consistent with limited in situ observations available for the region. Arp et al. (2015) reported ice-free periods of 80–110 days for instrumented lakes on the Arctic Coastal Plain during 2004–2013. Our results extend these findings across the full thermokarst lake population of roughly 30,000 lakes. We do note that there are a small number of statistical outlier lakes (outside ±2σ) that exhibit phenological events that are physically improbable compared to climate data. For the ice-off metrics, we find that there are 6,325 lake-date outliers (4.17% of total lake-dates). Likewise for ice-on, there are 8,641 lake-date outliers (5.27% of total lake-dates). Here, we note that these outliers comprise approximately 5% of data and are not representative of the total aggregate population of lake-dates.

The bias in offsets we observe in the ice-off and ice-on dates between the RF model and the ADD estimates are likely due to inherent assumptions made in the RF and ADD models, as well as spatial scale differences in data resolutions (~10 m/pixel for Sentinel-1 vs ~30 km/pixel for ERA5). Additionally, the loss of Sentinel-1B on December 23rd 2021 reduced the density of repeat passes and coarsened the granularity of the ice-on/off day-of-year estimates following its failure. However, the RF and ADD models have fair agreeance, but they also exhibit inter-model biases that manifest differently during thaw and freeze. We do not necessarily interpret the offsets between the RF and ADD models as errors, but rather as the expected lag between the onset of freezing or thawing and the lake reaching a fully ice-covered or ice-free state and inherent differences between the accuracy of both models.

The accuracy of the RF model implicitly depends upon the assumed polarimetric scattering model, wherein changes in radar backscatter within and between polarization channels are assumed to be driven by the systematic difference in polarized backscatter between ice and liquid water (Jeffries et al. 1994; Beaven 1995). In reality, dielectric permittivity changes due to partial surface melting, the presence of snow, changes in snow water equivalent, and ice metamorphism can all introduce changes to SAR backscatter that can complicate the interpretation of a simple water phase transition (Paul et al., 2017; Marin et al., 2020; Lund et al., 2022; Veijola et al., 2024). These confounding effects can be particularly apparent during the shoulder seasons making satellite observation of ice phenology difficult (Posch et al., 2024). Additionally, it is possible that the Sentinel-1 descending track (~61% of scenes) images, which are acquired between 7:00-9:00 AM AKST, may also introduce variability in predicted ice phenology with the RF model due to diurnal freeze/thaw processes during the shoulder seasons, as compared to the ascending track images acquired between 6:00-8:00 PM AKST (Fig. ~~S7~~S8).

Likewise, the use of ADD to model thermal diffusion through the ice cover implicitly assumes a purely one-dimensional diffusive heat transfer at the air-water surface interface, neglecting lateral heat transfer or wind-driven thermal advection. Additionally, the ADD model does not account for thermal mixing that occurs before ice formation begins. Larger and deeper lakes take longer to mix, which delays ice formation. We expect that the ADD model would likely estimate earlier ice-on dates than when ice formation may actually begin as a result. We further do not consider the effect of snowpack cover, which serves to delay ice-off timing, due to the lack of sufficiently accurate, high spatial resolution snow depth and snow water equivalent products. The 40 cm mean snowpack on the Arctic Coastal Plain reported in Stuefer et al. 2025 is significant enough to insulate the soil and delay ice-off timing.

Consequently, the RF model may be capturing this thermal insulation and relative delay in ice-off timing compared to the ADD model, which is consistent with the observed lag in RF model timing compared to the ADD model for ~~72~~60% of lake-year combinations. Our deliberate choice of a relatively simple empirical model might explain some of the observed offset between RF and ADD model ice-on and ice-off calls, particularly during ice thaw when the presence of snowpack can considerably impact the timing of ice-off (Pouw et al., 2023). This epistemic uncertainty propagates into both models' abilities to predict exact dates for ice phenology.

To minimize the potential for an early ice-off bias, our analysis relies on observations from the lake interior rather than the shoreline, where ice loss is expected to occur first because adjacent land surfaces warm more rapidly than the frozen lake. Although ice drifting toward the shore is possible, the interior edge buffer used in this study is only 10 m wide, corresponding to a single Sentinel-1/2 pixel. Given this spatial limitation, we cannot reliably quantify uncertainty or bias associated with potential ice drift within this single-pixel boundary zone. Nonetheless, any remaining risk of an early ice-off bias is further reduced by the temperature-based labelling criterion described in the methodology, which provides an additional constraint on ice-off classification.

The observed offset between the RF and ADD model phenology dates is likely also due to differences between the onset of freezing or thawing and when a lake becomes fully ice free or ice covered. The fact that ~~72~~60% of ice-off dates are predicted later by the RF model than by the ADD model may reflect the effects of thermal inertia during spring, where higher air

temperatures and ADDT suggest faster ice thinning than is captured by the RF model analyzing lake backscatter. Despite the simple radar scattering and thermodynamic models employed here, each model exhibits decent predictive power for ice phenology and timing, and the inter-model biases are physically consistent.

4.2 Latitudinal and morphometric effects on ice phenology

The primary control on thermokarst lake ice phenology on the Alaska North Slope is latitude, which implicitly includes temperature. Latitude also incorporates other factors such as coastal proximity, elevation, topography, and local weather patterns. As a result, these likely also contribute to ice timing variability in addition to temperature. Even after excluding the high residual lakes primarily associated with the Brooks Range foothills, latitude continues to inform the regression predictor variables (mean spring temperature and morphometric properties). The weak but non-zero latitudinal correlation with residuals ($\rho = -0.24$) suggests there are still processes that covary with latitude but are not represented in the regression, such as elevation, which may explain the remaining correlation. The strong latitudinal gradient in ice-free duration (23 days per degree northward for lakes with residuals within the 5th-95th percentile of the distribution; Fig. 5) is consistent with the heightened climate sensitivity of lakes near the northern limits of the seasonal ice zone. This sensitivity implies that continued Arctic warming could produce disproportionately large changes in ice phenology for these northernmost lakes, with consequently disproportionate effects on future green-house gas emissions (Walter Anthony et al., 2018).

Our statistical analysis suggests that the offset between the RF and ADD model ice-on and ice-off dates is due in part to latitudinal effects. This is particularly so for lakes for which the RF-ADD model offset is greater than 2 revisit gaps (> 12 days). Lakes at lower latitudes are more likely to have ice-off detected by the RF model earlier than the ADD estimates and ice-on detected later relative to the ADD estimates. Lakes at higher latitudes have ice-off detected by the RF model later than the ADD estimates and ice-on earlier relative to the ADD estimates. This indicates that the RF model detects a longer ice-free period for lower latitude lakes and a shorter ice-free period for higher latitudes relative to the ADD estimates, suggesting that the ADD model is more conservative in predicting ice-free duration at lower latitudes whereas at higher latitudes, the RF model is more conservative.

The latitudinal gradient in offsets between the models may be explained through ice thickness variation as a function of latitude, the RF model not accounting for ice thickness, topography and microclimate, and differences in the spatial resolution of data. Given that lower latitudes have higher temperatures and are located further inland, they may have thinner ice overall and more complex freezing and thawing dynamics than lakes at higher latitudes. Meanwhile, these higher latitude lakes experience cooler temperatures and are more exposed to the Arctic Ocean thus freezing more rapidly. Furthermore, it is likely that the topographic gradient from the Brooks Range in the south to the low-elevation ACP in the north of our study area has an effect on the thermal budgeting of the lakes, owing to high-relief, radiation shadowing, and aspect differences that affect SAR backscatter through the imaging geometry (e.g. layover and shadowing). Additionally, the higher resolution of the SAR observations (~10 m/pixel) compared to the ERA5 temperature data (~30 km/pixel) may cause the RF model to capture thawing

and freezing more precisely than the lower resolution ADD estimates of ice thickness by accounting for other thermal effects around lakes and the local microclimate and topography that are beyond the resolution capabilities of the ERA5 data.

The morphometric relationships we identified provide insight into the physical controls on ice phenology beyond geographic location. Morphology regulates heat transfer from a lake's surroundings to the lake itself (Basu et al., 2024; Brown and Dugay, 2010) ultimately modulating the coupled effects of air temperature and latitude for an individual lake.

Shoreline development index (SDI) and convexity showed the largest effects after spatial correction (~3 days across typical ranges), while lake area and circularity showed smaller but significant effects (~1–2 days) (Fig. 9). Higher SDI, indicating a more complex shoreline, generally suggests shorter ice-free periods, with the onset of freezing occurring earlier in the year and onset of thaw beginning later. Similarly, lower convexity, associated with higher boundary concavity or irregularity (Bozeman and Pilling, 2013), shortens the ice-free period due to ice-on occurring at an earlier date. However, this trend with lower convexity and shorter ice-free periods holds true for the full distribution of lakes in the dataset not accounting for spatial autocorrelation. Conversely, higher circularity lakes tend to remain ice-free longer because they freeze later and thaw earlier in the year.

These patterns may reflect thermal sheltering in inlets and reduced fetch, which promotes ice growth earlier in the year (Basu et al., 2024). Smaller fetch results in reduced wind speeds, which facilitates the growth of ice crystals, while higher wind speeds can break up the initially forming ice layer. (Brown and Dugay, 2010). Ice growth typically begins along sheltered parts of the lakes and along the shore (Ashton, 1986), due in part to the wind-effect but also because of the shallower depths, which are more sensitive to temperature changes (Kirillin, et al., 2012). Irregular lakes have more shoreline per area, helping to initiate ice growth earlier than more circular or compact lakes. Lower convexity may also be related to overall shallower depths (Zhan et al., 2023), suggesting that deeper lakes have longer ice-free periods than shallower lakes. This likely corresponds to the patterns we identify with lake area, where ice free period increases with lake area. Lake depths tend to scale with area (Cael and Seekell, 2022); thus, these larger and deeper lakes remain ice-free longer because they begin freezing later in the year likely due to a combination of longer periods of thermal mixing within the water column and their greater thermal inertia. Indeed, lake depth has been found to be a significant control on the timing of ice-off conditions for thermokarst lakes in this region (Arp et al., 2015).

Lake depth also serves as a control on which lakes have bedfast or floating ice, which may also influence ice-out timing (Grunblatt and Atwood 2014; Sellmann et al. 1975). We combined SAR-based lake classifications of bedfast and floating ice regimes (Engram 2018; Bergstedt and Jones 2025) with our SAR RF-detected ice phenology across 6,797 lakes overlapping between the combined datasets. We observe that the difference between bedfast and floating ice regime has little effect on ice phenology over this large geographic extent encompassing seven thousand lakes as shown by the relatively small values of Cohen's *d* separability (Fig. 10). However, we do observe that bedfast ice lakes in the aggregate thaw prior to floating ice lakes in the aggregate; we see a modest five-day difference in median values between bedfast and floating ice for ice off compared to the larger 16-day difference found in Arp et al. 2015.




Figure 10: Bedfast and floating ice regime data (Engram 2018; Bergstedt and Jones 2025) histograms and kernel density estimations for A) ice off, B) ice on, and C) total ice-free days across 6,797 spatially overlapping lakes with our SAR RF-detected ice phenology events.

While a highly correlated scaling between lake depth, area and ice-free duration remains outstanding--potentially due to the limited range of thermokarst lake depths, and the importance of the underlying substrate (Hinkel et al., 2012) -- some studies have shown moderate correlations between lake depth and area (Arp et al., 2011), with modelling suggesting physical mechanisms by which lake deepening and enlargement can occur (Ohara et al., 2022). We therefore suggest a very general correspondence between these morphometric properties, which allows for the observed variability we identify.

Overall, morphometry explained only 1.5–2% of variance beyond temperature, indicating that climate remains the dominant control on regional ice phenology. Nonetheless, morphology plays a significant role in regulating the climatic impact for thermokarst lakes on the Alaska North Slope.

4.3 Implications for GHG emissions and Arctic change

Based on our results, Arctic thermokarst lakes are likely to experience pronounced effects on ice phenology due to global climate change. The strong latitudinal-temperature dependence suggests that increased warming will likely increase the ice free-period, which will further contribute to greenhouse gas emissions (Guo et al. 2020; Zhuang et al., 2023). Because larger lakes have longer ice-free durations, their potential contribution to GHG emissions may increase. However, these effects may be dwarfed by the sheer number of smaller lakes, which likely have a higher potential contribution. Furthermore, if larger lakes are indeed deeper, than they may also be less efficient emitters of methane due to increased oxidation that occurs in deeper water columns. Recent work (Levenson et al., 2025) found that thermokarst lakes in previously unglaciated landscapes underlain with continuous permafrost, including our study area, are projected to decrease in overall total area under projected climate scenarios. This reduction in total area may have direct effects on average lake depth and consequently emission potential, as shallower lakes and ponds exhibit higher methane emissions than deeper lakes (Manasypov et al., 2023).

Other climatic processes that depend upon the timing of ice formation and breakup may also be affected by a changing global climate. Evaporative fluxes from ice-free lakes increase substantially following ice-out, enhancing the exchange of moisture and heat between lakes and the atmosphere. As demonstrated by Arp et al. (2015), variations in lake depth, ice thickness, and ice-out timing can lead to divergent hydrologic responses among lakes in this region, further highlighting the potential importance of lake morphology in modulating lake ice phenology and its impact of climate processes.

The phenology baseline established here provides a foundation for detecting future changes in thermokarst lake ice regimes that can be incorporated with other Arctic remote sensing studies of methane emissions from thermokarst lakes (Elder et al., 2020; Engram and Walter Anthony, 2024; Kyzivat and Smith, 2023). Future monitoring efforts can be leveraged in conjunction with these results to assess whether ice-free periods increase as predicted by climate models. Lastly, upcoming higher-resolution spaceborne methane measurements (e.g., MethaneSAT, Carbon Mapper, GHGSat (McLinden et al., 2024)) could be leveraged in conjunction with this ice phenology and timing observational method to directly observe the ice-phenology-emission relationship from the individual lake-scale to the regional-scale and determine the relative importance of numerous small lakes vs. fewer larger lakes in consideration of GHG emission budgeting.

5 Conclusion

This study presents the first comprehensive analysis of ice phenology across the Alaska North Slope thermokarst lake population. Using a RF classifier trained using labels derived from Sentinel-2 optical imagery and ERA5 temperature data and

applied to Sentinel-1 SAR imagery, we achieved 94% accuracy for ice state classification and retrieved phenology for 30,774 lakes over five years (2019–2023).

The mean ice-free period is 114 days, with ice-off occurring in mid-June (DOY 159) and ice-on in late September (DOY 273). Latitude exerts dominant control on ice timing, with ice-free duration decreasing by 23 days per degree northward. Morphometric predictors showed modest but significant effects after spatial correction, with shoreline development index and convexity each contributing ~3 days variation across typical lake ranges.

These results provide essential baseline data for monitoring how thermokarst lake dynamics respond to continued Arctic warming, with implications for regional carbon cycling and climate feedbacks. The lake-specific ice phenology dates also provide critical timing constraints for methane emission models and for planning targeted high-resolution observation campaigns.

Code and data availability

Sentinel-1 and Sentinel-2 imagery were accessed through Google Earth Engine (https://earthengine.google.com). ERA5 reanalysis data are available from the Copernicus Climate Data Store (https://cds.climate.copernicus.eu). The Alaska Lake and Pond Occurrence Dataset (ALPOD) is available from the ORNL DAAC (doi:10.3334/ORNLDAAC/2399). Processed ice phenology data and analysis code are archived at https://github.com/washu-eeps/thermokarst-lakes-alaska.

Interactive computing environment

All code used in this study is provided in the form of Jupyter Notebooks within the provided Zenodo and GitHub repositories. Each notebook lists the necessary Python packages and calls to Google Cloud storage buckets through Washington University in St. Louis. These Jupyter Notebooks can be directly downloaded to run locally or in the cloud, and the GitHub repository can be forked and cloned for execution and reproduction.

Author contributions

Conceptualization: all authors. Data curation: A.L.N., A.S.B., R.J.M. Formal analysis: all authors. Investigation: C.G.L., A.L.N., A.S.B., R.J.M. Methodology: all authors. Software: C.G.L., A.S.B. Writing: all authors. Supervision: R.J.M., C.C.M. A.L.N. and C.G.L contributed equally to the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We thank the Google Earth Engine team for providing computational resources and for open access to satellite data (Gorelick et al., 2017). Hersbach et al, (2017) was downloaded from the Copernicus Climate Change Service (C3S) Climate Data Store. We thank C3S for allowing us to redistribute the data. The results contain modified Copernicus Climate Change Service information 2022. Neither the European Commission nor ECMWF is responsible for any use that may be made of the Copernicus information or data it contains. We thank the Natural Earth for vector data provided through Amazon Web Services used in figure map generation. Cloud computing resources for satellite data processing and analysis were provided by the Fossett Laboratory at Washington University in St. Louis, which we gratefully acknowledge.

We would additionally like to thank Isabel Lopez for assistance in initial project conceptualization and planning and preliminary manual data analysis. Furthermore, we thank Alisa Oates and Amelia Mogul for assistance in preliminary work manually validating an earlier version of the RF model used in this study.

We used generative AI (gpt-5.2, opus-4.6, and gemini-3) to assist with computational code associated with this study. Initial code was produced and refined through iterative prompting and subsequently reviewed and manually revised by the authors.

Financial support

This research was supported by the Washington University Department of Earth, Environmental, and Planetary Sciences and by a National Science Foundation Graduate Research Fellowship and the Ann W. and Spencer T. Olin-Chancellor's fellowship to A.N.

This material is based upon work supported by the National Science Foundation's Graduate Research Fellowship Program (under Grant No. DGE 2139839). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

References

Arp, C. D., Jones, B. M., Urban, F. E., and Grosse, G.: Hydrogeomorphic processes of thermokarst lakes with grounded-ice and floating-ice regimes on the Arctic coastal plain, Alaska, Hydrol. Process., 25, 2422–2438, https://doi.org/10.1002/hyp.8019, 2011.

Arp, C. D., Jones, B. M., Liljedahl, A. K., Hinkel, K. M., and Welker, J. A.: Depth, ice thickness, and ice-out timing cause divergent hydrologic responses among Arctic lakes, Water Resour. Res., 51, 9379–9401, https://doi.org/10.1002/2015WR017362, 2015.

Ashton, G.D.: River and Lake Ice Engineering. Water Resources Publications, Littleton, CO, 1986.

Bartsch, A., Pointner, G., Leibman, M. O., Dvornikov, Y. A., Khomutov, A. V., and Trofaier, A. M.: Circumpolar mapping of ground-fast lake ice, Front. Earth Sci., 5, 12, https://doi.org/10.3389/feart.2017.00012, 2017.

Bartsch, A., Strozzi, T., and Nitze, I.: Permafrost Monitoring from Space, Surveys in Geophysics, https://doi.org/10.1007/s10712-023-09770-3, 2023.

Basu, A., Culpepper, J., Blagrave, K., & Sharma, S.: Phenological shifts in lake ice cover across the Northern Hemisphere: A glimpse into the past, present, and the future of lake ice phenology. Water Resour. Res., 60, https://doi.org/10.1029/2023WR036392, 2024.

Beaven, S. G.: Sea Ice Radar Backscatter Modeling, Measurements, and the Fusion of Active and Passive Microwave Data. Radar Remote Sensing Laboratory, The University of Kansas Center for Research, Inc., 1995.

Bergstedt, H., and Jones, B.: Integrated Field and Synthetic Aperture Radar-Based Dataset for Arctic Lake Ice and Under-Ice Water Salinity Classification, Northern Alaska. Arctic Data Center, https://doi:10.18739/A25M6288G, 2025.

Bozeman, J. R., & Pilling, M.: The Convexity Ratio and Applications, Scientiae Mathematicae Japonicae, 76(1), 47-53, https://doi.org/10.32219/isms.76.1_47, 2013.

Breiman, L.: Random Forests, Machine Learning, 45, 5-32, https://doi.org/10.1023/A:1010933404324, 2001.

Brown, L. C., & Duguay, C. R.: The response and role of ice cover in lake-climate interactions. Progress in Physical Geography: Earth and Environment, 34(5), 671-704, https://doi.org/10.1177/0309133310375 65, 2010.

Bryant, M. B., Borsa, A. A., Anderson, E. J., Masteller, C. C., Michaelides, R. J., Siegfried, M. R., and Young, A. P.: Multiple modes of shoreline change along the Alaskan Beaufort Sea observed using ICESat-2 altimetry and satellite imagery, The Cryosphere, 19, 1825–1847, https://doi.org/10.5194/tc-19-1825-2025, 2025.

Cael, B.B. and Seekell, D.: A theory for the relationship between lake surface area and maximum depth. Limnol. Oceanogr. Lett, 7, 527-533, https://doi.org/10.1002/lol2.10269, 2022.

Carmack, E. C., Vagle, S., Morrison, J., and Laval, B. E.: Space-for-time proxy for climate change in deep lakes in the Canadian cordillera: Seasonality along a latitudinal climate gradient, J. Great Lakes Res., 40, 608–617, https://doi.org/10.1016/j.jglr.2014.06.007, 2014.

Carson C.E., and Hussey K.M.: The oriented lakes of Arctic Alaska. Journal of Geology, 70, 417–439, 1962.

Carver, R. W., and Merose, A.: ARCO-ERA5: An Analysis-Ready Cloud-Optimized Reanalysis Dataset. 22nd Conference on AI for Environmental Science. American Meterological Society, 4A.1, 2023AMS...10315842C, 2023.

Detterman, R. L., Bowsher, A. L., & Dutro, J. T.: Glaciation on the Arctic Slope of the Brooks Range, Northern Alaska. Arctic, 11(1), 43–61, https://doi.org/10.14430/arctic3732, 1958.

Duguay, C.R., Flato, G.M., Jeffries, M.O., Ménard, P., Morris, K. and Rouse, W.R. (2003), Ice-cover variability on shallow lakes at high latitudes: model simulations and observations. Hydrol. Process., 17, 3465-3483, https://doi.org/10.1002/hyp.1394, 2003.

Elder, C. D., Thompson, D. R., Thrope, A. K., Hanke, P., Walter Anthony, K. M., Miller, C. E.: Airborne Mapping Reveals Emergent Power Law of Arctic Methane Emissions, Geophysical Research Letters, 47, 3, https://doi.org/10.1029/2019GL085707, 2020.

Engram, M.: Floating and bedfast lake ice regimes across Arctic Alaska using space-borne SAR imagery from 1992-2016. Arctic Data Center, https://doi:10.18739/A2CJ87K8H, 2018.

Engram, M., and Walter Anthony, K.: Synthetic aperture radar (SAR) detects large gas seeps in Alaska lakes. Environmental Research Letters, 19, 10.1088/1748-9326/ad2b2a, 2024.

Farquharson, L.M., Mann, D.H., Grosse, G., Jones, B.M., Romanovsky, V.E.: Spatial distribution of thermokarst terrain in Arctic Alaska. Geomorphology, 273, 116–133, 10.1016/j.geomorph.2016.08.007, 2016.

Gorelick, N., Hancher, M., Dixon, M., Ilyushchenko, S., Thau, D., and Moore, D.: Google Earth Engine: Planetary-scale geospatial analysis for everyone. Remote Sensing of Environment, 10.1016/j.rse.2017.06.031, 2017.

Grunblatt, J. and Atwood, D.: Mapping lakes for winter liquid water availability using SAR on the North Slope of Alaska, International Journal of Applied Earth Observation and Geoinformation, 27, 63-69, 10.1016/j.jag.2013.05.006, 2014 .

Guo M., Zhuang, W., Tan, Z., Shurpali, N., Juutinen, S., Kortelainen, P., and Martikainen, P. J.: Rising methane emissions from boreal lakes due to increasing ice-free days, Environmental Research Letters, 15, 6, 10.1088/1748-9326/ab8254, 2020.

Hall, D. K., Fagre, D. B., Klasner, F., Linebaugh, G., and Liston, G. E.: Analysis of ERS 1 synthetic aperture radar data of frozen lakes in northern Montana and implications for climate studies. Journal of Geophysica Research Oceans, 99, C11, 22473-22482, https://doi.org/10.1029/94JC01391, 1994.

Hall, D., Riggs, G.A., and Salomonson, V. V.: Development of methods for mapping global snow cover using moderate resolution imaging spectroradiometer data, Remote Sensing of Environment, 54, 2, 127-40, 10.1016/0034-4257(95)00137-P, 1995.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F.,Villaume, S., Thépaut, J-N.: Complete ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. Copernicus Climate Change Service (C3S) Data Store (CDS), 2017.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Jean-Noël Thépaut: The ERA5 global reanalysis, Q. J. R. Meteorol. Soc., 146, 1999–2049, https://doi.org/10.1002/qj.380, 2020.

Heslop, J. K., Walter Anthony, K. M., Winkel, M., Sepulveda-Jauregui, A., Martinez-Cruz, K., Bondurant, A., Gross, G., and Liebner, S.: A synthesis of methane dynamics in thermokarst lake environments, Earth-Science Reviews, https://doi.org/10.1016/j.earscirev.2020.103365, 2020.

Higgins, S. N., Desjardins, C. M., Drouin, H., Hrenchuk, L. E., & van der Sanden, J. J.: The role of climate and lake size in regulating the ice phenology of boreal lakes. Journal of Geophysical Research: Biogeosciences, 126, https://doi.org/10.1029/2020JG005898, 2021.

Hinkel, K. M., Sheng, Y., Lenters, J. D., Lyons, E. A., Beck, R. A., Eisner, W. R., and Wang, J.: Thermokarst Lakes on the Arctic Coastal Plain of Alaska: Geomorphic Controls on Bathymetry, Permafrost and Periglacial Processes, 23, 3, 218-230, https://doi.org/10.1002/ppp.1744, 2012.

Hoekstra, M., Jiang, M., Clausi, D. A., and Duguay, C.: Lake ice-water classification of RADARSAT-2 images by integrating IRGS segmentation with pixel-based random forest labeling, Remote Sens. (Basel), 12, 1425, https://doi.org/10.3390/rs12091425, 2020.

Hutchinson, G. E.: A Treatise on Limnology, Vol. 1: Geography, Physics, and Chemistry, John Wiley & Sons, New York, NY, 1957.

In' T Zandt, M. H., Liebner, S., and Welte, C. U.: Roles of Thermokarst Lakes in a Warming World, Trends in Microbiology, 28, 9, https://doi.org/10.1016/j.tim.2020.04.002, 2020.

Jeffries, M. O., Morris, K., Weeks, W. F., and Wakabayashi, H.: Structural and stratigraphie features and ERS 1 synthetic aperture radar backscatter characteristics of ice growing on shallow lakes in NW Alaska, winter 1991-1992, Journal of Geophysical Research Oceans, 99, C11, 22459-22471, https://doi.org/10.1029/94JC01479, 1994.

Jones, B.M., Grosse, G., Farquharson, L.M. et al. Lake and drained lake basin systems in lowland permafrost regions. Nat Rev Earth Environ 3, 85–98, https://doi.org/10.1038/s43017-021-00238-9, 2022.

Kirillin, G., Leppäranta, M., Terzhevik, A. et al.: Physics of seasonally ice-covered lakes: a review. Aquat Sci 74, 659–682, https://doi.org/10.1007/s00027-012-0279-y, 2012.

Latifovic, R., and Pouliot, D.: Analysis of climate change impacts on lake ice phenology in Canada using the historical satellite data record, Remote Sensing of Environment, 106, 4, 10.1016/j.rse.2006.09.015, 2007.

Lafrenière, M. J., and Lamoureux, S. F.: Effects of changing permafrost conditions on hydrological processes and fluvial fluxes, Earth-Science Reviews, 191, 212-223, https://doi.org/10.1016/j.earscirev.2019.02.018, 2019.

Levenson, E., Cooley, S. W., and Mullen, A.: ABoVE: Alaska Lake and Pond Occurrence, https://doi.org/10.3334/ORNLDAAC/2399, 2025.

Lund, J., Forster, R. R., Deeb, E. J., Liston, G. E., Skiles, S. M., and Marshall, H.: Interpreting Sentinel-1 SAR Backscatter Signals of Snowpack Surface Melt/Freeze, Warming, and Ripening, through Field Measurements and Physically-Based SnowModel, Remote Snsing, 14 (16), 4002, https://doi.org/10.3390/rs14164002, 2022.

Manasypov, R., Fan, L., Lim, A. G., Krickov, I. V., Pokrovsky, O. S., Kuzyakov, Y., and Dorodnikov, M.: Size matters: Aerobic methane oxidation in sediments of shallow thermokarst lakes, Global Change Biology, https://doi.org/10.1111/gcb.17120, 2023.

Matthews, E., Johnson, M. S., Genovese, V., Du, J., and Bastviken, D.: Methane emission from high latitude lakes: Methane-centric lake classification and satellite-driven annual cycle of emissions. Scientific Reports, 10, 1, https://doi.org/10.1038/s41598-020-68246-1, 2020.

Marin, C., Bertoldi, G., Premier, V., Callegari, M., Brida, C., Hürkamp, K., Tschiersch, J., Zebisch, M., and Notarnicola, C., Use of Sentinel-1 radar observations to evaluate snowmelt dynamics in alpine regions, The Cryosphere, 14, 935-956, https://doi.org/10.5194/tc-14-935-2020, 2020.

McLinden, C. A., Griffin, D., Davis, Z., Hempel, C., Smith, J., Sioris, C., Nassar, R., Moeini, O., Legault-Ouellet, E., and Malo, A.: An Independent Evaluation of GHGSat Methane Emissions: Performance Assessment, Journal of Geophysical Research: Atmospheres, 129, e2023JD039906, https://doi.org/10.1029/2023JD039906, 2024.

Mu, C., Lei, P., Mu, M., Zhang, C., Zhou, Z., Song, J., Jia, Y., Fan, C., Peng, X., Zhang, G., Yang, Y., Wang, L., Li, D., Song, C., Wang, G., and Zhang, Z.: Methane emissions from thermokarst lakes must emphasize the ice-melting impact on the Tibetan Plateau, Nature Communications, 16, 1, https://doi.org/10.1038/s41467-025-57745-2, 2025.

Mullen, A. L., Watts, J. D., Rogers, B. M., Carroll, M. L., Elder, C. D., Noomah, J., Williams, Z., Caraballo-Vega, J. A., Bredder, A., Rickenbaugh, E., Levenson, E., Cooley, S. W., Hung, J. K. Y., Fiske, G., Potter, S., Yang, Y., Miller, C. E., Natali, S. M., Douglas, T. A., and Kyzivat, E. D.: Using high-resolution satellite imagery and deep learning to track dynamic seasonality in small water bodies, Geophysical Research Letters, 50, 7, https://doi.org/10.1029/2022GL102327, 2023.

Mullissa, A.; Vollrath, A.; Odongo-Braun, C.; Slagter, B.; Balling, J.; Gou, Y.; Gorelick, N.; Reiche, J. Sentinel-1 SAR Backscatter Analysis Ready Data Preparation in Google Earth Engine. Remote Sensing, 13, 10, https://doi.org/10.3390/rs13101954, 2021.

Murfitt, J. and Duguay, C.: Assessing the Performance of Methods for Monitoring Ice Phenology of the World's Largest High Arctic Lake Using High-Density Time Series Analysis of Sentinel-1 Data. Remote Sensing, 12, 382, https://doi.org/10.3390/rs12030382, 2020.

Muster, S., Roth, K., Langer, M., Lange, S., Cresto Aleina, F., Bartsch, A., Morgenstern, A., Grosse, G., Jones, B., Sannel, A. B. K., Sjöberg, Y., Günther, F., Andresen, C., Veremeeva, A., Lindgren, P. R., Bouchard, F., Lara, M. J., Fortier, D., Charbonneau, S., Virtanen, T. A., Hugelius, G., Palmtag, J., Siewert, M. B., Riley, W. J., Koven, C. D., and Boike, J.: PeRL: a circum-Arctic Permafrost Region Pond and Lake database, Earth Syst. Sci. Data, 9, 317–348, https://doi.org/10.5194/essd-9-317-2017, 2017.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., and Duchesnay, E.: Scikit-learn: Machine learning in Python, J. Mach. Learn. Res., 12, 2825–2830, https://doi.org/10.48550/arXiv.1201.0490, 2011.

Posch, C., Abermann, J., and Silva, T.: Lake ice break-up in Greenland: timing and spatiotemporal variability, The Cryosphere, 18, 2035-2059, https://doi.org/10.5194/tc-18-2035-2024, 2024.

Pouw, A. F., Pour, H. K., and MacLean, A.: Mapping snow depth on Canadian sub-arctic lakes using ground-penetrating radar, The Cryosphere, 17, 2367-2385, https://doi.org/10.5194/tc-17-2367-2023, 2023.

Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, Commun. Earth Environ., 3, 168, https://doi.org/10.1038/s43247-022-00498-3, 2022.

Riggs, G. A., Hall, D. K., and Román, M. O.: Overview of NASA's MODIS and VIIRS snow-cover Earth System Data Records, Earth Syst. Sci. Data, 8, 215–226, https://doi.org/10.5194/essd-9-765-2017, 2017.

Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, Nature, 520, 7546, https://doi.org/10.1038/nature14338, 2015.

Seekell, D., Cael, B. B., Byström, P.: Problems With the Shoreline Development Index—A Widely Used Metric of Lake Shape, Geophysical Research Letters, 49, 10, https://doi.org/10.1029/2022GL098499, 2022.

Sellmann, P.V., Weeks, W.F., & Campbell, W.: Use of side–looking airborne radar to determine lake depth on the Alaskan North Slope, U.S. Cold Regions Research and Engineering Laboratory. Special Report 230, 1975

Sentinel-Hub: Sentinel-2 cloud detector, https://github.com/sentinel-hub/sentinel2-cloud-detector, 2025.

Serreze, M. C., and Barry, R. G.: Processes and impacts of Arctic Amplification: A research synthesis, Global and Planetary Change, 77, 1-2, https://doi.org/10.1016/j.gloplacha.2011.03.004, 2011.

Skakun, S., Wevers, J., Brockmann, C., Doxani, G., Aleksandrov, M., Batič, M., Frantz, D., Gascon, F., Gómez-Chova, L., Hagolle, O., López-Puigdollers, D., Louis, J., Lubej, M., Mateo-García, G., Osman, J., Peressutti, D., Pflug, B., Puc, J., Richter, R., Roger, J.-C., Scaramuzza, P., Vermote, E., Vesel, N., Zupanc, A., and Žust, L.: Cloud Mask Intercomparison eXercise (CMIX): An evaluation of cloud masking algorithms for 8 and Sentinel-2, Remote Sens. Environ., 274, 112990, 10.1016/j.rse.2022.112990, 2022.

Smits, A. P., Gomez, N. W., Dozier, J., & Sadro, S.: Winter climate and lake morphology control ice phenology and under-ice temperature and oxygen regimes in mountain lakes. Journal of Geophysical Research: Biogeosciences, 126, https://doi.org/10.1029/2021JG006277, 2021.

Stefan, J.: On the theory of ice formation, with particular regard to ice formation in the polar sea, Sitz. Ber. Kais. Akad. Wiss. Wien, 98, 965-983, 1890.

Streletskiy, D. A., Maslakov, A., Grosse, G., Shiklomanov, N. I., Farquharson, L., Zwieback, S., Iwahana, G., Bartsch, A., Liu, L., Strozzi, T., Lee, H., and Debolskiy, M. V.: Thawing permafrost is subsiding in the Northern Hemisphere-Reivew and perspectives, Environmental Research Letters, 20, 1, 10.1088/1748-9326/ada2ff, 2025.

Stuefer, S. L., Hale, K., May, L. D., Mason, M., Vuyovich, C., Marshall, H., Vas, D., and Elder, K.: Snow depth measurements from Arctic tundra and boreal forest collected during NASA SnowEx Aalaska campaign, Sci Data, 12, 919, https://doi.org/10.1038/s41597-025-05170-x, 2025.

Surdu, C. M., Duguay, C. R., Brown, L. C., and Fernández Prieto, D.: Response of ice cover on shallow lakes of the North Slope of Alaska to contemporary climate conditions (1950--2011): Radar remote-sensing and numerical modeling data analysis, Cryosphere, 8, 167–180, https://doi.org/10.5194/tc-8-167-2014, 2014.

Surdu, C. M., Duguay, C. R., Pour, H. K., and Brown, L. C.: Ice Freeze-up and Break-up Detection of Shallow Lakes in Northern Alaska with Spaceborne SAR, Remote Sensing, 7, 6133-6159, https://doi.org/10.3390/rs70506133, 2015.

Surdu, C. M., Duguay, C. R., and Fernández Prieto, D.: Evidence of recent changes in the ice regime of lakes in the Canadian High Arctic from spaceborne satellite observations, Cryosphere, 10, 941–960, https://doi.org/10.5194/tc-10-941-2016, 2016.

Tom, M., Aguilar, R., Imhof, P., Leinss, S., Baltsavias, E., and Schindler, K.: Lake ice detection from Sentinel-1 Sar with deep learning, ISPRS Ann. Photogramm. Remote Sens. Spat. Inf. Sci., V-3–2020, 409–416, 10.5194/isprs-annals-V-3-2020-409-2020, 2020.

Veijola, K., Cohen, J., Mäkynen, M., Lemmetyinen, J., Praks, J., and Cheng, B.: X- and Ku-Band SAR Backscattering Signatures of Snow-Covered Lake Ice and Sea Ice, Remote Sensing, 16 (2), 369, https://doi.org/10.3390/rs16020369, 2024.

Walter Anthony, K., Schneider von Deimling, T., Nitze, I., Frolking, S., Emond, A., Daanen, R., Anthony, P., Lindgren, P., Jones, B., and Grosse, G.: 21st-century modeled permafrost carbon emissions accelerated by abrupt thaw beneath lakes, Nat. Commun., 9, 3262, https://doi.org/10.1038/s41467-018-05738-9, 2018.

West, W. E., Creamer, K. P., and Jones, S. E.: Productivity and depth regulate lake contributions to atmospheric methane, Limnology and Oceanography, 61, S1, https://doi.org/10.1002/lno.10247, 2016.

Wik, M., Varner, R. K., Anthony, K. W., MacIntyre, S., and Bastviken, D.: Climate-sensitive northern lakes and ponds are critical components of methane release, Nat. Geosci., 9, 99–105, https://doi.org/10.1038/ngeo2578, 2016.

USACE.: Method to Estimate River Ice Thickness Based on Meteorological Cold Regions Research and Engineering Laboratory Data Report, 6 pp., U.S. Army Corps of Eng., Hanover, N. H, 2004.

Zhan, P., Song, C., Liu, K., Chen, T., Ke, L., Luo, S., and Fan, C.: Can we estimate the lake mean depth and volume from the deepest record and auxiliary geospatial parameters?, Journal of Hydrology, 617(A), 10.1016/j.jhydrol.2022.128958, 2023.

Zhuang, Q., Guo, M., Melack, J. M., Lan, X., Tan, Z., Oh, Y., and Leung, L. R.: Current and Future Global Lake Methane Emissions: A Process-Based Modeling Analysis, Journal of Geophysical Research: Biogeosciences, 128, 3, https://doi.org/10.1029/2022JG007137, 2023.