Dynamical precursors to summer temperature extremes on the Antarctic Peninsula

William J. Dow[1], Amanda C. Maycock[1], Andrew N. Ross[1], Ryan S. Williams[2], Thomas J. Bracegirdle[2]

1 - Institute for Climate and Atmospheric Sciences, School of Earth, Environment and Sustainability, University of Leeds, Leeds, UK
2 - British Antarctic Survey, Cambridge, UK

Correspondence to: William J. Dow (earwd@leeds.ac.uk)

Abstract: Extreme warm summer near-surface temperatures over the Antarctic Peninsula (AP) can lead to surface melting and the disintegration of ice shelves. While individual case studies have linked such events to anomalous large-scale circulation, a systematic assessment of the dynamical pathways leading to AP-wide extreme austral summer warm events remains limited. This study uses ERA5 reanalysis data to investigate the large-scale dynamical precursors associated with extreme warm days over the Antarctic Peninsula. We apply k-means clustering to mean sea level pressure anomalies during high temperature extremes and identify five distinct circulation patterns with different dominant zonal wavenumbers. We investigate the spatio-temporal evolution and persistence of near-surface wind, temperature and pressure for each cluster. Four clusters are associated with rapidly amplifying planetary-scale wave patterns, while a fifth resembles a negative Southern Annular Mode–like state with enhanced persistence prior to event onset. Despite these differing pathways, all regimes promote anomalous northerly flow toward the AP, driving strong meridional temperature advection and regional warming. We demonstrate that extreme Antarctic Peninsula warm events arise from distinct circulation pathways, reflecting diverse dynamical states likely influenced by hemispheric-scale teleconnections and planetary wave interactions.

1 Introduction

The Antarctic Peninsula (AP) has undergone marked multi-decadal variability in surface air temperature since the mid-20th century. Warming trends calculated over periods longer than approximately 30 years appear robust, particularly during austral summer and autumn (González and Fortuny, 2018), with late 20th century increases exceeding the global mean in several observational records (Marshall et al., 2006; Oliva et al., 2017; Wang et al., 2022a). Concurrently, the late 1990s marked the onset of a period of relative cooling consistent with internal variability (Carrasco et al., 2019; Turner et al., 2016). However, more recent analyses suggest that this early 21st century cooling may be weakening, with temperature

evolution closely tied to changes in large-scale circulation modes and regional pressure systems (Carrasco et al., 2019; Bozkurt et al., 2020).

During austral summer (December-February), episodes of extreme warm surface air temperatures over the AP — frequently exceeding 0 °C — are projected to become increasingly common under future scenarios that include continued increases in greenhouse gases (e.g. Siegert et al. 2019). Similarly, pronounced increases in high extremes (p90; 90th percentile compared to mean background climate) in precipitation and westerly wind anomalies are projected to occur on the AP over the 21st century compared to increases in background climatology (Bracegirdle et al., 2024). Such extreme weather events can drive surface melt (Gorodetskaya et al. 2023), firn air depletion (Dunmire et al. 2024), and, in some cases, contribute to the destabilisation and collapse of ice shelves (Scambos et al., 2009; Wille et al., 2022). Improving our understanding of the atmospheric drivers of extreme warm temperatures is therefore critical for assessing Antarctica's vulnerability to climate change and its broader implications for the Earth system.

Extreme warm events over the AP are typically associated with strong poleward meridional wind anomalies near the AP at synoptic timescales, which are often associated with atmospheric blocking, atmospheric rivers and enhanced moisture transport towards Antarctica (Banwell et al., 2021; Bozkurt et al., 2024; Gorodetskaya et al., 2023; Wille et al., 2022; Zou et al., 2023; Wang et al., 2022a; Wang et al., 2022b). The synoptic scale circulation anomalies can be linked with remote forcing, and their effects on temperature may be modulated by local scale processes, including föhn warming on the eastern Peninsula (Bevan et al., 2020; Bae et al., 2022; Guarino et al., 2026; Kirchgaessner et al., 2021; Xu et al., 2021; Zou et al., 2023; Zhu et al., 2026).

Anomalous tropical convection associated with the Madden–Julian Oscillation (MJO) and the El Niño-Southern Oscillation (ENSO) can initiate poleward propagating Rossby wave trains that favour blocking highs over the South Pacific and Drake Passage (Renwick and Revell, 1999; Tseng et al., 2019; Clem et al., 2022; Barrett et al., 2025). In addition, a Rossby wave train from the tropical Atlantic has been shown to modulate the intensity of the Amundsen–Bellingshausen Seas Low (ABSL) on decadal timescales, thereby priming the regional circulation to influence conditions across the AP (Li et al., 2014; Li et al., 2015). Furthermore, anomalous sea surface temperatures across the South Pacific Convergence Zone (SPCZ) have been linked to anticyclonic blocking over coastal West Antarctica through enhanced Rossby wave propagation (Deb et al., 2025). Together, these tropical-extratropical teleconnections can promote poleward transport of heat and moisture toward the Peninsula (e.g., Rondanelli et al., 2019; Clem et al., 2022).

Atmospheric circulation near the AP is also modified by the Southern Annular Mode (SAM) (e.g. Fogt et al., 2012; Clem and Fogt, 2013; Clem et al., 2016; Montano Bello and Villegas Bolanos, 2025). Post 1979, the SAM has become less zonally symmetric, with an increase in zonal wave-3 structure and an

eastward shift in phase, modulating the likelihood ~~resulting in~~ of more northerly flow anomalies over the AP during positive SAM conditions, consistent with an eastward displacement of the Amundsen–Bellingshausen Sea Low (Marshall et al., 2022). The SAM has also exhibited a positive trend in austral summer and autumn since the late 20th century, with the summer trend driven primarily by stratospheric ozone depletion (e.g. Thompson and Solomon, 2002; Marshall et al., 2003; Fogt and Marshall, 2020), alongside a deepening of the ABSL (England et al., 2016). The shift toward more positive summer SAM conditions has contributed to long-term warming over the AP through the intensification of high-latitude cyclones, strengthening of meridional flow onto the AP via amplification of the ABSL, and enhancement of lee-side föhn winds along the eastern AP (Marshall, 2007; Orr et al., 2008; Fogt et al., 2011, 2012; Turner et al., 2013; Hosking et al., 2013). Importantly, SAM variability has been linked to changes in the frequency of Peninsula-centred pressure dipole patterns and ABSL-dominated regimes (Hosking et al., 2013; Raphael et al., 2016), including an increased occurrence of an AP-centred dipole during summer between 1979 and 2016 (González et al., 2018)

While the typical circulation features necessary to drive anomalous high temperatures over the AP have been described in the literature (e.g. Balay and Renom, 2026; Clem et al., 2022; Gorodetskaya et al., 2023; Wang et al., 2022a; Wille et al., 2022), there has been relatively little investigation into the diversity in circulation features amongst extremes. In particular: are all dynamically-driven AP extremes made equal? Gonzalez et al. (2018) performed daily clustering of circulation patterns around the AP during the entire year, and therefore did not isolate circulation features associated with extreme high temperatures. Nielsen et al. (2025) applied clustering to surface air temperature trends over the entire Antarctic continent ~~which~~ and identified three~~identified a single~~ clusters ~~representing~~ that encompassed the AP and analysed their ~~with its~~ associated extremes. However, they did not specifically examine ~~, and therefore did not consider~~ the diversity of circulation features amongst AP temperature extremes. As a result, a systematic characterisation of the circulation pathways producing summer high temperatures over the AP is lacking.

This study uses ERA5 reanalysis data to examine the synoptic and large-scale circulation patterns linked to extreme austral summer temperature events across the AP. We define extremes over the AP and apply k-means clustering (cf. Nielsen et al., 2025) to mean sea level pressure (MSLP) anomalies at the onset of the extreme temperatures to identify their characteristic clusters which are then studied. The remainder of the paper is structured as follows: Section 2 outlines the methods, Section 3 presents the results, and Section 4 discusses the implications for understanding and anticipating future extreme events in a warming climate.

2 Data and Methods

2.1 Data

Daily mean near-surface air temperature, 10 metre wind fields, sea level pressure and 500 hPa geopotential height data were obtained from ERA5 for the period 1979-2024 on a regular 0.25° x 0.25° grid (Hersbach et al., 2020). Hourly zonal and meridional winds and temperature fields at 300 hPa were utilised in the meridional temperature advection analysis. ERA5 exhibits regionally varying near-surface biases at coastal stations and a general warm bias (DJF mean = +0.96 K) (Zhu et al. 2021) over the AP; however, it demonstrates generally high agreement in temporal variability with Antarctic station observations (DJF seasonal mean correlation coefficient = 0.82) (Zhu et al., 2021) and is considered to effectively capture the spatial structure and temporal variability of near surface temperature and circulation climate across the AP (Tetzner et al., 2019; Hillebrand et al., 2020; Zhu et. al., 2021). This makes ERA5 well suited for analysing synoptic scale circulation associated with austral summer temperature extremes across the AP.

2.2 Methods

2.2.1 Definition of temperature extremes

Extreme high temperature days are identified based on 2-m temperature anomalies calculated relative to a 21-day running mean climatology. Alternative climatological window lengths (7, 10, 14, 30 days) were tested (Supplementary Fig. S1), with 21 days chosen to best capture the seasonal cycle without oversmoothing. The AP temperature anomalies are then calculated as the area-weighted average over the whole region [80°W-50°W; 60°S-75°S]. Extreme summer temperature anomalies are defined as days when the temperature anomaly exceeds the DJF 95th percentile of departures from the 21-day running mean. This approach aims to capture coherent high temperature events which have the potential to impinge on ice shelf surface melting, and are most likely driven by large-scale regional circulation anomalies (e.g. Wille et al., 2022). The seasonal DJF temperature trend over 1979-2019~~24~~ is not significant across the AP region in ERA5 (Zhu et al., 2021), and we find no systematic trend in the frequency of extreme high temperature days during the analysis period (Supplementary Fig S2).

Extreme high temperature days are grouped into events allowing up to 2 consecutive non-extreme days within a sequence (cf. Baldwin et al., 2019; Ruosteenoja & Jylhä, 2023). Sequences separated by >2 days were treated as independent events. This definition captures 87 events (mean duration 2.4 days), with most lasting one day (n = 35, 40%) and 82% lasting three days or fewer.

2.2.2 K-means clustering

K-means clustering has been applied to understand Antarctic climate in previous work (Coggins et al., 2014; Nielsen et al., 2025; Gonzalez et al., 2018). Here, we apply k-means clustering using Lloyd's algorithm (Lloyd, 1982), to the daily MSLP anomaly fields over the region [50°S - 90°S, 180°W - 0°W] for days -3 to +1 relative to the event onset (day 0). The −3 to +1 day window is chosen to capture the immediate synoptic evolution preceding and during extreme high temperature events, while limiting the inclusion of unrelated background variability. Each day within this window is treated as an independent sample in the clustering procedure. That is, daily MSLP anomaly fields from all events spanning days −3 to +1 are input to the clustering. This allows the temporal evolution of circulation patterns leading up to event onset to be assessed through the event duration. For event-based analyses each event was assigned to a single cluster according to the cluster membership of the day 0 circulation pattern. Lead-lag composites outside the clustering window were grouped according to the cluster assignment determined during the clustering stage and were not independently clustered.

To assess whether the clustering window constrains circulation states associated with warm extremes, we computed the Euclidean distance between daily MSLP fields and their associated cluster centroids as a function of lag. We tested for discontinuities at the clustering window boundaries (lag −3 to −4 and +1 to +2), which would indicate a mismatch between clustered and unclustered regimes. Instead, distances vary smoothly with lag, with boundary changes comparable to typical day-to-day variability (not shown).

Lloyd's algorithm is iterative and centroid-based, partitioning the dataset X containing n observations into K distinct groups C_K, such that each day is assigned to the cluster with the nearest centroid U_K. The goal is to reduce the total within-cluster variance by minimizing the sum of the squared distance between each data point and the center of its assigned cluster. Mathematically, this is expressed as:

$$\Sigma_{k=1}^{K} \Sigma_{x_n \epsilon C_k} \left|| x_n - \mu_k \right||^2 \quad (1)$$

Starting from an initial set of centroids, the algorithm first assigns each data point to the cluster whose centroid is nearest in terms of distance:

$$C_K = \left\{ x_n : \left|| x_n - \mu_k \right|| \leq all \left|| x_n - \mu_l \right|| \right\} \quad (2)$$

Then, for the given clusters, the centroids are repositioned to the mean of all points belonging to each cluster:

$$u_k = \frac{1}{C_k} \Sigma_{x_n \epsilon C_k} x_n \quad (3)$$

This iterative process continues until they converge, minimising the sum of squares in the clustering arrangement. We determined the optimal number of clusters ($k = 5$) ~~using~~ based on the peak in the silhouette coefficient (Supplementary Figure S3) (Rousseeuw, 1987). We further tested ($k = 4$) and ($k = 6$) observing that the centroids derived from the ($k = 4$) represents a coarser partitioning of the ($k = 5$) regime space. In contrast, ~~two of the five clusters (~~$k = 5$) ~~at (merged~~ ($k = 4$) ~~single cluster at while one cluster from (split into two distinct clusters at~~the ($k = 6$) solution yields a reduction in the silhouette score and primarily results in subdivision of the existing regimes. (Supplementary Figs. S4 and S5). Importantly, the ($k = 5$) solution retains an additional zonal-like circulation regime that is not resolved at($k = 4$) and which is dynamically distinct from the other clusters.

2.2.3 Horizontal thermal advection

Horizontal thermal advection was calculated using ERA5 reanalysis data at hourly resolution (see Section 2.1) following the formula:

$$-v.\nabla_p T = -\left(u\left(\frac{\delta T}{\delta x}\right)_p + v\left(\frac{\delta T}{\delta y}\right)_p\right) \quad (4)$$

Where ∇_p denotes the horizontal gradient on a constant pressure surface, u and v represent the zonal and meridional components of wind and T represents air temperature.

Temperature gradients are calculated using centred finite differences on the native grid, with grid spacing in units derived from spherical geometry assuming an Earth radius of 6371km. The results were converted to units of K day^{-1}. To minimise the influence of orographic effects, the analysis was performed on a constant pressure level of 300 hPa. Sensitivity test using alternative tropospheric pressure levels including 500hPa produced qualitatively similar results, although lower levels exhibited a greater influence of topography. The horizontal thermal advection diagnostic should not be interpreted as a complete temperature tendency budget, as it does not account for adiabatic processes associated with vertical motion or diabatic heating and latent heat release, which have been shown to contribute to observed temperature extremes (e.g. Röthlisberger and Papritz, 2023).

3 Results

3.1 Spatial patterns of extreme high temperature events

The composite 2-m temperature anomalies for high temperature extremes show spatially coherent positive temperature anomalies across the AP (Figure 1a), with typical magnitudes exceeding 4 K. Figure 1b shows the extreme events identified based on the 95th percentile of AP average 2-m temperature coincide with local temperature anomalies above the 95th percentile on average at least 50% of the time at most locations. Coherence is strongest across the central and southern AP, with somewhat weaker correspondence toward the northern tip. The presence, on average, of high temperature anomalies across the entire AP in the set of extreme events confirms they are typically large-scale events that are likely to have a primary origin from synoptic scale atmospheric circulation drivers.




Figure 1: (a) Composite daily mean 2m temperature anomalies over the Antarctic Peninsula on summer days where the temperature exceeds the 95th percentile. (b) Percentage of extreme events (see Methods) from (a) during which the local 2m temperature exceeds the local 95th percentile threshold at least once.

3.2 Circulation clusters for high temperature events

To motivate the applicability of k-means clustering to distinguish circulation patterns associated with extreme high temperature events over the AP, we first review three well-documented case studies. Figure 2a–c shows daily 2-m temperature anomalies for: 6-11th February 2020 - the record breaking heatwave over AP triggered by deep convection in the central tropical Pacific (González-Herrero et al., 2022; Clem et al., 2022); 31st December 1988; and 25th January 1985 - corresponding to extreme warm days

recorded at the Faraday/Vernadsky station on the west of the Peninsula (Turner et. al., 2020). The temperature anomalies during the three events exhibit pronounced spatial heterogeneity and duration. Coincident with the February 2020 event (Fig. 2a), positive anomalies exceeding +5 K were observed across Southwest and West Antarctica (WA), whereas over Dronning Maud Land (DML) and East Antarctica (EA) temperatures remained near climatology. The AP warming coincided with an anomalous ridge positioned to the west of the AP and associated westerly wind anomalies of approximately 3 m s^{-1} (Fig. 2d). In contrast, the extreme warm event in December 1988 (Fig. 2b) was more confined to the AP, with coincident negative temperature anomalies across central and EA and weaker warm anomalies over Queen Mary Land and the Ross Ice Shelf. This event is characterised by a dipole MSLP anomaly straddling the AP, with a ridge extending along its axis and anomalous northwesterly flow exceeding 5 m s^{-1}. The January 1985 event (Fig. 2c) coincided with widespread positive temperature anomalies that extended across EA. The corresponding MSLP and near-surface wind anomalies (Fig. 2f) resemble those of the December 1988 event, although the high-pressure centre is displaced closer to the AP and shifted westward. The contrasting spatial footprints and circulation anomalies associated with these events serve to highlight the diversity of regional circulation patterns driving extreme high temperatures on the AP and their related anomalies across the continent.




Figure 2: Three extreme warm summer events recorded across AP. (a, d) 6th-11th February 2020, (b, e) 31st December 1988, (c, f) 25th January 1985. (a-c) Shading: daily near-surface temperature anomalies; contours: MSLP anomalies (interval 2 hPa, solid = positive, dashed = negative). (d-f) Shading: 500 hPa geopotential height anomalies; vectors: near-surface winds [domain 280°E - 320°E, 60°S - 90°S].

The MSLP anomalies associated with the five k-means cluster centroids are presented in Figure 3a. Cluster 1 is characterised by a high pressure anomaly centred over the Drake Passage and a low pressure anomaly centred between the Amundsen Sea and Ross Sea, with peaks in the mean power spectrum at zonal wavenumbers 1 and 3. (Fig. 3ai, Fig. 3b). Cluster 2 exhibits a dipole structure straddling the AP, consistent with a zonal wavenumber-2 pattern (Fig. 3aii). Cluster 3 features a broad low pressure anomaly centred over the Amundsen Sea with negative anomalies extending over continental WA, with a peak in the mean power spectrum at zonal wavenumber 1 (Fig. 3aiii, Fig. 3b). Cluster 4 shows a peak in the mean power spectrum at zonal wavenumber 2 and is associated with high pressure across WA and the eastern Bellingshausen Sea and a pronounced anticyclone centred northeast of the AP (Fig. 3aiv, Fig. 3b). ~~Cluster 5 displays a zonal wavenumber 3 pattern including a dipole flanking the AP (Fig. 3av).~~ Cluster 5 exhibits a wave-like circulation pattern with a dipole structure

flanking the Antarctic Peninsula. The zonal wavenumber spectrum indicates broadly enhanced power across low wavenumbers (ZW2–ZW4), consistent with a broadband wave structure rather than a single dominant zonal mode (Fig. 3av and Fig. 3b).




Figure 3: (a) Centroids (in hPa) and their associated wind vectors (in m s^{-1}) for the 5 clusters shown across the hemisphere over which they were defined [50°S - 90°S, 180°E - 360°E]. (b)

Composite average zonal wavenumber power spectrum for each cluster calculated over the hemipshere [50°S - 90°S, 180°E - 360°E]. Shading represents ±1 standard error of the mean.

3.3 Temporal behaviour of high temperature events by cluster




Figure 4: (a) Histogram showing the duration of each extreme event separated by each k-means cluster. (b) Histogram showing the area-weighted mean 2-m temperature anomaly over the Antarctic Peninsula across all the days of the extreme high temperature events.

Figure 4 summarises the differences between clusters in terms of event duration and the associated AP temperature anomalies, with the accompanying statistics presented in Table 1. Clear distinctions emerge in both the persistence and intensity of extreme high temperature events between the clusters. Cluster 1 comprises the longest and most intense events with durations ranging from 1 to 7 days and the largest mean AP temperature anomaly. In contrast, Clusters 2–4 represent intermediate-duration events with weaker mean anomalies compared to those from Cluster 1. Cluster 5, which contains the second largest number of events, is characterised by the shortest durations and the weakest mean anomalies. To quantify within-cluster variability, we calculated the mean Euclidean distance between individual samples and their assigned centroid. Larger clusters generally exhibit lower mean distances (not shown), indicating that the most frequently occurring circulation regimes are also among the most dynamically coherent.

	Number of events	Mean length (days)	Mean temperature

	(Percentage of total)		anomaly (K)
Cluster 1	11 (13%)	3.3±1.2	5.5±0.6
Cluster 2	26 (30%)	2.6±0.8	4.6±0.2
Cluster 3	11 (13%)	2.1±1.0	4.5±0.4
Cluster 4	15 (17%)	2.7±0.8	4.4±0.2
Cluster 5	24 (28%)	1.9±0.6	4.3±0.2

Table 1: Summary of event statistics by cluster, including event duration and associated temperature anomalies, highlighting differences in cluster characteristics. Uncertainty values represent 95% confidence range.

Figure 5a shows the temporal evolution of AP-mean temperature anomalies during extreme high temperature events for each cluster, with the corresponding temperature growth rates, calculated using centred finite differences, shown in Fig. 5b. Temperature anomalies remain below 2 K for all clusters until two days prior to onset. At this point, Clusters 3–5 exhibit steadily increasing anomalies, reaching peaks of approximately 4.0–4.5 K at day 0. By contrast, Cluster 1 shows weaker anomalies during days −2 to −1, followed by a rapid intensification in the final day before onset, with growth rates approaching ~2 K day^{-1} and peak anomalies near 5 K. Cluster 2 also exhibits rapid pre-onset growth but attains a slightly lower peak (~4.3 K), representing, on average, a fast-developing yet more moderate extreme event.




Figure 5: (a) Mean temperature anomaly across the Antarctic Peninsula for the events in each cluster between days -5 and +3. (b) As in (a) but for average temperature growth rate (dT/dt), calculated using a centred finite-difference scheme method. Shading represents ±1 standard error of the mean. (c) Fraction of events in each cluster whose circulation patterns are assigned to all clusters as a function of lag. Percentages shown in the bars give the proportion of days classified to each cluster. Bars are ordered by decreasing fraction.

Figure 5c shows the fraction of events assigned to each cluster at day 0 that are assigned to all clusters at different lead and lag times. It should not be interpreted as the evolution of individual events, but rather as an aggregate measure of cluster coherence through time. It therefore quantifies how consistently each synoptic regime is retained relative to its day-0 classification. By construction, 100% of events are assigned to their respective clusters at day 0. Only 55% of Cluster 1 events are assigned to the same cluster at day −5, indicating substantial variability in circulation in the lead-up to the extreme event. Cluster coherence on day −5 is lower for Clusters 2 (28%) and 3 (27%), but higher for Cluster 4 (53%). Cluster 3 exhibits higher coherence at day −3 (78%), although this fraction declines at days −2 and −1, suggesting that the circulation patterns associated with many events are located between centroids prior to onset. Similar temporal evolution is observed for Clusters 1, 4, and 5 in the days preceding onset, with Cluster 5 showing the highest persistence, as 70% of events retain their cluster assignment at day −3. Together, these results indicate that some clusters represent relatively stable, long-lived large-scale circulation anomalies (e.g., Cluster 5), whereas others are characterised by rapidly evolving or transitional synoptic conditions leading up to extreme high temperature events.

To further explore this temporal evolution, Figure 6 shows composites of MSLP and 2-m temperature anomalies for the days preceding, during and following the onset of the extreme high temperature events for each cluster. The composite MSLP anomalies closely resemble the respective centroids, reinforcing the robustness of the k-means classification. The associated temperature anomaly patterns display substantial differences in spatio-temporal structure between the clusters.




Figure 6: Time-averaged lagged composites of events in each cluster for (a-e) days -5 to -3; (f-j) days –2 to 0 and (k-o) days 1 to 3. Day 0 = first day of the extreme event. Shading: near-surface temperature anomalies; contours: MSLP anomalies (interval 2 hPa, solid = positive, dashed = negative). Non-significant anomalies (p > 0.05) are not plotted.

Fig. 6a-e shows composite anomalies over days −5 to −3 prior to the onset of extreme warm conditions across AP. In Cluster 1, the warming over WA evident at event onset is already established, though warm anomalies are absent from AP (Fig. 6a). During Cluster 2 events, anomalies remain close to climatological values over days -5 to -3 prior to onset (Fig. 6b). For Cluster 3, there is warming evident in the Bellingshausen Sea and cooling across the Weddell Sea evident in the lead-up period (Fig. 6c). In Cluster 4, there is continental wide warm anomalies with the exception of parts of WA prior to the onset of

extreme high temperatures on the AP (Fig. 6d). Temperature anomalies associated with Cluster 5 are broadly close to climatological values prior to the onset of extreme high temperatures on the AP (Fig. 6e).

During the event onset phase (days -2 to 0), Cluster 1 exhibits widespread warm anomalies across the AP, extending into WA and across the Ross Ice Shelf, while cold anomalies are present over Dronning Maud Land (Fig. 6f). Following the event onset (days 1-3), the warm temperature anomalies extend further across to East Antarctica (Fig. 6k) while the cold anomalies near the Dronning Maud Land return towards climatology. Cluster 2 is characterised by warming over the AP and the Bellingshausen Sea, accompanied by cooling at the South Pole and across coastal East Antarctica including Dronning Maud Land (Fig. 6g). The warm anomalies expand into the Ross Ice Shelf area in the period after event onset, with areas of EA characterised by an intensification of their cold anomalies (Fig. 6l). In Cluster 3, the warm anomalies are largely restricted to the AP and WA, while cooling is observed over Wilkes Land (Fig. 6h). After the event onset, warming is largely confined to the same region whilst the cold anomalies found over Wilkes Land return towards climatology (Fig. 6m). Cluster 4 exhibits extensive warm anomalies spanning most of the Antarctic continent (Fig. 5i), which persist in the days following event onset (Fig. 6n). Cluster 5 shows warming over the AP, Weddell Sea and Bellingshausen Sea and cooling over much of EA (Fig. 6j). Following the event, the warm anomalies grow across WA whilst cold anomalies persist over much of EA (Fig. 6o).

Figure 7 shows composite 500 hPa geopotential height anomalies and 10-m wind anomalies over the AP for days −5 to −3, −2 to 0, and 1 to 3 relative to the onset of high AP temperature events. Across all clusters, there are positive mid-tropospheric height anomalies over or near the AP, indicating the presence of an anomalous ridge, although the amplitude and position of the ridge varies between clusters.




Figure 7: As in Figure 6, but for 500 hPa geopotential height anomalies (shading) and near-surface wind speed anomalies (vectors) for the sector 280°E - 320°E, 60°S - 90°S. Non-significant anomalies (p > 0.05) are not plotted.

~~During the preconditioning phase (days −5 to −3), an anomalous ridge is established close to the AP, with the high pressure centre located to the west and migrating eastward toward the AP by the onset phase (days −2 to 0). An exception occurs in Clusters 4 and 5, where the high pressure centre is positioned to the east of the AP during preconditioning, placing the AP beneath the western flank of the ridge. 10-m wind anomalies exhibit substantial inter-cluster differences in both magnitude and direction. Cluster 1 is characterised by strong (~5 m s^{-1}) and statistically significant northerly anomalies across the Bellingshausen Sea, Weddell Sea, and northern AP during event onset (Fig. 7l). These anomalies are approximately twice the magnitude of the climatological winds in this region, consistent with enhanced meridional warm-air advection. The wind anomalies weaken in the days following onset, while the ridge remains quasi-stationary. Clusters 3 and 4 also exhibit significant northerly anomalies of comparable magnitude, centred primarily over the Bellingshausen Sea (Cluster 3) and the Weddell Sea (Cluster 4) (Fig. 7h,m). Following onset, northerly anomalies in Cluster 4 intensify and expand to encompass much of the AP, whereas in Cluster 3 the ridge weakens and northerly anomalies largely dissipate. In contrast, Cluster 2 displays anomalous southerly near-surface winds offshore to the east of the AP (Weddell Sea) during event onset (Fig. 7c), and the associated ridge migrates northward away from the AP in the post-onset period. Cluster 5 exhibits relatively weak and spatially limited northerly anomalies at event onset (Fig. 7d), but strong northerly anomalies during the preconditioning phase (Fig. 7g), indicating a different~~

~~temporal phasing between circulation anomalies and peak warming. These patterns suggest that while elevated mid-tropospheric geopotential heights are a common precursor to extreme warm events, differences in the strength, location, and timing of low-level meridional flow likely modulate the magnitude and spatial footprint of surface warming through thermal advection and interactions with AP topography that favour föhn-like warming in some regions.~~ During the preconditioning phase (days −5 to −3), an anomalous ridge is established close to the AP, with the high-pressure centre located to the west and migrating eastward towards the AP by the onset phase (days −2 to 0). An exception occurs in Clusters 4 and 5, where the high-pressure centre is positioned to the east of the AP during preconditioning, placing the AP beneath the western flank of the ridge. The 10-m wind anomalies exhibit substantial inter-cluster differences in both magnitude and direction. Cluster 1 is characterised by strong (~5 m s^{-1}) and statistically significant anticyclonic anomalies across the Bellingshausen Sea and AP during event onset, with southerlies evident across the Weddell Sea (Fig. 7f). These anomalies are approximately twice the magnitude of the climatological winds in this region, consistent with enhanced meridional warm-air advection. The wind anomalies persist in the days following onset, while the ridge which remains quasi-stationary. Clusters 3 and 4 also exhibit significant anticyclonic wind anomalies of comparable magnitude, with ridges centred primarily over the Bellingshausen Sea (Cluster 3) and the northern AP Sea (Cluster 4) (Fig. 7h,i). Following onset, the ridge in Cluster 4 persists over the AP, albeit with reduced intensity, whereas the anomalous ridge in Cluster 3 dissipates. Cluster 2 displays anomalous northerly near-surface winds across the AP and to the east of the AP (Weddell Sea) during event onset (Fig. 7c), and the associated ridge migrates northward away from the AP in the post-onset period. Cluster 5 exhibits a ridge centred over the Weddell Sea during event onset, with associated anticyclonic wind anomalies, with the ridge migrating further northeast after onset. These patterns suggest that while elevated mid-tropospheric geopotential heights are a common precursor to extreme warm events, differences in the strength, location, and timing of low-level meridional flow likely modulate the magnitude and spatial footprint of surface warming through thermal advection and interactions with AP topography that favour föhn-like warming in some regions.

3.4 Role of thermal advection for high temperature events

Figure 8 quantifies the anomalous upper tropospheric horizontal thermal advection associated with extreme high AP temperature events for each cluster. In the days immediately preceding event onset, all clusters exhibit pronounced positive horizontal thermal advection to the west of the AP, locally exceeding 4 K day^{-1} (Fig. 8f–j). These strong warming tendencies represent a marked intensification from the preconditioning period, during which comparable advection magnitudes occur only in Cluster 3, across the Bellingshausen Sea and centred near ~50°S west of South America (Fig. 8c). Although all clusters show anomalous warming tendencies in the region of the AP, as anticipated, their spatial structures differ.

Cluster 5 displays weaker and more geographically confined warming, with anomalies of ~3 K day^{-1} largely restricted to the AP region. In contrast, Clusters 1–4 exhibit broader horizontal advection patterns extending from further west across the Bellingshausen Sea. Across all clusters, anomalous negative meridional advection (cooling) occurs over the Weddell Sea, consistent with anomalous southerly flow in that region.




Figure 8: As in Figure 6, but for anomalies in 300 hPa horizontal thermal advection (shading) and wind speed (vectors) for the sector 105ºW - 45ºW , 45ºS - 80ºS.

Following event onset (Fig. 8k–o), more pronounced cluster-dependent behaviour emerges. Cluster 2 shows widespread positive advection across much of the AP, whereas Clusters 1 and 4 continue to be dominated by warming originating from the lower-latitude Bellingshausen Sea. In Cluster 3, the advection signal weakens across the Bellingshausen Sea region substantially after onset whilst showing warming in the Weddell Sea and north of AP; this is also seen in Cluster 5.

A decomposition of the anomalous upper tropospheric thermal advection for all high temperature events reveals that the meridional component is dominant and this is largely explained by anomalous meridional wind anomalies (Fig. S6). In the zonal component, zonal wind anomalies contribute more than variations in the zonal temperature gradient. These results indicate that anomalous warming during extreme events is driven primarily by changes in flow strength and direction, rather than by modulation of the background temperature gradients.

Together, these results provide further evidence that enhanced poleward heat transport via anomalous meridional flow is a fundamental mechanism underpinning extreme warm events across all clusters, while differences in the location, magnitude, and persistence of this advection help put into context the contrasting growth rates, spatial footprints, and lifetimes of temperature anomalies.

4 Discussion

González et al. (2018) identified five recurrent synoptic-scale circulation patterns over the Drake Passage and AP region using a similar method applied over the full annual cycle. These included patterns characterised by low pressure in the Weddell Sea, Amundsen–Bellingshausen Seas, and Drake Passage, as well as zonal flow and a ridge over the AP. Our results suggest that extreme high AP temperatures in austral summer are associated with a subset of the broad circulation regimes identified by González et al. (2018), particularly those featuring ridging over or immediately upstream of the AP. To draw further comparison, they find that a cluster featuring a low over the Drake Passage is the most frequent pattern during austral summer, followed by a more zonally oriented flow, which qualitatively agrees with our findings where Cluster 5 is the most abundant cluster, albeit with its low centre of action located further north. Furthermore, we identify circulation patterns specifically associated with extreme high AP temperature events that were not separated in González et al. (2018); for example conditioning on temperature extremes can reveal additional nuances in the patterns of synoptic scale flow during extreme events such as between clusters 2 and 4, which might otherwise be grouped into a single broad cluster. In this sense, the present study extends González et al. (2018) by demonstrating how specific synoptic patterns are selectively associated with temperature extremes, and by documenting the distinct dynamical signatures associated with each extreme-favouring regime.

The hemispheric-scale atmospheric circulation anomalies show signatures consistent with zonal wavenumber-3 and wavenumber-2 patterns, demonstrated in cluster 1 and clusters 2 and 4, respectively, indicating a hemispheric-scale connection through planetary-scale Rossby wave trains between the tropics, midlatitudes and polar regions (e.g., Ding and Steig 2013; Boschat et al., 2023; Goyal et al., 2021). Such wave trains are frequently initiated by anomalous tropical convection associated with ENSO and the MJO (e.g. Lee and Seo, 2019; Schneider et al., 2012), and can promote persistent blocking highs over the South Pacific and Drake Passage that favour warm air advection toward the AP (Rondanelli et al., 2019; Clem et al., 2022). Nevertheless, there is evidence to suggest that there is a spatial dependency regarding the impacts of the different modes of climate variability on regions of the AP (e.g. Clem and Fogt., 2013; Clem et al., 2016). Our results suggest a role for planetary scale modes in driving high AP temperature events. In particular, Cluster 4 exhibits a more persistent and extensive high-pressure anomaly over Antarctica in the lead-up to the onset of extreme warm temperatures across AP, suggestive of a negative SAM-like state, consistent with circulation patterns that are known to weaken the circumpolar westerlies and shift the storm track equatorward~~which is known to weaken the circumpolar~~

~~westerlies and shift the storm track equatorward~~, ~~thereby promoting~~ which can promote warm air intrusions into the AP (Turner et al., 2013; Raphael et al., 2016). This is consistent with well established evidence that variations in SAM strongly modulate AP temperature extremes (Fogt et al., 2012; Clem et al., 2016; Turner et al., 2020).

The spatial footprint of warming differs markedly between clusters, with some exhibiting continent-wide warming while others show warming largely confined to West Antarctica or the AP alone. This behaviour mirrors continental-scale analyses showing that West Antarctic temperature extremes tend to be spatially confined and closely linked to meridional flow anomalies, whereas East Antarctic temperature extremes are often associated with inland high-pressure systems and katabatic wind enhancement (Nielsen et al., 2025). The present results support this evidence and suggest that the AP occupies a dynamical regime in which both large-scale advection and local topographic effects strongly influence extreme temperature expression.

Our results have shown that large-scale thermal advection plays a key role in driving high temperature events on the AP. However, the thermal advection diagnostic presented here does not constitute a complete temperature tendency budget. Adiabatic temperature changes associated with vertical motion and diabatic heating will also contribute in part to Antarctic temperature extremes. Furthermore, latent heat release associated with moist heat fluxes has been identified as an important contributor to Antarctic heat extremes (e.g. Blanchard-Wrigglesworth et al., 2023) but is not explicitly quantified here. Station-based analyses indicate that the majority of record local temperature extremes on the AP occur following flow across steep orography, consistent with an important contribution from föhn warming (Turner et al., 2020). In addition, the presence of high pressure and near-surface wind anomalies extending for several days over the AP, particularly in Clusters 1 and 5, supports a mechanism whereby anomalous anticyclonic flow drives warm, dry northwesterly winds onto the upwind side of the AP, producing föhn-like conditions on the leeward side. ERA5 does not fully resolve the dynamics of föhn flow due to its coarse spatial resolution, however, the presence of northwesterly low-level wind anomalies in several clusters suggests there would be a central role of föhn processes in the high AP temperature events identified as found in various studies (Elvidge et al., 2014; Laffin et al. 2021; Turton et al., 2018). These dynamics are well-established contributors to surface melt and firn air depletion on the AP (e.g. Cape et al., 2015; Zou et al., 2023; Kirchgaessner et al., 2021), and have been implicated in major melt events over ice shelves such as Larsen C (Elvidge et al., 2014). Case study events pertaining to each cluster, using observations and high-resolution modelling tools such as dynamically downscaled regional climate model outputs, should be looked at in future to help understand the relative role of local-scale (föhn wind) from background forcing due to large-scale drivers.

5 Conclusions

This study has identified five distinct circulation patterns associated with extreme high summer temperatures across AP in austral summer using k-means clustering applied to daily MSLP on days when the 2-m AP averaged temperature exceeds the 95th percentile. The resulting clusters exhibit substantial diversity in the patterns of continental 2-m temperature, 10-m wind, and mid-tropospheric geopotential height anomalies, demonstrating that high temperature extremes on the AP are driven by a range of synoptic circulation states. While Cluster 4 shows continent-wide warming, Clusters 1 and 3 show warming confined to WA. The warming in Cluster 2 and 5 is more confined to the AP with coincident cooling across parts of EA. Unsurprisingly, all clusters are associated with a ridge over the northern AP, though the magnitude, orientation and persistence of this feature varies significantly.

The classification of dynamical patterns associated with similar high AP summer temperatures extends prior work on circulation patterns throughout the seasonal cycle (Gonzalez et al., 2018) and continental trends and extremes (Turner et al., 2020; Nielsen et al., 2025) and reveals that extreme high AP temperatures arise not from a single canonical pattern, but from a range of distinct synoptic states. Future work could examine the connection of these states to remote large-scale drivers such as tropical and subtropical convection (e.g., Turner et al., 2016; Pezza et al. 2012), the SAM and the stratospheric polar vortex (e.g., Lim et al., 2018), as well as assess their representation in climate models in order to better understand their predictability across future timescales.

While this study has applied k-means clustering to circulation fields in order to focus on the role of atmospheric dynamics, it may be interesting in the future to include other information in the clustering related to thermodynamic preconditioning and radiative effects, which have been shown to contribute to extreme melt and temperature events (e.g. Gorodetskaya et al., 2023; Wille et al., 2022), and to quantify thermal tendencies from adiabatic descent, turbulent mixing, and cloud-radiative effects.

Data Availability Statement

The ERA5 reanalysis data sets (Hersbach et al., 2020) can be obtained from the Copernicus Climate Data Store (https://cds.climate.copernicus.eu/). All figures are prepared using Python (version 3.11).

Author Contributions

WJD, ACM and ANR designed the study. WJD performed the analysis and produced the figures. WJD wrote the manuscript draft; ACM and ANR edited the manuscript with comments from RSW, TJB.

Competing Interests

The contact author has declared that none of the authors have any competing interests.

Acknowledgements

We are grateful to the wider NERC-funded ExtAnt project team for their contributions to the discussion of the results. We thank Tony Phillips for assistance with data handling. We are also grateful to NERC's JASMIN facility for providing resources to conduct the analysis.

Financial Support

All authors were funded as part of the ExtAnt project (NERC grants NE/Y503307/1 and NE/Y503319/1).

References

Baba, K., & Renwick, J. (2017). Aspects of intraseasonal variability of Antarctic sea ice in austral winter related to ENSO and SAM events. *Journal of Glaciology*, *63*(241), 838–846. https://doi.org/10.1017/JOG.2017.49

Bae, H. J., Kim, S. J., Kim, B. M., & Kwon, H. (2022). Causes of the Extreme Hot Event on February 9, 2020, in Seymour Island, Antarctic Peninsula. *Frontiers in Environmental Science*, *10*, 865775. https://doi.org/10.3389/FENVS.2022.865775

Balay, F., & Renom, M. (2026). Characterization of extreme warm temperature events (1998–2016) on King George Island, Antarctic Peninsula. *Antarctic Science*, *00*, 1–15. https://doi.org/10.1017/S0954102025100552

Baldwin, J. W., Dessy, J. B., Vecchi, G. A., & Oppenheimer, M. (2019). Temporally Compound Heat Wave Events and Global Warming: An Emerging Hazard. *Earth's Future*, *7*(4), 411–427. https://doi.org/10.1029/2018EF000989

Banwell, A. F., Tri Datta, R., Dell, R. L., Moussavi, M., Brucker, L., Picard, G., Shuman, C. A., & Stevens, L. A. (2021). The 32-year record-high surface melt in 2019/2020 on the northern George VI Ice Shelf, Antarctic Peninsula. *Cryosphere*, *15*(2), 909–925. https://doi.org/10.5194/TC-15-909-2021,

Barrett, B. S., Lafleur, D. M., & Henderson, G. R. (2025). Madden–Julian Oscillation Modulation of Antarctic Sea Ice. *Glacies 2025, Vol. 2*, *2*(4), 16. https://doi.org/10.3390/GLACIES2040016

Bevan, S., Luckman, A., Hendon, H., & Wang, G. (2020). The 2020 Larsen C Ice Shelf surface melt is a 40-year record high. *Cryosphere*, *14*(10), 3551–3564. https://doi.org/10.5194/TC-14-3551-2020,

Boschat, G., Purich, A., Rudeva, I., & Arblaster, J. (2023). Impact of Zonal and Meridional Atmospheric Flow on Surface Climate and Extremes in the Southern Hemisphere. *Journal of Climate*, *36*(15), 5041–5061. https://doi.org/10.1175/JCLI-D-22-0251.1

Bozkurt, D., Bromwich, D. H., Carrasco, J., Hines, K. M., Maureira, J. C., & Rondanelli, R. (2020). Recent Near-surface Temperature Trends in the Antarctic Peninsula from Observed, Reanalysis and Regional Climate Model Data. *Advances in Atmospheric Sciences*, *37*(5), 477–493. https://doi.org/10.1007/S00376-020-9183-X/

Bozkurt, D., Marín, J. C., Verdugo, C., & Barrett, B. S. (2024). Atmospheric blocking and temperatures in the Antarctic Peninsula. *Science of The Total Environment*, *931*, 172852. https://doi.org/10.1016/J.SCITOTENV.2024.172852

Bracegirdle, T. J., Caton Harrison, T., Holmes, C. R., Lu, H., Martineau, P., & Phillips, T. (2024). Antarctic extreme seasons under 20th and 21st century climate change. *Npj Climate and Atmospheric Science*, *7*(1), 1–12. https://doi.org/10.1038/S41612-024-00822-Y

Cape, M. R., Vernet, M., Skvarca, P., Marinsek, S., Scambos, T., & Domack, E. (2015). Foehn winds link climate-driven warming to ice shelf evolution in Antarctica. *Journal of Geophysical Research: Atmospheres*, *120*(21), 11,037-11,057. https://doi.org/10.1002/2015JD023465

Carrasco, J. F., Bozkurt, D., & Cordero, R. R. (2021). A review of the observed air temperature in the Antarctic Peninsula. Did the warming trend come back after the early 21st hiatus? *Polar Science*, *28*, 100653. https://doi.org/10.1016/j.polar.2021.100653

Clem, K. R., Renwick, J. A., McGregor, J., & Fogt, R. L. (2016). The relative influence of ENSO and SAM on Antarctic Peninsula climate. *Journal of Geophysical Research: Atmospheres*, *121*(16), 9324–9341.https://doi.org/10.1002/2016JD025305

Clem, K.R., Bozkurt, D., Kennett, D. et al. Central tropical Pacific convection drives extreme high temperatures and surface melt on the Larsen C Ice Shelf, Antarctic Peninsula. *Nat. Commun* 13, 3906 (2022). https://doi.org/10.1038/s41467-022-31119-4)

Clem, K. R., & Fogt, R. L. (2013). Varying roles of ENSO and SAM on the Antarctic Peninsula climate in austral spring. *Journal of Geophysical Research Atmospheres*, *118*(20), 11,481-11,492. https://doi.org/10.1002/JGRD.50860

Clem, K. R., Renwick, J. A., McGregor, J., & Fogt, R. L. (2016). The relative influence of ENSO and SAM on Antarctic Peninsula climate. *Journal of Geophysical Research*, *121*(16), 9324–9341. https://doi.org/10.1002/2016JD025305;PAGE:STRING:ARTICLE/CHAPTER

Coggins, J. H. J., McDonald, A. J., & Jolly, B. (2014). Synoptic climatology of the Ross Ice Shelf and Ross Sea region of Antarctica: K-means clustering and validation. *International Journal of Climatology*, *34*(7), 2330–2348. https://doi.org/10.1002/JOC.3842

DeConto, R. M., & Pollard, D. (2016). Contribution of Antarctica to past and future sea-level rise. *Nature*, *531*(7596), 591–597. https://doi.org/10.1038/NATURE17145

Deb, P., Bromwich, D., Orr, A., Sen, A., & Clem, K. R. (2025). Recent increase in surface melting of West Antarctic ice shelves linked to Interdecadal Pacific Oscillation. *Communications Earth & Environment 2025 6:1*, *6*(1), 99–. https://doi.org/10.1038/s43247-025-02077-8

Ding, Q., & Steig, E. J. (2013). Temperature Change on the Antarctic Peninsula Linked to the Tropical Pacific. *Journal of Climate*, *26*(19), 7570–7585. https://doi.org/10.1175/JCLI-D-12-00729.1

Ding, Q., Steig, E., Battisti, D., & Küttel, M. (2011). Winter warming in West Antarctica caused by central tropical Pacific warming. *Nature Geoscience*. https://doi.org/10.1038/NGEO1129

Dunmire, D., Wever, N., Banwell, A. F., & Lenaerts, J. T. M. (2024). Antarctic-wide ice-shelf firn emulation reveals robust future firn air depletion signal for the Antarctic Peninsula. *Communications Earth and Environment*, *5*(1), 1–13. https://doi.org/10.1038/S43247-024-01255-4

Elvidge, A. D., Renfrew, I. A., King, J. C., Orr, A., Lachlan-Cope, T. A., Weeks, M., & Gray, S. L. (2014). Foehn jets over the Larsen C Ice Shelf, Antarctica. *Quarterly Journal of the Royal Meteorological Society*, *141*(688), 698–713. https://doi.org/10.1002/QJ.2382

England, M. R., Polvani, L. M., Smith, K. L., Landrum, L., & Holland, M. M. (2016). Robust response of the Amundsen Sea Low to stratospheric ozone depletion. *Geophysical Research Letters*, *43*(15), 8207–8213. https://doi.org/10.1002/2016GL070055

Feron, S., Cordero, R. R., Damiani, A., Malhotra, A., Seckmeyer, G., & Llanillo, P. (2021). Warming events projected to become more frequent and last longer across Antarctica. *Scientific Reports*, *11*(1), 1–9. https://doi.org/10.1038/S41598-021-98619-Z

Fogt, R. L., Bromwich, D. H., & Hines, K. M. (2010). Understanding the SAM influence on the South Pacific ENSO teleconnection. *Climate Dynamics 2010 36:7*, *36*(7), 1555–1576. https://doi.org/10.1007/S00382-010-0905-0

Fogt, R. L., Jones, J. M., & Renwick, J. (2012). Seasonal Zonal Asymmetries in the Southern Annular Mode and Their Impact on Regional Temperature Anomalies. *Journal of Climate*, *25*(18), 6253–6270. https://doi.org/10.1175/JCLI-D-11-00474.1

Fogt, R. L., & Marshall, G. J. (2020). The Southern Annular Mode: Variability, trends, and climate impacts across the Southern Hemisphere. *Wiley Interdisciplinary Reviews: Climate Change*, *11*(4), e652. https://doi.org/10.1002/WCC.652;ISSUE:ISSUE:DOI

Hosking, J. S., Orr, A., Marshall, G. J., Turner, J., & Phillips, T. (2013). The Influence of the Amundsen–Bellingshausen Seas Low on the Climate of West Antarctica and Its Representation in Coupled Climate Model Simulations. *Journal of Climate*, *26*(17), 6633–6648. https://doi.org/10.1175/JCLI-D-12-00813.1

Gonzalez, S., & Fortuny, D. (2018). How robust are the temperature trends on the Antarctic Peninsula? *Antarctic Science*, *30*(5), 322–328. https://doi.org/10.1017/S0954102018000251

Gonzalez, S., Vasallo, F., Recio-Blitz, C., Guijarro, J. A., & Riesco, J. (2018). Atmospheric Patterns over the Antarctic Peninsula. *Journal of Climate*, *31*(9), 3597–3608. https://doi.org/10.1175/JCLI-D-17-0598.1

González-Herrero, S., Barriopedro, D., Trigo, R. M., López-Bustins, J. A., & Oliva, M. (2022). Climate warming amplified the 2020 record-breaking heatwave in the Antarctic Peninsula. *Communications Earth & Environment 2022 3:1*, *3*(1), 122 . https://doi.org/10.1038/s43247-022-00450-5

González-Herrero, S., Vasallo, F., Bech, J., Gorodetskaya, I., Elvira, B., & Justel, A. (2023). Extreme precipitation records in Antarctica. *International Journal of Climatology*, *43*(7), 3125–3138. https://doi.org/10.1002/JOC.8020

Gorodetskaya, I. v, Durán-Alarcón, C., González-Herrero, S., Clem, K. R., Zou, X., Rowe, P., Rodriguez-Imazio, P., Campos, D., Leroy, C., Santos, -Dos, Dutrievoz, N., Wille, J. D., Chyhareva, A., Favier, V.,

Blanchet, J., Pohl, B., Cordero, R. R., Park, S.-J., Colwell, S., … Picard, G. (2023). Record-high Antarctic Peninsula temperatures and surface melt in February 2022: a compound event with an intense atmospheric river. *Npj Climate and Atmospheric Science 2023 6:1*, *6*(1), 1–18. https://doi.org/10.1038/s41612-023-00529-6

Goyal, R., Jucker, M., sen Gupta, A., Hendon, H. H., & England, M. H. (2021). Zonal wave 3 pattern in the Southern Hemisphere generated by tropical convection. *Nature Geoscience 2021 14:10*, *14*(10), 732–738. https://doi.org/10.1038/s41561-021-00811-3

Guarino, M.-V., Ridley, J. K., Colwell, S., Farneti, R., Giuliani, G., Hindley, N., King, J., Kucharski, F., Polichtchouk, I., Tompkins, A. M., Vignon, É., Wright, C., Guarino, M.-V., Ridley, J. K., Colwell, S., Farneti, R., Giuliani, G., Hindley, N., King, J., … Wright, C. (2026). A Long-Term Shift in Flow Regimes over the Antarctic Peninsula. *Journal of Climate*, *39*(2), 749–767. https://doi.org/10.1175/jcli-d-25-0330.1

Hillebrand, F. L., Bremer, U. F., Arigony-Neto, J., da Rosa, C. N., Mendes, C. W., Costi, J., de Freitas, M. W. D., & Schardong, F. (2021). Comparison between Atmospheric Reanalysis Models ERA5 and ERA-Interim at the North Antarctic Peninsula Region. *Annals of the American Association of Geographers*, *111*(4), 1147–1159. https://doi.org/10.1080/24694452.2020.1807308

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., … Thépaut, J. N. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, *146*(730), 1999–2049. https://doi.org/10.1002/qj.3803

Jones, M. E., Bromwich, D. H., Nicolas, J. P., Carrasco, J., Plavcová, E., Zou, X., & Wang, S. H. (2019). Sixty Years of Widespread Warming in the Southern Middle and High Latitudes (1957–2016). *Journal of Climate*, *32*(20), 6875–6898. https://doi.org/10.1175/JCLI-D-18-0565.1

Kirchgaessner, A., King, J. C., & Anderson, P. S. (2021). The Impact of Föhn Conditions Across the Antarctic Peninsula on Local Meteorology Based on AWS Measurements. *Journal of Geophysical Research: Atmospheres*, *126*(4), e2020JD033748. https://doi.org/10.1029/2020JD033748

Laffin, M. K., Zender, C. S., Singh, S., van Wessem, J. M., Smeets, C. J. P. P., & Reijmer, C. H. (2021). Climatology and Evolution of the Antarctic Peninsula Föhn Wind-Induced Melt Regime From 1979–2018. *Journal of Geophysical Research: Atmospheres*, *126*(4), e2020JD033682. https://doi.org/10.1029/2020JD033682

Lee, H. J., & Seo, K. H. (2019). Impact of the Madden-Julian oscillation on Antarctic sea ice and its dynamical mechanism. *Scientific Reports 2019 9:1*, *9*(1), 10761–. https://doi.org/10.1038/s41598-019-47150-3

Li, X., Gerber, E. P., Holland, D. M., & Yoo, C. (2015). A Rossby Wave Bridge from the Tropical Atlantic to West Antarctica. *Journal of Climate*, *28*(6), 2256–2273. https://doi.org/10.1175/JCLI-D-14-00450.1

Li, X., Holland, D. M., Gerber, E. P., & Yoo, C. (2014). Impacts of the north and tropical Atlantic Ocean on the Antarctic Peninsula and sea ice. *Nature*, *505*(7484), 538–542. https://doi.org/10.1038/NATURE12945;TECHMETA

Lim, E. P., Hendon, H. H., & Thompson, D. W. J. (2018). Seasonal Evolution of Stratosphere-Troposphere Coupling in the Southern Hemisphere and Implications for the Predictability of Surface Climate. *Journal of Geophysical Research: Atmospheres*, *123*(21), 12,002-12,016. https://doi.org/10.1029/2018JD029321

Marshall, G. J. (2003). Trends in the southern annular mode from observations and reanalyses. Journal of Climate, 16(24), 4134–4143. https://doi.org/10.1175/1520- 0442(2003)016<4134:TITSAM>2.0.CO;2

Marshall, G. J. (2007). Half-century seasonal relationships between the Southern Annular Mode and Antarctic temperatures. *International Journal of Climatology*, *27*(3), 373–383. https://doi.org/10.1002/JOC.1407

Marshall, G. J., Fogt, R. L., Turner, J., & Clem, K. R. (2022). Can current reanalyses accurately portray changes in Southern Annular Mode structure prior to 1979? *Climate Dynamics 2022 59:11*, *59*(11), 3717–3740. https://doi.org/10.1007/s00382-022-06292-3

Marshall, G. J., Orr, A., van Lipzig, N. P. M., & King, J. C. (2006). The Impact of a Changing Southern Hemisphere Annular Mode on Antarctic Peninsula Summer Temperatures. *Journal of Climate*, *19*(20), 5388–5404. https://doi.org/10.1175/JCLI3844.1

Montaño Bello, D., Villegas Bolaños, N. L., (2025). Influence of ENSO and SAM on the occurrence of absolute extremes and behavior of Ta, SST, and SIC anomalies in the western Antarctic Peninsula. *Atmósfera*, *39*, 437–450. https://doi.org/10.20937/ATM.53423

Nielsen, E. B., Katurji, M., Zawar-Reza, P., & Cullen, N. (2025). Air Temperature Trends and Extreme Warming Events Across Regions of Antarctica for the Period 2003–2021. *Journal of Geophysical Research: Atmospheres*, *130*(9), e2024JD043042. https://doi.org/10.1029/2024JD043042

Oliva, M., Navarro, F., Hrbáček, F., Hernández, A., Nývlt, D., Pereira, P., Ruiz-Fernández, J., & Trigo, R. (2017). Recent regional climate cooling on the Antarctic Peninsula and associated impacts on the cryosphere. *Science of The Total Environment*, *580*, 210–223. https://doi.org/10.1016/J.SCITOTENV.2016.12.030

Orr, A., Marshall, G. J., Hunt, J. C. R., Sommeria, J., Wang, C. G., van Lipzig, N. P. M., Cresswell, D., & King, J. C. (2008). Characteristics of Summer Airflow over the Antarctic Peninsula in Response to Recent Strengthening of Westerly Circumpolar Winds. *Journal of the Atmospheric Sciences*, *65*(4), 1396–1413. https://doi.org/10.1175/2007JAS2498.1

Pezza, A. B., Rashid, H. A., & Simmonds, I. (2012). Climate links and recent extremes in antarctic sea ice, high-latitude cyclones, Southern Annular Mode and ENSO. *Climate Dynamics*, *38*(1–2), 57–73. https://doi.org/10.1007/S00382-011-1044-Y/

Raphael, M. N., Marshall, G. J., Turner, J., Fogt, R. L., Schneider, D., Dixon, D. A., Hosking, J. S., Jones, J. M., & Hobbs, W. R. (2016). The Amundsen Sea Low: Variability, Change, and Impact on Antarctic Climate. *Bulletin of the American Meteorological Society*, *97*(1), 111–121. https://doi.org/10.1175/BAMS-D-14-00018.1

Renwick JA, Revell MJ (1999) Blocking over the South Pacific and Rossby wave propagation. Mon Weather Rev 127(10):2233–2247. https://doi.org/10.1175/1520-0493(1999)127<2233:BOTSPA>2.0.CO;2

Rondanelli, R., Hatchett, B., Rutllant, J., Bozkurt, D., & Garreaud, R. (2019). Strongest MJO on record triggers extreme Atacama rainfall and warmth in Antarctica. Geophysical Research Letters, 46, 3482-3491. https://doi.org/10.1029/2018GL081475

Rousseeuw, P. J. (1987). Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. *Journal of Computational and Applied Mathematics*, *20*(C), 53–65. https://doi.org/10.1016/0377-0427(87)90125-7

Ruosteenoja, K., & Jylhä, K. (2023). Average and extreme heatwaves in Europe at 0.5–2.0 °C global warming levels in CMIP6 model simulations. *Climate Dynamics*, *61*(9–10), 4259–4281. https://doi.org/10.1007/S00382-023-06798-4/

Sadai, S., Condron, A., DeConto, R., & Pollard, D. (2020). Future climate response to Antarctic Ice Sheet melt caused by anthropogenic warming. *Science Advances*, *6*(39). https://doi.org/10.1126/SCIADV.AAZ1169/SUPPL_FILE/AAZ1169_SM.PDF

Scambos, T., Fricker, H. A., Liu, C. C., Bohlander, J., Fastook, J., Sargent, A., Massom, R., & Wu, A. M. (2009). Ice shelf disintegration by plate bending and hydro-fracture: Satellite observations and model results of the 2008 Wilkins ice shelf break-ups. *Earth and Planetary Science Letters*, *280*(1–4), 51–60. https://doi.org/10.1016/J.EPSL.2008.12.027

Schneider, D. P., Deser, C., & Okumura, Y. (2012). An assessment and interpretation of the observed warming of West Antarctica in the austral spring. *SpringerDP Schneider, C Deser, Y OkumuraClimate Dynamics, 2012•Springer*, *38*(1–2), 323–347. https://doi.org/10.1007/S00382-010-0985-X

Siegert, M., Atkinson, A., Banwell, A., Brandon, M., Convey, P., Davies, B., Downie, R., Edwards, T., Hubbard, B., Marshall, G., Rogelj, J., Rumble, J., Stroeve, J., & Vaughan, D. (2019). The Antarctic Peninsula under a 1.5°C global warming scenario. *Frontiers in Environmental Science*, *7*(JUN), 465700. https://doi.org/10.3389/FENVS.2019.00102

Tetzner, D., Thomas, E., & Allen, C. (2019). A Validation of ERA5 Reanalysis Data in the Southern Antarctic Peninsula—Ellsworth Land Region, and Its Implications for Ice Core Studies. *Geosciences 2019, Vol. 9, Page 289*, *9*(7), 289. https://doi.org/10.3390/GEOSCIENCES9070289

Tewari, K., Mishra, S. K., Salunke, P., & Dewan, A. (2022). Future projections of temperature and precipitation for Antarctica. *Environmental Research Letters*, *17*(1), 014029. https://doi.org/10.1088/1748-9326/AC43E2

Thompson, D. W. J., & Solomon, S. (2002). Interpretation of recent Southern Hemisphere climate change. *Science.orgDWJ Thompson, S SolomonScience, 2002•science.Org*, *296*(5569), 895–899. https://doi.org/10.1126/SCIENCE.1069270

Tseng, K. C., Maloney, E., & Barnes, E. (2019). The Consistency of MJO Teleconnection Patterns: An Explanation Using Linear Rossby Wave Theory. *Journal of Climate*, *32*(2), 531–548. https://doi.org/10.1175/JCLI-D-18-0211.1

Turner, J., Barrand, N. E., Bracegirdle, T. J., Convey, P., Hodgson, D. A., Jarvis, M., Jenkins, A., Marshall, G., Meredith, M. P., Roscoe, H., Shanklin, J., French, J., Goosse, H., Guglielmin, M., Gutt, J., Jacobs, S., Kennicutt, M. C., Masson-Delmotte, V., Mayewski, P., … Klepikov, A. (2014). Antarctic climate change and the environment: an update. *Polar Record*, *50*(3), 237–259. https://doi.org/10.1017/S0032247413000296

Turner, J., Colwell, S. R., Marshall, G. J., Lachlan-Cope, T. A., Carleton, A. M., Jones, P. D., Lagun, V., Reid, P. A., & Iagovkina, S. (2005). Antarctic climate change during the last 50 years. *International Journal of Climatology*, *25*(3), 279–294. https://doi.org/10.1002/JOC.1130

Turner, J., Lu, H., White, I., King, J. C., Phillips, T., Hosking, J. S., Bracegirdle, T. J., Marshall, G. J., Mulvaney, R., & Deb, P. (2016). Absence of 21st century warming on Antarctic Peninsula consistent with natural variability. *Nature*, *535*(7612), 411–415. https://doi.org/10.1038/NATURE18645

Turner, J., Maksym, T., Phillips, T., Marshall, G. J., & Meredith, M. P. (2013). The impact of changes in sea ice advance on the large winter warming on the western Antarctic Peninsula. *International Journal of Climatology*, *33*(4), 852–861. https://doi.org/10.1002/JOC.3474

Turner, J., Marshall, G. J., Clem, K., Colwell, S., Phillips, T., & Lu, H. (2020). Antarctic temperature variability and change from station data. *International Journal of Climatology*, *40*(6), 2986–3007. https://doi.org/10.1002/joc.6378

Turton, J. V., Kirchgaessner, A., Ross, A. N., King, J. C., & Munneke, P. K. (2020). The influence of föhn winds on annual and seasonal surface melt on the Larsen C Ice Shelf, Antarctica. *Cryosphere*, *14*(11), 4165–4180. https://doi.org/10.5194/TC-14-4165-2020

Wang, J., Luo, H., Yu, L., Li, X., Holland, P. R., & Yang, Q. (2023). The Impacts of Combined SAM and ENSO on Seasonal Antarctic Sea Ice Changes. *Journal of Climate*, *36*(11), 3553–3569. https://doi.org/10.1175/JCLI-D-22-0679.1

Wang, S., Ding, M., Liu, G., Wei, T., Zhang, W., Chen, W., Dou, T., & Xiao, C. (2022a). On the drivers of temperature extremes on the Antarctic Peninsula during austral summer. *Climate Dynamics*, *59*(7–8), 2275–2291. https://doi.org/10.1007/S00382-022-06209-0/

Wang, S., Ding, M., Liu, G., & Chen, W. (2022b). Processes and Mechanisms of Persistent Extreme Rainfall Events in the Antarctic Peninsula during Austral Summer. *Journal of Climate*, *35*(12), 3643–3657. https://doi.org/10.1175/JCLI-D-21-0834.1

Wille, J. D., Favier, V., Jourdain, N. C., Kittel, C., Turton, J. V., Agosta, C., Gorodetskaya, I. v., Picard, G., Codron, F., Santos, C. L. dos, Amory, C., Fettweis, X., Blanchet, J., Jomelli, V., & Berchet, A. (2022). Intense atmospheric rivers can weaken ice shelf stability at the Antarctic Peninsula. *Communications Earth and Environment*, *3*(1), 1–14. https://doi.org/10.1038/S43247-022-00422-9

Xie, A., Wang, S., Xiao, C., Kang, S., Gong, J., Ding, M., Li, C., Dou, T., Ren, J., & Qin, D. (2018). *Arctic, Antarctic, and Alpine Research Can Temperature Extremes in East Antarctica be Replicated from ERA Interim Reanalysis?* https://doi.org/10.1657/AAAR0015-048

Xu, M., Yu, L., Liang, K., Vihma, T., Bozkurt, D., Hu, X., & Yang, Q. (2021). Dominant role of vertical air flows in the unprecedented warming on the Antarctic Peninsula in February 2020. *Communications Earth and Environment*, *2*(1), 1–9. https://doi.org/10.1038/S43247-021-00203

Zhang, L., Delworth, T. L., Yang, X., Zeng, F., Lu, F., Morioka, Y., & Bushuk, M. (2022). The relative role of the subsurface Southern Ocean in driving negative Antarctic Sea ice extent anomalies in 2016–2021. *Communications Earth & Environment 2022 3:1*, *3*(1), 302–. https://doi.org/10.1038/s43247-022-00624-1

Zhu, J., Xie, A., Qin, X., Wang, Y., Xu, B., & Wang, Y. (2021). An Assessment of ERA5 Reanalysis for Antarctic Near Surface Air Temperature. *Atmosphere 2021, Vol. 12, Page 217*, *12*(2), 217. https://doi.org/10.3390/ATMOS12020217

Zou, X., Rowe, P. M., Gorodetskaya, I., Bromwich, D. H., Lazzara, M. A., Cordero, R. R., Zhang, Z., Kawzenuk, B., Cordeira, J. M., Wille, J. D., Ralph, F. M., & Bai, L. S. (2023). Strong Warming Over the Antarctic Peninsula During Combined Atmospheric River and Foehn Events: Contribution of Shortwave Radiation and Turbulence. *Journal of Geophysical Research: Atmospheres*, *128*(16), e2022JD038138. https://doi.org/10.1029/2022JD038138

Baba, K. and Renwick, J.: Aspects of intraseasonal variability of Antarctic sea ice in austral winter related to ENSO and SAM events, Journal of Glaciology, 63, 838–846, https://doi.org/10.1017/JOG.2017.49, 2017.

Bae, H. J., Kim, S. J., Kim, B. M., and Kwon, H.: Causes of the Extreme Hot Event on February 9, 2020, in Seymour Island, Antarctic Peninsula, Front Environ Sci, 10, 865775, https://doi.org/10.3389/FENVS.2022.865775/TEXT, 2022.

Balay, F. and Renom, M.: Characterization of extreme warm temperature events (1998–2016) on King George Island, Antarctic Peninsula, Antarct Sci, 38, 146–160, https://doi.org/10.1017/S0954102025100552, 2026.

Baldwin, J. W., Dessy, J. B., Vecchi, G. A., and Oppenheimer, M.: Temporally Compound Heat Wave Events and Global Warming: An Emerging Hazard, Earths Future, 7, 411–427, https://doi.org/10.1029/2018EF000989, 2019.

Banwell, A. F., Tri Datta, R., Dell, R. L., Moussavi, M., Brucker, L., Picard, G., Shuman, C. A., and Stevens, L. A.: The 32-year record-high surface melt in 2019/2020 on the northern George VI Ice Shelf, Antarctic Peninsula, Cryosphere, 15, 909–925, https://doi.org/10.5194/TC-15-909-2021, 2021.

Barrett, B. S., Lafleur, D. M., and Henderson, G. R.: Madden–Julian Oscillation Modulation of Antarctic Sea Ice, Glacies 2025, Vol. 2, 2, 16, https://doi.org/10.3390/GLACIES2040016, 2025.

Bevan, S., Luckman, A., Hendon, H., and Wang, G.: The 2020 Larsen C Ice Shelf surface melt is a 40-year record high, Cryosphere, 14, 3551–3564, https://doi.org/10.5194/TC-14-3551-2020, 2020.

Blanchard-Wrigglesworth, E., Cox, T., Espinosa, Z. I., and Donohoe, A.: The Largest Ever Recorded Heatwave—Characteristics and Attribution of the Antarctic Heatwave of March 2022, Geophys Res Lett, 50, e2023GL104910, https://doi.org/10.1029/2023GL104910, 2023.

Boschat, G., Purich, A., Rudeva, I., Arblaster, J., Boschat, G., Purich, A., Rudeva, I., and Arblaster, J.: Impact of Zonal and Meridional Atmospheric Flow on Surface Climate and Extremes in the Southern Hemisphere, J Clim, 36, 5041–5061, https://doi.org/10.1175/JCLI-D-22-0251.1, 2023.

Bozkurt, D., Bromwich, D. H., Carrasco, J., Hines, K. M., Maureira, J. C., Rondanelli, R., Bozkurt, C. :, Bromwich, D. H., Carrasco, J., Hines, K. M., Maureira, J. C., and Rondanelli, R.: Recent Near-surface Temperature Trends in the Antarctic Peninsula from Observed, Reanalysis and Regional Climate Model Data, Advances in Atmospheric Sciences 2020 37:5, 37, 477–493, https://doi.org/10.1007/s00376-020-9183-x, 2020.

Bozkurt, D., Marín, J. C., Verdugo, C., and Barrett, B. S.: Atmospheric blocking and temperatures in the Antarctic Peninsula, Science of The Total Environment, 931, 172852, https://doi.org/10.1016/J.SCITOTENV.2024.172852, 2024.

Bracegirdle, T. J., Caton Harrison, T., Holmes, C. R., Lu, H., Martineau, P., and Phillips, T.: Antarctic extreme seasons under 20th and 21st century climate change, npj Climate and Atmospheric Science 2024 7:1, 7, 276-, https://doi.org/10.1038/s41612-024-00822-y, 2024.

Cape, M. R., Vernet, M., Skvarca, P., Marinsek, S., Scambos, T., and Domack, E.: Foehn winds link climate-driven warming to ice shelf evolution in Antarctica, Journal of Geophysical Research: Atmospheres, 120, 11,037-11,057, https://doi.org/10.1002/2015JD023465, 2015.

Carrasco, J. F., Bozkurt, D., and Cordero, R. R.: A review of the observed air temperature in the Antarctic Peninsula. Did the warming trend come back after the early 21st hiatus?, Polar Sci, 28, 100653, https://doi.org/10.1016/j.polar.2021.100653, 2021.

Clem, K. R. and Fogt, R. L.: Varying roles of ENSO and SAM on the Antarctic Peninsula climate in austral spring, Journal of Geophysical Research Atmospheres, 118, 11,481-11,492, https://doi.org/10.1002/JGRD.50860, 2013.

Clem, K. R., Bozkurt, D., Kennett, D., King, J. C., and Turner, J.: Central tropical Pacific convection drives extreme high temperatures and surface melt on the Larsen C Ice Shelf, Antarctic Peninsula, Nature Communications 2022 13:1, 13, 3906-, https://doi.org/10.1038/s41467-022-31119-4, 2022.

Coggins, J. H. J., McDonald, A. J., and Jolly, B.: Synoptic climatology of the Ross Ice Shelf and Ross Sea region of Antarctica: K-means clustering and validation, International Journal of Climatology, 34, 2330–2348, https://doi.org/10.1002/JOC.3842, 2014.

Deb, P., Bromwich, D., Orr, A., Sen, A., and Clem, K. R.: Recent increase in surface melting of West Antarctic ice shelves linked to Interdecadal Pacific Oscillation, Communications Earth & Environment 2025 6:1, 6, 99-, https://doi.org/10.1038/s43247-025-02077-8, 2025.

DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, Nature 2016 531:7596, 531, 591–597, https://doi.org/10.1038/nature17145, 2016.

Ding, Q., Steig, E. J., Battisti, D. S., and Küttel, M.: Winter warming in West Antarctica caused by central tropical Pacific warming, Nature Geoscience 2011 4:6, 4, 398–403, https://doi.org/10.1038/ngeo1129, 2011.

Ding, Q., Steig, E. J., Ding, Q., and Steig, E. J.: Temperature Change on the Antarctic Peninsula Linked to the Tropical Pacific, J Clim, 26, 7570–7585, https://doi.org/10.1175/JCLI-D-12-00729.1, 2013.

Dunmire, D., Wever, N., Banwell, A. F., and Lenaerts, J. T. M.: Antarctic-wide ice-shelf firn emulation reveals robust future firn air depletion signal for the Antarctic Peninsula, Communications Earth & Environment 2024 5:1, 5, 100-, https://doi.org/10.1038/s43247-024-01255-4, 2024.

Elvidge, A. D., Renfrew, I. A., King, J. C., Orr, A., Lachlan-Cope, T. A., Weeks, M., and Gray, S. L.: Foehn jets over the Larsen C Ice Shelf, Antarctica, Quarterly Journal of the Royal Meteorological Society, 141, 698–713, https://doi.org/10.1002/QJ.2382,2015.

England, M. R., Polvani, L. M., Smith, K. L., Landrum, L., and Holland, M. M.: Robust response of the Amundsen Sea Low to stratospheric ozone depletion, Geophys Res Lett, 43, 8207–8213, https://doi.org/10.1002/2016GL070055, 2016.

Feron, S., Cordero, R. R., Damiani, A., Malhotra, A., Seckmeyer, G., and Llanillo, P.: Warming events projected to become more frequent and last longer across Antarctica, Scientific Reports 2021 11:1, 11, 19564-, https://doi.org/10.1038/s41598-021-98619-z, 2021.

Fogt, R. L. and Marshall, G. J.: The Southern Annular Mode: Variability, trends, and climate impacts across the Southern Hemisphere, Wiley Interdiscip Rev Clim Change, 11, e652, https://doi.org/10.1002/WCC.652, 2020.

Fogt, R. L., Bromwich, D. H., and Hines, K. M.: Understanding the SAM influence on the South Pacific ENSO teleconnection, Climate Dynamics 2010 36:7, 36, 1555–1576, https://doi.org/10.1007/S00382-010-0905-0, 2010.

Fogt, R. L., Jones, J. M., Renwick, J., Fogt, R. L., Jones, J. M., and Renwick, J.: Seasonal Zonal Asymmetries in the Southern Annular Mode and Their Impact on Regional Temperature Anomalies, J Clim, 25, 6253–6270, https://doi.org/10.1175/JCLI-D-11-00474.1, 2012.

Gonzalez, S. and Fortuny, D.: How robust are the temperature trends on the Antarctic Peninsula?, Antarct Sci, 30, 322–328, https://doi.org/10.1017/S0954102018000251, 2018.

Gonzalez, S., Vasallo, F., Recio-Blitz, C., Guijarro, J. A., Riesco, J., Gonzalez, S., Vasallo, F., Recio-Blitz, C., Guijarro, J. A., and Riesco, J.: Atmospheric Patterns over the Antarctic Peninsula, J Clim, 31, 3597–3608, https://doi.org/10.1175/JCLI-D-17-0598.1, 2018.

González-Herrero, S., Barriopedro, D., Trigo, R. M., López-Bustins, J. A., and Oliva, M.: Climate warming amplified the 2020 record-breaking heatwave in the Antarctic Peninsula, Communications Earth & Environment 2022 3:1, 3, 122-, https://doi.org/10.1038/s43247-022-00450-5, 2022.

Gonzalez-Herrero, S., Vasallo, F., Bech, J., Gorodetskaya, I., Elvira, B., and Justel, A.: Extreme precipitation records in Antarctica, Int J Climatol, 1, 0, https://doi.org/10.1002/joc.8020, 2023.

Gorodetskaya, I. v., Durán-Alarcón, C., González-Herrero, S., Clem, K. R., Zou, X., Rowe, P., Rodriguez Imazio, P., Campos, D., Leroy-Dos Santos, C., Dutrievoz, N., Wille, J. D., Chyhareva, A., Favier, V., Blanchet, J., Pohl, B., Cordero, R. R., Park, S. J., Colwell, S., Lazzara, M. A., Carrasco, J., Gulisano, A. M., Krakovska, S., Ralph, F. M., Dethinne, T., and Picard, G.: Record-high Antarctic Peninsula temperatures and surface melt in February 2022: a compound event with an intense atmospheric river, npj Climate and Atmospheric Science 2023 6:1, 6, 202-, https://doi.org/10.1038/s41612-023-00529-6, 2023.

Goyal, R., Jucker, M., sen Gupta, A., Hendon, H. H., and England, M. H.: Zonal wave 3 pattern in the Southern Hemisphere generated by tropical convection, Nature Geoscience 2021 14:10, 14, 732–738, https://doi.org/10.1038/s41561-021-00811-3, 2021.

Guarino, M.-V., Ridley, J. K., Colwell, S., Farneti, R., Giuliani, G., Hindley, N., King, J., Kucharski, F., Polichtchouk, I., Tompkins, A. M., Vignon, É., Wright, C., Guarino, M.-V., Ridley, J. K., Colwell, S., Farneti, R., Giuliani, G., Hindley, N., King, J., Kucharski, F., Polichtchouk, I., Tompkins, A. M., Vignon, É., and Wright, C.: A Long-Term Shift in Flow Regimes over the Antarctic Peninsula, J Clim, 39, 749–767, https://doi.org/10.1175/jcli-d-25-0330.1, 2026.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., de Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I.,

Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, Quarterly Journal of the Royal Meteorological Society, 146, 1999–2049, https://doi.org/10.1002/QJ.3803, 2020.

Hillebrand, F. L., Bremer, U. F., Arigony-Neto, J., da Rosa, C. N., Mendes, C. W., Costi, J., de Freitas, M. W. D., and Schardong, F.: Comparison between Atmospheric Reanalysis Models ERA5 and ERA-Interim at the North Antarctic Peninsula Region, Ann Am Assoc Geogr, 111, 1147–1159, https://doi.org/10.1080/24694452.2020.1807308, 2021.

Hosking, J. S., Orr, A., Marshall, G. J., Turner, J., and Phillips, T.: The Influence of the Amundsen–Bellingshausen Seas Low on the Climate of West Antarctica and Its Representation in Coupled Climate Model Simulations, J Clim, 26, 6633–6648, https://doi.org/10.1175/JCLI-D-12-00813.1, 2013.

Jones, M. E., Bromwich, D. H., Nicolas, J. P., Carrasco, J., Plavcová, E., Zou, X., Wang, S.-H., Jones, M. E., Bromwich, D. H., Nicolas, J. P., Carrasco, J., Plavcová, E., Zou, X., and Wang, S.-H.: Sixty Years of Widespread Warming in the Southern Middle and High Latitudes (1957–2016), J Clim, 32, 6875–6898, https://doi.org/10.1175/JCLI-D-18-0565.1, 2019.

Kirchgaessner, A., King, J. C., and Anderson, P. S.: The Impact of Föhn Conditions Across the Antarctic Peninsula on Local Meteorology Based on AWS Measurements, Journal of Geophysical Research: Atmospheres, 126, e2020JD033748, https://doi.org/10.1029/2020JD033748,2021.

Laffin, M. K., Zender, C. S., Singh, S., van Wessem, J. M., Smeets, C. J. P. P., and Reijmer, C. H.: Climatology and Evolution of the Antarctic Peninsula Föhn Wind-Induced Melt Regime From 1979–2018, Journal of Geophysical Research: Atmospheres, 126, e2020JD033682, https://doi.org/10.1029/2020JD033682, 2021.

Lee, H. J. and Seo, K. H.: Impact of the Madden-Julian oscillation on Antarctic sea ice and its dynamical mechanism, Scientific Reports 2019 9:1, 9, 10761-, https://doi.org/10.1038/s41598-019-47150-3, 2019.

Li, X., Holland, D. M., Gerber, E. P., and Yoo, C.: Impacts of the north and tropical Atlantic Ocean on the Antarctic Peninsula and sea ice, Nature, 505, 538–542, https://doi.org/10.1038/NATURE12945, 2014.

Li, X., Gerber, E. P., Holland, D. M., Yoo, C., Li, X., Gerber, E. P., Holland, D. M., and Yoo, C.: A Rossby Wave Bridge from the Tropical Atlantic to West Antarctica, J Clim, 28, 2256–2273, https://doi.org/10.1175/JCLI-D-14-00450.1, 2015.

Lim, E. P., Hendon, H. H., and Thompson, D. W. J.: Seasonal Evolution of Stratosphere-Troposphere Coupling in the Southern Hemisphere and Implications for the Predictability of Surface Climate, Journal of Geophysical Research: Atmospheres, 123, 12,002-12,016, https://doi.org/10.1029/2018JD029321, 2018.

Lloyd, S. P.: Least Squares Quantization in PCM, IEEE Trans Inf Theory, 28, 129–137, https://doi.org/10.1109/TIT.1982.1056489, 1982.

Marshall, G. J.: Trends in the Southern Annular Mode from Observations and Reanalyses, J Clim, 2003.

Marshall, G. J.: Half-century seasonal relationships between the Southern Annular Mode and Antarctic temperatures, International Journal of Climatology, 27, 373–383, https://doi.org/10.1002/JOC.1407, 2007.

Marshall, G. J., Orr, A., Lipzig, N. P. M. van, King, J. C., Marshall, G. J., Orr, A., Lipzig, N. P. M. van, and King, J. C.: The Impact of a Changing Southern Hemisphere Annular Mode on Antarctic Peninsula Summer Temperatures, J Clim, 19, 5388–5404, https://doi.org/10.1175/JCLI3844.1, 2006.

Marshall, G. J., Fogt, R. L., Turner, J., and Clem, K. R.: Can current reanalyses accurately portray changes in Southern Annular Mode structure prior to 1979?, Climate Dynamics 2022 59:11, 59, 3717–3740, https://doi.org/10.1007/s00382-022-06292-3, 2022.

Montaño Bello, D., Villegas Bolaños, N. L., Montaño Bello, D., and Villegas Bolaños, N. L.: Influence of ENSO and SAM on the occurrence of absolute extremes and behavior of Ta, SST, and SIC anomalies in the western Antarctic Peninsula, Atmósfera, 39, 437–450, https://doi.org/10.20937/ATM.53423, 2025.

Nielsen, E. B., Katurji, M., Zawar-Reza, P., and Cullen, N.: Air Temperature Trends and Extreme Warming Events Across Regions of Antarctica for the Period 2003–2021, Journal of Geophysical Research: Atmospheres, 130, e2024JD043042, https://doi.org/10.1029/2024JD043042, 2025.

Oliva, M., Navarro, F., Hrbáček, F., Hernández, A., Nývlt, D., Pereira, P., Ruiz-Fernández, J., and Trigo, R.: Recent regional climate cooling on the Antarctic Peninsula and associated impacts on the cryosphere, Science of The Total Environment, 580, 210–223, https://doi.org/10.1016/J.SCITOTENV.2016.12.030, 2017.

Orr, A., Marshall, G. J., Hunt, J. C. R., Sommeria, J., Wang, C. G., van Lipzig, N. P. M., Cresswell, D., and King, J. C.: Characteristics of Summer Airflow over the Antarctic Peninsula in Response to Recent

Strengthening of Westerly Circumpolar Winds, J Atmos Sci, 65, 1396–1413, https://doi.org/10.1175/2007JAS2498.1, 2008.

Pezza, A. B., Rashid, H. A., Simmonds, I., Pezza, A. B., Rashid, H. A., Simmonds, I., Pezza, Bernardes, A., Rashid, A., H., Simmonds, and Ian: Climate links and recent extremes in antarctic sea ice, high-latitude cyclones, Southern Annular Mode and ENSO, ClDy, 38, 57–73, https://doi.org/10.1007/S00382-011-1044-Y, 2012.

Raphael, M. N., Marshall, G. J., Turner, J., Fogt, R. L., Schneider, D., Dixon, D. A., Hosking, J. S., Jones, J. M., and Hobbs, W. R.: The Amundsen Sea Low: Variability, Change, and Impact on Antarctic Climate, Bull Am Meteorol Soc, 97, 111–121, https://doi.org/10.1175/BAMS-D-14-00018.1, 2016.

Renwick, J. A. and Revell, M. J.: Blocking over the South Pacific and Rossby Wave Propagation, Mon Weather Rev, 1999.

Rondanelli, R., Hatchett, B., Rutllant, J., Bozkurt, D., and Garreaud, R.: Strongest MJO on Record Triggers Extreme Atacama Rainfall and Warmth in Antarctica, Geophys Res Lett, 46, 3482–3491, https://doi.org/10.1029/2018GL081475, 2019.

Röthlisberger, M. and Papritz, L.: Quantifying the physical processes leading to atmospheric hot extremes at a global scale, Nature Geoscience 2023 16:3, 16, 210–216, https://doi.org/10.1038/s41561-023-01126-1, 2023.

Rousseeuw, P. J.: Silhouettes: A graphical aid to the interpretation and validation of cluster analysis, J Comput Appl Math, 20, 53–65, https://doi.org/10.1016/0377-0427(87)90125-7, 1987.

Ruosteenoja, K. and Jylhä, K.: Average and extreme heatwaves in Europe at 0.5–2.0 °C global warming levels in CMIP6 model simulations, Climate Dynamics 2023 61:9, 61, 4259–4281, https://doi.org/10.1007/S00382-023-06798-4, 2023.

Sadai, S., Condron, A., DeConto, R., and Pollard, D.: Future climate response to Antarctic Ice Sheet melt caused by anthropogenic warming, Sci Adv, 6, https://doi.org/10.1126/SCIADV.AAZ1169, 2020.

Scambos, T., Fricker, H. A., Liu, C. C., Bohlander, J., Fastook, J., Sargent, A., Massom, R., and Wu, A. M.: Ice shelf disintegration by plate bending and hydro-fracture: Satellite observations and model results

of the 2008 Wilkins ice shelf break-ups, Earth Planet Sci Lett, 280, 51–60, https://doi.org/10.1016/J.EPSL.2008.12.027, 2009.

Schneider, D. P., Deser, C., and Okumura, Y.: An assessment and interpretation of the observed warming of West Antarctica in the austral spring, SpringerDP Schneider, C Deser, Y OkumuraClimate Dynamics, 2012•Springer, 38, 323–347, https://doi.org/10.1007/S00382-010-0985-X, 2012.

Siegert, M., Atkinson, A., Banwell, A., Brandon, M., Convey, P., Davies, B., Downie, R., Edwards, T., Hubbard, B., Marshall, G., Rogelj, J., Rumble, J., Stroeve, J., and Vaughan, D.: The Antarctic Peninsula under a 1.5°C global warming scenario, Front Environ Sci, 7, 465700, https://doi.org/10.3389/FENVS.2019.00102, 2019.

Tetzner, D., Thomas, E., and Allen, C.: A Validation of ERA5 Reanalysis Data in the Southern Antarctic Peninsula—Ellsworth Land Region, and Its Implications for Ice Core Studies, Geosciences 2019, Vol. 9, Page 289, 9, 289, https://doi.org/10.3390/GEOSCIENCES9070289, 2019.

Tewari, K., Mishra, S. K., Salunke, P., and Dewan, A.: Future projections of temperature and precipitation for Antarctica, Environmental Research Letters, 17, 014029, https://doi.org/10.1088/1748-9326/AC43E2, 2022.

Thompson, D. W. J. and Solomon, S.: Interpretation of recent Southern Hemisphere climate change, science.orgDWJ Thompson, S SolomonScience, 2002•science.org, 296, 895–899, https://doi.org/10.1126/SCIENCE.1069270, 2002.

Tseng, K.-C., Maloney, E., Barnes, E., Tseng, K.-C., Maloney, E., and Barnes, E.: The Consistency of MJO Teleconnection Patterns: An Explanation Using Linear Rossby Wave Theory, J Clim, 32, 531–548, https://doi.org/10.1175/JCLI-D-18-0211.1, 2019.

Turner, J., Colwell, S. R., Marshall, G. J., Lachlan-Cope, T. A., Carleton, A. M., Jones, P. D., Lagun, V., Reid, P. A., and Iagovkina, S.: Antarctic climate change during the last 50 years, International Journal of Climatology, 25, 279–294, https://doi.org/10.1002/JOC.1130, 2005.

Turner, J., Maksym, T., Phillips, T., Marshall, G. J., and Meredith, M. P.: The impact of changes in sea ice advance on the large winter warming on the western Antarctic Peninsula, International Journal of Climatology, 33, 852–861, https://doi.org/10.1002/JOC.3474, 2013.

Turner, J., Lu, H., White, I., King, J. C., Phillips, T., Hosking, J. S., Bracegirdle, T. J., Marshall, G. J., Mulvaney, R., and Deb, P.: Absence of 21st century warming on Antarctic Peninsula consistent with natural variability, Nature 2016 535:7612, 535, 411–415, https://doi.org/10.1038/nature18645, 2016.

Turner, J., Marshall, G. J., Clem, K., Colwell, S., Phillips, T., and Lu, H.: Antarctic temperature variability and change from station data, International Journal of Climatology, 40, 2986–3007, https://doi.org/10.1002/JOC.6378, 2020.

Turton, J. v., Kirchgaessner, A., Ross, A. N., King, J. C., and Munneke, P. K.: The influence of föhn winds on annual and seasonal surface melt on the Larsen C Ice Shelf, Antarctica, Cryosphere, 14, 4165–4180, https://doi.org/10.5194/TC-14-4165-2020, 2020.

Wang, S., Ding, M., Liu, G., Wei, T., Zhang, W., Chen, W., Dou, T., and Xiao, C.: On the drivers of temperature extremes on the Antarctic Peninsula during austral summer, Clim Dyn, 59, 2275–2291, https://doi.org/10.1007/S00382-022-06209-0, 2022a.

Wang, S., Ding, M., Liu, G., Chen, W., Wang, S., Ding, M., Liu, G., and Chen, W.: Processes and Mechanisms of Persistent Extreme Rainfall Events in the Antarctic Peninsula during Austral Summer, J Clim, 35, 3643–3657, https://doi.org/10.1175/JCLI-D-21-0834.1, 2022b.

Wille, J. D., Favier, V., Jourdain, N. C., Kittel, C., Turton, J. v., Agosta, C., Gorodetskaya, I. v., Picard, G., Codron, F., Santos, C. L. dos, Amory, C., Fettweis, X., Blanchet, J., Jomelli, V., and Berchet, A.: Intense atmospheric rivers can weaken ice shelf stability at the Antarctic Peninsula, Commun Earth Environ, 3, 1–14, https://doi.org/10.1038/S43247-022-00422-9;SUBJMETA=106,125,35,704,823, 2022.

Xie, A., Wang, S., Xiao, C., Kang, S., Gong, J., Ding, M., Li, C., Dou, T., Ren, J., and Qin, D.: Can Temperature Extremes in East Antarctica be Replicated from ERA Interim Reanalysis?, Arct Antarct Alp Res, 48, 603–621, https://doi.org/10.1657/AAAR0015-048, 2016.

Xu, M., Yu, L., Liang, K., Vihma, T., Bozkurt, D., Hu, X., and Yang, Q.: Dominant role of vertical air flows in the unprecedented warming on the Antarctic Peninsula in February 2020, Communications Earth & Environment 2021 2:1, 2, 133-, https://doi.org/10.1038/s43247-021-00203-w, 2021.

Zhang, L., Delworth, T. L., Yang, X., Zeng, F., Lu, F., Morioka, Y., and Bushuk, M.: The relative role of the subsurface Southern Ocean in driving negative Antarctic Sea ice extent anomalies in 2016–2021,

Communications Earth & Environment 2022 3:1, 3, 302-, https://doi.org/10.1038/s43247-022-00624-1, 2022.

Zhu, J., Xie, A., Qin, X., Wang, Y., Xu, B., and Wang, Y.: An Assessment of ERA5 Reanalysis for Antarctic Near-Surface Air Temperature, Atmosphere 2021, Vol. 12, Page 217, 12, 217, https://doi.org/10.3390/ATMOS12020217, 2021.

Zou, X., Rowe, P. M., Gorodetskaya, I., Bromwich, D. H., Lazzara, M. A., Cordero, R. R., Zhang, Z., Kawzenuk, B., Cordeira, J. M., Wille, J. D., Ralph, F. M., and Bai, L. S.: Strong Warming Over the Antarctic Peninsula During Combined Atmospheric River and Foehn Events: Contribution of Shortwave Radiation and Turbulence, Journal of Geophysical Research: Atmospheres, 128, e2022JD038138, https://doi.org/10.1029/2022JD038138, 2023.

Zou, X., Rowe, P. M., Gorodetskaya, I. v., Orr, A., Bromwich, D. H., Lubin, D., Lazzara, M. A., Zhang, Z., Brian, K., Wille, J. D., Cordeira, J. M., Hansen, N., Li, J., Gan, P., and Ralph, F. M.: Föhn-induced melting over Larsen C modulated by atmospheric river shape, direction and landfall location, Nature Communications 2026 17:1, 17, 4788-, https://doi.org/10.1038/s41467-026-71359-2, 2026.