

Decadal predictions of wind, solar and compound power indicators to support the European renewable energy sector

Sara Moreno-Montes¹, Carlos Delgado-Torres¹, Matías Olmo¹, Sushovan Ghosh¹, Verónica Torralba¹, and Albert Soret¹

¹Barcelona Supercomputing Center (BSC), Barcelona, Spain

Correspondence: Sara Moreno-Montes (sara.moreno@bsc.es)

Abstract. Renewable energy production is strongly influenced by climate variability and change, making the energy sector sensitive to fluctuations on decadal timescales. Decadal climate predictions, which aim to forecast climate variability over the next few years, therefore offer potential value for anticipating near-term changes in wind and solar resources and supporting climate-informed energy planning. However, the predictive skill of decadal forecasts for energy-relevant indicators remains poorly quantified, which is crucial to know the potential usability of any forecast product.

This study evaluates the skill of decadal climate predictions over Europe for forecast years 1–3 using a multi-model ensemble from the Coupled Model Intercomparison Project Phase 6 (CMIP6) Decadal Climate Prediction Project (DCPP). We assess three energy-relevant indicators: photovoltaic potential (PVpot), wind capacity factor (WCF), and a compound indicator describing the number of energy drought days (NED), defined as days with inefficient production from both wind and solar resources. The skill is evaluated against the ERA5 reanalysis, and the added value of the model initialization is estimated by comparing the decadal predictions against the non-initialized historical forcing simulations. PVpot exhibits the highest and most spatially homogeneous skill for annual, spring and summer aggregations, closely reflecting the high predictability of surface solar radiation. WCF shows low and spatially heterogeneous skill, consistent with the high intrinsic variability of wind. The compound NED indicator displays strong seasonal dependence: its predictability is largely controlled by solar conditions in high-radiation seasons and by wind in winter and autumn. Model initialization generally provides added value where historical simulations already show some skill, especially for PVpot, while its impact is lower for WCF. This work shows the specific seasons, regions and energy indicators for which decadal predictions can provide actionable climate information to support renewable energy applications.

1 Introduction

The renewable energy sector has gained increasing importance in Europe in recent years, with wind and solar representing a growing fraction of total electricity generation (European Environment Agency, 2022). Continued policy support and decarbonisation targets are expected to further accelerate this expansion in the coming decades (International Energy Agency, 2024). Renewable energy production depends strongly on variations in key climate variables such as near-surface air temperature, solar radiation or surface wind speed (Solomon, 2007), and ambitious deployment targets across Europe further highlight the need

25 for climate-informed planning in the energy sector (e.g., Ely et al., 2013; Gonzalez et al., 2019). While long-term climate projections are essential for strategic planning, and short-term forecasts support daily operations, the intermediate horizon of next years remains comparatively underexploited (Smith et al., 2019). Decadal climate predictions, which bridge the gap between seasonal forecasts and long-term projections, are emerging as a valuable tool for sectors that must adapt to both climate variability and climate change (Meehl et al., 2009; Boer et al., 2016). In the energy sector, such information could support infrastructure development, system resilience, and risk management decisions (~~Challinor et al., 2017; European Commission, 2020~~) (European Commission, 2020). Despite this potential, the explicit integration of decadal climate information into wind and solar energy assessments remains limited, highlighting a gap between advances in decadal prediction science and their application in the renewable energy sector.

Previous studies have evaluated the forecast skill of decadal prediction systems for essential climate variables and extreme indices (~~Association et al., 2023; Delgado-Torres et al., 2022, 2023; World Meteorological Organization, 2024~~) (Delgado-Torres et al., 2022), and have also explored the role of statistical downscaling in improving regional skill (Moreno-Montes et al., 2026). In parallel, decadal forecasts have been applied to sector-specific climate services, including agriculture (Solaraju-Murali et al., 2022; Delgado-Torres et al., 2025), water management (Paxian et al., 2022), marine fisheries (Payne et al., 2022), and hydropower (Tsartsali et al., 2023), as well as multi-sector studies demonstrating the application of decadal predictions (Dunstone et al., 2022; Done et al., 2021). However, research on decadal climate services explicitly focused on the energy sector remains limited. Early assessments have reported regionally dependent skill for decadal predictions of wind-related variables, including near-surface wind speed and wind energy output, with skill often enhanced through downscaling approaches (Haas et al., 2015; Moemken et al., 2016). More recent work has highlighted relatively high skill for surface solar radiation, while wind speed skill remains generally low over Europe (Hutchins et al., 2025). By comparison, climate services for the energy sector have been more extensively explored at sub-seasonal to seasonal timescales (Clark et al., 2017; Torralba et al., 2017b; Bloomfield et al., 2021; Bett et al., 2022; Cionni et al., 2022; Lledó et al., 2022; Soret et al., 2026) and with long-term climate projections (~~Carvalho et al., 2021; Hou et al., 2021; Olmo et al., 2026~~) (Carvalho et al., 2021; Hou et al., 2021).

In this study, we assess the skill of decadal predictions for energy-relevant indicators derived separately for solar and wind energy over forecast years 1–3, and we further explore compound indicators that combine information from both sources. ~~Compound events are particularly relevant because simultaneous deficits in multiple energy resources can exacerbate the risk of energy shortages and stress the energy system (Zscheischler et al., 2018; Otero et al., 2022)~~ This 3-year forecast period is particularly relevant for short- to medium-term energy planning applications, including maintenance scheduling, estimation of expected renewable energy production, and management of seasonal energy mixes. Both annual and seasonal indicators are considered: annual means highlight low-frequency signals relevant for year-ahead planning, while seasonal means allow the identification of season-dependent predictability. Analyzing each energy source independently allows us to characterize source-specific predictability, while the combined indicators capture the co-variability of wind and solar resources and its implications for energy system operations. Compound events are particularly relevant because simultaneous deficits in multiple energy resources can exacerbate the risk of energy shortages and stress the energy system (Zscheischler et al., 2018; Otero et al., 2022). Previous studies have shown that combining ~~multiple renewable energy sources, particularly~~ wind and solar resources can

60 enhance system robustness ~~by exploiting through~~ their temporal and spatial complementarity ~~, and can support planning and~~
~~balancing strategies in highly renewable energy systems~~ (Heide et al., 2010; Monforti et al., 2014; Jerez et al., 2019; Kapica
et al., 2024). Decadal predictions of compound hot-dry extreme events have ~~been recently~~ recently been explored by Aranyossy
et al. (2025). However, to the best of our knowledge, an assessment of the forecast quality of decadal predictions for compound
power indicators remains lacking, limiting the understanding of their utility for energy ~~systems~~ system planning and risk
65 management over multi-year timescales.

This study is organized as follows. Section 2 describes the datasets used to compute the indicators including observation-
based reference datasets, decadal predictions and historical simulations. Section 3 presents the methodology to compute the
solar- and wind-based indicators, and compound indicators. Section 4 presents the results together with the discussion and a
comparison with previous studies. Finally, Section 5 summarizes the main findings, discussing their implications and suggest-
70 ing directions for future research.

2 Data

In this study, five different decadal forecast systems contributing to the Decadal Climate Prediction Project (DCPP; Boer
et al., 2016) of the Coupled Model Intercomparison Project Phase 6 (CMIP6; Eyring et al., 2016), are used. These forecast
systems are the EC-Earth3 (with three different initialization strategies), IPSL-CM6A-LR and MPI-ESM1.2-HR. In addition,
75 to analyze the impact of initialization, the indicators are computed for the three corresponding CMIP6 non-initialized historical
forcing simulation models (7 EC-Earth3 members, 11 IPSL-CM6A-LR members, and 2 MPI-ESM1.2-HR members). The main
characteristics of the DCPD forecast systems and the historical simulation models, including ensemble size, spatial resolutions,
initialization months (for the DCPD-A), and references, are detailed in Table S1 of the Supplementary Material. Additionally,
the ERA5 reanalysis (Hersbach et al., 2020) dataset is used as the observation-based reference. This observational reference
80 has been selected taking into account its temporal and spatial coverage and because it is widely used by the energy community
(Ramon et al., 2024).

Depending on the energy source, different temporal resolutions and variables are used. For solar energy indicators, daily
values of surface downwelling shortwave radiation (RSDS), near-surface air temperature (TAS), and surface wind speed
(SFCWIND) are employed. For wind energy indicators, instantaneous 6-hourly surface wind (6h-SFCWIND) data are used.
85 As sub-daily data are required to compute the wind indicators (Lledó et al., 2022), the number of available models is reduced.

Start dates from 1960 to 2016 are used for decadal predictions, focusing the analysis on forecast years 1–3. Therefore, to
match the period, we use the ERA5 data, as well as the historical forcing simulations concatenated with the scenario ssp245
(O'Neill et al., 2016), during 1961–2019. Historical simulations provide data until 2014, and ssp245 is used for the rest of the
years (2015–2019).

90 The analysis focuses on Europe, defined by the spatial domain spanning longitudes 30°W–12°E and latitudes 35°N–65°N.

3 Methodology

To ensure consistency across datasets, variables from all model simulations are first interpolated provide information at a spatial scale relevant for regional applications, a statistical downscaling approach is applied to model simulations to increase the horizontal resolution, bringing them to the ERA5 grid (0.25°). To calculate wind energy indicators, since decadal prediction systems and historical simulations do not provide wind speeds at hub height, This downscaling procedure consists of a combination of spatial interpolation and calibration using Empirical Quantile Mapping (EQM), described below. This method is adopted, as previous studies have shown that more complex downscaling techniques often provide limited added value in terms of predictive skill compared to simpler approaches (e.g., Manzanas et al. (2018); Gutiérrez et al. (2019); Moreno-Montes et al. (2026)), while minimising the risk of introducing artificial skill. To assess the sensitivity of the results to the spatial resolution used for the analysis, all indicators and skill metrics are additionally computed on the native spatial resolution of the coarsest prediction system considered (IPSL-CM6A-LR: $2.5^\circ \times 1.25^\circ$).

After interpolation and before calibration, wind energy indicators are derived from 6h-SFCWIND data from both the reference dataset and the model simulations are data converted to wind speeds at 100m-100 m (6-hWIND). A representative hub height of 100m-100 m is assumed for modern wind turbines (Wiser et al., 2023). To estimate hub-height hub-height winds from surface winds, a power law is applied (Brower, 2012), using a shear exponent of $\alpha = 0.143$ (Touma, 1977) over land and $\alpha = 0.11$ over water (Hsu et al., 1994), under the assumption of neutral atmospheric stability.

In addition, ERA5 wind speeds are adjusted using the Global Wind Atlas (GWA) to better represent long-term mean conditions at hub height. Figure S1 shows the climatological ratio between ERA5 and GWA mean 6-hWIND over 1961–2019, which is applied as a pointwise correction to the multiplicative pointwise correction factor (GWA/ERA5 data. This ensures a more realistic representation of wind resources) to adjust the mean wind speed while preserving the temporal variability from of the ERA5 data (Lledó et al., 2022).

After interpolation and Once wind-specific corrections, all have been applied, model outputs are adjusted to the observed climatology by removing the model mean and replacing it with the corresponding observed mean. These climatologies are computed over a common reference period (1963–2017), defined by the availability of all forecast years included in this study (i.e. 1–3) across the selected start dates (1960–2016). To ensure consistency with the cross-validation framework, the climatological means are estimated excluding the target year being calibrated. This adjustment reduces the impact of model drift while preserving the temporal variability of the simulations (Boer et al., 2016).

The adjusted decadal predictions and historical simulations are then calibrated against ERA5 using Empirical Quantile Mapping (EQM; EQM (Panofsky and Brier, 1958). EQM maps the quantiles (percentiles) of the modelled data to the observed ones without assuming any underlying distribution and then it corrects the model biases By adjusting empirical quantiles at 1% intervals (from the 1st to the 99th percentile), EQM corrects the distribution of the variable, including the mean, variance, and higher-order moments. Calibration is performed independently at each grid point, per forecast year and member for decadal predictions, and per calendar year and member for historical simulations. In both cases, empirical distributions are estimated using a 30-day moving window centered on the target date is applied to remove short-term fluctuations and to pre-

125 serve seasonality while ensuring a sufficient sample size. A leave-one-year-out cross-validation is ~~used to prevent overfitting~~
~~and applied~~. For each target year, the climatological adjustment and EQM transfer function are estimated using all other years
in the calibration period. This prevents information from the target year from being used in the calibration and avoids artificial skill inflation (Elsner and Schmertmann, 1994). For 6-hWIND, the calibration is performed separately for each time of
the day, since wind distributions differ markedly across the diurnal cycle (~~Kalverla et al., 2019; Weide Luiz and Fiedler, 2022~~)
130 (Kalverla et al., 2019; Weide Luiz and Fiedler, 2022).

The calibrated climate variables are then used to compute the energy indicators. All indicators are calculated separately for
each climatological season and for the whole year. We define three indicators: photovoltaic potential (PVpot) for solar energy,
capacity factor for wind energy (WCF), and the Number of Energy Droughts (NED). The intermediate metric Number of
Effective days (Neff) is first computed for each resource and subsequently combined to derive NED. Indicators are computed
135 for each forecast system and for the ERA5 reanalysis, which we use as a reference to assess predictive performance. Results
for wind-related indicators are analyzed over land and over sea-adjacent regions, as both onshore and offshore wind resources
are of practical interest for current and future wind energy deployment.

Following previous studies (Mavromatakis et al., 2010; Jerez et al., 2015), PVpot is computed as a dimensionless metric
that combines RSDS, TAS and SFCWIND to represent the performance of photovoltaic cells under ambient environment. PV
140 power generation at a specific location is determined by multiplying PVpot with the nominal installed capacity of that loca-
tion. The PVpot is mainly driven by RSDS, with adjustments for TAS, which lowers module efficiency at higher temperatures
(Radziemska, 2003), and SFCWIND, which can partially offset thermal losses through convective cooling. Although the equa-
tion S1 provides as a unitless quantity, in this study they are converted to percentage values to facilitate the visual interpretation.
Even though daily inputs are less precise than hourly due to aggregation, they keep seasonal and annual errors small and avoid
145 the much larger biases seen with monthly data (Müller et al., 2019; Bett and Thornton, 2016).

WCF is defined as the ratio between the energy actually produced and the energy that would be generated if the turbine
operated continuously at its rated power. WCF depends on the turbine's power curve, which varies by turbine type and site
conditions; the standard power curves defined ~~in international~~ by the International Electrotechnical Commission (IEC) 61400-1
are shown in Figure S2. In this study IEC Class I turbine is adopted as a common reference, as a sensitivity analysis presented
150 no notable differences among the skill performance of the different turbines (not shown). Given the high temporal resolution
of turbine power curves and their strong nonlinearity, the use of 6-hourly wind data reduces aggregation bias. Although daily
means are often adequate (MacLeod et al., 2018), sub-daily data better preserves variability and yields more accurate WCF
estimates (Lledó et al., 2022). WCF is derived using the R-based CSIndicators package (Pérez-Zanón et al., 2023).

PVpot and WCF are defined differently, so their absolute values should not be compared across energy sources. To enable a
155 meaningful cross-resource comparison, Neff is defined as the number of days meeting efficiency-related thresholds, providing
a common performance indicator with direct operational relevance.

For solar energy production, Neff efficiency thresholds are defined using a minimum RSDS and a maximum cell temperature
(Tcell: Chenni et al. (2007)). The RSDS threshold is based on the concept of peak sun hours (PSH), defined as the equivalent
number of hours per day with an irradiance of 1000 W m^{-2} (Duffie and Beckman, 1980). A commonly used lower limit for

160 meaningful PV generation is 5 PSH per day, which corresponds to a daily mean RSDS of 208 W m^{-2} (Ghosh et al., 2022). Since higher TAS increases Tcell and reduces PVpot, an upper operating threshold of Tcell = 45°C is applied (see equation S2), at which the PV module works at 90% of its rated-efficiency and above the threshold it progressively declines PV generation (Ghosh et al., 2024).

For wind energy, Neff is defined using both operational and performance criteria. Although IEC Class I turbines operate
165 between cut-in (2 ms^{-1}) and cut-out (25 ms^{-1}) wind speeds, winds just above cut-in typically yield little energy production. Therefore, effectiveness is defined using a minimum threshold of $\text{WCF} \geq 25\%$, which excludes periods of low-efficiency production and can be associated with economically relevant wind power generation (Boccard, 2009; Kealy et al., 2015). This threshold is translated into an equivalent minimum wind speed using the turbine power curve, and effective time steps are counted when wind speeds exceed this value while remaining below cut-out. To ensure consistency with solar indicators, 6-
170 hourly wind data are aggregated to daily values, defining a day as effective when at least 3 of 4 time steps meet the threshold (Figure S3).

Based on the effective number of days in which any energy source can operate efficiently, we define the combined indicator NED, which corresponds to the number of days when neither source produces efficiently. Besides, the Neff for each energy source (solar-Neff and wind-Neff) ~~and for both (both-Neff)~~ are calculated.

175 Once the corresponding indicators are calculated for each forecast year, the forecast years 1-3 are averaged in order to obtain a mean value over the entire forecast period for each forecast system. To ensure a consistent comparison with decadal predictions, a rolling mean with a window length equal to the number of forecast years (3) is applied to the historical simulations and ERA5 time series. Subsequently, multi-model ensembles are constructed for decadal predictions (DCPP) and historical simulations (HIST) following the multi-model mean approach (Delgado-Torres et al., 2022).

180 The performance of the DCPP is evaluated against the ERA5 reanalysis using the Anomaly Correlation Coefficient (ACC; Wilks, 2011). ACC ranges from -1 (perfect inverse correlation) to 1 (perfect correlation). A one-sided t-test is applied to determine if ACC values differ significantly from zero (Wilks, 2011). Following Storch and Zwiers (1999), the effective number of degrees of freedom is used to account for autocorrelation in the time series. To assess the impact of initialization, the residual correlation (ResCorr; Smith et al., 2019) between the DCPP and ERA5 is calculated relative to the HIST. Positive values of
185 ResCorr indicate that decadal predictions can capture more observed variability that is not already captured by the historical simulations.

Since the DCPP and HIST ensembles have different numbers of members, we perform an additional sensitivity analysis to assess the impact of ensemble size on the estimated added value of initialization. The analysis is applied to the multi-model mean used throughout the study; therefore, ensemble-size adjustments are performed independently for each model before
190 constructing the multi-model average. For each forecast system, the same number of DCPP and HIST members is selected, corresponding to the minimum ensemble size available in both systems (7 for EC-Earth3, 10 for IPSL-CM6A-LR, and 2 for MPI-ESM1.2-HR). For this analysis, the 7 EC-Earth3 decadal prediction members are taken from the three different initializations for their comparison with the non-initialized EC-Earth3 historical simulations. For each resampling, an equal-member multi-model ensemble is constructed and used to estimate ResCorr. The procedure is repeated 5000 times, and the sensitivity

195 to ensemble size is quantified as the difference in residual correlation:

$$\Delta\text{ResCorr} = \text{ResCorr}_{\text{equal-members}} - \text{ResCorr}_{\text{full-ensemble}}.$$

The median of the resulting $\Delta\text{ResCorr}$ distribution is used as the central estimate at each grid point, while the 2.5th and 97.5th percentiles define a 95% bootstrap interval. Regions where this interval does not include zero are considered robustly sensitive to ensemble size.

200 Finally, after computing the indicators for each grid point, a regionalization is applied to analyze trends and correlations at the regional scale. For each sub-region, indicators are first spatially area-weighted averaged to obtain a single representative time series, from which regional trends, correlations and ResCorr are calculated. This step is motivated by the known contribution of externally forced trends and low-frequency variability to decadal prediction skill in some variables (e.g. van Oldenborgh et al., 2012), and allows us to assess how long-term changes influence the estimated predictability. Trends are estimated
205 using a linear fit, while statistical significance is assessed independently using a modified Mann-Kendall test that accounts for temporal autocorrelation following Hamed and Rao (1998). Europe is divided in seven sub-regions (Figure S4), following the regionalization of Priestley et al. (2024): Iberia, Western Europe, Mediterranean, Central Europe, Eastern Europe, Scandinavia, Great Britain and Ireland.

4 Results and discussion

210 The results are presented in three separate sections for each indicator: PVpot, WCF, and NED. ~~In each section, the ACC between the DCP and ERA5, the ACC between HIST and ERA5, and the ResCorr between the DCP and ERA5 relative to HIST are shown for the annual mean and individual seasons. The annual means provide an integrated measure of performance and reduce high-frequency variability, helping to highlight low-frequency signals relevant at decadal timescales. This information is usually required by energy stakeholders for planning the year ahead. Seasonal means allow the identification of season-dependent predictability which can be relevant for energy deployment. The ACC between the DCP~~ For each indicator, the climatology and multiannual standard deviation are presented to provide context on the mean state and variability of the indicator (Figures S5, S10 and S13). The ACC of the indicator between DCP and ERA5, the ACC between HIST and ERA5, and the ResCorr between the DCP and ERA5 relative to HIST are shown for the annual mean and individual seasons. Equivalent results obtained on the native resolution of the coarsest prediction system are shown in Figures S6, S11 and S15. Additionally, the ACC between DCP and ERA5 and the regional trends of the variables used to calculate the indicators (RSDS, TAS and SFCWIND) and ERA5 are shown
220 in Figures S7 and S9. Finally, the sensitivity of the ResCorr results to differences in ensemble size between DCP and HIST, assessed through the bootstrap resampling procedure described in Section 2, are presented in Figure S5 Figures S8, S12 and S17.

4.1 Photovoltaic potential (PVpot)

The climatology of PVpot (~~S6a~~S5a–e) shows higher values over southern Europe than at higher latitudes, reflecting the merid-
225 ional gradient in solar irradiance, in line with previous studies (e.g. Šúri et al., 2007; Castillo et al., 2016). The associated

~~interannual~~multiannual standard deviation is low across the domain (~~S6f~~S5f–j), indicating that PVpot variability is small and largely controlled by mean climatological conditions.

Figure 1a-e shows the ACC between DCPD and ERA5 of PVpot for the annual mean and each season. The annual mean (Figure 1a) and JJA (Figure 1d) show the highest fractions of significant area, with positive skill over most of the region, except in parts of the UK, southern Europe and Scandinavia. In MAM (Figure 1c), the ACC is not significantly positive in northern Europe and Iberia and the percentage of significant area is lower than in the annual mean and JJA, although the skill patterns are similar. ~~This reduction is mainly due to northern Europe and parts of Iberia, where MAM skill is generally non-significant.~~ On the other hand, DJF (Figure 1b) shows the lowest fraction of significant points, with significant skill only over parts of Iberia and Italy. ACC values during ~~the SON season~~SON (Figure 1e) are significant in regions along the Atlantic coast and parts of Poland and the Baltic countries. Across all seasons, skill tends to be lower in mountainous regions, particularly over the Alps.

~~First row: ACC of DCPD of solar PVpot for the annual mean (a) and seasonal means (b–e) for forecast years 1–3. The percentages of significant grid points are indicated in brackets, and hatched regions denote non-significant correlations. Second row: ResCorr between DCPD and ERA5 relative to the HIST for the annual mean (f) and seasonal means (g–k). Percentages indicate the fractions of significantly positive (red) and significantly negative (blue) values. Positive (negative) ResCorr indicates higher (lower) skill for DCPD compared to HIST. Correlations are computed against ERA5 during the 1961–2019 period, and statistical significance is assessed using a one-sided t-test at the 95% confidence level, accounting for time-series autocorrelation. Similar skill patterns are obtained at the coarser resolution (Figure S6a–e), indicating that the main results are robust to spatial resolution.~~

Although the percentage of significant grid points is slightly lower for PVpot than for its main driver RSDS (Figure ~~S5a–e~~S7a–e), both variables show very similar spatial skill patterns. The reduction in PVpot skill relative to RSDS is more noticeable over southern Europe in the annual mean (Figure ~~S5a–S7a~~ and Figure 1a), over parts of central Europe in MAM (Figure ~~S5c–S7c~~ and Figure 1c), and over Iberia in SON (Figure ~~S5e–S7e~~ and Figure 1e). This loss of skill reflects the fact that PVpot is not only driven by RSDS but is also modulated by TAS and, with a smaller contribution, by SFCWIND. While TAS generally shows high and spatially homogeneous skill across most seasons (except in DJF; Figure ~~S5g–S7g~~), and SFCWIND exhibits weaker and more heterogeneous skill (Figure ~~S5k–S7k–o~~), their combined effect slightly decreases the overall PVpot predictability compared to RSDS alone.

The seasonal contrast in skill suggests a link with large-scale atmospheric conditions. Skill is higher in JJA and MAM, when more persistent radiative conditions typically dominate, and lower in DJF and in mountainous regions, when variability is stronger. However, these relationships are not spatially uniform and should be interpreted cautiously. Besides, the similarity between the annual and JJA skill patterns reflects the dominant contribution of high-radiation seasons to the annual PVpot signal. Since PVpot is strongly controlled by RSDS, and solar irradiance peaks during spring and summer, the annual mean largely integrates the predictability of these seasons. In addition, annual averaging reduces short-term variability and increases signal-to-noise ratios, resulting in higher predictability and thus higher skill values.




Figure 1. First row: ACC between DCP and ERA5 for PVpot for the annual mean (a) and seasonal means (b–e) for forecast years 1–3. Second row: ACC between HIST and ERA5 for the annual mean (f) and seasonal means (g–j). The percentages of significant grid points are indicated in brackets, and hatched regions denote non-significant correlations. Third row: ResCorr between DCP and ERA5 relative to HIST for the annual mean (k) and seasonal means (l–o). Percentages indicate the fractions of significantly positive (red) and significantly negative (blue) values. Positive (negative) ResCorr indicates higher (lower) skill for DCP compared to HIST. Correlations are computed against ERA5 during the 1961–2019 period, and statistical significance is assessed using a one-sided t-test at the 95% confidence level, accounting for time-series autocorrelation.

260 Few studies have assessed decadal predictability for solar-energy indicators. Hutchins et al. (2025) have reported high summer skill for RSDS over southern and central Europe, broadly consistent with our results. They have also identified additional winter predictability using a NAO-based approach, which is not captured when evaluating PVpot from the multi-model ensemble. Focusing on seasonal predictions, Bett et al. (2022) have found higher predictability for RSDS in summer and lower in winter, in line with the seasonal contrast observed here for decadal predictions.

265 The ACC between HIST and ERA5 is shown in Figure 1f-j and the added value of the DCPD relative to the HIST is HIST assessed using the ResCorr (Figure 1f-j), ~~and the ACC between HIST and ERA5 is shown in Figure S7. In 1k-o. HIST reproduces the main spatial patterns identified in DCPD, although with a lower fraction of significant area, particularly in the annual mean (Figure 1f) and JJA (Figure 1i), ResCorr is significantly positive over large parts of the domain, indicating that model initialization enhances PVpot skill relative to historical simulations. In these seasons, HIST skill is already generally~~
270 ~~high (Figure S7a,d), so initialization mainly amplifies existing positive skill. In DJF (Figure 1g), ResCorr exhibits localized negative values, indicating reduced skill in some regions. On the other hand, DCPD skill (Figure 1b) is lower than HIST skill (Figure S7b1g) in areas such as southern Iberia or the Alps. During the transition seasons, . In the annual mean (Figure 1k) and JJA (Figure 1n), ResCorr is significantly positive over large parts of the domain, indicating that model initialization enhances PVpot skill relative to historical simulations. During MAM and SON (Figure 1h,j,m,o), positive ResCorr values~~
275 ~~appear only in are restricted to~~ specific regions, mainly over eastern Europe and parts of the Atlantic coast ~~for in~~ MAM and over northern Atlantic regions ~~for SON, again corresponding to areas where HIST already exhibits positive skill (Figure S7c,e). Overall, initialization mainly enhances PVpot skill where historical simulations already exhibit predictability in SON. Similar conclusions are obtained at the native resolution of the coarsest prediction system (Figure S6f-o), where both the ACC of HIST and the ResCorr show broadly consistent patterns. Figure S8 shows that the median ResCorr differences obtained when~~
280 ~~subsampling the ensembles are generally close to zero and non-significant across Europe. This indicates that the added value of initialization identified in Figure 1k-o is largely insensitive to the unequal ensemble sizes of the prediction systems.~~

Figure 2 summarizes the regional behaviour of PVpot long-term trends across the seven European sub-regions (Figure S2S4), showing ERA5 trends (a), DCPD trends (b), their correlations (c), and their ResCorr ~~respect the relative to~~ HIST (d) for the annual mean and each season. For Iberia, Western Europe, the Mediterranean, Central Europe ~~and~~ Eastern Europe, at ERA5
285 and DCPD generally show positive significant trends in the annual mean ~~and during~~, MAM and JJA, ~~both datasets show positive significant trends, which are accompanied by significantly resulting in significant~~ positive correlations. In DJF and SON, ERA5 trends in these regions are often weak or non-significant, while DCPD trends remain significantly positive, leading to generally ~~low contrast, correlations are mostly weak~~ and non-significant ~~correlations. Some exceptions to this behaviour occur (Iberia in DJF, Western Europe in SON, and Eastern Europe in SON), where better agreement between trends results in significant~~
290 ~~correlations. For in DJF and SON, when observed trends are small or absent. Scandinavia and Great Britain and Ireland exhibit little skill overall, with significant correlations only in a few cases. ResCorr values are generally positive where correlations are significant, particularly in JJA and over Western and Eastern Europe during MAM, trends generally disagree and correlations are low, with a few exceptions (Scandinavia in the annual mean and SON and Great Britain and Ireland in MAM), although significant correlations are limited to the the annual mean in Scandinavia and MAM in Great Britain and Ireland. Overall,~~




Figure 2. PVpot trends derived from ERA5 for years 1961-2019 with a three years rolling mean (a) and from the DCP for the start dates 1960-2016 for over forecast years 1–3 (b), the correlation between ERA5 and DCP (c), and the ResCorr between DCP and ERA5 respect HIST (d) shown for the annual mean and for each season across the seven european sub-regions (Figure S4). Statistically significant trends and correlations at the 95% confidence level are marked with an asterisk.

these results highlight the strong link between trend-agreement and the skill of decadal predictions and in Eastern Europe and Scandinavia during annual mean, indicating that initialization contributes additional skill beyond the common trend signal. Overall, PVpot skill is closely linked to the ability of DCP to reproduce the observed long-term trends.

PVpot trends derived from ERA5 for years 1961-2019 with a three years rolling mean (a) and from the DCP for the start dates 1960-2016 for over forecast years 1–3 (b), the correlation between ERA5 and DCP (c), and the ResCorr between DCP and ERA5 respect HIST (d) shown for the annual mean and for each season across the seven european sub-regions (Figure S4). Statistically significant trends and correlations at the 95% confidence level are marked with an asterisk.

PVpot trends follow similar patterns to those from RSDS (Figure S8a-bS9a-b) for both ERA5 and DCP, showing again that the variable is the main driver of the indicator. TAS trends (Figure S8e-dS9c-d) are significantly positive in all the seasons and regions for both datasets, which are not aligned with the PVpot values.

ResCorr regional patterns (Figure 2d) show that in regions and seasons where correlations are significantly positive (Figure 2e), ResCorr values are generally also positive, although they are only significant in JJA for most regions, in Western and

Eastern Europe during MAM, and in Eastern Europe and Scandinavia for the annual mean. This indicates that not all the skill shown in Figure 2c is explained by the similar trends of ERA5 (Figure 2a) and DCP (Figure 2b) alone. Initialization adds additional skill beyond the externally forced signal in these cases. In DJF, ResCorr values are not significant in any region, indicating that initialization does not add skill beyond what is captured by the HIST. In SON, ResCorr values are mostly positive but small, suggesting that initialization provides a limited additional contribution compared to the forced trend. In summary, when ERA5 and DCP trends agree, part of the skill is linked to the common trend, but initialization can still enhance predictability in all the regions, particularly in JJA. On the other hand, when trends have opposite signs and correlations are non-significantly positive, ResCorr values are also non-significant.

The strong contribution of long-term trends to significant skill has been identified in several studies for decadal predictions (Meehl et al., 2009; Suckling et al., 2017) and historical simulations (Donat et al., 2023). Some studies have applied approaches to reduce model drift or systematic errors (Kharin et al., 2012), and others have shown that the skill of TAS is substantially reduced after removing the forced trend, whereas precipitation skill is less affected (Bellucci et al., 2015; Delgado-Torres et al., 2022). However, removing the trend may also influence forecast reliability and modify the balance between externally forced and internally driven variability (Corti et al., 2012; Smith et al., 2012). Rather than adopting a single approach, we explicitly present both perspectives. The ACC quantifies the total predictive skill, including the contribution from externally forced trends, which are themselves relevant for decision-making in a climate services context (Donat et al., 2023; Vaughan and Dessai, 2014; Delgado-Torres et al., 2025). The ResCorr isolates the added value of initialization beyond the historical simulations, providing insight into ~~the role of internally generated variability~~ climate variations not captured by the historical simulations. Together, these two metrics offer a more complete assessment of decadal prediction performance for energy-relevant indicators.

The predominantly positive PVpot trends over southern and central Europe are consistent with recent historical analyses. Segado-Moreno et al. (2026) have reported weak to positive RSDS trends over southern and central Europe when comparing observations from 1994–2004 and 2004–2023, in line with the patterns found here for the longer period 1961–2019. By contrast, studies focusing on future climate change suggest different responses, with projected decreases over northern Europe and weaker or near-neutral changes in the south (Jerez et al., 2015; Hou et al., 2021). Although these future-oriented results are not directly comparable to the present analysis, they provide context suggesting that the positive PVpot trends identified here over southern Europe are consistent with behaviours projected to strengthen in the future, whereas the future negative signals over northern Europe are not yet clearly expressed in the historical record.

4.2 Wind energy capacity factor (WCF)

Figure S9–S10 shows the climatology (Figure S9a–S10a–e) and ~~interannual–multiannual~~ standard deviation (Figure S9f–S10f–j) of WCF for the annual and seasonal means. The climatological patterns follow a distribution of wind resources over Europe, with higher WCF values over the ~~North Atlantic and~~ northern Europe and lower values over southern and Mediterranean regions, together with a clear seasonal cycle (Bett and Thornton, 2016) and higher values offshore than onshore. Unlike PVpot, WCF exhibits a non-negligible ~~interannual–multiannual~~ standard deviation, with the largest values (up to 5–6 percentage points) occurring mainly in DJF and over northern Europe, reflecting stronger year-to-year wind variability. This reflects the

intrinsically high variability of wind and the non-linear turbine power curve, which translates wind fluctuations into larger variations in WCF.

Figure 3a-e shows the ACC of WCF between DCP and ERA5 for the annual mean and each season. Overall, the percentage of grid points with significant skill is ~~generally~~ low across Europe, with the highest values ~~occurring~~ in JJA (Figure 3d), followed by the annual mean (Figure 3a) and MAM (Figure 3c). Eastern Europe consistently exhibits ~~higher~~ high skill, although the spatial patterns ~~differ with season~~ vary seasonally. In the annual mean ~~and DJF~~, DJF and MAM (Figure 3a-b), ~~there are some points with positive skill values over southern Iberia and southern France; the signal over southern France is weakly visible in MAM~~ a-c), significant positive skill is present over parts of southern France, and over Iberia in DJF. In MAM, ~~additionally~~, northern Germany and northern Poland ~~also~~ show areas of ~~significantly positive~~ significant skill. In JJA, ~~the high skill is present in many parts of central and eastern Europe and in points of the UK. In~~ skill extends across much of central Europe, whereas in SON (Figure 3e), ~~there are significant ACC values only in some points~~ it is limited to parts of Poland and the Baltics. ~~Outside these regions, ACC values are generally low~~ These regions of significant skill are also present in the coarsest-resolution results (Figure S11a-e).

The spatial distribution and the percentage of significant grid points for WCF closely resemble those obtained for SFCWIND (Figure ~~S5k-o~~ S9k-o), with only minor higher values for WCF. This similarity is expected as wind speed is the main driver of WCF.

Low predictive skill for wind-related variables is physically expected given the intrinsic characteristics of atmospheric circulation. Climate predictability partly depends on the nature and variability of the variable considered (Alizadeh, 2022). Variables that are strongly constrained by large-scale energy balances and evolve more slowly, such as TAS and RSDS, tend to be more predictable (Trenberth et al., 2009). In contrast, variables with higher variability and stronger dependence on regional and synoptic processes, such as SFCWIND or precipitation (PR), generally exhibit larger uncertainty (Gettelman and Rood, 2016; Slingo and Palmer, 2011). This behaviour is consistent with previous studies showing comparatively higher skill for decadal predictions for TAS than for PR (e.g. Smith et al., 2019; Delgado-Torres et al., 2022; Moreno-Montes et al., 2026).

Previous studies have examined the predictability of wind-related indicators for decadal predictions over Europe. Haas et al. (2015) and Moemken et al. (2016) have reported significant skill for regional peak winds and wind energy output (Eout) over central Europe, particularly for annual means and short lead times, using a statistical-dynamical downscaling approach applied to a single model. However, these results are not directly comparable to the present assessment of WCF, which relies on ~~large-scale model winds without regional downscaling~~ a simpler statistical downscaling approach based on interpolation and calibration, and on a different wind-energy metric. More recently, Hutchins et al. (2025) have reported limited and spatially heterogeneous skill for surface wind over Europe, with significant predictability mainly confined to parts of central and eastern Europe, consistent with the patterns identified here.

The ~~impact of model initialization on WCF skill is assessed using the ResCorr (Figure 3f~~ ACC between HIST and ERA5 is shown in Figure 3f-j). For all seasons and the annual mean, ResCorr exhibits both positive and negative values, indicating ~~that initialization locally enhances skill in some regions while reducing it in others. For the annual mean and j,~~ while the impact of initialization is assessed through the ResCorr in Figure 3k-o. HIST exhibits higher skill than DCP in DJF (Figure




Figure 3. Same as Figure 1 but First row: ACC between DCP and ERA5 for WCF for the annual mean (a) and seasonal means (b–e) for forecast years 1–3. Second row: ACC between HIST and ERA5 for the annual mean (f) and seasonal means (g–j). The percentages of significant grid points are indicated in brackets, and hatched regions denote non-significant correlations. Third row: ResCorr between DCP and ERA5 relative to HIST for the annual mean (k) and seasonal means (l–o). Percentages indicate the fractions of significantly positive (red) and significantly negative (blue) values. Positive (negative) ResCorr indicates higher (lower) skill for DCP compared to HIST. Correlations are computed against ERA5 during the 1961–2019 period, and statistical significance is assessed using a one-sided t-test at the 95% confidence level, accounting for time-series autocorrelation.

3f–g), initialization reduces WCF skill over Scandinavia and northern Europe, although HIST skill in these regions is already non-significant (Figure S10a–b), and in some parts of central Europe. The skill reduction may be related to the high interannual variability of WCF in these regions (Figure S9f–g), which can make the forecasts more sensitive to initialization errors. By contrast, initialization increases skill in the Atlantic coast of Iberia and Ireland for the annual mean, where HIST does not show

380 ~~significant skill, and particularly over Mediterranean regions. In contrast, HIST shows slightly lower skill than DCPD in MAM and JJA (Figure 3h-i). The ResCorr indicates that initialization enhances skill over eastern Europe for in the annual mean and DJF, where HIST does show it. In DJF, although ResCorr values are mostly non-significant over southern Europe, some regions with significantly positive HIST skill (Figure S10b) lose significance in the DCPD (Figure 3b), leading to a strong reduction in the overall percentage of significant area. In MAM (Figure 3h), positive ResCorr values are present in parts of northern~~
385 ~~France and Germany, and the DCPD (Figure 3c) has higher skill values than HIST (Figure S10h) over points of Germany but lower over points of the UK, resulting in a similar percentage of significant points. During JJA k), over northern Germany and western France in MAM (Figure 3i), initialization increases skill over parts of eastern Europe and central Germany, while reducing it over Iberia, where HIST skill is already low (Figure S10d), resulting in a higher fraction of significant area for the DCPD than for the HIST. In SON m), and over eastern Europe in JJA (Figure 3j)n). Outside these regions, ResCorr values~~
390 ~~remain close to zero and DCPD skill (Figure 3e) largely mirrors that of the HIST (Figure S10e). Overall, model initialization affects WCF skill mainly in winter and summer: it leads to a marked reduction of skill in DJF, while it enhances skill in JJA. In the remaining seasons, initialization has a limited impact and largely preserves the skill already present in the historical simulations are generally weak and spatially incoherent. Similar conclusions are obtained at the native resolution of the coarsest prediction system (Figure S11), which shows the same large-scale patterns, supporting the robustness of these results. Figure~~
395 ~~S12 further shows that the ResCorr patterns are largely insensitive to ensemble size, with generally small and non-significant differences across Europe.~~

Applying the same regionalization used for PVpot, regional trends of WCF differ markedly between ERA5 and the DCPD (Figure 4a-b). ERA5 shows no uniform behaviour, with both significantly positive and negative trends depending on region and season, whereas the DCPD exhibits predominantly significant negative trends across all regions and seasons. WCF trends
400 follow similar patterns to those of SFCWIND (Figure S8e-fS9e-f) for both ERA5 and the DCPD.

Consequently, regional correlations between ERA5 and the DCPD (Figure 4c) are significantly positive mainly where both datasets display coherent negative trends. This is most evident in Eastern Europe, with significant correlations for the annual mean, JJA and SON, and weaker agreement in DJF and MAM, consistent with the regions showing skill in Figure 3. Similar behaviour appears in Central Europe in JJA and SON, with additional agreement in Western Europe in MAM and weaker in
405 Iberia in DJF. Elsewhere, differences in trend sign or magnitude lead to weak or negative correlations. Overall, these results emphasize the key role of trend consistency in explaining regional skill.

ResCorr regional values (Figure 4d) are generally non-significant, with a few isolated exceptions. A significantly positive ResCorr appears in Eastern Europe in JJA, aligned with the agreement in negative trends between ERA5 and the DCPD (Figure 4a-b) and the significantly positive correlation in Figure 4c. In addition, ResCorr is ~~significantly-significantly~~ positive in Central
410 Europe in MAM, ~~even though the correlation is not significant and trend agreement is weak, indicating the increase of skill due to the initialization in this case; however, this signal is driven by localized positive values (Figure 3h) and may be influenced by sampling variability.~~ Other regions show non-significant but positive ResCorr values, suggesting local improvements due to initialization. In ~~some of these cases, trends agree and correlations are positive (e.g. Western Europe in MAM). In others, trends have opposite signs and correlations are~~ regions with weak or negative, ~~indicating that the limited skill comes mainly~~




Figure 4. Same as Figure 2 but for WCF trends derived from ERA5 for years 1961-2019 with a three years rolling mean (a) and from the DCPP for the start dates 1960-2016 for over forecast years 1–3 (b), the correlation between ERA5 and DCPP (c), and the ResCorr between DCPP and ERA5 respect HIST (d) shown for the annual mean and for each season across the seven european sub-regions (Figure S4). Statistically significant trends and correlations at the 95% confidence level are marked with an asterisk.

415 from initialization (e.g. Western Europe in JJA; Mediterranean in JJA and SON; Great Britain and Ireland in SON and annual). Conversely, there are regions where trends have the same sign and correlations are significantly positive, but ResCorr is close to zero (e.g. Eastern Europe in Annual and SON; Central Europe in JJA and SON). This suggests that most of the skill in those cases is explained by the externally forced trend rather than by initialization. Taken together, these results show that initialization can enhance skill in some regions, sometimes reinforcing existing trend agreement and in other cases providing additional skill independently of it. correlations, however, they mainly reflect a reduction in negative skill rather than a transition to meaningful predictive skill.

The WCF trends found are consistent with signals reported in previous studies. Using climate projections, Pryor et al. (2005) have identified slight decreases in wind speed and wind energy density over northern Europe. CMIP6 historical simulations, Miao et al. (2023) identified negative historical wind speed trends over Europe across a broad CMIP6 ensemble, including the three CMIP6 models analysed in the present study. Similarly, Olmo et al. (2026) Pryor and Barthelmie (2010), analysing future

climate scenarios, reported a projected weakening of near-surface winds over Europe, particularly towards the late 21st century. Although these projections are not directly comparable with the present analysis, they point to a tendency towards declining wind resources that may already be weakly emerging in current conditions. Observational ~~analyses by Vautard et al. (2010) and Torralba et al. (2017a) have revealed~~ and reanalysis-based studies (e.g. Vautard et al. (2010); Torralba et al. (2017a)) have
430 ~~reported~~ spatially heterogeneous near-surface wind trends across Europe, ~~with predominantly negative signals in several regions.~~ Our results are broadly consistent in showing negative trends over central and eastern Europe and weak signals over southern regions. However, differences remain over northern Europe and the British Isles, where these previous studies reported neutral or negative trends while positive trends are found here, likely due to differences in datasets, periods and methodologies. Together, these studies, based on different datasets and periods, indicate that declining wind signals have been
435 reported in both modelling and observational contexts, while also highlighting the substantial uncertainty in the magnitude and spatial structure of historical wind changes.

4.3 Number of energy droughts (NED)

Figure ~~S11–S13~~ shows the climatology (Figure ~~S11a–e~~) and interannual standard deviation (Figure ~~S11f–j~~) of NED for the annual mean and each season. Overall, NED values are relatively high across Europe, reaching around 50% for the annual
440 mean (Figure ~~S11a~~). Seasonal patterns reflect the relative availability of each resource. In DJF and SON (Figure ~~S11b,e~~), the NED values are generally high (between 40 and 100% in all the regions) and the highest ones occur particularly in mountainous regions ($>80\%$), where low multiannual standard deviation of NED, while Figure S14 shows the climatology of solar-Neff (Figure ~~S14a–e~~) and wind-Neff (Figure ~~S14f–j~~). NED is generally highest in DJF (Figure ~~S13b~~) and SON (Figure ~~S13e~~) and lowest in JJA (Figure ~~S13d~~), reflecting the strong seasonal cycle of solar-Neff ($<5\%$) (Figure ~~S12b,e~~) and low
445 wind-Neff ($<10\%$) (Figure ~~S12g,j~~) are found. Wind-Neff has values $<10\%$ in mountainous regions for all the seasons. In MAM (Figure ~~S11e~~). Spatially, higher NED values appear over parts of central Europe and some mountainous regions, mainly associated with low wind-Neff (Figure ~~S12h~~), are found in mountainous regions and, during summer, over northern Europe. Wind-Neff is consistently low over mountainous areas and largely explains several regional NED features, while solar-Neff (Figure ~~S12e~~) decreases primarily with latitude. In JJA (Figure ~~S11d~~), although being low (between 0 and 20%), higher NED
450 values are largely confined to northern Europe, reflecting the strong latitudinal gradient in solar-Neff (Figure ~~S12d~~), despite generally favourable solar conditions elsewhere. The standard deviation values are particularly high mainly follows a latitudinal gradient. The largest NED variability occurs in DJF (Figure ~~S11g~~) and over mountainous regions ~~S13g~~, in central and eastern Europe in JJA (Figure ~~S11i~~). The high standard deviation observed over northern Poland and the Baltics in almost all seasons closely resembles the patterns found for the WCF standard deviation (Figure ~~S8f–j~~), highlighting the strong influence of wind
455 variability on NED ~~S13i~~) and in some regions of northern Europe in SON (Figure ~~S13j~~).

The relatively high climatological values of the NED indicator mainly reflect the use of fixed, efficiency-based thresholds to define effective energy production. Previous studies have analyzed compound energy droughts using percentile-based metrics (e.g. Otero et al., 2022), which characterise relative extremes within each region. Despite these different definitions, the main seasonal patterns they report of higher drought frequencies in winter over southern Europe and in summer over northern

460 regions are broadly consistent with our results. Here, fixed thresholds motivated by operational efficiency are adopted, an approach used in previous studies like Richardson et al. (2023). In our case, the prior calibration of all climate variables against ERA5 ensures consistent magnitudes, supporting the use of absolute, efficiency-related thresholds to assess compound energy droughts reduces systematic biases relative to the observational reference, including biases in mean values and variability. This helps avoid artificial shifts in the frequency of threshold exceedances caused by biases in the raw model data.

465 Figure 5a-e shows the ACC between the DCP and ERA5 of NED. The highest fraction of significant ACC values occurs in JJA (Figure 5d), when most of eastern Europe and parts of central Europe and Iberia exhibit significant skill. The annual mean (Figure 5a) and MAM (Figure 5c) show significant ACC values over regions including also show significant values over parts of eastern Europe and the Atlantic coast of Iberia; however, MAM displays a slightly higher percentage of significant area, mainly due to additional significance over Italy. By contrast, skill is generally limited in DJF (Figure 5b), where significant ACC appears only in small areas of eastern Europe and Iberia, and in Iberia. In SON (Figure 5e), where skill is skill is generally weak and confined to parts of eastern Europe and the Baltics. Overall, the skill patterns are broadly consistent with those obtained at the coarsest resolution (Figure S15a-e).

Figure S13 shows the ACC of the DCP S16 presents the corresponding ACC for solar-Neff (Figure S13a-S16a-e), and wind-Neff (Figure S13f-S16f-j), and both-Neff (Figure S13k-o). The skill patterns of solar-Neff closely resemble those of solar PVpot (Figure 1a-e), while wind-Neff shows patterns similar to those of WCF (Figure 3a-e), as expected given that all three indicators reflect energy-production efficiency. By contrast, both-Neff exhibits more spatially fragmented skill, with isolated regions of significant values. In solar-Neff and both-Neff, some areas appear as missing values because correlations cannot be computed when the indicator remains equal to zero for all years, and solar energy droughts are expectable when the amount of daily sun hours is low (Figure S13b, e).

480 The. Consequently, the spatial distribution of NED skill generally reflects the combined, but not necessarily simultaneous, reflects the seasonally varying contribution of solar- and wind-Neff predictability wind-related conditions. For the annual mean (Figure 5a), significant NED skill over parts of eastern Europe and Iberia broadly coincides with regions where both solar-Neff (Figure S13a-S16a) and wind-Neff (Figure S13f) are skillful. By contrast, over parts of the Alps and southeastern region, NED exhibits significant skill despite wind-Neff being non-significant, indicating a dominant contribution from S16f show positive skill. Over parts of southeastern Europe, including the Carpathian region and Greece, localized NED skill is more consistent with solar-Neff. In DJF (Figure 5b), NED skill over Iberia and eastern Europe aligns mainly than with wind-Neff skill (Figure S13g). In MAM (Figure 5c), NED skill over southern and southeastern Europe Europe generally follows the solar-Neff signal, although NED remains non-significant in many-several regions where solar-Neff shows skill. In During JJA (Figure 5d), the widespread NED skill closely follows-resembles the solar-Neff pattern (Figure S13d), only showing, with lower values over parts of Western Europe. Finally, in SON (Figure 5e) western Europe. Over most of Europe in DJF and in northern regions in SON, as solar-Neff remains close to zero (Figure S14b,e), NED skill is confined to parts of Poland and the Baltic region, closely matching ACC can not be computed (Figure S16b,e) and NED ACC is directly the wind-Neff skill pattern (Figure S13j) and in points of southeastern Europe, which are skillful at solar-Neff (Figure S13e).




Figure 5. Same as Figure 1 but First row: ACC between DCP and ERA5 for NED for the annual mean (a) and seasonal means (b–e) for forecast years 1–3. Second row: ACC between HIST and ERA5 for the annual mean (f) and seasonal means (g–j). The percentages of significant grid points are indicated in brackets, and hatched regions denote non-significant correlations. Third row: ResCorr between DCP and ERA5 relative to HIST for the annual mean (k) and seasonal means (l–o). Percentages indicate the fractions of significantly positive (red) and significantly negative (blue) values. Positive (negative) ResCorr indicates higher (lower) skill for DCP compared to HIST. Correlations are computed against ERA5 during the 1961–2019 period, and statistical significance is assessed using a one-sided t-test at the 95% confidence level, accounting for time-series autocorrelation.

Overall, these results may indicate that NED predictability is strongly seasonally modulated by the dominant energy source. During periods of high solar availability (JJA and, in the southern region, MAM), one (Figure S16g,j). In SON, outside these regions, the weak NED skill is largely driven by solar-Neff, whereas in low-radiation seasons (DJF and SON) more consistent

with wind-Neff becomes the main modulator (Figure S16j), although some contribution from solar-Neff (Figure S16e) is also apparent.

The heterogeneous ResCorr patterns (ACC obtained with HIST is shown in Figure 5f–j), with both positive and negative values in all seasons, indicate that, while the impact of initialization on this indicator varies across seasons and regions. In the annual mean is assessed through ResCorr (Figure 5f) k–o). Overall, HIST reproduces the main spatial patterns found in DCP, positive ResCorr values appear along the Atlantic coast of Iberia and in parts of eastern Europe, while negative values dominate along the rest of the Atlantic coast. Although the DCP skill pattern (Figure 5a) resembles that of the HIST (Figure S14a) and ResCorr is not significant there, the skill of DCP is lower than that of HIST in eastern Iberia and over a large part of Italy, as well as in the fraction of significant grid points. In DJF although the relative skill varies by season, HIST shows slightly lower skill than DCP in MAM (Figure 5g), ResCorr is positive in parts of eastern and Southwestern Europe but negative in parts of northern Europe, resulting in a lower skill for DCP (Figure 5b) than for HIST (Figure S14b). In MAM h) and particularly in JJA (Figure 5h), initialization enhances skill over parts of southern and eastern Europe while reducing it over Iberia, leading to a modest improvement of DCP skill relative to HIST i), whereas it has higher skill in DJF (Figure 5c vs Figure S14c). During g), JJA (Figure 5i), g) exhibits widespread positive ResCorr values extend across large areas of central and southern Europe, and DCP shows clearly higher skill than HIST (Figure 5d vs Figure S14d). By contrast, in SON (Figure 5j), ResCorr is predominantly negative with only isolated positive values, resulting in a lower fraction of significant skill in DCP (Figure 5e) compared to HIST across central, southern and eastern Europe, consistent with the substantially larger fraction of significant ACC values in DCP (Figure S14e). For this indicator, initialization generally increases predictability under conditions of higher sunshine duration (JJA and MAM) and reduces it under the opposite conditions (DJF and SON) 38.8% than in HIST (24.1%). ResCorr values are generally weak and spatially heterogeneous in the rest of the seasons. Similar conclusions are obtained at the native resolution of the coarsest prediction system (Figure S16f–o), with broadly consistent HIST ACC and ResCorr patterns. The main exception is DJF (Figure S16g), where HIST exhibits substantially fewer significant ACC values at the coarser resolution. This suggests that part of the additional skill identified in the high-resolution results during this season may reflect spatial noise rather than robust large-scale predictability. Figure S17 shows that the ResCorr differences obtained when subsampling the ensembles are generally close to zero and non-significant across Europe, indicating that the NED results are largely insensitive to the unequal ensemble sizes of the prediction systems.

Figure 6 shows the regional trends of NED for ERA5 (a), for the DCP (b), the corresponding correlations (c), and the ResCorr (d) across the seven sub-regions defined above. In some regions and seasons, ERA5 and DCP exhibit similar trends, while in others they show opposite signs. In JJA, Iberia, Western Europe, Mediterranean, Central and Eastern Europe display significant negative trends in both datasets, which results in significantly positive correlations (Figure 6c). A similar behaviour is found for the annual mean and MAM in Iberia, Western Europe and Mediterranean. In addition, there are cases where both ERA5 and DCP show significantly positive trends, leading to positive, though non-significant, correlations, such as in DJF for Iberia, in SON for Central Europe, and in DJF and SON for Eastern Europe. In the remaining regions and seasons, ERA5 and DCP trends generally differ in sign, typically negative in ERA5 and positive in the DCP, resulting in correlations that




Figure 6. Same as Figure 2 but for NED trends derived from ERA5 for years 1961-2019 with a three years rolling mean (a) and from the DCPP for the start dates 1960-2016 for over forecast years 1–3 (b), the correlation between ERA5 and DCPP (c), and the ResCorr between DCPP and ERA5 respect HIST (d) shown for the annual mean and for each season across the seven european sub-regions (Figure S4). Statistically significant trends and correlations at the 95% confidence level are marked with an asterisk.

are weak or negative. Overall, these results again underline the importance of consistently reproducing the trend for obtaining high correlation values.

ResCorr (Figure 6d) partly mirrors the correlation patterns shown in Figure 6c. Regions and seasons with significantly positive ResCorr values (notably Iberia, Mediterranean, Central and Eastern Europe in JJA; and more weakly Iberia in DJF; Mediterranean in MAM,) or high but not significant ones (Mediterranean and Eastern Europe in SON) Annual generally correspond to cases where correlations are also significantly positive. This indicates that, in these cases, initialization enhances skill beyond the externally forced signal already contributing to the correlation. However, there are also cases with significantly positive correlations but low or negative ResCorr values (e.g. Iberia in MAM and Annual; Mediterranean in Annual). In these situations, the agreement in trends between ERA5 and the DCPP explains most of the skill, while initialization adds little or even decreases predictive value.

NED variability is largely controlled by the energy source that dominates effective production in each season: wind in DJF and SON, To better understand the contribution of solar-Neff and solar in JJA and, secondarily, in MAM. As a result, wind-Neff to NED variability and trends, solar-Neff and wind-Neff trends are analysed separately (Figure S18). The results show a clear seasonal dependence. In JJA, the strong negative NED trends in DJF and SON tend to be anticorrelated with WCF trends (Figure 4). In the DCP, WCF trends are consistently negative across regions, while NED trends are generally positive, leading to opposite signs. In ERA5, WCF trends are more heterogeneous, and this spatial variability is reflected in the NED trends; for example, positive WCF trends in DJF over Scandinavia and Great Britain and Ireland and negative trends over Iberia correspond to opposite sign trends in NED. A similar, though weaker, relationship is found between NED and PVpot in JJA and in southern regions during MAM. In these cases, positive trends in PVpot (Figure 2) for both ERA5 and DCP are primarily associated with marked increases in solar-Neff (Figure S18a-b), while wind-Neff changes are weaker (Figure S18c-d). In MAM, solar-Neff also drives NED trends in ERA5, although with smaller magnitude, whereas in DCP this signal is less clear due to strong negative trends in wind-Neff. In DJF, solar-Neff shows negligible variability over most regions, and NED trends are therefore directly anticorrelated with wind-Neff. In SON, both components exhibit mixed signals in ERA5 and the DCP are associated with generally negative, while in DCP negative wind-Neff trends tend to dominate, leading to positive NED trends. In the annual means, the strong negative trends in NED trends, particularly in JJA and in southern regions during MAM. This link is clearer in ERA5 and less pronounced in DCP seem to be dominated by positive solar-Neff trends, while the mixed DCP signals are influenced by both energy sources.

The negative NED trends identified in ERA5 for the recent past are consistent with the ERA5-based analysis of (Meng et al., 2025) Meng et al. (2025), which have reported a decrease in the frequency of renewable energy droughts under present-day conditions, despite differences in indicator definition. Additionally, DCP trends show seasonal patterns that broadly resemble some qualitative similarities with the future evolution of hybrid energy droughts projected by (Kapica et al., 2024) Kapica et al. (2024), particularly in DJF and SON. Although their results refer to future climate scenarios and are based on a different definition of compound events, the similarities suggest that some of the changes projected for the future may already be emerging in the decadal predictions analyzed here MAM and over parts of northern and eastern Europe in DJF. However, Kapica et al. (2024) have used a percentile-based definition of droughts, whereas the NED metric employed here is based on fixed physically motivated thresholds. The comparison is therefore limited to broad spatial patterns, and important regional differences remain, particularly in SON.

5 Conclusions

The renewable energy sector is strongly influenced by variability and long-term changes in key climate variables such as temperature, solar radiation and wind speed. However, the intermediate timescale of the coming years remains comparatively underexplored, and there is a clear need for tailored climate information to bridge the gap between decadal prediction advances and their application in renewable energy planning.

This study evaluates the predictive skill of a multi-model ensemble of decadal predictions (DCPP) for energy-relevant indicators over Europe, focusing on photovoltaic potential (PVpot), wind capacity factor (WCF), and the compound indicator number of energy droughts (NED) for forecast years 1–3. Skill is assessed using the ACC between the DCPP and ERA5, and the impact of model initialization is estimated with the ResCorr relative to historical simulations. In addition to grid-point analyses, sub-regional mean trends and correlations are examined to explore the relationship between long-term changes and predictive skill.

PVpot shows the highest and most spatially coherent skill among the indicators analyzed. Skill is significantly positive over large parts of the region at the annual scale and during MAM and JJA, while it is weaker and more spatially limited in DJF and SON. ~~This strong seasonal dependence indicates that PVpot predictability is highest during periods of higher and persistent solar radiation, while such predictability is more limited under weaker and more variable radiative conditions.~~ The spatial and seasonal patterns of PVpot skill closely follow those of surface downwelling shortwave radiation (RSDS), its main driver. PVpot skill is slightly lower than RSDS skill because the indicator is also modulated by near-surface air temperature (TAS) and surface wind speed (SFCWIND), which act as secondary constraints on predictability.

Initialization ~~clearly improves~~ enhances PVpot skill in the annual mean and particularly in JJA, mainly by ~~enhancing already positive skill values of the~~ strengthening regions that already exhibit skill in multi-model of historical simulations (HIST). In DJF and SON, the added value of initialization is limited and spatially heterogeneous, with localized improvements and degradations. ~~Overall, initialization tends to amplify existing PVpot predictability in HIST.~~

WCF exhibits lower and more spatially heterogeneous skill than PVpot, reflecting the higher intrinsic variability of wind and its strong sensitivity to synoptic-scale circulation. Significant skill is consistently found over parts of eastern Europe across seasons, while other regions show ~~skill only seasonally~~ weaker and more seasonally dependent skill, including central Europe in JJA ~~, southern Europe in DJF,~~ and northern Europe in MAM and SON. The spatial distribution of WCF skill closely mirrors that of SFCWIND, confirming wind variability as the primary control on WCF predictability.

~~Initialization tends to enhance skill in summer and reduce it in winter, while its impact is generally small in the transition seasons.~~ The impact of initialization is regionally dependent and generally limited. The clearest skill enhancements are found in JJA over parts of eastern Europe, while in other seasons the impact of initialization is weaker and spatially heterogeneous. As for PVpot, initialization mainly improves skill where historical simulations already show some skill. The ~~overall low skill for WCF is partly associated with opposite trend signs~~ generally low WCF skill is consistent with the weak agreement between ERA5 and ~~the DCPP~~ DCPP trends in many regions and seasons, together with the high intrinsic variability of wind.

The compound indicator NED displays predictability characteristics that reflect the combined influence of solar and wind resources. NED skill is highest in JJA, followed by the annual mean and MAM, and is generally limited in DJF and SON. NED predictability does not require simultaneous skill in both components: in some regions it emerges where both solar-Neff and wind-Neff are predictable, while in others it is dominated by a single source. This highlights the ~~non-linear nature of compound energy droughts and the~~ importance of ~~interactions between resources~~ considering the relative contribution of both resources when assessing compound energy drought predictability.

Seasonal differences in NED skill reflect the ~~dominant energy source. During high-radiation periods (JJA and partly MAM) relative contribution of wind and solar components. In JJA, NED predictability is mainly driven by solar-Neff, whereas during low-radiation seasons (DJF and SON) wind-Neff becomes the main contributor. The~~. In DJF, variability over much of Europe is largely driven by wind conditions, while in MAM and SON the relative contribution of both components varies regionally. As a result, the impact of initialization on NED ~~follows this seasonal dependence~~ is also seasonally dependent, enhancing skill when the dominant source is more predictable and reducing it otherwise. These results show that energy-relevant indicators derived from decadal predictions can energy source is predictable, while providing little added value otherwise. This is consistent with the broader finding that initialization generally enhances existing predictability rather than creating skill where historical simulations show none. These results demonstrate the potential of decadal predictions to provide actionable information ~~to anticipate energy droughts. However, predictability is for~~ anticipating energy droughts, although forecast quality remains strongly dependent on season, region and indicator. Initialization generally enhances skill where some degree of predictability already exists but rarely creates new skill when historical simulations do not have it and region.

By evaluating compound energy indicators, explicitly separating total skill from the added value of initialization, and analysing results at the regional scale, this study advances beyond variable-based assessments and provides a more application-oriented understanding of decadal predictability for renewable energy. These findings highlight both the potential and the limitations of decadal predictions for energy-related applications and underscore the need to clearly communicate where and when such information is reliable. Besides, the main conclusions were robust across spatial resolutions and were largely insensitive to differences in ensemble size between DCP and HIST.

Several limitations should be noted. ERA5 is the only reference dataset used, and reanalysis uncertainties, especially for wind, may affect skill estimates. However, ERA5 has been identified as ~~the best reanalysis for wind~~ one of the most reliable reanalysis datasets for near-surface wind variability, particularly in terms of correlation with observations and representation of temporal variability at turbine-relevant heights (Ramon et al., 2019) and Global Wind Atlas has been used to correct it for a more accurate representation. Secondly, fixed, efficiency-based thresholds, although operationally motivated, may limit sensitivity and transferability across regions or technologies. Besides, temporal aggregation choices influence the indicators, particularly for wind, given the finer temporal resolution of turbine power curves. In addition, the power-law extrapolation used to estimate 100 m wind speeds assumes neutral stability and may be less representative under non-neutral atmospheric conditions, which could affect the reliability of local-scale decadal predictions. Finally, the ACC and ResCorr metrics capture correlation-based skill but do not fully describe other aspects relevant for operational decision-making, such as reliability or the representation of extreme events. Other metrics might be selected depending on the climate forecast products required by the energy users.

Future work could assess how sensitive the indicators are to the choice of thresholds (e.g. different RSDS or WCF limits), compare results using alternative reanalyses or observation-based datasets. The framework could also be extended to include electricity demand, which would allow a more comprehensive assessment of supply–demand imbalances and increase the relevance of the indicators for system operators and energy planners. ~~In addition, hybrid approaches that combine percentile-based metrics with physically motivated thresholds may help balance climatological comparability and operational~~

~~relevance when assessing compound renewable energy risks~~ Furthermore, as decadal prediction systems are known to be affected by signal-to-noise problems (Scaife and Smith, 2018), future studies could further investigate the impact of signal-to-noise characteristics of the calibrated ensemble or apply correction methods such as the NAO-matching proposed by Smith et al. (2020)

645 .

Code availability. We acknowledge the use of the startR, s2dv, CStools, multiApply, and CSIndicators R-language-based software packages, all of them available on the Comprehensive R Archive Network (CRAN; <https://cran.r-project.org/>, last access: 2 March 2026). The code used during the study is available from the corresponding author on reasonable request.

650 *Data availability.* All datasets used in the study are publicly available. SEAS5 predictions and ERA5 reanalysis data are available from the Copernicus Climate Data Store (CDS; <https://cds.climate.copernicus.eu/>, last access: 2 March 2026). The decadal predictions, historical simulations and climate projections are available on the Earth System Grid Federation system (ESGF; <https://esgf.github.io/>, last access: 2 March 2026).

Author contributions. SMM and CDT designed the study. SMM carried out the analysis and wrote the first draft. All authors contributed equally to the interpretation of results and writing thereafter.

655 *Competing interests.* The contact author has declared that none of the authors has any competing interests.

660 *Acknowledgements.* This study has been supported by the Spanish national project BOREAS (PID2022-140673OA-I00) funded by MICI-U/AEI/10.13039/501100011033 and by ERDF, EU. SMM gratefully acknowledges financial support from the Spanish Ministry for Science and Innovation (FPI PREP2022-000684, funded by MCIN/AEI/10.13039/501100011033). MEO is funded by the AI4Science PN070500 fellowship within the “Generación D” initiative, Red.es, Ministerio para la Transformación Digital y de la Función Pública, for talent attraction (C005/24-ED CV1). Funded by the European Union NextGenerationEU funds, through PRTR. VT has received funding from the EU Horizon 2020 Marie Skłodowska-Curie grant 101152499 (SINFONIA).

References

- Alizadeh, O.: Advances and challenges in climate modeling, *Climatic Change*, 170, 18, 2022.
- Aranyosy, A., De Luca, P., Delgado-Torres, C., Solaraju-Murali, B., Samso Cabre, M., and Donat, M. G.: Multi-annual predictions of hot,
665 dry and hot-dry compound extremes, *Earth System Dynamics*, 16, 2225–2251, 2025.
- Association, W. M. et al.: WMO global annual to decadal climate update, 2023.
- Bellucci, A., Haarsma, R., Gualdi, S., Athanasiadis, P., Caian, M., Cassou, C., Fernandez, E., Germe, A., Jungclaus, J., Kröger, J., et al.: An
assessment of a multi-model ensemble of decadal climate predictions, *Climate Dynamics*, 44, 2787–2806, 2015.
- Bett, P. E. and Thornton, H. E.: The climatological relationships between wind and solar energy supply in Britain, *Renewable Energy*, 87,
670 96–110, 2016.
- Bett, P. E., Thornton, H. E., Troccoli, A., De Felice, M., Suckling, E., Dubus, L., Saint-Drenan, Y.-M., and Brayshaw, D. J.: A simplified
seasonal forecasting strategy, applied to wind and solar power in Europe, *Climate services*, 27, 100 318, 2022.
- Bloomfield, H. C., Brayshaw, D. J., Gonzalez, P. L., and Charlton-Perez, A.: Sub-seasonal forecasts of demand and wind power and solar
power generation for 28 European countries, *Earth System Science Data*, 13, 2259–2274, 2021.
- 675 Boccard, N.: Capacity factor of wind power realized values vs. estimates, *energy policy*, 37, 2679–2688, 2009.
- Boer, G. J., Smith, D. M., Cassou, C., Doblas-Reyes, F., Danabasoglu, G., Kirtman, B., Kushnir, Y., Kimoto, M., Meehl, G. A., Msadek, R.,
et al.: The decadal climate prediction project (DCPP) contribution to CMIP6, *Geoscientific Model Development*, 9, 3751–3777, 2016.
- Brower, M.: Wind resource assessment: a practical guide to developing a wind project, John Wiley & Sons, 2012.
- Carvalho, D., Rocha, A., Costoya, X., DeCastro, M., and Gómez-Gesteira, M.: Wind energy resource over Europe under CMIP6 future
680 climate projections: What changes from CMIP5 to CMIP6, *Renewable and Sustainable Energy Reviews*, 151, 111 594, 2021.
- Castillo, C. P., e Silva, F. B., and Lavalle, C.: An assessment of the regional potential for solar power generation in EU-28, *Energy policy*,
88, 86–99, 2016.
- Challinor, A. J., Adger, W. N., and Benton, T. G.: Climate risks across borders and scales, *Nature Climate Change*, 7, 621–623, 2017.
- Chenni, R., Makhoulouf, M., Kerbache, T., and Bouzid, A.: A detailed modeling method for photovoltaic cells, *Energy*, 32, 1724–1730, 2007.
- 685 Cionni, I., Lledo, L., Torralba, V., and Dell’Aquila, A.: Seasonal predictions of energy-relevant climate variables through Euro-Atlantic
Teleconnections, *Climate Services*, 26, 100 294, 2022.
- Clark, R. T., Bett, P. E., Thornton, H. E., and Scaife, A. A.: Skilful seasonal predictions for the European energy industry, *Environmental
Research Letters*, 12, 024 002, 2017.
- Corti, S., Weisheimer, A., Palmer, T., Doblas-Reyes, F., and Magnusson, L.: Reliability of decadal predictions, *Geophysical Research Letters*,
690 39, 2012.
- Delgado-Torres, C., Donat, M. G., Gonzalez-Reviriego, N., Caron, L.-P., Athanasiadis, P. J., Bretonnière, P.-A., Dunstone, N. J., Ho, A.-C.,
Nicoli, D., Pankatz, K., et al.: Multi-model forecast quality assessment of CMIP6 decadal predictions, *Journal of Climate*, 35, 4363–4382,
2022.
- Delgado-Torres, C., Donat, M. G., Soret, A., González-Reviriego, N., Bretonnière, P.-A., Ho, A.-C., Pérez-Zanón, N., Samso Cabre, M., and
695 Doblas-Reyes, F. J.: Multi-annual predictions of the frequency and intensity of daily temperature and precipitation extremes, *Environmen-
tal Research Letters*, 18, 034 031, 2023.

- Delgado-Torres, C., Octenjak, S., Marcos-Matamoros, R., Pérez-Zanón, N., Baulenas, E., Doblas-Reyes, F. J., Donat, M. G., Lwiza, L. M., Milders, N., Soret, A., et al.: Supporting food security with multi-annual climate information: Co-production of climate services for the Southern African Development Community, *Science of The Total Environment*, 975, 179 259, 2025.
- 700 Donat, M. G., Delgado-Torres, C., De Luca, P., Mahmood, R., Ortega, P., and Doblas-Reyes, F. J.: How credibly do CMIP6 simulations capture historical mean and extreme precipitation changes?, *Geophysical Research Letters*, 50, e2022GL102 466, 2023.
- Done, J. M., Morss, R. E., Lazrus, H., Towler, E., Tye, M. R., Ge, M., Das, T., Munévar, A., Hewitt, J., and Hoeting, J. A.: Toward usable predictive climate information at decadal timescales, *One Earth*, 4, 1297–1309, 2021.
- Duffie, J. A. and Beckman, W. A.: *Solar engineering of thermal processes*, Wiley New York, 1980.
- 705 Dunstone, N., Lockwood, J., Solaraju-Murali, B., Reinhardt, K., Tsartsali, E. E., Athanasiadis, P. J., Bellucci, A., Brookshaw, A., Caron, L.-P., Doblas-Reyes, F. J., et al.: Towards useful decadal climate services, *Bulletin of the American Meteorological Society*, 103, E1705–E1719, 2022.
- Elsner, J. B. and Schmertmann, C.: Assessing forecast skill through cross validation, *Weather and Forecasting*, 9, 619–624, 1994.
- Ely, C. R., Brayshaw, D. J., Methven, J., Cox, J., and Pearce, O.: Implications of the North Atlantic Oscillation for a UK–Norway renewable power system, *Energy Policy*, 62, 1420–1427, 2013.
- 710 European Commission: *Powering a climate-neutral economy: An EU strategy for energy system integration*, 2020.
- European Environment Agency: *Share of energy consumption from renewable sources in Europe*, Published: 26.10. 022, 2022.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, 2016.
- 715 Gettelman, A. and Rood, R. B.: *Demystifying climate models: A users guide to earth system models*, Springer, 2016.
- Ghosh, S., Dey, S., Ganguly, D., Baidya Roy, S., and Bali, K.: Cleaner air would enhance India’s annual solar energy production by 6–28 TWh, *Environmental Research Letters*, 17, 054 007, 2022.
- Ghosh, S., Ganguly, D., Dey, S., and Chowdhury, S. G.: Future photovoltaic potential in India: navigating the interplay between air pollution control and climate change mitigation, *Environmental Research Letters*, 19, 124 030, 2024.
- 720 Gonzalez, P. L., Brayshaw, D. J., and Zappa, G.: The contribution of North Atlantic atmospheric circulation shifts to future wind speed projections for wind power over Europe, *Climate Dynamics*, 53, 4095–4113, 2019.
- Gutiérrez, J. M., Maraun, D., Widmann, M., Huth, R., Hertig, E., Benestad, R., Roessler, O., Wibig, J., Wilcke, R., Kotlarski, S., et al.: An intercomparison of a large ensemble of statistical downscaling methods over Europe: Results from the VALUE perfect predictor cross-validation experiment, *International journal of climatology*, 39, 3750–3785, 2019.
- 725 Haas, R., Reyers, M., and Pinto, J. G.: Decadal predictability of regional-scale peak winds over Europe using the Earth System Model of the Max-Planck-Institute for Meteorology, *Meteorol. Z*, 25, 739–752, 2015.
- Hamed, K. H. and Rao, A. R.: A modified Mann-Kendall trend test for autocorrelated data, *Journal of hydrology*, 204, 182–196, 1998.
- Heide, D., Von Bremen, L., Greiner, M., Hoffmann, C., Speckmann, M., and Bofinger, S.: Seasonal optimal mix of wind and solar power in a future, highly renewable Europe, *Renewable Energy*, 35, 2483–2489, 2010.
- 730 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., et al.: The ERA5 global reanalysis, *quarterly journal of the royal meteorological society*, *Quarterly Journal of the Royal Meteorological Society*, 2020.
- Hou, X., Wild, M., Folini, D., Kazadzis, S., and Wohland, J.: Climate change impacts on solar power generation and its spatial variability in Europe based on CMIP6, *Earth System Dynamics*, 12, 1099–1113, 2021.

- 735 Hsu, S., Meindl, E. A., and Gilhousen, D. B.: Determining the power-law wind-profile exponent under near-neutral stability conditions at sea, *Journal of Applied Meteorology* (1988-2005), pp. 757–765, 1994.
- Hutchins, B. W., Brayshaw, D. J., Shaffrey, L. C., Thornton, H. E., and Smith, D. M.: Decadal prediction for the European energy sector, *Meteorological Applications*, 32, e70 054, 2025.
- International Energy Agency: Renewables 2023, 2024.
- 740 Jerez, S., Tobin, I., Vautard, R., Montávez, J. P., López-Romero, J. M., Thais, F., Bartok, B., Christensen, O. B., Colette, A., Déqué, M., et al.: The impact of climate change on photovoltaic power generation in Europe, *Nature communications*, 6, 10 014, 2015.
- Jerez, S., Tobin, I., Turco, M., Jiménez-Guerrero, P., Vautard, R., and Montávez, J. P.: Future changes, or lack thereof, in the temporal variability of the combined wind-plus-solar power production in Europe, *Renewable energy*, 139, 251–260, 2019.
- Kalverla, P. C., Duncan Jr, J. B., Steeneveld, G.-J., and Holtslag, A. A.: Low-level jets over the North Sea based on ERA5 and observations: together they do better, *Wind Energy Science*, 4, 193–209, 2019.
- 745 Kapica, J., Jurasz, J., Canales, F. A., Bloomfield, H., Guezgouz, M., De Felice, M., and Kobus, Z.: The potential impact of climate change on European renewable energy droughts, *Renewable and Sustainable Energy Reviews*, 189, 114 011, 2024.
- Kealy, T., Barrett, M., and Kearney, D.: How profitable are wind turbine projects? An empirical analysis of a 3.5 MW wind farm in Ireland, *International journal on recent technologies in mechanical and electrical engineering*, 2015.
- 750 Kharin, V., Boer, G., Merryfield, W., Scinocca, J., and Lee, W.-S.: Statistical adjustment of decadal predictions in a changing climate, *Geophysical Research Letters*, 39, 2012.
- Lledó, L., Ramon, J., Soret, A., and Doblas-Reyes, F.-J.: Seasonal prediction of renewable energy generation in Europe based on four teleconnection indices, *Renewable Energy*, 186, 420–430, 2022.
- MacLeod, D., Torralba, V., Davis, M., and Doblas-Reyes, F.: Transforming climate model output to forecasts of wind power production: how much resolution is enough?, *Meteorological Applications*, 25, 1–10, 2018.
- 755 Manzanas, R., Gutiérrez Llorente, J. M., van Meijgaard, E., Calmanti, S., Magariño Manero, M. E., Cofiño González, A. S., Herrera García, S., et al.: Dynamical and statistical downscaling of seasonal temperature forecasts in Europe: Added value for user applications, 2018.
- Mavromatakis, F., Makrides, G., Georgiou, G., Pothrakis, A., Franghiadakis, Y., Drakakis, E., and Koudoumas, E.: Modeling the photovoltaic potential of a site, *Renewable energy*, 35, 1387–1390, 2010.
- 760 Meehl, G. A., Goddard, L., Murphy, J., Stouffer, R. J., Boer, G., Danabasoglu, G., Dixon, K., Giorgetta, M. A., Greene, A. M., Hawkins, E., et al.: Decadal prediction: can it be skillful?, *Bulletin of the American Meteorological Society*, 90, 1467–1486, 2009.
- Meng, Y., Schmidt, J., Zscheischler, J., and Bevacqua, E.: Climate-driven compounding effects and historical trends in renewable electricity droughts in Europe, *Applied Energy*, 401, 126 623, 2025.
- Miao, H., Xu, H., Huang, G., and Yang, K.: Evaluation and future projections of wind energy resources over the Northern Hemisphere in CMIP5 and CMIP6 models, *Renewable energy*, 211, 809–821, 2023.
- 765 Moemken, J., Reyers, M., Buldmann, B., and Pinto, J. G.: Decadal predictability of regional scale wind speed and wind energy potentials over Central Europe, *Tellus A: Dynamic Meteorology and Oceanography*, 68, 29 199, 2016.
- Monforti, F., Huld, T., Bódis, K., Vitali, L., D’isidoro, M., and Lacal-Arántegui, R.: Assessing complementarity of wind and solar resources for energy production in Italy. A Monte Carlo approach, *Renewable Energy*, 63, 576–586, 2014.
- 770 Moreno-Montes, S., Delgado-Torres, C., Duzenli, E., Pérez-Zanón, N., Marcos-Matamoros, R., and Soret, A.: Comparative analysis of statistical downscaling methods for multi-model decadal climate predictions over Western Europe, *Climate Services*, 42, 100 639, <https://doi.org/https://doi.org/10.1016/j.cliser.2026.100639>, 2026.

- Müller, J., Folini, D., Wild, M., and Pfenninger, S.: CMIP-5 models project photovoltaics are a no-regrets investment in Europe irrespective of climate change, *Energy*, 171, 135–148, 2019.
- 775 Olmo, M., Moreno-Montes, S., Trascasa-Castro, P., Balmaceda-Huarte, R., Delgado-Torres, C., Torralba, V., and Soret, A.: Disentangling the spread in wind projections over Europe and the impacts on energy generation., *Earth's future*, under review, 2026.
- O'Neill, B. C., Tebaldi, C., Van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, J., et al.: The scenario model intercomparison project (ScenarioMIP) for CMIP6, *Geoscientific Model Development*, 9, 3461–3482, 2016.
- Otero, N., Martius, O., Allen, S., Bloomfield, H., and Schaeffli, B.: A copula-based assessment of renewable energy droughts across Europe, 780 *Renewable Energy*, 201, 667–677, 2022.
- Panofsky, H. A. and Brier, G. W.: Some applications of statistics to meteorology, Mineral Industries Extension Services, College of Mineral Industries . . . , 1958.
- Paxian, A., Reinhardt, K., Pankatz, K., Pasternack, A., Lorza-Villegas, M. P., Scheibel, M., Hoff, A., Mannig, B., Lorenz, P., and Früh, B.: High-resolution decadal drought predictions for German water boards: a case study for the Wupper catchment, *Frontiers in Climate*, 4, 785 867 814, 2022.
- Payne, M. R., Danabasoglu, G., Keenlyside, N., Matei, D., Miesner, A. K., Yang, S., and Yeager, S. G.: Skilful decadal-scale prediction of fish habitat and distribution shifts, *Nature Communications*, 13, 2660, 2022.
- Pérez-Zanón, N., Ho, A.-C., Chou, C., Lledó, L., Marcos-Matamoros, R., Rifà, E., and González-Reviriego, N.: CSIndicators: Get tailored climate indicators for applications in your sector, 2023.
- 790 Priestley, M. D., Stephenson, D. B., Scaife, A. A., Bannister, D., Allen, C. J., and Wilkie, D.: Forced trends and internal variability in climate change projections of extreme European windstorm frequency and severity, *Quarterly Journal of the Royal Meteorological Society*, 150, 4933–4950, 2024.
- Pryor, S., Schoof, J., and Barthelmie, R.: Climate change impacts on wind speeds and wind energy density in northern Europe: empirical downscaling of multiple AOGCMs, *Climate Research*, 29, 183–198, 2005.
- 795 Pryor, S. C. and Barthelmie, R. J.: Climate change impacts on wind energy: A review, *Renewable and sustainable energy reviews*, 14, 430–437, 2010.
- Radziemska, E.: The effect of temperature on the power drop in crystalline silicon solar cells, *Renewable energy*, 28, 1–12, 2003.
- Ramon, J., Lledó, L., Torralba, V., Soret, A., and Doblas-Reyes, F. J.: What global reanalysis best represents near-surface winds?, *Quarterly Journal of the Royal Meteorological Society*, 145, 3236–3251, 2019.
- 800 Ramon, J., Lledó, L., Ferro, C. A., and Doblas-Reyes, F. J.: Uncertainties in the observational reference: Implications in skill assessment and model ranking of seasonal predictions, *Quarterly Journal of the Royal Meteorological Society*, 150, 897–910, 2024.
- Richardson, D., Pitman, A., and Ridder, N.: Climate influence on compound solar and wind droughts in Australia, *npj Climate and Atmospheric Science*, 6, 184, 2023.
- Scaife, A. A. and Smith, D.: A signal-to-noise paradox in climate science, *npj Climate and Atmospheric Science*, 1, 28, 2018.
- 805 Segado-Moreno, L. C., Ruiz-Arias, J. A., Montávez, J. P., and Betak, J.: Past, current and future solar radiation trends in Europe: Multi-source assessment of the role of clouds and aerosols, *Remote Sensing of Environment*, 333, 115 122, 2026.
- Slingo, J. and Palmer, T.: Uncertainty in weather and climate prediction, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369, 4751–4767, 2011.
- Smith, D., Eade, R., Scaife, A., Caron, L.-P., Danabasoglu, G., DelSole, T., Delworth, T., Doblas-Reyes, F., Dunstone, N., Hermanson, L., 810 et al.: Robust skill of decadal climate predictions, *Npj Climate and Atmospheric Science*, 2, 13, 2019.

- Smith, D. M., Scaife, A. A., and Kirtman, B. P.: What is the current state of scientific knowledge with regard to seasonal and decadal forecasting?, *Environmental Research Letters*, 7, 015 602, 2012.
- Smith, D. M., Scaife, A. A., Eade, R., Athanasiadis, P., Bellucci, A., Bethke, I., Bilbao, R., Borchert, L. F., Caron, L.-P., Counillon, F., et al.: North Atlantic climate far more predictable than models imply, *Nature*, 583, 796–800, 2020.
- 815 Solaraju-Murali, B., Bojovic, D., Gonzalez-Reviriego, N., Nicodemou, A., Terrado, M., Caron, L.-P., and Doblas-Reyes, F. J.: How decadal predictions entered the climate services arena: an example from the agriculture sector, *Climate services*, 27, 100 303, 2022.
- Solomon, S.: *Climate change 2007-the physical science basis: Working group I contribution to the fourth assessment report of the IPCC*, vol. 4, Cambridge university press, 2007.
- Soret, A., Gonzalez, P., Bloomfield, H., Masato, G., Goutham, N., Plougonven, R., and Torralba, V.: Subseasonal-to-seasonal climate predic-
820 tions for energy, in: *Sub-seasonal to Seasonal Prediction*, pp. 779–807, Elsevier, 2026.
- Storch, H. v. and Zwiers, F. W.: *Statistical Analysis in Climate Research*, Cambridge University Press, 1999.
- Suckling, E. B., van Oldenborgh, G. J., Eden, J. M., and Hawkins, E.: An empirical model for probabilistic decadal prediction: global attribution and regional hindcasts, *Climate Dynamics*, 48, 3115–3138, 2017.
- Šúri, M., Huld, T. A., Dunlop, E. D., and Ossenbrink, H. A.: Potential of solar electricity generation in the European Union member states
825 and candidate countries, *Solar energy*, 81, 1295–1305, 2007.
- Torralba, V., Doblas-Reyes, F. J., and Gonzalez-Reviriego, N.: Uncertainty in recent near-surface wind speed trends: a global reanalysis intercomparison, *Environmental Research Letters*, 12, 114 019, 2017a.
- Torralba, V., Doblas-Reyes, F. J., MacLeod, D., Christel, I., and Davis, M.: Seasonal climate prediction: a new source of information for the management of wind energy resources, *Journal of Applied Meteorology and Climatology*, 56, 1231–1247, 2017b.
- 830 Touma, J. S.: Dependence of the wind profile power law on stability for various locations, *Journal of the Air Pollution Control Association*, 27, 863–866, 1977.
- Trenberth, K. E., Fasullo, J. T., and Kiehl, J.: Earth’s global energy budget, *Bulletin of the american meteorological society*, 90, 311–324, 2009.
- Tsartsali, E., Athanasiadis, P., Materia, S., Bellucci, A., Nicolì, D., and Gualdi, S.: Predicting precipitation on the decadal timescale: A
835 prototype climate service for the hydropower sector, *Climate Services*, 32, 100 422, 2023.
- van Oldenborgh, G. J., Doblas-Reyes, F. J., Wouters, B., and Hazeleger, W.: Decadal prediction skill in a multi-model ensemble, *Climate dynamics*, 38, 1263–1280, 2012.
- Vaughan, C. and Dessai, S.: *Climate services for society: origins, institutional arrangements, and design elements for an evaluation framework*, Wiley Interdisciplinary Reviews: Climate Change, 5, 587–603, 2014.
- 840 Vautard, R., Cattiaux, J., Yiou, P., Thépaut, J.-N., and Ciais, P.: Northern Hemisphere atmospheric stilling partly attributed to an increase in surface roughness, *Nature geoscience*, 3, 756–761, 2010.
- Weide Luiz, E. and Fiedler, S.: Spatiotemporal observations of nocturnal low-level jets and impacts on wind power production, *Wind Energy Science*, 7, 1575–1591, 2022.
- Wilks, D. S.: Forecast verification, in: *International geophysics*, vol. 100, pp. 301–394, Elsevier, 2011.
- 845 Wiser, R., Bolinger, M., Hoen, B., Millstein, D., Rand, J., Barbose, G., Darghouth, N., Gorman, W., Jeong, S., O’Shaughnessy, E., et al.: *Land-based wind market report: 2023 edition*, Tellus A: Dynamic Meteorology and Oceanography, 2023.
- World Meteorological Organization: *WMO Global Annual to Decadal Climate Update, 2024–2028*, Tech. rep., World Meteorological Organization, Geneva, Switzerland, <https://wmo.int/publication-series/wmo-global-annual-decadal-climate-update-2024-2028>, 2024.

Zscheischler, J., Westra, S., Van Den Hurk, B. J., Seneviratne, S. I., Ward, P. J., Pitman, A., AghaKouchak, A., Bresch, D. N., Leonard, M.,
850 Wahl, T., et al.: Future climate risk from compound events, *Nature climate change*, 8, 469–477, 2018.