

Co-seismic infrasound in the ionosphere over Central Europe from the M8.8 Kamchatka 2025 earthquake observed by Doppler sounding at record heights

5 Jaroslav Chum¹, Zbyšek Mošna¹, Jiří Baše¹, Jan Zedník², Carsten Schmidt³, Patrick Hannawald³, Jan Ruzs¹, Jaroslav Urbár¹, Šimon Mackovjak⁴

¹Institute of Atmospheric Physics of the Czech Academy of Sciences, Prague, 141 00, Czech Republic

²Institute of Geophysics of the Czech Academy of Sciences, Prague, 141 00, Czech Republic

10 ³German Remote Sensing Data Center of the German Aerospace Center (DLR-DFD), Oberpfaffenhofen, 82234, Germany

⁴Institute of Experimental Physics, Slovak Academy of Sciences, Košice, 040 01, Slovakia

Correspondence to: Jaroslav Chum (jachu@ufa.cas.cz)

15 **Abstract**

Unique observations of co-seismic infrasound waves and disturbances in the ionosphere recorded by continuous Doppler sounding systems (CDSS) in Czechia and Slovakia during geomagnetically quiet period and associated with the Kamchatka M8.8 earthquake on July 29, 2025, are analysed and discussed. It is shown by simultaneous ionospheric sounding by a digisonde that the co-seismic infrasound waves were detected by the CDSS at a record height of about 340 km over Czechia, which is much higher than in previous observations, at distances greater than 8000 km from the epicentre. The Doppler shift oscillations caused by ionospheric plasma quasiperiodic movement induced by the infrasound waves had a frequency around 0.005 Hz and were observed approximately 12 min after the arrival of causative Long period surface seismic waves in Czechia. The frequency spectrum of the vertical ground surface motion that generated the infrasound waves was much broader, including more intense fluctuations with frequencies around 0.05 Hz. However, the higher frequency infrasound waves were attenuated during their propagation upward and did not reach the observation altitude, which is confirmed by numerical simulation that is in a good agreement with the CDSS observation. The numerical simulation also proves that it is necessary to consider air/plasma compression when calculating air particle velocities from the measured Doppler shift values. Combination of the numerical simulation with measurements using the CDSS and ionosonde is therefore a useful tool for better understanding and monitoring of co-seismic infrasound that propagates up to the altitudes of ionospheric F2 layer.

35

1 Introduction

On July 29, 2025, at 23:24:52 UT, an earthquake with a magnitude of 8.8 occurred east of the Kamchatka peninsula, about 120 km from the city of Petropavlovsk-Kamchatsky with the epicentre at 52.495°N 160.240°E and at the depth of about 35 km. The earthquake was the

40 result of reverse faulting on the subduction zone plate interface of the Kuril-Kamchatka arc and
the affected fault area was about 390×140 km (<https://earthquake.usgs.gov>). It was the third
largest earthquake in the 21th century after the 2004 Sumatra and the 2011 Great Tohoku
earthquake. The purpose of this paper is to investigate co-seismic ionospheric perturbations
associated with the Kamchatka earthquake in the ionosphere over Central Europe, about
45 8100 km away from the epicentre.

An investigation of atmospheric and ionospheric responses to earthquakes started already in
sixties of the 20th century by the analysis of data recorded during the 1964 Alaskan earthquake
(Bolt 1964; Donn and Posmentier 1964; Davies and Baker 1965). Various observation
techniques have been used to detect the co-seismic perturbations in the ionosphere. One of the
50 first and still used technique is continuous HF Doppler sounding, which allows obtaining time
series of the co-seismic perturbations at reflection heights of the HF signals (Davies and Baker,
1965; Artru et al., 2004; Chum et al., 2012). An observation technique probably most widely
used today are measurements of co-seismic changes in the total electron content (TEC) using
the dual-frequency receivers of Global Navigation Satellite Systems (GNSS) and analysis of
55 maps of TEC changes, which is possible due to a large number of GNSS receivers (Calais and
Minster, 1995; Heki and Ping, 2005; Otsuka et al., 2006). A basic instrument for ionospheric
research, the ionospheric sounder, is not frequently used to observe the co-seismic signatures
in the ionosphere due the relatively low time resolution. However, multiple cusp signatures
were observed in ionograms during suitable conditions, providing an important information
60 about the vertical extent of the perturbations (Maruyama and Shinagawa, 2014; Haralambous
et al., 2023). Multiple cusp signatures were also observed for the 29 July Kamchatka 2025
earthquake in the region from the western Pacific to the American west coast (Paul et al., 2026).
In addition, the ionospheric sounders provide important complementary measurements to
continuous HF Doppler sounding by providing an information about the reflection height of
65 Doppler sounding signals.

It was confirmed both theoretically and experimentally that vertical motion of the ground
surface, caused by seismic waves that propagate at supersonic speeds, generates infrasound
waves that propagate quasi-vertically to the upper atmosphere and ionosphere and cause the co-
seismic ionospheric disturbances owing to collisions between neutral particles and ionospheric
70 plasma, detectable by techniques mentioned above (Watada et al., 2006; Rolland et al., 2011;
Chum et al., 2012; Chum et al., 2016a,b; Haralambous et al., 2023). Recent reviews of the
detection and analysis of ionospheric signatures generated by earthquakes and other natural
hazards such as volcano eruption and tsunamis were given by Astafyeva (2019) and Wüst et al.
(2026).

75 The GNSS TEC observations can be used to create maps displaying the co-seismic
disturbances and their propagation projected into the horizontal plane at a height, in which a

dense ionospheric plasma occurs and/or at a height in which perturbations are assumed (Astafyeva and Heki, 2009; Liu et al., 2011; Zettergren and Snively, 2019). The GNSS TEC values and their changes represent integral values along the line of sight to the GNSS satellite and do not provide information about the phase and amplitude of waves in a specific/known altitude. However, it is possible to do a three-dimensional reconstruction (tomography) of ionospheric perturbations using a dense network of GNSS receivers (Song et al., 2025), nevertheless, the vertical and time resolution needed for a successful reconstruction is relatively low, about 30 km and 2 min, respectively.

On the other hand, continuous HF Doppler sounding operating close to an ionospheric sounder can provide the waveform of the co-seismic perturbation, infrasound wave, at a specific altitude (given by the sounding frequency and determined from ionograms) with a high time resolution (up to several seconds). So far, the co-seismic infrasound waves were observed using Doppler sounding at altitudes around 200 km or lower in Central Europe (Chum et al., 2012; Chum et al., 2016a; Haralambous et al., 2023) and, to the best of our knowledge, also in the world. In this paper we present an observation of co-seismic infrasound by Doppler sounding in record breaking height of about 340 km. It is also shown that only waves with longer periods were able to reach such altitudes; the shorter periods were attenuated below. The observation is consistent with numerical simulation of infrasound propagation.

2 Data and methods

The Czech multipoint Continuous Doppler Sounding System (CDSS) operates at frequencies of 3.59, 4.65 and 7.04 MHz (Laštovička and Chum, 2017). Its main application is monitoring and analysis of medium-scale ionospheric travelling disturbances associated with atmospheric gravity waves (Chum et al., 2021), but due to its high time resolution (several seconds) it can also be used to detect and analyse co-seismic infrasound and transient phenomena such as sudden frequency deviations caused by X-ray and extreme ultraviolet radiation produced during solar flares and sudden storm commencements (Chum et al., 2016a,b; Chum et al., 2018a,b, Kikuchi et al., 2021, 2022). There are three transmitting sites (Tx), labelled PVC, PRU and DLO in Figure 1, which operate at all three frequencies in Czechia. The receiver (Rx) used in this study is located in Prague (50.04°N, 14.48°E), about 8150 km away from the epicentre and about 7 km away from the digisonde DPS-4D in Pruhonice, PQ052 (Reinisch et al., 2005), which is about 200 m away from the PRU Tx site. To exclude a local effect, data from CDSS (operating at 3.59 MHz) installed in Eastern Slovakia with receiver in Kolonica (48.93°N, 22.27°E), about 8060 km away from the epicentre and marked KOL in Figure 1, are also investigated. The Slovak transmitters are labelled KOS, ROZ and ZAH.




Fig. 1 Distribution of CDSS transmitters (Tx, blue dots) and receivers (Rx, red dots) in Czechia and Slovakia. Digisonde DPS-4D is collocated with the Tx in Pruhonice (PRU), and the seismometer is collocated with the Tx in Panská Ves (PVC) in Czechia.

Reflection points of the Doppler signals, if projected to the ground, are assumed in the midpoints between each transmitter and receiver (Figure 1). The half distances between the transmitters and receiver (few tens of km) are much smaller than the reflection heights (more than 300 km in this case). So, the sounding is quasi-vertical, which means that azimuthal angles α of sounding radio waves are small, $\cos(\alpha) \approx 1$, and the sounding radio waves (ordinary mode) reflect at heights where their frequency matches the local plasma frequency. If the ionospheric plasma in the reflection region moves (radially relative to the CDSS) or if the electron density changes, then the reflected radio wave experiences a Doppler shift. Assuming a sharp electron density gradient, the radial/vertical plasma velocity w_p can be computed from the obtained Doppler shift f_D as

$$w_p = -f_D \cdot \frac{c}{2f_0} \quad , \quad (1)$$

where c is the speed of light and f_0 is the frequency of the sounding radio wave (Davies et al., 1962; Jacobs and Watanabe, 1966).

In the ionospheric F region, plasma is fully magnetized and moves freely only along magnetic field lines. Therefore, the vertical component of plasma motion w_p induced by vertically moving air particles with velocity w due to the infrasound waves is given by

$$w_p = w \cdot \sin^2(I) \quad , \quad (2)$$

where I is the inclination of the magnetic field ($\sim 66^\circ$ in Czechia).

Because the plasma frequency and refractive index of the radio waves depend on the electron density, the Doppler shift is related to the electron density changes, especially at the region of reflection. Several authors (e.g., Sutcliffe and Poole, 1989 and references therein) used the

equation of continuity and argued that alternating plasma compression caused by compressional
 140 magnetohydrodynamic waves should also contribute to the Doppler shift. However, the
 experimental results were inconclusive in the case of magnetohydrodynamic waves. Chum et
 al. (2012, 2016a) applied their approach and using the equation of continuity derived equation
 (3) which relates the observed Doppler shift f_D with the air particle vertical velocity w induced
 by vertically propagating infrasound waves.

$$145 \quad w = -f_D \cdot \frac{c}{2f_0 \sin^2(I)} \cdot \frac{\frac{\partial N}{\partial z}}{\sqrt{(\frac{\partial N}{\partial z})^2 + (N \frac{2\pi f_{IS}}{c_s})^2}}, \quad (3)$$

where f_{IS} is the infrasound frequency, c_s is the sound speed at the reflection height, N is the
 electron density and $\partial N/\partial z$ is the vertical gradient of the electron density at the reflection height
 estimated from the true height profile of the plasma frequency f_p obtained from the ionogram,
 e.g., by SAO Explorer software (Reinisch et al. 2005). A definition of plasma frequency f_p is
 150 used to relate f_p and N . The term $N \cdot (2\pi f_{IS})/c_s$ in the denominator of equation (3) results from
 the air/plasma compression due to the infrasound waves. See Chum et al., (2016a) for more
 details and derivation. Note that if $(\partial N/\partial z)^2 \gg (N \cdot (2\pi f_{IS})/c_s)^2$, then equation (3) reduces to
 equation (1) with usage of equation (2). Chum et al. (2012, 2016a) also showed that not
 considering the compressional term $N \cdot (2\pi f_{IS})/c_s$ can lead to results that are inconsistent with
 155 energy conservation for specific cases of the co-seismic infrasound observations.

The near surface air particle vertical oscillation velocities w_0 are determined by the vertical
 velocity of Earth surface motion, v_z , (Watada et al., 2006), which is measured by a seismometer
 co-located with PVC transmitter in the Panská Ves observatory (50.53° N, 14.57° E). The
 infrasound waves generated by seismic waves propagating far away from the epicentre
 160 propagate nearly vertically owing to the supersonic speed of seismic waves. Specifically, the
 expected zenith angle δ of the generated infrasound waves above the surface is given by
 equation (4)

$$\sin(\delta) = \frac{c_{S0}}{c_G}, \quad (4)$$

where c_{S0} is the sound speed above the ground and c_G is the speed of seismic waves (Rolland et
 165 al., 2011; Chum et al., 2016a). As c_G is about 10 times larger than c_{S0} , the deviation of initial
 infrasound wave vector from the zenith direction is around 6°.

The observed time delays between the ground surface velocities and co-seismic infrasound
 waves recorded by the CDSS are also compared with ray tracing simulation, including
 attenuation along the ray tracing trajectories due to the viscosity and thermal conductivity. The
 air temperature and neutral species densities at specific locations and time needed for sound
 170 speed calculation and attenuation estimate are obtained from the NRLMSIS model (Emmert *et al.*
 2020) with a possibility to include also neutral winds using the horizontal wind model
 HWM14 (Drob *et al.* 2015). A more detailed description of the ray tracing software features

175 can be found in Chum et al. (2023) and references therein. In addition, one-dimensional full-
wave simulation of vertical infrasound propagation based on the numerical solution of the
equation of continuity, the equation of motion/momentum (Navier-Stokes equation) and the
heat equation using an implicit finite difference method, including non-linear terms, is
performed to investigate the expected evolution of the wave spectrum with height and to
180 estimate the expected air particle velocities at specific heights (see Chum et al., 2016b, 2018a
for a more detailed description of the simulation method).

We also check other available measurements that could provide information on atmospheric
waves at specific altitudes, such as airglow measurements of the hydroxyl (OH) layer in the
mesopause region around 86 km using the OH temperature measurements by the Ground-based
Infrared P-branch Spectrometer (GRIPS) and wave detection by the Fast Airglow Imager
185 (FAIM) installed in Panská Ves (Schmidt et al., 2013; Hannawald et al., 2016). Both airglow
instruments were setup to observe the OH layer directly below the reflection point of the CDSS
(the green point between PVC and PRG in Figure 1).

The indices of geomagnetic activity, searched to verify the geomagnetic activity, are taken
from the World Data Center for Geomagnetism, Kyoto.

190

3 Results

Despite the maximum of solar cycle 25, the M8.8 Kamchatka earthquake occurred during a
period when ionosphere was not significantly disturbed by space weather drivers. There were
no M or X-class flares and neither strong geoeffective coronal mass ejections nor high-speed
195 streams of solar wind plasma. Therefore, the geomagnetic activity was very low, with the
following values of geomagnetic indices: $Dst \sim 12$ nT and $Kp \sim 2$. The conditions were therefore
favourable to study the ionospheric response to acoustic-gravity waves of known origin and
amplitude propagating from below. As the space weather influence was minimal (Mackovjak
et al., 2026), the interpretation of the origin of the detected ionospheric variations is relatively
200 straightforward.

Figure 2a shows the vertical velocity v_z of ground surface motion measured in Panská Ves,
Czechia, in the time interval of 70 min from the onset of the earthquake on July 29, 2025, at
23:24:52 UT. A detection of the Primary body (P) waves in Panská Ves started after 23:36 UT
and continued until about 23:40 UT, with relatively small amplitudes in v_z (less than 0.1 mm/s).
205 Secondary shear body waves once reflected from the Earth's surface (SS) were detected in the
vertical component after 23:50 UT and had similar amplitudes as the P waves. The surface
waves reached much larger amplitudes and were observed on July 30, 2025, approximately
between 00:00 and 00:20 UT. Specifically, Long period surface (L) waves with amplitudes
reaching up to around 0.4 mm/s were observed approximately between 00:00 and 00:10 UT,

210 and Rayleigh (R) waves with shorter periods and amplitude exceeding 1 mm/s were recorded between 00:10 and 00:20 UT.


Following the activity of seismic waves, fluctuations in Doppler shifts f_D with a frequency about 0.005 Hz (period about 200 s) were observed in the CDSS records around 00:20 UT at sounding frequencies 4.65 and 3.59 MHz in Czechia (sounding frequency 7.04 MHz was above
215 the maximum plasma frequency in the ionosphere, the critical frequency, so the 7.04 MHz signals did not reflect from the ionosphere). There were no time delays (within the time resolution) between the infrasound signals on individual sounding paths (Tx-Rx pairs), so the signals from all three transmitters were averaged at each time point. The values of the observed Doppler shifts were converted to the vertical plasma velocities w_p and vertical air particle
220 velocities w using equations (1) and (3), respectively, and after filtering out longer period fluctuations of different origin (larger than about 5 min) displayed in Figure 2b (4.65 MHz) and Figure 2c (3.59 MHz). The w_p and w velocities obtained from the Czech CDSS are by blue and red solid, respectively. Note different scales on the axes for the w_p (left axis) and w (right axis) velocities. The w_p velocity measured by the CDSS in Slovakia is shown by blue dashed line in
225 Figure 2c (the w is not calculated because of the absence of a nearby ionosonde that could provide the $\partial N/\partial z$ gradient). The signal (dominant wave packet) observed by the Czech CDSS at 3.59 MHz precedes by about 40 seconds the signal recorded at 4.65 MHz. The w_p amplitudes (equation 1) reached about 9 and 15 m/s at 3.59 and 4.65 MHz, respectively, and the w amplitudes (equation 3) were about 5.5 and 4.8 m/s at 3.59 and 4.65 MHz. The w_p amplitudes
230 were larger at 4.65 MHz than at 3.59 MHz, whereas the w amplitudes were higher for 3.59 MHz. The reason for this is a difference in the values of electron density gradient, $\partial N/\partial z$, at reflection heights for 3.59 and 4.65 MHz waves, as will be shown later.

It is reminded that application of equation (3) also requires a knowledge of a dominant frequency at the reflection height to calculate w . An insight in a spectral content of the observed
235 waves and their evolution with time is shown in dynamic spectra displayed in Figure 3. It is important to note that the seismic wave packet labelled L (Figure 2a and Figure 3a) contains a very broad frequency spectrum, including frequencies around 0.005 Hz, which are detected also in the Doppler shift records (Figures 3b and 3c), approximately 11-13 min later. The frequency of 0.005 Hz was used to calculate w shown in Figure 2b and 2c.

240 The reflection heights of the ordinary mode of the sounding radio waves with frequencies 3.59 and 4.65 MHz are about 305 and 340 km, respectively, as shows the height profile of plasma frequency (true height profile) obtained from the ionogram recorded in Pruhonice on July 30 at 00:25 UT, which is displayed in Figure 4. The average reflection heights for the 3.59 and 4.65 MHz waves in the time interval from 00:10 to 00:30 UT obtained from a sequence of
245 ionograms are 303 and 336 km, respectively. Figure 4 also shows the estimated inverse of the

plasma density gradients at the reflection heights, from which the electron density gradient $\partial N/\partial z$ is obtained using the definition of plasma frequency f_p . The obtained values are $\partial N/\partial z = 3.56 \cdot 10^6$ and $2.75 \cdot 10^6 \text{ m}^{-4}$ at the reflection heights of 3.59 and 4.65 MHz waves, respectively. These values were used to calculate vertical air particle velocities w plotted in Figures 2b and 2c.


To verify that the observed time delays correspond to the propagation of infrasound waves, which are a probable cause of the ionospheric disturbances, we performed a ray tracing simulation. Figure 5a shows the simulated propagation time (orange) and sound speed (black) of the infrasound waves launched with the zenith angle 6° from the Panská Ves location on July 30, 2025 at 00:06 UT. The propagation times to altitudes 305 and 340 km are 680 and 724 s, respectively. The sound speeds at these altitudes are 807 and 837 m/s, respectively. The modelled propagation times are consistent with the observed time delays between the 0.005 Hz signal recorded on the ground by the seismometer and in the ionosphere by the CDSS at about 305 and 340 km altitudes.



260

Fig. 2 a) Vertical velocity v_z of the ground surface motion in Panská Ves on the night of July 29 to 30, 2025 b) and c) Vertical velocity of plasma motion w_p (blue solid) directly obtained from the observed Doppler shift by equation (1) and the calculated vertical velocity of air particle oscillation w (red) obtained using equation (3) for measurements at $f=4.65$ and 3.59 MHz, respectively. Black dashed lines show the simulated vertical air particle oscillation velocity at altitudes 340 and 305 km, respectively, scaled to the right axis. The blue dashed line in the plot c) displays the w_p velocity obtained in Slovakia.

265



270 **Fig. 3** Dynamic spectra (power spectral densities) of vertical velocity v_z of the ground surface motion on the night of July 29 to 30, 2025 (a), and of the Doppler shifts measured at $f=4.65$ MHz (b) and 3.59 MHz (c). The dynamic spectra correspond to signals presented in Figure 2.

Dashed lines in Figure 5b show the ratios w/w_0 , which represent the expected/simulated vertical air particle velocities w at a specific altitude normalized by their near surface value w_0 , including the frequency dependent attenuation along the ray trajectories for infrasound frequencies 0.05 (red) and 0.005 (orange) Hz. The solid line displays the expected ratio for an unrealistic lossless (infra)sound plane wave propagation for which the infrasound power flux is conserved. It means that the quantity $c_s \rho w^2$ is constant, where ρ is the air density, decreasing exponentially with height (obtained from the NRLMSIS model). Figure 5b shows that waves with a frequency of 0.05 Hz begin to be significantly attenuated above an altitude of approximately 150 km. On the other hand, the 0.005 Hz waves can reach the height of 300 km without a significant attenuation. The ratio w/w_0 (normalized velocity w) for 0.005 Hz waves at the observation heights 305 and 340 km is $1.26 \cdot 10^5$ and $1.19 \cdot 10^5$, respectively. For idealized lossless propagation it is $1.89 \cdot 10^5$ and $2.74 \cdot 10^5$, respectively, at these heights.

285




Fig. 4 True height profile of plasma frequency derived from the measurements by the digisonde located in Průhonice, PQ052, on July 30, 2025 at 00:25 UT. The CDSS sounding frequencies 3.59 and 4.65 MHz and the vertical plasma density gradients are marked by red.

290




Fig. 5 a) Simulated infrasound propagation time (orange) and sound speed (black) as a function of height. b) Simulated vertical air particle oscillation velocities w normalized to the initial value w_0 above the ground surface for frequencies 0.05 Hz (red) and 0.005 Hz (orange).


295 The solid orange line represents an idealized case of lossless propagation of a plane wave.

It is important to stress that a comparison of the observed values w (w_p) displayed in Figure 2 with the simulated values w shown in Figure 5b is not straightforward. The reason is that the calculated normalized values w/w_0 correspond to monochromatic waves. The wave attenuation with height strongly depends on the wave frequency. The higher frequency, the lower is the altitude at which the wave energy dissipates (Figure 5b). Figure 3a shows that the seismic wave packet L contained a very wide range of frequencies and therefore generated infrasound in a broad frequency spectrum. However, only the waves corresponding to frequencies around 0.005 Hz - a small energy fraction could reach the heights above 300 km, where the induced plasma fluctuations were detected by the CDSS. Therefore, for the wave packet L it makes no sense to compare the observed amplitudes w (w_p) normalized by the v_z amplitude (Figure 2) with the simulated ratio w/w_0 (Figure 5b).

To compare the observed amplitudes obtained from the CDSS measurements with the expected values, we performed full wave one-dimensional numerical simulation of infrasound propagation in the vertical direction, started with initial perturbations above the Earth's surface defined by the vertical motion of the ground surface, $w_0=v_z$. The assumption of this initial perturbation is based on theoretical work (Watada et al., 2006) and was successfully applied in previous studies (Chum et al., 2018a and references therein). The w values obtained by the full-wave numerical simulation at the heights 305 and 340 km are plotted in Figure 2c and 2b, respectively, by the dashed black lines. The values are scaled to the right (w) axis and the maximum amplitudes are about 6.8 m/s at 305 km and 6.2 at 340 km. It means that they are about 30% larger than the w values estimated from the measurement. The main fluctuations of the simulated waveforms occur at approximately similar times as the w (w_p) values derived from the observed Doppler shifts and have also similar frequencies. The simulated signals only

320 slightly advance the observation. Therefore, the simulation agrees reasonably well with the
observation. It should be noted in this respect, that the real trajectory of infrasound waves will
be somewhat longer than the vertical distance due to the non-zero zenith angle of propagation
and due to the neutral winds, which may affect the trajectory. Consequently, the location of the
seismometer in Panská Ves (PVC in Figure 1) used for the simulation may not be the exact
325 source location of the waves that reach the observation points.


Figure 6 shows the height evolution of the air particle velocity fluctuation spectra obtained
by the one-dimensional full-wave numerical simulation started with the fluctuations defined by
the seismic record, $w_0=v_z$. The power spectral densities of w are normalized to their maximum
value above the ground surface and their colour-coded common logarithm is displayed in Figure
330 6. The height-dependent spectra in Figure 6 confirm that only waves with the lowest frequencies
(about 0.005 Hz) can reach the observation altitudes, although they carry just a small fraction
of the total wave energy at altitudes lower than about 200 km.



335 **Fig. 6** Expected power spectral densities (colour coded common logarithm) at different
altitudes, normalized to the ground level.

We also investigated measurements that could provide information about wave propagation
at lower heights, namely OH airglow emissions originating from an about 10 km thick layer
340 around 86 km, which may be modulated by co-seismic atmospheric waves (Snively, 2013; Wüst

et al., 2023). Figure 7 shows that no significant changes in the OH temperature related to the infrasound waves were detected over Czechia, despite intensity drops in the used OH(3-1)- and OH(4-2)-Q-branches, approximately in the interval from 00:05 to 00:35 UT, which roughly coincides with the occurrence of the co-seismic infrasound waves. On the one hand the ratio of these two emissions serves as a cloud indicator (OH(3-1)-Q at 1.505 μm is more strongly attenuated by water and ice droplets compared to OH(4-2)-Q at 1.585 μm). On the other hand, both emissions originate at slightly different centroid altitudes, separated by 0.5-1.0 km (e.g., von Savigny et al., 2012). Atmospheric waves with short vertical wavelengths ($\ll 100$ km) passing through the OH layer will therefore leave a clear signature in the OH(3-1)/OH(4-2)-ratio, while waves with large vertical wavelengths (affecting both emissions at the same time) will not (Schmidt et al., 2018). This drop in intensity without a significant change in the OH(3-1)/OH(4-2)-ratio indicates the presence of an atmospheric structure with large vertical scales, which is in contrast with the expected wavelength of the co-seismic infrasound (about 6 km for the dominant 0.05 Hz waves at this altitude and about 60 km for the 0.005 Hz waves). Model results by Inchin et al. (2020, 2022) predicted a drop in intensity by about 15%, but only close to the epicenter. Data from the OH imager (FAIM) show the presence of a varying number of small clouds in the fields of view of both instruments at this time. A similar drop in OH intensities were not observed in other European stations equipped with the OH imagers and spectrometers in this time interval. Thus, the co-seismic origin of the observed intensity drop remains questionable. The time resolution of OH temperature measurement by GRIPS is 15 s. This sampling frequency does not allow for the analysis/detection of waves with a dominant frequency of approximately 0.05 Hz, which is expected at the height of the OH layer (Figure 6). The waves with frequency around 0.005 Hz were not detected in the OH temperature time series at the expected time (around 00:10 UT on July 29, 2025) using the continuous Wavelet Transform (not shown). The reason may be an insufficient signal-to-noise ratio (SNR), as the model predictions by Inchin et al. (2020, 2022) show that co-seismic airglow perturbations range between 1-5% and are weaker in temperature than in intensity requiring an SNR that wasn't reached during this particular night (Fig. 7).



370 **Fig. 7** OH temperature (blue) and OH(4-2) radiance (solid red) on the night of July 29 to 30,
 2025. The nearly constant ratio (dashed red) between OH(3-1)- and OH(4-2)-emissions
 indicates a structure with large vertical wavelengths/scales being associated with the drop in
 intensity. See text for further details.

375 **4 Discussion and Conclusions**

The simulated air particle vertical velocities w shown in Figure 2 by black dashed lines are
 about 30% larger than the values obtained from observation using equation (3). This is a
 reasonable agreement considering the uncertainties in determination of $\partial N/\partial z$ gradients (around
 10%). Importantly, it is reasonable to expect that the observed w are smaller, than the simulated
 values for several reasons. First, the simulation is based on the plane wave propagation (one-
 dimensional simulation) and does not take into account a possible divergence of infrasound ray
 trajectories, and therefore does not consider a decrease of the infrasound power flux due to the
 geometrical spreading of the waves. Second, the simulation only considers attenuation due to
 the viscosity and thermal conductivity (Chum et al., 2016b), but does not simulate other
 possible losses, for example, rotational relaxation losses, turbulent losses etc. (Bass et al., 1984).
 Third, the actual trajectory of infrasound waves may be longer than the pure vertical distance,
 and consequently the attenuation larger, due to the non-zero zenith angle and due to the neutral
 winds. It is useful to stress that both the plasma velocities w_p and the air particle velocities
 obtained directly from w_p by using equation (2), without including the compressional term, give

390 values roughly 100% larger than the simulated values, which is highly improbable for the reasons mentioned above, and indicates the need to use the compressional term in the quantitative analysis of infrasound amplitudes using Doppler shift measurements.

The observed time delay of about 12 min between the 0.005 Hz waves in the seismic record and the waves of the same frequency in the Doppler shift (air particle velocity fluctuations) is
395 consistent with the numerical simulation of infrasound propagating in the (quasi)vertical direction. In addition, the 3.59 MHz CDSS in Czechia detects the co-seismic signal approximately 40 s sooner than the 4.65 MHz CDSS, the signal of which is reflected at the altitude about 35 km higher.

The Slovak 3.59 MHz CDSS detects the co-seismic infrasound waves at a similar time as
400 the Czech CDSS, with a phase shift locating the main fluctuations roughly between the 3.59 and 4.65 MHz signals in Czechia (Figure 2c). This indicates that the reflection height for the 3.59 MHz radio waves is somewhat larger in Slovakia than in Czechia, considering (quasi)vertical propagation of infrasound and similar horizontal distances from the epicentre (about 8150 km the Czech CDSS and about 8060 km the Slovak CDSS). The vertical plasma
405 velocities w_p obtained without considering the compressional term (equation 1) are about 100% higher in Slovakia than in Czechia. It is reminded that there is no nearby ionosonde in Slovakia that would provide reliable $\partial N/\partial z$ values for calculating the more realistic velocities w using equation (3). However, if we assume that the air particle velocities w are approximately the same over Czechia and Slovakia, then it follows from equation (3) that the electron density
410 gradient $\partial N/\partial z$ at the corresponding reflection heights of 3.59 MHz signals must be smaller in Slovakia, than in Czechia. This is consistent with the higher reflection height in Slovakia deduced from the timing mentioned above (note that $\partial N/\partial z$ usually decreases with height in the F2 layer, when approaching the peak of the layer).

Unlike the previous observation of co-seismic infrasound waves by CDSS in Czechia,
415 triggered by strong earthquakes, namely M9.0 Tohoku 2011, M7.8 Nepal 2015, and M7.7 Turkey 2023 earthquakes, when the shape and spectral content of the wave packets in the co-seismic Doppler shift fluctuations was very similar to that recorded in the vertical velocity v_z of ground surface motion (Chum et al., 2012; Chum et al., 2016a; Haralambous et al., 2023), the spectral content of the detected co-seismic ionospheric waves associated with the M8.8
420 Kamchatka earthquake differed significantly from v_z . The main reason for that is the high altitude of observation and filtering out of waves with higher frequencies during the infrasound propagation to such a high altitude, including waves with frequencies that are dominant at lower altitudes. This is a novel observation, which contributes to a better understanding of propagation of the co-seismic infrasound to high altitudes, approximately up to the peak of the ionosphere.

425 It should be stressed that the change of the wave packet shape and absence of waves with
higher frequencies at the CDSS observation height in this case is not caused by nonlinear effects,
which are a consequence of large amplitude of infrasound waves in the upper atmosphere that
may occur for sufficiently strong initial perturbations $w_0=v_z$ (Chum et al., 2016b; Chum et al.,
2018a), but purely by filtering out of waves with higher frequencies due to the wave attenuation,
430 which strongly depends on wave frequency. Unlike the case of nonlinear effects (Chum et al.,
2016b), a nonlinear simulation started with an initial perturbation $w_0=v_z/10$ provides in this case
practically the same wave packet (spectral content) at the observation altitude, only with ~ 0.1
amplitude, when compared with the simulation started for $w_0=v_z$.

The co-seismic infrasound waves have never been observed at ~ 340 km height by the CDSS
435 in Czechia before. All the previous Czech observations of co-seismic infrasound were from an
altitude range approximately 150-220 km. To the best of our knowledge, we are not aware of
any Doppler observation of co-seismic infrasound in such a height in other countries. It is
reminded that although TEC perturbations detected by GNSS receivers are usually mapped to
the peak of F2 layer, they are usually observed with a 8-9 min delay with respect to the seismic
440 record (Astafyeva et al., 2019; Wüst et al., 2026 and references therein), which indicates
dominant perturbations at ~ 250 km or below when considering the expected sound speed height
profile (e.g., Figure 5a). There is actually no direct information about the altitude of the TEC
disturbance due to the height integrated nature of the TEC GNSS measurements. In addition,
we have not identified any TEC perturbations over the Central Europe that could be reliably
445 associated with the co-seismic infrasound. On the other hand, ionospheric disturbances
associated with tsunami waves were with a high probability detected at the altitudes above 400
km by Low Earth Orbit satellites (Yang et al., 2022; Alfonsi et al., 2024), but these perturbations
are related to the gravity waves, rather than to infrasound. Therefore, the detection of infrasound
waves in the ionosphere by the CDSS at the height of around 340 km and about 12 min after
450 the arrival of causative Long period surface seismic waves is a unique observation, which
experimentally proves that low frequency infrasound waves can reach such a high altitude. It
should also be remembered that the detailed analysis of the co-seismic infrasound observed by
the CDSS was possible due to the operation of a nearby digisonde which provided the reflection
heights of the CDSS signals and electron density gradients there.

455

Appendix A: List of abbreviations

CDSS continuous Doppler sounding system

GNSS global navigation satellite system

OH hydroxyl

460 Rx receiver

TEC total electron content

Tx transmitter
UT universal time

465 **Data availability**

The CDSS data are available at the <http://datacenter.ufa.cas.cz/> (use link to spectrogram archive). Specifically, time series of Doppler shifts related to Kamchatka earthquake which are used in this study (including detailed Doppler shift spectrograms), can be obtained using the link http://datacenter.ufa.cas.cz/publicdata/archive/three_paths/ (select the corresponding frequency and date).

470

The digisonde data can be found at <https://giro.uml.edu/ionoweb/> (select PQ052 station). The seismic data are available, e.g., at <https://www.fdsn.org/networks/detail/CZ/>. The OH airglow data can be obtained at <https://zenodo.org/records/18788838>.

475 **Authors contribution**

JC designed the article, wrote the texts and performed most of the analysis. ZM analysed the ionograms, JB engineered and maintained the CDSS, JZ provided the seismic data and their analysis, CS was responsible for the OH temperature measurements, PH was responsible for the OH imager data, JR helped with Figures, JU checked the GNSS data, and SM was responsible for the CDSS in Slovakia. All authors read and approved the submitted version.

480

Competing Interest

The authors declare that they have no conflict of interest.

485 **Acknowledgement**

Community Coordinated Modeling Center at Goddard Space Flight Center and their publicly available simulation services (<https://ccmc.gsfc.nasa.gov>) are acknowledged for the NRLMSIS model used in our ray tracing code and full wave numerical simulation. The World Data Center for Geomagnetism, Kyoto (<https://wdc.kugi.kyoto-u.ac.jp/>) is acknowledged for the geomagnetic indices.

490

Funding

The support by the ESA project Swarm QUID-REGIS ESA, contract No. 4000143632/24/I-EB is acknowledged.

495

References

- Alfonsi, L., Cesaroni, C., Hernandez - Pajares, M., Astafyeva, E., Bufferal, S., Elias, P., et al.: Ionospheric response to the 2020 Samos earthquake and tsunami, *Earth Planet. Space*, 76(1), 13, <https://doi.org/10.1186/s40623-023-01940-2>, 2024.
- 500 Artru, J., Farges, T., and Lognonné, P.: Acoustic waves generated from seismic surface waves: propagation properties determined from Doppler sounding observations and normal-mode modeling, *Geophys J. Int.*, 158:1067–1077, 2004.
- Astafyeva, E.: Ionospheric detection of natural hazards, *Reviews of Geophysics* 57(4), 1265-1288, <https://doi.org/10.1029/2019RG000668>, 2019.
- 505 Astafyeva, E. and Heki, K.: Dependence of waveform of near-field co-seismic ionospheric disturbances on focal mechanisms, *Earth Planet. Space*, 61(7), 939-943. <https://doi.org/10.1186/bf03353206>, 2009.
- Bass, H., Sutherland, ELC., Piercy, J., Evans, L.: Absorption of sound by the atmosphere, in *Physical Acoustics*, Vol. XVII, Chap. 3., edited by W. P. Mason and R. N. Thurston, pp.145-232, Academic Press, Inc, 1984.
- 510 Bolt, BA.: Seismic air waves from the great 1964 Alaskan earthquake. *Nature* 202:1095-1096. doi:10.1038/2021095a0, 1964.
- Calais, E. and Minster, J. B.: GPS detection of ionospheric perturbations following the January 17, 1994, Northridge earthquake, *Geophys. Res. Lett.*, 22, 1045–1048, doi:10.1029/95GL00168, 1995.
- 515 Chum, J., Hruska, F., Zednik, J., and Lastovicka, J.: Ionospheric disturbances (infrasound waves) over the Czech Republic excited by the 2011 Tohoku earthquake, *J. Geophys. Res.*, 117, A08319, doi:10.1029/2012JA017767, 2012
- Chum, J., Liu, J.Y., Laštovička, J., Fišer, J., Mošna, Z., Baše, J., and Sun, Y.Y.: Ionospheric signatures of the April 25, 2015 Nepal earthquake and the relative role of compression and advection for Doppler sounding of infrasound in the ionosphere, *Earth Planet. Space*, 68:24, DOI 10.1186/s40623-016-0401-9, 2016a
- 520 Chum, J., Cabrera, M. A., Mošna, Z., Fagre, M., Baše, J., and Fišer, J.: Nonlinear acoustic waves in the viscous thermosphere and ionosphere above earthquake, *J. Geophys. Res. Space Physics*, 121, doi:10.1002/2016JA023450, 2016b
- 525 Chum, J., Liu, J.Y., Podolská, K., and Šindelářová, T.: Infrasound in the ionosphere from earthquakes and typhoons, *J. Atmos. Sol. Terr. Phys.*, 171, 72-82, doi:/10.1016/j.jastp.2017.07.022, 2018a
- Chum, J., Urbář, J., Laštovička, J., Cabrera, M. A., Liu, J.Y., Bonomi, F., Fagre, M., Fišer, J., and Mošna, Z.: Continuous Doppler sounding of the ionosphere during solar flares, *Earth, Planet. Space*, 2018, 70: 198, <https://doi.org/10.1186/s40623-018-0976-4>, 2018b
- 530

- Chum, J., Podolská, K., Rusz, J., Baše, J., and Tedoradze, N.: Statistical investigation of gravity wave characteristics in the ionosphere, *Earth Planet. Space*, **73**, 60, <https://doi.org/10.1186/s40623-021-01379-3>, 2021.
- 535 Chum, J., Šindelářová, T., Koucká Knížová, P., Podolská, K., Rusz, J., Baše, J. et al.: Atmospheric and ionospheric waves induced by the Hunga eruption on 15 January 2022; Doppler sounding and infrasound, *Geophys. J. Int.*, **233**, 1429–1443, <https://doi.org/10.1093/gji/ggac517>, 2023.
- Davies, K., Watts, J., and Zacharisen, D.: A study of F2-layer effects as observed with a
540 Doppler technique, *J. Geophys. Res.*, **67**:2, doi:10.1029/JZ067i002p00601, 1962.
- Davies, K. and Baker, D.M.: Ionospheric effects observed around the time of the Alaskan earthquake of March 28, 1964, *J. Geophys. Res.*, **70**(9), 2251–2253, doi:10.1029/JZ070i009p02251, 1965.
- Donn, W.L., and Posmentier, E.S.: Ground-coupled air waves from the great Alaskan
545 earthquake. *J. Geophys. Res.*, **69**:5357–536, 1964.
- Drob, D. P., et al.: An update to the Horizontal Wind Model (HWM): The quiet time thermosphere, *Earth Space Sci.*, **2**, 301–319, doi:10.1002/2014EA000089, 2015.
- Emmert, J.T., Drob, D.P., Picone, J.M., Siskind, D.E., Jones, M., Jr., Mlynchak, M.G. et al.:
550 NRLMSIS 2.0: a whole-atmosphere empirical model of temperature and neutral species densities, *Earth Space Sci.*, **8**, e2020EA001321, doi:10.1029/2020EA001321, 2020.
- Hannawald, P., Schmidt, C., Wüst, S., and Bittner, M.: A fast SWIR imager for observations of transient features in OH airglow, *Atmos. Meas. Tech.*, **9**, 1461–1472. doi:10.5194/amt-9-1461-2016, 2016.
- Haralambous, H., Guerra, M., Chum, J., Verhulst, T. G. W., Barta, V., Altadill, D., et al.:
555 Multi-instrument observations of various ionospheric disturbances caused by the 6 February 2023 Turkey earthquake, *J. Geophys. Res.- Space*, **128**, e2023JA031691. <https://doi.org/10.1029/2023JA031691>, 2023.
- Heki, K. and Ping, J.: Directivity and apparent velocity of the coseismic ionospheric disturbances observed with a dense GPS array, *Earth Planet. Sci. Lett.*, **236**, 845–855,
560 doi:10.1016/j.epsl.2005.06.010, 2005.
- Inchin, P. A., Snively, J. B., Williamson, A., Melgar, D., Aguilar Guerrero, J., and Zettergren, M. D.: Mesopause airglow disturbances driven by nonlinear infrasonic acoustic waves generated by large earthquakes, *J. Geophys. Res.- Space*, **125**, e2019JA027628, <https://doi.org/10.1029/2019JA027628>, 2020.
- 565 Inchin, P. A., Guerrero, J. A., Snively, J. B., and Kaneko, Y.: Simulation of infrasonic acoustic wave imprints on airglow layers during the 2016 M7.8 Kaikoura earthquake, *J.*

- Geophys. Res.- Space, 127, e2021JA029529. <https://doi.org/10.1029/2021JA029529>, 2022.
- 570 Jacobs, J.A. and Watanabe, T.: Doppler frequency changes in radio waves propagating through a moving ionosphere, *Radio Science*, 1(3):257–264, 1966.
- Kikuchi, T., Chum, J., Tomizawa, I., Kumiko, K., Hosokawa, K., Ebihara, Y., Hozumi, K., Supnithi, P.: Penetration of the electric fields of the geomagnetic sudden commencement over the globe as observed with the HF Doppler sounders and magnetometers, *Earth Planet. Space*, 73:10, <https://doi.org/10.1186/s40623-020-01350-8>, 2021.
- 575 Kikuchi, T., Araki, T., Hashimoto, K. K., Ebihara, Y., Tanaka, T., Nishimura, Y., Vichare, G., Sinha, A. K., Chum J., Hosokawa, K., Tomizawa, I., Tanaka, Y., Kadokura, A.: Instantaneous Achievement of the Hall and Pedersen–Cowling Current Circuits in Northern and Southern Hemispheres During the Geomagnetic Sudden Commencement on 12 May 2021, *Front. Astron. Space Sci.*, 9: 879314, 580 <https://doi.org/10.3389/fspas.2022.879314>, 2022.
- Laštovička, J. and Chum, J.: A review of results of the international ionospheric Doppler sounder network, *Adv. Space Res.*, 60, 1629-1643, <http://dx.doi.org/10.1016/j.asr.2017.01.032>, 2017.
- Liu, J.Y., Chen, C.H., Lin, C.H., Tsai, H.F., Chen, C.H., Kamogawa, M.: Ionospheric 585 disturbances triggered by the 11 March 2011 M9.0 Tohoku earthquake. *J. Geophys. Res.*, 116, <http://dx.doi.org/10.1029/2011JA016761>. A06319, 2011.
- Mackovjak, Š., Kostárová, S., Chum, J., Schmidt, C., Hannawald, P., Wüst, S., Küchelbacher, L., Truhlík, V., Koucká Knížová, P., Urbář, J., Randuška, L., Štempel, F., and Kubančák, J.: Dataset of Quiet Space Weather Periods for Ionospheric Studies, *Earth Space Sci.*, 13, 590 e2025EA004537, <https://doi.org/10.1029/2025EA004537>, 2026.
- Maruyama, T. and Shinagawa, H.: Infrasonic sounds excited by seismic waves of the 2011 Tohoku-Oki earthquake as visualized in ionograms, *J. Geophys. Res.- Space*, 119(5), 4094–4108, <https://doi.org/10.1002/2013JA019707>, 2014.
- Otsuka, Y., Kotake, N., Tsugawa, T. *et al.*: GPS detection of total electron content variations 595 over Indonesia and Thailand following the 26 December 2004 earthquake, *Earth Planet. Space*, 58, 159–165, <https://doi.org/10.1186/BF03353373>, 2006.
- Paul, K. S., Biswas, T., Haralambous, H., and Moses, M.: Ionospheric disturbances induced by the 29 July 2025 Kuril-Kamchatka Earthquake: Multi - instrument observations and analysis, *J. Geophys. Res.- Space*, 131, <https://doi.org/10.1029/2025JA034646>, 2026.
- 600 Reinisch, B. W., Huang, X., Galkin, I. A., Paznukhov, V., and Kozlov, A.: Recent advances in real-time analysis of ionograms and ionospheric drift measurements with digisondes, *J. Atmos. Sol. Terr. Phys.*, 67 (12), 1054-1062, doi:10.1016/j.jastp.2005.01.009, 2005.

- Rolland, L.M., Lognonné, P., and Munekane, H.: Detection and modeling of Rayleigh wave induced patterns in the ionosphere. *J. Geophys. Res.*, 116: A05320, doi:10.1029/2010JA016060, 2011.
- 605 Schmidt, C., Höppner, K., and Bittner, M.: A ground-based spectrometer equipped with an InGaAs array for routine observations of OH(3-1) rotational temperatures in the mesopause region, *J. Atmos. Sol. Terr. Phys.*, 102, 125–139, <https://doi.org/10.1016/j.jastp.2013.05.001>, 2013.
- 610 Schmidt, C., Dunker, T., Lichtenstern, S., Scheer, J., Wüst, S., Hoppe, U.-P., and Bittner, M.: Derivation of vertical wavelengths of gravity waves in the MLT-region from multispectral airglow observations, *J. Atmos. Sol. Terr. Phys.*, 173: 119-127, <https://doi.org/10.1016/j.jastp.2018.03.002>, 2018.
- Snively, J.B.: Mesospheric hydroxyl airglow signatures of acoustic and gravity waves generated by transient tropospheric forcing, *Geophys. Res. Lett.*, 40(17):4533-4537, <https://doi.org/10.1002/grl.50886>, 2013.
- 615 Song, R., Hattori, K., Zhang, X. *et al.*: A case study of the three-dimensional co-seismic ionospheric disturbance evolution, *Sci. Rep.*, 15, 42209, <https://doi.org/10.1038/s41598-025-26074-1>, 2025.
- 620 Sutcliffe, P.R. and Poole, A.W.V.: Ionospheric Doppler and electron velocities in the presence of ULF waves, *J. Geophys. Res.*, 94(A10):13505-13514, <https://doi.org/10.1029/JA094iA10p13505>, 1989.
- von Savigny, Ch., McDade, I.C., Eichmann, K.U., and Burrows, J.P.: On the dependence of the OH* Meinel emission altitude on vibrational level: SCIAMACHY observations and model simulations, *Atmos. Chem. Phys.* 12 (18), 8813–8828, <https://doi.org/10.5194/acp-12-8813-2012>, 2012.
- 625 Watada, S., Kunugi, T., Hirata, K., Sugioka, H., Nishida, K., Sekiguchi, S., Oikawa, J., Tsuji, Y., and Kanamori, H.: Atmospheric pressure chase associated with the 2003 Tokachi-Oki earthquake, *Geophys. Res. Lett.*, 33, L24306, doi:10.1029/2006GL027967, 2006.
- 630 Wüst, S., Bittner, M., Espy, P.J., French, W.J.R., and Mulligan, F.J.: Hydroxyl airglow observations for investigating atmospheric dynamics: results and challenges, *Atmos. Chem. Phys.*, 23(2):1599-1618, <https://doi.org/10.5194/acp-23-1599-2023>, 2023.
- Wüst, S., Guerra, M., Chum, J., and Gasque, L.C.: Using atmospheric waves for the detection and early warning of natural hazards: a review combining results from the neutral atmosphere and the ionosphere, *Surv. Geophys.*, <https://doi.org/10.1007/s10712-025-09909-4>, 2026.
- 635 Yang, H., Hernandez - Pajares, M., Jarmolowski, W., Wielgosz, P., Vadas, S. L., Colombo, L., et al.: Systematic detection of anomalous ionospheric perturbations above LEOs from

640 GNSS POD data including possible tsunami signatures, IEEE T. Geosci. Remote, 60, 1–
23, <https://doi.org/10.1109/TGRS.2022.3182885>, 2022.

Zettergren, M. D. and Snively, J. B. (2019). Latitude and longitude dependence of ionospheric
TEC and magnetic perturbations from infrasonic-acoustic waves generated by strong
seismic events, Geophys. Res. Lett., 46(3), 1132-1140,
<https://doi.org/10.1029/2018GL081569>, 2019.

645