

On thin glacial ice: New Austrian Glacier Inventory shows accelerating glacier shrinkage and 31 % area loss within two decades

Lea Hartl¹, Jakob Abermann², Ayla Akgün¹, Giulia Bertolotti¹, Tobias Bolch³, Svenja Conzelmann¹, Codrut-Andrei Diaconu⁴, Iris Hansche², Anne Hartig¹, Anna Haut⁵, Kay Helfricht⁶, Bernhard Hynek⁷, Marie Sophie Kaucher¹, Andreas Kellerer-Pirklbauer², Ann Christin Kogel⁸, Julie Krippes⁵, Marcela Violeta Lauria¹, Christoph Mayer⁹, Jan-Christoph Otto⁵, Rainer Prinz⁸, Sina Pröhl¹⁰, Lorenzo Rieg¹¹, Lea Schönleber¹, Gabriele Schwaizer¹², Bernd Seiser¹, Martin Stocker-Waldhuber¹, Markus Strudl¹³, Martin Verhounik³, and Harald Zandler²

¹Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences, 6020 Innsbruck, Austria

²Department of Geography and Regional Science, University of Graz, 8010 Graz, Austria

³Institute of Geodesy, Graz University of Technology, 8010 Graz, Austria

⁴Chair of Remote Sensing Technology, Technical University of Munich, 80333 Munich, Germany

⁵Department of Environment and Biodiversity, University of Salzburg, 5020 Salzburg, Austria

⁶Hydrographic Service Tyrol, Office of the Tyrolean Government, 6020 Innsbruck, Austria

⁷Department Climate Impact Research, Geosphere Austria, 1190 Vienna, Austria

⁸Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, 6020 Innsbruck, Austria

⁹Geodesy and Glaciology, Bavarian Academy of Sciences and Humanities, 80539 Munich, Germany

¹⁰Hochgebirgsnaturpark Zillertaler Alpen, 6295 Ginzling, Austria

¹¹University of Innsbruck, 6020 Innsbruck, Austria

¹²ENVEO-Environmental Earth Observation IT GmbH, 6020 Innsbruck, Austria

¹³Independent researcher, 6460 Imst, Austria

Correspondence: Lea Hartl (lea.hartl@oeaw.ac.at)

Abstract. We present the new Austrian glacier inventory, AGI5. Glacier outlines were manually digitized from high-resolution orthoimagery and digital elevation models, using older inventories as a baseline. The delineation of debris-covered ice was supported by visual analysis of multi-temporal imagery and elevation model differencing, depending on data availability. Differences in interpretation between analysts were assessed using a round robin experiment (mapping of selected glaciers by several analysts). The updated inventory reflects glacier extent in 2023 (55 % of total glacier area in Austria), 2022 (43 %), and 2021 (2 %). The total glacier area in AGI5 is $285 \pm 12 \text{ km}^2$. Most glaciers in Austria (87 %) are smaller than 0.5 km^2 . These "very small" glaciers comprise 22 % of the total glacier area. Nine glaciers remain larger than 5 km^2 and account for more than a quarter of Austria's glacierized area. Area losses since the previous inventory (2004-2012) amount to $129 \pm 23 \text{ km}^2$, corresponding to 31% of the total glacier area. Median area loss rates differ between regions, ranging from 2-3 % per year in more heavily glacierized regions to almost 7 % per year in regions with predominantly smaller glaciers. Of 894 glaciers listed in the previous inventory, 95 have disappeared completely or were no longer mappable. Compared to other glacierized regions, Austria's glacier recession since the Little Ice Age (LIA) maximum is well constrained with a LIA inventory, four high-resolution, consistent AGIs from 1969 to 2022/23, and additional coverage in complementary inventories using different data sources. As glacier loss accelerates, more frequent updates to the AGIs are needed to keep pace with rapid changes.

1 Introduction

Glaciers in the European Alps lost nearly 40 % of their mass between 2000 and 2023 (The GlaMBIE Team, 2025). During recent extreme years, in particular the record breaking summer of 2022, glaciers thinned across all elevation ranges and unprecedented mass loss was recorded throughout the Alps (Berthier et al., 2024; Voordendag et al., 2023; Hartl et al., 2025a; WGMS, 2025; van Tiel et al., 2026). Regional studies in Austria indicate the same trends, with the complete disappearance of several small glaciers since the mid-2000s, volume change patterns indicative of strong disequilibrium, and progressing disintegration processes of increasingly debris-covered glacier remnants (Kellerer-Pirklbauer and Kulmer, 2019; Fischer et al., 2021a; Hartl et al., 2022, 2025b; Conzelmann et al., 2026). Given the magnitude of the current changes, it is essential that local and regional glacier monitoring continues to keep pace with ongoing ice loss.

Multi-temporal glacier outlines are a key part of glacier monitoring and form an important prerequisite for assessing glacier volume and mass change (e.g., Paul et al., 2007; Zemp et al., 2014). They support assessments of the impacts of glacier shrinkage on local hydrological systems, runoff patterns, and natural hazards, aid the calibration and validation of glacier evolution models, and ultimately foster sustainable mountain development (Nussbaumer et al., 2017; Gärtner-Roer et al., 2019). To maintain their usefulness for such applications, inventories need to be updated regularly. Paul et al. (2020) suggested decadal updates of global glacier inventories, noting that more frequent updates are required in regions with more rapid change, such as the European Alps. In Switzerland, Linsbauer et al. (2021) stated 6-year repeat inventories as a goal to appropriately monitor regional glacier area evolution.

In Austria, national-scale glacier inventories (AGI) were compiled for glacier state during the Little Ice Age maximum (AGI LIA, Fischer et al., 2015b), the late 1960s (AGI1, Patzelt, 1980; Groß, 1987), late 1990s (AGI2, Lambrecht and Kuhn, 2007), and the mid-2000s (AGI3, Fischer et al., 2015b). The most recent glacier inventories covering all of Austria reflect glacier state in 2015-2016 (AGI4, Buckel et al., 2018; Paul et al., 2020; Sommer et al., 2020) but used differing methodological approaches compared to the earlier AGI. Table 1 summarizes the available glacier inventories.

A comprehensive inventory update for glaciers in Austria is urgently needed to account for the rapid area losses and progressing glacier disappearance in recent years. This study presents results of a community effort to compile a new Austrian Glacier Inventory along with area change metrics derived from the resulting dataset. The Fifth Austrian Glacier Inventory (AGI5) follows the approach of prior national inventories (Table 1, AGI 1-3, Fischer et al., 2015b) and is based on manual mapping of glacier outlines from high-resolution orthoimagery and digital elevation models (DEM) derived from airborne laser scanning data. Most glaciers in Austria are small and many are fragmented, partially debris-covered, and no longer have persistent accumulation zones. Despite their relatively small size and limited contribution to total glacier area, data on the distribution of such features provides important context for catchment scale applications related to, for example, hydrology, touristic infrastructure, cartography, or potential hazards (e.g., Intergovernmental Panel on Climate Change (IPCC), 2023; Van Tiel et al., 2021; Mani et al., 2023).

The aims of this study are to 1) delineate glacier ice in Austria for the target years 2021-2023 and enable direct comparisons with prior AGIs by adhering to established ID-numbering systems and definitions 2) quantify glacier area changes in Austria since the last inventories 3) discuss the main challenges and sources of uncertainties in compiling regional inventories of rapidly receding mountain glaciers.

2 Methods and data

2.1 Compilation of the fifth Austrian Glacier Inventory (AGI5)

Efforts to compile a new national glacier inventory, AGI5, evolved from discussions within the Austrian glacier monitoring community and were implemented by the same community. In keeping with the prior AGIs, glaciers were grouped by mountain ranges into 20 inventory subregions (Fig. 1), which were assigned to analysts or groups of analysts for outline mapping. Where possible, the mapping of a given subregion was carried out by analysts who were involved in the prior AGIs for that region and/or otherwise familiar with the area.

AGI5 applies the same naming conventions and ID number system as AGI1, 2 and 3 (Table 1). An ID number can be associated with multiple separate glacier fragments if the fragments were previously connected and listed under the same ID. In the following, "glacier" refers to all glacier fragments associated with the same ID even if they are no longer connected. This is in line with the prior AGIs and allows consistent counting of glaciers (the overall number of glaciers does not increase if a glacier splits into two fragments) and per-glacier area change assessments through the AGI time series. In keeping with prior AGI and in contrast to the Swiss national glacier inventories (SGI) and many larger-scale inventories (e.g., Linsbauer et al., 2021; Paul et al., 2020), we use the term "glacier" to refer to any glacial ice identified in the inventory, regardless of feature size or other characteristics, and no minimum feature size is applied in the AGI.

2.1.1 Data basis and general mapping procedure

Glacier outlines were mapped manually using Geographical Information System programs (ArcGIS, QGIS). The AGI3 outlines and, where available, consistent intermediate regional inventories (Table 1) were used as a starting point for the new outlines. The primary data type used for the mapping process was high-resolution regional orthoimagery (spatial resolution: 10-25 cm). Where available, high-resolution DEMs (0.5-1 m) and derived products (hillshades from different illumination angles, elevation change rasters) were used as supporting information or instead of orthophotos depending on data coverage. The georeferenced aerial imagery and DEMs were accessed through governmental open data services. The DEMs are derived from airborne laserscanning surveys and provided by Austrian regional authorities. High-resolution satellite imagery (Pleiades images, 0.5 m; Planet Scope images, 3 m (PBC, 2022)) was used in some cases to compensate for image quality issues in the orthophotos related to snow cover or shading. Table S1 in the supplement provides an overview of the main data sources and acquisition years per subregion. Fig. 2 shows the different available data types for the examples of Großelend Kees in the Ankogel-Hochalmspitze Group and Nördlicher Schalf Ferner in the Ötztal Alps. Großelend Kees (Fig. 2a, b) was mapped

Table 1. Overview of inventories covering Austria’s glaciers at global, Alps-wide, national or regional level.

Name	Inventory Year	Data basis	Coverage	References
AGI LIA	1850	Moraines, historical data	Austria	Fischer et al. (2015b); Groß and Patzelt (2015)
AGI 1	1969	Orthoimagery, DEMs	Austria	Patzelt (1980); Groß (1987); Patzelt (2015)
AGI 2	1996-2002	Orthoimagery, DEMs	Austria	Eder et al. (2000); Lambrecht and Kuhn (2007); Kuhn et al. (2015)
Sommer 2000	1999-2001	L5 TM, L7 ETM+	Alps	Sommer et al. (2020)
RGI 6, 7	2003	L5 TM	Alps	Paul et al. (2011); Pfeffer et al. (2014); RGI Consortium (2023)
AGI 3	2004-2012	Orthoimagery, DEMs	Austria	Fischer et al. (2015b, a)
Sommer 2011	2011	L5 TM	Alps	Sommer et al. (2020)
Sommer 2014	2013-2015	L8 OLI	Alps	Sommer et al. (2020)
AGI 4	2015	Google Earth Imagery	Austria	Buckel et al. (2018); Buckel and Otto (2018)
Paul 2020	2015-2016	Sentinel-2	Alps	Paul et al. (2020)
Salzburg	2008-2018	Orthoimagery, DEMs	Federal Province (Austria)	Bertolotti et al. (2025)
Vorarlberg	2017, 2020, 2021, 2022, 2023	Orthoimagery, DEMs	Federal Province (Austria)	Conzelmann et al. (2026); Conzelmann et al. (2026)
Stubai Alps	2017-2018	Orthoimagery, DEMs	Subregion (Austria)	Helfricht et al. (2024b)
Ötztal Alps	2017	Orthoimagery, DEMs	Subregion (Austria)	Helfricht et al. (2024a)
Silvretta Group	2017	Orthoimagery, DEMs	Subregion (Austria)	Fischer et al. (2021b)
AGI 5	2021-2023	Orthoimagery, DEMs	Austria	This study




Figure 1. a: The black box indicates the location of the 20 subregions and glacierized terrain shown in b in Austria. b: Colors indicate acquisition years of the AGI5 source data for the respective glaciers and regions. Background terrain map: Government of Austria (CC BY-SA 2.0 AU), data.bev.gv.at, region outlines: open street map data.

using mainly information derived from a 2023 DEM, whereas Nördlicher Schalf Ferner (Fig. 2c, d) was mapped based on multi-temporal orthoimagery.

Data availability and acquisition years vary between subregions and Austrian federal provinces. The target year for AGI5 glacier outlines was 2023. Data acquired in 2021 or 2022 was used if no suitable coverage for 2023 was available. Outlines for 2023 were produced for 60 % of glaciers in Austria (55 % of total glacier area). Outlines for 2022 were produced for 39 % of glaciers (43 % of total area). The remaining 1 % of glaciers (2 % of area) were mapped with data acquired in 2021. For the glaciers in the federal province of Vorarlberg (Fig.1), outlines for 2023 produced for a province-level inventory (Conzelmann et al., 2026) were incorporated into AGI5.

About 95 % of the total glacier area in Austria was mapped using mainly aerial images. The remaining area was mapped mainly based on DEM derived products (approx. 4 % of area), or high resolution satellite imagery (1 % of area), typically with orthophotos as auxiliary data. In most cases, analysts reported using more than one data type. This mainly refers to using hillshades or elevation change rasters in combination with aerial or satellite imagery, using satellite imagery instead of orthophotos if snow conditions were more favourable in the former, or using imagery from multiple years to assess glacier evolution. As a general guideline and following the approach of AGI3, analysts were asked to use a maximum zoom scale of 1:3000 in both aerial imagery and DEM derivatives as a starting point for mapping. Most analysts mentioned using a higher zoom level for very small features and cases they considered challenging.




Figure 2. a: Großelend Kees (centroid coordinates: Lon. 13.315°, Lat. 47.025°; Ankogel-Hochalmspitz Group, region 2) as seen in a hillshade generated from a 2023 DEM (September 2023; Province of Carinthia, CC BY 4.0) b) Großelend Kees elevation change 2023-2010 overlayed on the hillshade (DoD: DEM of Differences). c, d: Multitemporal orthophotos of Nördlicher Schalf Ferner (centroid coordinates: Lon. 10.956°, Lat. 46.796°; Ötztal Alps, region 11; orthophotos: 28. July 2020, 19. August 2023; Province of Tyrol; CC BY 4.0 AT). a, b: EPSG: 31258; c, d: EPSG: 31287. Grid in meters. For additional information on data sources see Table S1 in the supplement.

95 Information from field surveys of the glacier margins and relevant local knowledge were incorporated at the discretion of the analysts. Such terrain knowledge was available mainly for glaciers with in situ monitoring programs and a subregion of the Ötztal Alps where an analyst documented debris-covered ice outside of older AGI outlines during site visits over multiple years. In some cases, analysts found that the imagery and data used to delineate AGI5 outlines showed ice where none was mapped in prior outlines due to varying image quality, snow cover, and/or differences in interpretation between analysts. The
100 prior outlines were not modified in these instances. The AGI5 mapping process also revealed several errors in prior inventory attributes, mainly related to wrongly assigned or mistyped ID numbers. Corrections for these cases were issued as updates to the existing AGI3 data publication (Fischer et al., 2015a).

2.1.2 Metadata and quality metrics

For each glacier ID and corresponding outline, the AGI5 attribute tables (Tab. 2) list information related to the source data (acquisition date, data type, image identifiers). Categorical flags indicate potentially detrimental image characteristics (e.g., snow cover or shadows), a qualitative level of uncertainty of the outline, and a debris cover score. In addition, the presence of visible crevasses was flagged. Crevasses have been used as an indicator of past or present flow (e.g., Leigh et al., 2019) and, hence, a criterion that allows distinguishing "glaciers" from "ice bodies" as defined by Cogley (2011). AGI5 does not make such a distinction but including the crevasse flag allows users to filter the data accordingly, for example when comparing AGI5 with datasets that distinguish between these classes, or only include glaciers that show signs of flow (e.g., the SGI). The attribute table also includes a flag indicating overlap of AGI5 with the outlines of the most recent Austrian Rock Glacier Inventory (RoGI, Wagner et al., 2020) to identify potential cases of misclassifications or ambiguous landforms.

The AGI5 attribute table further contains the area of each glacier, the number of fragments per glacier, and elevation statistics. Minimum, maximum, and median elevation of all pixels within a given outline were extracted from a 10 m x 10 m resolution DEM provided by the Austrian Federal Office of Metrology and Surveying (BEV). This DEM is a resampled product from original high-resolution DEM data (0.5 - 1 m) derived from airborne laserscanning produced for the individual provinces of Austria. It includes data from different undisclosed epochs around 2015 and is available through the Austrian open government data platform (Geoland.at, 2015). Accordingly, the national DEM does not exactly match the outline years in AGI5. It was chosen as the most recent product available across all of Austria to derive consistent per-glacier elevation statistics.

2.1.3 Vanishing glaciers

In keeping with the previous AGI, no minimum size threshold for glaciers or fragments of glaciers was applied in AGI5. Analysts were asked to map the remaining glacial ice patches in their subregions to the best of their abilities regardless of feature size. Experience with recent regional inventories showed that very small, highly debris-covered features can be difficult to map even with very high-resolution imagery and auxiliary DEM-derived information (Fischer et al., 2021a; Conzelmann et al., 2026). To account for this, analysts had the option to flag glaciers or glacier fragments that they considered impossible to map but assumed may still contain ice as "vanishing features".

Two possibilities for further handling of such "vanishing features" were incorporated in the processing workflow:

- Case 1: A single glacier ID was associated only with "vanishing features". That is, there were no longer any "mappable" fragments of a given glacier but it was assumed that ice may still be present, for example under debris cover. In this case, the respective glacier was considered a "vanishing glacier" close to disappearance. As the outlines of these features are not mappable with the given source data, they were not included in the main inventory dataset. Centroid coordinates were recorded in a separate file.
- Case 2: A glacier ID was associated with fragments that could still be mapped as well as with "vanishing fragments", which may still contain ice but could not be outlined. In such cases, the outlines of the remaining "mappable" fragments

Table 2. Categorical scores included in the data attributes of each AGI5 glacier outline, indicating potential image quality issues, estimated outline quality, debris cover, the presence of crevasses, and overlap with a nation-wide rock glacier inventory (Wagner et al., 2020). A full list of attributes is provided in the supplement (Table S3 and Fig. S2).

Image quality flag	Categorical value indicating whether snow, shading, or image errors/artifacts affect outline mapping
0 ("good")	No snow cover and no shading over any part of the glacier, no relevant image errors/artifacts.
1 ("medium")	Snow cover and shading allow mapping but parts of the feature are affected by shadows, snow, or image errors/artifacts.
2 ("poor")	Snow cover, shading, or image errors/artifacts do not allow good confidence assessment of the glacier outline.
Outline quality flag	Categorical value indicating issues not related to the quality of the images that affect outline mapping, typically related to debris cover over very small features
0 ("good")	The outline of the feature can generally be mapped with good confidence (e.g., debris cover may be present in some parts but does not substantially obscure the feature boundary, or the boundary can be determined despite debris cover)
1 ("medium")	Some fragments of the feature (single polygons with the same id number) cannot be mapped with confidence (e.g., debris cover obscures part of the boundary)
2 ("uncertain")	The entire feature (all polygons associated with the id number) cannot be mapped with confidence (e.g., fully debris-covered or otherwise obscured boundary)
3 ("very uncertain")	The feature probably still contains some ice (e.g. a small ice body under debris cover) but it is not possible to determine this with certainty and the extent of the feature cannot be mapped with confidence
Debris flag	Categorical value to approximately indicate the amount of debris cover
0	No debris cover (usage: no continuous debris cover on any part of the glacier. Individual rocks are not considered debris cover)
1	Partial debris cover (for example: sections of the terminus are debris-covered)
2	Mostly debris-covered (most of the feature is debris-covered)
3	Fully debris-covered (the entire feature is debris-covered)
4	Not possible to determine (e.g. due to snow cover)
Crevasse flag	Categorical value to indicate whether there are visible crevasses
0	No visible crevasses
1	Visible crevasses
2	Not possible to determine (e.g., due to snow cover)
RoGI	Indicates overlap of the feature with the rock glacier inventory
0	No overlap
1	Up to 20 % of the feature area overlaps with a feature in the rock glacier inventory
2	More than 20 % of the feature area overlaps with a feature in the rock glacier inventory

135 were included in the inventory dataset and the possible existence of "vanishing fragments" was flagged in the dataset attribute table.

Analysts categorized glaciers as "vanished" if they found no evidence of remaining ice within the previous glacier outline. The centroid coordinates of the glaciers that were categorized as "vanishing" or "vanished" are provided as an extra file in the AGI5 dataset publication and the two categories can be distinguished in the data file based on the attributes. To improve the
140 readability of this manuscript, we refer to both categories as "vanishing" in the following.

2.2 Special cases

2.2.1 Glacier ski resorts

The AGI5 region contains nine ski resorts located in glacierized terrain. The resorts apply white reflective coverings to locally reduce melt and preserve particular patches of ice and snow relevant to resort operations (for example, to maintain lift tracks
145 or for snow storage). Such patches of covered ice and snow were included in the outline mapping if they were connected to a remaining uncovered feature. Covered patches not connected to uncovered ice were not included. Fig. 3a and b show examples of coverings and snow management used to locally reduce ablation on Wurten Kees, Carinthia. Fig. 3c and d highlight a narrow connection between a covered remnant of the tongue of Tiefenbach Ferner, Tyrol, and the main glacier.

2.2.2 Ice divides and country borders

150 We maintained the AGI3 ice divides in AGI5 to enable consistent area change assessments and due to the challenges associated with accurately determining the current location of the divides. The ice divides were initially defined for AGI1 (Patzelt, 1980; Groß, 1987) and largely applied in the same way in AGI2 and AGI3. We note that the true location of the ice divides may have shifted since AGI1 due to ice losses.

Some Austrian glaciers border on neighboring glaciers in Italy. Country borders along the main chain of the Alps are
155 generally defined based on drainage divides. That is, the country border mostly follows the ice divides in the glacierized regions of the Ötztal, Stubai, and Zillertal Alps. The main exceptions to this are Hochjoch Ferner and Niederjoch Ferner in the Ötztal Alps. Here, the country border locally deviates from the drainage divides and the glaciers extend across the border. For consistency with AGI3, we continue to include the Italian parts of the respective glaciers in AGI5.

2.3 Area uncertainty estimation

160 Mapping glacier outlines is subject to various sources of uncertainty, which result in uncertainties in the derived glacier area. Building on approaches by previous studies (Abermann et al., 2010; Fischer et al., 2015b; Conzelmann et al., 2026), we applied different area uncertainty estimates to glaciers of different size classes and outline quality categories. Regional uncertainty values were calculated as the sum of the individual glacier uncertainties (that is, uncertainties always cumulate), in keeping with AGI3 (Fischer et al., 2015b).




Figure 3. a: Wurten Kees in the Goldberg Group with the Mölltaler Gletscher ski resort (Carinthia). The red box marks the subset shown in b. b: Zoomed in view of Wurten Kees; reflective coverings (yellow arrow) are applied to reduce ice melt in this sector in the vicinity of resort infrastructure. The orange arrow indicates a narrow connection between two sections of the glacier which is maintained by the resort operator through snow management measures. c: The ski resort of Sölden (Tyrol) operates infrastructure and ski runs on glacierized terrain in the central Ötztal Alps, mainly on Tiefenbach Ferner (T) and Rettenbach Ferner (R). Yellow arrows indicate coverings near the glacier margins. The red box marks the subset shown in d. d: Close-up view of reflective coverings on the terminus of Tiefenbach Ferner and a narrow band of connecting ice between the two coverings. Blue lines indicate AGI5 outlines in all panels. Background image a, b: 2023 UAV orthophoto, courtesy of Geosphere Austria. c, d: 2023 orthophoto mosaic, Province of Tyrol (CC BY 4.0 AT).

165 The relative uncertainties based on glacier size categories correspond to uncertainties applied by Conzelmann et al. (2026), who used uncertainties determined by Abermann et al. (2010) as a starting point for their assessment (± 1.5 % for glaciers larger than 1 km^2 , ± 5 % for smaller glaciers). The values of Abermann et al. (2010) were also applied in the uncertainty estimates of AGI3 (Fischer et al., 2015b). However, Conzelmann et al. (2026) found higher uncertainties for very small, highly debris-covered features based on multi-analyst mapping comparisons (referred to as "Round Robin" experiments in the following),
170 similar to the results of Paul et al. (2013). Accordingly, they adjusted the uncertainty estimates of Abermann et al. (2010), adding smaller size categories with higher relative uncertainties (± 10 % and ± 25 % for features smaller than 0.1 km^2 and 0.05 km^2 , respectively). We used the same approach for AGI5 and additionally accounted for cases flagged by the analysts as highly uncertain by incorporating the outline quality attribute.

For outline quality 0 or 1 ("good" and "medium"), relative area uncertainties were assigned solely based on size categories:

- 175
- Glacier area $\geq 1 \text{ km}^2$: ± 1.5 %
 - $0.1 \text{ km}^2 \leq \text{glacier area} < 1 \text{ km}^2$: ± 5 %
 - $0.05 \text{ km}^2 \leq \text{glacier area} < 0.1 \text{ km}^2$: ± 10 %
 - Glacier area $< 0.05 \text{ km}^2$: ± 25 %

For outline quality 2 and 3 ("uncertain", "very uncertain") the following relative uncertainties were applied regardless of glacier
180 size:

- Outline quality 2: ± 25 %
- Outline quality 3: ± 50 %

To assess differences in interpretation between analysts and enable comparisons of uncertainty estimates between AGI5 and similar studies, we carried out a Round Robin (RR) experiment. The outlines of six glaciers of different sizes and characteristics
185 were digitized independently by 15 analysts using the AGI5 imagery and the AGI3 outlines as a starting point. The selection of glaciers for the RR reflects the abundance of small ($< 1 \text{ km}^2$) and very small ($< 0.5 \text{ km}^2$) glaciers in Austria and intentionally included cases that were considered challenging due to debris cover, vanishing status, and discrepancies between AGI3 and AGI5. Section 3 of the supplementary material provides figures showing the glaciers considered in the RR experiment.

The outlines generated in the RR experiment enable an alternative approach to uncertainty estimation similar to the "buffer
190 method" applied by, e.g., Granshaw and Fountain (2006); Bolch et al. (2010); Mölg et al. (2018); Paul et al. (2020), in which glacier area is computed with buffers of varying sizes applied to the outlines to obtain a likely area range. In large-scale inventories derived from 10 m to 30 m resolution satellite imagery, typical buffer sizes are in the range of 0.5 pixels for clean ice and 1 to 2 pixels for debris-covered ice (e.g., Mölg et al., 2018; Paul et al., 2020). We explored how well the AGI5 RR outlines are aligned by determining the fraction of outlines that fall within a set of buffers (± 2 m, ± 2 -5 m, ± 5 -10 m, ± 10 -20
195 m, ± 20 -40 m, ...) around the main AGI5 outline (supplement Section S4). This yields a buffer distance for clean ice glacier margins with a derived range of likely glacier area, and a larger buffer and area range for debris-covered margins. We applied

the smaller buffer (± 2 m) to glaciers with low debris cover scores (0 or 1, Table 2) and the larger buffer (± 20 m) to mostly or fully debris-covered glaciers (debris score 2 or higher, Table 2) to generate an alternative area uncertainty estimate.

In a separate experiment, we additionally compared two sets of outlines (A, B) for a subset of glaciers in the Ötztal Alps. Outline dataset A was mapped based on aerial imagery and extensive local terrain knowledge, whereas dataset B was mapped exclusively from aerial imagery by analysts without detailed field experience at these sites. Statistics computed from the comparison of these datasets serve as a first-order assessment of potential biases in identification of debris-covered ice without ground truth (e.g., Racoviteanu et al., 2009; Paul et al., 2013).

2.4 Change analysis

Area change since AGI3 was assessed for all glaciers included in both inventories. Average annual change rates were computed for each glacier based on the ID numbering system of the AGIs, and for the 20 subregions of the inventory. To account for the variability in inventory years in both AGI3 and AGI5, change rates were computed on a per-glacier basis and regional values are given as the median change rate of the sample. Vanishing glaciers were included in the change analysis with an area of zero in 2023 (AGI5). That is, a "vanishing year" of 2023 was assumed for the computation. The true vanishing years are generally not known. Uncertainties in area change were computed as the sum of the respective AGI3 and AGI5 uncertainties.

In subregions where regional inventories represent a consistent intermediate time step between AGI3 and AGI5, additional change rates for individual glaciers were computed. This refers to the province-level inventories of Vorarlberg and Salzburg and regional inventories for the Silvretta Group and the Ötztal and Stubai Alps (see references in Table 1).

Per-glacier change rates for AGI1 and AGI2 were computed to present a complete overview of the AGI time series. Regional glacier area change since 1850 was assessed for the subregions contained in all AGIs (some of the smaller regions were omitted in AGI LIA and AGI1).

3 Results

3.1 Glacier distribution in the Austrian Alps

The updated inventory (AGI5) contains 799 glaciers covering a total area of 285 ± 12 km². Five subregions contain 88 % of Austria's glacier area, namely the Ötztal Alps (35 %), the Venediger Group (18 %), the Glockner Group (14 %), the Stubai Alps (11 %), and the Zillertal Alps (10 %, Table 3). Most glaciers in Austria (87 %) are smaller than 0.5 km². These "very small" glaciers (Huss and Fischer, 2016) comprise 22 % of the total glacier area (Fig. 4a, b). The largest size class (> 5 km², 9 glaciers) accounts for more than a quarter of Austria's glacier covered area. Fig. 4c additionally highlights the 10 largest glaciers in Austria. The smallest glacier of this group (Schalf Ferner 4.98 km²) falls outside of the > 5 km² category by only a slight margin and is larger than all other glaciers by more than 1 km².

Most of Austria's glacier area is located between about 2800 and 3200 m a.s.l. (60 % of total area ; Fig. 5a). The AGI5 glacier area extends from a minimum altitude of 1880 m.a.s.l. (Boggenei Kees, Glockner Group) to a maximum altitude of




Figure 4. a: Area of all glaciers in the given size classes as percentage of the total glacier area in AGI5. b: Number of glaciers per size class. c: Glacier area (log scale), median elevation (circular markers) and elevation range (max. and min. elevation, vertical lines) for all 799 glaciers in Austria. The names and sizes (with outline years) of the 10 largest glaciers are indicated in the legend.

3755 m (Rofenkar Ferner, Ötztal Alps). Figure S1 in the supplement provides further visualizations of glacier distribution by area and aspect.

Analysts identified 82 completely debris-covered glaciers and an additional 158 "mostly" debris-covered glaciers (Table 2). These glaciers comprise about 1 % and 4 % of the total glacier area, respectively. The fully and mostly debris-covered glaciers are substantially smaller on average (median area 0.02 and 0.04 km^2 , respectively) than the glaciers with no or only partial debris cover (median area 0.08 and 0.10 km^2 , respectively; supplementary Table S3). Crevasses were visible on 404 glaciers. These glaciers are roughly an order of magnitude larger (median area 0.19 km^2) than glaciers without visible crevasses (363 glaciers, median area 0.02 km^2). We estimate that the area covered by visible reflective geotextiles in ski resorts amounts to <0.2% of total glacier area in Austria (that is, in the imagery used to map AGI5 outlines - the distribution of the coverings changes as they are applied and removed depending on resort operations).

Image quality was mostly considered "good" or "medium", indicating that snow and shadows caused only minor issues in the mapping process (see flag descriptions in Table 2; supplementary Fig. S2). About a third of all glaciers (32 %) were classified as "uncertain" or "very uncertain", mainly due to debris cover (outline quality 2 and 3, Table 2). These glaciers account for 5 % of the total glacier area (supplementary Table S3).

Table 3. Glacier area mapped in AGI5 by region in km² and as a percentage of total area. Percentage change refers to the difference in area between AGI5 and AGI3 relative to AGI3 total area. Loss rates were computed per glacier and the median value is reported for each region. The number of glaciers in AGI5 is exclusive of the vanished and vanishing features.

Region	Glacier area (km ²)	Percent of total area (%)	Area loss (km ²)	Percentage loss (%)	Loss rate (% yr ⁻¹)	Number of glaciers	Number of vanishing glaciers	AGI3 outline years	AGI5 outline years
Ötztal Alps	99.812±3.821	34.96	-37.54±7.067	-27.3	-2.5	185	20	2006	2023
Venediger Group	51.591±1.454	18.07	-17.67±2.951	-25.5	-3.3	88	11	2007-2009	2022
Glockner Group	41.246±1.331	14.45	-10.39±2.577	-20.1	-3.2	74	4	2009	2022
Stubai Alps	30.108±1.454	10.55	-19.03±2.959	-38.7	-3.4	108	9	2006	2023
Zillertal Alps	28.346±1.518	9.93	-16.85±2.93	-37.3	-5.3	126	10	2007-2011	2022-2023
Silvretta Group	10.606±0.504	3.72	-7.87±1.101	-42.6	-3.2	40	5	2004-2006	2023
Ankogel Group	7.127±0.293	2.50	-4.79±0.639	-40.2	-4.6	43	1	2009	2023
Dachstein	4.439±0.379	1.55	-0.64±0.493	-12.5	-2.0	8	0	2012	2021
Goldberg Group	4.403±0.34	1.54	-3.49±0.735	-44.2	-3.7	30	0	2009	2022-2023
Granatspitz Group	2.727±0.314	0.96	-2.75±0.545	-50.2	-3.9	23	7	2009	2023
Verwall Group	2.148±0.264	0.75	-1.93±0.468	-47.3	-2.6	27	8	2006	2023
Rieserferner Group	1.18±0.072	0.41	-1.56±0.167	-57.0	-6.9	5	5	2009	2023
Schober Group	0.629±0.095	0.22	-1.94±0.223	-75.5	-5.8	16	10	2007-2009	2022-2023
Lechtaler Alps	0.315±0.087	0.11	-0.27±0.116	-46.1	-2.5	11	3	2006	2023
Rätikon	0.303±0.017	0.11	-1.25±0.042	-80.50	-4.7	2	0	2006	2023
Hochkönig Group	0.2±0.019	0.07	-0.96±0.077	-82.8	-6.6	2	1	2009	2022
Deferegger Group	0.142±0.065	0.05	-0.16±0.08	-53.0	-3.3	6	1	2009	2023
Carnic Alps	0.126±0.006	0.04	0.03±0.011	35.70	2.7	1	0	2009	2022
Allgäu Alps	0.02±0.005	0.01	-0.05±0.008	-69.50	-4.1	1	0	2006	2023
Samnaun Group	0.018±0.008	0.01	-0.05±0.011	-73.50	-3.2	3	0	2006	2023
Total	285.486±12.046	100	-129.162±23.2	-31.15	-3.4	799	95	2004-2012	2021-2023




Figure 5. a: Glacier area per 50 m elevation bands in AGI3 and AGI5, and distribution of area loss per elevation band (contribution to total loss per elevation band, red line). b: Median elevation of the 20 subregions (vertical axis) from west to east (horizontal axis) with regional area loss (marker color indicates percentage of AGI3 area lost; positive change due to different interpretation of debris covers in the Carnic Alps shown in grey). Marker size indicates regional glacier area in AGI5.

3.2 Glacier area change since AGI3

AGI5 indicates a reduction of glacier area in the Austrian Alps by $129 \pm 23 \text{ km}^2$ compared to AGI3. This corresponds to an area loss of 31 % (relative to AGI3) within a period of about 15 years (area weighted inventory year for AGI3: 2007.5; AGI5: 2022.5), or roughly $2.1 \text{ \% yr}^{-1}$. The median per-glacier area loss rate is $3.4 \text{ \% yr}^{-1}$ (Table 3). 72 % of area loss occurred in an altitudinal range between 2500 and 3000 m. Altitudes above 3000 m contributed 22 % of area loss (Fig. 5a).

Regionally, the greatest absolute losses since AGI3 occurred in the Ötztal Alps with an area reduction of $37.5 \pm 7.1 \text{ km}^2$ followed by the Stubai Alps with $19.0 \pm 3.0 \text{ km}^2$. In relative terms, two regions (Hochkönig Group and Rätikon) lost over 80 % of their AGI3 area. This amounts to absolute losses of $0.96 \pm 0.08 \text{ km}^2$ and $1.25 \pm 0.04 \text{ km}^2$, respectively. An additional six regions lost more than 50 % of their AGI3 area (Schober Group, Samnaun Group, Allgäu Alps, Rieserferner Group, Deferegger Group, Granatspitz Group; Fig. 5b, Table 3). One region (Carnic Alps) showed a slight area increase (Fig. 5b). This region consists of a single glacier (Eiskar Ferner, $0.126 \pm 0.006 \text{ km}^2$ in AGI5, $0.093 \pm 0.005 \text{ km}^2$ in AGI3). The positive area change here is due to differences in interpretation of debris-covered areas in the AGI3 and AGI5 outlines and does not represent actual glacier growth.

255 The most negative median area change rates were found in the Rieserferner Group with $-6.9 \text{ \% yr}^{-1}$ (Table 3). The Schober Group, Hochkönig Group, and Zillertal Alps also had median change rates exceeding $-5 \text{ \% yr}^{-1}$. The least negative median change rates were found in the Dachstein Group with $-2.0 \text{ \% yr}^{-1}$ (AGI5 inventory year: 2021).

Median glacier area decreased from 0.10 km^2 in AGI3 to 0.06 km^2 in AGI5, and median glacier elevation increased from 2849 to 2882 m a.s.l. (Fig. 6a, b). Grouping glaciers by size, the most negative median change rates occurred in the smallest category ($\leq 0.01 \text{ km}^2$) with $-5.9 \text{ \% yr}^{-1}$. Absolute median change rates decreased with increasing glacier size, dropping to losses of less than $2 \text{ \% yr}^{-1}$ for glaciers larger than 1 km^2 (Fig. 6c).

3.3 Vanishing glaciers

AGI5 contains 799 individual glaciers compared to 894 in AGI3. The 95 "missing" glaciers were classified as vanishing between AGI3 and AGI5. That is, analysts found no remaining evidence of ice or considered it impossible to determine an outline for potential remnants of debris-covered ice. The vanishing glaciers had a median size of 0.028 km^2 in AGI3, which is about 70 % smaller than the AGI3 median (Fig. 6a). Of 95 vanishing glaciers, 80 were between 0.01 and 0.1 km^2 in AGI3 (Fig. 6 c). The area loss from the vanishing glaciers amounts to 3.702 km^2 , equivalent to 2.9 % of total area loss between AGI3 and AGI5. Glaciers have vanished across most of the altitudinal range of glacierized terrain in Austria (median elevation in AGI3 between 2363 m and 3356 m). The median elevation of the vanishing glaciers was 44 m (77 m) lower than the AGI3 (AGI5) median (Fig. 6b) and their median vertical extent was considerably smaller (127 m) than the AGI5 median (206 m, Fig. 6 d, e).

The disappearance of 95 glaciers between AGI3 and AGI5 represents a substantial increase in disappearances compared to AGI1 to AGI2 (5) and AGI2 to AGI3 (11). Fig. 7a indicates the location of the vanishing glaciers for the three inventory time steps and their size class prior to being classified as vanishing.

3.4 Area change rates since AGI LIA

275 Comparing AGI LIA with AGI5 shows an area reduction of 68 % for the subregions that were included in all AGI (supplementary material Fig. S3). Median area change rates of Austrian glaciers have become increasingly negative since AGI1 (Table 4). Median change rates amounted to approximately $-1 \text{ \% yr}^{-1}$ from 1969 (AGI1) to the late 1990s (AGI2), increased to $-1.6 \text{ \% yr}^{-1}$ between the late '90s and mid-2000s (AGI3) and to $-3.4 \text{ \% yr}^{-1}$ from AGI3 to AGI5. The intermediate regional inventories available for the time frame around 2017/18 show even more negative median change rates of $-4.8 \text{ \% yr}^{-1}$ for the most recent subperiod. The histograms of per-glacier change rates for AGI1-2, AGI2-3, and AGI3-5 in Fig. 7b - d indicate increasing variability and more frequent occurrences of strongly negative change rates in the most recent periods, in addition to the higher median losses. In absolute terms, losses increased from $3.1 \text{ km}^2 \text{ yr}^{-1}$ (AGI1-AGI2) to $5.8 \text{ km}^2 \text{ yr}^{-1}$ (AGI2-AGI3) and $8.8 \text{ km}^2 \text{ yr}^{-1}$ since AGI3 (Table 4).

285 There are 13, 12, and 17 instances of positive change rates in the three timesteps, respectively. These are caused by differences in interpretation of outlines, imagery revealing ice that was previously not visible, or inclusion of knowledge gained from in situ surveys (Table 4, supplementary material Section S5). For the subregions with an intermediate inventory for 2017/18, 49 of 511 glaciers show positive change rates, some of which are substantially larger than typical positive change

rates associated with subjective differences in interpretation of images between analysts. These cases are primarily related to the inclusion of debris-covered areas based on local knowledge in AGI5 for glaciers in the Ötztal Alps (refer to supplementary material Section S5 for more information on this subset of glaciers).




Figure 6. a: Violin plots for glacier area in AGI5, AGI3, and the AGI3 area of the vanishing glaciers. b: Median glacier elevation in AGI5, AGI3, and the vanishing glaciers (AGI3 elevation). c: Median area change rates for glaciers grouped by size categories (left axis) and number of vanishing glaciers per size category (right axis). Vanishing glaciers are grouped by size bins using their AGI3 area. These values are provided in tabular form in the supplement (Table S4) d: Glacier elevation range (maximum - minimum elevation) plotted against glacier area (log-scale) for all AGI5 glaciers. Colors indicate area change rates. Black stars show AGI3 area and elevation of the vanishing glaciers. e: Subset of d; scale and marker transparency adjusted to highlight the vanishing glaciers.




Figure 7. a: Vanishing glaciers as identified in comparisons of AGI1 and AGI2 (yellow markers), AGI2 and AGI3 (orange), and AGI3 and AGI5 (red). Marker size is scaled to glacier size categories in the respective prior inventories. Background terrain map: basemap.at. Labels indicate Austrian provinces (black border around label text) and neighbouring countries (grey border). Sources for the borders: BEV (2026), Federal Office of Topography swisstopo. b, c, d: Histograms of relative area change for AGI1-AGI2 (b), AGI2-AGI3 (c), and AGI3-AGI5 (d). The legend indicates the number (n) of glaciers assessed in each time period.

Table 4. Change rates since AGI1 expressed as percentages per year and in km² per year. The median and total values are computed from the glacier-wise change rates. That is, different inventory years at individual glaciers are taken into account.

Inventory period	Median change rate (% yr ⁻¹)	Total change rate (km ² yr ⁻¹)	Number of glaciers with positive area change (total number of glaciers)
AGI1-AGI2	-0.88	-3.065	13 (793)
AGI2-AGI3	-1.58	-5.789	12 (905)
AGI3-AGI5	-3.36	-8.760	17 (894)
AGI3-intermediate*	-2.62	-4.461	0 (511)
intermediate*-AGI5	-4.75	-6.440	49 (511)

* intermediate inventories available for some of the subregions, see Table 1

3.5 Round Robin experiment

Table 5 lists the glaciers used for the RR experiment with median glacier area and standard deviation (SD) of the outlines produced by the analysts. Overview maps of all test cases and summarized commentary by the analysts are provided in the supplementary material (Section S3). The SD of the RR outlines ranges from 0.006 to 0.269 km², with generally larger SD
295 with increasing glacier size (Table 5). The SD reaches up to 50 % of the median for the smallest test case not classified as vanishing by the majority of analysts. SD decreases with glacier size and drops to less than 2 % of the median for the largest test case. In absolute terms, the difference between the smallest and largest area values derived from the analysts’ outlines (Fig. 8 a-f) ranges from 0.15 km² at Seekarles Ferner to more than 1 km² for the largest test case (Pasterze), where analysts had different interpretations of the lake-terminating, partly debris-covered glacier tongue. At Arvental Kees, 9 of 15 analysts
300 identified ice beyond the AGI3 outlines, leading to an increase in mapped area between AGI3 and AGI5.

Three of the six RR glaciers were classified as vanishing by at least one analyst. Madlein Ferner, the smallest of the RR glaciers in AGI3, was classified as vanishing by 14 of 15 analysts (Fig. 8a). The second-smallest glacier was classified as vanishing by two analysts, and the third-smallest by one analyst (Fig. 8b, c). That is, most analysts agree on the status of these glaciers but there are individual outlier opinions in both directions. The largest area estimates at all six test glaciers were
305 produced by either analyst 4 or analyst 11 (Fig. 8), who included larger debris-covered areas. Analysts 7 and 13 produced the lowest estimates for the three larger RR glaciers (Fig. 8d-f).

In general, agreement between the analysts is high in sections of the outlines that can clearly be identified as debris-free. Fig. 9 shows examples from Seekarles Ferner to illustrate the decrease in alignment of the identified ice margins in ambiguous or debris-covered sections. The glacier outline can clearly be delineated with little variation among the 15 analysts where it
310 borders a bedrock outcrop, whereas there are substantial discrepancies in a neighbouring section where loose rocks obscure the margin (Fig. 9b). Similarly, there is good agreement between the analysts along the largely debris-free upper margin of the

Table 5. Glaciers used in the Round Robin (RR) experiment, with regions, size in AGI3, standard deviation (SD), SD divided by median area, and notes on characteristics relevant to the mapping process. All listed glaciers were digitized by at least 14 analysts. Figures showing each glacier and the RR outlines are provided in the supplementary material (Section S3).

Name (ID)	Region	Lon, Lat (centroid)	Size GI3 km ² (year)	Median size (km ²)	SD (km ²)	SD/median (%)	Outlines within ±2 m (%)	Comments
Madlein Ferner (13029)	Verwall	10.250, 47.020	0.036 (2006)	0.0	0.009	-	-	Highly debris-covered in AGI5 imagery; snow patches but no visible ice. Overlap with rock glacier inventory.
NN* (3028)	Stubai Alps	11.203, 46.978	0.044 (2006)	0.012	0.006	50.0	20	Very small feature, mostly clean-ice in AGI5 imagery.
Arvental** Kees S (6013)	Venediger Group	12.158, 47.01	0.050 (2007)	0.064	0.024	37.5	45	Fragmented and partially debris-covered. AGI5 outline mapped beyond the AGI3 extent based on 2023 imagery.
Wurten Kees (4038)	Goldberg Group	13.01, 47.037	0.916 (2009)	0.41	0.061	14.9	59	Multiple fragments, mostly clean-ice.
Seekarles Ferner (14033)	Ötztal Alps	10.809, 46.976	1.106 (2006)	0.814	0.045	5.5	57	Debris-covered sector near the terminus. One analyst identified a "vanishing fragment" that may still contain ice but cannot be delineated with confidence (Fig. 9a).
Pasterzen Kees (4027)	Glockner Group	12.695, 47.101	16.316 (2009)	14.568	0.269	1.8	46	Largest glacier in Austria; partly debris-covered tongue; lake-terminating.

* nameless glacier, ** also known as Affental Kees




Figure 8. Glacier area derived from the outlines digitized by the analysts (n =number of analysts) for the six test cases in the RR experiment. The boxplots indicate the first and third quartiles (black boxes) and the median (grey line); grey shading denotes the corresponding violin plots. In panel c, the AGI3 area (dashed black line) is included for comparison because the AGI5 outlines were mapped beyond the extent of the AGI3 outlines in this case. The AGI3 area was larger than all RR results for the other examples shown.

glacier despite shading and snow cover. Agreement decreases along the lower margin, which is partially debris-covered and shadowed (Fig. 9c).

Applying distance buffers to the outlines confirms good alignment of the RR outlines with each other and with the main
 315 AGI5 outline in clean ice sections of the glacier margins, and heterogeneous results in debris-covered sections (Fig. 9b, c; supplementary material Sections S3, S4). At Seekarles Ferner and Wurten Kees, 57 % and 59 % of RR outlines are within ± 2 m of each other (Table 5). At Pasterze, outlines deviate along the large, partly debris-covered and lake-terminating glacier tongue so that only 46 % of the RR outlines are aligned within ± 2 m. At the small, nameless glacier (NN) this value drops to 20 %. A buffer size of ± 40 m covers 90 % of the RR outlines at this glacier (supplementary material Section 4).

320 For all AGI5 glaciers (total area 285.5 km²), applying a ± 2 m buffer yields an area range of 279.7 - 292.7 km², or an approximate uncertainty of ± 6.5 km² (± 2.3 %). With a ± 20 m buffer, the range increases to 231.2 km² - 347.7 km² (± 58 km², or ± 20 %). A ± 40 m buffer corresponds to a ± 37 % area uncertainty. Accounting for differing amounts of debris cover by incorporating the debris attributes (Table 2) in the calculation results in a regional area uncertainty of ± 13.9 km² (± 4.8 %) using a ± 20 m buffer for mostly or fully debris-covered glaciers and ± 2 m for the mostly or entirely clean ice cases. With a

325 ± 40 m buffer the uncertainty increases to ± 20.5 km² (± 5.9 %).




Figure 9. a: The outlines of Seekarles Ferner produced for the RR experiment by 15 analysts. The star marks a "vanishing fragment" placed by one analyst on the debris-covered sector of the tongue (may contain ice, cannot be mapped with confidence). Two analysts excluded this sector entirely, whereas the remaining 12 analysts included it. The red boxes indicate the close-up views shown in panels b and c. Panels b and c: close up views of Seekarles Ferner outlines highlighting good alignment (blue arrows) for bare ice margins and reduced agreement (orange arrows) between analysts for debris-covered sections. The solid and dashed red lines respectively indicate a ± 2 m and ± 20 m buffer around the AGI5 outline.

4 Discussion

4.1 Uncertainties and inherent limitations

4.1.1 Uncertainties in outline mapping

In general terms, we follow an epistemic approach to uncertainties. That is, "the best value is that which is most credible, from the current perspective of knowledge" (Bokulich and Parker, 2025). Uncertainty estimates can be interpreted as a measure of the extent that our knowledge of glacier area - at the glacier-level and regionally - remains inexact. Considering the RR experiment, we assume the most credible value for each glacier is the median of the sample. However, the RR approach is a measure of precision rather than accuracy in the sense that we do not know the "true" area value. Rather, we are assessing how closely the analysts' interpretations of the source data are aligned (see Section 3.5).

The AGI5 uncertainties should be understood as fallible epistemic products that can and should be iteratively improved (Bokulich and Parker, 2025). Notably, improvement does not necessarily mean a decrease in uncertainty. The AGI5 uncertainties provide an estimate of potential errors introduced by differences in interpretation between analysts, but they do not account for potential systematic biases, for example related to debris cover identification, and are hence fallible for glaciers that may be affected by such biases. The AGI5 uncertainty estimates build upon prior work, adapting the approaches of Abermann et al. (2010) and Conzelmann et al. (2026), and have been iteratively improved by incorporating feedback from analysts, who are arguably in the best position to estimate how inexact their outlines may be for a given glacier. The uncertainty estimates could be further improved in the future by explicitly including additional sources of uncertainty as related knowledge improves, for example regarding quantitative bias-estimates for debris cover identification.

The "outline quality" and debris attributes assigned by the analysts (Table 2), indicate that about a third (259) of all glaciers in the study area have high outline uncertainties due to debris cover or otherwise obscured ice margins. Although the attributes are also affected by subjective interpretations (e.g., partially vs. mostly debris-covered), this clearly reflects the most common challenges in manually mapping glacier outlines from orthoimagery and is aligned with expectations given the numerous very small glaciers in Austria. Findings by other studies similarly indicate that surface conditions, timing of image acquisition, quality and resolution strongly affect the visibility and delineation of glacier margins (e.g., Paul et al., 2016; Prinz et al., 2018; Leigh et al., 2019; Fischer et al., 2021a; Andreassen et al., 2022; Hinzmann et al., 2024; Conzelmann et al., 2026).

Discrepancies in manual mapping between different analysts (or the same analyst mapping a glacier multiple times) are to be expected and it is assumed that they cannot be avoided completely. RR experiments are a standard approach to estimating the general magnitude of the uncertainties introduced by such discrepancies (e.g., Abermann et al., 2010; Paul et al., 2013; Fischer et al., 2015b; Paul et al., 2020; Linsbauer et al., 2021; Diaconu et al., 2025). Measures of variability in the area derived from RR outlines are typically used as uncertainty metrics. For example, the 2010 Swiss Glacier Inventory (SGI 2010, Fischer et al. (2014)) found uncertainties of ± 7.6 % for glaciers smaller than 1 km² and uncertainties between ± 3 % and ± 5 % for larger glaciers. For SGI 2016, Linsbauer et al. (2021) reported a standard deviation of glacier area between multiple analysts of 23.8 % for a "very small glacier in a shadowed, snow-covered north face" and values between 0.3 % and 7.1 % in other cases.

In their glacier inventory for Vorarlberg, Conzelmann et al. (2026) found uncertainties of over $\pm 50\%$ in very challenging (small, mostly debris-covered) cases, which is generally in line with studies assessing outlines of debris-covered or otherwise challenging glaciers mapped from satellite imagery (e.g., Paul et al., 2013; Mölg et al., 2018).

Once uncertainty estimates for individual glaciers are determined (often derived from RR comparisons), they are frequently used to estimate region-wide uncertainties in a second step. For example, Linsbauer et al. (2021) applied an area weighted average uncertainty derived from the RR experiment to obtain the total regional uncertainty of SGI 2010. For AGI3, Fischer et al. (2015b), applied relative area uncertainty estimates to individual glaciers (following Abermann et al. (2010), see section 2.3) and summed uncertainties to produce a region-wide uncertainty value.

The regional uncertainties given for AGI3 (Fischer et al., 2015b) and SGI2016 (Linsbauer et al., 2021) are in the range of 2-3 % of glacier area in Austria and Switzerland, respectively (Austria: $415.11 \pm 11.18 \text{ km}^2$, Switzerland: $961 \pm 22 \text{ km}^2$, as per the above studies). Uncertainty estimates for AGI5 are slightly higher with $\pm 4.2\%$ ($285.5 \pm 12.0 \text{ km}^2$). This increase compared to AGI3 can be explained with the higher relative uncertainties applied to highly debris-covered or otherwise very uncertain glaciers. In AGI5, the glaciers with the highest assumed uncertainties account for about 5 % of total area.

The buffer approach derived from the RR outlines yields uncertainty estimates comparable to the uncertainties computed from the tiered system based on glacier size and outline quality scores (Section 2.3), with higher values depending on how debris cover is treated. Outlines of clean ice margins in our RR experiment are generally aligned within $\pm 2 \text{ m}$ of each other. Extrapolating this to the AGI5 region yields a regional area uncertainty of around $\pm 2\%$ (i.e., similar values to AGI3 and SGI2016). In debris-covered sections of the glacier margins, relative uncertainties derived from the buffer approach reach up to around $\pm 40\%$ (i.e., similar values as in Table 5 and as found by Linsbauer et al. (2021) and Conzelmann et al. (2026) for challenging cases).

4.1.2 Takeaways from the AGI5 Round Robin experiment

The results of the AGI5 RR experiment (Fig. 8, Table 5) are broadly in line with the uncertainty magnitudes reported by the studies discussed above and confirm that uncertainties increase with debris cover and decreasing glacier size. Generalizing the comments provided by the analysts regarding the RR experiment (supplement Section S3), differing interpretations of debris cover are the main source of discrepancies. Most of the 15 analysts incorporated geomorphological process understanding in their mapping approach and relied on visible surface features (e.g., changes in surface structure, presence of snow or water) that they considered indicative of subsurface ice to delineate debris-covered ice margins. Other analysts focused more on ice that could be visually identified in the imagery and incorporated less detailed process thinking. The RR experiment showed a general tendency towards smaller glacier area for the latter approach but this is not universally true for all test cases and analysts.

In some cases, analysts provided somewhat contrasting process-based interpretations (e.g., Arvental Kees), or had generally similar interpretations of the geomorphology that nonetheless lead to different conclusions regarding the outline or glacier status (e.g., Madlein Ferner, supplement Section S3). In practice, most of the AGI5 glacier outlines were checked by multiple people and analysts discussed challenging cases among each other. Some of the low outlier cases in the RR (e.g., lowest area

estimate for Pasterze, "vanishing" status for Arvental Kees) would likely have been identified as anomalous and revised in such quality control settings. The high outlier cases are all related to interpretations of debris cover and are difficult to refute or verify. Improved mapping consistency might be achieved with agreed upon, detailed guidelines regarding identification of debris-covered areas. However, improved consistency does not necessarily mean improved accuracy, for example if the majority opinion is subject to unknown or unquantifiable systematic biases.

In discussions prior to the main AGI5 mapping effort, it was agreed that all available information, including relevant process understanding, should be used, as opposed to only mapping visible ice. The latter would likely have produced more consistent results in the RR (and the overall AGI outlines), but would have underestimated the debris-covered area. Similarly, setting an initial zoom-scale of 1:3000 was a measure intended to improve consistency and reduce effort spent on very small-scale details. Most analysts reported deviating from this scale to check for evidence of subsurface ice in challenging cases and considered closer zoom levels to result in improved accuracy. Aside from inter-analyst discrepancies, there is also some amount of variability in outlines mapped by the same analyst on different days (sometimes referred to as "digitization uncertainty"). In our experience, this variability is substantially lower than the inter-analyst variability and we assume it to be covered by our general uncertainty estimates (see also Section S3 in the supplement).

4.1.3 Likely underestimation of debris-covered area without ground truth

As evidenced above, accurately mapping debris-covered ice from remote sensing is challenging even with very high-resolution imagery, and independent ground truth is rarely available (e.g., Racoviteanu et al., 2009; Zandler et al., 2025). Including local terrain knowledge (where available) in the compilation of AGI5 caused area change discrepancies at individual glaciers due to the greater inclusion of debris-covered area compared to AGI3 (supplement Section S5). However, we consider the approach more accurate and, as above, found this more important than maintaining "consistency" with the previous data by omitting the field observations. The same principle (accuracy over consistency with prior outlines) was applied when AGI5 imagery showed evidence of ice beyond the extent of AGI3 (e.g., Eiskar Ferner in the Carnic Alps, Table 3; or Arvental Kees in the Venediger Group Fig. 8).

For the subset of glaciers for which detailed local knowledge was available in AGI5 (supplementary material Section S5), the comparison with outlines mapped exclusively from aerial imagery suggests a systematic underestimation of debris-covered area, similar to previous work in the same region (Strudl, 2024). However, the magnitude of the bias is difficult to quantify at larger scales and it is unknown how often overestimations occur. The tendency towards underestimation of debris-covered area is in line with findings from neighboring South Tyrol (Alto Adige, Northern Italy), where Galos et al. (2025) reported an underestimation of glacier area by 2.3 % in their 2017 glacier inventory compared to a subsequent inventory for 2023. They suggest the underestimation is mainly related to debris cover and assume a similar underestimation is present in outlines produced with 2023 imagery. Given the comparable source data and mapping approaches, AGI5 underestimations of this type can be assumed to be of similar magnitudes. We note that the opposite scenario (overestimation of debris-covered area) may also occur but this is hard to verify and generally cannot be determined from optical imagery alone (e.g., Paul et al., 2013).

Incorporating DEM-derived products or other auxiliary information in glacier outline mapping can support the identification of debris-covered ice margins (e.g., Strudl, 2024; Diaconu et al., 2025; Zandler et al., 2025). Discussing specifically manual mapping with very high-resolution DEM-derived and optical information at the scale of individual mountain ranges (as applied in this study), Abermann et al. (2010) and Fischer et al. (2021a) highlighted that surface elevation change and geometric information improve confidence in outline detection under debris cover. Inventory studies from Switzerland have similarly noted that surface elevation change information reduces debris-related uncertainties (Fischer et al., 2014; Linsbauer et al., 2021).

However, even with very high-resolution data (sub-meter pixels), challenges remain. Snow cover during DEM acquisition, avalanches, or rock fall can create locally positive elevation change signals even if glacier ice was lost over the same time period, making such data difficult to interpret especially for very small glaciers close to disappearance (examples in Conzelmann et al. (2026)). In AGI5, analysts used surface elevation change data where available, usually in addition to optical imagery. Feedback on this was mixed - some found the additional information helpful, but others noted that data were inconclusive in their regions, for example because it was not possible to distinguish loss of ground ice from loss of glacier ice, or due to ambiguities in areas where ice cover was lost completely during the elevation change epoch (i.e., an elevation change signal is present but no ice remains at the end of the epoch). In practical terms, on-site inspections of glacier margins would often be beneficial for accurate mapping but are not feasible at larger scales. Lake-terminating glaciers such as Pasterze (Supplement Section S3, Fig. S10) can pose an additional mapping challenge if ice extends below the water level (Kellerer-Pirklbauer et al., 2021).

In summary, it seems likely that debris-covered area is systematically underestimated in glacier inventories compiled from optical imagery, including AGI5, but this remains hard to quantify exactly. Contextual information (surface elevation change, other data types) has the potential to improve delineation of debris-covered glacier margins but needs to be assessed within the specific geomorphological setting. Geophysical investigations would be needed to clearly identify the ice margins beneath thick debris cover.

4.1.4 Classification challenges and conceptual uncertainties

In addition to differences in interpretation and approach between analysts, "classification errors" and "conceptual errors" are two main sources of uncertainties in remote sensing of glacier outlines (Racoviteanu et al., 2009). The former refers to "misidentified features" and the latter is used for a variety of challenges, e.g., "glacier definition issues such as ice divides, perennial snowfields, minimum size, and fragmentation" (Racoviteanu et al., 2009).

Applying these ideas to AGI5, the conceptual issues related to glacier definitions were addressed in the same way as in prior AGIs. That is, there is no explicit distinction between glaciers, ice patches or perennial snow patches; ice divides were maintained since AGI1; and glacier fragments retain their initial ID number.

The imagery and source data used for mapping of the AGI was largely acquired during favourable, mostly snow-free conditions. However, it cannot be ruled out that off-glacier seasonal snow was erroneously included in glacier outlines in some instances across the AGI time series. If these snow patches then disappear and are not included in the subsequent inventory, derived loss rates are overestimated. Due to the mostly good quality of imagery in AGI5 and based on the available information

460 on source data of older AGI, we do not consider this to be a systematic issue in the dataset but acknowledge that the effect may be present in individual cases.

A further relevant classification error may be found in the distinction between glaciers, rock glaciers, and other periglacial landforms. Comparing the AGI5 outlines with the Austrian Rock Glacier inventory (Wagner et al., 2020) shows 29 cases where AGI5 and the rock glacier inventory outlines overlap. About half of these have only minimal overlap indicative of different
465 landform types existing in close spatial proximity (i.e., the outlines "touch"). The remaining cases have more than 20 % area overlap and may represent classification errors of rock glaciers misidentified as glaciers or vice versa. The distinction between debris-covered glaciers and rock glaciers can be challenging, especially for landform-sequences where both occur in close proximity (e.g., Kellerer-Pirklbauer and Kaufmann, 2018; Haeberli et al., 2024). Nonetheless, the low number of overlapping cases between AGI5 and the rock glacier inventory suggests that this is a relatively minor error source for both data sets.

470 Whether a glacier can and should be classified as "vanished" or "vanishing" might be considered a classification issue as well as a conceptual one. Classification requires consistent inventories or other forms of record keeping that allow a comparison over time to determine if a glacier that was previously present has disappeared. A conceptual definition of a "vanished glacier" requires a common understanding of the term "glacier" that is consistent at least across the inventories used for the comparisons (e.g., Raup et al., 2025; Linsbauer et al., 2025; Pope, 2025; Carlson et al., 2026; Paul, 2026)).

475 In their recent compilation of vanished glaciers in Switzerland, Linsbauer et al. (2025) and GLAMOS (2025) consider glaciers that were larger than 0.01 km² in the SGI of 1979 and were no longer included in the SGI2016, or the forthcoming SGI2023 (GLAMOS, 2025). That is, they classify glaciers as "vanished" if they no longer meet the size requirements for inclusion in the SGIs. In AGI5, we allow a "fuzzy" classification as "vanishing" for cases that may still contain ice but cannot reasonably be mapped with the AGI methodology. This approach was based on feedback by analysts, who indicated that
480 uncertainties in mapping the outlines of very small, debris-covered features can effectively be as large as the remaining area of said features. Of the 95 vanishing glaciers in AGI5, 85 were larger than 0.01 km² in AGI3.

Our RR experiment showed that analysts mostly but not always agree whether a glacier has disappeared. Considering the example of Madlein Ferner (Table 5, Supplement Section 3), we suggest that phrasing such as "14 of 15 analysts agree this glacier is vanishing" or "this glacier has disappeared with over 90 % confidence" can serve as an adequate expression of
485 uncertainty (Bokulich and Parker, 2025) depending on the application. Such an approach would benefit from more extensive RR-type experiments designed specifically to assess vanishing glaciers and the limits of their detection.

In addition to the minimum size requirement, the recent SGIs (Linsbauer et al., 2025; GLAMOS, 2025) apply criteria related to "evidence of flow" such as crevasses and deformation features to exclude ice bodies that do not meet the definition of a glacier as per Cogley (2011). Similar approaches are described in, e.g., Leigh et al. (2019) and Fountain et al. (2023), although not
490 all types of crevasses necessarily indicate ice flow (Kellerer-Pirklbauer and Kulmer, 2019). In AGI5, filtering by size (>0.01 km²) and the "visible crevasses" attribute, 396 glaciers covering an area of 265 km² remain. That is, almost half of the features included in AGI5 would not meet the criteria for inclusion in the SGI. Although the impact of the filter on glacier count is large, the impact on total area is limited with excluded features accounting for only 7 % of total AGI5 area. Despite the differing

criteria, AGI3 and AGI5 and SGI2016 and SGI2023 (forthcoming, (GLAMOS, 2025)) both indicate the disappearance of about
495 11 % of the total glacier count in the respective inventories.

How to treat disappeared or vanishing glaciers in inventories, how to define when a glacier has disappeared, and what to
call glacial remnants that may no longer meet common definitions of a "glacier" are matters of current discussion within the
glaciological community (e.g., Pope, 2025; Paul, 2026). Following the approach of previous AGI, AGI5 includes all glacier
ice identified in Austria, regardless of size or other criteria. Comparability with other approaches, such as that of the SGI, is
500 achieved with the various data attributes that allow users to filter depending on their use-cases. We concur with Pope (2025),
who points out that when a glacier has disappeared depends on "who is asking and why", and hope that AGI5 can contribute
to answering this question in Austria for different types of users and applications.

4.2 Comparison with other inventories and outlook

Comparing total glacier area in Austria in the AGI time series and other inventories, overall area loss trends are consistent
505 although absolute area can vary substantially. For example, AGI2 and AGI3 indicate a larger total glacier area than the roughly
contemporary RGI 7 (RGI Consortium (2023) - Austrian RGI glacier outlines are mainly based on Paul et al. (2011)) and the
inventories of Sommer et al. (2020). This is likely due in large part to the comparatively coarse resolution of the Landsat source
imagery. Paul et al. (2020) noted improved mapping of small features in their 2015-16 Sentinel-2 inventory, which led to the
inclusion of "new" glaciers compared to the RGI. This is reflected by a glacier count of over 1000 in Austria in Paul et al.
510 (2020) versus 800 in the RGI and an improved alignment of the outlines in Paul et al. (2020) with AGI 2.

The number of individual glaciers counted in different inventories generally varies widely, highlighting the influence of
different approaches to fragmentation and exclusion of glaciers based on size criteria. Excluding glaciers $<0.01 \text{ km}^2$ from the
AGIs only marginally affects total glacier area but has noticeable impacts on glacier counts. This calls for pursuing a uniform
and consistent inventory strategy, at least at the national level. Arguably, the number of glaciers in a given region is not essential
515 for typical applications focusing on hydrology or regional glacier area change. However, such numbers are often included in
public communication for general audiences, in particular regarding the number of glaciers that have disappeared in a given
time period (e.g., Boyer and Howe, 2025). Depending on the use-case, care should be taken to contextualize glacier counts by
explaining specifically what was counted and how "glaciers" are defined (e.g., Pope, 2025; Huss et al., 2025).

Glaciers in Austria are expected to largely disappear well before the end of the century under current warming trajectories,
520 with the greatest losses expected in the coming two to three decades (Hartl et al., 2025b). This overall trend is apparent from all
available observational data and from projected future glacier evolution in regional and global modeling studies (e.g., Hanzer
et al., 2018; Zekollari et al., 2019; Rounce et al., 2023; Zekollari et al., 2024; Hartl et al., 2025a; Tricht et al., 2025). Since
AGI2, updates to the AGIs were compiled in roughly decadal intervals. With recent median area change rates of near $-5 \text{ \%}$
 yr^{-1} and losses expected to accelerate as deglaciation progresses, more frequent updates are needed to provide adequate input
525 for hydrological modeling at catchment scales, biotic succession studies, cartographic material, and other applications that rely
on up to date information about local to regional glacier coverage. As has been noted in Alps-wide studies (Paul et al., 2020)
and is evident from the effort to compile AGI5 and previous AGI, variable data availability means that inventories often span

multiple years and it is rarely possible to obtain complete coverage of a larger region from one year. In Austria, new high-resolution orthoimagery and DEMs become available at irregular intervals, usually on a province-level rather than nationally.

530 Targeted airborne campaigns that combine optical and laserscanning acquisitions during low-snow conditions and centralized data processing and digitization efforts would be very beneficial for systematic inventory updates, but require considerable financial resources.

Besides the availability of adequate source data, the work required for manual digitization of glacier outlines is a key challenge for increasing the update frequency of glacier inventories. Combining manual calibration and quality control with

535 automatic approaches has the potential to substantially reduce the workload. Object based image analysis and deep learning approaches (Strudl, 2024; Diaconu et al., 2025; Maslov et al., 2025) can incorporate auxiliary information in addition to optical imagery, potentially improving the detection of debris-covered ice. Glacier outlines produced for 2015/16 and 2023 in an automated, deep learning-based approach (Diaconu et al., 2025) indicate area loss rates in a similar range as for the AGI3 to AGI5 period in Austria. However, data coverage is limited and model reliability appears to decrease for very small,

540 debris-covered features.

For further method development aimed at automation, independent reference inventories compiled from high-resolution data sources, such as the AGIs, can serve to quantify potential biases introduced by methodological differences and support model evaluation. Debris cover is likely to remain a central and difficult to quantify source of uncertainty in manual mapping as well as any automated approaches that might be operationalized in the future.

545 **5 Conclusions**

AGI5 provides an updated status report on Austria's glaciers in 2021-2023, highlighting ongoing glacier recession in all regions. Since AGI3, over 30 % of glacier area has been lost and 95 glaciers have disappeared completely or are no longer mappable. Area change in the highest altitudinal zones above 3000 m is substantial and contributed 22 % of the total losses, indicating that many former accumulation zones have reached a stage of rapid recession and mass loss. The main source of uncertainty in

550 delineating glacier area for AGI5 was the identification of debris-covered glacier ice. The resulting uncertainties for very small glaciers can be large ($> \pm 50$ %) but the glaciers with the greatest uncertainties contribute only marginally to total glacier area (approx. 5 %).

With accelerating disintegration processes and glacier disappearance, mapping challenges related to very small, debris-covered features will gain in relative importance for estimating how much glacial ice remains in individual catchments and at

555 regional scales. Inventories compiled from high-resolution source data, ideally taking into account both optical imagery and elevation change information, can help track changes as well as constrain uncertainties.

Observational data documenting ongoing, progressing regional deglaciation are essential for model development and local and regional planning and adaptation processes. We recommend more frequent updates to Austrian glacier inventories than in the past to match the accelerating rates of change. Ideally, such inventories would extend beyond national borders. This poses

560 challenges related to data availability, which might be mitigated by greater integration of high resolution satellite imagery. The

compilation of AGI5 showed the potential of community initiatives to coordinate inventory updates. In the future, such efforts - at national or Alps-wide scales - would benefit from targeted data acquisition campaigns, improved automation in glacier detection, and systematic support from established monitoring structures.

. The AGI5 outlines have been submitted to the Pangaea data repository and will be freely available at the following DOI once the data
565 curation and publication process is complete: <https://doi.pangaea.de/10.1594/PANGAEA.991106>
The data will also be made available in GLIMS.
Code to produce the figures and analyses in this manuscript is available at: <https://github.com/LeaHartl/inventories>

. Conceptualization: JA, TB, SC, AHg, LH, KH, BH, AKP, JCO, RP, GS, BS, MSW, MS, HZ
Data curation: LH, AHg, MSW, BS
570 Formal analysis: AA, GB, TB, SC, AHg, LH, AHt, KH, BH, MK, AKP, AK, JK, MV, CM, JCO, RP, SP, LR, LS, GS, BS, MSW, MS, MV, HZ
Investigation: LH, AHg
Methodology: LH, AHg, JCO
Validation: CD
575 Visualization: LH, IH, AHg, SC, MS
Writing (original draft preparation): LH
Writing (review and editing): All co-authors

. One co-author is an editor at TC. We have no other competing interests.

. We gratefully acknowledge the departments for geodata of the Federal Provinces of Vorarlberg, Tyrol, Salzburg, and Carinthia, and the
580 BEV for providing essential source datasets and data support. We thank Nina Kirchner, Frank Paul, and an anonymous reviewer for their constructive comments during the review process, as well as Mauro Fischer and Andreas Linsbauer for interesting exchange on the SGI approach and the forthcoming SGI 2023, and Christian Sommer for information on their inventory dataset. R. Prinz and A.C. Kogel are grateful for funding from the University of Innsbruck. This work was partially funded by the Earth System Sciences program of the Austrian Academy of Sciences and the FFG FEMtech program. L. Hartl acknowledges that this research was funded in whole or in part by the Austrian
585 Science Fund (FWF) [10.55776/PAT2089925].

References

- Abermann, J., Fischer, A., Lambrecht, A., and Geist, T.: On the potential of very high-resolution repeat DEMs in glacial and periglacial environments, *The Cryosphere*, 4, 53–65, <https://doi.org/10.5194/tc-4-53-2010>, 2010.
- Andreassen, L. M., Nagy, T., Kjølmoen, B., and Leigh, J. R.: An inventory of Norway's glaciers and ice-marginal lakes from 2018–19
 590 Sentinel-2 data., *Journal of Glaciology*, 68, <https://doi.org/https://doi.org/10.1017/jog.2022.20>, 2022.
- Berthier, E., Vincent, C., and Six, D.: Exceptional thinning through the entire altitudinal range of Mont-Blanc glaciers during the 2021/22 mass balance year, *Journal of Glaciology*, 70, e30, <https://doi.org/10.1017/jog.2023.100>, 2024.
- Bertolotti, G., Wiesenegger, H., Seiser, B., and Fischer, A.: Area and volume change of glaciers in the Salzburg region, Austria: a new inventory (2008–2018), <https://doi.pangaea.de/10.1594/PANGAEA.984878>, 2025.
- 595 BEV: Verwaltungsgrenzen (VGD) BEV 1:250 000 Stichtag 01.04.2026, <https://doi.org/https://doi.org/10.48677/be1d6a26-965a-4548-a6c7-c005e87129d7>, 2026.
- Bokulich, A. and Parker, W. S.: Understanding data uncertainty, *Studies in History and Philosophy of Science*, 112, 90–101, <https://doi.org/https://doi.org/10.1016/j.shpsa.2025.06.003>, 2025.
- Bolch, T., Menounos, B., and Wheate, R.: Landsat-based inventory of glaciers in western Canada, 1985–2005, *Remote sensing of*
 600 *Environment*, 114, 127–137, <https://doi.org/10.1016/j.rse.2009.08.015>, 2010.
- Boyer, D. and Howe, C.: Communication efforts to educate the public about vanishing glaciers, 1958–2025, *Annals of Glaciology*, 67, e2, <https://doi.org/10.1017/aog.2025.10033>, 2025.
- Buckel, J. and Otto, J.-C.: The Austrian Glacier Inventory GI 4 (2015) in ArcGis (shapefile) format, <https://doi.org/10.1594/PANGAEA.887415>, 2018.
- 605 Buckel, J., Otto, J.-C., Prasicek, G., and Keuschnig, M.: Glacial lakes in Austria-Distribution and formation since the Little Ice Age, *Global and Planetary Change*, 164, 39–51, <https://doi.org/10.1016/j.gloplacha.2018.03.003>, 2018.
- Carlson, A. E., Bakken-French, N., Thayne, M., Pappas, S., Molnar, D., and Rood, D.: Disappearing glaciers of the Oregon Cascades, USA, *Annals of Glaciology*, 67, e1, <https://doi.org/10.1017/aog.2025.10037>, 2026.
- Cogley, J.: others (2011) Glossary of glacier mass balance and related terms, 2011.
- 610 Conzelmann, S., Seiser, B., Fischer, A., Lauria, M. V., Stocker-Waldhuber, M., Bertolotti, G., and Hartl, L.: Mapping vanishing glaciers in Vorarlberg, Austria, *Annals of Glaciology*, p. 1–27, <https://doi.org/10.1017/aog.2026.10044>, 2026.
- Conzelmann, S., Seiser, B., Fischer, A., Lauria, M. V., Stocker-Waldhuber, M., Bertolotti, G., and Hartl, L.: Vorarlberg (Austria) Glacier outlines, 2017–2020–2022–2023, <https://doi.org/10.1594/PANGAEA.984116>, 2026.
- Diaconu, C.-A., Zekollari, H., and Bamber, J. L.: DLAGAM: A multi-modal deep learning-based framework for glacier area monitoring,
 615 trained and validated on the European Alps, *Earth and Space Science*, 12, e2025EA004 197, 2025.
- Eder, K., Würlländer, R., and Rentsch, H.: Digital photogrammetry for the new glacier inventory of Austria, *International archives of photogrammetry and remote sensing*, 33, 254–261, 2000.
- Fischer, A., Seiser, B., Stocker-Waldhuber, M., and Abermann, J.: The Austrian Glacier Inventory GI 3, 2006, in ArcGIS (shapefile) format, <https://doi.org/10.1594/PANGAEA.844985>, 2015a.
- 620 Fischer, A., Seiser, B., Stocker Waldhuber, M., Mitterer, C., and Abermann, J.: Tracing glacier changes in Austria from the Little Ice Age to the present using a lidar-based high-resolution glacier inventory in Austria, *The Cryosphere*, 9, 753–766, <https://doi.org/10.5194/tc-9-753-2015>, 2015b.

- Fischer, A., Schwaizer, G., Seiser, B., Helfricht, K., and Stocker-Waldhuber, M.: High-resolution inventory to capture glacier disintegration in the Austrian Silvretta, *The Cryosphere*, 15, 4637–4654, <https://doi.org/10.5194/tc-15-4637-2021>, 2021a.
- 625 Fischer, A., Stocker-Waldhuber, M., Seiser, B., Helfricht, K., and Schwaizer, G.: Glacier inventory Austrian Silvretta 2017/2018, <https://doi.org/10.1594/PANGAEA.936109>, 2021b.
- Fischer, M., Huss, M., Barboux, C., and Hoelzle, M.: The new Swiss Glacier Inventory SGI2010: relevance of using high-resolution source data in areas dominated by very small glaciers, *Arctic, Antarctic, and Alpine Research*, 46, 933–945, <https://doi.org/https://doi.org/10.1657/1938-4246-46.4.933>, 2014.
- 630 Fountain, A. G., Glenn, B., and Mcneil, C.: Inventory of glaciers and perennial snowfields of the conterminous USA, *Earth System Science Data*, 15, 4077–4104, <https://doi.org/https://doi.org/10.5194/essd-15-4077-2023>, 2023.
- Galos, S., klug, C., and Dinale, R.: Short Communication: The 2023 Glacier Inventory for South Tyrol in Context of the Inventories 1997, 2005 and 2017, *Geografia Fisica e Dinamica Quaternaria*, 47, 381–388, <https://doi.org/10.4454/5agvnb15>, 2025.
- Gärtner-Roer, I., Nussbaumer, S. U., Hüsler, F., and Zemp, M.: Worldwide assessment of national glacier monitoring and future perspectives, *Mountain Research and Development*, 39, A1–A11, <https://doi.org/https://doi.org/10.1659/MRD-JOURNAL-D-19-00021.1>, 2019.
- 635 Geoland.at: Digitales Geländemodell (DGM) Österreich, <https://www.data.gv.at/katalog/datasets/d88a1246-9684-480b-a480-ff63286b35b7>, 2015.
- GLAMOS: Swiss Glacier Bulletin 2025, Glacier Monitoring Switzerland, Annual report 146 on glacier observation in Switzerland, Tech. rep., GLAMOS - Glacier Monitoring Switzerland, https://doi.org/10.18752/glbulletin_2025, 2025.
- 640 Granshaw, F. D. and Fountain, A. G.: Glacier change (1958–1998) in the north Cascades national park complex, Washington, USA, *Journal of Glaciology*, 52, 251–256, <https://doi.org/10.3189/172756506781828782>, 2006.
- Groß, G. and Patzelt, G.: The Austrian Glacier Inventory for the Little Ice Age Maximum (GI LIA) in ArcGIS (shapefile) format, <https://doi.org/10.1594/PANGAEA.844987>, 2015.
- Groß, G.: Der Flächenverlust der Gletscher in Österreich 1850–1920–1969, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 23, 131–141, 1987.
- 645 Haeberli, W., Arenson, L. U., Wee, J., Hauck, C., and Mölg, N.: Discriminating viscous-creep features (rock glaciers) in mountain permafrost from debris-covered glaciers – a commented test at the Gruben and Yerba Loca sites, Swiss Alps and Chilean Andes, *The Cryosphere*, 18, 1669–1683, <https://doi.org/10.5194/tc-18-1669-2024>, 2024.
- Hanzer, F., Förster, K., Nemec, J., and Strasser, U.: Projected cryospheric and hydrological impacts of 21st century climate change in the Ötztal Alps (Austria) simulated using a physically based approach, *Hydrology and Earth System Sciences*, 22, 1593–1614, 2018.
- 650 Hartl, L., Helfricht, K., Stocker-Waldhuber, M., Seiser, B., and Fischer, A.: Classifying disequilibrium of small mountain glaciers from patterns of surface elevation change distributions, *Journal of Glaciology*, 68, 253–268, <https://doi.org/10.1017/jog.2021.90>, 2022.
- Hartl, L., Covi, F., Stocker-Waldhuber, M., Baldo, A., Fugazza, D., Di Mauro, B., and Naegeli, K.: Loss of accumulation zone exposes dark ice and drives increased ablation at Weißseespitze, Austria, *The Cryosphere*, 19, 3329–3353, <https://doi.org/10.5194/tc-19-3329-2025>, 2025a.
- 655 Hartl, L., Schmitt, P., Schuster, L., Helfricht, K., Abermann, J., and Maussion, F.: Recent observations and glacier modeling point towards near complete glacier loss in western Austria (Ötztal and Stubai mountain range) if 1.5° C is not met, *The Cryosphere*, 19, 1431–1452, <https://doi.org/https://doi.org/10.5194/tc-19-1431-2025>, 2025b.
- Helfricht, K., Hartl, L., Stocker-Waldhuber, M., Seiser, B., and Fischer, A.: Glacier inventory Ötztal Alps 2017 [dataset], 660 <https://doi.org/10.1594/PANGAEA.965798>, 2024a.

- Helfricht, K., Hartl, L., Stocker-Waldhuber, M., Seiser, B., and Fischer, A.: Glacier inventory Stubai Alps 2017/2018 [dataset], <https://doi.pangaea.de/10.1594/PANGAEA.965791>, 2024b.
- Hinzmann, A., Mölg, T., Braun, M., Cullen, N. J., Hardy, D. R., Kaser, G., and Prinz, R.: Tropical glacier loss in East Africa: recent areal extents on Kilimanjaro, Mount Kenya, and in the Rwenzori Range from high-resolution remote sensing data, *Environmental Research: Climate*, 3, 011 003, <https://doi.org/10.1088/2752-5295/ad1fd7>, 2024.
- Huss, M. and Fischer, M.: Sensitivity of very small glaciers in the Swiss Alps to future climate change, *Frontiers in earth science*, 4, 34, <https://doi.org/https://doi.org/10.3389/feart.2016.00034>, 2016.
- Huss, M., Fischer, M., Linsbauer, A., and Bauder, A.: Continuous monitoring of a glacier’s extinction, *Annals of Glaciology*, 66, e27, <https://doi.org/10.1017/aog.2025.10024>, 2025.
- Intergovernmental Panel on Climate Change (IPCC): Climate Change 2021 – The Physical Science Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, 2023.
- Kellerer-Pirklbauer, A. and Kaufmann, V.: Deglaciation and its impact on permafrost and rock glacier evolution: New insight from two adjacent cirques in Austria, *Science of The Total Environment*, 621, 1397–1414, <https://doi.org/https://doi.org/10.1016/j.scitotenv.2017.10.087>, 2018.
- Kellerer-Pirklbauer, A. and Kulmer, B.: The evolution of brittle and ductile structures at the surface of a partly debris-covered, rapidly thinning and slowly moving glacier in 1998–2012 (Pasterze Glacier, Austria), *Earth Surface Processes and Landforms*, 44, 1034–1049, <https://doi.org/https://doi.org/10.1002/esp.4552>, 2019.
- Kellerer-Pirklbauer, A., Avian, M., Benn, D. I., Bernsteiner, F., Krisch, P., and Ziesler, C.: Buoyant calving and ice-contact lake evolution at Pasterze Glacier (Austria) in the period 1998–2019, *The Cryosphere*, 15, 1237–1258, <https://doi.org/10.5194/tc-15-1237-2021>, 2021.
- Kuhn, M., Lambrecht, A., and Abermann, J.: The Austrian glacier inventory GI 2, 1998, in ArcGIS (shapefile) format, <https://doi.org/10.1594/PANGAEA.844984>, 2015.
- Lambrecht, A. and Kuhn, M.: Glacier changes in the Austrian Alps during the last three decades, derived from the new Austrian glacier inventory, *Annals of Glaciology*, 46, 177–184, <https://doi.org/10.3189/172756407782871341>, 2007.
- Leigh, J., Stokes, C., Carr, R., Evans, I., Andreassen, L., and Evans, D.: Identifying and mapping very small (< 0.5 km²) mountain glaciers on coarse to high-resolution imagery, *Journal of Glaciology*, 65, 873–888, 2019.
- Linsbauer, A., Huss, M., Hodel, E., Bauder, A., Fischer, M., Weidmann, Y., Bärtschi, H., and Schmassmann, E.: The New Swiss Glacier Inventory SGI2016: from a topographical to a glaciological dataset, *Frontiers in Earth Science*, 9, 704 189, 2021.
- Linsbauer, A., Huss, M., Hodel, E., Bauder, A., and Barandun, M.: Vanished glaciers of the Swiss Alps: An inventory-based assessment from 1973 to 2016, *Annals of Glaciology*, p. 1–24, <https://doi.org/10.1017/aog.2025.10031>, 2025.
- Mani, P., Allen, S., Evans, S. G., Kargel, J. S., Mergili, M., Petrakov, D., and Stoffel, M.: Geomorphic Process Chains in High-Mountain Regions—A Review and Classification Approach for Natural Hazards Assessment, *Reviews of Geophysics*, 61, e2022RG000 791, <https://doi.org/https://doi.org/10.1029/2022RG000791>, e2022RG000791 2022RG000791, 2023.
- Maslov, K. A., Persello, C., Schellenberger, T., and Stein, A.: Globally scalable glacier mapping by deep learning matches expert delineation accuracy, *Nature communications*, 16, 43, 2025.
- Mölg, N., Bolch, T., Rastner, P., Strozzi, T., and Paul, F.: A consistent glacier inventory for Karakoram and Pamir derived from Landsat data: distribution of debris cover and mapping challenges, *Earth System Science Data*, 10, 1807–1827, <https://doi.org/https://doi.org/10.5194/essd-10-1807-2018>, 2018.

Nussbaumer, S. U., Hoelzle, M., Hüsler, F., Huggel, C., Salzmann, N., and Zemp, M.: Glacier monitoring and capacity building: Important ingredients for sustainable mountain development, *Mountain Research and Development*, 37, 141–152, <https://doi.org/https://doi.org/10.1659/MRD-JOURNAL-D-15-00038.1>, 2017.

Patzelt, G.: The Austrian glacier inventory: status and first results, 1980.

Patzelt, G.: The Austrian glacier inventory GI 1, 1969, in ArcGIS (shapefile) format, <https://doi.org/10.1594/PANGAEA.844983>, in supplement to: Fischer, Andrea; Seiser, Bernd; Stocker-Waldhuber, Martin; Mitterer, Christian; Abermann, Jakob (2015): Tracing glacier changes in Austria from the Little Ice Age to the present using a lidar-based high-resolution glacier inventory in Austria. *The Cryosphere*, 9(2), 753–766, <https://doi.org/10.5194/tc-9-753-2015>, 2015.

Paul, F.: When has a glacier disappeared?, *Annals of Glaciology*, 67, e18, <https://doi.org/10.1017/aog.2026.10048>, 2026.

Paul, F., Kääb, A., and Haeberli, W.: Recent glacier changes in the Alps observed by satellite: Consequences for future monitoring strategies, *Global and Planetary Change*, 56, 111–122, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2006.07.007>, 2007.

Paul, F., Frey, H., and Le Bris, R.: A new glacier inventory for the European Alps from Landsat TM scenes of 2003: challenges and results, *Annals of Glaciology*, 52, 144–152, <https://doi.org/10.3189/172756411799096295>, 2011.

Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S., Konovalov, V., Le Bris, R., et al.: On the accuracy of glacier outlines derived from remote-sensing data, *Annals of Glaciology*, 54, 171–182, 2013.

Paul, F., Winsvold, S. H., Kääb, A., Nagler, T., and Schwaizer, G.: Glacier remote sensing using Sentinel-2. Part II: Mapping glacier extents and surface facies, and comparison to Landsat 8, *Remote Sensing*, 8, 575, <https://doi.org/https://doi.org/10.3390/rs8070575>, 2016.

Paul, F., Rastner, P., Azzoni, R. S., Diolaiuti, G., Fugazza, D., Le Bris, R., Nemec, J., Rabatel, A., Ramusovic, M., Schwaizer, G., et al.: Glacier shrinkage in the Alps continues unabated as revealed by a new glacier inventory from Sentinel-2, *Earth System Science Data*, 12, 1805–1821, <https://doi.org/10.5194/essd-12-1805-2020>, 2020.

PBC, P. L.: Planet Application Program Interface: In Space for Life on Earth, <https://api.planet.com>, 2022.

Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., et al.: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *Journal of glaciology*, 60, 537–552, 2014.

Pope, A.: Glacier or not? The importance of nuance in definitions of vanishing glaciers, *Annals of Glaciology*, 66, e32, <https://doi.org/10.1017/aog.2025.10030>, 2025.

Prinz, R., Heller, A., Ladner, M., Nicholson, L. I., and Kaser, G.: Mapping the loss of Mt. Kenya’s glaciers: an example of the challenges of satellite monitoring of very small glaciers, *Geosciences*, 8, 174, <https://doi.org/https://doi.org/10.3390/geosciences8050174>, 2018.

Racoviteanu, A. E., Paul, F., Raup, B., Khalsa, S. J. S., and Armstrong, R.: Challenges and recommendations in mapping of glacier parameters from space: results of the 2008 Global Land Ice Measurements from Space (GLIMS) workshop, Boulder, Colorado, USA, *Annals of Glaciology*, 50, 53–69, 2009.

Raup, B. H., Andreassen, L. M., Boyer, D., Howe, C., Pelto, M., and Rabatel, A.: Tracking extinct glaciers in GLIMS, *Annals of Glaciology*, 66, e35, <https://doi.org/10.1017/aog.2025.10027>, 2025.

RGI Consortium: Randolph Glacier Inventory - A Dataset of Global Glacier Outlines, Version 7, <https://doi.org/10.5067/F6JMOVY5NAVZ>, 2023.

Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno, L., Copland, L., et al.: Global glacier change in the 21st century: Every increase in temperature matters, *Science*, 379, 78–83, <https://doi.org/10.1126/science.abo1324>, 2023.

- 735 Sommer, C., Malz, P., Seehaus, T. C., Lippl, S., Zemp, M., and Braun, M. H.: Rapid glacier retreat and downwasting throughout the European Alps in the early 21st century, *Nature communications*, 11, 3209, <https://doi.org/10.1038/s41467-020-16818-0>, 2020.
- Strudl, M.: “Veränderung der Gletscher und deren Gesteinsbedeckung am Kaunergrat in 75 Jahren” Klassifizierung der Luftbilder von 1945 bis 2020 mittels OBIA, Master thesis, Paris Lodron-Universität Salzburg, <https://unigis.at/files/Masterthesen/Full/106918.pdf>, 2024.
- The GlaMBIE Team: Community estimate of global glacier mass changes from 2000 to 2023, *Nature*, pp. 1–7, <https://doi.org/https://doi.org/10.1038/s41586-024-08545-z>, 2025.
- 740 Tricht, L. V., Zekollari, H., Huss, M., Rounce, D. R., Schuster, L., Aguayo, R., Schmitt, P., Maussion, F., Tober, B., and Farinotti, D.: Peak glacier extinction in the mid-twenty-first century, *Nature Climate Change*, <https://doi.org/10.1038/s41558-025-02513-9>, 2025.
- Van Tiel, M., Van Loon, A. F., Seibert, J., and Stahl, K.: Hydrological response to warm and dry weather: do glaciers compensate?, *Hydrology and Earth System Sciences*, 25, 3245–3265, <https://doi.org/10.5194/hess-25-3245-2021>, 2021.
- 745 van Tiel, M., Huss, M., Zappa, M., Jonas, T., and Farinotti, D.: Swiss glacier mass loss during the 2022 drought: persistent streamflow contributions amid declining melt water volumes, *Hydrology and Earth System Sciences*, 30, 23–43, <https://doi.org/10.5194/hess-30-23-2026>, 2026.
- Voordendag, A., Prinz, R., Schuster, L., and Kaser, G.: Brief communication: The Glacier Loss Day as an indicator of a record-breaking negative glacier mass balance in 2022, *The Cryosphere*, 17, 3661–3665, <https://doi.org/https://doi.org/10.5194/tc-17-3661-2023>, 2023.
- 750 Wagner, T., Pleschberger, R., Kainz, S., Ribis, M., Kellerer-Pirklbauer, A., Krainer, K., Philippitsch, R., and Winkler, G.: The first consistent inventory of rock glaciers and their hydrological catchments of the Austrian Alps., *Austrian Journal of Earth Sciences*, 113, 2020.
- Wagner, T., Ribis, M., Kellerer-Pirklbauer, A., Krainer, K., and Winkler, G.: The Austrian rock glacier inventory RGI_1 and the related rock glacier catchment inventory RGCI_1 in ArcGis (shapefile) format, <https://doi.org/10.1594/PANGAEA.921629>, 2020.
- WGMS: Fluctuations of Glaciers (FoG) Database, <https://doi.org/https://doi.org/10.5904/wgms-fog-2025-02>, 2025.
- 755 Zandler, H., Sulzer, W., and Kaufmann, V.: Delineation of debris-covered glaciers with multi-temporal UAV images (Gössnitzkees, Schober Group/Austria), *Advances in Geodesy and Geoinformation*, 74, e66, 1–17, <https://doi.org/10.24425/agg.2025.154150>, 2025.
- Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-CORDEX RCM ensemble, *The Cryosphere*, 13, 1125–1146, <https://doi.org/https://doi.org/10.5194/tc-13-1125-2019>, 2019.
- Zekollari, H., Huss, M., Schuster, L., Maussion, F., Rounce, D. R., Aguayo, R., Champollion, N., Compagno, L., Hugonnet, R., Marzeion, B., Mojtavavi, S., and Farinotti, D.: Twenty-first century global glacier evolution under CMIP6 scenarios and the role of glacier-specific observations, *The Cryosphere*, 18, 5045–5066, <https://doi.org/10.5194/tc-18-5045-2024>, 2024.
- 760 Zemp, M., Armstrong, R., Gärtner-Roer, I., Haeberli, W., Hoelzle, M., Kääb, A., Kargel, J. S., Khalsa, S. J. S., Leonard, G. J., Paul, F., and Raup, B. H.: Introduction: Global Glacier Monitoring—a Long-Term Task Integrating in Situ Observations and Remote Sensing, pp. 1–21, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-540-79818-7_1, 2014.