

CoRSEER: The Calculator of Rock Surface Exposure Age and Erosion Rates for Luminescence Rock Surface Exposure Dating

Arbaz N. Pathan^{1*}, Rabiul H. Biswas^{1*}, Devender Kumar^{2,3}

¹ Department of Earth Sciences, Indian Institute of Technology Kanpur, Kanpur, 208016, India

² CSIR-National Geophysical Research Institute (NGRI), Uppal Road, Hyderabad, Telangana 500007, India

³ Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, Uttar Pradesh 201002, India

*Correspondence to: Arbaz N. Pathan (akp20@iitk.ac.in) and Rabiul H. Biswas (rabiul@iitk.ac.in)

Abstract.

Luminescence rock surface exposure dating (LRSED) estimates the duration of rock surface exposure and the rate of erosion by analysing changes in luminescence with depth below the surface. The wider use of this method has been slowed by limitations in mathematical analysis, such as the subjective selection of the deep plateau for profile normalisation, inconsistent modelling choices across studies, and the lack of a unified inverse-modelling workflow that delivers both parameter estimates and uncertainty bounds. We introduce the Calculator of Rock Surface Exposure Age and Erosion Rates (CoRSEER), an open-source MATLAB application that standardises the full workflow from luminescence depth profiles to exposure and erosion history. CoRSEER first normalises profiles objectively using weighted three-parameter logistic sigmoidal fitting to identify the saturation level and profile shape. It then simulates profile evolution with a finite-difference forward model that includes ambient-radiation-induced signal growth, depth-dependent bleaching described by a surface bleaching rate, attenuation coefficient, and advection caused by erosion. Finally, CoRSEER performs Monte Carlo inversion to estimate calibration parameters from known-age calibration samples, apparent exposure ages for unknown samples, and steady or stepwise erosion histories while reporting the best-fit and uncertainty ranges from likelihood-based filtering. We tested all three modules by reanalysing published datasets from multiple regions, including 23 calibration samples from the original dataset. For a representative calibration profile, CoRSEER-I improved the fit (coefficient of determination of 0.928 compared with 0.868 for the literature dataset) and substantially tightened the uncertainty bounds after objective renormalisation. Across the compiled studies, the CoRSEER-derived apparent ages cluster close to the one-to-one line relative to the published values and span approximately 1 year to 8.5 thousand years, while erosion estimates for a benchmark case remain comparable to the literature results. These results show that objective normalisation plus a transparent, standardised inversion framework can materially improve the reproducibility of luminescence rock surface exposure dating and support consistent inter-study comparisons of exposure and erosion history.

Short Summary:

We developed an open-source program that estimates how long a rock surface has been exposed and how rapidly it has eroded from changes in luminescence with depth. The program objectively normalises measured profiles, simulates signal bleaching and erosion, and reports uncertainty. Tests using published data produced results consistent with previous studies and improved reproducibility, supporting more reliable reconstruction of recent landscape change.

Keywords: Rock Surface Exposure Dating, Infrared Stimulated Luminescence, Rock surface exposure and erosion rate modelling, CoRSEER

Reconstructing the timing of rock surface exposure and erosion history is crucial in geomorphology because it links climate, tectonics, and surface processes that shape the landforms observed today (Molnar and England, 1990). Generally, there are two major reconstruction scenarios: (i) Establishing the exposure ages of host rocks (e.g., valley walls, bedrock, or strath surface) that can predict the age of the natural (e.g., deglaciation) and anthropogenic (e.g., stripping of regolith and quarrying) events that create a freshly exposed surface (Owen and Dortch, 2014; Gliganic et al., 2019). (ii) The transportation, deposition, and reworking histories of transportable clasts, such as cobbles and boulders (Luo et al., 2018). Furthermore, establishing the erosion rates of these rock surfaces provides information on the interactions between climate and landforms (Burbank et al., 1996). Evaluating the exposure-erosion history provides chronological and quantitative constraints on landscape evolution (Colman, 1981).

Terrestrial cosmogenic nuclide (TCN) dating has been extensively used to quantify rock surface exposure age and erosion rate (Lal, 1991; Balco et al., 2008; Dunai, 2010; Schaefer et al., 2022). ^{10}Be dating, the most commonly used technique, is sensitive only above the millennial timescale (Schaefer et al., 2022). Furthermore, the unique concentration of ^{10}Be can provide infinitely many pairs of exposure–erosion histories (Lehmann et al., 2019a) when the growth of ^{10}Be is in a transient state; lower exposure–lower erosion may have a similar effect to higher exposure–higher erosion. Thus, an independent chronometer and/or the use of an additional nuclide is required to constrain both the exposure and erosion histories. Furthermore, the choice of different scaling models has a high impact on the derived exposure ages, and because of the unavailability of a production rate calibration site in most geologically significant sites (e.g., the Himalayan-Tibetan orogen), this problem becomes more critical (Owen and Dortch, 2014).

The introduction of the novel luminescence rock surface exposure dating (LRSED) has shown promising possibilities for overcoming some of these limitations and providing an additional independent chronometer. When rock is shielded from sunlight (part of the host rock or covered by opaque materials) and exposed to ambient radiation over Myr timescales, the ionising radiation fills electron traps to field saturation in minerals such as quartz and feldspar (Wintle and Murray, 2006). When exposed to sunlight, it penetrates the rock surface, causing instantaneous detrapping of electrons, that is, bleaching of the luminescence signal on the surface. As the photon flux attenuates with depth, the bleaching rate decreases, causing a sigmoidal distribution of luminescence signals, also called the luminescence depth profile (LDP) (Sohbati et al., 2011; Sohbati et al., 2012a). The LDP is highly sensitive to exposure and erosion but weakly sensitive to ambient dose rates. The exposure causes deeper propagation of the LDP, erosion cause advection towards the surface and ambient-dose-driven signal growth modifies its shape and position (Sohbati et al., 2018; Lehmann et al., 2019a). Even exposure of <1 year can be significant enough to cause a sigmoidal distribution. Conversely, erosion at extremely low erosion rates (order of 10^{-5} mm/a) can cause measurable advection of the LDP (Pathan et al., 2024). We argue that LRSED has the potential to act as a complementary method to TCN dating, and the LRSED-TCN coupled framework has been introduced in a few studies (Sohbati et al., 2018; Lehmann et al., 2019a; Smedley et al., 2021; Sohbati and Hippe, 2023). A comparison of TCN and LRSED dating is shown in Table 1.

Comparative parameter	Terrestrial cosmogenic nuclide dating using ^{10}Be	Luminescence rock surface exposure dating
Signal principle	In-situ nuclide production in the top few meters by secondary cosmic rays; concentration accumulates with exposure and competes with decay and erosion.	Sunlight bleaches trapped charge in the outer few cm of rock; post-exposure, a depth-dependent luminescence profile develops whose shape reflects exposure time, erosion, and light penetration.
Exposure-age Range	Generally, from 10^3 to 10^6 years, but it is mainly affected by erosion. A lower rate of erosion ($< 10^{-4}$ mm/a) will have no effect, but a higher rate of erosion ($>10^{-2}$ mm/a) can significantly reduce the maximum dateable age (Lal, 1991).	The minimum dateable age is ~ 1 year (often months) given adequate light, and the maximum dateable age is around 50 kyr to 100 kyr, which is highly sensitive to even lower erosion rate on the order of 10^{-4} mm/a, causing a reduction in maximum dateable age (Pathan et al., 2024).
Calibration	Requires production-rate calibration and scaling to site conditions (Schaefer et al., 2022). Location and sample-specific calibration are difficult and generally not performed.	Requires calibration of optical bleaching kinetic parameters (i.e., $\overline{\sigma\varphi_0}$, μ) to site conditions (Sohbati et al., 2012a). Site- and sample-specific calibration is logistically easier.
Scaling/model dependence	Multiple scaling schemes for estimating production rates, such as Lal/Stone (Stone, 2000), the Dunai scaling model (Dunai, 2000), the Lifton-Sato-Dunai (LSD) model (Lifton et al., 2014), the Desilets and Zreda model (Desilets and Zreda, 2003), are available. Predicted exposure age is greatly affected by the choice of model (Owen and Dortch, 2014; Schaefer et al., 2022).	Various LRSED models are available, such as the first-order kinetic model (FOK) (Sohbati et al., 2012b), fading model (Jain et al., 2015; Freiesleben et al., 2023) and general-order kinetic model (GOK) (Biswas et al., 2023; Freiesleben et al., 2023; Pathan et al., 2024). The derived exposure-erosion history highly depends on the choice of models, and the superiority of the GOK model is established, though (Pathan et al., 2024).
Advantages	Robust for multi-kyr exposures; simple sampling; extensive literature; pairing of nuclides provides erosion information.	Resolves short exposures; broad erosion-rate sensitivity; field calibration is logistically straightforward.
Limitations	Weak for decadal–centennial exposures; high erosion biases ages; inheritance can over-age clasts; scaling/PR uncertainties; cost/AMS access (Owen and Dortch, 2014; Schaefer et al., 2022).	Strong dependence on modelling aspect and derived calibration parameters; greater uncertainty for older exposure ages; different kinetics model for feldspar IRSL; community-standard codes/workflows still emerging.
Best suited materials/settings	Quartz-bearing lithologies; stable bedrock; moraines/erosional surfaces with low–moderate erosion (Schaefer et al., 2022).	Feldspar and/or quartz-bearing rocks with light-exposed surfaces; settings with recent exposure or erosion (e.g., deglaciated bedrock, rockfall blocks, anthropogenic cuts).

72 **2 Model**

73 The choice of the incorporated model significantly affects the chronology derived using the LRSED. Thus, it is crucial to include
74 the most accurate, peer-reviewed, simple, and flexible model that can encompass all the geological scenarios for the most robust
75 chronology. This section reviews the LRSED models and discusses the model adopted in this study.

76 **Table 2: Symbology, units, and descriptions of parameters used.**

Symbol	Unit	Description
n	<i>Dimensionless</i>	Number of trapped electrons
N		Maximum possible number of trapped electrons
$\bar{n}$	<i>dimensionless</i>	Normalised electron trap population
$\bar{n}_{sim}$		Simulated normalised luminescence signal
$\bar{n}_{obs}$		Natural normalised luminescence signal
σ	mm^2	Luminescence photoionisation cross section
φ_o	$mm^{-2} s^{-1}$	Photon flux on the surface of the rock
$\overline{\sigma\varphi_0}$	s^{-1}	Wavelength-dependent luminescence decay constant of rocks
μ	mm^{-1}	Attenuation coefficient

Symbol	Unit	Description
$\dot{D}$	Gy/a	Environmental dose rate
D_o	Gy	Characteristic dose of saturation (Dose that fills ~63% of the traps) (Wintle and Murray, 2006)
r	<i>dimensionless</i>	Order of kinetics (Biswas et al., 2023; Freiesleben et al., 2023)
L_x	$counts\ s^{-1}$	The luminescence intensity of the natural sample
T_x	$counts\ s^{-1}$	The luminescence intensity of the laboratory-dosed sample
L_x/T_x	<i>dimensionless</i>	The laboratory-dose normalised luminescence intensity for the natural sample
a	<i>dimensionless</i>	Upper asymptote, the maximum value the function can attain when x approaches infinity
b	mm^{-1}	Steepness of the curve or growth rate of the curve
c	mm	Depth of the inflection point, also referred to as $x_{0.5}$
χ^2	<i>dimensionless</i>	Chi-square
P_k		Likelihood value associated with the synthetic LDP numbered as k
P_{max}		Maximum likelihood, which is associated with the best-fitting synthetic LDP
P_{norm}^k		Normalised likelihood value associated with the synthetic LDP numbered as k
δ		Standard deviation
α		Fractional error, which is equal to $(L_x/T_x)/(L_x/T_x)_{error}$
R^2		Coefficient of determination
x		Depth from surface
$x_{0.5}$	mm	Depth of half saturation from the surface
$x_{0.5,max}$		Forward modelled half saturation depth associated with the maximum detectable age limit.
$x_{0.5,ss}$		Forward-modelled half-saturation depth when the LDP is in a steady state and purely acts as an erosion meter.
$x_{0.5,natural}$		Half saturation depth for natural LDP also equals the c value.
t		Exposure age
t_{max}	a	Maximum age limit for the samples
t_i		Independent exposure age by any other method
t_{cal}		Known exposure age of calibration sample
t_s		Onset time of erosion
t_{ss}		Steady-state erosion time
t_{app}		Apparent exposure age
$\dot{\epsilon}$	mm/a	Erosion rate
$\dot{\epsilon}_{ss}$		Steady-state erosion rate

2.1 Laskaris and Liritzis (2011)

This is the first parametric fit function presented for luminescence rock surface exposure dating (Laskaris and Liritzis, 2011). This was suggested for both OSL and TL.

$$L(x; a_1, b_1, c_1, d_1) = a_1 + \frac{b_1}{2} \operatorname{erfc} \left(\frac{\ln \left(\frac{x}{c_1} \right)}{d_1 \sqrt{2}} \right) \quad (1)$$

where L is the distribution of luminescence signal with respect to depth, and it depends on the fitting parameters as a_1 , b_1 , c_1 , and d_1 . While this empirical model enables precise fitting to the luminescence depth profile, it does not explain the luminescence mechanism

inside the rock, and the derived fitting parameters are not linked to the natural kinetic parameters. Additionally, this model assumes zero erosion conditions, which are not always achievable. Thus, this model was not incorporated into any further LRSED studies.

2.2 Sohbati et al. (2018)

Sohbati et al. (2018) proposed a model whereby the instantaneous concentration of trapped electrons n at depth x is:

$$\frac{dn}{dt} = (N - n)F(x) - nE(x) \quad (2)$$

where,

$$E(x) = \overline{\sigma\varphi_0}e^{-\mu x} \quad (3)$$

$$F = \dot{D}(x)/D_0 \quad (4)$$

With the boundary conditions $n = N$ at $t = 0$, the analytical solution was evaluated as

$$\frac{n(x, t)}{N} = \frac{E(x)e^{-t[E(x)+F]} + F}{E(x) + F} \quad (5)$$

$$\frac{n(x, t)}{N} = \frac{\overline{\sigma\varphi_0}e^{-\mu x}e^{-t\left(\overline{\sigma\varphi_0}e^{-\mu x} + \frac{\dot{D}}{D_0}\right)} + \frac{\dot{D}}{D_0}}{\overline{\sigma\varphi_0}e^{-\mu x} + \frac{\dot{D}}{D_0}} \quad (6)$$

The model was further improved by incorporating an erosion scenario (Sohbati et al., 2018).

$$x(t) = x_0 - \dot{\epsilon}t \quad (7)$$

$$E(x(t)) = \overline{\sigma\varphi_0}e^{-\mu(x_0 - \dot{\epsilon}t)} = (\overline{\sigma\varphi_0}e^{-\mu x_0})e^{\mu\dot{\epsilon}t} = E_0e^{\mu\dot{\epsilon}t} \quad (8)$$

$$\frac{dn}{dt} = (N - n)F - nE_0e^{\mu\dot{\epsilon}t} \quad (9)$$

where the analytical solution, expressed as a confluent hypergeometric function M , is:

$$\frac{n(x, \dot{\epsilon})}{N} = M\left(1, 1 + \frac{F}{\mu\dot{\epsilon}}, -\frac{E(x)}{\mu\dot{\epsilon}}\right) \quad (10)$$

The analytical solution improves the feasibility of modelling by incorporating the effects of dose rates, first-order kinetics (FOK) decay behaviour and erosion. However, in the case of polymineral or feldspar samples, where the infrared stimulated luminescence (IRSL) of feldspar minerals exhibits non-first-order decay, the general-order kinetic (GOK) model is more suitable for these samples.

2.3 Freiesleben et al. (2023) model

Freiesleben et al. (2023) developed this model to incorporate GOK:

$$\frac{dn}{dt} = -E_{\dot{\epsilon}}(x, t)n^r \quad (11)$$

where r is the order of kinetics.

$$E_{\dot{\epsilon}}(x, t) = \overline{\sigma\varphi_0}e^{-\mu(x_0 - \dot{\epsilon}t)} \quad (12)$$

$$n(x, t_{\dot{\epsilon}}) = [(r - 1)t_{\dot{\epsilon}}E(x) + n^{1-r}]^{\frac{1}{1-r}} \quad (13)$$

where the apparent exposure time is given as

$$t_{\dot{\epsilon}} = \frac{1}{\mu\dot{\epsilon}}(1 - e^{-\mu\dot{\epsilon}t}) \quad (14)$$

This model considers the order of kinetics and erosion scenario but ignores the effect of the dose rate on LDP propagation. However, in natural environments, the dose rate considerably affects the propagation of LDP over a longer timescale ($>10^3$ years), and this effect is amplified with an increase in the order of kinetics (Pathan et al., 2024). This model also assumes a constant average rate of erosion, which may not be applicable in different erosion scenarios, such as step-function or stochastic histories (Ganti et al., 2016; Brown and Moon, 2019; Lehmann et al., 2019a).

2.4 Model framework after Lehmann et al. (2019a), Biswas et al. (2023) and Pathan et al. (2024)

The modelling framework adopted in this study brings together the current understanding of rock-surface luminescence signal evolution within a single computational framework. It includes: (1) general-order kinetics, (2) advection through the natural mechanism of erosion, and (3) ambient dose-rate-driven signal accumulation. This formulation was required to simulate the luminescence-depth profile more accurately and builds upon earlier work by Lehmann et al. (2019a), Biswas et al. (2023), and Pathan et al. (2024). Lehmann et al. (2019a) introduced the advection term in response to erosion as $\varepsilon \frac{d\bar{n}}{dx}$ but for the FOK ($r = 1$) in addition to earlier models. Thus, the instantaneous change in normalised luminescence $\bar{n}$ (see Section 3.1 for normalisation procedure) with respect to time t at depth x , is expressed as:

$$\frac{d\bar{n}}{dt} = \left\{ \frac{\dot{D}}{D_0} (1 - \bar{n}) \right\}_{growth} - \left\{ \bar{n}^r \overline{\sigma\varphi} e^{-\mu x} \right\}_{decay} + \left\{ \varepsilon \frac{d\bar{n}}{dx} \right\}_{erosion} \quad (15)$$

This model can be solved using a finite-difference method with a second-order upwind scheme for the advection term rather than altering the depth term (Sohbati et al., 2018). Furthermore, Biswas et al. (2023) modified this model for a zero erosion scenario by incorporating GOK ($r > 1$) in the decay term and demonstrated the severe underestimation of the exposure age if FOK was used. Pathan et al. (2024) solved the GOK model (Eq. 15) for a non-zero erosion scenario and reported an overestimation of the erosion rate when FOK was used. Thus, we incorporate the most recent model, as in Eq. 15, into the mathematical framework of CoRSEER. The model accounts for signal growth due to environmental dose rate, depth-dependent light attenuation, nonlinear bleaching behaviour of IRSL signals, and advective loss of material due to surface erosion. Therefore, it provides a physically consistent framework for simulating LDP evolution under both exposure and erosion. This class of model has been used in several studies. Lehmann et al. (2019a) and Elkadi et al. (2022) used this model assuming FOK ($r = 1$) to jointly invert ^{10}Be cosmogenic nuclide data and LRSED profiles, allowing erosion histories and erosion-corrected exposure ages to be constrained through complementary dating systems. Smedley et al. (2021), Brill et al. (2021), and other studies also applied this framework for reconstructing erosion histories, generally assuming FOK.

The incorporation of general-order kinetics is particularly important for feldspar-based IRSL signals, where slower and non-exponential bleaching behaviour is commonly observed. The FOK model assumes simple exponential decay with ($r = 1$), whereas the GOK model introduces a kinetic-shape parameter (r), which can represent slower bleaching behaviour observed in feldspar signals (Duller, 1997; Chen and Leung, 2003; Guralnik et al., 2015). Previous studies have shown that the use of FOK may lead to underestimation of exposure ages and overestimation of erosion rates, in some cases by up to an order of magnitude (Biswas et al., 2023; Pathan et al., 2024; Luo et al., 2026). By embedding this model within CoRSEER, we provide a reproducible platform for forward and inverse modelling of LRSED profiles, enabling more consistent reconstruction of exposure–erosion histories. The prerequisites for CoRSEER, sampling strategy, and sample-processing methods for LRSED are provided in Supplementary Sections S1, S2, and S3, respectively.

3 Methodology

CoRSEER is a MATLAB-based app designed to streamline the quantification of rock surface exposure age (for uneroded surfaces) and erosion histories (for eroded surfaces) through inverse modelling of luminescence depth profiles. The workflows are as follows.

3.1 Normalisation of experimental data

To eliminate the effect of intersample variability (mass and luminescence production efficiency), the test-dose normalised luminescence (L_x/T_x) was measured at different depths to generate the LDP. The LDP is further adjusted (or normalised) to plateau at 1 to facilitate further mathematical operations (Sohbati et al., 2012a). Traditionally, users select a depth beyond which the

experimental LDP is assumed to be saturated through visual inspection and compute the mean L_x/T_x of all points thereafter to normalise the entire profile. However, the high scatter in the saturation region often leads to subjective depth selection, introducing human error and bias. The incorrectly identified plateau can systematically distort the normalised signals and skew the kinetic parameter estimation. To overcome this, we present an empirical normalisation using the sigmoidal curve fitting (SCF) method. We chose the following three-parameter logistic sigmoidal function (Kaufmann, 1981) to model the luminescence-depth profile:

$$f(x) = \frac{a}{1 + \exp[-b(x - c)]} \quad (16)$$

Here, a is the upper asymptote, the maximum value the function can attain as $x \sim \infty$, denotes the saturation luminescence intensity (or L_x/T_x), and b is the growth rate, which controls the steepness of the curve. A higher b indicates a steeper transition of L_x/T_x with depth. c represents the inflection point or $x_{0.5}$, that is, the depth at which half of the saturation signal is reached (Fig. 1). To derive a , b , and c , we fitted Eq. 16 to the experimental LDP using weighted nonlinear least-squares regression in MATLAB. The initial guesses for fitting parameters were set to $a_0 = \max\left(\frac{L_x}{T_x}\right)$, $b_0 = 1$ and $c_0 = \text{median}(x)$. We enforced bounds to ensure physically meaningful solutions: (a) lower bounds: $[0, 0, 0]$, (b) upper bounds: $[\infty, \infty, \max(x)]$, and provided the weight factors (w_i) as $w_i = \frac{1}{\alpha}$ where α is the fractional error. The experimentally observed LDP (L_x/T_x vs depth) was normalised with the respective a value for each sample and referred as SCF-normalised LDP. This approach was used to identify the plateau and normalise the experimental data. This method removes bias, improves reproducibility, and ensures a consistent LDP normalisation algorithm across diverse datasets by eliminating manual intervention.




Figure 1: The luminescence-depth profile (LDP) sigmoidal shape consists of three distinct regions: (i) the near-surface bleaching plateau, (ii) the middle slope ($\tan \theta = \left(\frac{ab}{4}\right)$ at $c = x_{0.5}$) of sigmoidal shape, and (iii) the deep saturation plateau of magnitude ' a '.

3.2 Inverse modelling in CoRSEER

In inverse modelling, multiple LDPs are simulated by solving Eq. 15 using the finite difference method for two main scenarios: (a) varying kinetic parameters, $\overline{\sigma\varphi_0}$, and μ (keeping the exposure-erosion history as fixed) or (b) by varying the exposure-erosion history (keeping $\overline{\sigma\varphi_0}$, and μ as fixed). For a calibration sample (known exposure-erosion history), kinetic parameters ($\overline{\sigma\varphi_0}$, and μ) are randomly varied, whereas for natural samples (known $\overline{\sigma\varphi_0}$, and μ), the exposure-erosion history is randomly varied. The other kinetic parameters, $\dot{D}$, D_0 , and r are constrained independently using DRAC (Durcan et al., 2015; Sohbati et al., 2018), IRSL dose-response curve (Wintle and Murray, 2006) and fitting decay of IRSL with light exposure (Biswas et al., 2023), respectively. These parameters are user-defined in CoRSEER. The most probable, or best-fitting, LDPs and corresponding parameters are constrained

by comparing the simulated LDPs ($\bar{n}_{sim}$) and experimentally observed SCF normalised LDP ($\bar{n}_{obs}$) and by minimising the chi-square misfit (χ^2) and maximising the *likelihood*, following the approach by Lehmann et al. (2019a).

$$\chi^2 = \sum_{i=1}^l \frac{((\bar{n})_{obs}^{(i)} - (\bar{n})_{sim}^{(i)})^2}{\delta^2} \quad (17)$$

$$likelihood = P = e^{-\chi^2/2} \quad (18)$$

$$P_{norm}^k = \frac{P_k}{P_{max}} \quad (19)$$

where l is the number of experimental data points, δ is the standard deviation of the residual between the observed experimental profile and the best-fitting sigmoidal curve, divided by ‘ a ’ factor, P_k is the likelihood value associated with the simulated LDP numbered as k , P_{max} is maximum-likelihood, which is associated with the best-fitting synthetic LDP and P_{norm}^k is the normalised likelihood value associated with the simulated LDP numbered as k . The best LDP and corresponding parameters were selected by applying the Monte Carlo acceptance criterion as $P_{norm}^k > U$, where U is a random number between zero and one. Finally, CoRSEER establishes the values of best-fit, median, $\pm 1\delta$, and $\pm 2\delta$ for all required parameters with confidence intervals.

3.2.1 Calibration

CoRSEER-I module constrains $\overline{\sigma\varphi_0}$ and μ from the SCF normalised LDP of calibration sample (known exposure age and known erosion rate). The inverse modelling (described above) is performed with fixed input parameters $\dot{D}$, D_0 , r , t , and ε and random selection of $\overline{\sigma\varphi_0}$ and μ from a user-defined range for a large number of iterations. It must be mentioned that erosion is assumed to be negligible for the young calibration sample. However, for the older calibration sample, erosion might not be zero, which may bias the parameter estimates. CoRSEER-I has the option to include erosion rate ($\varepsilon > 0$), if known.

The parameter pair ($\overline{\sigma\varphi_0}$ and μ) associated with the lowest normalised chi-square is designated the best-fit bleaching parameter. To assess uncertainty, CoRSEER-I filters the entire chi-square field using a probabilistic threshold: all solutions where $P_{norm}^k > \text{rand}(m, m)$ (following the method of Pathan et al. (2024)) are retained, and their median and $\pm\delta$ uncertainty bounds are reported. This approach ensures the statistical robustness and physical interpretability of the calibrated parameters. The details of the input, mechanism, and output of CoRSEER-I are discussed in Supplementary Section S4.1. These location- and rock-type-specific calibration parameters ($\overline{\sigma\varphi_0}$ and μ) are used as inputs in exposure age and erosion rate calculation.

A representative analysis was performed on the LDP of the available literature data ROAD02 (Smedley et al., 2021). As we are solving for only one LDP corresponding to the IRSL50 signal, the LDP is termed IR50_ROAD02. The LDP was normalised to 1 by dividing the mean deep-plateau L_x/T_x . This road-cut sample has an exposure age of 57 a (when the LDP was generated). The other parameters are $r = 1$, $\dot{D} = 4.76 \pm 0.15 \text{ Gy/kyr}$, and $D_0 = 500 \text{ Gy}$. The inverse modelling is performed for 500×500 pairs of $\overline{\sigma\varphi_0}$ (ranging from 10^{-10} to 10^0 s^{-1} , spacing in log-scale) and μ (ranging from 0 to 4 mm^{-1}), and considering no erosion. After the calibration is completed, CoRSEER generates three diagnostic plots, as shown in Fig. 2a-c.

The normalisation factor in the SCF analysis was estimated as $a = 1.085$ (Fig. 2a), which shows the presence of under-normalisation, and the profiles were further renormalised (SCF normalisation) using this factor for accuracy. Fig. 2b and 2c show the plausible fitting using CoRSEER-I comparable to the fitting seen in the literature (Fig. 6g of Smedley et al. (2021)). The resulting best-fitting $\overline{\sigma\varphi_0}$ and μ values are $1 \times 10^{-7} \text{ s}^{-1}$ and 1.08 mm^{-1} which vary significantly from reported values of $6.67 \times 10^{-6} \text{ s}^{-1}$ (overestimated approximately by an order of 2) and 2.1 mm^{-1} (overestimated by factor of ~ 2) by Smedley et al. (2021). Furthermore, the $\pm 1\delta$ range for $\overline{\sigma\varphi_0}$ is reduced from $3.50 \times 10^{-7} - 1.27 \times 10^{-4} \text{ s}^{-1}$ to $6.39 \times 10^{-8} - 3.11 \times 10^{-7} \text{ s}^{-1}$, and μ is reduced from $1.4 - 2.6$ to 1.00 to 1.31

mm⁻¹, showing the narrowing of the uncertainty interval. We investigated the reason for this discrepancy between the results by comparing the manually normalised LDP (literature) from the SCF-normalised LDP (factor $a = 1.085$). We plotted the simulated profiles using the calibration dataset associated with both LDPs (literature and CoRSEER-I, respectively). Fig. 2d shows that there is no tangible depth difference between the literature and CoRSEER-I generated profiles, but we see a noticeable difference in the slope of the LDPs. The estimated R^2 values were 0.868 for the literature dataset and 0.928 for CoRSEER-I, indicating greater precision in fitting with CoRSEER, as shown in Fig. 2e.

Despite the slight difference in the R^2 values, the evaluated kinetic parameters varied significantly. The apparent reason for this discrepancy likely results from (1) SCF normalisation of the LDP and (2) the Fitting Strategy.




Figure 2: Reevaluation of the IR50 calibration parameters for sample ROAD02 (IR50_ROAD02) (Smedley et al., 2021). (a) The natural IR50 L_x/T_x depth profiles (hollow circles) and associated sigmoidal fit (teal solid line). Although the normalisation was already performed by Smedley et al. (2021), the fitted amplitude showed $a = 1.085$, indicating $\sim 8.5\%$ under-normalisation. Parameters b and c indicate the steepness and $x_{0.5}$, respectively. (b) The simulated LDPs for best-fit (dashed blue) and median (solid orange) calibration parameters associated with the SCF-normalised LDP (hollow circles). The precise overlap between both simulated LDPs shows a strong agreement between the best-fit and median values, indicating plausible fitting strategies. (c) Probability likelihood distribution in $\sigma\phi_0$ and μ space from Bayesian inversion. The dashed lines represent the best-fitting values, and the solid line represents the median of the filtered values using the criteria described in Section 3.2.1. (d) Direct comparison of the literature-normalised profile (black circles) and SCF-normalised profile (cyan squares) using CoRSEER-I alongside their corresponding best-fit simulated luminescence depth profile (black and cyan lines). Both datasets show comparable $x_{0.5}$, but there is a slight change in the slope after renormalisation. (e) Residuals (data – model) as a function of depth for both fits, highlighting the improved agreement for CoRSEER-I. The coefficient of determination increases from $R^2 = 0.868$ (literature handling) to $R^2 = 0.928$ (CoRSEER-I), indicating a tighter, more precise fit to the same underlying measurements.

3.2.2 Age estimation

This module evaluates the exposure age from the LDP of an unknown sample for a no-erosion scenario. If the sample had an erosion history, the derived age would be the apparent exposure age (t_{app}). The inverse modelling is performed with fixed input parameters, $\dot{D}$, D_0 , r , $\sigma\phi_0$, μ and ϵ and random selection of t from a user-defined range for a large number of iterations. A probabilistic threshold filter is applied (e.g., $P_{norm}^k > rand(m,1)$) to extract accepted LDPs from which the best-fitting age, median age and $\pm 1\delta$ age

uncertainty are derived. The algorithm also calculates t_{max} , the maximum possible exposure age, which is associated with depth beyond which the LDP does not hold any information about exposure history ($x_{0.5,max}$). Here, we define $x_{0.5,max} < x_{0.5,ss} - 1/\mu$, as described by Sohbati et al. (2018). We simulated the evolution of $x_{0.5}$, the depth at which the luminescence signal reaches 50% of the saturation signal. The simulation continued until $x_{0.5}$ asymptotically approached a steady-state depth of $x_{0.5,ss}$. The input, mechanism, and output of CoRSEER-II are discussed in Supplementary Section S4.2.

Here, we evaluated the exposure age of the rock avalanche boulder BALL02 (Smedley et al., 2021) (hereafter IR50 BALL02) using CoRSEER-II. The boulder was deposited and exposed due to the rock avalanche event dated to $t_i = 4.52 \pm 0.27$ ka. Smedley et al. (2021) estimated the erosion-uncorrected exposure age (t_{app}) as 8 ± 2 a, which was evaluated using $\dot{D} = 3.32 \pm 0.12$ Gy/ka, $D_0 = 500$ Gy, $\overline{\sigma\varphi_0} = 6.67 \times 10^{-6}$ s $^{-1}$, $\mu = 2.1$ mm $^{-1}$, $\varepsilon = 0$ and $r = 1$ (The calibration parameters were evaluated from ROAD02). Here, we replace the calibration parameters, $\overline{\sigma\varphi_0}$, and μ with the values obtained in CoRSEER-I, $\overline{\sigma\varphi_0} = 1 \times 10^{-7}$ s $^{-1}$, $\mu = 1.08$ mm $^{-1}$, and keeping the other parameters ($\dot{D}$, D_0 , r) same. CoRSEER-II generates three diagnostic plots, as shown in Fig. 3a-c.

The fitted value $a = 1.032$ shows (Fig. 3a) the presence of mild inaccurate normalisation, and the unknown age LDP was renormalised. The impressive fitting of the SCF-normalised LDP with the accepted paths shows a successful inversion, as shown in Fig. 3b. The apparent exposure age calculated as 9^{+2}_{-1} a using CoRSEER-II is comparable to 8 ± 2 a as derived by Smedley et al. (2021). Despite the significant difference in calibration parameters derived from CoRSEER-I and the literature, we observed almost the same ages. This can be explained by the fact that while performing a single calibration (Sohbati et al., 2012b), the $\overline{\sigma\varphi_0}$ and μ can covary (Fig. 5c), and based on fitting strategies, both parameters can change. The calibration profile of higher $\log(\overline{\sigma\varphi_0})$ and μ will overlap to lower $\log(\overline{\sigma\varphi_0})$ and μ for the same age sample, but the magnitude of the slope of the sigmoid will be less for lower μ (Sohbati et al., 2018). Finally, this is affected by the precision of the model and fitting strategy used to account for the slope of the curvature.

Using the kinetic parameters given earlier, we simulate the propagation of the LDP by tracking the $x_{0.5}$ with a simulated exposure time ($> 10^5$ a) for different constant erosion rate scenarios ($\dot{\varepsilon} = 10^{-4}, 10^{-3}, 10^{-2}, 10^{-1}, 10^0, 0$ mm/a), as shown in Fig. 3c. We plot two points on the no erosion $x_{0.5}$ line: (i) The $x_{0.5,natural}$ (blue filled circle in Fig. 3c) corresponds to the apparent exposure age. (ii) $x_{0.5,max}$ (red filled circle in Fig. 3c), depth of maximum exposure age ($x_{0.5,max} < x_{0.5,ss} - 1/\mu$). Thus, for a no-erosion scenario, the current sample can be dated up to an age of ~ 52 ka, which is far higher than the actual exposure age (4.52 ± 0.27 ka) of the boulder. The highly underestimated apparent exposure age of 9 ± 1 a indicates that this sample underwent significant erosion. Assuming that the LDP is in a steady-state equilibrium, for current $x_{0.5,natural}$, the erosion rates may lie between 10^{-2} and 10^0 mm/a. This represents the plausible performance and application of CoRSEER-II to derive and diagnose the apparent exposure ages and provide insight into the possible erosion rate.




Figure 3: CoRSEER-II exposure-age inversion and erosion-diagnostic modelling for rock avalanche boulder IR50 BALL02. (a) Unknown natural IR50 Lx/Tx depth profiles (hollow circles) and their sigmoidal fit (teal solid line). The fitted $a = 1.032$ shows 3.2% under normalisation, which was further used for SCF normalisation of the natural LDP. The b -value of 3.906 indicates a very steep slope. (b) The renormalised LDP (black hollow circles) and its corresponding Bayesian inversion fit resulting in the probability likelihood patch of

$\pm \delta$ exposure age LDP. Convergence towards the centre indicates an excellent fit to the model. (c) The evolution of $x_{0.5}$ for long-term exposure and constant erosion-rate ($0, 10^{-4}, 10^{-3}, 10^{-2}, 10^{-1}, 10^0$, mm/a) scenarios using $\dot{D} = 3.32 \pm 0.12$ Gy/kyr, $D_0 = 500$ Gy, $\overline{\sigma\varphi_0} = 1 \times 10^{-7}$ s $^{-1}$, $\mu = 1.08$ mm $^{-1}$ and $r = 1$. The blue-filled circle denotes the $x_{0.5, natural}$, taken as the c value of the current profile from panel (a). The red-filled marker plotted on the zero-erosion $x_{0.5}$ line indicates the no-erosion chronometer limit ($x_{0.5, max} < x_{0.5, ss} - 1/\mu$), showing that the profile could, in principle, record ages up to ~ 52 ka if no ongoing erosion of the surface occurred.

3.2.3 Erosion estimation

Here, we evaluated the erosion history of a rock surface undergoing step-function erosion. Although the natural erosion pattern is more stochastic, we prefer the step-function history for computational simplicity, as used in previous studies (Lehmann et al., 2019a,b; Smedley et al., 2021; Elkadi et al., 2022; Pathan et al., 2024). Using user-provided logarithmic spacing of exposure-erosion history, the CoRSEER-III creates two vectors: erosion rates ($\dot{\epsilon}$) and exposure durations (t). If the erosion-corrected exposure age is known by any other independent method, the t can be replaced with its value. For each ($\dot{\epsilon}$, t) pair, it simulates LDPs using a two-phase approach: Phase 1 with no erosion (from 0 to zero erosion exposure period (t_o)), and Phase 2 with constant erosion, $\dot{\epsilon}$ (for the period of t_s) ($t = t_o + t_s$, total exposure age). This forward modelling solves Equation (15) using the finite difference method and produces the final simulated LDPs. We plot $\dot{\epsilon}$ - t_s matrix and its associated P_{norm}^k and interpret the exposure-erosion conditions. Furthermore, we perform a Gaussian distribution operation on $\dot{\epsilon}$ column associated with steady erosion onset exposure age (t_{ss}). The input, mechanism, and output of CoRSEER-III are discussed in Supplementary Section S4.3.

We evaluated the exposure-erosion history of IR50_BALL02, which was suspected to face significant erosion, as discussed in Section 3.2.2. Smedley et al. (2021) estimated the steady-state erosion rate ($\dot{\epsilon}_{ss}$) and erosion onset exposure age (t_{ss}) as 0.066 mm/a and 73 a, which was evaluated using $\dot{D} = 3.32 \pm 0.12$ Gy/kyr, $D_0 = 500$ Gy, $\overline{\sigma\varphi_0} = 6.67 \times 10^{-6}$ s $^{-1}$, $\mu = 2.1$ mm $^{-1}$. If the t_{app} of a sample is much lower than the expected age as observed for IR50_BALL02, the LDP is in a steady state. To derive the erosion rates, the model should run for t_{max} . However, if the true exposure age is known by independent means, we suggest running the model for that exposure age. As the boulder was previously dated to 4.52 ± 0.27 ka, we solved Eq. 15 for t of 5 ka (t_s value varied from 0 to 5 ka in log-scale) and the erosion rate range as 10^{-6} to 10^1 mm/a (spacing in log-scale) as a 100×100 matrix using $\dot{D} = 3.32 \pm 0.12$ Gy/kyr, $D_0 = 500$ Gy, $\overline{\sigma\varphi_0} = 1 \times 10^{-7}$ s $^{-1}$, $\mu = 1.08$ mm $^{-1}$ and $r = 1$. After completion, the CoRSEER-III generates the three diagnostic plots: (1) Sigmoidal fitting (already shown in Fig. 3a), (2) the probability likelihood surface plot of $\dot{\epsilon}$ - t_s pairs as shown in Fig. 4, and (3) a Gaussian distribution of erosion rates for onset exposure age. We derive the $\dot{\epsilon}_{ss}$ as $0.107_{-0.033}^{+0.048}$ mm/a and t_{ss} as 38 a using CoRSEER-III, which are comparable to the previously established values of 0.066 mm/a and 73 a as shown in Fig. 4. This established the applicability of CoRSEER-III for predicting erosion rates.




Figure 4: (a) The normalised probability likelihood surface plot of $\dot{\epsilon}$ - t_s 10^4 pairs in log-space for the sample IR50_BALL02. The light greenish-blue shaded sloped portion points towards a mildly fitting possible transient-state solution. The yellow-shaded patch represents the possible steady-state solutions. The white-filled circle defines the minimum t_{ss} . (b) The Gaussian distribution of $\dot{\epsilon}_{ss}$ for $t_{ss} = 38$ a. The blue circles define the normalised values of probability likelihood of the erosion rate column of t_{ss} and fitted with a Gaussian distribution function. The solid, dotted and dashed black line represent the mean, $+\delta$ and $-\delta$, respectively.

4 Performance of CoRSEER

We tested all three modules of CoRSEER for the natural data available in the literature (Lehmann et al., 2019a,b; Brill et al., 2021; Smedley et al., 2021; Luo et al., 2022; Balco et al., 2023) to: (i) benchmark the app's in-built parameters for performance, accuracy, and flexibility, and (ii) to understand how CoRSEER's mathematical framework and algorithmic implementation influence the final result. We derive three sets of outputs: (i) calibration parameters ($\overline{\sigma\varphi_0}$ and μ) for the calibration samples of Lehmann et al. (2019a,b), Brill et al. (2021), Smedley et al. (2021), Luo et al. (2022), and Balco et al. (2023) using CoRSEER-I; (ii) apparent ages (t_{app}) for Smedley et al. (2021), Brill et al. (2021), Lehmann et al. (2019b), and Luo et al. (2022) using CoRSEER-II, because no usable unknown age LDP was available for the remaining datasets; and (iii) erosion rates for Smedley et al. (2021) and Lehmann et al. (2019b), using CoRSEER-III, because both studies have a well-established dataset of erosion history. The complete methodology for deriving the required parameters is provided in Section 3. We must note that all operations were performed using the FOK model ($r = 1$), as the literature data were previously analysed using FOK. However, CoRSEER has the flexibility to use $r > 1$, which is important when luminescence depth profiles are derived using feldspar IRSI, which follows non-FOK behaviour.

4.1 Calibration

Across the published calibration datasets, CoRSEER-I generally reproduces literature calibration parameters, with the best agreement observed for Lehmann et al. (2019a) and Luo et al. (2022), where the CoRSEER-I-derived $\log_{10}(\overline{\sigma\varphi_0})$ and μ values are broadly comparable to reported estimates. This agreement is consistent with a similar normalisation choice for the L_x/T_x depth profiles. In contrast, several datasets show systematic offsets between CoRSEER-I and the literature values. For example, in Balco et al. (2023), the sample IR50_KAY_110, the calibration parameters shift to higher values, from $\log_{10}(\overline{\sigma\varphi_0}) = -7.30$ and $\mu = 0.34$ mm⁻¹ to -6.58 and 0.51 mm⁻¹ whereas for the sample pIRIR225_KAY_110 shifts to lower values, from -7.68 and 0.54 mm⁻¹ to -7.77 and 0.36 mm⁻¹. In Brill et al. (2021), CoRSEER-I predicted decreased calibration parameters for IR50_HAR1-1_CAL but increased parameters for IR50_TEM3-1_CAL and IR50_VAL4-1_CAL1. For the two calibration samples for which Brill et al. (2021) did not report the final parameters (IR50_RAB5-1_CAL and IR50_VAL4-1_CAL2), CoRSEER-I yielded -7.38 , 0.76 mm⁻¹ and -7.50 , 0.37 mm⁻¹, respectively. In Smedley et al. (2021), the ROAD set was dominated by a decrease in both parameters, with two clear increased cases (IR50_ROAD01 and IR50_ROAD03). The Lehmann et al. (2018) dataset shows predominantly decreased parameters for MBMB samples, with one increased case (IR50_MBMV7) and one near-balanced case (IR50_MBMV1). We attribute these discrepancies primarily to (1) differences in the fitting strategy, (2) differences in the L_x/T_x normalisation (manual plateau selection versus automated normalisation) and (3) both the $\overline{\sigma\varphi_0}$ and μ may slightly covary during optimisation of the best fitting. The boundary ranges of $\log_{10}(\overline{\sigma\varphi_0})$ and μ values predicted by CoRSEER-I are -8.41 to -4.82 and 0.36 to 3.36 mm⁻¹ (Table 3), showing the plausible flexibility of the app for a wide variety of exposure conditions and lithologies.

We plotted CoRSEER-I vs the literature dataset and evaluated the R^2 value with respect to the 1:1 line to summarise the compilation-scale performance for 23 samples. In Fig. 5a, μ_{CoRSEER} versus $\mu_{\text{literature}}$ gives $R^2_{1:1} = 0.562$ for $n = 23$, meaning CoRSEER-I tracks published μ values moderately well, even though a large fraction of points falls below the 1:1 line. In Fig. 5b, $\overline{\sigma\varphi_0}_{\text{CoRSEER}}$ versus $\overline{\sigma\varphi_0}_{\text{literature}}$ gives $R^2_{1:1} = 0.316$ for $n = 23$, showing weaker agreement and stronger scatter. It is well established that the $\overline{\sigma\varphi_0}$ and μ covary, and we present the relationship between them in Supplementary Section S5. To investigate the offset between the established calibration parameters and the underlying reason behind it, we plot $(\log_{10}(\overline{\sigma\varphi_0})/\mu)_{\text{CoRSEER}}$ vs $(\log_{10}(\overline{\sigma\varphi_0})/\mu)_{\text{literature}}$ in Fig. 5c which gives $R^2_{1:1} = 0.379$ showing weaker agreement. This suggests that covariance contributes only weakly to the offset, whereas normalisation and fitting strategy are the dominant contributors. Visually, most points fall below the 1:1 line in both panels, so the dominant overall tendency in this compilation is that CoRSEER estimates lower μ and $\overline{\sigma\varphi_0}$ than those reported in the literature for many samples, while a smaller set of samples shows the opposite paired behaviour.




Figure 5: The 1:1 comparison between CoRSEER-I and literature calibration parameters. The round filled circles show the median/best fitting values, and the solid lines represent the respective $\pm \delta$ values, which are plotted along with the black dotted 1:1 correlation line. (a) Attenuation coefficients (μ) (b) bleaching rates ($\sigma\phi_0$) and (c) the ratio $\log_{10}(\sigma\phi_0)/\mu$ for all compiled calibration samples ($n = 23$).

Table 3: Comparison of model parameters of samples in the selected studies and estimated using CoRSEER-I

Author	Samples	D (Gy /kyr)	D_o (Gy)	Location	Rock type	Sampling type	Known age (a)	$\log_{10}(\sigma\bar{\Phi}_0, \text{CoRSEER})$			$\log_{10}(\sigma\bar{\Phi}_0, \text{literature})$			$\mu_{\text{CoRSEER}} (\text{mm}^2)$			$\mu_{\text{literature}} (\text{mm}^2)$		
								-1 σ	BF/M*	+1 σ	-1 σ	BF/M	+1 σ	-1 σ	BF/M	+1 σ	-1 σ	BF/M	+1 σ
Balco et al. (2023)	IR50_KAY_110	-	-	Kay Peak Ridge, West Antarctica	Biotite gneiss	Surface bedrock in the ice-sheet-free portion	4	-	-	-	-	-	-	-	-	-	-	-	-
	pIRIR225_KAY_110	-	-					6.97	-6.58	6.21	*	-7.30	*	0.40	0.51	0.61	*	0.34	*
Brill et al. (2021)	IR50_HAR 1-1 CAL	0.53	500	Harhoura, Morocco	Calcarenite, carbonate limestone	Coastal boulders	2.15	-	-	-	-	-	-	-	-	-	-	-	-
	IR50_RAB 5-1 CAL	-		Rabat, Morocco				7.02	-6.59	6.20	-6.64	-6.52	-6.43	0.87	1.25	1.60	1.24	1.39	1.54
	IR50_TEM 3-1 CAL	0.51		Temara, Morocco				7.60	-7.38	7.16	*	*	*	0.52	0.76	0.97	*	*	*
	IR50_VAL 4-1 CAL 1	-		Val d'Or, Morocco				4.82	-4.54	4.30	-5.68	-5.58	-5.50	1.88	2.04	2.18	1.24	1.39	1.54
	IR50_VAL 4-1 CAL 2	-		Val d'Or Morocco				6.50	-6.16	5.84	-6.64	-6.52	-6.43	1.29	1.59	1.85	1.24	1.39	1.54
								7.57	-7.50	7.44	*	*	*	0.33	0.37	0.41	*	*	*
	IR50_MBMV1	-	-	Mer de Glace, Mont Blanc Massif, France	Gneiss	Bedrock outcrop	137	7.94	-7.80	7.60	-7.89	-7.85	-7.82	0.95	1.04	1.13	1.05	1.07	1.08
	IR50_MBMV6	-	-		Gneiss		2	6.33	-6.13	5.89	-5.74	-5.70	-5.66	0.71	0.79	0.89	0.92	0.92	0.95
	IR50_MBMV7	-	-		Granite		69	5.32	-4.98	4.64	-5.44	-5.40	-5.30	2.55	2.72	2.95	2.46	2.52	2.56
	IR50_MBMV8	-	-		Gneiss		3	7.50	-7.34	7.18	-6.82	-6.66	-6.62	0.89	1.07	1.28	1.70	1.89	1.98
	IR50_MBMV10	-	-		Gneiss		18	7.43	-7.30	7.14	-6.96	-6.82	-6.82	1.20	1.31	1.42	1.75	1.82	1.87
	IR50_MBMV11	-	-		Gneiss		30	7.72	-7.57	7.40	-7.41	-7.38	-7.28	0.96	1.05	1.15	1.13	1.21	1.22
Lehmann et al. (2019a)	IR50_MBTP7_CAL	-	-		Granite	Bedrock	2	5.57	-5.33	5.00	-5.71	-5.65	-4.84	0.55	0.61	0.68	0.51	0.53	0.69
	IR50_MBTP8_CAL	-	-		Granite		11	6.40	-6.19	5.94	-6.07	-5.92	-3.53	0.44	0.48	0.53	0.53	0.53	0.97
Luo et al. (2022)	pIRIR175_LS_IK_CAL	3.3	506	Langshan Normal Fault, Inner Mongolia, China	Granite	Fault scarp	3	7.13	-7.05	6.94	7.081	7.078	7.076	1.06	1.16	1.25	0.91	1.10	1.29
Smedley et al. (2021)	IR50_ROAD01	4.81	-	NW Scotland, roadcut	Torridonian sandstone	In situ bedrock	0.01	3.72	-3.37	3.09	-4.33	-3.55	-3.08	2.82	3.36	3.78	2.50	3.20	3.80
	IR50_ROAD02	4.76	-				57	7.40	-6.88	6.36	-6.46	-5.18	-3.90	0.91	1.14	1.36	1.40	2.10	2.60
	IR50_ROAD03	4.00	-				44	5.15	-4.04	2.88	-5.83	-4.81	-3.79	2.32	2.95	3.66	2.00	2.70	3.20
	IR150_ROAD01	4.81	-				0.01	5.20	-5.17	5.13	-4.67	-4.49	-3.94	1.60	1.74	1.89	2.20	3.10	3.70
	IR150_ROAD02	4.76	-				57	8.02	-7.77	7.49	-8.06	-7.76	-7.16	1.32	1.55	1.81	1.50	1.90	2.30
	IR150_ROAD03	4.00	-				44	8.21	-8.01	7.84	-7.95	-7.42	-7.36	0.96	1.11	1.28	1.30	1.90	2.50
	IR225_ROAD01	4.81	-				0.01	5.16	-5.11	5.06	-4.82	-4.54	-4.40	1.97	2.16	2.40	2.30	3.00	3.60
	IR225_ROAD02	4.76	-				57	7.70	-7.45	7.11	-7.64	-7.05	-6.26	1.94	2.24	2.61	1.80	2.80	3.60
	IR225_ROAD03	4.00	-				44	8.51	-8.41	8.27	-8.33	-7.77	-6.93	0.75	0.84	0.96	0.90	1.40	2.50

4.2 Exposure age

We evaluated the apparent exposure ages (t_{app}) of the unknown age LDPs of the literature dataset using CoRSEER-II for 41 samples. As the evaluated age exclusively depends on the chosen calibration samples, dose rate, and characteristic dose, we follow the same pattern as given in the literature dataset (unknown-age LDPs), except for a few, as mentioned further. Lehmann et al. (2019b) used a combined calibration parameter of IR50_MBTP7_CAL (2 a) and IR50_MBTP8_CAL (11 a), whereas the CoRSEER-I supports only a single calibration sample. Thus, we used IR50_MBTP7_CAL, as it closely resembles the calibration parameters in the literature. For Brill et al. (2021), we used CoRSEER-I to evaluate the parameters of IR50_RAB 5-1 CAL for the samples of IR50_RAB 1-1, IR50_RAB 1-2, and IR50_RAB 5-1 (these samples were evaluated using IR50_HAR 1-1 CAL in the literature due to the large uncertainty in IR50_RAB 5-1 CAL). We plotted the CoRSEER-II derived versus published literature apparent age dataset and evaluated the $R^2_{1:1}$ value with respect to the 1:1 line to summarise the compilation-scale performance. Interestingly, the evaluated t_{app} using CoRSEER-II is comparable to the literature and falls closely around the 1:1 line (Fig. 6) despite a significant difference in the calibration parameters used in this study. The derived $R^2_{1:1} = 0.889$ represents the CoRSEER-II track, which has apparent exposure ages with high certainty. There was a significant underestimation by CoRSEER-II for the IR50_RAB 1-1, IR50_RAB 5-1 and IR50_TEM 2-1. This is due to the following reasons: (1) a change in the calibration sample, as mentioned

earlier. (2) The high SCF normalisation factors $a = 1.1$ (up to 10%) of the sigmoidal fitting for the IR50_TEM 2-1, which might have changed the position of $x_{0.5}$.

The range of t_{app} values predicted by CoRSEER-II extends from 1 a to 8.5 ka (Table 4), showing its potential as a decadal-to-millennial chronometric mathematical tool for LRSed.




Figure 6: The 1:1 comparison between CoRSEER-II and literature t_{app} data. The round filled circles show the median/best fitting values, and the solid lines represent the respective $\pm \delta$ values, which are plotted along with the black dotted 1:1 correlation line.

Table 4: Representation of exposure ages for samples evaluated from both the literature and CoRSEER.

Author	Unknown age samples	t_{app} literature (a)			t_{app} CoRSEER (a)		
		-1δ	BF/M	$+1\delta$	-1δ	BF/M	$+1\delta$
Brill et al. (2021)	IR50 HAR 1-1	5	5	5	4	4	4
	IR50 HAR 2-1	11	17	23	8	9	11
	IR50 HAR 3-1	25	33	41	15	18	21
	IR50 RAB 1-1	141	162	183	23	26	30
	IR50 RAB 1-2	9	12	15	8	9	10
	IR50 RAB 5-1	1022	1180	1338	59	74	93
	IR50 TEM 2-1	14	18	22	3	4	4
	IR50 TEM 3-1	1	1	2	2	2	3
	IR50 TEM 4-1	1	1	2	1	2	2
	IR50 VAL 1-1	15	22	29	26	33	42
	IR50 VAL 1-2	7	11	15	9	11	13
	IR50 VAL 4-1	44	68	92	106	130	162
	IR50 VAL 6-1	25	33	41	32	37	42
	IR50 VAL 6-1	25	33	41	32	37	42
Lehmann et al. (2019b)	IR50 MBAM1	26	34	42	31	36	43
	IR50 MBAM2	51	67	83	61	64	67
	IR50 MBAM3	10	12	14	12	13	13
	IR50 MBTP1	471	618	765	690	813	962
	IR50 MBTP2	462	606	750	537	612	697
	IR50 MBTP5	4	5	5	14	15	16
	IR50 MBTP9	40	54	68	49	54	60
	IR50 MBTP11	3	3	4	4	4	5
Luo et al. (2022)	pIRIR175 LS0 10	13	13	13	12	13	14
	pIRIR175 LS30 40	14	15	16	16	17	18
	pIRIR175 LS70 80	31	34	37	37	39	41
	pIRIR175 LS90 100	32	35	38	39	41	43
	pIRIR175 LS140 150	5	6	7	7	8	9
	pIRIR175 LS160 170	69	84	99	81	90	98
	pIRIR175 LS210 220	21	25	29	26	30	34
	pIRIR175 LS290 300	17	19	21	21	21	22
	pIRIR175 LS310 320	600	700	800	881	978	1082
	pIRIR175 LS370 380	800	900	1000	951	1003	1057
	pIRIR175 LS390 400	400	500	600	533	581	635
	pIRIR175 LS420 430	4200	5000	5800	7591	8503	9508
	pIRIR175 LS450 460	3000	3500	4000	3800	4299	4866
	pIRIR175 LS590 600	1700	2000	2300	3520	3745	3985
Smedley et al. (2021)	IR50 BALL02	6	8	10	8	9	11
	IR50 BALL03	284	387	490	1165	1174	1183
	IR150 BALL02	50	66	82	77	85	92
	IR150 BALL03	242	296	350	303	337	374
	IR225 BALL02	233	263	293	149	162	176
	IR225 BALL03	313	362	411	201	223	246

4.3 Erosion rates

Here, we establish the steady-state erosion rate ($\dot{\epsilon}_{ss}$) and corresponding onset time of erosion (t_{ss}) of literature dataset ($n = 11$) using CoRSEER-III. We carefully chose the calibration and dose parameters, as mentioned in the earlier sections. The choice of sample for analysis assumes that these samples are in a steady state, as described in the literature (Lehmann et al., 2019a; Smedley et al., 2021). Comparison of the literature and CoRSEER-derived erosion rates is important because, unlike the previous studies, CoRSEER-III derives erosion rates directly from the $\dot{\epsilon}_{ss} - t_{ss}$ matrix. Therefore, the present result is not only a comparison with previously reported values, but also the first systematic demonstration of an algorithmic method for extracting erosion-rate solutions from this matrix. The results are shown in Table 5.

The CoRSEER-III-derived erosion rates broadly agree with the literature erosion-rate values. The literature erosion rates range from 0.0035 to 4.3 mm/a, whereas the CoRSEER-III-derived erosion rates range from 0.003 to 4.327 mm/a. Using the literature values as reference, the absolute percentage difference ranges from 0.6% to 86.7%. The minimum difference occurs for IR50_MBTP6, where CoRSEER-III gives 4.327 mm/a compared with the literature value of 4.3 mm/a. The largest difference occurs for IR50_MBTP5, where CoRSEER-III gives 0.560 mm/a compared with the literature value of 0.300 mm/a.

The uncertainty associated with CoRSEER-III-derived erosion rates remains within a reasonable range for most samples. The relative upper error in the CoRSEER-III erosion rates varies from 10.8% to 44.9%, whereas the relative lower error varies from 10.8% to 33.3%. The lowest relative uncertainty occurs for IR50_MBAM2, where the erosion rate is 0.037 mm/a with an uncertainty of approximately $\pm 10.8\%$. The highest upper uncertainty occurs for IR50_BALL02, where the upper error reaches approximately 44.9%. These values show that the algorithm provides bounded erosion-rate estimates rather than single manually selected solutions. In contrast, the erosion-onset time, t_{ss} , shows a considerably larger departure from the literature values. The literature t_{ss} values range from 4 to 3146 a, whereas the CoRSEER-III-derived t_{ss} values range from 3 to 3080 a. Using the literature values as reference, the absolute percentage difference ranges from 18.0% to 64.1%. In most samples, CoRSEER-III estimates a younger erosion-onset time than the literature value. The only exception is IR50_MBTP1, where CoRSEER-III gives a larger onset time of 3080 a compared with the literature value of 2610 a, corresponding to an increase of approximately 18.0%. The maximum difference occurs for IR50_MBTP5, where CoRSEER-III estimates 23 a compared with the literature value of 64 a, corresponding to a reduction of approximately 64.1%.

Overall, the results indicate that CoRSEER-III reproduces the erosion-rate field reasonably well, as presented in Table 5, but it estimates erosion-onset time differently from the literature. This difference is expected because the CoRSEER-III estimate is derived from an automated best-fitting erosion-time slope detection algorithm, whereas previous studies relied more strongly on manual or interpretative selection of model solutions.

When the sample is in a transient state, one can visually identify the solution that generally lies in the sloped portion of the $\dot{\epsilon}_{ss}$ vs t_{ss} likelihood plot. It is mathematically challenging to produce an objective method to constrain a pair of possible solutions. Thus, we do not present any transient-state condition scenarios. But it is possible to derive a transient state solution from the results of CoRSEER-III if the onset of erosion (t_s) is known by some means.

Table 5: Representation of steady-state onset exposure age and erosion rates for samples evaluated from both the literature and CoRSEER-III.

Author	Samples	$\dot{\epsilon}_{ss, literature} (mm/a)$	$\dot{\epsilon}_{ss, CoRSEER-III} (mm/a)$	$t_{ss, literature} (a)$	$t_{ss, CoRSEER-III} (a)$
Lehmann et al. (2019b)	IR50_MBAM1	0.057	$0.068^{+0.027}_{-0.019}$	285	178
	IR50_MBAM2	0.028	$0.037^{+0.004}_{-0.004}$	461	354
	IR50_MBAM3	0.2	$0.277^{+0.059}_{-0.048}$	56	40
	IR50_MBTP1	0.0035	$0.003^{+0.001}_{-0.001}$	2610	3080
	IR50_MBTP2	0.0035	$0.005^{+0.001}_{-0.001}$	3146	2466
	IR50_MBTP5	0.3	$0.560^{+0.227}_{-0.161}$	64	23
	IR50_MBTP6	4.3	$4.327^{+0.645}_{-0.561}$	4	3
	IR50_MBTP9	0.032	$0.048^{+0.007}_{-0.006}$	486	206
	IR50_MBTP11	0.53	$0.618^{+0.079}_{-0.070}$	42	23
Smedley et al. (2021)	IR50_BALL02	0.066	$0.107^{+0.048}_{-0.033}$	73	38
	IR150_BALL02	0.009	$0.012^{+0.003}_{-0.002}$	593	327

5 Discussion

The LRSED is a promising chronometer of geological processes which occur under sufficient sunlight. Rigorous methodological development and precise field application of LRSED can cause a paradigm shift in surface exposure chronometry by enabling us to date decadal to millennial exposure events. Furthermore, it can accurately establish steady-state in situ erosion rates. Although this method is affected by broader challenges such as the availability of calibration samples, representative luminescence depth-profile selection, lithological suitability, and independent exposure-age constraints, additional methodological uncertainty can arise from subjective plateau selection and the use of non-standardised modelling approaches. Thus, CoRSEER attempts to improve the application, standardisation, and intercomparison of results among several studies.

5.1 Preliminary precautions

Large scatter in the luminescence depth profile (LDP) and high fractional uncertainty in L_x/T_x make plateau identification unreliable and reduce the stability of any inversion (i.e., the SCF normalisation, estimation of calibration parameters, exposure age, erosion rates). We can reduce L_x/T_x uncertainty by adopting a modified post-infrared infrared stimulated luminescence protocol with lower heating rates and longer isothermal holding times (Jenkins et al., 2018), and by using metal cups (Elkadi et al., 2021). For each sample, we recommend measuring two or more LDPs and maintain millimetre-scale sampling density with a substantial fraction of points located on the sigmoidal transition. Importantly, the fractional error structure should be similar or random with depth; if the fractional error at lower depths is systematically higher than at higher depths, the fitting becomes biased toward the plateau, and the inferred parameters become unreliable. Although we cannot yet provide a statistical criterion for the required fraction of points on the sigmoidal part of the curve, based on our observation in Fig. 2a and Fig. 3a, we recommend that more than 50% of all L_x/T_x measurement points should lie before the final plateau. If the measured profile is dominated by plateau points, we recommend reducing the measurement depth such that the total number of points on the initial plateau and sigmoidal region exceeds 50% of the total.

In the case of incomplete LDP, the maximum measurement depth is shallower than the steady-state depth, the plateau cannot be reconstructed. We therefore recommend measuring L_x/T_x up to the equilibrium depth (which is the stable plateau). Similarly, for the large scatter in experimental data, it is difficult to identify the plateau manually. The SCF and parameter estimation to normalise the LDP is a major step forward as it resolves both issues and removes the manual intervention from mathematical inversion, thereby improving the accuracy of fitting (Fig. 2e).

5.2 Calibration

While using CoRSEER-I, we suggest that there are a few precautions we must take to estimate $\overline{\sigma\varphi_0}$ and μ accurately. If the IRSL signal from feldspar minerals is used for LDP measurements, we suggest evaluating the order of kinetics (r) beforehand and incorporating it into the calculation. The application of FOK can cause severe underestimation and overestimation of the exposure age and erosion rate (Biswas et al., 2023; Pathan et al., 2024). The different methods for evaluating the r value are thoroughly discussed in Biswas et al. (2023) and Pathan et al. (2024). Furthermore, if the calibration sample is young (say 1-10 a), for example, a known-age deglaciated surface or manually exposed surface created by chipping off a boulder face, erosion is assumed to have no effect on the LDP. In contrast, when using older natural calibration samples ($t > 100$ a), erosion is expected to affect LDP significantly (Lehmann et al., 2018; Smedley et al., 2021), and this effect becomes stronger in high-erosion environments. This can lead to substantial underestimation of $\overline{\sigma\varphi_0}$ and smaller bias in μ . This is inferred from the samples IR50_MBMV1 ($t = 137$ a) and IR50_MBMV6 ($t = 2$ a), which were samples from the same site with similar lithology, but the $\log \overline{\sigma\varphi_0}$ was estimated as -7.8 and -6.13, respectively, which is a difference of about two orders of magnitude, whereas μ was estimated at 1.04 and 0.8 mm^{-1} . Thus, the erosion rate must be explicitly incorporated into CoRSEER-I. To avoid this complexity and reduce model dependence on uncertain erosion histories, it is recommended to use young calibration surfaces at sampling sites where there is no erosion over the period of interest. This allows for the precise calibration of kinetic parameters under well-constrained exposure conditions. This will further remove any ambiguity due to changes in decay kinetics, lithology, surface orientation, shielding, and light penetration.

When selecting a calibration sample, it is critical to recognise that the light attenuation coefficient (μ) and the effective detrapping rate at the surface ($\overline{\sigma\varphi_0}$) of the calibration samples are the best representatives of the actual sample. Recent empirical studies demonstrate that photon penetration and attenuation are highly sensitive to micro-mineralogical heterogeneity, such as the spatial distribution of opaque and translucent minerals, which can cause localised 'light piping' or 'shadowing' (Meyer et al., 2018; Cui et al., 2024). Furthermore, μ is strongly wavelength-dependent and can be drastically altered over time by dynamic surficial boundary conditions, including the accretion of rock varnish (Luo et al., 2018) or localised iron-hydroxide precipitation due to moisture (Meyer et al., 2018; Smedley et al., 2021). The effective detrapping rate ($\overline{\sigma\varphi_0}$) is similarly sensitive to external environmental

geometries, fluctuating significantly based on the target surface's topographic aspect, inclination, and seasonal solar incidence angles (Fuhrmann et al., 2022). High-resolution spatial analyses reveal that these optical properties can vary significantly even on a sub-millimetre scale within contiguous rock formations (Ou et al., 2018; Smedley et al., 2025). Consequently, to avoid severe systematic chronological errors, it is imperative to ensure that the optical characteristics, dynamic surficial coatings, and spatial orientation governing μ and $\overline{\sigma\varphi_0}$ in the calibration sample, the measurements are meticulously matched to those of the unknown target sample. Although our present modelling framework focuses on single calibration (SC), it is worth noting that multi-calibration strategies (MC) can be highly effective under certain conditions. When multiple calibration samples of different known ages, but from the same lithology and sunlight exposure conditions, are available, it becomes possible to constrain $\overline{\sigma\varphi_0}$, μ and r simultaneously (Pathan et al., 2024). However, the accuracy of such multi-calibration depends strongly on the number of calibration samples and their age spread. A wider temporal distribution across well-characterised surfaces enhances the resolution of the kinetic behaviour. While we recognise the potential of this methodology, it lies beyond the scope of the current work and is not implemented in the present version of CoRSEER.

5.3 Apparent age estimation

After estimation of kinetic parameters, the estimation of exposure and erosion history appears to be highly dependent on the position of $x_{0.5}$. There are two possible scenarios:

Case 1: For non-eroding exposure conditions, if the depth of half saturation of the LDP, $x_{0.5}$ is smaller than $x_{0.5,max}$, then the apparent exposure age is equal to the actual exposure age of the sample, and if $x_{0.5,max} < x_{0.5} \leq x_{0.5,ss}$ then the sample reaches steady-state, and the exposure age is greater than the t_{max} , maximum dateable exposure age.

Case 2: If the derived apparent ages are substantially lower than the expected age (derived using an independent method), this indicates the presence of erosion, which was seen in the sample IR50_BALL02 (Smedley et al., 2021). CoRSEER-III should be used to further re-establish the exposure-erosion history. Furthermore, when using IRSL to establish the LDP, the r value should be the same for calibration and unknown samples (Pathan et al., 2024). We incorporated all these modelling aspects, diagnostics, and interpretation into the input and output GUI plots (Fig. 3).

5.4 Erosion rate estimation

The fundamental assumptions and modelling aspects underpinning CoRSEER-III were taken from Lehmann et al. (2019a), where the authors coupled ^{10}Be concentration and the LDP to evaluate the erosion-corrected ^{10}Be exposure age range. The main difference is that CoRSEER-III does not incorporate ^{10}Be concentration; rather, it incorporates independent exposure age (i) which can be derived from an independent dating method or (ii) by incorporating t_{max} from forward modelling of $x_{0.5}$ vs t .

One of the most important observations is that CoRSEER-III captures the same broad erosion-rate scale as the literature values. This indicates that the erosion-rate extraction algorithm is able to preserve the first-order structure of the erosion solution space. In practical terms, this is encouraging because the algorithm avoids subjective manual picking while still recovering erosion-rate estimates comparable to previous studies. This is also a possible reason for the large relative difference between CoRSEER and literature results in a few cases. Thus, for maximum accuracy and precision, we recommend following the guidelines. Users should not run the model once over a very large erosion-exposure window and treat the result as final. A better strategy is to run the model in multiple stages. First, use a broad search window to identify the approximate high-likelihood region. Then shrink the exposure-erosion window around that region. Finally, increase the matrix density so that the best-fitting slope is resolved with higher numerical precision. This stepwise refinement should reduce grid artefacts and improve the stability of both erosion-rate (and errors) and onset-time estimates.

6 Conclusion

In this study, we present CoRSEER as an integrated, reproducible workflow for LRSED that converts the LDP into physically interpretable exposure and erosion information. CoRSEER reduces user-dependent choices by automating key steps that vary across studies. CoRSEER comprises three linked modules: (i) calibration of site- and lithology-specific bleaching parameters, (ii) apparent exposure age estimation, and (iii) erosion rate estimation under steady-state scenarios. Reanalysis of published datasets shows that CoRSEER-I generally reproduces plausible calibration parameter estimates and captures literature trends, while also presenting systematic offsets in some datasets that mainly arise from differences in normalisation and fitting strategy. Despite these calibration differences, CoRSEER-II yields apparent exposure ages that closely track published values with strong 1:1 agreement, demonstrating that the platform provides robust chronometric outputs when calibration inputs are selected consistently with the reference studies. Finally, CoRSEER-III reproduces steady-state erosion histories with excellent agreement for well-constrained erosion datasets, supporting its use as a quantitative erosion-rate estimator when exposure age constraints are available, and the profile indicates steady-state behaviour. These results support the central conclusion that objective normalisation and standardised inversion materially improve reproducibility in LRSED without reducing model flexibility. CoRSEER enables consistent decadal-to-millennial exposure reconstructions and strengthens its practical use, but there are a few limitations which must be addressed. The CoRSEER's performance depends on data quality; that is, a significant number of L_x/T_x samples across the sigmoidal slope and lesser L_x/T_x error estimates remain essential, and shallow or plateau-dominated profiles can bias sigmoidal fitting. The current release prioritises single-sample calibration and does not yet implement multi-calibration optimisation, and the objective selection of transient-state erosion solutions remains mathematically challenging and therefore user-guided. Future versions should (i) implement multi-calibration strategies to constrain kinetic parameters jointly, (ii) expand diagnostics that flag profiles likely to be poorly constrained, and (iii) develop a similar GUI framework for luminescence rock surface burial dating.

7 Code availability

CoRSEER (Calculator of Rock Surface Exposure Age and Erosion Rates) was released as open-source software and archived on Zenodo. The exact version used in this study is CoRSEER v7 (Zenodo version DOI: <https://doi.org/10.5281/zenodo.20796442>). The source repository is hosted on GitHub at <https://github.com/arbazNP/CoRSEERv1>.

8 Video Supplement

The explanatory video is posted on TIB AV-Portal as <https://doi.org/10.5446/72689>.

9 Author contributions

AP contributed to conceptualisation, data curation, formal analysis, investigation, methodology, project administration, software, validation, visualisation, writing of the original draft, and review and editing of the manuscript. RHB contributed to conceptualisation, methodology, project administration, resources, supervision, writing of the original draft, and review and editing of the manuscript. DK contributed to the resources and to the review and editing of the manuscript.

10 Competing interests

The authors declare no conflicts of interest.

11 Acknowledgements:

The authors thank the editor and reviewer for their constructive comments, which helped improve the clarity and presentation of the manuscript. Arbaz and Rabiul acknowledges the Department of Earth Sciences, Indian Institute of Technology Kanpur, for providing institutional and academic support during this work.

12 Financial Support:

This research received no external funding.

13 References

- Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J.: A complete and easily accessible means of calculating surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements, *Quaternary Geochronology*, 3, 174–195, <https://doi.org/10.1016/j.quageo.2007.12.001>, 2008.
- Balco, G., Brown, N., Nichols, K., Venturelli, R. A., Adams, J., Braddock, S., Campbell, S., Goehring, B., Johnson, J. S., Rood, D. H., Wilcken, K., Hall, B., and Woodward, J.: Reversible ice sheet thinning in the Amundsen Sea Embayment during the Late Holocene, *The Cryosphere*, 17, 1787–1801, <https://doi.org/10.5194/tc-17-1787-2023>, 2023.
- Biswas, R. H., Pathan, A. N., and Malik, J. N.: General-order kinetics model for OSL rock surface exposure dating, *Proceedings of the Indian National Science Academy*, 89, 644–654, <https://doi.org/10.1007/s43538-023-00172-y>, 2023.
- Brill, D., May, S. M., Mhammedi, N., King, G., Lehmann, B., Burow, C., Wolf, D., Zander, A., and Brückner, H.: Evaluating optically stimulated luminescence rock surface exposure dating as a novel approach for reconstructing coastal boulder movement on decadal to centennial timescales, *Earth Surface Dynamics*, 9, 205–234, <https://doi.org/10.5194/esurf-9-205-2021>, 2021.
- Brown, N. D. and Moon, S.: Revisiting erosion rate estimates from luminescence profiles in exposed bedrock surfaces using stochastic erosion simulations, *Earth and Planetary Science Letters*, 528, 115842, <https://doi.org/10.1016/j.epsl.2019.115842>, 2019.
- Burbank, D. W., Leland, J., Fielding, E., Anderson, R. S., Brozovic, N., Reid, M. R., and Duncan, C.: Bedrock incision, rock uplift and threshold hillslopes in the northwestern Himalayas, *Nature*, 379, 505–510, <https://doi.org/10.1038/379505a0>, 1996.
- Chen, R. and Leung, P. L.: The decay of OSL signals as stretched-exponential functions, *Radiation Measurements*, 37, 519–526, [https://doi.org/10.1016/S1350-4487\(03\)00007-6](https://doi.org/10.1016/S1350-4487(03)00007-6), 2003.
- Colman, S. M.: Rock-weathering rates as functions of time, *Quaternary Research*, 15, 250–264, [https://doi.org/10.1016/0033-5894\(81\)90029-6](https://doi.org/10.1016/0033-5894(81)90029-6), 1981.
- Cui, F., Kook, M., Murray, A. S., Qin, J., Liu, J., and Jain, M.: Do attenuation coefficients based on luminescence bleaching fronts reflect true light attenuation in rocks?, *Radiation Measurements*, 174, 107129, <https://doi.org/10.1016/j.radmeas.2024.107129>, 2024.
- Desilets, D. and Zreda, M.: Spatial and temporal distribution of secondary cosmic-ray nucleon intensities and applications to in situ cosmogenic dating, *Earth and Planetary Science Letters*, 206, 21–42, [https://doi.org/10.1016/S0012-821X\(02\)01088-9](https://doi.org/10.1016/S0012-821X(02)01088-9), 2003.
- Duller, G. A. T.: Behavioural studies of stimulated luminescence from feldspars, *Radiation Measurements*, 27, 663–694, [https://doi.org/10.1016/S1350-4487\(97\)00216-3](https://doi.org/10.1016/S1350-4487(97)00216-3), 1997.
- Dunai, T. J.: Scaling factors for production rates of in situ produced cosmogenic nuclides: a critical reevaluation, *Earth and Planetary Science Letters*, 176, 157–169, [https://doi.org/10.1016/S0012-821X\(99\)00310-6](https://doi.org/10.1016/S0012-821X(99)00310-6), 2000.
- Dunai, T. J.: *Cosmogenic nuclides: principles, concepts and applications in the earth surface sciences*, Cambridge University Press, Cambridge, 2010.
- Durcan, J. A., King, G. E., and Duller, G. A.: DRAC: Dose Rate and Age Calculator for trapped charge dating, *Quaternary Geochronology*, 28, 54–61, <https://doi.org/10.1016/j.quageo.2015.03.012>, 2015.
- Elkadi, J., King, G. E., Lehmann, B., and Herman, F.: Reducing variability in OSL rock surface dating profiles, *Quaternary Geochronology*, 64, 101169, <https://doi.org/10.1016/j.quageo.2021.101169>, 2021.
- Elkadi, J., Lehmann, B., King, G. E., Steinemann, O., Ivy-Ochs, S., Christl, M., and Herman, F.: Quantification of post-glacier bedrock surface erosion in the European Alps using ^{10}Be and optically stimulated luminescence exposure dating, *Earth Surface Dynamics*, 10, 909–928, <https://doi.org/10.5194/esurf-10-909-2022>, 2022.
- Freiesleben, T. H., Thomsen, K. J., and Jain, M.: Novel luminescence kinetic models for rock surface exposure dating, *Radiation Measurements*, 160, 106877, <https://doi.org/10.1016/j.radmeas.2022.106877>, 2023.
- Fuhrmann, S., Meyer, M. C., Gliganic, L. A., and Obleitner, F.: Testing the effects of aspect and total insolation on luminescence depth profiles for rock surface exposure dating, *Radiation Measurements*, 153, 106732, <https://doi.org/10.1016/j.radmeas.2022.106732>, 2022.
- Ganti, V., Von Hagke, C., Scherler, D., Lamb, M. P., Fischer, W. W., and Avouac, J.-P.: Time scale bias in erosion rates of glaciated landscapes, *Science Advances*, 2, e1600204, <https://doi.org/10.1126/sciadv.1600204>, 2016.
- Gliganic, L. A., Meyer, M. C., Sohhati, R., Jain, M., and Barrett, S.: OSL surface exposure dating of a lithic quarry in Tibet: Laboratory validation and application, *Quaternary Geochronology*, 49, 199–204, <https://doi.org/10.1016/j.quageo.2018.04.012>, 2019.

580 Guralnik, B., Li, B., Jain, M., Chen, R., Paris, R. B., Murray, A. S., Li, S.-H., Pagonis, V., Valla, P. G., and Herman, F.: Radiation-
581 induced growth and isothermal decay of infrared-stimulated luminescence from feldspar, *Radiation Measurements*, 81, 224–231,
582 <https://doi.org/10.1016/j.radmeas.2015.02.011>, 2015.

583 Jain, M., Sohbati, R., Guralnik, B., Murray, A. S., Kook, M., Lapp, T., Prasad, A. K., Thomsen, K. J., and Buylaert, J. P.: Kinetics
584 of infrared stimulated luminescence from feldspars, *Radiation Measurements*, 81, 242–250,
585 <https://doi.org/10.1016/j.radmeas.2015.02.006>, 2015.

586 Jenkins, G. T. H., Duller, G. A. T., Roberts, H. M., Chiverrell, R. C., and Glasser, N. F.: A new approach for luminescence dating
587 glaciofluvial deposits - High precision optical dating of cobbles, *Quaternary Science Reviews*, 192, 263–273,
588 <https://doi.org/10.1016/j.quascirev.2018.05.036>, 2018.

589 Kaufmann, K. W.: Fitting and using growth curves, *Oecologia*, 49, 293–299, <https://doi.org/10.1007/BF00347588>, 1981.

590 Lal, D.: Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models, *Earth and Planetary Science*
591 *Letters*, 104, 424–439, [https://doi.org/10.1016/0012-821X\(91\)90220-C](https://doi.org/10.1016/0012-821X(91)90220-C), 1991.

592 Laskaris, N. and Liritzis, I.: A new mathematical approximation of sunlight attenuation in rocks for surface luminescence dating,
593 *Journal of Luminescence*, 131, 1874–1884, <https://doi.org/10.1016/j.jlumin.2011.04.052>, 2011.

594 Lehmann, B., Valla, P. G., King, G. E., and Herman, F.: Investigation of OSL surface exposure dating to reconstruct post-LIA
595 glacier fluctuations in the French Alps (Mer de Glace, Mont Blanc massif), *Quaternary Geochronology*, 44, 63–74,
596 <https://doi.org/10.1016/j.quageo.2017.12.002>, 2018.

597 Lehmann, B., Herman, F., Valla, P. G., King, G. E., and Biswas, R. H.: Evaluating post-glacial bedrock erosion and surface exposure
598 duration by coupling in situ optically stimulated luminescence and ^{10}Be dating, *Earth Surface Dynamics*, 7, 633–662,
599 <https://doi.org/10.5194/esurf-7-633-2019>, 2019a.

600 Lehmann, B., Herman, F., Valla, P. G., King, G. E., Biswas, R. H., Ivy-Ochs, S., Steinemann, O., and Christl, M.: Postglacial erosion
601 of bedrock surfaces and deglaciation timing: New insights from the Mont Blanc massif (western Alps), *Geology*, 48, 139–144,
602 <https://doi.org/10.1130/g46585.1>, 2019b.

603 Lifton, N., Sato, T., and Dunai, T. J.: Scaling in situ cosmogenic nuclide production rates using analytical approximations to
604 atmospheric cosmic-ray fluxes, *Earth and Planetary Science Letters*, 386, 149–160, <https://doi.org/10.1016/j.epsl.2013.10.052>,
605 2014.

606 Luo, M., Chen, J., Owen, L. A., Qin, J., Yin, J., Yang, H., Liu, J., Gong, Z., and Luo, J.: A novel approach for reconstructing slip
607 histories for bedrock fault scarps using rock surface luminescence dating, *Geophysical Research Letters*, 49, e2022GL099526,
608 <https://doi.org/10.1029/2022GL099526>, 2022.

609 Luo, M., Chen, J., Gong, Z., Owen, L. A., Yang, H., Jiang, X., Yin, J., Xu, W., and Wang, H.: A new surface rupture of the 1556
610 Huaxian M $\sim$ 8.5 earthquake revealed by rock surface luminescence dating of a bedrock fault scarp and colluvial wedges,
611 *Geophysical Research Letters*, 53, e2025GL119878, <https://doi.org/10.1029/2025GL119878>, 2026.

612 Luo, M., Chen, J., Liu, J., Qin, J., Owen, L. A., Han, F., Yang, H., Wang, H., Zhang, B., Yin, J., and Li, Y.: A test of rock surface
613 luminescence dating using glaciofluvial boulders from the Chinese Pamir, *Radiation Measurements*, 120, 290–297,
614 <https://doi.org/10.1016/j.radmeas.2018.07.017>, 2018.

615 Meyer, M. C., Gliganic, L. A., Jain, M., Sohbati, R., and Schmidmair, D.: Lithological controls on light penetration into rock
616 surfaces – Implications for OSL and IRSL surface exposure dating, *Radiation Measurements*, 120, 298–304,
617 <https://doi.org/10.1016/j.radmeas.2018.03.004>, 2018.

618 Molnar, P. and England, P.: Late Cenozoic uplift of mountain ranges and global climate change: chicken or egg?, *Nature*, 346, 29–
619 34, <https://doi.org/10.1038/346029a0>, 1990.

620 Ou, X. J., Roberts, H. M., Duller, G. A. T., Gunn, M. D., and Perkins, W. T.: Attenuation of light in different rock types and
621 implications for rock surface luminescence dating, *Radiation Measurements*, 120, 305–311,
622 <https://doi.org/10.1016/j.radmeas.2018.06.027>, 2018.

623 Owen, L. A. and Dortch, J. M.: Nature and timing of Quaternary glaciation in the Himalayan–Tibetan orogen, *Quaternary Science*
624 *Reviews*, 88, 14–54, <https://doi.org/10.1016/j.quascirev.2013.11.016>, 2014.

625 Pathan, A. N., Biswas, R. H., Lehmann, B., King, G. E., and Herman, F.: Towards accurate modelling of rock surface exposure
626 dating using luminescence to estimate post-exposure erosion rate, *Quaternary Geochronology*, 85, 101634,
627 <https://doi.org/10.1016/j.quageo.2024.101634>, 2024.

628 Pathan, A., Biswas, R. H., & Kumar, D. (2026). CoRSEER: The Calculator of Rock Surface Exposure Age and Erosion Rates for
629 Rock Surface Luminescence dating [Computer software]. Zenodo. <https://doi.org/10.5281/zenodo.20796442>.

630 Schaefer, J. M., Codilean, A. T., Willenbring, J. K., Lu, Z.-T., Keisling, B., Fülöp, R.-H., and Val, P.: Cosmogenic nuclide
631 techniques, *Nature Reviews Methods Primers*, 2, 18, <https://doi.org/10.1038/s43586-022-00096-9>, 2022.

632 Smedley, R. K., Small, D., Jones, R. S., Brough, S., Bradley, J., and Jenkins, G. T. H.: Erosion rates in a wet, temperate climate
633 derived from rock luminescence techniques, *Geochronology*, 3, 525–543, <https://doi.org/10.5194/gchron-3-525-2021>, 2021.

634 Smedley, R. K., Fletcher, M., Honnold, A., Kane, A., and Sevestre, H.: Highly variable light attenuation across a gneiss rock wall
635 in eastern Greenland, *Radiation Measurements*, 190, 107568, <https://doi.org/10.1016/j.radmeas.2025.107568>, 2025.

636 Sohbati, R. and Hippe, K.: OSL- ^{14}C - ^{10}Be : A novel composite geochronometer for simultaneous quantification of timing and
637 magnitude of change in bedrock outcrop erosion rate, *Earth Surface Processes and Landforms*, 48, 322–331,
638 <https://doi.org/10.1002/esp.5487>, 2023.

639 Sohbati, R., Jain, M., and Murray, A.: Surface exposure dating of non-terrestrial bodies using optically stimulated luminescence: A
640 new method, *Icarus*, 221, 160–166, <https://doi.org/10.1016/j.icarus.2012.07.017>, 2012a.

641 Sohbati, R., Murray, A. S., Chapot, M. S., Jain, M., and Pederson, J.: Optically stimulated luminescence (OSL) as a chronometer
642 for surface exposure dating, *Journal of Geophysical Research: Solid Earth*, 117, <https://doi.org/10.1029/2012jb009383>, 2012b.

Sohbati, R., Murray, A. S., Jain, M., Buylaert, J.-P., and Thomsen, K. J.: Investigating the resetting of OSL signals in rock surfaces, *Geochronometria*, 38, 249–258, <https://doi.org/10.2478/s13386-011-0029-2>, 2011.

Sohbati, R., Liu, J., Jain, M., Murray, A., Egholm, D., Paris, R., and Guralnik, B.: Centennial- to millennial-scale hard rock erosion rates deduced from luminescence-depth profiles, *Earth and Planetary Science Letters*, 493, 218–230, <https://doi.org/10.1016/j.epsl.2018.04.017>, 2018.

Stone, J. O.: Air pressure and cosmogenic isotope production, *Journal of Geophysical Research: Solid Earth*, 105, 23753–23759, <https://doi.org/10.1029/2000JB900181>, 2000.

Wintle, A. G. and Murray, A. S.: A review of quartz optically stimulated luminescence characteristics and their relevance in single-aliquot regeneration dating protocols, *Radiation Measurements*, 41, 369–391, <https://doi.org/10.1016/j.radmeas.2005.11.001>, 2006.