

Paleoenvironmental and paleoclimatic reconstruction in the Western Mediterranean during the Late Early Pleistocene

Maé Catrain¹, Séverine Fauquette², Odile Peyron², Nathalie Combourieu-Nebout¹, Vincent Lebreton¹, Morgane Fischer-Fries³, Mary Robles², Viviane Bout-Roumazeilles⁴, Patricia Richard³, Marion Delattre⁴, Lionel Dubost¹, Sébastien Joannin², Amandine Viala², Yul Altolaguirre⁵, Jean-Pierre Suc⁶, Jeanne Lepelletier¹, Marie-Hélène Moncel¹

¹ Histoire Naturelle des Humanités Préhistoriques, UMR 7194 HNHP, MNHN-CNRS-UPVD, Muséum National d'Histoire Naturelle, Paris, France

² Institut des Sciences de l'Evolution de Montpellier, UMR 5554, Université de Montpellier-CNRS-IRD-EPHE, Montpellier, France

³ Laboratoire des Sciences du Climat et de l'Environnement, UMR 8212 LSCE/IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, Gif-sur-Yvette, France

⁴ Laboratoire d'Océanologie et de Géosciences, UMR 8187 LOG. CNRS, Université de Lille, Université du Littoral-Côte d'Opale, IRD, F-59000 Lille, France

⁵ ROCEEH Research Centre 'The Role of Culture in Early Expansions of Humans' of the Heidelberg Academy of Sciences, Senckenberg Research Institute, Senckenberganlage 25, 60325, Frankfurt am Main, Germany

⁶ Institut des Sciences de la Terre Paris, UMR 7193 ISTeP, CNRS-INSU, Sorbonne Université, Paris, France

Correspondence to: Maé Catrain (mae.catrain@edu.mnhn.fr)

Abstract. The major climate change recorded during the Early-Middle Pleistocene Transition (1.4-0.4 Ma) is characterized by an increase in the length of climatic cycles from 41 ka to 100 ka. During this period, the Mediterranean climate underwent aridification associated with a drop in temperatures. This period also corresponds to the arrival of the first hominins in Western Europe. This study aims to establish the climatic framework at the end of the Early Pleistocene between MIS 37-31 (~1.25 to 1.06 Ma). To this end, a multiproxy approach was applied to assemblages of planktonic foraminifera, pollen and clay mineralogy, constituting a multi-method approach to climate reconstruction. Comparisons with other sequences from the Central and Western Mediterranean show major differences between the Iberian Peninsula and southern Italy. Indeed, the MIS 31 appears cooler and dryer in the south of the Iberian Peninsula than in central Mediterranean. Clay mineralogy analysis highlights a sudden change in oceanic and atmospheric circulation in the Alboran Basin around 1140 ka. Climate reconstructions follow climatic cycles, with temperatures that appear to be underestimated compared to temperature reconstructions based on fauna from southern Spain. Precipitation reconstructions, on the other hand, are more consistent, suggesting that the climate and vegetation of this region are more likely to be influenced by variations in precipitation. These new data enhance our understanding of the climate of the South-western Mediterranean at the beginning of the Early-Middle Pleistocene Transition.

1. Introduction

The boundaries of the Early Middle Pleistocene Transition (EMPT), also known as the Mid-Pleistocene Transition, have been set at 1.25 to 0.75 Ma by Clark et al. (2006). This chronological window has been widely discussed and extended to 1.4 to 0.4 Ma by Head and Gibbard (2015). At that time, the Earth's climatic cycles lengthened from 41 kyr to 100 kyr (Pisias and Moore, 1981; Lisiecki and Raymo, 2005), and cycles became asymmetrical, with gradual onsets of glaciation and abrupt glacial terminations (Lisiecki and Raymo, 2005). This difference in structure is linked to a combination of changes in the influence of orbital parameters on the Earth's climate system, and internal changes involving feedback loops such as the long-term cooling and a decrease in CO₂, the removal of regolith and ice dynamics (Chalk et al., 2017; Head and Gibbard, 2015; Berends et al., 2021; Herbert, 2023). Before the EMPT, climate cycles were controlled by obliquity, whereas after the EMPT, they are often presented as being controlled by eccentricity. However, this cyclicity is much more complex, involving the influence of obliquity (41 ka), precession (23 ka) and an eccentricity cycle (100 ka) (Maslin and Brierley, 2015; Herbert, 2023). During the EMPT, the cooling trend that began in the Late Pliocene continued and accelerated in parallel with a decrease in the concentration of atmospheric CO₂ (Martínez-Botí et al., 2015). Glacial periods became more intense and longer (Herbert, 2023). The increase in cycle duration associated with the 100-ka cyclicity is recorded towards the end of the Early Pleistocene, between Marine Isotope Stages (MIS) 37 to 31 (1.24 to 1.06 Ma), comprising the particularly long MIS 36-35 cycle (~75 ka; Barker et al., 2022; Hodell et al., 2023). It is also marked by the MIS 31 'super interglacial' (DeConto et al., 2012; Melles et al., 2012) and the glacial MIS 34, which has recently been characterized as extreme cooling (Margari et al., 2023).

The start of the EMPT coincides with the arrival of the first hominins in Western Europe, as shown by evidence of occupation dating to around 1.4-1.2 Ma in Italy and on the Iberian Peninsula (Peretto, 2006; Carbonell et al., 2008; Moyano et al., 2011; Duval et al., 2024; Cucart-Mora et al., 2026). The Mediterranean is a key region for studying the spread of hominids, because the presence of environmental studies associated with archaeological evidence (Leroy et al., 2011; Messenger et al., 2011), and the pathways they took to reach Europe are particularly concentrated in this area (Abbate and Sagri, 2012). In addition, the region is highly sensitive to climate change and is thus appropriate for cross-checking climate results from the high- and mid-latitude systems of the Northern Hemisphere and the African monsoon regimes. The impact of global climate change during the EMPT on the Mediterranean is still poorly understood due to the scarcity of reliable sedimentary archives for this region. Some previous studies have shown that aridification in the Mediterranean was a major parameter throughout this period, leading to changes in vegetation, including forest regression and expansion and the progressive disappearance of relict taxa with affinities for warm and humid environments (Joannin et al., 2011; Pross et al., 2015; Magri et al., 2017; Donders et al., 2021). Foraminifera assemblages reveal that changes in Mediterranean circulation, and particularly variations in Mediterranean outflow water (MOW), resulted in a modulation of Atlantic oceanic circulation, heat transport (Bahr et al., 2018, 2022) and ventilation (Guo et al., 2020). To date, changes in the atmospheric system based on the mineralogy of clays in the Mediterranean at the end of the Early Pleistocene are only continuously documented at ODP Site 964 (Ionian Sea) and MD 964 (Levantine Basin), in the Central and Eastern Mediterranean, respectively. Both records highlight changes in sedimentary sources, varying between North Africa and the southern Apennine Peninsula (Zhao et al., 2016), and changes in precipitation in the Nile source areas linked to monsoon activity (Zhao et al., 2012). Climate reconstruction efforts for this period are still in their infancy. Based on pollen assemblages, only three sequences have been used for climate reconstruction at the end of the Early Pleistocene: Lake Acigöl in Türkiye (Robles et al., 2026), which covers the entire EMPT

period, previous data from ODP 976 (Joannin et al., 2011) and the Palominas site (Altolaguirre et al., 2020), which covers the beginning of the transition. Other climate reconstructions based on faunal assemblages from Spanish archaeological and palaeontological sites also provide information on specific time windows, depending on site chronology: Gran Dolina (Blain et al., 2009), Fuente Nueva 3, Barranco Leon (Agustí et al., 2009; Blain et al., 2011, 2016; Saarinen et al., 2021; Sánchez-Bandera et al., 2023), and Almena-Casablanca 3 (Blain et al., 2007; Marquina-Blasco et al., 2025). Comparing different types of climate reconstruction (based on pollen and fauna) can yield a more complete picture of the climate. It is therefore essential to provide new long-term climate data in order to understand global and regional differences in the Mediterranean climate during the EMPT.

Despite the widespread use of marine sequences, the possibilities for studying climate change throughout this period with a multiproxy and integrated approach are limited by the restricted number of long sequences covering the entirety of the EMPT. Thus, our knowledge of the EMPT climate is fragmentary, and questions persist as to the impact of the EMPT on hydroclimatic dynamics, terrigenous sources and vegetation changes. How did the different basins of the Mediterranean respond to climatic changes during the EMPT? How did this transition begin in the Mediterranean? In this paper, we aim to better constrain the land–sea climate dynamics across different Mediterranean basins, with a particular focus on the Alboran Sea, during the onset of the EMPT.

This study provides the first multiproxy and multi-method analysis covering the period ranging from MIS 37 to 31 (1.24 to 1.06 Ma) in the Western Mediterranean using the ODP 976 marine sequence (Alboran Sea). It connects isotope records, pollen data from previous studies (Joannin et al., 2011; Catrain et al., 2025), with new reconstructed Mean Annual Air Temperatures (MAAT) and Mean Annual Precipitation (MAP) based on this pollen data, planktonic foraminifera assemblages, Sea Surface Temperatures (SSTs) based on foraminifera data and clay mineralogy. For this time period, comparisons with the sites of Montalbano Jonico in south Italy (Joannin et al., 2008; Girone et al., 2013), Monte San Giorgio in Sicily (Catrain et al., 2025) and Palominas in southern Spain (Altolaguirre et al., 2020) provide the first land-sea comparison between the central and western Mediterranean showing that differences in vegetation already existed (Catrain et al., 2025). This approach aims to enhance our understanding of climate system variability through: i) examining links between sea and continent, ii) a regional focus and comparison of climate response in the Central and Western Mediterranean, and iii) a methodological discussion of pollen-based climate reconstruction methods.

2. Environmental setting

According to the Köppen classification, the Mediterranean climate is a temperate climate characterized by dry summers (Cs) with precipitation mainly concentrated in winter. Some areas are characterized by more arid steppe (BS) and desert (BW) climates (Köppen, 1936; Daget, 1977; Quézel and Médail, 2003; Peel et al., 2007; Beck et al., 2023). Current annual rainfall around the Alboran Sea varies between 200 mm and 1100 mm and annual temperatures between 12 and 19°C, depending on areas and altitudes (Fig.1). The surface circulation of the Alboran Sea is linked to the entry of the Atlantic jet (AJ) and two main surface gyres: the Western Alboran Gyre (WAG) and the Eastern Alboran Gyre (EAG), both with an anticyclonic circulation. The WAG and EAG are connected to each other by a smaller gyre; the Central Cyclonic Gyre (CCG) (Fig.1.A) (Sánchez-Garrido and Nadal, 2022). The surface waters from the Atlantic Ocean entering through the Strait of Gibraltar move towards the Almeria Oran Front (AOF) where they mix with saltier waters. From the east, the Levantine Intermediate

Water (LIW) and the colder and more saline Western Mediterranean Deep Water (WMDW) move west before leaving through the strait. The mixing of these two water masses upon arrival in the Atlantic Ocean is called Mediterranean Outflowing Water (MOW) (Fig.1.B) (Fabres et al., 2002; Macías et al., 2008; Moreno et al., 2005; Pérez-Folgado et al., 2003; Tintore et al., 1988).




Figure 1. ODP site 976 climate and location maps. Climate data from ERA5 reanalysis between 1960-2022 with a lat-lon grid of 0.25 degrees. A) Mean annual precipitation with sea surface circulation. AJ: Atlantic Jet, WAG: Western Alboran Gyre, CCG: Central Cyclonic Gyre and EAG: Eastern Alboran Gyre. B) Mean annual temperature with sea deep circulation. MOW: Mediterranean Outflowing Water, LIW: Levantine Intermediate Water and WMDW: Western Mediterranean Deep Water. C) Atmospheric circulation. NSAL: Northern Saharan Air Layer.

In the Western Mediterranean region, atmospheric circulation is influenced by the westerlies and variations in the positioning of the intertropical convergence zone and therefore the monsoon system. The Western Mediterranean is subject to winds from Africa, such as Northern Saharan pluri-directional winds, which originate in the Sahara. On the Atlantic side, the Saharan Air Layer (SAL) originating from the monsoon system is deflected by the Azores high pressure system, thus creating the northern branch of the SAL (NSAL), which goes up to the Western Mediterranean (Bout-Roumazelles et al., 2007; Leblanc, 2022). The Alboran basin is subject to westerly and south-westerly winds, the Poniente and the Vendaval, and warm easterly winds such as the Levanter.

Around the Alboran Sea, Mediterranean vegetation is organized altitudinally according to temperature and precipitation distribution. The Thermo-Mediterranean belt (0 to 300-700 m) is mainly composed of Mediterranean species, such as olive trees (*Olea*), pistachio trees (*Pistacia*) and carob trees (*Ceratonia*). The Meso-Mediterranean (300-700 to 1000-1200 m) and Supra-Mediterranean (1000-1200 and 1500-1700 m) belts consist of evergreen mixed forest and deciduous oak (*Quercus*) and pine (*Pinus*) forests respectively. Above these altitudes, forests are composed of conifers such as pine (*Pinus*), fir (*Abies*) and cedar (*Cedrus*) in Morocco (Benabid, 1985; Ozenda, 1994; Quézel and Médail, 2003; Médail, 2022).

3. Site location and chronology

ODP Leg 161 site 976 (36°12.30'N, 4°18.8'W, 1108 m b.s.l. (below sea level); Fig. 1) is located in the northwest of the Alboran Sea. The sediment core records a continuous lithostratigraphy from the Upper Miocene to the Holocene (Comas et al., 1996). Our study focuses on the depth interval of the master core from 281.23 to 256.53 mcd (metre composite depth), corresponding to the MIS 37-31 interval, with an age model based on nannofossil events, sapropel and the correlation between ODP site 976 (Catrain et al., 2025) and global stack oxygen isotope curves (Lisiecki and Raymo, 2005). The sedimentary sequence consists of homogeneous clays rich in nannofossils and silty nannofossils. The lithogenic particles in the Alboran Sea come from aeolian and fluvial sources from Spain and Morocco.

4. Materials and Methods

This paper uses the ODP Site 976 palynological data obtained by Catrain et al. (2025) which followed a standard protocol of chemical treatment. For every sample a minimum of 150 grains excluding *Pinus* (overrepresented) and a minimum of 15 taxa were recorded to have a reliable representation of the vegetation.

Additional proxies and statistical analyses were used to cross environmental information and to better constrain the land-sea climate dynamics.

4.1. Planktonic foraminifera analysis and isotope analysis

Planktonic foraminifera analysis was carried out on one hundred and sixteen samples with a spacing of 10 cm to 80 cm, with an average of every 23 cm. The sediments were sieved to 150 µm. Foraminifera were counted using a binocular magnifying glass with a minimum of 450-500 tests per sample to be representative. Identification is based on publications (Orbigny, 1839) and a book (Schiebel and Hemleben, 2017). No evidence of diagenesis was identified.

Between eight and 12 individuals of the foraminifera *Globigerina bulloides* were selected from 56 of these same samples to conduct $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ analyses. The samples were cleaned with methanol before being inserted into the dual-entry Isoprime isotope ratio mass spectrometer. Analyses were performed at the Laboratory for Sciences of Climate and Environment (LSCE, France) on the analytical platform PANOPLY.

4.2. Clay mineral analysis

Clay mineral associations were studied using X-ray diffraction following the protocol of Bout-Roumazeilles et al. (1999). In this study, 'clay minerals' refer to the main phyllosilicate minerals in the clay-size fraction

(generally less than 2- μ m particles). All samples were first decalcified with 0.2 M HCl, before clay deflocculation by successive washing with distilled water. The clay-size fraction was separated by settling according to Stokes's law, concentrated by centrifugation, and oriented by wet smearing on glass slides. X-ray diagrams were obtained using a Bruker D4 Endeavor diffractometer with CuK α radiation and a Ni filter, coupled with a Lynxeye detector. A tube voltage of 30 kV and a tube current of 35 mA were utilised. Three X-ray diagrams were performed: air-dried sample (normal run), ethylene-glycol vapour saturation for 12 h (glycol run) and heating at 490°C during 2 h (heating run). The goniometer scanned from 2.49° to 32.49° 2 θ for normal and glycol run and from 2.5 to 14.5° 2 θ heating run. Each clay mineral is characterized by its basal layer plus interlayer interval (d), as revealed by XRD analysis (Brown and Brindley, 1980). Smectite (S) is characterized by a peak at 14 Å on the normal run, which expands to 17 Å after saturation by ethylene-glycol and retracts to 10 Å after heating. Illite (I) presents a basal peak at 10 Å on the three runs. Palygorskite (P) is characterized by the presence of a peak at 10.34 Å on the normal run. Chlorite (C) is characterized by peaks at 14 Å (001), 7 Å (002), 4.75 Å (003) and 3.53 Å (004) on the three runs. Kaolinite (K) is characterized by peaks at 7 Å (001) and 3.58 Å (002) on the normal and glycol runs. Both peaks disappear or are strongly reduced after heating. To distinguish kaolinite from chlorite, the portion of the spectrum containing the basal peaks of kaolinite (002) and chlorite (004) around 3.55 Å is step-scanned in a high-resolution mode following standard procedures described in detail by Petschick et al. (1996). Semi-quantitative estimation of clay mineral abundance is based on peak areas and totalled to 100% (S + I + K + C + P = 100%). Peak area measurements were taken in the glycol runs using the Macintosh MacDiff® 4.2.5 software (Petschick, 2000). The error on measurement reproducibility is estimated to be $\pm$ 5% for each clay mineral.

4.3. Sea surface temperature reconstruction from planktonic foraminifera

Sea surface temperatures (SST) were reconstructed based on planktonic foraminifera assemblages, using PaleoAnalog software (version 3.0). This software uses the Modern Analogue Technique (MAT), which compares a database of modern assemblages with fossil assemblages to find the closest modern samples (Theron et al., 2003). The degree of dissimilarity between samples is calculated using the Squared-Chord-Distance. A minimum of ten temperature analogues were selected and then averaged to estimate those of the fossil samples. Reconstructions were based on a North Atlantic Mediterranean database (Siccha and Kucera, 2017).

4.4. Pollen-inferred climate reconstructions

For climate reconstructions, pollen counts were used to reconstruct four climatic parameters: Mean Annual Air Temperature (MAAT), Mean Temperature of the Coldest month (MTCO), Mean Temperature of the Warmest month (MTWA), and Mean Annual Precipitation (PANN). A multi-method approach was adopted for greater reliability (Peyron et al., 2011, 2013; Salonen et al., 2019; Chevalier et al., 2020), and taxa that have now disappeared from the Mediterranean (relict taxa) were included in certain methods (Fauquette et al., 1998). Five methods with different mathematical bases were selected: the modern analogue technique (MAT; (Guiot, 1990)), weighted averaging partial least-squares regression (WAPLS; (Ter Braak et al., 1993)), random forest (RF; (Breiman, 2001; Prasad et al., 2006)), boosted regression trees (BRT; (De'ath, 2007; Elith et al., 2008)) and the climatic amplitude method (CAM; (Fauquette et al., 1998)). To discuss the climate of the Western Mediterranean

Basin with a multi-site approach, additional pollen-based climate reconstructions were carried out using data from Monte San Giorgio and Palominas, following the same multi-method approach (CAM, MAT, WAPLS, RF and BRT).

The CAM is based on the overlap of the climatic requirements of taxa and takes relict taxa into account. The climatic requirements of ~135 taxa were defined on the basis of 8000 modern spectra and the literature. The past climate is estimated by transposing the climatic requirements of the maximum number of modern taxa to the fossil data. The most probable climate corresponds to the suitable climatic interval for the highest number of taxa and to a ‘Most Likely Value’ corresponding to a weighted mean (Fauquette et al., 1998). Some taxa represented by a single grain in the record (*Lygeum*, *Croton*, *Rhus*, *Myrthus*, *Calligonum*, *Taxodium*, and Sapotaceae) and Cichorioideae were not taken into account in the climate reconstruction as they generate noise. For the CAM, Poaceae were not taken into account for the reconstruction of the Palominas site for the same reason as for Cichorioideae in ODP site 976.

RF and BRT methods are based on machine learning and have recently been used for palaeoclimatology in Northern Europe (Salonen et al., 2019) and the Mediterranean Basin (d’Oliveira et al., 2023, 2025; Robles et al., 2023, 2025; Sassoon et al., 2025; Charton et al., 2025). For RF, each tree is estimated from a random set of different subsets defined separately from the modern data assembly. This is called bootstrapping and each sample has the same probability of being selected (Breiman, 2001; Prasad et al., 2006). For BRT, the random creation of sets from modern assemblages means that samples that were not sufficiently taken into account in the previous tree have a higher probability of being selected in the next training sessions. This is known as boosting (De’ath, 2007; Elith et al., 2008). WAPLS is a non-linear regression technique that creates relationships between climate parameters and pollen taxa within the modern database, then applies them to fossil assemblages (Ter Braak et al., 1993; Chevalier et al., 2020). MAT method works in a similar way to the one use for reconstruct SST based on foraminifera. The MAT and WAPLS methods were applied with the R *rioja* package (Juggins, 2024), RF with *randomForest* package (Breiman et al., 2024) and BRT with *dismo* package (Hijmans et al., 2024).

The Eurasian and Mediterranean Current Surface Sample Database (EAPDB; n = 3373 sites), compiled by Peyron et al. (1998, 2011, 2013), and completed by Dugerdil et al. (2021) and Robles et al. (2023), was used for the first four methods. In this dataset, each sample is associated with a biome (Tarasov et al., 1998; Elenga et al., 2000). Some biomes were selected and others were not taken into account in order to better constrain the database. Here the biomes selected: cold deciduous forest (CLDE), cold desert (CODE), cool mixed forest (COMX), cold steppe (COST), hot desert (HODE), pioneer (PION), temperate deciduous forest (TEDE), warm mixed forest (WAMX), warm steppe (WAST) and xerophilous (XERO). Anthropogenic (ANTH), cold mixed forest (CLMX), cold conifer forest (COCO), taiga (TAIG) and tundra (TUND) biomes were not taken into account because they are too far from the study area. One hundred and seven sites from the Eurasian modern database (EMPD) (Davis et al., 2020) were added to the EAPDB for a more reliable representation of certain taxa (Ericaceae). Particular attention was paid to these new samples to avoid selecting anthropized sites. *Pinus*, kept in the modern database, was not included in the fossil sequence due to its overrepresentation, except for the Palominas site which corresponds to a continental basin.

It is difficult to rank climate reconstruction methods, as each has its own advantages and disadvantages. The order in which the methods appear in this section reflects our subjective ranking. CAM offers a key advantage for

this time period because, unlike other methods, it accounts for relict taxa. Furthermore, it considers taxa individually, allowing them to be distinguished within pollen assemblages. Machine learning methods such as BRT and RF do not rely on ecological theories but are based on statistical methods, which helps limit ecological biases in the database. MAT appears to be the least suitable method for this study period, as there is no known true analogue of past vegetation in the Mediterranean region, hence the use of a database covering Eurasia to identify the closest possible assemblages. However, the dominance of *Quercus* in the arboreal component of the pollen signal makes it difficult to constrain the reconstructions effectively, regardless of the method used, since trees exert a stronger constraining influence on climate reconstructions than grasses due to their higher taxonomic resolution.

4.5. PERMANOVA

The use of PERMANOVA in this study aims to compare different methods of climate reconstruction with one another, taking into account different time periods and climate parameters. PERMANOVA is a non-parametric test based on the F-test. The p-value of PERMANOVA is estimated by permutation, which makes it particularly well adapted to data that do not follow multivariate normality or homogeneity of variances, as is the case with climate reconstruction data. These two criteria are important for the application of traditional analyses of variance methods (ANOVA, MANOVA), as confirmed using a Mardia test. Whilst ANOVA compares group means to verify whether or not they come from the same population, PERMANOVA projects data points into multivariate space, and therefore compares group centroids instead of means. The null hypothesis (H0) of the PERMANOVA test is that the reconstructed values belong to the same centroids and that their dispersion is equivalent across all groups. In other words, reconstructed values are considered as permutable or “equivalent” under the null hypothesis. When the p-value of PERMANOVA is less than or equal to 0.05, the test is significant and rejects H0. The centroids of reconstructed values and their spread is different for each group, which are considered not permutable in multivariate space and therefore, “different”. This test was applied to reconstructed temperature and precipitation data, which were centred and reduced to avoid the influence of scale and give equal weight to each variable and to compare trends.

5. Results

As a reminder, the pollen data presented by Catrain et al. (2025) are composed of 31 families and 72 genera distributed over the different belts of Mediterranean vegetation. The temperate taxa curves, mainly represented by the *Quercus* deciduous type, *Isoetes* and elements of steppe environments (*Artemisia*, *Amaranthaceae*, *Ephedra*) highlight alternating warm/ humid periods and cold/dry periods between MIS 37 and 31. The high percentages of high-altitude forest elements, such as *Cedrus* and *Abies*, and of Ericaceae, point to the onset of colder but still humid climatic conditions at the end of the interglacial. The constant values of Asteroideae, Cichorioideae and Poaceae show a continuous presence of an open vegetation environment (Fig.S1).

5.1. Late Early Pleistocene planktonic foraminifera assemblages and SST reconstruction

Altogether, ninety-seven samples were analysed. However, fourteen samples could not be used due to the low number of foraminifera. Thirty-nine species were identified, dominated by *Neoglobobulimina incompta* (on average 30%), *Globigerina bulloides*, *Globobulimina inflata*, *Neoglobobulimina pachyderma*, and *Globigerinoides*

ruber white (Fig. 2). *Globigernella calida* (around 1%), *Globigerinita glutinata* (4%), and *Globigerina falconensis* (3%) are consistently present. Taxa with affinity for warm waters, such as *Orbulina*, *Globorotalia crassaformis*, and *G. calida*, are poorly represented in the assemblage. Warm taxa, *Globigerinoides ruber* white type, and taxa with an affinity for temperate waters, such as *Globigerina bulloides* and *Globorotalia inflata*, are found in interglacial periods. Two peaks of *G. Globigerinoides ruber* white are recorded at 1229 ka (39%) and 1084 ka (49%). *Turborotalita quinqueloba* develops significantly (up to 30%) between 1215 and 1200 ka. The high *Neogloboquadrina pachyderma* percentages, with peaks between 40 and 70%, are synchronized with glacial periods.




Figure 2. Planktonic foraminifera assemblages from ODP site 976 and SST reconstructed with the Modern Analog Technique (MAT) associated with MIS chronology. The red curves correspond to foraminifera with warm affinities and the blue curves to cold affinities. A map of the analog distribution (Fig. S2) and a table of analog's distances (Tab. S1) are available in supplementary data.

Mean annual sea surface temperature values reconstructed from planktonic foraminifera assemblages vary between 5 and 19.6 °C. The values show marked variation with a trend of cooling during glacial periods and warming during interglacial periods (Fig. 2).

5.2. Late Early Pleistocene clay analysis in the Alboran Sea

Over the studied interval, the clay assemblage is mainly composed of illite (35%) and smectite (30%) associated to 15% chlorite and 10% kaolinite, palygorskite being less abundant (6%). This composition is very similar to the composition of recent sediments deposited in the Alboran Sea (MIS1-MIS3), consisting mainly of illite and smectite, which account on average for 35% and 30% of the clay assemblage respectively (Bout-


Roumazeilles et al., 2007; Hassouni and El Hmaidi, 2007), with kaolinite and chlorite at around 16%, whilst palygorskite accounts for less than 5% (Bout-Roumazeilles et al., 2007; Hassouni and El Hmaidi, 2007). The clay mineral signals can be divided into four main phases of variation between stages 37 to 31 (Fig. 3):

- Phase I, corresponding to MIS 37 is slightly enriched in illite (40%) and depleted in smectite (20%), with a I/K ratio around 2.4. The end of stage 37 is marked by an increase in kaolinite (~20-21%) and chlorite (~19%) and a decrease in smectite (~12%) and the I/K ratio (~2).

- Phase II is the longest phase, covering MIS 36-35, and records fairly constant values for each mineral, very similar to those for phase I.

- Phase III, which begins with the transition from MIS 35 to 34, shows the greatest change in clay composition. A general decreasing trend is observed for illite (~31%), kaolinite (~10%), chlorite (~11%) and palygorskite (~6%), although the latter peaks at around 1114, 1103, 1001 and 1088 ka. On the other hand, during this phase, smectite increases significantly, reaching average values of around 42%. The I/K ratio fluctuates more regularly, with values rising slightly to 3.69.

- Phase IV corresponding to MIS 31 presents a return to the values of phases II and I.



325 **Figure 3. Clay mineralogy from ODP site 976 associated with MIS chronology. Solid lines represent the mean and dotted**
326 **lines the standard deviation.**

5.3. Climate reconstruction during the Late Early Pleistocene

Reconstructions from pollen data at ODP site 976 show similar trends between all the different methods and for each parameter (Fig 4). BRT and MAT have the best R^2 and RMSE, while WAPLS yield the poorest for all climatic parameters (Tab. 1). MAT and WAPLS show greater variation amplitude than BRT, RF and CAM, where differences between warm, wet phases and cold, dry phases are less pronounced. The RF shows the same variations as the CAM for the MAAT and PANN; for the MTWA, the CAM exhibits greater amplitudes, whereas for the MTCO the RF has greater amplitudes. Parameter variations follow those of interglacial and glacial periods. Interglacial phases are characterized by higher temperatures and precipitation than for glacial periods. RF appears colder than other methods, with a general mean value of around 10 °C in MAAT and 0 °C for MTCO. WAPLS and MAT present more humid conditions.

All methods except BRT show that MIS 37 (1245–1215 ka) was the warmest period of the study, based on MAAT, MTWA and MTCO. Precipitation levels were high, but were nevertheless similar to those reconstructed for MIS 31 (1081–1062 ka), which had lower temperatures. MIS 35 (1190–1141 ka), meanwhile, shows temperature values close to those of MIS 31 but with slightly lower precipitation. For this stage, the BRT is the only method that shows a decrease in annual temperatures and in the temperature of the warmest month. The RF, MAT and WAPLS methods indicate that MIS 33 (1114–1104 ka) was the coolest and driest interglacial of the period.

According to all methods, MIS 36 (1215–1190 ka) was characterised by cold and dry conditions, although these were less severe than during the subsequent glacial phases; the only exception is the BRT method, which shows no decrease in temperature or precipitation relative to the flanking interglacial periods. According to the MAT and WAPLS methods, the MIS 34 glacial phase (1141–1114 ka) is the coldest and driest of the cold phases of the period for the MAAT and MTCO. The other methods show that the end of MIS 32 (1104–1081 ka) was the most extreme period. However, all methods unanimously indicate that MIS 34 is the stage with the driest conditions.




Figure 4. ODP site 976 pollen climate reconstruction. In green, RF (Random Forest), in blue BRT (Boosted Regression Trees), in yellow WAPLS (Weighted Averaging Partial Least Squares regression), in red MAT (Modern Analogue Technique) and in purple CAM (Climatic Amplitude Method). Four climatic parameters: MAAT (Mean Annual air Temperature), MTWA (Mean Temperature of the Warmest month), MTCO (Mean Temperature of the Coldest month) and PANN (mean annual precipitation). Grey bands correspond to interglacial periods and the thick coloured lines correspond to loess-smoothed curves and shaded areas to the 95 % confidence interval.

Table 1. R^2 (Pearson correlation coefficient) and RMSE (Root Mean Square Error) values for climatic parameters. The lower the RMSE and the higher the R^2 , the closer the method is to the target value from the modern database. R^2 and RMSE are based on bootstrapping for MAT and WAPLS and on leave-one-out validation for BRT and RF. The calibration range associated is given in supplementary data in the Tab. S2 and the WAPLS has 3 components. Fig S3 presenting the observation vs predicted plots. For CAM, R^2 and RMSE are obtained by applying the method to an independent validation dataset and comparing the predictions with the observed data.

R^2	MAT	WAPLS	RF	BRT	CAM
MAAT	0.83	0.65	0.71	0.81	0.87
MTWA	0.79	0.54	0.62	0.70	0.68
MTCO	0.87	0.70	0.77	0.87	0.86
PANN	0.72	0.46	0.60	0.71	0.45
RMSE					
MAAT	2.53	3.30	2.92	2.36	3.4
MTWA	2.38	3.17	2.81	2.24	2.9
MTCO	3.30	4.53	3.88	2.95	5.5
PANN	194.25	241.41	204.44	176.96	257

369


370 **5.4. Statistical analysis of climate reconstruction**

371 The test comparing the methods for each MIS (Fig. 5) shows that, for most of them, the values reconstructed
372 by the methods belong to the same centroid. Indeed, for PANN, H0 is not rejected for any methods from MIS 37
373 to MIS 33 (Fig. 5). For MAAT, H0 cannot be rejected for any of the methods (Fig. 5). Therefore, the various tests
374 carried out with PERMANOVA on the centred and reduced climate reconstruction data suggest that the methods
375 can be considered as consistent between each other. H0 is only rejected for certain MIS or for PANNs. For MIS
376 32, only CAM has a p-value < 0.05 and is significantly different to MAT (p-value = 0.004), WAPLS (p-value =
377 0.004) and BRT (p-value = 0.005). RF differs from MAT (p-value = 0.039) and BRT (p-value = 0.049) during
378 MIS 31. In those cases, reconstructed values are considered to be significantly different between each other (Fig.
379 5)

380

381 PERMANOVA tests that compare the glacial and interglacial periods with each other for each method,
382 however, all reject H0 (p-value < 0.05; Fig.6). All interglacial and glacial phases can therefore be considered to
383 be different from the others, regardless of the methods. .

384



385 Figure 5. PERMANOVA p-values between climate reconstruction methods for each isotopic stage and for Mean Annual
386 Air Temperature (MAAT) and Mean of Annual Precipitation (PANN). A p-value less than or equal to 0.05 is significant
387 and indicates that the two methods are different from each other.




Figure 7. PERMANOVA p-values between the same types of periods (glacial-glacial and interglacial-interglacial) for each climate reconstruction method on centred-reduced data. A p-value less than or equal to 0.05 is significant and indicates that the two periods are different from each other.

6. Discussion

6.1. Land-sea responses from the Western to Central Mediterranean

Planktonic foraminifera data from the same marine sequence as the pollen data highlight simultaneous changes in oceanic characteristics and in the floristic composition of terrestrial ecosystems. The Mediterranean Basin, and in particular the Western Mediterranean, only comprises rare sites combining these two proxies. This foraminifera/pollen data approach is recorded in the Montalbano Jonico site over a large EMPT time window (Joannin et al., 2008; Girone et al., 2013).

Between MIS 37 and MIS 31 in ODP site 976 (Western Mediterranean basin), all the proxies highlight the typical interglacial and glacial variations recorded during this period, except clay mineralogy, which shows longer cycles. The SST estimates should be interpreted with caution, as the high abundance of *Neogloboquadrina pachyderma* is likely to introduce a cold bias, as already observed in a previous study (Mega et al., 2025). However, the SST variations are of great interest and may be compared to the other proxies. The comparison between pollen data and planktonic foraminifera assemblages highlights a synchronization of steppe elements with the polar affinity foraminifera *Neogloboquadrina pachyderma*. A similar pattern is observed during MIS 20-19 (Toti et al., 2020). This correlation emphasizes that the steppes from the Iberian Peninsula and/or North Africa are not only linked to aridity but also to low temperatures (Fig. 8.a). The constant dominance of the subpolar affinity *Neogloboquadrina incompta* (~30%) throughout the period also attests to relatively cool surface waters (average SST of 12 °C). Air and ocean temperatures (MAAT and SST) reconstructed from pollen and planktonic foraminifera also show these glacial-interglacial variations (Fig. 8.a). The SST signal is highly influenced by the quantity of *Globigerinoides ruber* of subtropical affinity during interglacial phases, and *Neogloboquadrina*

pachyderma of polar affinity, during glacial phases. For both air and ocean temperatures, these are lower than modern values. The current SST is 17.7 °C between 1981-2010 (36°N,03°W; (Guijarro et al., 2015; Sánchez-Laulhé et al., 2021)) and the current mean air annual temperature in a perimeter of 400 km around ODP 976 is 16.8°C between 1960-2022 (calculated with ERA5 reanalysis). Furthermore, multiproxy analyses from ODP site 976 show that the warmest phase is MIS 37 (1245–1215 ka). During this phase, temperate forests reached their maximum development, averaging 35%, while steppes were reduced to 10%. This stage also has the highest percentages of *Globigerinoides ruber* (~20%), a warm-water foraminifera, and SSTs around 16.5 °C, as well as low $\delta^{18}\text{O}$ values (~-0.56‰; Catrain et al., 2025) (Fig. 8).




Figure 8. Land-sea variability comparison at ODP site 976 (left), compared to the site of Montalbano Jonico (right) in southern Italy. Isotope data for the ODP site 976 are from Catrain et al. (2025). For the Montalbano Jonico site, pollen and oxygen isotope data come from Joannin et al. (2008), and planktonic foraminifera data from Girone et al. (2013).

This observation counters current knowledge, which presents MIS 31 (1081-1062 ka) as the warmest interglacial during this period. It is generally referred to as the ‘Super Interglacial’, associated with maxima in obliquity, eccentricity and insolation (DeConto et al., 2012; Melles et al., 2012). However, this result is consistent with interpretations made at sites off the Iberian Peninsula. Indeed cores U1385 and U1387 show that MIS 31 is not associated with particularly high temperature conditions compared to those recorded at high latitudes (Voelker

et al., 2015; Oliveira et al., 2017). Finally, in ODP site 976, MIS 31 is less marked than MIS 37 in terms of isotope values, pollen and foraminifera assemblages, and is closer to MIS 33 (1114-1104 ka) and MIS 35 (1190-1141 ka).

Concerning glacial phases, MIS 34 (1141-1114 ka) seems to be the most pronounced, with significant steppe expansion (~28%) and forest regression (10%). *Neogloboquadrina pachyderma* reached 40% and reconstructions estimate an annual air temperature of ~9°C, which is corroborated by high $\delta^{18}\text{O}$ values of up to 1.62‰. However, these values remain very close to those observed during MIS 36 (1215-1190 ka). Reconstructed SSTs show colder values for MIS 36 (9.5°C) than MIS 34 (9.8°C), even though they remain within the same range of values. This slight difference can be explained by the combined presence of *Neogloboquadrina pachyderma* and the subpolar taxa *Turborotalita quinqueloba* in MIS 36, whereas the latter is almost absent in MIS 34.

6.2. Provenance of clay minerals in the Western Mediterranean

The variations in the mineralogical composition of the sediments with smectite ranging from 20% to 60%, illite ranging from 20 to 45% and lower amounts of kaolinite (up to 20%) suggest changes in the provenance and/or transport patterns of clay particles during the studied interval.

However, the origin and interpretation of smectite in the Alboran basin is still debated (Holeman, 1968; Milliman and Syvitski, 1992; Stanley et al., 1992; Krom et al., 1999; Weldeab et al., 2002; Garzanti et al., 2006; Ehrmann et al., 2007; Hassouni and El Hmaidi, 2007; Hamann et al., 2009; Zhao et al., 2012; Bout-Roumzeilles et al., 2013). Smectite is ubiquitous in various types of environments, as it is easily transported over long-range distances due to its specific buoyancy characteristics. There are several hypotheses regarding the potential sources of the smectite collected in the Alboran Basin, as well as for illite and kaolinite : (1) fluvial origin (the Nile, major European rivers, coastal rivers of the Iberian Peninsula, Moroccan wadis, etc.), (2) eolian origin, and (3) advection/redistribution by oceanic water masses (turbidites, Suspended Particulate Matter (SPM)) from the central and eastern Mediterranean, or from the Gulf of Cádiz and the North Atlantic via the Gibraltar Strait.

The Nile, according to its average discharge (from 120 to 230.10⁶ t yr⁻¹) is one of the primary fluvial source of smectite in the eastern Mediterranean basin (Venkatarathnam and Ryan, 1971; Foucault and Mélières, 2000; Bout-Roumzeilles et al., 2007; Hamann et al., 2009; Zhao et al., 2012) but previous studies have suggested that the dispersion of clays originating from the Nile is negligible to the west of the Mediterranean Ridge, (Krom et al., 1999; Weldeab et al., 2002).

The river systems in southern Spain are unlikely to influence clay sedimentation in the Alboran Sea. Indeed, the Guadalquivir flows into the Gulf of Cádiz, on the other side of Gibraltar, and the Segura, Júcar and Ebro river systems have no outlet into the Alboran Sea, as they flow further north.

The Po River, with an annual discharge of 17.10⁶ t yr⁻¹, as well as most SE European rivers are mainly supplying illite toward central Mediterranean flows via the Adriatic Sea and the Aegean Sea (Venkatarathnam and Ryan, 1971; Dominik and Stoffers, 1979; Chamley, 1989; Alonso and Maldonado, 1990; Tomadin, 2000). The Rhone River may supply illite and chlorite to the Gulf of Lion, but its outlet is located at distance from the studied area, reducing its direct influence on the Alboran Sea sedimentation.

Riverine contribution to the central and western Mediterranean from areas bordering the southern shores is generally enriched in illite and is severely reduced due to the narrow drainage basin and sparse rainfall (Martin and Milliman, 1997). Nevertheless, the Medjerda and Miliane wadis, discharging into the Gulf of Tunis, may influence the sedimentation in the study area. Smectite is a dominant component of suspended particulate material

in the Alboran Sea (Pierce and Stanley, 1975; Grousset et al., 1988; Baringer and Price, 1999), especially within the Western Alboran Gyre (Bulian, 2022), resulting from the interaction between the smectite-rich entering surface Atlantic water, and smectite-rich Levantine Intermediate Water and Western Mediterranean Deep Water (Ambar and Howe, 1979; Bryden and Stommel, 1984; Millot, 2014).

Given the physiographic context of the Alboran Sea, part of the terrigenous fraction may be transported from the South Iberian Plateau to ODP Site 976 via submarine canyon/delta systems such as the Baños, Calahonda or Fuengirola systems, which allow sediments to be transported to depths of over 1000 m (Ercilla et al., 1992; Alonso et al., 1999; Vázquez et al., 2015). Nevertheless, the sedimentological description of ODP 976 (Comas et al., 1996) indicates that sedimentation over the studied interval is predominantly hemipelagic, with no major turbiditic intervals. This contrasts with the older Pliocene section, characterised by numerous gravitational episodes, reflecting reduced sediment transport through these canyons during the interval studied. These observations suggest that gravitational transport processes did not play a major role in sedimentation. Sea-level changes may also affect the clay composition through modifications of the source-to-sink transfers. During glacial period, low sea-level would promote the transfer from South Spain riverine systems enriched in illite and kaolinite and depleted in smectite, which is however not supported by the clay mineral assemblages at site ODP 976. Similarly, during deglaciations, sea level rise would favour the rapid remobilization of sediments from the shelf. By contrast, deglaciations are not associated with major modifications of the clay mineral fraction, given low confidence in this hypothesis.

The advection of smectite-rich sediments from the Gulf of Cádiz via surface water masses entering the Mediterranean is a potential candidate for smectite supply. A recent study evidenced that sediment drifts from the Gulf of Cádiz likely formed under the influence of the MOW are enriched in smectite (Moal-Darrigade et al., 2022). As a consequence, we should not rule out the possibility that sediments passing through Gibraltar are returned to the Mediterranean by ocean circulation, thereby allowing accumulation in the Alboran Basin via the WAG gyre.

Finally, the generally low direct fluvial input and potential eolian contribution, as well as the overall presence of palygorskite suggest that a significant proportion of the clay sediments may be of eolian origin. Previous studies evidenced the dominance of illite and kaolinite, associated with palygorskite in eolian dust from northern Africa, and highlighted the scarcity of smectite in Saharan dust (Guerzoni et al., 1999; Bout-Roumazielles et al., 2007, 2013).

6.3. Atmospheric and oceanographic implications

In order to highlight the potential eolian contribution to marine clay sedimentation, comparison between palygorskite content and I/K ratio may be used to refine the identification of the various eolian sources. Physical erosion and chemical weathering are responsible for the diversity of clay assemblages, but previous studies (Thiry, 2000), on the paleoclimatic interpretation of clay mineral in marine deposits, shown that direct interpretation of variations in the composition of the clay fraction may be tricky. In particular, processes as chemical weathering and hydrolysis, which are responsible for the transformation of clays in soils, are slow-acting mechanisms, which raises issues of synchronisation and time lag between the moment when hydrolysis conditions change, the moment when the nature of the minerals formed changes, and the moment when they are finally transported and then deposited in the oceanic environment. Based on these assumptions, any direct link between modifications of

weathering conditions on the continent and contemporaneous variations in clay composition within the Alboran sea is difficult to assess and is not supported by the whole dataset.

Based on these findings, the clay mineralogy analysis likely record change in atmospheric and oceanographic conditions in the Western Mediterranean with reduced impact of gravity transfer processes. Considering the almost permanent presence of palygorskite, and based upon previous studies in the central and western Mediterranean including the Alboran sea (Bout-Roumzeilles et al., 2007, 2013), the illite/kaolinite (I/K) ratio will be used in order to tentatively retrace variations in dust sources from North Africa. Currently, illite decreases from 60% of clay content in Northern Algeria to 30% in the Sahelian area (Paquet et al., 1984), whereas kaolinite, scarce in the northern Sahara, increases progressively southward and eastward and becomes dominant in the Sahel (Caquineau et al., 1998; Guerzoni et al., 1999; Bout-Roumzeilles et al., 2007; Stumpf et al., 2011). A high I/K ratio therefore suggests inputs preferentially from the Sahara, while lower values indicate a provenance associated with the Sahel. In ODP 976, the I/K ratio strongly increases after the transition between MIS 35 and 34 (around 1140 ka), reflecting a dominant Saharan origin for aeolian particles (Fig.8). This suggests the development of arid conditions with an enhanced supply of Saharan vs Sahelian origin at that time (Bout-Roumzeilles et al., 2007; Skonieczny et al., 2011).

While variations in the I/K ratio are commonly used to retrace aeolian supplies, the interpretation of the individual clay mineral as smectite is less direct and more complex. Given the various scenarios outlined above, the variations observed in the smectite from ODP 976 can be interpreted as reflecting the oceanic transport of terrestrial material into the Alboran Sea, whereas the I/K ratio may partly retrace aeolian supply. As a consequence, the shift in the smectite signal is observed at ~1140ka, concomitant to the observed shift in the I/K ratio, suggesting a change in terrigenous advection and thus in the oceanic configuration in the Alboran Sea around 1140 ka.

In the Central Mediterranean, the vegetation signal from Montalbano Jonico shows extensive steppe developments during glacial periods and different interglacial periods compared to the western basin, more pronounced in MIS 31 and 35 than MIS 37 (Fig. 8.b). This shows the influence of the Atlantic mitigated forest development during the MIS 31 super interglacial period, compared to the Eastern Mediterranean. In the marine environment, the signal also differs from ODP site 976 with the dominance of *Turborotalita quinqueloba* and higher values for *Globigerinoides ruber* and lesser values for *Neogloboquadrina pachyderma* and *N. incompta*. *T. quinqueloba* is a cold-water taxon which represents very productive waters and reflects the distribution of water masses and the stratification of the water column (Capotondi et al., 2004; Girone et al., 2013; Schiebel and Hemleben, 2017; Bazzicalupo et al., 2018). The central Mediterranean waters therefore appear warmer than in the western basin, but still relatively cool as attested by the presence of *T. quinqueloba*. This is consistent with a lesser influence of the input of cold Atlantic waters through the Gibraltar Strait compare to the Alboran Sea. Throughout the period, *Turborotalita quinqueloba* variations appear to be partly linked to interglacial-glacial variations. At the end of MIS 35, it experienced a drastic decline (from 80 % to 20%). This major decline coincides with changes in the percentage of smectite in the Alboran basin (Fig.8). However, this event is not found in the clay mineralogy from ODP Site 964 in the Ionian Sea (Zhao et al., 2016), nor further east at Site MD90-964 in the Levantine Basin (Zhao et al., 2011). This suggests that the change in marine circulation in the Alboran Sea is likely associated with Atlantic circulation and attenuates eastwards in the Mediterranean.

6.4. Late Early Pleistocene climate in the Southwest Mediterranean

Among the small number of continuous palynological sequences from the Early-Middle Pleistocene period in the Mediterranean, only two provide climate reconstruction data; Palominas site in southern Spain (Altolaquirre et al., 2020) with the Coexistence approach (CA) and Acigöl site in Türkiye (Robles et al., 2026), with a multi-method approach (CAM, MAT, WAPLS, RF and BRT).

The three Western Mediterranean sites present consistent temperature variations (MAAT), which mirror the alternation of glacial and interglacial phases (Fig. 9). The MAT, WAPLS, BRT, and CAM methods show average annual temperatures of around 11 to 12°C for both marine sites (ODP site 976 and Monte San Giorgio) over the entire period, while the RF drops to an average of 10°C (Fig. 9.a.c.). In the continental sequence of Palominas, average temperature varies depending on the method used: 15°C for CA, 12°C for CAM, 10°C for MAT and BRT, 9.5°C for RF and 8°C for WAPLS (Fig.9.b). The lower values appear to be consistent with the altitude of the site, around 780 m above sea level. However, these values seem very cold for annual temperatures in this Southern Mediterranean region, where current values are around 16 to 19°C (Brun et al., 2022). The MAT method shows the greatest fluctuations over time across all sites. For precipitation (PANN), the signal from ODP site 976 mirrors the interglacial-glacial variations with large amplitudes for MAT and WAPLS (Fig. 9.a). The other two sites show wetter conditions for certain glacial phases: MIS 34 at Monte San Giorgio, except for CAM, and MIS 36 at Palominas (Fig. 9.b.c). In addition, MIS 35 appears drier at Palominas than at ODP 976 and Monte San Giorgio. MIS 34 appears quite marked at ODP site 976 and Monte San Giorgio with temperature decreases down to 9°C depending on the method. However, precipitation is reversed at Monte San Giorgio with more rainfall during this cold period. Even though this phase seems to be the most intense, it remains close to the conditions obtained for MIS 36, which are dry at Monte San Giorgio and ODP site 976, but wet at Palominas. This observation is consistent with current knowledge presenting MIS 34 as a very cold phase (Hodell et al., 2023; Margari et al., 2023), but we cannot exclude the possibility that conditions were less extreme in other Mediterranean areas.

CAM reconstruction shows higher precipitation and temperature fluctuations for Monte San Giorgio and Palominas than for ODP 976. Two hypotheses based on pollen data may explain this difference: i) forest diversity at the ODP 976 site is lower than at the other sites as the signal is almost exclusively composed of deciduous *Quercus*; ii) the slightly higher presence of relict taxa, among them *Tsuga*, *Zelkova* or *Cathaya*. The deciduous *Quercus* pollen type encompasses a large number of species that develop within a very wide temperature range (MAAT: 5 and 23°C; Fauquette et al., 1998), which does not allow for effective constraints on climate reconstruction. The other temperate deciduous forest taxa, which are present within narrower temperature ranges, are not present in sufficient quantities to exceed the presence and abundance thresholds, and therefore to constrain the climate estimate. For the other two sites, *Quercus* also dominates the assemblage, but other taxa are sufficiently represented to constrain the model, such as *Ulmus* or *Ostrya* and *Celtis* for the Spanish site, and *Carpinus* for the Italian site. Monte San Giorgio also presents sufficient quantities of Arecaceae to better constrain the reconstruction.

The methods applied to the Palominas dataset (CAM, MAT, BRT, RF, and WAPLS) are consistently colder than the results provided by the Coexistence Approach (CA) (Altolaquirre et al., 2020). They rely on different modern databases: PALEOFLORE for the CA (www.paleoflora.de), the database of modern pollen spectra used by Fauquette et al. (1998) for the CAM and the Eurasian and Mediterranean Current Surface Sample Database (EAPDB; Peyron et al., 1998, 2011, 2013) for the other methods. The difference between the CAM and CA is all

the more striking since they are based on similar concept and parameterization choices (i.e., the removal of taxa present only once, [Altolaguirre et al., 2020](#)). However, unlike the CA, CAM takes into account not only the presence/absence criteria but also the pollen percentages of the taxa, that certainly refines the climatic estimates.




Figure 9. Mean Annual Air Temperature (MAAT) and Annual Precipitation (PANN) reconstructions of three sites in the Western Mediterranean basin between 36 and 38°N: ODP site 976, Palominas and Monte San Giorgio. Five methods were used: Weighted Averaging Partial Least Square (WAPLS), Modern Analog Technique (MAT), Random Forest (RF), Boosted Regression Tree (BRT) and Climatic Amplitude Method (CAM). For Palominas, the Coexistence Approach (CA) method was performed by [Altolaguirre et al. \(2020\)](#). m.b.s.l: metres below sea level; m.a.s.l: metres above sea level. The dotted lines correspond to modern values: for the ODP site 976 the mean value for a 400 km radius is taken from the ERA5 reanalysis dataset (1960–2022), for Palominas and Monte Sangiorgio the values are taken from www.climate-date.org respectively for the cities of Baza and Caltagirone.

6.5. Comparison with climate reconstruction from archaeological/palaeontological sites in South-western Spain

In southern Spain, archaeological faunal data is an additional proxy for climate reconstructions and comparisons with pollen-based climate reconstructions from Palominas and ODP 976. The faunal assemblages come from archaeological sites (Fuente Nueva 3, Barranco Leon and Cueva Victoria) and palaeontological sites (Quibas and El Chaparral) (Fig. 10). Faunal-based climate reconstructions follow three different methods in terms of concepts and types of fauna. All these sites have higher temperature estimates (Fig. 10) than the temperature reconstructions based on pollen assemblages (Fig. 9). Only the results of the Coexistence Approach from Palominas pollen data (Fig. 9) are in the same range of values (15.5°C). On the other hand, pollen-based reconstructions of precipitation are more consistent with results from faunal data (Fig. 10). The WAPLS values in ODP 976 and Palominas (~750 and ~700 mm/year) are close to those reconstructed at Fuente Nueva (738-753 mm/year; Agustí et al. (2009); Blain et al. (2011, 2016)), Barranco León (732-741 mm/year; Agustí et al. (2009); Blain et al. (2011)), Quibas (723 mm/year; Agustí et al. (2009)) and Cueva Victoria (716 mm/year; Agustí et al. (2009)). BRT values (~475 mm/year) in ODP 976 are close to the lowest reconstructed precipitation values at Quibas (417 mm/year; Hernández Fernández et al. (2007)). This probably suggests that climate and vegetation in the southern Mediterranean region were more strongly influenced by, and sensitive to, variations in rainfall and drought than temperatures. Underestimations of temperatures using pollen-based reconstruction methods raise questions about the methodology, the modern databases used, the functioning of each method, and the application of these methods to marine pollen records and to the ancient Quaternary period (except for CAM which was developed for Neogene and Early Quaternary climate reconstruction). In the case of the two marine sites ODP 976 and Monte San Giorgio, temperatures appear to be underestimated. However, this same trend is observed at the continental site of Palominas. Unlike faunal data, ODP 976 records plants in the Alboran Sea from a large area stretching from the coasts to the mountains of both Morocco and southern Spain. The reconstructed climate is therefore an average climate over a large geographical area and not a local one.




Figure 10. Climate reconstruction based on faunal assemblages from archaeological and palaeontological sites in south-eastern Spain. Palaeontological sites are represented by the image of a horse, and archaeological sites are represented by a stone tool. MAAT, MTCO and MTWA are in degrees and PANN in mm per year.

629

630 **Conclusion**

631

632 This high-resolution study of the ODP 976 site provides multiproxy reconstructions of climate change in
633 the South-western Mediterranean at the end of the Early Pleistocene. The sequence continuously documents
634 marine and continental environments during the EMPT, when climate cycles began to lengthen from 41 kyr to 100
635 kyr. This multiproxy approach highlights the synchronicity between the continental and marine response to glacial-
636 interglacial cycles in the Western Mediterranean between MIS 37 and MIS 31 (~1235-1062 ka). Comparisons with
637 other sequences from the Central and Western Mediterranean show major differences between the Iberian
638 Peninsula and southern Italy. Results highlight a marked climatic gradient between the Western and Central
639 Mediterranean during MIS 31, constraining forest development under the influence of the Atlantic climate system,
640 compared to southern Italy, which was less affected by this dynamic. Clay mineralogy shows an abrupt change in
641 smectite and I/K signals, reflecting atmospheric and marine conditions in the Alboran Sea around 1140 ka, i.e., at
642 the transition between MIS 35 and 34. Multi-method climate reconstructions based on foraminifera (MAT) and
643 pollen assemblages (CAM, MAT, WAPLS, FR and BRT) show variations in temperature and precipitation
644 mirroring climate cycles. However, some glacial periods appear wetter than some interglacial periods, depending
645 on the method used. MIS 34 reconstructions appear more pronounced in terms of temperature and precipitation,
646 compared to other glacial phases. Pollen-based reconstructions systematically show colder temperatures than those
647 based on faunal assemblages from palaeontological and archaeological sites in southern Spain, which could be
648 explained by local vs regional climate reconstruction. On the other hand, precipitation reconstruction seems more
649 consistent for the two types of archives, suggesting that vegetation changes are driven more by precipitation than
650 temperature fluctuations in this part of the Southern Mediterranean. These results enhance our understanding of
651 the environmental changes that occurred in the Western Mediterranean during the period between MIS 37-31,
652 providing land-sea information. This type of research must now be extended to the entire EMPT in order to better
653 understand the climate changes that took place at a time of major modification of the Earth's climate system, and
654 to comprehend the environmental context in which the first hominins settled in Western Europe.

655 **Competing interests**

656 One of the authors (O. Peyron) is a member of the editorial board of *Climate of the Past*. The other authors declare
657 that they have no competing interests.

658 **Data availability**

659 Data will be made available to the corresponding author upon request before being made available on Pangea..

660 **Author contribution**

661 MC, VL, NCN, OP, SF designed the project. MC performed the palynological analyses and MF and JL the
662 foraminifera assemblages. PR and MF conducted the isotope analyses used for the age model and VBR and MD
663 the clay mineralogy analysis. LD conducted all the palynological HF attack treatment. MC, MR, OP, AM and SF
664 performed the climate reconstruction on pollen data and the statistical tests. MF applied the SST reconstruction

method to foraminifera assemblages. SJ, YA, JPS gave permission to use their data and contributed to the interpretation of results. All the co-authors participated in writing this manuscript and in the discussion of results.

Acknowledgments

We are grateful to the ERC LATEUROPE project (n°101052653), the Centre National de la Recherche Scientifique (CNRS) and the Museum national d'Histoire naturelle (MNHN) for their financial support. This research was also supported by the Franco-Italian PHC Galileo project BioMed (project FR n°48933VE/project IT n°2023-G23_53). We thank the PANOPLY analytical platform at LSCE for facilities for carrying out isotopic measurements and the study of the foraminifera assemblages, and the Ocean Drilling Program (ODP) for making available material from the core ODP leg 161 site 976. We thank the CARMIN analytical platform at the University of Lille for facilities for carrying out clay mineralogical measurements. We would also like to thank Angela Girone for sharing and giving us permission to use her data. Finally, we thank Lucas Dugerdil and Léa d'Oliveira (ISEM) for providing us their code to performed climate reconstructions (MAT, WAPLS, BRT and RF). We thank the editor Antje Voelker, Timme Donders and an anonymous reviewer whose relevant comments helped significantly improve the manuscript quality. This is an ISEM publication n° XXXX.

Financial support

This project was financially supported by the ERC LATEUROPE project (n°101052653), by the Centre National de la Recherche Scientifique (CNRS) and by the Museum National d'Histoire Naturelle (MNHN)

References

- Abbate, E. and Sagri, M.: Early to Middle Pleistocene *Homo* dispersals from Africa to Eurasia: Geological, climatic and environmental constraints, *Quaternary International*, 267, 3–19, <https://doi.org/10.1016/j.quaint.2011.02.043>, 2012.
- Agustí, J., Blain, H.-A., Cuenca-Bescós, G., and Bailon, S.: Climate forcing of first hominid dispersal in Western Europe, *Journal of Human Evolution*, 57, 815–821, <https://doi.org/10.1016/j.jhevol.2009.06.005>, 2009.
- Alonso, B. and Maldonado, A.: Late Quaternary sedimentation patterns of the Ebro turbidite systems (northwestern Mediterranean): Two styles of deep-sea deposition, *Marine Geology*, 95, 353–377, [https://doi.org/10.1016/0025-3227\(90\)90124-3](https://doi.org/10.1016/0025-3227(90)90124-3), 1990.
- Alonso, B., Ercilla, G., Martínez-Ruiz, F., Baraza, J., and Galimont, A.: Pliocene-Pleistocene sedimentary facies at site 976: Depositional history in the northwestern Alboran sea, in: *Proceedings of the Ocean Drilling Program, 161 Scientific Results*, vol. 161, edited by: Zahn, R., Comas, M. C., and Klaus, A., Ocean Drilling Program, <https://doi.org/10.2973/odp.proc.sr.161.1999>, 1999.
- Altolaguirre, Y., Bruch, A. A., and Gibert, L.: A long Early Pleistocene pollen record from Baza Basin (SE Spain): Major contributions to the palaeoclimate and palaeovegetation of Southern Europe, *Quaternary Science Reviews*, 231, 106199, <https://doi.org/10.1016/j.quascirev.2020.106199>, 2020.
- Ambar, I. and Howe, M. R.: Observations of the Mediterranean outflow—I mixing in the Mediterranean outflow, *Deep Sea Research Part A. Oceanographic Research Papers*, 26, 535–554, [https://doi.org/10.1016/0198-0149\(79\)90095-5](https://doi.org/10.1016/0198-0149(79)90095-5), 1979.

Bahr, A., Kaboth, S., Hodell, D., Zeeden, C., Fiebig, J., and Friedrich, O.: Oceanic heat pulses fueling moisture transport towards continental Europe across the mid-Pleistocene transition, *Quaternary Science Reviews*, 179, 48–58, <https://doi.org/10.1016/j.quascirev.2017.11.009>, 2018.

Bahr, A., Catunda, M. C. A., and Friedrich, O.: The first appearance of Glacial North Atlantic Intermediate Water (GNAIW) during the Mid-Pleistocene transition, *Global and Planetary Change*, 218, 103977, <https://doi.org/10.1016/j.gloplacha.2022.103977>, 2022.

Baringer, M. O. and Price, J. F.: A review of the physical oceanography of the Mediterranean outflow, *Marine Geology*, 155, 63–82, [https://doi.org/10.1016/S0025-3227\(98\)00141-8](https://doi.org/10.1016/S0025-3227(98)00141-8), 1999.

Barker, S., Starr, A., Conn, S., Lordsmith, S., Owen, L., Nederbragt, A., Hemming, S., Hall, I., and Levay, L.: Persistent influence of precession on northern ice sheet variability since the early Pleistocene, *Science*, 376, 961–967, <https://doi.org/10.1126/science.abm4033>, 2022.

Bazzicalupo, P., Maiorano, P., Girone, A., Marino, M., Combourieu-Nebout, N., and Incarbona, A.: High-frequency climate fluctuations over the last deglaciation in the Alboran Sea, Western Mediterranean: Evidence from calcareous plankton assemblages, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 506, 226–241, <https://doi.org/10.1016/j.palaeo.2018.06.042>, 2018.

Beck, H. E., McVicar, T. R., Vergopolan, N., Berg, A., Lutsko, N. J., Dufour, A., Zeng, Z., Jiang, X., Van Dijk, A. I. J. M., and Miralles, D. G.: High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections, *Sci Data*, 10, 724, <https://doi.org/10.1038/s41597-023-02549-6>, 2023.

Benabid, A.: Les écosystèmes forestiers, préforestiers et presteppiques du Maroc : diversité, répartition biogéographique et problèmes posés par leur aménagement, *Forêt Méditerranéenne*, VII, 53–64, 1985.

Berends, C. J., Köhler, P., Lourens, L. J., and van de Wal, R. S. W.: On the Cause of the Mid-Pleistocene Transition, *Reviews of Geophysics*, 59, e2020RG000727, <https://doi.org/10.1029/2020RG000727>, 2021.

Blain, H.-A., Bailon, S., and Agustí, J.: Anurans and squamate reptiles from the latest early Pleistocene of Almenara-Casablanca-3 (Castellón, East of Spain). Systematic, climatic and environmental considerations, 2007.

Blain, H.-A., Bailon, S., Cuenca-Bescós, G., Arsuaga, J. L., Bermúdez De Castro, J. M., and Carbonell, E.: Long-term climate record inferred from early-middle Pleistocene amphibian and squamate reptile assemblages at the Gran Dolina Cave, Atapuerca, Spain, *Journal of Human Evolution*, 56, 55–65, <https://doi.org/10.1016/j.jhevol.2008.08.020>, 2009.

Blain, H.-A., Bailon, S., Agustí, J., Martínez-Navarro, B., and Toro, I.: Paleoenvironmental and paleoclimatic proxies to the Early Pleistocene hominids of Barranco León D and Fuente Nueva 3 (Granada, Spain) by means of their amphibian and reptile assemblages, *Quaternary International*, 243, 44–53, <https://doi.org/10.1016/j.quaint.2010.12.031>, 2011.

Blain, H.-A., Lozano-Fernández, I., Agustí, J., Bailon, S., Menéndez Granda, L., Espígares Ortiz, M. P., Ros-Montoya, S., Jiménez Arenas, J. M., Toro-Moyano, I., Martínez-Navarro, B., and Sala, R.: Refining upon the climatic background of the Early Pleistocene hominid settlement in western Europe: Barranco León and Fuente Nueva-3 (Guadix-Baza Basin, SE Spain), *Quaternary Science Reviews*, 144, 132–144, <https://doi.org/10.1016/j.quascirev.2016.05.020>, 2016.

Bout-Roumazeilles, V., Cortijo, E., Labeyrie, L., and Debrabant, P.: Clay mineral evidence of nepheloid layer contributions to the Heinrich layers in the northwest Atlantic, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 146, 211–228, [https://doi.org/10.1016/S0031-0182\(98\)00137-0](https://doi.org/10.1016/S0031-0182(98)00137-0), 1999.

Bout-Roumazeilles, V., Combourieu Nebout, N., Peyron, O., Cortijo, E., Landais, A., and Masson-Delmotte, V.: Connection between South Mediterranean climate and North African atmospheric circulation during the last 50,000yrBP North Atlantic cold events, *Quaternary Science Reviews*, 26, 3197–3215, <https://doi.org/10.1016/j.quascirev.2007.07.015>, 2007.

Bout-Roumazeilles, V., Combourieu-Nebout, N., Desprat, S., Siani, G., Turon, J.-L., and Essallami, L.: Tracking atmospheric and riverine terrigenous supplies variability during the last glacial and the Holocene in central Mediterranean, *Climate of the Past*, 9, 1065–1087, <https://doi.org/10.5194/cp-9-1065-2013>, 2013.

Breiman, L.: Random Forests, *Machine Learning*, 45, 5–32, <https://doi.org/10.1023/A:1010933404324>, 2001.

Breiman, L., Cutler, A., Liaw, A., and Wiener, M.: Package ‘randomForest’, 4.7–1.2, 2024.

Brown, G. and Brindley, G. W.: X-ray Diffraction Procedures for Clay Mineral Identification, in: *Crystal Structures of Clay Minerals and their X-Ray Identification*, edited by: Brown, G. and Brindley, G. W., Mineralogical Society, London, 305–359, 1980.

Brun, P., Zimmermann, N. E., Hari, C., and Karger, D. N.: CHELSA-BIOCLIM+ A novel set of global climate-related predictors at kilometre-resolution, <https://doi.org/https://www.doi.org/10.16904/envdat.332>, 2022.

Bryden, H. L. and Stommel, H. M.: Limiting processes that determine basic features of the circulation in the Mediterranean Sea, *Oceanologia acta*, 7, 289–296, 1984.

Bulian, F.: The late Miocene – early Pliocene offshore onshore sedimentary records in the vicinity of Gibraltar, Universidad de Salamanca, 2022.

Capotondi, L., Soroldoni, E., and Principato, M. S.: Late Quaternary planktonic foraminiferal distributions: problems related to size fraction, *Proceedings of the First Italian Meeting on Environmental*, 2004.

Caquineau, S., Gaudichet, A., Gomes, L., Magonthier, M.-C., and Chatenet, B.: Saharan dust: Clay ratio as a relevant tracer to assess the origin of soil-derived aerosols, *Geophysical Research Letters*, 25, 983–986, <https://doi.org/10.1029/98GL00569>, 1998.

Carbonell, E., Bermúdez de Castro, J. M., Parés, J. M., Pérez-González, A., Cuenca-Bescós, G., Ollé, A., Mosquera, M., Huguet, R., van der Made, J., Rosas, A., Sala, R., Vallverdú, J., García, N., Granger, D. E., Martínón-Torres, M., Rodríguez, X. P., Stock, G. M., Vergès, J. M., Allué, E., Burjachs, F., Cáceres, I., Canals, A., Benito, A., Díez, C., Lozano, M., Mateos, A., Navazo, M., Rodríguez, J., Rosell, J., and Arsuaga, J. L.: The first hominin of Europe, *Nature*, 452, 465–469, <https://doi.org/10.1038/nature06815>, 2008.

Catrain, M., Combourieu Nebout, N., Lebreton, V., Fauquette, S., Peyron, O., Fries, M., Richard, P., Dubost, L., Joannin, S., Suc, J.-P., Paquier Comas, E., Lepelletier, J., and Moncel, M.-H.: Vegetation and climate dynamics in the south-western mediterranean during MIS 37–31 (~1.25 - ~1.06 Ma): Insights from the marine core ODP site 976, *Quaternary Science Reviews*, 369, <https://doi.org/https://doi.org/10.1016/j.quascirev.2025.109635>, 2025.

Chalk, T. B., Hain, M. P., Foster, G. L., Rohling, E. J., Sexton, P. F., Hendry, K. R., Badger, M. P. S., Cherry, S. G., Hasenfratz, A. P., Haug, G. H., Jaccard, S. L., Martínez-García, A., Pälike, H., Pancost, R. D., and Wilson, P. A.: Causes of ice age intensification across the Mid-Pleistocene Transition, *Proceedings of the National Academy of Sciences of the United States of America*, 114, 13114–13119, <https://doi.org/10.1073/pnas.1702143114>, 2017.

Chamley, H.: *Clay Sedimentology*, Springer Berlin, Heidelberg., 1989.

Charton, L., Combourieu-Nebout, N., Bertini, A., Lebreton, V., Peyron, O., Robles, M., Sassoon, D., and Moncel, M.-H.: Vegetation and climate changes during the Middle to Upper Palaeolithic transition in the southwestern

781 Mediterranean: What happened to the last Neanderthals during Heinrich stadial 4?, *Quaternary Science Reviews*,
782 359, 109345, <https://doi.org/10.1016/j.quascirev.2025.109345>, 2025.

783 Chevalier, M., Davis, B. A. S., Heiri, O., Seppä, H., Chase, B. M., Gajewski, K., Lacourse, T., Telford, R. J.,
784 Finsinger, W., Guiot, J., Kühl, N., Maezumi, S. Y., Tipton, J. R., Carter, V. A., Brussel, T., Phelps, L. N., Dawson,
785 A., Zanon, M., Vallé, F., Nolan, C., Mauri, A., de Vernal, A., Izumi, K., Holmström, L., Marsicek, J., Goring, S.,
786 Sommer, P. S., Chaput, M., and Kupriyanov, D.: Pollen-based climate reconstruction techniques for late
787 Quaternary studies, *Earth-Science Reviews*, 210, 103384, <https://doi.org/10.1016/j.earscirev.2020.103384>, 2020.

788 Clark, P. U., Archer, D., Pollard, D., Blum, J. D., Rial, J. A., Brovkin, V., Mix, A. C., Pisias, N. G., and Roy, M.:
789 The middle Pleistocene transition: characteristics, mechanisms, and implications for long-term changes in
790 atmospheric pCO₂, *Quaternary Science Reviews*, 25, 3150–3184,
791 <https://doi.org/10.1016/j.quascirev.2006.07.008>, 2006.

792 Comas, M. C., Zahn, R., and Klaus, A. (Eds.): SITE 976, in: *Proceedings of the Ocean Drilling Program, Initial*
793 *Reports*, 179–297, 1996.

794 Cucart-Mora, C., Lombao, D., Pineda, A., Boemke, B., Mecozzi, B., and Moncel, M.-H.: The First Occupations
795 of Western Europe: Dispersals and Population Dynamics in the Early to Middle Pleistocene, *J Archaeol Method*
796 *Theory*, 33, 19, <https://doi.org/10.1007/s10816-025-09752-2>, 2026.

797 Daget, P.: Le bioclimat méditerranéen: Caracteres generaux, modes de caracterisation, *Vegetatio*, 34, 1–20,
798 <https://doi.org/10.1007/BF00119883>, 1977.

799 Davis, B. A. S., Chevalier, M., Sommer, P., Carter, V. A., Finsinger, W., Mauri, A., Phelps, L. N., Zanon, M.,
800 Abegglen, R., Åkesson, C. M., Alba-Sánchez, F., Anderson, R. S., Antipina, T. G., Atanassova, J. R., Beer, R.,
801 Belyanina, N. I., Blyakharchuk, T. A., Borisova, O. K., Bozilova, E., Bukreeva, G., Bunting, M. J., Clò, E.,
802 Colombaroli, D., Combourieu-Nebout, N., Desprat, S., Di Rita, F., Djamali, M., Edwards, K. J., Fall, P. L.,
803 Feurdean, A., Fletcher, W., Florenzano, A., Furlanetto, G., Gaceur, E., Galimov, A. T., Galka, M., García-
804 Moreiras, I., Giesecke, T., Grindean, R., Guido, M. A., Gvozdeva, I. G., Herzsuh, U., Hjelle, K. L., Ivanov, S.,
805 Jahns, S., Jankovska, V., Jiménez-Moreno, G., Karpińska-Kołaczek, M., Kitaba, I., Kołaczek, P., Lapteva, E. G.,
806 Latałowa, M., Lebreton, V., Leroy, S., Leydet, M., Lopatina, D. A., López-Sáez, J. A., Lotter, A. F., Magri, D.,
807 Marinova, E., Matthias, I., Mavridou, A., Mercuri, A. M., Mesa-Fernández, J. M., Mikishin, Y. A., Milecka, K.,
808 Montanari, C., Morales-Molino, C., Mrotzek, A., Muñoz Sobrino, C., Naidina, O. D., Nakagawa, T., Nielsen, A.
809 B., Novenko, E. Y., Panajiotidis, S., Panova, N. K., Papadopoulou, M., Pardoe, H. S., Pędziszewska, A., Petrenko,
810 T. I., Ramos-Román, M. J., Ravazzi, C., Rösch, M., Ryabogina, N., Sabariego Ruiz, S., Salonen, J. S., Sapelko,
811 T. V., Schofield, J. E., Seppä, H., Shumilovskikh, L., Stivrins, N., Stojakowits, P., Svobodova Svitavská, H.,
812 Święta-Musznicka, J., Tantau, I., Tinner, W., Tobolski, K., Tonkov, S., Tsakiridou, M., et al.: The Eurasian
813 Modern Pollen Database (EMPD), version 2, *Earth System Science Data*, 12, 2423–2445,
814 <https://doi.org/10.5194/essd-12-2423-2020>, 2020.

815 De'ath, G.: Boosted Trees for Ecological Modeling and Prediction, *Ecology*, 88, 243–251,
816 [https://doi.org/10.1890/0012-9658\(2007\)88%5B243:BTFEMA%5D2.0.CO;2](https://doi.org/10.1890/0012-9658(2007)88%5B243:BTFEMA%5D2.0.CO;2), 2007.

817 DeConto, R. M., Pollard, D., and Kowalewski, D.: Modeling Antarctic ice sheet and climate variations during
818 Marine Isotope Stage 31, *Global and Planetary Change*, 88–89, 45–52,
819 <https://doi.org/https://doi.org/10.1016/j.gloplacha.2012.03.003>, 2012.

820 Dominik, J. and Stoffers, P.: The influence of late Quaternary stagnations on clay sedimentation in the eastern
 821 Mediterranean sea, *Geol Rundsch*, 68, 302–317, <https://doi.org/10.1007/BF01821133>, 1979.
 822 Donders, T., Panagiotopoulos, K., Koutsodendris, A., Bertini, A., Mercuri, A. M., Masi, A., Combourieu-Nebout,
 823 N., Joannin, S., Kouli, K., Kousis, I., Peyron, O., Torri, P., Florenzano, A., Francke, A., Wagner, B., and Sadori,
 824 L.: 1.36 million years of Mediterranean forest refugium dynamics in response to glacial–interglacial cycle strength,
 825 *Proceedings of the National Academy of Sciences*, 118, e2026111118, <https://doi.org/10.1073/pnas.2026111118>,
 826 2021.
 827 Dugerdil, L., Joannin, S., Peyron, O., Jouffroy-Bapicot, I., Vannière, B., Boldgiv, B., Unkelbach, J., Behling, H.,
 828 and Ménot, G.: Climate reconstructions based on GDGT and pollen surface datasets from Mongolia and Baikal
 829 area: calibrations and applicability to extremely cold–dry environments over the Late Holocene, *Clim. Past*, 17,
 830 1199–1226, <https://doi.org/10.5194/cp-17-1199-2021>, 2021.
 831 Duval, M., Arnold, L. J., Bahain, J.-J., Parés, J. M., Demuro, M., Falguères, C., Shao, Q., Voinchet, P., Arnaud,
 832 J., Berto, C., Berruti, G. L. F., Daffara, S., Sala, B., and Arzarello, M.: Re-examining the earliest evidence of
 833 human presence in western Europe: New dating results from Pirro Nord (Italy), *Quaternary Geochronology*, 82,
 834 101519, <https://doi.org/10.1016/j.quageo.2024.101519>, 2024.
 835 Ehrmann, W., Schmiedl, G., Hamann, Y., Kuhnt, T., Hemleben, C., and Siebel, W.: Clay minerals in late glacial
 836 and Holocene sediments of the northern and southern Aegean Sea, *Palaeogeography, Palaeoclimatology,*
 837 *Palaeoecology*, 249, 36–57, <https://doi.org/10.1016/j.palaeo.2007.01.004>, 2007.
 838 Elenga, H., Peyron, O., Bonnefille, R., Jolly, D., Cheddadi, R., Guiot, J., Andrieu, V., Bottema, S., Buchet, G., De
 839 Beaulieu, J. -L., Hamilton, A. C., Maley, J., Marchant, R., Perez-Obiol, R., Reille, M., Riollet, G., Scott, L., Straka,
 840 H., Taylor, D., Van Campo, E., Vincens, A., Laarif, F., and Jonson, H.: Pollen-based biome reconstruction for
 841 southern Europe and Africa 18,000 yr BP, *Journal of Biogeography*, 27, 621–634, <https://doi.org/10.1046/j.1365-2699.2000.00430.x>, 2000.
 842
 843 Elith, J., Leathwick, J. R., and Hastie, T.: A working guide to boosted regression trees, *Journal of Animal Ecology*,
 844 77, 802–813, <https://doi.org/10.1111/j.1365-2656.2008.01390.x>, 2008.
 845 Ercilla, G., Alonso, B., and Baraza, J.: Sedimentary evolution of the northwestern Alboran Sea during the
 846 Quaternary, *Geo-Marine Letters*, 12, 144–149, <https://doi.org/10.1007/BF02084925>, 1992.
 847 Fabres, J., Calafat, A., Sanchez-Vidal, A., Canals, M., and Heussner, S.: Composition and spatio-temporal
 848 variability of particle fluxes in the Western Alboran Gyre, Mediterranean Sea, *Journal of Marine Systems*, 33–34,
 849 431–456, [https://doi.org/10.1016/S0924-7963\(02\)00070-2](https://doi.org/10.1016/S0924-7963(02)00070-2), 2002.
 850 Fauquette, S., Guiot, J., and Suc, J.-P.: A method for climatic reconstruction of the Mediterranean Pliocene using
 851 pollen data, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 144, 183–201, [https://doi.org/10.1016/S0031-0182\(98\)00083-2](https://doi.org/10.1016/S0031-0182(98)00083-2), 1998.
 852
 853 Foucault, A. and Mélières, F.: Palaeoclimatic cyclicity in central Mediterranean Pliocene sediments: the
 854 mineralogical signal, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 158, 311–323,
 855 [https://doi.org/10.1016/S0031-0182\(00\)00056-0](https://doi.org/10.1016/S0031-0182(00)00056-0), 2000.
 856 Garzanti, E., Andò, S., Vezzoli, G., Megid, A., and Kammar, A.: Petrology of Nile River sands (Ethiopia and
 857 Sudan): Sediment budgets and erosion patterns, *Earth and Planetary Science Letters*, 252, 327–341,
 858 <https://doi.org/10.1016/j.epsl.2006.10.001>, 2006.

859 Girone, A., Capotondi, L., Ciaranfi, N., Di Leo, P., Lirer, F., Maiorano, P., Marino, M., Pelosi, N., and Pulice, I.:
860 Paleoenvironmental changes at the lower Pleistocene Montalbano Jonico section (southern Italy): Global versus
861 regional signals, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 371, 62–79,
862 <https://doi.org/10.1016/j.palaeo.2012.12.017>, 2013.

863 Grousset, F. E., Joron, J. L., Biscaye, P. E., Latouche, C., Treuil, M., Maillet, N., Faugères, J. C., and Gonthier,
864 E.: Mediterranean outflow through the Strait of Gibraltar since 18,000 Years B.P.: Mineralogical and geochemical
865 arguments, *Geo-Marine Letters*, 8, 25–34, <https://doi.org/10.1007/BF02238003>, 1988.

866 Guerzoni, S., Molinaroli, E., Rossini, P., Rampazzo, G., Quarantotto, G., De Falco, G., and Cristini, S.: Role of
867 desert aerosol in metal fluxes in the Mediterranean area, *Chemosphere*, 39, 229–246,
868 [https://doi.org/10.1016/S0045-6535\(99\)00105-8](https://doi.org/10.1016/S0045-6535(99)00105-8), 1999.

869 Guijarro, J. A., Conde, J., Campins, J., Orro, M. L., and Picornell, M. Á.: Atlas de clima marítimo (0-52°N, 35°W-
870 12°E; 1981-2010), Agencia Estatal de Meteorología, <https://doi.org/10.31978/281-15-028-8>, 2015.

871 Guiot, J.: Methodology of palaeoclimatic reconstruction from pollen in France: *Palaeogeography,*
872 *Palaeogeography, Palaeoclimatology, Palaeoecology*, 80, 49–69, [https://doi.org/10.1016/0031-0182\(90\)90033-4](https://doi.org/10.1016/0031-0182(90)90033-4),
873 1990.

874 Guo, Q., Li, B., Voelker, A. H. L., and Kim, J.-K.: Mediterranean Outflow Water dynamics across the middle
875 Pleistocene transition based on a 1.3 million-year benthic foraminiferal record off the Portuguese margin,
876 *Quaternary Science Reviews*, 247, 106567, <https://doi.org/10.1016/j.quascirev.2020.106567>, 2020.

877 Hamann, Y., Ehrmann, W., Schmiedl, G., and Kuhnt, T.: Modern and late Quaternary clay mineral distribution in
878 the area of the SE Mediterranean Sea, *Quat. res.*, 71, 453–464, <https://doi.org/10.1016/j.yqres.2009.01.001>, 2009.

879 Hassouni, B. and El Hmadi, A.: Intérêts de l'étude des minéraux argileux en mer d'Alboran, *Notes et Mém. Serv.*
880 *géol. Maroc*, 113–120, 2007.

881 Head, M. J. and Gibbard, P. L.: Early–Middle Pleistocene transitions: Linking terrestrial and marine realms,
882 *Quaternary International*, 389, 7–46, <https://doi.org/10.1016/j.quaint.2015.09.042>, 2015.

883 Herbert, T. D.: The Mid-Pleistocene Climate Transition, *Annual Review of Earth and Planetary Sciences*, 51, 389–
884 418, <https://doi.org/10.1146/annurev-earth-032320-104209>, 2023.

885 Hijmans, R. J., Phillips, S., Leathwick, J., and Elith, J.: Package ‘dismo,’ , 1.3-16, 2024.

886 Hodell, D. A., Crowhurst, S. J., Lourens, L., Margari, V., Nicolson, J., Rolfe, J. E., Skinner, L. C., Thomas, N. C.,
887 Tzedakis, P. C., Mleneck-Vautravers, M. J., and Wolff, E. W.: A 1.5-million-year record of orbital and millennial
888 climate variability in the North Atlantic, *Climate of the Past*, 19, 607–636, [https://doi.org/10.5194/cp-19-607-](https://doi.org/10.5194/cp-19-607-2023)
889 2023, 2023.

890 Holeman, J.: The Sediment Yield of Major Rivers of the World, *Water resources research*, 4, 1968.

891 Joannin, S., Ciaranfi, N., and Stefanelli, S.: Vegetation changes during the late Early Pleistocene at Montalbano
892 Jonico (Province of Matera, southern Italy) based on pollen analysis, *Palaeogeography, Palaeoclimatology,*
893 *Palaeoecology*, 270, 92–101, <https://doi.org/10.1016/j.palaeo.2008.08.017>, 2008.

894 Joannin, S., Bassinot, F., Nebout, N. C., Peyron, O., and Beaudouin, C.: Vegetation response to obliquity and
895 precession forcing during the Mid-Pleistocene Transition in Western Mediterranean region (ODP site 976),
896 *Quaternary Science Reviews*, 30, 280–297, <https://doi.org/10.1016/j.quascirev.2010.11.009>, 2011.

897 Juggins, S.: Package ‘rioja,’ , 1.0-7, 2024.

898 Köppen, W.: Das geographische System der Klimate, in: Handbuch der Klimatologie, vol. I, edited by: Köppen,
899 W. and Geiger, R., Verlag von Gebrüder Borntraeger, Berlin, 44, 1936.

900 Krom, M. D., Michard, A., Cliff, R. A., and Strohle, K.: Sources of sediment to the Ionian Sea and western
901 Levantine basin of the Eastern Mediterranean during S-1 sapropel times, *Marine Geology*, 160, 45–61,
902 [https://doi.org/10.1016/S0025-3227\(99\)00015-8](https://doi.org/10.1016/S0025-3227(99)00015-8), 1999.

903 Leblanc, M.: Étude des périodes humides Africaines enregistrées au cours des deux derniers cycles climatiques à
904 partir de la caractérisation de poussières Sahariennes déposées en Méditerranée Occidentale et dans l’océan
905 Atlantique Tropical Nord-Est, Université Paris-Saclay, 2022.

906 Leroy, S. A. G., Arpe, K., and Mikolajewicz, U.: Vegetation context and climatic limits of the Early Pleistocene
907 hominin dispersal in Europe, *Quaternary Science Reviews*, 30, 1448–1463,
908 <https://doi.org/10.1016/j.quascirev.2010.01.017>, 2011.

909 Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records,
910 *Paleoceanography*, 20, <https://doi.org/10.1029/2004PA001071>, 2005.

911 Macías, D., Bruno, M., Echevarría, F., Vázquez, A., and García, C. M.: Meteorologically-induced mesoscale
912 variability of the North-western Alboran Sea (southern Spain) and related biological patterns, *Estuarine, Coastal
913 and Shelf Science*, 78, 250–266, <https://doi.org/10.1016/j.ecss.2007.12.008>, 2008.

914 Magri, D., Di Rita, F., Aranbarri, J., Fletcher, W., and González-Sampériz, P.: Quaternary disappearance of tree
915 taxa from Southern Europe: Timing and trends, *Quaternary Science Reviews*, 163, 23–55,
916 <https://doi.org/10.1016/j.quascirev.2017.02.014>, 2017.

917 Margari, V., Hodell, D. A., Parfitt, S. A., Ashton, N. M., Grimalt, J. O., Kim, H., Yun, K.-S., Gibbard, P. L.,
918 Stringer, C. B., Timmermann, A., and Tzedakis, P. C.: Extreme glacial cooling likely led to hominin depopulation
919 of Europe in the Early Pleistocene, *Science*, 381, 693–699, <https://doi.org/10.1126/science.adf4445>, 2023.

920 Marquina-Blasco, R., Sánchez-Bandera, C., López-García, J. M., Martínez-Ortí, A., Sevilla, P., Ruiz-Sánchez, F.
921 J., and Blain, H.-A.: Palaeoecological characterization of the Pliocene-Pleistocene transition on the Mediterranean
922 littoral area: Almenara-Casablanca (Castellón, eastern Spain), *Quaternary Science Reviews*, 352, 109154,
923 <https://doi.org/10.1016/j.quascirev.2024.109154>, 2025.

924 Martin, J.-M. and Milliman, J. D.: EROS 2000 (European River Ocean System). The western Mediterranean : an
925 introduction, *Deep-sea res., Part 2, Top. stud. oceanogr*, 44, 521–529, 1997.

926 Martínez-Botí, M. A., Foster, G. L., Chalk, T. B., Rohling, E. J., Sexton, P. F., Lunt, D. J., Pancost, R. D., Badger,
927 M. P. S., and Schmidt, D. N.: Plio-Pleistocene climate sensitivity evaluated using high-resolution CO₂ records,
928 *Nature*, 518, 49–54, <https://doi.org/10.1038/nature14145>, 2015.

929 Maslin, M. A. and Brierley, C. M.: The role of orbital forcing in the Early Middle Pleistocene Transition,
930 *Quaternary International*, 47–55, <https://doi.org/http://dx.doi.org/10.1016/j.quaint.2015.01.047>, 2015.

931 Médail, F.: Plant Biogeography and Vegetation Patterns of the Mediterranean Islands, *Bot. Rev.*, 88, 63–129,
932 <https://doi.org/10.1007/s12229-021-09245-3>, 2022.

933 Mega, A., Rodrigues, T., Salgueiro, E., Padilha, M., Kuhnert, H., and Voelker, A. H. L.: The Early–Middle
934 Pleistocene Transition in the Gulf of Cadiz (NE Atlantic) – an interplay between subtropical gyre and extremely
935 cold surface waters, *Clim. Past*, 21, 919–939, <https://doi.org/10.5194/cp-21-919-2025>, 2025.

936 Melles, M., Brigham-Grette, J., Minyuk, P. S., Nowaczyk, N. R., Wennrich, V., DeConto, R. M., Anderson, P.
937 M., Andreev, A. A., Coletti, A., Cook, T. L., Haltia-Hovi, E., Kukkonen, M., Lozhkin, A. V., Rosén, P., Tarasov,

P., Vogel, H., and Wagner, B.: 2.8 Million Years of Arctic Climate Change from Lake El'gygytgyn, NE Russia, *Science*, 337, 315–320, <https://doi.org/10.1126/science.1222135>, 2012.

Messenger, E., Lebreton, V., Marquer, L., Russo-Ermolli, E., Orain, R., Renault-Miskovsky, J., Lordkipanidze, D., Despriée, J., Peretto, C., and Arzarello, M.: Palaeoenvironments of early hominins in temperate and Mediterranean Eurasia: new palaeobotanical data from Palaeolithic key-sites and synchronous natural sequences, *Quaternary Science Reviews*, 30, 1439–1447, <https://doi.org/10.1016/j.quascirev.2010.09.008>, 2011.

Milliman, J. D. and Syvitski, J. P. M.: Geomorphic/Tectonic Control of Sediment Discharge to the Ocean: The Importance of Small Mountainous Rivers, *The Journal of Geology*, 100, 525–544, <https://doi.org/10.1086/629606>, 1992.

Millot, C.: Heterogeneities of in- and out-flows in the Mediterranean Sea, *Progress in Oceanography*, 120, 254–278, <https://doi.org/10.1016/j.pocean.2013.09.007>, 2014.

Moal-Darrigade, P., Ducassou, E., Bout-Roumazeilles, V., Hanquiez, V., Perello, M.-C., Mulder, T., and Giraudeau, J.: Source-to-sink pathways of clay minerals in the cadiz contourite system over the last 25 kyrs: The segregational role of mediterranean outflow water, *Marine Geology*, 443, 106697, <https://doi.org/10.1016/j.margeo.2021.106697>, 2022.

Moreno, A., Cacho, I., Canals, M., Grimalt, J. O., Sánchez-Goñi, M. F., Shackleton, N., and Sierro, F. J.: Links between marine and atmospheric processes oscillating on a millennial time-scale. A multi-proxy study of the last 50,000yr from the Alboran Sea (Western Mediterranean Sea), *Quaternary Science Reviews*, 24, 1623–1636, <https://doi.org/10.1016/j.quascirev.2004.06.018>, 2005.

Moyano, I. T., Barsky, D., Cauche, D., Celiberti, V., Grégoire, S., Lebegue, F., Moncel, M. H., and De Lumley, H.: The archaic stone tool industry from Barranco León and Fuente Nueva 3, (Orce, Spain): Evidence of the earliest hominin presence in southern Europe, *Quaternary International*, 243, 80–91, <https://doi.org/10.1016/j.quaint.2010.12.011>, 2011.

Oliveira, D., Sánchez Goñi, M. F., Naughton, F., Polanco-Martínez, J. M., Jimenez-Espejo, F. J., Grimalt, J. O., Martrat, B., Voelker, A. H. L., Trigo, R., Hodell, D., Abrantes, F., and Desprat, S.: Unexpected weak seasonal climate in the western Mediterranean region during MIS 31, a high-insolation forced interglacial, *Quaternary Science Reviews*, 161, 1–17, <https://doi.org/10.1016/j.quascirev.2017.02.013>, 2017.

d'Oliveira, L., Dugerdil, L., Ménot, G., Evin, A., Muller, S. D., Ansanay-Alex, S., Azuara, J., Bonnet, C., Bremond, L., Shah, M., and Peyron, O.: Reconstructing 15 000 years of southern France temperatures from coupled pollen and molecular (branched glycerol dialkyl glycerol tetraether) markers (Canroute, Massif Central), *Climate of the Past*, 19, 2127–2156, <https://doi.org/10.5194/cp-19-2127-2023>, 2023.

d'Oliveira, L., Joannin, S., Ménot, G., Combourieu-Nebout, N., Dugerdil, L., Blache, M., Robles, M., Florenzano, A., Masi, A., Mercuri, A. M., Sadori, L., Balasse, M., and Peyron, O.: Holocene climate dynamics in the central Mediterranean inferred from pollen data, *Climate of the Past*, 21, 2331–2359, <https://doi.org/10.5194/cp-21-2331-2025>, 2025.

Orbigny, A.: Foraminifères, in de la Sagra, Bertrand, 1–224 pp., 1839.

Ozenda, P.: Végétation du continent européen, Delachaux et Niestlé., 272 pp., 1994.

Paquet, H., Coudé-Gaussen, G., and Rognon, P.: Etude minéralogique de poussière sahariennes le long d'un transect entre 19 et 35° de latitude Nord, *Revue de Géographie Physique et de Géologie Dynamique*, 25, 257–265, 1984.

978 Peel, M. C., Finlayson, B. L., and McMahon, T. A.: Updated world map of the Köppen-Geiger climate
 979 classification, *Hydrol. Earth Syst. Sci.*, 11, 1633–1644, 2007.

980 Peretto, C.: The first peopling of southern Europe: the Italian case, *Comptes Rendus Palevol*, 5, 283–290,
 981 <https://doi.org/10.1016/j.crpv.2005.11.006>, 2006.

982 Pérez-Folgado, M., Sierro, F. J., Flores, J. A., Cacho, I., Grimalt, J. O., Zahn, R., and Shackleton, N.: Western
 983 Mediterranean planktonic foraminifera events and millennial climatic variability during the last 70 kyr, *Marine*
 984 *Micropaleontology*, 48, 49–70, [https://doi.org/10.1016/S0377-8398\(02\)00160-3](https://doi.org/10.1016/S0377-8398(02)00160-3), 2003.

985 Petschick, R.: MacDiff Ver.4.2.5, Manual, 2000.

986 Petschick, R., Kuhn, G., and Gingele, F.: Clay mineral distribution in surface sediments of the South Atlantic:
 987 sources, transport, and relation to oceanography, *Marine Geology*, 130, 203–229, [https://doi.org/10.1016/0025-](https://doi.org/10.1016/0025-3227(95)00148-4)
 988 [3227\(95\)00148-4](https://doi.org/10.1016/0025-3227(95)00148-4), 1996.

989 Peyron, O., Guiot, J., Cheddadi, R., Tarasov, P., Reille, M., De Beaulieu, J.-L., Bottema, S., and Andrieu, V.:
 990 Climatic Reconstruction in Europe for 18,000 YR B.P. from Pollen Data, *Quat. res.*, 49, 183–196,
 991 <https://doi.org/10.1006/qres.1997.1961>, 1998.

992 Peyron, O., Goring, S., Dormoy, I., Kotthoff, U., Pross, J., De Beaulieu, J.-L., Drescher-Schneider, R., Vannière,
 993 B., and Magny, M.: Holocene seasonality changes in the central Mediterranean region reconstructed from the
 994 pollen sequences of Lake Accesa (Italy) and Tenaghi Philippon (Greece), *The Holocene*, 21, 131–146,
 995 <https://doi.org/10.1177/0959683610384162>, 2011.

996 Peyron, O., Magny, M., Goring, S., Joannin, S., de Beaulieu, J. L., Brugiapaglia, E., Sadori, L., Garfi, G., Kouli,
 997 K., Ioakim, C., and Combourieu-Nebout, N.: Contrasting patterns of climatic changes during the Holocene across
 998 the Italian Peninsula reconstructed from pollen data, *Clim. Past*, 2013.

999 Pierce, J. W. and Stanley, D. J.: Suspended-sediment concentration and mineralogy in the central and western
 1000 Mediterranean and mineralogic comparison with bottom sediment, *Marine Geology*, 19, M15–M25,
 1001 [https://doi.org/10.1016/0025-3227\(75\)90053-5](https://doi.org/10.1016/0025-3227(75)90053-5), 1975.

1002 Pisias, N. G. and Moore, T. C.: The evolution of Pleistocene climate: A time series approach, *Earth and Planetary*
 1003 *Science Letters*, 52, 450–458, [https://doi.org/10.1016/0012-821X\(81\)90197-7](https://doi.org/10.1016/0012-821X(81)90197-7), 1981.

1004 Prasad, A. M., Iverson, L. R., and Liaw, A.: Newer Classification and Regression Tree Techniques: Bagging and
 1005 Random Forests for Ecological Prediction, *Ecosystems*, 9, 181–199, <https://doi.org/10.1007/s10021-005-0054-1>,
 1006 2006.

1007 Pross, J., Koutsodendris, A., Christanis, K., Fischer, T., Fletcher, W. J., Hardiman, M., Kalaitzidis, S., Knipping,
 1008 M., Kotthoff, U., Milner, A. M., Müller, U. C., Schmiedl, G., Siavalas, G., Tzedakis, P. C., and Wulf, S.: The 1.35-
 1009 Ma-long terrestrial climate archive of Tenaghi Philippon, northeastern Greece: Evolution, exploration, and
 1010 perspectives for future research, *nos*, 48, 253–276, <https://doi.org/10.1127/nos/2015/0063>, 2015.

1011 Quézel, P. and Médail, F.: *Écologie et biogéographie des forêts du bassins méditerranéen*, Elsevier., Paris, 571
 1012 pp., 2003.

1013 Robles, M., Peyron, O., Ménot, G., Brugiapaglia, E., Wulf, S., Appelt, O., Blache, M., Vannière, B., Dugerdil, L.,
 1014 Paura, B., Ansanay-Alex, S., Cromartie, A., Charlet, L., Guédron, S., De Beaulieu, J.-L., and Joannin, S.: Climate
 1015 changes during the Late Glacial in southern Europe: new insights based on pollen and brGDGTs of Lake Matese
 1016 in Italy, *Clim. Past*, 19, 493–515, <https://doi.org/10.5194/cp-19-493-2023>, 2023.

Robles, M., Andrieu, V., Rochette, P., Fauquette, S., Peyron, O., Demory, F., Parlak, O., Charrat, E., Gambin, B., and Alçiçek, M. C.: Vegetation and climate changes at the Early-Late Pliocene Transition around the Mediterranean basin: A case from the Burdur Basin in Southwestern Anatolia, *Climate of the Past*, 1–33, <https://doi.org/10.5194/egusphere-2025-174>, 2025.

Robles, M., Andrieu, V., Fauquette, S., Peyron, O., Joannin, S., Azuara, J., Rochette, P., Demory, F., Helvacı, C., Gambin, B., Pross, J., Koutsodendris, A., and Alçiçek, M. C.: Vegetation and climate changes during the Early–Middle Pleistocene transition: evidence from Acıgöl Basin (Southwestern Anatolia, Türkiye), *Global and Planetary Change*, 262, <https://doi.org/https://doi.org/10.1016/j.gloplacha.2026.105409>, 2026.

Saarinen, J., Oksanen, O., Žliobaitė, I., Fortelius, M., DeMiguel, D., Azanza, B., Bocherens, H., Luzón, C., Solano-García, J., Yravedra, J., Courtenay, L. A., Blain, H.-A., Sánchez-Bandera, C., Serrano-Ramos, A., Rodríguez-Alba, J. J., Viranta, S., Barsky, D., Tallavaara, M., Oms, O., Agustí, J., Ochando, J., Carrión, J. S., and Jiménez-Arenas, J. M.: Pliocene to Middle Pleistocene climate history in the Guadix-Baza Basin, and the environmental conditions of early *Homo* dispersal in Europe, *Quaternary Science Reviews*, 268, 107132, <https://doi.org/10.1016/j.quascirev.2021.107132>, 2021.

Salonen, J. S., Korpela, M., Williams, J. W., and Luoto, M.: Machine-learning based reconstructions of primary and secondary climate variables from North American and European fossil pollen data, *Sci Rep*, 9, 15805, <https://doi.org/10.1038/s41598-019-52293-4>, 2019.

Sánchez Goñi, M. F., Llave, E., Oliveira, D., Naughton, F., Desprat, S., Ducassou, E., Hodell, D. A., and Hernández-Molina, F. J.: Climate changes in south western Iberia and Mediterranean Outflow variations during two contrasting cycles of the last 1Myrs: MIS 31–MIS 30 and MIS 12–MIS 11, *Global and Planetary Change*, 136, 18–29, <https://doi.org/10.1016/j.gloplacha.2015.11.006>, 2016.

Sánchez-Bandera, C., Fagoaga, A., Serrano-Ramos, A., Solano-García, J., Barsky, D., DeMiguel, D., Ochando, J., Saarinen, J., Piñero, P., Lozano-Fernández, I., Courtenay, L. A., Tilton, S., Luzón, C., Bocherens, H., Yravedra, J., Fortelius, M., Agustí, J., Carrión, J. S., Oms, O., Blain, H.-A., and Jiménez-Arenas, J. M.: Glacial/interglacial climate variability in southern Spain during the late Early Pleistocene and climate backdrop for early *Homo* in Europe, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 625, 16, <https://doi.org/10.1016/j.palaeo.2023.111688>, 2023.

Sánchez-Garrido, J. C. and Nadal, I.: The Alboran Sea circulation and its biological response: A review, *Front. Mar. Sci.*, 9, 933390, <https://doi.org/10.3389/fmars.2022.933390>, 2022.

Sánchez-Laulhé, J. M., Jansa, A., and Jiménez, C.: Alboran Sea Area Climate and Weather, in: *Alboran Sea - Ecosystems and Marine Resources*, edited by: Báez, J. C., Vázquez, J.-T., Camiñas, J. A., and Malouli Idrissi, M., 939, 2021.

Sassoon, D., Combourieu-Nebout, N., Peyron, O., Bertini, A., Toti, F., Lebreton, V., and Moncel, M.-H.: Pollen-based climatic reconstructions for the interglacial analogues of MIS 1 (MIS 19, 11, and 5) in the southwestern Mediterranean: insights from ODP Site 976, *Climate of the Past*, 21, 489–515, <https://doi.org/10.5194/cp-21-489-2025>, 2025.

Schiebel, R. and Hemleben, C.: *Planktic Foraminifers in the Modern Ocean*, Springer Berlin Heidelberg, Berlin, Heidelberg, <https://doi.org/10.1007/978-3-662-50297-6>, 2017.

Siccha, M. and Kucera, M.: ForCenS, a curated database of planktonic foraminifera census counts in marine surface sediment samples, *Sci Data*, 4, 170109, <https://doi.org/10.1038/sdata.2017.109>, 2017.

Skonieczny, C., Bory, A., Bout-Roumazeilles, V., Abouchami, W., Galer, S. J. G., Crosta, X., Stuut, J.-B., Meyer, I., Chiapello, I., Podvin, T., Chatenet, B., Diallo, A., and Ndiaye, T.: The 7–13 March 2006 major Saharan outbreak: Multiproxy characterization of mineral dust deposited on the West African margin, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2011JD016173>, 2011.

Stanley, D. J., Warne, A. G., Davis, H. R., Bernasconi, S. M., and Chen, Z.: Nile delta, *National Geographic Research Exploration*, 8, 22–51, 1992.

Stumpf, R., Frank, M., Schönfeld, J., and Haley, B. A.: Climatically driven changes in sediment supply on the SW Iberian shelf since the Last Glacial Maximum, *Earth and Planetary Science Letters*, 312, 80–90, <https://doi.org/10.1016/j.epsl.2011.10.002>, 2011.

Tarasov, P. E., Cheddadi, R., Guiot, J., Bottema, S., Peyron, O., Belmonte, J., Ruiz-Sanchez, V., Saadi, F., and Brewer, S.: A method to determine warm and cool steppe biomes from pollen data; application to the Mediterranean and Kazakhstan regions, *J. Quaternary Sci.*, 13, 335–344, [https://doi.org/10.1002/\(SICI\)1099-1417\(199807/08\)13:4%3C335::AID-JQS375%3E3.0.CO;2-A](https://doi.org/10.1002/(SICI)1099-1417(199807/08)13:4%3C335::AID-JQS375%3E3.0.CO;2-A), 1998.

Ter Braak, C. J. F., Juggins, S., Birks, H. J. B., and Van der Voet, H.: Weighted Averaging Partial Least Squares regression (WA-PLS): definition and comparison with other methods for species-environment calibration, in: *Multivariate Environmental Statistics*, edited by: Patil, G. P. and Rao, C. R., Amsterdam, 525–560, 1993.

Theron, R., Paillard, D., Cortijo, E., Flores, J. A., Vaquero, M., Sierro, F. J., and Waelbroeck, C.: Data-mining the past environment, in: *IGARSS 2003. 2003 IEEE International Geoscience and Remote Sensing Symposium. Proceedings (IEEE Cat. No.03CH37477)*, IGARSS 2003. 2003 IEEE International Geoscience and Remote Sensing Symposium., 3688–3690, <https://doi.org/10.1109/IGARSS.2003.1295238>, 2003.

Thiry, M.: Palaeoclimatic interpretation of clay minerals in marine deposits: an outlook from the continental origin, *Earth-Science Reviews*, 49, 201–221, [https://doi.org/10.1016/S0012-8252\(99\)00054-9](https://doi.org/10.1016/S0012-8252(99)00054-9), 2000.

Tintore, J., Violette, P. E. L., Blade, I., and Cruzado, A.: A Study of an Intense Density Front in the Eastern Alboran Sea: The Almeria–Oran Front, *Journal of Physical Oceanography*, 18, 1384–1397, [https://doi.org/10.1175/1520-0485\(1988\)018%3C1384:ASOAIID%3E2.0.CO;2](https://doi.org/10.1175/1520-0485(1988)018%3C1384:ASOAIID%3E2.0.CO;2), 1988.

Tomadin, L.: Sedimentary fluxes and different dispersion mechanisms of the clay sediments in the Adriatic Basin, *Rend. Fis. Acc. Lincei*, 11, 161–174, <https://doi.org/10.1007/BF02904649>, 2000.

Toti, F., Bertini, A., Girone, A., Marino, M., Maiorano, P., Bassinot, F., Combourieu-Nebout, N., Nomade, S., and Buccianti, A.: Marine and terrestrial climate variability in the western Mediterranean Sea during marine isotope stages 20 and 19, *Quaternary Science Reviews*, 243, 106486, <https://doi.org/10.1016/j.quascirev.2020.106486>, 2020.

Vázquez, J. T., Alonso, B., Fernández-Puga, M. C., Gómez-Ballesteros, M., Iglesias, J., Palomino, D., Roque, C., Ercilla, G., and Díaz-del-Río, V.: Seamounts along the Iberian continental margins, *Boletín Geológico y Minero*, 126, 483–514, 2015.

Venkatarathnam, K. and Ryan, W. B. F.: Dispersal patterns of clay minerals in the sediments of the eastern Mediterranean Sea, *Marine Geology*, 11, 261–282, [https://doi.org/10.1016/0025-3227\(71\)90028-4](https://doi.org/10.1016/0025-3227(71)90028-4), 1971.

Voelker, A. H. L., Salgueiro, E., Rodrigues, T., Jimenez-Espejo, F. J., Bahr, A., Alberto, A., Loureiro, I., Padilha, M., Rebotim, A., and Röhl, U.: Mediterranean Outflow and surface water variability off southern Portugal during the early Pleistocene: A snapshot at Marine Isotope Stages 29 to 34 (1020–1135ka), *Global and Planetary Change*, 133, 223–237, <https://doi.org/10.1016/j.gloplacha.2015.08.015>, 2015.

1097 Weldeab, S., Emeis, K.-C., Hemleben, C., Vennemann, T. W., and Schulz, H.: Sr and Nd isotope composition of
 1098 Late Pleistocene sapropels and nonsapropelic sediments from the Eastern Mediterranean Sea: Implications for
 1099 detrital influx and climatic conditions in the source areas, *Geochimica et Cosmochimica Acta*, 66, 3585–3598,
 1100 2002.

1101 Zhao, Y., Liu, Z., Colin, C., Paterne, M., Siani, G., Cheng, X., Duchamp-Alphonse, S., and Xie, X.: Variations of
 1102 the Nile suspended discharges during the last 1.75Myr, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 311,
 1103 230–241, <https://doi.org/10.1016/j.palaeo.2011.09.001>, 2011.

1104 Zhao, Y., Colin, C., Liu, Z., Paterne, M., Siani, G., and Xie, X.: Reconstructing precipitation changes in
 1105 northeastern Africa during the Quaternary by clay mineralogical and geochemical investigations of Nile deep-sea
 1106 fan sediments, *Quaternary Science Reviews*, 57, 58–70, <https://doi.org/10.1016/j.quascirev.2012.10.009>, 2012.

1107 Zhao, Y., Colin, C., Liu, Z., Bonneau, L., and Siani, G.: Climate forcing of terrigenous sediment input to the central
 1108 Mediterranean Sea since the early Pleistocene, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 442, 23–35,
 1109 <https://doi.org/10.1016/j.palaeo.2015.11.006>, 2016.

1110

