

Supplemental Material

Energetic near-inertial waves induced by winter storms and mesoscale eddies in the subtropical Northwestern Pacific Ocean

5 Hongkai Wang^{1,2}, Zifei Chen¹, Xinyuan Diao^{1,2,3}, Fei Yu^{1,2,3}, Xingchuan Liu¹, Qiang Ren¹, Feng Nan^{1,2,3}

¹ Key Laboratory of Ocean Observation and Forecasting & Laboratory of Ocean Circulation and Waves, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, 266071, China


² Laboratory for Ocean Dynamics and Climate, Qingdao Marine Science and Technology Center, Qingdao, 266071, China

³ College of Marine Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

10 *Correspondence to:* Zifei Chen (chenzifei@qdio.ac.cn), Fei Yu (yuf@qdio.ac.cn)

Text S1

Data comparisons between the CPLDA reanalysis product and the subsurface mooring observations are shown in Fig. S1. The results suggest that the reanalysis data and observations have nearly the same patterns in mean flows and potential density. Therefore, the CPLDA data can be used to analysis the variation of stratification background velocities in this study.



15

Figure S1. Comparisons of the velocity (a and b) and density (c and d), between the reanalysis product and subsurface mooring observations from 1 November, 2017 to 28 February, 2018.