

Severe convective weather in Italy: Current understanding and Research Priorities in the TIM campaign

Mario Marcello Miglietta^{1,2}, Elenio Avolio¹, Alessandro Bracci¹, Massimiliano Burlando³, Francesco Cairo¹, Federico Canepa³, Vincenzo Capozzi⁴, Sebastiano Carpentari⁵, Federico Cassola⁶, Alessandro Ceppi^{7,8}, Silvio Davolio^{1,9}, Francesco De Martin¹⁰, Giorgio Doglioni⁵, Costanza Di Felice Fabrizi^{11,12}, Luca Di Liberto¹, Francesco Domenichini¹³, Stefano Federico¹, Massimiliano Enrico Ferrario¹³, Federico Grazzini¹⁴, Antonio Iengo⁵, Sante Laviola¹, Agostino Manzato¹⁵, Paolo Paganini¹⁶, Antonio Parodi¹⁷, Alessandro Pavan¹⁵, Andrea Piazza¹⁸, Arturo Pucillo¹⁵, Giovanni Ravezzani⁷, Francesco Sioni¹⁵, Barbara Turato⁶, Gianfranco Vulpiani¹⁹, Marco Zanatta¹, Dino Zardi^{5,11}

¹CNR-ISAC, Italy

²Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria

³Department of Civil, Chemical and Environmental Engineering, University of Genoa, Genoa, Italy

⁴Department of Science and Technology, University of Naples “Parthenope”, Centro Direzionale di Napoli, Isola C4, CAP 80143

⁵Department of Civil Environmental and Mechanical Engineering (DICAM), University of Trento, Trento, Italy

⁶Regional Agency for Environmental Protection of Liguria (ARPAL), Genoa, Italy

⁷Department of Civil and Environmental Engineering (D.I.C.A.), Politecnico di Milano, Milan, Italy

⁸Pegaso Telematic University, Faculty of Engineering and Computer Science, Department of Engineering, Naples, Italy

⁹Dipartimento di Scienze della Terra, Università degli studi di Milano, Milan, Italy

¹⁰School of Meteorology, University of Oklahoma, Norman, Oklahoma

¹¹Center Agriculture Food Environment (C3A), University of Trento, Trento, Italy

¹²National Doctoral School in Polar Sciences, University of Venice “Ca’ Foscari”, Venice, Italy

¹³Regional Agency for Environmental Protection of Veneto (ARPAV), Teolo, Italy

¹⁴Regional Agency for Prevention, Environment and Energy of Emilia-Romagna (ARPAE-SIMC), Bologna, Italy

¹⁵ARPA FVG–OSMER, Palmanova, Italy

¹⁶Section of Geophysics, National Institute of Oceanography and Applied Geophysics, Sgonico, Italy

¹⁷CIMA Research Foundation, Savona, Italy

¹⁸Provincia Autonoma di Trento, Servizio Prevenzione e Rischi e CUE, Ufficio Previsioni e Pianificazione, Trento, Italy

¹⁹Department of civil protection, Rome, Italy

Correspondence to: Mario Marcello Miglietta (m.miglietta@isac.cnr.it)

Abstract. The status and priorities of research on severe convective storms in Italy are outlined here, with particular attention to the upcoming Thunderstorm Intensification from Mountains to Plains (TIM) observation campaign. The increased intensity of events induced by climate change is attracting growing attention from the Italian scientific community on this topic. While northern Italy is the most studied, with numerous papers analyzing intense events such as tornadoes, hailstorms, and flash floods, there are still many open questions. The central and southern regions are even less explored, even though they are also occasionally subject to intense events modulated by mesoscale circulations, sea-land interactions, and complex orography.

The TIM campaign represents a unique opportunity to improve the understanding, monitoring, and forecasting of severe storms. The campaign, led by the European Severe Storm Laboratory, represents a first-of-this-kind pan-European campaign aimed at obtaining coordinated data on severe convective storms, and is a key step toward improving warnings, forecasts, climate change impact estimation, and adaptation measures. The value of the campaign for Italy is indicated by the participation of several institutions, both from the academic and the operational community. Among the planned initiatives, the use of two Italian airborne platforms will allow for a more complete characterization of the environments associated with convective storms, including an improvement in our understanding of the role of aerosols and storm-scale modeling.

1. Introduction

Convective storms result from a complex interplay between thermodynamics and dynamical processes at different scales, ranging from the synoptic scale to the microscale. Therefore, their forecast is still challenging due to the complexity of the physical processes, which are still only approximately accounted for in numerical weather prediction models, and the scarcity of high-resolution observations within the pre-storm environment, which limits the representativeness of the initial conditions in model simulations. Improving our understanding of these processes requires a coordinated strategy that combines in-situ microphysics observations, thermodynamic profiling, remote sensing of clouds, precipitation, winds and aerosols, turbulence measurements, together with modelling activities.

Severe convective storms are responsible for injuries and even fatalities (e.g., De Martin et al., 2025b), as well as for a substantial amount of damage across Europe. Their growing impact (Battaglioli et al., 2026) highlights the importance of research and a better understanding of their effects to adopt appropriate loss mitigation measures.

Unfortunately, focused field campaigns in Europe were relatively small (Kunz et al., 2022). Conversely, several observational campaigns on severe convection have been performed for several decades in the USA. Considering the complex morphology of the Italian territory, where land-sea transitions and complex orography may especially affect the meso-gamma scale circulation, the conceptual models developed for the USA should be at least modified or adapted to the unique Mediterranean environment. This was shown, for example, in Miglietta et al. (2016) or, more recently, in De Martin et al. (2024), in which the triple point configuration typical of tornadogenesis in US Midwest was observed in the Po Valley, but with very different time and spatial scales. In particular, the role of the increasingly warmer Mediterranean Sea and the presence of a very long and complex coastline should be properly included in this framework, together with the complex orography of the region, as demonstrated in recent studies of extreme convective weather (e.g. De Martin et al., 2025a).

In the Mediterranean region, only a limited number of extensive observational campaigns was performed, but none was focused on small-scale convective storms. In northern Italy (and partly central Italy), severe weather was investigated during the Mesoscale Alpine Program (Bougeault et al., 2001) and the HyMeX program (Ducrocq et al., 2012), but both projects were mainly focused on the large scale and mesoscale conditions conducive to severe weather, heavy rain in particular (Miglietta and Davolio, 2022).

In recent years, one major observational effort was the observational campaign on Monte Baldo (Italian Alps), managed by the University of Trento under the DECIPHER project. This activity was part of the summer Extended Observation Period of TEAMx (Serafin et al., 2020; Rotach et al., 2022), during which nine research teams from various European countries provided a remarkable number of different instruments and personnel to the observational activities from June to October 2025. The Adige valley target area and its experimental setup are shown in Figure 1. Although the project focused on planetary boundary layer (PBL) processes in a complex Alpine environment, some attention was posed to air-surface exchange processes and ventilation mechanisms towards the free atmosphere, possibly conducive to the initiation of atmospheric convection. Through in situ observations-including flux towers, Doppler and Raman Lidar, ceilometer, aerosol sensors and disdrometers-, the TEAMx-DECIPHER campaign was a great opportunity to consolidate a high level of logistical and organizational expertise in the management of complex observation campaigns with many partners involved. Various turbulence regimes, slope and valley circulations were observed, allowing to investigate their role in transport and mixing, and indirectly their potential influence on convective initiation through the distribution of humidity, aerosols and atmospheric tracers, leading to a rich dataset in the Alpine area.

Together with convective initiation, another important issue is the predictability of severe thunderstorms. In the Po valley, it is known that predictability is generally high in the presence of a strong synoptic forcing (e.g., a cold front advancing from the northwest), but it is much lower during the convective season, when severe storms may locally occur without any substantial large-scale driver. To make the picture more complex, storms over the Alps and Prealps follow a strong diurnal cycle, associated with diurnal heating, while those near the coasts or offshore are mostly associated with the annual cycle of sea surface temperature (SST; Manzato et al., 2022a). Consequently, the distribution of thunderstorms is not uniform in the area, showing a strong maximum in the northeastern Prealps (Fig. 2).

Any approach based uniquely on synoptic analysis (e.g., a simple “weather-type” classification) is insufficient for severe weather forecasting in the Po Valley. Even any approach based on simple mesoscale “environmental conditions” would not be very helpful, because potential instability is present on most summer days in the Po Valley (Manzato, 2003; Manzato, 2023); hence, it is not obvious that the instability (e.g., estimated by CAPE) has a direct proportionality to the storm probability or even to the storm intensity (Manzato et al., 2025).

Finally, severe convective weather events, particularly supercell thunderstorms and hailstorms, constitute a significant hazard and have accounted for the costliest natural disasters in recent years (Bowen et al., 2023). These phenomena pose substantial risks to populations, infrastructure, and sectors such as agriculture and transport. These issues illustrate how timely is the planning of the field campaign on Thunderstorm Intensification from Mountains to plains (TIM; see Fischer et al., 2025 for further details on the motivation of the campaign and its research topics). The campaign, led by the European Severe Storm Laboratory (ESSL), represents a first-of-its-kind pan-European campaign on severe convective storms aimed at obtaining coordinated and dense data on orographically driven storms, and is a key step toward improving warnings, forecasts, climate change impact estimation, and adaptation measures. The value of the campaign for Italy is indicated by the participation of several institutions, both from the academic and the operational community.

This paper represents a summary of the contribution that the Italian community could and is willing to provide to the forthcoming TIM campaign, planned in the 2028-2030 years. The northern regions are the most studied in Italy, as they are affected by the most intense convective phenomena, so this article focuses on these regions (and partly also on the central regions). Section 2 presents the state of the art of severe convective weather research in Italy. Issues worth studying during the campaign are reported in Section 3. Section 4 describes which tools are available or would be necessary for better understanding the dynamics and microphysics of severe convective storms. Conclusions are drawn in Section 5.




Figure 1: Geographical location of Monte Baldo (left; Imagery © 2025 NASA, Map data © 2025 Google); sketch of the experimental setup on the Monte Baldo slope during the TEAMx campaign (right).




Figure 2: Spatial distribution of Cloud-to-Ground lightning (per km² yr) based on EUCLID data from 2005 to 2019 (adapted from Manzato et al., 2022a). The location of northern Italian regions is also shown.

2. State of the Art

2.1 Northern Italy

The Liguria region, characterized by complex orography and proximity to the sea, every year experiences intense meteorological events, such as flash floods, severe thunderstorms, hail, downbursts and strong winds. These events are generally localized and typically develop in correspondence with the convergence between the southerly flow over the Ligurian Sea with the northerly cold outbreak from the Po valley across the gaps in the orography (Buzzi et al., 2014; Fiori et al., 2017). They can cause significant socio-economic impacts, occasionally even casualties, while their predictability is still quite limited. The available observations from the Regional Agency for the Environmental Protection of Liguria region (ARPAL), including a dense network of weather stations, enriched by open hardware ACRONETWORK stations (Loglisci et al. 2024), operational radars, and continuous satellite monitoring, combined with numerical simulation analysis, have highlighted the key role of orography, sea-breeze interactions and SST patterns in enhancing convective activity (e.g., Cassola et al., 2016, Meroni et al. 2018), but the understanding of small-scale processes remains limited. In particular situations to be further investigated, storms originating from mesoscale convergence over the Ligurian Sea or northern Tyrrhenian Sea can move further inland in the Tuscan hinterland and, more rarely even downwind the main orographic barrier, even affecting the Emilia-Romagna region, as happened on 13-14 September 2015 during the devastating flash floods in Val Trebbia and Val di Nure (Segadelli et al., 2020).

In the Veneto and Friuli Venezia Giulia (FVG) regions, a series of particularly intense phenomena, such as supercells and intense convective systems, have occurred in the last decades. Remarkable examples are shown in Fig. 3: the F4 (category 4 of the Fujita scale) Mira-Dolo tornado of July 2015 (Zanini et al., 2017); the violent downburst in Verona in August 2020; the recurrent summer hailstorms, which peaked in July 2023, when the European record for hailstone size was broken (De Martin et al., 2025a). Other severe weather events were directly or indirectly related to cyclonic activity, such

184 as the famed Vaia storm of 2018 (Davolio et al., 2020; Sioni et al., 2023), the impressive “acqua alta” in Venice on
 185 November 12, 2019 (the second highest level in the historical record; Ferrarin et al., 2021; Miglietta et al., 2023), the
 186 intermediate-duration floods (12-24 hours long) over the pre-Alpine area near Vicenza that tested the efficiency of the
 187 recently implemented operational detention basins, and the back-building storms responsible for flash floods in eastern
 188 FVG in November 2025. A statistical study by the Regional Agency for the Environmental Protection of Veneto region
 189 (ARPAV) has identified a marked increase in high precipitation threshold exceedances during the last 30 years for what
 190 concerns the short and very short accumulation timescales (5 min - 12 hours), while no significant trend was found in
 191 extreme precipitations in FVG (Manzato et al., 2025). The increased interest in intense events has brought ARPAV to
 192 maintain a register of intense rainfall events and tornadoes in the region, as part of its weather forecasting practice.
 193 ARPAV continuously monitors these episodes with its ground-based high-resolution observational network, including
 194 over 300 hydro-meteorological stations, 2 C-band radars, 8 disdrometers, a 10-km temperature and humidity profiler, and
 195 a historical archive of images from installed webcams.
 196 Overall, the Alpine region appears to exhibit increased hydrogeological sensitivity, emerging during several heavy
 197 downpours during the summer season in the form of recurrent debris flows. In fact, the increased frequency and intensity
 198 of extreme weather events (e.g., Dallan et al., 2022) resulted in a rise in hydrogeological risk and in the vulnerability of
 199 ecosystems and infrastructure. These phenomena, often characterized by strong spatial and temporal variability and very
 200 difficult to predict, are particularly impactful in urban areas (Gambini et al., 2025). For example, the Milan hydraulic
 201 node represents an emblematic case, where the Seveso, Olona, and Lambro rivers cause recurrent inundations in the
 202 metropolitan area, leading to widespread damage to infrastructure and urban mobility (Ceppi et al., 2022). Furthermore,
 203 wide urban areas (such as Milan) may alter the dynamics of local severe storms, by intensifying updrafts and shifting
 204 storm tracks (De Martin et al., 2025c), thus adding an additional level of complexity.




206 **Figure 3: Photos of recent severe weather events in northern Italian regions: supercell with downburst in Verona – hundreds**
 207 **of trees fallen in urban areas (August 23, 2020; top left); Mira-Dolo F4 tornado (July 8, 2015; top right; Photo: Alberto Gobbi);**
 208

209 hailstorm (hailstone diameter: 16 cm) in Carmignano del Brenta (July 19, 2023; bottom left; Photo: “Associazione Meteo in
 210 Veneto”); flash flood in Meda due to severe rainfall in Brianza province (September 22, 2025; bottom right).

211 2.2 Central and Southern Apennines

212 While northern Italy has been the subject of several studies and some conceptual models are now available to identify
 213 conditions conducive to severe weather, the central and southern Apennine range remains largely under-researched,
 214 although they constitute a steep, coastal-proximal mountain range occasionally subject to heavy precipitation, strongly
 215 modulated in terms of timing and persistence by mesoscale circulations, land-sea contrasts and complex topography. In
 216 the Apennine setting, modest but steep relief modulates convective initiation by: (i) channeling low-level flow through
 217 gaps and valleys, (ii) forcing ascent on the windward flanks, and (iii) focusing near-shore convergence where the marine
 218 boundary layer impinges on the coastal range. The efficacy of these pathways depends on the synoptic flow, boundary-
 219 layer stability, shear, and sea-land thermal contrasts, which together set diurnal phasing and storm organization. The
 220 available literature clearly documents how the interaction between coastal orography and marine inflows over Italy
 221 prolongs event duration and shifts precipitation peaks relative to the upstream forcing (see Miglietta and Davolio, 2022,
 222 for a review), leading to intense inland flooding (Rosso and Ceppi, 2022).

223 Across the Apennines, orographic precipitation encompasses both cold-season events, driven by moisture transported
 224 from the sea (stratiform or embedded convection in the presence of long-fetch advection) and warm-season events,
 225 characterized by convection initiated by the orography, including hail-producing storms. Capozzi et al. (2023a) identified
 226 a limited set of recurrent heavy-precipitation spatial patterns for Campania region (southern Italy) that are systematically
 227 associated with synoptic types and the regional orography; importantly, such patterns occur in autumn-winter under long-
 228 fetch maritime moisture transport impinging on the mountain ranges, highlighting the key role of moisture-fluxes in
 229 organizing and sustaining heavy rainfall events (Fig. 4).



230
 231 **Figure 4: Schematic illustration of two of the six large-scale configurations associated with heavy-precipitation events in the**
 232 **Campania Region (southern Italy) identified by Capozzi et al. (2023a). Panels (a)–(b) show Pattern 1 (warm and moist**
 233 **southwesterly flow and orographic enhancement of precipitation along the main Apennine ranges); panels (c)–(d) represent**
 234 **Pattern 6 (convective activity over the Tyrrhenian Sea and the northern sector under local low-level convergence). In (a) and**
 235 **(c), the black dashed line marks the approximate position of the 552-dam 500-hPa geopotential-height contour, while the orange**

236 box in (a) locates the Campania region, zoomed in b) and in d). Images credit: © Google Earth, Data Sio, NOAA, U.S. Navy,
237 NGA, GEBCO.

238 2.3 Atmospheric Rivers

239 Atmospheric rivers (ARs) have emerged as one of the most relevant global drivers for extreme hydro-meteorological
240 events in many areas of the globe. Defined as narrow corridors of enhanced horizontal transport of low-level moisture,
241 ARs typically form over midlatitude regions pulling up water vapor charge also from subtropics and feeding precipitation
242 systems in the areas where they make landfall and are forced to rise over the orography. Whilst the science of ARs is
243 well developed in the US, where they have been recognized as responsible for extreme precipitation especially along the
244 Pacific coast, only in the last decade the role of ARs has become evident over Europe (Lavers and Villarini, 2015). Over
245 the Mediterranean basin, ARs were found relevant for recent extreme rainfall and major floods over southern Spain
246 (Lorente-Plazas et al., 2019), southern France (Francis et al., 2022), the Dinaric Alps (Martinković et al., 2017), the
247 eastern basin (Bozkurt et al., 2019) and over northern and central Italy (Davolio et al., 2020; Davolio et al., 2023; Sioni
248 et al., 2023).

249 Focusing on the Alpine chain southern slopes, the presence of moisture transport, which may originate from either remote
250 or local sources, is critical for the development of orographic rainfall and convection. Heavy precipitation often develops
251 eastward of an advancing upper-level disturbance, a trough or a cyclone deepening over the basin. The latter is responsible
252 for steering warm and moist low-level (unstable) air from the sea towards the steep slopes of the Alps and the Apennines
253 (Miglietta and Davolio, 2022). At the mesoscale, precipitation systems are likely triggered by the direct orographic uplift
254 or by the deflection or blocking action exerted by the orography (Miglietta and Buzzi, 2004), which can also renew
255 convection triggering at the same location.

256 The role of ARs in heavy precipitation has been recently disclosed in a climatological framework over northern Italy
257 (Davolio et al., 2026), as well as in case study analyses (Davolio et al., 2020, Sioni et al., 2023), providing evidence of a
258 strict connection between extreme organized water vapor transport and extreme events affecting both the Alps and the
259 Apennines (Grazzini et al., 2020). Conversely, it is well known that the thermodynamic characteristics of the impinging
260 moist flow are fundamental to determine the interaction with the orography (Miglietta and Rotunno, 2014, among others),
261 thus the intensity and distribution of rainfall. In this context, ARs have been so far described as characterized by moist-
262 neutral stratification in the lowest layers (~ 3 km) of the atmosphere, as observed from dropsondes released during many
263 field campaigns over the Pacific (Ralph et al. 2020). This weak stability makes AR prone to rise over mountain barriers,
264 producing orographic precipitation where convection is likely embedded, which can greatly amplify the amounts of
265 precipitation.

266 2.4 Tornadoes and downbursts

267 Italy is among the European countries with the highest tornado occurrence and intensity. Most Italian tornadoes fall within
268 the categories of damage F0 and F2 of the Fujita scale, but intense cases (F3, F4 or even F5) have also been reported, as
269 documented by the European Severe Weather Database (ESWD; Dotzek et al., 2009). Some climatological studies
270 (Miglietta and Matsangouras, 2018; Avolio and Miglietta, 2022) have shown that the Italian areas most prone to tornado
271 development include some southern areas (Ionian regions and Sicily), the Po Valley (in particular, its eastern part), and
272 the Tyrrhenian coasts (Fig. 5). These events occur predominantly in autumn (in the southern regions) and summer (in the
273 northern regions).

274 Some recent studies have improved our understanding of tornadic environments in Italy. The conditions typically
 275 identified include the presence of strong wind shear (especially in the low-level) and significant Convective Available
 276 Potential Energy (CAPE) (Ingrosso et al., 2020). In southern Italy, SST anomalies also play a key role (Miglietta et al.,
 277 2017a; Bagaglini et al., 2021; Avolio and Miglietta, 2021) as well as the convective initiation by the orography (Miglietta
 278 et al., 2017b). More recently, a conceptual model based on the proximity of three air masses with different characteristics
 279 has been proposed for the Po valley (De Martin et al., 2024), showing some similarities with the US Midwest tornadic
 280 environments, although at a much smaller spatial scale. Due to its small-scale, this feature was not identified earlier in
 281 reanalysis studies. A detailed analysis of a recent IF3 (category 3 of the International Fujita scale) tornado in the same
 282 area reaffirmed the applicability of this conceptual model (De Martin et al., 2025d).

283 Significant field campaigns in Italy, focused on observations of high-resolution surface wind fields during thunderstorms,
 284 have been conducted using a wind monitoring network, which integrates ultrasonic anemometers, three LiDAR profilers,
 285 and one scanning Lidar, installed in the commercial ports of the Ligurian Sea (Solari et al., 2012; Repetto et al., 2018).
 286 This system has captured numerous thunderstorm-related wind events over the past few decades (e.g., Burlando et al.,
 287 2017, 2018, 2020; Canepa et al., 2020; Solari et al., 2015; Zhang et al., 2018). Canepa et al. (2024) include a database of
 288 twenty-nine thunderstorm wind records acquired by ultrasonic anemometers within the monitoring network. Several of
 289 these events were simultaneously captured by multiple instruments located within the same area, enabling spatiotemporal
 290 cross-correlation analyses and providing a depiction of different thunderstorm outflows. This level of spatial and temporal
 291 resolution makes the dataset unique in the national context.

292 In terms of direct measurements of thunderstorm outflows and their potentially destructive impacts on the ground, full-
 293 scale wind observations remain scarce due to the limited spatial and temporal extent of the phenomenon. When available,
 294 high-sampling anemometric records capture only a single, event-specific realization, shaped by both the intrinsic
 295 characteristics of the downburst (e.g., size, intensity, duration, and interaction with background flow) and the relative
 296 position of the measuring instrument with respect to the evolving phenomenon. Consequently, field data seldom provide
 297 a complete description of the spatial flow structure or of its temporal evolution, particularly when the parent storm moves
 298 relative to the sensor. In-situ measurements alone do not permit a quantitative reconstruction of the interplay between the
 299 various flow components that collectively define the downburst system. As a result, much of the research on downburst
 300 dynamics relies on laboratory-scale physical modelling and numerical simulations. Recent large-scale experimental
 301 campaigns have produced valuable datasets, enabling statistical analyses of the spatiotemporal evolution of downburst
 302 flows and their interaction with the near-surface PBL and the translating storm systems. Subsequent experiments
 303 incorporated thermodynamic effects by imposing temperature differentials between the downdraft and ambient air,
 304 highlighting the key role of thermal gradients in shaping the downburst structure and dynamics (Canepa et al., 2025).
 305 Nevertheless, the quantitative influence of thermal processes in natural conditions remains largely unresolved.




Figure 5. Map of selected Italian tornadoes crossing "land" for the period 1990–2024; different colors represent the seasons (data derived from ESWD).

2.5 Role of aerosols

Aerosols are the seeds upon which cloud droplets and ice crystals form. Their ability to act as ice-nucleating particles (INPs) or cloud condensation nuclei (CCN) regulate microphysical processes, e.g. increasing the number of liquid droplets (Twomey, 1977) or initiating cloud glaciation (Murray et al., 2021). Aerosol particles play an important role in the climate by controlling the radiative budget of the atmosphere (aerosol-radiation interaction; ARI) and by directly modifying the microphysical properties of clouds (aerosol-cloud interaction; ACI).

In fact, variations in *aerosol properties* can dramatically affect convective storms as they may: modify droplet activation and warm-rain formation; shift the height and temperature of primary glaciation; influence secondary ice processes such as rime-splintering; determine the number and type of ice particles, graupel, and hail embryos; affect latent heating profiles, updraft strength, and storm intensity; control the rate and efficiency of precipitation development; modulate storm electrification and lightning activity.

Despite limited observational evidence, coarse-mode aerosol particles, particularly mineral dust, have recently been linked to hail formation through their ability to act as efficient ice-nucleating particles. Several studies reported positive associations between hail occurrence and enhanced Saharan dust loadings (Brennan and Wilhelm, 2025), which is extremely important for Italy, given its frequent exposure to Saharan dust outbreaks (Vogel et al., 2025).

2.6 Hail

Currently, there is considerable interest within the research community in hail over Italy for several reasons. While Italy has emerged as a hail hotspot in Europe over the last decade (e.g., Punge et al., 2017) and Northern Italy has shown the largest global increase in very-large hail frequency (up to +0.03 events per decade; Battaglioli et al., 2026), the most recent hail climatologies show remarkable differences across the country. Some studies indicate a frequency peak over the foothills of Northern Italy (e.g., Giordani et al., 2023; Cui, 2025; Manzato et al., 2022b), while others show a maximum over the sea (e.g., Laviola et al., 2022; Kahraman et al., 2024). Furthermore, there has been growing awareness of the

substantial economic damage caused by hail to crops, buildings, and vehicles (Púčík et al., 2019; Panosetti and Tomassetti, 2023). A record of US\$6 billion loss has been experienced in Northern Italy in 2023 (Battaglioli et al., 2026).

Generally, hail is not explicitly forecast by regional and national weather services and is generally treated as one of several possible hazards associated with severe storms. While this approach is partly justified by the uncertainty in hail prediction, other weather services (e.g., NOAA in the United States) issue daily hail outlooks, suggesting that such forecasts are feasible and could be tested during the TIM campaign. Notably, OSMER-ARPA FVG mentioned in their forecast for 24 July 2023 the possible occurrence of severe hailstorms, a few hours before a record-breaking hail event occurred.

2.7 Numerical weather prediction models and data assimilation

Despite the progress made in the model resolution and physical parameterizations, the forecast of convective events is challenging for numerical weather prediction (NWP) models because of the multitude of physical processes and scales involved (Hu and Franzke, 2020) and the intrinsic limited predictability of convection.

Current weather and climate models are progressing towards finer spatiotemporal scales, which is largely due to the increased computational capabilities that have become available. Improved model performance can be also achieved by steering the model using observations, i.e. applying data assimilation (DA). Several operational systems around the world use convection permitting DA (e.g. Ballard et al., 2016), that is regional-scale DA with a horizontal grid spacing of approximately 1-4 km, where convection is explicitly resolved rather than parametrized. These systems provide improved nowcasts and short-term (0-24h) forecasts particularly suitable for convective storms (Milan et al., 2020).

Several studies have been conducted over Italy focusing on the assimilation of different data sources, often at convection resolving resolution, as radar (Gastaldo et al., 2021; Maiello et al., 2017; Lagasio et al., 2019b; Mazzarella et al., 2020), GNSS (Lagasio et al., 2019a; Torcasio et al., 2023; Faccani et al., 2005), lightning (Federico et al., 2024; 2026), weather stations (Maggioni et al., 2023), rainrate (Davolio et al., 2017; Torcasio et al., 2024), among others. These studies used different DA techniques namely 3DVar, 4DVar, nudging, and Ensemble Kalman filter applied to different models used in Italy (e.g., WRF, MOLOCH, COSMO).

3. Open Issues

Observational networks in Italy are mostly managed at regional scale. Although some fields (e.g. precipitation), especially in northern Italy, are densely monitored, several challenges and significant limitations persist, due to sparse and heterogeneous observational coverage and reporting (e.g., for tornadoes and hail) and the complex topography.

For example, flash floods in small catchments are often poorly observed because of their rapid and highly localized nature. Existing observational networks may miss peak showers because measurements are either insufficiently frequent or too sparse to adequately cover the affected areas, and the accuracy of numerical models is not always satisfactory (Ceppi et al., 2023). These limitations, together with the lack of specific measurements needed to better understand convective initiation and development, make the need for a dedicated field campaign urgent. Some key issues are outlined below.

3.1 Convective initiation

The difficulty in predicting localized convective storms requires a better understanding of the local conditions leading to *convective initiation*. Although some recurrent patterns for heavy rain have been recently identified in Italy (e.g., Capozzi et al., 2023b; Grazzini et al., 2020, 2021; Iacomino et al., 2025), and the role of equilibrium and non-equilibrium conditions has been analyzed (Molini et al. 2011), the characteristics of the synoptic and mesoscale conditions associated

with other severe storms should be clarified, identifying the observational/model-based precursors for intense convection in the different regions. This investigation will allow improving the capacity of forecasting severe thunderstorms and combined with observational analysis, should help identify the typical trajectories followed by the storms in their evolution.

Within the framework of orographic convection in the Alpine areas, the experience gained during the TEAMx–DECIPHER campaign has emphasized the need to better understand the PBL mechanisms leading to convective initiation and, in particular, any mechanism that may produce vertical motion sufficient to lift high- θ_e air mass to the level of free convection. In this framework, Manzato et al. (2022a) provided the first climatology of convective initiation over northern Italy based on lightning strikes. The campaign stimulated several scientific questions, both methodological and operational: which dynamic and thermodynamic preconditions govern the initiation and lifecycle of *orographic cumulus clouds*? which atmospheric and terrain factors govern their location and persistence? how do local slope/valley circulations, atmospheric stability, and direction of the synoptic flow interact to determine the stationarity or propagation of convective clouds?

Different initiation mechanisms may favor the development of convective systems, such as convergence induced by local breeze systems, land-coast gradient boundaries, sunrise-sunset induced thermal gradients, moisture gradients due to advection (e.g., from the Adriatic Sea). In addition, cold-pools, bores and gravity waves may trigger new convective events (the so-called “secondary convection”). However, it is difficult to discriminate which initiation mechanism is effective. This issue is related to a key question: why on days with similar instability and “weather-type” conditions, do severe storms sometimes occur and sometimes not? This point should be analyzed for heavy rain events both in the convective season (summer), in the period characterized by “strong flux of rain” (autumn), and in the transition between the two regimes (Manzato, 2007a; Davolio et al., 2016).

In the past, Manzato (2007b) and Manzato et al. (2019) developed a neural network-based scheme to perform a multivariate analysis for forecasting the occurrence and intensity of rainfall events using multiple sounding-derived indices in the Friuli-Venezia-Giulia region. Building on this work, the TIM campaign aims also to better understand whether a relationship between instability indices and severe precipitation that can lead to flash flood episodes may exist. Given the limitations identified in this kind of studies, new aspects should probably be incorporated in research, in particular the role of aerosol concentration and of warm rain in enhancing storm dynamics and rain efficiency. At a later stage of development, another issue is the transition of convective storms from mountains to plains, which seems neither well understood from a storm-dynamics perspective nor generally well predicted by numerical models (Fischer et al., 2025).

The presence of the *Mediterranean Sea* introduces an additional degree of complexity, making the prediction of convective initiation and evolution particularly challenging, especially in the case of stationary or back-building convective systems, as identified in some heavy rain events in the Liguria region (Cassola et al., 2016). A better understanding of the role of SST anomalies has implications not only for a better detection of observed events, but also in terms of reduction of false alarms.

3.2 Role of aerosols

Extreme convective storms often form in environments characterized by strong horizontal and vertical aerosol gradients—for example, dust intrusions, biomass-burning plumes, polluted boundary layers, or marine aerosol inflow. The ability to detect these gradients over large areas is essential for interpreting storm initiation and spatial evolution; in fact, depending on their distribution within the atmosphere, aerosols may either invigorate or suppress convective development (Stier et

al., 2024). Unfortunately, direct observations in convective environments remain scarce. Limited airborne measurements reveal substantial variability in aerosol concentrations and properties throughout the cloud column, suggesting that aerosol–cloud interactions are highly sensitive to local microphysical, dynamical and thermodynamic conditions (e.g. Zanatta et al., 2023).

Moreover, the understanding of *microphysical processes* leading to hail formation and extreme precipitation is still incomplete and leads to a limited consensus on the impacts of aerosol particles on precipitation, making microphysics parametrization in numerical models a critical issue (Stier et al., 2024). Also, the impact of ACI on cloud formation and precipitation is nonlinear (Zhao et al., 2024), leading to contrasting effects, i.e. invigorating or weakening microphysics effects at cloud base that accelerate or delay precipitation. Lastly, aerosol-mediated effects on hail development remain a potentially important but poorly constrained process.

Despite recent progress, wide observational gaps limit our knowledge of aerosol-cloud interaction and their influence on Mediterranean severe storms: do aerosols act as CCN and/or as INP? Do they promote the formation of rain and/or graupel, and if so, through which mechanisms and relationships? Recently, the role of aerosols was identified in modulating the precipitation in Mediterranean areas exposed to Saharian dust advection (e.g., Ferrari et al., 2024); however, the generality of these results is still uncertain and requires further investigation.

Within the Mediterranean, climate-change induced desertification and heatwaves modify the local concentration of aerosols (e.g. Cristofanelli et al., 2013; Vogel et al., 2025), hence the availability of CCN and INPs. Even though numerical modelling studies correlate these aerosol perturbations with changes in convective systems, the inability of reconciling observations and modelling data complicates the attribution of observed changes to aerosol-cloud interactions (Stier et al., 2024).

Along similar lines, Napoli et al. (2022) showed through cloud-permitting simulations over the Great Alpine Region (GAR) that the lifetime of clouds is generally increased at high levels of aerosols, while convective clouds (typical of the summer season) can decrease at high levels of pollution as a result of the reduction in strong updrafts associated with an increased air column stability. All in all, it is crucial to study observations related to the 3D distribution of aerosol in connection with convective initiation (e.g., Brennan and Wilhelm, 2025 have identified for hail in Switzerland an “optimal concentration” of aerosol to maximize hail probability).

A central challenge is to disentangle aerosol-driven microphysical effects from dynamic forcing. Accordingly, open questions are: what are the background aerosol states (composition, CCN/INP abundance, vertical structure) that do not promote storm intensification, and how do they differ from perturbations associated with Mediterranean severe convective weather? Under which dynamical and thermodynamic conditions (CAPE, shear, humidity, synoptic forcing) do Mediterranean aerosol perturbations transition from weakening to invigorating convective storms? How can in-situ observations be used to attribute observed storm variability to aerosol–cloud interactions rather than to meteorological dynamics alone?

3.3 Convective storm changes in future climate

The potential changes in storm frequency and intensity under future climate scenarios is an open question that is still subject of strong debate in the scientific community. While there are strong evidences that potential instability increased in recent years (Cavalleri et al., 2026), and will be enhanced in a warmer climate, the signal in rainfall observations from regional network is not so clear (e.g. Manzato et al., 2025 found no significant trend of precipitation during the convective season), emphasizing the need for the analysis of multi-decadal trends in different data to detect possible shifts in precipitation regimes (Mazzoglio et al., 2025). Conversely, attribution studies based on pseudo-global warming

simulations have clearly identified the role of a warmer atmosphere and sea in enhancing the intensity of severe convective events in the Mediterranean (Gonzalez-Aleman et al., 2023; Calvo-Sanchez et al., 2026). The fact that the observed super-Clausius–Clapeyron scaling of extreme precipitation may be explained by a shift from stratiform to convective rain type (Da Silva and Haerter, 2025; Haslinger et al., 2025) opens new questions on the role of climate change in modifying the precipitation formation pathways (Hawkins et al., 2023). In this framework, and following recent evidence for the Apennine areas that the intensity and frequency of heavy precipitation events are increasing, particularly during autumn (Capozzi et al., 2023b), it is worth investigating how this result applies to different types of precipitation (e.g., autumn-winter orographic rainfall episodes driven by long-fetch maritime moisture transport in the Apennines). Some studies report a marked increase in large hail events in the Po Valley (Battaglioli et al., 2023), others over the Alps (Cui et al., 2025), while in the plain of Friuli Venezia Giulia region, located in north-eastern Italy, no significant hail trends were observed from 29 years of data collected by a network of about 300 hailpad stations, except for hail size (Manzato et al., 2025). These inconsistencies may ultimately be related to the scarcity of systematic ground-based hail observations, a limitation that could only be addressed through an extensive and dense network of automated hailpads (e.g., Kopp et al., 2023) and disdrometers (Jameson et al., 2015).

3.4 Conceptual models for central Italy

While conceptual models for precipitation have been developed in the Alpine areas, for other regions specific conceptual models need to be developed. The open issues include: (i) better investigating the convective initiation along the Adriatic/Tyrrhenian flanks, discriminating between terrain-forced convergence versus synoptically-forced cases; (ii) identifying under which combinations of SST anomalies, stability and shear, orographic channeling most effectively trigger convection and favors hail growth in central Italy; (iii) assessing the transferability of hail proxies calibrated in other Mediterranean regions, such as the Alps and northeastern Italy (Manzato, 2012; Giordani et al., 2024), to the interior Apennine basins with different relief scales and coastal geometry.

3.5 Atmospheric rivers and severe convection

ARs associated with severe organized convective storms, over or at some distance from the orography, have been rarely documented. For example, Moore et al. (2012) analyzed a heavy rainfall event that occurred over the Central US, where an AR fed two quasi-stationary mesoscale convective systems (MCSs), linearly organized and exhibiting back-building mechanisms. Recent events affecting the Mediterranean basin revealed the presence of intense convection associated with ARs, as in the case of the dramatic flood in Valencia in October 2024 or the record-breaking rainfall over Liguria and Piedmont in October 2020 (Davolio et al., 2023). Therefore, given the availability of various AR datasets, both global and Mediterranean-specific, it is timely and worthy to investigate whether, and to what extent, ARs are associated with severe convective storms that develop near the Alpine chain during different seasons.

3.6 Tornadoes and downbursts

Significant limitations persist in monitoring and predicting rapidly evolving storms, such as tornadoes or downbursts, hence raising key challenges. These include: i) sparse and heterogeneous observational coverage, including radar gaps and non-uniform scanning strategies, that limit the monitoring of mesocyclones and small-scale storms; ii) underreporting and inconsistent intensity classification; iii) the intricate geomorphology of Italy and the Mediterranean, which can enhance or hinder small-scale circulations, including sea–land and orographic interactions; iv) the absence of tornado-specific forecasting or warning systems in Italy, as discussed in Miglietta and Rotunno (2016).

While some progress has been made in identifying the tornadic environment (Bagaglini et al., 2021; De Martin et al., 2024), more attention should be paid to radar patterns associated with severe convection. A climatology and a systematic

analysis of the environments conducive to downbursts is also missing, although they represent a significant cause of damage in some Italian regions (Burlando et al., 2025). Future advances in downburst understanding will require integrated observational networks combining high-frequency anemometry, Doppler Lidar and radar systems, and thermodynamic profilers capable of resolving temperature, humidity, and pressure gradients at fine spatial and temporal scales. The coupling of these measurements with numerical weather prediction (NWP) models and Computational Fluid Dynamics (CFD) simulations could enable real-time reconstruction and nowcasting of downburst structures.

3.7 Hail

Understanding of the physical processes leading to hail formation remains limited, particularly for giant hail events (Allen et al., 2020). In Italy, this issue is further complicated by complex topography, as recently demonstrated by De Martin et al. (2025a) for northeastern Italy, and Ricchi et al. (2023a, 2023b) for a giant hail event in Central Italy, where the Apennine topography, together with SST anomalies, guided the low-level flow and focused coastal convergence, effectively preconditioning convective initiation. This uncertainty has important implications. For instance, why does large hail appear to be a rare phenomenon along the Ligurian and Tyrrhenian coasts, despite the very high frequency of severe thunderstorms? Conversely, why is large hail so common along the Alpine foothills? Uncertainty in the mechanisms leading to large hail production also affects hail forecasting. For example, how important are environmental kinematics with respect to thermodynamics?

Beyond direct hail observation and forecasting, substantial effort is currently devoted to estimating hail size using remote sensing, like radar data (Allen et al., 2020; Aregger et al, 2025). An effective hail size detection algorithm, combined with a tracking algorithm, could represent a powerful tool for early warnings of severe hailstorms. Regional weather services in Emilia-Romagna and Piedmont are testing several algorithms for this purpose (Fornasiero et al., 2024), and the data collected during the TIM campaign could significantly support their calibration.

Further work is needed to clarify the spatial distribution of hailstorms over Italy. Extending hail climatologies to include economic hail damage —potentially involving insurance companies— should improve hail risk maps and lay the groundwork for impact-oriented hail warnings.

3.8 Numerical weather prediction models and data assimilation

As stated, there are some studies over Italy addressing the problem of DA at the convective scale. However, very high-resolution simulations (i.e. with a grid spacing of less than 1 km) will be necessary to study local low-level circulations (e.g., up-valley and down-valley flows, mountain-valley breeze, ...) that are relevant for the initiation of convection. Furthermore, the sensitivity of the model to parameterization schemes (e.g., convection, turbulence, microphysics) should clarify the role of convection-relevant and not yet fully studied physical processes, such as the impact of aerosols on cloud microphysics. Therefore, there are many challenges requiring further work and the TIM campaign can help to cope with some of them: i) quantifying the observation impact of different data sources to guide future observing network design for the prediction of convection; b) quantifying the impact of a background correlation matrix, which is aware of the “error of the day”, compared to climatological approaches in order to better designing DA system for the convective scale; c) using different observations at high spatio-temporal resolution to consider explicitly the correlation among observations; d) refining the use of currently available observations, and assimilating new observation types.

4. Required Tools

Observational tools and modelling techniques available nowadays make it possible to investigate severe convective weather features at unprecedented resolution.

4.1 Satellite data

Satellite observations are limited by their relatively coarse space and time resolution; however, the integration of geostationary and polar data remain a privileged tool to explore and monitor storm evolution, from pre-storm conditions and convection initiation to the mature and decaying stages of the systems. In addition, flash observations from the Lightning Imager onboard MTG are very useful to identify, track and forecast convective storms.

New satellite techniques are available to identify and monitor hailstorms. The Multi-sensor Approach for Satellite Hail Advection (MASHA) is a new multi-instrument blended technique conceived for real-time detecting of hail-bearing clouds. MASHA (Fig. 6) identifies hail clouds from satellite measurements and monitors the evolution of hail-bearing systems every 5 min by combining the strength of the MicroWave Cloud Classification-Hail (MWCC-H) method, which detects hail through the whole Global Precipitation Measurement (GPM) sensor constellation (Laviola et al., 2020a,b), with the high temporal rate of the Meteosat Rapid Scan Service (MSG-RSS). MASHA has been applied to various case studies demonstrating strong capability in identifying severe events and high robustness to detect small-scaled hailstorms in mountain catchments. These results open new perspectives for investigating hailstorms and hydro-meteorological events in mountain areas, whereas traditional methodologies often miss events or underestimate their severity. Although further experiments are needed, MASHA's performance paves the way towards pseudo-operational applications in support to hailstorm nowcasting and assimilation in regional numerical weather predictions and appears mature for application during the TIM campaign.

Finally, the data provided by the new satellite constellations (e.g., Meteosat Third Generation, MTG), including vertical profiling, will represent a tremendous opportunity to better characterize the environment where convective storms develop and to track their evolution.




Figure 6: MASHA conceptual scheme (top images; Credit: NASA and EUMETSAT) and highlights of hailstorm evolution affecting Northern Italy on 26 July 2021 (bottom bracketed images).

4.2 Radar

Investigating convective initiation requires high-temporal resolution observations of the vertical velocity and moisture distribution in the lowest part of the troposphere (first 2-3 km). *Doppler radar* wind data with overlapping scan domains (at least every 5 minutes) would be necessary to estimate the *vertical* velocity spatial distribution using the classical

continuity equation method (Dual-Doppler Analysis, DDA) or more modern techniques (e.g. Lopez-Carrillo et al. 2011, Tsai et al. 2025). For northeastern Italy, these techniques can be applied to the available Loncon and Fossalon C-band Doppler radars. In the Emilia-Romagna region, an X-band radar will be soon installed and two C-band radar sites updated. The Italian Civil Protection Department (DPC) expressed its willingness to join the TIM campaign, being particularly interested in analyzing the ground impact of severe convective storms and in integrating the damage surveys with radar data. A mobile X-band radar (2m high antenna) is available for the campaign but requires power supply stations in the target areas and some funds to assure the transport of the instrument. Also relevant for the campaign is a radar situated on mount Zoufplan at 2000 m height, at the border with Austria.

The recent installation of two X-band dual-polarization radars in the Lombardy Region (Desio and Flero) aims to strengthen the regional monitoring network and enhance nowcasting capabilities using machine learning models (Franch et al., 2025). In addition to supporting operational forecasts (Xu et al., 2025), these radar-based estimates will serve as inputs for hydrological models, whether physically based and spatially distributed (hybrid systems) or entirely AI-driven (full AI systems). Consequently, integrating AI-based radar nowcasting with hydrological models will represent a significant advancement in improving the effectiveness of operational warning systems and the dynamic management of structural mitigation measures, providing reliable and accurate short-term forecasts, particularly for summer convective events (Ceppi et al., 2025).

AEROLAB (AERosol mOBile LABoratory) is a CNR-ISAC mobile platform designed for deployment in remote and non-instrumented areas for the study and characterization of atmospheric particulate matter through in situ sampling. AEROLAB is housed in an ISO10 container, certified for operation also on-board research vessels, temperature-controlled, and equipped with dedicated sampling lines for multiple instruments. The remote sensing operational observational capability of the mobile platform consists of a co-located Doppler wind lidar, a 35-GHz cloud radar, and a lidar, enabling an integrated characterization of convection-relevant processes from the surface layer to the cloud column. Together, the suite links aerosol structure, inflow kinematics, and near-cloud vertical motions to the cloud and precipitation response, enabling process studies and model evaluation of triggering, microphysics–dynamics feedbacks, and aerosol-mediated convective intensification.

4.3 Integration of remote sensing and surface data

A possible correspondence between severe storms and ARs can be identified based on the exploitation of satellite analysis and ground-based observations. Being characterized by large-scale moisture transport, ARs can be identified using analysis fields and detection algorithms. However, to describe precisely the low-level thermodynamic profiles of the impinging moist flow and infer the precipitation characteristics, the availability of on-demand *radio-soundings*, *dropsondes* and vertical moisture/wind measurements, including weather *drones*, especially upstream of convection, is hoped for. Additional instruments, like *optical lidar* and powerful *wind profilers*, which have been extensively employed during the TEAMx campaign, should monitor the PBL at high-resolution, and the low-level wind, including its vertical component.

A substantial upgrade of in-situ observations is also desirable, mainly in areas which have been excluded from the mainstream scientific interest, such as the Apennines. Dense, elevation-stratified *raingauges* with ≤ 10 -min logging along coastal-to-ridge transects and windward crests may complement the available disdrometer network (Adirosi et al., 2023) to resolve drop size distributions, phase and fall speeds in mixed-phase and convective events. An extensive *hail monitoring network* is also desirable to document occurrence, size spectra and kinetic energy, ideally co-located with disdrometers for cross-validation.

4.4 Aerosol and cloud in-situ observations

The available ACI in-situ observations are rarely coordinated with large scale dynamic campaigns (Kunz et al., 2022). The Italian contribution to TIM aims to provide systematic observations of both aerosol properties and cloud microphysics at the WMO/GAW observatory of Monte Cimone (CNR-ISAC), a facility representative of the Mediterranean troposphere. However, without vertical profiles of wind, stability, humidity, and aerosol distribution (Fig. 7), it remains impossible to separate microphysical aerosol effects from dynamical forcing. Implementing remote-sensing and profiling capabilities in parallel to in-situ aerosol-microphysical observations during the TIM campaign is therefore desirable to enable robust attribution of aerosol impacts on Mediterranean severe storms.

Airborne platforms are the only way of providing vertically and horizontally resolved observations of aerosol properties, thermodynamic structure, and cloud development in the vicinity of convective systems. To address this need, we propose the integrated deployment of two available Italian research aircrafts: the SkyArrow ERA and the Piper Seneca III, in coordination with aerosol remote sensing (Lufft CHM – Nimbus 15k). Used in combination, the Cimone observatory and the two mobile platforms may characterize the environmental conditions from the surface to the mid-upper troposphere with complementary measurements.

Resolving aerosol concentrations in 3D is crucial considering that convective storms source their inflow differently, depending on their mode (supercell, MCS, etc.) and on the fact they are elevated or surface based. For supercells, different flanks of the storm even have different inflow origins, which has implications for hail growth (e.g., Lin and Kumjian, 2022). Near the orography, inflows might be even more complex and changing over time. Thus, monitoring of aerosols, moisture, and winds would be a unique and impactful observation set, yielding a 3D, multi-scale picture of the storm environment over a wide geographic domain, capturing horizontal gradients in aerosol concentration, shifts in aerosol types, thermodynamic stability, moisture and temperature gradients, all of which are critical for convective initiation and strongly modulate the spatial evolution and intensity of convective storms.

4.5 Campaign coordination challenges

A multi-institutional, multi-platform field campaign entails practical and coordination challenges, which require rigorous operational frameworks, such as instrument interoperability, data-sharing protocols, real-time decision-making during IOPs, and the logistical complexity of deploying airborne platforms alongside ground networks. In fact, different institutions bring instruments with different hardware, data format, and calibration standards, which will require establishing compliance profiles and creating centralized preprocessing scripts to align all outputs, while centralising multiple observations requires the transition to federated, cloud-based data repositories that enforce interoperable formatting like netCDF for sharing. In addition, mission-planning strategies to rapidly evolving conditions will be required to ensure multi-aircraft coordination and to allow simultaneous data collection in the same atmospheric column from fixed-point ground networks and fast-moving airborne platforms.

5. Conclusions

The TIM campaign offers a unique opportunity to improve our understanding of severe weather in Italy. This objective needs, however, a fine data coverage, which includes: high-resolution dual-polarization radars, with short scan times, and improved overlapping coverage for storm detection and tracking; integration of high-frequency satellite and radar data to support effective nowcasting; dense networks of rain gauges and hydrological sensors for real-time monitoring of flash floods; radiosondes and weather drones to better understand the local conditions favorable to deep convective initiation; convection-permitting numerical limited-area models, either deterministic or probabilistic, potentially combined with advanced convective-scale data assimilation methods, providing severe convection indices as outputs to give sufficient guidance in forecasting localized severe convection and in concentrating the observational capabilities in specific areas;

637 post-event damage survey analysis techniques for correlating storm dynamics and ground impacts; development and
638 improvement of nowcasting algorithms in identifying and reproducing the dynamics of propagating, stationary and back-
639 building systems, including the use of data-driven AI algorithms.


640 The national radar reflectivity mosaic, of which we will encourage the full integration in the OPERA European mosaic,
641 may be combined with the data from the available mobile radar and (hopefully) from weather drones (Leuenberger et al.,
642 2020), representing a great opportunity to discover signatures and monitor the detailed evolution of severe weather
643 directly in the areas affected by the most intense events, solving the weaknesses over complex terrain, due for example
644 to radar beam blocking. The recent development of AI methods paves the way to new applications, in terms of both
645 nowcasting and forecasting capabilities, and of pattern-recognition tools trained on radar, satellite, and reanalysis datasets.
646 Finally, the campaign can be crowdsourced, using the platforms where weather amateurs already share the information
647 in real-time (e.g., PRETEMP, De Martin et al., 2023, and Meteonetwork, Giazzi et al., 2022).

648 The proposed dual-aircraft strategy will allow, for the first time in Italy, to deliver one of the most comprehensive
649 environmental characterizations ever obtained for European deep convective storms. This combination will allow to
650 directly link aerosol spatial gradients to microphysical evolution (glaciation, hail, precipitation); to improve storm-scale
651 modelling, especially microphysical parameterizations; to advance the understanding of aerosol–convection coupling, a
652 major climate uncertainty. The dedicated airborne measurements, combined with the data from the observatory of Monte
653 Cimone, offers a unique opportunity to better investigate the troposphere during severe convective events and to improve
654 our understanding of the role of aerosols in enhancing or reducing severe storm activity.

655 Therefore, we believe that the TIM campaign, under the coordination of ESSL and the involvement of many institutions
656 across Europe, may represent a tremendous boost for: (i) moving ahead the understanding of severe weather over Italy;
657 (ii) improving the monitoring and forecasting capabilities; (iii) creating more comprehensive, and generate detailed and
658 (iv) accurate statistics for the intensity and distribution of severe convective events, which is partially missing over Italy
659 (e.g., downbursts); enhancing resilience of the population.

660 Finally, the communication activities planned aside the campaign should be able to increase the awareness about severe
661 weather in the framework of climate change and to explain to the population how to deal with alerts. While the recurring
662 floods over Italy in the last few years have surely improved the situation and made clear the need to adopt more
663 precautionary behavior, unfortunately public education on severe convection storms is still limited, and there is a long
664 way to go to reach an acceptable level.

665



666

667

668

669

670

Figure 7. Schematic representing the aerosol cloud interactions (ACI) under the warming scenario in the Mediterranean. The cloud condensation nuclei (CCN) and ice nucleating particles (INPs) trigger different processes during severe storm development. To isolate the ACI contribution, observations of aerosol and cloud microphysics must be coordinated with vertically resolved dynamics observations.

671

Data availability

672

Not relevant for the present paper.

673

Authors contributions

674

All Authors contributed to the writing of the paper.

675

Conflicts of interest

676

The contact author declares that none of the authors has any competing interests.

677

Acknowledgements

678

679

The authors gratefully acknowledge the European Severe Storms Laboratory (ESSL) for the ESWD extreme weather reports (Fig. 5), Alberto Gobbi and “Associazione Meteo in Veneto” for the photos (Fig. 3).

680

Financial support

681

682

683

MMM and EA acknowledge financial support from the project “REDUCE METEO HAZARDS (RMH)”, BRiC-2024 call. SD received the support of the project "Geosciences for society: resources and their evolution" supported by the Italian Ministry of University and Research (MUR) through the funds "Dipartimenti di Eccellenza 2023/2027".

684

References

685

686

687

688

689

690

- Adirosi, E., Porcù, F., Montopoli, M., Baldini, L., Bracci, A., Capozzi, V., Annella, C., Budillon, G., Bucchignani, E., Zollo, A. L., Cazzuli, O., Camisani, G., Bechini, R., Cremonini, R., Antonini, A., Ortolani, A., Melani, S., Valisa, P., and Scapin, S.: Database of the Italian disdrometer network, Earth Syst. Sci. Data, 15, 2417–2429, <https://doi.org/10.5194/essd-15-2417-2023>, 2023.
- Allen, J. T., Giammanco, I. M., Kumjian, M. R., Jurgen Punge, H., Zhang, Q., Groenemeijer, P., and Ortega, K.: Understanding hail in the earth system, Reviews of Geophysics, 58, 2019 000665, 2020.

691 Areggeret, M. and Germann, H.: Differential reflectivity columns and hail: Linking C-band radar-based estimated column
 692 characteristics to crowdsourced hail observations in Switzerland, *Quarterly Journal of the Royal Meteorological Society*,
 693 151, 5003, 2025.
 694 Avolio, E. and Miglietta, M. M.: Multiple tornadoes in the Italian Ionian regions: Observations, sensitivity tests and
 695 mesoscale analysis of convective storm environmental parameters, *Atmospheric Research*, 263, 105800, 2021.
 696 Avolio, E. and Miglietta, M. M.: Tornadoes in the Tyrrhenian regions of the Italian peninsula: The case study of 28 July
 697 2019, *Atmospheric Research*, 278, 106285, 2022.
 698 Bagaglini, L., Ingrosso, R., and Miglietta, M. M.: Synoptic patterns and mesoscale precursors of Italian tornadoes,
 699 *Atmospheric Research*, 253, 105503, 2021.
 700 Ballard, S. P., Li, Z., Simonin, D., and Caron, J.-F.: Performance of 4D-Var NWP-based nowcasting of precipitation at
 701 the Met Office for summer 2012, *Quarterly Journal of the Royal Meteorological Society*, 142, 472–487,
 702 <https://doi.org/10.1002/qj.2665>, 2016.
 703 Battaglioli, F., Groenemeijer, P., Púčík, T., Taszarek, M., Ulbrich, U., and Rust, H.: Modeled multidecadal trends of
 704 lightning and (very) large hail in Europe and North America (1950–2021), *Journal of Applied Meteorology and*
 705 *Climatology*, 62, 1627–1653, 2023.
 706 Battaglioli, F., Taszarek, M., Groenemeijer, P. *et al.* Contrasting trends in very large hail events and related economic
 707 losses across the globe. *Nat. Geosci.* 19, 52–58, <https://doi.org/10.1038/s41561-025-01868-0>, 2026.
 708 Bougeault, P. and Coauthors: The MAP special observing period, *Bull. Am. Meteor. Soc.*, 82, 433–462, 2001.
 709 Bowen, S., Kerschner, B., and Zheng, J.: Natural Catastrophe and Climate Report (Gallagher Re);
 710 www.ajg.com/gallagher/news-and-insights/2024/january/2023-natural-catastrophe-and-climate-report/, 2024.
 711 Bozkurt, D., Ezber, Y., and Sen, O. L.: Role of the East Asian Trough on the eastern Mediterranean temperature variability
 712 in early spring and the extreme case of 2004 warm spell, *Clim. Dynam.*, 53, 2309–2326, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-019-04847-5)
 713 [019-04847-5](https://doi.org/10.1007/s00382-019-04847-5), 2019.
 714 Brennan, K. P. and Wilhelm, L.: Saharan dust linked to European hail events, *Atmos. Chem. Phys.*, 25, 10823–10836,
 715 <https://doi.org/10.5194/acp-25-10823-2025>, 2025.
 716 Burlando, M., Romanić, D., Solari, G., Hangan, H., and Zhang, S.: Field Data Analysis and Weather Scenario of a
 717 Downburst Event in Livorno, Italy, on 1 October 2012, *Mon. Wea. Rev.*, 145, 3507–3527, [https://doi.org/10.1175/MWR-](https://doi.org/10.1175/MWR-D-17-0018.1)
 718 [D-17-0018.1](https://doi.org/10.1175/MWR-D-17-0018.1), 2017.
 719 Burlando, M., Zhang, S., and Solari, G.: Monitoring, cataloguing, and weather scenarios of thunderstorm outflows in the
 720 northern Mediterranean, *Nat. Hazards Earth Syst. Sci.*, 18, 2309–2330, <https://doi.org/10.5194/nhess-18-2309-2018>, 2018.
 721 Burlando, M., Romanic, D., Boni, G., Lagasio, M., and Parodi, A.: Investigation of the Weather Conditions During the
 722 Collapse of the Morandi Bridge in Genoa on 14 August 2018 Using Field Observations and WRF Model, *Atmosphere*,
 723 11, 724, <https://doi.org/10.3390/atmos11070724>, 2020.
 724 Buzzi, A., Davolio, S., Malguzzi, P., Drofa, O., and Mastrangelo, D.: Heavy rainfall episodes over Liguria in autumn
 725 2011: numerical forecasting experiments, *Nat. Hazards Earth Syst. Sci.*, 14, 1325–1340, [https://doi.org/10.5194/nhess-14-](https://doi.org/10.5194/nhess-14-1325-2014)
 726 [1325-2014](https://doi.org/10.5194/nhess-14-1325-2014), 2014.
 727 Calvo-Sancho, C., Díaz-Fernández, J., González-Alemán, J. J., Halifa-Marín, A., Miglietta, M. M., Azorin-Molina, C.,
 728 Prein, A. F., Montoro-Mendoza, A., Bolgiani, P., Morata, A., and Martín, M. L.: Human-induced climate change
 729 amplification on storm dynamics in Valencia’s 2024 catastrophic flash flood, *Nature Communications*, 17, 1492,
 730 <https://doi.org/10.1038/s41467-026-68929-9>, 2026.

Canepa, F., Burlando, M., Solari, G., F., R., M.P., and Burlando, M.: Vertical profile characteristics of thunderstorm outflows, *Journal of Wind Engineering and Industrial Aerodynamics*, 206, 104332, <https://doi.org/10.1016/j.jweia.2020.104332>, 2020.

Canepa, F., Repetto, M. P., and Burlando, M.: Full-scale measurements of thunderstorm outflows in the Northern Mediterranean. *Geoscience Data Journal* gdj3.247, <https://doi.org/10.1002/gdj3.247>, 2024.

Canepa, F., Guibert, A., Xhelaj, A., Žužul, J., Romanic, D., Ricci, A., Hangan, H., Bouchet, J.-P., Delpech, P., Flamand, O., and Burlando, M.: CLIMATHUNDERR: Experimental database of buoyancy-driven downbursts, *Earth Syst. Sci. Data Discuss.* [preprint], <https://doi.org/10.5194/essd-2025-11>, 2025.

Capozzi, V., Annella, C., and Budillon, G.: Classification of daily heavy precipitation patterns and associated synoptic types in the Campania Region (southern Italy), *Atmos. Res*, 289, <https://doi.org/10.1016/j.atmosres.2023.106781>, 2023a.

Capozzi, V., Rocco, A., Annella, C., Cretella, V., Fusco, G., and Budillon, G.: Signals of change in the Campania region rainfall regime: An analysis of extreme precipitation indices (2002–2021), *Meteorological Applications*, 30, 2168, <https://doi.org/10.1002/met.2168>, 2023b.

Cassola, F., Ferrari, F., Mazzino, A., and Miglietta, M. M.: The role of the sea on the flash floods events over Liguria (northwestern Italy), *Geophys. Res. Lett.*, 43, 3534–3542, <https://doi.org/10.1002/2016GL068265>, 2016.

Cavalleri, F., Lussana, C., Viterbo, F., Brunetti, M., Bonanno, R., Manara, V., Lacavalla, M., and Maugeri, M.: Hourly Precipitation Patterns and Extremization over Italy using convection-permitting reanalysis data, *Nat. Hazards Earth Syst. Sci.*, 26, 279–297, <https://doi.org/10.5194/nhess-26-279-2026>, 2026.

Ceppi, A., Gambini, E., Lombardi, G., Ravazzani, G., and Mancini, M.: SOL40: Forty Years of Simulations under Climate and Land Use Change, *Water*, 14, <https://doi.org/10.3390/w14060837>, 2022.

Ceppi, A., Chaves González, N. A., Davolio, S., and Ravazzani, G.: Can meteorological model forecasts initialize hydrological simulations rather than observed data in ungauged basins?, *Meteorological Applications*, 30, 2165, <https://doi.org/10.1002/met.2165>, 2023.

Ceppi, A., Gambini, E., Ravazzani, G., Lombardi, G., Cerri, L., Meucci, S., and Mancini, M.: Attribution of Flood Forecasting Errors From a Multi-Model Perspective in Milan Urbanized River Basins, *J Flood Risk Management*, 18, 70023, <https://doi.org/10.1111/jfr3.70023>, 2025.

Cristofanelli, P., Brattich, E., Decesari, S., Landi, T. C., Maione, M., Putero, D., Tositti, L., and Bonasoni, P.: Aerosol Chemical Composition at the Mt. Cimone WMO/GAW Global Station, in: *High-Mountain Atmospheric Research : The Italian Mt. Cimone WMO/GAW Global Station (2165 m a.s.l, edited by: Cristofanelli, P., Brattich, E., Decesari, S., Landi, T. C., Maione, M., Putero, D., Tositti, L., and Bonasoni, P., Springer International Publishing, Cham, 99–118, pp.,* https://doi.org/10.1007/978-3-319-61127-3_5, 2018.

Cui, R.: A European hail and lightning climatology from an 11-year kilometer-scale regional climate simulation, *Journal of Geophysical Research: Atmospheres*, 130, 2024 042828, 2025.

Dallan, E., Borga, M., Zaramella, M., and Marra, F.: Enhanced summer convection explains observed trends in extreme subdaily precipitation in the eastern Italian Alps, *Geophysical Research Letters*, 49, e202GL096727, <https://doi.org/10.1029/2021GL096727>, 2022.

Da Silva, N. A. and Haerter, J. O.: Super-Clausius–Clapeyron scaling of extreme precipitation explained by shift from stratiform to convective rain type, *Nat. Geosci.*, 18, 382–388, 2025.

Davolio, S., Volonté, A., Manzato, A., Pucillo, A., Cicogna, A., and Ferrario, M. E.: Mechanisms producing different precipitation patterns over north-eastern Italy: insights from HyMeX-SOP1 and previous events, *Quarterly Journal of the Royal Meteorological Society*, 142, 188–205, 2016.

772 Davolio, S., Silvestro, F., and Gastaldo, T.: Impact of rainfall assimilation on high-resolution hydrometeorological
773 forecasts over Liguria, Italy, *J. Hydrometeorol*, 18, 2659–2680, 2017.

774 Davolio, S., Della Fera, S., Laviola, S., Miglietta, M. M., and Levizzani, V.: Heavy precipitation over Italy from the
775 Mediterranean storm “Vaia” in October 2018: Assessing the role of an atmospheric river, *Mon. Wea. Rev.*, 148, 3571-
776 3588, 2020.

777 Davolio, S., Vercellino, M., Miglietta, M. M., Drago Pitura, L., Laviola, S., and Levizzani, V.: The influence of an
778 atmospheric river on a heavy precipitation event over the Alps, *Wea. Clim. Extremes*, 39, 100542,
779 <https://doi.org/10.1016/j.wace.2022.100542>, 2023.

780 Davolio, S., Sala, I., Comunian, A., Mastrangelo, D., Miglietta, M. M., Drago Pitura, L., and Grazzini, F.: Detection of
781 atmospheric rivers affecting the western Mediterranean and producing extreme rainfall over northern-central Italy.
782 *Nat. Hazards Earth Syst. Sci.*, 26, 2291-2304, <https://doi.org/10.5194/nhess-26-2291-2026>, 2026.

783 De Martin F., Carlon N., Pavan F., Carpentari S., Giazzi M., Peressutti G., Miglietta M. M., Davolio S.: Toward a
784 dedicated warning system of severe storms in Italy: the PRETEMP project, *METMED2023*, 22-24 May 2023, Genoa
785 (Italy), 2023.

786 De Martin, F., Davolio, S., Miglietta, M. M., and Levizzani, V.: A Conceptual Model for the Development of Tornadoes
787 in the Complex Orography of the Po Valley, *Monthly Weather Review*, 152, 1357–1377, 2024.

788 De Martin, F., Manzato, A., Carlon, N., Setvák, M., and Miglietta, M. M.: Dynamic and statistical analysis of giant hail
789 environments in northeast Italy, *Quarterly Journal of the Royal Meteorological Society*, 151, 4945,
790 <https://doi.org/10.1002/qj.4945>, 2025a.

791 De Martin, F., Zonato, A., and Di Sabatino, S.: Effects of the urban land use on a severe convective windstorm, *Quarterly*
792 *Journal of the Royal Meteorological Society*, 151, 5058, <https://doi.org/10.1002/qj.5058>, 2025b.

793 De Martin, F., Miglietta, M. M., Gastaldo, T., Martinazzo, M., Pavan, F., Siena, M., and Sabatino, S.: The Bayesian
794 sinking in Porticello: a predictable convective windstorm?, *Weather*, 80, 365–373, <https://doi.org/10.1002/wea.7715>,
795 2025c.

796 De Martin, F., Pavan, F., Carlon, N., Cioni, G., Rozoff, C., Poli, V., Carpentari, S., and Miglietta, M. M.: A Significant
797 Tornado Event near a Dryline Bulge in Northern Italy, *Wea. Forecasting*, 40, 2293-2315, [https://doi.org/10.1175/WAF-](https://doi.org/10.1175/WAF-D-25-0071.1)
798 [D-25-0071.1](https://doi.org/10.1175/WAF-D-25-0071.1), 2025d.

799 Dotzek, N., Groenemeijer, P., Feuerstein, B., and Holzer, A. M.: Overview of ESSL’s severe convective storms research
800 using the European Severe Weather Database (ESWD, *Atmospheric Research*, 93, 575–586, 2009.

801 Ducrocq, V., and Coauthors: HyMeX SOP1, the field campaign dedicated to heavy precipitation and flash flooding in the
802 northwestern Mediterranean, *B. Am. Meteorol. Soc.*, 95, 1083-1100, <https://doi.org/10.1175/BAMS-D-12-00244.1>, 2014.

803 Faccani, C., Ferretti, R., Pacione, R., Paolucci, T., Vespe, F., and Cucurull, L.: Impact of a high density GPS network on
804 the operational forecast, *Adv. Geosci*, 2, 73-79, <https://doi.org/10.5194/adgeo-2-73-2005>, 2005.

805 Federico, S., Torcasio, R. C., Popova, J., Sokol, Z., Pop, L., Lagasio, M., Lynn, B. H., Puca, S., and Dietrich, S.: Improving
806 the lightning forecast with the WRF model and lightning data assimilation: Results of a two-seasons numerical experiment
807 over Italy, *Atmospheric Research*, 304, 107 382, <https://doi.org/10.1016/j.atmosres.2024.107382>, 2024.

808 Federico, S., Torcasio, R. C., Transerici, C., Realini, E., Song, X., and Venuti, G.: Forecasting convective precipitation
809 over northern Italy: A comparison of lightning and GNSS-ZTD data assimilation, *Atmospheric Research*, 331, 108687,
810 <https://doi.org/10.1016/j.atmosres.2025.108687>, 2026.

811 Ferrari, F., Rizza, U., Morichetti, M., Cassola, F., Miglietta, M. M., and Mazzino, A.: The role of atmospheric aerosols
 812 on severe convective precipitation in a Mediterranean coastal region, *Atmospheric Research*, 305, 107421,
 813 <https://doi.org/10.1016/j.atmosres.2024.107421>, 2024.
 814 Ferrarin, C., Bajo, M., Benetazzo, A., Cavaleri, L., Chiggiato, J., Davison, S., Davolio, S., Lionello, P., Orlić, M., and
 815 Umgiesser, G.: Local and large-scale controls of the exceptional Venice floods of November 2019, *Progress in*
 816 *Oceanography*, 197, 102628, <https://doi.org/10.1016/j.pocean.2021.102628>, 2021.
 817 Fiori, E., Ferraris, L., Molini, L., Siccardi, F., Kranzlmüller, D., and Parodi, A.: Triggering and evolution of a deep
 818 convective system in the Mediterranean Sea: modelling and observations at a very fine scale, *Q. J. Roy. Meteor. Soc.*, 143,
 819 927–941, <https://doi.org/10.1002/qj.2977>, 2017.
 820 Fischer, J., Groenemeijer, P., Holzer, A., Feldmann, M., Schröer, K., Battaglioli, F., Schielicke, L., Púčik, T., Antonescu,
 821 B., Gatzen, C., and Partners, T. I. M.: Invited perspectives: Thunderstorm intensification from mountains to plains, *Nat.*
 822 *Hazards Earth Syst. Sci.*, 25, 2629–2656, <https://doi.org/10.5194/nhess-25-2629-2025>, 2025.
 823 Fornasiero, A., Campana, V., Cremonini, R., Alberoni, P. P., and Vulpiani, G.: Comparative Analysis of two algorithms
 824 for estimating large hail occurrence using radar data, ERAD24, Rome (Italy), 2024.
 825 Franch, G., Tomasi, E., Wanjari, R., Poli, V., Cardinali, C., Alberoni, P. P., and Cristoforetti, M.: GPTCast: a weather
 826 language model for precipitation nowcasting, *Geosci. Model Dev.*, 18, 5351–5371, [https://doi.org/10.5194/gmd-18-5351-](https://doi.org/10.5194/gmd-18-5351-2025)
 827 2025, n.d.
 828 Francis, D., Fonseca, R., Nelli, N., Bozkurt, D., Picard, G., and Guan, B.: Atmospheric river drive exceptional Saharan
 829 dust transport towards Europe, *Atmos. Res.*, 266, 105959, <https://doi.org/10.1016/j.atmosres.2021.105959>, 2025.
 830 Gambini, E. and Coauthors: Uncertainty Quantification and Spatial Biases Assessment in Precipitation Forecasts: A
 831 Methodology for Real-Time Flood Forecasting Applications, *J. Hydrometeor.*, 26, 1423–1436,
 832 <https://doi.org/10.1175/JHM-D-24-0140.1>, 2025.
 833 Gastaldo, T., Poli, V., Marsigli, C., Cesari, D., Alberoni, P. P., and Paccagnella, T.: Assimilation of radar reflectivity
 834 volumes in a pre-operational framework, *Q J R Meteorol Soc.*, 147, 1031–1054, <https://doi.org/10.1002/qj.3957>, 2021.
 835 Giazzi, M., Peressutti, G., Cerri, L., Fumi, M., Riva, I. F., Chini, A., Ferrari, G., Cioni, G., Franch, G., Tartari, G., Galbiati,
 836 F., Condemi, V., and Ceppi, A.: Meteonetwork, An Open Crowdsourced Weather Data System. *Atmosphere*, 13, 928,
 837 <https://doi.org/10.3390/atmos13060928>, 2022.
 838 Giordani, A., Kunz, M., Bedka, K. M., Punge, H. J., Paccagnella, T., Pavan, V., Cerenzia, I. M. L., and Di Sabatino, S.:
 839 Characterizing hail-prone environments using convection-permitting reanalysis and overshooting top detections over
 840 south-central Europe, *Nat. Hazards Earth Syst. Sci.*, 24, 2331–2357, <https://doi.org/10.5194/nhess-24-2331-2024>, 2024.
 841 González-Alemán, J. J., and Coauthors: Anthropogenic warming had a crucial role in triggering the historic and
 842 destructive Mediterranean Derecho in summer 2022, *Bull. Am. Meteorol. Soc.*, 104, 1526–1532, 2023.
 843 Grazzini, F., Craig, G. C., Keil, C., Antolini, G., and Pavan, V.: Extreme precipitation events over northern Italy. Part I:
 844 A systematic classification with machine-learning techniques, *Q. J. Roy. Meteorol. Soc.*, 146, 69–85, 2020.
 845 Grazzini, F., Fragkoulidis, G., Teubler, F., Wirth, V., and Craig, G. C.: Extreme precipitation events over northern Italy.
 846 Part II: Dynamical precursors, *Q. J. Roy. Meteorol. Soc.*, 147, 1237–1257, <https://doi.org/10.1002/qj.3969>, 2021.
 847 Haslinger, K., Breinl, K., and Pavlin, L.: Increasing hourly heavy rainfall in Austria reflected in flood changes, *Nature*,
 848 639, 667–672, <https://doi.org/10.1038/s41586-025-08647-2>, 2025.
 849 Hawkins, E., Compo, G. P., and Sardeshmukh, P.: D.: ESD Ideas: Translating historical extreme weather events into a
 850 warmer world, *Earth Syst. Dynam.*, 14, 1081–1084, <https://doi.org/10.5194/esd-14-1081-2023>, 2023.

851 Hu, G. and Franzke, C. L. E.: Evaluation of daily precipitation extremes in reanalysis and gridded observation-based data
 852 sets over Germany, *Geophysical Research Letters*, 47, 2020 089624, <https://doi.org/10.1029/2020GL089624>, 2020.
 853 Iacomino, C., Pascale, S., and Zappa, G.: A Classification of High-Risk Atmospheric Circulation Patterns for Italian
 854 Precipitation Extremes, *International Journal of Climatology*, 45, 70118, <https://doi.org/10.1002/joc.70118>, 2025.
 855 Ingrosso, R., Lionello, P., Miglietta, M. M., and Salvadori, G.: A statistical investigation of mesoscale precursors of
 856 significant tornadoes: The Italian case study, *Atmosphere*, 11, 301, 2020.
 857 Jameson, A. R., Larsen, M. L., and Kostinski, A. B.: Disdrometer Network Observations of Finescale Spatial–Temporal
 858 Clustering in Rain, *J. Atmos. Sci.*, 72, 1648–1666, <https://doi.org/10.1175/JAS-D-14-0136.1>, 2015.
 859 Kahraman, A., Kendon, E. J., and Fowler, H. J.: Climatology of severe hail potential in Europe based on a convection-
 860 permitting simulation, *Climate Dynamics*, 62, 6625–6642, 2024.
 861 Kopp, J., Manzato, A., Hering, A., Germann, U., and Martius, O.: How observations from automatic hail sensors in
 862 Switzerland shed light on local hailfall duration and compare with hailpad measurements, *Atmos. Meas. Tech.*, 16, 3487-
 863 3503, <https://doi.org/10.5194/amt-16-3487-2023>, 2023.
 864 Kunz, M.: An interdisciplinary field campaign for investigating convective storms and their event chains, *Front. Earth*
 865 *Sci.*, 10, <https://doi.org/10.3389/feart.2022.999593>, 2021.
 866 Lagasio, M., Parodi, A., Pulvirenti, L., Meroni, A. N., Boni, G., Pierdicca, N., Marzano, F. S., Luini, L., Venuti, G.,
 867 Realini, E., Gatti, A., Tagliaferro, G., Barindelli, S., Monti Guarnieri, A., Goga, K., Terzo, O., Rucci, A., Passera, E.,
 868 Kranzlmüller, D., and Rommen, B.: A synergistic use of a high-resolution numerical weather prediction model and high-
 869 resolution earth observation products to improve precipitation forecast, *Remote Sensing*, 11, 2387, 2019a.
 870 Lagasio, M., Silvestro, F., Campo, L., and Parodi, A.: Predictive Capability of a High-Resolution Hydrometeorological
 871 Forecasting Framework Coupling WRF Cycling 3DVAR and Continuum, *J. Hydrometeorol.*, 20, 1307–1337,
 872 <https://doi.org/10.1175/JHM-D-18-0219.1>, 2019b.
 873 Lavers, D. A. and Villarini, G.: The contribution of atmospheric rivers to precipitation in Europe and the United States,
 874 *Journal of Hydrology*, 522, 382–390, 2015.
 875 Laviola S., V. Levizzani, R. R. Ferraro, and J. Beauchamp: Hailstorm Detection by Satellite Microwave Radiometers,
 876 *Remote Sens.*, 12, 621, <https://doi.org/10.3390/rs12040621>, 2020a.
 877 Laviola S., G. Monte, V. Levizzani, R. R. Ferraro, and J. Beauchamp: A new method for hail detection from the GPM
 878 constellation. A prospective for a global hailstorm climatology, *Remote Sens.*, 12, 3553,
 879 <https://doi.org/10.3390/rs12213553>, 2020b.
 880 Laviola, S., Monte, G., Cattani, E., and Levizzani, V.: Hail climatology in the Mediterranean basin using the GPM
 881 constellation (1999–2021), *Remote Sensing*, 14, 4320, 2022.
 882 Leuenberger, D., Haefele, A., Omanovic, N., Fengler, M., Martucci, G., Calpini, B., Fuhrer, O., and Rossa, A.: Improving
 883 High-Impact Numerical Weather Prediction with Lidar and Drone Observations, *Bull. Amer. Meteor. Soc.*, 101, 1036-
 884 1051, <https://doi.org/10.1175/BAMS-D-19-0119.1>, 2020.
 885 Lin, Y. and Kumjian, M. R.: Influences of CAPE on Hail Production in Simulated Supercell Storms, *J. Atmos. Sci.*, 79,
 886 179–204, <https://doi.org/10.1175/JAS-D-21-0054.1>, 2022.
 887 Loglisci, N., Boni, G., Cauteruccio, A., Faccini, F., Milelli, M., Paliaga, G., and Parodi, A.: The role of citizen science in
 888 assessing the spatiotemporal pattern of rainfall events in urban areas: a case study in the city of Genoa, Italy, *Natural*
 889 *Hazards and Earth System Sciences*, 24, 2495–2510, 2024.
 890 López Carrillo, C. and Raymond, D. J.: Retrieval of three-dimensional wind fields from Doppler radar data using an
 891 efficient two-step approach, *Atmos. Meas. Tech.*, 4, 2717–2733, <https://doi.org/10.5194/amt-4-2717-2011>, 2011.

892 Lorente-Plazas, R., Montavez, J. P., Ramos, A. M., Jerez, S., Trigo, R. M., and Jimenez-Guerrero, P.: Unusual
 893 atmospheric-river-like structures coming from Africa induced extreme precipitation over the western Mediterranean Sea,
 894 *J. Geophys. Res. (Atmosphere)*, 125, e2019JD031280, <https://doi.org/10.1029/2019JD031280>, 2019.
 895 Maggioni, E. C., Manzoni, T., and Perotto, A.: WRF data assimilation of weather stations and lightning data for a
 896 convective event in northern Italy, *Bull. of Atmos. Sci. & Technol.*, 4, 8, <https://doi.org/10.1007/s42865-023-00061-8>,
 897 2023.
 898 Maiello, I., Gentile, S., Ferretti, R., Baldini, L., Roberto, N., Picciotti, E., Alberoni, P. P., and Marzano, F.: S.: Impact of
 899 multiple radar reflectivity data assimilation on the numerical simulation of a flash flood event during the HyMeX
 900 campaign, *Hydrol. Earth Syst. Sci.*, 21, 5459-5476, <https://doi.org/10.5194/hess-21-5459-2017>, 2017.
 901 Manzato, A.: A climatology of instability indices derived from Friuli Venezia Giulia soundings, using three different
 902 methods, *Atmos. Res.*, 67–68, 417–454, 2003.
 903 Manzato, A.: The 6 Hours Climatology of Thunderstorms and Rainfalls in the Friuli Venezia Giulia Plain, *Atmos. Res.*
 904 *ECSS 2004 special issue*, 83, 336–348, 2007a.
 905 Manzato, A.: Sounding-Derived Indices for Neural Network Based Short-Term Thunderstorm and Rainfall Forecasts,
 906 *Atmos. Res.*, 83, 349–365, <https://doi.org/10.1016/j.atmosres.2005.10.021>, 2007b.
 907 Manzato, A.: Hail in northeast Italy: Climatology and bivariate analysis with the sounding-derived indices, *J. Appl.*
 908 *Meteorol. Clim.*, 51, 449-467, 2012.
 909 Manzato, A., Pucillo, A., and Cicogna, A.: Improving ECMWF-based 6-hours maximum rain using instability indices
 910 and neural networks, *Atmospheric Research*, 217, 184-197, <https://doi.org/10.1016/j.atmosres.2018.10.020>, 2019.
 911 Manzato, A., Serafin, S., Miglietta, M. M., Kirshbaum, D., and Schulz, D. W.: A pan-Alpine climatology of lightning and
 912 convective initiation, *Monthly Weather Review*, 150, 2213-2230, <https://doi.org/10.1175/MWR-D-21-0149.1>, 2022a.
 913 Manzato, A., Cicogna, A., Centore, M., Battistutta, P., and Trevisan, M.: Hailstone Characteristics in Northeast Italy from
 914 29 Years of Hailpad Data, *J. Appl. Meteor. Climatol.*, 61, 1779-1795, <https://doi.org/10.1175/JAMC-D-21-0251.1>, 2022b.
 915 Manzato, A.: Heavy rain, hailstorms, and lightning in Alpine areas, <https://doi.org/10.13140/RG.2.2.33321.98403>, 2023.
 916 Manzato, A., Fasano, G., Cicogna, A., Sioni, F., and Pucillo, A.: Relationships between Environmental Parameters and
 917 Storm Observations in Po Valley: Are They Climate Change Invariant?, *J. Appl. Meteor. Climatol.*, 64, 267-298,
 918 <https://doi.org/10.1175/JAMC-D-24-0034.1>, 2025.
 919 Martinković, M., Strelec Mahović, N., Mikuš Jurković, P., Renko, T., Pelajić, I., and Smiljanić, I.: Relationship between
 920 atmospheric rivers and extreme precipitation events. In: Karacostas, T., Bais, A., and Nastos, P. (Eds.): Relationship
 921 between atmospheric rivers and extreme precipitation events, in: *Perspectives on Atmospheric Sciences*, Springer
 922 *Atmospheric Sciences*. Springer, Cham, https://doi.org/10.1007/978-3-319-35095-0_54, 2017.
 923 Mazzearella, V., Maiello, I., Ferretti, R., Capozzi, V., Picciotti, E., Alberoni, P. P., Marzano, F. S., and Budillon, G.:
 924 Reflectivity and velocity radar data assimilation for two flash flood events in central Italy: A comparison between 3D and
 925 4D variational methods, *Q. J. Roy. Meteorol. Soc.*, 146, 348-366, <https://doi.org/10.1002/qj.3679>, 2020.
 926 Mazzoglio, P., Viglione, A., Ganora, D., and Claps, P.: Mapping the Uneven Temporal Changes in Ordinary and
 927 Extraordinary Rainfall Extremes in Italy, *Journal of Hydrology: Regional Studies*, 58, 102287,
 928 <https://doi.org/10.1016/j.ejrh.2025.102287>, 2025.
 929 Meroni, A. N., Parodi, A., and Pasquero, C.: Role of SST patterns on surface wind modulation of a heavy midlatitude
 930 precipitation event, *Journal of Geophysical Research: Atmospheres*, 123, 9081–9096, 2018.
 931 Miglietta, M. M. and Buzzi, A.: A numerical study of moist stratified flow regimes over isolated topography, *Q. J. Roy.*
 932 *Meteor. Soc.*, 130, 1749–1770, <https://doi.org/10.1256/qj.02.225>, 2004.

933 Miglietta, M. M. and Rotunno, R.: Numerical simulations of sheared conditionally unstable flows over a mountain ridge,
934 J. Atmos. Sci, 71, 1747–1762, <https://doi.org/10.1175/JAS-D-13-0297.1>, 2014.

935 Miglietta, M. M., Manzato, A., and Rotunno, R.: Characteristics and predictability of a supercell during HyMeX SOP1,
936 Quart. J. Roy. Meteor. Soc, 142, 2839–2853, <https://doi.org/10.1002/qj.2872>, 2016.

937 Miglietta, M. M., Mazon, J., Motola, V., and Pasini, A.: Effect of a positive sea surface temperature anomaly on a
938 Mediterranean tornadic supercell, Scientific Reports, 7, 1–8, 2017a.

939 Miglietta, M. M., Mazon, J., and Rotunno, R.: Numerical simulations of a Tornadic supercell over the Mediterranean,
940 Weather Forecast, 32, 1209–1226, <https://doi.org/10.1175/WAF-D-16-0223.1>, 2017b.

941 Miglietta, M. M. and Matsangouras, I. T.: An updated climatology of tornadoes and waterspouts in Italy, International
942 Journal of Climatology, 1–17, 2018.

943 Miglietta, M. M. and Davolio, S.: Dynamical forcings in heavy precipitation events over Italy: lessons from the HyMeX
944 SOP1 campaign, Hydrol. Earth Syst. Sci, 26, 627–646, <https://doi.org/10.5194/hess-26-627-2022>, 2022.

945 Miglietta, M. M., Buscemi, F., Dafis, S., Papa, A., Tiesi, A., and Conte, D.: A high-impact meso-beta vortex in the
946 Adriatic Sea, Quarterly Journal of the Royal Meteorological Society, 149, 637–656, <https://doi.org/10.1002/qj.4432>,
947 2023.

948 Milan, M., Macpherson, B., Tubbs, R., Dow, G., Inverarity, G., and Mittermaier, M.: Hourly 4D-Var in the Met Office
949 UKV operational forecast model, Quarterly Journal of the Royal Meteorological Society, 146, 1281–1301, 2020.

950 Molini, L., Parodi, A., Rebora, N., and Craig, G. C.: Classifying severe rainfall events over Italy by hydrometeorological
951 and dynamical criteria, Quarterly Journal of the Royal Meteorological Society, 137, 148–154, 2011.

952 Moore, B. J., Neiman, P. J., Ralph, F. M., and Barthold, F. E.: Physical Processes Associated with Heavy Flooding
953 Rainfall in Nashville, Tennessee, and Vicinity during 1–2 May 2010: The Role of an Atmospheric River and Mesoscale
954 Convective Systems, Mon. Wea. Rev, 140, 358–378, <https://doi.org/10.1175/MWR-D-11-00126.1>, 2012.

955 Murray, B. J., Carslaw, K. S., and Field, P.: R.: Opinion: Cloud-phase climate feedback and the importance of ice-
956 nucleating particles, Atmospheric Chemistry and Physics, 21, 665–679, <https://doi.org/10.5194/acp-21-665-2021>, 2021.

957 Napoli, A., Desbiolles, F., Parodi, A., and Pasquero, C.: Aerosol indirect effects in complex-orography areas: a numerical
958 study over the Great Alpine Region, Atmos. Chem. Phys, 22, 3901–3909, <https://doi.org/10.5194/acp-22-3901-2022>,
959 2022.

960 Panosetti, D., and Tomassetti, U.: The July 2023 Northern Italy hailstorms from a climatological and (re) insurance market
961 perspective, EGU General Assembly, 3901–3916, 2024.

962 Púčik, T., Castellano, C., Groenemeijer, P., Kühne, T., Rädler, A. T., Antonescu, B., Faust, E., J., H., and Kunz, M.: Large
963 hail incidence and its economic and societal impacts across Europe, Monthly Weather Review, 147, 159–184, 2019.

964 Punge, H. J., Bedka, K. M., Kunz, M., and Reinbold, A.: Hail frequency estimation across Europe based on a combination
965 of overshooting top detections and the ERA-INTERIM reanalysis, Atmospheric Research, 198, 34–43,
966 <https://doi.org/10.1016/j.atmosres.2017.07.025>, 2017.

967 Ralph, F. M. and Coauthors: Atmospheric Rivers, Springer Nature, 252pp, ISBN 978-3-030-28905-8, 2020.

968 Repetto, M. P., Burlando, M., Solari, G., Gaetano, P., Pizzo, M., and Tizzi, M.: A web-based GIS platform for the safe
969 management and risk assessment of complex structural and infrastructural systems exposed to wind, Advances in
970 Engineering Software, 117, 29–45, <https://doi.org/10.1016/j.advengsoft.2017.03.002>, 2018.

971 Ricchi, A., Rotunno, R., Miglietta, M. M., Picciotti, E., Montopoli, M., Marzano, F. S., Baldini, L., Vulpiani, G., Tiesi,
972 A., and Ferretti, R.: Analysis of the development mechanisms of a large-hail storm event on the Adriatic Sea, Atmospheric
973 Research, 296, 107079, <https://doi.org/10.1016/j.atmosres.2023.107079>, 2023a.

974 Ricchi, A., Sangelantoni, L., Redaelli, G., Mazzarella, V., Montopoli, M., Miglietta, M. M., Tiesi, A., Mazzà, S., Rotunno,
 975 R., and Ferretti, R.: Impact of the SST and topography on the development of a large-hail storm event, on the Adriatic
 976 Sea, *Atmospheric Research*, 296, 107078, <https://doi.org/10.1016/j.atmosres.2023.107078>, 2023b.
 977 Rosso, R. and Ceppi, A.: Land–Sea Distribution of Ground Precipitation in Mediterranean Storms, *Water*, 15,
 978 <https://doi.org/10.3390/w15101894>, 2023.
 979 Rotach, M. W.: A collaborative effort to better understand, measure and model atmospheric exchange processes over
 980 mountains, *Bulletin of the American Meteorological Society*, 103, E1282-E1295, [https://doi.org/10.1175/BAMS-D-21-](https://doi.org/10.1175/BAMS-D-21-0232.1)
 981 0232.1, 2022.
 982 Segadelli, S., Grazzini, F., Adorni, M., Nardo, M. T., Fornasiero, A., Chelli, A., and Cantonati, M.: Predicting Extreme-
 983 Precipitation Effects on the Geomorphology of Small Mountain Catchments: Towards an Improved Understanding of the
 984 Consequences for Freshwater Biodiversity and Ecosystems, *Water*, 12, 79, <https://doi.org/10.3390/w12010079>, 2020.
 985 Serafin, S.: Multi-scale transport and exchange processes in the atmosphere over mountains - Programme and experiment,
 986 Innsbruck University Press, 42 pp., <https://doi.org/10.15203/99106-003-1.>, 2020.
 987 Sioni, F., Davolio, S., Grazzini, F., and Giovannini, L.: Revisiting the atmospheric dynamics of the two century floods
 988 over north-eastern Italy, *Atm. Res*, 286, 106662, <https://doi.org/10.1016/j.atmosres.2023.106662>, 2023.
 989 Solari, G., Repetto, M. P., Burlando, M., Gaetano, P., Pizzo, M., Tizzi, M., and Parodi, M.: The wind forecast for safety
 990 management of port areas, *Journal of Wind Engineering and Industrial Aerodynamics*, 104–106, 266-277,
 991 <https://doi.org/10.1016/j.jweia.2012.03.029>, 2012.
 992 Solari, G., Burlando, M., Gaetano, P., and Repetto, M. P.: Characteristics of thunderstorms relevant to the wind loading
 993 of structures, *Wind and Structures*, 20, 763-791, <https://doi.org/10.12989/WAS.2015.20.6.763>, 2015.
 994 Stier, P.: Multifaceted aerosol effects on precipitation, *Nat. Geosci.*, 17, 719-732, [https://doi.org/10.1038/s41561-024-](https://doi.org/10.1038/s41561-024-01482-6)
 995 01482-6, 2024.
 996 Torcasio, R. C., Mascitelli, A., Realini, E., Barindelli, S., Tagliaferro, G., Puca, S., Dietrich, S., and Federico, S.: The
 997 impact of global navigation satellite system (GNSS) zenith total delay data assimilation on the short-term precipitable
 998 water vapor and precipitation forecast over Italy using the Weather Research and forecasting (WRF) model, *Nat. Hazards*
 999 *Earth Syst. Sci*, 23, 3319–3336, 2023.
 1000 Torcasio, R. C., Papa, M., Del Frate, F., Mascitelli, A., Dietrich, S., Panegrossi, G., and Federico, S.: Data assimilation
 1001 of satellite-derived rain rates estimated by neural network in convective environments: a study over Italy, *Remote Sens.*,
 1002 16, 1769, <https://doi.org/10.3390/rs16101769>, 2024.
 1003 Tsai, C.-L., Kim, K., Liou, Y.-C., and Lee, G.: Advantages of using multiple Doppler radars with different wavelengths
 1004 for three dimensional wind retrieval, *Atmos. Meas. Tech*, 18, 6371-6392, <https://doi.org/10.5194/amt-18-6371-2025>,
 1005 2025.
 1006 Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of the Atmospheric Sciences*, 34,
 1007 1149-1152, 1977.
 1008 Vogel, F., Putero, D., Bonasoni, P., Cristofanelli, P., Zanatta, M., and Marinoni, A.: Saharan dust transport event
 1009 characterization in the Mediterranean atmosphere using 21 years of in-situ observations, *Atmos. Chem. Phys*, 25, 15453-
 1010 15468, <https://doi.org/10.5194/acp-25-15453-2025>, 2025.
 1011 Xu, C., Zhong, P.-A., Davolio, S., Drofa, O., Gambini, E., Ravazzani, G., and Ceppi, A.: Testing machine learning
 1012 algorithms as post-processing tools for hydro-meteorological modelling over a small river basin, *Environ. Model. Softw.*,
 1013 [https://doi.org/10.1016/j.envsoft.2025.106592.](https://doi.org/10.1016/j.envsoft.2025.106592), 2025.

1014 Zanatta, M., Mertes, S., Jourdan, O., Dupuy, R., Järvinen, E., Schnaiter, M., Eppers, O., Schneider, J., Jurányi, Z., and
 1015 Herber, A.: Airborne investigation of black carbon interaction with low-level, persistent, mixed-phase clouds in the Arctic
 1016 summer, *Atmospheric Chemistry and Physics*, 23, 7955–7973, <https://doi.org/10.5194/acp-23-7955-2023>, 2023.
 1017 Zanini, M. A., Hoferl, L., Faleschinil, F., and Pellegrino, C.: Building damage assessment after the Riviera del Brenta
 1018 tornado, northeast Italy, *Nat. Hazards*, 86, 1247–1273, <https://doi.org/10.1007/s11069-017-2741-6>, 2017.
 1019 Zhang, S., Solari, G., Gaetano, P., Burlando, M., and Repetto, M. P.: A refined analysis of thunderstorm outflow
 1020 characteristics relevant to the wind loading of structures, *Probabilistic Engineering Mechanics*, 54, 9-24,
 1021 <https://doi.org/10.1016/j.pro bengmech.2017.06.003>, 2018.
 1022 Zhao, C., Sun, Y., Yang, J., Li, J., Zhou, Y., Yang, Y., Fan, H., and Zhao, X.: Observational evidence and mechanisms
 1023 of aerosol effects on precipitation, *Science Bulletin*, 69, 1569-1580, <https://doi.org/10.1016/j.scib.2024.03.014>, 2024.