

The pH and Phosphorus Availability as Primary Drivers of Compost-Induced CO₂ Emissions from Malaysian Tropical Soil: Empirical Relationships

Xingxing Cheng¹, Faridah Othman¹, Rosazlin Abdullah², Chiu Chuen Onn^{1*}


¹Department of Civil Engineering, Faculty of Engineering, Universiti Malaya, 50603 Kuala Lumpur, Malaysia

²Institute of Biological Sciences, Faculty of Science, Universiti Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence to: Chiu Chuen Onn (onnchiuchuen@um.edu.my)

Abstract. Confronting the global need for climate-smart agriculture, this study investigated the major factors controlling CO₂ emissions from a Malaysian tropical soil amended with four composts. Multiple regression analyses identified soil available phosphorus (AP) and pH as the key interactive drivers of CO₂ emissions, in the following order:: chicken dung compost (CDCS) > sludge compost (SLS) > goat manure-leaf compost (GLCS) > food waste compost (FWCS).The significantly higher emissions from CDCS were primarily due to its pronounced elevation of soil pH, likely stimulating microbial activity. The positive correlation with AP indicated that enhanced phosphorus availability further promoted microbial carbon mineralization. The findings demonstrate that compost is not a carbon-neutral amendment; its net climate impact depends on the specific physic-chemical changes it induces in the soil. This provides a scientific basis for optimizing compost selection to reconcile soil fertility improvement with greenhouse gas mitigation in tropical agroecosystems.

Graphical abstract



20 In this experiment, four composts (FWC, GLC, CDC, SL) were applied at 2.5% and 10% rates to typical Malaysian tropical rainforest soils and incubated for 90 days. The CO₂ emission characteristics were measured in different soil types. To determine key drivers, the study examined correlations between soil physic-chemical properties and CO₂ emissions and evaluated the relative impact of each factor's impact.

25 **Keywords:** Soil; Compost; CO₂ emission; Available phosphoru; pH

1 Introduction

Confronting global climate change and the strategic goals of “carbon peak and carbon neutrality” (IPCC, 2022), agricultural soils play a pivotal dual role in the global carbon cycle due to their vast carbon storage capacity and active biogeochemical processes serving as both a crucial organic carbon sink (Bolinder et al., 2020; Shu et al., 2022; Rumpel et al., 2022; Gu et al., 2024) and a potential significant source of CO₂ emissions (Lal, 2004; Paustian et al., 2016). The stability of this massive carbon pool, estimated to exceed the carbon in vegetation and the atmosphere combined, is highly sensitive to land management practices (Amelung et al., 2020; Sánchez-Moreno et al., 2025), with its persistence now understood to be driven by functional complexity beyond mere input quantity (Lehmann et al., 2020). Consequently, strategies that can concurrently enhance soil carbon sequestration (SOC) and support agricultural productivity are critical for climate-resilient food systems.

In pursuit of sustainable agriculture, organic amendment practices, particularly compost application, have emerged as a cornerstone strategy. Composting transforms organic waste into a stable soil conditioner, closing nutrient loops and reducing reliance on synthetic fertilizers (Bong et al., 2018). Beyond waste management benefits, compost application is widely regarded as a core pathway for enhancing soil fertility by improving nutrient supply, water retention, and aggregate stability, and for actively promoting long-term carbon sequestration (Cotrufo et al., 2013; Oldfield et al., 2019). Recent meta-analyses confirm that organic amendments can significantly increase SOC stocks, with the magnitude of increase dependent on climate, soil type, and application duration (Bai et al., 2023; Huang et al., 2022; Mao et al., 2024). However, integrating this practice into climate-smart frameworks requires a nuanced understanding of its full carbon budget, accounting not only for gains in stable carbon but also for potential losses.

However, this practice entails a fundamental scientific dilemma: while the input of exogenous organic matter enhances the soil's carbon sequestration potential, it may also lead to substantial initial carbon loss by priming microbial metabolism (Bai et al., 2018; Li et al., 2023; Kuzyakov et al., 2000; Bernard et al., 2022). Therefore, precisely quantifying the dynamics of soil CO₂ emissions following compost application and deeply deciphering the underlying biogeochemical drivers have become indispensable scientific foundations for accurately assessing the net climate impact of agricultural management practices (Smith et al., 2020) and formulating truly “climate-smart” land management strategies (Lipper et al., 2024).

It is well established that adding compost to soil typically causes a short-term increase in CO₂ emissions. This occurs because compost introduces readily decomposable organic carbon, which rapidly stimulates microbial metabolism and accelerates carbon turnover (Sarker et al., 2017). Consequently, a large body of research has focused on monitoring CO₂ flux dynamics following compost application and on establishing its empirical relationships with environmental factors such as temperature and moisture. However, a critical limitation persists in the current understanding. There is a lack of systematic investigation into the key factors through which compost alters the soil's internal state to drive these emissions (Bardgett et al., 2021). Specifically, compost input triggers a cascade of changes in key soil properties, including the dynamics of labile carbon pools (Chen et al., 2019; Laffely et al., 2020), nitrogen availability (Liu et al., 2019; Li et al., 2021), pH (Aye et al., 2017; Xiao et al., 2021), and microbial biomass with its stoichiometry (Rosinger et al., 2022; Gurmessa et al., 2024; Razavi et al., 2016). Yet, studies that systematically and quantitatively link these specific physic-chemical changes to CO₂ emission dynamics remain insufficient (Sarkar et al., 2018; Chen et al., 2018). This gap in process-based understanding hinders our ability to accurately predict the net carbon balance of different compost management practices (Niu et al., 2024; Wang et al., 2024a; Tang et al., 2023), as the ultimate outcome depends on complex interactions among these multiple factors (Cho et al., 2024; Yang et al., 2023; Ma et al., 2023) and is highly influenced by initial soil conditions (McFarlane et al., 2024; Oldfield et al., 2024). This limitation, in turn, constrains the development of optimized strategies to maximize soil carbon sequestration potential (Xiao et al., 2021).

To address the research gaps identified above, we propose the following hypotheses. First, compost application is known to induce a short term pulse of soil CO₂ emissions. We therefore hypothesize that the magnitude and duration of these emissions will vary significantly among different compost types due to their distinct physicochemical properties. Second, compost simultaneously alters multiple soil properties, including labile carbon pools, nitrogen availability, pH, etc. We hypothesize that changes in these physicochemical parameters will collectively drive the dynamics of post compost CO₂ emissions. Third, many tropical soils are acidic and limited in phosphorus. We therefore hypothesize that compost induced increases in pH and available phosphorus may exert particularly strong influences on CO₂ emissions. Multiple regression analysis will be used to quantitatively assess the relative contribution of each factor and to identify the dominant drivers governing variations in CO₂ emissions.

To address the current gaps in process-based understanding and quantitative prediction, this study focuses on a Malaysian typical soil and aims to precisely quantify the soil carbon release process following compost input and to decipher its key driving factors. Through a controlled incubation experiment, this research will systematically achieve the following objectives: (1) to monitor the continuous dynamics of soil CO₂ emissions under different compost treatments; (2) to simultaneously determine changes in key soil physic-chemical parameters (including water-soluble organic carbon, microbial biomass carbon, mineral nitrogen, and pH, among others) post-incubation; and (3) to employ correlation analysis and multiple regression modeling to quantitatively assess the independent influence of each physic-chemical parameter on CO₂ flux and to calculate their relative contribution weights, thereby identifying the core driving factors governing emission variations.

85 Consequently, by quantifying the relationships between CO₂ emissions and multiple intrinsic soil properties (e.g., labile
carbon pools, nitrogen availability) following compost addition and identifying the dominant driving factors, this study
provides concrete mechanistic evidence for understanding the carbon cycle response of tropical soils at a regional scale. The
research findings can directly inform localized practices for agricultural soil carbon management in Malaysia, offering direct
data support and decision-making references for assessing the short-term carbon loss risk associated with compost
90 application and for optimizing application strategies to reconcile the conflict between soil improvement and carbon
emissions.

2 Materials and Methods

2.1 Soil and Compost Materials

Using a five-point random sampling strategy, a composite soil sample (10-20 cm depth) was collected from an Agricultural
95 Research Center of the University of Malaya and named ARC-soil. The soil is a highly weathered, acidic Ultisol typical of
the humid tropics, with low natural fertility, a clayey texture, and a mineralogy dominated by kaolinite and oxides. The
sample was air-dried, disaggregated, sieved (<2 mm), and stored airtight at 3°C until use. The methods for testing soil
physic-chemical properties are described in Supporting Information **Text 1**.

Four types of compost were obtained from the composting facility at the Faculty of Science, University of Malaya. They
100 were produced from distinct feedstocks and designated as: FWC, primarily derived from food waste; GLC, produced from a
mixture of goat manure and shredded leaves; CDC, originating from chicken dung and amended with sawdust and rice bran
as bulking agents; and SL, generated from dewatered food and beverage wastewater sludge. The four composts (FWC, GLC,
CDC, and SL) used in this study were purchased from the Faculty of Science, University of Malaya. Detailed information on
their production processes is not disclosed by the supplier. However, it is known that CDC and SL are commercially
105 available products, and additional phosphorus and potassium were supplemented during their production. To characterize the
compost quality and maturity, the following physicochemical properties were determined: Total phosphorus (TP) (0.22%,
0.42%, 0.68%, and 1.94%, respectively), total potassium (TK) (0.076%, 0.33%, 0.76%, and 4.43%), TOC (18.5%, 24.9%,
14.5%, and 15.8%), and pH (6.63, 6.12, 7.90, and 6.0). These parameters indicate that all four composts were well-stabilized
and suitable for the subsequent experiments.

110 2.2 Soil Incubation Experiment for CO₂ Emission Monitoring

The experiment involved preparing soil-compost mixtures at two amendment ratios (1:9 and 1:39, compost-to-soil w/w),
both adjusted to a total weight of 200 g per incubation unit. The two amendment ratios (2.5% and 10%, w/w) were selected
based on the following: (1) the 2.5% ratio corresponds to a typical field application rate of approximately 50 t/ha, which is
commonly used in local agricultural practice; (2) the 10% ratio was chosen to amplify the treatment effects, allowing for
115 clearer mechanistic observation of compost-induced changes in soil properties and CO₂ emissions under controlled

incubation conditions. Each of the four composts was tested at both ratios, creating eight distinct amended mixtures. These, along with a pure soil control (200 g), constituted the nine core treatments, each replicated three times for a total of 33 units. The soils were incubated at a constant temperature of 28 ± 2 °C. CO₂ efflux was measured on 14 occasions over the 90 day incubation period, specifically on days 1, 5, 7, 14, 28, 40, 47, 52, 60, 70, 80, and 90.

CO₂ efflux monitoring followed a pre-determined schedule of 14 samplings spanning 90 days. The measurement protocol required the temporary placement of a CO₂ trap (20 mL of 1 mol/L NaOH in an open container) in each vessel one day before a scheduled sampling. Following a 24-h accumulation period, the trap solution was removed, and its carbonate content was determined via back-titration to quantify the CO₂-C produced during that interval. The CO₂ emissions from soil were captured with NaOH and subsequently quantified by back-titration. This established method involved precipitating the trapped CO₂ as barium carbonate before titration with standardized acid, enabling the calculation of CO₂-C evolution, and was performed as described by Cheng et al. (2025).

2.3 Characterization of Soil and Compost Incubated Soil

Baseline soil properties were analyzed. Soil pH and electrical conductivity (EC) were measured in a 1:2.5 soil-to-water suspension. Total organic carbon (TOC) was determined by wet oxidation. AP was extracted with 0.5 M NaHCO₃ and measured spectrophotometrically. Ammonium (NH₄⁺-N) and nitrate (NO₃⁻-N) were extracted with 2mol/L KCl and analyzed by a continuous flow analyzer. Available potassium (AK) was extracted with 1 M NH₄OAc and quantified via flame atomic absorption spectroscopy (AAS). Cation exchange capacity (CEC) was determined by the ammonium acetate (NH₄OAc) saturation method at pH 7.0.

2.4 Statistical Analysis

One way analysis of variance (ANOVA) was performed to compare all soil physicochemical parameters (pH, Moisture, EC, CEC, TOC, TC, OM, NH₄⁺-N, NO₃⁻-N, TN, AP, and AK) among the nine treatments (ARC- soil control, FWCS 2.5%, FWCS 10%, GLCS 2.5%, GLCS 10%, CDCS 2.5%, CDCS 10%, SLS 2.5%, and SLS 10%). Tukey's honestly significant difference (HSD) post hoc test was used for multiple comparisons. Homogeneity of variances was assessed using Levene's test. A significance level of $p < 0.05$ was considered statistically significant. All statistical analyses were conducted using SPSS version 26.0 .

A standard multiple linear regression model (Montgomery et al., 2021) was employed to evaluate the effects of soil properties on the response variable, following the general form:

$$Y = \xi_0 + \sum \xi_i K_i + \epsilon \quad (1)$$

In this formulation, Y corresponds to the response or outcome variable. Each K_i corresponds to an explanatory or predictor variable included in the model. The parameters ξ_i quantify the partial effect of their respective K_i and Y, holding other variables constant. Finally, ϵ captures all unexplained variation, commonly known as the model residual.


Coefficients were estimated via ordinary least squares, which minimizes the sum of squared residuals. Model diagnostics ensured: (1) significant correlations between predictors and the response, (2) independent, normally distributed and

homoscedastic residuals, and (3) absence of severe multicollinearity among predictors. Only after satisfying these criteria
150 were the regression outputs deemed reliable.

3 Results


3.1 Changes in Soil Physic-chemical Properties Following Incubation

155 **Figure 1** presents physical photographs of soils incubated for 90 days with different compost types and application rates. The original ARC-soil appeared visually more aggregated than the compost-treated soils (Figure 1). . This observation clearly indicates that compost amendment alleviates soil compaction, with the mitigation effect becoming more pronounced as the compost application rate increases.



160 **Figure 1: Images illustrating soil appearance after incubation under different compost treatments (types: FWC, GLC, CDC, SL; application rates: 2.5% and 10% by mass).** (Quantitative measurements of bulk density were not performed because the visual difference was so pronounced that sample processing for other analyses was prioritized. Nevertheless, the visual observation suggests a qualitative improvement in soil structure following compost addition)

The detailed ANOVA results and Tukey HSD grouping letters are provided in Supporting Information Tables S4 and S5. One-way ANOVA followed by Tukey's HSD post hoc test ($p < 0.05$) was used to compare treatments. Compared to the unamended control, compost application significantly elevated soil TN content by 1.5- to 16-fold, and all compost-amended soils showed significantly higher TN than the control ($p < 0.05$). Both $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ levels were also markedly higher in amended soils, with GLCS-10% exhibiting the highest values among all treatments ($p < 0.05$). Furthermore, increasing the compost addition rate from 2.5% to 10% consistently enhanced the concentrations of all three nitrogen forms, and for each compost type, the 10% addition gave significantly higher values than the 2.5% addition ($p < 0.05$). Notably, while the original soil showed no significant difference between $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$, compost-treated soils exhibited consistently lower $\text{NH}_4^+\text{-N}$ than $\text{NO}_3^-\text{-N}$ across all application rates, indicating that compost addition promoted the nitrification process in the ARC-soil system. As shown in Figure 2(b), the ARC-soil had an AP content below the instrument's detection limit. However, as the compost addition increased from 2.5% to 10%, both AP and AK contents increased significantly. According to Tukey's HSD test, SLS-10% and SLS-2.5% showed the highest AP levels with no significant difference between them, while SLS-10% had the highest AK among all treatments ($p < 0.05$). Notably, the high AP and AK contents in CDCS and SLS were due to additional external additions during composting, whereas FWCS and GLCS did not receive external additions of AP and AK. Figure 2(c) demonstrates that incubation with compost generally increased soil TOC relative to the non-incubated control. Tukey HSD comparisons revealed that FWCS-10% and GLCS-10% gave significantly higher TOC than the control and all 2.5% treatments ($p < 0.05$). Among the amendments, soils receiving FWC and GLC exhibited higher TOC levels than those treated with CDC or SL. It should be noted that lower TOC in a given amendment does not directly translate to reduced CO_2 emissions, as the stability of the carbon present also plays a determining role.






Figure 2: A comparison between the original ARC-soil and compost-incubated soils is shown for: (a) nitrogen species ($\text{NH}_4^+\text{-N}$, $\text{NO}_3^-\text{-N}$, TN); (b) AP and AK; (c) carbon and organic matter (TOC, TC, OM); (d) pH; (e) electrical conductivity (EC); (f) moisture content; and (g) cation exchange capacity (CEC). Data are presented as mean $\pm$ SD (n = 3). Different lowercase letters above the bars indicate significant differences among treatments according to one-way ANOVA followed by Tukey's HSD post hoc test (p < 0.05). Bars sharing the same letter are not significantly different.

As shown in Figure 2(d), after adding 10% compost, the pH values of FWCS and GLCS were slightly lower than that of the original ARC-soil, with values of 6.17 and 5.86 (the ARC soil pH was 6.23), while the pH values of CDCS and SLS significantly increased. As shown in Figure 2(e), after adding 10% compost, the EC values of FWCS, GLCS, and SLS were approximately 2-8 times higher than that of the original soil, while the EC values of SLS-10% and SLS-2.5% increased to 18.02 and 7.07, which were approximately 34-13 times higher than that of the original ARC-Soil, and SLS-10% was significantly higher than all other treatments (p < 0.05). This indicates that the sludge composting from food and beverage wastewater contains a high level of soluble salts.

As shown in Figure 2(f), after adding 10% compost, the moisture contents of FWCS, GLCS, CDCS, and SLS after 90 days were 1.12%, 1.57%, 3.03%, and 2.63%, respectively, all of which were higher than that of the original ARC-soil (0.5%). Moreover, as the compost addition increased from 2.5% to 10%, the moisture content of FWCS, GLCS, CDCS, and SLS all increased to varying degrees with CDCS-10% being significantly higher than other treatments (p < 0.05). During the incubation experiment, irrigation time and water volume were kept constant with all samples, indicating that CDCS and SLS exhibited higher water retention than FWCS and GLCS. During the incubation experiment, the irrigation time and water

volume were kept consistent for all samples, indicating that CDCS and SLS exhibited higher water retention compared to FWCS and GLCS. As shown in Figure 2(g), after adding 10% compost, the CEC of the soil increased 3-5 times overall, with SLS-10% and CDCS-10% showing significantly higher CEC than FWCS-10% and GLCS-10% ($p < 0.05$). A higher CEC value indicates a stronger ability of the soil to retain fertilizer components, better soil fertility, and the potential to mitigate fluctuations in soil pH and EC. Therefore, the results of this study show that CDCS and SLS had higher soil fertility than FWCS and GLCS. Additionally, CDCS and SLS had better regulation of soil pH and EC fluctuations.

3.2 The dynamics of CO₂ flux from the soil were monitored throughout the incubation period

Throughout the first two weeks of incubation, CO₂ emissions from FWCS, GLCS, CDCS, and SLS remained consistently higher than those from ARC-soil, despite some instability during this period (Figure 3a). Over time, the CO₂ emissions from FWCS-10% / FWCS-2.5% and GLCS-10% / GLCS-2.5% significantly increased, reaching peak values around day 52 (0.264/0.245 and 0.266/0.242 g/(kg-soil·d), respectively). CO₂ emissions declined briefly from days 52 to 60, then stabilised thereafter. As shown in Figure 3(b), with the addition of 10%, CO₂ emissions from CDCS and SLS also exhibited instability, but remained higher than those of the original soil. After two weeks, CDCS emissions gradually increased, peaking around day 30, then decreasing over the next 20 days before stabilizing. In contrast, the CO₂ emissions from SLS showed a downward trend after 10 days and stabilized only after day 50. CO₂ emission trends for CDCS and SLS were consistent across the 2.5% and 10% application rates.




Figure 3: (a) Comparative CO₂ efflux in ARC-soil and FWCS (10%), FWCS 2.5% GLCS 10% and GLCS 2.5%; and (b) for ARC-soil, CDCS 10%, CDCS 2.5%; SLS 10% and SLS 2.5% during the 90-day period.

Figure 4 presents the cumulative CO₂ emissions from soils amended with four distinct compost types (FWCS, GLCS, CDCS, SLS) at two application rates, totalling of nine treatments. The emission patterns varied considerably depending on the compost type. As illustrated in Figure 4(a), soils treated with GLCS consistently exhibited slightly higher cumulative emissions than those receiving FWCS. Figure 4(b) further demonstrates that CDC-amended soils produced the highest

225 cumulative emissions throughout the experiment, significantly surpassing the levels observed in soils treated with SL, GLC, and FWC. Overall, the compost types can be ranked in descending order of their effect on cumulative soil CO₂ emissions as follows: CDCS > SLS > GLCS > FWCS.




Figure 4: (a) Cumulative CO₂ emission of ARC-soil, FWCS 10%, FWCS 2.5%, GLCS 10% and GLCS 2.5%; (b) for ARC-soil, CDCS 10%, CDCS 2.5%; SLS 10% and SLS 2.5% throughout the 90-d incubation period.

230 **3.3 Multiple Linear Regression Analysis (MLRA)**

Figure 5a and b present the correlation and significance matrices for twelve soil physic-chemical properties and CO₂ emissions. The analysis reveals that CO₂ efflux was closely linked to several factors, showing strong positive correlations with available phosphorus (0.79), pH (0.76), moisture content (0.74), available potassium (0.73), and electrical conductivity (0.62). Based on the magnitude of correlation coefficients, the relative influence of these key drivers on CO₂ emissions decreased in the order: AP > pH > moisture content > AK > EC.




Figure 5: Analysis of key drivers of CO₂ emission: (a) inter-variable correlations and (b) their statistical significance ($P < 0.05$) for compost-incubated soil.

Validating the absence of residual autocorrelation is critical for confirming that regression model assumptions hold. The Durbin-Watson statistic, which ranges from 0 to 4, reflects residual independence when near 2. Values deviating toward 0 or 4 imply possible autocorrelation. In this analysis, the obtained statistic of 1.851 (Table S1) closely approaches 2, indicating no substantial autocorrelation and thus satisfying the independence assumption.

The histogram of standardized residuals (Figure 6a) displays a shape consistent with normality, centered near zero with a standard deviation of about 0.5. Correspondingly, the close alignment of points along the reference line in the first quadrant of the normal probability plot (Figure 6b) also supports the normality of the residual distribution.




Figure 6: (a) Distribution Histogram; (b) P-P Plot for Normality Check; (c) Ridge Regression Trace; (d) Trade-off Curve: k vs. R².

Collinearity diagnostics based on tolerance and variance inflation factor (VIF) are summarized in [Table S3](#). Conventionally, tolerance < 0.2 or VIF > 10 indicates potential multicollinearity. In this dataset, total carbon (TC), moisture content, available phosphorus (AP), and available potassium (AK) all exhibited tolerance values below 0.2 and VIF values exceeding 10, confirming multicollinearity among these predictors.

Table 1. Coefficients

Model		Variables in the Equation				P	R ²	Adj-R ²
		B	SE (B)	Beta	B/SE (B)			
1	(Constant)	-1.561	0.677	0.000	-2.303			
	pH	0.726	0.115	1.331	0.651			
	EC	0.123	0.189	1.620	0.154			
	Moisture	-0.661	0.140	-1.261	-4.711	0.072	0.975	0.902
	AP	5.815	3.927	1.399	1.481			
	AK	-1.583	2.25	-2.144	-0.702			
	OM	0.0619	0.047	0.332	1.301			

Y=5.815×AP-1.583×AK+0.726×pH-0.661×Moisture+0.123×EC+0.062×OM

To addressed this problem, the study used ridge regression analysis. As depicted in [Figure 6c](#), which illustrates the trajectories of standardized coefficients across increasing values of the ridge parameter (k). Each coefficient curve descends rapidly before stabilizing, and all curves converge to a common range. Based on the stabilization patterns in [Figure 6c](#) and the corresponding MSE profile in [Figure 6c](#), a ridge parameter of k = 0.05 was selected. The resulting ridge-regression coefficients are reported in [Table 1](#). The established regression equation is:

$$Y = 5.815 \times AP - 1.583 \times AK + 0.726 \times pH - 0.661 \times Moisture + 0.123 \times EC + 0.062 \times OM$$

The regression model's overall validity was confirmed by analysis of variance (ANOVA), which yielded a statistically significant result (*p* < 0.1). The model demonstrated strong explanatory power, as evidenced by an adjusted R² value of 0.902, indicating that approximately 90.2% of the variance in the response variable is accounted for by the selected

predictors. Within the fitted equation, the estimated coefficient for each independent variable quantifies its individual effect size and direction on the dependent variable.

4 Discussion

4.1 Mechanisms of Compost in Improving Soil Properties

The application of compost significantly enhances key soil physic-chemical properties. Firstly, it stimulates microbial processes related to nitrogen cycling. As reported by [Zhang et al. \(2015\)](#), the addition of compost to lateritic soils increased the abundance of ammonia-oxidising bacteria, thereby promoting nitrification. In this study, the CDCS treatment exhibited the strongest nitrification activity, likely due to its highest pH among the treatments. This aligns with the finding that low pH is detrimental to NH_3 retention in soil, as it promotes the ionization of NH_3 to NH_4^+ , potentially creating a substrate shortage for nitrification and thus inhibiting the process ([Ying et al., 2010](#)). Furthermore, compost amendment markedly improves the bioavailability of phosphorus and potassium. Phosphorus in compost primarily exists in the form of mineral phosphates, such as calcium phosphate ($\text{Ca}_3(\text{PO}_4)_2$) and iron phosphate (FePO_4). Microbial activity during and after incorporation can solubilize these compounds, converting them into plant-available forms such as H_2PO_4^- and HPO_4^{2-} ([Lanno et al., 2021](#)). Potassium in compost remains relatively stable and bioavailable as it exists mainly in ionic form (K^+). During the composting process and subsequent soil incubation, these potassium ions are released into the soil solution, directly increasing the soil's readily AK content. The final levels of AP and AK contributed by compost depend on the initial nutrient content of the raw materials on any losses during the composting process.

4.2 Key Drivers of CO_2 Emissions: pH and Phosphorus as Primary Controls

The cumulative CO_2 emissions from the four compost-incubated soils followed a distinct descending order: CDCS > SLS > GLCS > FWCS. This pattern demonstrates that the specific type of compost significantly influences the magnitude of soil heterotrophic respiration. Notably, the CDCS treatment, characterized by the highest pH and substantial nutrient availability, also exhibited the greatest CO_2 efflux. This correlation suggests a strong link between compost-induced soil conditions and carbon mineralization rates.

Statistical analysis of influencing factors revealed that soil AP and pH were the primary drivers of CO_2 emissions in this compost-amended system, followed by moisture content and organic matter (OM). This hierarchy underscores the complex interplay between compost-derived nutrients, soil abiotic conditions, and microbial activity in controlling carbon dynamics.

The dominant role of pH stems from its comprehensive control over soil biogeochemistry. It is a master variable regulating microbial community composition and metabolic activity. For instance, bacterial growth rates are typically optimized in neutral to slightly alkaline conditions, and specific microbial taxa dominate at pH extremes (e.g., Actinobacteria in alkaline soils) ([Rousk et al., 2009](#); [Fierer et al., 2006](#)). Most extracellular enzymes involved in decomposition also exhibit peak activity between pH 6 and 8. Crucially, pH influences the stability of soil organic matter (SOM). Lower pH increases the positive surface charge on iron and aluminium oxides, enhancing their capacity to adsorb negatively charged organic molecules and thereby promoting SOM stabilization ([Wang et al., 2024b](#); [Oliver et al., 2019](#)). Conversely, a rise in pH, as

seen in CDCS, can accelerate the decomposition of particulate organic matter (Hayashi et al., 2024), explaining the higher CO₂ emissions from this treatment.

Phosphorus availability emerged as another critical factor. In many ecosystems, especially tropical ones, phosphorus is a key limiting nutrient for biological activity (Yang et al., 2014). The addition of compost, which increased soil AP, likely alleviated this limitation. As suggested by Antonio et al. (2024), phosphorus addition can indirectly stimulate CO₂ emissions by promoting the desorption of organic carbon compounds from mineral surfaces, thereby increasing substrate availability for microbes. This mechanism may explain the positive correlation between AP and CO₂ flux observed here. Soil moisture content, a secondary yet significant driver, primarily regulates the diffusion of oxygen within soil pores. This, in turn, governs the dominant mode of microbial respiration (aerobic vs. anaerobic) and overall microbial activity, making it a fundamental regulator for the soil biota responsible for decomposition (Lacroix et al., 2021).

5 Conclusion

This study examined the effects of adding four types of compost (FWCS, GLCS, CDCS, SLS) at two rates (2.5% and 10%) on ARC-soil properties and CO₂ emissions. The main findings are:

(1) Compost addition improved soil physical properties (e.g., reduced compaction, increased moisture retention) and significantly enhanced soil fertility. Total nitrogen, ammonium, nitrate, available phosphorus, and available potassium all increased markedly. Organic carbon increased significantly only under FWCS-10% and GLCS-10%. Notably, CDCS and SLS treatments resulted in higher pH, electrical conductivity (EC), and nutrient availability compared to FWCS and GLCS.

(2) CO₂ emissions from amended soils were consistently higher than from the control, with distinct temporal patterns. Cumulative emissions over the 90-day incubation followed the order: CDCS > SLS > GLCS > FWCS, indicating that compost type strongly influenced heterotrophic respiration.

(3) Statistical analysis identified available phosphorus (AP) and soil pH as the primary drivers of CO₂ emissions, followed by moisture content and available potassium (AK).

(4) The observed CO₂ emission patterns are linked to compost-induced changes in soil conditions. Higher pH likely stimulated microbial activity and organic matter decomposition, while increased AP alleviated phosphorus limitation. Moisture content further modulated microbial respiration dynamics.

These findings advocate for precision compost management. The core strategy is to select or formulate mature composts with near-neutral pH and controlled phosphorus release, guided by soil testing. This approach mitigates carbon loss risks while delivering agronomic benefits, advancing climate-smart agriculture.

We acknowledge that this study was conducted under controlled laboratory incubation conditions, which may not fully capture the complexity of field environments (e.g., temperature fluctuations, plant root activity, and longer-term carbon dynamics). Future field-scale studies with multi-year monitoring are needed to validate these findings under real agricultural conditions.

Code and data availability

The code and data are available from the corresponding author upon reasonable request.

330 **Author contributions**

Xingxing Cheng conceived and designed the study, developed the methodology, performed the formal analysis and data curation, conducted the validation, and wrote the original draft. Faridah Othman and Rosazlin Abdullah contributed to the investigation and reviewed and edited the manuscript. Chiu Chuen Onn acquired funding, supervised the project, contributed to the methodology and formal analysis, and reviewed and edited the manuscript. The final manuscript was prepared with
335 contributions from all co-authors..

Competing interests

The authors declare the following financial interests/personal relationships, which may be considered as potential competing interests: Chiu Chuen, Onn reports financial support was provided by Malaysia Ministry of Higher Education. If there are other authors, they declare that they have no known competing financial interests or personal relationships that could have
340 appeared to influence the work reported in this paper.

Disclaimer

Copernicus Publications adds a standard disclaimer: “Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the
345 authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.”
Please feel free to add disclaimer text at your choice, if applicable.

Acknowledgements

The author would like to thank [Universiti of malaya] for their support and assistance..

Financial support

350 This work was supported by the Ministry of Higher Education Malaysia under the Fundamental Research Grant Scheme (FRGS) [No. FRGS/1/2022/TK01/UM/02/2], and the Young Innovative Talent Project-Guangdong Scientific Research Platform and Projects for the Higher-educational Institution & Education Science Planning Scheme [NO. KY2022036401].

Review statement

The review statement will be added by Copernicus Publications listing the handling editor as well as all contributing referees
355 according to their status anonymous or identified.

References

- Amelung, W., Bossio, D., de Vries, W., Kögel-Knabner, I., Lehmann, J., Amundson, R., Bol, R., Collins, C., Lal, R.,
Leifeld, J., Minasny, B., Pan, G., Paustian, K., Rumpel, C., Sanderman, J., van Groenigen, J. W., Mooney, S., van
Wesemael, B., Wander, M., and Chabbi, A.: Towards a global-scale soil climate mitigation strategy, *Nat. Commun.*, 11,
360 5427, <https://doi.org/10.1038/s41467-020-18887-7>, 2020.
- Antonio, R. S., María, C. D. C., José, T., Cooledge, E. C., Chadwick, D. R., and Jones, D. L.: Phosphorus fertilization
promotes carbon cycling and negatively affects microbial carbon use efficiency in agricultural soils: Laboratory
incubation experiments, *Geoderma*, 450, 117038, <https://doi.org/10.1016/j.geoderma.2024.117038>, 2024.
- Bai, Z., Caspari, T., Gonzalez, M. R., Batjes, N. H., Mäder, P., Bünemann, E. K., de Goede, R., Brussaard, L., Xu, M.,
365 Ferreira, C. S. S., Reintam, E., Fan, H., Mihelič, R., Glavan, M., and Tóth, Z.: Effects of agricultural management
practices on soil organic carbon and greenhouse gas emissions: A synthesis, *Adv. Agron.*, 151, 163–
277, <https://doi.org/10.1016/bs.agron.2018.05.001>, 2018.
- Bardgett, R. D., Bullock, J. M., Lavorel, S., Manning, P., Schaffner, U., Ostle, N., Chomel, M., Durigan, G., Fry, E. L.,
Johnson, D., Lavalley, J. M., Le Provost, G., Luo, S., Png, K., Sankaran, M., Hou, X., Zhou, H., Ma, L., Ren, W., and
370 Shi, H.: Combatting global grassland degradation, *Nat. Rev. Earth Environ.*, 2, 720–
735, <https://doi.org/10.1038/s43017-021-00207-2>, 2021.
- Bolinder, M. A., Crotty, F., Elsen, A., Frac, M., Kismányoky, T., Lipiec, J., Tits, M., Tóth, Z., and Kätterer, T.: The effect
of crop residues, cover crops, manures and nitrogen fertilization on soil organic carbon changes in agroecosystems: A
synthesis of reviews, *Nutr. Cycl. Agroecosyst.*, 118, 1–22, <https://doi.org/10.1007/s11027-020-09916-3>, 2020.
- 375 Bong, C. P. C., Lim, L. Y., Lee, C. T., Klemeš, J. J., Ho, C. S., and Ho, W. S.: The characterisation and treatment of food
waste for improvement of biogas production during anaerobic digestion – A review, *J. Clean. Prod.*, 172, 1545–1558,
<https://doi.org/10.1016/j.jclepro.2017.10.199>, 2018
- Cho, H. N., Shin, M., Lee, I., Ryoo, H., Acharya, B. S., Park, J. H., Cheong, Y. H., Cho, J. S., and Kang, S. W.: Correction:
Impact of biochar and compost amendment on corn yield and greenhouse gas emissions under waterlogged conditions,
380 *Appl. Biol. Chem.*, 67, 14, <https://doi.org/10.1186/s13765-024-00868-9>, 2024.
- Chen, J., Luo, Y., van Groenigen, K. J., Hungate, B. A., Cao, J., Zhou, X., and Wang, R.: A keystone microbial enzyme for
nitrogen control of soil carbon storage, *Sci. Adv.*, 4, eaaq1689, <https://doi.org/10.1126/sciadv.aaq1689>, 2018.

- Chen, J., Seven, J., Zilla, T., Dippold, M. A., Blagodatskaya, E., and Kuzyakov, Y.: Microbial C:N:P stoichiometry and turnover depend on nutrients availability in soil: A ^{14}C , ^{15}N and ^{33}P triple labelling study, *Soil Biol. Biochem.*, 131, 206–216, <https://doi.org/10.1016/j.soilbio.2019.01.017>, 2019.
- Cheng, X., Onn, C. C., Othman, F., Abdullah, R., Guo, W., Gunn, P. F. E., and Yue, L.: Impact of biochars on CO_2 emissions and physic-chemical properties in Malaysian tropical soils: Factor analysis and weighted contribution, *J. Environ. Sci.*, 158, 84–97, <https://doi.org/10.1016/j.jes.2025.03.002>, 2025.
- Shu, X. Y., He, J., Zhou, Z. H., Xia, L. L., Hu, Y. F., Zhang, Y. L., Zhang, Y. Y., Luo, Y. Q., Chu, H. Y., Liu, W. J., Yuan, S., Gao, X. S., and Wang, C. Q.: Organic amendments enhance soil microbial diversity, microbial functionality and crop yields: A meta-analysis, *Sci. Total Environ.*, 829, 154627, <https://doi.org/10.1016/j.scitotenv.2022.154627>, 2022.
- Cotrufo, M. F., Wallenstein, M. D., Boot, C. M., Deneff, K., and Paul, E. A.: The Microbial Efficiency-Matrix Stabilization (MEMS) framework integrates plant litter decomposition with soil organic matter stabilization: do labile plant inputs form stable soil organic matter?, *Glob. Change Biol.*, 19(4), 988–995, <https://doi.org/10.1111/gcb.12113>, 2013.
- McFarlane, K. J., Cusack, D. F., Dietterich, L. H., Hedgpeth, A. L., Finstad, K. M., and Nottingham, A. T.: Experimental warming and drying increase older carbon contributions to soil respiration in lowland tropical forests, *Nat. Commun.*, 15, 7084, <https://doi.org/10.1038/s41467-024-51422-6>, 2024.
- Niu, J., Saeed, Q., Wang, W., Zhang, R., Liu, L., Lv, F., Xu, J., Han, Y., Zhang, P., Hu, C., Xu, H., Sun, B., Yang, X., and Zhang, S.: Manure replacing synthetic fertilizer improves crop yield sustainability and reduces carbon footprint under winter wheat–summer maize cropping system, *J. Environ. Manage.*, 358, 120936, <https://doi.org/10.1016/j.jenvman.2024.120936>, 2024.
- Yang, Y., Cheng, S. L., Fang, H. J., Guo, Y. F., Li, Y. N., and Zhou, Y.: Interactions between soil organic matter chemical structure and microbial communities determine the spatial variation of soil basal respiration in boreal forests, *Appl. Soil Ecol.*, 183, 104743, <https://doi.org/10.1016/j.apsoil.2022.104743>, 2023.
- Liu, H. M., Zhou, G. Y., Zhou, X. H., Zhou, Y. S., Jiang, Q., and Guo, D. L.: Differential response of soil respiration to nitrogen and phosphorus addition in a highly phosphorus-limited subtropical forest, China, *For. Ecol. Manage.*, 448, 499–508, <https://doi.org/10.1016/j.foreco.2019.06.033>, 2019.
- Fierer, N. and Jackson, R. B.: The diversity and biogeography of soil bacterial communities, *Proc. Natl. Acad. Sci. USA*, 103, 626–631, <https://doi.org/10.1073/pnas.0507535103>, 2006.
- Rumpel, C., Amiraslani, F., Bossio, D., Chenu, C., Henry, B., Fuentes Espinoza, A., Koutika, L.-S., Ladha, J., Madari, B., Minasny, B., Olaleye, A. O., Shirato, Y., Sall, S. N., Soussana, J.-F., and Varela-Ortega, C.: The role of soil carbon sequestration in enhancing human resilience in tackling global crises including pandemics, *Soil Secur.*, 8, 100069, <https://doi.org/10.1016/j.soisec.2022.100069>, 2022.
- Li, Y., Wu, L., Tang, L., Ren, F., Wang, X., Zhu, P., Sun, N., and Xu, M.: Manure application decreases soil organic carbon priming by increasing mineral protection and nitrogen availability, *Geoderma*, 439, 116676, <https://doi.org/10.1016/j.geoderma.2023.116676>, 2023.

- Aye, N. S., Butterly, C. R., Sale, P. W. G., and Tang, C.: Residue addition and liming history interactively enhance mineralization of native organic carbon in acid soils, *Biol. Fertil. Soils*, 53, 61–75, <https://doi.org/10.1007/s00374-016-1156-y>, 2017.
- 420 Hayashi, R., Maie, N., Wagai, R., Hirano, Y., Matsuda, Y., Okamoto, T., Wada, R., and Tanikawa, T.: Soil acidity accelerates soil organic matter decomposition in *Cryptomeria japonica* stands and *Chamaecyparis obtusa* stands, *Plant Soil*, 494, 627–649, <https://doi.org/10.1007/s11104-023-06308-9>, 2024.
- Bai, X., Tang, J., Wang, W., Ma, C., and Shi, L.: Organic amendment effects on cropland soil organic carbon and its implications: A global synthesis, *CATENA*, 231, 107343, <https://doi.org/10.1016/j.catena.2023.107343>, 2023.
- 425 IPCC: Climate Change 2022: Mitigation of Climate Change. Contribution of Working Group III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157926>, 2022.
- Gu, B., Qin, T., Qiu, M., Yu, J., Zhang, L., Li, Y., and Hatano, R.: Addition of exogenous organic ameliorants mediates soil bacteriome and microbial community carbon source utilization pattern in coastal saline-alkaline soil, *Agriculture-Basel*, 430 14, 44, <https://doi.org/10.3390/agriculture14010044>, 2024.
- Kuzyakov, Y., Friedel, J. K., and Stahr, K.: Review of mechanisms and quantification of priming effects, *Soil Biol. Biochem.*, 32, 1485–1498, [https://doi.org/10.1016/S0038-0717\(00\)00084-5](https://doi.org/10.1016/S0038-0717(00)00084-5), 2000.
- Lacroix, E. M., Rossi, R. J., Bossio, D., and Fendorf, S. E.: Effects of moisture and physical disturbance on pore-scale oxygen content and anaerobic metabolisms in upland soils, *Sci. Total Environ.*, 780, 435 146572, <https://doi.org/10.1016/j.scitotenv.2021.146572>, 2021.
- Lal, R.: Soil carbon sequestration impacts on global climate change and food security, *Science*, 304, 1623–1627, <https://doi.org/10.1126/science.1097396>, 2004.
- Lanno, M., Kriipsalu, M., Shanskiy, M., Silm, M., and Kisand, A.: Distribution of Phosphorus Forms Depends on Compost Source Material, *Resources*, 10, 102, <https://doi.org/10.3390/resources10100102>, 2021.
- 440 Lehmann, J., Hansel, C. M., Kaiser, C., Kleber, M., Maher, K., Manzoni, S., Nunan, N., Reichstein, M., Schimel, J. P., Torn, M. S., Wieder, W. R., and Kögel-Knabner, I.: Persistence of soil organic carbon caused by functional complexity, *Nat. Geosci.*, 13, 529–534, <https://doi.org/10.1038/s41561-020-0612-3>, 2020.
- Huang, X. L., Jia, Z. X., Jiao, X. Y., Wang, J. L., and Huang, X. F.: Long-term manure applications to increase carbon sequestration and macroaggregate-stabilized carbon, *Soil Biol. Biochem.*, 174, 108827, 445 <https://doi.org/10.1016/j.soilbio.2022.108827>, 2022.
- Li, J., Pei, J. M., Pendall, E., Fang, C. M., and Nie, M.: Stoichiometric imbalance and microbial community regulate microbial elements use efficiencies under nitrogen addition, *Soil Biol. Biochem.*, 156, 108207, <https://doi.org/10.1016/j.soilbio.2021.108207>, 2021.
- Lipper, L., Thornton, P., Campbell, B. M., Baedeker, T., Braimoh, A., Bwalya, M., Caron, P., Cattaneo, A., Garrity, D., 450 Henry, K., Hottle, R., Jackson, L., Jarvis, A., Kossam, F., Mann, W., McCarthy, N., Meybeck, A., Neufeldt, H.,

- Remington, T., Sen, P. T., Sessa, R., Shula, R., Tibu, A., and Torquebiau, E. F.: Climate-smart agriculture for food security, *Nat. Clim. Change*, 4, 1068–1072, <https://doi.org/10.1038/nclimate2437>, 2024.
- Gurmessa, B., Cocco, S., Ashworth, A. J., Udawatta, R. P., Cardelli, V., Serrani, D., Ilari, A., Foppa Pedretti, E., Fornasier, F., and Corti, G.: Enzyme activities and microbial nutrient limitations in response to digestate and compost additions in organic matter poor soils in the Marches, Italy, *Soil Till. Res.*, 242, 106136, <https://doi.org/10.1016/j.still.2024.106136>, 2024.
- Sánchez-Moreno, S., Ros, M. B. H., Walder, F., Peixoto, L., Gómez-Gallego, T., Jeanbille, M., Juhanson, J., Feiza, V., Aponte, C., Pulido-Moncada, M., Parasotas, I., Kochiieru, M., Amaleviciute-Volunge, K., Dammie, N., van der Heijden, M. G. A., Ramos, J. L., Feizene, D., Philippot, L., Munkholm, L., Velthof, G. L., Petersen, S. O., Hallin, S., and Goberna, M.: A global dataset of experimental agricultural management on soil carbon accrual, its synergies and trade-offs, *Sci. Data*, 12, 929, <https://doi.org/10.1038/s41597-025-05238-8>, 2025.
- Ma, L., Wang, Y., Liu, H., Wang, F., Zhang, X., and Li, Z.: Differential effects of compost and biochar on CO₂ and CH₄ emissions from paddy soil: Insight into dissolved organic matter composition and microbial function, *J. Clean. Prod.*, 397, 136547, <https://doi.org/10.1016/j.jclepro.2023.136547>, 2023.
- Montgomery, D. C., Peck, E. A., and Vining, G. G.: *Introduction to Linear Regression Analysis*, 6th Ed., John Wiley & Sons, Hoboken, NJ, USA, <https://www.wiley.com/en-us/Introduction+to+Linear+Regression+Analysis%2C+6th+Edition-p-9781119578727>, 2021.
- Oldfield, E. E., Lavallee, J. M., Blesh, J., Bradford, M. A., Cameron-Harp, M., Cotrufo, M. F., Eagle, A. J., Eash, L., Even, R. J., Kuebbing, S. E., Kort, E. A., Lark, T. J., Latka, C., Lin, Y., Machmuller, M. B., O'Neill, B., Raffeld, A. M., RoyChowdhury, T., Rudek, J., Sanderman, J., Sprunger, C. D., Toombs, T. P., Uludere Aragon, N., Vidal, M., Woolf, D., Zelikova, T. J., and Gordon, D. R.: Greenhouse gas mitigation on croplands: clarifying the debate on knowns, unknowns and risks to move forward with effective management interventions, *Carbon Manag.*, 15, <https://doi.org/10.1080/17583004.2024.2365896>, 2024.
- Oldfield, E. E., Bradford, M. A., and Wood, S. A.: Global meta-analysis of the relationship between soil organic matter and crop yields, *SOIL*, 5, 15–32, <https://doi.org/10.5194/soil-5-15-2019>, 2019.
- Oliver, D. P., Li, Y., Orr, R., Nelson, P., Barnes, M., McLaughlin, M. J., and Kookana, R. S.: The role of surface charge and pH changes in tropical soils on sorption behaviour of per- and polyfluoroalkyl substances (PFASs), *Sci. Total Environ.*, 673, 197–206, <https://doi.org/10.1016/j.scitotenv.2019.04.055>, 2019.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., and Smith, P.: Climate-smart soils, *Nature*, 532, 49–57, <https://doi.org/10.1038/nature17174>, 2016.
- Rumpel, C., Amiraslani, F., Bossio, D., Chenu, C., Henry, B., Fuentes Espinoza, A., Koutika, L.-S., Ladha, J., Madari, B., Minasny, B., Olaleye, A. O., Shirato, Y., Sall, S. N., Soussana, J.-F., and Varela-Ortega, C.: The role of soil carbon sequestration in enhancing human resilience in tackling global crises including pandemics, *Soil Secur.*, 8, 100069, <https://doi.org/10.1016/j.soisec.2022.100069>, 2022.

- 485 Razavi, B. S., Zarebanadkouki, M., Blagodatskaya, E., and Kuzyakov, Y.: Rhizosphere shape of lentil and maize: spatial distribution of enzyme activities, *Soil Biol. Biochem.*, 96, 229–237, <https://doi.org/10.1016/j.soilbio.2016.02.020>, 2016.
- Rosinger, C., Keiblinger, K. M., Rousk, J., and Sandén, H.: Shifts in microbial stoichiometry upon nutrient addition do not capture growth-limiting nutrients for soil microorganisms in two subtropical soils, *Biogeochemistry*, 159, 33–43, <https://doi.org/10.1007/s10533-022-00911-1>, 2022.
- 490 Rousk, J., Brookes, P. C., and Bååth, E.: Contrasting soil pH effects on fungal and bacterial growth suggest functional redundancy in carbon mineralization, *Appl. Environ. Microbiol.*, 75, 1589–1596, <https://doi.org/10.1128/AEM.02753-08>, 2009.
- Sarker, J. R., Singh, B. P., He, X., Fang, Y., Li, G. D., Collins, D., and Cowie, A. L.: Tillage and nitrogen fertilization enhanced belowground carbon allocation and plant nitrogen uptake in a semi-arid canola crop–soil system, *Sci. Rep.*, 7, 10726, <https://doi.org/10.1038/s41598-017-11190-4>, 2017.
- 495 Mao, X., Sun, T., Zhu, L., Wanek, W., Cheng, Q., Wang, X., Zhou, J., Liu, X., Ma, Q., Wu, L., and Jones, D. L.: Microbial adaption to stoichiometric imbalances regulated the size of soil mineral-associated organic carbon pool under continuous organic amendments, *Geoderma*, 445, 116883, <https://doi.org/10.1016/j.geoderma.2024.116883>, 2024.
- Sarkar, B., Singh, M., Mandal, S., Churchman, G. J., and Bolan, N. S.: Clay minerals-organic matter interactions in relation to carbon stabilization in soils, *The Future of Soil Carbon: Its Conservation and Formation*, Elsevier, 71 – 86, <https://doi.org/10.1016/B978-0-12-811687-6.00003-1>, 2018.
- 500 Smith, P., Soussana, J.-F., Angers, D., Schipper, L., Chenu, C., Rasse, D. P., Batjes, N. H., van Egmond, F., McNeill, S., Kuhnert, M., Arias-Navarro, C., Olesen, J. E., Chirinda, N., Fornara, D., Wollenberg, E., Álvaro-Fuentes, J., Sanz-Cobena, A., and Klumpp, K.: How to measure, report and verify soil carbon change to realize the potential of soil carbon sequestration for atmospheric greenhouse gas removal, *Glob. Change Biol.*, 26, 219–241, <https://doi.org/10.1111/gcb.14815>, 2020.
- Tang, R., Liu, Y., Ma, R., Zhang, L., Li, Y., Li, G., Wang, D., Lin, J., Li, Q., and Yuan, J.: Effect of moisture content, aeration rate, and C/N on maturity and gaseous emissions during kitchen waste rapid composting, *J. Environ. Manage.*, 326, 116662, <https://doi.org/10.1016/j.jenvman.2022.116662>, 2023.
- 510 Bernard, L., Basile-Doelsch, I., Derrien, D., Fanin, N., Fontaine, S., Guenet, B., Karimi, B., Marsden, C., and Maron, P. A.: Advancing the mechanistic understanding of the priming effect on soil organic matter mineralisation, *Funct. Ecol.*, 36, 1355–1377, <https://doi.org/10.1111/1365-2435.14038>, 2022.
- Wang, H., Dong, W., Shao, D., Liu, L., Liao, B., Gu, W., Tang, C., Liu, J., Hu, W., Feng, J., and Huang, J.: Biochar enhances paddy productivity, carbon sequestration, and reduces greenhouse gas emissions in the Middle Yangtze River region, *Agronomy*, 14, 3067, <https://doi.org/10.3390/agronomy14123067>, 2024a.
- 515 Wang, X., Wang, T., Huang, Y., Liu, A., Li, Q., Wang, Y., Li, M., Fan, F., and Tang, Z.: Effect of biochars on the immobilization and form of Cadmium (Cd) in simulated Cd deposition of iron rich soils, *Ecotoxicol. Environ. Saf.*, 272, 116045, <https://doi.org/10.1016/j.ecoenv.2024.116045>, 2024b.

- Yang, X., Thornton, P., Ricciuto, D., and Post, W.: The role of phosphorus dynamics in tropical forests—A modeling study
520 using CLM-CNP, *Biogeosciences*, 11, 1667–1681, <https://doi.org/10.5194/bg-11-1667-2014>, 2014.
- Ying, J. Y., Zhang, L. M., and He, J. Z.: Putative ammonia-oxidizing bacteria and archaea in an acidic red soil with different
land utilization patterns, *Environ. Microbiol. Rep.*, 2, 304–312, <https://doi.org/10.1111/j.1758-2229.2009.00130.x>,
2010.
- Zhang, M. M., Wang, B. R., Li, D. C., He, J. Z., and Zhang, L. M.: Effects of long-term N fertilizer application and liming
525 on nitrification and ammonia oxidizers in acidic soils, *Acta Ecol. Sin.*, 35, 6362–
6370, <https://doi.org/10.5846/stxb201404150722>, 2015.
- Laffely, A., Erich, M. S., and Ohno, T.: Dissolved organic carbon chemical composition controls the rate of CO₂ release
from rewetted soil, *Soil Sci. Soc. Am. J.*, 84, 483–493, <https://doi.org/10.1002/saj2.20035>, 2020.
- Xiao, Q., Huang, Y., Wu, L., Tian, Y., Wang, Q., Wang, B., Xu, M., and Zhang, W.: Long-term manuring increases
530 microbial carbon use efficiency and mitigates priming effect via alleviated soil acidification and resource limitation,
Biol. Fertil. Soils, 57, 925–934, <https://doi.org/10.1007/s00374-021-01583-z>, 2021.