

## Contents

|                                     |           |
|-------------------------------------|-----------|
| <b>Response to reviewer 1 .....</b> | <b>1</b>  |
| <b>Response to reviewer 2 .....</b> | <b>11</b> |
| <b>Response to reviewer 3 .....</b> | <b>25</b> |

### **Response to reviewer 1**

#### **Reviewer comment:**

Overall, I found this study well written, straight to the point and very interesting. The EFAST decomposition is a simple yet robust method to explore whether an individual parameter contribute more or less than its coupling with other parameter in a qualitative AND quantitative way. The paper does a really good job describing the coupling and non-coupling effect of the different snow parameters on the TBs and the different GRs. Given the fact that this paper relies on MOSAiC snow data to estimate the range of variability of the snow and ice parameters, I think it would be valuable to have results of this parametrization that fits the MOSAiC TBs to confirm that the proposed configuration still falls within the observations range. I think this study is well suited for publication in The Cryosphere as it fits very well its scope, but some minor adjustments are needed prior publication.

Thank you for your positive and valuable comments. We have added additional experiments and enhanced the clarity of the methodology section in the revised manuscript. Please see our response (blue text) below

#### **Specific comments**

Given the fact that this paper relies on MOSAiC snow data to estimate the range of variability of the snow and ice parameters, I think it would be valuable to have results of this parametrization that fits the MOSAiC TBs to confirm that the proposed configuration still falls within the observations range.

We have added an additional TB simulation experiment using MOSAiC measurements together with collocated AMSR2 TBs. In this experiment, we mainly used the single-layer snow configuration corresponding to Sect. 4.1.1 of the sensitivity analysis.

Because direct measurements of snow liquid water content during MOSAiC were not publicly available, we therefore restricted this comparison to the winter period, when dry snow conditions are expected to dominate. Values for the snow depth, density, temperature and radius were taken from the MOSAiC measurements. For each snow observation site, profile measurements were averaged to obtain a representative value for the corresponding snow column.

To make the modelled TBs more comparable with AMSR2 observations, we also included sea ice and atmospheric datasets in the forward simulation. Specifically, the SMRT model was first used to simulate the microwave emission from the snow-covered sea ice surface. The simulated surface TBs were then propagated through the atmosphere using the Wentz atmospheric radiative transfer model (Wentz and Meissner 2000).

**Tab. R1 Snow, sea ice and atmosphere configurations used in the forward simulation**

| Layer      | Parameter                    | Setting or Source                                            |
|------------|------------------------------|--------------------------------------------------------------|
| Snow       | Depth                        | MOSAiC measurements                                          |
|            | Density                      |                                                              |
|            | Temperature                  |                                                              |
|            | Radius                       |                                                              |
|            | Stickiness                   | 0.15                                                         |
| Sea ice    | Thickness                    | Same as that in the original<br>Manuscript Tab. 4            |
|            | Density                      |                                                              |
|            | Temperature                  |                                                              |
|            | Radius                       |                                                              |
|            | Brine volume fraction        | Dataset from University of Bremen<br>(Ye and Heygster, 2015) |
|            | Porosity                     |                                                              |
|            | Multi-year ice concentration |                                                              |
| Atmosphere | Total water vapour           | ERA5 reanalysis                                              |
|            | Total cloud liquid water     | (Hersbach et al, 2018)                                       |

Overall, this SMRT configuration can reproduce the general magnitude and temporal variability of the AMSR2 observed TB, particularly at the intermediate-

frequency channels from 18 to 36 GHz (See Fig. R1). At the low-frequency channels (6 and 10 GHz), the simulated TBs are systematically lower than the AMSR2 observations. This discrepancy is likely related to the stronger sensitivity to sea ice temperature and brine volume fraction. The best agreement is found at 18, 23 and 36 GHz. At 89 GHz, both the observed and simulated TBs exhibit much stronger short-term variability. The model captures part of the observed TB fluctuation and the increase in late winter. Overall, the comparison supports the use of the adopted snow and sea ice configurations as a physically plausible basis for the sensitivity analysis.

Following your suggestion, we have included this analysis in the Supplementary Material.




**Fig. R1 Time-series comparison of SMRT-simulated TBs and AMSR2-observed TBs during the MOSAiC winter period.**

Moreover, following the suggestions of the other reviewers, we conducted a sensitivity analysis using the exponential microstructure model instead of the sticky hard sphere (SHS) microstructure model. The results are broadly consistent with the main conclusions of this study, and detailed experimental results are also provided in

the Supplementary Material.

Although the authors mention in the text that the “global” term refers to the full parameter space, having it in the title can be quite confusing as it mostly focuses on Arctic snow. In general, the paper does a very good job describing the theoretical coupling between the ASMR2-3 frequencies, but given the title there is this feeling it would be more connected to observations. The Section 3.2 “Microwave radiometer configuration” is in data yet just describes the AMSR2 frequencies. If no observational TBs are used, this should be moved to the Method section.

The Global Sensitivity Analysis (GSA) is a specific terminology and we would still prefer to retain it in the title.

As for the Section “Microwave radiometer configuration”, although the comparison of satellite measurements and model-based simulations is not presented in the main manuscript. Following the reviewer’s suggestion (see previous comment), we have added a model–observation comparison experiment in the Supplementary Material. We therefore would like to keep this section in the Data section and add more detailed information regarding the configuration of AMSR2 (Also see reply to the next comment).

Still in the section 3.2 or later in the text, I am surprised nothing is said about the major footprint’s size difference between the 6GHz and the higher frequency channels (36 and 89 GHz. For snow depth retrieval from satellite measurement, this cannot be ignored as the 6GHz footprint is tens of km wider than the 36GHz, so the GR36/6 may also show something artificial due to the heterogeneity between the two footprints.

In the revised manuscript, we have added a description of the AMSR2 spatial resolutions in Section 3.2, Page 7:

*“AMSR2 measures upwelling microwave TBs at six frequency bands centred at 6.9, 10.7, 18.7, 23.8, 36, and 89 GHz (henceforth denoted as 6, 10, 18, 23, 36 and 89 GHz for simplicity), operating at a conical incidence angle of approximately 55°. The corresponding footprint sizes are approximately  $62 \times 35$ ,  $42 \times 24$ ,  $22 \times 14$ ,  $26 \times 15$ ,  $12 \times 7$ , and  $5 \times 3$  km, respectively, from the lowest to the highest frequency.”*

The present SMRT-based retrievability analysis assumes a horizontally homogeneous snow and ice scenario and the same footprint across channels. We have

added the description in Conclusion Section (Page 29):

*“It should also be noted that the present SMRT-based retrievability analysis assumes a homogeneous surface and the same footprint across channels. In practical applications, footprint mismatch between low- and high-frequency channels, e.g. GR(36/6), could potentially introduce additional artefacts to snow depth retrieval. The resolution-harmonized Level-1R (L1R) TB data from JAXA may be useful here (Maeda et al. 2016).”*

L255-260 and Tables 2-3: how is snow ice and superimposed ice modelled in SMRT? Are you using fresh ice layer? Snow with such high density ( $>500$ ) can be tricky to configure in SMRT as some permittivity models can't characterize snow with density that high. I would maybe suggest to better describe the permittivity model used for each layer (snow, ice, fresh, dry, wet etc..)

We are aware that existing snow microstructural assumptions and scattering formulation (Sticky hard spheres + Improved born approximation) may become unstable in the intermediate- to high-density regime. In addition, in response to comments from the other reviewers, we revised the description of the “snow ice” layer in the manuscript. Because “**snow ice**” layer is very rare in the Arctic and the term may therefore cause ambiguity, we have replaced it with “**saline interlayer**” layer. Apart from this change in terminology and the corresponding revisions throughout the manuscript, the parameter settings used in the experiments remain unchanged.

We took three precautions in our study:

- 1) In the SMRT simulations, the saline interlayer is treated as a thin sea ice layer rather than a snow layer. The difference is in the physical parameterisation of the layer: when treated as sea ice, the dense saline layer is represented as an ice-rich medium with brine inclusions, rather than as a high-density snow layer composed of ice grains embedded in an air matrix.
- 2) We limited the maximum density of snow layers (wind slab and depth hoar) to  $500 \text{ kg m}^{-3}$ . Although the MOSAiC observations indicate that locally higher snow densities may occur, a density of  $500 \text{ kg m}^{-3}$  already corresponds to an ice volume fraction of approximately 0.55, which is slightly above the upper practical limit for applying the IBA+SHS simulation (0.5 given in Picard et al. 2022). We checked the simulated TB over the density range used for the snow

layers. The 3D TB figures (Fig.6 – 9 in manuscript) show smooth and continuous variations over 200–500 kg m<sup>-3</sup>, without abrupt fluctuations or discontinuities. This gives us confidence that the IBA+SHS configuration behaves stably within the snow density range.

- 3) We used the dense snow correction available in SMRT. When the ice volume fraction exceeds 0.5, this option treats the medium as air inclusions embedded in an ice matrix rather than ice particles embedded in an air matrix.

To describe this more clearly, we have added a model setup table to the Supplemental Material (Tab. R2).

Tab. R2 SMRT model setups

| Topic                                             | Snow                       | First year ice                                                                            |                                                     | Multi-year ice |
|---------------------------------------------------|----------------------------|-------------------------------------------------------------------------------------------|-----------------------------------------------------|----------------|
|                                                   |                            | (including saline interlayer and superimposed ice)                                        |                                                     |                |
| Scattering theory                                 |                            | Improved born approximation (with dense snow correction)                                  |                                                     |                |
| Microstructure Representation                     |                            | Sticky hard spheres                                                                       |                                                     |                |
| Surface/interface scattering models and substrate |                            | Default flat/Fresnel interfaces were used; a water substrate was added below the sea ice. |                                                     |                |
| Permittivity model                                | ice_permittivity_matzler87 | ice_permittivity_matzler06<br>brine_permittivity_stogryn85                                | ice_permittivity_matzler06<br>polder_van_stat<br>en |                |
| Radiative transfer solver                         |                            | Discrete Ordinate Radiative Transfer solver                                               |                                                     |                |
| SMRT version                                      |                            | 1.5.1                                                                                     |                                                     |                |

How saline was the snow? A 5% of brine volume fraction isn't something we can easily represent, as BVF is not something we directly measure on the field. It could be better to explicitly say the salinity value set in SMRT to get this BVF of 5%, as Figure 2 also

shows the salinity of the snow in PSU.

In this study, we didn't treat ordinary snow layers as saline. The salinity of the snow layers, including wind slab and depth hoar, is set to 0 PSU in the SMRT simulations. The reason is that the MOSAiC measurements show that most of the snowpack has very low or negligible salinity, while enhanced salinity is mainly observed near the snow–ice interface. This near-interface layer is also characterised by higher density and is physically different from the overlying snow. Therefore, in our model configuration, this saline and dense transitional layer is represented as a “saline interlayer” layer (“snow ice” layer in the previous version).

The BVF=5% in Table 4 is used for basal first year ice. It is based on MOSAiC measurements, we found that BVF generally fluctuates around 5% (Fig. R2). Although BVF increases to about 10% at deeper depths, around 1.5–2 m, these deeper layers are expected to have limited influence on the passive microwave signal because the penetration depth is less than 1 m at 6 GHz. Therefore, we used BVF = 5% for first year ice.




Fig. R2 Brine volume fraction measurements (First year ice) during MOSAiC

Figure 3, 4 and 5: the author's choice of underlying the TSI value is quite confusing and sometime makes the results not readable. It would be better to just write the numbers (such as for Figure 3-b 18H where the author forgot to put the underline)

We have revised Figs. 3, 4, and 5.

Figure 4 is mostly empty, I wonder if there would be a better way to present these sparse results.

Reply: Thank you for this suggestion. Fig. 4 is sparse because many parameters have

very low sensitivity at several AMSR2 channels. We considered alternative ways to present the results, including compressing or narrowing columns with generally low sensitivity, such as some temperature columns. However, we found that changing the column width would alter the visual meaning of the matrix and could unintentionally imply that these parameters are less important a priori than the others, rather than simply having low sensitivity under the tested conditions and channels.

Therefore, we decided to retain the full matrix structure with equal column widths. The empty cells are meaningful in this figure because they indicate parameters or channels for which the sensitivity is below the display threshold, and their contrast with the filled cells highlights the channel- and layer-dependent distribution of sensitivity.

For the figures showing the GR as function of snow depth and snow density (Figures 6, 7, 8 and 9), I would suggest removing the X-Y projection as it takes a lot of space and decrease the readability of the results and in a lot of case, they are showing no variability at all. They could be added as Supplementary material, according to the author's decision.

We have moved the X–Y projection panels from Figs. 8 and 9 to the Supplementary Material. We retain the X–Y projections in Figs. 6 and 7 because, for readers who are less familiar with the concept of retrievable range, the projections help illustrate how the averaging kernel diagnostics are converted into the retrievable domain.

L446: “The 89 GHz TBs are also susceptible to atmospheric effects...” This is also true for the 36GHz and in a lower extent the 18GHz.

We have revised the sentence. The new text reads (in Section 4.2.1, Page 20):

*“Atmospheric effects can affect the measured TBs at several AMSR2 channels. This effect is strongest at 89 GHz because of its higher sensitivity to water vapour and cloud liquid water, but it is also non-negligible at 36 GHz and, to a lesser extent, at 18 GHz. The water vapor influence is similar there, just slightly lower than at 89 GHz. These atmospheric contributions can introduce additional uncertainty into snow-depth retrievals, particularly for retrievals or GRs involving the higher-frequency channels.”*

## References

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1979 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2018.

Maeda, T., Taniguchi, Y., and Imaoka, K.: GCOM-W1 AMSR2 Level 1R Product: Dataset of Brightness Temperature Modified Using the Antenna Pattern Matching Technique, *IEEE T. Geosci. Remote Sens.*, 54, 770–782, doi:10.1109/TGRS.2015.2465170, 2016.

Picard, G., Löwe, H., and Mätzler, C.: Brief communication: A continuous formulation of microwave scattering from fresh snow to bubbly ice from first principles, *The Cryosphere*, 16, 3861–3866, <https://doi.org/10.5194/tc-16-3861-2022>, 2022.

Wentz, F. J. and Meissner, T.: AMSR ocean algorithm, Algorithm Theoretical Basis Document (ATBD), Version 2, CA, USA, 2000.

Ye, Y. and Heygster, G.: Arctic multiyear ice concentration retrieval from SSM/I data using the NASA Team algorithm with dynamic tie points, in: *Towards an Interdisciplinary Approach in Earth System Science*, Springer, Cham, 99–108, [https://doi.org/10.1007/978-3-319-13865-7\\_12](https://doi.org/10.1007/978-3-319-13865-7_12), 2015.

## Response to reviewer 2

Their study evaluates the retrievability of snow depth over Arctic sea ice for the AMSR2 frequency channels. I appreciate the effort of taking a more fundamental view on the question: instead of trying out different snow depth retrievals the authors use radiative transfer modeling to test sensitivities depending on background conditions.

The introduction is well written. The methods are clearly described but could benefit from a bit of additional information (see below) and more caution regarding the use of snow ice. The Data section describes the MOSAiC data but remains a bit too vague as to how the MOSAiC data is actually used later in the model setup. The Results section is well structured, describing first the results of the different snow setups and then the retrievable ranges, see some comments for improvement below. The conclusions provide nice guidance for further developments, highlighting also the importance of snow grain sizes. The figures are well done and I congratulate the authors for visualizing the information/coupling effects of such high dimensions, without oversimplifying it.

Reply: Thank you for your positive and valuable comments. We have added additional experiments and enhanced the clarity of the methodology section in the revised manuscript. Please see our response (blue text) below

### Reviewer comment:

How do the authors ensure representative sampling? From the method section I understand that the parameters are uniformly sampled, thus unrealistic input cases, that is unrealistic combinations of layer parameters or values that are rare, such as very deep snow, are possible and could influence the sensitivity. Is that right? Or is there a mechanism to ensure that the observed gradients of, e.g., temperature and density, are preserved or that the output is weighted according to the likelihood of occurrence? The use of the MOSAiC data here is not clear to me.

In our original setup, the normalised EFAST samples were mapped uniformly onto the prescribed physical ranges of each parameter.

To evaluate whether this sampling choice affects our main conclusions, we performed an additional sensitivity test using bounded Gaussian sampling. Specifically, for parameters with available MOSAiC-derived means and standard deviations, we

generated EFAST samples from bounded Gaussian distributions parameterised by these statistics (see Tab. R1), while retaining the parameter range constraints specified in the original study. This allows the samples to be more concentrated around the MOSAiC-observed mean states while remaining within the physically plausible intervals.


Tab. R1 Means and standard deviations (SDs) of snow parameters derived from MOSAiC measurements and used in an additional sensitivity test with bounded Gaussian sampling.

| Parameter | Depth | Density           | Temperature | Radius |
|-----------|-------|-------------------|-------------|--------|
| Unit      | cm    | kg/m <sup>3</sup> | K           | mm     |
| Mean      | 13.2  | 313.0             | 257.2       | 0.37   |
| SD        | 11.7  | 137.7             | 7.2         | 0.38   |

The results of this additional sensitivity test show no fundamental change in the main conclusions. The most sensitive parameter in each channel remains consistent with the original uniform-sampling sensitivity test, and the frequency-dependent coupling patterns among snow depth, density and grain radius are also preserved. In particular, the dominant role of snow microstructure at higher frequencies and the strong parameter-coupling behaviour identified in the original analysis remain evident under the MOSAiC-constrained Gaussian sampling.

The EFAST framework requires a consistent sampling treatment across all input parameters in the normalised parameter space. We did not adopt Gaussian sampling as the primary EFAST sampling strategy because reliable MOSAiC-based means and standard deviations are not available for all parameters. For example, MOSAiC does not provide summer observations of liquid water content in snow, which is a key parameter in the wet snow simulations. In addition, some summer snow states observed during MOSAiC, especially very high density and large grain wet snow, exceed the validity limits of the SMRT simulation. The MOSAiC observations were used to define the physically plausible parameter ranges. Although we calculated MOSAiC-based means and standard deviations and used them in the additional robustness experiment, we did not use them in the primary EFAST sampling because of the above limitations and because the primary goal was to examine the full bounded response space of the radiative transfer model.

(a) Sampling: uniform



(b) Sampling: Gaussian distribution




Fig. R1 EFAST sensitivity result of single-layer Arctic snow parameters under dry snow conditions. Solid segments indicate the SSI of each parameter, while dotted extensions show the coupling effects captured by the TSI. For each channel (column), the strongest single sensitivity and the total sensitivity are also given numerically.

We have added supplementary explanations in the manuscript (in Section 3.3, Page 10):  
*“Notably, the sampling strategy used in the global sensitivity analysis assumes that all parameters are uniformly distributed within their prescribed ranges, which may differ to some extent from the actual distributions of snow parameters. We therefore also considered an alternative sampling strategy based entirely on Gaussian distributions derived from the MOSAiC observations. A comparison of the sensitivity analysis results obtained using the two sampling strategies shows that they are largely consistent. The results of this comparison are provided in the Supplementary*

### *Material.”*

Another general remark related to that: I think the confidence in the modeling results/snow configuration could be increased by a direct comparison, possibly in the supplemental material, of the modelled brightness temperature distributions to pan-Arctic satellite-measured brightness temperature distributions (possibly corrected for atmosphere), to see how modelled and measured TB variabilities compare (with no perfect agreement expected).

Following the reviewer’s suggestion, a pan-Arctic comparison requires spatially continuous information on key snow microstructural properties, especially snow density and grain radius. These two parameters are critical for the SMRT simulations, as demonstrated by our sensitivity analysis, but they are currently difficult to obtain at the pan-Arctic scale with sufficient accuracy and consistency. Therefore, rather than performing a pan-Arctic comparison based on poorly constrained snow microphysical inputs, we conducted an additional TB simulation experiment based on the MOSAiC observations, where snow depth, density, temperature, and grain radius are directly measured.

In this additional analysis, we mainly used the single-layer snow configuration corresponding to Sect. 4.1.1 of the sensitivity analysis. Because direct measurements of snow liquid water content were not available in the MOSAiC dataset used here, we restricted this comparison to the winter period, when dry snow conditions are expected to dominate. Values for the snow depth, density, temperature and radius were taken from the MOSAiC measurements. For each snow observation site, profile measurements were averaged to obtain a representative value for the corresponding snow column.

To make the modelled TBs more comparable with AMSR2 observations, we also included sea ice and atmospheric datasets in the forward simulation. Specifically, the SMRT model was first used to simulate the microwave emission from the snow-covered sea ice surface. The simulated surface TBs were then propagated through the atmosphere using the Wentz atmospheric radiative transfer model (Wentz 2000).

Tab. R2 Snow, sea ice and atmosphere configurations used in the forward simulation

| Layer | Parameter | Setting or Source   |
|-------|-----------|---------------------|
| Snow  | Depth     | MOSAiC measurements |

|            |                              |                                                              |
|------------|------------------------------|--------------------------------------------------------------|
|            | Density                      |                                                              |
|            | Temperature                  |                                                              |
|            | Radius                       |                                                              |
|            | Stickiness                   | 0.15                                                         |
|            | Thickness                    |                                                              |
|            | Density                      |                                                              |
|            | Temperature                  | Same setting as that in the original                         |
|            | Radius                       | Manuscript Tab. 4                                            |
| Sea ice    | Brine volume fraction        |                                                              |
|            | Porosity                     |                                                              |
|            | Multi-year ice concentration | Dataset from University of Bremen<br>(Ye and Heygster, 2015) |
|            | Total water vapour           | Use ERA5 reanalysis                                          |
| Atmosphere | Total cloud liquid water     | (Hersbach et al, 2018)                                       |

Overall, this SMRT configuration can reproduce the general magnitude and temporal variability of the AMSR2 observed TB, particularly at the intermediate-frequency channels from 18 to 36 GHz (See Fig. R2). At the low-frequency channels (6 and 10 GHz), the simulated TBs are systematically lower than the AMSR2 observations. This discrepancy is likely related to the stronger sensitivity to sea ice background properties. The best agreement is found at 18, 23 and 36 GHz. At 89 GHz, both the observed and simulated TBs exhibit much stronger short-term variability. The model captures part of the observed fluctuation and the increase in late winter. Overall, the comparison supports the use of the adopted snow and sea ice configurations as a physically plausible basis for the sensitivity analysis.

Following your suggestion, we have included this analysis in the Supplementary Material.



Fig. R2 Time-series comparison of SMRT-simulated TBs and AMSR2-observed TBs during the MOSAiC winter period.

Section 2.1: I recommend that the authors follow the guidelines from the official SMRT documentation: <https://smrt.readthedocs.io/en/v1.5.1/publish.html> providing a table with additional information about the model setup (in the supplemental material) including the surface roughness assumptions, the version number of the model and the exact way the salty snow (snow ice in Table 2) is modelled (what permittivity formulation is used)

We have added a model setup table to the supplemental material (Also see Tab.R3).

Tab. R3 SMRT model setups

| Topic                                             | Snow                       | First year ice                                                                               |                                                     |
|---------------------------------------------------|----------------------------|----------------------------------------------------------------------------------------------|-----------------------------------------------------|
|                                                   |                            | (including saline interlayer and superimposed ice)                                           | Multi-year ice                                      |
| Scattering theory                                 |                            | Improved born approximation<br>(with dense snow correction)                                  |                                                     |
| Microstructure Representation                     |                            | Sticky hard spheres                                                                          |                                                     |
| Surface/interface scattering models and substrate |                            | Default flat/Fresnel interfaces were used;<br>a water substrate was added below the sea ice. |                                                     |
| Permittivity model                                | ice_permittivity_matzler87 | ice_permittivity_matzler06<br>brine_permittivity_stogryn85                                   | ice_permittivity_matzler06<br>polder_van_stat<br>en |
| Radiative transfer solver                         |                            | Discrete Ordinate Radiative Transfer solver                                                  |                                                     |
| SMRT version                                      |                            | 1.5.1                                                                                        |                                                     |

- L. 116 : is  $i$  going from 1 to  $N$  or from 1 to  $n$  (as suggested by line 113)?

The index  $i$  denotes the input variable index rather than the sample index. Therefore,  $i$  should range from 1 to  $n$ , where  $n$  is the number of input variables. We have corrected this mistake.

- Section 2.3: While the general description of the averaging kernel as a matrix is of course correct, you could add that is 1D in your case, am I correct?

Yes, the averaging kernel matrix is one-dimensional because snow depth is the only target parameter considered.

- Section 3.1., Line 190: how are these measurements used exactly? To constrain the parameter ranges? Or to sustain realistic parameter combinations?

The MOSAiC measurements were used mainly to define variation ranges for the input parameters in the EFAST sensitivity analysis. We have added an additional sensitivity test in the Supplementary Material. In this additional test, the parameter variations were defined using the means and standard deviations derived from the MOSAiC measurements (see the reply to General Comment 1).

- Line 201/Table 4: you write about BVF for temperatures warmer than  $-2^{\circ}\text{C}$ , but you only model 265 K ice. How did you come up with BVF of 5% in Table 4?

The sentence referring to Cox and Weeks (1983) (For cold ice) and Leppäranta and Manninen (1988) (for ice warmer than  $-2^{\circ}\text{C}$ ) describes how BVF was derived in the MOSAiC ice-core data set from measured salinity, temperature, and density. The BVF value of 5% in Table 4 was used as an idealised representative value for cold ice, rather than being calculated from the warm ice formulation from Leppäranta and Manninen (1988). To avoid ambiguity, we remove the description of “*Leppäranta and Manninen (1988) for ice warmer than  $-2^{\circ}\text{C}$* ” from the original sentence.

- Section 3.2.: I suggest to add AMSR2 footprint sizes

Agreed and added

- Section 3.3.: snow ice formation: I suggest a bit more caution with using this term in the context of Arctic sea ice. To my knowledge flooding is common in the Antarctic (the Maksym and Jeffries paper quoted in line 217 and line 239 is also about Antarctica, which should be clarified), but the occurrence in the Arctic is still not clear. Also, the Merkouriadi study quoted later in line 360 talks about potential and not observations at large scale, please formulate accordingly. It is not clear to me whether the use of snow ice is backed by the MOSAiC observations. That is not to say that there is no salty snow

(commonly observed e.g. on fast ice) but to be careful with the flooding hypothesis. Finally, the density values listed in Table 2 go up very high, and scattering formulations tend to break down at intermediate/high densities, see Picard, G., Löwe, H., and Mätzler, C.: Brief communication: A continuous formulation of microwave scattering from fresh snow to bubbly ice from first principles, *The Cryosphere*, 16, 3861–3866, <https://doi.org/10.5194/tc-16-3861-2022>, 2022. Please have a look whether any issues are observed with regard to that.

In the revised manuscript, we have replaced the term “**snow ice**” with “**saline interlayer**” throughout the manuscript, figures, tables, and Supplementary Material. We have revised the description of saline interlayer in Sect. 3.3 (Page 8) and now use more cautious wording.

*“We include an idealised saline intermediate layer near the snow–ice interface. This layer represents a possible dense and saline transition between the snowpack and the underlying sea ice. For simplicity, we refer to this idealised intermediate layer as saline interlayer in the present study. The formation of saline interlayer may involve several processes. In Arctic sea ice, enhanced basal snow salinity may also arise from upward brine migration from the ice surface into the basal snow (Domine et al. 2004; Nandan et al. 2017; Mallett et al. 2024). Therefore, we treat this layer as an idealised sensitivity scenario to examine the potential radiometric impact of a dense saline layer at the snow-ice interface. Given the limitations of the SHS model in simulating high-density snow (Picard et al. 2022), the saline interlayer is treated as a thin sea ice layer rather than a snow layer in the SMRT simulation. In this parameterisation, the saline interlayer is represented as an ice-rich medium with brine inclusions, rather than as a high-density snow layer composed of ice grains embedded in an air matrix.”*

During the design of our SMRT simulation, we were aware of the issue discussed by Picard et al. 2022, that existing snow microstructural assumptions and scattering simulation (Sticky hard spheres + Improved born approximation) may become unstable in the intermediate- to high-density regime. We took three precautions:

- 4) In the SMRT simulations, the saline interlayer is treated as a thin sea ice layer rather than a snow layer (See Tab. R3). The difference is in the physical parameterisation of the layer: when treated as sea ice, the dense saline layer is represented as an ice-rich medium with brine inclusions, rather than as a high-

density snow layer composed of ice grains embedded in an air matrix.

- 5) We limited the maximum density of snow layers (wind slab and depth hoar) to  $500 \text{ kg m}^{-3}$ . Although the MOSAiC observations indicate that locally higher snow densities may occur, a density of  $500 \text{ kg m}^{-3}$  already corresponds to an ice volume fraction of approximately 0.55, which is slightly above the upper practical limit for applying the IBA+SHS simulation (0.5 given in Picard et al. 2022). We checked the simulated TB over the density range used for the snow layers. The 3D TB figures (Fig.6 – 9 in manuscript) show smooth and continuous variations over  $200\text{--}500 \text{ kg m}^{-3}$ , without abrupt fluctuations or discontinuities. This gives us confidence that the IBA+SHS configuration behaves stably within the snow density range.
- 6) We used the dense snow correction available in SMRT (See Tab.R3). When the ice volume fraction exceeds 0.5, this option treats the medium as air inclusions embedded in an ice matrix rather than ice particles embedded in an air matrix. . This medium inversion approach is also mentioned by Picard et al. (2022) as a possible treatment for dense snow.

-Section 3.4/ Table 4: what is the source of e.g. the radius values and the BVF values?  
Is it based on literature or MOSAiC data?

The radius values were derived from MOSAiC. The BVF value is also based on MOSAiC ice-core data. BVF generally fluctuates around 5% (Fig. R4). Although BVF increases to about 10% at deeper depths, around 1.5–2 m, these deeper layers are expected to have limited influence on the passive microwave signal because the penetration depth at 6 GHz is below 1 m. Therefore, we used  $\text{BVF} = 5\%$ .




Fig. R4 Brine volume fraction measurements (First year ice) during MOSAiC

- Figure 1: You could consider adding a panel giving the number of observations per snow depth/horizontal bars.

We have revised Fig.1 (See Fig. R5).




Fig. R5 Revised Fig. 1 in the manuscript.

- Section 4, Line 307: why is the temperature not shown for wet snow? I would think it is the biggest contributor with wet snow being more similar to a black body?

In our wet snow sensitivity analysis, snow temperature was not included as a varying parameter because it was fixed at 273.15 K. This choice was based on the MOSAiC observations, which show that wet snow temperature varies only weakly during the

melt season, with a mean value of 272.4 K and a standard deviation of 1.9 K (See Fig. R5). However, the snow temperature is not allowed to be lower than 273.15 K when snow liquid water is simulated in the SMRT model. Therefore, we set the wet snow temperature to a fixed value of 273.15 K and did not include it in the sensitivity analysis.

- Line 383: “finite depth” please be more specific, I assume snow is always of finite depth

We thank the reviewer for pointing this out. What we intended to express is that, when the wet snow layer is not thick enough to completely attenuate the contribution from the underlying interface, reflections from the superimposed ice layer can still affect the simulated TBs. We have revised the sentence accordingly in Section 4.1.3, Page 17.

*“When the wet snow layer is thin, e.g. <1 cm, it may not fully attenuate radiation from the lower boundary, so, the reflections from the underlying superimposed ice and residual microstructural scattering can still contribute to the observed signal.”*

- Section 4.2: I could not follow on which simulation setup this section is based. Are these extra simulations? Is the single-layer snow used? But then, how can we get snow depths up to 1 m when the range was limited to 60 cm in the previous section? Is wet snow excluded? Please clarify.

Sect. 4.2 is based on additional simulations, rather than on the EFAST sample sets used in Sect. 4.1. Specifically, the analyses in Sects. 4.2.1 and 4.2.2 use an idealised dry single-layer snowpack over typical Arctic FYI. Snow temperature and stickiness are fixed at 260 K and 0.1, respectively, and only snow depth, density, and grain radius are varied. The snow depth range in Sect. 4.2 was intentionally extended to 0–1.0 m to diagnose the saturation behaviour of single-channel TBs and GRs and to define the retrievable envelope. Wet snow is excluded from Sect. 4.2. This choice was made because most existing passive microwave snow depth retrieval algorithms over sea ice have been developed and applied primarily for dry snow conditions. We have clarified this simulation setup in the revised manuscript, Section 4.2, Page 19:

*“This Section is based on additional simulations, rather than on the EFAST sample sets used for the global sensitivity analysis in Sect. 4.1. Specifically, the analyses in Sects. 4.2.1 and 4.2.2 use an idealised dry single-layer snowpack over typical Arctic FYI. Snow temperature and stickiness are fixed at 260 K and 0.1, respectively, and*

*only snow depth, density, and grain radius are varied. The snow depth range is intentionally extended to 0–1.0 m in Sect. 4.2 in order to diagnose the saturation behaviour of single-channel TBs and GRs and to define the retrievable range.”*

- Figure 6: I am surprised to see no sharper gradient between 0 cm snow and the rest, is 0 snow actually modelled or is there always a thin snow layer present?

Yes, we used a very thin snow layer (0.001m). This very thin layer was introduced to avoid an abrupt numerical transition between a snow-covered surface and a bare ice surface. It is negligible compared with the 0 – 1 m snow depth range in Sect. 4.2 and does not affect the diagnosed retrievable ranges.

- Line 35: “most weather and all-sky conditions”: redundant

Deleted

## References

Domine, F., Sparapani, R., Ianniello, A., and Beine, H. J.: The origin of sea salt in snow on Arctic sea ice and in coastal regions, *Atmos. Chem. Phys.*, 4, 2259–2271, <https://doi.org/10.5194/acp-4-2259-2004>, 2004.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1979 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2018.

Mallett, R., Nandan, V., Stroeve, J., Willatt, R., Saha, M., Yackel, J., Veysière, G., and Wilkinson, J.: Dye tracing of upward brine migration in snow, *Ann. Glaciol.*, 65, e26, <https://doi.org/10.1017/aog.2024.27>, 2024.

Nandan, V., Geldsetzer, T., Yackel, J., Mahmud, M., Scharien, R., Howell, S., King, J., Ricker, R., and Else, B.: Effect of snow salinity on CryoSat-2 Arctic first-year sea ice freeboard measurements, *Geophys. Res. Lett.*, 44, 10419–10426, <https://doi.org/10.1002/2017GL074506>, 2017.

Picard, G., Löwe, H., and Mätzler, C.: Brief communication: A continuous formulation of microwave scattering from fresh snow to bubbly ice from first principles, *The Cryosphere*, 16, 3861–3866, <https://doi.org/10.5194/tc-16-3861-2022>, 2022.

Wentz, F. J. and Meissner, T.: AMSR ocean algorithm, Algorithm Theoretical Basis Document (ATBD), Version 2, CA, USA, 2000.

Ye, Y. and Heygster, G.: Arctic multiyear ice concentration retrieval from SSM/I data using the NASA Team algorithm with dynamic tie points, in: *Towards an Interdisciplinary Approach in Earth System Science*, Springer, Cham, 99–108, [https://doi.org/10.1007/978-3-319-13865-7\\_12](https://doi.org/10.1007/978-3-319-13865-7_12), 2015.

## Response to reviewer 3

### Reviewer comment:

I find this manuscript to be well written and the methodology (linking SMRT with EFAST and an averaging kernel to quantify retrievability) to be well chosen for such study. The rigor of the analysis while working in a multi-dimensional parameter space is commendable. Overall, this manuscript delivers valuable insights into the complexity surrounding retrievals of snow depth on sea ice that is of great interest to the remote sensing community and has the potential to influence future snow depth algorithms. For this, I recommend this manuscript for publication in The Cryosphere following some revisions. In particular, I find that some specific details of the methodology require clarification and that there is little to no discussion placing the study into the context of existing retrieval algorithms or modelling studies, or discussion regarding the limitations of this study. Furthermore, I find the authors' discussion around "snow ice" in the Arctic to be questionable. Please see below my comments and suggestions. These are divided into overarching, general comments and line-specific comments.

Thank you for your recommendation and valuable comments, and we are pleased to hear it. We have added additional experiments and added solid discussions in the revised manuscript. Please see our response (blue text) below

### 1: SMRT Configuration and Surface/Interface Roughness

The SMRT model is undoubtedly suitable for this study. However, by design, it is a highly configurable model. As such, user choices during model setup can influence the output. For example, the authors chose to represent the snowpack microstructure as sticky hard spheres (SHS). Several recent studies instead represent the snowpack using an exponential microstructure (e.g., Murfitt et al., 2024; Meloche, Sandells et al., 2024; Wivell et al., 2023; Meloche, Royer et al., 2024) as defined by a correlation length which can be derived using snow and ice densities and SSA. The cited literature on L95, Vargel et al. (2020), notably compares different SMRT snow microstructure configurations in reproducing observed brightness temperatures (Tbs). They found that a combination of the IBA model with snow layers defined by an exponential microstructure more accurately reproduced the Tbs than IBA with a SHS snowpack.

Because this study focuses on relative Tb differences, the absolute observation accuracy

may be secondary. However, if Tb errors scale non-linearly, then these results may be a reflection of this specific model configuration. To ensure robustness, I recommend running a subset of the experiment using an exponential snowpack microstructure and checking consistency. This could be added to the supplementary material and briefly referred to in the main text.

We thank the reviewer for this important comment. To evaluate whether our conclusions depend on the use of the sticky hard sphere (SHS) microstructure model, we repeated the sensitivity analysis using the exponential microstructure model while retaining the same SMRT configurations, parameter ranges, and sampling strategy. Because our objective was to isolate the influence of the snow microstructure model, snow depth, density, temperature, and effective grain radius were retained as the input parameters in the sensitivity analysis. For the exponential model, the correlation length ( $l_{ex}$ ) was derived from the sampled radius ( $r$ ) through the corresponding conversion based on snow density ( $\rho_{snow}$ ):

$$SSA = \frac{3}{\rho_{ice}r}$$

$$l_{ex} = \frac{4(1 - \rho_{snow}/\rho_{ice})}{\rho_{ice}SSA}$$

where  $\rho_{ice}$  denotes the density of pure ice (910 kg m<sup>-3</sup>),  $SSA$  denotes specific surface area. This conversion ensures consistency between the parameters prescribed in the two microstructure models. Because the exponential model does not use the stickiness parameter, stickiness was fixed at 0.2 in the SHS model and was not included as a variable in the sensitivity analysis.

The comparison, presented in the Supplementary Material (Also see Fig.R1), shows that the principal frequency-dependent sensitivity patterns are generally preserved between the two microstructure models. At 6 and 10 GHz, snow density remains the dominant parameter for both polarisations. At higher frequencies, particularly at 36 and 89 GHz, the snow grain radius remains dominant. Its total sensitivity indices are 0.66–0.71 at 36 GHz for both models and 0.60–0.67 and 0.53–0.59 at 89 GHz for the SHS and exponential model, respectively. Snow temperature has a consistently small influence in both models, whereas snow depth generally makes a secondary contribution. At 18 GHz V-polarisation and 23 GHz H-polarisation, the dominant parameter changes from density in the SHS model to the radius in the exponential model. At 18 GHz H-polarisation, density remains dominant, although its

total sensitivity decreases from 0.92 to 0.76. At 23 GHz V-polarisation, the radius remains dominant, with total sensitivity of 0.59 and 0.66 for the SHS and exponential model, respectively.




Fig.R1 Comparison of EFAST sensitivity indices obtained using (a) the SHS and (b) the exponential microstructure models. The values above each channel indicate the largest total sensitivity index among the evaluated parameters.

Furthermore, the strong coupling effects associated with snow grain radius, snow depth, and snow density, which are a central focus of this study, are clearly evident under both microstructure models and exhibit highly consistent frequency- and polarisation-dependent patterns (Fig. R1). In both the SHS and exponential model, the coupling contributions of these parameters become particularly pronounced at the 18 – 89 GHz. This consistency further supports our conclusion that the TB response is controlled not only by the individual snow properties but also by their nonlinear interactions, and that this key finding is robust to the choice of snow microstructure model. We have therefore added this comparison to the Supplementary Material and briefly summarised the results in the new Discussion Section (Pages 27-28):

*“The principal conclusions of this study were obtained using the SHS*

*microstructure model. SMRT provides several alternative representations of snow microstructure, and the selected configuration may influence both the magnitude of the simulated TBs and their sensitivity to snow properties. In several recent studies, snow microstructure has instead been represented using the exponential model (Wivell et al. 2023; Meloche et al. 2024; Meloche et al. 2026; Murfitt et al. 2024). In contrast to the SHS model, which idealises the snow microstructure as spherical ice particles characterised by an effective radius and a stickiness parameter, the exponential model describes the spatial organisation of the ice and air phases using an exponentially decaying autocorrelation function characterised by a correlation length. It therefore does not explicitly assume spherical grains and does not include a parameter equivalent to SHS stickiness. Some studies indicate that the suitability of a microstructure model may depend on snow type, microwave frequency, and the electromagnetic theory employed. The SHS model can provide a physically meaningful description of snow as an assembly of clustered ice particles and has shown performance comparable to, or slightly better than, the exponential model for some weakly metamorphosed or rounded-grain snow (Löwe and Picard, 2015). However, in a direct evaluation using 119 Arctic and sub-Arctic snow pits, Vargel et al. (2020) found that the exponential model produced lower errors than the SHS model, reducing the RMSE by up to 24 % for Arctic snow.*

*To assess the dependence of our results on this modelling choice, we repeated the sensitivity analysis using the exponential microstructure model for single-layer dry snow. Snow depth, density, temperature, and grain radius were retained as the parameters, with the exponential correlation length derived from the sampled SHS radius through the conversion given by Debye et al. (1957) and Mätzler (2002). Stickiness was fixed at 0.2 in the SHS model because it has no equivalent in the exponential model. Details of the experimental design and complete results are provided in the Supplementary Material. The principal frequency-dependent sensitivity characteristics are generally preserved between the two configurations. Snow density dominates the TBs at 6 – 10 GHz, whereas the grain radius becomes increasingly important at the higher frequencies, particularly at 36 and 89 GHz. Differences occur mainly within the 18 – 23 GHz channels, where the dominant parameter and the magnitude of the sensitivity indices vary between the SHS and exponential model. Importantly, the strong coupling effects associated with snow*

*radius, depth, and density, which are a central focus of this study, are evident under both microstructure models and exhibit highly consistent frequency- and polarisation-dependent patterns.”*

Additionally, SMRT permits adding roughness to interfaces. Surface roughness is known to affect the Tb as observed by satellites (e.g., Stroeve et al., 2006; Lee et al., 2018); however, this is not mentioned in the manuscript. As there is no mention of this, I assume the study chose not to add any rough interfaces in the SMRT simulations, meaning the retrievability conclusions all assume ice and snow layers are flat. While I think the added complexity of accounting for surface roughness is likely outside of the scope of this paper, which focuses on snow microstructure, I have 2 recommendations based on this omission. 1) Discuss neglecting snow and ice roughness as a limitation of the method, and 2) Flatten the interfaces of the ice layers in Fig. 1 and Fig. 3., so that the reader is not misled into thinking rough interfaces were modelled (if indeed they were not).

Snow and ice roughness were not included in the SMRT simulations. We have therefore revised Figs.1 and 3 by replacing the wavy boundaries with flat interfaces.

Following the reviewer’s suggestion, we added a discussion to fulfil the reviewer’s recommendations in the new Discussion section of the revised manuscript (Pages 27-28):

*“An additional limitation of this study is that all snow and ice interfaces were treated as smooth boundaries. This simplification does not fully represent natural snow-covered sea ice, where roughness may occur between snow layers and at the snow-ice boundary. SMRT can explicitly represent rough interfaces, and this capability has been applied in several previous studies (Picard et al. 2018; Meloche et al. 2024; Meloche et al. 2026; Murfitt et al. 2024). These studies demonstrate that roughness can modify microwave reflection, transmission, and scattering, thereby affecting the magnitude and polarisation characteristics of the simulated microwave signal. Surface and interfacial roughness may consequently influence the retrieval of snow and sea ice properties. (Stroeve et al., 2006; Lee et al., 2018). However, explicitly accounting for roughness would require additional parameters and substantially expand the multi-dimensional parameter space, which is beyond the scope of the present study, which focuses primarily on the effects of snow microstructure. Future*

*work should incorporate surface and interfacial roughness into the SMRT framework to assess whether the sensitivity rankings and snow depth retrievability identified here remain robust under more realistic interface conditions.”*

## **2: “Snow Ice” or simply Saline Snow?**

The authors note relatively high salinity in the bottom ~10 cm of the snowpacks sampled in the MOSAiC dataset and deem this to be “snow ice” on the basis that salinity is assumed to be negligible for Arctic snow (L234). I disagree with this conclusion and the associated terminology. Firstly, I think it is generally accepted that snow at the base of the snowpack on FYI is often saline; one cause being the upward wicking of brine rejected to the ice surface during sea ice growth (e.g., Domine et al., 2004; Nandan et al., 2017; Mallett et al., 2024). Secondly, widespread snow ice formation due to flooding and subsequent refreezing of the snowpack is generally considered to be an Antarctic phenomenon due to the Antarctic’s higher precipitation amount and higher fraction of seasonal ice. Although snow ice formation has been observed in the Arctic, I do not believe it is known to be widespread. The references provided on L259 (Merkouriadi et al., 2017; Merkouriadi et al., 2020) state that their results are confined to the Atlantic sector of the Arctic which is heavily influenced by storms.

Additionally, snow ice in this manuscript is treated as a snow layer, but in its conventional definition, it is a solid layer of ice that is often indistinguishable from sea ice without isotopic analysis (e.g., Jeffries et al., 1997; Granskog et al., 2017; Arndt et al., 2021). To avoid confusion with the conventional definition, I recommend renaming this layer to “Saline Snow,” or similar, and removing the suggestion that the properties of the basal snow layers in these profiles were a product of flooding, unless there is clear evidence to support it.

We would like to clarify that the snow ice layer in our study is treated as ice rather than snow in the SMRT model. In the revised manuscript, we have replaced the term “**snow ice**” with “**saline interlayer**” throughout the manuscript, figures, tables, and Supplementary Material. We have revised the description in Sect. 3.3 to make this modelling choice explicit and to use more cautious wording regarding Arctic saline interlayer formation (in Section 3.3, Page 8).

*“We include an idealised saline intermediate layer near the snow–ice interface. This layer represents a possible dense and saline transition between the snowpack and the*

*underlying sea ice. For simplicity, we refer to this idealised intermediate layer as “saline interlayer” in the present study. The formation of saline interlayer may involve several processes. In Arctic sea ice, enhanced basal snow salinity may also arise from upward brine migration from the ice surface into the basal snow (Domine et al. 2004; Nandan et al. 2017; Mallett et al. 2024). Therefore, we treat this layer as an idealised sensitivity scenario to examine the potential radiometric impact of a dense saline layer at the snow-ice interface. Given the limitations of the SHS model in simulating high-density snow (Picard et al. 2022), the saline interlayer is treated as a thin sea ice layer rather than a snow layer in the SMRT simulation. In this parameterisation, the saline interlayer is represented as an ice-rich medium with brine inclusions, rather than as a high-density snow layer composed of ice grains embedded in an air matrix.”*

In the SMRT simulations, the saline interlayer is treated as a thin ice layer rather than a snow layer. The difference lies in the physical parameterisation of the layer: when treated as sea ice, the dense saline layer is represented as an ice-rich medium with brine inclusions, rather than as a high-density snow layer composed of ice grains embedded in an air matrix. To describe this more clearly, we have added a model setup table to the Supplemental Material

### **3: Limitations**

Although this study follows a thorough methodology, it is not without its limitations. However, the manuscript currently lacks a real discussion surrounding this. I strongly recommend adding a dedicated limitations paragraph or subsection.

For example, the MOSAiC dataset represents a spatially limited sample of Arctic floes due to the nature of the expedition being “locked-in” to the drifting ice. Additionally, the simulations are specific to observations at AMSR2 frequencies and 55-degree look angle. Furthermore, as mentioned in GC1, SMRT configuration choices can impact such simulations, and this study does not assess the quality of the absolute simulated Tb values. Perhaps another SMRT configuration (e.g., exponential snow microstructure) could be used to check for consistency, or the configuration could be evaluated by also simulating the Tbs observed by the ground-based radiometers on MOSAiC.

Finally, in the context of simulating AMSR2 data, it is important to note that observed Tbs do not come from a single, homogeneous snowpack. While the study investigates

different ice types within a footprint, it does not investigate the potential spatial heterogeneity of the snowpack within a footprint. Because this is a fundamental study, it is acceptable to assume a homogeneous snowpack; however, it should perhaps be noted as a limitation when drawing conclusions for practical remote sensing applications

We have added a new Discussion section to state the limitations of our study. As described in our response to GC1, we have added a discussion of the influence of SMRT configuration choices on the Discussion section. We have further expanded the Discussion section to the restricted spatial representativeness of the MOSAiC dataset and the omission of sub-footprint spatial heterogeneity in the simulated snow cover (Pages 27-28):

*“Although this study combines SMRT with EFAST to conduct sensitivity and snow depth retrievability analyses, several limitations should be considered when interpreting the results. First, the MOSAiC measurements represent a spatially and temporally restricted sample of Arctic snow and sea ice conditions rather than the full variability of the Arctic Ocean. Consequently, our results may not capture the full range of conditions across the Arctic, particularly in marginal ice zones, where the snow-covered sea ice is often thinner, more fragmented, and more spatially heterogeneous. However, for studies of microwave radiative transfer from snow-covered sea ice, the MOSAiC measurements remain the most suitable choice because it provides invaluable integrated observations of snow depth, snow microstructure, and sea ice conditions. Second, the simulations were performed specifically at the AMSR2 frequencies and a fixed incidence angle of 55°, and the resulting sensitivity patterns may have to be adapted to other sensor configurations. Finally, the simulations assume horizontally homogeneous snow and ice conditions. Although variations among sea ice types within a satellite footprint are considered, sub-footprint heterogeneity in snow depth, density, temperature, and microstructure is not explicitly represented. Because the microwave response to these properties may be nonlinear, additional footprint-scale simulations and observational validation may be required before directly applying these results to operational retrievals.”*

We have added an additional TB simulation experiment using MOSAiC measurements together with collocated AMSR2 TBs. In this additional analysis, we mainly used the single-layer snow configuration corresponding to Sect. 4.1.1 of the

sensitivity analysis. Because direct measurements of snow liquid water content were not available in the MOSAiC dataset used here, we restricted this comparison to the winter period, when dry snow conditions are expected to dominate. Values for snow depth, density, temperature and radius were taken from the MOSAiC measurements. For each snow observation site, profile measurements were averaged to obtain a representative value for the corresponding snow column.

To make the modelled TBs more comparable with AMSR2 observations, we also included sea ice and atmospheric datasets in the forward simulation (See Table R2). Specifically, the SMRT model was first used to simulate the microwave emission from the snow-covered sea ice surface. The simulated surface TBs were then propagated through the atmosphere using the Wentz atmospheric radiative transfer model (Wentz 2000).

Tab. R2 Snow, sea ice and atmosphere configurations used in the forward simulation

| Layer      | Parameter                    | Setting or Source                                            |
|------------|------------------------------|--------------------------------------------------------------|
| Snow       | Depth                        | MOSAiC measurements                                          |
|            | Density                      |                                                              |
|            | Temperature                  |                                                              |
|            | Radius                       |                                                              |
|            | Stickiness                   | 0.15                                                         |
| Sea ice    | Thickness                    | Same as that in the original<br>Manuscript Tab. 4            |
|            | Density                      |                                                              |
|            | Temperature                  |                                                              |
|            | Radius                       |                                                              |
|            | Brine volume fraction        | Dataset from University of Bremen<br>(Ye and Heygster, 2015) |
|            | Porosity                     |                                                              |
| Atmosphere | Multi-year ice concentration |                                                              |
|            | Total water vapour           | ERA5 reanalysis                                              |
|            | Total cloud liquid water     | (Hersbach et al, 2018)                                       |



Fig. R2 Time-series comparison of SMRT-simulated TBs and AMSR2-observed TBs during the MOSAiC winter period.

Overall, this SMRT configuration can reproduce the general magnitude and temporal variability of the AMSR2 observed TBs, particularly at the intermediate-frequency channels from 18 to 36 GHz (See Fig. R2). At the low-frequency channels (6 and 10 GHz), the simulated TBs are systematically lower than the AMSR2 observations. This discrepancy is likely related to the stronger sensitivity of these channels to sea ice background properties. The best agreement is found at 18, 23 and 36 GHz. At 89 GHz, both the observed and simulated TBs exhibit much stronger short-term variability. The model captures part of the observed fluctuation and the increase in late winter. Overall, the comparison supports the use of the adopted snow and sea ice configurations as a physically plausible basis for the sensitivity analysis. Following the reviewers, we have included this analysis in the Supplementary Material.

#### **4: Contextualisation of Study**

I think the reader would appreciate some more context surrounding existing snow depth retrieval algorithms in the introduction and why they are limited, i.e., that they are typically parametric equations or neural networks trained on airborne or ground-based springtime data (often biased to the Western Arctic, or even Antarctica), therefore representing a limited snapshot of the metamorphic state of the snowpack and making their year-round applicability uncertain. Furthermore, there should be some introduction and discussion regarding prior modelling works that have sought to provide some understanding of the influence of snow properties on the retrievability of snow depth. For example, Markus et al. (2006), Powell et al. (2006), and Rostosky et al. (2020).

We have revised the introduction to describe existing snow depth retrieval algorithms and their main limitations (in Section 1, Page 2):

*“Existing passive microwave snow depth retrieval algorithms rely mainly on empirical regressions or neural networks trained using airborne or ground-based observation collected primarily in spring, with many datasets concentrated in the western Arctic or derived from Antarctica (Markus and Cavalieri, 1998; Braakmann-Folgmann and Donlon 2019; Kilic et al., 2019; Rostosky et al. 2020; Rösel et al. 2021; He et al., 2024). The training datasets capture a limited range of regional metamorphic snow conditions, despite substantial seasonal and spatial variations in snow density and grain size. Consequently, snow depth retrievals may exhibit*

*seasonally and regionally varying biases when the actual snow microstructure differs from that represented in the training data or assumed by the retrieval algorithm (Yang et al., 2021; Gao et al., 2023; He et al., 2024).”*

Regarding the prior modelling works, we have added the following sentences to the Introduction (in Section 1, Pages 2-3): D

*“Previous studies have combined radiative transfer models with local sensitivity or Monte Carlo uncertainty analyses to quantify how uncertainties in the snow and ice properties propagate into snow depth retrieval (Markus et al. 2006; Powell et al. 2006; Rostosky et al. 2020).”*

## 5: Quoting Results

I find the results section to be well done and interesting. However, I do suggest integrating more specific quantitative values (e.g., of TSI and SSI values) into the text to support your points and save the reader from needing to continuously scroll back to the figures.

We have added specific TSI and SSI values to the relevant paragraphs where they directly support the main findings.

## Line-specific Comments:

L30-31: The snow not only restricts oceanic heat transfer through the ice but also insulates the ice and ocean from atmospheric heat fluxes, both of which influence the sea ice mass balance as per the following sentence.

Agreed and added, the new text reads (in Introduction, Page 1-2):

*“Its high albedo reduces the absorption of solar radiation, while its low thermal conductivity provides an insulating layer that suppresses heat exchange between the ocean and the atmosphere, thereby affecting both sea ice growth and melt”*

L36: I think “all-sky” here refers to day and night / regardless of the solar zenith angle. In which case, I do not think it should be hyphenated.

Agreed and removed.

L38-39: Penetration depth is itself a function of frequency-dependent absorption and scattering within the system and perhaps listing it individually is redundant.

Agreed and changed, The new text reads (in Introduction, Page 2): *“most notably the penetration, absorption, and scattering of radiation within the snowpack and the underlying sea ice”*

L43-45: Such variability does not inherently show that the algorithms are problematic, as snow depth is expected to vary in space and time. Consider rephrasing this sentence to highlight that biases caused by unaccounted for snow microstructural properties can lead to unrealistic variability, using appropriate references.

Done. The new text reads (in Introduction, Page 2):

*“As a result, unaccounted snow microstructural properties within snow retrieval algorithms can lead to large biases or unrealistic variability of snow depth (Yang et al., 2021; Gao et al., 2023; He et al., 2024).”*

L45: I think further context on the “limited application ranges of existing snow depth algorithms” would be appreciated here (in line with GC4); explaining why current algorithms are limited, e.g., perhaps due to their simplistic parametric form that was derived using airborne campaign data that is spatially and temporally limited, reflecting a metamorphic state of the snowpack on Arctic sea ice that is unrepresentative of the entire winter. Overall, there is little explanation of how current algorithms “work” or were derived in this paragraph.

We have revised this sentence in Introduction (Page 2):

*“Existing passive microwave snow depth retrieval algorithms rely mainly on empirical regressions or neural networks trained using airborne or ground-based observation collected primarily in spring, with many datasets concentrated in the western Arctic or derived from Antarctica (Markus and Cavalieri, 1998; Braakmann-Folgmann and Donlon 2019; Kilic et al., 2019; Rostosky et al. 2020; Rösel et al. 2021; He et al., 2024). The training datasets capture a limited range of regional metamorphic snow conditions, despite substantial seasonal and spatial variations in snow density and grain size. As a result, unaccounted snow microstructural properties within snow retrieval algorithms can lead to large biases or unrealistic variability of snow depth (Yang et al., 2021; Gao et al., 2023; He et al., 2024).”*

L64-65: Although this work may provide the most comprehensive study into the impact

of microstructural properties on the retrievability of snow depth on Arctic sea ice and may be the first to consider coupling effects into the analysis, I do not agree there has not been “similar work done before.” Others have also modelled Tb as a function of snow and ice microstructure to assess uncertainty in snow depth algorithms, and I think these should be noted in this manuscript. For example, Markus et al. (2006), Powell et al.,(2006), and Rostosky et al. (2020), per GC4.

We modified our description accordingly (in Section 1, Page 2):

*“Previous studies have combined radiative transfer models with local sensitivity or Monte Carlo uncertainty analyses to quantify how uncertainties in snow and ice properties propagate into snow depth retrieval (Markus et al. 2006; Powell et al. 2006; Rostosky et al. 2020).”*

L90-91: I am a bit unsure of the phrasing here regarding “coupled snow and sea ice stratification” configuration. Presumably this is referring to the experiment in which the fraction of each ice type is varied? However, it makes it sound like the single- and multi-layer snow configurations are not coupled to a sea ice substrate. Consider rephrasing.

Done, the new text reads (in Section 2.1):

*“In this study, we considered both single-layer and multi-layer snow over a representative first year ice. In addition, a representative multi-year ice was included to assess the influence of sea ice type on snow depth retrieval.”*

L93-95: Please see my GC1 regarding the SMRT setup and possible extension to ensure robustness using another model representation of snow microstructure.

We have added the corresponding experiment and a new Discussion section.

L193: There is a minor inconsistency in this list. The instrumentation used is listed for all parameters but salinity, in which the location the sample was measured (the lab) is listed. Listing the lab is redundant and it would be better to list the instrument (presumably salinometer?) to match the rest of the list.

We have modified the sentence accordingly. The new text reads (in Section 3.2, Page 8): “... .. and snow salinity profiles from snow pit samples measured using a salinity, conductivity and temperature sensor .”

L202/Section 3.1 and 3.2: Consider swapping the order of these two sections. The MOSAiC observation section flowing into Section 3.3, how it was used to create the layered snow scenarios, seems more natural.

[Agreed and done.](#)

L210: Perhaps reference Table 1 in Fuhrhop et al. (1998) directly to help guide the reader.

[Done.](#)

L216-217: While the description of snow ice formation is true, in accordance with GC2, it may not be relevant here.

[Please see our reply to GC2.](#)

L219: I recommend changing “freezes in the snow-ice surface” to “freezes at the snow-ice interface.”

[Accepted and done.](#)

Figure 2: Check punctuation in the panel descriptions (inconsistencies with colons and semicolons). Also, consider changing “the horizontal bars are the standard deviations” to “the horizontal bars represent the standard deviation”.

[Done.](#)

L221/Fig. 2 caption/306/419. British English “colour” is used, however the rest of the manuscript is consistent with American English (e.g., “utilizing”, “summarised”, “parameterized”). I recommend changing to the American English “color”.

[Thank you, we have applied British English throughout the entire manuscript.](#)

L233-239: Please refer to GC2.

[Please see reply to GC2.](#)

L244: The choice of parameter space is key to this study. However, it is not clear exactly

how the “physically plausible ranges” were derived. Was there a quantitative selection process, or were they selected based on the authors’ judgement? Furthermore, how were the MOSAiC measurements combined with the snow properties listed in Fuhrhop et al.(1998) to define the study’s ranges? Please add a sentence or two detailing the range selection process more clearly.

We have clarified this issue in the revised manuscript. We have added this explanation to the revised manuscript (in Section 3.3, Pages 9-10):

*“The parameter ranges were primarily constrained by the MOSAiC observations, while also accounting for the applicable range of the SMRT configuration and the ranges adopted in Fuhrhop et al. (1998). For example, the snow density upper limit was set to  $500 \text{ kg m}^{-3}$ , rather than including MOSAiC measurements reaching  $700\text{--}900 \text{ kg m}^{-3}$  (Figs.2b and 2c), because the performance of the IBA+SHS model deteriorates at high snow densities (Picard et al. 2022). This is also broadly consistent with the snow density upper range of  $546 \text{ kg m}^{-3}$  used by Fuhrhop et al. (1998). ”*

L283-286: The MYIF study is a great addition due to the known challenge of PM snow depth retrievals over MYI but the unknown nature of how this may couple with snow microstructure properties and the fraction of ice types within a footprint. However, please consider adding a brief 1-2 sentences on why the MYI bias is perceived to exist (i.e., scattering within the porous upper ice layer that mimics the scattering of a snowpack). Additionally, it should be stated what snowpack is assumed for the MYI here – I assume it is the same as the FYI dry snow single layer?

We analysed the influence of the fraction of MYI on snow depth retrieval in Section 4.2.3 using the same single-layer dry snow configurations as that used for FYI. We have added an explanation of why the MYI bias is thought to occur and clarified the snowpack configuration in the revised manuscript (in Section 3.4, Page 11):

*“The porous, air-rich upper layer of MYI can produce volume scattering similar to that of snow, thereby introducing a bias in snow-depth retrievals based on GR algorithms (Markus and Cavalieri, 1998; Rostosky et al., 2018).”*

In Section 4.2.3 (Page 24), we also added:

*“In this analysis, the same single-layer dry snow configuration was used over FYI and MYI, while the fraction of MYI was varied.”*

L302: I suggest changing the section header to “Results and Discussion.” Discussion points are interwoven into this section and there is no stand-alone discussion section.

We added a new “Discussion section” in the revised manuscript.

L359-361: See GC2 regarding Arctic snow ice and its occurrence.

Please see response to GC2.

Figure 4: What determines when TSI and SSI values are overlaid on the figure? There are clearly valid instances for a given channel, layer, and physical parameter that do not have a corresponding number set overlaid. Are TSI and SSI values over a certain threshold deemed to be of interest?

The TSI and SSI values were overlaid only for parameters with TSI values greater than 0.1. We have clarified this criterion in the revised Figs 4 and 5 captions.

L413: Why “based on V-pol TBs”? I assume this is because V-pol is the typical choice for existing snow depth algorithms, however the reader may not know this without context (see GC4).

The reviewer is correct, and we have clarified this in the revised manuscript (in Section 4.2, Page 18):

*“We evaluate the retrievable range of snow depth using V-pol TBs and V-pol GRs, because V-pol TBs are widely used in existing retrieval algorithms and exhibit greater sensitivity than H-pol TBs (Markus and Cavalieri, 1998; Rostosky et al., 2018; He et al. 2024).”*

L495: Consider adding an “e.g.,” before Markus and Cavalieri, 1998 as it is not the only study demonstrating this.

Done.

L496: Perhaps an opportunity to relate this result (GR(18/6) extending the snow depth retrievable range under large grain snow conditions) to the studies of Powell et al. (2006) and Rostosky et al., (2018).

Agreed and done.

L522-525: It is my understanding that the lower frequency is typically considered to be the ‘anchor’ for GR-based snow depth retrievals because it is conventionally regarded to be less sensitive to volume scattering within the snowpack. Check terminology.

We have revised the sentence and checked the terminology throughout the manuscript.

L525-528 and L566-568: The discussion surrounding replacing the 36 GHz (L525-529) and 18 GHz (L566-568) channel with 23 GHz for these applications is interesting. But it may be worth noting the ~23 GHz channel is seldom used for sea ice applications, I believe due to its strong sensitivity to atmospheric water vapour. If this channel were to be used operationally, it would likely require an atmospheric correction using a radiative transfer model. On this topic, I appreciate the supplementary material demonstrating that the higher frequency channel observations from AMSR2 can correlate better with in situ data following an atmospheric correction (depending on the atmospheric state) – perhaps this operational requirement could be reiterated in your conclusions?.

We have added more descriptions linking the result in the supplementary material to our conclusions (in Section 4.2.2, Page 24):

*“The potential benefits of channels at and above 23 GHz should, however, be interpreted with caution because observations at these frequencies are more susceptible to atmospheric effects. The supplementary experiments show that applying atmospheric correction using an atmospheric radiative transfer model generally improves the correlation between GRs incorporating high frequency channels and in-situ snow depth observations (See Supplementary Material). Nevertheless, the magnitude of this improvement depends on the atmospheric state, and further validation using a broader range of in-suit observation is required before these channel combinations can be applied operationally.”*

L543: Reiterating that Sect. 3.4. lacks clear information of the snowpack set up for this ice type experiment. Were both types considered to have the same snowpack here?

Yes, we used the same snowpack. We have added description in Section 4.2.3, Page 24:

*“In this analysis, the same single-layer dry snow configuration was used over FYI and MYI, while the fraction of MYI was varied.”*

L544-545: It may be worth noting that the spatial response of the AMSR2 antenna is not uniform. Because the antenna gain function is approximately Gaussian, a floe's actual contribution to the measured Tb depends on its relative position in the footprint. Therefore, a simple linear weighting is a simplification.

We have clarified this assumption in the revised manuscript and noted that a more realistic application would require convolution of the spatially distributed surface TBs with the AMSR2 antenna gain pattern in Section 4.2.3, Page 25:

*“This represents a first-order approximation because it assumes a uniform spatial response within the footprint. In practice, the AMSR2 antenna gain is approximately Gaussian, and the contribution of each ice type therefore also depends on its position within the footprint.”*

## References

Braakmann-Folgmann, A. and Donlon, C.: Estimating snow depth on Arctic sea ice using satellite microwave radiometry and a neural network, *The Cryosphere*, 13, 2421–2438, <https://doi.org/10.5194/tc-13-2421-2019>, 2019.

Debye, P., Anderson, J., and Brumberger, H.: Scattering by an Inhomogeneous Solid, II. The Correlation Function and Its Application, *J. Appl. Phys.*, 28, 679 – 683, <https://doi.org/10.1063/1.1722830>, 1957.

Domine, F., Sparapani, R., Ianniello, A., and Beine, H. J.: The origin of sea salt in snow on Arctic sea ice and in coastal regions, *Atmos. Chem. Phys.*, 4, 2259–2271, <https://doi.org/10.5194/acp-4-2259-2004>, 2004.

Fuhrhop, R., Grenfell, T. C., Heygster, G., Johnsen, K. P., Schlüssel, P., Schrader, M., and Simmer, C.: A combined radiative transfer model for sea ice, open ocean, and atmosphere, *Radio Science*, 33, 303–316, 1998, <https://doi.org/10.1029/97RS03020>.

Gao, S., Li, Z., Zhang, P., Chen, Q., Huang, L., Zhou, J., Zhao C., Qiao H. and Zheng, Z.: A novel global snow depth retrieval method considering snow metamorphism and forest influence, *Remote Sensing of Environment*, 295, 113712, 2023, <https://doi.org/10.1016/j.rse.2023.113712>.

He, L., Xue, B., Hui, F., Xu, S., Chen, Z., and Cheng, X.: Toward daily snow depth estimation on Arctic sea ice during the whole winter season from passive microwave radiometer data, *IEEE Transactions on Geoscience and Remote Sensing*, 62, 1–15, 2024, <https://doi.org/10.1109/TGRS.2024.3358340>.

Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 hourly data on single levels from 1979 to present, Copernicus Climate Change Service (C3S) Climate Data Store (CDS) [data set], <https://doi.org/10.24381/cds.adbb2d47>, 2018.

Kilic, L., Tonboe, R. T., Prigent, C., and Heygster, G.: Estimating the snow depth, the snow–ice interface temperature, and the effective temperature of Arctic sea ice using Advanced Microwave Scanning Radiometer 2 and ice mass balance buoy data, *The Cryosphere*, 13, 1283–1296, <https://doi.org/10.5194/tc-13-1283-2019>, 2019.

Lee, S., Sohn, B., and Shi, H.: Impact of ice surface and volume scatterings on the microwave sea ice apparent emissivity, *J. Geophys. Res.-Atmos.*, 123, 9220–9237, <https://doi.org/10.1029/2018JD028688>, 2018.

Löwe, H. and Picard, G.: Microwave scattering coefficient of snow in MEMLS and DMRT-ML revisited: the relevance of sticky hard spheres and tomography-based estimates of stickiness, *The Cryosphere*, 9, 2101 – 2117, <https://doi.org/10.5194/tc-9-2101-2015>, 2015.

Mallett, R., Nandan, V., Stroeve, J., Willatt, R., Saha, M., Yackel, J., Veysi re, G., and Wilkinson, J.: Dye tracing of upward brine migration in snow, *Ann. Glaciol.*, 65, e26, <https://doi.org/10.1017/aog.2024.27>, 2024.

Markus, T. and Cavalieri, D. J.: Snow depth distribution over sea ice in the Southern Ocean from satellite passive microwave data, in: *Antarctic Sea Ice: Physical Processes, Interactions and Variability*, 74, 19–39, 1998, <https://doi.org/10.1029/AR074p0019>

Markus, T., Powell, D. C., and Wang, J. R.: Sensitivity of passive microwave snow depth retrievals to weather effects and snow evolution, *IEEE T. Geosci. Remote*, 44, 68–77, <https://doi.org/10.1109/TGRS.2005.860208>, 2006.

Macfarlane, Amy R., Schneebeil, Martin, Dadic, Ruzica, Wagner, David N., Arndt, Stefanie, Clemens-Sewall, David, Hämmerle, Stefan, Hannula, Henna-Reetta, Jaggi, Matthias, Kolabutin, Nikolai, Krampe, Daniela, Lehning, Michael, Matero, Ilkka, Nicolaus, Marcel, Oggier, Marc, Pirazzini, Roberta, Polashenski, Chris, Raphael, Ian, Regnery, Julia, Shimanchuck, Egor, Smith, Madison M., and Tavri, Aikaterini: Snowpit raw data collected during the MOSAiC expedition, PANGAEA [data set], <https://doi.org/10.1594/PANGAEA.935934>, 2021.

Meloche, J., Royer, A., Roy, A., Langlois, A., and Picard, G.: Improvement of polar snow microwave brightness temperature simulations for dense wind slab and large grain, *IEEE T. Geosci. Remote*, 62, 1 – 10, 2024, <https://doi.org/10.1109/TGRS.2024.3428394>

Meloche, J., Sandells, M., Löwe, H., Rutter, N., Picard, G., Essery, R., Scharien, R. K., Langlois, A., Jaggi, M., King, J., Toose, P., Scagliola, M., Bouffard, J., Di Bella, A., and Haas, C.: Altimetric Ku-band Radar Observations of Snow on Sea Ice Simulated with SMRT, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2026-3615>, 2026.

Murfitt, J., Duguay, C., Picard, G., and Lemmetyinen, J.: Forward modelling of synthetic-aperture radar (SAR) backscatter during lake ice melt conditions using the Snow Microwave Radiative Transfer (SMRT) model, *The Cryosphere*, 18, 869–888, <https://doi.org/10.5194/tc-18-869-2024>, 2024.

Nandan, V., Geldsetzer, T., Yackel, J., Mahmud, M., Scharien, R., Howell, S., King, J., Ricker, R., and Else, B.: Effect of snow salinity on CryoSat-2 Arctic first-year sea ice freeboard measurements, *Geophys. Res. Lett.*, 44, 10419–10426, <https://doi.org/10.1002/2017GL074506>, 2017.

Picard, G., Sandells, M., and Löwe, H.: SMRT: An active–passive microwave radiative transfer model for snow with multiple microstructure and scattering formulations (v1.0), *Geoscientific Model Development*, 11, 2763–2788, 2018, <https://doi.org/10.5194/gmd-11-2763-2018>.

Powell, D. C., Markus, T., Cavalieri, D. J., Gasiewski, A. J., Klein, M., Maslanik, J. A., Stroeve, J. C., and Sturm, M.: Microwave signatures of snow on sea ice: Modeling, *IEEE T. Geosci. Remote*, 44, 3091–3102, <https://doi.org/10.1109/TGRS.2006.882139>, 2006.

Picard, G., Löwe, H., and Mätzler, C.: Brief communication: A continuous formulation of microwave scattering from fresh snow to bubbly ice from first principles, *The Cryosphere*, 16, 3861–3866, <https://doi.org/10.5194/tc-16-3861-2022>, 2022.

Rostosky, P., Spreen, G., Gerland, S., Huntemann, M., and Mech, M.: Modeling the microwave emission of snow on Arctic sea ice for estimating the uncertainty of satellite retrievals, *J. Geophys. Res.-Oceans*, 125, e2019JC015465, <https://doi.org/10.1029/2019JC015465>, 2020.

Rostosky, P., Spreen, G., Farrell, S. L., Frost, T., Heygster, G., and Melsheimer, C.: Snow depth retrieval on Arctic sea ice from passive microwave radiometers—Improvements and extensions to multiyear ice using lower frequencies, *Journal of Geophysical Research: Oceans*, 123, 7120–7138, 2018, <https://doi.org/10.1029/2018JC014028>.

Rösel, A., Farrell, S. L., Nandan, V., Richter-Menge, J., Spreen, G., Divine, D. V., Steer, A., Gallet, J.-C., and Gerland, S.: Implications of surface flooding on airborne estimates of snow depth on sea ice, *The Cryosphere*, 15, 2819–2833, <https://doi.org/10.5194/tc-15-2819-2021>, 2021.

Stroeve, J. C., Markus, T., Maslanik, J. A., Cavalieri, D. J., Gasiewski, A. J., Heinrichs, J. F., Holmgren, J., Perovich, D. K., and Sturm, M.: Impact of surface roughness on AMSR-E sea ice products, *IEEE T. Geosci. Remote*, 44, 3103–3117,

<https://doi.org/10.1109/TGRS.2006.880619>, 2006.

Vargel, C., Royer, A., St-Jean-Rondeau, O., Picard, G., Roy, A., Sasseville, V., & Langlois, A. (2020). Arctic and subarctic snow microstructure analysis for microwave brightness temperature simulations. *Remote Sensing of Environment*, 242, 111754. <https://doi.org/10.1016/j.rse.2020.111754>

Wentz, F. J. and Meissner, T.: AMSR ocean algorithm, Algorithm Theoretical Basis Document (ATBD), Version 2, CA, USA, 2000.

Wivell, K., Fox, S., Sandells, M., Harlow, C., Essery, R., and Rutter, N.: Evaluating Snow Microwave Radiative Transfer (SMRT) model emissivities with 89 to 243 GHz observations of Arctic tundra snow, *The Cryosphere*, 17, 4325–4341, 2023, <https://doi.org/10.5194/tc-17-4325-2023>

Yang, J. W., Jiang, L. M., Lemmetyinen, J., Pan, J. M., Luojus, K., and Takala, M.: Improving snow depth estimation by coupling HUT-optimized effective snow grain size parameters with the random forest approach, *Remote Sensing of Environment*, 264, 112630, 2021, <https://doi.org/10.1016/j.rse.2021.112630>.

Ye, Y. and Heygster, G.: Arctic multiyear ice concentration retrieval from SSM/I data using the NASA Team algorithm with dynamic tie points, in: *Towards an Interdisciplinary Approach in Earth System Science*, Springer, Cham, 99–108, [https://doi.org/10.1007/978-3-319-13865-7\\_12](https://doi.org/10.1007/978-3-319-13865-7_12), 2015.