

Title Page

Biochar addition enhances soil aggregate stability by modulating soil internal forces: The critical role of readily oxidized organic carbon and particulate organic carbon

Xiaoshuai Song^{a,b,c}, Chenyang Xu^{d,e}, Huanhuan Zhao^d, Jiangwen Li^{a,b,f}, Wei Du^{d,e}, Wuquan Ding^g, Zhenghong Yu^h, Fengbao Zhang^{a,b,f}, Feinan Hu^{a,b,f,*}

^a State Key Laboratory of Soil and Water Conservation and Desertification Control, the Research Center of Soil and Water Conservation and Ecological Environment, Chinese Academy of Sciences and Ministry of Education, Yangling, Shaanxi 712100, China;

^b Institute of Soil and Water Conservation, Chinese Academy of Sciences and Ministry of Water Resources, Yangling, Shaanxi 712100, China;

^c University of Chinese Academy of Sciences, Beijing 100049, China;

^d College of Natural Resources and Environment, Northwest A&F University, Yangling, Shaanxi 712100, China;

^e Key Laboratory of Plant Nutrition and Agri-environment in Northwest China, Ministry of Agriculture and Rural Affairs, Yangling, Shaanxi 712100, China;

^f College of Soil and Water Conservation Science and Engineering, Northwest A&F University, Yangling, Shaanxi 712100, China;

^g Chongqing Key Laboratory of Environmental Materials & Remediation Technologies, Chongqing University of Arts and Science, Chongqing 402168, China;

^h College of Desert Management, Inner Mongolia Agricultural University, Hohhot 010000, China.

***Corresponding author:** Feinan Hu, address: No. 26, Xi'nong Road, State Key

Laboratory of Soil and Water Conservation and Desertification Control, Northwest
A&F University, Yangling, Shaanxi 712100, China; email: hufn@nwafu.edu.cn;
telephone: +86-029-87011863; fax: +86-029-87010623.

Abstract: Degradation of soil structure and function is a significant obstacle to sustainable agriculture, yet using biochar is known to be an effective way of increasing soil organic carbon and boosting structure. Aggregate stability is greatly influenced by soil internal forces (SIFs), such as van der Waals, hydration, and electrostatic forces. However, it is still not clear which specific organic carbon components play a significant role in the stability of aggregates and the action of soil internal forces. In this study, a field simulation experiment was conducted using maize straw and wheat straw biochars to investigate how biochar addition affects soil aggregate stability. The findings showed that as the rate of biochar application increased from 0% to 10% over 180 days and 365 days, the contents of soil light fraction organic carbon (LFOC), particulate organic carbon (POC), dissolved organic carbon (DOC), readily oxidized organic carbon (ROC), as well as soil specific surface area and surface charge number, increased. The content of different carbon fractions decreased with prolonged incubation time, while soil electrochemical parameters continued to increase throughout the incubation period. Pearson correlation and redundancy analyses showed that soil active organic carbon fractions were strongly correlated with soil surface electrochemical properties, with ROC and POC being the primary factors. Soil aggregate stability increased with higher biochar application rates, and the beneficial effect of biochar on aggregate stability became increasingly pronounced over time. Based on the quantitative estimates of SIFs, the addition of biochar facilitated soil aggregates stability by increasing van der Waals attractive pressure and decreasing electrostatic repulsive pressure, and finally lowering net

repulsive pressure between soil particles. The soil aggregate breaking strength increased quickly at initially and then leveled out as the electrolyte content decreased; $10^{-2} \text{ mol L}^{-1}$ marked the tipping point. The study identified the active organic carbon (ROC and POC) fractions which could mediate the soil surface electrochemical properties and soil internal forces, finally enhance soil aggregate stability. Our research provides a theoretical basis for the targeted use and modifying of biochar to efficiently improve soil structure.

Keywords: Biochar; Active organic carbon fraction; Soil internal forces; Aggregate stability

1. Introduction

Soil aggregate formation and stabilization play a fundamental role in shaping soil architecture and underpin a wide range of soil functions (Batista et al., 2023; Chen et al., 2025). Physical characteristics like porosity, aeration, compaction, and water retention, as well as biochemical characteristics like organic carbon fixation, conversion, and microbial activity, are all impacted by aggregate stability (Liu et al., 2022; Tian et al., 2022; Hashimi et al., 2026). Furthermore, both intrinsic soil characteristics—such as particle composition, organic matter content, and clay mineralogy—and external factors—such as weathering, tillage methods, and erosion—have an impact on aggregate stability (Zhang et al., 2021; Siedt et al., 2021; Li et al., 2025). Therefore, improving soil aggregate stability is essential for reducing soil erodibility, improving soil health and sustaining agricultural productivity.

The application of biochar as a soil amendment is primarily attributed to its ability to boost organic carbon content and contribute to better soil functionality (Ji et al., 2024; Zhang et al., 2026a). Numerous studies have demonstrated that biochar have successfully raised soil organic carbon levels and improved soil aggregate stability (Sheng et al., 2023; Sun et al., 2021; Jia et al., 2024). However, Blair et al. (1995) concluded that anthropogenic management practices in agricultural systems primarily affect the reactive organic carbon fraction of soil organic carbon. Soil organic matter has been found to degrade slowly, making short-term changes in its content of limited practical significance. As an early and quick sign of ecosystem management techniques, soil active organic carbon, on the other hand, is less plentiful but more sensitive to environmental changes than soil organic matter (Stevenson, 1994; Wander et al., 1994).

According to Wei et al. (2014), soil active organic carbon represents the most dynamic fraction of soil organic carbon that has the greatest direct impact on plant nutrient supply; it is highly reactive, has a certain solubility in the soil, and is readily oxidized and mineralized by soil microorganisms. Light fraction organic carbon (LFOC), dissolved organic carbon (DOC), particulate organic carbon (POC), and readily oxidized organic carbon (ROC) are the primary forms of soil active organic carbon (Zhao et al., 2018; Yang et al., 2026). Despite making up a small fraction of soil organic carbon, soil active carbon has a direct role in soil biochemical transformation processes and can indicate slight variations in soil organic carbon before changes in total soil carbon (Coleman et al., 1983).

Accordingly, variations in the amount of organic carbon's active fraction in agricultural soils significantly impact on the soil's nutrient cycling and overall quality (Liang et al., 2011; Zhang et al., 2026b).

Overall, increased soil active organic carbon enhances soil aggregate stability (Bronick and Lal, 2005; Chen et al., 2026). It has been demonstrated that soil active organic carbon fractions can affect the nutrient content and bioavailability of the soil and have a significant and positive correlation with the formation and stability of soil macroaggregates (Wang et al., 2018). Aggregate stability was shown to increase positively with organic carbon content, mainly in the light organic matter component (Yagüe et al., 2016). In comparison to soils treated with nitrogen, phosphorus, and potassium (NPK) in combination with green manure or NPK alone, Li et al. (2012) found that soils treated with NPK and animal manure had significantly higher levels of biologically active soil organic carbon and aggregate stability. Microaggregates and silt + clay fractions showed a significant negative association with the mean weight diameter and geometric mean diameter of soil aggregates, while DOC, macroaggregates, macroaggregates-C, and microaggregates-C showed a significant positive correlation (Cao et al., 2024). In trials with simulated rainfall, soil aggregate stability rose when soil organic carbon and LFOC levels increased (Li et al., 2021). Although the role of active organic carbon in enhancing soil aggregate stability has been widely studied, the composition and characteristics of its various forms differ substantially, potentially leading to divergent effects on aggregate stability. Therefore, further research is needed to

elucidate how different fractions of soil active organic carbon influence the formation and stabilization of soil aggregates.

The stability of soil aggregates in water is mostly due to soil internal forces (SIFs), which include electrostatic, hydration, and van der Waals attraction (Ma et al., 2021b; Li et al., 2026). Theoretically, internal forces may reach hundreds of thousands of atmospheric pressures between two adjacent soil particles (Ding et al., 2019; Ma et al., 2022). The classical double layer theory states that the characteristics of the soil, including its specific surface area (SSA), surface charge density, and cation exchange capacity, as well as external environmental factors (moisture content, solution concentration, etc.), primarily determine the function of SIFs in mesoscale soils (Quirk, 1994; Li et al., 2013). The addition of organic matter modifies SIFs by altering the interfacial properties of soil particles. In a four-year field experiment, the use of weathered coal and biochar had a significant impact on the SSA and surface charge density of soil particles, as well as improving van der Waals attraction, decreasing net soil pressure, and decreasing aggregate fragmentation strength and splash erosion rate (Wang et al., 2024). Changes in the electrochemical characteristics of the soil surface were mostly caused by an increase in the amount of soil organic matter after vegetation restoration (Liu et al., 2020). Nevertheless, few research have examined how various active organic carbon fractions affect SIFs and aggregate stability by altering the surface properties of the soil particles when biochar is applied.

In this study, maize straw and wheat straw biochars were applied at different rates in a field simulation experiment to systematically evaluate their effects on soil

internal forces. Specifically, we aimed to (i) quantify the changes in SIFs under different biochar types and application rates, (ii) identify the active organic carbon fractions that most strongly regulate soil interfacial electrochemical properties and thereby influence SIFs, and (iii) elucidate the mechanisms and time effects by which biochar-mediated modifications of SIFs contribute to soil aggregate formation and stabilization. This work provides new mechanistic insights into biochar–soil interactions from the perspective of soil internal forces and offers a scientific basis for targeted soil structural improvement and sustainable agricultural management.

2. Materials and methods

2.1 Study area

This study was conducted in the Yangling Demonstration Zone, Shaanxi Province, China (108°08'47"E, 34°27'21"N). The region has a temperate continental monsoon semi-humid climate, with an average of 2167 hours of sunlight per year, temperatures between 11 and 13 °C, and 500 to 700 mm of rainfall. According to the FAO soil classification, the soil under study was Lou soil, which is categorized as calcareous Cambisol. The basic physicochemical properties of the soil are shown in Table 1.

Table 1 Basic physicochemical properties of the soil

	Bulk	Organic	Total	Available	Available
pH	density	carbon	nitrogen	phosphorus	potassium
	(g cm ³)	(g kg ⁻¹)	(g kg ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)
7.53	1.33	9.02	0.82	20.79	165.52

2.2 Experimental design

The biochar made from wheat and maize straw that was utilized in this

investigation was acquired from Zhengzhou Jinbang Environmental Protection Technology Co. An elemental analyzer (Elementar Vario EL III, Germany) was used to determine the elemental composition of the biochar samples. A fully automated specific surface and porosity analyzer (Micromeritics ASAP 2460, USA) was used to measure the SSA of biochar. A Mettler pH meter with a solid-liquid ratio of 1: 20 was used to measure the pH. The amount of ash was measured by placing the biochar in a crucible and burning it for two hours at 800 °C. Table 2 displays the fundamental physical and chemical characteristics of biochar.

Table 2 Basic properties of biochar

Biochar	pH	C (%)	H (%)	O (%)	N (%)	H/C	O/C	SSA (m ² g ⁻¹)	Ash (%)
Maize	6.30	32.67	0.92	11.72	0.48	0.03	0.36	14.48	7.51
Wheat	6.08	76.10	2.18	24.21	0.67	0.03	0.32	15.17	3.52

In mid-December 2021, biochar was added to the top layer of soil (0–20 cm) and tilled lightly until it was incorporated into the soil. In a control treatment (CK), no biochar was used. Two varieties of biochar, manufactured from maize and wheat straw, were treated at five different dose levels: 1%, 3%, 5%, and 10% by mass (designated M1, M3, M5, and M10 for maize straw biochar; W1, W3, W5, and W10 for wheat straw biochar). There were 27 plots in all, each measuring 1.5 m by 1.5 m, and a completely randomized design with three replications was employed (Figure 1). At the end of the 180-day and 365-day experimental periods, five soil auger samples were collected from the 0–20 cm layer of each plot and combined to form a composite soil sample. The samples were then allowed to dry naturally and were ready for use after being cleared of stones, plant roots, and other contaminants. Before determining

the soil properties, the unbound biochar particles in the soil samples were floated off in distilled water.




Fig. 1. Experimental plot design. Note: W is wheat straw biochar and M is maize straw biochar; 0, 1, 3, 5, and 10 represent 0%, 1%, 3%, 5%, and 10% biochar additions, respectively; A, B, and C represent three replicates, respectively.

2.3 Determination of soil active organic carbon and electrochemical properties

A carbon and nitrogen analyzer (Multi N/C 3000, Jena, Germany) was used to measure DOC (Qiu et al., 2023). By using potassium permanganate for oxidation, ROC was determined (Zhang et al., 2022). Sulfuric acid-potassium dichromate oxidation with external heating and sodium iodide heavy liquid separation were used to calculate LFOC (Janzen et al., 1992). Sulfuric acid-potassium dichromate oxidation using an external heating technique and sodium hexametaphosphate dispersion were used to determine POC (Ma et al., 2021a).

The combined method of surface property determination provided by Li et al. (2011) was used to measure the soil surface charge number (SCN), surface charge density (σ_0), and SSA. In short, soil samples were subjected to exchange equilibrium testing using a combination of NaOH and $\text{Ca}(\text{OH})_2$ after being pre-saturated with H^+ . After measuring the amounts of Na^+ and Ca^{2+} in the supernatant, the double-layer

theory was used to compute the cation exchange capacity, SSA, and σ_0 ; the detailed procedures can be found in our previous work (Li et al., 2026).

2.4 Quantification of soil internal forces

At the mesoscopic level, the primary SIF interactions among soil particles are van der Waals attractive pressure (P_{vdw}), surface hydration repulsive pressure (P_{h}), and electrostatic repulsive pressure (P_{E}). The net inter-particle force (P_{net}) is the combined force of the three, which can be calculated from the soil surface property parameter to obtain the SIFs (Liu et al., 2021; Li and Xu, 2008).

2.5 Soil aggregate stability evaluation

In this experiment, soils were prepared as Na^+ -saturated samples to quantitatively characterize the intrinsic link between SIFs and aggregate stability. Soil aggregate breaking strength (SABS) was used to evaluate the aggregate stability (Xu et al., 2015). It was defined as the proportion of mass released as small particles (less than 5 and less than 10 μm) when macroaggregates broke down. As the percentage mass release increases, soil aggregate stability decreases. The following was the preparation procedure: A 5 L beaker containing 2 kg of air-dried soil was filled with 5 L of NaCl electrolyte solution at a concentration of 0.5 mol L^{-1} . The sample was centrifuged and the supernatant was disposed of after 24 hours of stirring and equilibration. Following three repetitions of the aforementioned procedure, the extra Na^+ was removed using distilled water and dried through a 1–5 mm sieve. Different concentrations of NaCl solutions (10^{-5} , 10^{-3} , 10^{-2} , 10^{-1} , 1 mol L^{-1}) were used in the experiment to simulate soil electrolyte solutions. The time required for <

10 μm and $< 5 \mu\text{m}$ soil particles to fall at a specific distance was first calculated according to Stokes formula, respectively. About 450 ml of a NaCl solution with concentrations of 1, 10^{-1} , 10^{-2} , 10^{-3} , and $10^{-5} \text{ mol L}^{-1}$ was put into a 500 ml measuring cylinder. Weigh 15 g of Na^+ -saturated aggregates and transfer them into a measuring cylinder. Add the same solution to bring the total volume to 500 mL, then seal the cylinder. Allow it to stand for 1 minute. Afterward, hold the cylinder horizontally in your hand and rotate it slowly, inverting it once and allowing it to stand for approximately 30 seconds. Repeat this process four times. Finally, place the cylinder on the tabletop, record the start time, and calculate the end time. A U-shaped tube was used to aspirate the suspension at the conclusion of the allotted period, and it was then moved into an aluminum box. The sample was then weighed after drying at 105°C , following the method of Hu et al. (2015).

2.6 Statistical analysis

Microsoft Excel was used to arrange and compute the data, while SPSS Statistics 26 was used to conduct the statistical analysis. Statistical significance was determined using Duncan's test ($P < 0.05$). The relationship between soil surface electrochemical characteristics and soil active carbon fractions was assessed using the Pearson's coefficient test. Soil carbon fractions and electrochemical properties were analyzed redundantly using Canoco 5.0. Origin 2023 software was applied for data plotting.

3. Results

3.1 Effect of biochar addition on soil active organic carbon fractions

After applying maize straw biochar and wheat straw biochar after 180 days, the

overall amount of soil active organic carbon rose as the amount of biochar increased (Figure 2). Both biochar treatments significantly increased soil LFOC concentrations ($P < 0.05$), although slight, non-significant decreases were observed in the M5 and W5 treatments compared to M3 and W3. The M10 and W10 treatments showed the highest increase, 147.9 and 127.0 times compared to CK, respectively. The LFOC under each treatment for the two kinds of straw biochar was only substantially different at the 10% addition rate; at the 3% and 5% biochar application rates, the difference was not significant (Figure 2a). Both biochar treatments significantly enhanced soil POC content ($P < 0.05$). In terms of magnitude, M10 and W10 exhibited the largest increases, with POC contents 84.2 and 78.4 times higher than that of the CK, respectively. With the exception of the 5% application rate (Figure 2b), no appreciable variations in the soil POC concentration were noted in relation to the impacts of the two straw biochar treatments at the same application rate. The M10 and W10 treatments increased DOC by 18.6 and 18.1 times, respectively, compared to CK. At the same application rate, there were no discernible variations in DOC between the two biochars (Figure 2c). The M10 and W10 treatments increased ROC by 5.1 and 5.3 times, respectively, compared to CK. At the same application rate, there were no discernible variations in ROC between the two biochars (Figure 2d). The contents of different carbon fractions under each treatment decreased with the extension of incubation time, and the data for soil active carbon fractions at 365 days are provided in Supplementary Figure S1.




Fig. 2. Effect of biochar addition on soil active organic carbon fractions after 180 days. Note: M1, M3, M5, M10, add 1%, 3%, 5%, 10% maize straw biochar; W1, W3, W5, W10, add 1%, 3%, 5%, 10% wheat straw biochar.

3.2 Effect of biochar addition on soil surface electrochemical properties

Table 3 shows that as the amount of biochar rose from 0% to 10%, SCN increased considerably ($P < 0.05$). Compared with CK, SCN increased by 23.3%, 33.3%, 39.8%, and 42.7% under each treatment of maize straw biochar, and 27.1%, 33.3%, 39.4%, and 46.4% under the wheat straw biochar treatment, respectively. The difference between the two biochars at the same addition level was not significant. The soil SSA increased significantly ($P < 0.05$) with increasing biochar application rates; the M1 treatment and CK did not differ significantly, and at 1%, 3%, and 5% application rates, the differences between the treatments were not significant, and

M10 and W10 increased by 91.06% and 91.53%, respectively, in comparison to the CK treatment. Surface charge density σ_0 showed no significant change at lower application rates (1%, 3%, 5%) compared to CK. However, σ_0 tended to decline as biochar application rate increased. After 365 days of incubation with biochar, the trends of soil electrochemical parameters were consistent with those observed at 180 days, and the values of these parameters were slightly higher than those at 180 days. The data for soil surface electrochemical properties at 365 days are provided in Supplementary Table S1.

Table 3 Soil electrochemical properties under biochar treatment for 180 days

Treatment	SCN (cmol(c) kg ⁻¹)	SSA (m ² g ⁻¹)	σ_0 (c m ⁻²)
CK	13.12 ± 0.36e	32.23 ± 2.42c	0.39 ± 0.04a
M1	16.18 ± 0.72d	40.63 ± 7.67bc	0.39 ± 0.08a
M3	17.50 ± 0.14c	45.69 ± 3.56b	0.37 ± 0.03abc
M5	18.35 ± 0.08b	49.94 ± 0.57b	0.35 ± 0.01abc
M10	18.72 ± 0.03ab	61.58 ± 7.24a	0.30 ± 0.03c
W1	16.67 ± 0.36d	42.26 ± 7.04b	0.39 ± 0.07ab
W3	17.50 ± 0.09c	45.70 ± 3.39b	0.37 ± 0.03abc
W5	18.29 ± 0.40b	47.85 ± 5.43b	0.37 ± 0.05abc
W10	19.21 ± 0.16a	61.73 ± 7.40a	0.30 ± 0.04bc

Note: M1, M3, M5, M10, add 1%, 3%, 5%, 10% maize straw biochar; W1, W3, W5, W10, add 1%, 3%, 5%, 10% wheat straw biochar; CEC: cation exchange capacity, SSA: specific surface area, σ_0 : surface charge density; Values are means ± standard deviation (n = 3).

3.3 Relationship between soil electrochemical properties and active carbon fractions

After 180 days of biochar application, the correlation between soil

electrochemical properties and active organic carbon fractions is presented in Figure 3a. SCN exhibited a strong positive association with LFOC and POC ($P < 0.05$) and a highly significant positive correlation with DOC and ROC ($P < 0.01$). SSA and DOC ($P < 0.01$), LFOC, POC, and ROC ($P < 0.001$) all showed a substantial positive connection. In contrast, σ_0 was strongly negatively correlated with LFOC, POC, and ROC ($P < 0.001$), and with DOC ($P < 0.01$). The results at 365 days (Fig. 3b) showed that SCN and SSA were significantly and positively correlated with LFOC, POC, DOC, and ROC ($P < 0.001$). In contrast, σ_0 was negatively correlated with LFOC, POC, and ROC ($P < 0.05$), and showed a highly significant negative correlation with DOC ($P < 0.001$).




Fig. 3. Pearson correlation analysis between soil active carbon fractions and soil electrochemical properties after 180 and 365 days. Note: * indicates significant correlation at the 0.05 level; ** indicates significant correlation at the 0.01 level; and *** indicates significant correlation at the 0.001 level.

Figure 4 displays the results of redundancy analysis. The results showed that after 180 days of biochar addition, ROC explained 63.6% of the variation in soil electrochemical properties, whereas after 365 days, POC accounted for 66.1% of the

variation (Table 4). Therefore, ROC and POC are the primary factors influencing the electrochemical properties of the soil surface. The acute angle formed between the SCN, SSA, and soil active carbon component arrows indicates a strong positive correlation between them (the smaller the acute angle, the stronger the relationship), consistent with the results in Figure 3.




Fig. 4. Redundancy analysis of soil active carbon fractions and soil electrochemical properties after 180 and 365 days.

Table 4 Redundancy analysis of soil surface electrochemical properties and active carbon fractions after 180 and 365 days.

	Soil active carbon fractions	Explanation rate (%)	<i>F</i>	<i>P</i>
180 d	ROC	63.6	43.7	0.002
	DOC	1.7	1.2	0.322
	LFOC	1.5	1	0.37
	POC	1.7	1.2	0.348
365 d	POC	66.1	48.7	0.002
	DOC	0.5	0.4	0.614

LFOC	0.4	0.3	0.762
ROC	0.1	0.1	0.878

Note: ROC: readily oxidized organic carbon; DOC: dissolved organic carbon; LFOC: Light fraction organic carbon; POC: particulate organic carbon.

3.4 Effect of biochar addition on soil internal forces

3.4.1 Electrostatic repulsive pressure between soil particles

The electrostatic repulsive pressure at various electrolyte concentrations at a distance of 2 nm between soil particles is depicted in Figure 5. The findings demonstrated that as the amount of maize straw and wheat straw biochar applied rose from 1% to 10%, the electrostatic repulsive pressure gradually reduced at the same electrolyte concentration. At a constant biochar application rate, electrostatic repulsive pressure increased sharply under all treatments as the electrolyte concentration decreased progressively from 1 mol L⁻¹ to 10⁻² mol L⁻¹. However, when the electrolyte concentration is further decreased below 10⁻² mol L⁻¹, the rate of growth of electrostatic repulsive pressure slows down and stabilizes. After 365 days of incubation, the trends in electrostatic repulsive pressure for each treatment were consistent with those at 180 days, and increased compared to the soil after 180 days of incubation.




Fig. 5. Electrostatic repulsive pressure between soil particles at an interparticle distance of 2 nm under different electrolyte concentrations after 180 and 365 days.

Note: M1, M3, M5, M10, add 1%, 3%, 5%, 10% maize straw biochar; W1, W3, W5, W10, add 1%, 3%, 5%, 10% wheat straw biochar.

3.4.2 Van der Waals attractive pressure and surface hydration repulsive pressure

Table 5 shows that all of the Hamaker constant values were higher than CK and rose as the amount of biochar applied increased. The strength of the van der Waals attractive pressure between soil particles is directly determined by the Hamaker constant, a crucial element in the computation of this pressure. The amount and distribution of the van der Waals attractive pressure and surface hydration repulsive pressure between particles with biochar addition after 180 and 365 days (as seen in Figure 6) may be determined using the formula, in conjunction with the acquired

Hamaker constant. The van der Waals attractive pressure (absolute value) and the surface hydration repulsive pressure both sharply decline with increasing soil particle distance. When the spacing between soil particles was smaller than 1.3 nm, the surface hydration repulsive pressure was substantially larger in all treatments than the van der Waals attractive pressure. For instance, under various treatments, the interparticle van der Waals attractive pressure (in absolute value) is less than 100 atm and the interparticle surface hydration repulsive pressure is around 591 atm when the interparticle distance is 0.7 nm. The van der Waals attractive pressure (absolute values) is all below 200 atm, but the surface hydration repulsive pressure is around 1051 atm at an interparticle distance of 0.6 nm. This suggests that the difference between the two rises quickly as the distance decreases. Van der Waals attractive pressure (absolute value) rose with increasing application rate in both biochar treatments. After 365 days of incubation, the trends in the Hamaker constant and van der Waals attraction (in absolute value) for each treatment were consistent with those at 180 days, and increased compared to the soil after 180 days of incubation.

Table 5 Changes in soil Hamaker constant under different treatments after 180 and 365 days.

Treatment	Hamaker (J) 180 d	R^2	Hamaker (J) 365 d	R^2
CK	$4.78 \times 10^{-20} \pm 5.91 \times 10^{-21}$	0.82	$4.79 \times 10^{-20} \pm 4.63 \times 10^{-21}$	0.90
M1	$6.01 \times 10^{-20} \pm 6.05 \times 10^{-21}$	0.82	$8.89 \times 10^{-20} \pm 7.27 \times 10^{-21}$	0.85
M3	$6.08 \times 10^{-20} \pm 7.99 \times 10^{-21}$	0.73	$9.47 \times 10^{-20} \pm 8.29 \times 10^{-21}$	0.83
M5	$6.06 \times 10^{-20} \pm 7.71 \times 10^{-21}$	0.78	$9.78 \times 10^{-20} \pm 6.28 \times 10^{-21}$	0.91
M10	$6.15 \times 10^{-20} \pm 6.90 \times 10^{-21}$	0.78	$10.60 \times 10^{-20} \pm 3.44 \times 10^{-21}$	0.98
W1	$4.96 \times 10^{-20} \pm 5.95 \times 10^{-21}$	0.81	$7.38 \times 10^{-20} \pm 7.57 \times 10^{-21}$	0.75

W3	$5.02 \times 10^{-20} \pm 6.67 \times 10^{-21}$	0.75	$8.39 \times 10^{-20} \pm 7.80 \times 10^{-21}$	0.79
W5	$5.19 \times 10^{-20} \pm 6.08 \times 10^{-21}$	0.82	$8.73 \times 10^{-20} \pm 3.78 \times 10^{-21}$	0.95
W10	$5.22 \times 10^{-20} \pm 5.55 \times 10^{-21}$	0.84	$8.85 \times 10^{-20} \pm 2.69 \times 10^{-21}$	0.98

Note: M1, M3, M5, M10, add 1%, 3%, 5%, 10% maize straw biochar; W1, W3, W5, W10, add 1%, 3%, 5%, 10% wheat straw biochar; Values are means $\pm$ standard deviation (n = 3).




Fig. 6. Distributions of van der Waals attractive pressure and surface hydration repulsive pressure under biochar addition after 180 and 365 days. Note: M1, M3, M5, M10, add 1%, 3%, 5%, 10% maize straw biochar; W1, W3, W5, W10, add 1%, 3%, 5%, 10% wheat straw biochar.

3.4.3 Net pressure between soil particles

Figure 7 shows the net pressure of soil particles at 2 nm for different electrolyte concentrations under biochar addition after 180 and 365 days. Under both biochar

types, net pressure increased as electrolyte concentration decreased. From 1 mol L^{-1} to $10^{-2} \text{ mol L}^{-1}$, net pressure rose markedly; below $10^{-2} \text{ mol L}^{-1}$, the rate of increase plateaued. At the same electrolyte concentration, net pressure decreased as biochar application increased. After 365 days of incubation, the trends in the net interparticle force were consistent with those at 180 days, and decreased compared to the soil after 180 days of incubation.




Fig. 7. Net pressure of soil particles at 2 nm for different electrolyte concentrations under biochar addition after 180 and 365 days. Note: M1, M3, M5, M10, add 1%, 3%, 5%, 10% maize straw biochar; W1, W3, W5, W10, add 1%, 3%, 5%, 10% wheat straw biochar.

3.5 Effect of biochar addition on soil aggregate stability

Figure 8 shows SABS ($< 5 \mu\text{m}$, $< 10 \mu\text{m}$) at different electrolyte concentrations

under biochar addition after 180 days. As the electrolyte concentration increased, the SABS decreased. The change in SABS was relatively smooth when the electrolyte concentration was less than 10^{-2} mol L⁻¹, but it significantly decreased for all treatments when the electrolyte concentration went from 10^{-2} to 1 mol L⁻¹. A concentration of 10^{-2} mol L⁻¹ represents the critical threshold for aggregate disintegration. SABS dropped as biochar application increased under the same electrolyte concentration, suggesting that biochar application improved aggregate stability. After 365 days of incubation, the trends in aggregate breaking strength for each treatment were consistent with those at 180 days, and the values decreased compared to the soil after 180 days of incubation. The data for soil aggregate breaking strength at 365 days are provided in Supplementary Figure S2.




Fig. 8. Soil aggregate breaking strength (< 5 μ m, < 10 μ m) at different electrolyte

concentrations under biochar addition after 180 days. Note: M1, M3, M5, M10, add 1%, 3%, 5%, 10% maize straw biochar; W1, W3, W5, W10, add 1%, 3%, 5%, 10% wheat straw biochar.

3.6 Relationship between soil internal forces and aggregate stability

Figure 9 depicts the relationship between the net pressure at a 2 nm distance between soil particles and SABS under biochar addition after 180 and 365 days. The figure reveals a significant positive correlation between SABS and net pressure across all treatments, with SABS exhibiting an exponential increase in response to rising net pressure.




Fig. 9. Relationships between net interparticle pressure at a separation distance of 2 nm and soil aggregate breaking strength (SABS) for the < 5 μm and < 10 μm particle fractions under biochar addition after 180 and 365 days. Note: Panels (a) and (b)

represent maize straw biochar and wheat straw biochar treatments after 180 days, respectively, while panels (c) and (d) represent the corresponding treatments after 365 days. M1, M3, M5, and M10 indicate the addition of 1%, 3%, 5%, and 10% maize straw biochar, respectively; W1, W3, W5, and W10 indicate the addition of 1%, 3%, 5%, and 10% wheat straw biochar, respectively. Blue and black lines indicate the fitting curves for the $< 5 \mu\text{m}$ and $< 10 \mu\text{m}$ particle fractions, respectively.

4. Discussion

4.1 Response of soil active carbon fraction and electrochemical properties to biochar application

The application of biochar enhanced soil active carbon fractions and had a considerable impact on soil organic carbon (Li et al., 2024; Han et al., 2020; Lu et al., 2021). According to our findings, the soil active carbon percentage rose when both types of biochar were added (Figure 2 and Figure S1). The reason can be explained: the biochar itself contains more carbon, especially more resistant humic acid and condensed aromatic carbon structures. Secondly, biochar possesses a large surface area, which may facilitate the accumulation of specific organic carbon fractions in the soil by adsorbing them onto its surface. Biochar has a highly porous structure that also protects some organic carbon components from microbial degradation (He et al., 2023; Feng et al., 2021). In a field experiment, Qiu et al. (2023) applied biochar to 0%, 1%, 2%, and 4% of the dry weight of topsoil. After 8 and 12 months, they observed a positive correlation between the amount of particulate organic carbon and the amount of biochar applied, as well as significant increases in organic carbon, particulate

organic carbon, and DOC content. In comparison to the control, soil organic carbon, ROC, and POC levels rose by 7.09–38.08%, 1.62–39.80%, and 9.52–62.30%, respectively, when Jiang et al. (2022) applied 0, 5, 15, 20, and 40 t ha⁻¹ of biochar to the soil. As a result, soil active carbon fractions were significantly impacted by biochar additions.

With the extension of incubation time from 180 days to 365 days, the content of soil active carbon fractions decreased, which may be attributed to several factors (Stewart et al., 2009; Yang et al., 2017; Zhao et al., 2018). Firstly, as the incubation period increased, organic carbon in the soil was gradually degraded and mineralized by microorganisms, leading to a decrease in the content of ROC and DOC. Secondly, the long-term application of biochar may have transformed some organic carbon into more stable forms, which are less susceptible to microbial decomposition, resulting in a decline in the active organic carbon fractions. Additionally, changes in the soil environment, such as fluctuations in moisture, temperature, and pH, may have affected the availability and turnover rate of organic carbon, further contributing to the reduction in active organic carbon content.

Both biochar applications considerably raised the SSA and SCN of soil particles in this investigation (Table 2 and Table S1). According to Liang et al. (2006), this phenomenon is caused by the numerous negative charges that are carried on the surface of biochar. These charges come from the oxygen-containing functional groups, like -COOH, -OH, etc., that are widely distributed on the surface of biochar and which effectively improve the cation exchange of the soil. The high SSA of biochar

itself directly contributes to an increase in soil SSA, improves cation adsorption, and thus enhances exchange capacity (Lone et al., 2015). Consistent with Hu et al. (2021), our findings show that increasing the biochar application rate enhances soil SSA and SCN (Table 2 and Table S1). However, the increase in soil organic carbon (particularly the active carbon component) brought about by the addition of biochar may have contributed to the rise in SSA and SCN. The study's findings demonstrated that adding two different kinds of biochar raised the amount of active carbon in the soil and that there was a significant relationship between soil active carbon and soil electrochemical characteristics, with ROC and POC being the primary determinant of the soil surface's electrochemical characteristics (Table 4; Figures 3–4). According to Meyer et al. (1994) and Pennell et al. (1995), soil organic matter, particularly humic acid, has a cation exchange capacity of up to 100–700 cmol(c) kg⁻¹ and an SSA of up to 500–800 m² g⁻¹. Vertisol soil SSA (about 35.3%) and SCN (approximately 29.3%) were shown to decrease with the removal of soil organic matter (Yu et al., 2020). Compared with more stable organic carbon pools, ROC represents the relatively amorphous, chemically reactive fraction that is enriched in oxygen-containing functional groups (e.g. –COOH, –OH, –C=O) and is preferentially adsorbed onto or co-precipitated with mineral surfaces. Such functional groups provide abundant negatively charged sites and enhance the development of the diffuse double layer, thereby increasing soil SSA, SCN and surface charge density and modifying the balance between electrostatic repulsion and van der Waals attraction at the particle–water interface (He et al., 2021). In contrast, POC is typically found in smaller soil

particles and influences the electrochemical properties of the soil surface through physical adsorption. The role of POC primarily involves its interaction and stability with minerals, which increases the charge and physical surface characteristics of the soil, further regulating the soil's electrochemical behavior. ROC and POC jointly influence the electrochemical properties of the soil, with ROC being more sensitive to changes in soil electrochemical properties in the short term, while POC affects the soil's electrochemical reactions over a longer period through stable mineral-organic carbon complexes. These organic carbon fractions have been repeatedly identified as a sensitive indicator of short-term management effects and is closely linked to key fertility parameters, including effective CEC and other electrochemical properties (Culman et al., 2012). Therefore, soil active organic carbon (especially ROC and POC) is closely related to the electrochemical properties of the soil surface.

4.2 Response of soil internal forces to biochar application

Soil particles in close proximity experience mutual repulsion due to like charges, generating electrostatic repulsive pressure. Because soil texture, organic matter content, and mineral composition all affect the particles' surface charge, variations in these characteristics will result in variations in electrostatic repulsive pressure (Hu et al., 2015; Liu et al., 2021). At the same electrolyte concentration, soil electrostatic repulsive pressure decreased under both biochar treatments as biochar application increased from 1% to 10% (Figure 5). This phenomenon is caused by the substantial alterations in the electrochemical characteristics of the soil surface, which are shown by the increase in SCN and SSA, the decrease in the intensity of the surface electric

field and charge density, and the effective control of electrostatic repulsive pressure. Both forms of biochar reduced electrostatic repulsive pressure similarly, which was essentially in line with the trend of soil surface electrochemical parameters under varying rates of biochar application, which is in line with Wang et al. (2024) findings.

Biochar application also significantly increased van der Waals attractive pressure between soil particles, particularly at higher application rates (Figure 6). Van der Waals pressure is directly related to the Hamaker constant, which itself depends on the properties of the materials and the intervening medium (Bergström, 1997; Hu et al., 2021; Yu et al., 2020). The observed increases in Hamaker constants (Table 4) suggest that biochar incorporation introduced more organic complexes into the soil (Resurreccion et al., 2011). Additionally, our results revealed that net interparticle pressure decreased as biochar application increased (Figure 7), consistent with the findings of Yu et al. (2017). These results suggest that the influence of biochar on net pressure is primarily through increased van der Waals attraction and reduced electrostatic repulsion.

4.3 Response of soil aggregate stability to biochar application

In this study, SABS values decreased with increasing biochar application (Figure 8 and Figure S2), indicating improved aggregate stability. This aligns with previous findings that biochar enhances soil aggregation (Fu et al., 2021; Wang et al., 2022). The observed decline in net pressure with increasing biochar (Figure 7) supports this conclusion, as lower internal forces correspond to greater structural stability. This is consistent with prior reports by Yu et al. (2020) and Hu et al. (2021), who also found

that biochar reduces internal particle forces and enhances aggregate stability.

We also observed that SABS increased with decreasing electrolyte concentration and then plateaued, with a threshold near 10^{-2} mol L⁻¹, indicating a critical transition in aggregate disintegration. This turning point corresponded to a similar trend in net interparticle pressure, reinforcing the connection between soil internal forces and aggregate stability. Regression analysis confirmed a significant positive correlation between SABS and net pressure (Figure 9), highlighting the central role of interparticle forces in determining aggregate stability.

The effectiveness of biochar in improving soil aggregation depends on environmental conditions, soil type, and feedstock characteristics (Yang and Lu, 2021; Pituello et al., 2018). Overall, there were no significant differences in SIF distribution and aggregate stability between the two biochar treatments in this study. This may be attributed to the fact that the basic properties of the two types of straw biochar are relatively similar (Table 2). Additionally, there were no significant changes in the active carbon components and surface electrochemical properties of the soil after the two types of biochar were added. In addition, our study found that the improvement in soil aggregate stability was greater at 365 days compared to 180 days following the application of both types of biochar. This is because, compared to 180 days, biochar addition at 365 days increased both soil charge density and the Hamaker constant, which in turn enhanced both the electrostatic repulsive force and molecular attraction at the mesoscale between soil particles. However, the increase in molecular attraction was much greater than that of electrostatic repulsion, resulting in a lower net force.

Therefore, soil aggregate stability was stronger after 365 days of biochar addition.

Although this study focused on short-term (365-day) effects, biochar is known for its long-term stability in soil environments. Previous research has shown that biochar can enhance soil aggregation, nutrient retention, and carbon sequestration for years to decades (Jiang et al., 2022; Sun et al., 2021). However, these long-term benefits may be affected by aging processes, environmental variability, and management practices. Future long-term field experiments are needed to better understand the persistence and dynamics of biochar effects in diverse agroecosystems.

It should be noted that the Na^+ -saturation treatment used in this study represents a simplified condition compared with natural soil environments, where multiple exchangeable cations and complex electrolyte compositions coexist. Such differences may influence ionic strength, electrical double-layer characteristics, and particle interaction forces, thereby affecting soil aggregate stability. Therefore, the SIFs quantified in this study should be interpreted as a mechanistic assessment of biochar-induced changes in particle interactions rather than a direct representation of field conditions. Nevertheless, this controlled approach enables the quantitative evaluation of intrinsic changes in soil interparticle forces and provides insights into the mechanisms underlying biochar-mediated aggregate stabilization. Future research integrating mixed electrolyte systems, contrasting soil types, and long-term field experiments is required to further validate and extend these findings.

5. Conclusions

After 180 and 365 days of maize straw biochar and wheat straw biochar addition,

the contents of soil active carbon fractions increased with higher biochar application rates. Soil active carbon fractions were closely related to soil electrochemical properties, with ROC and POC being the primary influencing factors. The application of biochar markedly modified the SSA and SCN of soil particles, thereby increasing van der Waals attractive forces, reducing net repulsive pressure, and ultimately enhancing aggregate stability. These beneficial effects were further amplified as the application rate increased. As the electrolyte concentration decreased, the SABS first increased and then tended to level off, and $10^{-2} \text{ mol L}^{-1}$ was the turning point for the change. Although this study utilized Na^+ -saturated soil samples, which do not fully reflect natural soil conditions, the findings quantitatively assessed the intensity and distribution of soil internal forces (SIFs) and elucidated the mechanistic pathway by which biochar-induced increases in ROC and POC, along with associated changes in soil surface electrochemistry, enhance aggregate stability. Collectively, these findings offer a theoretical basis for the targeted use of biochar to manipulate soil internal forces, strengthen aggregate structure, and mitigate soil erosion in agricultural systems.

Author contributions.

The idea for this paper was conceptualized by XSS, HHZ, and FNH. XSS wrote the initial draft of the manuscript. All authors were involved in the review and editing the manuscript.

Competing interests.

The contact author has declared that none of the authors has any competing interests.

Code and data availability.

No custom code was used in this study. The experimental data that support the findings of this study are available from the corresponding author upon reasonable request. The data are not publicly available because they form part of an ongoing research project.

Acknowledgements.

This work was supported by the National Natural Science Foundation of China (42277311 and 42361144707), the Open Fund from the State Key Laboratory of Soil and Water Conservation and Desertification Control (F2010121002-202419), Natural Science Foundation of Shaanxi Province (2023-JC-YB-263) and the Natural Science Foundation of Chongqing, China (CSTB2024NSCQ-LZX0020).

References

Batista, A. M., Nunes, M. R., Pessoa, T. N., and Libardi, P. L.: Seasonal variation of the rhizosphere soil aggregation in an Oxisol, *Soil Tillage Res.*, 231, <https://doi.org/10.1016/j.still.2023.105741>, 2023.

Bergström, L.: Hamaker constants of inorganic materials, *Adv. Colloid Interface Sci.*,

- 70, 125-169, [https://doi.org/10.1016/s0001-8686\(97\)00003-1](https://doi.org/10.1016/s0001-8686(97)00003-1), 1997.
- Blair, G. J., Lefroy, R. D. B., and Lisle, L.: Soil carbon fractions based on their degree of oxidation, and the development of a carbon management index for agricultural systems, *Aust. J. Agric. Res.*, 46, <https://doi.org/10.1071/ar9951459>, 1995.
- Bronick, C. J. and Lal, R.: Soil structure and management: a review, *Geoderma*, 124, 3-22, <https://doi.org/10.1016/j.geoderma.2004.03.005>, 2005.
- Cao, X., Xu, Y., Wang, F., Zhang, Z., and Xu, X.: Changes of soil organic carbon and aggregate stability along elevation gradient in *Cunninghamia lanceolata* plantations, *Sci Rep*, 14, 31778, <https://doi.org/10.1038/s41598-024-81847-4>, 2024.
- Chen, Y., Huang, L., Huang, S., Guo, T., Zhang, S., Guo, D., Song, X., Ding, S., Mehran, M., Yang, Y., Yue, K., Su, S., Geng, M., and Zhang, H.: Long-term green manure incorporation increases soil carbon sequestration and improves aggregate stability by changing organic carbon components, *Soil Tillage Res.*, 258, <https://doi.org/10.1016/j.still.2025.107024>, 2026.
- Chen, Y., Kuzyakov, Y., Ma, Q., Du, Z., Sun, K., Xiao, K., Liang, X., Li, Y., Zhang, Y., Lai, X., Fu, W., Gao, B., Wang, F., Zhu, S., Gao, Q., and Rillig, M. C.: Aggregate size mediates the stability and temperature sensitivity of soil organic carbon in response to decadal biochar and straw amendments, *Soil Biology and Biochemistry*, 211, <https://doi.org/10.1016/j.soilbio.2025.109969>, 2025.
- Coleman, D. C., Reid, C. P. P., and Cole, C. V.: Biological Strategies of Nutrient Cycling in Soil Systems, in: *Adv. Ecol. Res.*, 1-55,

10.1016/s0065-2504(08)60107-5, 1983.

Culman, S. W., Snapp, S. S., Freeman, M. A., Schipanski, M. E., Beniston, J., Lal, R., Drinkwater, L. E., Franzluebbers, A. J., Glover, J. D., Grandy, A. S., Lee, J., Six, J., Maul, J. E., Mirksy, S. B., Spargo, J. T., and Wander, M. M.: Permanganate Oxidizable Carbon Reflects a Processed Soil Fraction that is Sensitive to Management, *Soil Sci. Soc. Am. J.*, 76, 494-504, <https://doi.org/10.2136/sssaj2011.0286>, 2012.

Ding, W., Liu, X., Hu, F., Zhu, H., Luo, Y., Li, S., and Li, H.: How the particle interaction forces determine soil water infiltration: Specific ion effects, *J. Hydrol.*, 568, 492-500, <https://doi.org/10.1016/j.jhydrol.2018.11.017>, 2019.

Feng, Z., Fan, Z., Song, H., Li, K., Lu, H., Liu, Y., and Cheng, F.: Biochar induced changes of soil dissolved organic matter: The release and adsorption of dissolved organic matter by biochar and soil, *Sci. Total Environ.*, 783, 147091, <https://doi.org/10.1016/j.scitotenv.2021.147091>, 2021.

Fu, G. Q., Qiu, X. N., Xu, X. Y., Zhang, W., Zang, F., and Zhao, C. Y.: The role of biochar particle size and application rate in promoting the hydraulic and physical properties of sandy desert soil, *Catena*, 207, <https://doi.org/10.1016/j.catena.2021.105607>, 2021.

Han, L., Sun, K., Yang, Y., Xia, X., Li, F., Yang, Z., and Xing, B.: Biochar's stability and effect on the content, composition and turnover of soil organic carbon, *Geoderma*, 364, <https://doi.org/10.1016/j.geoderma.2020.114184>, 2020.

Hashimi, R., Ganjegunte, G. K., Suthar, J. D., Kumar, S., and Palmate, S. S.: Soil

- carbon fractions and aggregate stability across management zones under flood-irrigated pecan orchards in an arid region, *J. Clean Prod.*, 549, <https://doi.org/10.1016/j.jclepro.2026.147871>, 2026.
- He, M. R., Jiang, Y., Han, Y. H., Zhu, W. H., Meng, D., Li, C. L., Cai, H. G., and Zhang, J. J.: Rice Straw Biochar is More Beneficial to Soil Organic Carbon Accumulation and Stabilization than Rice Straw and Rice Straw Ash, *J. Soil Sci. Plant Nutr.*, 23, 3023-3033, <https://doi.org/10.1007/s42729-023-01256-w>, 2023.
- He, Y., Yang, M., Huang, R., Wang, Y., and Ali, W.: Soil organic matter and clay zeta potential influence aggregation of a clayey red soil (Ultisol) under long-term fertilization, *Sci Rep*, 11, 20498, <https://doi.org/10.1038/s41598-021-99769-w>, 2021.
- Hu, F., Xu, C., Li, H., Li, S., Yu, Z., Li, Y., and He, X.: Particles interaction forces and their effects on soil aggregates breakdown, *Soil Tillage Res.*, 147, 1-9, <https://doi.org/10.1016/j.still.2014.11.006>, 2015.
- Hu, F., Xu, C., Ma, R., Tu, K., Yang, J., Zhao, S., Yang, M., and Zhang, F.: Biochar application driven change in soil internal forces improves aggregate stability: Based on a two-year field study, *Geoderma*, 403, <https://doi.org/10.1016/j.geoderma.2021.115276>, 2021.
- Janzen, H. H., Campbell, C. A., Brandt, S. A., Lafond, G. P., and Townley-Smith, L.: Light-Fraction Organic Matter in Soils from Long-Term Crop Rotations, *Soil Sci. Soc. Am. J.*, 56, 1799-1806, <https://doi.org/10.2136/sssaj1992.03615995005600060025x>, 1992.

- Ji, D. C., Ge, L. W., Van Zwieten, L., An, T. T., Li, S. Y., Kuzyakov, Y., Ding, F., and Wang, J. K.: Contrasting effects of maize straw and its biochar on aggregation and soil organic matter stabilization, *Plant Soil*, 495, 221-233, <https://doi.org/10.1007/s11104-023-06313-y>, 2024.
- Jia, A., Song, X., Li, S., Liu, Z., Liu, X., Han, Z., Gao, H., Gao, Q., Zha, Y., Liu, Y., Wu, X., and Wang, G.: Biochar enhances soil hydrological function by improving the pore structure of saline soil, *Agric. Water Manage.*, 306, <https://doi.org/10.1016/j.agwat.2024.109170>, 2024.
- Jiang, M. H., Li, C. B., Gao, W. C., Cai, K., Tang, Y., and Cheng, J. Z.: Comparison of long-term effects of biochar application on soil organic carbon and its fractions in two ecological sites in karst regions, *Geoderma Reg.*, 28, <https://doi.org/10.1016/j.geodrs.2021.e00477>, 2022.
- Li, C. L., Xu, J. B., He, Y. Q., Liu, Y. L., and Fan, J. B.: Dynamic Relationship Between Biologically Active Soil Organic Carbon and Aggregate Stability in Long-Term Organically Fertilized Soils, *Pedosphere*, 22, 616-622, [https://doi.org/10.1016/S1002-0160\(12\)60046-0](https://doi.org/10.1016/S1002-0160(12)60046-0), 2012.
- Li, H., Hou, J., Liu, X. M., Li, R., Zhu, H. L., and Wu, L. S.: Combined Determination of Specific Surface Area and Surface Charge Properties of Charged Particles from a Single Experiment, *Soil Sci. Soc. Am. J.*, 75, 2128-2135, <https://doi.org/10.2136/sssaj2010.0301>, 2011.
- Li, H., Liu, G., Dan, C., Zhou, Z., Liu, Y., Liu, C., Shen, E., Zhang, Q., Xia, X., Liu, D., Guo, Z., and Zhao, X.: Effects of soil porosity on water stability of

- aggregates, *Soil Tillage Res.*, 254, <https://doi.org/10.1016/j.still.2025.106741>, 2025.
- Li, H. R., Liu, G., Gu, J., Chen, H., Shi, H. Q., Abd Elbasit, M. A. M., and Hu, F. N.: Response of soil aggregate disintegration to the different content of organic carbon and its fractions during splash erosion, *Hydrol. Process.*, 35, <https://doi.org/10.1002/hyp.14060>, 2021.
- Li, J., Song, S., Xu, C., Du, W., Liu, X., Ding, W., and Hu, F.: Combined effects of soil internal and external forces on aggregate fragments detachment and displacement during splash erosion, *Catena*, 266, <https://doi.org/10.1016/j.catena.2026.109961>, 2026.
- Li, Q., Zhang, J., Ye, J., Liu, Y., Lin, Y., Yi, Z., and Wang, Y.: Biochar affects organic carbon composition and stability in highly acidic tea plantation soil, *J. Environ. Manage.*, 370, 122803, <https://doi.org/10.1016/j.jenvman.2024.122803>, 2024.
- Li, S.-z. and Xu, R.-k.: Electrical double layers' interaction between oppositely charged particles as related to surface charge density and ionic strength, *Colloid Surf. A-Physicochem. Eng. Asp.*, 326, 157-161, <https://doi.org/10.1016/j.colsurfa.2008.05.023>, 2008.
- Li, S., Li, H., Xu, C. Y., Huang, X. R., Xie, D. T., and Ni, J. P.: Particle Interaction Forces Induce Soil Particle Transport during Rainfall, *Soil Sci. Soc. Am. J.*, 77, 1563-1571, <https://doi.org/10.2136/sssaj2013.01.0009>, 2013.
- Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O., Thies, J., Luizao, F. J., Petersen, J., and Neves, E. G.: Black

- Carbon increases cation exchange capacity in soils, *Soil Sci. Soc. Am. J.*, 70, 1719-1730, <https://doi.org/10.2136/sssaj2005.0383>, 2006.
- Liang, Q., Chen, H., Gong, Y., Fan, M., Yang, H., Lal, R., and Kuzyakov, Y.: Effects of 15 years of manure and inorganic fertilizers on soil organic carbon fractions in a wheat-maize system in the North China Plain, *Nutr. Cycl. Agroecosyst.*, 92, 21-33, <https://doi.org/10.1007/s10705-011-9469-6>, 2011.
- Liu, J., Wang, Z., Hu, F., Xu, C., Ma, R., and Zhao, S.: Soil organic matter and silt contents determine soil particle surface electrochemical properties across a long-term natural restoration grassland, *Catena*, 190, <https://doi.org/10.1016/j.catena.2020.104526>, 2020.
- Liu, J., Hu, F., Xu, C., Wang, Z., Ma, R., Zhao, S., and Liu, G.: Comparison of different methods for assessing effects of soil interparticle forces on aggregate stability, *Geoderma*, 385, <https://doi.org/10.1016/j.geoderma.2020.114834>, 2021.
- Liu, J. Y., Yang, Y. A., Zheng, Q. W., Su, X. M., and Zhou, Z. C.: Response of soil aggregate stability and rill erodibility to soil electric field, *Catena*, 215, <https://doi.org/10.1016/j.catena.2022.106338>, 2022.
- Lone, A. H., Najar, G. R., Ganie, M. A., Sofi, J. A., and Ali, T.: Biochar for Sustainable Soil Health: A Review of Prospects and Concerns, *Pedosphere*, 25, 639-653, [https://doi.org/10.1016/s1002-0160\(15\)30045-x](https://doi.org/10.1016/s1002-0160(15)30045-x), 2015.
- Lu, T., Wang, X., Du, Z., and Wu, L.: Impacts of continuous biochar application on major carbon fractions in soil profile of North China Plain's cropland: In comparison with straw incorporation, *Agric. Ecosyst. Environ.*, 315,

<https://doi.org/10.1016/j.agee.2021.107445>, 2021.

Ma, L. J., Lv, X. B., Cao, N., Wang, Z., Zhou, Z. G., and Meng, Y. L.: Alterations of soil labile organic carbon fractions and biological properties under different residue-management methods with equivalent carbon input, *Appl. Soil Ecol.*, 161, <https://doi.org/10.1016/j.apsoil.2020.103821>, 2021a.

Ma, R., Hu, F., Liu, J., and Zhao, S.: Evaluating the effect of soil internal forces on the stability of natural soil aggregates during vegetation restoration, *J. Soils Sediments*, 21, 3034-3043, <https://doi.org/10.1007/s11368-021-03011-z>, 2021b.

Ma, R. t., Guo, W., Hu, F. N., Xu, C. Y., Liu, G., Zhao, S. w., and Zheng, F. l.: Effects of soil internal forces on splash detachment and transport of aggregate fragments in Mollisols of Northeast China, *Eur. J. Soil Sci.*, 73, <https://doi.org/10.1111/ejss.13273>, 2022.

Meyer, W. L., Arp, P. A., and Marsh, M.: Cation exchange capacities of upland soils in eastern Canada, *Can. J. Soil Sci.*, 74, 393-408, <https://doi.org/10.4141/cjss94-053>, 1994.

Pennell, K. D., Abriola, L. M., and Boyd, S. A.: Surface Area of Soil Organic Matter Reexamined, *Soil Sci. Soc. Am. J.*, 59, 1012-1018, <https://doi.org/10.2136/sssaj1995.03615995005900040008x>, 1995.

Pituello, C., Dal Ferro, N., Francioso, O., Simonetti, G., Berti, A., Piccoli, I., Pisi, A., and Morari, F.: Effects of biochar on the dynamics of aggregate stability in clay and sandy loam soils, *Eur. J. Soil Sci.*, 69, 827-842, <https://doi.org/10.1111/ejss.12676>, 2018.

- Qiu, H., Liu, J., Boorboori, M. R., Li, D., Chen, S., Ma, X., Cheng, P., and Zhang, H.: Effect of biochar application rate on changes in soil labile organic carbon fractions and the association between bacterial community assembly and carbon metabolism with time, *Sci. Total Environ.*, 855, 158876, <https://doi.org/10.1016/j.scitotenv.2022.158876>, 2023.
- Quirk, J. P.: Interparticle Forces: A Basis for the Interpretation of Soil Physical Behavior, in: *Adv. Agron.*, 121-183, 10.1016/s0065-2113(08)60614-8, 1994.
- Resurreccion, A. C., Moldrup, P., Tuller, M., Ferr é T. P. A., Kawamoto, K., Komatsu, T., and De Jonge, L. W.: Relationship between specific surface area and the dry end of the water retention curve for soils with varying clay and organic carbon contents, *Water Resour. Res.*, 47, <https://doi.org/10.1029/2010wr010229>, 2011.
- Sheng, M. H., Ai, X. Y., Huang, B. C., Zhu, M. K., Liu, Z. Y., and Ai, Y. W.: Effects of biochar additions on the mechanical stability of soil aggregates and their role in the dynamic renewal of aggregates in slope ecological restoration, *Sci. Total Environ.*, 898, 165478, <https://doi.org/10.1016/j.scitotenv.2023.165478>, 2023.
- Siedt, M., Schaffer, A., Smith, K. E. C., Nabel, M., Ross-Nickoll, M., and van Dongen, J. T.: Comparing straw, compost, and biochar regarding their suitability as agricultural soil amendments to affect soil structure, nutrient leaching, microbial communities, and the fate of pesticides, *Sci. Total Environ.*, 751, 141607, <https://doi.org/10.1016/j.scitotenv.2020.141607>, 2021.
- Stevenson, F. J.: *Humus chemistry: Genesis, composition, reactions*, John Wiley, New York 1994.

- Stewart, C. E., Paustian, K., Conant, R. T., Plante, A. F., and Six, J.: Soil carbon saturation: Implications for measurable carbon pool dynamics in long-term incubations, *Soil Biol. Biochem.*, 41, 357-366, <https://doi.org/10.1016/j.soilbio.2008.11.011>, 2009.
- Sun, Q., Meng, J., Lan, Y., Shi, G. H., Yang, X., Cao, D. A. Y., Chen, W. F., and Han, X. R.: Long-term effects of biochar amendment on soil aggregate stability and biological binding agents in brown earth, *Catena*, 205, <https://doi.org/10.1016/j.catena.2021.105460>, 2021.
- Tian, S., Zhu, B., Yin, R., Wang, M., Jiang, Y., Zhang, C., Li, D., Chen, X., Kardol, P., and Liu, M.: Organic fertilization promotes crop productivity through changes in soil aggregation, *Soil Biol. Biochem.*, 165, <https://doi.org/10.1016/j.soilbio.2021.108533>, 2022.
- Wander, M. M., Traina, S. J., Stinner, B. R., and Peters, S. E.: Organic and Conventional Management Effects on Biologically Active Soil Organic Matter Pools, *Soil Sci. Soc. Am. J.*, 58, 1130-1139, <https://doi.org/10.2136/sssaj1994.03615995005800040018x>, 1994.
- Wang, J., Fu, X., Sainju, U. M., and Zhao, F.: Soil carbon fractions in response to straw mulching in the Loess Plateau of China, *Biol. Fertil. Soils*, 54, 423-436, <https://doi.org/10.1007/s00374-018-1271-z>, 2018.
- Wang, K., Zhang, X., Zheng, J., Zhang, W., Yang, Z., Zhang, Q., Cai, J., and Wang, X.: Amended soils with weathered coal exhibited greater resistance to aggregate breakdown than those with biochar: From the viewpoint of soil internal forces,

- Soil Tillage Res., 244, <https://doi.org/10.1016/j.still.2024.106244>, 2024.
- Wang, L. W., Ok, Y. S., Tsang, D. C. W., Alessi, D. S., Rinklebe, J., Masek, O., Bolan, N. S., and Hou, D. Y.: Biochar composites: Emerging trends, field successes and sustainability implications, *Soil Use Manage.*, 38, 14-38, <https://doi.org/10.1111/sum.12731>, 2022.
- Wei, Z., Zhao, X., Zhu, C., Xi, B., Zhao, Y., and Yu, X.: Assessment of humification degree of dissolved organic matter from different composts using fluorescence spectroscopy technology, *Chemosphere*, 95, 261-267, <https://doi.org/10.1016/j.chemosphere.2013.08.087>, 2014.
- Xu, C. Y., Li, H., Hu, F. N., Li, S., Liu, X. M., and Li, Y.: Non-classical polarization of cations increases the stability of clay aggregates: specific ion effects on the stability of aggregates, *Eur. J. Soil Sci.*, 66, 615-623, <https://doi.org/10.1111/ejss.12252>, 2015.
- Yagüe, M. R., Domingo-Olivé, F., Bosch-Serra, À. D., Poch, R. M., and Boixadera, J.: Dairy Cattle Manure Effects on Soil Quality: Porosity, Earthworms, Aggregates and Soil Organic Carbon Fractions, *Land Degrad. Dev.*, 27, 1753-1762, <https://doi.org/10.1002/ldr.2477>, 2016.
- Yang, C. D. and Lu, S. G.: Effects of five different biochars on aggregation, water retention and mechanical properties of paddy soil: A field experiment of three-season crops, *Soil Tillage Res.*, 205, <https://doi.org/10.1016/j.still.2020.104798>, 2021.
- Yang, P., Liu, K., Wang, J., Yang, H., Gao, D., Chen, G., Liu, W., Tong, C., Sun, D.,

- Xu, Y., and Tang, K. W.: Shifts in organic carbon fractions, chemical functional groups, and mineralization enhance soil carbon storage after restoring earthen aquaculture ponds to wetlands, *Catena*, 272, <https://doi.org/10.1016/j.catena.2026.110370>, 2026.
- Yang, X., Meng, J., Lan, Y., Chen, W., Yang, T., Yuan, J., Liu, S., and Han, J.: Effects of maize stover and its biochar on soil CO₂ emissions and labile organic carbon fractions in Northeast China, *Agric. Ecosyst. Environ.*, 240, 24-31, <https://doi.org/10.1016/j.agee.2017.02.001>, 2017.
- Yu, Z., Zhang, J., Zhang, C., Xin, X., and Li, H.: The coupling effects of soil organic matter and particle interaction forces on soil aggregate stability, *Soil Tillage Res.*, 174, 251-260, <https://doi.org/10.1016/j.still.2017.08.004>, 2017.
- Yu, Z., Zheng, Y., Zhang, J., Zhang, C., Ma, D., Chen, L., and Cai, T.: Importance of soil interparticle forces and organic matter for aggregate stability in a temperate soil and a subtropical soil, *Geoderma*, 362, <https://doi.org/10.1016/j.geoderma.2019.114088>, 2020.
- Zhang, H., Zhang, R., Li, Y., Zhang, Z., Gao, Y., Dong, B., and Ma, J.: Biochar amendment coupled with deficit irrigation enhance silage maize yield by modulating soil physico-chemical properties and enzyme activities, *Industrial Crops and Products*, 246, <https://doi.org/10.1016/j.indcrop.2026.123354>, 2026a.
- Zhang, L., Chen, X., Liang, A., Huang, D., Huang, W., Liu, Y., Jiao, S., Ye, W., Wang, Y., and Wei, B.: Straw return depth and tillage intensity shape organic carbon fractions and their stability in black soil, *Agric. Ecosyst. Environ.*, 408,

<https://doi.org/10.1016/j.agee.2026.110478>, 2026b.

Zhang, Y., Shengzhe, E., Wang, Y. A., Su, S. M., Bai, L. Y., Wu, C. X., and Zeng, X.

B.: Long-term manure application enhances the stability of aggregates and aggregate-associated carbon by regulating soil physicochemical characteristics, *Catena*, 203, <https://doi.org/10.1016/j.catena.2021.105342>, 2021.

Zhang, Y. F., Dou, S., Ndzelu, B. S., Ma, R., Zhang, D. D., Zhang, X. W., Ye, S. F., and Wang, H. R.: Effects of returning corn straw and fermented corn straw to fields on the soil organic carbon pools and humus composition, *Soil*, 8, 605-619, <https://doi.org/10.5194/soil-8-605-2022>, 2022.

Zhao, S. X., Ta, N., Li, Z. H., Yang, Y., Zhang, X., Liu, D., Zhang, A., and Wang, X. D.: Varying pyrolysis temperature impacts application effects of biochar on soil labile organic carbon and humic fractions, *Appl. Soil Ecol.*, 123, 484-493, <https://doi.org/10.1016/j.apsoil.2017.09.007>, 2018.