Addressing systematic underestimation in global ship emission inventories through improved fleet representation and fuel compliance characterization


Weiwei Zhang[1,2], Zhaofeng Lv[1], Wen Yi[1,2], Tingkun He[1,2], Bensheng Xiao[1], Feng Sun[4], Qiang Zhang[3], Huan Liu[1,2]

[1] State Key Laboratory of Regional Environment and Sustainability, School of Environment, Tsinghua University, Beijing 100084, China
[2] Institute for Carbon Neutrality, Tsinghua University, Beijing 100084, China
[3] Ministry of Education Key Laboratory for Earth System Modeling, Department of Earth System Science, Tsinghua University, Beijing 100084, China
[4] COSCO SHIPPING Heavy Industry (Dalian) Co.,Ltd…, Dalian 116113, China

Correspondence to: Huan Liu (liu_env@tsinghua.edu.cn)

Abstract. As a sector with high energy demand, long-lived assets, and limited near-term low-carbon alternatives, global shipping is considered difficult to decarbonize and is subject to stringent emission controls. To evaluate emission control effects and conduct rapid response air quality simulations, accurate and timely ship emission inventories are indispensable. However, existing inventories face challenges from incomplete fleet representation and large uncertainties in $PM_{2.5}$ emissions after the global sulfur regulation. Here, we established a dynamically updated Ship Emission Inventory Model that allows near-real-time emission calculation within several days after AIS data acquisition. Ship activity and technical database are updated daily instead of yearly to obtain a more complete and rapid description of the global ship fleet. Fleet's multiple choices to comply with fuel sulfur regulation were considered, including switching to very low sulfur fuel and utilizing scrubbers to keep consuming heavy fuel oil. The daily expansion of ship technical database uncovered 8% and 6.2% of the total gross tonnage of active bulk carrier and container fleets, unveiling up to 5.4% of global ship CO_2 emissions. The resulting emission discrepancies gradually accumulated over 2024, demonstrating the persistent impact of delayed fleet information updates on ship emission estimates. Meanwhile, assuming that all ships switch to low-sulfur fuels after 2020 would lead to underestimation of $PM_{2.5}$ and BC emissions by approximately 43% and 15%. Although South China Ocean had the most absolute underestimation, the Indian Ocean had the highest underestimated portion, reaching 56% and 15% of its total $PM_{2.5}$ and BC emissions.

删除了: **Addressing systemic underestimation in global ship emissions from fleet growth and fuel compliance…**
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: As a hard-to-abate sector, global shipping is under international and regional emission control regulations. …
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: Without the expansion, the daily underestimation would enlarge over time from about 0.20 Mt CO_2 /d to 0.29 Mt CO_2 /d throughout 2024. …
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: 55
删除了: 27
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: 75
删除了: 39
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)



Systemic emission underestimat
Multiple compliance ch
• Switch to VLSF(
• HFO + scrubber
Incomplete fleet cover

删除了:

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

1 Introduction

As one of the hard-to-abate sectors (Sharmina et al., 2021), global shipping experienced a threefold increase in cargo turnover and a 1.7-fold increase in carbon dioxide (CO_2) emissions from 1970 to 2021 (Wang et al., 2025). Growing ship emissions have brought adverse impacts on the air quality and public health of coastal regions (Liu et al., 2016; Liu et al., 2019; Anenberg et al., 2019; Zhang et al., 2021). In 2018, global shipping was responsible for 2.9% of anthropogenic CO_2 emissions, 11% of sulfur oxides (SO_x), and 15% of nitrogen oxides (NO_x) (IMO, 2020). Under a business-as-usual scenario without additional mitigation measures, international shipping was projected to account for approximately 17% of global CO_2 emissions by 2050 (Cames et al., 2015), which reflects both the future development of international shipping emissions and the emission trajectories of other sectors. To restrain ship emissions, international and regional regulations have been implemented. Previous research showed that after the International Maritime Organization (IMO) low sulfur regulation came into force in 2020, ship SO_2 emissions dropped by 81.3% globally (Yi et al., 2024a) and by 71% in Southern China (Feng et al., 2023). Studies evaluating regional sulfur regulations have demonstrated their effectiveness in reducing ship-related SO_2 emissions. The sulfur emission control measures implemented in the Marmara Sea and Turkish Straits since 2006 reduced SO_2 concentrations by 90% in Istanbul (Viana et al., 2015) while China's Domestic Emission Control Area (DECA) policy, implemented progressively during 2016–2019, achieved an emission reduction by 33% within 200 nautical miles away from the Chinese mainland's territorial sea baseline (Wang et al., 2021).

删除了: If unregulated, the contribution to total CO_2 emissions would surge to 17% in 2050 (Cames et al., 2015)….

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: Studies evaluating regional regulations found that sulfur emission control areas (ECA) in Marmara Sea and Turkish Straits reduced SO_2 concentration by 90% in Istanbul…

设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗

Accurate and timely emission inventories are key to evaluating air quality and climate impacts and facilitating future policymaking (Im et al., 2018; Luo et al., 2024; Liu et al., 2024; Goldsworthy et al., 2019; Huang et al., 2020; Li et al., 2024). Bottom-up models using Automatic Identification System (AIS) have substantially improved the spatial and temporal

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

representation of shipping emissions by combining ship activity with ship-specific technical characteristics, engine-load calculations, and emission factors. However, converting ship activity into emissions requires information beyond vessel movements, including fuel consumption, technical specifications, and voyage-specific compliance pathways, which are not directly provided by AIS records. Therefore, existing models such as STEAM, MariTEAM, and MoSES generally rely on vessel characteristics, fleet statistics, regulatory assumptions, or predefined scenarios to estimate fuel use and emission-control practices (Schwarzkopf et al., 2021; Jalkanen et al., 2009; Kramel et al., 2021). These assumptions introduce two major uncertainties in global ship emission inventories: incomplete representation of the evolving fleet and insufficient characterization of fuel compliance pathways. In particular, distinguishing Heavy Fuel Oil (HFO) use with scrubbers from low-sulfur fuel switching remains challenging at the global ship level, despite their substantially different implications for $PM_{2.5}$, and BC emissions.

Among these uncertainties, the representation of sulfur compliance pathways became particularly important after the implementation of the IMO 2020 sulfur regulation. From January 1, 2020, the global sulfur limit for ship fuel oil was reduced from 3.50% to 0.50%, while a stricter limit of 0.10% applies within ECAs. Ships can comply with this regulation either by switching to low-sulfur fuels such as Very Low Sulfur Fuel oil (VLSFO) and Marine Gas Oil (MGO) or by continuing to use HFO with scrubber systems. However, globally complete ship-level information on scrubber installation is rarely available, limiting the ability of emission inventories to explicitly represent HFO-with-scrubber pathways. Some previous shipping emission inventories assumed that ships previously using HFO switched to VLSFO or MGO after 2020 without explicitly considering scrubber-based compliance (Yi et al., 2024a; Yi et al., 2024b).

Nevertheless, multi-source evidence indicates that HFO-based compliance pathways remained important after 2020. IMO fuel statistics showed that the broad HFO category still accounted for approximately 65% of ship fuel consumption after 2020 (IMO, 2024), while regional bunker data indicated that HFO remained an important fuel category in major maritime hubs, with reported shares ranging from 21.4% to 37.2% in Singapore and Rotterdam. Moreover, Lunde Hermansson et al. (2024) identified more than 800 scrubber-equipped ships operating in the Baltic Sea after 2020, demonstrating the continued use of scrubbers in regional fleets. However, these statistics provide indirect evidence of HFO consumption or regional scrubber presence rather than globally representative ship-level compliance information. Therefore, neglecting HFO-with-scrubber pathways may introduce substantial uncertainty into post-2020 estimates of $PM_{2.5}$, and BC emissions.

Another challenge facing ship emission inventories is the timely incorporation of newly identified ships into ship technical databases. In this study, newly identified ships include both newly delivered vessels and existing vessels absent from the original static database. Ships built after 2016 contributed 10.2% of global ship emissions in 2021 (Yi et al., 2024a), indicating that recently delivered vessels already represent a non-negligible fraction of shipping activity. Unlike ship activity data that can be processed at daily resolution, ship technical databases are generally updated with substantial delays (Yi et al., 2024a). Newly identified ships often lack key technical specifications and are represented by proxy values, which may affect engine power estimation, load-factor correction, and emission factor assignment. Previous studies have extracted ship parameters from AIS data (Wang et al., 2021; Huang et al., 2020; Schwarzkopf et al., 2021) and supplement missing

设置了格式: 默认段落字体, 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 英语 (英国)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: However, the accuracy of existing emission inventories is systemically constrained by the incomplete and delayed technical data (Hu et al., 2025; Huang et al., 2021; Ye et al., 2025), especially for the global maritime transportation sector (Liu et al., 2022; Baek et al., 2024). First, the fuel choice assumption after the implementation of IMO low sulfur regulation by existing ship emission inventory model do not align with the actual situation. IMO low sulfur regulation requires that, from January 1, 2020, the global cap on the sulfur content of ship fuel oil be reduced from 3.50% to 0.50%. Within ECAs, the sulfur content cap is 0.10%. Previous ship emission inventory models assumed that after 2020, all ships using heavy fuel oil (HFO) had switched to very low sulfur fuel oil (VLSFO) or marine gas oil (MGO) (Yi et al., 2024a; Yi et al., 2024b). Yet, multi-source data suggest that using HFO with scrubber to comply with IMO regulation is a mainstream choice for global ship fleets. IMO fuel statistics showed that HFO still accounts for 65% of the ship fuel consumption even after 2020 (IMO, 2024). HFO also constituted of 21.4% - 37.2% of the marine bunker fuel in Singapore and Rotterdam (https://shipandbunker.com/). Lunde Hermansson et al. (2024) found that after 2020 there were over 800 ships in the Baltic Sea equipped with scrubbers to comply with IMO regulation. As the choice of HFO with installation of scrubber would be cost-competitive to the choice of VLSFO for over 95% of the ships within five years after installation(Lunde Hermansson et al., 2024), ship scrubber use would maintain the mainstream choice for ship fleets in the short or middle term.
删除了: (Eirik Ovrum, 2024)to include newly built or newly identified ships into ship technical database. Ships built after 2016 contributed to 10.2% of global ship emissions in 2021 (Yi et al., 2024a). With the number and gross tonnage of ships on order r ... [1]
删除了: Unlike ship activity data, which can be updated daily, ship technical database are generally supplemented at least one year
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: As a result, newly built ships often lack key technical specifications and have to be represented by proxy values, whi ... [2]
删除了: Automatic identification systems (
删除了:)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)

parameters with machine learning methods (Yang et al., 2022; Huang et al., 2024; Yang et al., 2024; Raman et al., 2025). However, an integrated framework that simultaneously updates ship activity and technical information with quality control remains lacking.

Here, we developed SEIMv2.3, a dynamically updated ship emission inventory model based on Shipping Emission Inventory Model (SEIM) (Liu et al., 2016; Wang et al., 2021; Yi et al., 2024a). SEIMv2.3 improves the representation of ship activity, fleet composition, and sulfur compliance pathways through three methodological developments. First, AIS activity records are processed at a daily temporal resolution with an enhanced preprocessing framework to support near-real-time emission updates. Second, newly identified ships are dynamically detected from AIS activity records and incorporated into the static ship technical database, with missing specifications supplemented through AIS-derived information and machine-learning-based reconstruction. Third, the fleet low sulfur compliance choice module (FLSCCM) is introduced to reconstruct post-2020 sulfur compliance pathways by integrating ship-level scrubber installation records with constrained gap filling. Using SEIMv2.3, we calculated global ship emissions from 2022 to 2024 at daily resolution and quantified the effects of improved fleet representation and sulfur compliance pathway characterization on emission estimates.

2 Method

SEIM was first established by (Liu et al., 2016) based on the idea of the disaggregated dynamic method. Driven by AIS data and ship technical information, SEIM realized vessel-by-vessel ship emission simulations, suitable for the establishment of multi-scale shipping emission inventories with applications in regions (Liu et al., 2016; Wang et al., 2021; Zhang et al., 2025b) and ports (Fu et al., 2017). The specific formula used in this study to calculate ship emissions are identical to those reported by (Yi et al., 2024a) and provided in Supplementary Information. Emission factors for different ship types and fuels are listed in Table S2-S4, primarily based on the IMO Fourth Greenhouse Gas Study (IMO, 2020) and U.S. Environmental Protection Agency reports (Starcrest Consulting Group, 2022). Engine-load effects were explicitly considered by estimating the main-engine load for each AIS record and applying pollutant-specific low-load adjustment factors when the load was below 20%. Detailed factors are provided in Fig. S4.

Similar to other AIS-based bottom-up models, SEIM reconstructs ship activity at high spatial and temporal resolution but cannot directly observe fuel consumption or compliance technologies used during each voyage. Therefore, emission estimation depends on both accurate fleet representation and appropriate characterization of fuel compliance pathways. To improve the timeliness and completeness of these key inputs, SEIMv2.3 introduces three methodological improvements: (1) Daily AIS-based activity processing. AIS records are processed at a daily temporal resolution rather than annual aggregation, enabling ship activity reconstruction and emission estimation with a short delay and providing the technical basis for near-real-time emission updates. Detailed methods are provided in Sect. 2.1. (2) Daily identification of previously missing ships. In SEIMv2.2, ship activity records were matched with our Ship Technical Specifications Database (STSD), which may become incomplete as new ships enter operation or previously unrecorded ships become active. SEIMv2.3 uses daily AIS

设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: , there is currently no established framework where activity data and technical data are updated simultaneously with quality control.…
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: SEIMv2.3 updates both activity data and ship technical data on a daily basis. On one hand, ship activity data went through pre-processing framework specifically designed to handle anomalies in daily signals. On the other hand, ship technical parameters of newly identified ships were extracted from AIS data to supplement the ship technical database. The fleet low sulfur compliance choice module was added to identify HFO and scrubber use. We calculated global ship emissions from 2022 to 2024 day-by-day and evaluated the improvement by SEIMv2.3 on global fleet description, and daily and annual emission calculation.…
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: and Table S3
删除了: (IMO, 2020)
删除了: ,
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: , and the National Standard for General Diesel Fuel of the People's Republic of China (GB 252-2015)…
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: 6
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗
设置了格式: 字体: (默认) +西文正文 (Times New Roman)

activity records to identify previously missing ships and update the STSD, with detailed procedures described in Sect. 2.2. (3) Fleet low sulfur compliance choice module (FLSCCM). SEIMv2.3 incorporates multiple sulfur compliance pathways, including low-sulfur fuel switching and HFO use with scrubbers, to better represent post-2020 sulfur compliance choices. Detailed methods are provided in Sect. 2.3.

2.1 From daily AIS to near-real-time emissions

This study established a daily AIS pre-processing framework specifically designed to handle anomalies in daily signals. AIS data consist of both dynamic data and static data. Dynamic AIS offers ship activity information encompassing timestamps and latitude-longitude coordinates, and are subject to multi-stage cleaning to establish a well-cleaned daily ship activity database. Static AIS contains ship technical information including ship type, length, width, and draft and are used for daily identification and characterization of new ships. In this study, we collected global daily AIS data from 2022 to 2024, comprising approximately 113 million dynamic signals and 14 million static signals per day. The daily AIS processing resulted in approximately 45 billion processed signals annually. AIS data contain anomalies and inaccuracies due to various reasons. Therefore, rigorous pre-processing was necessary especially for daily near-real-time emission calculation.

For the daily AIS dataset, firstly, to ensure the stability of daily input data, the model reads AIS signals from the three days preceding and following the target date when calculating daily emissions. Unlike annual AIS preprocessing, the daily framework explicitly accounts for transmission delays because daily AIS data received on a given day may contain signals with timestamps from both the current and previous days. As demonstrated in Fig. 1 (a), the total AIS signals received in one day generally comprised of about 90% of signals with timestamp of the current day and 10% of signals with timestamp of the previous day. Secondly, three-stage cleaning was performed to remove low-quality signals, temporal anomaly, and spatial anomaly. Fig. 1(c) and (d) present the spatial anomaly which leads to the abnormal AIS signal surge during February 21 and 29, 2024 in Fig. 1 (b), when the signal count exceeded 200 million per day. If unprocessed, the abnormal surge would result in up to 2 times of emission overestimation. The cleaning criteria in this study is the same as the work by (Yi et al., 2024a). Signals with invalid geographic coordinates, excessively long-time intervals, or ground speed exceeding 50 knots were discarded in this process. Thirdly, linear interpolation and route restoration was employed to reconstruct the actual sailing trajectory of ships. In the previous step, we removed signals located on land. Additionally, although some signals are not located over land themselves, the connecting lines with adjacent signals may cross over land. In this step, we reconstructed the sailing trajectory between the starting and ending points of those signals based on a global shipping route network using the Dijkstra shortest path algorithm (Wang et al., 2021; Aulinger et al., 2016; Johansson et al., 2017; Cherkassky et al., 1996). Interpolation was performed segment by segment at fixed time intervals, and emissions are calculated accordingly. Besides, as shown in Fig. 1 (e) and (f), the delayed signals contributed to a considerable part of the entire shipping routes, indicating the necessity of three-day AIS processing to the reconstruction of actual sailing trajectories. This preprocessing window is required for trajectory reconstruction from delayed AIS transmissions, while SEIMv2.3 enables near-real-time emission updates with a short delay after AIS data acquisition. The overall effect of pre-processing

设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: ↵
Compared to previous versions (Wang et al., 2021; Yi et al., 2024a), **this dynamically updated version, SEIMv2.3, features three key improvements: (1) AIS data are collected, preprocessed, and calculated daily instead of yearly. Emission inventories with a rapid response capability are crucial for pollution control and air quality simulation. In SEIMv2.3, we collected daily AIS data and enhanced the pre-process accordingly to enable near-real-time ship emission calculation. Compared with collecting AIS annually, this operation does not change the emission results, but only changes the temporal resolution of the results. Detailed methods are provided in Sect. 2.1. (2) Daily identification and characterization of new ships. The number of active ships identified by SEIMv2.2 dropped markedly in recent years, from 108,000 in 2021 to 104,000 in 2023 while global fleet data from UNCTAD showed an increase by 9.2 % from 99,800 ships 2021 to 109,000 ships in 2023(UNCTAD, 2021, 2024). To obtain a more complete description of global ship fleet, we promptly updated the Ship Technical Specification Database (STSD) with daily AIS data, which is discussed in detail in Sect. 2.2.** (3) The fleet low sulfur compliance choice module. We simulated the multiple choices of fleet to comply with IMO low sulfur regulation to generate more accurate emission calculation. Detailed methods are provided in Sect. 2.3. …
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 字体颜色: 文字 1
删除了: daily AIS from 2022 to 2024, which comprised of ... [3]
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: Annual AIS data used in previous SEIM were also ... [4]
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)

framework on global AIS signals from 2022 to 2024 is illustrated in Fig. S1. Figure 1 (g) and (h) display the improvement in coastal regions along China and western Europe on February 22, 2024. The framework managed to reduce inland signals by 46% and 51% in the two regions, where spatial anomalies were removed while signals on inland rivers were kept.




Figure 1: Anomalies and transmission delays of daily AIS signals. Timestamp of dynamic and static AIS signals received on Day n (a) and their spatial distribution in coastal areas along China (e) and western Europe (f). Example of daily signal count anomaly during February 21 to 29, 2024, Day 52 to Day 60 of 2024 (b). Spatial distribution of AIS signals received on Day 53 of 2024 in coastal areas along China (c) and western Europe (d). Spatial distribution of AIS signals of Day 53 after pre-processing in coastal areas along China (g) and western Europe (h).

2.2 Daily identification and characterization of previously missing ships

The STSD in SEIM describes ship attributes such as ship type, deadweight tonnage (DWT), and main engine power, and is originally updated once every two years. In our previous work, we updated the STSD with new-built ships from 2019 to 2021 obtained from Lloyd's Register (Yi et al., 2024a). To keep up with the update frequency of ship activity database and mitigate emission underestimation brought by delayed STSD update, a dynamic framework utilizing daily static AIS to promptly update the STSD has been established.

First, we identified newly detected ships by filtering IMO number that were not yet included in STSD from daily static AIS signals. Before utilizing static AIS to update the STSD, two major challenges need to be solved: (1) the classification of cargo ships in static AIS is overly broad, which does not align with the 14-type ship classification required by SEIMv2.3. While static AIS assign values 70–79 uniformly to cargo ships, SEIMv2.3 further subdivides cargo ships into auto carriers, bulk carriers, container ships, and general cargo ships, each with distinct emission factors. (2) Static AIS lacks main engine power and DWT information, which is also critical for matching emission factors. Without additional parameter completion, these missing specifications would have to be represented by proxy values, which may introduce considerable uncertainties in engine power estimation and subsequent emission factor assignment. Through extensive literature review, we selected

设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: 1
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: new ships
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: Ship Technical Specification Database (
删除了:)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)

machine learning algorithms and feature sets that have demonstrated strong fitting performance to supplement and refine these missing or coarse parameters (Baek et al., 2024; Huang et al., 2024; Raman et al., 2025; Yang et al., 2024). The final models, features, and hyperparameters used are listed in Table S1. Cargo ships were further subdivided into auto carriers, bulk carriers, container ships, and general cargo ships based on geometric features such as ship length and width using the XGBoost, achieving an accuracy of 0.82. XGBoost and Random Forest were applied to predict DWT (r^2 = 0.92, RMSE = 4.9 ktons, MAE = 1.9 ktons) and main engine power (r^2 = 0.80, RMSE = 1.7 kW, MAE = 0.9 kW), significantly improving data completeness. Figure S2 demonstrates the performance of key ship parameter prediction. Ultimately, the reconstructed technical information of newly identified ships was merged with the original STSD, forming a dynamically updated STSD as shown in Fig. 2.




Figure 2: The composition of ship types (a), deadweight tonnage (b), and main engine power (c) for daily newly detected ships. The ship categories shown here represent the dominant categories identified from processed AIS static information for newly identified ships and do not correspond directly to the 14 emission categories used in SEIMv2.3.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: the supplemented information of newly detected ship was merged with the original STSD, forming a daily updated STSD as shown in Fig. 2. Daily static AIS data from 2022 to 2024 were used to enhance our description of the global ship fleet.…
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman), 非加粗

2.3 Fleet low sulfur compliance choice module (FLSCCM)

In response to the IMO low sulfur regulation, ships primarily have three compliance options (Chu Van et al., 2019; Deng et al., 2021; Ji and El-Halwagi, 2020), with different implications for fuel consumption, SO_2 emissions, $PM_{2.5}$, and BC emissions: (1) Switching to low-sulfur fuels (e.g., MGO, VLSFO). This straightforward option directly lowers fuel sulfur content, resulting in sharp reduction in SO_2, $PM_{2.5}$ and BC emissions. As shown in Table S2 and Table S4, the emission factors for HFO (2.43% sulfur) are 0.051 gSO_2/g fuel and 1.39 $gPM_{2.5}$/kWh for the main engine, whereas for VLSFO (0.5% sulfur) they are 0.0098 gSO_2/g fuel and 0.31 $gPM_{2.5}$/kWh. Assuming other operational factors remain constant, this switch reduces SO_2 by 81% and reduces $PM_{2.5}$ and BC emissions by approximately 77%. (2) Continuing to use cheap HFO with a scrubber. Scrubbers remove SO_2 from exhaust but are less effective against $PM_{2.5}$. Studies reported the $PM_{2.5}$ removal efficiency ranging only from about 16% to 37% and no significant BC removal efficiency for ship scrubbers (Yang et al., 2021; Kuittinen et al., 2024). Therefore, compared to option (1), this option yields similar SO_2 emissions but results in relatively higher $PM_{2.5}$ and BC emissions. (3) Using alternative fuels such as LNG, which can substantially reduce emissions but currently represent a limited share of global marine fuel consumption. Therefore, FLSCCM focuses on the allocation of the two dominant sulfur compliance pathways: low-sulfur fuel switching and HFO with scrubbers.

Because globally complete ship-level compliance information is unavailable, FLSCCM reconstructs fleet sulfur compliance pathways by combining ship-level scrubber installation records with constrained statistical gap filling. First, scrubber-equipped ships recorded in the Clarksons Research database are identified and matched with the SEIM fleet using ship MMSI. Ships were considered to have observed scrubber capability only when they were identified as fitted with scrubbers and the installation was effective before the corresponding emission calculation period. These records indicate scrubber installation availability rather than voyage-level operation. In this study, ships with effective installations were assumed to follow the HFO-with-scrubber pathway during the corresponding period.

For ships without available scrubber installation records, FLSCCM applies structurally constrained gap filling based on characteristics derived from scrubber-equipped vessels identified from the installation database. Analysis of the scrubber installation database in Supplementary Information showed that scrubber adoption was concentrated among major cargo ship categories and larger vessels. Bulk carriers, containers, and tankers together accounted for approximately 79.5% of identified installations (Fig. S8). Therefore, gap filling was restricted to bulk carriers, containers, oil tankers, and chemical tankers. In addition, ship size and construction year were introduced as screening criteria because scrubber installation is economically favoured for ships with higher fuel consumption and sufficient remaining service life. Database-unmatched ships were considered eligible candidates only when they had a DWT of at least 50,000 tonnes and a construction year later than 2006 when reliable information was available. These criteria were applied only to ships without scrubber installation records during the residual gap-filling procedure and did not affect ships with confirmed scrubber installations. The threshold settings were determined based on the characteristics of the observed scrubber-equipped fleet. For vessels identified or assigned as following the HFO-with-scrubber pathway, SEIMv2.3 applies a baseline $PM_{2.5}$ removal efficiency of 37% and a

设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: 3
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: . This option drastically reduces most air pollutants, as the emission factors are negligible compared to HFO. However, the current ship LNG consumption is minimal relative to HFO, VLSFO, or MGO.…
删除了: Consequently, the fleet low sulfur compliance choice module only considers the allocation of option (1) and (2). …

BC removal efficiency of 30% in the emission calculation. Uncertainties associated with scrubber assignment criteria and removal efficiencies are further evaluated through sensitivity analyses in the Supplementary Information and the Discussion section.

The reconstructed fleet-level fuel structure was evaluated against IMO fuel consumption statistics for the post-2020 period (Fig. 3). SEIMv2.3 reproduced the major transition from HFO-dominated fuel use before IMO 2020 to increased adoption of low-sulfur fuels and HFO-with-scrubber pathways afterward. The remaining differences between modelled and IMO-reported fuel shares are likely attributable to differences in fuel classification, fleet coverage, and the explicit separation of HFO-with-scrubber pathways in SEIMv2.3. Therefore, IMO statistics were used as an independent fleet-level reference to assess the plausibility of reconstructed compliance pathways rather than as a direct constraint for model calibration.




Figure 3: Fuel consumption validation with IMO statistics

3 Results

3.1 Enhanced description of global shipping fleet

The daily identification and characterization mechanism enables the model to effectively capture newly-detected ship activity. An average of 20 ships (identified by MMSI number) previously absent from the STSD are detected per day, their characteristics are presented in Fig. 2. Bulk carriers constitute the primary category among these newly added ships, accounting for approximately 57% of the total. Regarding DWT range, newly identified vessels were predominantly distributed within the 10,000–100,000 DWT range, accounting for about 59% of newly identified vessels by number. Meanwhile, main engine power was largely concentrated in the 1,000–10,000 kW range, which included approximately 58% of newly identified vessels by number. The daily identification and characterization mechanism also led to a continuous

设置了格式: 字体: (默认) +西文正文 (Times New Roman), 英语 (英国)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

上移了 [1]: The module was validated by comparing its fuel consumption results with IMO statistics of 2020, 2022 and 2023. SEIMv2.3 successfully reproduced the dramatic fuel shift: the HFO share of total consumption fell sharply from 82.8% in 2019 to 52.2% in 2020, matching the IMO-reported values of 81.2% and 50.7%, respectively (Fig. 3). The absolute HFO consumption estimates for the post-2020 period also align with the statistics.

删除了: The module was validated by comparing its fuel consumption results with IMO statistics of 2020, 2022 and 2023. SEIMv2.3 successfully reproduced the dramatic fuel shift: the HFO share of total consumption fell sharply from 82.8% in 2019 to 52.2% in 2020, matching the IMO-reported values of 81.2% and 50.7%, respectively (Fig. 3). The absolute HFO consumption estimates for the post-2020 period also align with the statistics.

Due to the lack of global, ship-specific data on actual compliance choices, this study developed a proxy method based on vessel type and size to assign the compliance choice to each ship. Analysis of IMO fuel consumption statistics by ship type and DWT revealed that bulk carriers, container ships, and tankers accounted for nearly 90% of global HFO consumption (IMO, 2024). Within these categories, the proportion of HFO use increases significantly with vessel size (Fig. S3). Based on this pattern, a DWT threshold was established for these three key ship types. Ships exceeding the threshold are assigned to option (2) as the economic incentive for scrubber installation is stronger for larger, higher-fuel-consumption ships. Ships at or below the threshold are assigned to option (1). The DWT threshold was calibrated by testing a range of values (e.g., 40,000–60,000 DWT) against the IMO fuel consumption data. The threshold that minimized the difference between the modeled and reported HFO consumption share in 2021 was selected. For this study, a threshold of 40,000 DWT was applied (Fig. S4). The DWT threshold is used as a proxy for scrubber adoption among major HFO-consuming cargo ship types rather than as a ship-type screening criterion. Therefore, it does not explicitly capture scrubber adoption by passenger ships and ser ... [5]

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman), (中文) 宋体, 非加粗, (中文) 中文(中国), (其他) 英语(美国)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman), (中文) 中文(中国), (其他) 英语(美国)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)


设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: Regarding DWT range, the newly detected ships are predominantly distributed within the 10k–100k ton range, representing about 59% of the total. Meanwhile, the main engi ... [6]

expansion of our STSD, as illustrated in Fig. S5. Over the study period, the number of ships (identified by unique MMSI) in the STSD increased from 418,000 to 440,000, representing a growth of 5.3%. If we identified ship number by unique IMO number (primarily for oceangoing ships), the STSD exhibited more significant growth, expanding from 105,000 to 126,000 ships—a notable increase of 20.0%, underlining that dynamic update of STSD effectively enhanced the coverage of oceangoing ships.

The expansion of the STSD have enabled SEIMv2.3 to provide a more comprehensive characterization of the global active fleet. In 2024, SEIMv2.3 captured activities from 132,000 ships, including 68,000 IMO-identified ships. Compared with SEIMv2.2, the number of active bulk carriers, container ships, general cargo ships, and fishing ships increased by 22.9%, 6.8%, 8.0%, and 14.6%, respectively. Bulk carriers and container ships showed the largest increases in fleet tonnage, with gross tonnage increases of 8.0% and 6.2%, respectively. Newly identified ships were primarily constructed between 2000 and 2010, indicating that the update mainly supplemented missing information for existing vessels (Fig. 4). DWT distribution reveals that while the ship number increase in SEIMv2.3 was dominated by ship whose DWT was less than 10 kt, the growth in total fleet tonnage was driven predominantly by ship DWT between 60 to 80 kt. The newly identified ships accounted for approximately 10.3% of all active ships annually and contributed about 11.3% of the total annual sailing distance in 2024. Their activities exhibited the highest signal density in major maritime hubs, notably in East Asia and Europe, with their activity widely distributed across global shipping routes, as shown in Fig. 5. This geographical pattern reflects the higher rate of new ship deployment in these key shipbuilding and operational regions, such as China, South Korea, and the major ports of Northern Europe.



设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: Using 2024 as an example, SEIMv2.3 captured the activities of 132,000 ships, including 68,000 ships identified by IMO number. As illustrated in Fig. 4, compared to SEIMv2.2, the number of active bulk carriers, container ships, general cargo ships, and fishing ships identified by SEIMv2.3 increased by 22.9%, 6.8%, 8.0%, and 14.6%, respectively. Regarding transport capacity, bulk carriers remained the primary contributor to total fleet tonnage, with SEIMv2.3 reporting an 8.0% increase in gross tonnage for bulk carriers, while container ship tonnage grew by 6.2%. In terms of age distribution, SEIMv2.3 primarily supplemented activities of ships constructed between 2000 and 2010, both in ship number and gross tonnage. …

设置了格式: 字体: (默认) +西文正文 (Times New Roman)


删除了: The newly detected ships represent a significant and active component of global shipping activity. …

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: Without the dynamic update of the STSD, this substantial portion of maritime activity would either be omitted or be inaccurately estimated using generalized default emission factors. Spatially, the newly detected ship…

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: ↵



删除了:

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

Figure 4: The composition of ship number and total weight of the active global fleet by SEIMv2.3 and SEIMv2.2. Panels (a–c) show fleet number by ship type, build period, and weight bin; panels (d–f) show the corresponding distributions of total fleet weight. The inner ring distinguishes ships contained in the original STSD (light grey) from ships newly identified through the dynamic STSD updating procedure (dark grey). The outer coloured ring shows the category composition of the original STSD fleet, and the linked stacked bar shows the composition of newly identified vessels using the same colour scheme.




Figure 5: Spatial distribution of newly detected ship signals.

3.2 Annual ship emissions and comparison with other studies

Following our previous global ship emission estimates for 2016–2021 (Yi et al., 2024a), SEIMv2.3 extends the inventory to 2022–2024. To improve the representation of emission changes before and after the IMO 2020 sulfur regulation, historical emissions from 2019 to 2021 were recalculated using the revised SEIMv2.3 framework. Figure 6 shows the global ship emissions of nine atmospheric pollutants and greenhouse gases (GHG), including CO_2, CH_4, and N_2O, calculated using different models from 2019 to 2024. For comparison, available international shipping emission estimates from the Emissions Database for Global Atmospheric Research (EDGAR) and the Community Emissions Data System (CEDS) were used as independent global references to evaluate the temporal consistency of SEIMv2.3. For EDGAR, international shipping emissions were derived using the Ship Traffic Emission Assessment Model (STEAM3) framework (Crippa et al., 2024). Comparisons were limited to pollutants with available and comparable estimates.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: **For the Fleet number by weight bin subplot, the radial axis is displayed on a logarithmic scale to improve the visualization of categories with relatively small fleet sizes.…**
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: Building on our previous work of ship emission from 2016 to 2021…
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: this study further presents global ship emissions from 2022 to 2024. …
删除了: To obtain more accurate emission results before and after IMO low sulfur regulation, we calculated historical ship emissions from 2019 to 2021 using SEIMv2.3. …
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
设置了格式: 字体: (默认) +西文正文 (Times New Roman)
删除了: For comparison, we used international shipping emissions from the Emissions Database for Global Atmospheric Research (EDGAR), a global sector-resolved inventory of anthropogenic greenhouse gases and air pollutants, and the Community Emissions Data System (CEDS), a consistent historical inventory of sectoral anthropogenic emissions. These datasets provide independent global estimates against which the temporal trends in SEIMv2.3 can be evaluated.…
设置了格式: 字体: (默认) +西文正文 (Times New Roman)

From 2019 to 2020, ship SO_2 emissions dropped sharply from 12,215 kt to 2,212 kt, consistent with the results from EDGAR and CEDS. $PM_{2.5}$ emissions showed a substantial but less pronounced reduction than SO_2 because the implementation of sulfur regulations affected not only fuel sulfur content but also the selection of compliance pathways. In SEIMv2.3, global ship $PM_{2.5}$ emissions decreased by 60.2%, from 1,725 kt in 2019 to 686 kt in 2020, and remained relatively stable at 703–768 kt during 2021–2024. In comparison, SEIMv2.2, which did not represent HFO-with-scrubber pathways after 2020, estimated only 393 kt $PM_{2.5}$ emissions in 2020, corresponding to a 42.7% underestimation relative to SEIMv2.3. BC emissions exhibited a comparable but weaker response. SEIMv2.3 estimated that global ship BC emissions decreased by 30.3%, from 98 kt in 2019 to 68.3 kt in 2020, followed by relatively stable emissions between 67.2 and 70.6 kt during 2021–2024. In contrast, SEIMv2.2 estimated 58 kt BC emissions in 2020, corresponding to a 15.1% underestimation relative to SEIMv2.3 and indicating the importance of representing HFO-with-scrubber pathways for BC emission estimation. For other atmospheric pollutants and GHGs, the improvement mainly results from daily identification and characterization of newly observed ships. Comparison between SEIMv2.3 and SEIMv2.2 shows that the absence of daily ship technical database updates would underestimate CO_2 emissions by 5.4% in 2023 and 2.6% in 2024. Although the proportion of underestimated emissions relative to total shipping emissions was relatively low, the inclusion of newly identified ships enabled us to capture emission changes caused by fluctuations in maritime trade and provided a more complete basis for monitoring shipping decarbonization and evaluating mitigation policies.

To distinguish the contributions of daily STSD updates and the FLSCCM, numerical experiments were conducted for the 2024 inventory. Experiment (i) applied SEIMv2.2 without daily STSD updates or FLSCCM; experiment (ii) applied the updated STSD without FLSCCM; and experiment (iii) applied the complete SEIMv2.3 framework. The difference between experiments (iii) and (ii) represents the contribution of FLSCCM, whereas the difference between experiments (ii) and (i) represents the contribution of daily STSD updates. The results indicate that emission underestimation sources differ among pollutants: STSD updates mainly improve CO_2 and gaseous pollutant estimates by increasing fleet completeness, while FLSCCM remains particularly important for $PM_{2.5}$ and BC because these pollutants are directly affected by fuel sulfur content and scrubber-related removal efficiencies.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: For $PM_{2.5}$ emissions, the results from SEIMv2.3 show that from 2019 to 2020, ship $PM_{2.5}$ emissions decreased from 1,725 kt to 893 kt, a reduction of approximately 48.2%, and then fluctuated slightly between 856 kt and 932 kt from 2021 to 2024. SEIMv2.2 did not consider HFO use after 2020 and underestimated $PM_{2.5}$ emissions to be 393 kt in 2020, representing a decrease of about 77.2% compared to 2019. EDGAR results do not reflect the control effect of IMO's sulfur regulations on $PM_{2.5}$ emissions. Although there was also a decrease by 8.6% in ship $PM_{2.5}$ emissions from 2019 to 2020 in EDGAR results, this can be attributed to reduced ship activity due to the pandemic in 2020. BC emissions show a similar pattern to $PM_{2.5}$ emissions. Results SEIMv2.3 show that from 2019 to 2020, ship BC emissions decreased from 98 kt to 83 kt, a reduction of approximately 15.3%, and then fluctuated between 83 kt and 88 kt from 2021 to 2024, highly consistent with CEDS results. SEIMv2.2 underestimated BC emissions to be 58 kt in 2020, representing a decrease of about 40.8% compared to 2019. Comparison between SEIMv2.3 and SEIMv2.2 shows that the ignorance of HFO use after 2020 would lead to an underestimation of $PM_{2.5}$ and BC emissions by approximately 55% and 27%, 97% of which can be mitigated by fleet low sulfur compliance choice module. For other atmospheric pollutants and GHGs, the emission underestimation was mitigated by daily identification and characterization of new ships.
Comparison between SEIMv2.3 and SEIMv2.2 shows that the absence of daily identification and characterization of new ships would lead to CO_2 emission underestimation by 5.4% in 2023 and 2.6% in 2024. Although the proportion of underestimated emissions relative to total shipping emissions was relatively low, the inclusion of newly identified ships enabled us to capture emission changes caused by fluctuations in maritime trade and provided a more complete basis for monitoring shipping decarbonization and evaluating mitigation policies. To quantitatively distinguish the contributions of daily identification and characterization of new ships (denoted as STSD-2024 in Fig. 6) and the fleet low sulfur compliance choice module (denoted as FLSCCM-2024 in Figure 6), numerical experiments were conducted for the 2024 emission inventory. (i) SEIMv2.2 was applied without daily STSD updates or the FLSCCM; (ii) SEIMv2.3 was applied with the updated STSD but with the FLSCCM disabled; and (iii) the complete SEIMv2.3 framework was applied. Consequently, the difference between experiments (iii) and (ii) represents the contribution of the FLSCCM, whereas the difference between experiments (ii) and (i) represents the contribution of the daily STSD update. The results indicate that the dominant source of emission underestimation varies among pollutants. The daily STSD update contributes most to the reduction in underestimation for CO_2 and most gaseous pollutants by improving fleet completeness, whereas the FLSCCM plays a larger role for $PM_{2.5}$ and BC because these pollutants are highly sensitive to post-2020 fuel compliance pathways.





Figure 6: Comparison of annual global ship emissions results from SEIMv2.3, SEIMv2.2, and other emission inventories from 2019 to 2024. SEIMv2.2 results were sourced from (Yi et al., 2024a). SEIMv2.3 represents the method presented in this paper. STEAM3 represents the emission results from EDGAR (Crippa M., 2025), whose shipping emission results are based on STEAM3. CEDS represents the emission results from Community Earth atmospheric Data Systems, sourced from (Hoesly, 2025). STSD-2024 and FLSCCM-2024 denote configurations incorporating updated fleet information and reconstructed sulfur compliance pathways, respectively.

3.3 Spatiotemporal heterogeneity in ship emission underestimation

Emission underestimation showed distinct spatial patterns among pollutants (Fig. 7). The major discrepancies were concentrated in the South China Sea, North Atlantic Ocean, Indian Ocean, and North Pacific Ocean. Figure 7 compares regional emission discrepancies with sailing distances of newly identified ships and scrubber-equipped ships to identify potential sources of underestimation. For CO_2 and NOx, the largest discrepancies were primarily associated with incomplete



删除了:

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

带格式的: 制表位: 2.24 字符, 左对齐 + 不在 3.17 字符

删除了: **studies**

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: **EDGAR**

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: ↵

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: For different air pollutants and GHGs, the underestimated emissions were concentrated in the South China Sea, North Atlantic Ocean, Indian Ocean, and North Pacific Ocean, consistent with the spatial distribution of newly added ship signals shown in Fig. 5. Figure 7 illustrates the ocean regions with the top 20 underestimated emissions derived by subtracting SEIMv2.2 results from the SEIMv2.3 results, most of which appeared in South China Sea, accounting for 13.4% to 15.2% of the total underestimation. However, in terms of the percentage of underestimated emissions relative to the total emissions of each region, the Indian Ocean and the South Atlantic Ocean exhibited the highest proportions of underestimation. For CO_2 and NOx emissions, the underestimation in the Indian Ocean and the South Atlantic Ocean accounted for 8.1% and 7.5% of ... [7]

fleet representation. The South China Sea contributed the largest shares of global CO_2 and NOx underestimation, accounting for 14.2% and 14.4%, respectively, followed by the Indian Ocean and North Atlantic Ocean. These regions also exhibited relatively high STSD-added vessel sailing distances, indicating the important contribution of incomplete fleet representation to CO_2 and NOx discrepancies. In contrast, $PM_{2.5}$ and BC underestimations were more strongly associated with incomplete representation of sulfur compliance pathways. The Indian Ocean showed the largest contributions to global $PM_{2.5}$ and BC underestimation, accounting for 15.1% and 14.7%, respectively. These regions also showed substantial scrubber-associated sailing distances, indicating that incomplete representation of HFO-with-scrubber pathways was a major contributor to particulate emission discrepancies.

The relative impact of underestimation on individual regions was further evaluated by normalizing emission differences against regional total emissions. For CO_2, the relative differences were relatively small across major shipping regions, reaching 3.0% in the South China Sea, 2.9% in the Indian Ocean, and 2.8% in the South Atlantic Ocean. Similarly, NOx underestimation ratios ranged from 2.7% to 4.6%, with the highest value observed in the South China Sea. In contrast, particulate emissions exhibited substantially larger relative differences. The relative underestimation of $PM_{2.5}$ reached 56.1% in the Indian Ocean, 53.7% in the South Atlantic Ocean, and 48.5% in the South China Sea. For BC, the corresponding values were 14.5%, 12.8%, and 10.8%, respectively. These results highlight the stronger influence of compliance pathway assumptions than fleet completeness on regional $PM_{2.5}$ and BC estimates.

The differences between SEIMv2.3 and SEIMv2.2 accumulated throughout 2024 due to differences in fleet representation and sulfur compliance pathway characterization. The daily emission discrepancies remained consistently positive throughout 2024. As shown in Fig. 8, the daily emission discrepancies between SEIMv2.3 and SEIMv2.2 increased from 52.8 kt CO_2 d^{-1}, 1.52 kt NOx d^{-1}, 0.93 kt $PM_{2.5}$ d^{-1}, and 0.017 kt BC d^{-1} at the beginning of 2024 to 62.1 kt CO_2 d^{-1}, 1.79 kt NOx d^{-1}, 0.95 kt $PM_{2.5}$ d^{-1}, and 0.018 kt BC d^{-1} at the end of 2024. The corresponding cumulative emission discrepancies reached 21.3 Mt CO_2, 615.8 kt NOx, 350.3 kt $PM_{2.5}$, and 6.5 kt BC by the end of 2024.





Figure 7. Ship $PM_{2.5}$, BC, CO_2, and NOx emission underestimation mitigated by SEIMv2.3 in different ocean regions in 2024.



删除了：↵
By increasing the update frequency of the STSD from yearly to daily, we managed to include emissions previously omitted or miscalculated due to delayed update of ship parameters. The average supplemented daily emissions are 1.52 kt $PM_{2.5}$, 6.07 kt NOx, 0.62 kt SO_2, 0.22 kt CO, 0.24 kt HC, 0.25 Mt CO_2, 0.01 kt N_2O, 0.01 CH_4, and 0.07 kt BC. Those emissions account for 58% of daily $PM_{2.5}$ emissions and 28% of daily BC emissions. For the remaining pollutants, the underestimated proportion was 9.1% for SO_2, 12.6% for NOx, 9.5% for CO, 11.2% for HC, 10.0% for CO_2, 1.9% for CH_4, and 7.6% for N_2O. Figure 8 shows the temporal variations in calculated ship $PM_{2.5}$ and CO_2 emissions throughout 2024, derived by subtracting SEIMv2.2 results from the SEIMv2.3 results. Despite some fluctuations, the daily emission differences show an increasing trend over time. The daily $PM_{2.5}$ emission difference climbed from about 1.45 kt/d at the start of 2024 to 1.58 kt/d at the end of 2024. The same was true with other emissions. The daily CO_2 emission difference climbed from about 0.20 Mt/d at the start of 2024 to 0.29 Mt/d at the end of 2024. The findings highlight the major advantage of SEIMv2.3 in its accuracy in daily emission calculation.↵
↵
↵
... [8]

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman), (中文) 宋体, 13 磅, 非加粗

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)


设置了格式: 字体: (默认) +西文正文 (Times New Roman)

Figure 8. Temporal accumulation of emission discrepancies between SEIMv2.3 and SEIMv2.2 during 2024.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

4 Summary

In this study, we introduced SEIMv2.3, which features daily updated ship activity database and STSD to support near-real-time ship emission calculation. With fleet low sulfur compliance choice module, it can simulate fleet's multiple choices to comply with emission control regulations, which forms the fuel consumption structure that aligns well with the actual situation. On average, 23 newly detected ships were supplemented to the STSD daily. From 2022 to 2024, ships included in our STSD rose from 105,000 to 126,000, an increase of 20.0%. The daily expansion of the STSD unveiled an important proportion of the global maritime transport capacity, about 8.0% and 6.2% of the total gross tonnage of the active bulk carrier and container fleets. Those newly-detected ships contributed to 11.3% of the total sailing distance and 10.3% of active ship fleet, uncovering 5.4% and 2.6% of the CO_2 emissions in 2023 and 2024. For $PM_{2.5}$ and BC emissions, the neglect of HFO-with-scrubber pathways after 2020 would lead to underestimations of approximately 43% and 15%, respectively. Our results suggest a decrease of 60.2% in $PM_{2.5}$ emissions from 2019 to 2020, compared with the 77.2% reduction estimated by SEIMv2.2. The South China Sea contributed the largest shares of global CO_2 and NOx underestimation, accounting for 14.2% and 14.4%, respectively, while the Indian Ocean showed the largest contributions for $PM_{2.5}$ and BC underestimation, accounting for 15.1% and 14.7%, respectively. When normalized by regional total emissions, the Indian Ocean and South Atlantic Ocean exhibited the largest relative underestimations. For $PM_{2.5}$ and BC emissions, the underestimated portion reached 56.1% and 14.5% of regional emissions in the Indian Ocean, respectively, compared with 48.5% and 10.8% in the South China Sea. Without the methodological improvements in SEIMv2.3, emission differences accumulated throughout 2024. The daily CO_2 emission difference increased from about 0.053 Mt/d to 0.062 Mt/d, while the daily $PM_{2.5}$ emission difference increased from about 0.93 kt/d to 0.95 kt/d.



删除了:

Figure 8. Temporal variations in daily ship $PM_{2.5}$, BC, CO_2, and NOx emission underestimation.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: For $PM_{2.5}$ and BC emissions, the ignorance of HFO use after 2020 would lead to an underestimation of $PM_{2.5}$ and BC emissions by approximately 55% and 27%. Our results suggest a decrease by 48.2% in $PM_{2.5}$ emission from 2020 to 2019, instead of by 77.2% by previous bottom-up method.…

5 Discussion

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

Overall, our study establishes a dynamically updated framework for global ship emission estimation and reveals substantial emission underestimation associated with incomplete fleet representation and simplified sulfur compliance assumptions in previous SEIM configurations. Near-real-time emission inventories enable tracking of short-term emission changes and provide a data foundation for emission reduction policies (Xu et al., 2023). The underestimation addressed in this study may affect assessments of the environmental and health impacts of global shipping. Underestimated $PM_{2.5}$ and SO_2 emissions may bias estimates of shipping-induced aerosol–cloud interactions and radiative forcing (Chen et al., 2017; Goldsworthy et al., 2019; Yuan et al., 2024; Zhang et al., 2025a). Such underestimation may also affect assessments of atmospheric deposition impacts in coastal and marine environments (Shi et al., 2023). The additional contribution to population exposure

删除了: Overall, our study pioneers in its near-real-time framework to calculate global ship emissions and its discovery of the systemic emission underestimation in previous emission inventory studies.…

删除了: systemic

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: The deposition of sulfur and nitrogen compounds may also affect surface-water acidity and marine biogeochemistry, particularly along heavily trafficked routes and in coastal or semi-enclosed

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

and health burdens may be limited, because the additional emissions identified in this study were mainly in open-ocean regions, distant from populated coastal areas.

Our conclusions are also supported by independent evidence from IMO fuel consumption statistics, scrubber installation records, and ambient particle observations. IMO fuel statistics indicate that HFO-based pathways remained important after 2020 (IMO, 2024), while (Lunde Hermansson et al., 2024) identified continued operation of scrubber-equipped ships after the sulfur regulation, supporting the necessity of explicitly representing HFO-with-scrubber pathways. (Song et al., 2022) analyzed $PM_{2.5}$ concentrations near Busan Port before and after the IMO 2020 sulfur regulation and found limited changes in ambient $PM_{2.5}$, highlighting the complexity of evaluating shipping impacts based solely on particle mass concentrations.

In addition to $PM_{2.5}$ mass emissions, particle number (PN) emissions deserve further attention because ship exhaust contains large numbers of ultrafine particles that are not fully represented by particle mass. Previous studies demonstrated that shipping remains an important source of anthropogenic PN emissions after 2020 (Kuittinen et al., 2021). Measurements in the Baltic Sea sulfur emission control area showed that reducing fuel sulfur content from 1.5% to 0.1% decreased plume PN concentrations by 27% and ambient PN concentrations by 32% (Seppälä et al., 2021). However, the influence of scrubbers on PN is complex and depends on particle size and scrubber configuration. Onboard measurements of a modern cruise ship showed that a hybrid scrubber reduced total PN and nonvolatile PN, whereas particles larger than 50 nm, which may include BC and metal-containing particles, were not significantly reduced (Kuittinen et al., 2024). These findings indicate that future ship emission inventories should extend beyond $PM_{2.5}$ mass emissions and incorporate PN size distributions and scrubber-specific particle removal characteristics.

Despite the improvements introduced in SEIMv2.3, several uncertainties remain. First, missing vessel specifications supplemented through machine learning may introduce errors at the individual-ship level, although these uncertainties are expected to decrease after aggregation at regional and global scales. Second, FLSCCM reconstructs sulfur compliance pathways using vessel-level scrubber records combined with constrained statistical allocation because global ship-specific fuel-use observations are unavailable. Third, FLSCCM does not distinguish ULSFO and MGO, and fuels of 0.1% sulfur are represented using MGO emission factors. This simplification is expected to have limited influence on SO_2, but it may lead to some underestimation of $PM_{2.5}$ and BC emissions where residual-type ULSFO is used. Therefore, uncertainties may arise from incomplete scrubber records, assumed scrubber operation, and simplified representation of ULSFO using MGO emission factors. These uncertainties mainly affect $PM_{2.5}$ and BC estimates, while their influence on SO_2 remains limited.

To quantify the influence of scrubber-related uncertainty on particulate emission estimates, we conducted sensitivity analyses using different scrubber removal efficiency assumptions. Under the low-, baseline-, and high-removal scenarios, corresponding to $PM_{2.5}$ removal efficiencies of 10%, 37%, and 70% and BC removal efficiencies of 10%, 30%, and 60%, global $PM_{2.5}$ emissions varied by +26.6% and −32.6% relative to the baseline case, while BC emissions varied by +11.5% and −17.3%. Despite this uncertainty, all scenarios consistently show higher particulate emissions than inventories that neglect or incompletely represent HFO-with-scrubber pathways, indicating that the identified underestimation is robust to plausible variations in scrubber removal performance.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

删除了: Our conclusions align well with IMO fuel consumption statistics, fleet installation of scrubbers by (Lunde Hermansson et al., 2024), and $PM_{2.5}$ concentration observational research. (Song et al., 2022) analyzed the $PM_{2.5}$ concentrations near Busan Port, South Korea, before and after 2020. His findings indicated that the IMO low sulfur regulation did not affect $PM_{2.5}$ emissions. In addition to $PM_{2.5}$ mass emissions, PN emissions deserve further attention because ship exhaust contains large numbers of ultrafine particles that are not well represented by particle mass alone. (Kuittinen et al., 2021) argued that ship emissions remain a significant source of anthropogenic PN emissions after 2020. Measurements in the Baltic Sea SECA showed that reducing fuel sulfur content from 1.5% to 0.1% decreased plume PN concentrations by 27% and ambient PN concentrations by 32% (Seppälä et al., 2021). Meanwhile, ship plumes can still contribute up to 19% of PN concentrations on days affected by shipping lanes (Kivekäs et al., 2014). The effect of scrubbers on PN is also size- and configuration-dependent. Onboard measurements on a modern cruise ship showed that a hybrid scrubber reduced total PN and nonvolatile PN, but no significant reduction was observed for particles larger than 50 nm, which commonly include BC and metal-containing particles (Kuittinen et al., 2024).↵
Admittedly, certain limitations exist in our methodology. On one hand, although SEIMv2.3 improves the representation of shipping $PM_{2.5}$ emissions, the present inventory does not include PN emissions. Future development should incorporate PN emission factors and size-resolved particle information to better assess the climatic and health impacts of shipping emissions. On the other hand, uncertainties arise from multiple aspects. First, missing ship specifications supplemented by machine learning may introduce errors for individual vessels, although such errors are expected to decrease after aggregation to regional and global scales. Second, FLSCCM relies on statistical allocation of fuel consumption because actual ship-level fuel choices are generally unavailable. Third, FLSCCM does not distinguish ULSFO and MGO, and fuels of 0.1% sulfur are represented using MGO emission factors. This simplification is expected to have limited influence on SO_2, but it may lead to some underestimation of $PM_{2.5}$ and BC emissions where ULSFO is used. Finally, incomplete scrubber installation records may bias $PM_{2.5}$, BC, and PN emission calculation because unidentified scrubber-equipped vessels may be treated as using low-sulfur fuels. Besides, installation records alone do not fully indicate whether scrubbers are actually operated during the whole voyages, which introduces further uncertainty.

删除了: ↵

Future reductions in uncertainty require improved ship-specific technical databases, broader sharing of scrubber installation records and fuel-use data, and additional validation datasets such as onboard measurements, EU MRV, or ship bunkering records. Direct single-ship validation remains challenging because such data are rarely publicly available. Nevertheless, comparisons with independent inventories and sensitivity analyses indicate that these uncertainties are unlikely to alter the main conclusions regarding the magnitude and characteristics of global shipping emissions.

Code and data availability

Python codes used during the current study are available from the corresponding author on reasonable request.

Author contributions

Weiwei Zhang developed the model, performed the simulations, and prepared the manuscript. Zhaofeng Lv managed the project and edited the manuscript. Bensheng Xiao collected the data. Wen Yi, Tingkun He, and Feng Sun validated the results and reviewed the manuscript. Huan Liu and Qiang Zhang conceptualized the research aim, supervised the project and acquired the financial support.

Competing interests

One of the co-authors is member of the editorial board of Atmospheric Chemistry and Physics.

Acknowledgements

This work was supported by the National Key R&D Program of China (No. 2023YFC3705604) and COSCO SHIPPING Co., Ltd. (No. 2025-4-Z001-03).

References

Anenberg, S. C., Miller, J., Henze, D. K., Minjares, R., and Achakulwisut, P.: The global burden of transportation tailpipe emissions on air pollution-related mortality in 2010 and 2015, Environmental Research Letters, 14, 094012, 10.1088/1748-9326/ab35fc, 2019.

Aulinger, A., Matthias, V., Zeretzke, M., Bieser, J., Quante, M., and Backes, A.: The impact of shipping emissions on air pollution in the greater North Sea region – Part 1: Current emissions and concentrations, Atmos. Chem. Phys., 16, 739-758, 10.5194/acp-16-739-2016, 2016.

删除了：These uncertainties can be constrained at the inventory level through complete data curation and systematic sensitivity analyses, while future improvements in ship-specific technical databases and broader sharing of ship technical specifications, scrubber installation records, and fuel use data would further reduce uncertainties at the individual-ship level and facilitate the use of shipping emission inventories in future climate and impact models. Direct validation at the single-ship level remains difficult because such data are rarely publicly available. Future validation could use onboard measurements, EU MRV, or ship bunkering data. Nevertheless, comparisons with independent inventories and sensitivity analyses indicate that these uncertainties are unlikely to affect the main conclusions on the magnitude, temporal evolution, and pollutant characteristics of global shipping emissions.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

Baek, W.-K., Kim, E., Jeon, H.-K., Lee, K.-J., Kim, S.-W., Lee, Y.-K., and Ryu, J.-H.: Monitoring Maritime Ship Characteristics Using Satellite Remote Sensing Data from Different Sensors, Ocean Science Journal, 59, 8, 10.1007/s12601-023-00131-0, 2024.

Cames, M., Graichen, J., Siemons, A., and Cook, V.: Emission Reduction Targets for International Aviation and Shipping European Parliament, 2015.

Chen, D., Wang, X., Nelson, P., Li, Y., Zhao, N., Zhao, Y., Lang, J., Zhou, Y., and Guo, X.: Ship emission inventory and its impact on the PM2.5 air pollution in Qingdao Port, North China, Atmos. Environ., 166, 351-361, https://doi.org/10.1016/j.atmosenv.2017.07.021, 2017.

Cherkassky, B. V., Goldberg, A. V., and Radzik, T.: Shortest paths algorithms: Theory and experimental evaluation, Mathematical Programming, 73, 129-174, 10.1007/BF02592101, 1996.

Chu Van, T., Ramirez, J., Rainey, T., Ristovski, Z., and Brown, R. J.: Global impacts of recent IMO regulations on marine fuel oil refining processes and ship emissions, Transportation Research Part D: Transport and Environment, 70, 123-134, https://doi.org/10.1016/j.trd.2019.04.001, 2019.

Crippa, M., Guizzardi, D., Pagani, F., Schiavina, M., Melchiorri, M., Pisoni, E., Graziosi, F., Muntean, M., Maes, J., Dijkstra, L., Van Damme, M., Clarisse, L., and Coheur, P.: Insights into the spatial distribution of global, national, and subnational greenhouse gas emissions in the Emissions Database for Global Atmospheric Research (EDGAR v8.0), Earth Syst. Sci. Data, 16, 2811-2830, 10.5194/essd-16-2811-2024, 2024.

Crippa M., G. D., Pagani F., Banja M., Muntean M., Schaaf, E., Quadrelli, R., Risquez Martin, A., Taghavi-Moharamli, P., Grassi, G., Rossi, S., Melo, J., Oom, D., Branco, A., Suarez Moreno, M., Sedano, F. San-Miguel, J., Manca, G., Pisoni, E., Pekar, F.: GHG emissions of all world countries – JRC/IEA 2025 Report, LuxembourgJRC143227, https://data.europa.eu/doi/10.2760/9816914, 2025.

Deng, J., Wang, X., Wei, Z., Wang, L., Wang, C., and Chen, Z.: A review of NOx and SOx emission reduction technologies for marine diesel engines and the potential evaluation of liquefied natural gas fuelled vessels, Science of The Total Environment, 766, 144319, https://doi.org/10.1016/j.scitotenv.2020.144319, 2021.

Feng, X., Ma, Y., Lin, H., Fu, T.-M., Zhang, Y., Wang, X., Zhang, A., Yuan, Y., Han, Z., Mao, J., Wang, D., Zhu, L., Wu, Y., Li, Y., and Yang, X.: Impacts of Ship Emissions on Air Quality in Southern China: Opportunistic Insights from the Abrupt Emission Changes in Early 2020, Environmental Science & Technology, 57, 16999-17010, 10.1021/acs.est.3c04155, 2023.

Fu, M., Liu, H., Jin, X., and He, K.: National- to port-level inventories of shipping emissions in China, Environmental Research Letters, 12, 114024, 10.1088/1748-9326/aa897a, 2017.

Goldsworthy, B., Enshaei, H., and Jayasinghe, S.: Comparison of large-scale ship exhaust emissions across multiple resolutions: From annual to hourly data, Atmospheric Environment, 214, 116829, https://doi.org/10.1016/j.atmosenv.2019.116829, 2019.

Hoesly, R., Smith, S. J., Ahsan, H., Prime, N., O'Rourke, P., Crippa, M., Klimont, Z., Guizzardi, D., Feng, L., Harkins, C., MCDONALD, B., & Wang, S.: CEDS v_2025_03_18 Aggregate Data (v_2025_03_18) https://doi.org/10.5281/zenodo.15059443, 2025.

Huang, I.-L., Lee, M.-C., Nieh, C.-Y., and Huang, J.-C.: Ship Classification Based on AIS Data and Machine Learning Methods, Electronics, 13, 98, 2024.

Huang, L., Wen, Y., Zhang, Y., Zhou, C., Zhang, F., and Yang, T.: Dynamic calculation of ship exhaust emissions based on real-time AIS data, Transportation Research Part D: Transport and Environment, 80, 102277, https://doi.org/10.1016/j.trd.2020.102277, 2020.

Im, U., Christensen, J. H., Geels, C., Hansen, K. M., Brandt, J., Solazzo, E., Alyuz, U., Balzarini, A., Baro, R., Bellasio, R., Bianconi, R., Bieser, J., Colette, A., Curci, G., Farrow, A., Flemming, J., Fraser, A., Jimenez-Guerrero, P., Kitwiroon, N., Liu, P., Nopmongcol, U., Palacios-Peña, L., Pirovano, G., Pozzoli, L., Prank, M., Rose, R., Sokhi, R., Tuccella, P., Unal, A., Vivanco, M. G., Yarwood, G., Hogrefe, C., and Galmarini, S.: Influence of anthropogenic emissions and boundary conditions on multi-model simulations of major air pollutants over Europe and North America in the framework of AQMEII3, Atmos. Chem. Phys., 18, 8929-8952, 10.5194/acp-18-8929-2018, 2018.

IMO: Fourth IMO GHG Study – Final Report, 2020.

IMO: Report of fuel oil consumption data submitted to the IMO Ship Fuel Oil Consumption Database in GISIS (Reporting year: 2023), MEPC 82/6/38, 2024.

Jalkanen, J. P., Brink, A., Kalli, J., Pettersson, H., Kukkonen, J., and Stipa, T.: A modelling system for the exhaust emissions of marine traffic and its application in the Baltic Sea area, Atmos. Chem. Phys., 9, 9209-9223, DOI 10.5194/acp-9-9209-2009, 2009.

Ji, C. and El-Halwagi, M. M.: A data-driven study of IMO compliant fuel emissions with consideration of black carbon aerosols, Ocean Engineering, 218, 108241, https://doi.org/10.1016/j.oceaneng.2020.108241, 2020.

Johansson, L., Jalkanen, J.-P., and Kukkonen, J.: Global assessment of shipping emissions in 2015 on a high spatial and temporal resolution, Atmospheric Environment, 167, 403-415, https://doi.org/10.1016/j.atmosenv.2017.08.042, 2017.

Kramel, D., Muri, H., Kim, Y., Lonka, R., Nielsen, J. B., Ringvold, A. L., Bouman, E. A., Steen, S., and Strømman, A. H.: Global Shipping Emissions from a Well-to-Wake Perspective: The MariTEAM Model, Environmental Science & Technology, 55, 15040-15050, 10.1021/acs.est.1c03937, 2021.

Kuittinen, N., Timonen, H., Karjalainen, P., Murtonen, T., Vesala, H., Bloss, M., Honkanen, M., Lehtoranta, K., Aakko-Saksa, P., and Rönkkö, T.: In-depth characterization of exhaust particles performed on-board a modern cruise ship applying a scrubber, Science of The Total Environment, 946, 174052, https://doi.org/10.1016/j.scitotenv.2024.174052, 2024.

Kuittinen, N., Jalkanen, J.-P., Alanen, J., Ntziachristos, L., Hannuniemi, H., Johansson, L., Karjalainen, P., Saukko, E., Isotalo, M., Aakko-Saksa, P., Lehtoranta, K., Keskinen, J., Simonen, P., Saarikoski, S., Asmi, E., Laurila, T., Hillamo, R., Mylläri, F., Lihavainen, H., Timonen, H., and Rönkkö, T.: Shipping Remains a Globally Significant Source of Anthropogenic PN Emissions Even after 2020 Sulfur Regulation, Environmental Science & Technology, 55, 129-138, 10.1021/acs.est.0c03627, 2021.

Li, Z., Fei, J., Du, Y., Ong, K.-L., and Arisian, S.: A near real-time carbon accounting framework for the decarbonization of maritime transport, Transportation Research Part E: Logistics and Transportation Review, 191, 103724, https://doi.org/10.1016/j.tre.2024.103724, 2024.

Liu, H., Fu, M., Jin, X., Shang, Y., Shindell, D., Faluvegi, G., Shindell, C., and He, K.: Health and climate impacts of ocean-going vessels in East Asia, Nat. Clim. Chang., 6, 1037-1041, 10.1038/nclimate3083, 2016.

Liu, H., Yi, W., Jalkanen, J.-P., Luo, Z., Majamäki, E., Matthias, V., Moldanová, J., Shi, Z., and He, K.: Atmospheric impacts and regulation framework of shipping emissions: achievements, challenges and frontiers, Fundamental Research, https://doi.org/10.1016/j.fmre.2024.02.013, 2024.

Liu, H., Meng, Z.-H., Lv, Z.-F., Wang, X.-T., Deng, F.-Y., Liu, Y., Zhang, Y.-N., Shi, M.-S., Zhang, Q., and He, K.-B.: Emissions and health impacts from global shipping embodied in US–China bilateral trade, Nat. Sustain., 2, 1027-1033, 10.1038/s41893-019-0414-z, 2019.

Lunde Hermansson, A., Hassellöv, I.-M., Grönholm, T., Jalkanen, J.-P., Fridell, E., Parsmo, R., Hassellöv, J., and Ytreberg, E.: Strong economic incentives of ship scrubbers promoting pollution, Nature Sustainability, 7, 812-822, 10.1038/s41893-024-01347-1, 2024.

Luo, Z., Lv, Z., Zhao, J., Sun, H., He, T., Yi, W., Zhang, Z., He, K., and Liu, H.: Shipping-related pollution decreased but mortality increased in Chinese port cities, Nat. Cities, 1, 295-304, 10.1038/s44284-024-00050-8, 2024.

Raman, R., Jethava, G., Chandrasekaran, H., and Tiwari, V.: Utilizing AIS Data and Machine Learning Techniques for Ship Type Classification, 2025 International Conference on Knowledge Engineering and Communication Systems (ICKECS), 28-29 April 2025, 1-6, 10.1109/ICKECS65700.2025.11034805,

Schwarzkopf, D. A., Petrik, R., Matthias, V., Quante, M., Majamäki, E., and Jalkanen, J.-P.: A ship emission modeling system with scenario capabilities, Atmospheric Environment: X, 12, 100132, https://doi.org/10.1016/j.aeaoa.2021.100132, 2021.

Seppälä, S. D., Kuula, J., Hyvärinen, A. P., Saarikoski, S., Rönkkö, T., Keskinen, J., Jalkanen, J. P., and Timonen, H.: Effects of marine fuel sulfur restrictions on particle number concentrations and size distributions in ship plumes in the Baltic Sea, Atmos. Chem. Phys., 21, 3215-3234, 10.5194/acp-21-3215-2021, 2021.

Sharmina, M., Y., E. O., C., W., R., F., J., G. D. E. H., P., G., A., L., W., L. E., M., T., P., v. V. D., E., V. N., R., W. F., and and Le Quéré, C.: Decarbonising the critical sectors of aviation, shipping, road freight and industry to limit warming to 1.5–2°C, Climate Policy, 21, 455-474, 10.1080/14693062.2020.1831430, 2021.

Song, S.-K., Shon, Z.-H., Moon, S.-H., Lee, T.-H., Kim, H.-S., Kang, S.-H., Park, G.-H., and Yoo, E.-C.: Impact of international Maritime Organization 2020 sulfur content regulations on port air quality at international hub port, Journal of Cleaner Production, 347, 131298, https://doi.org/10.1016/j.jclepro.2022.131298, 2022.

Starcrest Consulting Group, L.: PORT OF LOS ANGELES Inventory of Air Emission 2021, 2022.

Viana, M., Fann, N., Tobías, A., Querol, X., Rojas-Rueda, D., Plaza, A., Aynos, G., Conde, J. A., Fernández, L., and Fernández, C.: Environmental and Health Benefits from Designating the Marmara Sea and the Turkish Straits as an Emission Control Area (ECA), Environmental Science & Technology, 49, 3304-3313, 10.1021/es5049946, 2015.
Wang, X., Yi, W., Lv, Z., Deng, F., Zheng, S., Xu, H., Zhao, J., Liu, H., and He, K.: Ship emissions around China under gradually promoted control policies from 2016 to 2019, Atmos. Chem. Phys., 21, 13835-13853, 10.5194/acp-21-13835-2021, 2021.
Wang, X., Liu, H., Zhang, J., Fu, X., Chen, D., Zhang, W., Yi, W., Lv, Z., Zhang, Q., and He, K.: Global shipping emissions from 1970 to 2021: Structural and spatial change driven by trade dynamics, One Earth, 10.1016/j.oneear.2025.101243, 2025.
Xu, Y., Huang, Z., Ye, J., and Zheng, J.: Hourly emissions of air pollutants and greenhouse gases from open biomass burning in China during 2016–2020, Scientific Data, 10, 629, 10.1038/s41597-023-02541-0, 2023.
Yang, J., Tang, T., Jiang, Y., Karavalakis, G., Durbin, T. D., Wayne Miller, J., Cocker, D. R., and Johnson, K. C.: Controlling emissions from an ocean-going container vessel with a wet scrubber system, Fuel, 304, 121323, https://doi.org/10.1016/j.fuel.2021.121323, 2021.
Yang, T., Wang, X., and Liu, Z.: Ship Type Recognition Based on Ship Navigating Trajectory and Convolutional Neural Network, Journal of Marine Science and Engineering, 10, 84, 2022.
Yang, Y., Liu, Y., Li, G., Zhang, Z., and Liu, Y.: Harnessing the power of Machine learning for AIS Data-Driven maritime Research: A comprehensive review, Transportation Research Part E: Logistics and Transportation Review, 183, 103426, https://doi.org/10.1016/j.tre.2024.103426, 2024.
Yi, W., Wang, X., He, T., Liu, H., Luo, Z., Lv, Z., and He, K.: High-resolution global shipping emission inventory by Shipping Emission Inventory Model (SEIM), Earth Syst. Sci. Data Discuss., 2024, 1-31, 10.5194/essd-2024-258, 2024a.
Yi, W., He, T., Wang, X., Soo, Y. H., Luo, Z., Xie, Y., Peng, X., Zhang, W., Wang, Y., Lv, Z., He, K., and Liu, H.: Ship emission variations during the COVID-19 from global and continental perspectives, Science of The Total Environment, 176633, https://doi.org/10.1016/j.scitotenv.2024.176633, 2024b.
Yuan, T., Song, H., Oreopoulos, L., Wood, R., Bian, H., Breen, K., Chin, M., Yu, H., Barahona, D., Meyer, K., and Platnick, S.: Abrupt reduction in shipping emission as an inadvertent geoengineering termination shock produces substantial radiative warming, Communications Earth & Environment, 5, 281, 10.1038/s43247-024-01442-3, 2024.
Zhang, J., Chen, Y.-S., Gryspeerdt, E., Yamaguchi, T., and Feingold, G.: Radiative forcing from the 2020 shipping fuel regulation is large but hard to detect, Communications Earth & Environment, 6, 18, 10.1038/s43247-024-01911-9, 2025a.
Zhang, W., Yi, W., He, T., Peng, X., Xing, Q., Wu, X., Yang, Z., Xie, R., Xian, A., Liu, H., and Hao, J.: Ship emissions and reduction potential under domestic policies and international cooperation: A case study in Hainan, China, Journal of Environmental Sciences, 156, 253-266, https://doi.org/10.1016/j.jes.2025.02.006, 2025b.
Zhang, Y., Eastham, S. D., Lau, A. K. H., Fung, J. C. H., and Selin, N. E.: Global air quality and health impacts of domestic and international shipping, Environ. Res. Lett., 16, 084055, 10.1088/1748-9326/ac146b, 2021.

设置了格式: 字体: (默认) +西文正文 (Times New Roman)

第 3 页: [1] 删除了 唯唯 张 2026/8/23 15:44:00

第 3 页: [2] 删除了 唯唯 张 2026/8/23 15:43:00

第 5 页: [3] 删除了 唯唯 张 2026/8/23 14:06:00

第 5 页: [4] 删除了 唯唯 张 2026/8/23 14:08:00

第 9 页: [5] 删除了 唯唯 张 2026/8/18 13:34:00

第 9 页: [6] 删除了 唯唯 张 2026/8/22 21:52:00

第 13 页: [7] 删除了 唯唯 张 2026/8/23 16:26:00

第 15 页: [8] 删除了 唯唯 张 2026/8/23 15:10:00