

Opinion: Status, Plans and Needs of Southern Ocean Modelling

Torge Martin¹, Carolina O. Dufour², Andrew J. S. Meijers³ and Alyce M. Hancock⁴

¹ GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

² University of Brest, CNRS, Ifremer, IRD, Laboratoire d'Océanographie Physique et Spatiale (LOPS), IUEM, F29280, Plouzané, France

³ British Antarctic Survey, Cambridge, United Kingdom

⁴ Southern Ocean Observing System, International Project Office, Institute for Marine and Antarctic Studies, University of Tasmania, Australia

ORCID:

T. Martin: 0000-0002-0882-8780

C.O. Dufour: 0000-0002-1441-3880

A.J.S. Meijers: 0000-0003-3876-7736

A.M. Hancock: 0000-0001-6049-5592

Correspondence to: Torge Martin (tomartin@geomar.de)

Abstract. In preparation for the *SOOS/OCEAN:ICE Workshop on ice-ocean observation harmonization and future priorities agenda*, a survey targeting the modelling community was conducted to assess research priorities for the Southern Ocean and Antarctica. While this initiative was tailored mostly towards physical ocean and ice modelling, its outcome specifically supports the design of field activities from the open Southern Ocean to the Antarctic shelf for the forthcoming *Antarctica InSync* campaign and is aligned with broader strategic planning efforts ahead of the next International Polar Year (IPY). The survey results are a useful basis to further communication between modeling and observing science communities. We believe this is crucial for optimizing campaign planning, achieving enhanced data usage and improving numerical experiments.

1 Background

The Southern Ocean is responsible for 83% of the global ocean heat content increase over the historical period and more than 40% of global ocean anthropogenic carbon uptake (Frölicher et al., 2015; Williams et al., 2024). Ocean heat is a major driver of the current Antarctic ice sheet mass imbalance (Adusumilli et al., 2020; Bell & Seroussi, 2020; Noble et al., 2020). Nevertheless, this part of the world ocean features some of the most severe and long-standing biases present in state-of-the-art climate models with far reaching implications for climate projections (Stouffer et al., 2017; Beadling et al., 2020; Moreno-Chamarro et al., 2022; Zhang et al., 2023).

Full understanding of the processes and feedbacks of climate change in the Southern Ocean and Antarctica can only be gained by combining in-situ observations, satellite reconnaissance and numerical modelling. The region is difficult to access, especially where ice covered, and direct observations thus remain sparse. Remote sensing is limited to the surface, at least for the ocean. And models are valuable tools but never perfect. The upcoming Antarctica InSync and the IPY programs are centered around field campaigns, which is natural considering the need for strategically planning ship schedules and equipment acquisition years in advance.

50 Satellite missions and model development follow again different cycles and routines, partly
51 on even longer schedules. Dedicated numerical experiments can be accomplished on a much
52 shorter time scale though. Nevertheless, coordinating efforts across these science
53 communities and intensifying exchange between them from early on in these major programs
54 will be crucial for turning advanced process understanding into improved projections of the
55 future climate in and beyond the southern high latitudes. Therefore, the survey initiative
56 documented here supports a push for early integration of and engagement by the modeling
57 community in these observation-driven efforts.

58

59 As a rather spontaneous effort, the survey was of ad hoc design. Despite being launched just
60 before the northern hemisphere summer break, it received a relatively large number of
61 completed responses (98), representing a broad cross-section of the ocean and climate
62 modeling community. This great turnout is also owed to the endorsement by [SOOS](https://www.soos.aq/)
63 (<https://www.soos.aq/>), the Southern Ocean Observing System and [CLIVAR](https://www.clivar.org/)
64 (<https://www.clivar.org/>), Climate and Ocean Variability, Predictability, and Change offices
65 and Scientific Steering Groups, who spread the call in the SOOS Update (Issue 31) and the
66 CLIVAR Bulletin, respectively, in August 2025 as well as the sharing of the call across
67 mailing lists of [Antarctica InSync](https://www.antarctica-insync.org/) (<https://www.antarctica-insync.org/>), [APECS](https://www.apecs.is/)
68 (<https://www.apecs.is/>), [ASPeCt](https://aspectsouth.org/) (<https://aspectsouth.org/>), [BEPSII](https://sites.google.com/site/bepsiiwg140/home)
69 (<https://sites.google.com/site/bepsiiwg140/home>), [BioEcoOcean](https://bioecocean.org/) (<https://bioecocean.org/>),
70 [Polar-CORDEX](https://climate-cryosphere.org/polar-cordex/about/) (<https://climate-cryosphere.org/polar-cordex/about/>), [Cryolist](https://lists.cryolist.org/mailman/listinfo/cryolist)
71 (<https://lists.cryolist.org/mailman/listinfo/cryolist>), the [EU Polar Cluster](https://polarcluster.eu/)
72 (<https://polarcluster.eu/>), [ICED](https://www.iced.ac.uk/) (<https://www.iced.ac.uk/>), [IMBeR](https://imber.info/) (<https://imber.info/>),
73 [IMECaN](http://imecan/) (<http://imecan/>), [MISOMIP2](https://misomip.github.io/misomip2/) (<https://misomip.github.io/misomip2/>), [ObsSea4Clim](https://obssea4clim.eu/)
74 (<https://obssea4clim.eu/>), [Ocean & Carbon Biogeochemistry](https://www.us-ocb.org/) (<https://www.us-ocb.org/>),
75 [OCEAN:ICE](https://ocean-ice.eu/) (<https://ocean-ice.eu/>), CLIVAR's [Ocean Modeling Development Panel](https://www.clivar.org/clivar-panels/omdp)
76 (<https://www.clivar.org/clivar-panels/omdp>) and [Southern Ocean Region Panel](https://www.clivar.org/clivar-panels/southern)
77 (<https://www.clivar.org/clivar-panels/southern>), [POGO](https://pogo-ocean.org/) (<https://pogo-ocean.org/>), [SCAR](https://scar.org/)
78 (<https://scar.org/>), [SCOR](https://scor-int.org/) (<https://scor-int.org/>), [SOCCOM](https://socom.org/) (<https://socom.org/>), the
79 CLIVAR task team and Community-MIP [SOFIA](https://sofiamip.github.io/) (<https://sofiamip.github.io/>), and [TipESM](https://tipesm.eu/)
80 (<https://tipesm.eu/>). This positive engagement yielded a valuable and unprecedented dataset
81 that offers quantitative insights into current priorities and gaps in Southern Ocean research
82 and modeling. It provides a robust foundation for ongoing and future strategic discussions
83 regarding the alignment of modeling and observational efforts.

84

85 The survey was designed with primarily the physical ocean modeling community in mind and
86 thus pre-defined answers often highlight physical oceanographic processes (see Appendix).
87 Although leaning towards such processes, coupled interactions with other components of the
88 climate system, such as sea ice, ice shelves, and the atmosphere were considered as well.
89 Contributions related to biogeochemical processes and ecosystem modeling were also
90 encouraged though not covered comprehensively. It turns out that a more careful selection of
91 the pre-defined answers would have been advantageous and likely beneficial for a broader
92 coverage of these coupled processes. Relatively little use was made of the “Other” free text
93 option by the respondents. Similarly, the survey started out with an emphasis on probing
94 modelling groups using realistic regional Southern Ocean configurations and CMIP-class
95 global climate and Earth system models to study the historical period and maybe 21st century
96 projections. But it evolved into covering a much broader range of spatial and temporal scales,
97 and model complexities, for which the respondents did make use of the free text comment
98 fields. For pragmatic reasons, we limited the geographical region and defined the Southern
99 Ocean as the area south of approximately 50°S in the survey context.

100
101 The following summary presents key findings of the survey. While the [dataset](https://doi.org/10.5281/zenodo.17289776)
102 (<https://doi.org/10.5281/zenodo.17289776>) (Martin et al., 2025) can be further explored, a
103 first look already holds significant potential for informing cross-disciplinary planning and
104 collaborative program development in the Southern Ocean and Antarctic research landscape.
105

106 **2 Who participated?**


107
108 About half of the 98 survey participants [48%] identified as oceanographers, others see
109 themselves as experts in coupled climate [13%], sea ice [12%], ice shelf cavity [10%] and
110 land ice [8.2%] modelling. Colleagues studying processes at basin to global scales and from
111 annual to centennial scales contributed two-thirds of the replies; less than a quarter indicated
112 a research focus on mesoscale (10-100 km) processes with periods of months to seasons.
113

114 **3 Survey results and discussion**

116 **3.1 Model status and evolution**

117
118 We asked the participants for the most problematic ocean model bias allowing a single choice
119 only aiming for a clear emergence of the most pressing issue from the survey. About a
120 quarter identified processes of the Antarctic continental margin (shelf seas, slope current, ice
121 shelf cavities) as requiring most attention (Figure 1). Further, open ocean deep convection
122 and water mass transformation [both 14.3%] and mixing were highlighted. Individual free
123 text answers mentioned deep ocean circulation, modeling of biogeochemical cycles and the
124 carbon pump, planetary boundary layer of ice-covered seas, and impacts on benthic
125 ecosystems as other major model biases, which are not displayed in Figure 1. The
126 respondents related such biases in particular to global coupled climate [38%] and ocean
127 models [28%] in general such as those used for the Climate Modelling Intercomparison
128 Project (CMIP); this may reflect the dominant area of expertise of the participants however.
129 Moreover, the respondents identified a dozen specific ocean models and state estimates as
130 well as specifically high resolution model versions that include some of these major biases
131 (see published survey data for details; Martin et al., 2025). We emphasize that moving
132 towards finer grid resolution alone may yield individual improvements but will not solve all
133 the biases—as is documented in Moreno-Chamarro et al. (2022).
134

ocean model biases by expert group




135
 136 *Figure 1: “What is the most problematic ocean model bias?” Eight well-known issues were listed as pre-*
 137 *defined, single-choice answers to ensure emergence of the most pressing problems. In this distribution of the*
 138 *responses we merged seasonality of surface fluxes and air-sea CO₂ flux into surface fluxes. The 91 responses (of*
 139 *98 in total) are color-coded with respect to the area of expertise provided by the respective respondents. Here,*
 140 *“other” refers to all areas of expertise not explicitly listed, e.g. atmosphere and ecosystem.*

141
 142 It is important to understand that open ocean deep convection—while physically not
 143 unrealistic (Gordon, 1978)—is dramatically overestimated in many climate models,
 144 specifically those of coarse resolution, with consequences for ocean to atmosphere heat
 145 redistribution, sea ice coverage, bottom water characteristics, and eventually also internal
 146 climate variability (Reintges et al., 2017; Heuzé et al., 2021). In reality, Antarctic bottom
 147 water (AABW) is formed on the continental shelf where sea ice formation and ice shelf melt
 148 play key roles in the transformation of upwelled deep water (Silvano et al., 2023) and so does
 149 mixing for its transformation into Antarctic intermediate water (AAIW) further equatorward
 150 (Li et al., 2022). To this end, nearly three-quarters of all participating experts pointed out
 151 biases that are inter-connected and play an imminent role in the formation of water masses,
 152 such as AABW and AAIW, crucial for the global overturning circulation and for the natural
 153 sequestration of heat and anthropogenic carbon.

154
 155 Hence, it is no surprise that implementation of ice shelf cavities [20%], convection
 156 parameterization [18%], scale aware (mixing) parameterizations [15%] and overflow
 157 parameterization [13%] were listed as most urgent model development targets. There has
 158 been remarkable advancement in these directions over the past two decades (e.g., De Rydt et
 159 al., 2024; Legg et al., 2009; Bruciaferri et al., 2024). And it has been demonstrated that these
 160 new developments can mitigate model biases in the Southern Ocean even at relatively coarse
 161 resolution despite remaining issues (e.g. for ice shelf cavities see Hutchinson et al., 2023).
 162 However, we believe there is often significant delay or inaction in implementing such
 163 advancements, as model development is rarely funded directly by dedicated research projects.
 164 The push for rapid research outcomes tends to favor easily implemented targets — the 'low-
 165 hanging fruit' — over more complex, long-term efforts.

166



167
 168 *Figure 2: “What is in your view the singular key science topics in the Southern Ocean?” The five most frequent*
 169 *answers were provided as part of seven examples and could be simply ticked. Pre-defined answers less picked*
 170 *were air-sea exchange and extreme events. Additional topics were given by the respondents as free text input.*
 171 *These include, amongst others, air-sea exchange, cloud-radiation processes, extreme events, nutrient*
 172 *redistribution and cross-disciplinary topics, and are collated as ‘other’.*
 173

174 This behavior is evident in the responses on near-term model evolution. Among those,
 175 increasing model complexity [27%] and spatial resolution [24%] stand out. Other goals such
 176 as improving or developing novel parameterizations and including artificial intelligence
 177 based modules are only considered by 11-13% of the participants. In this case multiple
 178 answers were possible and participants ticked or listed 2-3 responses on average. It appears
 179 that preference is given to model complexity—evolving climate models into Earth system
 180 models by coupling more components, for example, ice sheets or biogeochemistry modules—
 181 over improving model physics. However, we think this could also be a sign of a more
 182 diversified, cross-disciplinary science landscape. As compute power keeps growing,
 183 resolving model issues by enhancing grid resolution appears to be a possible avenue to reduce
 184 biases (e.g., Rackow et al., 2022). But this is a costly option and impractical for applications
 185 on centennial time scales since proper representation of mesoscale dynamics in the high
 186 latitudes of the Southern Ocean requires grid spacing of $1/8^{\circ}$ - $1/20^{\circ}$ (and finer on the
 187 continental shelf), to properly resolve the Rossby radius (Hallberg, 2013, their Fig. 1). We
 188 suggest that new observations supporting model development could thus lead to improved
 189 and yet affordable simulations on a large range of spatial and temporal scales.
 190

191 3.2 Scientific focus

192
 193 Reducing the major model biases and advancing ocean and climate models as laid out above
 194 will be essential to address the key research topics identified in the survey responses.
 195 Freshwater, heat and carbon budgets are high on the scientific agenda of the modelling
 196 community (Figure 2). Questions on process understanding and future evolution of the
 197 Antarctic ice sheet, its ice shelves and their interaction with the ocean through heat and
 198 meltwater dominate the results [40%]. While this is research at the continental margin, heat
 199 and carbon uptake where the low latitude Southern Ocean plays a major role was named
 200 second [19%], followed by interest in the recent and future sea ice trends [14%]. On the one
 201 hand, these results are somewhat biased by the research areas of the participants. On the other
 202 hand, scientific interest has migrated poleward in the Southern Ocean, where major
 203 challenges have been identified, such as knowledge gaps in ice-ocean interaction affecting

204 global sea level rise projections, and where new observational techniques for under-ice
205 sampling and mesoscale ocean simulations have become available.

206

207 The results suggest that oceanic processes themselves, such as dynamics from mesoscale
208 eddies to large-scale circulation, tides, waves and mixing are not “big questions” by
209 themselves anymore despite remaining issues and their important role in current “grand
210 challenges” like ice-ocean interaction, warm water intrusion onto the continental shelf and
211 biogeochemical modeling. Further, we assume that the underrepresentation of
212 biogeochemical and ecosystem research as well as atmospheric process understanding, most
213 prominently clouds and aerosols, is likely a consequence of questionnaire design and the
214 focus group addressed.

215

216 **3.3 Observations used and needed**

217


218 Before diving into a discussion on the observational needs of the modelling community, we
219 would like to call attention to the data sources actually used. There is an unbroken preference
220 by modellers to use gridded data products [28%], i.e. statistically interpolated fusions of
221 observations from various sources, and reanalysis or state estimates [21%], which are based
222 on a numerical model and incorporate observations through, e.g. assimilation techniques.
223 Likely also due to their extensive spatial coverage, satellite-borne remote sensing products
224 are favored as well [15%]. Data from ship-borne instruments, moorings, and floats appear to
225 be used less often directly [10-13%]. We acknowledge that observational data of all kinds, in
226 particular including in-situ data, feed into the gridded products though. Nevertheless, it is
227 important to note that modelers tend to validate their simulations against these kind of
228 “observations”, which in fact are advanced data products and rather not viewed as actual
229 observations by the observing, sea-going science community. Further, any in-situ data that is
230 not included in such gridded products is likely less used or even overlooked by modelers and
231 thus does not contribute as much to the improvement of models.

232

233 Based on our personal experience, modelers tend to lean on derived products, such as
234 reanalyses and state estimates, because (1) observations are typically sparse and direct
235 comparisons require complex subsampling of model output, (2) formats and platforms used
236 to share observational data are still not unified and easily accessible to users despite ongoing
237 efforts, and (3) original measurements require advanced background knowledge and
238 interpretation for use in model validation. These issues can be mitigated by, for instance,
239 providing open-source validation packages along with the measurements and the
240 development of virtual instruments, i.e. tool boxes allowing for the subsampling of model
241 output in ways emulating specific observational instruments. Further, we suggest that
242 communication between the science communities should be strengthened and educational
243 programs, such as summer schools, could introduce instruments, proper data handling and
244 aforementioned tool boxes to the next generation of modelers.

245

246



247
248
249
250
251
252
253
254
255
256
257

Figure 3: (a) “Which kind of observations would further this [model] development?” (without pre-defined answers). Free text responses were grouped by key words (right hand side bar labels) and sorted into groups of temporal coverage, instrumentation/sensors/platform and Earth system components (left hand side labels). In total 130 responses were identified. (b) Linking the two questions “What is the most problematic ocean model bias?” and “Which observations could help understanding biases or further the process understanding?” by the same respondent. In both cases pre-defined answers were provided but free text replies were also possible; multiple choice was allowed. Overall 200 responses were cross-linked. The number of responses on the left side are normalized, see Fig. 1 for specific numbers.

258
259
260
261
262
263
264
265
266
267
268
269

For reducing observational gaps, especially with mitigating model biases in mind, modelers should be given and take the opportunity to provide input to evolving observational programs early in the planning phase. This would ensure multiple use of the data collected in the end. Figure 3a shows observations desired for bias mitigation with in-situ observations clearly dominating over remote sensing data. This preference can be interpreted as a need for better process understanding in which in-situ data are considered actual “ground truth” and often provide higher resolution in space and time, which is useful in several aspects, such as model validation, identifying of processes resolved at a given grid spacing, improving model parameterizations, etc. Another interpretation is that modellers are well aware of in-situ measurements being crucial for better quality gridded products. And while such products are preferred in the actual validation process, the dire need for more ground truth data in a changing climate is acknowledged and its collection valued.

270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288

Moreover, the scientific goals and observational plans of Antarctica InSync appear to be very much in line with the needs of the modelling community wishing for year-round data especially in ice-covered seas and combining physical and biogeochemical measurements (Figure 3a). The strong desire for winter observations and year-round monitoring in the Southern Ocean [29%] is linked nearly equally to all major biases identified (Figure 3b). As is already discussed by the Antarctica InSync community, building capacity for sustained monitoring in preparation for the International Polar Year in 2032/33 and beyond would also strongly support model improvement and advancement [25%]. Observations of the upper ocean stratification and surface fluxes would enable a better process understanding of the mixed layer and help to constrain vertical/diapycnal mixing parameterizations in models. Interestingly, the role of other climate system components causing biases in the ocean, for example, sea ice and snow, clouds and radiative processes, was not highly considered. This is likely owed to the limited choice of pre-defined answers we provided and the behavioral bias of the respondents preferring to tick one of those rather than entering individual answers. Nevertheless, we take this as an opportunity to point out the need for improved fundamental understanding and acknowledgement of coupled mechanisms and feedbacks within and beyond the focus group. Having better records of magnitude and variability of Southern Ocean surface fluxes of both physical and chemical quantities will help this significantly.

289 Similarly, any observations in support of heat and freshwater budgets especially with a focus
290 on ice-ocean interaction will be instrumental in advancing models and improving climate
291 projections. And last but not least, high-resolution bathymetry data of the Southern Ocean
292 from the Antarctic Circumpolar Current to the continental margin and into the ice shelf
293 cavities is urgently needed. Topography is a key ingredient for realistic simulations of the
294 ocean circulation, specifically the import of warm deep water and the export of dense bottom
295 waters, and therefore crucial for reliable projections of ice shelf melting.

296 **4 Conclusions and Outlook**

297 In conclusion, surveys like this provide a valuable overview of the current status, plans, and
298 data needs not only for the Southern Ocean but also for the global modeling community.
299 There are also some experiences to take away from this exercise, in particular how the pool
300 of respondents shapes the usefulness of the survey, how to ask targeted questions without
301 being exclusive, how pre-defined multiple-choice answers simplify the analysis but reduce
302 the variety of responses, and which meta-information really is instrumental for interpreting the
303 responses. All of this is well known and demonstrates the importance to involve experts on
304 questionnaire design rather than constructing an ad hoc survey. Nevertheless, the feedback by
305 the community to our survey has been very positive indicating that such surveys can be a
306 valuable tool for future international program planning.

307 With the Antarctica InSync program in active planning and IPY approaching, we hope the
308 results presented—with additional data available (Martin et al., 2025)—will inform both the
309 scientific community and stakeholders to advance observations and models. The findings
310 already contributed to the *SOOS/OCEAN:ICE Workshop* discussions and conclusions.
311 Research priorities include ice–ocean interactions, Southern Ocean heat and carbon uptake,
312 and the recent major changes in sea ice. Addressing these challenges requires model
313 developments such as ice-shelf–ocean coupling, biogeochemistry, and higher resolution,
314 alongside improved understanding of continental shelf processes and upper-ocean
315 stratification. This, in turn, requires new observations in key regions with ice shelves most
316 vulnerable to warm water intrusions and ocean circulation choke points. Further, the survey
317 results call for stronger communication between the modeling and observing communities
318 and dedicated data-use training for modelers. Antarctica InSync offers a major opportunity to
319 advance such efforts.

320 **Data availability**

321 The survey results are available through Zenodo (Martin et al., 2025),
322 <https://doi.org/10.5281/zenodo.17289777>.

323 **Author Contributions**

324 TM conceived the original idea of conducting the survey, led the analysis and produced all
325 figures. All authors contributed to the survey design, its content and writing of this
326 manuscript. AMH explored different survey platforms and distributed the survey call.

327 **Competing Interests**

328 The authors declare to have no competing interests.

329
330
331
332
333
334
335
336
337
338

339 **Acknowledgement**

340 The authors thank three anonymous reviewers for their positive, useful comments leading to a
341 more careful interpretation of the survey. We specifically thank SOOS and CLIVAR for their
342 support and endorsement of the survey facilitating this study. We are grateful to all the
343 scientific networks and coordinators for widely spreading the survey call and all the
344 participants for responding yielding such great turnout. Helpful comments on the
345 questionnaire design by Chris Danek, Ivy Frenger, Alexander Haumann, Qian Li, Lavinia
346 Patara, Ariaan Purich, David Stevens were much appreciated. The survey was conducted
347 using Google Forms and we made use of the initial visualization of the statistics provided by
348 this platform in material provided by Martin et al. (2025). AJSM was supported by Ocean
349 Cryosphere Exchanges in Antarctica: Impacts on Climate and the Earth system, OCEAN
350 ICE, which is funded by the European Union, Horizon Europe Funding Programme for
351 research and innovation under grant agreement Nr. 101060452, 10.3030/101060452; OCEAN
352 ICE contribution number 50; he was also funded by UK Research and Innovation (UKRI)
353 under the UK government's Horizon Europe funding Guarantee [grant number 10048443].
354 SOOS is a joint initiative of the Scientific Committee on Oceanic Research (SCOR) and the
355 Scientific Committee on Antarctic Research (SCAR). SOOS sponsors include SCAR and
356 SCOR, the Tasmanian State Government, CSIRO, the Australian Antarctica Program
357 Partnership (AAPP), the Alfred Wegener Institute (AWI), OCEAN:ICE, the Swedish Polar
358 Research Secretariat and the TÜBİTAK MAM Polar Research Institute. The SOOS
359 International Project Office is hosted by the Institute for Marine and Antarctica Studies,
360 University of Tasmania.

361

362 **References**

363 Adusumilli, S., H. A. Fricker, B. Medley, L. Padman & M. R. Siegfried (2020). Interannual
364 variations in meltwater input to the Southern Ocean from Antarctic ice shelves. *Nat.*
365 *Geosci.* 13, 616–620, <https://doi.org/10.1038/s41561-020-0616-z>.

366 Beadling, R. L., Russell, J. L., Stouffer, R. J., Mazloff, M., Talley, L. D., Goodman, P. J.,
367 Sallée, J. B., Hewitt, H. T., Hyder, P., & Pandde, A. (2020). Representation of Southern
368 Ocean properties across coupled model intercomparison project generations: CMIP3 to
369 CMIP6. *J. Climate*, 33, 6555–6581, <https://doi.org/10.1175/JCLI-D-19-0970.1>.

370 Bell, R. E. & H. Seroussi (2020). History, mass loss, structure, and dynamic behavior of the
371 Antarctic Ice Sheet. *Science* 367, 1321–1325, <https://doi.org/10.1126/science.aaz5489>.

372 Bruciaferri, D., Guiavarc'h, C., Hewitt, H. T., Harle, J., Almansi, M., Mathiot, P., &
373 Colombo, P. (2024). Localized general vertical coordinates for quasi-Eulerian ocean
374 models: The Nordic overflows test-case. *Journal of Advances in Modeling Earth Systems*,
375 16, e2023MS003893, <https://doi.org/10.1029/2023MS003893>.

376 De Rydt, J., Jourdain, N. C., Nakayama, Y., van Caspel, M., Timmermann, R., Mathiot, P.,
377 Asay-Davis, X. S., Seroussi, H., Dutrieux, P., Galton-Fenzi, B., Holland, D., & Reese, R.
378 (2024). Experimental design for the Marine Ice Sheet–Ocean Model Intercomparison
379 Project – phase 2 (MISOMIP2), *Geosci. Model Dev.*, 17, 7105–7139,
380 <https://doi.org/10.5194/gmd-17-7105-2024>.

381 Frölicher, T. L., J. L. Sarmiento, D. J. Paynter, J. P. Dunne, J. P. Krasting, & M. Winton
382 (2015). Dominance of the Southern Ocean in Anthropogenic Carbon and Heat Uptake in
383 CMIP5 Models. *J. Climate*, 28, 862–886, <https://doi.org/10.1175/JCLI-D-14-00117.1>.

384 Gordon AL (1978). Deep Antarctic convection west of Maud rise. *J. Phys. Oceanogr.*,
385 8(4):600–612, [https://doi.org/10.1175/1520-0485\(1978\)008<0600:DACWOM>2.0.CO;2](https://doi.org/10.1175/1520-0485(1978)008<0600:DACWOM>2.0.CO;2)

386 Hallberg, R. (2013). Using a resolution function to regulate parameterizations of oceanic
387 mesoscale eddy effects, *Ocean Modelling*, 72, 92-103,
388 <https://doi.org/10.1016/j.ocemod.2013.08.007>.

389 Heuzé, C. (2021). Antarctic Bottom Water and North Atlantic Deep Water in CMIP6 models,
390 *Ocean Sci.*, 17, 59–90, <https://doi.org/10.5194/os-17-59-2021>.

391 Hutchinson, K., Deshayes, J., Éthé, C., Rousset, C., de Lavergne, C., Vancoppenolle, M.,
392 Jourdain, N. C., & Mathiot, P. (2023). Improving Antarctic Bottom Water precursors in
393 NEMO for climate applications, *Geosci. Model Dev.*, 16, 3629–3650,
394 <https://doi.org/10.5194/gmd-16-3629-2023>.

395 Legg, S., B. Briegleb, Y. Chang, E. P. Chassignet, G. Danabasoglu et al. (2009). Improving
396 Oceanic Overflow Representation in Climate Models: The Gravity Current Entrainment
397 Climate Process Team. *Bull. Amer. Meteor. Soc.*, 90, 657–670,
398 <https://doi.org/10.1175/2008BAMS2667.1>.

399 Li, Z., S. Groeskamp, I. Cerovečki, and M. H. England (2022). The Origin and Fate of
400 Antarctic Intermediate Water in the Southern Ocean, *J. Phys. Oceanogr.*, 52, 2873–2890,
401 <https://doi.org/10.1175/JPO-D-21-0221.1>.

402 Martin, T., Dufour, C., Meijers, A., & Hancock, A. (2025). Southern Ocean modelling survey
403 (1.0) [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.17289777>.

404 Moreno-Chamarro, E., Caron, L.-P., Loosveldt Tomas, S., Vegas-Regidor, J., Gutjahr, O.,
405 Moine, M.-P., Putrasahan, D., Roberts, C. D., Roberts, M. J., Senan, R., Terray, L.,
406 Tourigny, E., & Vidale, P. L. (2022) Impact of increased resolution on long-standing biases
407 in HighResMIP-PRIMAVERA climate models, *Geosci. Model Dev.*, 15, 269–289,
408 <https://doi.org/10.5194/gmd-15-269-2022>.

409 Noble, T. L., Rohling, E. J., Aitken, A. R. A., Bostock, H. C., Chase, Z., Gomez, N., et al.
410 (2020). The Sensitivity of the Antarctic Ice Sheet to a Changing Climate: Past, Present, and
411 Future. *Reviews of Geophysics*, 58, e2019RG000663,
412 <https://doi.org/10.1029/2019RG000663>.

413 Rackow, T., S. Danilov, H. F. Goessling, H. H. Hellmer, D. V. Sein, T. Semmler, D.
414 Sidorenko & T. Jung (2022). Delayed Antarctic sea-ice decline in high-resolution climate
415 change simulations. *Nat Commun* 13, 637, <https://doi.org/10.1038/s41467-022-28259-y>.

416 Reintges, A., T. Martin, M. Latif, & W. Park (2017). Physical controls of Southern Ocean
417 deep-convection variability in CMIP5 models and the Kiel Climate Model, *Geophys. Res.*
418 *Let.*, 44, <https://doi.org/10.1002/2017GL074087>.

419 Silvano A, Purkey S, Gordon AL, Castagno P, Stewart AL, Rintoul SR, Foppert A, Gunn KL,
420 Herraiz-Borreguero L, Aoki S, Nakayama Y, Naveira Garabato AC, Spingys C, Akhoudas
421 CH, Sallée J-B, de Lavergne C, Abrahamsen EP, Meijers AJS, Meredith MP, Zhou S,
422 Tamura T, Yamazaki K, Ohshima KI, Falco P, Budillon G, Hattermann T, Janout MA,
423 Llanillo P, Bowen MM, Darelius E, Østerhus S, Nicholls KW, Stevens C, Fernandez D,
424 Cimoli L, Jacobs SS, Morrison AK, Hogg AM, Haumann FA, Mashayek A, Wang Z, Kerr
425 R, Williams GD and Lee WS (2023). Observing Antarctic Bottom Water in the Southern
426 Ocean. *Front. Mar. Sci.*, 10:1221701, <https://doi.org/10.3389/fmars.2023.1221701>.

427 Stouffer, R. J., V. Eyring, G. A. Meehl, S. Bony, C. Senior, B. Stevens, & K. E. Taylor
428 (2017). CMIP5 scientific gaps and recommendations for CMIP6. *Bull. Amer. Meteor. Soc.*,
429 98, 95–105, <https://doi.org/10.1175/BAMS-D-15-00013.1>.

430 Williams, R.G., Meijers, A.J.S., Roussenov, V.M. et al. (2024). Asymmetries in the Southern
431 Ocean contribution to global heat and carbon uptake. *Nat. Clim. Chang.* 14, 823–831,
432 <https://doi.org/10.1038/s41558-024-02066-3>.

433 Zhang, Q., Liu, B., Li, S., & Zhou, T. (2023). Understanding models' global sea surface
434 temperature bias in mean state: From CMIP5 to CMIP6. *Geophysical Research Letters*, 50,
435 e2022GL100888. <https://doi.org/10.1029/2022GL100888>.

436

437

438

439

440 **Appendix**

441 Here we repeat for convenience the survey introduction and questions along with the
442 response options. A screenshot of the original questionnaire website is available at
443 <https://doi.org/10.5281/zenodo.17289777>.

444

445 **Southern Ocean modelling: status, plans and needs**

446 In preparation of the *SOOS/OCEAN:ICE/Antarctica InSync workshop* on ice-ocean
447 observation harmonisation and future priorities agenda, which will be held in Copenhagen in
448 September this year and also serves for the planning of [Antarctica InSync](#) activities, we
449 kindly ask for your input.

450 All of the 12 questions below focus on the Southern Ocean region (south of ~50°S), physical
451 ocean processes are prioritised but immediate interaction with other system components can
452 be addressed (atmosphere, sea ice, ice shelves, biogeochemistry, ecosystem etc.), and the
453 applications we have in mind range from realistic regional models in the Southern Ocean to
454 CMIP-type global climate models (with or without further coupled components like ice
455 shelves or biogeochemistry).

456 **Please select from the multiple choice items or add an alternative item as "other".**
457 **This should not take more than 10 minutes.**

458 Note, the first three questions on expertise and scales may help to "set the stage" for the
459 answers of the other questions. If you work on very different scales, you may consider
460 answering the questionnaire twice for different cases. A link for another response is provided
461 after submitting the first one.

462 ***Please respond until September 15. (deadline extended)***

463 Thank you for your time and effort.

464 Torge Martin (OMDP), Carolina Dufour (SORP), Andrew Meijers (SOOS), Alyce Hancock
465 (SOOS)

466 *This questionnaire is supported by SOOS and CLIVAR panels SORP and OMDP. The*
467 *anonymous(!) data may be used by these panels for planning of further activities beyond the*
468 *above mentioned workshop.*

469 <https://www.soos.aq>, <https://www.clivar.org/clivar-panels/southern>,
470 <https://www.clivar.org/clivar-panels/omdp>

471

472 **1) What is your primary area of expertise?**

473 This is independent of a focus on physical, biological or other processes.

- 474 ○ Ocean
- 475 ○ Sea ice
- 476 ○ Ice-shelf cavity
- 477 ○ Ice sheet/ice shelf/icebergs
- 478 ○ Atmosphere
- 479 ○ Coupled climate modelling
- 480 ○ Other: _____

481

482 **2) What spatial scales do you work on?**

- 483 ○ Submesoscale (<10 km)
- 484 ○ Mesoscale (10-100 km)
- 485 ○ Basin-scale (>100 km)
- 486 ○ Global impacts

487

488 **3) What temporal scales do you work on?**

- 489 ○ Days to weeks
- 490 ○ Months to seasons
- 491 ○ Years to decades
- 492 ○ Decades to centuries
- 493 ○ Beyond centuries

494

495 **4) What is in your view the singular key science topic in the SO?**

- 496 ○ Ocean dynamics (sub-/mesoscale eddies to large-scale circulation)
- 497 ○ Ice sheet/shelf–ocean interaction, fresh water
- 498 ○ Heat and/or carbon uptake and redistribution
- 499 ○ Sea-ice change and drivers/consequences thereof
- 500 ○ Air-sea exchange, incl. carbon/trace gasses
- 501 ○ Marine biogeochemistry and ecosystems
- 502 ○ Extreme events
- 503 ○ Other: _____

504

505 **5) What kind of observational data do you preferably use?**

506 *[multiple answers allowed]*

- 507 ○ Campaign based data (e.g. ship transects, gliders, ...)
- 508 ○ Stationary, year-round data (e.g. moorings)
- 509 ○ Varied spatial and temporal coverage (e.g. Argo-Floats)
- 510 ○ Gridded data products integrating various data sources
- 511 ○ Reanalysis/ocean state estimate products
- 512 ○ Remote sensing data (please specify below)
- 513 ○ Other: _____

514

515 **If you selected “Remote sensing data” above, please specify the variable and**
516 **product here:**

517 _____

518

519 **6) What is the most problematic ocean model bias?**

520

○ Surface warm bias, too small sea ice extent

521

○ Deep warm bias

522

○ Mixing

523

○ Seasonality of surface fluxes

524

○ Air-sea CO₂ flux

525

○ Water mass formation and characteristics (AABW, AAIW, ...)

526

○ Open-ocean convection (polynyas)

527

○ Shelf processes (e.g. slope current, ice shelf cavities, overflows, ...)

528

○ Other: _____

529

530 **7) Which type of model did you have in mind when answering the last question**
531 **on biases?**

532

For example, list model complexity, grid resolution, application case, or anything else.

533

534

535

536 **8) Which observations could help understanding biases or further the process**
537 **understanding? [multiple answers allowed]**

538

○ Winter data (year-round observations)

539

○ Upper ocean stratification

540

○ Surface fluxes

541

○ Cloud observations

542

○ Sustained monitoring

543

○ Other: _____

544

545 **9) Where do mismatches exist between traditionally observed quantities and**
546 **model diagnostics (e.g. mixed layer depth)?**

547

548

549 **10) What is the most pressing ocean model development?**

550

[multiple answers allowed]

551

○ Implementation of ice shelf cavities

552

○ Overflow parameterization, vertical coordinates

553

○ Convection parameterization, vertical mixing

554

○ Scale-aware parameterization, e.g. for eddies

555

○ Biogeochemistry module

556

○ Coupled ecosystem module

557

○ Surface waves and/or tides

558

○ Sea ice dynamics

559

○ Other: _____

560

561

562 **11) Where will you or your collaborators evolve ocean models towards over the**
563 **next decade?**

564 *[multiple answers allowed]*

- 565 ○ Coupling more Earth system components (ice sheets, biogeochemistry, ...)
- 566 ○ Including more processes (tides, waves...)
- 567 ○ Novel parameterizations (submesoscale, scale-aware...)
- 568 ○ Higher resolution (horizontal and/or vertical)
- 569 ○ Unstructured grids, variable topography (wetting/drying, ice shelf cavity
- 570 shape)
- 571 ○ AI-based model components (parameterizations, emulators, ...)
- 572 ○ Other: _____

573

574 **12) Which kind of observations would further this development?**

575 _____

576

577

578 **Voluntary option:** provide your contact details for further discussion regarding Antarctica
579 InSync modeling activities: **name, affiliation, e-mail** (comma delimited please).

580 _____

581

582