

Hydrological implications of vegetation-associated precipitation recycling during peak growing season over the Loess Plateau

Jiaxiang Deng¹, Quan Quan¹, Shuangcheng Tang², Hanbo Yang³, and Xiaoyu Song¹

¹State Key Laboratory of Water Engineering Ecology and Environment in Arid Area, Xi'an University of Technology, Xi'an, China

²College of Hydraulic Science and Engineering, Yangzhou University, Yangzhou, China

³Tsinghua University, Beijing, China

Correspondence to: Quan Quan (qq@xaut.edu.cn)

Abstract. Vegetation restoration on the Loess Plateau has ~~raised an important question of~~ ~~led to continued debate over~~ whether enhanced land-atmosphere coupling can ~~alleviate regional water scarcity in this water-limited environment~~ ~~meaningfully alleviate water scarcity in this water-limited region~~. In this study, we ~~integrate~~ ~~combined~~ the WAM-2layers atmospheric moisture-tracking model with a random forest-enhanced Budyko framework and introduced a vegetation-weighted leaf area index (LAI_w) to examine vegetation-associated precipitation recycling and its hydrological ~~effects~~ ~~implications~~ during the peak growing season (July–August) ~~from during 2000–to 2022~~. Results show that precipitation is dominated by terrestrial moisture sources (86.5%), while oceanic contributions account for 13.5%. ~~Precipitation over the Loess Plateau is dominated by land-sourced moisture, which accounts for 86.5% of total precipitation.~~ Internal moisture recycling contributes 14.4% of total precipitation ~~but exhibits a~~ ~~and shows a declining~~ trend of $-0.073 \text{ mm yr}^{-1}$. ~~Surface water availability also declines significantly, at a rate of -0.34 mm yr^{-1} , mainly because evapotranspiration increases by 0.31 mm yr^{-1} .~~ At the regional mean scale, vegetation-associated recycled precipitation ~~makes only~~ ~~has~~ a limited ~~effect~~ ~~contribution~~ ~~onto~~ precipitation, evapotranspiration, and surface water availability, ~~accounting for~~ (0.059%, 0.02%, and 0.107%, respectively), ~~reflecting strong compensation between positive and negative vegetation effects and the limited fraction of internal recycling.~~ ~~This low net contribution partly reflects the fact that internal moisture recycling itself represents only a limited fraction of total precipitation, and that positive and negative vegetation-associated effects partly offset each other.~~ However, ~~a clear regime shift emerges~~ ~~the hydrological effect varies clearly~~ along the LAI_w gradient, ~~from~~ ~~with~~ weakly positive ~~contribution~~ ~~effects~~ under low vegetation density ~~to~~ ~~and~~ increasingly negative ~~effect~~ ~~contributions~~ under high vegetation density. These findings suggest that under water-limited conditions, enhanced vegetation-atmosphere coupling does not necessarily ~~lead to meaningful gains in~~ ~~increase~~ surface water availability, ~~but instead reflects a constrained redistribution of water within a monsoon-continental transition system characterized by nonlinear vegetation density-dependent regime shifts.~~ ~~This seasonal focus is intended to capture the period of strongest vegetation-atmosphere coupling rather than the full annual water balance.~~

Water scarcity and ecological fragility are major constraints on sustainable development in arid and semi-arid regions worldwide (Kummu et al., 2016; He et al., 2021). Climate warming has intensified the hydrological cycle by altering evaporation and precipitation processes, thereby reshaping regional water availability and increasing hydroclimatic risks, especially in water-limited environments (Giorgi et al., 2014; Huang et al., 2016; Wang et al., 2023; Zhang et al., 2023).

35 Although enhanced moisture recycling may partly mitigate water stress by returning evapotranspired water to the land surface as precipitation, the extent to which this process can effectively alleviate regional water limitation remains insufficiently understood.

Atmospheric moisture transport plays a fundamental role in regulating regional precipitation and water availability (Dominguez et al., 2020; He et al., 2021). Precipitation over a given region may originate either from evapotranspiration within
40 the region through internal moisture recycling or from moisture advected from upwind source areas (van der Ent et al., 2010; Tuinenburg et al., 2022; Cui et al., 2022). The balance between these internal and external moisture sources is particularly important in arid and semi-arid regions, where water availability is highly sensitive to both land–atmosphere coupling and atmospheric circulation (Te Wierik et al., 2022; Li et al., 2023; Zhang et al., 2022). Therefore, quantifying the contribution of local evapotranspiration to regional precipitation is essential for understanding whether land-surface changes can meaningfully
45 influence regional water resources.

The Loess Plateau (LP) of China is a representative water-limited region that has experienced severe ecological degradation and long-term water stress (Huang et al., 2016; Huang et al., 2017). Since the implementation of large-scale ecological restoration programs around 2000, vegetation cover and ecosystem productivity have increased substantially across the region (Li et al., 2021; Li et al., 2022). However, the hydrological consequences of vegetation restoration remain debated. On the one
50 hand, vegetation recovery can increase evapotranspiration, intensify soil moisture depletion, and reduce runoff, thereby aggravating water scarcity (Feng et al., 2016; Deng et al., 2020; Zhang et al., 2018). On the other hand, stronger evapotranspiration may enhance land–atmosphere coupling, promote atmospheric moisture recycling, and potentially increase precipitation (Yosef et al., 2018; Zhang et al., 2022a; Wei et al., 2019). These processes occur within a monsoon-dominated climatic system, where large-scale moisture transport and land–atmosphere interactions jointly regulate the evolution of
55 atmospheric water vapor along its pathway. These competing effects raise a key but unresolved question: under water-limited conditions, can vegetation-associated precipitation recycling provide a meaningful hydrological benefit, or is it largely offset by the additional water consumption associated with vegetation growth?

The seasonal timing of moisture recycling analysis requires careful consideration in dryland regions. While the full growing season on the Loess Plateau spans June to September, previous studies indicate that vegetation–atmosphere coupling intensity
60 varies substantially within this period. July and August represent the peak growing season, when leaf area index reaches its annual maximum, solar radiation is strongest, and soil moisture remains favorable for active evapotranspiration (Wang et al., 2019; Li et al., 2024). During this period, transpiration accounts for the dominant fraction of evapotranspiration, and land–

atmosphere moisture exchange is most pronounced (Guo et al., 2025). Consequently, focusing on July–August allows targeted assessment of vegetation-associated hydrological effects when they are expected to be strongest. This approach follows established practices in moisture recycling research where seasonal subsets are analyzed to isolate specific process periods (Zemp et al., 2014).

Addressing this question requires linking vegetation dynamics, evapotranspiration, atmospheric moisture tracking, and regional water balance within a unified framework. Existing studies have provided important insights into either the hydrological costs of revegetation or the climatic effects of land–atmosphere feedbacks, but quantitative assessments that explicitly connect vegetation-associated moisture recycling to regional water availability remain limited. In particular, few studies have integrated atmospheric moisture tracking with water-balance analysis to evaluate whether the precipitation returned through vegetation-associated recycling is large enough to compensate for the accompanying evapotranspiration increase in dryland ecosystems.

The Budyko framework provides a widely used basis for evaluating climatic controls on water partitioning between evapotranspiration and runoff (Budyko, 1974). However, its conventional form generally assumes long-term equilibrium conditions and fixed catchment characteristics, which may be insufficient for regions such as the Loess Plateau, where hydroclimatic conditions are spatially heterogeneous and land-surface properties have changed substantially under ecological restoration (Yang et al., 2009; Yokoo et al., 2008; Woods, 2003; Yang et al., 2015). To better capture these variations, it is necessary to allow the effective catchment parameter to vary with vegetation, soil moisture, and meteorological conditions.

In this study, we develop an integrated framework that combines the WAM-2layers atmospheric moisture tracking model with a random forest-enhanced Budyko approach to quantify vegetation-associated precipitation recycling and its hydrological implications over the Loess Plateau during the peak growing season (July–August) from 2000 to 2022. We further introduce a vegetation-weighted indicator, LAI_w , as a diagnostic measure that combines vegetation structure with relative evapotranspiration intensity in order to represent vegetation conditions more closely related to land–atmosphere moisture exchange. Specifically, this study aims to (1) characterize the spatiotemporal patterns of growing-season moisture recycling and water availability over the Loess Plateau, (2) quantify the vegetation-associated component of recycled precipitation, and (3) evaluate whether this recycled precipitation meaningfully offsets evapotranspiration-induced water losses under different vegetation-density conditions. By doing so, this study seeks to clarify the hydrological limits of vegetation-associated precipitation recycling in a water-constrained dryland region.


2 Data and methods

2.1 Study area and climatic characteristics

The Loess Plateau (LP) is located in the middle reaches of the Yellow River basin (33°43′–41°16′N, 104°54′–114°33′E) and represents a typical arid and semi-arid region in northern China. The region covers an area of approximately 640,000 km² and is characterized by complex topography, with elevation generally decreasing from northwest to southeast and an overall

95 elevation range exceeding 3000 m. The Loess Plateau is dominated by a temperate continental monsoon climate. The mean annual air temperature and precipitation are approximately 9.4 °C and 593.7 mm, respectively. Precipitation exhibits strong seasonality, with about 60–70% of the annual total occurring between June and September, whereas winter precipitation is limited and contributes negligibly to the annual water budget (Wang et al., 2023). Under the influence of the East Asian monsoon system, precipitation is primarily driven by large-scale moisture transport, which is subsequently modified by
 100 regional land-atmosphere recycling processes. Precipitation over the Loess Plateau also shows pronounced interannual variability, with wet years receiving several times more precipitation than dry years and extreme rainfall events occurring frequently (Liu et al., 2025).

Given the strong seasonality of both precipitation and vegetation activity, this study focuses on July and August as the primary analysis period. These two months correspond to the peak growing season on the Loess Plateau, during which leaf area index (LAI) and fractional vegetation cover generally reach their annual maxima, solar radiation is relatively strong, and soil moisture conditions remain favorable for active land-atmosphere exchange. As a result, evapotranspiration and vegetation-atmosphere interactions are most pronounced during this period (Wang et al., 2019; Li et al., 2024; Guo et al., 2025). Focusing on July and August therefore allows this study to better capture the period when vegetation-associatedrelated moisture recycling is



expected to be strongest, providing a targeted basis for evaluating vegetation-associated hydrological effects during the peak growing season in this water-limited region. This seasonal focus is intended to isolate the period of strongest vegetation–atmosphere coupling rather than to represent the full annual water balance.

Figure 1. Study area and key characteristics, including (a) topography and river network, (b) land cover, (c) evaporation, (d) precipitation, and (e) land-cover conversions.

2.2 Data

Monthly evapotranspiration (ET) and potential evapotranspiration (ET_0) data for July and August during 2000–2022 were obtained from the Global Land Evaporation Amsterdam Model (GLEAM, v3.6a) dataset, with a spatial resolution of 0.25° (Martens et al., 2017). GLEAM ET has been extensively evaluated against eddy covariance (EC) observations across China and shows good agreement at the monthly scale, with an average correlation coefficient of 0.86 (Yang et al., 2017).

Monthly precipitation data for July and August during 2000–2022 were derived from the ERA5 reanalysis dataset at a spatial resolution of 0.25° (Hersbach et al., 2020). ERA5 precipitation has been shown to capture the annual and seasonal variability of observed precipitation over China reasonably well (Jiao et al., 2021) and exhibits a strong correlation ($r = 0.92$) with official national statistics at the 99% confidence level (An et al., 2025). Before subsequent analyses, the ERA5 precipitation and GLEAM ET datasets were temporally aligned for July–August during 2000–2022 and resampled to a common 0.25° grid using nearest-neighbor interpolation to ensure spatial consistency.

For atmospheric moisture tracking, hourly meteorological variables from the ERA5 dataset were collected for June–August during 2000–2022. These variables included air temperature, precipitation, specific humidity, wind components, surface pressure, dew-point temperature, total column water vapor, and surface evaporation. Specific humidity and wind fields were extracted for 17 vertical pressure levels ranging from 1000 to 100 hPa, all at a spatial resolution of 0.25° . ERA5 data have been widely used in previous moisture-tracking studies over China and have shown robust performance (An et al., 2025; Xie et al., 2024; Zhang et al., 2024). Considering the atmospheric residence time of water vapor (van der Ent et al., 2013), meteorological data from June were included as auxiliary input to enable more complete tracking of the moisture contributing to precipitation in July and August.

Vegetation conditions were characterized using the High-Quality Leaf Area Index (HiQ-LAI) dataset, a globally consistent LAI product derived from MODIS LAI Collection 6.1 using a spatiotemporal information compositing algorithm (STICA) (Yan et al., 2024). Compared with the original MODIS LAI product, HiQ-LAI improves temporal stability and spatial consistency by integrating pixel-level quality information and spatiotemporal correlations while preserving the underlying radiative transfer framework. The dataset provides global coverage at an 8-day temporal resolution and a spatial resolution of 5 km for 2000–2022, and has been validated against ground-based observations, showing reduced noise and improved phenological consistency relative to the original MODIS LAI . In this study, HiQ-LAI data for July and August were extracted and aggregated to characterize vegetation conditions during the peak growing season over the Loess Plateau. The HiQ-LAI dataset is publicly available via Zenodo (Yan et al., 2023).

Additional variables used in the random forest modeling of the Budyko parameter n were derived from the ERA5-Land and ERA5 reanalysis datasets. Four-layer soil water content (swvl1–swvl4), near-surface wind components (u and v), and surface temperature were obtained from ERA5-Land at a spatial resolution of 0.1° . Surface runoff (src) and surface shortwave radiation (ssr) were obtained from the ERA5 hourly single-level dataset at a spatial resolution of 0.25° . Vegetation cover (VC) was characterized using the China regional fractional vegetation cover (FVC) dataset at 250 m resolution (Gao et al., 2024). All variables were spatially aggregated to a common 0.25° grid to maintain consistency with the Budyko framework.

2.3 Methodology

This study employs an integrated framework combining atmospheric moisture tracking, vegetation-based diagnostics, and hydrological modeling to examine vegetation-associated precipitation recycling and its hydrological implications over the Loess Plateau during the peak growing season (Fig. 2a).




Figure 2: Methodological framework and model validation results, including (a) schematic diagram of the methodological framework, (b) scatter plot of predicted n values from the random forest model versus observed n values in the test set, and (c) scatter plot of evapotranspiration estimated by the Budyko–RF framework versus observed evapotranspiration.

2.3.1 Moisture tracking and precipitation source attribution

To identify the moisture sources of precipitation over the Loess Plateau (LP) and to separate the contributions from internal recycling and external advection, we employed the Eulerian moisture-tracking model WAM-2layers v3 (Kalverla et al., 2025; van der Ent et al., 2013, 2014). The model partitions the atmosphere into two layers, namely a lower layer ($k=1$) below

approximately 812 hPa and an upper layer ($k=0$) representing the free atmosphere, and assumes that moisture is well mixed within each layer. In the model, tagged moisture is defined as $S^*=cS$, where c is the tagged moisture concentration. Its spatiotemporal evolution is governed by the tagged moisture budget equation:

$$\frac{\Delta_t S^*}{\Delta t} + \frac{1}{A} (\Delta_x F_x^* + \Delta_y F_y^* + \Delta_p F_p^*) = E^* - P^* \quad (1)$$

where S^* is tagged moisture storage, F_x^* , F_y^* , and F_p^* are the tagged moisture fluxes in the zonal, meridional, and vertical directions, respectively, and E^* and P^* are the tagged evaporation and precipitation terms. In WAM-2layers, tagged horizontal fluxes are assumed to be proportional to total moisture fluxes, whereas the vertical tagged flux includes an additional dispersion term to enhance interlayer mixing. The vertical transport term is treated as a closure term representing exchange at the interface between the two layers.

For the two-layer implementation, evaporation is assigned entirely to the lower layer, whereas precipitation is partitioned between the upper and lower layers according to their relative moisture contents. The resulting upper- and lower-layer equations can be written as:

$$S_{ij,0}^{t+1} = S_{ij,0}^t + \Delta t \left[\frac{\left(F_{x,i+\frac{1}{2},j,0}^* - F_{x,i-\frac{1}{2},j,0}^* \right)}{A} - \frac{\left(F_{y,i,j+\frac{1}{2},0}^* - F_{y,i,j-\frac{1}{2},0}^* \right)}{A} - \frac{\left(F_{p,ij,\frac{1}{2}}^* + k_{vf} \left| F_{p,ij,\frac{1}{2}} \right| (c_{ij,1}^t - c_{ij,0}^t) \right)}{A} - c_{ij,0} \frac{S_{ij,0}}{S_{ijT}} P_{ij}^{t+\frac{1}{2}} \right] \quad (2)$$

and

$$S_{ij,1}^{t+1} = S_{ij,1}^t + \Delta t \left[\frac{\left(F_{x,i+\frac{1}{2},j,1}^* - F_{x,i-\frac{1}{2},j,1}^* \right)}{A} - \frac{\left(F_{y,i,j+\frac{1}{2},1}^* - F_{y,i,j-\frac{1}{2},1}^* \right)}{A} + \frac{\left(F_{p,ij,\frac{1}{2}}^* + k_{vf} \left| F_{p,ij,\frac{1}{2}} \right| (c_{ij,1}^t - c_{ij,0}^t) \right)}{A} + \delta_e E_{ij}^{t+\frac{1}{2}} - c_{ij,1} \frac{S_{ij,1}}{S_{ijT}} P_{ij}^{t+\frac{1}{2}} \right] \quad (3)$$

where A is grid-cell area, S_{ijT} is total column moisture, and k_{vf} is the vertical dispersion coefficient. This two-layer formulation improves the representation of vertical moisture exchange relative to single-layer models, particularly under strong vertical wind shear.

In this study, WAM-2layers was run in backward-tracking mode to trace the sources of precipitation over the LP during July–August for 2000–2022. Considering the atmospheric residence time of water vapor, June meteorological inputs were also included, and the tracking period was set to July–August to ensure more complete tracing of moisture contributing to July–August precipitation. Hourly ERA5 variables were temporally interpolated to a 10 min time step and simulated at 0.25° spatial resolution. This configuration follows the model recommendation that moisture transport within a time step should not exceed one grid cell, thereby maintaining numerical stability under the Courant-Friedrichs-Lewy criterion.

Based on the tracking results, precipitation over the LP was decomposed into internally recycled and externally advected components, and the relative contributions of land and ocean source regions were quantified. The recycling ratio α is defined as (van der Ent et al., 2010):

$$\alpha = \frac{P_{v0}}{P_v} \quad (4)$$

where P_v is the total precipitation over the LP during the tracking period, and P_{v0} is the amount of precipitation over the LP that originates from evaporation within the LP and subsequently returns to the LP through atmospheric moisture transport.

2.3.2 Construction of LAI_w and vegetation-associated component of recycled precipitation

To characterize vegetation conditions that are more directly related to land–atmosphere moisture exchange, a vegetation-weighted index, denoted as LAI_w , was constructed in this study. The purpose of this index is not to define a new physical state variable, but to provide a diagnostic indicator that better links vegetation structure to the evaporative conditions under which atmospheric moisture recycling occurs.

Leaf area index (LAI) is widely used to represent vegetation canopy structure, but LAI alone does not fully reflect the extent to which vegetation is actively coupled to atmospheric moisture exchange. Under dryland conditions, the hydrological role of vegetation depends not only on canopy density, but also on the contemporaneous evaporative intensity associated with available water and energy. Grid cells with similar LAI may therefore contribute differently to moisture recycling if their evapotranspiration rates differ substantially. To account for this combined control, LAI was weighted by normalized actual evapotranspiration, so that vegetation in grid cells with stronger evapotranspiration receives greater emphasis in the subsequent diagnosis of recycled precipitation.

For each grid cell, the mean leaf area index during the peak growing season (July–August) was derived from the HiQ-LAI dataset. Actual evapotranspiration during the same period, denoted as E_{obs} , was used to represent the evaporative strength associated with local vegetation and hydroclimatic conditions. The vegetation-weighted index is defined as:

$$LAI_w = LAI \cdot \frac{E_{obs}}{\bar{E}_{obs}} \quad (5)$$

where E_{obs} is the observed evapotranspiration at the grid scale, and $\bar{E}_{obs}$ is the spatial mean of E_{obs} over the Loess Plateau, used for normalization. This formulation preserves the structural information represented by LAI while moderately increasing the relative weight of vegetation in grid cells with stronger evapotranspiration. Accordingly, LAI_w is intended as a diagnostic measure of vegetation conditions that are more closely associated with land–atmosphere moisture exchange than LAI alone.

It should be noted that LAI_w is a derived indicator rather than a directly observed physical quantity. Its role in this study is therefore diagnostic rather than strictly mechanistic. The weighting does not imply that evapotranspiration is solely controlled by vegetation, nor that LAI_w should be interpreted as an independent biophysical property. Instead, it is used to describe

vegetation conditions under which structural canopy effects and evaporative activity jointly influence the likelihood of moisture returning to the study region as recycled precipitation.

215 At the grid scale, a multiple linear regression was then applied to estimate the vegetation-associated component of recycled precipitation over the Loess Plateau:

$$P_{rec} = d + a \cdot LAI_w + b \cdot T + c \cdot SSR \quad (6)$$

where P_{rec} represents the grid-based recycled precipitation, defined as the amount of precipitation over the Loess Plateau originating from evaporation at a given grid cell and returning to the plateau through atmospheric moisture transport. T and SSR denote near-surface air temperature and surface shortwave radiation, which are included to account for major thermodynamic controls on moisture recycling. The coefficient a represents the statistical sensitivity of recycled precipitation to LAI_w , conditional on T and SSR .

Based on the estimated coefficient a , the vegetation-associated contribution to recycled precipitation is defined as:

$$P_0 = a \cdot LAI_w \quad (7)$$

225 where P_0 represents the component of recycled precipitation statistically associated with vegetation conditions as characterized by LAI_w . Here, P_0 should be interpreted as a vegetation-associated diagnostic term rather than as a strictly causal estimate derived from process-based perturbation experiments. This term is subsequently used to construct precipitation scenarios with the vegetation-associated recycling effect removed.

The use of LAI_w instead of the original LAI is intended to better capture the joint role of vegetation structure and evapotranspiration intensity in regulating recycled precipitation during the peak growing season. However, to avoid over-interpreting the weighting scheme, the robustness of this diagnostic formulation was further evaluated by repeating the key analytical steps using unweighted LAI as an alternative vegetation predictor. This comparison was used to assess whether the major results depend strongly on the specific construction of LAI_w , or whether they primarily reflect the broader vegetation pattern itself.

235 2.3.3 Budyko–RF framework for assessing hydrological impacts of vegetation-associated precipitation recycling

The Budyko framework was originally developed for long-term water balance analysis, but recent advances have extended its application to sub-annual timescales by incorporating dynamic storage effects and time-varying parameters (Yang et al., 2026; Liu et al., 2025). At the monthly scale, explicitly accounting for soil water storage change can substantially improve the performance of Budyko-type equations, thereby supporting the feasibility of such approaches beyond traditional annual mean applications (Yang et al., 2026).

In this study, the Budyko framework is applied as a comparative diagnostic tool for the peak growing season (July–August), rather than as a process-based simulator of the full land–atmosphere system. Unlike climate model (GCM/RCM) approaches, which explicitly resolve coupled vegetation–atmosphere interactions through dynamic representations of radiation, convection, and moisture transport, the Budyko framework provides a constraint-based diagnostic description of surface water balance under observed hydroclimatic conditions. This distinction implies that vegetation–precipitation feedbacks in this study are not

simulated dynamically, but are instead quantified through their contribution to the observed precipitation–evapotranspiration balance, a strict representation of the full seasonal or annual water balance.

The framework is used to isolate the first-order hydrological effects of vegetation-associated precipitation recycling under consistent land surface and atmospheric conditions. This seasonal application is conceptually consistent with previous monthly
250 Budyko studies that emphasize the importance of parameter variability under changing hydroclimatic conditions (Chen et al., 2013).

To better represent sub-annual variability in land-surface and hydroclimatic conditions, the key Budyko parameter n was allowed to vary across grid cells and years through a Random Forest (RF) model (Breiman, 2001). This treatment is conceptually consistent with previous monthly Budyko studies that emphasize the importance of storage effects and parameter
255 variability under changing hydroclimatic and land-surface conditions. By linking n to vegetation, soil moisture, and meteorological conditions, this Budyko–RF framework was used to evaluate the hydrological implications of vegetation-associated precipitation recycling.

Actual evapotranspiration (E) simulated by the Budyko framework was calculated using the Mezentsev–Choudhury–Yang equation:

$$E = \frac{ET_0 \cdot P}{(P^n + ET_0^n)^{1/n}} \quad (8)$$

where P is precipitation, ET_0 is potential evapotranspiration, and n is a dimensionless parameter characterizing integrated land surface properties.

Because n is affected by vegetation conditions, soil water status, and atmospheric forcing, an RF regression model was used to estimate n for each grid cell:

$$n = RF(VC, swvl1, swvl2, swvl3, swvl4, u, v, T, src, ssr) \quad (9)$$

where VC is vegetation cover; $swvl1$ – $swvl4$ represent soil moisture in four layers; u and v are wind components; T is near-surface air temperature; src is surface runoff; and ssr is surface shortwave radiation.

To evaluate the performance of the Budyko–RF framework at the analysis scale used in this study, a two-step validation was performed. First, 80% of the samples were used for model training and the remaining 20% for testing. The RF model showed
270 good predictive skill for n in the test set (Fig. 2b, $R^2=0.82$), indicating that the effective Budyko parameter can be reasonably estimated from vegetation associated, hydrological, and meteorological predictors. Second, evapotranspiration estimated by the Budyko–RF framework was compared with observed evapotranspiration at the grid scale. The comparison showed strong agreement (Fig. 2c, $R^2=0.93$), suggesting that the framework can satisfactorily reproduce actual evapotranspiration during the study period.

To further quantify the hydrological impacts of vegetation-associated precipitation recycling, a counterfactual precipitation
275 scenario was constructed by removing the vegetation-associated recycled component from regional mean precipitation while keeping other climatic controls unchanged. For each year t , the study domain contains N_t valid grid cells, and regional mean

precipitation and potential evapotranspiration are denoted by $\bar{P}_t$ and $\overline{ET}_{0t}$, respectively. Based on the grid-scale vegetation-associated recycled precipitation $P_{0,i,t}$, the total contribution associated with a given set of grid cells S is expressed as:

$$P_{0,t}^{(S)} = \sum_{i \in S} P_{0,i,t} \quad (10)$$

The precipitation scenario without vegetation-associated recycling is then defined at the regional mean scale as:

$$P_{\text{clim},t}^{(S)} = \bar{P}_t - \frac{P_{0,t}^{(S)}}{N_t} \quad (11)$$

This formulation represents the removal of the vegetation-associated recycling component as a uniform perturbation to regional mean precipitation, allowing direct comparison among years and among vegetation subsets.

Hydrological responses to the counterfactual precipitation were evaluated within the Budyko framework. For each year t , the Budyko parameter n_t was held constant when comparing the baseline and counterfactual scenarios, so that changes in evapotranspiration could be attributed solely to differences in precipitation. Accordingly, regional mean evapotranspiration under baseline and counterfactual conditions is given by:

$$\bar{E}_t = \beta(\bar{P}_t, \overline{ET}_{0t}, n_t), \quad E_{\text{clim},t}^{(S)} = \beta(P_{\text{clim},t}^{(S)}, \overline{ET}_{0t}, n_t) \quad (12)$$

where $\beta(\cdot)$ denotes the Budyko mapping.

The vegetation-associated component of regional evapotranspiration associated with precipitation recycling is quantified by scenario differencing:

$$E_{0,t}^{(S)} = \bar{E}_t - E_{\text{clim},t}^{(S)} \quad (13)$$

The rRegional aridity index is introduced to further characterize the regional hydroclimatic conditions, defined as:

$$AI_t = \frac{\bar{P}_t}{\bar{E}_t} \quad (14)$$

Here, AI is used as a diagnostic hydroclimatic index defined as P/E , rather than the conventional PET-based aridity index.

The rRegional surface water availability is defined as the difference between precipitation and evapotranspiration:

$$SWA_t = \bar{P}_t - \bar{E}_t \quad (15)$$

with the corresponding value under the counterfactual precipitation scenario given by:

$$SWA_{\text{clim},t}^{(S)} = P_{\text{clim},t}^{(S)} - E_{\text{clim},t}^{(S)} \quad (16)$$

The relative contribution of vegetation-associated precipitation recycling to regional surface water availability is finally expressed as:

$$C_{SWA,t}^{(S)} = \frac{SWA_t - SWA_{\text{clim},t}^{(S)}}{SWA_t} \times 100\% \quad (17)$$

This framework provides a consistent attribution of vegetation-associated precipitation recycling effects on regional evapotranspiration and surface water availability, and can be applied uniformly to the entire study domain as well as to vegetation subsets defined by LAI_w magnitude or long-term LAI_w trends.

3 Results

3.1 Spatiotemporal characteristics and sources of recycled precipitation over the Loess Plateau

Atmospheric moisture tracking based on WAM-2layers for the peak growing season (July–August) during 2000–2022 shows that precipitation over the Loess Plateau (LP) is sustained by both external moisture advection and internal moisture recycling. The mean amount of precipitation originating from evaporation at individual grid cells and returning to the LP (P_{v0}) is 2.05 mm, indicating that terrestrial evaporation contributes measurably to regional precipitation. The dominant moisture transport pathways supplying precipitation to the LP exhibit a clear bimodal pattern (Fig. 3a). One pathway is associated with the mid-latitude westerlies, through which moisture from Central and Western Asia is transported eastward toward the LP. The other is related to southerly transport driven by the southwest and southeast monsoons, which carries moisture from the Indian and Pacific Oceans northward into the study region.

Quantitative source attribution further shows that land-sourced moisture dominates the precipitation supply to the LP. On average, terrestrial moisture contributes approximately 71.0 mm of precipitation, accounting for 86.5% of total growing-season precipitation, whereas oceanic moisture contributes about 11.1 mm, corresponding to 13.5% (Fig. 3b). This land-sourced moisture signal reflects moisture that has undergone substantial transformation through repeated evaporation–precipitation cycles during inland transport, rather than exclusively locally generated evaporation. Such a pattern suggests progressive modification of monsoon-derived moisture along continental transport pathways within a monsoon–inland transition zone, where ocean-origin vapor is continuously mixed with terrestrial evapotranspiration inputs before reaching the Loess Plateau.

In terms of recycling pathways, externally advected precipitation accounts for 70.3 mm (85.6%), while internally recycled precipitation contributes 11.8 mm, representing 14.4% of total precipitation over the LP. During the study period, internally recycled precipitation shows a declining trend of $-0.073 \text{ mm yr}^{-1}$ (Fig. 3c). This result indicates a gradual weakening of the contribution of internal moisture recycling to growing-season precipitation over the LP.





Figure 3. Recycled precipitation over the Loess Plateau during the growing season (July–August, 2000–2022). (a) Spatial distribution of mean recycled precipitation and moisture transport. (b) Contributions of oceanic and terrestrial moisture sources and the partitioning between external and internal recycling. (c) Interannual variation of internally recycled precipitation.

3.2 Vegetation-weighted patterns and contributions to recycled precipitation

To characterize the spatial patterns of vegetation conditions relevant to moisture recycling over the Loess Plateau, the vegetation-weighted index LAI_w was analyzed at the grid scale. The spatial distribution and temporal trends of LAI_w are shown in Fig. 4a–b. The multi-year mean LAI_w over the Loess Plateau is 0.45, and approximately 90.6% of grid cells exhibit LAI_w values lower than 1, mainly in the northwestern part of the plateau where vegetation cover is relatively sparse. In contrast, grid cells with LAI_w values greater than 4 account for only 2.9% of the region and are concentrated in localized areas of the southeastern Loess Plateau. At the regional scale, 48.7% of grid cells show increasing trends in LAI_w , whereas 26.5% show decreasing trends. Spatially, increases in LAI_w are mainly located in areas that already exhibit relatively high LAI_w values (Fig. 4b).

340 To further evaluate the statistical association between vegetation conditions and recycled precipitation, a grid-based multiple linear regression was performed with recycled precipitation (P_{rec}) as the dependent variable and LAI_w , near-surface air temperature, and surface shortwave radiation as explanatory variables. Here, P_{rec} represents the amount of precipitation over the Loess Plateau that originates from evaporation at a given grid cell and subsequently returns to the plateau through atmospheric moisture transport. The regional mean of the sensitivity coefficient of LAI_w (a) is 2.34, and 57.5% of grid cells exhibit positive a -values, indicating that higher LAI_w is statistically associated with greater recycled precipitation across much of the study area (Fig. 4c–d).



350 **Figure 4.** Spatial patterns and effects of vegetation-weighted index LAI_w over the Loess Plateau. (a) Spatial distribution of mean LAI_w during the growing season. (b) Spatial patterns of linear trends in LAI_w . (c) Spatial distribution of the regression coefficient a . (d) Relationship between mean LAI_w and the regression coefficient a . (e) Interannual variation of vegetation contributions to the evaporation–precipitation cycle.

Fig. 4e shows the interannual evolution of the vegetation-associated contribution to the evaporation–precipitation (E–P) cycle, decomposed into positive, negative, and net components. Over the study period, the mean positive contribution is 1.16%, with a weak increasing trend of $0.007\% \text{ yr}^{-1}$. In contrast, the mean negative contribution is -0.97% , with a ~~decreasing~~ trend of $-0.026\% \text{ yr}^{-1}$. As a combined result of these opposing components, the net vegetation-associated contribution to the E–P cycle remains small, with a mean value of 0.12% and a slight ~~negative~~ trend of $-0.019\% \text{ yr}^{-1}$. Overall, these results show that positive and negative vegetation-associated influences on internal moisture recycling largely offset each other at the regional scale.

3.3 Declining surface water availability and hydrological implications of vegetation-associated precipitation recycling

A pronounced drying tendency is observed over the Loess Plateau during the growing season from 2000 to 2022 (Fig. 5a). Interannual variations in precipitation (P), evapotranspiration (E), surface water availability (SWA), and the aridity index (AI) show that regional precipitation exhibits a non-significant ~~declining~~ trend of $-0.0043 \text{ mm yr}^{-1}$, whereas evapotranspiration shows a significant increasing trend of 0.31 mm yr^{-1} . The contrasting changes in precipitation and evapotranspiration correspond to increasingly dry hydroclimatic conditions over the study period. Although the interannual variations in SWA and AI generally follow the temporal pattern of precipitation, both indices exhibit significant downward trends under increasing evapotranspiration, with SWA decreasing at a rate of -0.34 mm yr^{-1} and AI decreasing at a rate of -0.008 yr^{-1} .

The spatial distribution of mean SWA and its temporal trends also shows clear heterogeneity across the Loess Plateau (Fig. 5b–c). Higher SWA values are mainly distributed in the southeastern part of the region, whereas persistently low SWA values characterize the northwestern arid areas. Declining SWA trends occur in both relatively high- SWA and low- SWA regions, whereas increasing trends are mainly concentrated in the southwestern Loess Plateau. Although 51% of grid cells show increasing trends in SWA and 49% show decreasing trends, the pattern becomes asymmetric when statistical significance is considered. Among the 56.6% of grid cells with significant SWA trends ($P < 0.05$), 40% exhibit significant declines, whereas only 16.6% show significant increases. This indicates that the most robust regional signal is a widespread decline in SWA .

To quantify the hydrological implications of vegetation-associated recycled precipitation (P_0), the P_0 term estimated in Section 3.2 was subtracted from total precipitation and the resulting water-balance components were evaluated within the Budyko–RF framework. The results (Fig. 5d) show that the mean relative contributions of vegetation-associated recycled precipitation to P , E , and SWA are 0.059%, 0.02%, and 0.107%, respectively. A consistent response pattern is observed across the study region: grid cells with positive P_0 contributions to precipitation also show higher E and SWA , whereas negative P_0 contributions are accompanied by concurrent reductions in both E and SWA . Overall, the positive hydrological contribution of P_0 is relatively more evident in drier areas with lower SWA .




Figure 5: Vegetation-associated impacts on precipitation, evapotranspiration, and surface water availability over the Loess Plateau. (a) Interannual variations of growing-season P , E , SWA , and AI . (b) Spatial distribution of SWA . (c) Spatial patterns of SWA trends. (d) Interannual contribution rates of vegetation to P , E , and SWA . (e) Contribution rates of vegetation to SWA under different LAI_w classes.

Fig. 5e further shows a systematic dependence of the SWA contribution associated with vegetation-associated recycled precipitation on LAI_w . Under low- LAI_w conditions ($LAI_w < 1$), the contribution to SWA remains consistently positive, with a

385 mean value of 0.2855% and no negative values across all years, although it shows a ~~weakening~~ trend of $-0.0023\% \text{ yr}^{-1}$. As LAI_w increases, this positive contribution diminishes and gradually shifts toward negative values. For intermediate LAI_w conditions ($1 \leq LAI_w \leq 2.5$), the mean contribution is close to zero (0.02565%) and shows substantial interannual variability, with negative values occurring in 11 years; however, a modest positive trend ($0.0035\% \text{ yr}^{-1}$) is still observed. In contrast, high-
390 for $2.5 \leq LAI_w \leq 4$ and -0.1495% for $LAI_w > 4$, and the strongest negative trend ($-0.0092\% \text{ yr}^{-1}$) occurs ~~ring~~ in the highest- LAI_w class. Overall, these results indicate a systematic shift from weakly positive to increasingly negative SWA contributions along the LAI_w gradient.

4 Discussion

4.1 Hydroclimatic constraints and precipitation recycling over the Loess Plateau

395 Recent global ~~studies show assessments have shown~~ that surface water availability (SWA) has increased over ~~much large parts~~ of the terrestrial surface. ~~This is mainly, mainly~~ because precipitation has ~~risen increased~~ faster than evapotranspiration across more than 53% of global land areas (Cui et al., 2022). In contrast, our results ~~indicate show~~ a decline in SWA over the Loess Plateau during the study period (Fig. 5a). In this arid and semi-arid region, evapotranspiration increased faster than precipitation, ~~reducing leading to a reduction in~~ available surface water. Similar drying trends have been widely reported in
400 dryland ~~regions, where e~~Climate warming is expected to increase drought frequency and severity (Stevenson et al., 2022; Tabari et al., 2021). ~~In addition, and where regional water resources often decline under~~ limited precipitation and enhanced evaporation ~~often lead to declining regional water resources~~ (Lai et al., 2022). ~~Together, t~~These ~~findings indicate results together suggest~~ that water availability in in drylands is strongly constrained by low precipitation and high evaporative demand. ~~arid and semi-arid systems remains strongly constrained by limited precipitation and high evaporative demand.~~

405 Under this hydroclimatic background, precipitation over the Loess Plateau is strongly controlled by large-scale monsoon moisture transport. It is continuously modified by land-atmosphere exchanges along its inland pathway. The WAM-2layers results show that Central China is the predominantly continental-influenced moisture source for precipitation over the Loess Plateau (Fig. 3a). This is consistent with previous Lagrangian studies. These studies indicate that summer moisture mainly comes from Central-Eastern China and northwestern continental Asia. In contrast, direct oceanic contributions from the East China Sea, South China Sea, Bay of Bengal, and Arabian Sea are minor, together accounting for only about 10% of total precipitation (Hu et al., 2016).

410 The dominance of continental moisture sources reflects progressive changes in monsoon vapor during inland transport. Along this pathway, repeated evaporation-precipitation cycles and boundary layer recycling gradually reshape the original oceanic moisture signal. Recent studies also show that vegetation greening enhances evapotranspiration and strengthens regional moisture recycling (Cao et al., 2024). In this process, the land surface acts both as a moisture source and an intermediate

contributor to atmospheric moisture through evapotranspiration and recycling processes. This reinforces the continental component in the monsoon–land coupled system. Therefore, the high fraction of land-sourced precipitation (86.5%) should not be interpreted as a weak oceanic influence. Instead, it reflects the combined effects of monsoon moisture transport and terrestrial recycling. The East Asian monsoon provides the initial moisture input, while continental recycling determines the final precipitation composition over the Loess Plateau. Under this hydroclimatic setting, precipitation over the Loess Plateau still depends strongly on external moisture supply. The WAM-2layers results show that Central China is the dominant external moisture source for precipitation over the Loess Plateau (Fig. 3a). This pattern is broadly consistent with previous studies based on other moisture-tracking approaches, including FLEXPART-based analyses (Hu et al., 2018) and WAM-2layers simulations (Cao et al., 2024). This consistency suggests that the large-scale circulation pattern controlling moisture transport to the region is relatively stable.

Within this externally controlled system, internal precipitation recycling contributes about 14.4% of total precipitation over the Loess Plateau during the study period (Fig. 3b–c). This estimate value is higher than those reported in values reported in some previous studies, such as 7% from the UTrack Lagrangian moisture-tracking model (Wang, Liu et al., 2023) and $9.16 \pm 0.35\%$ to $10.18 \pm 0.28\%$ from the Dynamic Precipitation Recycling Method (Tian et al., 2022). These differences mainly reflect methodological approaches and climatic context rather than physical inconsistency. Most previous studies use annual or long-term averages. In contrast, this study focuses on July–August, when vegetation activity, evapotranspiration, and land–atmosphere coupling are strongest over the Loess Plateau. One likely reason is the seasonal focus of this study. Our analysis is limited to July and August, when vegetation activity, evapotranspiration, and land–atmosphere coupling are strongest over the Loess Plateau (Li et al., 2024; Guo et al., 2025).

This seasonal amplification agrees with recent findings that vegetation and land use changes regulate atmospheric moisture recycling by affecting evapotranspiration and atmospheric moisture residence time (Wang et al., 2023). In this framework, precipitation recycling is not a fixed ratio. It responds dynamically to land surface conditions and atmospheric moisture supply. Similarly, afforestation has been shown to strengthen land–atmosphere coupling and increase both external and internal moisture contributions to precipitation (Tian et al., 2022). Therefore, the relatively high recycling ratio in this study reflects enhanced seasonal coupling during the peak growing season, rather than overestimation. These results indicate that precipitation over the Loess Plateau is jointly controlled by monsoon moisture transport and terrestrial recycling processes. During this period, land derived moisture accounts for about 86.5% of the total precipitation supply (Fig. 3b). Stronger transpiration and soil evaporation may therefore increase the fraction of precipitation recycled from regional land surfaces. In this sense, the relatively high internal recycling ratio reported here is more likely to reflect seasonal amplification during the peak growing season than a uniformly high contribution throughout the year.

4.2 Conditional and nonlinear vegetation-associated effects on surface water availability

Under water-limited conditions, vegetation ~~affects~~~~influencees~~ regional water cycling through two competing pathways. ~~First~~
450 ~~On the one hand~~, enhanced vegetation activity increases evapotranspiration and strengthens land–atmosphere coupling, ~~which~~
~~may~~~~This can~~ promote the return of locally evaporated moisture as precipitation (Sterling et al., 2013). ~~Second~~~~On the other~~
~~hand~~, vegetation growth ~~also~~ increases terrestrial water consumption. ~~This reduces soil moisture and limits water availability~~
~~for surface and subsurface hydrological processes which can reduce soil moisture and limit the water available for surface and~~
~~subsurface hydrological processes~~ (Piao et al., 2020). ~~The coexistence of t~~These two pathways coexist. As a result, vegetation-
455 ~~associated effects on internal moisture recycling do not translate into a uniformly positive impact on surface water~~
~~availability helps explain why vegetation-associated effects on internal moisture recycling do not lead to a uniformly positive~~
~~effect on surface water availability.~~

Recent work by Tian et al. (2026) provides a complementary system-level view. They show that vegetation greening can
enhance precipitation efficiency through albedo-driven energy feedbacks. These feedbacks increase surface radiation,
460 atmospheric instability, and moisture-to-precipitation conversion efficiency. Their results also indicate that greening can
simultaneously increase evapotranspiration and streamflow in semi-arid regions. This suggests a coupled energy–moisture
mechanism that intensifies the atmospheric water cycle. The results are broadly consistent with the energy–moisture coupling
perspective proposed by Zhang (2025), particularly in terms of enhanced atmospheric moisture recycling under greening
conditions. We observe strengthened land–atmosphere coupling and enhanced internally recycled moisture. However, the
465 effect on surface water availability depends on temporal scale and hydrological closure assumptions.

At the regional scale, ~~the long-term~~ vegetation-associated ~~influence effects~~ on the evaporation–precipitation cycle ~~appears to~~
largely cancel out ~~in net effect over the long term~~. Vegetation-related moisture recycling can enhance precipitation during
periods of active growth periods. However, ~~but~~ this ~~is effect is~~ accompanied by ~~an~~ increased ~~in~~ evapotranspiration. As also
suggested by Tian et al. (2026), this represents a redistribution of water within the system rather than a net gain in water
470 resources. Consequently, As a result, the net effect on regional water availability is very small ~~contribution to regional water~~
~~availability remains very small~~ (Fig. 4e; Fig. 5d). This small ~~value net effect~~ should be interpreted ~~carefully in the context of~~
~~the process boundary considered here~~. Specifically ~~Here, we only quantify the vegetation contribution to precipitation generated~~
~~from local evaporation and recycled within the Loess Plateau. this study quantifies only the vegetation-associated contribution~~
~~to precipitation generated from evaporation within the Loess Plateau and recycled back to the plateau itself, whereas internally~~
475 recycled precipitation accounts for only 14.4% of total growing-season precipitation (Fig. 3c). ~~Therefore, Consequently~~, even
a ~~detectable measurable~~ vegetation effect on this ~~internally recycled~~ component becomes ~~strongly substantially~~ diluted when
expressed relative to total precipitation and, ~~therefore~~, total surface water availability.

In addition, the regional mean signal reflects strong compensation between positive and negative vegetation-associated effects.
The strongest positive and negative ~~effects correspond contributions to reach~~ 12.2% and –32.2% of the internally recycled
480 precipitation ~~component~~, respectively (Fig. 3c; Fig. 6a). ~~Thus, Therefore~~, the ~~small low net value arises effect results~~ not only


from the limited share of internal recycling ~~in the total precipitation supply~~, but also from ~~substantial-strong~~ cancellation ~~between opposing responses~~ across vegetation conditions. ~~This compensation is consistent with the nonlinear energy–moisture coupling described by Tian et al. (2026). In that framework, enhanced moisture conversion efficiency coexists with increased evaporative demand. This leads to spatially heterogeneous hydrological responses rather than uniform changes.~~

485 ~~This~~ A small net effect does not imply that vegetation–atmosphere feedbacks are negligible. ~~Instead~~ Rather, it indicates that under the dry ~~background~~ conditions ~~in~~ the Loess Plateau, ~~additional precipitation generated through recycling is generally insufficient to offset increased evapotranspiration caused by vegetation growth. As a result, vegetation does not substantially increase net surface water availability. Instead, it primarily reflects a redistribution of water within a tightly constrained land–atmosphere system.~~

490 ~~the additional precipitation returned through recycling is generally too limited to compensate for the additional water consumption associated with vegetation growth. Therefore, the main implication of our results is not that vegetation substantially increases available water, but that the hydrological benefit of vegetation-related recycling is strongly constrained by background hydroclimatic conditions. In this context, the hydrological benefit of vegetation-associated recycling is strongly limited by background hydroclimatic conditions. This contrasts with different from~~ humid or energy-limited regions, where
495 ~~vegetation-associated-associated moisture-recycling can more readily easily translates into effective-net~~ water gains (Cui et al., 2022; Meier et al., 2021). The comparison ~~highlights the key role of climate setting in regulating vegetation–water coupling efficiency. It also suggests that small mean effects should not be interpreted as hydrological irrelevance. Rather, they reflect strong compensatory feedbacks under water-limited conditions. further highlights the importance of climatic context in shaping vegetation–water interactions.~~

500 More importantly, ~~the influence of~~ vegetation ~~effects~~ on surface water availability shows a clear nonlinear pattern along the LAI_w gradient. Two ~~lines of evidence~~ results support this. ~~interpretation.~~ First, the contribution of vegetation-associated recycled precipitation ~~to SWA~~ shifts from positive under low- LAI_w ~~conditions~~ to negative under high- LAI_w ~~conditions~~. This ~~suggests indicates~~ that the hydrological benefit of ~~moisture~~ recycling is mainly ~~confined to relatively limited to~~ sparse vegetation. ~~It cover and~~ weakens ~~when one~~ vegetation density ~~increases beyond exceeds~~ a moderate level (Fig. 5e; Fig. 6a).
505 Second, the temporal evolution of ~~the~~ positive and negative components shows ~~an asymmetric compensation pattern behavior.~~ Positive contributions ~~remain are~~ small and intermittent, ~~while whereas~~ negative contributions become more persistent with increasing LAI_w (Fig. 4e; Fig. 5e; Fig. 6b). This pattern is consistent with the limited water storage capacity and strong evaporative sensitivity ~~of in~~ the Loess Plateau.

~~Within the broader framework of Tian et al. (2026), this nonlinear behavior reflects a trade-off between enhanced precipitation efficiency and increased evaporative demand under higher vegetation density. Under such conditions, further vegetation growth is more likely to intensify water consumption than to producegenerate sustained gainsincreases in surface water availability. Therefore, the key scientific value of the small regional mean effect therefore lies notless in its magnitude, but than in what it revealings about the hydrological conditions under which vegetation-associatedrelated precipitation recycling~~




shifts from weakly beneficial to increasingly negative under higher LAI_w conditions. shifts from weakly beneficial to persistently adverse.

Figure 6: Vegetation-associated variations in recycled precipitation and surface water availability: (a) recycled precipitation along the LAI_w gradient, (b) Contribution rates of vegetation to SWA under increasing and decreasing LAI_w conditions.

4.3 Robustness of the LAI_w diagnostic framework

Because LAI_w is a derived diagnostic indicator rather than a directly observed physical variable, its formulation may raise concerns about the question of whether the identified hydrological patterns depend strongly on the weighting scheme itself. In particular, weighting LAI by normalized actual evapotranspiration could, in principle, influence the apparent strength of vegetation-hydrological coupling enhance the apparent association with recycled precipitation and thereby affect the inferred threshold-like behavior.

To test this possibility, we repeated the full analytical workflow using the original, unweighted peak-growing-season LAI instead place of LAI_w . This parallel analysis included both the multiple linear regression for P_{rec} and the subsequent-Budyko-based attribution of hydrological effects. We then compared The resulting vegetation-associated contributions to surface water availability (C_{SWA}) were then compared between the LAI -based and LAI_w -based frameworks at both the regional mean scale and across the four vegetation classes defined in Section 3.3.



530 **Figure 7: Vegetation-associated related variations in surface water availability and vegetation contribution under LAI and LAI_w conditions: (a) interannual variations in vegetation-associated related C_{SWA} estimated from LAI and LAI_w , (b) contribution rates of vegetation to SWA under promotive and inhibitive conditions along the mean LAI gradient.**

535 The result comparison shows that the main findings major results are highly robust to the choice of vegetation indicator this substitution. At the regional mean scale, the interannual variations in P_0 and C_{SWA} are nearly identical between derived from the two frameworks are nearly identical, with Pearson correlation coefficients of 0.99996 and 0.99995, respectively (Fig. 7). Their multi-year mean values are likewise very close also show only minor differences. For example, the mean C_{SWA} is 0.107% in the LAI_w -based framework and 0.094% in the LAI -based framework, (with an mean absolute difference of only 0.013 percentage points%). This high level of consistency indicates that the identified observed hydrological response This close agreement suggests that the threshold-like hydrological response identified in this study is not an artifact of the LAI_w weighting scheme formulation. Instead Rather, LAI_w primarily serves mainly acts as a diagnostic rescaling of the underlying vegetation gradient under different evaporative conditions. The while the dominant signal is still controlled governed by the vegetation structure itself mainly acts as a diagnostic rescaling of the original LAI pattern, while the dominant signal appears to be rooted in the underlying vegetation gradient itself. In this sense, the use of LAI_w does not change the qualitative conclusions, but instead provides a slightly more process oriented representation of vegetation conditions under different evaporative backgrounds.

545 Importantly, the observed threshold-like behavior should be interpreted as an empirical real regime transition driven by the interaction between vegetation density and evaporative demand. It is not produced, rather than an artifact of by the LAI_w formulation. Rather, LAI_w helps to does not generate this pattern but serves as a diagnostic tool that reveal how vegetation effects on moisture recycling and surface water availability change under different atmospheric conditions. This suggests that vegetation-water interactions in the Loess Plateau operate within a constrained hydrological regime. The system shifting from a weakly coupled state with slight positive effects to a strongly evaporative-dominated state with negative effects. This transition reflects limited water storage capacity and high evaporative sensitivity in the region.

From a management perspective, this shift implies an implicit water-limited carrying capacity for vegetation restoration. Beyond this level, further increases in vegetation density are more likely to increase evapotranspiration than to improve surface water availability. However, this should not be interpreted as a fixed restoration threshold for restoration. Instead, it represents but as a context-dependent response controlled by hydroclimatic conditions in dryland regions.

4.4 Limitations and Research Prospects

This study provides a quantitative assessment of vegetation-associated precipitation recycling and its contribution to surface water availability during the peak growing season over the Loess Plateau. Several limitations should be acknowledged, but several limitations should be noted.

First, the seasonal focus on July–August. This period represents a peak-coupling regime stage within the monsoon-dominated growing season from (June to September, The growing season), which accounts for 60–70% of annual precipitation and coincides with the annual maxima of precipitation, evapotranspiration, and land–atmosphere coupling (Liang et al., 2015; Zhang et al., 2016). During July–August, This period is characterized by fully developed vegetation is fully developed and enhanced atmospheric moisture exchange is strong. In contrast, whereas coupling is weaker in early summer due to incomplete because vegetation is not yet fully developed. It also weakens and in late summer because of due to soil moisture depletion (Feng et al., 2016; Liang et al., 2015). As a result, Therefore, the present analysis captures the period of strongest vegetation–atmosphere interaction rather than a full-season or annual mean state. Consequently, The identified recycling signals should thus be interpreted as representing the upper bound of seasonal coupling intensity, while a Annual-scale water balance assessments, by contrast, reflect integrated hydrological responses across multiple seasonal phases (Wang et al., 2016). First, the seasonal focus of this study should be explicitly acknowledged. By concentrating on July and August, the analysis targets the period when vegetation activity and land–atmosphere coupling are strongest over the Loess Plateau, but it does not represent the full growing season (June–September) or annual water balance. This seasonal restriction is methodologically necessitated by the WAM-2layers tracking configuration, which requires computational trade-offs between temporal coverage and process resolution. The two-month focus captures the peak of vegetation precipitation recycling interactions but excludes the early growing season (June) when vegetation is still developing and late season (September) when soil moisture depletion may limit evapotranspiration. Consequently, the results should be interpreted as reflecting vegetation-associated hydrological effects during the period of maximum coupling intensity, rather than as representative of the entire growing season. Future work should extend this framework across the full growing season to test the robustness of the threshold-like behavior identified here.

Second, the analysis is limited to local hydrological the analysis focuses on hydroclimatic responses within the Loess Plateau. It itself. It does not explicitly consider the possible effects of vegetation-related moisture export on downwind regions,

nor does it examine how changes in external source areas may influence local precipitation and water availability, and does not explicitly account for consider downwind effects of caused by vegetation-induced moisture export. Vegetation-enhanced evapotranspiration may reduce local water availability. At the same time, it may while also contributing to precipitation in downstream regions elsewhere through atmospheric moisture transport processes (Greve et al., 2025). A more complete assessment would require a coupled local-downwind framework. Such a framework could to quantify the broader full hydroclimatic footprint of vegetation-induced moisture redistribution. evaluation would therefore require an integrated local-downwind framework. In this sense, the present results should be interpreted as reflecting the local hydrological consequences of vegetation-associated recycling within the Loess Plateau, rather than the full hydroclimatic footprint of vegetation-induced moisture redistribution.

Third, uncertainty arises from both the input datasets and the attribution framework. Recycled precipitation and hydrological responses were derived from multiple reanalysis and remote-sensing products, estimated using multiple reanalysis and remote-sensing datasets, and These associated uncertainties uncertainties in these products may propagate through both the moisture-tracking analysis and the Budyko-RF attribution framework analyses. In addition, LAI_w is a diagnostic indicator rather than a directly observed physical variable, and the P0 term also represents reflects a statistical attribution of vegetation-associated recycled precipitation, rather than a strictly causal estimate from process-based perturbation experiments. Therefore, the counterfactual analysis therefore isolates a first-order hydrological effect, but does not resolve secondary feedbacks involving associated with soil moisture, vegetation dynamics, or atmospheric circulation responses. adjustment, or atmospheric circulation.

Future work should address these limitations in three ways. First, identified in this study by extending the analysis should be extended from the peak growing season to full seasonal and annual scales. Second, explicitly incorporating downwind moisture transport should be explicitly incorporated to capture the broader hydroclimatic footprint of vegetation. Third, and improving uncertainty characterization should be improved for in both input datasets and diagnostic attribution frameworks. Together, these efforts would help clarify to better resolve how vegetation-atmosphere feedbacks evolve across spatial gradients and temporal scales in water-limited environments.

5 Conclusions

The Loess Plateau is a water-limited region in arid and semi-arid northern China, where Large-scale vegetation restoration has raised an important question: can vegetation-associated related precipitation recycling meaningfully alleviate regional water scarcity? To address this question, this study combined an WAM-2layers Eulerian atmospheric moisture-tracking model with a random forest-enhanced Budyko framework. We examined peak growing season water cycling over the Loess Plateau in during July and August during from 2000 to 2022, at the period when vegetation-atmosphere coupling is expected to be strongest.

The results show that growing-season precipitation over the Loess Plateau is dominated by land-derived moisture. Terrestrial evaporation contributes 86.5% of total precipitation, while oceanic sources contribute 13.5%. Internal moisture recycling contributes 11.83 mm on average, accounting for 14.4% of total precipitation. It also shows a declining trend of $-0.073 \text{ mm yr}^{-1}$ during the study period. Vegetation-associated processes strengthen land-atmosphere coupling, but their net contribution to the evaporation-precipitation cycle remains small, at about 0.12%. Positive effects (1.16%) are largely offset by negative effects (-0.97%).

During the same period, the Loess Plateau shows a clear drying tendency. Evapotranspiration increased at 0.31 mm yr^{-1} , while surface water availability decreased at -0.34 mm yr^{-1} . At the regional mean scale, the hydrological contribution of vegetation-associated recycled precipitation is limited. On average, it accounts for only 0.059% of precipitation and 0.107% of surface water availability. This weak mean signal is partly explained by two factors. The first is that only the vegetation-associated component of internally recycled precipitation is isolated, which is itself a small fraction of total growing-season precipitation. The second is the partial cancellation between positive and negative vegetation effects. This small regional mean contribution partly reflects the fact that the analysis isolates only the vegetation-associated component of internally recycled precipitation, which itself represents a limited fraction of total growing-season precipitation, and that positive and negative vegetation-associated effects partly offset each other. However, this small regional mean effect the response is not spatially uniform. A threshold-like pattern emerges along the LAI_w gradient. Along the LAI_w gradient, the contribution shows a threshold-like pattern. Under low- LAI_w conditions ($LAI_w < 1$), vegetation-associated recycling contributes shows a weakly but consistently positive contribution to surface water availability, with a mean contribution of about 0.29%. Under high- LAI_w conditions ($LAI_w \geq 2.5$), the contribution becomes persistently negative, with mean values reaching about -0.15% on average and tending to become more negative over time.

Overall, these results suggest that under the water-limited conditions of the Loess Plateau, stronger vegetation-atmosphere coupling does not necessarily lead to increased greater surface water availability. Instead, vegetation mainly reshapes the internal partitioning of water fluxes within a constrained land-atmosphere system governed by monsoon-continental transition dynamics. The core contribution of this work lies not in quantifying mean changes in water availability, but in revealing the hydroclimatic conditions under which vegetation effects shift from weakly positive to persistently negative. These findings emphasize that the hydrological consequences of dryland vegetation restoration must be evaluated in the context of climatic constraints, nonlinear feedbacks, and vegetation density-dependent regime shifts, with the caveat that these conclusions are specific to the peak growing season and that seasonal hydroclimatic boundaries should be factored into restoration assessments. The additional precipitation returned through recycling is generally too small to offset the additional terrestrial water consumption associated with vegetation growth. The hydrological significance of vegetation-related precipitation recycling in this dryland region therefore appears to lie less in increasing mean water availability than in revealing the hydroclimatic conditions under which vegetation effects shift from weakly positive to persistently negative. More broadly, these findings suggest that the hydrological consequences of vegetation restoration in drylands need to be evaluated within the context of background climatic constraints, process boundaries, and vegetation thresholds. While these findings are specific to the peak

~~growing season, they highlight the importance of evaluating vegetation restoration effects within the context of seasonal hydroclimatic constraints and vegetation density thresholds.~~

655 **Code and data availability**

The ERA5 and ERA5-Land reanalysis data used in this study are publicly available from the European Centre for Medium-Range Weather Forecasts (ECMWF) Climate Data Store. The GLEAM v3.6a evapotranspiration data are publicly available from the GLEAM project. The HiQ-LAI dataset is publicly available via Zenodo, and the fractional vegetation cover dataset is available from the National Tibetan Plateau Data Center. The WAM-2layers model used for atmospheric moisture tracking
660 is available from the corresponding published source. The code used for data processing, statistical analysis, and figure generation in this study is available from the corresponding author upon reasonable request.

Supplement link

Author contributions

Jiaxiang Deng conducted the formal analysis, processed the data, and prepared the original manuscript draft. Quan Quan
665 conceived and designed the study, supervised the research, and revised the manuscript. Shuangcheng Tang contributed to the methodology, interpretation of the results, and manuscript revision. Hanbo Yang contributed to the methodology and interpretation of the results. Xiaoyu Song contributed to data curation and manuscript revision. All authors discussed the results and approved the final manuscript.

Competing interests

670 The authors declare that they have no conflict of interest.

Disclaimer

Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and
675 do not necessarily reflect the views of the publisher.

Acknowledgements

The authors would like to express their gratitude to the National Key Research and Development Program of China for their financial support under Grant No. 2024YFF1306901.

Financial support

680 The authors would like to express their gratitude to the National Key Research and Development Program of China for their financial support under Grant No. 2024YFF1306901.

Review statement

The review statement will be added by Copernicus Publications listing the handling editor as well as all contributing referees according to their status anonymous or identified.

685 **References**

An, Q., Liu, L., Cheng, Y., Liu, J., Staal, A., and Huang, G.: Unraveling the role of moisture recycling in water consumption across different ecosystems in China, *J. Hydrol.*, 662, 134020, <https://doi.org/10.1016/j.jhydrol.2025.134020>, 2025.

Cao, M., Wang, W., Wei, J., Forzieri, G., Fetzer, I., and Wang-Erlandsson, L.: Revegetation Impacts on Moisture Recycling and Precipitation Trends in the Chinese Loess Plateau, *Water Resour. Res.*, 60, e2024WR038199, <https://doi.org/10.1029/2024WR038199>, 2024.

690 Cui, J., Lian, X., Huntingford, C., Gimeno, L., Wang, T., Ding, J., He, M., Xu, H., Chen, A., Gentile, P., and Piao, S.: Global water availability boosted by vegetation-driven changes in atmospheric moisture transport, *Nat. Geosci.*, 15, 982–988, <https://doi.org/10.1038/s41561-022-01061-7>, 2022.

Deng, Y., Wang, S., Bai, X., Luo, G., Wu, L., Chen, F., Wang, J., Li, C., Yang, Y., Hu, Z., Tian, S., and Lu, Q.: Vegetation greening intensified soil drying in some semi-arid and arid areas of the world, *Agric. For. Meteorol.*, 292–293, 108103, <https://doi.org/10.1016/j.agrformet.2020.108103>, 2020.

695 Dominguez, F., Hu, H., and Martinez, J. A.: Two-Layer Dynamic Recycling Model (2L-DRM): Learning from Moisture Tracking Models of Different Complexity, *J. Hydrometeorol.*, 21, 3–16, <https://doi.org/10.1175/JHM-D-19-0101.1>, 2020.

van der Ent, R. J., Savenije, H. H. G., Schaeffli, B., and Steele-Dunne, S. C.: Origin and fate of atmospheric moisture over continents, *Water Resour. Res.*, 46, <https://doi.org/10.1029/2010WR009127>, 2010.

700 van der Ent, R. J., Tuinenburg, O. A., Knoche, H.-R., Kunstmann, H., and Savenije, H. H. G.: Should we use a simple or complex model for moisture recycling and atmospheric moisture tracking?, *Hydrol. Earth Syst. Sci.*, 17, 4869–4884, <https://doi.org/10.5194/hess-17-4869-2013>, 2013.

- van der Ent, R. J., Wang-Erlandsson, L., Keys, P. W., and Savenije, H. H. G.: Contrasting roles of interception and transpiration in the hydrological cycle – Part 2: Moisture recycling, *Earth Syst. Dynam.*, 5, 471–489, <https://doi.org/10.5194/esd-5-471-2014>, 2014.
- Feng, X., Fu, B., Piao, S., Wang, S., Ciais, P., Zeng, Z., Lü, Y., Zeng, Y., Li, Y., Jiang, X., and Wu, B.: Revegetation in China’s Loess Plateau is approaching sustainable water resource limits, *Nat. Clim. Change*, 6, 1019–1022, <https://doi.org/10.1038/nclimate3092>, 2016.
- GAO, J., SHI, Y., ZHANG, H., CHEN, X., ZHANG, W., SHEN, W., XIAO, T., and ZHANG, Y.: China regional 250m fractional vegetation cover data set (2000–2024), <https://doi.org/10.11888/Terre.tpd.300330>, 2025.
- Giorgi, F., Coppola, E., and Raffaele, F.: A consistent picture of the hydroclimatic response to global warming from multiple indices: Models and observations, *J. Geophys. Res.-Atmos.*, 119, 11,695–11,708, <https://doi.org/10.1002/2014JD022238>, 2014.
- Greve, P., Schmitt, A. U., Miralles, D. G., McDermid, S., Findell, K. L., García-García, A., and Peng, J.: Observational evidence of increased afternoon rainfall downwind of irrigated areas, *Nat. Commun.*, 16, 3415, <https://doi.org/10.1038/s41467-025-58729-y>, 2025.
- Guo, J., Fan, L., Feng, P., Sun, X., and Xue, S.: Response of vegetation evapotranspiration to landscape pattern changes in an arid region: A case study of the Loess Plateau, China, *CATENA*, 252, 108878, <https://doi.org/10.1016/j.catena.2025.108878>, 2025.
- He, C., Liu, Z., Wu, J., Pan, X., Fang, Z., Li, J., and Bryan, B. A.: Future global urban water scarcity and potential solutions, *Nat. Commun.*, 12, 4667, <https://doi.org/10.1038/s41467-021-25026-3>, 2021a.
- He, Y., Tian, W., Huang, J., Wang, G., Ren, Y., Yan, H., Yu, H., Guan, X., and Hu, H.: The Mechanism of Increasing Summer Water Vapor Over the Tibetan Plateau, *J. Geophys. Res.-Atmos.*, 126, e2020JD034166, <https://doi.org/10.1029/2020JD034166>, 2021b.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- Hu, Q., Jiang, D., Lang, X., and Xu, B.: Moisture sources of the Chinese Loess Plateau during 1979–2009, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 509, 156–163, <https://doi.org/10.1016/j.palaeo.2016.12.030>, 2018.
- Huang, J., Yu, H., Guan, X., Wang, G., and Guo, R.: Accelerated dryland expansion under climate change, *Nat. Clim. Change*, 6, 166–171, <https://doi.org/10.1038/nclimate2837>, 2016.
- Huang, J., Li, Y., Fu, C., Chen, F., Fu, Q., Dai, A., Shinoda, M., Ma, Z., Guo, W., Li, Z., Zhang, L., Liu, Y., Yu, H., He, Y., Xie, Y., Guan, X., Ji, M., Lin, L., Wang, S., and Wang, G.: Dryland Climate Change: Recent Progress and Challenges: Dryland Climate Change, *Reviews of Geophysics*, 55, <https://doi.org/10.1002/2016RG000550>, 2017.

- Jiao, D., Xu, N., Yang, F., and Xu, K.: Evaluation of spatial-temporal variation performance of ERA5 precipitation data in China, *Sci. Rep.*, 11, 17956, <https://doi.org/10.1038/s41598-021-97432-y>, 2021.
- 740 Kalverla, P., Benedict, I., Weijenborg, C., and van der Ent, R. J.: Atmospheric moisture tracking with WAM2layers v3, *Geosci. Model Dev.*, 18, 4335–4352, <https://doi.org/10.5194/gmd-18-4335-2025>, 2025.
- Kummu, M., Guillaume, J. H. A., de Moel, H., Eisner, S., Flörke, M., Porkka, M., Siebert, S., Veldkamp, T. I. E., and Ward, P. J.: The world’s road to water scarcity: shortage and stress in the 20th century and pathways towards sustainability, *Sci. Rep.*, 6, 38495, <https://doi.org/10.1038/srep38495>, 2016.
- 745 Lai, J., Li, Y., Chen, J., Niu, G.-Y., Lin, P., Li, Q., Wang, L., Han, J., Luo, Z., and Sun, Y.: Massive crop expansion threatens agriculture and water sustainability in northwestern China, *Environ. Res. Lett.*, 17, 034003, <https://doi.org/10.1088/1748-9326/ac46e8>, 2022.
- Li, C., Fu, B., Wang, S., Stringer, L. C., Wang, Y., Li, Z., Liu, Y., and Zhou, W.: Drivers and impacts of changes in China’s drylands, *Nature Reviews Earth & Environment*, 2, 858–873, <https://doi.org/10.1038/s43017-021-00226-z>, 2021.
- 750 Li, R., Fan, B., Jin, Z., Hao, M., Liu, H., Zhang, Y., Jiang, Y., Yao, H., Lin, K., and Guo, L.: Divergent rainfall infiltration patterns on the Chinese Loess Plateau between growing and non-growing seasons after long-term revegetation, *J. Hydrol.*, 641, 131816, <https://doi.org/10.1016/j.jhydrol.2024.131816>, 2024.
- Li, Z., Wang, S., Li, C., Ye, C., Gao, D., and Chen, P.: The trend shift caused by ecological restoration accelerates the vegetation greening of China’s drylands since the 1980s, *Environ. Res. Lett.*, 17, <https://doi.org/10.1088/1748-9326/ac6002>, 2022.
- 755 Liu, R., Yang, L., Shi, Z., and Xing, M.: Assessment of the applicability of multi-satellite precipitation products on the Loess Plateau over the past four decades, *International Journal of Applied Earth Observation and Geoinformation*, 141, 104634, <https://doi.org/10.1016/j.jag.2025.104634>, 2025a.
- Liu, W., Liu, P., Cheng, L., Zhang, X., and Zhou, L.: Identification of time-varying parameters of a monthly Budyko function in US MOPEX catchments and its implications, *J. Hydrol.: Reg. Stud.*, 59, 102348, <https://doi.org/10.1016/j.ejrh.2025.102348>, 2025b.
- 760 Martens, B., Miralles, D. G., Lievens, H., van der Schalie, R., de Jeu, R. A. M., Fernández-Prieto, D., Beck, H. E., Dorigo, W. A., and Verhoest, N. E. C.: GLEAM v3: satellite-based land evaporation and root-zone soil moisture, *Geosci. Model Dev.*, 10, 1903–1925, <https://doi.org/10.5194/gmd-10-1903-2017>, 2017.
- 765 Meier, R., Schwaab, J., Seneviratne, S. I., Sprenger, M., Lewis, E., and Davin, E. L.: Empirical estimate of forestation-induced precipitation changes in Europe, *Nat. Geosci.*, 14, 473–478, <https://doi.org/10.1038/s41561-021-00773-6>, 2021.
- Piao, S., Wang, X., Park, T., Chen, C., Lian, X., He, Y., Bjerke, J. W., Chen, A., Ciais, P., Tømmervik, H., Nemani, R. R., and Myneni, R. B.: Characteristics, drivers and feedbacks of global greening, *Nature Reviews Earth & Environment*, 1, 14–27, <https://doi.org/10.1038/s43017-019-0001-x>, 2020.
- 770 Sterling, S. M., Ducharne, A., and Polcher, J.: The impact of global land-cover change on the terrestrial water cycle, *Nat. Clim. Change*, 3, 385–390, <https://doi.org/10.1038/nclimate1690>, 2013.

- Stevenson, S., Coats, S., Touma, D., Cole, J., Lehner, F., Fasullo, J., and Otto-Bliesner, B.: Twenty-first century hydroclimate: A continually changing baseline, with more frequent extremes, *Proceedings of the National Academy of Sciences*, 119, e2108124119, <https://doi.org/10.1073/pnas.2108124119>, 2022.
- 775 Tabari, H., Hosseinzadehtalaei, P., Thiery, W., and Willems, P.: Amplified Drought and Flood Risk Under Future Socioeconomic and Climatic Change, *Earth's Future*, 9, e2021EF002295, <https://doi.org/10.1029/2021EF002295>, 2021.
- Te Wierik, S. A., Keune, J., Miralles, D. G., Gupta, J., Artzy-Randrup, Y. A., Gimeno, L., Nieto, R., and Cammeraat, L. H.: The Contribution of Transpiration to Precipitation Over African Watersheds, *Water Resour. Res.*, 58, e2021WR031721, <https://doi.org/10.1029/2021WR031721>, 2022.
- 780 Tian, L., Yang, Y., Feng, J., Li, Y., and Zhang, B.: Energy-mediated feedbacks of vegetation greening enhance precipitation efficiency and sustain water yield in global semi-arid regions, *Nat. Water*, 4, 559–570, <https://doi.org/10.1038/s44221-026-00631-y>, 2026.
- Tian, L., Zhang, B., Chen, S., Wang, X., Ma, X., and Pan, B.: Large-Scale Afforestation Enhances Precipitation by Intensifying the Atmospheric Water Cycle Over the Chinese Loess Plateau, *J. Geophys. Res.-Atmos.*, 127, e2022JD036738, <https://doi.org/10.1029/2022JD036738>, 2022.
- 785 Tuinenburg, O. A., Bosmans, J. H. C., and Staal, A.: The global potential of forest restoration for drought mitigation, *Environ. Res. Lett.*, 17, 034045, <https://doi.org/10.1088/1748-9326/ac55b8>, 2022.
- Wang, X., Wang, B., and Xu, X.: Effects of large-scale climate anomalies on trends in seasonal precipitation over the Loess Plateau of China from 1961 to 2016, *Ecological Indicators*, 107, 105643, <https://doi.org/10.1016/j.ecolind.2019.105643>, 2019.
- 790 Wang, X., Zhang, Z., Zhang, B., Tian, L., Tian, J., Arnault, J., Kunstmann, H., and He, C.: Quantifying the Impact of Land Use and Land Cover Change on Moisture Recycling With Convection-Permitting WRF-Tagging Modeling in the Agro-Pastoral Ecotone of Northern China, *J. Geophys. Res.-Atmos.*, 128, e2022JD038421, <https://doi.org/10.1029/2022JD038421>, 2023.
- Wei, J. and Dirmeyer, P. A.: Sensitivity of land precipitation to surface evapotranspiration: a nonlocal perspective based on water vapor transport, *Geophysical Research Letters*, 46, 12588–12597, <https://doi.org/10.1029/2019GL085613>, 2019.
- 795 Woods, R.: The relative roles of climate, soil, vegetation and topography in determining seasonal and long-term catchment dynamics, *Advances in Water Resources*, 26, 295–309, [https://doi.org/10.1016/S0309-1708\(02\)00164-1](https://doi.org/10.1016/S0309-1708(02)00164-1), 2003.
- Xie, D., Zhang, Y., Zhang, M., Tian, Y., Cao, Y., Mei, Y., Liu, S., and Zhong, D.: Hydrological impacts of vegetation cover change in China through terrestrial moisture recycling, *Science of The Total Environment*, 915, 170015, <https://doi.org/10.1016/j.scitotenv.2024.170015>, 2024.
- 800 Yan, K., Wang, J., Peng, R., Yang, K., Chen, X., Yin, G., Dong, J., Weiss, M., Pu, J., and Myneni, R. B.: HiQ-LAI: a high-quality reprocessed MODIS leaf area index dataset with better spatiotemporal consistency from 2000 to 2022, *Earth System Science Data*, 16, 1601–1622, <https://doi.org/10.5194/essd-16-1601-2024>, 2024.
- Yang, D., Shao, W., Yeh, P. J.-F., Yang, H., Kanae, S., and Oki, T.: Impact of vegetation coverage on regional water balance in the nonhumid regions of China, *Water Resour. Res.*, 45, <https://doi.org/10.1029/2008WR006948>, 2009.
- 805

- Yang, X., Yong, B., Ren, L., Zhang, Y., and Long, D.: Multi-scale validation of GLEAM evapotranspiration products over China via ChinaFLUX ET measurements, *Int. J. Remote Sens.*, 38, 5688–5709, <https://doi.org/10.1080/01431161.2017.1346400>, 2017.
- 810 Yang, Y., Donohue, R. J., McVicar, T. R., and Roderick, M. L.: An analytical model for relating global terrestrial carbon assimilation with climate and surface conditions using a rate limitation framework, *Geophysical Research Letters*, 42, 9825–9835, <https://doi.org/10.1002/2015GL066835>, 2015.
- Yang, Z., Dong, Q., Zhang, X., Fang, L., Zhu, X., Chen, K., and Mendingo, E. M.: Interpretable Budyko-constrained machine learning framework for monthly runoff attribution in U.S. CAMELS basins, *J. Hydrol.*, 670, 135209, <https://doi.org/10.1016/j.jhydrol.2026.135209>, 2026.
- 815 Yokoo, Y., Sivapalan, M., and Oki, T.: Investigating the roles of climate seasonality and landscape characteristics on mean annual and monthly water balances, *J. Hydrol.*, 357, 255–269, <https://doi.org/10.1016/j.jhydrol.2008.05.010>, 2008.
- Yosef, G., Walko, R., Avisar, R., Tatarinov, F., Rotenberg, E., and Yakir, D.: Large-scale semi-arid afforestation can enhance precipitation and carbon sequestration potential, *Sci. Rep.*, 8, 996, <https://doi.org/10.1038/s41598-018-19265-6>, 2018.
- 820 Zhang, B.: Albedo-driven hydroclimatic impacts of large-scale vegetation restoration should not be overlooked, *Nat. Water*, 3, 358–359, <https://doi.org/10.1038/s44221-025-00422-x>, 2025.
- Zhang, B., Tian, L., Yang, Y., and He, X.: Revegetation Does Not Decrease Water Yield in the Loess Plateau of China, *Geophysical Research Letters*, 49, e2022GL098025, <https://doi.org/10.1029/2022GL098025>, 2022a.
- Zhang, J., Wang, S., He, Y., Ren, Y., and Huang, J.: Contribution of the Precipitation-Recycling Process to the Wetting Trend in Xinjiang, China, *J. Geophys. Res.-Atmos.*, 127, e2021JD036407, <https://doi.org/10.1029/2021JD036407>, 2022b.
- 825 Zhang, S., Yang, D., Yang, Y., Piao, S., Yang, H., Lei, H., and Fu, B.-J.: Excessive Afforestation and Soil Drying on China's Loess Plateau, *J. Geophys. Res.-Biogeosci.*, 123, <https://doi.org/10.1002/2017JG004038>, 2018.
- Zhang, Y., Li, C., Chiew, F. H. S., Post, D. A., Zhang, X., Ma, N., Tian, J., Kong, D., Leung, L. R., Yu, Q., Shi, J., and Liu, C.: Southern Hemisphere dominates recent decline in global water availability, *Science*, 382, 579–584, <https://doi.org/10.1126/science.adh0716>, 2023.
- 830 Zhang, Y., Zhao, X., Gong, J., Luo, F., and Pan, Y.: Effectiveness and driving mechanism of ecological restoration efforts in China from 2009 to 2019, *Science of The Total Environment*, 910, 168676, <https://doi.org/10.1016/j.scitotenv.2023.168676>, 2024.
- An, Q., Liu, L., Cheng, Y., Liu, J., Staal, A., Huang, G., 2025. Unraveling the role of moisture recycling in water consumption across different ecosystems in China. *Journal of Hydrology* 662, 134020. <https://doi.org/10.1016/j.jhydrol.2025.134020>
- 835 Budyko, M.I., 1974. *Climate and life*. Academic, San Diego, Calif.
- Cao, M., Wang, W., Wei, J., Forzieri, G., Fetzer, I., Wang Erlandsson, L., 2024. Revegetation impacts on moisture recycling and precipitation trends in the Chinese Loess Plateau. *Water Resources Research* 60, e2024WR038199. <https://doi.org/10.1029/2024WR038199>

- Cui, J., Lian, X., Huntingford, C., Gimeno, L., Wang, T., Ding, J., He, M., Xu, H., Chen, A., Gentile, P., Piao, S., 2022. Global water availability boosted by vegetation driven changes in atmospheric moisture transport. *Nature Geoscience* 15, 982–988. <https://doi.org/10.1038/s41561-022-01061-7>
- Deng, Y., Wang, S., Bai, X., Luo, G., Wu, L., Chen, F., Wang, J., Li, C., Yang, Y., Hu, Z., Tian, S., Lu, Q., 2020. Vegetation greening intensified soil drying in some semi-arid and arid areas of the world. *Agricultural and Forest Meteorology* 292–293, 108103. <https://doi.org/10.1016/j.agrformet.2020.108103>
- Dominguez, F., Hu, H., Martinez, J.A., 2020. Two-layer dynamic recycling model (2L-DRM): Learning from moisture tracking models of different complexity. *Journal of Hydrometeorology* 21, 3–16. <https://doi.org/10.1175/JHM-D-19-0101.1>
- Feng, X., Fu, B., Piao, S., Wang, S., Ciais, P., Zeng, Z., Lü, Y., Zeng, Y., Li, Y., Jiang, X., Wu, B., 2016. Revegetation in China's Loess Plateau is approaching sustainable water resource limits. *Nature Climate Change* 6, 1019–1022. <https://doi.org/10.1038/nclimate3092>
- Gao, J.X., S.Y., Zhang, H.W., Chen, X.H., Zhang, W.G., Shen, W.M., Xiao, T., Zhang, Y.H., 2024. China regional 250m fractional vegetation cover data set (2000–2023). National Tibetan Plateau Data Center. <https://doi.org/10.11888/Terre.tpdc.300330>
- Giorgi, F., Coppola, E., Raffaele, F., 2014. A consistent picture of the hydroclimatic response to global warming from multiple indices: Models and observations. *Journal of Geophysical Research: Atmospheres* 119, 11695–11708. <https://doi.org/10.1002/2014JD022238>
- Greve, P., Schmitt, A.U., Miralles, D.G., McDermid, S., Findell, K.L., García-García, A., Peng, J., 2025. Observational evidence of increased afternoon rainfall downwind of irrigated areas. *Nature Communications* 16, 3415. <https://doi.org/10.1038/s41467-025-58729-y>
- Guo, J., Fan, L., Feng, P., Sun, X., Xue, S., 2025. Response of vegetation evapotranspiration to landscape pattern changes in an arid region: A case study of the Loess Plateau, China. *CATENA* 252, 108878. <https://doi.org/10.1016/j.catena.2025.108878>
- He, C., Liu, Z., Wu, J., Pan, X., Fang, Z., Li, J., Bryan, B.A., 2021. Future global urban water scarcity and potential solutions. *Nature Communications* 12, 4667. <https://doi.org/10.1038/s41467-021-25026-3>
- He, Y., Tian, W., Huang, J., Wang, G., Ren, Y., Yan, H., Yu, H., Guan, X., Hu, H., 2021. The mechanism of increasing summer water vapor over the Tibetan Plateau. *Journal of Geophysical Research: Atmospheres* 126, e2020JD034166. <https://doi.org/10.1029/2020JD034166>
- Hersbach, H., et al., 2020. The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society* 146, 1999–2049. <https://doi.org/10.1002/qj.3803>
- Hu, Q., Jiang, D., Lang, X., Xu, B., 2018. Moisture sources of the Chinese Loess Plateau during 1979–2009. *Palaeogeography, Palaeoclimatology, Palaeoecology* 509, 156–163. <https://doi.org/10.1016/j.palaeo.2016.12.030>
- Huang, J., Li, Y., Fu, C., Chen, F., Fu, Q., Dai, A., Shinoda, M., Ma, Z., Guo, W., Li, Z., Zhang, L., Liu, Y., Yu, H., He, Y., Xie, Y., Guan, X., Ji, M., Lin, L., Wang, S., Wang, G., 2017. Dryland climate change: Recent progress and challenges: Dryland climate change. *Reviews of Geophysics* 55. <https://doi.org/10.1002/2016RG000550>

- Huang, J., Yu, H., Guan, X., Wang, G., Guo, R., 2016. Accelerated dryland expansion under climate change. *Nature Climate Change* 6, 166–171. <https://doi.org/10.1038/nclimate2837>
- 875 Jiao, D., Xu, N., Yang, F., Xu, K., 2021. Evaluation of spatial-temporal variation performance of ERA5 precipitation data in China. *Scientific Reports* 11, 17956. <https://doi.org/10.1038/s41598-021-97432-y>
- Kalverla, P., Benedict, I., Weijenborg, C., van der Ent, R.J., 2025. Atmospheric moisture tracking with WAM2layers v3. *Geoscientific Model Development* 18, 4335–4352. <https://doi.org/10.5194/gmd-18-4335-2025>
- Kummu, M., Guillaume, J.H.A., de Moel, H., Eisner, S., Flörke, M., Porkka, M., Siebert, S., Veldkamp, T.I.E., Ward, P.J.,
880 2016. The world's road to water scarcity: shortage and stress in the 20th century and pathways towards sustainability. *Scientific Reports* 6, 38495. <https://doi.org/10.1038/srep38495>
- Lai, J., Li, Y., Chen, J., Niu, G. Y., Lin, P., Li, Q., Wang, L., Han, J., Luo, Z., Sun, Y., 2022. Massive crop expansion threatens agriculture and water sustainability in northwestern China. *Environmental Research Letters* 17, 034003. <https://doi.org/10.1088/1748-9326/ac46e8>
- 885 Li, C., Fu, B., Wang, S., Stringer, L.C., Wang, Y., Li, Z., Liu, Y., Zhou, W., 2021. Drivers and impacts of changes in China's drylands. *Nature Reviews Earth & Environment* 2, 858–873. <https://doi.org/10.1038/s43017-021-00226-z>
- Li, R., Fan, B., Jin, Z., Hao, M., Liu, H., Zhang, Y., Jiang, Y., Yao, H., Lin, K., Guo, L., 2024. Divergent rainfall infiltration patterns on the Chinese Loess Plateau between growing and non-growing seasons after long-term revegetation. *Journal of Hydrology* 641, 131816. <https://doi.org/10.1016/j.jhydrol.2024.131816>
- 890 Li, Z., Wang, S., Li, C., Ye, C., Gao, D., Chen, P., 2022. The trend shift caused by ecological restoration accelerates the vegetation greening of China's drylands since the 1980s. *Environmental Research Letters* 17(4), 044062. <https://doi.org/10.1088/1748-9326/ac55b8>
- Liu, R., Yang, L., Shi, Z., Xing, M., 2025. Assessment of the applicability of multi-satellite precipitation products on the Loess Plateau over the past four decades. *International Journal of Applied Earth Observation and Geoinformation* 141, 104634. <https://doi.org/10.1016/j.jag.2025.104634>
- 895 Liu, W., Liu, P., Cheng, L., Zhang, X., Zhou, L., 2025. Identification of time-varying parameters of a monthly Budyko function in US-MOPEX catchments and its implications. *Journal of Hydrology: Regional Studies* 59, 102348. <https://doi.org/10.1016/j.ejrh.2025.102348>
- Martens, B., Miralles, D.G., Lievens, H., van der Schalie, R., de Jeu, R.A.M., Fernández-Prieto, D., Beck, H.E., Dorigo, W.A.,
900 Verhoest, N.E.C., 2017. GLEAM v3: satellite-based land evaporation and root zone soil moisture. *Geoscientific Model Development* 10, 1903–1925. <https://doi.org/10.5194/gmd-10-1903-2017>
- Meier, R., Schwaab, J., Seneviratne, S.I., Sprenger, M., Lewis, E., Davin, E.L., 2021. Empirical estimate of forestation-induced precipitation changes in Europe. *Nature Geoscience* 14, 473–478. <https://doi.org/10.1038/s41561-021-00773-6>
- Piao, S., Wang, X., Park, T., Chen, C., Lian, X., He, Y., Bjerke, J.W., Chen, A., Ciais, P., Tømmervik, H., Nemani, R.R.,
905 Myneni, R.B., 2020. Characteristics, drivers and feedbacks of global greening. *Nature Reviews Earth & Environment* 1, 14–27. <https://doi.org/10.1038/s43017-019-0001-x>

Sterling, S.M., Ducharme, A., Polcher, J., 2013. The impact of global land-cover change on the terrestrial water cycle. *Nature Climate Change* 3, 385–390. <https://doi.org/10.1038/nclimate1690>

910 Stevenson, S., Coats, S., Touma, D., Cole, J., Lehner, F., Fasullo, J., Otto-Bliesner, B., 2022. Twenty-first century hydroclimate: A continually changing baseline, with more frequent extremes. *Proceedings of the National Academy of Sciences* 119, e2108124119. <https://doi.org/10.1073/pnas.2108124119>

Tabari, H., Hosseinzadehtalaei, P., Thiery, W., Willems, P., 2021. Amplified drought and flood risk under future socioeconomic and climatic change. *Earth's Future* 9, e2021EF002295. <https://doi.org/10.1029/2021EF002295>

915 Te Wierik, S.A., Keune, J., Miralles, D.G., et al., 2022. The contribution of transpiration to precipitation over African watersheds. *Water Resources Research* 58(11), e2021WR031721. <https://doi.org/10.1029/2021WR031721>

Tian, L., Zhang, B., Chen, S., Wang, X., Ma, X., Pan, B., 2022. Large-scale afforestation enhances precipitation by intensifying the atmospheric water cycle over the Chinese Loess Plateau. *Journal of Geophysical Research: Atmospheres* 127, e2022JD036738. <https://doi.org/10.1029/2022JD036738>

920 Tuinenburg, O.A., Bosmans, J.H.C., Staal, A., 2022. The global potential of forest restoration for drought mitigation. *Environmental Research Letters* 17, 034045. <https://doi.org/10.1088/1748-9326/ac55b8>

van der Ent, R.J., Savenije, H.H.G., Schaeffli, B., Steele-Dunne, S.C., 2010. Origin and fate of atmospheric moisture over continents. *Water Resources Research* 46. <https://doi.org/10.1029/2010WR009127>

van der Ent, R.J., Tuinenburg, O.A., Knoche, H. R., Kunstmann, H., Savenije, H.H.G., 2013. Should we use a simple or complex model for moisture recycling and atmospheric moisture tracking? *Hydrology and Earth System Sciences* 17, 4869–4884. <https://doi.org/10.5194/hess-17-4869-2013>

925 van der Ent, R.J., Wang Erlandsson, L., Keys, P.W., Savenije, H.H.G., 2014. Contrasting roles of interception and transpiration in the hydrological cycle—Part 2: Moisture recycling. *Earth System Dynamics* 5, 471–489. <https://doi.org/10.5194/esd-5-471-2014>

Wang, X., Wang, B., Xu, X., 2019. Effects of large-scale climate anomalies on trends in seasonal precipitation over the Loess Plateau of China from 1961 to 2016. *Ecological Indicators* 107, 105643. <https://doi.org/10.1016/j.ecolind.2019.105643>

930 Wang, X., Zhang, Z., Zhang, B., Tian, L., Tian, J., Arnault, J., Kunstmann, H., He, C., 2023. Quantifying the impact of land use and land cover change on moisture recycling with convection-permitting WRF-tagging modeling in the agro-pastoral ecotone of northern China. *Journal of Geophysical Research: Atmospheres* 128, e2022JD038421. <https://doi.org/10.1029/2022JD038421>

935 Wei, J., Dirmeyer, P.A., 2019. Sensitivity of land precipitation to surface evapotranspiration: a nonlocal perspective based on water vapor transport. *Geophysical Research Letters* 46, 12588–12597. <https://doi.org/10.1029/2019GL085613>

Woods, R., 2003. The relative roles of climate, soil, vegetation and topography in determining seasonal and long-term catchment dynamics. *Advances in Water Resources* 26, 295–309. [https://doi.org/10.1016/S0309-1708\(02\)00164-1](https://doi.org/10.1016/S0309-1708(02)00164-1)

- Xie, D., Zhang, Y., Zhang, M., Tian, Y., Cao, Y., Mei, Y., Liu, S., Zhong, D., 2024. Hydrological impacts of vegetation cover change in China through terrestrial moisture recycling. *Science of The Total Environment* 915, 170015. <https://doi.org/10.1016/j.scitotenv.2024.170015>
- Yan, K., Wang, J., Peng, R., Yang, K., Chen, X., Yin, G., Dong, J., Weiss, M., Pu, J., Myneni, R.B., 2024. HiQ LAI: a high-quality reprocessed MODIS leaf area index dataset with better spatiotemporal consistency from 2000 to 2022. *Earth System Science Data* 16, 1601–1622. <https://doi.org/10.5194/essd-16-1601-2024>
- Yang, D., Shao, W., Yeh, P., Yang, H., Kanae, S., Oki, T., 2009. Impact of vegetation coverage on regional water balance in the nonhumid regions of China. *Water Resources Research* 45. <https://doi.org/10.1029/2008WR006948>
- Yang, Z., Dong, Q., Zhang, X., Fang, L., Zhu, X., Chen, K., Mendiondo, E.M., 2026. Interpretable Budyko-constrained machine learning framework for monthly runoff attribution in U.S. CAMELS basins. *Journal of Hydrology* 670, 135209. <https://doi.org/10.1016/j.jhydrol.2026.135209>
- Yang, X., Yong, B., Ren, L., Zhang, Y., Long, D., 2017. Multi-scale validation of GLEAM evapotranspiration products over China via ChinaFLUX ET measurements. *International Journal of Remote Sensing* 38, 5688–5709. <https://doi.org/10.1080/01431161.2017.1346400>
- Yang, Y., Donohue, R., McVicar, T., Roderick, M., 2015. An analytical model for relating global terrestrial carbon assimilation with climate and surface conditions using a rate limitation framework. *Geophysical Research Letters* 42. <https://doi.org/10.1002/2015GL066835>
- Yokoo, Y., Sivapalan, M., Oki, T., 2008. Investigating the roles of climate seasonality and landscape characteristics on mean annual and monthly water balances. *Journal of Hydrology* 357, 255–269. <https://doi.org/10.1016/j.jhydrol.2008.05.010>
- Yosef, G., Walko, R., Avisar, R., Tatarinov, F., Rotenberg, E., Yakir, D., 2018. Large-scale semi-arid afforestation can enhance precipitation and carbon sequestration potential. *Scientific Reports* 8, 996. <https://doi.org/10.1038/s41598-018-19265-6>
- Zhang, B., Tian, L., Yang, Y., He, X., 2022. Revegetation does not decrease water yield in the Loess Plateau of China. *Geophysical Research Letters* 49, e2022GL098025. <https://doi.org/10.1029/2022GL098025>
- Zhang, J., Wang, S., He, Y., Ren, Y., Huang, J., 2022. Contribution of the precipitation recycling process to the wetting trend in Xinjiang, China. *Journal of Geophysical Research: Atmospheres* 127, e2021JD036407. <https://doi.org/10.1029/2021JD036407>
- Zhang, S., Yang, D., Yang, Y., Piao, S., Yang, H., Lei, H., Fu, B. J., 2018. Excessive afforestation and soil drying on China's Loess Plateau. *Journal of Geophysical Research: Biogeosciences* 123. <https://doi.org/10.1002/2017JG004038>
- Zhang, Y., Li, C., Chiew, F.H.S., Post, D.A., Zhang, X., Ma, N., Tian, J., Kong, D., Leung, L.R., Yu, Q., Shi, J., Liu, C., 2023. Southern hemisphere dominates recent decline in global water availability. *Science* 382, 579–584. <https://doi.org/10.1126/science.adh0716>
- Zhang, Y., Zhao, X., Gong, J., Luo, F., Pan, Y., 2024. Effectiveness and driving mechanism of ecological restoration efforts in China from 2009 to 2019. *Science of The Total Environment* 910, 168676. <https://doi.org/10.1016/j.scitotenv.2023.168676>