



Synthesis of the tethered balloon system and other TRACER campaign measurements elucidates aerosol property profiles

Fan Mei¹, Jian Wang², Israel Silber¹, Nurun Nahar Lata³, Gregory W. Vandergrift³, Jing Li², Bo Chen^{4,7}, Sarah D. Brooks⁴, Michael P. Jensen⁵, Min Deng⁵, Damao Zhang¹, Darielle Dexheimer⁶, Beat Schmid¹, Zezhen Cheng³, Swarup China³

¹Earth and Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland, WA, 99352, USA

²Energy, Environmental & Chemical Engineering Department, Washington University in St. Louis, St. Louis, MO, 63130, USA

³Environmental Molecular Sciences Laboratory, Pacific Northwest National Laboratory, Richland, WA, 99352, USA

⁴Department of Atmospheric Sciences, Texas A&M University, College Station, 77843, USA

⁵Environmental and Climate Sciences Department, Brookhaven National Laboratory, Upton, NY, 11973, USA

⁶Sandia National Laboratories, Albuquerque, NM, 87185, USA

⁷Department of Atmospheric Science, Colorado State University, Fort Collins, CO, 80521, USA

Correspondence to: Fan Mei (fan.mei@pnnl.gov)

Abstract. Coastal urban environments exhibit strong vertical and horizontal heterogeneity in aerosol properties, complicating process-level understanding of aerosol–cloud interactions. This study analyzes tethered balloon system (TBS) measurements from 149 flights during summer over the greater Houston, Texas, region as part of the DOE Atmospheric Radiation Measurement (ARM) Tracking Aerosol Convection interactions ExpeRiment (TRACER) campaign. We characterize the vertical structure of aerosol number concentrations, size distributions, and inferred cloud condensation nuclei (CCN) concentrations. Air mass history was classified using back-trajectory analysis and k-means clustering into three clusters: (1) marine-influenced, (2) mixed marine and urban emissions, and (3) urban/anthropogenic and long-range transported aerosols. CCN concentrations are estimated from observed size distributions using κ -Köhler theory. The resulting profiles show pronounced vertical variability across clusters, strongly modulated by boundary-layer depth and coastal circulations, leading to substantial variability in the aerosol population available for cloud activation. The marine cluster showed the lowest concentrations, with CCN at 0.8% supersaturation below $1,000\text{ cm}^{-3}$, while urban and mixed clusters displayed higher concentrations and more complex layering. Profiles influenced by the mixed marine–urban cluster frequently exhibit decoupling between near-surface aerosol and elevated layers, including enhanced accumulation-mode number aloft, consistent with prior TBS-based compositional studies. A September 6–7, 2022 case study demonstrates that mesoscale transport can simultaneously transform both the thermodynamic environment and aerosol population, highlighting the importance of constraining boundary-layer dynamics and air mass origin before attributing cloud changes to aerosol effects in complex coastal environments.



1. Introduction

Atmospheric aerosols are tiny solid or liquid particles from both natural sources and human activities suspended in the atmosphere. These aerosols influence the Earth's system in multiple ways, and their direct radiative forcing can be positive (warming) or negative (cooling). For example, sulfate aerosols primarily scatter sunlight, leading to cooling, while black carbon aerosols absorb sunlight, causing warming. (Huang et al., 2014; Ervens et al., 2011; Andreae et al., 2005). The net effect of aerosols on radiative forcing is complex and also depends on their properties and interactions with clouds (Fuzzi et al., 2015; Lohmann and Feichter, 1997; Lohmann and Hoose, 2009; Twomey, 1977). This dual role makes aerosols a significant factor in atmospheric process modeling and weather prediction, and has also stimulated an enormous number of research publications (Fuzzi et al., 2015). Acting as cloud condensation nuclei (CCN), atmospheric aerosols facilitate a critical process in cloud formation by catalyzing the condensation of water vapor into cloud droplets. The efficiency of this process depends on the size, composition, mixing state, and concentration of the aerosol particles (Rosenfeld et al., 2019; Schmale et al., 2017; Rosenfeld et al., 2014). Overall, aerosols play a crucial role in Earth's systems and significantly influence severe weather monitoring (Tsimpidi et al., 2024; Mushtaq et al., 2022).

Due to the complexity of aerosol properties and their roles in the atmospheric processes, accurate vertical profiles of aerosols are essential for quantifying radiative forcing, assessing air quality and transport, and understanding aerosol–cloud interactions. Remote sensing techniques initially emerged as crucial solutions for measuring aerosol vertical distributions and have since become essential tools for applications across diverse geographic regions, facilitating advanced research and applications. Elastic-backscatter lidar inversions, pioneered by Klett and Fernald (Fernald, 1984; Klett, 1981), laid the foundation for aerosol profiling by retrieving range-dependent backscatter and extinction using an assumed lidar ratio. While computationally efficient and widely applied, these methods carry inherent uncertainties due to their dependence on external assumptions about the lidar ratio (Chen et al., 2025; Burton et al., 2012; Fernald, 1984; Klett, 1981). Raman (inelastic) lidar significantly improved retrievals by utilizing Raman-shifted molecular returns to directly derive aerosol extinction and backscatter, eliminating the need for an assumed lidar ratio and enhancing profile accuracy (Ansmann et al., 2012; Ansmann and Müller, 2005). High-Spectral-Resolution Lidar (HSRL) further advanced profiling capabilities by separating molecular and aerosol scattering components, enabling precise direct retrievals of extinction, backscatter, and the lidar ratio. These advancements have established lidar approaches as critical tools in field campaigns and satellite validation efforts, underpinning a deeper understanding of aerosol vertical structure and its global impacts (Dmitrovic et al., 2024; Burton et al., 2012; Hair et al., 2008).

While remote sensing provides broader spatial coverage and long-term monitoring, airborne measurements can offer high-resolution in-situ data that comprehensively characterize vertical profiles of aerosol properties, including concentration, size, and chemical composition. Airborne platforms such as piloted aircraft, uncrewed aerial systems (UAS), and tethered balloon systems (TBS) play a critical role in accurately measuring aerosol vertical profiles, providing essential insights into their spatial distribution, composition, transport, and atmospheric processing (Mei et al., 2020; Chyliński et al., 2018; Geiß et



65 al., 2017; Brady et al., 2016; Ferrero et al., 2016; Corrigan et al., 2008; Ramanathan et al., 2001; Anderson et al., 1996). Recently, studies like the Department of Energy (DOE) Atmosphere Radiation Measurement (ARM) TBS and UAS
70 deployments (Mei et al., 2025; Chen et al., 2025; Creamean et al., 2025; Creamean et al., 2021) and the ACTIVATE (Aerosol Cloud meTeorology Interactions oVer the western ATlantic Experiment) aircraft campaigns (Ajayi et al., 2024) collectively advanced the state of atmospheric science by coupling airborne in situ measurements, remote sensing, and new platform
75 innovations to more accurately profile aerosols with unprecedented vertical resolution, filling the gaps of ground-based data. Aircraft can access high altitudes and carry advanced sensors, making them ideal for extensive regional measurements and satellite validation, though they are costly and often limited to short-term campaigns. UAS are portable, cost-efficient, and well-suited for fine-scale aerosol profiling at low-to-mid altitudes, but their operational range, payload capacity, and sensitivity to weather conditions can restrict their applications. TBS can provide detailed profiles of lower atmospheric layers at specific
altitudes and are affordable for deployment in challenging environments, but are spatially constrained to their tethered location and limited to low altitudes due to operational constraints. Together, these airborne platforms and remote sensing techniques provide a synergistic approach to understanding aerosol distribution, composition, transport processes, and climatic effects across varying spatial and temporal scales.

Given its scientific importance, integrating multi-platform observations collected at the same location and time—such as
80 those from the Tracking Aerosol Convection interactions ExpeRiment (TRACER)—is highly desirable. TRACER was a U.S. DOE ARM field campaign in the Houston, Texas, coastal urban region during 2021–2022, with an intensive observing period in summer 2022 (Wang et al., 2025; Jensen et al., 2025; Chen et al., 2025; Rapp et al., 2024; Farley et al., 2024). It deployed the ARM Mobile Facility and the ARM TBS, both equipped with a comprehensive suite of in situ and remote sensors, to characterize aerosol properties, boundary-layer structure, and convective cloud evolution under the influence of the sea breeze
85 and urban pollution. The campaign’s main objective was to quantify how aerosol loading and composition modulate convective cloud initiation, microphysics, and precipitation, thereby providing process-level constraints for weather and Earth’s system models; TRACER was also coordinated with partner efforts to enable source attribution and multi-scale evaluation of aerosol–cloud–precipitation interactions (Jensen et al., 2025; Rapp et al., 2024).

One of the primary objectives of this study is to derive high-resolution vertical aerosol profiles from TBS measurements,
90 targeting the lower troposphere and the atmospheric layers up to the cloud base, where remote-sensing retrievals are most uncertain. Unlike aircraft, which can sample these altitudes but typically only intermittently and for limited durations, TBS platforms can provide sustained, high-frequency profiling at fixed locations, enabling more continuous characterization of aerosol structure in these critical regions. From these profiles, we will quantify vertical variability in aerosol number, size distribution, composition, CCN activity, and optical properties across altitudes and meteorological regimes. Using vertical
95 profiles of aerosol properties, researchers can better constrain boundary-layer structure and entrainment by directly leveraging observed vertical variability. This study can potentially lead to further validation and refinement of ground-based and satellite remote sensing retrievals, improve data assimilation and model process parameterizations, and provide actionable constraints for both observations and models.

2. Data & Methodology

2.1. Field Campaigns Overview

Conducted from June to September 2022 in the Houston, Texas region, the TRACER Intensive Observation Period (IOP) focused on investigating the complex interactions between aerosols and convective clouds in an environment with varied aerosol loading and meteorological conditions (Jensen et al., 2025). During the IOP months, the ARM TBS was deployed the first two weeks of each month at an ancillary (ANC) site in Guy, Texas (29.33°N, 95.74°W), southwest of Houston. Concurrently with TBS flights, a co-located microwave radiometer provided continuous retrievals of precipitable water vapor (PWV) and liquid water path (LWP) profiles. This location was selected to represent a rural background with less influence from a mix of aerosol sources (such as fresh aerosol emissions from the metropolitan Houston area).




Figure 1. Locations of Vertical Aerosol Observations during the TRACER IOP. Sources: Esri, TomTom, Garmin, SafeGraph, FAO, MET/NASA, USGS, EPA, NPS, USFWS | Powered by Esri



Complementing the stationary measurements conducted by the ARM TBS team, the collaborating Texas A&M University (TAMU) team undertook mobile sampling of the sea breeze. Their sampling transects, illustrated in Figure 1, typically originated on the coast in Galveston. From there, they followed the sea-breeze inland, concluding at either the AMF site in La Porte or at one of several sites west and northwest of Houston (Bear Creek, Hockley, or Hempstead) (Thompson et al, 2025a). Along this mapped corridor, TAMU mobile measurements frequently transitioned from marine boundary-layer air characterized by low particle concentration to continental air ahead of the sea-breeze front, which generally contained particles that were slightly more effective for cloud formation and exhibited substantial variability over tens of kilometers (Thompson et al., 2025a; Thompson et al., 2025b; Chen et al., 2025).

Additionally, the Next Generation Weather Radar (NEXRAD), designated KHGX and located at 29.47°N, 95.08°W, played a pivotal role in observational activities throughout the study period (Galfione et al., 2025). This advanced radar system conducted volume scans using Plan Position Indicator (PPI) techniques, systematically operating over elevation angles from 0.5° to 19.5°. By varying these elevation angles, the radar prioritized acquiring data essential for detecting and characterizing convective processes embedded within stratiform regions. This approach ensured comprehensive coverage and improved the accuracy of convective feature identification, which is critical for understanding aerosol–cloud interactions and precipitation patterns in the Houston area.

2.2. Vertical Observations

The TBS observations enable examination of vertical variations in aerosol properties, including changes in concentration and composition driven by sources such as urban pollution, industrial emissions, long-range transport, and marine influences. The ARM TBS (Dexheimer et al., 2026) uses winch-controlled, helium-filled aerostats to obtain in situ vertical profiles up to about 1.5 km above ground, operating below clouds. Typical payloads, as shown in Table 1, include sensors for pressure, temperature, and humidity (e.g., iMet-XQ2), an anemometer for wind speed and direction, and a portable optical particle spectrometer (POPS) and a condensation particle counter for aerosol size distributions and concentrations; a Size- and Time-resolved Aerosol Collector (STAC) collected the aerosol particles at various altitudes for offline chemical analysis. During the TRACER IOP, a subset of meteorological and aerosol measurements was collected from 149 ARM TBS flights, which were conducted in the first two weeks of each month from June to September. The monthly distribution included 46 flights on 11 days between 03 and 14 June (46.45 hours), 32 flights on 13 days between 02 and 14 July (41.12 hours), 43 flights on 13 days between 02 and 14 August (46.92 hours), and 28 flights on 12 days between 02 and 14 September (56.2 hours). Detailed TBS daily operation reports can be found in Appendix A of the campaign report (Jensen et al., 2023).

Table 1. TBS measurements during TRACER IOP

Instruments	Property Measured
-------------	-------------------



iMet RSB-1 and RSB-4 radiosondes	Pressure, temperature, relative humidity, 3D GPS
iMet XQ2 UAV Sensor	Pressure, temperature, relative humidity, 3D GPS
40C cup anemometers	1 Hz horizontal wind speed
VectorNav GNSS wind direction	1 Hz wind direction
Portable Optical Particle Spectrometer (POPS)	Aerosol size distribution from 135 to 3000 nm
Condensation Particle Counter (CPC) Model 3007	Total aerosol concentration from 10 nm to 1000 nm
Size- and Time-Resolved Aerosol Collector (STAC)	Size- and time-resolved chemical composition from 70 nm to 2300 nm

During the 2022 deployment, aerosol particles were collected using a 4-stage Sioutas impactor at ground level and a STAC (Cheng et al., 2022) on a TBS. The instrument collected size- and time-resolved samples directly onto substrates for two complementary analytical pathways at the Environmental Molecular Sciences Laboratory (EMSL, <https://www.emsl.pnnl.gov/>). For single-particle characterization, particles collected at stage D (50% cutoff size of 0.12 microns) of the STAC sampler were analyzed using Computer-Controlled Scanning Electron Microscopy with Energy-Dispersive X-ray spectroscopy (CCSEM-EDX). This technique yields individual particle size, morphology, mixing state, and elemental composition to help identify particle types and sources (Lata et al., 2023). For molecular-level characterization, samples were collected on Teflon filters for about 12 hours at ground level and analyzed by nanospray desorption electrospray ionization high-resolution mass spectrometry (nano-DESI-HRMS) (Gautam et al., 2024). The resulting mass spectra inform the constituent organic molecular formulas in the particle phase, from which we derive key parameters such as O:C ratios, carbon oxidation state, aromaticity indices, and organic aerosol volatility (Roach et al., 2010; Vandergrift et al., 2024; Vandergrift et al., 2022). The results discussed herein are from samples collected during the June and September 2022 flights.

A range of ground-based measurements was obtained from the ARM Mobile Facility (AMF) site at La Porte, TX. The AMF was equipped with advanced radar and lidar systems, aerosol-observing instruments, and meteorological sensors. ARM has developed a set of Planetary Boundary Layer Height (PBLHT) value-added products to help scientists evaluate and compare estimates. In this study, the Planetary Boundary Layer Height Radiosonde Retrievals with yearly output (PBLHTSONDEYR1MCFARL) VAP was used to provide the PBLHT (Sivaraman et al., 2013; Li et al., 2021; Liu and Liang, 2010).

Additionally, this study used aerosol vertical profiles retrieved from a micropulse lidar (MPL) using a newly developed technique by the Brooks group at Texas A&M University (TAMU) that combines ground-based aerosol measurements with lidar and radiosonde data to obtain vertical profiles of aerosol, CCN, and INP. (Chen et al., 2025). This technique assumes that surface measurements of aerosol size distribution, composition, and cloud-activating ability are representative of their vertical profiles, except in cases of elevated layers like smoke or dust. The approach involves determining the vertical profile of the



cloud-free aerosol backscatter coefficient using lidar and radiosonde data. The aerosol backscatter coefficient is then corrected for hygroscopic growth using the lidar hygroscopic growth correction factor $f(RH)$, which is derived from ground-based Scanning Mobility Particle Sizer (SMPS), POPS, and CCN counter measurements, along with relative humidity profiles from radiosondes. This process yields the dry aerosol backscatter coefficient profile, which serves as the basis for further analysis of aerosol, CCN, and INP distributions.

2.3. CCN concentration estimation

To estimate CCN concentration, our method combines the observed TBS aerosol size distribution with ground-based chemical composition measurements. Our approach includes two steps. The first step is to address the size-range limitation of the aerosol size-distribution data. We extrapolate the measured particle number size distribution from its original POPS range of 135–3000 nm to 10 nm. We employed a modal-fitting technique based on the algorithm detailed by Hussein et al. (Seinfeld and Pandis, 2016; Hussein et al., 2005). The complete size distribution was represented as the sum of up to two lognormal modes, each defined by its total number concentration (N_i), geometric mean diameter (D_{gi}), and geometric standard deviation (σ_{gi}). An iterative optimization was performed to determine the parameters for these modes by minimizing the sum of squared errors between the estimated and measured data points in the 135–1000 nm range. Crucially, this fitting process was constrained so that the integral of the bimodal distribution over the entire 10–1000 nm range matched the independently measured total number concentration measured by the CPC.

The second step is to determine the CCN concentration profile using the κ -Köhler theory. In this study, we assume that the aerosol chemical composition at the ground was representative of that at each TBS flight altitude when the same air mass influenced the TBS flight. Ground-level non-refractory PM_{10} was measured with a Time-of-Flight Aerosol Chemical Speciation Monitor (ToF-ACSM, Aerodyne Research) equipped with a standard vaporizer (Li et al., 2025; Fröhlich et al., 2013), using a $PM_{2.5}$ cyclone upstream of the inlet. The instrument provided 10-min-resolution mass concentrations of organics (Org), sulfate (SO_4), nitrate (NO_3), ammonium (NH_4), and chloride (Chl) to complement the TBS deployment. However, the ACSM only quantifies non-refractory submicron species and cannot detect refractory components such as black carbon, sea salt, and mineral dust. These missing species, together with the assumption of a vertically uniform chemical composition, can influence the bulk hygroscopicity and introduce additional uncertainty into CCN concentration estimates.

We averaged the volumetric fractions of aerosol chemical composition over the flight period. Aerosol particles were assumed as internal mixtures, and the particle hygroscopicity (κ_{CCN}) is therefore the volumetric average of the three participating species. More information can be found in our previous publication (Mei et al., 2024)

$$\kappa_{CCN} = \sum_i x_i \kappa_i = x_{org} \kappa_{org} + x_{(NH_4)_2SO_4} \kappa_{(NH_4)_2SO_4} + x_{NH_4NO_3} \kappa_{NH_4NO_3} \quad (1)$$

where x_i is the volume fraction of species i . The values of κ are 0.15, 0.67 and 0.61 for organic, $(NH_4)_2SO_4$ and NH_4NO_3 , respectively.

For a given supersaturation (SS), the κ -Köhler theory is used to determine a corresponding critical activation diameter D_{pc} ,



$$D_{pc}^3 = \frac{4A^3}{27\kappa_{CCN}(SS)^2} \quad (2)$$

where $A = \frac{4\sigma_w M_w}{RT\rho_w D}$, M_w the molecular weight of water, σ_w the surface tension of pure water, ρ_w the density of water, R the gas constant, and T the absolute temperature.

All particles in the measured size distribution larger than this critical diameter are assumed to activate into cloud droplets. The total CCN concentration is then found by integrating the particle size distribution from this critical diameter upwards.

3. Results & Discussion

3.1. Air Mass Trajectory and Clusters Analysis

The ARMTRAJ dataset (Silber et al., 2025b) is generated using a Lagrangian methodology, primarily employing the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT) model with the fifth-generation European Center for Medium-Range Weather Forecasts (ECMWF) atmospheric reanalysis. (Hersbach et al., 2020; Stein et al., 2015) This model is driven by meteorological inputs from the ERA5 reanalysis dataset at its highest spatial resolution (0.25 degrees) and 1-h temporal resolutions. The core value of ARMTRAJ is to bridge the gap between the fixed, or *Eulerian*, nature of ARM's ground-based measurements and the need to understand the air mass's history as it passes over the site. To quantify uncertainty, the dataset incorporates results from ensemble runs for each trajectory initialization; the variability within these ensembles serves as a valuable metric for assessing the consistency and robustness of the reported air-mass coordinates and state variables. (Silber et al., 2025b; Silber et al., 2025a). For the ARMTRAJ-TBS, the trajectories were estimated at 11-equally-distant altitudes between the lowest and highest altitudes probed by the balloon over 1-hour intervals (Silber et al., 2025a).

Using air-mass geospatial trajectory data derived from latitude and longitude coordinates, Figure 2 illustrates clustered atmospheric transport pathways over two time periods: 24 hours (Figure 2a) and 5 days (Figure 2b). K-means clustering was applied to classify trajectories into three groups, highlighting contrasting transport pathways and source regions. For the 24-hour back trajectories, Cluster 1 (blue) comprises air masses that travel almost exclusively over the Gulf of Mexico before reaching the ANC site, indicating a relatively direct marine inflow. In contrast, Cluster 2 (orange) follows more complex marine pathways: the trajectories initially traverse the Gulf, then curve inland and pass over land for part of their transport, indicating a mixed marine–continental influence and a greater potential for modification by boundary-layer processes over land. Cluster 3 (yellow) originates primarily from the north and west, reflecting stronger influence from continental sources, including anthropogenic emissions from the broader Houston metropolitan region and upwind areas. The 5-day analysis (Figure 2b) provides crucial insight into the long-range history of these air masses. It confirms that Clusters 1 and 2 spent the vast majority of their transit time over the ocean, with pathways extending deep into the Gulf of Mexico and the western Atlantic. Conversely, the 5-day trajectories for Cluster 3 reveal a significant, long-range continental influence, with air masses originating from the central and upper Midwest of the United States and Canada. This demonstrates that while Clusters 1 and



2 are primarily influenced by oceanic processes, Cluster 3 is shaped by prolonged transport over land, bringing distinctly continental air to the site.




Figure 2. ARMTRAJ-TBS back trajectory properties 2022 (a) for 24 hours, and (b) for five days. The illustrated trajectory clusters are characterized by the geophysical range during measurement periods in June, July, August, and September. The star is the ancillary site in Guy, TX. All figures are generated using Natural Earth by MATLAB®.

To identify the dominant patterns of meteorological variability within three clustered atmospheric regimes, we performed Principal Component Analysis (PCA) on the ERA5 reanalysis dataset. Before analysis, each variable was standardized to ensure that contributions were not biased by scale, thereby allowing the PCA to reveal more meaningful underlying structures. The first two principal components (PC1 and PC2) accounted for more than 80% of the total variance in each regime. The weights for these components reveal distinct physical processes. As shown in Figure 3, the component weights illustrate the influence of eight meteorological variables on PC1 and PC2. The color indicates whether a variable's contribution is positive (yellow) or negative (red). The variables analyzed include: air mass pressure (pres), potential temperature (theta), temperature (temp), planetary boundary layer height (pblh), relative humidity (rh), specific humidity (qv), equivalent potential temperature (theta_e), and the mean hourly air mass ascent rate (wvert).

In the marine-influenced regime (Cluster 1), the PCA cleanly separates the advective characteristics of the marine boundary layer from local diurnal forcing. The leading component (PC1) is strongly associated with planetary boundary layer height (pblh, -0.45), potential temperature (theta, -0.34), relative humidity (rh, 0.48), and specific humidity (qv, 0.46). We therefore interpret PC1 as an index of marine layer strength and intrusion. Positive PC1 scores correspond to a cool, moist, and shallow marine boundary layer characteristic of direct onshore flow, whereas negative scores indicate a deeper, warmer, and drier boundary layer associated with marine boundary layer erosion or increased continental influence. The second



250 component (PC2) is dominated by positive weights for pblh (0.63), temperature (temp, 0.46), and equivalent potential temperature (theta_e, 0.41). Because PC2 is orthogonal to the marine-layer signal, it is interpreted as representing the diurnal cycle of surface heating and boundary-layer growth.

For the regime influenced by anthropogenic emissions and transport (Cluster 3), the PCA identifies atmospheric dispersion capacity as the primary mode of variability. The leading component (PC1) is dominated by pblh (0.71) and is
255 strongly correlated with rh (-0.52). In this regime, PC1 functions as a ventilation index, quantifying the atmosphere’s capacity to dilute surface emissions. High positive scores correspond to a deep, well-mixed boundary layer and reduced near-surface humidity, consistent with vigorous vertical mixing. Negative scores indicate a shallow, stable boundary layer that traps pollutants and moisture near the surface, favoring episodes of poor air quality. The second component (PC2) is characterized by positive weights for surface pressure (pres, 0.57), temperature (temp, 0.47), and specific humidity (qv, 0.47), and likely
260 correspond to conditions within a warm, moist high-pressure system.

Under mixed-influence conditions (Cluster 2), the leading mode of variability (PC1) is overwhelmingly characterized by strong, co-varying positive weights for specific humidity (qv, 0.58) and equivalent potential temperature (theta_e, 0.58). This component serves as a robust proxy for the large-scale advection of moisture and thermodynamic energy into the region. The second component is almost exclusively defined by a large weight for pblh (0.84) and a strong correlation with rh (-0.42).
265 This signature suggests vertical mixing and entrainment, in which a growing boundary layer mixes with drier air from the free troposphere, thereby increasing pblh while reducing the layer-averaged rh.

In summary, the two-component PCA, stratified by regime, isolates the dominant atmospheric controls: marine-layer confinement versus diurnal heating (Cluster 1), moisture/energy advection versus entrainment-driven drying (Cluster 2), and boundary-layer ventilation capacity versus synoptic airmass forcing (Cluster 3).

270






Figure 3. Heatmaps of component weights from the Principal Component Analysis (PCA) for the three distinct atmospheric regimes: (a) Cluster 1 (Marine/Coastal Influence), (b) Cluster 2 (Regional/Marine Mixed Influence), and (c) Cluster 3 (Anthropogenic/transport Influence).

3.2. Vertical Profiles of Aerosol Properties

Air mass transport pathways, origins, and processes profoundly affect aerosol distribution, particularly vertical variability. To understand how different atmospheric regimes influence the vertical aerosol structure, we grouped the flight data into three distinct clusters, as discussed in section 3.1. Cluster 1 affected 28 days, Cluster 2 influenced 12 days, and Cluster 3 impacted 9 days. Figure 4 shows the mean vertical profiles of temperature (T), relative humidity (RH), and aerosol number concentrations ($N_{\text{pops}} > 135 \text{ nm}$ and $N_{\text{cpc}} > 10 \text{ nm}$) composited by three trajectory-based influence clusters: Cluster 1 (marine-influenced), Cluster 3 (anthropogenic/transport influenced), and Cluster 2 (mixed influence). The aerosol number concentrations were reported at the standard temperature and pressure (STP; 1013.15 hPa and 298.15 K). In Figure 4, plots represent daytime TBS profiles during the intensive period in 2022, and error bars denote one standard deviation about the mean in each 75-m altitude bin for each cluster.

The vertical profiles of temperature, relative humidity, and aerosol concentrations reveal clear distinctions between the three air mass clusters. The temperature profiles are broadly similar across all three clusters, with comparable lapse rates from the surface to approximately 1.5 km. This suggests that the local-scale thermal structure and boundary-layer depth were not strongly dependent on air-mass origin during the sampled period. However, there are subtle differences near the surface: the anthropogenic cluster (Cluster 3) is slightly cooler ($\sim 30^\circ\text{C}$) than the marine (Cluster 1, $\sim 32^\circ\text{C}$) and mixed (Cluster 2, $\sim 31^\circ\text{C}$) clusters. The relative humidity (RH) profiles show similar variation. Cluster 3 is slightly more humid throughout the lower atmosphere, with RH values sustained at $\sim 60\%$ below 1 km, consistent with a moist, coastal air mass. Notably, the RH in Cluster 2 shows a weak positive gradient: starting from around 60–65% near the surface, RH gradually rises with height, reaching its highest values ($\approx 70\text{--}75\%$) near 1.3–1.4 km. In Cluster 3, RH aloft behaves differently. It is comparable to the other two cluster-influenced conditions within the lowest 1 km, then declines above $\sim 1.2 \text{ km}$. This implies that the anthropogenic cluster is characterized by a moist, well-mixed lower boundary layer capped by relatively drier air aloft, which is typical of a polluted, coastal boundary layer capped by subsiding or advected drier free-tropospheric air.

The most dramatic differences between the clusters are evident in the aerosol concentrations. Cluster 1 (marine) exhibits low aerosol concentrations characteristic of clean background air. Both $N_{\text{cpc}} (> 10 \text{ nm})$ and $N_{\text{pops}} (> 135 \text{ nm})$ profiles show a uniform concentration up to $\sim 1.1 \text{ km}$. This vertical structure is consistent with a well-mixed marine boundary layer, in which the dominant aerosol sources and precursors originate from the ocean surface (e.g., sea spray and DMS). Furthermore, the gradual decrease with height at altitudes above 1000 m is likely due to vertical mixing and/or dilution from entrainment of even cleaner free-tropospheric air from above. Cluster 3 shows a similar uniform concentration in N_{cpc} and a gradual decrease in the aerosol concentration of N_{pops} as altitude increases from 100 m to 1.5 km. This indicates that, in both the clean marine (Cluster 1) and polluted anthropogenic (Cluster 3) cases, the dominant aerosol sources are located within the lower boundary



layer. However, Cluster 3 (anthropogenic), influenced by human activities such as industrial emissions, is also characterized by higher concentrations of larger particles (N_{pops}), while the total aerosol number concentrations (N_{cpc}) are comparable to those in Cluster 1. This combination suggests an aged, processed aerosol source.

Cluster 2 (mixed) presents a complex vertical structure. At lower altitudes, the aerosol concentration profiles are relatively constant below 800 m. However, above 1 km, the N_{cpc} and N_{pops} profiles show a notable increase, making it the most concentrated cluster aloft. This elevated enhancement is usually consistent with an advected, or long-range transported aerosol layer residing near or just above the boundary-layer top. Such layers often arise from regional or continental pollution, biomass burning, or other distant sources and can be decoupled from the local surface emission regime. The fact that Cluster 2 is intermediate in concentration near the surface but enhanced aloft supports the interpretation of a mixed air mass influenced simultaneously by local fresh emissions, new particle formations and an overlying transported pollution layer entrained into the boundary layer.




Figure 4. Atmospheric conditions observed during various cluster-influenced flights in the 2022 TRACER campaign. The subplots include: (a) ambient temperature; (b) ambient relative humidity; (c) total particle concentration measured by the condensation particle counter (CPC, particles greater than 10 nm); and (d) total particle concentration measured by the portable optical particle spectrometer (POPS, particles ranging from 135 to 3,000 nm). The square symbols represent the averaged mean values of the sampling periods, while the error bars denote the standard deviation of the measurements.



3.3. Summertime Planetary Boundary Characterization

To contextualize the summertime boundary-layer behavior, we characterized its diurnal evolution using radiosonde profiles launched from the AMF site and two established algorithms implemented in the ARM PBLHT VAP product (Sivaraman et al., 2013). Estimating the Planetary Boundary Layer Height (PBLHT) from sounding data is a classic approach to understanding the PBL structure. However, no single perfect method exists, and the optimal approach varies with the definition; therefore, a given approach may be a better fit depending on the goal. Data from the ARM program enable comparison of several radiosonde-based approaches, including the Liu-Liang and bulk Richardson number methods (Li et al., 2021; Stull, 2012; Seidel et al., 2010; Liu and Liang, 2010), which are particularly effective for studying the diurnal cycle. The bulk Richardson number method, which identifies the PBL top as the point at which turbulence is dynamically suppressed, is physically robust and well-suited to determining the height of the nocturnal Stable Boundary Layer (SBL) and the transition periods. In contrast, the Liu-Liang method defines the PBLHT as the height at which the maximum vertical gradient in thermodynamic variables occurs, a feature characteristic of the entrainment zone atop a convective boundary layer.




Figure 5. Diurnal variation in UTC time of mean planetary boundary layer (PBL) height for June (1–14), July (2–14), August (2–14), and September (2–14) 2022. Dots indicate the mean PBL height at each sampling time; error bars show the standard deviation of PBL height during each sampling period.



As shown in Figure 5, the mean diurnal evolution of the PBLHT during the summer and early autumn months (June-September) aligns with theoretical expectations – increasing over the course of daylight hours coupled with the diurnal cycle of surface temperature. The PBLHT determined by the Heffter method is systematically higher than that obtained by other methods. Because it identifies the PBLHT by locating the base of the elevated temperature inversion and using empirical thresholds that require tuning across regions, it might not be robust in this case for moist/marine regimes. A comparison between the Liu-Liang and Richardson number methods (with a critical value of 0.25) reveals strong agreement during daytime hours (05:30 (00:30), 11:30 (6:30), 17:30 (12:30), and 23:30 (18:30) UTC (local time). The convergent estimates during this period indicate that both methods effectively capture the same fundamental process: the growth of a convective mixed layer driven by surface heating. The Richardson method illustrates a pronounced collapse of the PBLHT, while the Liu-Liang method shows a much more gradual decrease. This divergence is physically instructive. The Richardson method, being a direct measure of turbulence stability, captures the rapid decay of buoyancy-driven mixing as solar heating ceases, signaling the swift transition to a shallower, shear-driven stable layer. The Liu-Liang method, by seeking the maximum gradient, is prone to identifying the top of the now-inactive, neutrally stratified residual layer left over from the afternoon, thus overestimating the height of the active nocturnal boundary layer.

Although the sounding data from the middle of the night were limited to fully characterize the developed SBL, the 05:30 data point is particularly instructive for inferring the preceding nocturnal conditions. The trend of a progressively shallower early-morning PBL from summer to autumn is evident. The mean PBLHT at 05:30 decreased from approximately 500–600 m in June and July to 300–400 m in August, and further to around 200–300 m in September. This implies that the nocturnal SBL becomes systematically shallower and more stable as the season progresses from summer to autumn, a trend consistent with longer nights and more extensive radiative cooling, which fosters the development of stronger, more vertically constrained surface inversions.

To assess how well the TBS profiled the summertime boundary layer, we overlaid flight altitudes with the mean PBLHT derived from the Liu–Liang algorithm. The overlay (Figure S1) shows that most daytime flights in June–August remained below the mean Liu–Liang PBLHT, whereas several September flights reached into and, at times, above the mean PBL top, as shown in Figure 6. This is primarily explained by the systematic seasonal decrease in PBL depth from mid-summer to early autumn. Operational factors also contributed: earlier in the summer, the requirement to operate below the cloud base often constrained the TBS to lower altitudes, whereas in September, clearer conditions allowed the balloon to ascend to its maximum altitude. As established in the previous analysis, the mean afternoon PBLHT is significantly deeper in June and July (often exceeding 1.5–2.0 km) compared to September, when it typically peaks around 1.0–1.3 km. This is a direct consequence of reduced solar insolation and shorter day lengths, which lessen the surface heating that drives convective turbulence and PBL growth. With a fixed or operationally limited TBS ceiling, flights were therefore sampled primarily within the well-mixed layer in JJA. Within this regime, temperature, humidity, and aerosol properties are vertically homogeneous to first order, and profiles are representative of surface conditions and daytime dilution. The data collected in JJA are thus well suited to characterizing mixed-layer chemistry and aerosol processing and to relating CCN to surface emissions.




Figure 6. Daily TBS flight altitudes during September 2022, overlaid with the mean planetary boundary layer height (solid blue line) and its variability (shaded region) derived from radiosonde profiles.

To link air-mass origin with boundary-layer dynamics, we combined back-trajectory cluster analysis with radiosonde-derived PBL heights (Liu–Liang and Ri=0.25) and TBS flight altitudes to diagnose how source regions are introduced into, and mixed within, the summertime boundary layer. During most JJA sampling periods, the air mass is consistently dominated by Cluster 1 below 500 m, while above 500 m it is dominated by Cluster 2. This indicates a layered structure—local/near-field air occupying the nocturnal/morning boundary layer and a distinct, advected layer aloft. As the convective boundary layer grows during the day (often exceeding 1–1.5 km in JJA), entrainment at the PBL top brings the Cluster-2 air downward, gradually diluting the near-surface signature of Cluster 1. Because most TBS flights in JJA remained below the mean PBLHT, they primarily sampled the well-mixed local layer. In September, the air mass above 500 m becomes more heterogeneous, with both Cluster 2 and Cluster 3 appearing aloft. The decline in PBLHT (shallower afternoon maxima and lower morning depths) lowers the entrainment zone. September flights often reached and exceeded the mean PBLHT, enabling direct sampling of the entrainment zone and the lower free troposphere, where the influences of Clusters 2 and 3 are strongest. With a shallower morning SBL (~200–300 m) and reduced daytime growth in September, the boundary layer is more vulnerable to changes in synoptic and mesoscale flow (e.g., shifting trajectories, sea-breeze passages, residual-layer interactions). Consequently, all three clusters can influence the air below 500 m, with the dominant cluster varying from day to day. This variability might be



390 caused by intermittent entrainment of the overlying Cluster 2 layers into a shallow mixed layer and the occasional persistence of the previous day's residual layer aloft.

3.4. Clusters-influenced CCN Profiles Estimation

Figure 7 presents vertical profiles of estimated CCN concentrations for three distinct air mass clusters, plotted as a function of altitude. These plots provide useful vertical context for the variation in CCN availability across airmass regimes. Another version of this figure was plotted against altitude normalized by the PBLHT (Altitude/PBLHT), as shown in Figure S2. This normalized version aims to reveal key differences in the vertical distribution of aerosols relative to the PBL structure across differing airmass influences. In Figures (a)-(c), the size of the circle symbols at each altitude level is proportional to the number of data points collected within that altitude range, providing a visual indication of the statistical robustness of measurements. A threshold of 2,000 data points is established as the confidence limit for data representativeness. Altitude bins containing fewer than 2,000 data points (represented by smaller circles) should be interpreted with caution, as they may be less representative of the true vertical CCN distribution. Figures 7(d) and S2(d) appropriately document the sampling limitations inherent in TBS operations. They show a strong reduction in the number of samples collected aloft, and the data density decreases substantially above ~800-1000 m altitude in Figure 7(d). In Figure S2 (d), Cluster 2 shows notably fewer samples at the high normalized ratio (Altitude/PBLHT>0.8) compared to Clusters 1 and 3. Because statistical confidence in the upper-level observations is correspondingly reduced, the figures are most suitable for discussing within-PBL gradients and airmass differences but provide less definitive information about the free-tropospheric tail.

The vertical structure in all three cluster cases generally shows a well-mixed boundary layer (<800 m), characterized by relatively uniform CCN concentrations. Cluster 1 (Marine-influenced) is characterized by the lowest overall CCN concentrations, consistent with a relatively clean marine air mass originating from the Gulf of Mexico. Within the lower PBL (Altitude/PBLHT $\leq$ 0.4), mean CCN concentrations at SS = 0.80% are on the order of 1,000 cm⁻³. The ratio of CCN at 0.80% to 0.13% supersaturation is around 3 in this region, indicating relatively homogeneous critical supersaturation among aerosol particles, likely dominated by sea salt and marine biogenic particles with similar hygroscopic properties. The low ratio also suggests fewer small Aitken-mode particles relative to the accumulation mode, which is typical of clean marine environments where new particle formation is suppressed or where coagulation/wet scavenging removes smaller particles.

Cluster 3 (Urban/Anthropogenic/Transport) exhibits the highest CCN concentrations and maintains elevated values through much of the sampled depth. Within the PBLHT, CCN concentrations at SS = 0.80% reach ~3,000 cm⁻³. The ratio (N_{0.80%}/N_{0.13%}) is about 5, suggesting a broader, but still relatively processed, size distribution. This is consistent with aged urban/industrial pollution, where the transport time from Houston likely allows for photochemical aging and coagulation, and secondary aerosol formation (sulfate, nitrate, SOA) has grown particles into the accumulation mode, but a fraction of smaller, less hygroscopic particles remains (Li et al., 2025; Farley et al., 2024). This interpretation is further supported by the characteristics of aerosol chemical composition. The presence of an "aged" or "processed" aerosol population was observed by chemical analysis of a June 3 sample representative of Cluster 3's influence. This sample was exceptionally complex, with



1020 molecular formulas (MFs) assigned (Figure S3). A large fraction of these were nitrogen- and sulfur-containing organics, including 340 organonitrates (CHON) and 104 organosulfates (CHOS). The prevalence of these compounds resulted in a distinct chemical signature: high oxygenation and low aromaticity, both hallmarks of extensive atmospheric processing. A detailed manuscript (under review) presents chemical analyses of organic aerosols collected during TBS flights on June 3, 5, 8, and 9, 2022, and reports significant day-to-day variability attributable to distinct air-mass influences.

Cluster 2 (Mixed Influence from marine and regional) represents air masses influenced by both marine and regional continental sources and sits between Clusters 1 and 3 in absolute CCN concentration but shows the highest spread ratio ($N_{0.80\%}/N_{0.13\%} \sim 6$).

Cluster 2 exhibits a distinct vertical structure compared with Clusters 1 and 3. In addition to being well-mixed in the lower 70% of PBL, CCN concentrations influenced by Cluster 2 increase significantly with altitude at the top of the PBLHT (above 1000m), culminating in a distinct peak near the PBL top (Altitude/PBLHT > 0.8) with values exceeding $5,000 \text{ cm}^{-3}$ at SS=0.80%. This accumulation of aerosols aloft is strong evidence of active in situ particle growth occurring preferentially at the top of the boundary layer. One possible explanation is that coastal recirculation enhanced new particle formation (NPF) by mixing marine DMS-derived sulfur with continental NO_x and organics, producing fresh Aitken-mode particles. The recirculation process at the PBL top may also bring in intermittent continental influence with fresher, less-processed particles that have not yet grown into the accumulation mode. These continental pollutants are trapped beneath a temperature inversion and interact with the high humidity and strong solar radiation characteristic of the coastal marine environment. These conditions are ideal for efficient aqueous-phase chemistry (cloud processing) and photochemical reactions, which rapidly convert precursor gases into particulate matter and add mass to existing small particles, thereby growing them into the CCN size range (Li et al., 2025; Farley et al., 2024). However, Figure 7(d) shows that the number of samples decreases markedly with altitude, so the upper-level enhancement in Cluster 2 should be interpreted with caution, and additional observations would be needed to confirm this as a systematic feature.




Figure 7. Estimated CCN concentration vertical profiles as a function of TBS altitude for each cluster (a - c), with the number of data points collected at the corresponding altitude for three clusters (d).

We noticed elevated CCN layers in Figure S2(c) ($1.5 < \text{Altitude/PBLHT} < 2.2$), which is due to the single-day event on September 9, might not represent the general trend in this airmass regime.

3.5. Why Observational Aerosol-Cloud Interaction Studies Are So Difficult – A Case Study

A central obstacle in observational aerosol–cloud interaction research is disentangling aerosol microphysical effects from the meteorological variability that simultaneously governs both aerosol distributions and cloud formation. The September 6–7, 2022, case study from the TRACER campaign near Houston provides a compelling, data-rich illustration of this challenge—demonstrating how mesoscale circulations, airmass advection, and differential transport can simultaneously transform the thermodynamic environment and the aerosol population, rendering causal attribution of cloud changes to aerosol effects deeply ambiguous from observations alone.

3.5.1. Aerosol Vertical Structure and Airmass Complexity: September 6

In most boundary-layer flights during the campaign, the TBS vertical profiles reflected a single, well-mixed air mass. However, several cases in September deviated significantly from this pattern. The case from September 6, 2022, presented in



Figure 8, is a prominent example of such a deviation. The vertical profile of estimated CCN concentration (Figure 8a) reveals a distinct vertical stratification: below approximately 1000 m, CCN concentrations are largely constant with altitude across all measured supersaturations, while above 1000 m, CCN concentrations increase with altitude. These trends strongly suggest the presence of two vertically decoupled air masses with different aerosol characteristics, likely separated at the top of the marine boundary layer.

An analysis of the 24-hour ARMTRAJ back-trajectory provides a compelling explanation for this observed stratification. Figure 8(b) shows the trajectories for air parcels arriving at altitudes higher than 1000 m. This airmass originated over the open Gulf of Mexico and partially underwent subsidence (as indicated by increasing pressure along the trajectory paths) as it advected towards the measurement site. This air mass history is characteristic of marine air (Cluster 1) from the free troposphere, consistent with the lower CCN concentrations ($<500 \text{ cm}^{-3}$) observed in Figure 8 (a). Conversely, the trajectories for air parcels arriving near the ground (Figure 8(c)) reveal a drastically different history. These air masses exhibit a complex, low-level recirculation pattern confined near the coast (Cluster 2). This low-level recirculation pattern is consistent with NEXRAD imagery (Fig. S8a), which shows an airmass boundary propagating from the Gulf Coast, reaching the site at approximately 17 UTC. Therefore, the September 6 case demonstrates a scenario of different airmass types inside and above the boundary layer. The TBS profile sampled two vertically stacked but historically distinct air masses: a recirculated continental/coastal layer near the surface, and a cleaner, subsided marine layer entraining from the free troposphere above it.





Figure 8 Estimated CCN concentration vertical profiles (a) using TBS measurements for both flights on September 6 at the supersaturations of 0.13%, 0.35%, 0.47%, 0.63%, and 0.8% with ARMTRAJ TBS 24-hour back-trajectories for the TBS flight altitude higher than 1000 m (b) and near ground (c).

Critically, the corresponding radiosonde from September 6 (launched at 23:30 UTC, right after the TBS flight was ending, Figure S4) indicates a thermodynamic environment that is fundamentally hostile to cloud formation. The skew-T profile shows a deep layer of extremely dry mid-level air (~800–400 hPa). The large temperature–dewpoint separation indicates that any nascent cloud penetrating the cap would be rapidly eroded by entrainment of this dry air. Above the moist boundary layer (which extends only to roughly 950–900 mb), the dewpoint drops, with a corresponding dewpoint depression of 20–30°C or more through the 800–400 hPa layer. Although the dewpoint trace shows some oscillations in the 700–600 mb range (suggesting thin, intermittent layers of slightly higher moisture, possibly associated with the marine airmass above 1000 m identified by the back-trajectories. No clouds were observed on this day. The combination of CCN profiles, back trajectories and the thermodynamic sounding collectively indicates an atmosphere in which cloud formation was suppressed by dynamical and thermodynamic barriers—regardless of aerosol loading.

Additionally, the size-resolved chemical compositions derived from CCSEM-EDX analysis show considerable differences between the ground and aloft samples, as shown in Figure 9. The CCSEM-EDX results were consistent with the back-trajectory



analysis - two distinct aerosol environments linked to different air masses at different heights. The ground-level sample is characterized by a high fraction of Carbonaceous (35%) and Na-rich (18% + 20% Na-rich Sulfate) particles, indicating a strong influence of marine-influenced, boundary-layer air. The abundance of Na-rich sulfate suggests active chemical processing of sea salt in a polluted marine boundary layer. Meanwhile, the sample collected at 1350 m exhibits a markedly different chemical profile, indicative of a distinct airmass history. The most striking feature is the massive increase in the fraction of Dust (30%) compared to the ground (9%). This, combined with the still-significant Carbonaceous (31%) fraction and Na-rich particles (17%), suggests that the marine air masses also contained long-range-transported aerosol. The presence of internally mixed particles, such as Sulfate+Dust (3%) and Carbonaceous+Dust (8%), indicates atmospheric aging during transport. Overall, the ground-level composition reflects locally and marine-influenced boundary-layer air, whereas the aloft composition reflects a marine air mass with aged, long-range transported airmass rich in dust and processed carbonaceous material.





Figure 9 Size-resolved particle classifications from CCSEM/EDX at (a) ground (750 particles analyzed) and (b) 1350 m altitude (515 particles analyzed) for flight on September 6, 2022. The inset shows the percentage of each CCSEM/EDX-derived particle class.

3.5.2. Dramatic Airmass Transformation: September 7

By September 7, both the aerosol environment and the thermodynamic structure had undergone a wholesale transformation. The vertical CCN profile showed exceptionally high concentrations, with values exceeding $3,000 \text{ cm}^{-3}$ at a supersaturation (SS) of 0.8%, extending throughout the boundary layer, as shown in Figure 10. This airmass, a processed mixture of continental pollution and humid marine air, was characterized by an abundance of small, hygroscopic particles (the $\text{CCN}(0.80\%)/\text{CCN}(0.13\%)$ ratio was about 10). The x-axis scale of the CCN profile expanded from $0\text{--}1,500 \text{ cm}^{-3}$ on September 6 to $0\text{--}10,000 \text{ cm}^{-3}$ on September 7, reflecting a $5\text{--}10\times$ increase in CCN across all supersaturations and altitudes.



515 Figure 10. Estimated CCN concentration vertical profiles (a) using TBS measurements for all flights on September 7 at the supersaturations of 0.13%, 0.35%, 0.47%, 0.63%, and 0.8% with ARMTRAJ TBS 24-hour back-trajectories for the TBS flight altitude higher than 1000 m (b) and near ground (c).

The radiosonde from September 7 (20:30 UTC, Figure S5) reveals an equally dramatic thermodynamic transformation. The lower troposphere (surface through ~700 mb) was substantially moister on September 7 than on September 6, which
 520 reduced the destructive entrainment of dry air during the critical early growth phase of convective clouds. However, above ~600–700 mb, significant drying persists. The mid-to-upper troposphere was not deeply moistened. Both days had large CAPE. On September 6, no lifting mechanism existed to release this instability, so no clouds formed. On September 7, a complicated combination of a bay-breeze boundary propagating from Galveston Bay to the west and multiple gust fronts, associated with
 525 a mesoscale convective system to the northeast of Houston, propagating toward the south and southwest (see Fig. S8b), provided the mesoscale convergence and lift necessary to push parcels through the cap. Once initiated, clouds benefited from the moister lower troposphere (which reduced entrainment-driven erosion during early growth). The dramatically higher CCN concentrations were a byproduct of the same air-mass transformation that improved moisture and provided the outflow trigger.

In addition, we examined the CCN profiles derived from the TAMU MPL at Hockley (see Figure 1) during the overlapping operational periods (as shown in Table S1). While the dataset in Figure 8 highlighted the importance of vertical variance in
 530 CCN, the vertical profiles of CCN show a dramatic shift in air-mass properties between September 6 and September 7, 2022. As shown in Figure S6, the TBS and TAMU MPL-estimated CCN profiles clearly indicate a significant horizontal transport



event. On September 6, the region was characterized by relatively clean conditions. TBS and TAMU profiles at both 0.2% and 0.8% supersaturation are relatively similar in magnitude and vertical structure, suggesting that both platforms were largely sampling the same regional-continental background aerosol, with only modest spatial gradients. By 7 September, however, the TBS and TAMU profiles diverge sharply: CCN concentrations from the TBS increase by a factor of 2 (especially at 0.8% SS) and exhibit a much more vertically diversity, with an elevated layer up to about 1–1.4 km, whereas TAMU values remain closer to the magnitude on September 6. This contrast implies that mesoscale horizontal transport—likely associated with a change in wind direction or the passage of a plume—dominated CCN variability on 7 September, introducing strong spatial heterogeneity in aerosol properties over relatively short horizontal distances in the region.

3.5.3. The Covariability Trap

At first glance, the September 6–7 contrast might appear to present a natural experiment for aerosol-induced convective invigoration: a clean day with few clouds or precipitation, followed by a polluted day more clouds and precipitation. Indeed, when ingested by the observed updrafts, the September 7 aerosol population may have led to the nucleation of a very high number of small cloud droplets, potentially suppressing warm rain formation and potentially invigorating convection through enhanced latent heat release (as shown in Figures S7). However, the radar imagery from NEXRAD (Figures S8 and S9) reveals that the ANC site was strongly influenced by mesoscale dynamical forcing including density current boundary (most likely a sea breeze) on September 6th, while on September 7th, strong convection propagating from the north with multiple cold pools that also interact with a bay breeze frontal boundary.

The combined skew-T, CCN analysis and radar imagery reveals that this interpretation is untenable as a simple causal narrative for three reasons. First, the clean day (September 6th) was meteorologically incapable of producing deep clouds due to a general lack of moisture and low-level instability. Second, the polluted day (September 7th) was meteorologically primed for convection regardless of aerosol loading. With enormous CAPE, deep moisture, and a mesoscale lifting mechanism (convective cold pools and a bay breeze front), the September 7 environment would have produced deep convection even with September 6's modest CCN of a few hundred cm⁻³. Cloud droplets would have been fewer and larger, and warm rain would have developed more efficiently, but convection would still have occurred. Third, the aerosol and meteorological transformations were not independent. The same mesoscale process—sea-/bay-breeze circulations, convective cold pools, and air mass advection—that delivered deep moisture, removed CIN, and enhanced CAPE also imported an entirely different aerosol population with dramatically higher CCN concentrations. The aerosol and thermodynamic changes are confounded at their source.

4. Conclusions

This study used TBS observations in the greater Houston coastal urban region to investigate how vertical thermodynamic structure and transport modulate aerosol and inferred CCN distributions. TBS data were collected during the summer over the



greater Houston, Texas, region as part of the DOE ARM User Facility's TRACER field campaign. To interpret these measurements and understand the influence of air mass source regions, the study utilizes the ARMTRAJ dataset. This dataset consists of Lagrangian air-mass back trajectories generated with the HYSPLIT model, driven by high-resolution ERA5 meteorological reanalysis data. Its purpose is to provide historical context for air masses arriving at fixed (Eulerian) measurement sites, with trajectory uncertainty quantified through an ensemble methodology. A k-means clustering analysis of 24-hour and 5-day back trajectories demonstrates the dataset's utility in differentiating air mass histories. The analysis identified three distinct transport pathways: 1) direct marine inflow from the Gulf of Mexico, 2) mixed marine-continental transport, and 3) urban/anthropogenic flow originating from the central U.S. and Canada. This approach effectively classifies air masses by their source regions, providing an essential framework for interpreting the observed variations in aerosol properties.

By grouping flight data into three trajectory-based clusters (marine, anthropogenic, and mixed), distinct vertical profiles were identified. While temperature profiles were broadly similar across clusters, RH profiles revealed structural differences: the mixed cluster showed increasing RH with altitude, whereas the anthropogenic cluster featured a moist layer capped by drier air. The most dramatic differences were observed in the aerosol profiles. The marine cluster (Cluster 1) exhibited low aerosol concentrations that decreased with altitude, characteristic of a clean, well-mixed marine boundary layer with surface-based sources. Urban/anthropogenic cluster (cluster 3) showed much higher concentrations that also decreased with altitude, indicating strong surface-based pollution within a well-mixed layer. The mixed cluster (Cluster 2) displayed a complex vertical structure with moderate surface concentrations and a significant increase in aerosol levels above 1 km. This elevated layer is interpreted as advected, long-range-transported pollution residing above the local boundary layer, thereby confirming the cluster's mixed-influence nature.

To contextualize the measurements, this study characterized the summertime diurnal evolution of the PBL using radiosonde data. Then, an analysis comparing TBS flight altitudes to the derived PBL height (PBLHT) revealed that summer (June-August) flights primarily sampled within the deep, well-mixed boundary layer. In contrast, September flights often reached or exceeded the shallower PBL top, enabling direct measurement of the entrainment zone. By integrating this analysis with air-mass trajectory clusters, a dynamic picture emerges. Especially in September, the shallower PBL and more heterogeneous air masses aloft (Clusters 2 and 3) led to greater variability, with TBS flights capturing the direct influence of entrainment and distinct synoptic flows on the boundary-layer structure.

An analysis of estimated CCN vertical profiles reveals distinct aerosol characteristics across three air mass clusters. The marine-influenced cluster (Cluster 1) shows the lowest CCN concentrations within a well-mixed boundary layer, consistent with a clean marine air mass. The urban/anthropogenic cluster (Cluster 3) displays the highest CCN concentrations, also confined to a well-mixed PBL, reflecting strong surface-based pollution from fresh urban and industrial emissions. The CCN profiles in this air mass regime has the highest spread ratio ($N_{0.80\%}/N_{0.13\%}$), pointing to a broader size distribution with a large population of smaller particles. The mixed-influence cluster (Cluster 2) exhibits a unique vertical structure, with the highest



CCN concentrations peaking near the top of the PBL. This elevated aerosol layer is attributed to in-situ particle growth, in which continental pollutants trapped at the PBL inversion react with high humidity and solar radiation.

Finally, we focused on two interesting flight days. Analysis of vertically resolved CCN profiles from September 6th and 7th, 2022, revealed significant vertical and horizontal aerosol heterogeneity. On September 6th, TBS profiles captured two
600 vertically decoupled air masses: a polluted, recirculated coastal boundary layer and an overlying, cleaner, subsided marine free-tropospheric layer containing long-range transported dust. This case demonstrates the critical role of differential advection in stratifying aerosol chemical and physical properties. On September 7th, a mesoscale advection event introduced an air mass with anomalously high CCN concentrations ($>3,000 \text{ cm}^{-3}$ at 0.8% SS), creating conditions highly conducive to aerosol-induced convective invigoration. The key scientific advance highlighted by these events is the synergistic use of in-situ TBS profiling
605 with ground-based remote sensing. The TBS provided detailed, process-level measurements of the pre-convective aerosol microphysical properties, while subsequent radar and lidar observations captured the macroscopic cloud dynamical response—namely, the initiation of deep convection.

While this study provides valuable insights into the vertical structure of CCN under different air-mass regimes, several limitations frame the conclusions and motivate a clear path for future research. First, the TBS sampling provides high-
610 resolution vertical structure but limited horizontal representativeness, capturing a single-column view of a spatially heterogeneous aerosol field. The observational period was confined to the summer months, precluding assessment of seasonal variability in aerosol-cloud interactions that may arise from shifts in source contributions, photochemical processing rates, and boundary-layer dynamics. Secondly, CCN concentrations were inferred from size distributions rather than measured directly, introducing uncertainty associated with assumed hygroscopicity (κ) and the internal versus external mixing state. The
615 activation ratios ($N_{0.8\%}/N_{0.13\%}$) presented here are bulk metrics that cannot fully disentangle the relative contributions of size distribution shape versus composition-dependent hygroscopicity. This is particularly relevant for Cluster 2 (coastal recirculation), where the high activation ratio (~ 6) could reflect either a broad size distribution with abundant Aitken-mode particles, compositional heterogeneity from mixed marine-continental sources, or both. Thirdly, back-trajectory clustering provides a robust statistical framework for air-mass classification, but not deterministic attribution of sources and
620 transformations. Trajectories capture transport pathways but cannot resolve sub-grid emission sources, vertical mixing during transport, or chemical transformations that modify aerosol properties en route. Fourthly, reduced sampling aloft due to TBS operational limitations means conclusions about free-tropospheric CCN or fine-scale features near the highest altitudes should be framed as indicative rather than definitive. This is particularly relevant for understanding the influences of entrainment and the contributions of the residual layer to boundary layer CCN budgets. Finally, cloud responses were constrained by the
625 availability of coincident cloud observations, limiting the ability to establish direct linkages between specific aerosol populations and cloud microphysical responses.

These limitations highlight critical, actionable priorities for future work to achieve process-level closure on aerosol-cloud interactions. The next generation of field studies should move beyond characterization toward targeted, hypothesis-driven experiments. For example, using advanced chemical analyses to constrain source contributions and mixing state, especially



630 for cluster 3-influenced aerosol particles. Extending observations beyond the summer period would capture seasonal shifts in:
(a) marine biogenic emissions (DMS-derived sulfate), (b) photochemical processing rates affecting secondary aerosol
formation, (c) boundary layer depth and mixing dynamics, and (d) synoptic transport patterns influencing the frequency of
each air-mass regime. Winter observations, in particular, could reveal how reduced photochemistry and shallower boundary
layers alter the vertical structure of CCN and their activation characteristics. Finally, the cluster-stratified CCN profiles
635 demonstrated here offer a valuable dataset for evaluating how well regional and global models reproduce: (a) the magnitude
of CCN differences among marine, coastal recirculation, and urban-transported regimes, (b) the vertical structure of CCN
within and above the boundary layer, and (c) the supersaturation dependence (spread ratio) as a proxy for size distribution and
composition representation. Such comparisons could identify specific model deficiencies in aerosol emission inventories,
microphysical process parameterizations, or transport/mixing schemes.

640 **Author contribution**

FM and JW conceptualized the TBS study. BS secured financial support for the project that led to this publication. FM, GWV, and NNL
formally analyzed the data and provided the figures. FM, MPJ, BC, SDB, DZ and DD participated in the field campaign and data collection.
FM, JW, IS, NNL, GWV, ZC, and SC interpreted the results. FM wrote the initial draft of the paper. FM, JW, IS, NNL, GWV, JL, BC, SDB,
MJP, MD, ZC, SC, and DZ proofread and edited the paper.

645 **Data availability.**

The datasets presented in this article are available upon registration at the ARM Data Discovery (<https://adc.arm.gov/discovery/#/>, last
accessed: 27 January 2026). The cloud condensation nuclei concentrations estimated in this study are currently undergoing the data ingest
process with the Atmospheric Radiation Measurement Data Center and will be publicly accessible upon completion of that process. In the
interim, these data are available from the corresponding author upon reasonable request by contacting Fan Mei (fan.mei@pnnl.gov). The
650 NEXRAD data can be requested by contacting Michael Jensen (mjensen@bnl.gov). NEXRAD Level II radar data are publicly available
through the Unidata Amazon Web Services (AWS) S3 archive with anonymous access. In this study, data were programmatically retrieved
using a Python interface to the NEXRAD Level II archive by querying scans for a given radar site (e.g., KHGX, Houston, TX) over specified
UTC time ranges, and the resulting file URLs were used for subsequent processing and analysis.

Competing interests

655 The contact author has declared that none of the authors has any competing interests.



Acknowledgments

This research was supported by the ARM user facility, a US DOE Office of Science user facility managed by the Biological and Environmental Research (BER) program. A portion of this research was performed under project awards
660 (10.46936/lser.proj.2020.51377/60000185 and 10.46936/expl.proj.2021.60186/60008210) as user projects at the Environmental Molecular Sciences Laboratory (EMSL). Battelle operates the Pacific Northwest National Laboratory (PNNL) for the DOE under contract DE-AC05-76RL01830 to support both EMSL and ARM user facilities. JF acknowledges support from ASR under the Integrated Cloud, Land-surface, and Aerosol System Study (ICLASS) Science Focus Area. SDB and BC were supported by the Department of Energy Atmospheric System Research (ASR) program grant DE-SC0021047 to collect and analyze data, and by the ARM field campaigns AFC07023 and AFC07065.
665 Contributions from MPJ were supported by the ASR program under the Process-level AdvancementS of Coupled Cloud and Aerosol LifecycleS (PASCCALS) Science Focus Area through DOE Contract No. DE-SC0012704. JW acknowledges support by the Atmospheric System Research (ASR) program of the U.S. Department of Energy (DOE) under Award No. DE-SC0021017. We acknowledge the use of AI Incubator (including ChatGPT 5.4, Claude-opus-4-6-v1, and gemini-2.5-pro) to assist in refining the language used in this document. Authors also want to express their sincere gratitude for Jerome Fast's contributions to help shape this work.

670

References

- 675 Ajayi, T., Choi, Y., Crosbie, E. C., DiGangi, J. P., Diskin, G. S., Fenn, M. A., Ferrare, R. A., Hair, J. W., Hilario, M. R. A., Hostetler, C. A., Kirschler, S., Moore, R. H., Shingler, T. J., Shook, M. A., Soloff, C., Thornhill, K. L., Voigt, C., Winstead, E. L., Ziemba, L. D., and Sorooshian, A.: Vertical variability of aerosol properties and trace gases over a remote marine region: a case study over Bermuda, *Atmos. Chem. Phys.*, 24, 9197–9218, 10.5194/acp-24-9197-2024, 2024.
- Anderson, B. E., Grant, W. B., Gregory, G. L., Browell, E. V., Collins Jr., J. E., Sachse, G. W., Bagwell, D. R., Hudgins, C. H., Blake, D. R., and Blake, N. J.: Aerosols from biomass burning over the tropical South Atlantic region: Distributions and
680 impacts, *Journal of Geophysical Research: Atmospheres*, 101, 24117–24137, <https://doi.org/10.1029/96JD00717>, 1996.
- Andreae, M. O., Jones, C. D., and Cox, P. M.: Strong present-day aerosol cooling implies a hot future, *Nature*, 435, 1187–1190, 10.1038/nature03671, 2005.
- Ansmann, A., and Müller, D.: Lidar and Atmospheric Aerosol Particles, in: *Lidar: Range-Resolved Optical Remote Sensing of the Atmosphere*, edited by: Weitkamp, C., Springer New York, New York, NY, 105–141, 2005.
- 685 Ansmann, A., Seifert, P., Tesche, M., and Wandinger, U.: Profiling of fine and coarse particle mass: case studies of Saharan dust and Eyjafjallajökull/Grimsvötn volcanic plumes, *Atmos. Chem. Phys.*, 12, 9399–9415, 10.5194/acp-12-9399-2012, 2012.
- Brady, J. M., Stokes, M. D., Bonnardel, J., and Bertram, T. H.: Characterization of a Quadrotor Unmanned Aircraft System for Aerosol-Particle-Concentration Measurements, *Environ Sci Technol*, 50, 1376–1383, 10.1021/acs.est.5b05320, 2016.



- 690 Burton, S. P., Ferrare, R. A., Hostetler, C. A., Hair, J. W., Rogers, R. R., Obland, M. D., Butler, C. F., Cook, A. L., Harper, D. B., and Froyd, K. D.: Aerosol classification using airborne High Spectral Resolution Lidar measurements – methodology and examples, *Atmos. Meas. Tech.*, 5, 73–98, 10.5194/amt-5-73-2012, 2012.
- Chen, B., Thompson, S. A., Matthews, B. H., Sharma, M., Li, R., Nowotarski, C. J., Rapp, A. D., and Brooks, S. D.: A new technique to retrieve aerosol vertical profiles using micropulse lidar and ground-based aerosol measurements, *Atmos. Meas. Tech.*, 18, 5841–5859, 10.5194/amt-18-5841-2025, 2025.
- 695 Cheng, Z., Liyu, A., Dexheimer, D., Lata, N. N., Kulkarni, G., Longbottom, C. M., Mei, F., and China, S.: An automated size and time-resolved aerosol collector platform integrated with environmental sensors to study the vertical profile of aerosols, *Environmental Science: Atmospheres*, 2, 1263–1276, 2022.
- Chiliński, M. T., Markowicz, K. M., and Kubicki, M.: UAS as a Support for Atmospheric Aerosols Research: Case Study, *Pure and Applied Geophysics*, 175, 3325–3342, 10.1007/s00024-018-1767-3, 2018.
- 700 Corrigan, C. E., Roberts, G. C., Ramana, M. V., Kim, D., and Ramanathan, V.: Capturing vertical profiles of aerosols and black carbon over the Indian Ocean using autonomous unmanned aerial vehicles, *Atmos. Chem. Phys.*, 8, 737–747, 10.5194/acp-8-737-2008, 2008.
- Creamean, J. M., de Boer, G., Telg, H., Mei, F., Dexheimer, D., Shupe, M. D., Solomon, A., and McComiskey, A.: Assessing the vertical structure of Arctic aerosols using balloon-borne measurements, *Atmos. Chem. Phys.*, 21, 1737–1757, 10.5194/acp-21-1737-2021, 2021.
- 705 Creamean, J. M., Dexheimer, D., Hume, C. C., Vazquez, M., Hess, B. T. M., Longbottom, C. M., Ruiz, C. A., and Theisen, A. K.: Reaching new heights: A vertically-resolved ice nucleating particle sampler operating on Atmospheric Radiation Measurement (ARM) tethered balloon systems, *EGUsphere*, 2025, 1–22, 10.5194/egusphere-2025-5000, 2025.
- Dexheimer, D., Creamean, J., Gaustad, K., Longbottom, C., Cheng, Z., Sammon, J., and Mei, F.: Tethered Balloon System (TBS) Instrument Handbook, ARM Climate Research Facility; DOE Office of Science Atmospheric Radiation Measurement (ARM) Program (United States), United States, 2026.
- 710 Dmitrovic, S., Hair, J. W., Collister, B. L., Crosbie, E., Fenn, M. A., Ferrare, R. A., Harper, D. B., Hostetler, C. A., Hu, Y., Reagan, J. A., Robinson, C. E., Seaman, S. T., Shingler, T. J., Thornhill, K. L., Vömel, H., Zeng, X., and Sorooshian, A.: High Spectral Resolution Lidar – generation 2 (HSRL-2) retrievals of ocean surface wind speed: methodology and evaluation, *Atmos. Meas. Tech.*, 17, 3515–3532, 10.5194/amt-17-3515-2024, 2024.
- 715 Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies, *Atmos. Chem. Phys.*, 11, 11069–11102, 10.5194/acp-11-11069-2011, 2011.
- 720 Farley, R. N., Lee, J. E., Rivellini, L. H., Lee, A. K. Y., Dal Porto, R., Cappa, C. D., Gorkowski, K., Shawon, A. S. M., Benedict, K. B., Aiken, A. C., Dubey, M. K., and Zhang, Q.: Chemical properties and single-particle mixing state of soot aerosol in Houston during the TRACER campaign, *Atmos. Chem. Phys.*, 24, 3953–3971, 10.5194/acp-24-3953-2024, 2024.



- Fernald, F. G.: Analysis of atmospheric lidar observations: some comments, *Appl. Opt.*, 23, 652–653, 10.1364/AO.23.000652, 1984.
- 725 Ferrero, L., Cappelletti, D., Busetto, M., Mazzola, M., Lupi, A., Lanconelli, C., Becagli, S., Traversi, R., Caiazzo, L., Giardi, F., Moroni, B., Crocchianti, S., Fierz, M., Močnik, G., Sangiorgi, G., Perrone, M. G., Maturilli, M., Vitale, V., Udisti, R., and Bolzacchini, E.: Vertical profiles of aerosol and black carbon in the Arctic: a seasonal phenomenology along 2 years (2011–2012) of field campaigns, *Atmos. Chem. Phys.*, 16, 12601–12629, 10.5194/acp-16-12601-2016, 2016.
- 730 Fröhlich, R., Cubison, M., Slowik, J., Bukowiecki, N., Prévôt, A., Baltensperger, U., Schneider, J., Kimmel, J., Gonin, M., and Rohner, U.: The ToF-ACSM: a portable aerosol chemical speciation monitor with TOFMS detection, *Atmos Meas Tech*, 6, 3225–3241, 2013.
- Fuzzi, S., Baltensperger, U., Carslaw, K., Decesari, S., Denier van der Gon, H., Facchini, M. C., Fowler, D., Koren, I., Langford, B., Lohmann, U., Nemitz, E., Pandis, S., Riipinen, I., Rudich, Y., Schaap, M., Slowik, J. G., Spracklen, D. V., Vignati, E., Wild, M., Williams, M., and Gilardoni, S.: Particulate matter, air quality and climate: lessons learned and future needs, *Atmos. Chem. Phys.*, 15, 8217–8299, 10.5194/acp-15-8217-2015, 2015.
- 735 Galfione, A., Battaglia, A., Oue, M., Cattani, E., and Kollias, P.: Comparison of GOES16 Data with the TRACER-ESCAPE Field Campaign Dataset for Convection Characterization: A Selection of Case Studies and Lessons Learnt, *Remote Sensing*, 17, 2621, 2025.
- 740 Gautam, T., Vandergrift, G. W., Lata, N. N., Cheng, Z., Rahman, A., Minke, A., Lai, Z., Dexheimer, D. N., Zhang, D., Marcus, M. A., Zawadowicz, M. A., Kuang, C., Zhao, R., Steiner, A. L., and China, S.: Chemical Insights into the Molecular Composition of Organic Aerosols in the Urban Region of Houston, Texas, *ACS ES&T Air*, 1, 1304–1316, 10.1021/acsestair.4c00141, 2024.
- Geiß, A., Wiegner, M., Bonn, B., Schäfer, K., Forkel, R., von Schneidmesser, E., Munkel, C., Chan, K. L., and Nothard, R.: Mixing layer height as an indicator for urban air quality?, *Atmos. Meas. Tech.*, 10, 2969–2988, 10.5194/amt-10-2969-2017, 2017.
- 745 Hair, J. W., Hostetler, C. A., Cook, A. L., Harper, D. B., Ferrare, R. A., Mack, T. L., Welch, W., Izquierdo, L. R., and Hovis, F. E.: Airborne High Spectral Resolution Lidar for profiling aerosol optical properties, *Appl. Opt.*, 47, 6734–6752, 10.1364/AO.47.006734, 2008.
- 750 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 755 Huang, R.-J., Zhang, Y., Bozzetti, C., Ho, K.-F., Cao, J.-J., Han, Y., Daellenbach, K. R., Slowik, J. G., Platt, S. M., Canonaco, F., Zotter, P., Wolf, R., Pieber, S. M., Bruns, E. A., Crippa, M., Ciarelli, G., Piazzalunga, A., Schwikowski, M., Abbaszade, G., Schnelle-Kreis, J., Zimmermann, R., An, Z., Szidat, S., Baltensperger, U., Haddad, I. E., and Prévôt, A. S. H.: High



secondary aerosol contribution to particulate pollution during haze events in China, *Nature*, 514, 218–222, 10.1038/nature13774, 2014.

760 Hussein, T., Dal Maso, M., Petäjä, T., Koponen, I. K., Paatero, P., Aalto, P. P., Hämeri, K., and Kulmala, M.: Evaluation of an automatic algorithm for fitting the particle number size distributions, *Boreal environment research*, 10, 337, 2005.

Jensen, M. P., Flynn, J. H., Kollias, P., Kuang, C., McFarquhar, G., Powers, H., Brooks, S., Bruning, E., Collins, D., Collis, S. M., Fan, J., Fridlind, A., Giangrande, S. E., Griffin, R., Hu, J., Jackson, R. C., Kumjian, M., Logan, T., Matsui, T., Nowotarski, C. J., Oue, M., Rapp, A. D., Rosenfeld, D., Ryzhkov, A., Sheesley, R., Snyder, J., Stier, P., Usenko, S., van den Heever, S. C., van Lier-Walqui, M., Varble, A., Wang, Y., Aiken, A., Deng, M., Dexheimer, D., Dubey, M., Feng, Y., Ghaté, V. P., Johnson, K. L., Lamer, K., Saleeby, S. M., Wang, D., Zawadowicz, M. A., and Zhou, A.: Tracking Aerosol Convection Interactions Experiment (TRACER) Field Campaign Report, Oak Ridge National Laboratory (ORNL), Oak Ridge, TN (United States). Atmospheric Radiation Measurement (ARM) Data Center, United States, 2023.

770 Jensen, M. P., Flynn, J. H., Gonzalez-Cruz, J. E., Judd, L. M., Kollias, P., Kuang, C., McFarquhar, G. M., Powers, H., Ramamurthy, P., Sullivan, J., Aiken, A. C., Alvarez, S. L., Argay, P., Argrow, B., Bell, T. M., Boyer, D., Brooks, S. D., Bruning, E. C., Brunner, K., Butterworth, B., Calmer, R., Cappa, C. D., Chakrabarty, R. K., Chandrasekar, V., Chao, C.-Y., Chen, B., China, S., Collins, D. R., Collis, S. M., Crowell, S., Porto, R. D., Boer, G. d., Deng, M., Dexheimer, D., Drager, A. J., Du, X., Dubey, M. K., Dzambo, A. M., Etten-Bohm, M., Fan, J., Farley, R., Feng, Y.-C., Feng, Y., Fenn, M., Ferrare, R. E., Flusche, S., Fridlind, A. M., Galewsky, J., Gamarro, H., Gardner, S., Ghaté, V. P., Giangrande, S. E., Griffin, R. J., Griggs, T., Gronoff, G. P., Grover, M., Gaugenti, M., Guo, F., Gupta, S., Hu, J., Huang, Y., Jackson, R. C., Hair, J. W., Johnson, K. L., Kasparoglu, S., Klein, P., Kotsakis, A. E., Kumar, J., Kumjian, M. R., Lamer, K., Lappin, F. M., Lei, Z., Li, J., Li, R., Li, Y., Logan, T., Lombardo, K., Luke, E. P., Mages, Z., Matthews, A. A., Matthews, B. H., Mayol-Bracero, O., Matsui, T., McKeown, K. E., Mehra, M., Mei, F., Meskidzhe, N., Nguyen, C., Nielsen, E. R., Nowotarski, C. J., Oaks, D., Oktem, R., Oue, M., Park, J. M., Partida, N., Patil, S., Pena, J. C., Petters, M. D., Phoenix, D. B., Puthserry, J. V., Rapp, A. D., Romps, D. M., Roots, M., Rosenfeld, D., Saleeby, S. M., Savala, P., Sedlacek, A. J., Sharma, M., Sheesley, R., Shingler, T. J., Shrestha, S., Singh, A., Smith, E. N., Smith, J. N., Smith, S., Snyder, J., Spicer, E., Spinei, E., Spsychala, M., Steir, P., Storm, M. R., Subba, T., Treserras, B. P., Trojanowski, R., Theisen, A., Thompson, S. A., Twigg, L., Uin, J., Ulinski, A. R., van den Heever, S., van Lier-Walqui, M., Varble, A. C., Wagner, T. J., Wakeen, J., Wales, N. A., Walter, P. J., Wang, D., Wang, J., Wood, L., Wang, Y., Wolde, M., Yoon, S., Young, M. H., Zawadowicz, M. A., Zhang, Q., Zhou, A., Zhu, Z., and Zhu, Z.: Studying Aerosol, Clouds, and Air Quality in the Coastal Urban Environment of Southeastern Texas, *B Am Meteorol Soc*, BAMS–D–23–0331.0331, <https://doi.org/10.1175/BAMS-D-23-0331.1>, 2025.

785 Klett, J. D.: Stable analytical inversion solution for processing lidar returns, *Appl. Opt.*, 20, 211–220, 10.1364/AO.20.000211, 1981.

Lata, N. N., Cheng, Z., Dexheimer, D., Zhang, D., Mei, F., and China, S.: Vertical Gradient of Size-Resolved Aerosol Compositions over the Arctic Reveals Cloud Processed Aerosol in-Cloud and above Cloud, *Environmental Science & Technology*, 57, 5821–5830, 10.1021/acs.est.2c09498, 2023.

790 Li, H., Liu, B., Ma, X., Jin, S., Ma, Y., Zhao, Y., and Gong, W.: Evaluation of retrieval methods for planetary boundary layer height based on radiosonde data, *Atmos. Meas. Tech.*, 14, 5977–5986, 10.5194/amt-14-5977-2021, 2021.

Li, J., Zhang, J., Gong, X., Spielman, S., Kuang, C., Singh, A., Zawadowicz, M. A., Xu, L., and Wang, J.: The sources and diurnal variations of submicron aerosols in a coastal-rural environment near Houston, US, *Atmos Chem Phys*, 2025, 13975–13993, 10.5194/acp-25-13975-2025, 2025.

795



- Liu, S., and Liang, X.-Z.: Observed Diurnal Cycle Climatology of Planetary Boundary Layer Height, *Journal of Climate*, 23, 5790–5809, <https://doi.org/10.1175/2010JCLI3552.1>, 2010.
- Lohmann, U., and Feichter, J.: Impact of sulfate aerosols on albedo and lifetime of clouds: A sensitivity study with the ECHAM4 GCM, *Journal of Geophysical Research: Atmospheres*, 102, 13685–13700, <https://doi.org/10.1029/97JD00631>, 1997.
- Lohmann, U., and Hoose, C.: Sensitivity studies of different aerosol indirect effects in mixed-phase clouds, *Atmos. Chem. Phys.*, 9, 8917–8934, 10.5194/acp-9-8917-2009, 2009.
- Mei, F., Wang, J., Comstock, J. M., Weigel, R., Krämer, M., Mahnke, C., Shilling, J. E., Schneider, J., Schulz, C., Long, C. N., Wendisch, M., Machado, L. A. T., Schmid, B., Krisna, T., Pekour, M., Hubbe, J., Giez, A., Weinzierl, B., Zoeger, M., Pöhlker, M. L., Schlager, H., Cecchini, M. A., Andreae, M. O., Martin, S. T., de Sá, S. S., Fan, J., Tomlinson, J., Springston, S., Pöschl, U., Artaxo, P., Pöhlker, C., Klimach, T., Minikin, A., Afchine, A., and Borrmann, S.: Comparison of aircraft measurements during GoAmazon2014/5 and ACRIDICON-CHUVA, *Atmos. Meas. Tech.*, 13, 661–684, 10.5194/amt-13-661-2020, 2020.
- Mei, F., Wang, H., Zhu, Z., Zhang, D., Zhang, Q., Fast, J. D., Gustafson, W. I., Li, X.-Y., Schmid, B., Niedeck, C., Tomlinson, J., and Flynn, C.: Bridging New Observational Capabilities and Process-Level Simulation: Insights into Aerosol Roles in the Earth System, *B Am Meteorol Soc*, 105, E709–E724, <https://doi.org/10.1175/BAMS-D-23-0110.1>, 2024.
- Mei, F., Zhang, Q., Zhang, D., Fast, J. D., Kulkarni, G., Pekour, M. S., Niedeck, C. R., Glienke, S., Silber, I., Schmid, B., Tomlinson, J. M., Mehta, H. S., Mansoura, X., Cheng, Z., Vandergrift, G. W., Lata, N. N., China, S., and Zhu, Z.: Measurement report: Vertically resolved atmospheric properties observed over the Southern Great Plains with the ArcticShark uncrewed aerial system, *Atmos. Chem. Phys.*, 25, 3425–3444, 10.5194/acp-25-3425-2025, 2025.
- Mushtaq, Z., Sharma, M., Bangotra, P., Gautam, A. S., and Gautam, S.: Atmospheric Aerosols: Some Highlights and Highlighters, Past to Recent Years, *Aerosol Science and Engineering*, 6, 135–145, 10.1007/s41810-022-00133-w, 2022.
- Ramanathan, V., Crutzen, P. J., Lelieveld, J., Mitra, A. P., Althausen, D., Anderson, J., Andreae, M. O., Cantrell, W., Cass, G. R., Chung, C. E., Clarke, A. D., Coakley, J. A., Collins, W. D., Conant, W. C., Dulac, F., Heintzenberg, J., Heymsfield, A. J., Holben, B., Howell, S., Hudson, J., Jayaraman, A., Kiehl, J. T., Krishnamurti, T. N., Lubin, D., McFarquhar, G., Novakov, T., Ogren, J. A., Podgorny, I. A., Prather, K., Priestley, K., Prospero, J. M., Quinn, P. K., Rajeev, K., Rasch, P., Rupert, S., Sadourny, R., Satheesh, S. K., Shaw, G. E., Sheridan, P., and Valero, F. P. J.: Indian Ocean Experiment: An integrated analysis of the climate forcing and effects of the great Indo-Asian haze, *Journal of Geophysical Research: Atmospheres*, 106, 28371–28398, <https://doi.org/10.1029/2001JD900133>, 2001.
- Rapp, A. D., Brooks, S. D., Nowotarski, C. J., Sharma, M., Thompson, S. A., Chen, B., Matthews, B. H., Etten-Bohm, M., Nielsen, E. R., and Li, R.: TAMU TRACER: Targeted Mobile Measurements to Isolate the Impacts of Aerosols and Meteorology on Deep Convection, *B Am Meteorol Soc*, 105, E1685–E1702, <https://doi.org/10.1175/BAMS-D-23-0218.1>, 2024.
- Roach, P. J., Laskin, J., and Laskin, A.: Molecular Characterization of Organic Aerosols Using Nanospray-Desorption/Electrospray Ionization-Mass Spectrometry, *Analytical Chemistry*, 82, 7979–7986, 10.1021/ac101449p, 2010.



- Rosenfeld, D., Andreae, M. O., Asmi, A., Chin, M., de Leeuw, G., Donovan, D. P., Kahn, R., Kinne, S., Kivekäs, N., Kulmala, M., Lau, W., Schmidt, K. S., Suni, T., Wagner, T., Wild, M., and Quaas, J.: Global observations of aerosol-cloud-precipitation-climate interactions, *Reviews of Geophysics*, 52, 750–808, <https://doi.org/10.1002/2013RG000441>, 2014.
- 835 Rosenfeld, D., Zhu, Y., Wang, M., Zheng, Y., Goren, T., and Yu, S.: Aerosol-driven droplet concentrations dominate coverage and water of oceanic low-level clouds, *Science*, 363, eaav0566, doi:10.1126/science.aav0566, 2019.
- Schmale, J., Henning, S., Henzing, B., Keskinen, H., Sellegri, K., Ovadnevaite, J., Bougiatioti, A., Kalivitis, N., Stavroulas, I., Jefferson, A., Park, M., Schlag, P., Kristensson, A., Iwamoto, Y., Pringle, K., Reddington, C., Aalto, P., Äijälä, M., Baltensperger, U., Bialek, J., Birmili, W., Bukowiecki, N., Ehn, M., Fjæraa, A. M., Fiebig, M., Frank, G., Fröhlich, R., Frumau, A., Furuya, M., Hammer, E., Heikkinen, L., Herrmann, E., Holzinger, R., Hyono, H., Kanakidou, M., Kiendler-Scharr, A., Kinouchi, K., Kos, G., Kulmala, M., Mihalopoulos, N., Motos, G., Nenes, A., O'Dowd, C., Paramonov, M., Petäjä, T., Picard, D., Poulain, L., Prévôt, A. S. H., Slowik, J., Sonntag, A., Swietlicki, E., Svenningsson, B., Tsurumaru, H., Wiedensohler, A., Wittbom, C., Ogren, J. A., Matsuki, A., Yum, S. S., Myhre, C. L., Carslaw, K., Stratmann, F., and Gysel, M.: Collocated observations of cloud condensation nuclei, particle size distributions, and chemical composition, *Scientific Data*, 4, 170003, 10.1038/sdata.2017.3, 2017.
- 840
- 845 Seidel, D. J., Ao, C. O., and Li, K.: Estimating climatological planetary boundary layer heights from radiosonde observations: Comparison of methods and uncertainty analysis, *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD013680>, 2010.
- Seinfeld, J. H., and Pandis, S. N.: *Atmospheric chemistry and physics: from air pollution to climate change*, John Wiley & Sons, 2016.
- 850 Silber, I., Comstock, J. M., Kiebertz, M. R., Gaustad, K. L., and Mei, F.: ARM Trajectories Data Set Value-Added Product Report, Oak Ridge National Laboratory (ORNL), Oak Ridge, TN (United States). Atmospheric Radiation Measurement (ARM) Data Center, United States, 2025a.
- Silber, I., Comstock, J. M., Kiebertz, M. R., and Russell, L. M.: ARMTRAJ: a set of multipurpose trajectory datasets augmenting the Atmospheric Radiation Measurement (ARM) user facility measurements, *Earth Syst. Sci. Data*, 17, 29–42, 10.5194/essd-17-29-2025, 2025b.
- 855
- Sivaraman, C., McFarlane, S., Chapman, E., Jensen, M., Liu, S., and Fischer, M.: Planetary boundary layer height (PBL) value added product (VAP): Radiosonde retrievals, DOE Office of Science Atmospheric Radiation Measurement (ARM) Program ..., 2013.
- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J. B., Cohen, M. D., and Ngan, F.: NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System, *B Am Meteorol Soc*, 96, 2059–2077, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 860
- Stull, R. B.: *An introduction to boundary layer meteorology*, Springer Science & Business Media, 2012.
- Thompson, S. A., Chen, B., Matthews, B. H., Li, R., Nowotarski, C. J., Rapp, A. D., and Brooks, S. D.: Characterizing Greater Houston's Aerosol by Air Mass During TRACER, *Journal of Geophysical Research: Atmospheres*, 130, e2025JD043353, <https://doi.org/10.1029/2025JD043353>, 2025a.
- 865



Thompson, S. A., Peña, T. A., Chen, B., Sharma, M., Matthews, B. H., Li, R., Nowotarski, C. J., Rapp, A. D., and Brooks, S. D.: Variability in Ice Nucleating Particles Across Greater Houston Texas, *Journal of Geophysical Research: Atmospheres*, 130, e2025JD044164, <https://doi.org/10.1029/2025JD044164>, 2025b.

870 Tsimpidi, A. P., Scholz, S. M. C., Milousis, A., Mihalopoulos, N., and Karydis, V. A.: Aerosol Composition Trends during 2000–2020: In depth insights from model predictions and multiple worldwide observation datasets, *EGUsphere*, 2024, 1–66, 10.5194/egusphere-2024-3590, 2024.

Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of Atmospheric Sciences*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.

875 Vandergrift, G. W., Shawon, A. S. M., Dexheimer, D. N., Zawadowicz, M. A., Mei, F., and China, S.: Molecular Characterization of Organosulfate-Dominated Aerosols over Agricultural Fields from the Southern Great Plains by High-Resolution Mass Spectrometry, *ACS Earth and Space Chemistry*, 6, 1733–1741, 10.1021/acsearthspacechem.2c00043, 2022.

880 Vandergrift, G. W., Dexheimer, D. N., Zhang, D., Cheng, Z., Lata, N. N., Rogers, M. M., Shrivastava, M., Zhang, J., Gaudet, B. J., Mei, F., and China, S.: Tethered balloon system and High-Resolution Mass Spectrometry Reveal Increased Organonitrates Aloft Compared to the Ground Level, *Environmental Science & Technology*, 58, 10060–10071, 10.1021/acs.est.4c02090, 2024.

Wang, D., Kobrosly, R., Zhang, T., Subba, T., van den Heever, S., Gupta, S., and Jensen, M.: Aerosol impacts on isolated deep convection: findings from TRACER, *Atmos. Chem. Phys.*, 25, 9295–9314, 10.5194/acp-25-9295-2025, 2025.