

High Yields of Formic Acid and Acetic Acid during Multi-generational Oxidation of Toluene

Hengqing Shen^{1,2}, Liubin Huang^{1*}, Yue Zhao³, Min Zhao¹, Yu Yang¹, Huan Li⁴, Huihui Wu²,
Zhongming Chen^{2*}

¹Environment Research Institute, Shandong University, Qingdao, Shandong 266237, China

²State Key Laboratory of Regional Environment and Sustainability, College of Environmental Sciences and Engineering, Peking University, Beijing 100871, China

³School of Environmental Science and Engineering, Shanghai Jiao Tong University, Shanghai 200240, China

⁴China Center for Information Industry Development, Beijing 100871, China

***Corresponding Authors**

Liubin Huang (hliubin@sdu.edu.cn) and Zhongming Chen (zmchen@pku.edu.cn).

ABSTRACT

Formic acid and acetic acid are the most abundant gas-phase organic acids in the atmosphere, yet their concentrations are substantially underestimated by both global and regional atmospheric models across diverse environments. In this study, we report unexpectedly high yields of formic acid and acetic acid formed during the multi-generational photooxidation of toluene, a canonical anthropogenic volatile organic compound. Their yields exhibit a strong dependence on hydroxyl radical ($\bullet\text{OH}$) exposure ($[\bullet\text{OH}] \times \text{residence time}$), increasing from 25% and 24% under low exposure (<0.2 equivalent days) to 74% and 40% under elevated exposure (1–3 equivalent days) for formic and acetic acid, respectively. The formation of these organic acids is not significantly affected by NO_x concentrations. A modified box model based on MCM v3.3.1 underestimates the peak concentrations of both acids by approximately a factor of five, indicating substantial gaps in current mechanistic understanding. Although both secondary aerosol formation and organic acid production increase with aging to a certain degree of oxidation, their distinct temporal evolutions suggest that particle photodegradation is not the dominant pathway. The contrasting $\bullet\text{OH}$ exposure dependence between organic acids and primary carbonyl compounds further implies that these acids are predominantly multi-generational oxidation products. These findings demonstrate that multi-generational oxidation of aromatic compounds is an important and previously underappreciated source of atmospheric organic acids. The omission of organic acid formation from aromatic oxidation in current chemical mechanisms likely contributes to their widespread underestimation in models, highlighting the need for detailed laboratory studies and updated chemical mechanisms.

1 INTRODUCTION

Organic acids are ubiquitous in the troposphere and are important contributors to precipitation acidity (Stavrakou et al., 2012). Among them, formic acid (HCOOH) and acetic acid (CH_3COOH) are the most abundant gas-phase organic acids in various atmospheric environments, with mixing ratios spanning from several pptv to tens ppbv (Chebbi and Carlier, 1996; Veres et al., 2011; Mungall et al., 2018; Ding et al., 2025). Formic acid alone can contribute to more than half of rainwater acidity in many continental regions (Stavrakou et al., 2012). In addition, these organic acids can also affect secondary aerosol formation by altering aerosol pH and gas-particle partitioning processes (Shen et al., 2018; Tao and Murphy, 2019).

On a global scale, the dominant source of formic acid and acetic acid is considered to be secondary production from the photooxidation of biogenic volatile organic compounds (VOCs), particularly isoprene and its oxidation products (Paulot et al., 2011; Khan et al., 2018; Link et al., 2020). However, rapid secondary production and high amounts of formic acid and acetic acid have also been observed in anthropogenic air masses and urban environments (Veres et al., 2011; Yuan et al., 2015; Millet et al., 2015; Liggio et al., 2017), suggesting that anthropogenic VOCs may play a significant role in the formation of formic acid and acetic acid in polluted areas. Despite this, the sources and detailed formation pathways of atmospheric organic acids remain poorly understood. This is evidenced by the fact that current atmospheric chemical models substantially underestimate (up to >10 times) their field-observed concentrations on both global (Paulot et al., 2011; Stavrakou et al., 2012; Chaliyakunnel et al., 2016; Khan et al., 2018) and regional scales (Le Breton et al., 2012; Yuan et al., 2015; Millet et al., 2015; Liggio et al., 2017) in the free troposphere (Paulot et al., 2011; Millet et al., 2015), the forest (Stavrakou et al., 2012; Schobesberger et al., 2016), the urban (Yuan et al., 2015; Bannan et al., 2017; Khan et al., 2018), the polar region (Paulot et al., 2011; Mungall et al., 2018), and the oil sands region (Yuan et al., 2015; Liggio et al., 2017). Such widespread underestimation implies the existence of large and pervasive missing or underestimated sources in current models (Yuan et al., 2015; Liggio et al., 2017). Numerous hypotheses have been proposed to reconcile this model–observation gap, including previously unrecognized contributions from biomass burning (Paulot et al., 2011; Chaliyakunnel et al., 2016), unmeasured anthropogenic intermediate VOCs (Le Breton et al., 2012; Liggio et al., 2017), acetaldehyde tautomerization (Shaw et al., 2018), cloud-mediated methanediol oxidation (Franco et al., 2021), and chemical aging of organic aerosols (Paulot et al., 2011; Malecha and Nizkorodov, 2016, 2017; Bates et al., 2023; Jiang et al., 2023). Nevertheless, none of these processes alone can fully account for the high ambient

concentrations of formic acid and acetic acid observed in the atmosphere (Yuan et al., 2015; Liggio et al., 2017; Franco et al., 2021).

Recent studies have suggested that resolving this discrepancy requires much higher effective molar yields of formic acid from the oxidation of monoterpenes or anthropogenic VOCs than currently assumed (Stavrakou et al., 2012; Liggio et al., 2017). For example, Liggio et al. (2017) estimated that ~50% of formic acid yield would be necessary to reconcile observations and modeling results in an oil sand region (Liggio et al., 2017). In contrast, laboratory-reported yields of formic acid and acetic acid from VOC photooxidation are typically below 15% (Berndt and Böge, 2001; Baltensperger et al., 2005; Paulot et al., 2011; Yuan et al., 2015). Notably, most of these yields were measured under relatively low $\bullet\text{OH}$ exposures ($[\bullet\text{OH}] \times \text{residence time}$), corresponding to less than one equivalent atmospheric photochemical day (Praplan et al., 2014). However, recent studies have revealed that molecular fragmentation reactions become increasingly important with oxidative aging (Lambe et al., 2012; Ortega et al., 2013; Isaacman-VanWertz et al., 2018), potentially enhancing the formation of small oxygenated species. Consistent with this hypothesis, substantial increases in gas-phase organic acids have been observed during the aging of biomass-burning emissions (Ortega et al., 2013; Bruns et al., 2017; Permar et al., 2023) and diesel exhaust (Friedman et al., 2017). For instance, Bruns et al. (2017) reported that formic acid concentrations increased by factors of ~5–50 with aging. Friedman and Farmer (2018) further demonstrated that the molar yield of organic acids strongly depends on $\bullet\text{OH}$ concentrations during the photooxidation of monoterpenes. These findings suggest that the persistent model–observation discrepancy may be owing to an incomplete mechanistic and quantitative understanding of organic acid formation during multi-generational oxidation under atmospherically relevant conditions.

Oxidation flow reactor (OFR) provides a powerful complement to traditional environmental chambers by enabling the simulation of oxidation equivalent to multiple days of atmospheric aging within residence times of seconds to minutes through the use of elevated oxidant concentrations (Lambe et al., 2015). A key advantage of OFR is its ability to achieve high photochemical exposure while minimizing wall losses of gases and particles, which can be substantial in Teflon chambers due to the typically long residence time employed (Zhang et al., 2014; Brune, 2019). The validity of OFR has been demonstrated in numerous studies (Lambe et al., 2015; Peng et al., 2015), and they are increasingly utilized to investigate the effects of photochemical aging on gas-phase chemistry (Lambe et al., 2015; Friedman and Farmer, 2018), secondary organic aerosol (SOA) formation (Brunns et al., 2015), and aerosol optical properties

(Lambe et al., 2013). In this study, we employ an OFR to investigate the formation of formic acid and acetic acid during the photooxidation of toluene, a canonical and highly abundant anthropogenic VOC, over a wide range of $\bullet\text{OH}$ exposures, and examine how these processes are affected by relative humidity (RH), initial toluene and NO_x concentrations. By the combination of laboratory experiments and model simulations, the underlying mechanisms for the formation of formic acid and acetic acid are further explored.

2 METHODS

2.1 Oxidation Flow Reactor

Toluene photooxidation experiments were conducted in a custom-built quartz oxidation flow reactor (OFR) with a total volume of 2 L (1 m in length and 5 cm inner diameter; Figure 1). The reactor temperature was maintained at 298 K by a continuously circulating water jacket. Low-pressure mercury lamps (primary emission at 254 nm) were evenly distributed around the reactor, with up to eight lamps operating simultaneously. The OH radicals in the OFR were generated via the reaction of $\text{O}(^1\text{D}) + \text{H}_2\text{O} \rightarrow 2 \bullet\text{OH}$, where $\text{O}(^1\text{D})$ was produced by the photolysis of externally introduced O_3 . O_3 was generated upstream of the OFR using a custom-built O_3 generator in which O_2 was irradiated at 185 nm. Photons at 185 nm were effectively filtered by the quartz lamp sleeves, the quartz reactor walls, and the water jacket, thereby minimizing in situ O_3 formation inside the OFR (<10 ppbv). A movable sampling port (6 mm outer diameter) enabled gas and particle sampling at different axial positions along the OFR. The $\bullet\text{OH}$ exposure within the reactor was determined using the box model described in Section 2.3, and the results are provided in Table S1 and Figure S1. At 20% RH, the modeled $\bullet\text{OH}$ exposures ranged from 2.03×10^{10} to 1.97×10^{12} molecules cm^{-3} s, corresponding to 0.07–6.34 days of equivalent atmospheric aging assuming a 24-h average $\bullet\text{OH}$ concentration of 3.6×10^6 molecules cm^{-3} in urban Beijing (Tan et al., 2018). At 70% RH, the equivalent atmospheric aging times in the experiments ranged from 0.20 to 15.64 days.




Figure 1. Schematic diagram of the experimental setup.

2.2 Experimental Procedures

Toluene gas with an initial concentration of ~ 94 ppbv was generated by passing a flow of N_2 (99.999%, Beijing Haipubeifen Gas) over liquid toluene (Merck, $\geq 99.9\%$) in a temperature-controlled diffusion tube. RH in the OFR was adjusted by introducing water vapor generated using a bubbler. The gas mixture of toluene, O_3 , and water vapor were continuously introduced into the OFR at a total flow rate of 2 L min^{-1} , and the initial concentration of O_3 was about 1580 ppbv. Samples were collected at axial positions corresponding to 5%, 25%, 45%, 65%, and 85% of the reactor length using the movable sampling port. Under laminar flow conditions, these positions corresponded to residence times of 3.9, 15.8, 27.5, 39.3, and 51.0 seconds, respectively. Additional experiments were conducted with 2, 4, 6, or 8 lamps to achieve higher $\bullet\text{OH}$ exposures. The corresponding $\bullet\text{OH}$ exposures under different experimental conditions are summarized in Table S1. The influence of NO_x on organic acid formation was investigated by adding N_2O to the OFR. NO and NO_2 were produced via the reaction $\text{O}(^1\text{D}) + \text{N}_2\text{O} \rightarrow 2\text{NO}$, followed by $\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$ (Lambe et al., 2017). The NO concentration was systematically varied by adjusting the N_2O flow rate (0, 20, 40, 80, 120, 160, and 200 mL min^{-1}). All NO_x experiments were conducted using a single lamp, with sampling at the reactor outlet corresponding to a residence time of 60 s. It should be noted that although both the O_3 concentration and the 254 nm photon flux in the OFR substantially exceed typical ambient levels, $\bullet\text{OH}$ concentration is enhanced to an even greater extent, ensuring that $\bullet\text{OH}$ -initiated reactions remain the dominant oxidation pathway.

Toluene concentrations were quantified using gas chromatography with a flame ionization detector (GC-FID, Agilent 7890A) (Wu et al., 2017). O_3 and SO_2 were measured using an O_3 analyzer (model 202, 2B Technologies, USA) and an SO_2 analyzer (model 43i-TLE, Thermo), respectively. Particle number size distributions were measured using a scanning mobility particle sizer (SMPS, DMA 3081 connected to CPC 3776, TSI Inc.). Carbonyl compounds were collected using the cold trap method (Huang et al., 2013) and analyzed by high-performance liquid chromatography (HPLC) with ultraviolet detection (Shen et al., 2018, 2024). All experiments were conducted in the absence of seed particles. Gas-phase organic acids were collected using a temperature-controlled scrubbing glass coil maintained at 277 K and extracted with ultrapure water as the stripping solution (Wu et al., 2017; Xing et al., 2018). Gas and stripping solution flow rates were 0.6 L min^{-1} and 0.2 mL min^{-1} , respectively. The sample collection efficiencies were measured as approximately 85% for acetic acid, 95% for formic acid, and 99% for pyruvic acid. The eluent was collected for 15 min, and then the sample was

analyzed immediately using ion chromatography (DIONEX ICS–2000) (Shen et al., 2018). The concentrations of organic acids were quantified using a mixed liquid standard solution containing formic acid, acetic acid, and pyruvic acid. To evaluate potential artifacts associated with aqueous-phase formation in the stripping solution, five additional gas-phase samples were collected using Horibe tubes with an ethanol cold trap maintained at 183 K. Concentrations measured using the scrubbing coil and Horibe tubes agreed within 10%, indicating that aqueous-phase artifacts had a negligible influence on measurements. The molar yields of formic acid and acetic acid were calculated relative to the amount of toluene consumed at each sampling position as show in equation E1:

$$\text{Yield (\%)} = (\Delta[\text{organic acid}] / \Delta[\text{toluene}]) \times 100\% \quad \text{E1}$$

where $\Delta[\text{organic acid}]$ is the concentration of formic acid or acetic acid (ppbv) produced at a given sampling position, and $\Delta[\text{toluene}]$ is the corresponding amount of toluene consumed (ppbv) at that position. It should be noted that the reported yields are apparent yields and were not corrected for the subsequent reactions of organic acids with $\bullet\text{OH}$.

A series of control experiments were conducted to ensure data quality and reproducibility. (1) Before each experiment, high concentrations of O_3 were introduced into the OFR with two lamps illuminated until particle concentrations measured by the SMPS were negligible ($< 50 \text{ cm}^{-3}$ or $0.1 \mu\text{g m}^{-3}$) and no impurities were detected by GC-FID. (2) O_3 photolysis experiments were performed prior to each toluene photooxidation experiment to verify consistent irradiation conditions based on comparable O_3 decay rates. (3) Blank experiments were conducted without toluene, with toluene but without O_3 , and with both toluene and O_3 but without water vapor. Formic and acetic acid concentrations observed in these blank experiments were low relative to those measured during toluene photooxidation ($< 15\%$). All reported data were corrected for blank contributions. In addition, the wall loss rates of formic acid and acetic acid were measured to be less than 5% of their respective $\bullet\text{OH}$ reaction rates under our experimental conditions and were therefore not corrected for in the yield calculations.

2.3 Model Description and Determination of Photon Flux

A zero-dimensional box model based on the Master Chemical Mechanism (MCM) v3.3.1 (<http://mcm.leeds.ac.uk/MCM/>) and modified according to previous work (Li et al., 2015) was employed to simulate toluene photooxidation and organic acid formation in the OFR (Jenkin et al., 2015). Rate coefficients for additional reactions were adopted from the latest JPL kinetic evaluation (Li et al., 2015). Photolysis cross sections and quantum yields were taken from

Keller-Rudek et al. (2013). The initial concentrations of toluene, O₃, and water vapor were set according to the respective experimental conditions.

Accurate quantification of the photon flux in the OFR is essential for determining photolysis rates in the model. Direct measurement of photon flux is challenging due to attenuation by the water jacket and quartz reactor walls. Therefore, photon flux in the OFR was determined by combining model simulations with direct measurements of O₃ decay from experiments conducted in the absence of toluene. The photon flux was iteratively adjusted in the model until the simulated O₃ decay rate matched the observed decay. Using this approach, the photon flux with one lamp illuminated was estimated to be approximately 3.0×10^{15} photons cm⁻³ s⁻¹, which is consistent with photon fluxes commonly used in OFR studies (Peng et al., 2016). Photon fluxes corresponding to different numbers of illuminated lamps (up to eight) were determined using the same method, and the results are shown in Figure S2. The validity of this approach for photon flux determination, as well as the applicability of the box model to the OFR system, was further evaluated by comparing simulated and measured SO₂ decay rates (Figure S3).

3 RESULTS AND DISCUSSION

3.1 Production of Organic Acids in Toluene Photooxidation

3.1.1 •OH Exposure Dependence

The major gas-phase organic acids observed during the photooxidation of toluene were formic acid, acetic acid, and pyruvic acid, with minor contributions from lactic acid and oxalic acid at both 20% and 70% RH. Because formic acid and acetic acid together can account for approximately 90% of the total measured gas-phase organic acids, the following analysis focuses primarily on these two species. Figure 2 illustrates the dependence of the concentrations and molar yields of formic acid and acetic acid on •OH exposure. As •OH exposure increased, their concentrations initially increased and subsequently decreased, reaching maximum at approximately 1–3 equivalent days under both RH conditions. The similar trends observed for formic acid and acetic acid suggest that their formation is governed by closely related chemical pathways during toluene photooxidation. **The decrease in organic acid concentrations at higher •OH exposure is likely due to kinetic competition between organic acid formation and their subsequent degradation (including direct •OH reaction and depletion of their precursors) as OH exposure increases.** At the highest •OH exposure investigated (15.64 equivalent days), the concentrations of both acids decreased to about 25% of their respective peak values. As shown in Figures 2a and 2c, the evolution of both formic acid and acetic acid concentrations with the increasing •OH exposure at 20% RH is similar to that at 70% RH. However, it is found that

organic acid concentrations were consistently higher at low RH (20%) than at high RH (70%) under identical $\bullet\text{OH}$ exposures, suggesting that reactions involving water vapor may play a limited role in organic acid formation in this study, in contrast to the substantial contribution of stabilized Criegee intermediate–water reactions commonly observed during alkene ozonolysis (Neeb et al., 1997; Orzechowska and Paulson, 2005; Kang et al., 2025).

Figures 2b and 2d show a strong dependence of the molar yields of formic acid and acetic acid on $\bullet\text{OH}$ exposure at both 20% and 70% RH. The yields increase from 25% (formic acid) and 24% (acetic acid) at low $\bullet\text{OH}$ exposures (<0.2 equivalent days) to maximum values of 74% and 40%, respectively, at intermediate $\bullet\text{OH}$ exposures (1–3 equivalent days). Previous studies have shown that molecular fragmentation reactions become increasingly important under conditions of elevated $\bullet\text{OH}$ exposure (Lambe et al., 2012; Ortega et al., 2013), which are difficult to achieve in traditional environmental chamber experiments that typically span less than one equivalent day of atmospheric aging. The pronounced increase in organic acid yields with $\bullet\text{OH}$ exposure suggests that molecular fragmentation processes likely play an important role in formic acid and acetic acid formation.




Figure 2. Production of formic acid and acetic acid during the photooxidation of toluene as a

function of •OH exposure. Panels (a) and (b) correspond to 20% RH, and panels (c) and (d) correspond to 70% RH. The initial toluene concentration was ~94 ppbv. Error bars represent standard deviations of repeated experiments under identical conditions. **Note that the first data point in each panel corresponds to a condition where measurable toluene consumption has already occurred, with •OH exposures of 0.07 equivalent days at 20% RH and 0.2 equivalent days at 70% RH, respectively.**


Furthermore, we also compared the yields of formic acid and acetic acid measured from the photooxidation of toluene with literature results. Seuwen and Warneck (1996) quantified the yield of formic acid during toluene oxidation at approximately 13%, which is ~5 times lower than the peak value observed in our study. In addition to toluene, the yields of formic and acetic acid produced from other aromatic compounds reported in previous studies were also lower than our results (Gery et al., 1985; Bandow and Washida, 1985; Bandow et al., 1985; Berndt et al., 1999; Wyche et al., 2009; Müller et al., 2012). For example, Berndt et al. (1999) reported a formic acid yield of ~13% from benzene oxidation, and Wyche et al. (2009) observed a formic acid yield of 14% from 1,3,5-trimethylbenzene under high-NO_x conditions (Berndt et al., 1999; Wyche et al., 2009). **This discrepancy may be ascribed to the differences in •OH exposure levels. Previous chamber studies were typically conducted under •OH exposures of <0.1 equivalent days (Seuwen and Warneck, 1996; Wyche et al., 2009), which are substantially lower than those (1–3 equivalent days) at which peak organic acid yields were observed in our study. Under such limited •OH exposures, primary oxidation products dominate, whereas formic and acetic acids likely form as secondary or later-generation products requiring higher •OH exposures (Lambe et al., 2012; Isaacman-VanWertz et al., 2018; Drozd et al., 2019).** Evidence supporting the role of higher •OH exposures in organic acid formation comes from two previous studies. Wyche et al. (2009) observed formic acid and acetic acid yields of 49% and 33%, respectively, during the photooxidation of 1,3,5-trimethylbenzene under low-NO_x conditions, where the aromatic precursor was nearly fully consumed. Similarly, Murschell and Farmer (2018) reported formic acid yields as high as 43% during the oxidation of chlorinated aromatic herbicide 2-methyl-4-chlorophenoxyacetic acid under very high •OH exposures. Consistent with these findings, combustion studies, which effectively accelerate atmospheric oxidation, also reported enhanced formation of formic acid and acetic acid from toluene relative to non-aromatic compounds such as propane and isooctane (Zervas, 2005; Battin-Leclerc et al., 2007). Additionally, previous experiments often used very high concentrations of aromatic compounds (hundreds of ppbv to ppmv levels), which may suppress •OH concentrations and limit effective oxidative aging. Our

experiments similarly verify that increasing toluene concentrations inhibit the formation of organic acids (Figure S4). At 20% RH, the yield of formic acid decreased from 74% to 48% as the initial toluene concentration increased from 94 to 381 ppbv. Together, these results highlight the critical role of •OH exposure in controlling organic acid formation during aromatic oxidation. Further discussion of these yield discrepancies is provided in Sections 3.2 and 3.3.

3.1.2 NO_x Dependence

Small organic acids, such as formic and acetic acid, are commonly thought to form via reactions of peroxyacyl radicals with HO₂/RO₂ radicals (HO₂•/RO₂•). The presence of NO_x is expected to suppress this pathway by preferentially reacting with peroxy radicals, diverting them toward NO reaction channels that favor nitrate and carbonyl formation over organic acid production. To investigate the influence of NO_x on organic acid formation, N₂O was introduced into the OFR to generate NO_x resulting in NO concentrations of 0–29 ppbv. It should be noted that NO and NO₂ were produced simultaneously in our experiments (via O(¹D) + N₂O → 2NO, followed by NO + O₃ → NO₂ + O₂), and model simulations indicate that their concentrations were strongly correlated (R² = 0.98). NO primarily reacts with RO₂ to form alkoxy radicals (RO•), which can divert the traditional organic acid formation pathway toward carbonyl products. NO₂ can react with peroxyacyl radicals to form peroxyacyl nitrates (PANs), potentially sequestering acyl peroxy intermediates, and can also react with •OH, thereby affecting •OH concentration. As shown in Figure 3, molar yields of formic acid and acetic acid were not significantly affected by NO concentration at either 20% or 70% RH. Even at the highest NO level (~29 ppbv), which the ratio of NO concentration to HO₂• (simulated by model) exceeded 1000, substantial production of organic acids (~40% for formic acid and ~20% for acetic acid at 20% RH) were still observed. If organic acid formation were dominated by the peroxy radical pathway, such high NO: •HO₂ ratios would be expected to substantially suppress their formation. However, at 70% RH, organic acid yields even increased slightly with increasing NO concentrations. (Crounse et al., 2013). These results indicate that neither the NO-mediated RO₂• diversion pathway nor the NO₂-related pathways significantly control organic acid production. The observed ignorable suppression therefore provides robust evidence that a NO_x-insensitive pathway dominates organic acid production during toluene photooxidation. This finding is consistent with field observations of rapid organic acid formation in polluted environments, where NO_x levels are typically elevated (Veres et al., 2011; Yuan et al., 2015; Millet et al., 2015; Liggitto et al., 2017), and offers a plausible explanation for the persistent underestimation of organic acid concentrations in atmospheric models under both pristine and polluted conditions



306 (Stavrakou et al., 2012; Yuan et al., 2015; Bannan et al., 2017; Khan et al., 2018).



307 **Figure 3.** NO_x dependence of formic acid and acetic acid yields under 20% RH (a) and 70%
 308 RH (b) conditions. Error bars represent standard deviations from repeated experiments under
 309 identical conditions.

310 3.2 Underestimated Organic Acid Yields in Chemical Models


311 To further assess the ability of current chemical mechanisms to reproduce observed organic
 312 acid formation, we compared our experimental results with simulations using the MCM v3.3.1
 313 model. For acetic acid, two formation pathways are included in MCM v3.3.1 (reactions R1 and
 314 R2):



315 Model simulations indicate that reaction R2 ($\text{CH}_3\text{CO}_3\bullet + \text{RO}_2\bullet$) dominates acetic acid
 316 production (>90%). While the model can reproduce the general trend of acetic acid
 317 concentrations, that initially increasing and then decreasing with $\bullet\text{OH}$ exposure, it substantially
 318 underestimates the observed values. The maximum modeled yield (~7% at ~3 equivalent days)
 319 is roughly five times lower than the measured yields under both RH conditions (Figure 4a–b).
 320 This discrepancy suggests that additional formation pathways, currently absent from the model,
 321 may significantly contribute to acetic acid production during toluene photooxidation.

322 It is noted that formic acid is not considered as the product of $\bullet\text{OH}$ oxidation of aromatic
 323 compounds including toluene in the current MCM v3.3.1 model. Therefore, prior to model
 324 simulation, we implemented four candidate pathways based on previous studies (Yetter et al.,
 325 1989; Archibald et al., 2007; Fittschen et al., 2014; Yuan et al., 2015; Shaw et al., 2018). (1)
 326 $\text{HOCH}_2\text{OO}\bullet$ chemistry: Analogous to the acetic acid mechanism, formic acid can also be
 327 formed from either the reaction of $\text{HOCH}_2\text{OO}\bullet$ with $\text{HO}_2\bullet$ or the bimolecular reaction of

328 HOCH₂OO• (Jenkin et al., 2008; Yuan et al., 2015). (2) Vinyl alcohol (CH₂=CHOH) oxidation:
 329 Formic acid formation via •OH-initiated oxidation of vinyl alcohol, which is generated through
 330 acetaldehyde tautomerization. Both photo-induced and organic-acid-catalyzed tautomerization
 331 processes were included (Archibald et al., 2007; Yuan et al., 2015; Shaw et al., 2018). (3) HCHO
 332 + •OH reaction: Direct formation through HCHO + •OH → HCOOH + H, with a rate coefficient
 333 of $2.0 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Yetter et al., 1989). (4) CH₃O₂• + •OH reaction: Formic acid
 334 formation via CH₃O₂• + •OH → HCOOH + H₂O, with a very high rate coefficient of 2.8×10^{-10}
 335 $\text{cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Fittschen et al., 2014). **It should be noted that the branching ratio for formic**
 336 **acid formation from this reaction remains uncertain, thus, the yield of 100% was assumed in**
 337 **our model, representing an upper-limit estimation.** Model evaluation results suggest that
 338 pathways 1–3 contribute minimally (<1%). Pathway 1 (HOCH₂OO• chemistry) exhibits much
 339 lower yield than the analogous acetic acid mechanism due to the low stability and consequently
 340 low concentration of HOCH₂OO• relative to CH₃CO₃ radicals. Pathways 2 and 3 involve
 341 aldehydes as precursors, which also limit formic acid production. Only pathway 4 (CH₃O₂• +
 342 •OH) can produce a substantial amount of formic acid, with a maximum modeled yield of ~13%
 343 at ~3 equivalent days. **Even under this upper-limit assumption, the modeled concentrations of**
 344 **formic acid are still approximately five times lower than the experimental measurements**
 345 **(Figures 4c–d).** It is noted that pathway 4 is highly sensitive to NO_x. Given that our experimental
 346 results show that organic acid formation is insensitive to NO_x levels, this pathway is unlikely
 347 to be the dominant mechanism for organic acid production. Overall, these results indicate that
 348 known pathways account for only a small fraction of the observed formic and acetic acid
 349 production, implying the existence of additional, yet unidentified, formation mechanisms.






Figure 4. Comparison of measured and modeled concentrations of formic acid and acetic acid at different OH exposures. (a) Acetic acid at 20% RH; (b) Acetic acid at 70% RH; (c) Formic acid at 20% RH; (d) Formic acid at 70% RH.

3.3 Organic Acids Formed from the Multi-generation oxidation

Previous modeling studies have suggested that the aging of SOA could significantly contribute to the atmospheric budgets of organic acids (Paulot et al., 2011; Ervens et al., 2011), and laboratory experiments have confirmed the production of organic acids during SOA photodegradation (Malecha and Nizkorodov, 2016, 2017; Jiang et al., 2023). In this study, we analyzed the relationship between SOA particle mass and organic acid production to evaluate this potential pathway. However, the distinctly different temporal profiles of gas-phase organic acid production and SOA formation (Figure 5) imply that SOA photodegradation is not the primary source of the observed high organic acid yields. First, SOA formation exhibited a delay relative to the rapid production of organic acids, aligning with the “incubation period” for aromatic SOA formation (Ng et al., 2007; Wyche et al., 2009). Substantial organic acid yields are observed even when the SOA mass concentration was negligible (Figure 5a), effectively ruling out SOA photodegradation as the primary source. Second, as gas-phase organic acid concentrations began to decline, SOA mass continued to increase (Figure 5), indicating decoupled formation and removal pathways for SOA and organic acids. At very high $\bullet$ OH exposures (Figure 5b), organic acid concentrations decreased while the SOA mass remained nearly unchanged. This suggests that gas-phase organic acids are more reactive toward gas-phase oxidants than the more persistent SOA, aligning with the generally longer lifetimes of aerosol-phase organics against oxidation (Isaacman-VanWertz et al., 2018). Although SOA photodegradation can produce small oxygenated VOCs (including formic acid and acetic acid), previous studies showed that aromatic-derived SOA generates much lower yields of such

products compared to biogenic SOA (e.g., derived from isoprene or α -pinene) (Malecha and Nizkorodov, 2016). It should be noted that this conclusion is specific to toluene-derived SOA under our experimental conditions. In the real atmosphere, SOA formed from other precursors (e.g., biogenic VOCs or biomass burning emissions) may exhibit substantially higher organic acid yields upon photodegradation, and the contribution of aerosol-phase chemistry to the overall organic acid budget cannot be excluded based on our results alone. Additionally, it is worth noting that a greater aerosol mass was formed at 70% RH than at 20% RH under comparable $\bullet$ OH exposures (Figure 5). This enhanced SOA formation at higher humidity may partially account for the lower organic acid yields observed under high-RH conditions. Specifically, the larger partitioning of water-soluble intermediates into the particle phase could reduce their availability for gas-phase fragmentation into organic acids, and enhanced peroxide formation may divert intermediates toward non-acid-producing pathways.




Figure 5. Correlations of particle mass concentration with gas-phase formic acid and acetic acid during the photooxidation of toluene at different $\bullet$ OH exposure. a: 20% RH; b: 70% RH. Color of marker represents different $\bullet$ OH exposure levels.

To further investigate the origin of the high organic acid yields observed during toluene photooxidation, we first examined key first-generation ring-opened products, namely glyoxal and methylglyoxal, which are well-known primary products in aromatic photooxidation (Volkamer et al., 2001). In our experiments, the molar yields of these carbonyl compounds at low $\bullet$ OH exposure were 11% and 10%, respectively, aligning with results reported in previous studies (Gery et al., 1985; Bandow et al., 1985). Figure 6 displays that the yields of glyoxal and methylglyoxal decreased rapidly with increasing $\bullet$ OH exposure, dropping to approximately 1% at higher exposures. In contrast, the yields of formic acid and acetic acid showed a distinct maximum value at intermediate $\bullet$ OH exposures (1–3 equivalent days). Previous studies

proposed that oxidation of glyoxal and methylglyoxal can contribute to formic acid and acetic acid formation (Paulot et al., 2011). However, these pathways are excluded in this study given the substantially lower yields of glyoxal and methylglyoxal observed. Moreover, the poor linear correlation between the yields of formic acid and acetic acid and the first-generation ring-opened products suggests that these acids are secondary or later-generation products formed through multistep oxidation processes instead of the primary oxidation products. Previous study elucidated that the fragmentation and further oxidation of five-membered oxygen-containing heterocyclic compounds, which are important intermediates in the oxidation of aromatic compounds, can yield acetic acid (Forstner et al., 1997; Bahreini et al., 2005; Walavalkar et al., 2017). However, this pathway is still controversial due to the lack of robust evidence. The major precursors and detailed mechanisms underlying the multigenerational oxidation that results in the high yields of formic acid and acetic acid during the photooxidation of toluene warrant further investigation.




Figure 6. Correlations of yields of carbonyl compounds (sum of glyoxal and methylglyoxal) with yields of gas-phase formic acid and acetic acid during the photooxidation of toluene at different $\bullet$ OH exposure (20% RH). Markers are color-coded to represent different $\bullet$ OH exposure levels.

4 ATMOSPHERIC IMPLICATIONS

This study reveals unexpectedly high yields of formic acid (up to 74%) and acetic acid (up to 40%) during the multi-generational photooxidation of toluene, achieved over equivalent atmospheric aging times of 1–3 days in the OFR experiments. The formation of these organic acids exhibited minimal sensitivity to NO_x concentrations and RH. The up-to-date MCM v3.3.1 model substantially underestimated the observed organic acid concentrations in the OFR.

Although both small gas-phase organic acids and low-volatility particulate matter increased with atmospheric aging (within a certain degree of oxidation), their differing production timescales suggest that photodegradation of SOA may not be the dominant formation pathway of organic acids. Combined with measurements of carbonyl compounds, we proposed that the observed organic acids predominantly arise from the multi-generational oxidation of toluene. A mechanistic understanding of this process, particularly the photooxidation of intermediate products such as maleic anhydride, requires further investigation (Lu et al., 2024; Wang et al., 2024).

Our findings also provide new insight into understanding organic acid budgets in urban atmosphere. Previous studies have observed rapid secondary production of organic acids in urban air masses (Veres et al., 2011; Friedman and Farmer, 2018), but model simulations have significantly underestimated their concentrations (Le Breton et al., 2012; Yuan et al., 2015). **If the high yields of organic acids observed during the multi-generational oxidation of toluene are broadly applicable to other aromatics, the elevated yields of formic and acetic acids at high •OH exposures reported in this study may help explain these discrepancies.** Liggio et al. (2017) found that the underestimation of organic acids increased with reaction time, while Yuan et al. (2015) reported that the peak concentrations of formic acid occurred 4–6 days after the peak benzene levels (Yuan et al., 2015; Liggio et al., 2017). These studies evaluated the contributions of aromatics to formic acid assuming a yield of 15%, concluding that aromatics can contribute about 11%–12% of the total formic acid production. If a yield of 74% is adopted, the contributions of aromatics would rise to as high as 55%–60%. This outcome may explain the remaining 50% of missing sources in the model after known sources are considered (Yuan et al., 2015; Liggio et al., 2017). The large amounts of small gas-phase organic acids produced from the photooxidation of aromatics in urban areas may further affect the aerosol pH and aerosol formation. **It is noted that biomass burning emissions also contain substantial amounts of aromatic compounds, and multi-generational oxidation within biomass burning plumes may similarly produce significant quantities of organic acids, as supported by both laboratory experiments and field observations (Bruns et al., 2017; Permar et al., 2023).** Additionally, while small organic acids such as formic and acetic acid have long been recognized as useful tracers for atmospheric VOC oxidation and SOA aging (Munger et al., 1986; Hansen et al., 2014), our results suggest that their utility as reliable tracers for SOA production is limited to a relatively narrow •OH exposure window (approximately 0.2–3 equivalent days). Beyond this range, their concentrations decouple from SOA mass due to differences in reactivity and removal pathways

(Isaacman-VanWertz et al., 2018).

Supporting information

The supplement related to this article is available online.

Author contributions

ZC, LH, and HS designed the research. HS performed research, analyzed the data, and drafted the initial manuscript. LH and ZC revised the manuscript. MZ and YY assisted with the model simulation. YZ, HL, and HW provided valuable comments and suggestions for the manuscript.

Competing interests

All authors declare they have no competing interests.

Data availability

The data used in this study is available upon request from the corresponding authors.

Acknowledgements

The authors gratefully thank the National Key Research and Development Program of China (2022YFC3701102) for financial support.

References

Archibald, A. T., McGillen, M. R., Taatjes, C. A., Percival, C. J., and Shallcross, D. E.: Atmospheric transformation of enols: A potential secondary source of carboxylic acids in the urban troposphere, *Geophys. Res. Lett.*, 34, 2007GL031032, 2007.

Bahreini, R., Keywood, M. D., Ng, N. L., Varutbangkul, V., Gao, S., Flagan, R. C., Seinfeld, J. H., Worsnop, D. R., and Jimenez, J. L.: Measurements of Secondary Organic Aerosol from Oxidation of Cycloalkenes, Terpenes, and *m*-Xylene Using an Aerodyne Aerosol Mass Spectrometer, *Environ. Sci. Technol.*, 39, 5674–5688, 2005.

Baltensperger, U., Kalberer, M., Dommen, J., Paulsen, D., Alfarra, M. R., Coe, H., Fisseha, R., Gascho, A., Gysel, M., Nyeki, S., Sax, M., Steinbacher, M., Prevot, A. S. H., Sjögren, S., Weingartner, E., and Zenobi, R.: Secondary organic aerosols from anthropogenic and biogenic precursors, *Faraday Discuss.*, 130, 265–278, 2005.

Bandow, H. and Washida, N.: Ring-cleavage Reactions of Aromatic Hydrocarbons Studied by FT–IR Spectroscopy. III. Photooxidation of 1,2,3-, 1,2,4-, and 1,3,5-Trimethylbenzenes in the NO_x–Air System, *Bull. Chem. Soc. Jpn.*, 58, 2549–2555, 1985.

Bandow, H., Washida, N., and Akimoto, H.: Ring-cleavage Reactions of Aromatic Hydrocarbons Studied by FT–IR Spectroscopy. I. Photooxidation of Toluene and Benzene in the NO_x–Air System, *Bull. Chem. Soc. Jpn.*, 58, 2531–2540, 1985.

487 Bannan, T. J., Murray Booth, A., Le Breton, M., Bacak, A., Muller, J. B. A., Leather, K. E.,
 488 Khan, M. A. H., Lee, J. D., Dunmore, R. E., Hopkins, J. R., Fleming, Z. L., Sheps, L., Taatjes,
 489 C. A., Shallcross, D. E., and Percival, C. J.: Seasonality of formic acid (HCOOH) in London
 490 during the ClearfLo campaign, *J. Geophys. Res. Atmospheres*, 122, 12488–12498, 2017.

491 Bates, K. H., Jacob, D. J., Cope, J. D., Chen, X., Millet, D. B., and Nguyen, T. B.: Emerging
 492 investigator series: aqueous oxidation of isoprene-derived organic aerosol species as a source
 493 of atmospheric formic and acetic acids, *Environ. Sci. Atmospheres*, 3, 1651–1664, 2023.

494 Battin-Leclerc, F., Konnov, A. A., Jaffrezo, J. L., and Legrand, M.: To Better Understand the
 495 Formation of Short-Chain Acids in Combustion Systems, *Combust. Sci. Technol.*, 180, 343–
 496 370, 2007.

497 Berndt, T. and Böge, O.: Gas-phase reaction of OH radicals with benzene: products and
 498 mechanism, *Phys. Chem. Chem. Phys.*, 3, 4946–4956, 2001.

499 Berndt, T., Böge, O., and Herrmann, H.: On the formation of benzene oxide/oxepin in the gas-
 500 phase reaction of OH radicals with benzene, *Chem. Phys. Lett.*, 314, 435–442, 1999.

501 Brune, W. H.: The Chamber Wall Index for Gas–Wall Interactions in Atmospheric
 502 Environmental Enclosures, *Environ. Sci. Technol.*, 53, 3645–3652, 2019.

503 Bruns, E. A., El Haddad, I., Keller, A., Klein, F., Kumar, N. K., Pieber, S. M., Corbin, J. C.,
 504 Slowik, J. G., Brune, W. H., Baltensperger, U., and Prévôt, A. S. H.: Inter-comparison of
 505 laboratory smog chamber and flow reactor systems on organic aerosol yield and composition,
 506 *Atmospheric Meas. Tech.*, 8, 2315–2332, 2015.

507 Bruns, E. A., Slowik, J. G., El Haddad, I., Kilic, D., Klein, F., Dommen, J., Temime-Roussel,
 508 B., Marchand, N., Baltensperger, U., and Prévôt, A. S. H.: Characterization of gas-phase
 509 organics using proton transfer reaction time-of-flight mass spectrometry: fresh and aged
 510 residential wood combustion emissions, *Atmospheric Chem. Phys.*, 17, 705–720, 2017.

511 Chaliyakunnel, S., Millet, D. B., Wells, K. C., Cady-Pereira, K. E., and Shephard, M. W.: A
 512 Large Underestimate of Formic Acid from Tropical Fires: Constraints from Space-Borne
 513 Measurements, *Environ. Sci. Technol.*, 50, 5631–5640, 2016.

514 Chebbi, A. and Carlier, P.: Carboxylic acids in the troposphere, occurrence, sources, and sinks:
 515 A review, *Atmos. Environ.*, 30, 4233–4249, 1996.

516 Crounse, J. D., Nielsen, L. B., Jørgensen, S., Kjaergaard, H. G., and Wennberg, P. O.:
 517 Autoxidation of Organic Compounds in the Atmosphere, *J. Phys. Chem. Lett.*, 4, 3513–3520,
 518 2013.

519 Ding, Z., Zhu, Y., Wang, G., Li, H., Xu, J., Tian, M., Liu, Z., Chen, J., Yun, L., Zheng, H., Gui,
 520 H., Liu, J., Li, R., Deng, C., and Huang, K.: Coastal eutrophic ecosystem as an overlooked pool

521 of atmospheric formic acid: disentangling biogenic and abiotic contributions, *Environ. Sci.*
 522 *Technol. Lett.*, 2025.

523 Drozd, G. T., Zhao, Y., Saliba, G., Frodin, B., Maddox, C., Oliver Chang, M.-C., Maldonado,
 524 H., Sardar, S., Weber, R. J., Robinson, A. L., and Goldstein, A. H.: Detailed Speciation of
 525 Intermediate Volatility and Semivolatile Organic Compound Emissions from Gasoline Vehicles:
 526 Effects of Cold-Starts and Implications for Secondary Organic Aerosol Formation, *Environ. Sci.*
 527 *Technol.*, 53, 1706–1714, 2019.

528 Ervens, B., Turpin, B. J., and Weber, R. J.: Secondary organic aerosol formation in cloud
 529 droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies,
 530 *Atmospheric Chem. Phys.*, 11, 11069–11102, 2011.

531 Fittschen, C., Whalley, L. K., and Heard, D. E.: The reaction of CH_3O_2 radicals with OH radicals:
 532 a neglected sink for CH_3O_2 in the remote atmosphere, *Environ. Sci. Technol.*, 48, 7700–7701,
 533 2014.

534 Forstner, H. J. L., Flagan, R. C., and Seinfeld, J. H.: Secondary Organic Aerosol from the
 535 Photooxidation of Aromatic Hydrocarbons: Molecular Composition, *Environ. Sci. Technol.*, 31,
 536 1345–1358, 1997.

537 Franco, B., Blumenstock, T., Cho, C., Clarisse, L., Clerbaux, C., Coheur, P.-F., De Mazière, M.,
 538 De Smedt, I., Dorn, H.-P., Emmerichs, T., Fuchs, H., Gkatzelis, G., Griffith, D. W. T., Gromov,
 539 S., Hannigan, J. W., Hase, F., Hohaus, T., Jones, N., Kerkweg, A., Kiendler-Scharr, A., Lutsch,
 540 E., Mahieu, E., Novelli, A., Ortega, I., Paton-Walsh, C., Pommier, M., Pozzer, A., Reimer, D.,
 541 Rosanka, S., Sander, R., Schneider, M., Strong, K., Tillmann, R., Van Roozendael, M.,
 542 Vereecken, L., Vigouroux, C., Wahner, A., and Taraborrelli, D.: Ubiquitous atmospheric
 543 production of organic acids mediated by cloud droplets, *Nature*, 593, 233–237, 2021.

544 Friedman, B. and Farmer, D. K.: SOA and gas phase organic acid yields from the sequential
 545 photooxidation of seven monoterpenes, *Atmos. Environ.*, 187, 335–345, 2018.

546 Friedman, B., Link, M. F., Fulgham, S. R., Brophy, P., Galang, A., Brune, W. H., Jathar, S. H.,
 547 and Farmer, D. K.: Primary and Secondary Sources of Gas-Phase Organic Acids from Diesel
 548 Exhaust, *Environ. Sci. Technol.*, 51, 10872–10880, 2017.

549 Gery, M. W., Fox, D. L., Jeffries, H. E., Stockburger, L., and Weathers, W. S.: A continuous
 550 stirred tank reactor investigation of the gas-phase reaction of hydroxyl radicals and toluene, *Int.*
 551 *J. Chem. Kinet.*, 17, 931–955, 1985.

552 Hansen, A. M. K., Kristensen, K., Nguyen, Q. T., Zare, A., Cozzi, F., Nøjgaard, J. K., Skov, H.,
 553 Brandt, J., Christensen, J. H., Ström, J., Tunved, P., Krejci, R., and Glasius, M.: Organosulfates
 554 and organic acids in Arctic aerosols: speciation, annual variation and concentration levels,
 555 *Atmospheric Chem. Phys.*, 14, 7807–7823, 2014.

556 Huang, D., Chen, Z. M., Zhao, Y., and Liang, H.: Newly observed peroxides and the water
 557 effect on the formation and removal of hydroxyalkyl hydroperoxides in the ozonolysis of
 558 isoprene, *Atmospheric Chem. Phys.*, 13, 5671–5683, 2013.

559 Isaacman-VanWertz, G., Massoli, P., O’Brien, R., Lim, C., Franklin, J. P., Moss, J. A., Hunter,
 560 J. F., Nowak, J. B., Canagaratna, M. R., Misztal, P. K., Arata, C., Roscioli, J. R., Herndon, S.
 561 T., Onasch, T. B., Lambe, A. T., Jayne, J. T., Su, L., Knopf, D. A., Goldstein, A. H., Worsnop,
 562 D. R., and Kroll, J. H.: Chemical evolution of atmospheric organic carbon over multiple
 563 generations of oxidation, *Nat. Chem.*, 10, 462–468, 2018.

564 Jenkin, M. E., Hurley, M. D., and Wallington, T. J.: Investigation of the radical product channel
 565 of the $\text{CH}_3\text{C}(\text{O})\text{CH}_2\text{O}_2 + \text{HO}_2$ reaction in the gas phase, *Phys. Chem. Chem. Phys.*, 10, 4274–
 566 4280, 2008.

567 Jenkin, M. E., Young, J. C., and Rickard, A. R.: The MCM v3.3.1 degradation scheme for
 568 isoprene, *Atmospheric Chem. Phys.*, 15, 11433–11459, 2015.

569 Jiang, Y., Xia, M., Wang, Z., Zheng, P., Chen, Y., and Wang, T.: Photochemical ageing of
 570 aerosols contributes significantly to the production of atmospheric formic acid, *Atmospheric*
 571 *Chem. Phys.*, 23, 14813–14828, 2023.

572 Kang, M., Zhang, H., and Ying, Q.: Sulfate or organic acids: the atmospheric fate of stabilized
 573 criegee intermediates, *J. Geophys. Res. Atmospheres*, 130, e2025JD044867, 2025.

574 Keller-Rudek, H., Moortgat, G. K., Sander, R., and Sörensen, R.: The MPI-Mainz UV/VIS
 575 Spectral Atlas of Gaseous Molecules of Atmospheric Interest, *Earth Syst. Sci. Data*, 5, 365–373,
 576 2013.

577 Khan, M. A. H., Lyons, K., Chhantyal-Pun, R., McGillen, M. R., Caravan, R. L., Taatjes, C. A.,
 578 Orr-Ewing, A. J., Percival, C. J., and Shallcross, D. E.: Investigating the tropospheric chemistry
 579 of acetic acid using the global 3-D chemistry transport model, STOCHEM-CRI, *J. Geophys.*
 580 *Res. Atmospheres*, 123, 6267–6281, 2018.

581 Lambe, A., Massoli, P., Zhang, X., Canagaratna, M., Nowak, J., Daube, C., Yan, C., Nie, W.,
 582 Onasch, T., Jayne, J., Kolb, C., Davidovits, P., Worsnop, D., and Brune, W.: Controlled nitric
 583 oxide production via $\text{O}(^1\text{D}) + \text{N}_2\text{O}$ reactions for use in oxidation flow reactor studies,
 584 *Atmospheric Meas. Tech.*, 10, 2283–2298, 2017.

585 Lambe, A. T., Onasch, T. B., Croasdale, D. R., Wright, J. P., Martin, A. T., Franklin, J. P.,
 586 Massoli, P., Kroll, J. H., Canagaratna, M. R., Brune, W. H., Worsnop, D. R., and Davidovits, P.:
 587 Transitions from Functionalization to Fragmentation Reactions of Laboratory Secondary
 588 Organic Aerosol (SOA) Generated from the OH Oxidation of Alkane Precursors, *Environ. Sci.*
 589 *Technol.*, 46, 5430–5437, 2012.

590 Lambe, A. T., Cappa, C. D., Massoli, P., Onasch, T. B., Forestieri, S. D., Martin, A. T.,

591 Cummings, M. J., Croasdale, D. R., Brune, W. H., Worsnop, D. R., and Davidovits, P.:
 592 Relationship between Oxidation Level and Optical Properties of Secondary Organic Aerosol,
 593 Environ. Sci. Technol., 47, 6349–6357, 2013.

594 Lambe, A. T., Chhabra, P. S., Onasch, T. B., Brune, W. H., Hunter, J. F., Kroll, J. H., Cummings,
 595 M. J., Brogan, J. F., Parmar, Y., Worsnop, D. R., Kolb, C. E., and Davidovits, P.: Effect of
 596 oxidant concentration, exposure time, and seed particles on secondary organic aerosol chemical
 597 composition and yield, Atmospheric Chem. Phys., 15, 3063–3075, 2015.

598 Le Breton, M., McGillen, M. R., Muller, J. B. A., Bacak, A., Shallcross, D. E., Xiao, P., Huey,
 599 L. G., Tanner, D., Coe, H., and Percival, C. J.: Airborne observations of formic acid using a
 600 chemical ionization mass spectrometer, Atmospheric Meas. Tech., 5, 3029–3039, 2012.

601 Li, R., Palm, B. B., Ortega, A. M., Hlywiak, J., Hu, W., Peng, Z., Day, D. A., Knote, C., Brune,
 602 W. H., De Gouw, J. A., and Jimenez, J. L.: Modeling the Radical Chemistry in an Oxidation
 603 Flow Reactor: Radical Formation and Recycling, Sensitivities, and the OH Exposure
 604 Estimation Equation, J. Phys. Chem. A, 119, 4418–4432, 2015.

605 Liggio, J., Moussa, S. G., Wentzell, J., Darlington, A., Liu, P., Leithead, A., Hayden, K.,
 606 O'Brien, J., Mittermeier, R. L., Staebler, R., Wolde, M., and Li, S.-M.: Understanding the
 607 primary emissions and secondary formation of gaseous organic acids in the oil sands region of
 608 Alberta, Canada, Atmospheric Chem. Phys., 17, 8411–8427, 2017.

609 Link, M. F., Nguyen, T. B., Bates, K., Müller, J.-F., and Farmer, D. K.: Can Isoprene Oxidation
 610 Explain High Concentrations of Atmospheric Formic and Acetic Acid over Forests?, ACS Earth
 611 Space Chem., 4, 730–740, 2020.

612 Lu, R., Zhou, P., Ma, F., Zhao, Q., Peng, X., Chen, J., and Xie, H.-B.: Multi-generation
 613 oxidation mechanism of *m*-xylene: unexpected implications for secondary organic aerosol
 614 formation, Atmos. Environ., 327, 120511, 2024.

615 Malecha, K. T. and Nizkorodov, S. A.: Photodegradation of Secondary Organic Aerosol
 616 Particles as a Source of Small, Oxygenated Volatile Organic Compounds, Environ. Sci.
 617 Technol., 50, 9990–9997, 2016.

618 Malecha, K. T. and Nizkorodov, S. A.: Feasibility of photosensitized reactions with secondary
 619 organic aerosol particles in the presence of volatile organic compounds, J. Phys. Chem. A, 121,
 620 4961–4967, 2017.

621 Millet, D. B., Baasandorj, M., Farmer, D. K., Thornton, J. A., Baumann, K., Brophy, P.,
 622 Chaliyakunnel, S., De Gouw, J. A., Graus, M., Hu, L., Koss, A., Lee, B. H., Lopez-Hilfiker, F.
 623 D., Neuman, J. A., Paulot, F., Peischl, J., Pollack, I. B., Ryerson, T. B., Warneke, C., Williams,
 624 B. J., and Xu, J.: A large and ubiquitous source of atmospheric formic acid, Atmospheric Chem.
 625 Phys., 15, 6283–6304, 2015.

626 Müller, M., Graus, M., Wisthaler, A., Hansel, A., Metzger, A., Dommen, J., and Baltensperger,
 627 U.: Analysis of high mass resolution PTR-TOF mass spectra from 1,3,5-trimethylbenzene
 628 (TMB) environmental chamber experiments, *Atmospheric Chem. Phys.*, 12, 829–843, 2012.

629 Mungall, E. L., Abbatt, J. P. D., Wentzell, J. J. B., Wentworth, G. R., Murphy, J. G., Kunkel, D.,
 630 Gute, E., Tarasick, D. W., Sharma, S., Cox, C. J., Uttal, T., and Liggio, J.: High gas-phase
 631 mixing ratios of formic and acetic acid in the High Arctic, *Atmospheric Chem. Phys.*, 18,
 632 10237–10254, 2018.

633 Munger, J. W., Tillier, C., and Hoffmann, M. R.: Identification of hydroxymethanesulfonate in
 634 fog water, *Science*, 231, 247–249, 1986.

635 Murschell, T. and Farmer, D. K.: Atmospheric OH oxidation of three chlorinated aromatic
 636 herbicides, *Environ. Sci. Technol.*, 52, 4583–4591, 2018.

637 Neeb, P., Sauer, F., Horie, O., and Moortgat, G. K.: Formation of hydroxymethyl hydroperoxide
 638 and formic acid in alkene ozonolysis in the presence of water vapour, *Atmos. Environ.*, 31,
 639 1417–1423, 1997.

640 Ng, N. L., Kroll, J. H., Chan, A. W. H., Chhabra, P. S., Flagan, R. C., and Seinfeld, J. H.:
 641 Secondary organic aerosol formation from m-xylene, toluene, and benzene, *Atmospheric Chem.*
 642 *Phys.*, 7, 3909–3922, 2007.

643 Ortega, A. M., Day, D. A., Cubison, M. J., Brune, W. H., Bon, D., De Gouw, J. A., and Jimenez,
 644 J. L.: Secondary organic aerosol formation and primary organic aerosol oxidation from
 645 biomass-burning smoke in a flow reactor during FLAME-3, *Atmospheric Chem. Phys.*, 13,
 646 11551–11571, 2013.

647 Orzechowska, G. E. and Paulson, S. E.: Photochemical sources of organic acids. 1. Reaction of
 648 ozone with isoprene, propene, and 2-butenes under dry and humid conditions using SPME, *J.*
 649 *Phys. Chem. A*, 109, 5358–5365, 2005.

650 Paulot, F., Wunch, D., Crounse, J. D., Toon, G. C., Millet, D. B., DeCarlo, P. F., Vigouroux, C.,
 651 Deutscher, N. M., González Abad, G., Notholt, J., Warneke, T., Hannigan, J. W., Warneke, C.,
 652 De Gouw, J. A., Dunlea, E. J., De Mazière, M., Griffith, D. W. T., Bernath, P., Jimenez, J. L.,
 653 and Wennberg, P. O.: Importance of secondary sources in the atmospheric budgets of formic
 654 and acetic acids, *Atmospheric Chem. Phys.*, 11, 1989–2013, 2011.

655 Peng, Z., Day, D. A., Stark, H., Li, R., Lee-Taylor, J., Palm, B. B., Brune, W. H., and Jimenez,
 656 J. L.: HOx radical chemistry in oxidation flow reactors with low-pressure mercury lamps
 657 systematically examined by modeling, *Atmospheric Meas. Tech.*, 8, 4863–4890, 2015.

658 Peng, Z., Day, D. A., Ortega, A. M., Palm, B. B., Hu, W., Stark, H., Li, R., Tsigaridis, K., Brune,
 659 W. H., and Jimenez, J. L.: Non-OH chemistry in oxidation flow reactors for the study of
 660 atmospheric chemistry systematically examined by modeling, *Atmospheric Chem. Phys.*, 16,

661 4283–4305, 2016.

662 Permar, W., Jin, L., Peng, Q., O’Dell, K., Lill, E., Selimovic, V., J. Yokelson, R., S. Hornbrook,
 663 R., J. Hills, A., C. Apel, E., Ku, I.-T., Zhou, Y., C. Sive, B., P. Sullivan, A., L. Collett, J.,
 664 B. Palm, B., A. Thornton, J., Flocke, F., V. Fischer, E., and Hu, L.: Atmospheric OH reactivity
 665 in the western United States determined from comprehensive gas-phase measurements during
 666 WE-CAN, *Environ. Sci. Atmospheres*, 3, 97–114, 2023.

667 Praplan, A. P., Hegyi-Gaeggeler, K., Barnet, P., Pfaffenberger, L., Dommen, J., and
 668 Baltensperger, U.: Online measurements of water-soluble organic acids in the gas and aerosol
 669 phase from the photooxidation of 1,3,5-trimethylbenzene, *Atmospheric Chem. Phys.*, 14, 8665–
 670 8677, 2014.

671 Schobesberger, S., Lopez-Hilfiker, F. D., Taipale, D., Millet, D. B., D’Ambro, E. L., Rantala,
 672 P., Mammarella, I., Zhou, P., Wolfe, G. M., Lee, B. H., Boy, M., and Thornton, J. A.: High
 673 upward fluxes of formic acid from a boreal forest canopy, *Geophys. Res. Lett.*, 43, 9342–9351,
 674 2016.

675 Seuwen, R. and Warneck, P.: Oxidation of toluene in NO_x free air: product distribution and
 676 mechanism, *Int. J. Chem. Kinet.*, 28, 315–332, 1996.

677 Shaw, M. F., Sztáray, B., Whalley, L. K., Heard, D. E., Millet, D. B., Jordan, M. J. T., Osborn,
 678 D. L., and Kable, S. H.: Photo-tautomerization of acetaldehyde as a photochemical source of
 679 formic acid in the troposphere, *Nat. Commun.*, 9, 2584, 2018.

680 Shen, H., Chen, Z., Li, H., Qian, X., Qin, X., and Shi, W.: Gas-particle partitioning of carbonyl
 681 compounds in the ambient atmosphere, *Environ. Sci. Technol.*, 52, 10997–11006, 2018.

682 Shen, H., Huang, L., Qian, X., Qin, X., and Chen, Z.: Positive Feedback between Partitioning
 683 of Carbonyl Compounds and Particulate Sulfur Formation during Haze Episodes, *Environ. Sci.*
 684 *Technol.*, 58, 21286–21294, 2024.

685 Stavrakou, T., Müller, J.-F., Peeters, J., Razavi, A., Clarisse, L., Clerbaux, C., Coheur, P.-F.,
 686 Hurtmans, D., De Mazière, M., Vigouroux, C., Deutscher, N. M., Griffith, D. W. T., Jones, N.,
 687 and Paton-Walsh, C.: Satellite evidence for a large source of formic acid from boreal and
 688 tropical forests, *Nat. Geosci.*, 5, 26–30, 2012.

689 Tan, Z., Rohrer, F., Lu, K., Ma, X., Bohn, B., Broch, S., Dong, H., Fuchs, H., Gkatzelis, G. I.,
 690 Hofzumahaus, A., Holland, F., Li, X., Liu, Y., Liu, Y., Novelli, A., Shao, M., Wang, H., Wu, Y.,
 691 Zeng, L., Hu, M., Kiendler-Scharr, A., Wahner, A., and Zhang, Y.: Wintertime photochemistry
 692 in Beijing: observations of RO_x radical concentrations in the North China Plain during the
 693 BEST-ONE campaign, *Atmospheric Chem. Phys.*, 18, 12391–12411, 2018.

694 Tao, Y. and Murphy, J. G.: Evidence for the Importance of Semivolatile Organic Ammonium
 695 Salts in Ambient Particulate Matter, *Environ. Sci. Technol.*, 53, 108–116, 2019.

696 Veres, P. R., Roberts, J. M., Cochran, A. K., Gilman, J. B., Kuster, W. C., Holloway, J. S., Graus,
697 M., Flynn, J., Lefer, B., Warneke, C., and de Gouw, J.: Evidence of rapid production of organic
698 acids in an urban air mass, *Geophys. Res. Lett.*, 38, 2011.

699 Volkamer, R., Platt, U., and Wirtz, K.: Primary and Secondary Glyoxal Formation from
700 Aromatics: Experimental Evidence for the Bicycloalkyl–Radical Pathway from Benzene,
701 Toluene, and *p*-Xylene, *J. Phys. Chem. A*, 105, 7865–7874, 2001.

702 Walavalkar, M. P., Sharma, A., Dhanya, S., and Naik, P. D.: Reactions of lactones with
703 tropospheric oxidants: A kinetics and products study, *Atmos. Environ.*, 161, 18–26, 2017.

704 Wang, Y., Li, C., Zhang, Y., Li, Y., Yang, G., Yang, X., Wu, Y., Yao, L., Zhang, H., and Wang,
705 L.: Secondary reactions of aromatics-derived oxygenated organic molecules lead to plentiful
706 highly oxygenated organic molecules within an intraday OH exposure, *Atmospheric Chem.*
707 *Phys.*, 24, 7961–7981, 2024.

708 Wu, H., Wang, Y., Li, H., Huang, L., Huang, D., Shen, H., Xing, Y., and Chen, Z.: The OH-
709 initiated oxidation of atmospheric peroxyacetic acid: Experimental and model studies, *Atmos.*
710 *Environ.*, 164, 61–70, 2017.

711 Wyche, K. P., Monks, P. S., Ellis, A. M., Cordell, R. L., Parker, A. E., Whyte, C., Metzger, A.,
712 Dommen, J., Duplissy, J., Prevot, A. S. H., Baltensperger, U., Rickard, A. R., and Wulfert, F.:
713 Gas phase precursors to anthropogenic secondary organic aerosol: detailed observations of
714 1,3,5-trimethylbenzene photooxidation, *Atmospheric Chem. Phys.*, 9, 635–665, 2009.

715 Xing, Y., Li, H., Huang, L., Wu, H., Shen, H., and Chen, Z.: The production of formaldehyde
716 and hydroxyacetone in methacrolein photooxidation: New insights into mechanism and effects
717 of water vapor, *J. Environ. Sci.*, 66, 1–11, 2018.

718 Yetter, R. A., Rabitz, H., Dryer, F. L., Maki, R. G., and Klemm, R. B.: Evaluation of the rate
719 constant for the reaction $\text{OH} + \text{H}_2\text{CO}$: Application of modeling and sensitivity analysis
720 techniques for determination of the product branching ratio, *J. Chem. Phys.*, 91, 4088–4097,
721 1989.

722 Yuan, B., Veres, P. R., Warneke, C., Roberts, J. M., Gilman, J. B., Koss, A., Edwards, P. M.,
723 Graus, M., Kuster, W. C., Li, S.-M., Wild, R. J., Brown, S. S., Dubé, W. P., Lerner, B. M.,
724 Williams, E. J., Johnson, J. E., Quinn, P. K., Bates, T. S., Lefer, B., Hayes, P. L., Jimenez, J. L.,
725 Weber, R. J., Zamora, R., Ervens, B., Millet, D. B., Rappenglück, B., and De Gouw, J. A.:
726 Investigation of secondary formation of formic acid: urban environment vs. oil and gas
727 producing region, *Atmospheric Chem. Phys.*, 15, 1975–1993, 2015.

728 Zervas, E.: Formation of organic acids from propane, isooctane and toluene/isooctane flames,
729 *Fuel*, 84, 691–700, 2005.

730 Zhang, X., Cappa, C. D., Jathar, S. H., McVay, R. C., Ensberg, J. J., Kleeman, M. J., and

731 Seinfeld, J. H.: Influence of vapor wall loss in laboratory chambers on yields of secondary
732 organic aerosol, *Proc. Natl. Acad. Sci.*, 111, 5802–5807, 2014.

733