Improved subglacial boundary conditions near Dome C, Antarctica, through rigorous integration of multi-campaign radar data and an ensemble stochastic simulation approach

Calvin Shackleton[1*], *Kenichi Matsuoka*[1], *Brett Fraysher*[2], *Joohan Lee*[3], *Jørgen Dall*[4], *Allain Rapadas*[2], *Prasad Gogineni*[2], *Drew Taylor*[2], *Hyeon Tae Ju*[3], *Changhyun Chung*[3], *Anders Kusk*[4]

[*]*calvin.shackleton@npolar.no*
[1]Norwegian Polar Institute, Fram Senteret, Tromsø, Norway
[2]University of Alabama, Tuscaloosa, Alabama, USA
[3]Korea Polar Research Institute, Incheon, Korea
[4]DTU Space, Technical University of Denmark, Lyngby, Denmark

Abstract

Subglacial bed topography is a fundamental boundary condition for ice-sheet flow, basal thermal regimes, and the preservation of palaeoclimate records, yet it remains highly uncertain across Antarctica. We use stochastic simulation methods to generate an ensemble of gridded (500 m) ice thickness and bed elevation near Dome C, and map bed elevation uncertainty accounting for impacts of data availability and basal roughness. New airborne radar measurements uncover previously undocumented sectors of the ice-bed interface and we integrate these with existing data using a rigorous approach to ensure nearby measurement compatibility. Our simulated bed shows improved representation of basal roughness between survey profiles, revealing regional roughness over 50% higher than previous interpolations. We apply the ensemble to investigate uncertainty in meltwater routing, identifying well-constrained drainage away from the Dome C Plateau but poorly constrained outflow from Concordia Lake and parts of the Belgica Highlands. Topographic focussing of geothermal heat ~~predicts~~ could lead to local anomalies up to ±30% relative to the regional value, and several ice core drill sites are in a zone ~~of~~ with potentially 12% locally-reduced geothermal heating which ~~is~~ could be favourable for deep ice layer preservation. This work demonstrates that the critical subglacial bed boundary condition is better represented by ensemble-based, stochastic simulations based on strictly cross-validated data rather than single deterministic interpolations, and simulated beds can improve the reliability of analyses with robust consideration of uncertainties in data-sparse regions.

1. Introduction

Kilometer-scale subglacial topography influences basal thermal state through locally variable pressure-melting (Cuffey & Paterson, 2010) and also modulates the effective area for geothermal heat conduction (van der Veen et al., 2007). While basal roughness can be constrained in high resolution in the along-track orientation of radar profiles, this information is missing between survey profiles. Widely-used interpolation methods such as kriging and splines do not preserve the spatial statistics of measurements in data gaps, leading to over-smoothing in gridded products and unrealistic patterns in basal roughness. Compared with single-grid, elevation-preserving products such as Bedmap3 (Pritchard et al., 2025) and Bedmachine v4 (Morlighem, 2026), texture-preserving bed gridding approaches can provide more realistic boundary conditions for model projections (McCormack et al., 2026). While requiring an ensemble-based framework which increases computational cost and methodological complexity, this approach generates spatially explicit uncertainty estimates to propagate into models and can be used to prioritise sites for future geophysical surveys. Sequential Gaussian Simulation (SGS) is one texture-preserving geostatistical method that simulates values between measurements based on the values and statistical properties of nearby data (Deutsch & Journel, 1997). This approach has been applied to generate ensembles of equally-likely bed elevation grids to investigate the basal environment, subglacial meltwater drainage, and geothermal heat flow (GHF) with robust consideration of the impacts of elevation uncertainty and realistic basal roughness (MacKie et al., 2020; MacKie et al., 2021; Shackleton et al., 2023).

Subglacial topography around Dome C is of particular interest in the context of paleoclimate investigations. Oxygen isotope records from marine sediment cores indicate that global glacial-interglacial cycles experienced a major shift in periodicity from 41 Ka to ca. 100 Ka between 1.4 Ma and 800 Ka BP (Lisiecki & Raymo, 2005). The underlying mechanisms for this Mid-Pleistocene Transition (MPT) are not yet fully constrained (Berends et al., 2021), but deep ice cores that preserve atmospheric gases from the palaeo-climate could provide continuous, high-resolution climate records. Ice layers that span the last 1.5 Ma could be located near the base of thick ice domes in the interior of Antarctica, though basal melting, ice flow, and accretion should be minimal to preserve ice layer stratigraphy (Fischer et al., 2013; Van Liefferinge & Pattyn, 2013). To locate basal ice with undisturbed stratigraphy and sufficient age resolution spanning the MPT, the ice core community ~~are~~ is trying to delineate sites with: 1) ice around 2500 m thick, sufficient to resolve the MPT without inducing significant basal-melting; 2) low surface accumulation to limit vertical strain-induced thinning of ice layers; 3) low basal roughness to prevent disruption of basal ice stratigraphy through lateral strain; 4) low GHF to restrict basal melting and preserve the oldest ice layers. The European Project for Ice ~~coring~~ Coring in Antarctica (EPICA) successfully retrieved a 3260 m long ice core near Dome C, Antarctica that provides climate records for the last 800 Ka (Jouzel et al., 2007; Lüthi et al., 2008).

Ice thickness, basal roughness, and GHF are critical boundary conditions for thermomechanical modelling which delineate promising drill sites in the Dome C region that could resolve the MPT (Chung et al., 2023b; Parrenin et al., 2017; Passalacqua et al., 2017; Van Liefferinge et al., 2018; Wang et al., 2023; Young et al., 2017). Subglacial topography and ice thickness variability near Dome C (Fig. 1a-c) is documented in radar surveys (Drewry et al., 1982; Forieri et al., 2004; Tabacco et al., 1998), with thinner ice (<2.5 km) observed over the Belgica Highlands and Concordia Ridge, and 2.5-3.5 km thick ice over the Dome C plateau. Ice thicknesses over 4 km are located above Concordia and Vincennes subglacial lakes and in the Concordia Trench, with up to 4.7 km reported in the Aurora Subglacial Basin. The Concordia Trench and Adventure Trench are tectonic features 20 – 50 km in width with over 1 km thicker ice than the surroundings, oriented approximately northeast - southwest (Cianfarra et al., 2009). Spatial coverage of surveys varies from the well-resolved topography over the Dome C P~~p~~lateau and surrounding Concordia Station, to large data gaps over Vincennes Basin and southwest Belgica Highlands (Fig.1).

A large database of Antarctic radar-derived ice thickness measurements from surveys conducted between 1966 and 2022 was compiled recently by the Bedmap project (Frémand et al., 2023), and subsequent radar campaigns provide new measurements across key sites in the Dome C region to improve characterization of the basal environment (Chung et al., 2023a; Mulvaney et al., 2023). In this work, we integrate existing radar-derived ice thicknesses reported to the Bedmap database with subsequently released data as well as new radar survey measurements over the Belgica Highlands conducted by the Korea Polar Research Institute (KOPRI) and University of Alabama (UA) as well as over the Dome C Plateau by the Technical University of Denmark (DTU). This data integration was carried out using a robust filtering approach to minimize locally conflicting measurements in the compiled data. We use measurements in a 400 x 400 km^2 region surrounding Dome C (Fig. 1a) to simulate an ensemble of 100 bed elevation and ice thickness grids using SGS. The results for simulation #001 out of the ensemble are shown in Figure 1b and 1c as an illustrative example. By calculating variability over the ensemble we also map out regional topographic uncertainty (Fig. 1d). The simulated bed topography is used to investigate basal roughness, subglacial meltwater drainage, local topographic modification of GHF, and we discuss the implications of findings for the preservation of deep ice layers.




Figure 1: *Region map, radar survey data, and simulation results.* ***a)*** Radar-derived ice thickness measurements in the Dome C region. ***b)*** *Simulated bed elevation #001 from the ensemble of 100, with Concordia Station (green triangle) and ice cores plotted (red square).* ***c)*** *Simulated ice thickness #001 from the ensemble of 100, with subglacial lakes (orange circle) inferred from compiled radio-echo sounding (RES) analyses (Livingstone et al., 2022).* ***d)*** *Median Absolute Deviation (MAD) over the ensemble of 100 simulated bed elevation grids. All maps are projected to EPSG:3031. Histograms alongside colour bars show the distribution of values.*

2. Data compilation, quality control, and methodology

2.1 Radar-derived ice thickness data

We compiled available radar measurements of two-way travel time (TWT) from the ice surface to ice-bed interface from surveys conducted near Dome C, totalling ca. 2.3 million data points (Figs. 1a; A1-18). For all compiled TWT data, ice thicknesses were calculated assuming a radar wave propagation speed in ice of 169 m μs^{-1} and no firn corrections. The data compilation included new airborne surveys by a 2023/2024 Korea Polar Research Institute (KOPRI) and University of Alabama (UA) collaboration as well as a 2024/2025 survey by The Technical University of Denmark (DTU), 2016-2018 and 2019-2020 ground surveys from the Beyond EPICA - Oldest Ice (BE-OI) project, and ice thickness measurements with multiple sources from the Bedmap3 database:

2.1.1 KOPRI/UA airborne survey

A new 2023-2024 airborne survey from a KOPRI-UA collaboration (Fig. A12) provided 960 km lines of georeferenced radar data used to pick ice-bed TWT. These data were collected with a multi-channel ultra-wideband (UWB) ice-penetrating radar designed to operate over a frequency range of 170-470 MHz. The system consists of 8 transmit channels and 8 receive channels. Each transmit channel is supplied with chirped pulses of about 200 W of peak power. The radar is operated with a short pulse of 1-3 µs duration to sound shallow ice and map internal layers to a depth of 1000 m, and a long pulse of 5-10 us duration to sound thicker ice and map deeper layers. The system consists of a 16-element transmit array mounted under the left wing and a 16-element receive array mounted under the right wing. Two transmit and two receive channels are combined using a power divider/combiner to operate with a single transmit and receive channel. A more detailed system description is available in Awasthi et al. (2022) and Rizvi et al. (2025).

During this airborne campaign in 2023, we could not operate the system with its full bandwidth and power. One of the two inverters supplying power to the radar system from the aircraft failed. This led us to limit the number of channels, the operating frequency range (170-230 MHz), and the pulse width, so the radar could operate with only one inverter. Despite this constraint, the system maintained sufficient power and capability to recover bed echoes and internal stratigraphy across the region, demonstrating its ability to reconfigure to meet field and logistical constraints.

Data processing consisted of six major steps: 1) reduction of any coherent noise caused by the system or external sources; 2) pulse compression to improve range resolution and improve signal-to-noise ratio (Skolnik, 2008); 3) motion compensation to reduce errors introduced by aircraft flight path deviations; 4) synthetic aperture processing using a frequency-wavenumber (f-k) migration algorithm (Stolt, 1978) to improve both along-track resolution and signal-to-noise ratio; 5) correction of amplitude and phase differences between channels, and 6) combination of data from all channels and reduction of speckle using a reduction filter to improve image interpretation (Lee et al., 1994). Steps 1-4 were applied to each channel of received data, and processed data are then equalized and combined, with the additional image processing applied to the final image.

2.1.2 DTU airborne survey

A new 2024 airborne survey by DTU (Fig. A6) using the POLarimetric Airborne Radar Ice Sounder (POLARIS) system (Dall et al., 2010), provided ca. 1 million latitude, longitude, ice surface elevation, and TWT data points that spanned ca. 772 line km of survey profiles (Kusk, 2026). Commissioned by ESA, DTU developed the POLARIS radar, which can be configured in-flight ~~configured~~ for acquisition of either sounder data or synthetic aperture radar (SAR) data (Dall et al., 2013) at a center frequency of 435 MHz (P-band). With a maximum bandwidth of 85 MHz, a vertical resolution down to about 1 m in ice can be

achieved, but in order to improve the sensitivity to vertically distributed reflectors (e.g. internal layers), the Dome C data were acquired in a shallow/deep sounding mode, where a reduced bandwidth (30 MHz), a larger pulse length, and a higher receiver gain was used for the lower of two overlapping depth windows. The two depth windows were covered in a pulse-interleaved fashion, and the shorter pulse length that was used for shallow sounding enabled a lower flight altitude, hence less surface clutter. The POLARIS configuration prioritized polarimetry over coherent clutter suppression (Nielsen and Dall, 2015).

GNSS data and IMU data were acquired with an iMAR iNAT-M200-SLN navigation unit, PPP-processed with Waypoint Inertial Explorer, and used for motion compensation and geocoding. Narrowband RFI was masked out, and wideband RFI was suppressed by replacing the impacted pulses with interpolated pulses computed from highly correlated neighbouring pulses. Upon range compression and along-track focusing with a time-domain backprojection algorithm (Kusk and Dall, 2010), the POLARIS data were calibrated using data from an internal calibration loop and external calibration adjustments derived from POLARIS SAR data acquired over a trihedral radar reflector deployed near Casey Station. In this way both intensity and time delay was calibrated. Speckle and a potential birefringence loss was reduced by computing the span of the polarimetric covariance matrix and multi-looking by a factor of 5 in the along-track direction. The ice surface and bed were traced in a semi-automatic manner with an algorithm finding the depth where the leading edge of the return waveform has the steepest intensity slope.

2.1.3 BE-OI ground surveys

Ice thickness measurements from two ground surveys: a 2019-2020 survey near Little Dome C (LDC) consisting of 12 profiles over an area of 5x8 km^2 using a very high frequency (VHF) multi-channel coherent radar depth sounder (Chung et al., 2023a); a 2016-2018 survey using the Deep Looking radio echo sounder (DELORES) from British Antarctic Survey (BAS) consisting of 116 radar profiles near LDC and 21 profiles at North Patch covering an area of 5x5 km^2 (Mulvaney et al., 2023).

2.1.4 Bedmap Database

Ice thickness data from the Bedmap Database comprise radar-derived ice thickness measurements from many surveys (*Table A1*) between 1966 and 2018 (Frémand et al., 2023) and provided 1.9 million data points in the Dome C region. If not provided in the data files we calculated TWT from ice surface to the ice-bed interface using wave propagation speed and firn corrections described in the survey source reference.

2.2 Measurement consistency and data filtering

Nearby radar-derived ice thickness measurements can be inconsistent due to a combination of factors, including off-nadir reflections from steep or rough bed topography and signal attenuation or scattering in warm or debris-rich ice. To robustly calculate and model spatial variability in ice thickness any inconsistencies between nearby measurements need to be minimized; large differences in ice thickness could lead to overestimated variance. However, it is important to preserve as many data points as possible when filtering, especially in data-sparse regions.

We conducted a thorough analysis of measurement consistency to identify surveys or locations with significant differences within 50 m, which is less than the width of the first Fresnel zone for the regional minimum ice thickness (60 m at ca. 2100 m depth). We identified 43,337 sites which had measurements within a 50 m window that were not from the same radar profile. A large portion of these sites were from nearby sub-parallel profiles, repeated survey tracks, or duplicated data in the Bedmap Database.

Appendices Figures A1-18 show the consistency between each of the 18 surveys and all other surveys in the Dome C region. Although the differences between nearby data points from different surveys had a median of -1 m and standard deviation of 60 m, large and widespread inconsistencies in measured ice thickness with absolute differences up to 1324 m were identified (*Fig. 2)*.

This analysis was used as a basis for filtering out a total of 275,677 incompatible data points. Overall, we aimed to have a strict filtering procedure to minimize nearby data with large inconsistencies that would impact the spatial statistics of ice thickness variability. If possible, we filtered out older data that could be lacking accurate GPS positioning where newer data was available. Details of the filtering steps applied to individual surveys are given in Table A1. The filtering was applied in four stages:

1. Filtering out locally inconsistent survey segments and duplicated data from specific surveys within 0.2 km, with inconsistent surveys identified through manual inspection (Fig. A1-18) and preferred survey chosen based on lowest mean absolute differences between all other surveys. [*removed 98,176 data points*]
2. Filtering data within a 0.6 km radius of large proximity errors (>50 m), performed sequentially for each survey and ordered based on averaged mean absolute differences between the survey and all other surveys. [*removed 143,707 data points*]
3. Filtering all data points from surveys before 1990 with no GPS positioning and potentially uncertain locations that are within 10 km of more recent surveys with reliable positioning. [*removed 7963 data points*]
4. Filtering out data points along manually supplied sections of specific surveys where nearby measurement inconsistencies are large but could not be removed using automated filters. [*removed 25,831 data points]*




Figure 2: *Measurement compatibility analysis.* ***a)*** *Difference in radar-derived ice thickness measurements (all data: small yellow points) between different surveys if available within a 50 m radius (Red-Blue coloured points show measurement difference). The colormap histogram shows the distribution of differences and is capped at ±150 m, though minimum difference is -655 m and maximum difference is 1324 m.* ***b)*** *Heatmap matrix of inconsistencies, with number of crossover sites (red) in upper left and mean absolute differences for each survey combination (blue) in lower right. Table A1 lists all radar profiles used in this study and filtering steps applied.*

2.3 Spatial variability of filtered ice thickness data

The distance to the nearest measurement calculated for each grid cell in the simulation domain had a median of 2.2 km and standard deviation of 3.9 km, with a maximum distance of 34.1 km. We therefore aimed to calculate and model spatial relationships between measurements for a maximum lag distance of 40 km. We first decimated the radar-derived ice thickness measurements using a median reduction filter and gridded data to uniform 500 m spacings aligned to the Antarctic stereographic EPSG: 3031 grid using tools from the *Verde* python package (Uieda, 2018). This reduced the number of data points from ca. 2.3 million to ca. 61.5 thousand. Our median-decimated ice thickness measurements show that large-scale structure varies across different azimuths (Fig. 1a). To emphasize the relationships at shorter lag distances and minimize the variogram being impacted by large-scale structures, we modelled and removed these using a gaussian filter with a smoothing radius of 20 km (Fig. 3a), proceeding with the residual values (Fig. 3b) and storing modelled large-scale structures to be later added back onto simulated values. To facilitate a gaussian simulation process we normalised residual values to give a mean of 0 and standard deviation of 1, then computed the experimental variogram for 40 lag intervals with a separation of 1 km (Fig. 3c).




Figure 3: *Large-scale, smoothed topography model and measurement variograms.* ***a)*** *Large-scale elevation trend modelled using a gaussian filter with 20 km radius,* ***b)*** *Residuals after removing the regional trend from data,* ***c)*** *Experimental and modelled variogram for normalised ice thickness values for four different azimuths relative to the Polar Stereographic projection. Each azimuth represents a band ±22.5 ° of the central value.*

Despite removing large-scale ice thickness features, directional differences in variability can still be observed in the detrended data (Fig. 3b). To more accurately model spatial variability accounting for azimuthal differences, we calculated directional variograms across grid azimuths centered at 0°, 45°, 90°, 135° (Fig. 3c) relative to Polar Stereographic projected axes.~~,~~ This range adequately captured azimuthal variability when compared to larger ranges examined during exploratory tests and ~~which~~ should be sufficient for data aligned on a grid. All azimuths referred to herein are grid based, not geographical. The directional variograms show measurement variance diverges even at short lag distances, rising sharply for azimuths of 45° and 90° and levelling off at a range of ca. 35 km. Variance for azimuths of 0° and 135° rise more slowly, and these also diverge for lag distances more than 10 km, with the 135° azimuth levelling off after 50 km. We used the linear-weighted residual sums from a least-squares function applied to exponential, spherical, and Gaussian model fits to programmatically identify a best-fitting model type and parameters that can characterise the experimental variograms, prioritizing shorter lag distances and azimuths with the least variance. The experimental variogram at an azimuth of 135° shows the strongest continuity and is characterised well by an exponential model with a range of 50 km and a sill of 1.0 (Fig. 3c). The least continuity is observed for azimuths of 45° and 90° which are

characterised by an exponential model with a range 35 - 39 km and sill of 1.1 - 1.2, especially at lag distances < 10 km and > 30 km (Fig. 3c).

2.4 Stochastic simulation of ice thickness

We used Sequential Gaussian Simulation (SGS) (Deutsch & Journel, 1997) to simulate ice thicknesses based on code adapted from the *gstatsim* python package (MacKie et al., 2023). The algorithm operated over a randomized path over our domain grid to sequentially simulate values based on nearby data and modelled variance. For each unsimulated cell the algorithm builds a conditioning set from nearby measurements, including any previously simulated values, and selects a maximum of 240 values within a radius of 50 km. We tested combinations of search radii from 30 – 100 km and maximum number of conditioning data between 60 and 400 during the developmental phase to find optimum parameters which balanced simulation stability and compute time. The algorithm samples evenly from each of 8 octants if measurements exist within the search radius and populates the set with the nearest data otherwise.

A local mean is estimated through kriging of distance-weighted nearby values, and variance is determined from the covariance matrix for nearby data and the covariance function with model type and parameters identified by the variogram model. Anisotropy is propagated into the simulation by inverse-scaling distances used in the covariance functions based on the variogram range for the least-variable and most-variable azimuths, respectively 135° and 45° with ranges 50 km and 35 km (Fig. 3c). A value is selected at random for each grid cell from a Gaussian distribution drawn with centre at the local mean and variance estimate for the most relevant lag distance. Once all grid cells were simulated, we transformed the values from normalized space back into ice thicknesses and added back the modelled large-scale ice thickness trends. An ensemble of 100 equally-likely ice thickness grids were simulated.

Methodological, model, and parameter choices may impact our simulation results, and during the development stage of this methodology we considered and experimentally explored the impact of these choices to ensure robustness of the results. Our simulation approach can be sensitive to the choice of variogram model used as well as and the model fitting step to determine the range and sill for each azimuth. While regional spatial patterns in simulated elevation were mostly unaffected by these in preliminary tests, model choice impacted local spatial patterns and model fit influences the scale of local variability in simulated grids. It is therefore crucial to conduct a thorough exploratory analysis of the input data to ensure that the model choice and parameters adequately represent the spatial statistics of observations before proceeding with simulations. We mostly implemented a programmatic approach to ensure objectivity in model and parameter selection and used the exploratory experiments to manually check parameter selections where appropriate. Also, modelling spatial variability using isotropic instead of anisotropic experimental variograms produced similar regional spatial patterns and range in simulated bed elevations, and we chose anisotropic to more closely follow the patterns observed in measurements (Fig. 3c). Representing anisotropy using 2 orthogonal directions rather than implementing a more complex approach using continuously varying directional ranges balances improved local heterogeneity and computational complexity.

2.5 Bed elevation grids, uncertainties, and meltwater routing

We estimated 100 bed elevation grids by subtracting each of the 100 simulated ice thickness grids from the Reference Elevation Model of Antarctica (REMA) ice surface elevations (Howat et al., 2019). To evaluate short-scale bed roughness in the Dome C region we calculated the standard deviation within a 2 km radius of each grid cell (Taylor et al., 2004) for each simulated bed, calculating the mean over the ensemble as well as for the Bedmap3 (Pritchard et al., 2025) and Bedmachine v4 (Morlighem, 2026)

interpolations. The median absolute deviation (MAD) over the ensemble of simulated bed elevation grids was used to estimate topographic uncertainty, accounting for both distance from measurements and bed roughness (Shackleton et al., 2023).

We also predicted subglacial meltwater flow directions and hydraulic basins using *pysheds* (Bartos, 2020) to fill sinks and calculate D8 flow directions for subglacial hydraulic potential, $\varphi = \rho_w g z + k \rho_i g h$ (Shreve, 1972), where ρ_w and ρ_i are the density of water and ice (kg m^{-3}), g is gravitational acceleration (m s^{-2}), z and h are ensemble simulated bed elevations and ice thicknesses. We assume that spatial patterns in subglacial water routing are dominantly controlled by topography and hydraulic pressure and not influenced by basal melting rate, and that the ratio between subglacial water pressure and ice overburden pressure, k, is equal to 1~~We assume uniform basal melting and that the ratio between subglacial water pressure and ice overburden pressure, k, is equal to 1~~. In reality, these properties are expected to vary spatially and temporally in response to drainage configurations, subglacial geology, and groundwater interactions which are highly uncertain beneath Antarctica. During sensitivity testing, alternative pressure scenarios of $0.6 \leq k \leq 1.0$ at 0.1 intervals resulted in local shifts in drainage pathways and basin boundaries particularly in regions with low hydraulic gradients, while the broader patterns remained unchanged. Using our assumptions, we quantify how uncertainty in bed topography propagates into subglacial hydrological predictions under commonly adopted assumptions rather than reproduce the full complexity of the basal drainage system. Stream and basin probability was calculated from averaged stream locations over the ensemble.

3. Results

3.1 Simulated ice thickness and bed elevation

We simulated an ensemble of 100 ice thickness and bed elevation grids for a 400x400 km^2 region surrounding Dome C from 1120 km to 1520 km Easting, and -1170 km to -720 km Northing on an EPSG: 3031 Antarctic Stereographic projection. The full ensemble of simulated surfaces can be viewed in *Appendix animation A*, which shows that values near measurements remain consistent across the ensemble, whereas away from measurements values vary between simulations, increasingly with both distance from measurements and local bed roughness. Simulation #001 is shown in *Figures 1b* and *1c*, which confirms that large-scale structures such as northwest-southeast oriented bands of thicker and thinner ice observed in radar data (e.g. Concordia Ridge; Fig. 1a) are represented well. Topographic features 10's of kilometres in length are also resolved and are consistent across the ensemble in regions with high data density, such as the 3 patches of low ice thickness near 1350 km Easting, -960 km Northing. Features of this scale can shift in location over the ensemble for low data density regions in the southwest and north of the domain. Small-scale topography below 5 km varies significantly over the ensemble in all regions except in the central region where data density is the highest surrounding the ice core sites (Fig. 1a). The simulations incorporate new measurements from a KOPRI-UA airborne radar survey over the Belgica Highlands to the southwest of Concordia Station, documenting the southeast extent of the Belgica Highlands and Concordia Ridge with increased detail and revealing bed elevations more than 300 m higher than estimated from sparse data in previous interpolations (Bedmap 3: Pritchard et al., 2025; Bedmachine v4: Morlighem, 2026).

3.2 Bed elevation uncertainty

The Median Absolute Deviation (MAD) over the ensemble of 100 simulated bed elevation grids quantifies spatial uncertainties attributed to the combined influence of measurement density and basal roughness in the Dome C region (Fig. 1d). The domain mean is 53 m with standard deviation of 32 m, and

lowest MAD near measurements, especially in high data density regions near Concordia Station. The highest MAD is observed far from data, reaching a maximum of 260 m to the grid southeast. The primary pattern in MAD follows the radar survey tracks, but secondary patterns are observed with grid northwest-southeast trending bands of higher MAD aligned with the 10's of km scale troughs in the grid southwest and northwest. Regions of higher MAD are also observed for relatively uniform radar survey spacings, for example the band near 1300 km Easting, -1040 km Northing which is 60 m higher than the surroundings, which also correlates with rougher terrain in simulated bed. Over the Dome C Plateau MAD reaches a maximum of 60 m despite relatively dense radar data with maximum 4 km spaced surveys. However, within several kilometres of the Million Year Ice Core (MYIC) and Beyond EPICA Little Dome C (BELDC) ice core sites data is dense enough to constrain topographic uncertainty to 0 at the 500 m grid cell scale of our simulations.

4. Discussion

Gridded representations of subglacial bed topography are fundamental boundary conditions for ice sheet models and underpin a wide range of glaciological analyses. For modelling purposes, adequate representation of basal roughness should be prioritised for realistic constraints on basal shear stress. Although the impact of shear heating is likely to be small beneath ice domes due to low flow velocities, the basal thermal state could be sensitive to small changes since much of the basal ice is at or near the pressure-melting point. Replacing smooth bed with realistically rough topography was shown to improve modelled basal ice motion and replicate temperate ice layer thicknesses observed in Greenland boreholes (Law et al., 2023). In the context of ice core investigations, rougher basal topography can also lead to deformation of basal ice layers which disrupts the continuity of deep ice stratigraphy. Provided that all other conditions for locating old ice layers with undisturbed stratigraphy are also met (cf. Fischer et al., 2013) it is therefore favourable to choose sites with relatively smooth basal topography when selecting potential drill sites for old ice cores. We discuss our simulated bed topography results in the context of basal roughness (Section 4.1), subglacial meltwater drainage (Section 4.2), and geothermal heat flow (Section 4.3).

4.1 Basal roughness

We calculated standard deviations within a 2 km radius of each grid cell, $\sigma_{2km,}$ as a measure of basal roughness (Taylor et al., 2004) for each of our ensemble simulated beds (Fig. 4a: Sim #001), as well as the Bedmap 3 interpolation (Fig. 4b) (Pritchard et al., 2025) and Bedmachine v4 (Fig. 4c) (Morlighem, 2026). The Bedmachine v4 also includes 2-30 km scale topography inverted from ice surface observations (Ockenden et al., 2026). Roughness is generally much higher in our simulated bed compared with previous interpolations, especially in between radar profiles (Fig. 4a-c). For the Bedmap3 and Bedmachine v4 grids, respective σ_{2km} domain means are 21 m and 17 m, while domain standard deviations are 21 m and 16 m. The domain mean σ_{2km} across our results (with ensemble 1σ uncertainty estimate) is more than double that of previous work at 53 ± 0.2 m, reaching maximums of 390 ± 7 m with domain standard deviation of 25 ± 0.2 m. Regionally averaged basal roughness is therefore underestimated by more than 50% in single-interpolation based approaches.

The spatial distribution of rougher terrain in the Bedmap3 interpolation is dominated by radar data availability, as high σ_{2km} is located along survey profiles reaching a maximum of 279 m with little variability in between. Roughness for the Bedmachine v4 grid is not impacted to the same extent by measurement availability, although some roughness patterns appear to reflect ice surface features over the western Belgica Highlands which probably originate from the surface inversion methodology of

Ockenden et al., (2026). Several radar profiles are also still observed in the roughness map (Fig. 4c), likely due to the inclusion of incompatible ice thickness measurements in the interpolation process. This emphasises the necessity of our rigorous measurement compatibility analyses and screening steps before starting interpolation or simulation work. Our simulated beds have improved spatial distribution of basal roughness which is more consistent with that expected of natural landscapes, with minimal impacts from measurement distribution on individual bed simulations (Fig. 4a).

The pronounced anisotropy in topography in the Dome C region also limits the reliability of ice thickness or bed surfaces derived using isotropic interpolation strategies, which could mask important topographic structure by applying uniform smoothing in all directions in data gaps. By explicitly modelling and incorporating anisotropy into our stochastic simulations we also increase the probability that realistic structure is simulated in unsurveyed areas. We suggest that future analyses relying on gridded representations of basal topography such as ice flow and basal conditions modelling should carefully assess sensitivity to basal roughness before committing to using smoothed interpolations with unrealistic bed roughness. Our stochastically simulated surfaces preserve natural roughness patterns and could be used as more realistic boundary conditions for modelling as well as providing a suite of equally probable inputs for sensitivity analyses to constrain the impacts of bed elevation uncertainties.




Figure 4: *Standard deviation within a 2 km window for gridded ice thickness:* ***a)*** *Simulation #001,* ***b)*** *Bedmap 3 interpolation,* ***c)*** *Bedmachine v4 interpolation.*

4.2 Subglacial meltwater drainage

There are 111 locations with radar-inferred subglacial lakes in our simulation domain (Livingstone et al., 2022), and these occur beneath ice ranging from 2601 m to 4326 m thick. Satellite observations show that large volumes of meltwater can transit through the subglacial drainage system over short timescales, with 1.8 km^3 of water travelling 290 km over a 16 month period between lakes in the Adventure Trench (Wingham et al., 2006). We modelled basal hydraulic potential across the ensemble to constrain the range of possible subglacial water flow directions, meltwater routing, and drainage basins assuming uniform basal melting. ~~We modelled basal hydraulic potential to provide a simplified assessment of subglacial water flow directions to predict meltwater routing and drainage basins assuming uniform basal melting.~~ ~~We used~~ The simulated bed ensemble ~~to~~ constrains drainage uncertainty associated with bed elevation uncertainty, which we compare ~~and compared this~~ to drainage predictions using the Bedmap 3 interpolation (Fig. 5a,b). Subglacial drainage off the Dome C Plateau is well-constrained, with high probability for water routing away from the plateau following ice drainage divides and splitting water flow in either direction of the Concordia Trench (Fig. 5a). Our

analyses reveal uncertain drainage divides east of Concordia Station and towards the western extent of the Belgica Highlands, associated with large gaps between radar survey profiles.

Concordia Lake is the largest subglacial lake within our study area, estimated from radar profiles to be minimum 45 km long and 16 km wide and containing water volumes of 200 ± 40 km^3 (Tikku et al., 2005). Water inflow to this lake is well-constrained by the Concordia Trench, but outflow and downstream routing of released meltwater is not well-constrained by radar data. Our simulations suggest that water could drain either towards Vincennes basin or towards Adventure Trench, and the most probable drainage basin divides are to the southwest of the lake in contrast to the divides predicted




Figure 5: *Subglacial drainage predicted using ensemble simulated beds. a) Stream probability calculated from averaged stream predictions over the ensemble (blue), with predicted streams using Bedmap3 bed (orange). b) Drainage divide probability calculated as averaged divides over the ensemble, with divides predicted using Bedmap3 (black). Ice cores also plotted (red diamonds) and radar-detected subglacial lakes (green points). Lakes with estimated minimum surface area >50 km^2 are labelled: AL = Aurora Lake (Tabacco et al., 2003); CL = Concordia Lake (Oswald and Robin, 1973); HL = Horseshoe Lake (Blankenship et al., 2009); ITL12+13 (Tabacco et al., 2003).*

using Bedmap3 grids (Fig. 5b). The lack of strong basal topographic control over downstream water routing suggested by limited available radar data indicates that downstream drainage could be sensitive to migrating ice divides, and more radar surveys are necessary to reduce the uncertainty in water routing.

The drainage basin constrained by the western Belgica Highlands and Concordia Ridge contains 11 subglacial lakes with a combined radar-estimated surface area of 147 km^2. Five of these lakes are located beneath thinner ice <3000 m along Concordia Ridge and 6 along the valley floor where ice can be over 3400 m thick. Our stochastic simulations constrain potential flow paths for a drainage system linking these lakes and indicate that large volumes of water draining from this basin could be routed either to the southwest similar to predictions using the Bedmap3 grid or 100 km further north (Fig. 5).

The preservation of deep ice layers in undisturbed stratigraphic sequence ~~are~~ is sensitive to basal melt, ice flow, and basal ice accretion (Fischer et al., 2013; Van Liefferinge & Pattyn, 2013). Widespread subglacial lakes and meltwater drainage activity evidence basal melting in the Dome C region and poor hydraulic connectivity between lakes could increase potential for basal ice accretion, which together disturb deep ice stratigraphy and limit the preservation of old ice layers. For example, at Lake Concordia

enhanced basal melting of 4.3 mm a^{-1} ± 27% is predicted over most of the lake, and 15.1 m ± 65% of accreted ice in the northeast corner (Thoma et al., 2009). Careful consideration of the subglacial environment is required to identify targets for ice core drill sites and our uncertainty constrained subglacial drainage analyses identify sites with low potential for water ponding or drainage activity (Fig. 5a,b).

4.3 Geothermal heat flow

The GHF beneath Antarctica is largely unknown, however locally elevated GHF could impact basal conditions and ice flow especially beneath slow-flowing ice dome regions in the absence of frictional heat from basal sliding (Pittard et al., 2016). Ice sheet modelling investigations to identify suitable ice core drilling sites are also sensitive to GHF boundary conditions (Sutter et al., 2019). At Dome C, region-wide means range from 28.6 ± 5.1 to 59.0 ± 3.3 mW m^{-2} in large-scale GHF models (An et al., 2015; Burton-Johnson et al., 2020; Hazzard & Richards, 2023; Lösing & Ebbing, 2021; Martos et al., 2017; Purucker, 2012; Shen et al., 2020; Stål et al., 2021), and 10's of kilometre scale spatial distributions differ significantly between models (Fig. 6). Local GHF estimates at the EPICA core site of 57.9 ± 6.4 m Wm^{-2} (Talalay et al., 2020) and 54.5 ± 3.5 (Passalacqua et al., 2017) have high uncertainties and might not reflect regional values.

Topographic focusing of geothermal heat can redistribute heat to local ridges or valleys, and the polarity of heat refraction patterns depends on the thermal conductivity contrast between ice and the underlying rock (Willcocks et al., 2021; Reading et al., 2022). ~~lowers the amount of heat received along ridges and increase it in valleys relative to the background value (van der Veen et al., 2007).~~ The thermal conductivity of ice at depths beyond 750 m is estimated between 2.0 – 2.8 W m^{-1} K^{-1} (Willcocks et al., 2021), whereas igneous Antarctic bedrock ranges 1.5 – 4.0 W m^{-1} K^{-1} in samples limited to exposed outcrops mostly near the ice margin (Willcocks et al., 2025). The geology and corresponding thermal conductivity are uncertain beneath Dome C, so as an illustrative example of how topography could impact geothermal heat distribution we calculated an adjustment factor to account for local topographic modifications of background GHF following the~~a~~ simplified geostatistical approach of Colgan et al., (2021). Since the geology is not known, we adopt their globally averaged empirically determined averaging radius of 5 km and characteristic height needed to induce 100% change in local GHF of 950 m, using their equation 2 to calculate potential magnitude and spatial patterns of heat redistribution. However, we recognise that the magnitude and polarity of such modifications could be substantially different in reality depending on the conductivity of the bed.

This calculation was done for each of the simulated beds in our ensemble to produce a topographic adjustment factor map, and Figure 7a shows the ensemble median values. Similar to the ensemble elevation uncertainty, topographic adjustments to GHF are more consistent close to radar profiles, and the map inherently displays lower confidence adjustment factors closer to zero since topographic features far from data vary in location over the ensemble. Note that our confidence in adjustment factor is based only on uncertainty in bed geometry, and neglects uncertainty associated with bedrock thermal conductivity and polarity of thermal refraction, or the averaging radius and characteristic height parameters. Using our adopted parameterization of GHF redistribution, we show that ~~L~~locally elevated GHF up to 30% more than the background value ~~can be expected~~could occur within the troughs (e.g. Adventure, Concordia), yet along ridges (e.g. Concordia) local GHF could be 30% lower (Fig. 7a,b). Beneath the Dome Fuji ice dome, topographic modification to background GHF was predicted to induce widespread local anomalies of ±20%, aggregating to raise the regional average by 0.6% (Shackleton et al., 2023). The Dome C regional mean values are low over the ensemble at 0.05 ± 0.04%, though regional standard deviations of 0.1 ± 0.0% indicate widespread potential for localised GHF adjustments of 3-7 mW m^{-2} based on modelled range ca. 30-70 mW m^{-2}.




Figure 6: *Geothermal heat flow (GHF) estimates near Dome C. Large-scale models plotted as native resolution in the upper 8 maps, with domain mean and standard deviation for each model provided in bottom right corner. Core sites plotted as EPICA: diamond, MYIC: triangle, BELDC: circle. Histograms for modelled GHF are plotted below, with associated value at core sites (diamonds, triangles, circles) together with other local GHF estimates and reported uncertainty at the EPICA core site.*

We find 80% of the radar-inferred subglacial lake sites are within a zone of locally elevated GHF (Fig. 7c), suggesting that topographic modification of geothermal heat could be a contributing factor to the spatial pattern of basal melting and lake distributions in regions where basal ice is close to the pressure melting point. ~~We find 80% of the radar-inferred subglacial lake sites are within a zone of locally elevated GHF (Fig. 7c), which in conjunction with increased ice thickness could contribute to the spatial pattern of basal melting and lake distributions.~~ Spatial variability in GHF at much larger scale of 10's of kilometres increased modelled basal melt rates in the Aurora Basin by up to 4.5% (McCormack et al., 2022). Smoothed bed in data sparse regions of previous interpolations therefore limits robust assessments of basal thermal conditions, and using realistically rough topography should be prioritized when delineating thawed/frozen basal ice and calculating melt rates, which will also improve the reliability of basin-wide thermodynamical and ice flow modelling. Given the large uncertainties and range of modelled GHF near Dome C and across Antarctica, the local topographic adjustments to GHF could be used instead to find suitable sites for future ice coring, prioritising sites with locally lowered GHF. ~~The~~ Our analysis shows that the EPICA core site ~~is~~ could be situated in locally elevated GHF by 2% (Fig. 7c), whereas the new sites BELDC and MYIC are situated in a region with potentially ~~locally~~ lowered GHF by 12 -13% which

~~should~~ could be favourable for deep ice layer preservation. Considering the impacts of uncertain bed geometry alone, ~~V~~variability over the ensemble is less than 0.08% for all ice core sites since they are generally close to radar data. Borehole GHF estimates should also be adjusted to account for the potential range of local topographic modifications before extrapolating over a larger region.




Figure 7: *Topographic adjustments to geothermal heat flow (GHF).* ***a)*** *Local topographic adjustments to GHF displayed as the median adjustments over the ensemble of 100 simulated bed elevation grids. Ice thickness contours at 500 m intervals are drawn in black.* ***b)*** *Enlarged view of the central plateau near the ice core sites, also with radar-inferred subglacial lakes drawn as green points.* ***c)*** *Histogram of topographic adjustments, with ensemble median sampled adjustments at the subglacial lake sites (green) and core sites (orange) with median absolute deviation error bar.*

5. Conclusions

New airborne radar surveys uncover previously unmapped sectors of the Belgica Highlands, and we used a rigorous approach to integrating new observations with available ice thickness data near Dome C. Our measurement compatibility analysis guided strict filtering procedures, and we find that rigorous quality screening and cross-validation to integrate new measurements with existing data is crucial for characterising basal roughness. This process is a necessary pre-requisite for reliable gridding or simulating of bed topography to minimize the impact that nearby inconsistent measurements have on spatial statistics. Our simulated bed has improved representation of basal roughness in data sparse regions, with region-wide means more than double those of previous work and spatial patterns free from the influence of radar survey geometry. The ensemble simulation approach for mapping elevation uncertainty accounts for the combined influence of data density and local bed roughness characteristics, producing higher uncertainties where measurements are sparse and the terrain is rough. Domain-mean bed elevation uncertainty is 53 m and reaches 260 m in data-sparse regions, however, near the BELDC and MYIC ice core sites dense radar coverage constrains elevation uncertainty effectively to zero at a 500 m grid scale.

Subglacial drainage analyses using ensemble simulated bed identify well-constrained drainage away from the Dome C Plateau, but poorly constrained outflow from Concordia Lake and western Belgica Highlands drainage basin. These uncertainties impact predictions of subglacial lake connectivity and downstream water transport, highlighting locations where additional radar surveys are required to adequately constrain subglacial drainage. Our work also reveals ~~the~~ potential patterns of local topographic modifications to regional GHF, which could be critical to delineating thawed/frozen bed, locating suitable sites for ice core drilling, and interpreting retrieved cores and borehole

measurements/modelling investigations. We identify locations where local bed topography could lower GHF by up to 30% of the regional value, and the MYIC and BELDC ice core drilling sites are within a zone of locally lowered GHF by 12% which ~~should~~ could be favourable for preserving deep ice layer continuity. Topographic focussing of geothermal heat ~~likely enhances~~could enhance basal melting in many trough locations and ~~could~~ exert a control over the distribution of subglacial lakes~~;~~, ~~with~~ our analysis shows most radar-inferred subglacial lakes occurring in a region ~~of~~ where we predict locally elevated GHF. Given the large spread among continental-scale GHF models, we suggest that locally resolved topographic adjustments to GHF provide an independent and complementary constraint on spatial variability in basal conditions.

Our study emphasizes that the physical realism of Antarctic subglacial boundary conditions depends on capturing the inherent variability and anisotropy of the bed. These properties are currently obscured in widely used Antarctic bed products due to smoothing from isotropic deterministic interpolation, which limits their reliability for use in ice-sheet and subglacial environment modelling. Survey design should capture directional anisotropy by acquiring profiles along multiple azimuths, extending beyond traditional along- and across-flow transects, while gridded products should prioritize simulation-based approaches that maintain realistic spatial continuity in data-sparse regions. By resolving variability at scales relevant to basal processes and adopting ensemble-based topography, the community can more effectively propagate measurement and interpolation uncertainties into subsequent ice-sheet and subglacial environment analyses.

6. Acknowledgements

This research was supported by funding from Korea Institute of Marine Science & Technology Promotion (KIMST) grant funded by the Ministry of Oceans and Fisheries (KIMST RS-2026-25521468 and PE26080), and Research Council of Norway (FRINATEK 315246). The Technical University of Denmark acquired POLARIS data under ESA Contract No. 4000144378/23/NL/CT, supported by the European Space Agency (ESA), Australian Antarctic Division, and Institut Polaire Français Paul Émile Victor. Computing resources from Sigma2 – the National Infrastructure for High-Performance Computing and Data Storage in Norway were utilised in this work through project NN10061K. We thank the handling editor Yuzhen Yan and reviewers Tobias Stål and Shuai Yan for their reviews of this work which helped to improve the final manuscript. We also acknowledge the many researchers, institutions, and national Antarctic programmes who have contributed ice thickness data to the Bedmap Database and appreciate the community commitment to open data and software sharing which is essential for advancing Antarctic science. The Norwegian Polar Institute contribution forms part of the EU Horizon 2020 Beyond EPICA Oldest Ice Core (BE-OIC) project (Grant 815384). This is BE-OI contribution number XX.

Code availability

Code used for geostatistical modelling and stochastic simulation is available at: https://github.com/calvinshackleton/DomeC-BedElevation.git (Shackleton, 2026), which are based on tools from open-source python software: Verde (Uieda, 2018), GeoStatsPy (Pyrcz et al., 2021), SciKit Gstat (Mälicke et al., 2021), GStatSim (MacKie et al., 2023).

Data availability

Appendix animation A, as well as ensemble grids of simulated ice thickness, bed elevation, and local topographic adjustments to geothermal heat flow are available as GeoTIFFs from the Norwegian Polar Data Centre via: https://doi.org/10.21334/NPOLAR.2026.D05BFE78 (Shackleton and Matsuoka, 2026). The ice penetrating radar data from the KOPRI-UA survey are available here:

https://dx.doi.org/doi:10.22663/KOPRI-KPDC-00003141. Ice thickness data from the DTU survey using POLARIS are available here: https://doi.org/10.11583/DTU.32685036 (Kusk, 2026).

Author contributions

CS and KM conceptualized the study, developed the methodology, and wrote the manuscript. CS carried out the analysis, wrote the workflow code, and performed the simulations. BF, JL, AR, PG, DT, HTJ, CC, and CS contributed to the collection and processing of the KOPRI/UA radar data. JD and AK contributed to the collection and processing of the DTU radar data. All authors contributed to the preparation of the final manuscript.

Competing interests

The contact author declares that none of the authors have any competing interests.

References

An, M., Wiens, D. A., Zhao, Y., Feng, M., Nyblade, A., Kanao, M., et al.: Temperature, lithosphere-asthenosphere boundary, and heat flux beneath the Antarctic Plate inferred from seismic velocities. *Journal of Geophysical Research: Solid Earth*, *120*(12), 8720–8742. https://doi.org/10.1002/2015JB011917, 2015.

Arthur, J. F., Shackleton, C., Moholdt, G., Matsuoka, K., and van Oostveen, J.: Evidence of active subglacial lakes under a slowly moving coastal region of the Antarctic Ice Sheet. *The Cryosphere*, *19*(1), 375–392. https://doi.org/10.5194/tc-19-375-2025, 2025.

Awasthi, A., Taylor, D., Kolpuke, S., Reyhanigalangashi, O., Vanderburgh, L., Elluru, D., O'Neill, C., Chung, C., Lee, L., and Gogineni, P.: High-sensitivity transceiver for airborne ultra-wideband radio echo sounding of polar regions. *2022 IEEE Microwaves, Antennas, and Propagation Conference (MAPCON)*, Bangalore, India, 2022.

Bartos, M. (2020). pysheds. Python. 10.5281/zenodo.3822494

Berends, C.J., Köhler, P., Lourens, L. J., and van de Wal, R. S. W.: On the Cause of the Mid-Pleistocene Transition. *Reviews of Geophysics*, *59*(2), e2020RG000727. https://doi.org/10.1029/2020RG000727, 2021.

Blankenship, D. D., Kempf, S. D., Young, D. A., Richter, T. G., Schroeder, D. M., Greenbaum, J. S., Holt, J. W., van Ommen, T., Warner, R. C., Roberts, J. L., Young, N. W., Lemeur, E. and Siegert, M. J.: IceBridge HiCARS 1 L2 Geolocated Ice Thickness. (IR1HI2, Version 1). [Data Set]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. https://doi.org/10.5067/F5FGUT9F5089, 2011.

Blankenship, D. D., Kempf, S. D., Young, D. A., Richter, T. G., Schroeder, D. M., Greenbaum, J. S., van Ommen, T., Warner, R. C., Roberts, J. L., Young, N. W., Lemeur, E., Siegert, M. J. and Holt, J. W.: IceBridge HiCARS 1 L1B Time-Tagged Echo Strength Profiles. (IR1HI1B, Version 1). [Data Set]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. https://doi.org/10.5067/W2KXX0MYNJ9G, 2017.

612 Blankenship, D. D., Kempf, S. D., and Young, D. A.: Support Office for Airborne Research 1 km sampled
613 ice thickness data. U.S. Antarctic Program (USAP) Data Center. https://doi.org/10.15784/601588,
614 2022.

615 Bianchi, C., Cafarella, L., De Michelis, P., Forieri, A., Frezzotti, M., Tabacco, I. E., and Zirizzotti, A.: Radio
616 Echo Sounding (RES) Investigations at Talos Dome (East Antarctica): Bedrock Topography and Ice
617 Thickness. *Ann. Geophys, 46* (6). https://doi.org/10.4401/ag-3471, 2009.

618 Burton-Johnson, A., Dziadek, R., and Martin, C.: Review article: Geothermal heat flow in Antarctica:
619 current and future directions. *The Cryosphere*, *14*(11), 3843–3873. https://doi.org/10.5194/tc-14-
620 3843-2020, 2020.

621 Chung, A., Parrenin, F., Steinhage, D., Mulvaney, R., Martín, C., Cavitte, M. G. P., et al.: Internal reflecting
622 horizons at Little Dome C, Antarctica [Application/zip]. PANGAEA.
623 https://doi.org/10.1594/PANGAEA.957176, 2023a.

624 Chung, A., Parrenin, F., Steinhage, D., Mulvaney, R., Martín, C., Cavitte, M. G. P., et al.: Stagnant ice and
625 age modelling in the Dome C region, Antarctica. *The Cryosphere*, *17*(8), 3461–3483.
626 https://doi.org/10.5194/tc-17-3461-2023, 2023b.

627 Chung, A., Parrenin, F., Steinhage, D., Mulvaney, R., Martín, C., Cavitte, M., Lilien, D., Helm, V., Taylor,
628 D., Gogineni, P., Ritz, C., Frezzotti, M., O'Neill, C., Miller, H., Dahl-Jensen, D., and Eisen,
629 O.: Bedrock at Little Dome C, Antarctica
630 [dataset]. *PANGAEA*, https://doi.org/10.1594/PANGAEA.957461, 2023c.

631 Cianfarra, P., Forieri, A., Salvini, F., Tabacco, I. E., and Zirizotti, A.: Geological setting of the Concordia
632 Trench-Lake system in East Antarctica. *Geophysical Journal International*, *177*(3), 1305–1314.
633 https://doi.org/10.1111/j.1365-246X.2009.04123.x, 2009.

634 Colgan, W., MacGregor, J. A., Mankoff, K. D., Haagenson, R., Rajaram, H., Martos, Y. M., et al.:
635 Topographic Correction of Geothermal Heat Flux in Greenland and Antarctica. *Journal of*
636 *Geophysical Research: Earth Surface*, *126*(2). https://doi.org/10.1029/2020JF005598, 2021.

637 Cuffey, K. M., and Paterson, W. S. B. *The Physics of Glaciers* (4th ed.). Elsevier, 2010.

638 Dall, J., Kristensen, S. S., Krozer, V., Hernández, C. C., Vidkjær, J., Kusk, A., et al.: ESA'S POLarimetric
639 Airborne Radar Ice Sounder (POLARIS): design and first results. *IET Radar, Sonar & Navigation*,
640 *4*(3), 488–496. https://doi.org/10.1049/iet-rsn.2009.0035, 2010.

641 Dall, J., Nielsen, U., Kusk, A., and van de Wal, R.S.W.: Ice flow mapping with P-band SAR, *in Proceedings*
642 *of the IEEE 2013 International Geoscience and Remote Sensing Symposium*, 4 p., Melbourne,
643 July 2013.

644 Deutsch, C. V., and Journel, A. G.: *GSLIB: Geostatistical software library and user guide* (Second Edition).
645 New York: Oxford University Press. http://claytonvdeutsch.com/wp-
646 content/uploads/2019/03/GSLIB-Book-Second-Edition, 1997.

647 Drewry, D. J., Jordan, S. R., and Jankowski, E.: Measured Properties of the Antarctic Ice Sheet: Surface
648 Configuration, Ice Thickness, Volume and Bedrock Characteristics. *Annals of Glaciology*, *3*, 83–
649 91. https://doi.org/10.3189/S0260305500002573, 1982.

650 Ferraccioli, F., Corr, H., Jordan, T., Robinson, C., Armadillo, E., and Armadillo, G.: Airborne radar bed
651 elevation picks across the Wilkes Subglacial Basin, 2005-2006 [Data set]. Polar Data Centre,
652 Natural Environment Research Council, UK. https://doi.org/10.5285/59e5a6f5-e67d-4a05-99af-
653 30f65656940, 2018.

Fischer, H., Severinghaus, J., Brook, E., Wolff, E., Albert, M., Alemany, O., et al.: Where to find 1.5 million yr old ice for the IPICS "Oldest-Ice" ice core. *Climate of the Past*, *9*(6), 2489–2505. https://doi.org/10.5194/cp-9-2489-2013, 2013.

Forieri, A., Zuccoli, L., Bini, A., Zirizzotti, A., Remy, F., and Tabacco, I. E.: New bedrock map of Dome C, Antarctica, and morphostructural interpretation of the area. *Annals of Glaciology*, *39*, 321–325. https://doi.org/10.3189/172756404781814456, 2004.

Frezzotti M, Bitelli G, De Michelis P, et al. Geophysical survey at Talos Dome, East Antarctica: the search for a new deep-drilling site. *Annals of Glaciology*, 39:423-432. https://doi:10.3189/172756404781814591, 2004.

Frémand, A. C., Fretwell, P., Bodart, J. A., Pritchard, H. D., Aitken, A., Bamber, J. L., et al.: Antarctic Bedmap data: Findable, Accessible, Interoperable, and Reusable (FAIR) sharing of 60 years of ice bed, surface, and thickness data. *Earth System Science Data*, *15*(7), 2695–2710. https://doi.org/10.5194/essd-15-2695-2023, 2023.

Hazzard, J. A. N., and Richards, F. D.: Antarctic Geothermal Heat Flow, Crustal Conductivity and Heat Production Inferred From Seismological Data. *Retrieved from* https://eartharxiv.org/repository/view/5939/, 2023.

Howat, I. M., Porter, C., Smith, B. E., Noh, M.-J., and Morin, P.: The Reference Elevation Model of Antarctica. *The Cryosphere*, *13*(2), 665–674. https://doi.org/10.5194/tc-13-665-2019, 2019.

Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., et al.: Orbital and Millennial Antarctic Climate Variability over the Past 800,000 Years. *Science*, *317*(5839), 793–796. https://doi.org/10.1126/science.1141038, 2007.

Kusk, A., and Dall, J.: SAR focusing of P-band ice sounding data using back-projection. *Proceedings of the IEEE 2010 International Geoscience and Remote Sensing Symposium*, pp. 4071-4074, Honolulu, July 2010.

Kusk, A.: Dome C ice thickness measurements from POLARIS. [Dataset]. https://doi.org/10.11583/DTU.32685036, 2026.

Law, R., Christoffersen, P., MacKie, E., Cook, S., Haseloff, M., and Gagliardini, O.: Complex motion of Greenland Ice Sheet outlet glaciers with basal temperate ice. *Science Advances*, *9*(6), eabq5180. https://doi.org/10.1126/sciadv.abq5180, 2023.

Lee, J. S., Jurkevich, L., Dewaele, P., Wambacq, P., and Oosterlinck, A.: Speckle filtering of synthetic aperture radar images: A review. *Remote Sensing Reviews*, *8*(4), 313–340. https://doi.org/10.1080/02757259409532206, 1994.

Lisiecki, L. E., and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic δ 18O records. *Paleoceanography*, *20*(1), 1–17. https://doi.org/10.1029/2004PA001071, 2005.

Livingstone, S. J., Li, Y., Rutishauser, A., Sanderson, R. J., Winter, K., Mikucki, J. A., et al.: Subglacial lakes and their changing role in a warming climate. *Nature Reviews Earth & Environment*, 1–19. https://doi.org/10.1038/s43017-021-00246-9, 2022.

Lösing, M., and Ebbing, J.: Predicting Geothermal Heat Flow in Antarctica with a Machine Learning Approach. *Journal of Geophysical Research: Solid Earth*, *126*(6), e2020JB021499. https://doi.org/10.1029/2020JB021499, 2021.

Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., et al.: High-resolution carbon dioxide concentration record 650,000–800,000 years before present. *Nature*, *453*(7193), 379–382. https://doi.org/10.1038/nature06949, 2008.

MacKie, E. J., Schroeder, D. M., Caers, J., Siegfried, M. R., and Scheidt, C.: Antarctic Topographic Realizations and Geostatistical Modeling Used to Map Subglacial Lakes. *Journal of Geophysical Research: Earth Surface*, *125*(3), e2019JF005420. https://doi.org/10.1029/2019JF005420, 2020.

MacKie, Emma J., Schroeder, D. M., Zuo, C., Yin, Z., and Caers, J.: Stochastic modeling of subglacial topography exposes uncertainty in water routing at Jakobshavn Glacier. *Journal of Glaciology*, *67*(261), 75–83. https://doi.org/10.1017/jog.2020.84, 2021.

MacKie, Emma J., Field, M., Wang, L., Yin, Z., Schoedl, N., Hibbs, M., and Zhang, A.: GStatSim V1.0: a Python package for geostatistical interpolation and conditional simulation. *Geoscientific Model Development*, *16*(13), 3765–3783. https://doi.org/10.5194/gmd-16-3765-2023, 2023.

MacGregor, J. A., Boisvert, L. N., Medley, B., Petty, A. A., Harbeck, J. P., Bell, R. E., et al.: The scientific legacy of NASA's Operation IceBridge. Reviews of Geophysics, 59, e2020RG000712. https://doi.org/10.1029/2020RG000712, 2021.

Mälicke, M., Möller, E., Schneider, H. D., and Müller, S.: mmaelicke/scikit-gstat: A scipy flavoured geostatistical variogram analysis toolbox. Zenodo. https://doi.org/10.5281/zenodo.4835779, May 28th, 2021.

Martos, Y. M., Catalán, M., Jordan, T. A., Golynsky, A., Golynsky, D., Eagles, G., and Vaughan, D. G.: Heat Flux Distribution of Antarctica Unveiled. *Geophysical Research Letters*, *44*(22), 11,417-11,426. https://doi.org/10.1002/2017GL075609, 2017.

McCormack, F. S., Roberts, J. L., Dow, C. F., Stål, T., Halpin, J. A., Reading, A. M., and Siegert, M. J.: Fine-Scale Geothermal Heat Flow in Antarctica Can Increase Simulated Subglacial Melt Estimates. *Geophysical Research Letters*, *49*(15), e2022GL098539. https://doi.org/10.1029/2022GL098539, 2022.

McCormack, F. S., Stål, T., Shao, N., MacKie, E., Fabela Hinojosa, A., Lösing, M., et al.: Synthetic bed topographies for Antarctica and their utility in ice sheet modelling. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, *384*(2319), 20240537. https://doi.org/10.1098/rsta.2024.0537, 2026.

Morlighem, M.: MEaSUREs BedMachine Antarctica, Version 4 [Data set]. NASA National Snow and Ice Data Center Distributed Active Archive Center. https://doi.org/10.5067/POJQI54A45HX, 2026.

Mulvaney, R., Frezzotti, M., Ritz, C., Martín, C., Rix, J., and Chung, A.: Bedrock in the Dome C region of Antarctica [Dataset]. PANGAEA. https://doi.org/10.1594/PANGAEA.963474, 2023.

Nielsen, U., and Dall, J.: Direction-of-Arrival estimation for radar ice sounding surface clutter suppression. *IEEE Transactions on Geoscience and Remote Sensing*, Vol. 53, No. 9, pp. 5170-5179, September 2015.

Ockenden, H., Bingham, R. G., Goldberg, D., Curtis, A., and Morlighem, M.: Complex mesoscale landscapes beneath Antarctica mapped from space. *Science*, *391*(6782), 314–319. https://doi.org/10.1126/science.ady2532, 2026.

Parrenin, F., Cavitte, M. G. P., Blankenship, D. D., Chappellaz, J., Fischer, H., Gagliardini, O., et al.: Is there 1.5-million-year-old ice near Dome C, Antarctica? *Cryosphere*, *11*(6), 2427–2437. https://doi.org/10.5194/tc-11-2427-2017, 2017.

Passalacqua, O., Ritz, C., Parrenin, F., Urbini, S., and Frezzotti, M.: Geothermal flux and basal melt rate in the Dome C region inferred from radar reflectivity and heat modelling. *Cryosphere*, *11*(5), 2231–2246. https://doi.org/10.5194/tc-11-2231-2017, 2017.

Pittard, M. L., Galton-Fenzi, B. K., Roberts, J. L., and Watson, C. S.: Organization of ice flow by localized regions of elevated geothermal heat flux. *Geophysical Research Letters*, *43*(7), 3342–3350. https://doi.org/10.1002/2016GL068436, 2016.

Pritchard, H. D., Fretwell, P. T., Fremand, A. C., Bodart, J. A., Kirkham, J. D., Aitken, A., et al.: Bedmap3 updated ice bed, surface and thickness gridded datasets for Antarctica. *Scientific Data*, *12*(1), 414. https://doi.org/10.1038/s41597-025-04672-y, 2025.

Purucker, M.: Geothermal heat flux data set based on low resolution observations collected by the CHAMP satellite between 2000 and 2010, and produced from the MF-6 model following the technique described in Fox Maule et al., (2005). Retrieved June 9, 2022, from http://websrv.cs.umt.edu/isis/index.php/Main_Page, 2012.

Pyrcz, M., Jo, H., Kupenko, A., Liu, W., Gigliotti, A. E., Salomaki, T., and Javier, S.: GeostatsPy: Geostatistical Library in Python (Version 1.0.0). https://doi.org/10.5281/zenodo, October, 2021.

Reading, A. M., Stål, T., Halpin, J.A. et al.: Antarctic Geothermal Heat Flow and Its Implications for Tectonics and Ice Sheets. *Nature Reviews Earth and Environment, no. 12: 814–31.* https://doi.org/10.1038/s43017-022-00348-y, 2022.

Rizvi, S., Thapa, M., Fraysher, B., Rapadas, A., Neshani, S., Taylor, D., Lee, L., Dahl-Jensen, D., Gogineni, P., Kolpuke, S., and Chung, C.: Fine-Resolution Mapping of Near-Bed Ice Layers with Ultra-Wideband Radars, in *AGU 2025*, New Orleans, LA, USA, 2025.

Shackleton, C., Matsuoka, K., Moholdt, G., Van Liefferinge, B., and Paden, J.: Stochastic Simulations of Bed Topography Constrain Geothermal Heat Flow and Subglacial Drainage Near Dome Fuji, East Antarctica. *Journal of Geophysical Research: Earth Surface*, *128*(11), e2023JF007269. https://doi.org/10.1029/2023JF007269, 2023.

Shackleton, C.: DomeC-BedElevation [software]. https://github.com/calvinshackleton/DomeC-BedElevation.git, 2026.

Shackleton, C., and Matsuoka, K.: Simulated ice thickness, bed elevation and local topographic adjustments to geothermal heat flow near Dome C, Antarctica [Dataset]. *Norwegian Polar Institute.* https://doi.org/10.21334/NPOLAR.2026.D05BFE78, 2026.

Shen, W., Wiens, D. A., Lloyd, A. J., and Nyblade, A. A.: A Geothermal Heat Flux Map of Antarctica Empirically Constrained by Seismic Structure. *Geophysical Research Letters*, *47*(14), e2020GL086955. https://doi.org/10.1029/2020GL086955, 2020.

Shreve, R. L.: The movement of water in glaciers. *Journal of Glaciology*, *11*(62), 205–214, 1972.

Skolnik, M. I.: *Radar handbook* (3rd ed). New York: McGraw-Hill, 2008.

Steinhage, D., Franke, S., Eisen, O., Miller, H., and Helm, V.: Ice Thickness from East Antarctica, over Dome C, Dome A, Talos Dome and Vostok recorded with the airborne AWI EMR radar system in 2007/2008 [dataset]. PANGAEA, https://doi.org/10.1594/PANGAEA.957066, 2023.

Stål, T., Reading, A. M., Halpin, J. A., and Whittaker, J. M.: Antarctic Geothermal Heat Flow Model: Aq1. *Geochemistry, Geophysics, Geosystems*, *22*(2), e2020GC009428. https://doi.org/10.1029/2020GC009428, 2021.

Stolt, R. H.: Migration by Fourier transform. *Geophysics*, *43*(1), 23–48. https://doi.org/10.1190/1.1440826, 1978.

Sutter, J., Fischer, H., Grosfeld, K., Karlsson, N. B., Kleiner, T., Van Liefferinge, B., and Eisen, O.: Modelling the Antarctic Ice Sheet across the mid-Pleistocene transition – implications for Oldest Ice. *The Cryosphere*, *13*(7), 2023–2041. https://doi.org/10.5194/tc-13-2023-2019, 2019.

Tabacco, I. E., Passerini, A., Corbelli, F., and Gorman, M.: Determination of the surface and bed topography at Dome C, East Antarctica. *Journal of Glaciology*, *44*(146), 185–191. https://doi.org/10.3189/S0022143000002501, 1998.

Taylor, J., Siegert, M. J., Payne, A. J., and Hubbard, B.: Regional-scale bed roughness beneath ice masses: measurement and analysis. *Computers & Geosciences*, *30*(8), 899–908. https://doi.org/10.1016/j.cageo.2004.06.007, 2004.

Thoma, M., Grosfeld, K., Filina, I., and Mayer, C.: Modelling flow and accreted ice in subglacial Lake Concordia, Antarctica. *Earth and Planetary Science Letters*, *286*(1), 278–284. https://doi.org/10.1016/j.epsl.2009.06.037, 2009.

Tikku, A. A., Bell, R. E., Studinger, M., Clarke, G. K. C., Tabacco, I., and Ferraccioli, F.: Influx of meltwater to subglacial Lake Concordia, East Antarctica. *Journal of Glaciology*, *51*(172), 96–104. https://doi.org/10.3189/172756505781829494, 2005.

Uieda, L.: Verde: Processing and gridding spatial data using Green's functions. *Journal of Open Source Software*, *3*(30), 957. https://doi.org/10.21105/joss.00957, 2018.

Van Liefferinge, B., and Pattyn, F.: Using ice-flow models to evaluate potential sites of million year-old ice in Antarctica. *Climate of the Past*, *9*, 2335–2345, 2013.

Van Liefferinge, B., Pattyn, F., Cavitte, M. G. P., Karlsson, N. B., Young, D. A., Sutter, J., and Eisen, O.: Promising oldest ice sites in east antarctica based on thermodynamical modelling. *Cryosphere*, *12*(8), 2773–2787. https://doi.org/10.5194/tc-12-2773-2018, 2018.

van der Veen, C. J., Leftwich, T., von Frese, R., Csatho, B. M., and Li, J.: Subglacial topography and geothermal heat flux: Potential interactions with drainage of the Greenland ice sheet. *Geophysical Research Letters*, *34*(12). https://doi.org/10.1029/2007GL030046, 2007.

Wang, Z., Chung, A., Steinhage, D., Parrenin, F., Freitag, J., and Eisen, O.: Mapping age and basal conditions of ice in the Dome Fuji region, Antarctica, by combining radar internal layer stratigraphy and flow modeling. *The Cryosphere Discussions*, 1–22. https://doi.org/10.5194/tc-2023-35, 2023.

Willcocks, S., D. Hasterok, and S. Jennings.: Thermal Refraction: Implications for Subglacial Heat Flux. *Journal of Glaciology* 67, no. 265: 875–84. https://doi.org/10.1017/jog.2021.38, 2021.

Wingham, D. J., Siegert, M. J., Shepherd, A., and Muir, A. S.: Rapid discharge connects Antarctic subglacial lakes. *Nature*, *440*(7087), 1033–1036. https://doi.org/10.1038/nature04660, 2006.

Winter, A., Steinhage, D., Creyts, T. T., Kleiner, T., and Eisen, O.: Age stratigraphy in the East Antarctic Ice Sheet inferred from radio-echo sounding horizons. *Earth Syst. Sci. Data*, 11, 1069–1081, https://doi.org/10.5194/essd-11-1069-2019, 2019.

Young, D. A., Roberts, J. L., Ritz, C., Frezzotti, M., Quartini, E., Cavitte, M. G. P., et al.: High-resolution boundary conditions of an old ice target near Dome C, Antarctica. *The Cryosphere*, *11*(4), 1897–1911. https://doi.org/10.5194/tc-11-1897-2017, 2017.

Young, D. A., Beem, L. H., Blankenship, D. D., Cavitte, M., Greenbaum, J., Habbal, F., et al.: ICECAP: High resolution survey of the Little Dome C region in support of the IPICS Old Ice goal. *U.S. Antarctic Program (USAP) Data Center*. doi: https://doi.org/10.15784/601355, 2020.

Xiangbin, C., Jeofry, H., Greenbaum, J., Roberts, J., Blankenship, D., Bo, S., and Siegert, M.: ICECAP-2 consortium processed airborne ice thickness data from the Princess Elizabeth Land, East Antarctica [Data set]. Zenodo. https://doi.org/10.5281/zenodo.4023393, 2020.

Appendix A

Table A1: *Radar-derived ice thickness datasets used. BAS = British Antarctic Survey, BE-OI = Beyond EPICA Oldest Ice, KOPRI = Korea Polar Research Institute, UA = University of Alabama, DTU = The Technical University of Denmark, CPH = University of Copenhagen, AWI = Alfred Wegener Institute, AAD = Australian Antarctic Division, INGV = Istituto Nazionale di Geofisica e Vulcanologia, UTIG = University of Texas Institute for Geophysics, NASA = National Aeronautics and Space Administration, PRIC = Polar Research Institute of China, LDEO = Lamont Doherty Earth Observatory, CReSIS = Center for Remote Sensing of Ice Sheets. *All surveys had data filtered out within 0.6 km of crossover sites if differences were > 50 m, with priority determined by mean absolute differences between all other surveys.*

#	Name	Institution	Survey/Project	System	Source/DOI/Key reference	Filtering Notes*
1	AWI_2007_ANTR_AIR_BM2	AWI	Dronning Maud Land; Princess Elizabeth Land; Dome A; Dome C	AWI EMR	https://doi.org/10.1594/PANGAEA.957066 https://doi.org/10.5194/essd-11-1069-2019	
2	BAS_2005_WISE-ISODYN_AIR_BM2	BAS	Wilkes Subglacial Basin (WISE-ISODYN)	PASIN	https://doi.org/10.5285/59e5a6f5-e67d-4a05-99af-30f656569401 https://doi.org/10.5194/essd-14-3379-2022	Filtered out data within 0.2 km of surveys #9 and #11 due to inconsistencies.
3	BEDMAP1_1966-2000_AIR_BM1	*Various*	*Various*	*Various*	https://doi.org/10.5194/essd-15-2695-2023	Filtered out data along 5 manually supplied lines due to inconsistencies with surrounding data. Also filtered out data within 10 km of more recent data.
4	BEOI_2016-2018_LDC_DELORES	BAS	BE-OI	DELORES	https://doi.org/10.1594/PANGAEA.963474	Filtered out data along 1 manually supplied line due to inconsistencies with surrounding data.
5	BEOI_2019_LDC_VHF	UA/CPH/AWI	BE-OI	VHF	https://doi.org/10.1594/PANGAEA.957461 https://zenodo.org/records/4687049	
6	DTU_2024	DTU	ICESAR2024 & ICECAP	POLARIS	-	
7	INGV_1997_ITASE_AIR_BM2	INGV	Talos Dome; Oats Land (ITASE)	INGV-IT GlacioRadar	https://doi.org/10.3189/172756404781814591	Filtered out data within 0.2 km of surveys #8, #9, #10 due to duplicated data.
8	INGV_1997_Talos-Dome_AIR_BM3	INGV	Talos Dome; Terre Adelie	INGV-IT GlacioRadar	https://doi.org/10.4401/ag-3471	
9	INGV_1999_Talos-Dome_AIR_BM3	INGV	Talos Dome; Terre Adelie	INGV-IT GlacioRadar	https://doi.org/10.4401/ag-3471	
10	INGV_2001_Talos-Dome_AIR_BM3	INGV	Talos Dome; Oats Land	INGV-IT GlacioRadar	https://doi.org/10.5194/essd-15-2695-2023	
11	INGV_2003_Talos-Dome_AIR_BM3	INGV	Talos Dome; Queen Mary Land; Terre Adelie	INGV-IT GlacioRadar	https://doi.org/10.5194/essd-15-2695-2023	
12	KOPRI_UA_2023	KOPRI/UA	Belgica Highlands	UWB IPR	-	Filtered out data along 1 manually supplied line due to inconsistencies with surrounding data.
13	NASA_2013_ICEBRIDGE_AIR_BM3	NASA/ CReSIS	Operation Icebridge	MCoRDS v3	https://doi.org/10.1029/2020RG000712	
14	PRIC_2018_CHA4_AIR_BM3	PRIC	Princess Elizabeth Land (CHA4)	HiCARS	https://doi.org/10.5281/zenodo.4023393 https://doi.org/10.5194/essd-12-2765-2020	
15	UTIG_1999_SOAR-LVS-WLK_AIR_BM2	UTIG/LDEO	Transantarctic Mountains; South Pole; Lake Vostok; Dome C	SOAR TUD IV (RADgh1 digitizer)	http://dx.doi.org/10.1016/S0012-821X(04)00066-4 https://doi.org/10.15784/601588	Filtered out data along 1 manually supplied line due to inconsistencies with surrounding data.
16	UTIG_2008_ICECAP_AIR_BM2	UTIG/AAD	ICECAP/ Aurora Subglacial Basin	HiCARS	https://www.usap-dc.org/view/dataset/601411 https://doi.org/10.5067/F5FGUT9F5089 http://dx.doi.org/10.1038/nature10114	
17	UTIG_2010_ICECAP_AIR_BM3	UTIG/AAD	ICECAP/Antarctic-wide	HiCARS	https://doi.org/10.5067/W2KXX0MYNJ9G https://doi.org/10.1038/nature10114	Filtered out data within 0.2 km of survey #16 due to inconsistencies. Also filtered out data along 1 manually supplied line due to inconsistencies with surrounding data.
18	UTIG_2016_OLDICE_AIR_BM3	UTIG/AAD	ICECAP/ Dome C	MARFA	https://doi.org/10.15784/601355 https://doi.org/10.1098/rsta.2014.0297	

Figures A1-18: *Consistency between individual radar surveys and all other surrounding data. Each figure shows a map with 50 m radius crossover points as an X coloured according to the difference in value (Red-Blue colormap). Survey track is shown as a grey line and all other survey tracks are shown as yellow lines. The adjacent histograms show the differences in value between each intersecting survey as a probability density function, with the upper plot showing detailed differences within ±150 m and lower plot showing the full range of crossover differences. A legend for the histograms is shown to the right with some summary statistics for overall differences between the survey and surrounding measurements*

Figure A1




Figure A2




Figure A3





Figure A4



1

Figure A5




Figure A6




Figure A7





Figure A8



2

Figure A9




Figure A10




Figure A11





Figure A12



3

Figure A13




Figure A14




Figure A15





Figure A16



4

Figure A17




Figure A18

