

Saturation effect of background temperature and aridity on vegetation phenological sensitivity to urban warming

Zhang Zhenzhen^{1#}, Hu Xinxin^{1#}, Zhang Yongqi¹, Cui Shufen², Sun Liheng³, Lin Xingwen^{1*}, Wu Chaofan¹, Zhang Zhaoyang¹, Chen Yuanjian¹, Wen Qingqing⁴

¹College of Geography and Environmental Sciences, Zhejiang Normal University, Jinhua 321004, China

²College of Business, Lishui University, Lishui 323200, China

³Faculty of Management, Universiti Teknologi Malaysia, Johor Bahru 81310, Malaysia

⁴The Administration Center of Jinhua Wucheng Nanshan Provincial Nature Reserve, Jinhua 321000, China

* Correspondence: Lin Xingwen (linxw@zjnu.edu.cn)

These authors contributed equally to this work

Abstract:

Quantifying how background temperature and aridity constrain phenological sensitivity to urban warming requires identifying critical thermal thresholds and hydrological boundaries driving spatial heterogeneity. This study assessed urban warming effects on vegetation phenology across 293 Chinese cities during 2010–2020. Urban-rural disparities (Δ SOS, Δ EOS) and temperature sensitivity (R_{t-SOS} , R_{t-EOS}) were quantified using satellite metrics. Results showed that urban heat islands advanced SOS by 12.06 days and delayed EOS by 9.86 days relative to rural areas. A non-linear saturation effect was detected: phenological sensitivity to urban warming peaked at 4 °C (spring) and 6 °C (autumn), and significantly weakened beyond sensitivity saturation thresholds of 12.5 °C (SOS) and 4 °C (EOS). Aridity acted as a

删除[胡欣欣]: *

删除[胡欣欣]: ,

删除[胡欣欣]: Zhang Zhenzhen

设置格式[胡欣欣]: 超链接, 字体: 小四, 加粗

删除[胡欣欣]: zhangzhen@zjnu.cn

删除[胡欣欣]: Societal Impact Statement

As cities grow and climates warm, the Urban Heat Island (UHI) effect increasingly alters urban ecosystems. This study analyzed 293 Chinese cities to reveal how baseline temperature and aridity shape vegetation’s response to UHI. We found a "saturation effect," where UHI’s impact weakens under certain background conditions. This helps explain why urban greening outcomes vary regionally. Our findings offer new insight into urban ecology and can support climate-smart planning, helping cities adapt more effectively and sustainably.

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C

negative regulator that weakened warming benefits across most aridity index (AI) ranges and reversed them within $1.4 < AI < 2.0$. Combined, LST and AI explained 75.05% and 76.21% of spatial variance in SOS and EOS responses, respectively. Conceptually, these findings challenge the linear paradigm of warming-driven phenological shifts by demonstrating finite physiological benefits of urban warming. Ecologically, this study highlights the coupled heat-aridity constraints on urban vegetation, providing key implications for climate-adaptive urban ecological planning.

Keywords: urbanization; heat island effect; vegetation phenological sensitivity; aridity index; land surface temperature; saturation effect

1. Introduction

Vegetation phenology, a critical aspect of ecosystem dynamics, exhibits significant sensitivity to the coupling of thermal and moisture variations (Badeck et al., 2004; Yu et al., 2003). While recent studies have shown that urban warming advances the start of the growing season (SOS) and delays the end of the growing season (EOS), it is increasingly recognized that these responses are not solely temperature-driven but are fundamentally constrained by the background aridity of the locale. Thus, the phenological "benefits" of the Urban Heat Island (UHI) effect must be evaluated through a dual lens of heat and water availability (Jeong et al., 2019). The cascading effects of these shifts extend beyond mere phenological alterations, contributing to a range of ecological and environmental challenges (Zhou et al., 2016; Qiu et al., 2017; Ding et al., 2020).

However, the degree to which warming benefits phenological changes has been widely debated (Chmielewski and Rötzer, 2001; Ding et al., 2020; Yao et al., 2017). Urban environments offer a unique "natural laboratory" for testing these phenological theories. Unlike experimental warming plots, the urban heat island (UHI) effect represents a sustained, large-scale warming scenario that has persisted for decades (Grimm et al., 2008). This allows researchers to observe long-term vegetation adaptation and identify thresholds that are difficult to capture in short-term

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四
删除[胡欣欣]: investigations have highlighted how
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四
删除[胡欣欣]: precipitates an earlier
删除[胡欣欣]: a
删除[胡欣欣]: ed
删除[胡欣欣]: '
删除[胡欣欣]: '
删除[胡欣欣]: from the outset
删除[胡欣欣]: spectrum
删除[胡欣欣]: &
删除[胡欣欣]: '
删除[胡欣欣]: '

manipulative experiments. Moreover, the vast geographic distribution of cities across climatic gradients provides a natural experimental framework to test how identical warming magnitudes (UHI) interact with varying background climates (Fig. 1). Empirical evidence from the Northern Hemisphere consistently shows that cities in cold high-latitude regions exhibit more advanced SOS and delayed EOS in response to urban warming (Dallimer et al., 2016; Gazal et al., 2008; Jia et al., 2021; Li et al., 2021; Meng et al., 2020). While these studies offer valuable evidence regarding spatial heterogeneity, the non-linear dynamics of these responses—specifically, the saturation thresholds where urban warming benefits diminish, and how background aridity modulates these thresholds—remain inadequately quantified across climatic gradients. Identifying such thresholds is critical for predicting urban ecosystem stability under continuous global warming scenarios.

删除[胡欣欣]: Figure

The magnitude of urban-induced phenological shifts is not uniform. Background climate heterogeneity constrains vegetation responses to additional urban warming: phenological advancements are strong in cold regions but weaken as baseline temperatures rise, implying that phenological sensitivity is a variable state conditioned by local thermal conditions. In ecology, the "saturation effect" describes that stimulus efficacy declines beyond a threshold, rooted in tolerance curve theory (Angilletta, 2009): organismal physiological functions increase with environmental drivers until reaching physiological saturation, beyond which no further enhancement occurs (Ketola and Kristensen, 2017). Plant physiological processes (photosynthesis, respiration, transpiration) do not increase linearly with temperature; they slow down or even decrease beyond a thermal threshold (Sage and Kubien, 2007; Ge et al., 2021). Meta-analysis has shown greater positive growth responses to warming in colder ecosystems (Rustad et al., 2001; Büntgen et al., 2019). We therefore hypothesize that urban warming benefits for phenology (advanced SOS, delayed EOS) gradually approach saturation in warmer regions, and extra UHI warming will no longer promote phenological shifts. This hypothesis remains to be tested in urban ecosystems. Drought disrupts temperature-driven phenological responses (Peng et al., 2019;

删除[胡欣欣]: &

删除[胡欣欣]: &

删除[胡欣欣]: A m

删除[胡欣欣]: has proved

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: This theory remains to be verified in urban phenology systems.

删除[胡欣欣]: It is noteworthy that d

删除[胡欣欣]: conditions adversely disrupt

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

85 Yuan et al., 2020). Urban vegetation is particularly vulnerable to moisture deficits due
86 to the "Urban Dry Island" effect (reduced soil moisture and elevated evaporation in
87 built environments) and impervious surfaces (Zhang et al., 2019). UHI warming
88 combined with high aridity elevates vapor pressure deficit and water demand,
89 triggering defensive mechanisms (such as ABA accumulation and protein degradation)
90 (Brodrigg and McAdam, 2013), that offset thermal advancements. Drought delays
91 spring SOS and advances autumn EOS by inducing water stress and early dormancy
92 (Brodrigg and McAdam, 2013; Čehulić et al., 2019). However, in urban environments,
93 this drought effect is compounded by the Urban Dry Island and impervious surfaces,
94 further attenuating phenological sensitivity to extra warming, diminishing the positive
95 impacts of UHI; this effect is stronger in warmer cities where plants approach thermal
96 limits.

97 Consequently, we hypothesize that background aridity (AI) acts as a critical
98 negative regulator of the thermal saturation threshold. Specifically, we predict that as
99 background aridity increases, the "saturation point" at which warming benefits (such
100 as advanced SOS or delayed EOS) plateau will occur at lower LST values. This
101 implies that the saturation effect is not a fixed thermal threshold but a dynamic state
102 conditioned by water stress, leading to a more rapid decline in phenological
103 sensitivity in arid urban environments.

104 Furthermore, phenological sensitivity to environmental drivers is species-specific
105 and modulated by functional traits. Broad-leaf and coniferous species show divergent
106 responses to warming and drought due to differences in leaf morphology and
107 hydraulic architecture (Zheng et al., 2022). However, whether the saturation effect
108 and "aridity-driven inhibition" are consistent across urban vegetation types remains
109 unclear.

110 As a rapidly urbanizing country with diverse climates, China faces severe UHI
111 challenges (He et al., 2017). UHI intensity averages 0.9 ± 1.1 °C but varies drastically
112 across regions (Li et al., 2021). Previous urban phenology studies focused on linear
113 temperature sensitivity, ignoring non-linear saturation effects and aridity interactions.
114 No study has quantified critical thermal/arid saturation thresholds or clarified the

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: more

删除[胡欣欣]: '

删除[胡欣欣]: '

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: e.g., ABA accumulation

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: McAdam

删除[胡欣欣]: , protein degradation)

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 自动设置

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 自动设置

删除[胡欣欣]: McAdam

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: operate

删除[胡欣欣]: ly

删除[胡欣欣]: the interactive regulation of

删除[胡欣欣]: , n

coupling mechanism between pre-season temperature (T, phenological sensitivity) and land surface temperature (LST, background climate) in urban ecosystems. China's large climatic gradients and rapid urbanization make it an ideal platform to test the saturation hypothesis and aridity modulation.

删除[胡欣欣]: ,

Thus, an investigation into the phenological response of plants to urbanization in China would enhance our understanding of their ability to acclimate to climate change across various environments. However, how background temperature and aridity interact to determine the thresholds and "saturation" of phenological sensitivity to urban warming remains largely unquantified at the macro-scale. Addressing this gap is essential for predicting the stability of urban carbon sinks under future climate scenarios. To this end, the primary objectives of this study were to: (1) explore the spatial distribution of plant phenological responses to the urban heat island effect throughout mainland China; (2) quantify the saturation effect caused by the spatial variability of phenological responses correlated with background temperature across the cold-hot gradient; and (3) assess whether the benefits of urban warming would be counteracted in drought-affected areas.

删除[胡欣欣]: s

删除[胡欣欣]: ively

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: a

删除[胡欣欣]: encompassed the followin

删除[胡欣欣]: g

2. Materials and Methods

2.1 Study area

In our study, we conducted an analysis covering a total of 293 prefecture-level cities in China (excluding Taiwan, Autonomous Prefecture, and regions), taking into account the data from the year 2020 (<http://xzqh.mca.gov.cn/map>) (Fig. 1). It was hypothesized that the urban warming effect is most pronounced in urban centers and gradually diminishes towards rural areas (Zhang et al., 2004). To determine the urbanization intensity of each city, nighttime light data from NPP-VIIRS (Table 1) were utilized (Elvidge et al., 2017), employing the reference comparison method. The optimal segmentation threshold was established based on the statistical yearbook (Fan et al., 2019). Moreover, urban built-up areas were identified using the comparison method and annual average data derived from NPP-VIIRS night lights between 2013 and 2020. All data processing procedures were performed using ArcGIS 10.2. We

删除[胡欣欣]: Figure

删除[胡欣欣]: as

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

created ten buffer zones extending to a radius of 20 km from each city center, with each zone representing a 2 km annulus (Fig. 1). This 20 km boundary was chosen because our preliminary analysis revealed that the urban warming footprint on phenological metrics (SOS and EOS) gradually diminishes to near zero beyond 15 km (Zhang et al., 2004; Figs. 1 and 2). Thus, defining the outermost 20 km buffer zone as the "Rural" baseline guarantees a background environment conservatively free from urban heat island interference. The subscript (10) refers to the 10th buffer zone at 20 km from the urban center. Equation notations use a standardized buffer index (0 for urban center, 10 for rural reference) to avoid confusion between distance and buffer number.




Figure 1. City centers (red area) and buffer zones (blue area) of the 293 prefecture-level cities in China. The map, generated using ArcGIS version 10.2 (<https://www.esri.com/about/newsroom/arcwatch/the-best-of-arcgis-10-2/>), features

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: A total of ten buffer zones were created within a radius of 20 kilometers from the city center, with each buffer zone set at a distance of 2 kilometers from the center

删除[胡欣欣]: Figure

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: This specific 20 km delineation was established

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: and approaches zero at distances of 15 km and beyond

删除[胡欣欣]: ure

删除[胡欣欣]: , Figure

删除[胡欣欣]: s

删除[胡欣欣]: "

删除[胡欣欣]: "

删除[胡欣欣]: used throughout this manuscript

删除[胡欣欣]: -

删除[胡欣欣]: located

删除[胡欣欣]: away

删除[胡欣欣]: , and all notations in equations have been standardized to use

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: misinterpretation

删除[胡欣欣]: -

administrative boundaries sourced from the Ministry of Civil Affairs of the People's Republic of China (<http://xzqh.mca.gov.cn/map>). Source: Esri, TomTom, FAO, USGS, and the GIS User Community; Ministry of Civil Affairs of the People's Republic of China | Powered by Esri.

2.2 Vegetation phenology extraction

Phenology data spanning the years 2010 to 2020 were derived from MODIS Enhanced Vegetation Index (EVI) 16-day composite data with a spatial resolution of 250 m (MOD13Q1) and processed ~~with~~ TIMESAT (Jönsson ~~and~~ Eklundh, 2004). To mitigate challenges such as cloud cover and snow, the raw EVI time series data were smoothed using the Savitzky-Golay (S-G) filter method (Cai et al., 2017). Subsequently, a dynamic threshold technique was employed to extract the phenology data, utilizing a seasonal amplitude of 20%, a parameter proven effective for phenology extraction in urban environments (Zhou et al., 2016; Buyantuyev ~~and~~ Wu, 2012; Cong et al., 2012; White et al., 2009). The datasets underwent preprocessing and quality control to eliminate pixels with minimal vegetation EVI (e.g., bodies of water and fully paved surfaces). To exclude unrealistic phenological values, we retained only SOS (30–180 days) and EOS (210–360 days) within valid ranges, consistent with previous studies (Zhang et al., 2006). Efforts were made to reduce the impact of outlier years by averaging the phenology data for each pixel across the timeframe from 2010 to 2020. After applying these quality control thresholds and land-cover filtering criteria, approximately 48.5% of the initial pixels within the study buffer zones were excluded.

Furthermore, we employed land cover data from the MCD12Q1 dataset, which has a 500-meter resolution, to mitigate the impact of agricultural practices on the mainland during our analysis. The MCD12Q1 dataset categorizes land into 17 distinct classes. These include various types of forested areas such as Evergreen Coniferous Forests, Deciduous Needle-leaf Forests, Evergreen Broad-leaf Forests, Deciduous Broad-leaf Forests, Mixed Forests, as well as Closed Shrub lands, Open Shrub lands, Savannas, Woody Savannas, Grasslands, and Croplands.

删除[胡欣欣]: utilizing

删除[胡欣欣]: &

删除[胡欣欣]: &

Given the frequent human-induced disturbances in rural croplands, and the regular manual pruning of urban grasslands, we excluded savannas, woody savannas, grasslands, and croplands from our analysis. The remaining land types were consolidated into broader categories: broad-leaf forests (BF, deciduous and evergreen broadleaf forests), coniferous forests (CF, deciduous and evergreen coniferous forests), mixed forests (MF), and shrubs (closed and open shrublands). This classification approach allowed us to focus our analysis on land types that are less influenced by human activities.

删除[胡欣欣]: C
删除[胡欣欣]: in the rural regions
删除[胡欣欣]: G
删除[胡欣欣]: in the urban center
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四
删除[胡欣欣]: Savannas, woody savannas, Grasslands, and Croplands were excluded from our study.

Table 1. Data inventory used in this study

Data type	Resolution	Layer	Data sources	Year
MOD13Q1	250m/16d	EVI	https://search.earthdata.nasa.gov	2010-2020
MOD11A2	1000m/8d	LST_Day LST_Night	https://search.earthdata.nasa.gov	2010-2020
MOD16A2	500m/8d	PET	https://search.earthdata.nasa.gov	2010-2020
MCD12Q1	500m	Land Cover	https://search.earthdata.nasa.gov	2010-2020
VIIRS	500m	yearly average	https://eogdata.mines.edu	2013-2020
Precipitation data	1000m	monthly average	http://www.geodata.cn	2010-2020
China's urban zoning data			https://www.resdc.cn	2020
Station climates		monthly average	http://data.cma.cn/en	2010-2020

2.3 Pre-season temperature and background climates

Building upon the methodology of a previous study (Piao et al., 2006), we defined pre-season temperature (T), as the mean temperature for the three months preceding phenological onset. For alignment with the phenology information, the average temperature dataset was resampled to a 250 m resolution using the nearest neighbor interpolation technique. To evaluate the performance of the downscaling procedure, the observed long-term monthly temperature and precipitation across China were obtained from the National Meteorological Information Center of China (<http://data.cma.cn/en>). This dataset included observations from 496 national weather

删除[胡欣欣]: was characterized
删除[胡欣欣]: the onset of the
删除[胡欣欣]: season

stations during 2010~~to~~ 2020. Upon conducting a fitting analysis with station data, we observed that the coefficient of determination (R^2) for monthly mean temperature and remote sensing data reached 0.93, while that for monthly precipitation and remote sensing data achieved 0.85. These results indicated that the downscaled meteorological data sufficiently met the requirements of this study.

删除[胡欣欣]: -

删除[胡欣欣]: e

The background climates were evaluated utilizing land surface temperature (LST) and the aridity index (AI) as indicators. The LST and AI data for each pixel were averaged over 2010~~to~~ 2020. Specifically, MODIS LST datasets exhibiting an absolute bias of less than 1_K were selected as optimal for our investigation, in line with previous studies (Gow et al., 2016; Pablos et al., 2016; Qiao et al., 2013; Wan, 2008). Collection 6 LST products from the Aqua satellite (MOD11A2, 8-day composite, 1 km spatial resolution, covering the period from 2010 to 2020) were used. Additionally, daily temperature (T) was calculated as the mean of daytime and nighttime temperatures, following the methodology outlined by Kang et al., (2016). AI was expressed as:

删除[胡欣欣]: -

删除[胡欣欣]: et al.

删除[胡欣欣]: Kang (

删除[胡欣欣]: ,

删除[胡欣欣]:

设置格式[胡欣欣]: 字体: (默认) Times New Roman

设置格式[胡欣欣]: 字体: (默认) Times New Roman

设置格式[胡欣欣]: 字体: (默认) Times New Roman

设置格式[胡欣欣]: 字体: (默认) Times New Roman

设置格式[胡欣欣]: 字体: (默认) Times New Roman

where PET denotes the annual potential evapotranspiration and P denotes the annual precipitation. The PET values spanning from 2010 to 2020 were extracted from the MOD16A2 monthly synthetic product, characterized by an 8-day temporal resolution and a spatial resolution of 500 m (Mu et al., 2011), as delineated in Table 1. The annual precipitation (P) dataset was derived from the monthly precipitation records of China encompassing the timeframe from 1901 to 2020, ~~at~~ a spatial resolution of 1 km. Subsequently, both PET and P datasets were resampled to a spatial resolution of 250 m to facilitate the ensuing spatial analysis (Peng, 2020).

删除[胡欣欣]: possessing

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: clarify

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: Pre-season air temperature (T) is the direct biological driver of plant physiology

To robustly quantify these responses, we ~~defined~~ the distinct analytical roles of two critical thermal dimensions: pre~~-~~season air temperature (T) and land surface temperature (LST). In this study, T and LST serve strictly distinct analytical roles. ~~As the direct biological driver of plant physiology~~ (e.g., chilling requirements), T ~~was~~ used exclusively to quantify the pure biological phenological sensitivity (R_d) and identify true physiological thresholds. ~~In contrast, as the standard metric for~~

删除[胡欣欣]: . Therefore

删除[胡欣欣]: i

设置格式[胡欣欣]: 下标

删除[胡欣欣]: Conversely

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

235 quantifying the urban heat island (UHI) effect, LST was used to characterize the
236 spatial "background climate" of each city across the continent.

237 Because pre-season air temperature (T) is the direct driver of phenological
238 sensitivity, we first derived the biological thresholds using T (Fig. 5). However, to
239 address our core objective of quantifying how background LST shapes these
240 phenological shifts, we defined the quantitative relationship between the background
241 LST and T. We fitted the relationship curve between LST and T across all cities. To
242 minimize data matching biases during this conversion, we used the spatiotemporally
243 aligned 10-year climatological means (2010–2020) for both variables at a unified 250
244 m resolution. We observed a highly robust linear correlation (Fig. S4). Based on this
245 high goodness-of-fit, we applied the empirical conversion formula between the two
246 variables to mathematically substitute T with LST in the sensitivity curves (Wan,
247 2008). This empirical translation allowed us to systematically convert the T-driven
248 physiological thresholds into the specific LST-determined background climate
249 thresholds (12.5 °C for SOS and 4 °C for EOS), explicitly projecting pure biological
250 limits onto the urban map. Although T and LST represent different physical properties,
251 they are highly correlated in urban settings ($R^2 = 0.93$ for monthly mean temperature),
252 confirming the reliability of this empirical conversion between physiological response
253 and climatic background.

254 2.4 Urbanization effect on phenology

255 The phenology dates of each vegetation type were extracted in each buffer zone.
256 Finally, the mean phenology in each buffer zone was calculated as a weighted mean of
257 the phenology of each vegetation type. To ensure comparability in phenological
258 analyses across buffer zones, we first identified the common vegetation types present
259 in all zones and excluded the non-shared types. We then calculated the phenological
260 means for these common types within each buffer zone and averaged these values to
261 determine the overall mean vegetation phenology for the entire buffer region, as
262 shown in Eqs. (2) and (3).

$$SOS_i = \sum SOS_{ij} / n \quad (2)$$

删除[胡欣欣]: Land Surface Temperature (LST) is the ...

删除[胡欣欣]: i

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: Figure

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: —understanding how the background ...

删除[胡欣欣]: strictly prevent

删除[胡欣欣]: utilized

删除[胡欣欣]: -

删除[胡欣欣]: Figure

删除[胡欣欣]: utilized

删除[胡欣欣]: et al.

删除[胡欣欣]: While

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: ensuring the reliability of this empirically ...

删除[胡欣欣]: was

删除[胡欣欣]: this

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: weighted by

删除[胡欣欣]: one

删除[胡欣欣]: below (e

删除[胡欣欣]: uation

删除[胡欣欣]: ,

删除[胡欣欣]:)

设置格式[胡欣欣]: 字体: 倾斜

设置格式[胡欣欣]: 字体: 倾斜

设置格式[胡欣欣]: 字体: (默认) Times New Roman

设置格式[胡欣欣]: 字体: (默认) Times New Roman, 倾斜

设置格式[胡欣欣]: 字体: (默认) Times New Roman, 倾斜

设置格式[胡欣欣]: 字体: (默认) Times New Roman, 倾斜

设置格式[胡欣欣]: 字体: (默认) Times New Roman

$$EOS_i = \sum EOS_{ij} / n \quad (3)$$

SOS_i and EOS_i are the mean SOS and EOS of the i-th buffer zone; SOS_{ij} and EOS_{ij} are the mean SOS and EOS of the j-th vegetation type in the i-th buffer zone; n is the number of shared vegetation types across all buffer zones.

To investigate the impact of urbanization on phenology, mean values of start of season (SOS) and end of season (EOS) in each buffer zone were analyzed during 2010–2020, and fitted with respect to their distance from the urban center. In this study, the outermost buffer zone (20 km from the urban center) was defined as the rural reference region, and the urban center was defined as the urban region. This delineation was justified by the observation that phenological change rates approached zero at 15 km and beyond (Fig. S1), and similar patterns were observed for all four vegetation types (Fig. S2). Following prior investigations, the phenological sensitivity (R_t) to temperature was quantified by calculating the partial correlation coefficient between T and the phenology dates (R_{t-SOS} and R_{t-EOS}) (Meng et al., 2020). The phenological difference between urban and rural regions was calculated to quantify the magnitude of the urban warming effect on phenology:

$$\Delta SOS = SOS_{(0)} - SOS_{(10)} \quad (4)$$

$$\Delta EOS = EOS_{(0)} - EOS_{(10)} \quad (5)$$

where SOS₍₀₎ and SOS₍₁₀₎ are the mean SOS of the urban center (buffer index 0) and the 10th buffer zone (located 20 km away), respectively. Similarly, EOS₍₀₎ and EOS₍₁₀₎ are the mean EOS of the urban center and the 10th buffer zone. Biologically, a negative ΔSOS indicates an advanced start of the season (spring) in the urban center relative to the rural area, while a positive ΔEOS indicates a delayed end of the season (autumn senescence) in the urban center.

Similarly, the difference in phenological sensitivity to temperature between urban and rural regions was defined as:

$$\Delta R_{t-SOS} = R_{t-SOS(0)} - R_{t-SOS(10)} \quad (6)$$

$$\Delta R_{t-EOS} = R_{t-EOS(0)} - R_{t-EOS(10)} \quad (7)$$

where R_{t-SOS(0)} and R_{t-SOS(10)} are the mean R_{t-SOS} of the urban center (buffer index 0)

287 and the 10th buffer zone (located 20 km away), respectively. Similarly, $R_{t-EOS(0)}$ and
288 $R_{t-EOS(10)}$ are the mean R_{t-EOS} of the urban center and the 10th buffer zone.

289 2.5 Background climates determined phenology response

290 ~~To~~ elucidate the distinct impacts of LST and AI on urban and rural components,
291 ~~we performed~~ a partial correlation analysis. Specifically, the average LST or AI values
292 ~~for~~ 2010 to 2020 for each of the 293 cities were correlated with their respective Δ SOS,
293 Δ EOS, ΔR_{t-SOS} , and ΔR_{t-EOS} values, while ~~we~~ ~~controlled~~ ~~ed~~ for the influence of the other
294 climatic factor.

295 ~~Path analysis (PA), (Doncaster, 2007) was used to~~ examine both the direct and
296 indirect effects of background climate variables (LST and AI) on phenological
297 responses (SOS and EOS), as well as their corresponding discrepancies (Δ SOS and
298 Δ EOS) and rates of change (ΔR_{t-SOS} and ΔR_{t-EOS}).

299 In the path analysis, ~~we first assessed model fit using fit indices,~~ including the
300 chi-square (χ^2) statistic, Comparative Fit Index (CFI), and Root Mean Square Error of
301 Approximation (RMSEA). ~~Then,~~ path coefficients were derived for both the direct
302 and indirect effects of each independent variable on the dependent variable from the
303 model. ~~Squared path coefficients were used to,~~ estimate the respective direct and
304 indirect contributions of each independent variable to the dependent variable. The
305 ~~sum~~ of these direct and indirect contributions ~~represents~~ the total contribution degree,
306 or relative contribution.

307 2.6 Validation of phenology "saturation effect"

308 To determine if there was a temperature saturation effect on vegetation
309 phenology, we conducted a ~~model comparison~~. Rather than assuming a single
310 relationship a priori, we evaluated the fit of phenology versus pre-season temperature
311 using competing models: a linear model, a quadratic model, a segmented (piecewise)
312 model, and a logistic model. We specifically selected the logistic function as our
313 primary model because quadratic models biologically imply a symmetric reversal (a
314 continuous steep decline) rather than a plateau, and piecewise models assume an

删除[胡欣欣]: In order to

删除[胡欣欣]: was executed

删除[胡欣欣]: recorded from

删除[胡欣欣]: ing

删除[胡欣欣]: This study employed p

删除[胡欣欣]: as a methodological approach

删除[胡欣欣]: to

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: we initially ascertained the model's
congruence with empirical data by evaluating fit indices,

删除[胡欣欣]: Subsequently

删除[胡欣欣]: The s

删除[胡欣欣]: for the direct and indirect

删除[胡欣欣]: paths

删除[胡欣欣]: d

删除[胡欣欣]: degrees

删除[胡欣欣]: aggregate

删除[胡欣欣]: degrees

删除[胡欣欣]: constitutes

删除[胡欣欣]: comparative

abrupt, angular change in biological rates, which is less physiologically realistic than the smooth, asymptotic transition captured by the logistic curve (Pinheiro and Bates, 2000). The asymptote of the logistic function mathematically represents the biological "saturation point"—the maximum physiological limit of thermal benefits.

删除[胡欣欣]: &

删除[胡欣欣]: '

删除[胡欣欣]: '

Statistically, the model comparison results (Table 2) confirm that the logistic model consistently outperforms the other three formulations for both spring SOS and autumn EOS, providing quantitative support for the non-linear saturation effect interpretation. For SOS, the logistic model yields an AIC value 2778 units lower than the linear model and 1424 units lower than the piecewise model, along with an R^2 of 0.37, which is approximately 19% higher than the second-best piecewise model. For EOS, the logistic model still achieves the lowest AIC and highest R^2 , but the performance improvement is more modest: its R^2 (0.40) is only marginally higher than that of the piecewise model (0.39), consistent with a weaker but stable saturation pattern.

Table 2. Performance comparison of four competing models describing phenological responses to pre-season temperature

Phase	Model	AIC	BIC	R^2	RMSE
Spring_SOS	Linear	104203.62	104218.97	0.25	26.52
	Quadratic	103636.33	103659.35	0.27	26.05
	Piecewise	102849.67	102880.36	0.31	25.41
	Logistic	101425.33	101456.02	0.37	24.29
Autumn_EOS	Linear	93252.94	93268.34	0.23	17.39
	Quadratic	90241.5	90264.6	0.36	15.86
	Piecewise	89309.4	89340.2	0.39	15.41
	Logistic	85678.49	85709.14	0.40	15.22

删除[胡欣欣]: R^2

We then extracted phenological information from all vegetation pixels across the country and generated scatter plots of phenology versus pre-season temperature. Next, we fitted logistic growth curves to these scatter plots and evaluated the assumption using the goodness-of-fit metrics, p-value and R^2 . Subsequently, we applied the same

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: plotted scatter

删除[胡欣欣]: diagrams

删除[胡欣欣]: R^2

logistic curve to characterize the phenological responses of urban and suburban vegetation in 293 different cities against their pre-season temperatures, testing whether the phenological response to temperature in urban areas also follows a logistic growth pattern. Finally, we replaced the pre-season temperature with the background temperature LST to determine how the phenological saturation effect is regulated by the background temperature.

3. Results

3.1 Spatial distribution of urban-rural phenological differences

Among the 293 cities analyzed, 84% (N = 246) exhibited an advanced SOS in urban areas compared to rural regions, while 82% (N = 241) showed a delayed EOS (Fig. S3). Spatially, cities located in high-latitude regions generally demonstrated more negative Δ SOS values than those in low-latitude regions (Fig. 2), indicating that urban warming advanced spring phenology most strongly, in northern and western China (Fig. 2a,b). This pattern is consistent with the conventional understanding of earlier SOS in colder regions. Additionally, cities in coastal provinces exhibited larger Δ SOS values than inland cities (Fig. 2).

A strong positive correlation was observed between Δ SOS and the rural rate of SOS change (R_{t-SOS}), suggesting that cities with more negative R_{t-SOS} values exhibited larger urban-rural disparities (Fig. 3). Notably, cities with lower temperatures tended to have more negative R_{t-SOS} and Δ SOS values.

Δ EOS also showed a significant increasing trend with latitude (Fig. 2), with northern cities displaying more positive Δ EOS values, indicating a delayed autumn phenology in urban areas (Fig. 2c,d). In contrast, the majority of cities in southern regions demonstrated negative Δ EOS values, signifying an advancement in autumn phenology. Furthermore, cities in the coastal provinces also exhibited larger Δ EOS values.

Similarly, a positive correlation was observed between Δ EOS and the rural rate of EOS change (R_{t-EOS}). Positive and negative Δ EOS values were predominantly found in cold and warm regions, respectively (Fig. 3). Moreover, when Δ EOS > 0,

- 删除[胡欣欣]: experienced
- 删除[胡欣欣]: Figure
- 删除[胡欣欣]: Figure
- 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四
- 删除[胡欣欣]: indicating more advanced spring phenology in urban areas of high-latitude cities, which suggests a stronger impact of urban warming on SOS
- 删除[胡欣欣]: N
- 删除[胡欣欣]: W
- 删除[胡欣欣]: Figure
- 删除[胡欣欣]:
- 删除[胡欣欣]: Figure
- 删除[胡欣欣]: Figure
- 删除[胡欣欣]: distributions
- 删除[胡欣欣]: gradient
- 删除[胡欣欣]: Figure
- 删除[胡欣欣]: Figure
- 删除[胡欣欣]:
- 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四
- 删除[胡欣欣]: when Δ EOS was analyzed in relation to
- 删除[胡欣欣]: , a positive correlation was observed.
- 删除[胡欣欣]: Figure

both the absolute value of ΔEOS and $\Delta\text{R}_{\text{t-EOS}}$ increased with temperature, while for $\Delta\text{EOS} < 0$, the absolute value of ΔEOS and $\Delta\text{R}_{\text{t-EOS}}$ decreased with temperature.




Figure 2. Spatial distributions of urban-rural disparities in SOS (ΔSOS) (a and b) and in EOS (ΔEOS) (c and d) across mainland China. Mean land surface temperature (LST) is shown in grayscale for the study area, and the data reflect the 10-year mean for 2010–2020 for each city. Panel (e) depicts the linear correlation between ΔSOS and latitude ($R^2=0.35$, $p<0.01$); panel (f) depicts the linear correlation between ΔEOS and latitude ($R^2=0.46$, $p<0.01$); panel (g) compares ΔSOS between coastal and non-coastal provinces ($*p<0.05$); panel (h) compares ΔEOS between coastal and non-coastal provinces ($**p<0.01$). The map, generated using ArcGIS version 10.2, features administrative boundaries

删除[胡欣欣]: The s
删除[胡欣欣]: for
删除[胡欣欣]: ,
删除[胡欣欣]: was analyzed
删除[胡欣欣]: was represented in black-and-white range
删除[胡欣欣]: entire continent
删除[胡欣欣]: in these figures
删除[胡欣欣]: ed
删除[胡欣欣]: from
删除[胡欣欣]: to
删除[胡欣欣]: –

sourced from the Ministry of Civil Affairs of the People's Republic of China
 (http://xzqh.mca.gov.cn/map). Source: Esri, TomTom, FAO, USGS, and the GIS User
 Community; Ministry of Civil Affairs of the People's Republic of China | Powered by
 Esri.




Figure 3. Relationship between urban-rural phenological difference and temperature sensitivity across 293 Chinese cities.
 (a) Urban-rural difference in start of growing season (Δ SOS) versus temperature sensitivity of SOS (R_{t-SOS}).
 (b) Urban-rural difference in end of growing season (Δ EOS) versus temperature sensitivity of EOS (R_{t-EOS}).
 Each point represents one of the 293 cities analyzed, colored by land surface temperature (LST, °C). Red solid lines indicate linear regression fits.

3.2 Phenology response to LST

As we compared urban and rural regions, logistic trends were observed for both SOS and EOS in relation to LST (Fig. 4). In urban areas where LST was below 12 °C, the SOS occurred notably earlier compared to rural areas, while urban areas exhibited a delayed EOS when LST was below 18.5 °C. As LST decreased, the SOS stabilized at 101.3 days for urban regions and 129.7 days for rural regions, while the EOS stabilized at 321.3 days for urban areas and 310.8 days for rural areas. In contrast,

删除[胡欣欣]: Figure

删除[胡欣欣]: regions

删除[胡欣欣]: C

删除[胡欣欣]: onversely

when LST exceeded the specific thresholds for urban-rural differences (LST = 18 °C for Δ SOS and LST = 18.5 °C for Δ EOS), the phenological disparities between urban and rural areas were no longer significant and sometimes even reversed (Fig. 4a,b). As LST increased, the urban and rural SOS stabilized at 71.7 days (Fig. 4a), and the urban and rural EOS stabilized at 327.4 days and 332.4 days, respectively. Fitted curves of Δ SOS and Δ EOS with LST revealed positive and negative relationship, respectively, intersecting at LST = 18 °C and 18.5 °C, indicating a shift in SOS and EOS dynamics with increasing temperatures beyond these thresholds.




Figure 4. Relationship between background temperature (LST) and phenology (SOS and EOS) in urban and rural areas (a, b) for 2010 to 2020, and between Δ SOS and LST and between Δ EOS and LST (c,d).




Figure 5. Relationships between pre-season temperature (T) and SOS/EOS (a,b), and the corresponding first-order derivatives with respect to temperature (c,d), representing phenological sensitivity to temperature.

To determine whether the temperature sensitivity influenced the logistic trends of phenology with LST, we conducted a nationwide fitting of phenology to T and derived logistic curves for SOS and EOS (Fig. 5). The first-order derivatives of the fitted curves, representing the phenological sensitivity to T, revealed warming benefits on SOS and EOS, ranging from -3.5 to 10 °C and 2.0 to 13.5 °C, respectively. The highest sensitivity was observed at T = 4 °C for SOS and T = 6 °C for EOS. Notably, a decreasing trend for EOS was observed when T > 15 °C, corresponding to the advanced EOS when T > 18.5 °C (Fig. 5d).

To assess the temperature sensitivity of phenological variations between urban and rural settings across vegetation types, we established regression models for Δ SOS, Δ EOS, and LST for each type (Fig. 6). Results showed that shrublands had the highest temperature sensitivity for both metrics, whereas CF displayed the lowest. The temperature response of MF and BF did not differ significantly.

删除[胡欣欣]: The r
删除[胡欣欣]:
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四
删除[胡欣欣]: as well as their first-order derivatives
(implying the phenological sensitivity to T) of the fitting curve.
删除[胡欣欣]: also
删除[胡欣欣]: Figure
删除[胡欣欣]: ing
删除[胡欣欣]: that
删除[胡欣欣]: ed
删除[胡欣欣]: -
删除[胡欣欣]: Figure
删除[胡欣欣]: SOS
设置格式[胡欣欣]: 字体: 倾斜
删除[胡欣欣]: Figure
删除[胡欣欣]: Findings revealed
删除[胡欣欣]: showed no
删除[胡欣欣]: variation




Figure 6. Relationships between LST and Δ SOS/ Δ EOS across the 293 cities for four vegetation types. The inset subplots compare the mean phenological differences among vegetation groups. BF: Broad-leaf forest, CF: Coniferous forest, MF: Mixed forest, Shrubs: Shrublands.

3.3 Quantify the LST determined phenological sensitivity

To obtain LST-determined phenological sensitivity, the T in the first-order derivative curves shown in Figs. 5c and 5d was converted to LST based on the strong correlation between T and LST (Fig. S4). Three distinct LST thresholds were identified: phenological date stabilization thresholds, urban-rural disparity reversal thresholds, and phenological sensitivity saturation thresholds. Specifically, phenological date stabilization and urban-rural disparity reversal occur at LST = 12 °C (SOS) / 18.5 °C (EOS), while phenological sensitivity saturates at LST = 12.5 °C (SOS) / 4 °C (EOS). Based on the first-order derivatives, we identified the specific thresholds for phenological sensitivity. We observed that when background LST exceeded 12.5 °C for spring and 4 °C for autumn, the temperature sensitivity (Rt) began to significantly weaken. This indicates a saturation of the physiological response rate, even though the absolute urban-rural difference (Δ SOS/ Δ EOS) might still persist until higher temperatures are reached.

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: The relation between Δ SOS and Δ EOS with the LST across the 293 cities for different vegetation types, insert figures presented the comparison of averaged phenology difference across the 293 cities.

删除[胡欣欣]: Bush wood

删除[胡欣欣]:

删除[胡欣欣]: In order t

删除[胡欣欣]:

删除[胡欣欣]: s

删除[胡欣欣]: ure

删除[胡欣欣]: translated

删除[胡欣欣]: considering the

删除[胡欣欣]: Figure

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C




Figure 7. Schematic diagram of the three scenarios that likely occurred in urban (red dots) and rural (blue dots) regions for SOS (I: $LST_{R1} < LST_{U1} < 12.5\text{ }^{\circ}\text{C}$; II: $12.5\text{ }^{\circ}\text{C} < LST_{R2} < LST_{U2}$; III: $LST_{R3} < 12.5\text{ }^{\circ}\text{C} < LST_{U3}$) and EOS (I: $LST_{R1} < LST_{U1} < 4\text{ }^{\circ}\text{C}$; II: $4\text{ }^{\circ}\text{C} < LST_{R2} < LST_{U2}$; III: $LST_{R3} < 4\text{ }^{\circ}\text{C} < LST_{U3}$).

Scenario I: when LST was less than $12.5\text{ }^{\circ}\text{C}$ ($4\text{ }^{\circ}\text{C}$) for SOS (EOS), higher LST was associated with heightened temperature sensitivity, resulting in urban regions (U_1) exhibiting higher sensitivity than rural regions (R_1) (Fig. 5).

Scenario II: when LST exceeded $12.5\text{ }^{\circ}\text{C}$ ($4\text{ }^{\circ}\text{C}$) for SOS (EOS), temperature sensitivity weakened as LST increased, yet urban regions (U_2) still exhibited higher sensitivity than rural regions (R_2) (Fig. 5).

Scenario III: when urban LST surpassed $12.5\text{ }^{\circ}\text{C}$ ($4\text{ }^{\circ}\text{C}$) while rural LST remained below $12.5\text{ }^{\circ}\text{C}$ ($4\text{ }^{\circ}\text{C}$), the difference in sensitivity between urban and rural regions was inconclusive.

3.4 Phenology response to aridity

When the phenological responses were fitted to AI, R_{t-SOS} and ΔSOS were positively related to AI (Fig. 8). When $AI < 1.7$, there was a decrease in absolute R_{t-SOS} with increasing AI, indicating that increasing aridity weakened the temperature sensitivity of SOS. In contrast, when $AI > 1.7$, most urban areas exhibited positive R_{t-SOS} (resulting in delayed SOS), with higher AI values corresponding to greater R_{t-SOS} , suggesting that severe drought conditions adversely delayed the onset of SOS.

删除[胡欣欣]: for
删除[胡欣欣]: the
删除[胡欣欣]: regions

删除[胡欣欣]: an increased

删除[胡欣欣]: Figure

删除[胡欣欣]: there was a weakened

删除[胡欣欣]: leading to

删除[胡欣欣]: demonstrating

删除[胡欣欣]:

删除[胡欣欣]: Figure

删除[胡欣欣]: the

删除[胡欣欣]: Figure

删除[胡欣欣]: It was noted that w

删除[胡欣欣]: heightened

删除[胡欣欣]: Conversely




Figure 8. Phenological temperature sensitivity and urban-rural phenological differences determined by the aridity index (AI). When the phenological responses were fitted to AI, R_{t-SOS} and ΔSOS were positively related to AI.

When R_{t-EOS} and ΔEOS were fitted to AI, a negative relationship was observed. For R_{t-EOS} , when $AI < 1.4$, most R_{t-EOS} values were positive, and decreased with AI, i.e., enhanced drought offset the benefit of urban warming on the delayed EOS. When $AI > 1.4$, most cities exhibited negative R_{t-EOS} , and higher AI values corresponded to more negative R_{t-EOS} . That is, extreme drought leads to passive warming effects on EOS. Notably, negative ΔEOS tended to appear when $AI > 2$.

AI was observed to advance ΔSOS and hasten ΔEOS across 293 cities, with this pattern holding true for all four vegetation types (Fig. 9). Notably, ΔSOS for CF showed less sensitivity to AI than the other vegetation types. Both BF and MF demonstrated lower sensitivities to AI for both ΔSOS and ΔEOS than shrublands.

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: The AI determined phenological sensitivity and urban-rural difference.

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四

删除[胡欣欣]: the

设置格式[胡欣欣]: 下标

删除[胡欣欣]: ile

删除[胡欣欣]: against

删除[胡欣欣]: of the

设置格式[胡欣欣]: 非上标/ 下标

删除[胡欣欣]: w

删除[胡欣欣]: as

删除[胡欣欣]: the

删除[胡欣欣]: will

删除[胡欣欣]: shared a

删除[胡欣欣]: the

删除[胡欣欣]: I.e.

删除[胡欣欣]: will

删除[胡欣欣]: It is noted that the

删除[胡欣欣]: has been

删除[胡欣欣]: Figure

删除[胡欣欣]: in

删除[胡欣欣]: influences

删除[胡欣欣]: , as compared to




Figure 9. Relationships between AI and urban-rural phenological differences (ΔSOS , ΔEOS) across different vegetation types.

Panel (a): Scatter plot of ΔSOS versus AI with linear regression fits for each vegetation type.

Panel (b): Scatter plot of ΔEOS versus AI with linear regression fits for each vegetation type.




Figure 10. Venn diagrams illustrating the relative contributions of AI and LST to ΔSOS , ΔEOS , $R_{t-\text{SOS}}$, and $R_{t-\text{EOS}}$ across 293 Chinese cities. Note: Values in each section represent the percentage of variance independently explained by AI or LST

and their shared explanatory power in the overlapping region, and the total variance explained by both factors is labeled below each panel.

To further quantify the individual and combined contributions of LST and AI to phenological variations, a variance decomposition analysis was conducted (Legendre and Legendre, 2012; Fig. 10). The Venn diagram revealed that AI and LST explained 75.05%, 76.21%, 71.72% and 82.14% of the variance in Δ SOS, Δ EOS, R_{t-SOS} , and R_{t-EOS} , respectively (Fig. 10), and LST always contributed more to these phenological indicators. Additionally, the interaction effect of LST and AI ranged from 3.29% to 5.66%.

4. Discussion

4.1 Overall urbanization effects on plant phenology in China

Our multi-city assessment confirms a predominant urban-rural phenological divergence across China, characterized by advanced SOS and delayed EOS within a concentrated 10 km urbanization footprint. This consistent spatial pattern validates that urban warming acts as a primary driver of phenological shifts. Beyond mere thermal forcing, this convergence across 293 cities reflects the profound impact of urban microclimates on plant life cycles. The observed 10 km threshold is particularly significant as it suggests a localized but intense ecological footprint that necessitates specialized urban greening management strategies, especially in high-density northern cities where these shifts are most pronounced.

Spatially, the pronounced north-south gradient in Δ SOS and Δ EOS across China, as shown in Fig. 2, closely mirrors the spatial pattern of LST, reflecting stronger temperature sensitivity of vegetation phenology in northern regions than in the south. This finding aligns with previous research documenting greater phenological shifts in northern areas (Jia et al., 2021; Yang et al., 2025), and further highlights that background climate acts as a key modulator of urban phenological impacts. The more pronounced urban warming effects observed in coastal provinces relative to inland

删除[胡欣欣]: &
删除[胡欣欣]: Figure
删除[胡欣欣]: 71.72%,
设置格式[胡欣欣]: 字体: 小四
删除[胡欣欣]: for
删除[胡欣欣]: Figure
删除[胡欣欣]: had more
删除[胡欣欣]: ion
删除[胡欣欣]: s
删除[胡欣欣]: for
删除[胡欣欣]: y
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四
删除[胡欣欣]: Besides
删除[胡欣欣]: -
删除[胡欣欣]: -
删除[胡欣欣]: ure
删除[胡欣欣]: the far
删除[胡欣欣]: relative to
删除[胡欣欣]: and supports

cities (Fig. 2) likely stem from the buffering effect of the maritime climate on seasonal temperature constraints, which interacts with urban heat islands to amplify urban-rural phenological divergence. Moreover, the strong correlations between Δ SOS/ Δ EOS, temperature sensitivity indicators (R_{t-SOS} and R_{t-EOS}), LST, and AI, as presented in Fig. 3, confirm that background temperature and aridity are critical drivers shaping the magnitude of urbanization-induced phenological shifts across the 293 cities, rather than just passive correlates, validating the core hypothesis of this study.

删除[胡欣欣]: Figure

删除[胡欣欣]: ure

Notably, the pattern of urban warming's benefit on vegetation phenology was consistent among the four vegetation types (Fig. S2). This consistency could be attributed to the ecological convergence of plants coexisting in shared local habitats within each city (Wang et al., 2016). Ecologically, the consistent 10 km effective footprint validates that urban environments function as distinct microclimatic islands. This spatial boundary conceptually highlights that urban ecological engineering and greening strategies must account for highly localized, rather than regional, thermal forcings.

删除[胡欣欣]: It is noted that

删除[胡欣欣]: the

删除[胡欣欣]: Figure

删除[胡欣欣]: -

删除[胡欣欣]: '

删除[胡欣欣]: .'

4.2 Background temperature related phenological response variance

When we quantified phenological responses to temperature across the climatic gradient, clear logistic trends were observed for both SOS and EOS (Fig. 5). Consistent with this logistic trajectory, we observed that in urban areas, the growth rate for SOS/EOS began to plateau when temperatures exceeded 4 °C/6 °C, respectively (Fig. 5). Given the significant correlation between LST and T (Fig. S4), we anticipated a diminished sensitivity in phenology with higher LST (Fig. 7). These findings lend credibility to the "saturation" effect hypothesis, suggesting that the accelerating influence of warmth on vegetation phenology reported in earlier studies has its limits (Badeck et al., 2004; Li et al., 2021). Indeed, plants in southern regions exhibit not just a dampened phenological response but also a decelerated growth rate when faced with increasing temperatures (Li et al., 2021; Büntgen et al., 2019). As

删除[胡欣欣]: Figure

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: Figure

删除[胡欣欣]: Taking into account

删除[胡欣欣]: Figure

删除[胡欣欣]: could also

删除[胡欣欣]: Figure

删除[胡欣欣]: the

547 illustrated by the first-order derivatives in Figs. 5c and 5d, the phenological sensitivity
548 to temperature does not increase linearly, but peaks at $T = 4\text{ }^{\circ}\text{C}$ for SOS and $T = 6\text{ }^{\circ}\text{C}$
549 for EOS before rapidly declining. Specifically, the stabilization of urban SOS at 101.3
550 days and rural SOS at 129.7 days when LST was low (Fig. 4a) further demonstrated
551 that plants reach a required physiological threshold that cannot be bypassed by
552 additional thermal forcing. Collectively, these results highlight that threshold-based
553 non-linear models rather than simple linear relationships are more suitable for
554 predicting phenological responses under scenarios of continuous environmental
555 warming, especially in urban settings. Conceptually, these saturation thresholds
556 challenge the traditional linear paradigm of "warming-driven phenological
557 advancement". By identifying explicit thermal boundaries, our framework establishes
558 that the physiological capacity of urban vegetation to capitalize on UHI warming is
559 strictly finite.

560 The observation of temperature saturation for the SOS in our study likely stems
561 from the chilling requirements for vernalization (Kim et al., 2009; Hanninen et al.,
562 2019). Typically, vegetation needs to undergo a period of exposure to low chilling
563 temperatures to break dormancy. This mechanism is consistent with our spatial
564 observations in Fig. 2a, where cities in lower latitudes (warmer regions) exhibited
565 significantly smaller ΔSOS values. This suggests that in these warmer regions, the
566 higher background LST may fail to satisfy the necessary chilling requirements,
567 leading to the diminished phenological sensitivity we quantified in Fig. 3. Moreover,
568 an earlier onset of greening increases the risk of damage from freezing events, posing
569 a survival threat to plants (Duan et al., 2011; Wang et al., 2015). This phenomenon
570 explains the delayed SOS in approximately 16% of the cities under study, mainly
571 distributed in low-latitude regions (Fig. 4).

572 Regarding the EOS, a moderate increase in temperature can enhance the activity
573 of photosynthetic enzymes (Shi et al., 2014), slow down the degradation of
574 chlorophyll (Fracheboud et al., 2009), reduce the risk of frost exposure in autumn, and
575 increase the potential for growth and photosynthetic production, ultimately promoting
576 the extension of the growing season. However, the benefits of increased temperature

删除[胡欣欣]: ure
删除[胡欣欣]: $^{\circ}\text{C}$
删除[胡欣欣]: $^{\circ}\text{C}$
删除[胡欣欣]: is
删除[胡欣欣]: Figure
删除[胡欣欣]: s
删除[胡欣欣]: mandatory
删除[胡欣欣]: -
删除[胡欣欣]: ‘
删除[胡欣欣]: ’
删除[胡欣欣]: ure
删除[胡欣欣]: territories
删除[胡欣欣]: ure
删除[胡欣欣]: d
删除[胡欣欣]: the
删除[胡欣欣]: Figure
删除[胡欣欣]: could
删除[胡欣欣]: ies
删除[胡欣欣]: While

577 can be offset by the negative impacts of heat waves and excessive thermal stress
578 (Mathur et al., 2014). Our data in Figs. 5b and 5d provide direct evidence for this:
579 when pre-season temperature (T) exceeded 15 °C, a decreasing trend for EOS was
580 observed. This aligns with the intersection points identified in Figs. 4c and 4d at LST
581 = 18 °C, where the urban-rural phenology difference reverses, confirming that
582 excessive heat in urban centers triggers premature leaf senescence rather than further
583 delaying it. In fact, when T > 15 °C, EOS tends to be reduced, with an increased
584 sensitivity to temperature changes (Estiarte and Peñuelas, 2015; Fig. 5b).

585 Despite the existence of "saturation" effects, lower background temperatures do
586 not necessarily correspond to higher phenological sensitivity (Fig. 4b, c). For SOS,
587 when T < 4 °C, the phenological sensitivity approached zero with decreased T (Fig.
588 4c), which led to a stable SOS of 120 days (Fig. 5a,c). In fact, phenological events
589 such as bud burst could not be triggered when the threshold for high forcing
590 temperatures was not reached (Hanninen et al., 2019). For EOS, when T > 6 °C, the
591 increased T sensitivity produces more EOS advancement in warmer regions. This
592 reversal may be associated with abscisic acid synthesis induced by low temperatures,
593 which accelerates the progress of defoliation.

594 Beyond thermal controls alone, our findings regarding the aridity-driven
595 modulation of phenological responses provide a critical geographic context that
596 complements previous observations. While earlier studies primarily emphasized
597 thermal forcing as the dominant driver of urban phenology (Meng et al., 2020), our
598 results demonstrate that AI acts as a significant "environmental filter" that can
599 override or even reverse warming benefits. As shown in Figs. 8 and 10, the high
600 explanatory power of AI (17.33% for ΔEOS) suggests that in water-limited regions of
601 northern and western China, the physiological stress induced by drought prevents
602 plants from utilizing the additional heat provided by the Urban Heat Island (UHI)
603 effect. This phenomenon is consistent with the "resource limitation" theory, where
604 plant development is constrained by the most limiting resource (water; Harpole et al.,
605 2011), regardless of optimal temperatures. Compared to coastal cities where water is
606 abundant, the weakened warming benefits in arid inland cities highlight the need for

regionally differentiated urban irrigation strategies to maintain ecological productivity. However, when $LST < 12.5\text{ }^{\circ}\text{C}$, warming regions exhibited higher urban-rural differences in ΔR_{t-SOS} , which contradicted the weakened urban-rural difference in ΔR_{t-SOS} reported in the previous study. For EOS, the warming benefits only occurred when $0 < LST < 18\text{ }^{\circ}\text{C}$. Moreover, the weakened sensitivity was only detected in the LST interval of $LST > 4\text{ }^{\circ}\text{C}$. The conclusions of weakened ΔR_{t-SOS} and ΔR_{t-EOS} in the USA may reflect that their LST ranges were mostly within the range of $8\text{ }^{\circ}\text{C}$ to $22\text{ }^{\circ}\text{C}$.

4.3 AI related phenological response variance

In addition to background temperature (LST), the aridity index (AI) drives significant spatial variation in urban-rural phenological disparities (ΔSOS , ΔEOS). Spatially, the coastal cities experienced more benefits from urban warming (Fig. 2). Our statistical decomposition in Fig. 10 further elucidates this aridity-driven modulation, showing that AI and LST together explain more than 75% of the variance in ΔSOS and ΔEOS . Notably, the fact that AI explains 17.33% of the variation in ΔEOS (Fig. 10c) supports our inference that water availability is a critical limiting factor in autumn. This is further substantiated by the positive correlation between ΔSOS and AI in Fig. 8b, which quantifies how increasing aridity progressively offsets the advanced green-up typically induced by urban warming. These results highlight that the warming benefits on vegetation phenology will be weakened in those drought-dominated regions. In fact, the delayed SOS has already been demonstrated in previous studies (Ji et al., 2021; Yu et al., 2003).

Water deficit inhibits leaf growth via chemical signals (Knauer et al., 2017), and urbanization exacerbates drought stress (Zhang et al., 2019; Pollastrini et al., 2019), further weakening or even reversing the positive effects of urban warming on phenology. Furthermore, in water-stressed regions, temperatures are already close to the optimum for photosynthesis, thus fewer benefits are expected (Peñuelas et al., 2004). The weakened temperature sensitivity in drought-prone cities is likely to lead to less extension of the growing season.

635 Notably, the influence of AI on autumn phenology response was higher than that
636 on spring phenology response (Fig. 10). During spring leaf development, plants may
637 use water stored in their bodies to support the growth of new leaves (Shi et al., 2023).
638 Supporting evidence comes from the synchrony between plant water storage and leaf
639 sprouting for the boreal and temperate forests (Tian et al., 2018). In this case, plants
640 may have relatively little demand for external water supply, especially in some wetter
641 areas. By comparison, in summer and autumn, plant transpiration is higher, and water
642 demand is greater (Wu et al., 2022), so the influence of drought severity on plant
643 phenology is more pronounced. The overarching ecological implication here is a
644 profound vulnerability: as background climates dry under global change, the
645 compounding stress of UHI and background aridity may lead to "phenological
646 compression". In arid urban ecosystems, this moisture deficit effectively neutralizes
647 the anticipated carbon sequestration benefits of thermally extended growing seasons,
648 rendering them highly sensitive to future climate extremes.

649 **4.4 Phenological response variation among different vegetation types**

650 The phenological response to LST and AI varies significantly across vegetation
651 types, reflecting the influence of distinct ecological strategies. Our results (Figs. 6 and
652 9) demonstrate that shrublands are the most sensitive to both warming and aridity,
653 likely due to their opportunistic resource utilization strategy and higher leaf turnover
654 rates (Xiong et al., 2024). In contrast, coniferous forests exhibited a muted response to
655 both drivers (Malla et al., 2023). While our analysis relies on broad land-cover
656 categories and does not directly measure species-specific functional traits, these
657 discrepancies likely reflect generalized divergent ecological strategies among the
658 vegetation groups.

659 This insensitivity is substantiated by our AI-sensitivity analysis (Fig. 9a),
660 suggesting that the conservative hydraulic traits of conifers — such as needle
661 morphology and lower stomatal conductance (Sperry et al., 2002; Brodribb and
662 McAdam, 2013) — act as a buffer against the "aridity-driven inhibition" that more

删除[胡欣欣]: While it is noted that

删除[胡欣欣]: the

删除[胡欣欣]: the

删除[胡欣欣]: Figure

删除[胡欣欣]: ous

删除[胡欣欣]:

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: the external water supply may have relativ ...

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: Comparatively

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: the transpiration of plants is larger

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: the demand for water is strong

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: degree

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: will be more prominent

删除[胡欣欣]: '

删除[胡欣欣]: '

删除[胡欣欣]: ure

删除[胡欣欣]: S

删除[胡欣欣]: C

删除[胡欣欣]: F

删除[胡欣欣]: Figure

设置格式[胡欣欣]: 字体: 小四

设置格式[胡欣欣]: 字体: 小四

删除[胡欣欣]: McAdam

severely affects broadleaf species. Based on generalized ecological principles, this might be associated with typical drought-resistant adaptations found in many conifers (e.g., needle morphology and stricter stomatal control), which could limit the potential positive impact of warming on their phenology. In contrast, the heightened sensitivity of shrublands observed in our dataset aligns with literature suggesting more opportunistic resource utilization strategies in shrubs (Xiong et al., 2024).

删除[胡欣欣]: -

By considering these functional differences, our study provides a more nuanced understanding of urban ecosystem resilience, suggesting that urban greening programs should prioritize drought-resistant coniferous or mixed forest structures in regions where the warming-saturation effect is most pronounced. However, we note that these functional interpretations remain broad generalizations; future studies incorporating field-based, species-level trait measurements are required to mechanistically validate these large-scale categorical patterns. Recognizing these trait-mediated responses is a crucial conceptual step for urban forestry. It indicates that building climate-resilient urban ecosystems requires shifting from generic greening to deploying specific functional traits—such as the conservative hydraulic strategies of conifers—to buffer against the coupled heat-aridity saturation effects.

5. Conclusion

Our study used remote sensing data to investigate the impact of urbanization on vegetation phenology across 293 Chinese cities for 2010 to 2020. Urban heat islands consistently advanced the start of the growing season (SOS) by 12.06 days and delayed the end of the growing season (EOS) by 9.86 days relative to rural areas. However, the magnitude of these shifts varied markedly with background land surface temperature (LST) and aridity index (AI), indicating that urban warming benefits are not universally distributed but are constrained by local climatic context.

删除[胡欣欣]: utilized

删除[胡欣欣]: from

删除[胡欣欣]: "

删除[胡欣欣]: "

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C

删除[胡欣欣]: °C

We identified a non-linear saturation effect in phenological sensitivity to urban warming. Temperature sensitivity peaked at 4 °C (spring) and 6 °C (autumn), and weakened significantly beyond saturation thresholds of 12.5 °C for SOS and 4 °C for

EOS. Within the effective ranges of 3.5 ~~to 17.5 °C~~ (SOS) and 0–18 ~~°C~~ (EOS), urban warming exerted pronounced positive effects on phenological shifts; outside these ranges, sensitivity approached zero. Notably, extreme high temperatures reversed the beneficial effects for EOS, causing weakly delayed or even advanced senescence in warm urban regions.

删除[胡欣欣]: –

删除[胡欣欣]: °C

删除[胡欣欣]: °C

Aridity emerged as a critical negative regulator of warming benefits. Across most AI ranges, increasing aridity attenuated the positive effects of UHI on both SOS and EOS. Within the intermediate aridity band of $1.4 < AI < 2.0$, warming benefits were entirely reversed. Consequently, coastal cities with lower AI exhibited substantially larger urban-rural phenological disparities than arid inland cities, where coupled heat-aridity stress compressed the effective growing season.

Collectively, these findings challenge the linear assumption that urban warming universally extends the growing season. Instead, they demonstrate that phenological responses to UHI are bounded by finite physiological limits and modulated by water availability. For urban ecological planning, our results imply that greening strategies must be climate-specific: drought-resistant vegetation structures should be prioritized in arid regions where saturation occurs at lower thermal thresholds, whereas coastal and northern cities can more effectively harness UHI benefits for growing season extension. Future research incorporating species-level functional traits and process-based phenology models will be essential to refine these thresholds and project urban ecosystem dynamics under continued global warming.

Data availability statement

The data that support the findings of this study are available in those public domain resources that listed in the table 1.

Author contributions

Lin Xingwen: Conceptualization, Methodology, Funding Acquisition;

Zhang Zhenzhen: Data curation, Writing - Original draft preparation;

设置格式[胡欣欣]: 缩进: 左侧: 0 毫米, 首行缩进: 0 毫米, 无项目符号或编号

删除[胡欣欣]: : CRediT

删除[胡欣欣]: Cui Shufen: Conceptualization, Methodology;

718	Hu Xinxin: Methodology, Formal analysis, Validation, Writing - review <u>and</u> editing;	删除[胡欣欣]: &
719	Zhang Yongqi: Visualization, Investigation;	
720	<u>Sun Liheng, Wu Chaofan and Cui Shufen</u> ; Supervision;	删除[胡欣欣]:
721	Zhang Zhaoyang, Chen Yuanjian; Software, Validation;	删除[胡欣欣]: Lin Xingwen
722	<u>Wen Qingqing: Funding Acquisition,</u>	删除[胡欣欣]: and Wen Qingqing
723	Competing interests	设置格式[胡欣欣]: 字体: (中文) 宋体, 小四 删除[胡欣欣]: .
724	The authors declared that the research was conducted in the absence of any	删除[胡欣欣]: Data availability statement
725	commercial or financial relationships that could be construed as a potential conflict of	设置格式[胡欣欣]: 缩进: 左侧: 0 毫米, 首行缩进: 0 毫米
726	interest.	
727	Acknowledgements	设置格式[胡欣欣]: 标题 2, 行距: 单倍行距
728	<u>We appreciate the valuable suggestions from the Associate and anonymous reviewers,</u>	设置格式[胡欣欣]: 字体: (默认) Times New Roman,
729	<u>which substantially improved this manuscript. We also acknowledge the computing</u>	设置格式[胡欣欣]: 字体: (默认) Times New Roman,
730	<u>support provided by our university and the members of our research group for their</u>	设置格式[胡欣欣]: 标题 2, 行距: 单倍行距
731	<u>assistance throughout this work.</u>	删除[胡欣欣]:
732	Funding	设置格式[胡欣欣]: 字体: (默认) Times New Roman, 设置格式[胡欣欣]: 字体: (默认) Times New Roman,
733	<u>This study was supported by “Jinhua scientific and technological projects (grant</u>	设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 非加粗
734	<u>number, 2026-3-108, 2024-3-002), Open Research Project of Innovation Center of</u>	设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 非加粗
735	<u>Yangtze River Delta, Zhejiang University (grant number: 2025KFKY004)”</u>	删除[胡欣欣]: This study was supported by "Department of Ecology and Environment, 设置格式[胡欣欣]: 字体: (默认) Times New Roman,
736	References	设置格式[胡欣欣]: 正文, 行距: 1.5 倍行距
737	<u>Angilletta, M. J.: Thermal Adaptation: A Theoretical and Empirical Synthesis, Oxford</u>	设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 黑色
738	<u>University Press, Oxford, https://doi.org/10.1093/acprof:oso/9780198570875.001.1,</u>	删除[胡欣欣]: Angilletta, M. J. (2009). Thermal Adaptation: A Theoretical and Empirical Synthesis, Oxford University Press, Oxford, https://doi.org/10.1093/acprof:oso/9780198570875.001.1 , 2009.
739	<u>2009.</u>	删除[胡欣欣]: (2004).
740	Badeck, F.-W., Bondeau, A., Böttcher, K., Doktor, D., Lucht, W., Schaber, J., and Sitch, S.;	删除[胡欣欣]: .
741	Responses of spring phenology to climate change, <u>New Phytol., 162, 295–309,</u>	设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 黑色

742 <https://doi.org/10.1111/j.1469-8137.2004.01059.x>, 2004.

743 [Brodrribb, T. J. and McAdam, S. A. M.: Absciscic acid mediates a divergence in the drought](#)

744 [response of two conifers, Plant Physiol., 162, 1370–1377,](#)

745 <https://doi.org/10.1104/pp.113.217877>, 2013.

746 Buyantuyev, A., [and](#) Wu, J.: Urbanization diversifies land surface phenology in arid

747 environments: Interactions among vegetation, climatic variation, and land use pattern

748 in the Phoenix metropolitan region, [USA, Landsc. Urban Plan., 105, 149–159,](#)

749 <https://doi.org/10.1016/j.landurbplan.2011.12.013>, 2012.

750 Büntgen, U., Krusic, P. J., Piermattei, A., Coomes, D. A., Esper, J., Myglan, V. S.,

751 Kirdyanov, A. V., Camarero, J. J., Crivellaro, A., [and](#) Körner, C.: Limited capacity of

752 tree growth to mitigate the global greenhouse effect under predicted warming. [Nat.](#)

753 [Commun., 10, 2171, https://doi.org/10.1038/s41467-019-10174-4](#), 2019.

754 Cai, Z., Jönsson, P., Jin, H., [and](#) Eklundh, L.: Performance of smoothing methods for

755 reconstructing NDVI time-Series and estimating vegetation phenology from MODIS

756 Data. [Remote Sens., 9, 1271, https://doi.org/10.3390/rs9121271](#), 2017.

757 Čehulić, I., Sever, K., Katičić Bogdan, I., Jazbec, A., Škvorc, Ž., [and](#) Bogdan, S.: Drought

758 Impact on Leaf Phenology and Spring Frost Susceptibility in a Quercus robur L.

759 Provenance Trial. [Forests, 10, 50, https://doi.org/10.3390/f10010050](#), 2019.

760 Chmielewski, F.-M., [and](#) Rötzer, T.: Response of tree phenology to climate change across

761 Europe. [Agric. For. Meteorol., 108, 101–112,](#)

762 [https://doi.org/10.1016/S0168-1923\(01\)00233-7](https://doi.org/10.1016/S0168-1923(01)00233-7), 2001.

763 Cong, N., Piao, S., Chen, A., Wang, X., Lin, X., Chen, S., Han, S., Zhou, G., [and](#) Zhang,

764 X.: Spring vegetation green-up date in China inferred from SPOT NDVI data: A

765 multiple model analysis. [Agr. Forest Meteorol., 165, 104–113,](#)

766 <https://doi.org/10.1016/j.agrformet.2012.06.009>, 2012.

767 Dallimer, M., Tang, Z., Gaston, K. J., [and](#) Davies, Z. G.: The extent of shifts in vegetation

768 phenology between rural and urban areas within a human-dominated region. [Ecol.](#)

769 [Evol., 6, 1942–1953, https://doi.org/10.1002/ece3.1990](#), 2016.

770 Ding, H., Xu, L., Elmore, A. J., [Ma, X., and Li, X.](#): Vegetation phenology influenced by

771 rapid urbanization of The Yangtze Delta region. [Remote Sens., 12, 1783,](#)

删除[胡欣欣]: New Phytologist. **162**, 295-309.

删除[胡欣欣]: ,

删除[胡欣欣]: &

删除[胡欣欣]: (2012).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

删除[胡欣欣]: USA. Landscape and Urban Planning. **105**

删除[胡欣欣]: (2019).

删除[胡欣欣]: .

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

删除[胡欣欣]: Nature Communications. **10**, 2171.

删除[胡欣欣]: &

删除[胡欣欣]: (2017).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

删除[胡欣欣]: . Remote Sensing. **9**, 1271.

删除[胡欣欣]: &

删除[胡欣欣]: (2019).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

删除[胡欣欣]: . Forests. **10**, 50.

删除[胡欣欣]: ,

删除[胡欣欣]: &

删除[胡欣欣]: (2001).

删除[胡欣欣]: .

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

删除[胡欣欣]: Agricultural and Forest Meteorology. **108**,

删除[胡欣欣]: (2012).

删除[胡欣欣]: .

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

删除[胡欣欣]: Agricultural and Forest Meteorology. **165**,

删除[胡欣欣]: &

删除[胡欣欣]: (2016).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

删除[胡欣欣]: . Ecology and Evolution. **6**, 1942-1953.

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜

772 <https://doi.org/10.3390/rs12111783>, 2020.

773 Doncaster, C. P.: Structural Equation Modeling and Natural Systems, *Fish Fish.*, **8**, 368 –

774 369, <https://doi.org/10.1111/j.1467-2979.2007.00260.x>, 2007.

775 Duan, J., Zhang, Q.-B., Lv, L., and Zhang, C.: Regional-scale winter-spring temperature

776 variability and chilling damage dynamics over the past two centuries in southeastern

777 China, *Clim. Dynam.*, **39**, 919–928, <https://doi.org/10.1007/s00382-011-1323-9>,

778 2011.

779 Elvidge, C. D., Baugh, K., Zhizhin, M., Hsu, F. C., and Ghosh, T.: VIIRS night-time lights,

780 *Int. J. Remote Sens.*, **38**, 5860 – 5879,

781 <https://doi.org/10.1080/01431161.2017.1342050>, 2017.

782 Estiarte, M., and Peñuelas, J.: Alteration of the phenology of leaf senescence and fall in

783 winter deciduous species by climate change: effects on nutrient proficiency, *Glob.*

784 *Change Biol.*, **21**, 1005–1017, <https://doi.org/10.1111/gcb.12804>, 2015.

785 Fan, J., He, H., Hu, T., Zhang, P., Yu, X., and Zhou, Y.: Estimation of landscape pattern

786 changes in BRICS from 1992 to 2013 using DMSP-OLS NTL images, *J. Indian Soc.*

787 *Remote Sens.*, **47**, 725–735, <https://doi.org/10.1007/s12524-019-00963-1>, 2019.

788 Fracheboud, Y., Luquez, V., Bjorken, L., Sjodin, A., Tuominen, H., and Jansson, S.: The

789 control of autumn senescence in European aspen, *Plant Physiol.*, **149**, 1982–1991,

790 <https://doi.org/10.1104/pp.108.133249>, 2009.

791 Gazal, R., White, M. A., Gillies, R., Rodemaker, E. L. I., Sparrow, E., and Gordon, L.:

792 GLOBE students, teachers, and scientists demonstrate variable differences between

793 urban and rural leaf phenology, *Glob. Change Biol.*, **14**, 1568–1580,

794 <https://doi.org/10.1111/j.1365-2486.2008.01602.x>, 2008.

795 Ge, W., Han, J., Zhang, D., and Wang, F.: Divergent impacts of droughts on vegetation

796 phenology and productivity in the Yungui Plateau, southwest China, *Ecol. Indic.*, **127**,

797 107743, <https://doi.org/10.1016/j.ecolind.2021.107743>, 2021.

798 Gow, L. J., Barrett, D. J., Renzullo, L. J., Phinn, S. R., and O'Grady, A. P.: Characterising

799 groundwater use by vegetation using a surface energy balance model and satellite

800 observations of land surface temperature, *Environ. Modell. Softw.*, **80**, 66–82,

801 <https://doi.org/10.1016/j.envsoft.2016.02.021>, 2016.

删除[胡欣欣]: . Remote Sensing. **12**, 1783.

删除[胡欣欣]: (2007).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Fish and Fisheries, 8(4), 368-369.

删除[胡欣欣]: &

删除[胡欣欣]: (2011).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Climate Dynamics. **39**, 919-928.

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: et al. (2017).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . International Journal of Remote Sensing, ...

删除[胡欣欣]: ,

删除[胡欣欣]: &

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: Penuelas

删除[胡欣欣]: (2015).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Global Change Biology. **21**, 1005-1017.

删除[胡欣欣]: &

删除[胡欣欣]: (2019).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Journal of the Indian Society of Remote ...

删除[胡欣欣]: &

删除[胡欣欣]: (2009).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Plant Physiology. **149**, 1982-1991.

删除[胡欣欣]: &

删除[胡欣欣]: (2008).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Global Change Biology. **14**, 1568-1580 .

删除[胡欣欣]: &

删除[胡欣欣]: (2021).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

802 Grimm, N. B., Faeth, S. H., Golubiewski, N. E., Redman, C. L., Wu, J., Bai, X., and
803 Briggs, J. M.; Global change and the ecology of cities, Science, 319, 756–760,
804 <https://doi.org/10.1126/science.1150195>, 2008,
805 Hanninen, H., Kramer, K., Tanino, K., Zhang, R., Wu, J., and Fu, Y. H.; Experiments are
806 necessary in process-based tree phenology modelling, Trends Plant Sci., 24, 199–209,
807 <https://doi.org/10.1016/j.tplants.2018.11.006>, 2019,
808 Harpole, W. S., Ngai, J. T., Cleland, E. E., Seabloom, E. W., Borer, E. T., Bracken, M. E.,
809 Elser, J. J., Gruner, D. S., Hillebrand, H., Shurin, J. B., and Smith, J. E.; Nutrient
810 co-limitation of primary producer communities, Ecol. Lett., 14, 852–862,
811 <https://doi.org/10.1111/j.1461-0248.2011.01651.x>, 2011,
812 He, C., Gao, B., Huang, Q., Ma, Q., and Dou, Y.; Environmental degradation in the urban
813 areas of China: Evidence from multi-source remote sensing data, Remote Sens.
814 Environ., 193, 65–75, <https://doi.org/10.1016/j.rse.2017.02.027>, 2017,
815 Jeong, S.-J., Park, H., Ho, C.-H., and Kim, J.; Impact of urbanization on spring and
816 autumn phenology of deciduous trees in the Seoul Capital Area, South Korea, Int. J.
817 Biometeorol., 63, 627–637, <https://doi.org/10.1007/s00484-018-1610-7>, 2019,
818 Ji, S., Ren, S., Li, Y., Dong, J., Wang, L., Quan, Q., and Liu, J.; Diverse responses of
819 spring phenology to preseason drought and warming under different biomes in the
820 North China Plain, Sci. Total Environ., 766, 144437,
821 <https://doi.org/10.1016/j.scitotenv.2020.144437>, 2021,
822 Jia, W., Zhao, S., Zhang, X., Liu, S., Henebry, G. M., and Liu, L.; Urbanization imprint on
823 land surface phenology: The urban-rural gradient analysis for Chinese cities, Glob.
824 Change Biol., 27, 2895–2904, <https://doi.org/10.1111/gcb.15602>, 2021,
825 Jönsson, P., and Eklundh, L.; TIMESAT—a program for analyzing time-series of satellite
826 sensor data, Comput. Geosci., 30, 833–845,
827 <https://doi.org/10.1016/j.cageo.2004.05.006>, 2004,
828 Kang, X., Hao, Y., Cui, X., Chen, H., Huang, S., Du, Y., Li, W., Kardol, P., Xiao, X., and
829 Cui, L.; Variability and changes in climate, phenology, and gross primary production
830 of an Alpine wetland ecosystem, Remote Sens., 8, 391,
831 <https://doi.org/10.3390/rs8050391>, 2016,

删除[胡欣欣]: &
删除[胡欣欣]: (2008).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Science, 319(5864), 756-760.
删除[胡欣欣]: &
删除[胡欣欣]: (2019).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Trends in Plant Science. **24**, 199-209.
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: et al. (2011).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Ecology Letters, 14(9), 852-862.
删除[胡欣欣]: &
删除[胡欣欣]: (2017).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Remote Sensing of Environment. **193**, 65-75.
删除[胡欣欣]:
删除[胡欣欣]:
删除[胡欣欣]: &
删除[胡欣欣]: (2019).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . International Journal of Biometeorology. ...
删除[胡欣欣]: &
删除[胡欣欣]: (2021).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Science of the Total Environment. **766**, ...
删除[胡欣欣]: &
删除[胡欣欣]: (2021).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Global Change Biology. **27**, 2895-2904.
删除[胡欣欣]: ,
删除[胡欣欣]: &
删除[胡欣欣]: (2004).

832 Ketola, T., and Kristensen, T. N.: Experimental approaches for testing if tolerance curves
833 are useful for predicting fitness in fluctuating environments, Front. Ecol. Evol., 5,
834 129, <https://doi.org/10.3389/fevo.2017.00129>, 2017,
835 Kim, D. H., Doyle, M. R., Sung, S., and Amasino, R. M.: Vernalization: winter and the
836 timing of flowering in plants, Annu. Rev. Cell Dev. Biol., 25, 277–299,
837 <https://doi.org/10.1146/annurev.cellbio.042308.113411>, 2009,
838 Knauer, J., Zaehle, S., Reichstein, M., Medlyn, B. E., Forkel, M., Hagemann, S., and
839 Werner, C.: The response of ecosystem water-use efficiency to rising atmospheric
840 CO₂ concentrations: sensitivity and large-scale biogeochemical implications, New
841 Phytol., 213, 1654–1666, <https://doi.org/10.1111/nph.14288>, 2017,
842 Legendre, P., and Legendre, L.: Numerical Ecology, 3rd ed., Elsevier, Amsterdam, ISBN
843 978-0-444-53869-7, 2012,
844 Li, K., Wang, C., Sun, Q., Rong, G., Tong, Z., Liu, X., and Zhang, J.: Spring phenological
845 sensitivity to climate Change in the northern hemisphere: comprehensive evaluation
846 and driving force analysis, Remote Sens., 13, 1972,
847 <https://doi.org/10.3390/rs13101972>, 2021,
848 Malla, R., Neupane, P. R., and Köhl, M.: Climate change impacts: Vegetation shift of
849 broad-leaved and coniferous forests, Trees For. People, 14, 100457,
850 <https://doi.org/10.1016/j.tfp.2023.100457>, 2023,
851 Mathur, S., Agrawal, D., and Jajoo, A.: Photosynthesis: response to high temperature
852 stress, J. Photochem. Photobiol. B, 137, 116–126,
853 <https://doi.org/10.1016/j.jphotobiol.2014.01.010>, 2014,
854 Meng, L., Mao, J., Zhou, Y., Richardson, A. D., Lee, X., Thornton, P. E., Ricciuto, D. M.,
855 Li, X., Dai, Y., Shi, X., and Jia, G.: Urban warming advances spring phenology but
856 reduces the response of phenology to temperature in the conterminous United States,
857 Proc. Natl. Acad. Sci. USA, 117, 4228–4233,
858 <https://doi.org/10.1073/pnas.1911117117>, 2020,
859 Mu, Q., Zhao, M., and Running, S. W.: Improvements to a MODIS global terrestrial
860 evapotranspiration algorithm, Remote Sens. Environ., 115, 1781 – 1800,
861 <https://doi.org/10.1016/j.rse.2011.02.019>, 2011,

删除[胡欣欣]: ,
删除[胡欣欣]: &
删除[胡欣欣]: (2017).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Frontiers in Ecology and Evolution. **5**, 129.
删除[胡欣欣]: &
删除[胡欣欣]: (2009).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Annual Review of Cell and Development...
删除[胡欣欣]: &
删除[胡欣欣]: (2017).
删除[胡欣欣]:
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . New Phytologist. **213**, 1654-1666.
删除[胡欣欣]: ,
删除[胡欣欣]: &
删除[胡欣欣]: (2012).
删除[胡欣欣]: (
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]:). Amsterdam: Elsevier.
删除[胡欣欣]: &
删除[胡欣欣]: (2021).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Remote Sensing. **13**, 1972.
删除[胡欣欣]: &
删除[胡欣欣]: (2023).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Trees, Forests and People. **14**, 100457 .
删除[胡欣欣]: (2014).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Journal of Photochemistry and Photobiol...
删除[胡欣欣]: MeAdam, B. S. A. M. J. P. P. (2013). Abse...
设置格式[胡欣欣]: 字体颜色: 蓝色, 删除线
删除[胡欣欣]: (2020).

862 Pablos, M., Martínez-Fernández, J., Piles, M., Sánchez, N., Vall-llossera, M., and Camps,
863 A.: Multi-temporal evaluation of soil moisture and land surface temperature
864 dynamics using in situ and satellite observations, Remote Sens., 8, 587,
865 <https://doi.org/10.3390/rs8070587>, 2016, 删除[胡欣欣]: &
866 Peng, S.: 1-km monthly precipitation dataset for China (1901-2021), National Tibetan
867 Plateau / Third Pole Environment Data Center [data set],
868 <https://doi.org/10.5281/zenodo.3185722>, 2020, 删除[胡欣欣]: (2016).
869 Peng, J., Wu, C., Zhang, X., Wang, X., and Gonsamo, A.: Satellite detection of cumulative
870 and lagged effects of drought on autumn leaf senescence over the Northern
871 Hemisphere, Glob. Change Biol., 25, 2174–2188, <https://doi.org/10.1111/gcb.14627>,
872 2019, 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
873 Peñuelas, J., Gordon, C., Llorens, L., Nielsen, T., Tietema, A., Beier, C., Bruna, P.,
874 Emmett, B., Estiarte, M., and Gorissen, A.: Nonintrusive field experiments show
875 different plant responses to warming and drought among sites, seasons, and species
876 in a north-south European gradient, Ecosystems, 7, 598–612,
877 <https://doi.org/10.1007/s10021-004-0179-7>, 2004, 删除[胡欣欣]: . Remote Sensing. **8**, 587.
878 Piao, S., Fang, J., Zhou, L., Ciais, P., and Zhu, B.: Variations in satellite-derived
879 phenology in China's temperate vegetation, Glob. Change Biol., 12, 672–685,
880 <https://doi.org/10.1111/j.1365-2486.2006.01123.x>, 2006, 删除[胡欣欣]: &
881 Pinheiro, J. C., and Bates, D. M.: Mixed-Effects Models in S and S-PLUS, Springer, New
882 York, ISBN 978-0-387-22747-4, https://doi.org/10.1007/b98882, 2000, 删除[胡欣欣]: (2006).
883 Pollastrini, M., Puletti, N., Selvi, F., Iacopetti, G., and Bussotti, F.: Widespread crown
884 defoliation after a drought and heat wave in the forests of Tuscany (Central Italy) and
885 their recovery—A case study from summer 2017, Front. For. Glob. Change, 2, 74,
886 <https://doi.org/10.3389/ffgc.2019.00074>, 2019, 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
887 Qiao, Z., Tian, G., and Xiao, L.: Diurnal and seasonal impacts of urbanization on the
888 urban thermal environment: A case study of Beijing using MODIS data, ISPRS J.
889 Photogramm. Remote Sens., 85, 93–101,
890 <https://doi.org/10.1016/j.isprsjprs.2013.08.010>, 2013, 删除[胡欣欣]: . Global Change Biology. **25**, 2174-2188. ...
891 Qiu, T., Song, C., and Li, J.: Impacts of urbanization on vegetation phenology over the
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: Penuelas, J., Gordon, C., Llorens, L., Niels...
删除[胡欣欣]: &
删除[胡欣欣]: (2006).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Global Change Biology. **12**, 672-685.
删除[胡欣欣]: ,
删除[胡欣欣]: &
删除[胡欣欣]: (2000).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Springer.
删除[胡欣欣]: &
删除[胡欣欣]: (2019).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . Frontiers in Forests and Global Change. **2**.
删除[胡欣欣]: &
删除[胡欣欣]: (2013).
设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...
删除[胡欣欣]: . ISPRS Journal of Photogrammetry and ...
删除[胡欣欣]: &
删除[胡欣欣]: (2017).

892 past three decades in Shanghai, China, *Remote Sens.*, **9**, 970, 删除[胡欣欣]: .

893 <https://doi.org/10.3390/rs9090970>, 2017, 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

894 Rustad, L., Campbell, J., Marion, G., Norby, R., Mitchell, M., Hartley, A., Cornelissen, J., 删除[胡欣欣]: . Remote Sensing. **9**, 970. ...

895 Gurevitch, J., and Gcte, N.; A meta-analysis of the response of soil respiration, net 删除[胡欣欣]: (2001).

896 nitrogen mineralization, and aboveground plant growth to experimental ecosystem 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

897 warming, *Oecologia*, **126**, 543–562, <https://doi.org/10.1007/s004420000544>, 2001, 删除[胡欣欣]: . Oecologia. **126**, 543–562.

898 Sage, R. F. and Kubien, D. S.; The temperature response of C3 and C4 photosynthesis, 删除[胡欣欣]: (2007).

899 *Plant Cell Environ.*, **30**, 1086–1106, 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

900 <https://doi.org/10.1111/j.1365-3040.2007.01682.x>, 2007, 删除[胡欣欣]: . Plant. Cell & Environment. **30**, 1086-1106.

901 Shi, C., Sun, G., Zhang, H., Xiao, B., Ze, B., Zhang, N., and Wu, N.; Effects of warming 删除[胡欣欣]: &

902 on chlorophyll degradation and carbohydrate accumulation of Alpine herbaceous 删除[胡欣欣]: (2014).

903 species during plant senescence on the Tibetan Plateau, *PLoS One*, **9**, e107874, 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

904 <https://doi.org/10.1371/journal.pone.0107874>, 2014, 删除[胡欣欣]: . PLoS OnE. **9**, e107874.

905 Shi, W., Li, J., Zhan, H., Yu, L., Wang, C., and Wang, S.; Relation between water storage 删除[胡欣欣]: &

906 and photoassimilate accumulation of neosinocalamus affinis with phenology, *Forests*, 删除[胡欣欣]: (2023).

907 **14**, 531, <https://doi.org/10.3390/f14030531>, 2023, 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

908 Sperry, J. S., Hacke, U. G., Oren, R., and Comstock, J. P.; Water deficits and hydraulic 删除[胡欣欣]: . Forests. **14**, 531.

909 limits to leaf water supply, *Plant Cell Environ.*, **25**, 251 – 263, 删除[胡欣欣]: &

910 <https://doi.org/10.1046/j.0016-8025.2001.00799.x>, 2002, 删除[胡欣欣]: (2002).

911 Tian, F., Wigneron, J.-P., Ciais, P., Chave, J., Ogée, J., Peñuelas, J., Ræbild, A., Domec, 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

912 J.-C., Tong, X., Brandt, M., Mialon, A., Rodriguez-Fernandez, N., Tagesson, T., 删除[胡欣欣]: . Plant, Cell & Environment, 25(2), 251-263. ...

913 Al-Yaari, A., Kerr, Y., Chen, C., Myneni, R. B., Zhang, W., Ardö, J., and Fensholt, R.; 删除[胡欣欣]: (2018).

914 Coupling of ecosystem-scale plant water storage and leaf phenology observed by 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

915 satellite, *Nat. Ecol. Evol.*, **2**, 1428–1435, <https://doi.org/10.1038/s41559-018-0630-3>, 删除[胡欣欣]: . Nature ecology & Evolution. **2**, 1428-1435.

916 **2018**, 设置格式[胡欣欣]: 字体: 四号

917 Wan, Z.; New refinements and validation of the MODIS Land-Surface 设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

918 Temperature/Emissivity products, *Remote Sens. Environ.*, **112**, 59–74, 删除[胡欣欣]: . Remote Sensing of Environment. **112**, 59-74 .

919 <https://doi.org/10.1016/j.rse.2006.06.026>, 2008, 删除[胡欣欣]: &

920 Wang, C., Cao, R., Chen, J., Rao, Y., and Tang, Y.; Temperature sensitivity of spring 删除[胡欣欣]: (2015).

921 vegetation phenology correlates to within-spring warming speed over the Northern

922

Hemisphere, [Ecol. Indic., 50, 62–68, https://doi.org/10.1016/j.ecolind.2014.11.004,](#)

923

[2015.](#)

924

Wang, C., Tang, Y., [and](#) Chen, J.: Plant phenological synchrony increases under rapid

925

within-spring warming, [Sci. Rep., 6, 25460, https://doi.org/10.1038/srep25460, 2016,](#)

926

White, M. A., De Beurs, K. M., Didan, K., Inouye, D. W., Richardson, A. D., Jensen, O. P.,

927

O’Keefe, J., Zhang, G., Nemani, R. R., Van Leeuwen, W. J. D., Brown, J. F., De Wit,

928

A., Schaepman, M., Lin, X., Dettinger, M., Bailey, A. S., Kimball, J., Schwartz, M.

929

D., Baldocchi, D. D., Lee, J. T., [and](#) Lauenroth, W. K.: Intercomparison,

930

interpretation, and assessment of spring phenology in North America estimated from

931

remote sensing for 1982–2006, [Glob. Change Biol., 15, 2335–2359,](#)

932

[https://doi.org/10.1111/j.1365-2486.2009.01910.x, 2009.](#)

933

[Wu, C., Peng, J., Ciais, P., Peñuelas, J., Wang, H., Beguería, S., Black, T. A., Jassal, R. S.,](#)

934

[Zhang, X., Yuan, W., Liang, E., Wang, X., Hua, H., Liu, R., Ju, W., Fu, Y. H., and Ge,](#)

935

[Q.: Increased drought effects on the phenology of autumn leaf senescence, Nat. Clim.](#)

936

[Change, 12, 943–949, https://doi.org/10.1038/s41558-022-01464-9, 2022.](#)

937

Xiong, X., Wu, H., Wei, X., [and](#) Jiang, M.: Contrasting temperature and light sensitivities

938

of spring leaf phenology between understory shrubs and canopy trees: Implications

939

for phenological escape, [Agric. For. Meteorol., 355, 110144,](#)

940

[https://doi.org/10.1016/j.agrformet.2024.110144, 2024.](#)

941

Yang, H., Zhang, Q., Zhang, S., [Wang, X., and Yu, H.:](#) Joint control of urban expansion

942

and climate change on urban-rural vegetation phenology gradient in 31 cities of

943

China, [Front. Ecol. Evol., 13, 1637210, https://doi.org/10.3389/fevo.2025.1637210,](#)

944

[2025.](#)

945

Yao, R., Wang, L., Huang, X., Guo, X., Niu, Z., [and](#) Liu, H.: Investigation of Urbanization

946

effects on land Surface phenology in Northeast China during 2001–2015, [Remote](#)

947

[Sens., 9, 66, https://doi.org/10.3390/rs9010066, 2017.](#)

948

Yu, F., Price, K. P., Ellis, J., [and](#) Shi, P.: Response of seasonal vegetation development to

949

climatic variations in eastern central Asia, [Remote Sens. Environ., 87, 42–54,](#)

950

[https://doi.org/10.1016/S0034-4257\(03\)00144-5, 2003.](#)

951

Yuan, M., Zhao, L., Lin, A., Wang, L., Li, Q., She, D., [and](#) Qu, S.: Impacts of preseason

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Ecological Indicators. **50**, 62-68.

删除[胡欣欣]: &

删除[胡欣欣]: (2016).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Scientific Reports. **6**, 25460.

删除[胡欣欣]: (2009).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Global Change Biology. **15**, 2335-2359 .

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: Wu, C., Peng, J., Ciais, P., Peñuelas, J., W...

删除[胡欣欣]: &

删除[胡欣欣]: (2024).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Agricultural and Forest Meteorology. **355**

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: et al. (2025).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Frontiers in Ecology and Evolution, 13,

删除[胡欣欣]: &

删除[胡欣欣]: (2017).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Remote Sensing. **9**, 66.

删除[胡欣欣]: &

删除[胡欣欣]: (2003).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜...

删除[胡欣欣]: . Remote Sensing of Environment. **87**, 42-54.

删除[胡欣欣]: &

删除[胡欣欣]: (2020).

38

952 drought on vegetation spring phenology across the Northeast China Transect, Sci.
953 Total Environ., 738, 140297, https://doi.org/10.1016/j.scitotenv.2020.140297, 2020.

954 Zhang, X., Friedl, M. A., Schaaf, C. B., Strahler, A. H., and Schneider, A.; The footprint of
955 urban climates on vegetation phenology, Geophys. Res. Lett., 31, L12209,
956 https://doi.org/10.1029/2004GL020137, 2004,

957 Zhang, X., Friedl, M. A., and Schaaf, C. B.; Global vegetation phenology from Moderate
958 Resolution Imaging Spectro radiometer (MODIS): Evaluation of global patterns and
959 comparison with in situ measurements, J. Geophys. Res.-Biogeo., 111, G04017,
960 https://doi.org/10.1029/2006JG000217, 2006,

961 Zhang, X., Chen, N., Sheng, H., Ip, C., Yang, L., Chen, Y., Sang, Z., Tadesse, T., Lim, T. P.
962 Y., Rajabifard, A., Bueti, C., Zeng, L., Wardlow, B., Wang, S., Tang, S., Xiong, Z., Li,
963 D., and Niyogi, D.: Urban drought challenge to 2030 sustainable development goals,
964 Sci. Total Environ., 693, 133536, https://doi.org/10.1016/j.scitotenv.2019.07.342,
965 2019,

966 Zheng, W., Liu, Y., Yang, X., and Fan, W.; Spatiotemporal variations of forest vegetation
967 phenology and its response to climate change in northeast China, Remote Sens., 14,
968 2909, https://doi.org/10.3390/rs14122909, 2022,

969 Zhou, D., Zhao, S., Zhang, L., and Liu, S.; Remotely sensed assessment of urbanization
970 effects on vegetation phenology in China's 32 major cities, Remote Sens. Environ.,
971 176, 272–281, https://doi.org/10.1016/j.rse.2016.02.010, 2016,

972

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 自动设置

删除[胡欣欣]: . Science of the Total Environment. **738**, 140297.

删除[胡欣欣]: &

删除[胡欣欣]: (2004).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 自动设置

删除[胡欣欣]: . Geophysical Research Letters. **31**, L12209.

删除[胡欣欣]: &

删除[胡欣欣]: (2006).

设置格式[胡欣欣]: 字体: 非加粗

设置格式[胡欣欣]: 字体: 非加粗

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 自动设置

删除[胡欣欣]: . Journal of Geophysical Research: Biogeosciences. **111**.

删除[胡欣欣]:

Zhang, X., Chen, N., Sheng, H., Ip, C., Yang, L., Chen, Y., ... & Niyogi, D. (2019). Urban drought challenge to 2030 ...

设置格式[胡欣欣]: 字体颜色: 蓝色

删除[胡欣欣]: &

删除[胡欣欣]: (2022).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 自动设置

删除[胡欣欣]: . Remote Sensing. **14**, 2909.

删除[胡欣欣]: &

删除[胡欣欣]: (2016).

设置格式[胡欣欣]: 字体: (中文) 宋体, 小四, 字体颜色: 自动设置

删除[胡欣欣]: . Remote Sensing of Environment. **176**, 272-281.