



Everyday weather in a warmer world

Rhidian Thomas¹, Gilbert P. Compo^{2,3}, Steve George¹, Gabriele C. Hegerl⁴, Andrew Schurer⁴, Theodore G. Shepherd⁵, Laura C. Slivinski³, Vikki Thompson⁴, and Ed Hawkins¹

¹National Centre for Atmospheric Science, Department of Meteorology, University of Reading, Reading, UK

²Cooperative Institute for Research in Environmental Sciences, University of Colorado at Boulder, Boulder, USA

³NOAA Physical Sciences Laboratory, Boulder, USA

⁴School of GeoSciences, University of Edinburgh, Edinburgh, UK

⁵Department of Meteorology, University of Reading, Reading, UK

Correspondence: Rhidian Thomas (r.h.thomas@reading.ac.uk)

Abstract. How would the weather of a year from history be experienced in a warmer world? We reconstruct the weather of 1903 using a reanalysis system that assimilates only surface pressure observations (20CRv3) with observed SSTs, and then reconstruct it again with increased SSTs and atmospheric CO₂ levels. By assimilating the same pressure observations, the reanalysis experiments produce the same weather patterns, and so we translate the weather of 1903 into a warmer context.

5 We focus on changes in the everyday weather of four regions with a high density of historical pressure observations, where the circulation is constrained and differences between the experiments are due to the thermodynamic component of climate change. In these regions, nearly all days are warmer in the warmer world experiments, but the largest increases occur on cold days (below freezing) and hot days (above 20°C). Daily rainfall becomes more variable, even in regions where total rainfall is reduced. Fewer days experience light rain while more days experience heavy rain, and rainfall only increases on less than one
10 day in 10. This single year pair of reanalysis experiments also recovers common patterns of observed and projected long-term changes. For example, Western Mediterranean precipitation declines outside winter, but shows a small increase in winter in the absence of storm track shifts. By anchoring our analysis in weather patterns that have actually occurred, the reanalysis experiments point to how our day-to-day experience of the weather may change in a warmer world.

1 Introduction

15 Climate change is affecting the frequency and intensity of extreme weather, with implications for infrastructure adaptation and human health (Seneviratne et al., 2021). Over the 21st century, progress has been made in attributing the influence of climate change on individual extreme weather events (Stott et al., 2004; Otto et al., 2012; Clarke et al., 2022). However, while much attention has focused on extreme weather, less focus has been given to how climate change will affect the experience of everyday, non extreme weather. Though extreme weather is impactful, and can contribute disproportionately to the mean (Pendergrass, 2018), by definition it represents weather on only a small number of days. Previous work has considered how climate
20 change has and will affect the occurrence of mild weather (van der Wiel et al., 2017; Lin et al., 2019; Zhang et al., 2022) and the closely related concept of “outdoor days” (Choi et al., 2024a, b), defined variously as days with comfortable near-surface temperature and/or clear skies and minimal precipitation. Model projections broadly show an increase in mild conditions in the



mid-high latitudes and a decrease in the tropics (van der Wiel et al., 2017; Choi et al., 2024b), albeit with substantial regional
25 variation. However, regional projections of temperature and precipitation depend on uncertain future changes in large-scale
weather patterns (Shepherd, 2014).

A different approach is to consider how the thermodynamic aspects of climate change — those related to rising temperature
and humidity — will affect day-to-day weather, given no change in weather patterns. Those effects represent the most robust
aspects of regional climate change. We adopt a storyline approach (Shepherd, 2016) to explore changes in the weather of a year
30 from history. Hawkins et al. (2023a) reconstructed an extreme event from 1903, Storm Ulysses, in an atmospheric reanalysis
system, and translated its impacts into a warmer world. Using the same experiments, as well as an additional novel reanalysis
experiment described below, we present a broader survey of changes across the 12 months of 1903. The weather from all
12 months of 1903 is first reconstructed using an atmospheric reanalysis system that assimilates observations of atmospheric
pressure (including rescued pressure observations not available in the production version of the reanalysis) and uses observed
35 SSTs and sea ice concentrations as lower boundary conditions (Slivinski et al., 2019). Then, the model and data assimilation
system are completely re-run while assimilating the same pressure observations, but altering the SST boundary conditions
and atmospheric CO₂ in order to translate the historical weather into a warmer world. As the same pressure observations are
assimilated in each experiment, the experiments experience the same weather patterns, minimising uncertainty due to projected
changes in atmospheric circulation. We then explore how the thermodynamic impacts of day-to-day weather differ in a warmer
40 and moister atmosphere. For example, in the absence of circulation changes, how does a given ocean warming affect the
distribution of daily temperature? Do cold days and hot days warm by the same amount? Does rainfall become more or less
frequent, and more or less variable from day to day?

Our approach is similar to the pseudo global warming (PGW) method, where the boundary conditions for regional climate
models (RCMs) are adjusted to reflect the effects of climate change (Schär et al., 1996; Brogli et al., 2023; Lenderink et al.,
45 2025). These adjustments, often referred to as PGW “deltas”, are chosen such that the imposed dynamical driving at the
RCM boundaries is unchanged in the warmer world simulation. Similarly to the reanalysis experiments, there is no change
in intraannual variability between the control and PGW simulations, which allows robust signals to be extracted from shorter
simulations than are typically required with free-running climate models (Sato et al., 2007). The main differences between
these two methods are in how the circulation is constrained: in PGW, typically the 3D dynamics are imposed only at the edge
50 of the RCM domain, whereas the assimilation of surface pressure observations may constrain the synoptic circulation at the
surface more strongly in our reanalysis experiments. The reanalysis experiments are also global, allowing multiple locations to
be studied in a single run. The reanalysis experiments in this paper thus offer an additional and computationally efficient line
of evidence regarding the thermodynamic effects of warming on regional weather and climate.

We begin by describing the reanalysis system and assimilated pressure data before presenting a global overview of changes
55 in the warmer world experiments. The available pressure observations from 1903 are not distributed evenly in space, which
restricts the regions we can study in detail using this approach for this particular year. We therefore focus on four regions with
a high density of pressure observations in 1903. Our results are summarised in the final section.




Figure 1. Available pressure observations for the 1903 reanalysis experiments. Red circles denote land stations, blue dots denote pressure observations from ships, and orange crosses denote tropical cyclone best-track pressures. Blue boxes show well-observed regions used for analysis in later sections. 97.1% of all pressure observations in 1903 were successfully assimilated in the 20CRv3⁺ experiment. Note that some land stations have multiple observations per day and some have fewer.

2 Reanalysis experiments

We reconstruct the weather of 1903 using the NOAA-CIRES-DOE 20th Century Reanalysis version 3 data assimilation system (20CRv3, Slivinski et al. (2019)). Using an Ensemble Kalman Filter, 20CRv3 assimilates observed pressure values with a NOAA atmospheric model to provide a best estimate of the atmospheric state at each point in time. An 80-member ensemble of 3-hourly atmospheric fields is produced on a 0.7° Gaussian grid. SSTs and sea ice concentrations are prescribed from observed values. The system assimilates pressure observations taken over land and ocean as well as tropical cyclone central pressures from best-track datasets (see Slivinski et al. (2019) for details of the boundary conditions and pressure observations).

We present results from three reanalysis experiments. The first experiment, denoted 20CRv3⁺, is identical to the improved reanalysis in Hawkins et al. (2023a). It consists of a reconstruction of 1903 using the same SST and radiative forcings as in the production version of 20CRv3, with two further changes: a refinement of the data assimilation scheme, and the inclusion of many additional rescued pressure observations over Europe. These changes sharpen features of the weather fields and improve the utility of the reconstructions; see the appendices of Hawkins et al. (2023a) for details. Fig. 1 shows the pressure observations used in the 20CRv3⁺ experiment. Some regions are much richer in historical pressure observations than others. We expect the atmospheric circulation to be better constrained in these regions than in regions with fewer pressure observations, and so we focus most of our analysis on four regions with a high density of observations in 1903, shown by the blue boxes in Fig. 1. These regions cover northwestern Europe (NW_Europe), the western Mediterranean (Western_Med), the East Coast of the US (US_East_Coast), and Southeastern Australia (SE_Aus).

In addition to 20CRv3⁺, two further experiments are performed to translate the weather of 1903 into a warmer world. The first, denoted +2K, is as in Hawkins et al. (2023b): a uniform +2K perturbation is added to the SSTs, with radiative forcings kept at 1903 values. In the second, denoted +2K+CO₂, in addition to the +2K SST perturbation, global atmospheric CO₂ concentration is increased to 530ppm. Differences between the two warmer world experiments therefore indicate the direct role




Figure 2. Time series of 3-hourly 2m-temperature (dotted line) and precipitation rate (solid line) averaged over central England (3-0°W, 51.5-54.0°N, see boxed region in Fig. 8). The 20CRv3⁺ experiment is in black and the +2K+CO₂ experiment is in red (see Fig. S1 for the +2K experiment). Lines show the ensemble means and shading denotes the 10–90% range across the ensemble. Blue markers show daily mean temperatures from the Central England Temperature (CET) series (Legg et al., 2025), plotted at 9am each day. Vertical bars on each marker indicate the CET daily maximum and minimum temperatures. The grey dashed line in the top panel indicates the air frost threshold (0°C).

of the CO₂ radiative forcing. The SST and CO₂ perturbations are idealised and do not reflect observed or projected patterns of change — they are chosen to illustrate the role of ocean warming and atmospheric composition on the weather of 1903 in a simple way. In both experiments, the same pressure observations are assimilated as in 20CRv3⁺, so that the atmospheric flow is similarly constrained in all three experiments; we thus isolate the thermodynamic impacts of the perturbations on the reconstructed weather of 1903. We analyse 12 months of data (from January to December 1903) in each experiment. Each warmer world experiment is spun up for 6 months prior to January 1903 to allow the atmosphere to adjust to the perturbed boundary conditions. The 20CRv3⁺ and +2K experiments are as in Hawkins et al. (2023a) and Hawkins et al. (2023b), while the +2K+CO₂ experiment is novel.

An example of everyday weather in a well observed region is shown in Fig. 2, which shows 3-hourly time series of 2m-temperature and precipitation in 20CRv3⁺ and the +2K+CO₂ experiment averaged over Central England (the boxed area in Figs. 8b–d) for January and July 1903. The equivalent plots for the +2K experiment are shown in Fig. S1. Temperatures in 20CRv3⁺ are in good agreement with observed daily values from the Central England Temperature series (CET) (Legg et al., 2025), which represents observed average temperatures over a similar region. As no temperature observations are assimilated, this comparison provides an independent verification of the reanalysis skill in reproducing observed weather in this region. The warmer world simulations show day-to-day fluctuations in temperature and precipitation in phase with those of 20CRv3⁺,



which indicates that the synoptic circulation is well constrained by the assimilated observations (the experiments experience
95 the “same” weather). Temperatures in the warmer world experiments are consistently warmer than in 20CRv3⁺, with little
overlap between the 20CRv3⁺ and warmer world ensemble shading, especially in January. The grey dashed line in the top
panel indicates the air frost threshold (0°C). Eight nights in the 20CRv3⁺ ensemble-mean dip below this threshold, while only
five do in the +2K+CO₂ experiment — a reduction of 60% in this small sample. For comparison, the annual number of air
frost days in the UK has declined by around a quarter since the 1980s (Kendon et al., 2025). Precipitation peaks in the warmer
100 world experiments generally line up with those in 20CRv3⁺, though rainfall is less well constrained by the assimilated pressure
observations than is near-surface temperature.

3 Overview of global changes

Before focusing on regional aspects, we first present a global overview of changes in the warmer world experiments. Fig. 3
shows globally and annually averaged 2m-temperature, precipitable water (PWAT) and precipitation rate in each experiment.
105 The distributions show the spread of values across the 80 ensemble members. The spatial patterns of the ensemble mean
temperature and precipitation changes are shown in Fig. 4. As seen in Fig 3a, globally, the +2K experiments are 2.1K warmer
than 20CRv3⁺ in the ensemble mean, with the additional 0.1K of warming arising due to enhanced warming over land (Fig. 4a).
This highlights the prominent role of oceanic warming in driving continental temperatures, even in the absence of changes in
CO₂ forcing, such as through the moistening of continental air and hence enhanced downwelling longwave (LW) radiation over
110 land (Compo and Sardeshmukh, 2009). Interestingly, however, we also find an important role for direct CO₂-induced warming,
with a further 0.2K of global warming in the +2K+CO₂ experiment due to additional enhanced warming over extratropical
land (Fig. 4b). This corresponds to a larger global land/ocean warming ratio (1.35 in +2K+CO₂, compared to 1.21 in +2K).
Both warming ratio values are notably smaller in the warmer world experiments than in climate models and observations (e.g.
Sutton et al., 2007; Compo and Sardeshmukh, 2009; Byrne and O’Gorman, 2018); the warmer world experiments are idealised
115 and differ from observed climate change in several ways, including the uniform SST increase and the lack of changes in other
forcing agents such as tropospheric aerosols (which are constant at modern day levels in 20CRv3), so we would not expect
the ratios to match the real world values precisely. The direct CO₂-induced warming is small over tropical land (Fig. 4b–c),
likely due to masking of its radiative forcing by high clouds and upper-tropospheric water vapour (Merlis, 2015). It has a
larger warming effect in the extratropics, particularly over arid regions such as the Sahara and the Western US. In the latter
120 region, additional warming of 0.6K or more is seen due to the higher CO₂ levels, which is around a fifth of the total warming
in the +2K+CO₂ runs. This is likely to be a purely radiative consequence of the CO₂ forcing, as the high density of pressure
observations in the Western US will strongly constrain the circulation in each experiment (Fig. 1). Thus, even in the absence
of circulation changes, the direct warming effect of atmospheric CO₂ varies regionally and can be large in the extratropics.

Higher surface temperatures are associated with increased atmospheric moisture content in the warmer world experiments, as
125 seen in globally averaged precipitable water (PWAT, Fig. 3b). The PWAT increases follow the same ordering as the temperature
distributions. Accounting for the additional warming in the +2K+CO₂ runs, the relative increase in PWAT is similar in both




Figure 3. Globally and annually averaged 2m-temperature (a), total-column precipitable water (PWAT) (b), and precipitation rate (c) in the reanalysis experiments. Curves show the density of values across the 80 ensemble members in each experiment.

experiments and is broadly in line with Clausius-Clapeyron scaling at 8.5%/K and 8.2%/K respectively (Trenberth et al., 2005; O’Gorman and Muller, 2010; Wan et al., 2024).

Global precipitation also increases in the warmer world simulations (Fig. 3c), but more slowly than atmospheric moisture.

130 Latent heating from precipitation is balanced globally by LW cooling of the atmosphere, which increases as the surface and atmosphere warm. This explains why a smaller precipitation increase is seen in the +2K+CO₂ experiment than in +2K, despite a larger increase in atmospheric moisture: additional CO₂ reduces LW emission to space and thus a smaller increase in precipitation is required to balance it. Hence, in isolation, the higher CO₂ concentration acts to blunt the precipitation increase that arises due to surface warming (Allen and Ingram, 2002; Pendergrass and Hartmann, 2014a). The increase of 2.4%/K in the




Figure 4. Annual mean changes in 2m-temperature (a–c) and precipitation (d–f) between the reanalysis experiments. Changes are shown as differences between experiments for temperature, and as percentage changes between experiments for precipitation. The value at the top right of each plot is the global (area-weighted) average of the changes at each gridpoint (note that this is different to the changes in the global mean shown in Fig. 3). Hatching obscures non-significant changes at the 5% level using a two-tailed Student’s t-test. The colour bar in panels (a)–(b) is centred on a change of +2K as this is the perturbation applied to the SSTs.

135 +2K+CO₂ experiment is notably similar to values seen in climate model experiments (Allen and Ingram, 2002), despite the lack of strict energy conservation in the reanalysis system due to the inclusion of analysis increments.

The spatial pattern of precipitation change is shown in Figs. 4d–e. In many respects, the pattern of change in the warmer world simulations resembles the canonical projected response to greenhouse warming in comprehensive climate models (Semenov and Bengtsson, 2002; Lee et al., 2021), despite the specificity of the 1903 SST boundary conditions and atmospheric
 140 circulation. Precipitation increases over much of the extratropics, while drying is seen over the Mediterranean and subtropical ocean basins. Precipitation intensity over land increases everywhere, but land areas including Africa, the western US, and Mediterranean experience fewer wet days (Fig. S2). Much of the drying over subtropical oceans is due to the inclusion of CO₂ (Fig. 4f), which may reflect a shift of convection from ocean to land in association with the enhanced land warming driven by increased CO₂ (He and Soden, 2017). The sparse observations over much of the tropics mean that the atmospheric circulation
 145 is less constrained in these regions; in the following sections, we therefore focus on changes in the four well-observed regions identified in Fig. 1.



Temperature increase in the warmer world experiments




Figure 5. Change in ensemble-mean, daily-mean 2m-temperature in the warmer world experiments versus 20CRv3⁺, as a function of temperature percentile in 20CRv3⁺. The pairwise difference between experiments is calculated for every (lat, lon, day) pair and is then binned according to the temperature value of that (lat, lon, day) in 20CRv3⁺. Solid lines show the mean difference in each bin. The blue dotted line shows the pairwise difference between the +2K+CO₂ and +2K experiments (again binned according to 20CRv3⁺ temperature). Only land gridcells are used. The horizontal dashed line indicates a change of +2°C and the vertical dotted lines show absolute temperatures in 20CRv3⁺.

4 Regional changes in everyday weather

4.1 Temperature changes

Annual mean temperatures in the warmer world experiments are consistently higher than in 20CRv3⁺. But do cool and warm
150 days change by the same amount? Fig. 5 shows the pairwise change in near-surface temperature in the warmer world experiments compared to 20CRv3⁺, as a function of the 20CRv3⁺ daily mean temperature, for the four well-observed regions indicated in Fig. 1.

Fewer than 1% of points in each bin become cooler in the +2K or +2K+CO₂ experiments compared to 20CRv3⁺ (not shown). In the +2K experiment, temperatures increase by 2–2.5°C on what could be considered typical days in each region (between
155 the 20th and 80th percentiles of daily temperature in 20CRv3⁺). Higher CO₂ in the +2K+CO₂ experiment leads to a roughly constant additional warming of 0.2–0.4°C on these days. However, three of the four regions show much larger temperature increases on cold days (particularly days below freezing; Figs. 5a–c) in both the +2K and +2K+CO₂ experiments. Analysis of the surface heat budget reveals a reduction in the fraction of solar radiation that is reflected by the land surface on days below 5°C in these regions (Fig. S3), likely due to reduced snow cover. Thus, in the warmer world simulations, proportionally less
160 sunlight is reflected by snow and more is absorbed by the darker land surface, leading to amplified warming of cold days. The lack of cold-day amplification in SE_Aus (Fig. 5d) is consistent with the low occurrence of freezing temperatures.



Amplified warming is also seen for the hottest days in the Western_Med, US_East_Coast, and SE_Aus regions, with increases of 3°C or more on days above 30°C (Figs. 5b–d). In all regions, an increase in evaporative cooling is seen in the +2K experiment for days below the 70th percentile (Fig. S3); however, for the hottest days, the evaporative cooling increase is smaller. In the US East Coast region, the sign reverses, with a sharp reduction in evaporative cooling seen in the +2K experiment for days above the 97th percentile of temperature (Fig. S3c). Negative soil moisture anomalies can amplify heat waves, as the land surface is less able to lose heat through evaporation (Seneviratne et al., 2006; Fischer et al., 2007). This suggests that the hottest days in the US_East_Coast region are days with low soil moisture and thus reduced ability for the land surface to cool evaporatively, leading to larger increases in near-surface air temperature. In the Western_Med, evaporative cooling is also reduced on days above approximately the 70th percentile of temperature, but not on the very hottest days (matching the weaker amplification of warming in the +2K experiment on these days). Changes in latent heat flux contribute less to amplified warming of hot extremes in NW_Europe — perhaps indicating that the hottest days are less limited by soil moisture — and in SE_Aus, where the amplification of hot extremes appears related to reduced cloud cover and increased warming from incident solar radiation (Compo and Sardeshmukh, 2009).

Heat waves and cold spells as experienced in nature often coincide with particular atmospheric circulation patterns, such as the well-known relationship between cold air outbreaks over Northern Europe and the negative phase of the North Atlantic Oscillation (Sui et al., 2020). Long-lived anticyclones in the summer months can similarly lead to heat waves in the extratropics (Pfahl and Wernli, 2012; Wehrli et al., 2019). Feedbacks between the atmospheric circulation and land surface can also precondition the land surface in ways that amplify hot extremes, such as by reducing soil moisture in the months prior to a heatwave (Fischer et al., 2007). Changes in the number of hot and cold days can, in general, be due to changes in the occurrence of such persistent weather patterns, in addition to the direct thermodynamic effects of increased radiative forcing. The reanalysis experiments suggest that, even if large-scale weather patterns remain unchanged, daily cold and hot extremes will experience amplified warming in a warmer world. This highlights the role of land surface feedbacks in addition to the atmospheric circulation in amplifying extreme heat events (Wehrli et al., 2019).

4.2 Precipitation changes

Next we analyse how the global precipitation changes in Figs. 3 and 4 compare to changes in the well-observed regions. Additionally, are changes in mean precipitation accompanied by changes in variability? Figs. 6a–d show the change in mean precipitation and daily precipitation variability in +2K+CO₂ compared to 20CRv3⁺ for our selected regions. The global increase in annual mean precipitation is also seen in the NW_Europe, US_East_Coast, and SE_Aus regions. The distribution spread is notably wider in SE_Aus than in the other regions, which may reflect the lower observation density (Fig. 1) and hence more weakly constrained circulation.

In contrast to the other regions, mean precipitation declines in Western_Med. This is consistent with climate model experiments which show a reduction in Mediterranean precipitation in response to anthropogenic forcing. Observations also show a decline in Mediterranean precipitation, although at present the forced signal may be difficult to distinguish from multidecadal variability (Vicente-Serrano et al., 2025). In models, precipitation reduction is often associated with poleward expansion of the




Figure 6. (a)–(d): percentage changes in mean and variability of daily precipitation in the +2K+CO₂ experiment compared to 20CRv3⁺ across the whole year. Variability is calculated as the standard deviation of daily precipitation. The dashed grey lines indicate the local Clausius-Clapeyron scaling at 7%/K. (e)–(h): Changes in mean Western_Med precipitation in each season compared to 20CRv3⁺. Each quantity is calculated at each grid cell before being averaged over the region, using all 12 months in each experiment. Only land grid cells are used. The violin bodies show the range of changes possible by resampling the 80 ensemble members in each experiment, and the horizontal violin bars indicate the median, 5th and 95th percentiles.

subtropical dry zones leading to fewer Mediterranean cyclones (Zappa et al., 2015b, a), which is not represented in our experiments. Changes in large-scale circulation are more important in winter (Brogli et al., 2019), whereas the summer precipitation response is due mainly to thermodynamic factors such as changes in humidity and the land/sea warming contrast (Brogli et al., 2019; Barcikowska et al., 2020). We find that the annual mean change in Fig. 6b is due to reduced rainfall outside winter (Figs. 200 6f–h), with little change in DJF (Fig. 6e). The reanalysis experiments therefore suggest that local thermodynamic factors can drive a reduction in annual mean precipitation over the Western Mediterranean, even in the absence of large-scale circulation changes. Figs. 6e–h also show that, while the SST warming causes a small increase in winter rainfall but a decrease in the other seasons, the additional CO₂ in the +2K+CO₂ experiments always reduces rainfall in our experiments regardless of season. As the SSTs are the same in both warmer world experiments, the main effect of the CO₂ is to enhance land warming at the expense 205 of the ocean, potentially leading to reduced relative humidity over land and reduced precipitation.

Despite a decline in total precipitation, daily precipitation variability increases in Western_Med as well as in all other regions, meaning the warmer world experiences larger swings in rainfall from day to day (Figs. 6a–d). An increase in precipitation variability is consistent with modelling and theory (Pendergrass et al., 2017; Zhang et al., 2021) and with trends emerging in observations (Zhang et al., 2024). Variability increases more than the mean in each region, but by less than the availability of 210 moisture as defined by the local Clausius-Clapeyron scaling (grey dashed lines in Figs. 6a–d). Precipitation is proportional to




Figure 7. Precipitation characteristics of the NW_Europe (a–c) and Western_Med (d–f) regions. (a, d): Number of dry days, defined as days experiencing less than 1mm of rain (here 1.04mm due to numerical rounding). The median is shown by the horizontal line and the violin shows the distribution of ensemble members. (b, e) Daily precipitation occurrence in 20CRv3⁺ and +2K+CO2. The blue line shows the percentage change in each bin in +2K+CO2. Error bars show the 10–90% range through sampling all combinations of ensemble members, and vertical dashed lines show percentiles of the precipitation distribution in 20CRv3⁺. (c, f) Ratio of large event cutoff s_L in +2K+CO2 vs 20CRv3⁺ (see text). Only land gridcells are used for all calculations. Bins in (b) and (e) are logarithmically distributed with the smallest nonzero bin centred at 1.11 mm/day and successive bin widths increasing by 14.7%.

the product of moisture and vertical motion (Sardeshmukh et al., 2015) and so, for fixed vertical motion, precipitation variability scales with moisture availability (Pendergrass et al., 2017). Thus, the lower rate of increase in the warmer world experiment indicates a change in the distribution of vertical motion (Pendergrass and Gerber, 2016). This suggests that, although the synoptic circulation in the reanalysis experiments is constrained by the pressure assimilation, there is some flexibility for the statistics of vertical velocity to change, which may explain why the precipitation amplification in Fig. 2 is less robust than the temperature response.

Daily precipitation densities in the 20CRv3⁺ and +2K+CO2 experiments are shown by the grey and red points, respectively, in Figs. 7b and 7e. The curves follow a similar shape in both NW_Europe and Western_Med (the other regions are shown in Fig. S4), which approximately follow a gamma distribution (Martinez-Villalobos and Neelin, 2018):

$$p_s(s) \propto s^{-\tau} \exp[s/s_L] \quad (1)$$

where s is daily precipitation size, τ is a power law exponent, and s_L is a cutoff scale beyond which occurrence falls exponentially. Warming leads to an increase in s_L and therefore larger relative increases in the occurrence of heavier precipitation (Neelin et al., 2017; Martinez-Villalobos and Neelin, 2018). We find s_L increases approximately in line with Clausius-Clapeyron scaling in our regions (Figs. 7c and 7f, and S3c and S3f). However, this does not mean that the number of days with



225 heavy precipitation also increases by 7%/K: instead, the increase is exponential with precipitation rate, as can be seen from the blue curves in Figs. 7b and 7e which show the percentage change for each precipitation bin. The number of days with moderate to heavy rainfall — above the 90–95th percentile of all days in 20CRv3⁺ — increases exponentially with precipitation rate in the warmer world.

On the other hand, the number of days with light to moderate rainfall (below the 90–95th percentile in 20CRv3⁺) decreases slightly in the +2K+CO₂ experiment. The magnitude of these changes is smaller than at heavy rain rates and the sign of change is more uncertain, however the ensemble-mean values show similar characteristics in all four regions (see also Figs. S3b and S3e). This highlights the counterintuitive point, made previously by Allen and Ingram (2002), that the time-mean precipitation increases seen in previous figures (Figs. 3, 4, 6) occur only on less than one day in 10, with most wet days actually becoming drier in the warmer world experiments.

235 The number of dry days in each experiment is shown in Figs. 7a, 7d, S4a and S4d, defined as days experiencing less than 1mm of rain as in previous studies (here 1.04mm due to numerical rounding; Polade et al., 2014; Douville et al., 2023). The number of dry days also increases in three of the four regions in +2K+CO₂, with the exception of SE_Australia. The additional CO₂ appears to be an important factor for dry days over NW_Europe and Western_Med, but not over US_East_Coast and SE_Australia. Previous studies have also found an increase in the number of dry days in warmer world experiments (Polade et al., 2014), as is also seen on the global scale in the reanalysis experiments (Figs. S2a–c). One possible mechanism for this is a reduction in relative humidity over land leading to increased convective inhibition (CIN) and hence longer intervals between convective events (Dai et al., 2020; Chadwick et al., 2022), although convective precipitation is likely to be less well constrained by the assimilated pressure observations than rain associated with frontal systems.

5 October 1903: an extreme month

245 An advantage of the counterfactual reanalysis experiments is the ability to look at changes in climate impacts over a variety of timescales, as the assimilation process means weather patterns in the warmer world counterfactuals remain close to the original (factual) reanalysis. While short, intense weather events can lead to acute damages, extreme months or seasons, or even sequences of seasons, can also be significant for sectors such as agriculture (Noone et al., 2025). Here we present an example of an event falling between the annual timescales discussed in previous sections and the synoptic timescales that are typically the subject of storyline attribution studies.

October 1903 is the wettest recorded calendar month for the UK (220.0mm, in a series back to 1836 (Hollis et al., 2019)) and in the longer England and Wales Precipitation series (218.1mm, in a series back to 1766 (Alexander and Jones, 2000)). Fig. 8a shows the October 1903 rainfall total in a gridded observational product that interpolates between in-situ measurements (Hollis et al., 2019). Large parts of Wales, Southwest England, and western Scotland recorded rainfall totals exceeding 200mm. 255 These regions also receive the most rain in 20CRv3⁺ (Fig. 8b), though the magnitude is smaller in the reanalysis than in the observations. We would not expect perfect quantitative agreement with the observations as the reanalysis grid is too coarse to fully resolve the orography in these regions; similar underestimates are seen over western regions of Great Britain in the




Figure 8. October rainfall in (a) observations (HadUKGrid at 60km resolution (Hollis et al., 2019)) and (b)–(d) the reanalysis experiments. The value in the top right of panel (b) is the average rainfall over the UK and Ireland in 20CRv3⁺. The corresponding increase compared to 20CRv3⁺ is shown in the top right of panels (c) and (d). Panel (e) shows daily rainfall over Central England (the red boxed region, also used in Fig. 2; using reanalysis data from 9am–9am UTC to match the observational definition). Shading shows the 10–90% range across the ensemble and grey dashed lines indicate all-year daily rainfall percentiles in 20CRv3⁺ for 1903. Note that the boxes used to define the Central England region differ by a small amount in the observations compared to the reanalyses due to differences in the grids used (cf. the boxes in panels a and b).

case of Storm Ulysses earlier in 1903 (Hawkins et al., 2023a). Output from the reanalysis could be downscaled using regional atmospheric models to better represent rainfall over mountainous regions if desired.

260 Over flatter terrain, the reanalysis is in better agreement with the observations, such as over a less mountainous region in Central England (Fig. 8e). As no rainfall data is assimilated, this represents an additional independent verification of the reanalysis. Daily rainfall is shown in Fig. 8e for comparison with the observations, but the higher frequency rainfall data in the reanalysis reveals several intense, sub-daily rainfall events during October that are obscured in the daily mean (e.g. the 3hr-rain rate centred at 6th Oct 15:00 UTC is 44mm/day in 20CRv3⁺, Fig. S5).

265 Both warmer world experiments see increased rainfall over the UK and Ireland (Figs. 8b and 8c). The increase of 17% in the +2K experiment is equivalent to 8.8% per K of local warming, while the increase of 12% in +2K+CO2 is somewhat smaller (equivalent to 6.0%/K). These values are substantially larger than the increase in annual mean precipitation of 3.0%/K over the larger NW_Europe region in the +2K+CO2 experiment (not shown). The increase in October rainfall over the UK and Ireland is closer to, or even above, the Clausius-Clapeyron scaling seen in extreme daily rainfall events (Fig. 7c). This likely reflects

270 the unusually high occurrence of heavy rainfall events in this month, with 5 distinct events above the all-year 95th percentile of



daily precipitation in 20CRv3⁺ (Fig. 8e). Rainfall totals in extreme months or seasons may therefore scale much more rapidly with warming than the annual mean.

On sub-daily time scales, the largest changes in October rainfall occur on wet days, whereas rain rates on drier days are comparable in both experiments (Fig. S5). This supports the finding in the previous section that substantial increases in time-
275 mean rainfall can result from increased rainfall in a small number of heavy events. Notably, the warmer world rainfall rate at 6th Oct 15:00 UTC is 58mm/day, an increase of 31% over the value in 20CRv3⁺ or around 15%/K — much greater than Clausius-Clapeyron scaling, as has previously been found for sub-daily rainfall extremes (Lenderink and van Meijgaard, 2008; Guerreiro et al., 2018).

6 Discussion and conclusions

280 How would the everyday weather of a year from history be experienced in a warmer world? The reanalysis experiments in this paper share the same weather patterns, but by perturbing the SSTs and atmospheric CO₂ we can explore how the weather of 1903 would be experienced in a warmer world by residents of four well-observed regions (the boxed regions in Fig. 1). We focus on changes in two key meteorological variables which affect day-to-day perception of the weather: temperature and rainfall. Residents of these regions would experience warmer near-surface temperatures nearly every day, with 99% of days
285 becoming warmer than in the 20CRv3⁺ reconstruction of 1903. On a typical day, land areas would be warmer by around 2–2.5°C for a 2K ocean surface warming. However, increases of 2.5–3°C are seen on hot days (temperatures around 20°C or above), and larger increases of 4–5°C on days below freezing. The cold day response appears to be due to enhanced absorption of solar radiation by the land surface due to a reduction in snow cover; the warmer world version of 1903 would also be one with substantially less snow and ice. Mechanisms for the hot day response are mixed: in the US_East_Coast region it is linked
290 to reduced evaporative cooling on the hottest days, while changes in cloud cover appear to be more important in the SE_Aus region.

The everyday experience of rainfall would also be different in the warmer world. Even without substantial changes in large-scale weather patterns, annual rainfall would increase in NW_Europe and SE_Australia, decrease in Western_Med, and show little change in US_East_Coast – but in each region, daily rainfall would become more variable, meaning larger rainfall swings
295 from day to day, with implications for infrastructure and adaptation. Even in regions where the total annual rainfall increases, the increase would occur on roughly 1 day in every 10 (that is, on days above the 90th percentile of rainfall). Instead, there would be fewer drizzly days, and in three of the four regions an overall reduction in the number of rainy days when the additional CO₂ is included. Conversely, the number of days in the heavier rainfall bins grows exponentially with increasing rain rate.

300 We find that the October 1903 rainfall total over the UK and Ireland, already the wettest month on record in England and Wales, would be 12–17% higher in a +2K warmer world. The reanalysis experiments presented here are idealised experiments rather than best estimates of the current climate, however they can give an indication of how much more rain the weather conditions of October 1903 might produce today. If we assume that global temperatures have warmed by 1.5°C since 1903



(as compared to 2.3°C in the +2K+CO₂ experiment, Fig. 3a) then we estimate that the weather conditions of October 1903 would produce an extreme UK average rainfall of around 240mm in the current climate. The next wettest observed month is December 2015 at 216.3mm (Hollis et al., 2019). Future planned work using patterned SST perturbations and consistent CO₂ concentrations will provide better estimates of how the impacts of historical extreme weather events would be different in the current climate. We will also perform additional experiments to translate events into a warmer future climate to inform adaptation planning.

We also find evidence of a direct CO₂ effect on everyday weather. Higher atmospheric CO₂ leads to enhanced warming of extratropical land than from SST warming alone, contributing 19% of the total warming over the Western US in the +2K+CO₂ experiment. Higher CO₂ levels act to suppress global precipitation and reduce the number of rainy days over NW_Europe, Western_Med, and US_East_Coast.

The conclusions above consider what would happen if the same weather patterns of 1903 were to be repeated. Changes in the atmospheric circulation could lead to much larger changes in extreme temperature and precipitation, for example if persistent weather patterns were to occur more frequently (Petoukhov et al., 2013; Mann et al., 2017). The reanalysis experiments can thus be considered a strongly conditioned form of attribution (Thompson et al., 2026). However, by isolating the thermodynamic component of climate change, the reanalysis experiments can offer insight into the robust physical processes behind regional climate signals. For example, Mediterranean precipitation declines in spring, summer, and autumn in the warmer world experiments, but increases in winter, presumably due to local changes in atmospheric stability and humidity. We also find that, in the absence of shifts in the Mediterranean storm track, seasonality in the Mediterranean precipitation response is driven entirely by SST warming, as the direct effect of increased atmospheric CO₂ is to reduce Mediterranean rainfall regardless of season. Other Mediterranean climates are also projected to dry, with atmospheric circulation contributing to the recent Chilean “megadrought” (Garreaud et al., 2017, 2020). Forthcoming experiments will consider the modern period where there are more pressure observations over Chile and other Mediterranean-like climates outside Europe.

This study has presented an exploration of a single year’s weather. Our conclusions are restricted to regions with a high density of historical pressure observations, and some may be sensitive to the reanalysis system used — for example, there are longstanding problems in the simulation of light rainfall in numerical models (Chen et al., 2021), which limits confidence in the reduction of light rainfall and overall rainfall occurrence seen in the warmer world experiments. However, it is notable that even in a single year, the reanalysis experiments support changes in rainfall characteristics that have been seen in multiple generations of coupled models (Hennessy et al., 1997; Lau et al., 2013; Pendergrass and Hartmann, 2014b).

Data availability. Ensemble-means and ensemble-standard deviations for 2m-temperature and precipitation rate are available for each of the three experiments at: <https://doi.org/10.5281/zenodo.19949356>. The repository also includes the pressure observation text files used to generate Fig. 1. Individual ensemble members can be provided upon request.



335 *Author contributions.* GPC, SG, and LCS assisted with the design and running of the reanalysis experiments. EH and RT ran the reanalysis experiments. RT analysed the experiments and prepared the paper, with contributions from all co-authors. EH conceived and supervised the project.

Competing interests. The authors declare that they have no conflict of interest.

340 *Acknowledgements.* This work was funded by UKRI grant number NE/Z000203/1 and supported by the UK National Centre for Atmospheric Science. Author EH was also supported by a Co-Centre award number 22/CC/11103, with funding from Research Ireland, Northern Ireland's Department of Agriculture, Environment and Rural Affairs (DAERA) and UK Research and Innovation (UKRI). Author GPC was supported in part by NOAA cooperative agreement NA22OAR4320151 and by the NOAA Physical Sciences Laboratory. Author LS was supported by the NOAA Physical Sciences Laboratory. This work used JASMIN, the UK's collaborative data analysis environment (<https://www.jasmin.ac.uk>; Lawrence et al., 2013).



345 References

- Alexander, L. V. and Jones, P. D.: Updated precipitation series for the U.K. and discussion of recent extremes, *Atmospheric Science Letters*, 1, 142–150, <https://doi.org/10.1006/asle.2000.0016>, 2000.
- Allen, M. R. and Ingram, W. J.: Constraints on future changes in climate and the hydrologic cycle, *Nature*, 419, 224–232, <https://doi.org/10.1038/nature01092>, 2002.
- 350 Barcikowska, M. J., Kapnick, S. B., Krishnamurty, L., Russo, S., Cherchi, A., and Folland, C. K.: Changes in the future summer Mediterranean climate: contribution of teleconnections and local factors, *Earth System Dynamics*, 11, 161–181, <https://doi.org/10.5194/esd-11-161-2020>, 2020.
- Brogli, R., Sørland, S. L., Kröner, N., and Schär, C.: Causes of future Mediterranean precipitation decline depend on the season, *Environmental Research Letters*, 14, 114 017, <https://doi.org/10.1088/1748-9326/ab4438>, 2019.
- 355 Brogli, R., Heim, C., Mensch, J., Sørland, S. L., and Schär, C.: The pseudo-global-warming (PGW) approach: methodology, software package PGW4ERA5 v1.1, validation, and sensitivity analyses, *Geoscientific Model Development*, 16, 907–926, <https://doi.org/10.5194/gmd-16-907-2023>, 2023.
- Byrne, M. P. and O’Gorman, P. A.: Trends in continental temperature and humidity directly linked to ocean warming, *Proceedings of the National Academy of Sciences*, 115, 4863–4868, <https://doi.org/10.1073/pnas.1722312115>, 2018.
- 360 Chadwick, R., Pendergrass, A. G., Alves, L. M., and Moise, A.: How Do Regional Distributions of Daily Precipitation Change under Warming?, *Journal of Climate*, 35, 3243–3260, <https://doi.org/10.1175/JCLI-D-20-0864.1>, 2022.
- Chen, D., Dai, A., and Hall, A.: The Convective-To-Total Precipitation Ratio and the “Drizzling” Bias in Climate Models, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD034 198, <https://doi.org/10.1029/2020JD034198>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2020JD034198>, 2021.
- 365 Choi, Y.-W., Khalifa, M., and Eltahir, E. A. B.: Climate Change Impact on “Outdoor Days” Over the United States, *Geophysical Research Letters*, 51, e2024GL111 607, <https://doi.org/10.1029/2024GL111607>, 2024a.
- Choi, Y.-W., Khalifa, M., and Eltahir, E. A. B.: North–South Disparity in Impact of Climate Change on “Outdoor Days”, *Journal of Climate*, 37, 3269–3282, <https://doi.org/10.1175/JCLI-D-23-0346.1>, 2024b.
- Clarke, B., Otto, F., Stuart-Smith, R., and Harrington, L.: Extreme weather impacts of climate change: an attribution perspective, *Environmental Research: Climate*, 1, 012 001, <https://doi.org/10.1088/2752-5295/ac6e7d>, 2022.
- 370 Compo, G. P. and Sardeshmukh, P. D.: Oceanic influences on recent continental warming, *Climate Dynamics*, 32, 333–342, <https://doi.org/10.1007/s00382-008-0448-9>, 2009.
- Dai, A., Rasmussen, R. M., Liu, C., Ikeda, K., and Prein, A. F.: A new mechanism for warm-season precipitation response to globalwarming based on convection-permitting simulations, *Climate Dynamics*, 55, 343–368, <https://doi.org/10.1007/s00382-017-3787-6>, 2020.
- 375 Douville, H., Chadwick, R., Saint-Lu, M., and Medeiros, B.: Drivers of Dry Day Sensitivity to Increased CO₂, *Geophysical Research Letters*, 50, e2023GL103 200, <https://doi.org/10.1029/2023GL103200>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2023GL103200>, 2023.
- Fischer, E. M., Seneviratne, S. I., Vidale, P. L., Lüthi, D., and Schär, C.: Soil Moisture–Atmosphere Interactions during the 2003 European Summer Heat Wave, *Journal of Climate*, 20, 5081–5099, <https://doi.org/10.1175/JCLI4288.1>, 2007.



- 380 Garreaud, R. D., Alvarez-Garreton, C., Barichivich, J., Boisier, J. P., Christie, D., Galleguillos, M., LeQuesne, C., McPhee, J., and Zambrano-Bigiarini, M.: The 2010–2015 megadrought in central Chile: impacts on regional hydroclimate and vegetation, *Hydrology and Earth System Sciences*, 21, 6307–6327, <https://doi.org/10.5194/hess-21-6307-2017>, 2017.
- Garreaud, R. D., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., and Veloso-Aguila, D.: The Central Chile Mega Drought (2010–2018): A climate dynamics perspective, *International Journal of Climatology*, 40, 421–439, <https://doi.org/10.1002/JOC.6219>,
385 2020.
- Guerreiro, S. B., Fowler, H. J., Barbero, R., Westra, S., Lenderink, G., Blenkinsop, S., Lewis, E., and Li, X.-F.: Detection of continental-scale intensification of hourly rainfall extremes, *Nature Climate Change*, 8, 803–807, <https://doi.org/10.1038/s41558-018-0245-3>, 2018.
- Hawkins, E., Brohan, P., Burgess, S. N., Burt, S., Compo, G. P., Gray, S. L., Haigh, I. D., Hersbach, H., Kuyper, K., Martínez-Alvarado, O., McColl, C., Schurer, A. P., Slivinski, L., and Williams, J.: Rescuing historical weather observations improves quantification of severe
390 windstorm risks, *Natural Hazards and Earth System Sciences*, 23, 1465–1482, <https://doi.org/10.5194/nhess-23-1465-2023>, 2023a.
- Hawkins, E., Compo, G. P., and Sardeshmukh, P. D.: ESD Ideas: Translating historical extreme weather events into a warmer world, *Earth System Dynamics*, 14, 1081–1084, <https://doi.org/10.5194/esd-14-1081-2023>, 2023b.
- He, J. and Soden, B. J.: A re-examination of the projected subtropical precipitation decline, *Nature Climate Change*, 7, 53–57, <https://doi.org/10.1038/nclimate3157>, 2017.
- 395 Hennessy, K. J., Gregory, J. M., and Mitchell, J. F. B.: Changes in daily precipitation under enhanced greenhouse conditions, *Climate Dynamics*, 13, 667–680, <https://doi.org/10.1007/s003820050189>, 1997.
- Hollis, D., McCarthy, M., Kendon, M., Legg, T., and Simpson, I.: HadUK-Grid—A new UK dataset of gridded climate observations, *Geoscience Data Journal*, 6, 151–159, <https://doi.org/10.1002/gdj3.78>, _eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/gdj3.78>, 2019.
- 400 Kendon, M., Doherty, A., Hollis, D., Carlisle, E., Packman, S., Jevrejeva, S., Matthews, A., Williams, J., Garforth, J., and Sparks, T.: State of the UK Climate in 2024, *International Journal of Climatology*, 45, e70010, <https://doi.org/10.1002/joc.70010>, _eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/joc.70010>, 2025.
- Lau, W. K.-M., Wu, H.-T., and Kim, K.-M.: A canonical response of precipitation characteristics to global warming from CMIP5 models, *Geophysical Research Letters*, 40, 3163–3169, <https://doi.org/10.1002/grl.50420>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/grl.50420>, 2013.
- 405 Lawrence, B. N., Bennett, V. L., Churchill, J., Jukes, M., Kershaw, P., Pascoe, S., Pepler, S., Pritchard, M., and Stephens, A.: Storing and manipulating environmental big data with JASMIN, in: 2013 IEEE International Conference on Big Data, pp. 68–75, <https://doi.org/10.1109/BigData.2013.6691556>, 2013.
- Lee, J.-Y., Marotzke, J., Bala, G., Cao, L., Corti, S., Dunne, J., Engelbrecht, F., Fischer, E., Fyfe, J., Jones, C., Maycock, A., Mutemi, J., Ndiaye, O., Panickal, S., and Zhou, T.: Future Global Climate: Scenario-Based Projections and Near-Term Information, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou, pp. 553–672, Cambridge University Press, <https://doi.org/10.1017/9781009157896.006>, 2021.
- 410 Legg, T., Packman, S., Caton Harrison, T., and McCarthy, M.: An Update to the Central England Temperature Series—HadCET v2.1, *Geoscience Data Journal*, 12, e284, <https://doi.org/10.1002/gdj3.284>, _eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/gdj3.284>, 2025.



- Lenderink, G. and van Meijgaard, E.: Increase in hourly precipitation extremes beyond expectations from temperature changes, *Nature Geoscience*, 1, 511–514, <https://doi.org/10.1038/ngeo262>, 2008.
- 420 Lenderink, G., de Vries, H., van Meijgaard, E., de Rooy, W., van Ulft, L., Thompson, V., Qiu, X., and Fowler, H. J.: A pseudo global warming based system to study how climate change affects high impact rainfall events, *Weather and Climate Extremes*, 49, 100781, <https://doi.org/10.1016/j.wace.2025.100781>, 2025.
- Lin, L., Ge, E., Chen, C., and Luo, M.: Mild weather changes over China during 1971–2014: Climatology, trends, and interannual variability, *Scientific Reports*, 9, 2419, <https://doi.org/10.1038/s41598-019-38845-8>, 2019.
- 425 Mann, M. E., Rahmstorf, S., Kornhuber, K., Steinman, B. A., Miller, S. K., and Coumou, D.: Influence of Anthropogenic Climate Change on Planetary Wave Resonance and Extreme Weather Events, *Scientific Reports*, 7, 45242, <https://doi.org/10.1038/srep45242>, 2017.
- Martinez-Villalobos, C. and Neelin, J. D.: Shifts in Precipitation Accumulation Extremes During the Warm Season Over the United States, *Geophysical Research Letters*, 45, 8586–8595, <https://doi.org/10.1029/2018GL078465>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018GL078465>, 2018.
- 430 Merlis, T. M.: Direct weakening of tropical circulations from masked CO₂ radiative forcing, *Proceedings of the National Academy of Sciences*, 112, 13167–13171, <https://doi.org/10.1073/pnas.1508268112>, 2015.
- Neelin, J. D., Sahany, S., Stechmann, S. N., and Bernstein, D. N.: Global warming precipitation accumulation increases above the current-climate cutoff scale, *Proceedings of the National Academy of Sciences*, 114, 1258–1263, <https://doi.org/10.1073/pnas.1615333114>, 2017.
- Noone, C., Noone, S., McClean, D., and Thorne, P.: The Beast from the East and drought of summer 2018: an example
435 of compound event impacts upon Ireland’s agricultural sector, *Weather*, 80, 56–61, <https://doi.org/10.1002/wea.4586>, _eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/wea.4586>, 2025.
- Otto, F. E. L., Massey, N., van Oldenborgh, G. J., Jones, R. G., and Allen, M. R.: Reconciling two approaches to attribution of the 2010 Russian heat wave, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2011GL050422>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2011GL050422>, 2012.
- 440 O’Gorman, P. A. and Muller, C. J.: How closely do changes in surface and column water vapor follow Clausius–Clapeyron scaling in climate change simulations?, *Environmental Research Letters*, 5, 025207, <https://doi.org/10.1088/1748-9326/5/2/025207>, 2010.
- Pendergrass, A. G.: What precipitation is extreme?, *Science*, 360, 1072–1073, <https://doi.org/10.1126/science.aat1871>, 2018.
- Pendergrass, A. G. and Gerber, E. P.: The Rain Is Askew: Two Idealized Models Relating Vertical Velocity and Precipitation Distributions in a Warming World, *Journal of Climate*, 29, 6445–6462, <https://doi.org/10.1175/JCLI-D-16-0097.1>, 2016.
- 445 Pendergrass, A. G. and Hartmann, D. L.: The Atmospheric Energy Constraint on Global-Mean Precipitation Change, *Journal of Climate*, 27, 757–768, <https://doi.org/10.1175/JCLI-D-13-00163.1>, 2014a.
- Pendergrass, A. G. and Hartmann, D. L.: Changes in the Distribution of Rain Frequency and Intensity in Response to Global Warming, *Journal of Climate*, 27, 8372–8383, <https://doi.org/10.1175/JCLI-D-14-00183.1>, 2014b.
- Pendergrass, A. G., Knutti, R., Lehner, F., Deser, C., and Sanderson, B. M.: Precipitation variability increases in a warmer climate, *Scientific*
450 *Reports*, 7, 17966, <https://doi.org/10.1038/s41598-017-17966-y>, 2017.
- Petoukhov, V., Rahmstorf, S., Petri, S., and Schellnhuber, H. J.: Quasiresonant amplification of planetary waves and recent Northern Hemisphere weather extremes, *Proceedings of the National Academy of Sciences*, 110, 5336–5341, <https://doi.org/10.1073/pnas.1222000110>, 2013.



- Pfahl, S. and Wernli, H.: Quantifying the relevance of atmospheric blocking for co-located temperature extremes in the North-
455 ern Hemisphere on (sub-)daily time scales, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL052261>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2012GL052261>, 2012.
- Polade, S. D., Pierce, D. W., Cayan, D. R., Gershunov, A., and Dettinger, M. D.: The key role of dry days in changing regional climate and
precipitation regimes, *Scientific Reports*, 4, 4364, <https://doi.org/10.1038/srep04364>, 2014.
- Sardeshmukh, P. D., Compo, G. P., and Penland, C.: Need for Caution in Interpreting Extreme Weather Statistics, *Journal of Climate*, 28,
460 9166–9187, <https://doi.org/10.1175/JCLI-D-15-0020.1>, 2015.
- Sato, T., Kimura, F., and Kitoh, A.: Projection of global warming onto regional precipitation over Mongolia using a regional climate model,
Journal of Hydrology, 333, 144–154, <https://doi.org/10.1016/j.jhydrol.2006.07.023>, 2007.
- Schär, C., Frei, C., Lüthi, D., and Davies, H. C.: Surrogate climate-change scenarios for regional climate models, *Geophysical Research
Letters*, 23, 669–672, <https://doi.org/10.1029/96GL00265>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/96GL00265>,
465 1996.
- Semenov, V. and Bengtsson, L.: Secular trends in daily precipitation characteristics: greenhouse gas simulation with a coupled AOGCM,
Climate Dynamics, 19, 123–140, <https://doi.org/10.1007/s00382-001-0218-4>, 2002.
- Seneviratne, S., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Di Luca, A., Ghosh, S., Iskandar, I., Kossin, J., Lewis, S., Otto, F.,
Pinto, I., Satoh, M., Vicente-Serrano, S., Wehner, M., and Zhou, B.: Weather and Climate Extreme Events in a Changing Climate, in:
470 *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovern-
mental Panel on Climate Change*, pp. 1513–1766, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA,
<https://doi.org/10.1017/9781009157896.013>, 2021.
- Seneviratne, S. I., Lüthi, D., Litschi, M., and Schär, C.: Land–atmosphere coupling and climate change in Europe, *Nature*, 443, 205–209,
<https://doi.org/10.1038/nature05095>, 2006.
- 475 Shepherd, T. G.: Atmospheric circulation as a source of uncertainty in climate change projections, *Nature Geoscience*, 7, 703–708,
<https://doi.org/10.1038/ngeo2253>, 2014.
- Shepherd, T. G.: A Common Framework for Approaches to Extreme Event Attribution, *Current Climate Change Reports*, 2, 28–38,
<https://doi.org/10.1007/s40641-016-0033-y>, 2016.
- Slivinski, L. C., Compo, G. P., Whitaker, J. S., Sardeshmukh, P. D., Giese, B. S., McColl, C., Allan, R., Yin, X., Vose, R., Titchner, H.,
480 Kennedy, J., Spencer, L. J., Ashcroft, L., Brönnimann, S., Brunet, M., Camuffo, D., Cornes, R., Cram, T. A., Crouthamel, R., Domínguez-
Castro, F., Freeman, J. E., Gergis, J., Hawkins, E., Jones, P. D., Jourdain, S., Kaplan, A., Kubota, H., Blancq, F. L., Lee, T.-C., Lorrey, A.,
Luterbacher, J., Maugeri, M., Mock, C. J., Moore, G. K., Przybylak, R., Pudmenzky, C., Reason, C., Slonosky, V. C., Smith, C. A., Tinz,
B., Trewin, B., Valente, M. A., Wang, X. L., Wilkinson, C., Wood, K., and Wyszyński, P.: Towards a more reliable historical reanalysis:
Improvements for version 3 of the Twentieth Century Reanalysis system, *Quarterly Journal of the Royal Meteorological Society*, 145,
485 2876–2908, <https://doi.org/10.1002/qj.3598>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/qj.3598>, 2019.
- Stott, P. A., Stone, D. A., and Allen, M. R.: Human contribution to the European heatwave of 2003, *Nature*, 432, 610–614,
<https://doi.org/10.1038/nature03089>, 2004.
- Sui, C., Yu, L., and Vihma, T.: Occurrence and drivers of wintertime temperature extremes in Northern Europe during
1979–2016, *Tellus A: Dynamic Meteorology and Oceanography*, 72, 1–19, <https://doi.org/10.1080/16000870.2020.1788368>, _eprint:
490 <https://doi.org/10.1080/16000870.2020.1788368>, 2020.



- Sutton, R. T., Dong, B., and Gregory, J. M.: Land/sea warming ratio in response to climate change: IPCC AR4 model results and comparison with observations, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2006GL028164>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006GL028164>, 2007.
- Thompson, V., Ermis, S., and Athanase, M.: The need for multi-method extreme event attribution, *Weather*, 81, 40–45, <https://doi.org/10.1002/wea.7779>, _eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/wea.7779>, 2026.
- 495 Trenberth, K. E., Fasullo, J., and Smith, L.: Trends and variability in column-integrated atmospheric water vapor, *Climate Dynamics*, 24, 741–758, <https://doi.org/10.1007/s00382-005-0017-4>, 2005.
- van der Wiel, K., Kapnick, S. B., and Vecchi, G. A.: Shifting patterns of mild weather in response to projected radiative forcing, *Climatic Change*, 140, 649–658, <https://doi.org/10.1007/s10584-016-1885-9>, 2017.
- 500 Vicente-Serrano, S. M., Tramblay, Y., Reig, F., González-Hidalgo, J. C., Beguería, S., Brunetti, M., Kalin, K. C., Patalen, L., Kržič, A., Lionello, P., Lima, M. M., Trigo, R. M., El-Kenawy, A. M., Eddénjal, A., Türkes, M., Koutroulis, A., Manara, V., Maugeri, M., Badi, W., Mathbout, S., Bertalanič, R., Bocheva, L., Dabanli, I., Dumitrescu, A., Dubuisson, B., Sahabi-Abed, S., Abdulla, F., Fayad, A., Hodzic, S., Ivanov, M., Radevski, I., Peña-Angulo, D., Lorenzo-Lacruz, J., Domínguez-Castro, F., Gimeno-Sotelo, L., García-Herrera, R., Franquesa, M., Halifa-Marín, A., Adell-Michavila, M., Noguera, I., Barriopedro, D., Garrido-Perez, J. M., Azorin-Molina, C., Andres-Martin, M.,
- 505 Gimeno, L., Nieto, R., Llasat, M. C., Markonis, Y., Selmi, R., Ben Rached, S., Radovanović, S., Soubeyroux, J.-M., Ribes, A., Saidi, M. E., Bataineh, S., El Khalki, E. M., Robaa, S., Boucetta, A., Alsafadi, K., Mamassis, N., Mohammed, S., Fernández-Duque, B., Cheval, S., Moutia, S., Stevkov, A., Stevkova, S., Luna, M. Y., and Potopová, V.: High temporal variability not trend dominates Mediterranean precipitation, *Nature*, 639, 658–666, <https://doi.org/10.1038/s41586-024-08576-6>, 2025.
- Wan, N., Lin, X., Pielke Sr., R. A., Zeng, X., and Nelson, A. M.: Global total precipitable water variations and trends over the period 1958–2021, *Hydrology and Earth System Sciences*, 28, 2123–2137, <https://doi.org/10.5194/hess-28-2123-2024>, 2024.
- 510 Wehrli, K., Guillod, B. P., Hauser, M., Leclair, M., and Seneviratne, S. I.: Identifying Key Driving Processes of Major Recent Heat Waves, *Journal of Geophysical Research: Atmospheres*, 124, 11 746–11 765, <https://doi.org/10.1029/2019JD030635>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019JD030635>, 2019.
- Zappa, G., Hawcroft, M. K., Shaffrey, L., Black, E., and Brayshaw, D. J.: Extratropical cyclones and the projected decline of winter Mediterranean precipitation in the CMIP5 models, *Climate Dynamics*, 45, 1727–1738, <https://doi.org/10.1007/s00382-014-2426-8>, 2015a.
- 515 Zappa, G., Hoskins, B. J., and Shepherd, T. G.: The dependence of wintertime Mediterranean precipitation on the atmospheric circulation response to climate change, *Environmental Research Letters*, 10, 104 012, <https://doi.org/10.1088/1748-9326/10/10/104012>, 2015b.
- Zhang, J., You, Q., Ren, G., and Ullah, S.: Projected changes in mild weather frequency over China under a warmer climate, *Environmental Research Letters*, 17, 114 042, <https://doi.org/10.1088/1748-9326/ac9c70>, 2022.
- 520 Zhang, W., Furtado, K., Wu, P., Zhou, T., Chadwick, R., Marzin, C., Rostron, J., and Sexton, D.: Increasing precipitation variability on daily-to-multiyear time scales in a warmer world, *Science Advances*, 7, eabf8021, <https://doi.org/10.1126/sciadv.abf8021>, 2021.
- Zhang, W., Zhou, T., and Wu, P.: Anthropogenic amplification of precipitation variability over the past century, *Science*, 385, 427–432, <https://doi.org/10.1126/science.adp0212>, 2024.