

Orbital-scale ~~hydroclimateprecipitation~~ variations in the western
Qaidam Basin during the Late Pliocene: evidence from magnetic
parameters and median grain-size

Zeng Luo^{1*}, Hansheng Wang^{1,2}, Xiaoli Yu¹, Zhongbao Zhang³, Zhifen Song¹, Jing
Yang¹, Wanfeng Chen¹, Qingda Su^{4*}, Yang Lu⁵, Shun Hua¹

¹College of Geography and Tourism, Hunan University of Arts and Science, Changde,
Hunan, China 415000.

²Key Laboratory of Western China's Environmental System (Ministry of Education),
College of Earth and Environmental Sciences, Lanzhou University, Lanzhou, Gansu,
China 730000

³College of Urban and Environmental Sciences, Hubei Normal University, Huangshi,
Hubei, China 435002.

~~⁴Shandong Provincial Key Laboratory of Water and Soil Conservation and
Environmental Protection, College~~⁴College of Resources and Environment, Linyi
University, Linyi, Shandong, China 276000.

⁵Meteorological Bureau of Shunde District, Foshan, Guangdong, China 528399.

**Correspondence to:* Zeng Luo (luozeng2020@foxmail.com); Qingda Su
(suqingda@foxmail.com).

设置了格式

Abstract

A recent study suggests that the precipitation records of the western Qaidam Basin during the middle Piacenzian warm period (mPWP, 3.264-3.025 Ma) are broadly consistent with those of the monsoon-influenced regions, and are highly sensitive to 20-kyr precessional cycles, implying that the East Asian summer monsoon (EASM) precipitation intensified and ~~migrated~~penetrated farther westward into the Asian interior during this warm interval. However, it remains unclear whether such a precession-dominated pattern persisted throughout the Late Pliocene or was restricted to specific intervals. To address this question, we present new magnetic parameter and median grain-size records from the Gansen (~~GS~~) section for ~3.6-3.25 Ma and 2.95-1.8 Ma, and integrate them with the previously published records for 3.25-2.95 Ma to assess orbital-scale precipitation variations in the western Qaidam Basin during the Late Pliocene. The results show that ~~the EASM-related summer moisture reached~~ extended its influence into the western Qaidam Basin during ~3.6-3.25 Ma, although ~~the associated~~ precipitation was weaker than during 3.25-2.95 Ma. Spectral analysis further reveals that precipitation variations during ~3.6-3.3 Ma were characterized mainly by dominant eccentricity-scale (~100-125-kyr cyclicity) variability rather than the previously reported precession-dominated pattern during 3.25-2.95 Ma, ~~suggesting which may reflect~~ a nonlinear response to insolation forcing. This contrast indicates that the precession-dominated pattern was not a persistent feature throughout the Late Pliocene in the western Qaidam Basin. Instead, our results suggest that under different climatic background states, the dominant processes ~~controlling~~influencing

the westward penetration of summer moisture into the western Qaidam Basin were also different, and that the 20-kyr precessional signal and ~~100-kyr periodicities~~eccentricity-scale variability mainly reflect these differences in forcing response. In addition, the ~~similar phase variations~~broad correspondence between these precipitation records and the benthic ~~oxygen isotope~~ $\delta^{18}\text{O}$ stack at the 40-kyr band ~~imply~~suggests that high-latitude ice-volume variability, potentially including Antarctic ice-sheets-sheet variability, may also have modulated ~~hydroclimatic precipitation~~ variations in the basin-during ~3.6-3.3 Ma. These findings provide new insight into how background climate conditions may shape ~~the orbital-scale response of precipitation, and more broadly~~ hydroclimatic ~~variations, responses~~ in the Asian interior during ~~warm periods~~the Late Pliocene.

Keywords: Late Pliocene; Qaidam Basin; Monsoon precipitation; Insolation forcing; Orbital cycles.

1 Introduction

With continued global warming, knowledge of climate variations during past warm periods has major implications for understanding future climate change. The middle Piacenzian warm period (mPWP: 3.264-3.025 Ma), a particularly well-documented warm interval during the Late Pliocene, has long been considered an ideal geological analog for interpreting the path of future anthropogenic warming (Ravelo et al., 2004; Raymo et al., 2006; Dowsett et al., 2010; Salzmann et al., 2011; Haywood et al., 2011; Lunt et al., 2012; Burke et al., 2018; Su et al., 2024; Luo et al., 2021, 2024; Zhou and Liu, 2025). Compared to the present, this interval was characterized by similar orbital configurations and geographic boundary conditions (Haywood et al., 2011; Su et al., 2024), higher atmospheric carbon dioxide concentrations (400-450 ppmv) (Bartoli et al., 2011; Badger et al., 2013; Haywood et al., 2016), small Arctic ice sheets (Brigham-Grette et al., 2013; Rahaman et al., 2020), elevated global temperatures (Burke et al., 2018), and increased sea levels (Hearty et al., 2020). Over the past two decades, numerous studies based on proxy reconstructions from loess deposits, lake sediments, and marine sediments, together with climate model simulations, have suggested that the mid-Piacenzian world was not only warmer but also wetter on most continents than today (Reguero et al., 2007; Salzmann et al., 2008, 2011; Brigham-Grette et al., 2013; Herbert et al., 2016; Yang et al., 2018; Vieira et al., 2018; Zhou and Liu, 2025). Similarly, in the Asian interior, the extent of arid regions decreased during the mPWP. For example, arid conditions in the Qaidam Basin, an important part of the Asian interior, were alleviated during the

mPWP, contributing to a regional contraction of the inland arid zone (Koutsodendris et al., 2019; Huang et al., 2019; Schwarz et al., 2022; Su et al., 2024). This occurred because the influence of the East Asian summer monsoon (EASM) ~~migrated~~extended westward toward the western Qaidam Basin, bringing relatively more water vapor to this area and changing the local moisture conditions (Huang et al., 2019, 2021; Schwarz et al., 2022).

At the orbital timescale, recent precipitation records from the Gansen (GS) section of the western Qaidam Basin reveal dominant 20-kyr precessional cycles during the mPWP (Luo et al., 2024). These cycles are consistent with records from the Chinese Loess Plateau and the South China Sea (Wehausen and Brumsack, 2002; Wan et al., 2010; Luo et al., 2021), further indicating that the EASM strongly influenced precipitation in the western Qaidam Basin during this warm interval. However, EASM influence ~~on precipitation in the Qaidam Basin~~ has also been ~~inferred~~suggested for other intervals of the Late Pliocene. For example, pollen records from the SG-1b core in the western Qaidam Basin show that annual mean precipitation during 3.5-3.0 Ma was about ten times higher than today and was associated with EASM (Koutsodendris et al., 2019; Schwarz et al., 2022). ~~In addition,~~ ~~hematite~~Hematite-content records from the HTG section of the western Qaidam Basin also suggest that the EASM ~~also~~ affected precipitation variations in this region during the Late Pliocene (Su et al., 2024). These studies ~~imply that EASM-related moisture may also have influenced precipitation during other Late Pliocene intervals, but whether this was accompanied by persistent precession-dominated cyclicity remains~~

unresolved. Owing to the lack of high-resolution and continuous precipitation records, it is still unclear whether precession-dominated cyclicity was restricted to the mPWP or persisted throughout the Late Pliocene, which limits our understanding of precipitation variations in the Asian interior during the Pliocene warmth. Existing studies have also reached different conclusions regarding the dominant orbital periodicity of Late Pliocene precipitation in the Qaidam Basin. For example, Su et al. (2024) proposed that precipitation variations were dominated by 40-kyr obliquity cyclicity, whereas Schwarz et al. (2022) argued for dominant 100-kyr eccentricity cyclicity. This discrepancy highlights the need for additional high-resolution records to determine whether precession-dominated precipitation variability was persistent throughout the Late Pliocene or restricted to specific warm intervals such as the mPWP, however, inferred different dominant orbital periodicities, with Schwarz et al. (2022) arguing for dominant 100-kyr eccentricity cycles and Su et al. (2024) proposing dominant 40-kyr obliquity cycles. Consequently, it remains unclear whether the precession-dominated precipitation pattern identified during the mPWP was restricted to that interval or persisted throughout the Late Pliocene. Resolving this question requires additional high-resolution precipitation records from other intervals of the Late Pliocene.

To determine whether the precession-dominated precipitation pattern previously identified during 3.25–2.95 Ma was restricted to that interval or persisted through other stages of the Late Pliocene, we present new magnetic-parameter and median grain-size records from the GS section in the western Qaidam

Basin for ~3.6-3.25 Ma and 2.95-1.8 Ma. By integrating these new data with the previously published records for 3.25-2.95 Ma, we construct a longer precipitation record spanning ~3.6-1.8 Ma and ~~examine howcompare the~~ orbital-scale ~~characteristics of precipitation variations in the western Qaidam Basin varied~~ across different Late Pliocene intervals.

2 Geologic setting

The Qaidam Basin is located in the northeastern part of the Qinghai-Tibet Plateau, extending approximately from 35.9° to 39.3°N and from 90.3° to 99.3°E (Fig. 1). It is surrounded by several mountain ranges, including the Kunlun Mountains to the south, the Altyn Tagh Mountains to the northwest, the Qilian Mountains to the northeast, and the Ela Mountains to the east (Fig. 1b). This mountain-surrounding geographical pattern renders the Qaidam Basin relatively enclosed, forming a unique climate and ecological environment. Because the basin is far from major moisture ~~sourcesources~~ and lies in the rain shadow of the Tibetan Plateau, mean annual precipitation is very low, ranging from only 50 to 200 mm, whereas mean annual evaporation is as high as ~3000 mm (Cai et al., 2012; Wang et al., 2013; Yang et al., 2020; Luo et al., 2024). As a result, the Qaidam Basin is characterized by a hyper-arid climate, with widespread deposition of evaporite minerals such as halite, carbonates, and gypsum (Li et al., 2010; Wang et al., 2013; Luo et al., 2018, 2024).




Figure 1. Locations of the study GS site. (a) The modern summer atmospheric circulation pattern of Asia and the location of the GS section labeled with a red spot. ISM: Indian summer monsoon. EASM: East Asian summer monsoon. (b) Detailed geological map of the Qaidam Basin and the location of the GS section in the Qaidam Basin. ATF: Altyn Tagh fault, KLF: Kunlun fault, NQF: North Qaidam fault. This figure is modified after Bush et al. (2016). (c) A closer look at the GS section with a satellite image showing the sample locations and their distribution between the lower segment (0-226 m in stratigraphic height) and the upper segment (201-478 m in stratigraphic height) of the stratigraphic section.




Figure 2. Lithology, age model and representative sediments in the GS section.

The studied GS section (37.717°N, 92.38°E to 37.73°N, 92.41°E) lies on the southern flank of the anticline which is close to Jianshi Shan on the northwestern margin of the Qaidam Basin (Fig. 1b). Owing to the presence of thick salt and weathering crusts in the upper part of the section, only the lower 478 m was sampled. However, because the GS section is a composite stratigraphic succession composed of lower and upper segments that overlap between 201 and 226 m, the effective stratigraphic thickness is 453 m (Fig. 1c). The strata consist mainly of massive and laminated siltstone, interbedded with gypsum-rich siltstone and occasional sandstone

beds (Fig. 2). Gypsum layers occur throughout the sequence, whereas halite layers are confined to the intervals between 0 and 105–95 m and between 262–244 and 453 m (Fig. 2). Previous paleomagnetic analyses indicate that the sampled 0–453 m interval of the GS section spans ~3.6–1.8 Ma (Luo et al., 2024).

3 Materials and methods

3.1 Chronology of the GS section

The chronology of the GS section is based on the previously published paleomagnetic framework (of Luo et al., 2024), in which). In that study, the observed polarity sequence was of the sampled 0–453 m interval was well correlated with the geomagnetic polarity time scale (GPTS), and 10 paleomagnetic age control points were used to establish the age model for the section (Supplementary Table S1, Supplementary Fig. S1a), indicating that the sampled 0–453 m interval spans ~3.6–1.8 Ma (Luo et al., 2024). This age model provides the basis for plotting the new paleoclimatic records and conducting subsequent spectral analyses. Ages between adjacent paleomagnetic tie points were obtained by linear interpolation, and the sedimentation rates calculated between adjacent tie points indicate non-uniform sedimentation throughout the GS section (Supplementary Table S1, Supplementary Fig. S1b). Because the age model is constrained by discrete paleomagnetic tie points, some uncertainty remains in the interpolated ages between adjacent control points. Nevertheless, the paleomagnetic chronology is broadly consistent with the Plio-Quaternary age assignment of the GS strata on the regional geological map (Luo et al., 2024), providing an independent geological framework for the studied interval.

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式

Furthermore, the same paleomagnetic framework has successfully resolved significant orbital cycles in the previously published high-resolution GS record for 3.25-2.95 Ma (Luo et al., 2024), supporting the use of this chronology for orbital-scale analysis in the studied interval.

3.2 Environmental magnetic and grain-size measurements

For magnetic ~~analyses~~measurements, low-frequency and high-frequency magnetic susceptibility (χ) of the 149 samples ~~was~~were measured at 976 Hz and 15616 Hz, respectively, using the AGICO multifunction spinner Kappabridge (MFK1-FA). ~~frequency~~Frequency-dependent magnetic susceptibility (χ_{fd}) was calculated as $\chi_{fd} = \chi_{976\text{Hz}} - \chi_{15616\text{Hz}}$. Isothermal remanent magnetization (IRM) was then acquired in applied fields of 1.2 T and -0.3 T, respectively. Hard isothermal remanent magnetization (HIRM) was subsequently calculated as $(\text{IRM}_{1.2\text{T}} + \text{IRM}_{-0.3\text{T}})/2$. Finally, the χ_{fd}/HIRM (the ratio of the frequency-dependent magnetic susceptibility to the intensity of the isothermal remanent magnetization) values of the 149 samples were calculated. All magnetic measurements were carried out at the Paleomagnetism Laboratory of the China University of Geosciences, Beijing, China.

For grain-size analysis, all 149 samples were pretreated before measurement. Approximately 0.3 g of each bulk sample was first treated with 10 mL of 30% H_2O_2 to remove organic matter and then with 10 mL of 10% HCl to remove carbonates. The samples were subsequently immersed in distilled water for 12 h. After removal of the supernatant, the samples were ultrasonically dispersed in 10 mL of 10% $(\text{NaPO}_3)_6$ solution for 10 minutes. Grain-size distributions were then measured using a Malvern

设置了格式: 字体颜色: 自动设置

Mastersizer 2000 laser grain-size analyzer. These measurements were conducted at the Key Laboratory of Western China's Environmental Systems, Lanzhou University.

3.3 Spectral analysis

The average temporal spacing between samples is ~4 kyr for the ~3.6-3.3 Ma interval and ~18 kyr for the 2.95~1.8 Ma interval. These values describe the spacing of the analyzed samples, rather than the chronological resolution or precision of the paleomagnetic age model. Because ages between adjacent paleomagnetic tie points were obtained by linear interpolation, chronological uncertainty between tie points may exceed the average sample spacing and may have a greater influence on the interpretation of the shorter 20- and 40-kyr components than on eccentricity-scale variability. Therefore, the spectral analysis was focused mainly on the densely sampled ~3.6-3.3 Ma interval. For this interval, the χ_{rd} /HIRM and median grain-size records were first converted from stratigraphic height to time based on the paleomagnetic age model of the GS section and then interpolated to evenly spaced 1-kyr time series. The 1-kyr interpolation was used only to regularize the time spacing of the records for spectral analysis and band-pass filtering, and does not add independent temporal information. Power spectra were calculated using the REDFIT procedure implemented in PAST 3 software, which estimates the red-noise background and tests spectral peaks against an AR(1) red-noise null model. Spectral peaks exceeding the 90% red-noise confidence level were regarded as statistically significant. No additional detrending was applied prior to the REDFIT analysis. To test whether the main eccentricity-scale spectral result was sensitive to the

interpolation procedure, we repeated the REDFIT analysis using a 2-kyr interpolation time step. The eccentricity-scale (~100-125-kyr) variability remained evident in both proxy records, indicating that this feature is not an artifact of the 1-kyr interpolation (Supplementary Fig. S2).

Band-pass filtering was further applied to extract the 20-kyr, 40-kyr, and 100-kyr components. The central frequencies and bandwidths used for filtering were 0.05 kyr⁻¹ and 0.012 kyr⁻¹ for the 20-kyr band, 0.024 kyr⁻¹ and 0.004 kyr⁻¹ for the 40-kyr band, and 0.01 kyr⁻¹ and 0.002 kyr⁻¹ for the 100-kyr band, respectively. The filtered GS records were then compared with the corresponding filtered insolation, eccentricity, and benthic $\delta^{18}\text{O}$ stack.

4 Results

Because the χ_{fd} /HIRM and median grain-size records for 3.25-2.95 Ma have been ~~previously~~ published ~~previously~~ (Luo et al., 2024), the present study focuses on the newly generated data for ~~~3.6-3.25 Ma and 2.95-~1.8 Ma~~ and ~~integrates them~~ with the published 3.25-2.95 Ma records to construct an integrated record spanning ~~~3.6-~1.8 Ma. combines them with the published record for comparison.~~ The ~~combined/reconstructed~~ χ_{fd} /HIRM and median grain-size records ~~for 3.6-3.25 Ma and 2.95-1.8 Ma, together with the previously published records for 3.25-2.95 Ma (Luo et al., 2024),~~ are shown in Fig. 3.

设置了格式

设置了格式





Figure 3. Variation trends of the χ_{fd}/HIRM and median grain size records during 3.6–1.8 Ma. The blue rectangles represent the halite layers.

Based on lithology and the variation patterns of the χ_{fd}/HIRM and median grain size records, the paleoclimatic sequence can be divided into three phases (Fig. 3). During ~3.6–3.25 Ma (0–117 m; Phase I), χ_{fd}/HIRM values show large-amplitude fluctuations, ranging from 41.39×10^{-6} to $105 \times 10^{-6} \text{ Am}^{-1}$, with an average of $37.63 \times 10^{-6} \text{ Am}^{-1}$ (Fig. 3). Median grain size values range from 8.3 to 12.4 μm , with an average of 10.48 μm (Fig. 3). These two records display similar variation trends. In addition, the three thick halite layers observed in the stratigraphic

sequence correspond well to three low-value intervals in both the χ_{fd}/HIRM and median grain-size records (Fig. 3). During 3.25-2.95 Ma (117-220 m; Phase II), the published χ_{fd}/HIRM record exhibits high-frequency fluctuations, with an average of $52.6 \times 10^{-6} \text{ Am}^{-1}$ (Fig. 3). The published median grain-size record shows similar fluctuations, with an average of $11.2 \mu\text{m}$ (Fig. 3). No thick halite layers occur in the stratigraphic sequence during this interval (Fig. 3). During 2.95-1.8 Ma (220-453 m; Phase III), χ_{fd}/HIRM values range from 16.6×10^{-6} to $6668 \times 10^{-6} \text{ Am}^{-1}$, with an average of $3635 \times 10^{-6} \text{ Am}^{-1}$ (Fig. 3). Median grain-size values range from 5.42 to $13.5 \mu\text{m}$, with an average of $8.3637 \mu\text{m}$ (Fig. 3). Lithological observations indicate that halite layers occur more frequently during 2.95-1.8 Ma than during 3.6-3.25 Ma (Fig. 3).



带格式的: 缩进: 首行缩进: 8.1 字符

Figure 3. Variation trends of the χ_{fd}/HIRM and median grain-size records during $\sim 3.6\text{--}1.8\text{ Ma}$. The blue rectangles represent the halite layers.




Figure 4. Results of spectral analyses of the reconstructed χ_{fd}/HIRM and median grain-size records during $\sim 3.6\text{--}3.3\text{ Ma}$ and the published χ_{fd}/HIRM and median grain-size records during $3.25\text{--}2.95\text{ Ma}$. The orbital cycles revealed by these two paleoclimatic records during $3.6\text{--}3.3\text{ Ma}$ passed the 90% confidence level.

Our sampling yields an average resolution of ~ 4 Spectral analysis was performed on the interpolated 1-kyr for the time series using the REDFIT procedure in PAST 3 software. The spectral analysis for $\sim 3.6\text{--}3.3\text{ Ma}$ interval (82 samples) and 19 kyr is based on the newly generated records, whereas those for the $3.25\text{--}2.95\text{--}1.8\text{ Ma}$ interval (62 samples), indicating that the sampling resolution for the $3.6\text{--}3.3\text{ Ma}$ interval is sufficient to detect precession, obliquity, are from Luo et al. (2024). The red-noise background and eccentricity-scale cycles, the 90% red-noise confidence level are shown, and spectral peaks above the 90% confidence level are considered significant.

Spectral analysis shows that the strongest peak in the χ_{fd}/HIRM record is dominated by occurs at $\sim 100\text{ kyr}$ cycles, with relatively weak 40 and 23 kyr cycles,

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 蓝色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

298 ~~and, whereas the strongest peak in the median grain-size record is characterized~~
299 ~~mainly by occurs closer to ~125-kyr cycles, together with weak 42- and 23-kyr cycles~~
300 ~~kyr (Fig. 4).~~ These results indicate prominent eccentricity-scale (~100-125-kyr)
301 variability during ~3.6-3.3 Ma, accompanied by weaker ~40- and ~23-kyr
302 components. In addition, several ~~non-orbital cycles~~ periodicities, including ~30-kyr
303 and semiprecessional (~12-kyr) ~~cycles~~ components, are also present and may reflect
304 beats or harmonics of ~~primary orbital cycles~~ cycles (Fig. 4). A sensitivity test based
305 on a 2-kyr interpolation time step yielded similar eccentricity-scale spectral peaks in
306 both proxy records, further supporting that the main eccentricity-scale variability is
307 not caused by the 1-kyr interpolation procedure (Supplementary Fig. S2).

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 红色


设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 红色

设置了格式: 字体颜色: 自动设置



带格式的：缩进：首行缩进： 4.72 字符

Figure 5. Comparison of the GS χ_{fd}/HIRM and median grain-size records with insolation, eccentricity and benthic $\delta^{18}\text{O}$ stack during ~3.6-2.95 Ma. (a) June insolation gradient (Laskar et al., 2004). (b) GS χ_{fd}/HIRM . (c) GS median grain-size. (d) Eccentricity (Laskar et al., 2004). (e) Benthic $\delta^{18}\text{O}$ stack (Lisiecki and Raymo, 2005). The χ_{fd}/HIRM and median grain-size data during 3.25-2.95 Ma have been published by Luo et al. (2024). The green, gray and pink bands emphasize the potential corresponding relationships between χ_{fd}/HIRM , median grain-size, benthic $\delta^{18}\text{O}$ stack, eccentricity and insolation.

The dominant eccentricity-scale (~100-125-kyr cyclicity) variability in the χ_{fd}/HIRM and median grain-size records during ~3.6-3.3 Ma differs from the

设置了格式：字体颜色：自动设置

设置了格式：字体颜色：自动设置

设置了格式：字体颜色：自动设置

设置了格式：字体颜色：自动设置

319 previously reported precession-dominated pattern during 3.25-2.95 Ma (Fig. 4).
 320 Comparison with eccentricity (Laskar et al., 2004) shows that high-eccentricity
 321 intervals ~~broadly correspond well~~ to peak intervals in both records between $\sim$ 3.6 and
 322 3.3 Ma (Fig. 5), consistent with ~~dominant 100-kyr cyclicity~~ an eccentricity-scale signal
 323 during this interval. This interpretation is further supported by ~~the~~ broadly similar
 324 ~~phase relationship~~ filtered variations between eccentricity and both records ~~at~~ in the
 325 ~~100-kyr eccentricity~~ band (Fig. 6a, e). In contrast, most high-insolation intervals
 326 shown by the June insolation gradient (30°N to 30°S) do not correspond to peak
 327 values in either record between $\sim$ 3.6 and 3.3 Ma ~~(Fig. 5)~~, unlike the relationship
 328 observed during 3.25-2.95 Ma ~~(Fig. 5)~~ Luo et al., 2024. This mismatch, together with
 329 the weak 20-kyr signal (Fig. 6d, h), suggests that insolation did not directly force GS
 330 paleoclimatic variations during $\sim$ 3.6-3.3 Ma. In addition, five high-value intervals in
 331 the χ_{fd} /HIRM and median grain-size records ~~broadly correspond well with~~ those in
 332 the benthic oxygen isotope $\delta^{18}O$ stack (Lisiecki and Raymo, 2005), similar to the
 333 relationship observed during 3.25-2.95 Ma (Fig. 5). Both records also show ~~a~~ a similar
 334 ~~phase relationship with~~ filtered variations to the benthic oxygen isotope $\delta^{18}O$ stack at
 335 the 40-kyr band (Fig. 6b, f), ~~but an inverse phase relationship do not show a clear~~
 336 correspondence with insolation at the same band (Fig. 6c, g), suggesting that ~~ice~~
 337 ~~sheets, rather than insolation, likely influenced~~ the 40-kyr signal in the GS
 338 ~~paleoclimatic~~ χ_{fd} /HIRM and median grain-size records ~~was more closely linked to~~
 339 high-latitude ice-volume variability than to insolation
 340 forcing.

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置


设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置

设置了格式: 字体颜色: 自动设置



设置了格式: 字体: 五号, 字体颜色: 自动设置




Figure 6. Comparison of the filtered χ_{fd}/HIRM and median grain-size records from the GS section with the eccentricity, the filtered June insolation gradient and the filtered benthic $\delta^{18}\text{O}$ stack during ~3.6-3.3 Ma. (a) and (e) χ_{fd}/HIRM and median grain-size records at the 100-kyr band with eccentricity. (b) and (f) χ_{fd}/HIRM , median grain-size and benthic $\delta^{18}\text{O}$ stack at the filtered 40-kyr band. (c) and (g) χ_{fd}/HIRM , median grain-size and June insolation gradient at the filtered 40-kyr band. (d) and (h) χ_{fd}/HIRM , median grain-size and June insolation gradient at the filtered 20-kyr band. The 20-kyr central frequency = 0.05 kyr^{-1} , and the bandwidth = 0.012 kyr^{-1} . The 40-kyr central frequency = 0.024 kyr^{-1} , and the bandwidth = 0.004 kyr^{-1} . The 100-kyr central frequency = 0.01 kyr^{-1} , and the bandwidth = 0.002 kyr^{-1} .

5 Discussion

5.1 Late Pliocene-~~Quaternary~~Early Pleistocene precipitation variations in the western Qaidam Basin

The χ_{fd} /HIRM proxy has been shown to be sensitive to precipitation variations in arid to semiarid environments because χ_{fd} reflects pedogenic nanometer-scale ferrimagnetic grains, whereas HIRM mainly reflects hematite formed under relatively dry conditions (Nie et al., 2017; Luo et al., 2024). Therefore, their ratio records the relative balance between wetter and drier weathering conditions. ~~Previous studies further indicate that this proxy can be applied in Qaidam fluvial-lacustrine sediments. However, to interpret χ_{fd} /HIRM as a precipitation proxy in the GS section, potential non-climatic influences, including post-depositional dissolution, evaporite dilution, and changes in sediment source, need to be considered. First, thermal demagnetization results reported by Luo et al. (2024) for the GS section indicate the preservation of magnetite-bearing remanence carriers, suggesting that pervasive post-depositional reductive dissolution was unlikely to have dominated the magnetic signal. Previous studies from Qaidam fluvial-lacustrine sediments further support the applicability of χ_{fd} /HIRM as a precipitation proxy in settings where magnetic minerals~~ were not significantly dissolved after deposition (Nie et al., 2017; Su et al., 2019; Ren et al., 2020~~–~~). Second, although evaporite-bearing lithologies occur in the section, the χ_{fd} /HIRM variations are difficult to explain by simple evaporite dilution alone, given their broad consistency with the median grain-size record (Fig. 5). Third, the GS section is dominated mainly by fine-grained lacustrine deposits with relatively

设置了格式

uniform lithology, suggesting that major changes in sediment provenance were unlikely to be the primary driver of the observed χ_{fd} /HIRM variations. These considerations support the use of χ_{fd} /HIRM as a precipitation-sensitive proxy in the GS section, although minor sedimentological or post-depositional influences, including selective alteration of fine magnetic particles, evaporite dilution, and small provenance changes, cannot be entirely excluded. Median grain size can also serve as an indirect precipitation proxy because the grain size of fluvial, deltaic or nearshore lacustrine sediments is closely related to hydrodynamic conditions, which, in turn, are strongly influenced by runoff and precipitation. In the Qaidam Basin, ~~Luo et al., 2024~~. The median grain size can also serve as an indirect precipitation proxy because coarser particles in closed and shallow lacustrine settings generally reflect stronger hydrodynamic conditions associated with increased runoff and precipitation (Su et al., 2019; Yang et al., 2020; Luo et al., 2024). ~~In the GS section, the similar variation patterns of~~ We therefore interpret χ_{fd} /HIRM and median grain size during 3.6–3.25 Ma suggest that these two primarily as proxies consistently recorded for precipitation variations in the GS section.

The deposition of halite is generally associated with intensified aridity in arid regions (Song et al., 2013; Zhang et al., 2013; Gu et al., 2015; Luo et al., 2018). Therefore, the absence of halite layers likely reflects relatively wetter conditions during 3.25–2.95 Ma. This interpretation is consistent with paleoclimatic records from the HTG section in the western Qaidam Basin, which also indicate climatic wetting during the mPWP (Su et al., 2024). ~~In addition, the~~ The χ_{fd} /HIRM and median grain

设置了格式

设置了格式

设置了格式


设置了格式

设置了格式

设置了格式

设置了格式

size records reported here, together with the previously published GS records (Luo et al., 2024), further show that precipitation during 3.25-2.95 Ma was higher than during ~3.6-3.25 Ma and 2.95-1.8 Ma. ~~Taken together, these~~ These observations suggest that the relatively wet climate recorded in the GS section during the mPWP was likely associated with enhanced precipitation under a strengthened EASM. Previously published records from the GS section further indicate that precipitation variations there were significantly influenced by the intensified EASM during the mPWP (Luo et al., 2024). A possible mechanism for this monsoon intensification is the intensification and westward extension of the western Pacific subtropical high (Huang et al., 2019). Climate model simulations suggest that enhanced land-sea thermal contrast between the East Asian continent and the equatorial western Pacific during the mPWP may have promoted the intensification and westward expansion of the western Pacific subtropical high, thereby enhancing the EASM (Huang et al., 2019). This inference may further imply that, under continued global warming, the Qaidam Basin could receive increased summer moisture, potentially associated with a westward extension of the western Pacific subtropical high that favors a northwestward shift in EASM precipitation.



带格式的：缩进：首行缩进： 0.67 字符

Figure 7. Comparison of the GS χ_{ld} /HIRM and median grain size records with the paleoclimatic records affected by the EASM precipitation during 3.6–2.95 Ma. (a) Magnetic susceptibility at low frequency (χ_{lf}) from the Chaona section (Song et al., 2001). (b) Magnetic susceptibility at low frequency (χ_{lf}) from the Shilou section (Ao et al., 2016). (c) The ratio of free iron ($\text{Fe}_2\text{O}_3(\text{f})$) to total iron ($\text{Fe}_2\text{O}_3(\text{t})$) content of the Dianguan section (Yang et al., 2018). (d) GS χ_{ld} /HIRM. (e) GS median grain size. (f) Annual mean precipitation variations based on the pollen records from the SG-1b core (Schwarz et al., 2022).

The precipitation variations in the GS section during ~ 3.6 –2.95 Ma are similar to those recorded in the Chinese Loess Plateau and the SG-1b core from the western Qaidam Basin, both of which were influenced by the EASM (Fig. 7). This similarity suggests that EASM-related summer moisture likely also reached the GS section, at least during ~ 3.6 –3.25 Ma. From 2.95–1.8 Ma, however, the relatively large temporal spacing between samples precludes identification of clear temporal trends in the precipitation records (Fig. 3) the low sampling resolution of the precipitation

records precludes identification of clear temporal trends (Fig. 3) and prevents detailed comparison with precipitation records from the monsoon region. Nevertheless, the more frequent occurrence of halite layers suggests intensified aridification during 2.95–1.8 Ma interval (Fig. 3). This result is supported by other paleoenvironmental records from the western Qaidam Basin. For example, pollen support this interpretation. Pollen records from the SG-3 core suggest that the Qaidam Basin was a drier environment no later than ~3 Ma (Miao et al., 2013). The oxygen isotope records from the HTG section indicate that the Qaidam Basin has been characterized by a generally drier climate in the basin since 2.9 Ma (Su et al., 2024). Taken together, these observations suggest Overall, this evidence suggests enhanced aridification in the western Qaidam Basin following the Late Pliocene warmth.




Figure 7. Comparison of the GS χ_{lf} /HIRM and median grain-size records with the paleoclimatic records affected by the EASM precipitation during ~3.6–2.95 Ma. (a) Magnetic susceptibility at low frequency (χ_{lf}) from the Chaona section (Song et al.,

设置了格式

设置了格式

设置了格式

带格式的：缩进：首行缩进： 0.67 字符

2001). (b) Magnetic susceptibility at low frequency (χ_{lf}) from the Shilou section (Ao et al., 2016). (c) The ratio of free iron ($\text{Fe}_2\text{O}_3(\text{f})$) to total iron ($\text{Fe}_2\text{O}_3(\text{t})$) content of the Pianguan section (Yang et al., 2018). (d) GS χ_{fd}/HIRM . (e) GS median grain-size. (f) Annual mean precipitation variations based on the pollen records from the SG-1b core (Schwarz et al., 2022).

5.2 Nonlinear response of precipitation to insolation forcing during 3.6–3.3 Ma

Ma

The published ~~precipitation-GS~~ records ~~of the GS section reveals~~ show that precipitation in the western Qaidam Basin was dominated by the 20-kyr ~~precession~~ ~~cyclic~~ precessional cycles during the mPWP, and suggest that the June insolation gradient (30°N minus 30°S) directly forced precipitation variations in the Qaidam Basin during this interval (Luo et al., 2024). In contrast, the reconstructed ~~precipitation records of the GS section reveal that precipitation in the Qaidam Basin was dominated by the 100-kyr eccentricity cyclic~~ GS records indicate prominent eccentricity-scale (~100-125-kyr) variability during ~3.6-3.3 Ma. Although the ~3.6-3.3 Ma interval spans only ~0.3 Myr and therefore contains only about three eccentricity-scale cycles (Fig. 5), both χ_{fd}/HIRM and median grain-size records show their strongest spectral power in the eccentricity band, with the χ_{fd}/HIRM peak at ~100 kyr and the median grain-size peak closer to ~125 kyr (Fig. 4). This interpretation is further supported by the similar long-term variations in the two precipitation records and by the broadly similar patterns of their filtered eccentricity-scale components (Fig. 3 and 6).

It is widely believed that the insolation variations associated with the eccentricity signal of the Earth are much weaker than those associated with the precession and obliquity signals, although eccentricity can affect the insolation variations through the modulation of precession (Luo et al., 2018; Wang et al., 2023). ~~Thus, our precipitation records~~ Therefore, the eccentricity-scale variability during ~3.6-3.3 Ma may ~~show~~reflect a nonlinear response of precipitation to the insolation variations during ~~3.6-3.3 Ma forcing~~. This ~~finding~~interpretation is in line with the magnetic susceptibility records from the YZ01 borehole in the Xinding Basin, Shanxi Graben, North China, revealing that precipitation was dominated by the 100-kyr eccentricity ~~cycles~~ before 3.3 Ma, and constituted a nonlinear response to insolation (Wang et al., 2023). Moreover, a similar nonlinear response to insolation was found in the Rb/Sr records from the Huangdigou section in the Sanmenxia Basin, North China (Zhang et al., 2022). In the western Qaidam Basin, the pollen records from the SG-1b core reveal that there was a strong link between the 100-kyr ~~eccentricity~~cycles in the precipitation records and the eccentricity forcing during 3.5-3.0 Ma (Schwarz et al., 2022).

Interestingly, precipitation was stronger during intervals dominated by a 20-kyr precessional signal, whereas it was relatively weaker during intervals characterized mainly by ~~a 100-kyr eccentricity signal~~scale variability. However, this contrast does not simply reflect differences in precipitation intensity itself. Rather, it suggests that the dominant processes controlling the westward penetration of summer moisture into the western Qaidam Basin varied under different climatic background states. During

490 the wetter interval of 3.25-2.95 Ma, ~~EASM-related~~summer moisture associated with
 491 the EASM could penetrate farther inland, allowing precipitation to respond more
 492 directly to insolation forcing. By contrast, during $\sim$ 3.6-3.3 Ma, when precipitation was
 493 weaker, the ~~dominant 100-kyr cyclicity~~eccentricity-scale variability suggests ~~a more~~
 494 ~~indirect control on that~~ summer moisture delivery, ~~likely involving was influenced~~
 495 more indirectly by changes in large-scale boundary conditions. This indirect control
 496 may reflect several plausible mechanisms through which eccentricity-scale forcing
 497 influenced precipitation in the GS section during $\sim$ 3.6-3.3 Ma. ~~First, the dominant~~
 498 ~~100-kyr cyclicity can be interpreted as linked to insolation driven Antarctic ice sheet~~
 499 ~~forcing of the monsoonal intensity. That~~One possible explanation is, ~~that~~ eccentricity
 500 ~~controls~~ scale forcing may have modulated high-latitude ice-volume variability,
 501 including Antarctic ice ~~sheets~~ sheet variability, in a nonlinear way, ~~thereby affecting~~
 502 ~~the expansion and contraction of Antarctic ice sheets and hence controlling the~~that
 503 these changes may in turn have influenced EASM intensity through ~~the modulation~~
 504 ~~of changes in~~ the thermal contrast between Eurasia and the surrounding oceans and/or
 505 ~~the increase~~through changes in Pacific and Atlantic meridional overturning circulation
 506 (Zhang et al., 2022; Wang et al., 2023). Because the Arctic ice sheets did not develop
 507 on a large scale before $\sim$ 2.7 Ma (Brigham-Grette et al., 2013; Rahaman et al., 2020;
 508 Bridges et al., 2023), ~~the benthic oxygen isotope stack mainly represented~~ Antarctic
 509 ice ~~sheet variations~~ variability was likely an important component of high-latitude
 510 ice-volume changes during the Late Pliocene. Thus, ~~at~~in the ~~100-kyr~~eccentricity band,
 511 the ~~comparison of these~~ broadly similar variations between the two GS precipitation

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

records, eccentricity, and the benthic $\delta^{18}\text{O}$ stack are consistent with the hypothesis that eccentricity-paced high-latitude ice-volume variability, potentially involving Antarctic ice-sheet changes, contributed to the observed eccentricity-scale signal in the GS section with the eccentricity and benthic oxygen isotope stack seems to support this hypothesis (Fig. 8). Second, the dominant 100-kyr cyclicity Another possibility is that the eccentricity-scale variability in the precipitation records could have been controlled by linked to annual insolation instead of summer insolation. Because the annual insolation integrates seasonable seasonal insolation variations caused by precession, it has more stronger power in the eccentricity bands band and weak weaker power in the precession bands (Laskar band (Laskar et al., 2004). However, more investigations are needed to validate this hypothesis.




Figure 8. Comparison of the χ_{fd} /HIRM and median grain-size records with the eccentricity and benthic oxygen isotope $\delta^{18}\text{O}$ stack data. The χ_{fd} /HIRM, median grain-size and benthic oxygen isotope $\delta^{18}\text{O}$ stack data were filtered at the 100-kyr band.

Additionally, at the 40-kyr band, our precipitation records show a similar phase

设置了格式

带格式的：缩进：首行缩进： 6.07 字符

设置了格式

设置了格式

设置了格式

设置了格式

relationship with filtered variations to the benthic ~~oxygen isotope~~ $\delta^{18}\text{O}$ stack, suggesting that high-latitude ice-volume variability, potentially including Antarctic ice-sheet variability, may have ~~directly influenced~~ contributed to precipitation changes in the Qaidam Basin during $\sim 3.6\text{--}3.3$ Ma, rather than acting only indirectly through ~~insolation-related forcing~~. This inference is consistent with the published GS precipitation records for $3.25\text{--}2.95$ Ma (Luo et al., 2024), ~~and together~~. Together, these results suggest that high-latitude ice-volume variability, potentially including Antarctic ice-sheet variability ~~directly modulated~~, may have acted as an important background modulator of precipitation in the Qaidam Basin during the Late Pliocene.

6 Conclusions

Integrated χ_{fd} /HIRM and median grain-size records from the GS section provide ~~a continuous~~ an extended reconstruction of precipitation variability in the western Qaidam Basin during $\sim 3.6\text{--}1.8$ Ma. The combined record shows that precipitation during $3.25\text{--}2.95$ Ma was higher than during $\sim 3.6\text{--}3.25$ Ma and $2.95\text{--}1.8$ Ma, supporting the interpretation of relatively wetter conditions in the western Qaidam Basin during the mPWP. Comparison with previously published regional precipitation records further suggests that EASM-related summer moisture continued to reach the western Qaidam Basin during $\sim 3.6\text{--}3.25$ Ma, although the associated precipitation was weaker than that during $3.25\text{--}2.95$ Ma.

At the orbital timescale, precipitation variations during $\sim 3.6\text{--}3.3$ Ma ~~was~~ were characterized mainly by eccentricity-scale ($\sim 100\text{--}125\text{-kyr}$ -~~eyelicity~~) variability, in contrast to the previously reported precession-dominated pattern during $3.25\text{--}2.95$ Ma.

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

This shift indicates that the precession-dominated pattern was not persistent throughout the Late Pliocene in the western Qaidam Basin. Instead, the dominant periodicity of precipitation changed through time, implying that the processes controlling the westward penetration of summer moisture into the basin varied under different climatic background states. In particular, the stronger ~20-kyr signal during 3.25-2.95 Ma likely reflects a more direct response of precipitation to precessional insolation forcing under wetter conditions, whereas the ~~dominant 100-kyr~~ ~~eccentricity~~ eccentricity-scale (~100-125-kyr) variability during ~3.6-3.3 Ma suggests a more indirect forcing response under relatively drier conditions. In addition, the ~~phase relationship~~ broad correspondence between the precipitation proxies and the ~~benthic oxygen isotope~~ $\delta^{18}\text{O}$ stack at the 40-kyr band suggests that high-latitude ice-volume variability, potentially including Antarctic ice-sheet variability, may also have modulated ~~hydroclimatic variability~~ precipitation variations in the basin. Overall, these findings highlight that Late Pliocene precipitation ~~variability~~ variations in the western Qaidam Basin ~~was~~ were shaped not only by changes in moisture availability, but also by shifts in the dominant climatic processes governing inland penetration of EASM-related summer moisture.

Data availability

~~All data for this paper are available upon request.~~

The data underlying this study are provided in the Supplement. These include the GS magnetic-parameter and median grain-size records, paleomagnetic age-control points and sedimentation-rate calculations.

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

设置了格式

Author contributions

All authors approved the manuscript and agreed on its submission. **ZL:** writing-original draft, methodology, conceptualization, formal analysis, data collection, funding acquisition, investigation, resources. **HW:** Writing-review & editing, investigation, data collection. **XY:** Writing-review & editing, investigation. **ZZ, ZS, JY and WC:** investigation, sample processing. **QS:** investigation, sample processing, funding acquisition. **YL and SH:** investigation.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

This work was financially supported by the National Natural Science Foundation of China (grants 42202215, 42302213 and 42401004), the Key Foundation of the Education Department of Hunan Province (21A0424, [24C0341](#)), the PhD Research Startup Foundation of Hunan University of Arts and Science (grants-~~23BSQD05~~, [24BSQD79](#), [25BSQD58](#)-~~23BSQD05~~), the Fund of Geography Subject ([2022]351) and the Open Foundation of MOE Key Laboratory of Western China's Environmental System, Lanzhou University and the Fundamental Research Funds for the Central Universities (lzujbky-2021-kb01).

References

Ao, H., Roberts, A.P., Dekkers, M.J., Liu, X., Rohling, E.J., Shi, Z., An, Z., and Zhao, X.: Late Miocene-Pliocene Asian monsoon intensification linked to Antarctic ice-sheet growth, Earth Planet. Sci. Lett., 444, 75-87, doi:<https://doi.org/10.1016/j.epsl.2016.03.028>, 2016.

带格式的: 缩进: 首行缩进: 2 字符

带格式的: 缩进: 首行缩进: 2 字符

带格式的: 缩进: 首行缩进: 2 字符

595 Bartoli, G., Honisch, B., and Zeebe, R.E.: Atmospheric CO₂ decline during the
 596 Pliocene intensification of Northern Hemisphere glaciations, *Paleoceanography*,
 597 26, PA4213, doi:<https://doi.org/10.1029/2010pa002055>, 2011.

598 Badger, M. P. S., Schmidt, D. N., Mackensen, A., and Pancost, R. D.: High resolution
 599 alkenone palaeobarometry indicates relatively stable pCO₂ during the Pliocene
 600 (3.3 to 2.8 Ma), *Philos. Trans. A Math. Phys. Eng. Sci.*, 371, 20130094,
 601 doi:<https://doi.org/10.1098/rsta.2013.0094>, 2013.

602 Bridges, J.D., Tarduno, J.A., Cottrell, R.D., and Herbert, T.D.: Rapid strengthening of
 603 westerlies accompanied intensification of Northern Hemisphere glaciation, *Nat.*
 604 *Commun.*, 14, 3905, doi:<https://doi.org/10.1038/s41467-023-39557-4>, 2023.

605 Brigham-Grette, J., Melles, M., Minyuk, P., Andreev, A., Tarasov, P., DeConto, R.,
 606 Koenig, S., Nowaczyk, N., Wennrich, V., and Rosén, P.: Pliocene warmth, polar
 607 amplification, and stepped Pleistocene cooling recorded in NE Arctic Russia,
 608 *Science*, 340(6139), 1421-1427, doi:<https://doi.org/10.1126/science.1233137>,
 609 2013.

610 Bush, M., Horton, B.K., Saylor, J.E., and Nie, J.S.: Growth of the Qaidam Basin
 611 during Cenozoic exhumation in the northern Tibetan Plateau: Inferences from
 612 depositional patterns and multiproxy detrital provenance signatures, *Lithosphere*,
 613 8(1), 58-82, doi:<https://doi.org/10.1130/L449.1>, 2016.

614 Burke, K.D., Williams, J.W., Chandler, M.A., Haywood, A.M., Lunt, D.J., and
 615 Ottobliesner, B.L.: Pliocene and Eocene provide best analogs for near-future
 616 climates, *Proc. Natl. Acad. Sci.*, 115(52), 13288-13293,
 617 doi:<https://doi.org/10.1073/pnas.1809600115>, 2018.

618 Cai, M.T., Fang, X.M., Wu, F.L., Miao, Y.F., and Appel, E.: Pliocene-Pleistocene
 619 stepwise drying of Central Asia: Evidence from paleomagnetism and sporopollen

620 record of the deep borehole SG-3 in the western Qaidam Basin, NE Tibetan
 621 Plateau, *Glob. Planet. Chang.*, 94-95, 72-81,
 622 <https://doi.org/10.1016/j.gloplacha.2012.07.002>, 2012.

623 Dowsett, H.J., Robinson, M.M., Haywood, A.M., Salzmann, U., Hill, D.J., Sohl, L.E.,
 624 Chandler, M., Williams, M., Foley, K., and Stoll, D.K.: The PRISM3D
 625 paleoenvironmental reconstruction, *Stratigraphy*, 7,
 626 <https://doi.org/10.1111/j.1475-4983.2010.00949.x>, 2010.

627 Gu, N., Jiang, W.Y., Wang, L., Zhang, E.L., Yang, S.L., and Xiong, S.F.: Rainfall
 628 thresholds for the precipitation of carbonate and evaporite minerals in modern
 629 lakes in northern China, *Geophys. Res. Lett.*, 42 (14), 5895-5901,
 630 <https://doi.org/10.1002/2015GL064340>, 2015.

631 Haywood, A.M., Ridgwell, A., Lunt, D.J., Hill, D.J., Pound, D.J., and Dowsett, H.J.:
 632 Are there pre-Quaternary geological analogues for a future greenhouse warming?
 633 *Philos. T. R. Soc. A*, 369 (1938), 933-956,
 634 <https://doi.org/10.1098/rsta.2010.0317>, 2011.

635 Haywood, A., Dowsett, H., and Dolan, A.: Integrating geological archives and climate
 636 models for the mid-Pliocene warm period, *Nat. Commun.*, 7, 10646,
 637 <https://doi.org/10.1038/ncomms10646>, 2016.

638 Hearty, P. J., Rovere, A., Sandstrom, M.R., O'Leary, M. J., Roberts, D., and Raymo,
 639 M. E.: Pliocene-Pleistocene stratigraphy and sea-level estimates, Republic of
 640 South Africa with implications for a 400 ppmv CO₂ world, *Paleoceanogr.*
 641 *Paleoclimatol.*, 35, e2019PA003835, <https://doi.org/10.1029/2019PA003835>,
 642 2020.

643 Herbert, T.D., Lawrence, K.T., Tzanova, A., Peterson, L.C., Caballero-Gill, R., and
 644 Kelly, C.S.: Late Miocene global cooling and the rise of modern ecosystems, *Nat.*
 645 *Geosci.*, 9(11), 843-847, <https://doi.org/10.1038/ngeo2813>, 2016.
 646 Huang, X.F., Jiang, D.B., Dong, X.X., Yang, S.L., Su, B.H., Li, X.Y., Tang, Z.H., and
 647 Wang, Y.D.: Northwestward Migration of the Northern Edge of the East Asian
 648 Summer Monsoon During the Mid-Pliocene Warm Period: Simulations and
 649 Reconstructions, *J. Geophys. Res.*, 124, 1392-1404,
 650 <https://doi.org/10.1029/2018jd028995>, 2019.
 651 Huang, X.F., Yang, S.L., Haywood, A., Jiang, D.B., Wang, Y.D., Sun, M.M., Tang,
 652 Z.H., and Ding, Z.L.: Warming-induced northwestward migration of the Asian
 653 summer monsoon in the geological past: Evidence from climate simulations and
 654 geological reconstructions, *J. Geophys. Res-Atmos.*, 126, e2021JD035190,
 655 <https://doi.org/10.1029/2021JD035190>, 2021.
 656 Koutsodendris, A., Allstadt, F.J., Kern, O.A., Kousis, I., Schwarz, F., Vannacci, M.,
 657 Woutersen, A., Appel, E., Berke, M.A., and Fang, X. M.: Late Pliocene
 658 vegetation turnover on the NE Tibetan Plateau (Central Asia) triggered by early
 659 Northern Hemisphere glaciation, *Glob. Planet. Chang.*, 180, 117-125,
 660 <https://doi.org/10.1016/j.gloplacha.2019.06.001>, 2019.
 661 Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A.C.M., and Levrard, B.: A
 662 long-term numerical solution for the insolation quantities of the Earth, *Astron.*
 663 *Astrophys.*, 428, 261-285, <https://doi.org/10.1051/0004-6361:20041335>, 2004.
 664 Li, M.H., Fang, X.M., Yi, C.L., Gao, S.P., Zhang, W.L., and Galy, A.: Evaporite
 665 minerals and geochemistry of the upper 400 m sediments in a core from the
 666 Western Qaidam Basin, Tibet, *Quatern. Int.*, 218(1-2), 176-189,
 667 <https://doi.org/10.1016/j.quaint.2009.12.013>, 2010.

668 Lisiecki, L.E., and Raymo, M.E.: A Plio-Pleistocene stack of 57 globally distributed
 669 benthic $\delta^{18}\text{O}$ records, *Paleoceanography*, 20, PA1003,
 670 <https://doi.org/10.1029/2004pa001071>, 2005.

671 Lunt, D.J., Haywood, A.M., Schmidt, G.A., Salzmann, U., Dowsett, H.J., and Loptson,
 672 C.A.: On the causes of mid-Pliocene warmth and polar amplification, *Earth*
 673 *Planet. Sci. Lett.*, 321-322, 128-138, <https://doi.org/10.1016/j.epsl.2011.12.042>,
 674 2012.

675 Luo, Z., Su, Q.D., Wang, Z., Heermance, R.V., Garzione, C.N., Li, M., Ren, X.P.,
 676 Song, Y.G., and Nie, J.S.: Orbital forcing of Plio-Pleistocene climate variation in
 677 a Qaidam Basin lake based on paleomagnetic and evaporite mineralogic analysis,
 678 *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 510, 31-39,
 679 <https://doi.org/10.1016/j.palaeo.2017.09.022>, 2018.

680 Luo, Z., Nie, J.S., Moe, A.E., Heermance, R.V., Garzione, C; Herbert, T.D., Wang, Z.,
 681 Li, H., Zhang, R., Zhao, X.M., and Salzmann, U.: Joint insolation and ice
 682 sheet/CO₂ forcing on northern China precipitation during Pliocene warmth, *Sci.*
 683 *Bull.*, 66(4), 319-322, <https://doi.org/10.1016/j.scib.2020.10.025>, 2021.

684 Luo, Z., Wang, H.S., Heermance, R.V., Nie, J.S., Yang, J., Yu, X.L., Wang, Z., Zhang,
 685 Z., and Su, Q.D.: June insolation gradient and ice sheet forcing on Qaidam
 686 precipitation during the middle Piacenzian warm period, *Palaeogeogr.*
 687 *Palaeoclimatol. Palaeoecol.*, 648, 112277,
 688 <https://doi.org/10.1016/j.palaeo.2024.112277>, 2024.

689 Miao, Y.F., Fang, X.M., Wu, F.L., Cai, M.T., Song, C.H., Meng, Q.Q., and Xu, L.:
 690 Late Cenozoic continuous aridification in the western Qaidam Basin: evidence
 691 from sporopollen records, *Clim. Past*, 9(4), 1863-1877,
 692 <https://doi.org/10.5194/cp-9-1863-2013>, 2013.

693 Nie, J.S., Garziona, C.N., Su, Q.D., Liu, Q.S., Zhang, R., Heslop, D., Necula, C.,
 694 Zhang, S.H., Song, Y.G., and Luo, Z.: Dominant 100,000-year precipitation
 695 cyclicity in a late Miocene lake from Northeast Tibet, *Sci. Adv.*, 3(3), e1600762,
 696 <https://doi.org/10.1126/sciadv.1600762>, 2017.
 697 Raymo, M.E., Lisiecki, L.E., and Nisancioglu, K.H.: Plio-Pleistocene ice volume,
 698 Antarctic climate, and the global $\delta^{18}\text{O}$ record, *Science*, 313, 492-495,
 699 <https://doi.org/10.1126/SCIENCE.1123296>, 2006.
 700 Ravelo, A.C., Andreasen, D.H., Lyle, M., Olivarez, L.A., and Wara, M.W.: Regional
 701 climate shifts caused by gradual global cooling in the Pliocene epoch, *Nature*,
 702 429, 263-267, <https://doi.org/10.1038/nature02567>, 2004.
 703 Rahaman, W., Smik, L., Köseoglu, D., N, L., Tarique, M., Thamban, M., Haywood,
 704 A., Belt, S.T., and Knies, J.: Reduced Arctic sea ice extent during the
 705 mid-Pliocene Warm Period concurrent with increased Atlantic-climate regime,
 706 *Earth Planet. Sci. Lett.*, 550, 116535, <https://doi.org/10.1016/j.epsl.2020.116535>,
 707 2020.
 708 Ren, X.P., Nie, J.S., Saylor, J.E., Wang, X.X., Liu, F.B., and Horton, B.K.:
 709 Temperature control on silicate weathering intensity and evolution of the
 710 Neogene East Asian summer monsoon, *Geophys. Res. Lett.*, 47(15),
 711 e2020GL088808, <https://doi.org/10.1029/2020gl088808>, 2020.
 712 Reguero, M.A., Candela, A.M., and Alonso, R.N.: Biochronology and biostratigraphy
 713 of the Uquía Formation (Pliocene-early Pleistocene, NW Argentina) and its
 714 significance in the Great American Biotic Interchange, *J. S. Am. Earth Sci.*,
 715 23(1), 1-16, <https://doi.org/10.1016/j.jsames.2006.09.005>, 2007.
 716 Salzmann, U., Haywood, A. M., Lunt, D. J., Valdes, P. J., and Hill, D. J.: A new
 717 global biome reconstruction and data-model comparison for the Middle Pliocene,

Global Ecol. Biogeogr., 17, 432-447,
<https://doi.org/10.1111/j.1466-8238.2008.00381.x>, 2008.

Salzmann, U., Williams, M., Haywood, A.M., Johnson, A.L.A., Kender, S., and
 Zalasiewicz, J.: Climate and environment of a Pliocene warm world, *Palaeogeogr.*
Palaeoclimatol. Palaeoecol., 309(1-2), 1-8,
<https://doi.org/10.1016/j.palaeo.2011.05.044>, 2011.

Schwarz, F., Salzmann, U., Koutsodendris, A., Nie, J.S., Friedrich, O., Ni, J.,
 Garziane, C., Fang, X.M., Wu, F.L., Woodward, J., Appel, E., and Prosser, J.:
 Controls of precipitation and vegetation variability on the NE Tibetan Plateau
 during the Late Pliocene warmth (~3.5-3.0 Ma), *Glob. Planet. Chang.*, 208,
 103707, <https://doi.org/10.1016/j.gloplacha.2021.103707>, 2022.

Song, C.H., Hu, S.H., Han, W.X., Zhang, T., Fang, X.M., Gao, J.P., and Wu, F.L.:
 Middle Miocene to earliest Pliocene sedimentological and geochemical records
 of climate change in the western Qaidam Basin on the NE Tibetan Plateau,
Palaeogeogr. Palaeoclimatol. Palaeoecol., 395(2), 67-76,
<https://doi.org/10.1016/j.palaeo.2013.12.022>, 2013.

Song, Y.G., Fang, X.M., Li, J.J., An, Z.S., and Miao, X.D.: The Late Cenozoic uplift
 of the Liupan Shan, China, *Sci. China Ser. D-Earth Sci.*, 44, 176-184,
<https://doi.org/10.1007/bf02911985>, 2001.

Su, Q.D., Nie, J.S., Meng, Q.Q., Heermance, R., Gong, L.S., Luo, Z., Wang, Z.,
 Zhang, R., and Garziane, C.: Central Asian Drying at 3.3 Ma Linked to Tropical
 Forcing? *Geophys. Res. Lett.*, 46(17-18), 10561-10567,
<https://doi.org/10.1029/2019GL084648>, 2019.

Su, Q.D., Nie, J.S., Han, W.X., Garziane, C.N., and Gao, P.: Obliquity forcing of Late
 Pliocene–Early Pleistocene aridification in the Qaidam Basin, NE Tibetan

- Plateau, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 635, 111951, <https://doi.org/10.1016/j.palaeo.2023.111951>, 2024.
- Vieira, M., Pound, M.J., and Pereira, D.I.: The Late Pliocene palaeoenvironments and palaeoclimates of the western Iberian Atlantic margin from the Rio Maior flora, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 495, 245-258, <https://doi.org/10.1016/j.palaeo.2018.01.018>, 2018.
- Wan, S.M., Tian, J., Steinke, S., Li, A.C., and Li, T.G.: Evolution and variability of the East Asian summer monsoon during the Pliocene: Evidence from clay mineral records of the South China Sea, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 293 (1-2), 237-247, <https://doi.org/10.1016/j.palaeo.2010.05.025>, 2010.
- Wang, J.Y., Fang, X.M., Appel, E., and Zhang, W.L.: Magnetostratigraphic and radiometric constraints on salt formation in the Qaidam Basin, NE Tibetan Plateau, *Quat. Sci. Rev.*, 78(11), 53-64, <https://doi.org/10.1016/j.quascirev.2013.07.017>, 2013.
- Wang, Z.X., Mao, Y.D., Zhang, R., Kemp, D. B., and Huang, C.J.: Links between orbital forcing, Antarctic ice volume, and the East Asian hydrological cycle over the Pliocene, *J. Geophys. Res. Atmos.*, 128, e2023JD039519, <https://doi.org/10.1029/2023jd039519>, 2023.
- Wehausen, R., and Brumsack, H.J.: Astronomical forcing of the East Asian monsoon mirrored by the composition of Pliocene South China Sea sediments, *Earth Planet. Sci. Lett.*, 201, 621-636, [https://doi.org/10.1016/S0012-821X\(02\)00746-X](https://doi.org/10.1016/S0012-821X(02)00746-X), 2002.
- Yang, S.L., Ding, Z.L., Feng, S.H., Jiang, W.Y., Huang, X.F., and Guo, L.C.: A strengthened East Asian Summer Monsoon during Pliocene warmth: Evidence

from 'red clay' sediments at Pianguan, northern China, *J. Asian Earth Sci.*, 155, 124-133, <https://doi.org/10.1016/j.jseaes.2017.10.020>, 2018.

Yang, L.Y., Zhang, W.L., Fang, X.M., Cai, M.T., and Lu, Y.: Aridification recorded by lithofacies and grain size in a continuous Pliocene-Quaternary lacustrine sediment record in the western Qaidam Basin, NE Tibetan Plateau, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 556, 109903, <https://doi.org/10.1016/j.palaeo.2020.109903>, 2020.

Zhang, Z.G., Han, W.X., Fang, X.M., Song, C.H., and Li, X.Y.: Late Miocene–Pleistocene aridification of Asian inland revealed by geochemical records of lacustrine-fan delta sediments from the western Tarim Basin, NW China, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 377(2), 52-61, <https://doi.org/10.1016/j.palaeo.2013.03.008>, 2013.

Zhang, Z., Licht, A., De Vleeschouwer, D.D., Wang, Z.X., Li, Y.Z., Kemp, D. B., Tan, L.C., Zhang, R., Qiang, X.K., and Huang, C.J.: East Asian monsoonal climate sensitivity changed in the Late Pliocene in response to Northern Hemisphere glaciations, *Geophys. Res. Lett.*, 49, e2022GL101280, <https://doi.org/10.1029/2022gl101280>, 2022.

Zhou, X.Q., and Liu, C.L.: Mid-Piacenzian and future changes in South Asian precipitation under global warming, *Glob. Planet. Chang.*, 248, 104760, <https://doi.org/10.1016/j.gloplacha.2025.104760>, 2025.