

Major comments

In this study Laemmel et al. presented ~5 years' worth of very high quality $\Delta^{14}\text{CH}_4$ and $\Delta^{14}\text{CO}_2$ measurements from Jungfraujoch, Switzerland. Excluding reconstruction from ice cores/firn air (e.g., Hmiel et al., 2020) and sporadic atmospheric $^{14}\text{CH}_4$ measurements conducted during limited duration measurement campaigns (e.g., Sparrow et al., 2018) this study represent the first $\Delta^{14}\text{CH}_4$ measurements from clean background sites for the purpose of long-term global atmospheric monitoring probably since the Barring Head measurements from New Zealand presented by Lassey et al. (2007) almost 20 years ago. Due to the lack of measurements, for years we've been using +350‰ as “background” $\Delta^{14}\text{CH}_4$. This study presents a very welcome update to that value, and very interestingly the authors found that atmospheric $\Delta^{14}\text{CH}_4$ have been increasing since 2019 probably due to the proliferation of PWR (pressurized water reactor) nuclear power plants.

Beyond the novelty of the $\Delta^{14}\text{CH}_4$ measurements, this study also shows that the Bern Radiocarbon laboratory (LARA) can produce state-of-the-art high quality atmospheric ^{14}C data that are comparable to the background measurements made within the ICOS network and other international laboratories that handle background air samples. The passive sampling system using a large volume container (JASS) is very innovative and provides a blueprint for large volume sampling that I hope can be adapted by other global monitoring observatories. Given the study site (in Central Europe), interference from nuclear power plants is unavoidable, and I think the authors did their best to correct for the influences of local nuclear power plants. The paper is very well written; I highly recommend the paper for publication and congratulate the authors on a great manuscript/study.

I just have several general comments out of genuine curiosity for additional details that I think will help other readers, as well as some slightly nitpicky comments regarding choice of references, mostly in the introduction section (see minor comments).

1. In line 346-348 the authors mentioned: “*The standard deviation of the $\Delta^{14}\text{CH}_4$ value for all 40 CH_4 measurements over 15 months for each PAB bottle (so about 80 CH_4 measurements in total) is 8 ‰ (Fig. S4a,b), which is lower than the instrumental uncertainty of a single $\Delta^{14}\text{CH}_4$ measurement (12 ‰) indicating the satisfactory temporal reliability of our $\Delta^{14}\text{CH}_4$ measurements.*” Similar mention is repeated for $\Delta^{14}\text{CO}_2$ in line 438-441. First, I would replace “temporal reliability” with “long-term reproducibility.” I’m also interested in how the instrumental uncertainty/single sample precision is determined. For this, hopefully the authors are willing to share the composition of samples/standards for the 39 position MICADAS “wheel” (I guess you don’t use wheel in MICADAS, so batch/magazine/cartridges). How many OX-2s are used for normalization and how many replicates of the surveillance gas (the PABs

cylinders) are put in each sample batch. What was the average stdev of however many (4,6,8,10) primary OX-2 standards used for normalization within a single sample batch, and whether the average stdev of primary standards was bigger/smaller than the single sample precision derived from counting statistics. Also, were the samples measured together in small amounts of 39 position sample batches/magazines (like within 6-7 batches at the minimum to account for the 130-something samples and 80 surveillance samples, or are they more spread out over larger amounts of sample batches. Knowing the composition of the 39 position sample within a single MICADAS batch will give us a good sense of the inter-batch/magazine reproducibility.

2. Moving to integrated and automated sampling with the JASS system is clearly very beneficial in terms of sampling consistency and reducing labor needed to do the grab samples. But I have a couple of follow-up questions regarding the JASS system. The grab samples clearly show some highly elevated $\Delta^{14}\text{CH}_4$ (>420‰ cutoff) – likely given the sporadic nature of $^{14}\text{CH}_4$ emissions from PWR venting. Would it be fair to say/acknowledge that with integrated sampling it is harder to correct for potential elevation from sporadic PWR venting? The footprint-based correction relies on annual, quarterly, and monthly NPP emissions – this might not capture the sporadic emissions from venting so there's still some upside to grab samples. Second, in line 132-134 the authors mentioned that *“air sampling should occur passively, i.e., without any mechanical pumps in contact with the sampled air, thereby minimizing the risk of contamination from membrane outgassing.”* This is a noble principle to strive for, but is it fair to also acknowledge that one still needs to transfer the sample from the large volume container into the transport container? Reading the manuscript, in line 169-171 the authors wrote: *“At this stage, the transfer pump was evacuating the tank and after ~30 s of line flushing, a bag was connected to the line for the transfer of the sampled air. After about one hour, enough air was transferred to the bag for further ^{14}C analyses.”* So, the sample air is still in contact with mechanical pump, arguably even longer (1hr) compared to the 22-55 min needed to compress enough air during the manual grab samples (line 124). As such, the benefit of JASS sampling scheme in terms of sample cleanliness is only valid if the KNFN022 pump used for grab samples is dirtier than the KNF N922 pump used to transfer the sample from the large volume reservoir to the transfer bag.
3. Between the $\Delta^{14}\text{CH}_4$ with >420‰ cutoff, Radon filter, and Radon filter + footprint based nuclear correction (Table 1) – I personally think that the Radon filter is the most straightforward/foolproof representative of a well-mixed background mid-latitude $\Delta^{14}\text{CH}_4$, given the sporadic nature of $^{14}\text{CH}_4$ emissions from PWR that is not captured by emission inventories. The authors take a more agnostic approach to this, which is

fine, but I think at least the Radon filtered data should also be on Figure 4. I will also welcome an additional discussion if the authors have any inclination one way or another regarding which of the 3 annual averages they think is most representative of a true mid-latitude background value for $^{14}\text{CH}_4$.

4. Since the footprints are already made, it would be interesting if the authors could show comparison of footprints/catchment area for the grab samples vs. integrated JASS sampling, maybe as a supplementary figure. Either pick two footprints that are somewhat representative of the average for the two sampling schemes, or sum the footprints and provide a contour of influence for the two sampling schemes.

Minor comments

Line 30: “*biweekly*” is ambiguous as it can literally mean twice a week or once every two weeks. Be more specific and change it to either “*once every two weeks*” or “*fortnightly*”.

Line 34: Add “*continued*” before “*dilution*.” This is just to draw the contrast that the continued decrease in $^{14}\text{CO}_2$ is pretty much expected and has been a trend that’s been observed in all global sites, unlike the increase in $^{14}\text{CH}_4$, which is a novel discovery of this study.

Line 37: “*.. the mean nuclear ^{14}C contribution to our individual measurements was estimated to be $7 \pm 9 \%$...*” This is slightly misleading, as from my understanding not all measurements are corrected for local nuclear contribution, only measurements with high Radon. I would instead say something along the line of “*To assess the nuclear ^{14}C contribution to our individual measurements, we use a combination of in situ Radon-222 measurements and Lagrangian particle dispersion model convolved with bottom-up inventory of ^{14}C emissions from nuclear power plants.*” Probably also put this sentence before line 31 “*Over this period, $\Delta^{14}\text{CH}_4$ values showed an increase...*”

Line 47: The Hmiel et al. (2025) citation here is not appropriate. Please cite the original Law Dome ice core CH_4 mole fraction paper (MacFarling Meure et al., 2006).

Line 52: A more appropriate citation for the ^{14}C half-life of 5700 years is probably a combination of (Kutschera, 2013) – where the compilation of previous half-life measurements were made, (National Nuclear Data Center, Brookhaven National Laboratory, www.nndc.bnl.gov), and the most “recent” effort by Roberts and Southon (2007).

Line 61-62: “*Atmospheric $^{14}\text{CO}_2$ have a long history. Since the first measurements focused on the atmospheric $^{14}\text{CO}_2$ bomb peak in the 1950-60s...*” The most appropriate citation for this statement, even before Levin et al. (1985) is obviously the original Rafter and Fergusson (1957) paper.

Line 69-72: “*Since the bomb peak in the middle of the last century, $\Delta^{14}\text{CO}_2$ values have been declining mostly due to the emissions of ^{14}C -free fossil fuel CO_2 , which depletes the atmospheric $\Delta^{14}\text{CO}_2$ signal (Levin et al., 2010).*” I understand what the authors meant to say, but this is not 100% correct, atmospheric $^{14}\text{CO}_2$ is coming down from the bomb peak also (or even mostly) because of uptake by plants and the surface ocean. Maybe cite something like the Oeschger et al. (1975) paper or something more recent regarding the partitioning of land/ocean sink and revise the sentence accordingly.

Line 85: Consider giving acknowledgements to the very first pioneering atmospheric $^{14}\text{CH}_4$ measurements by Lowe et al. (1988) and Wahlen et al. (1989).

Line 94-95: “*Fujita et al. (2025) estimated 30% lower global CH_4 emissions from the fossil-fuel industry compared to previous isotope-based studies.*” Clarify that the “previous isotope-based studies” specifically referred in this sentence are $\delta^{13}\text{C}$ - CH_4 stable isotope only inversion like Basu et al. (2022) which does not include ^{14}C consideration.

Line 128: I don’t quite understand the use of UTC throughout the manuscript - I think this can be a bit confusing because JFJ local time is not UTC, and when trying to sample the free troposphere local time matters more and absolute time. I would recommend just using local time and mention (UTC+1/UTC+2).

Line 213: “... *and in equation 29 in Stenstrom et al. (2011).*” I’m a big fan of Stenstrom et al. (2011), but unfortunately it is not appropriate to cite as this was an internal report that was not peer-reviewed.

Line 227-229: “*In this study, R_n was used as a proxy to distinguish atmospheric conditions mostly influenced by free tropospheric conditions from conditions mostly influenced by the planetary boundary layer (i.e., with recent land contact).*” This sentence is a bit awkward, I would just say something along the lines of “*In this study, R_n was used as a proxy to distinguish free tropospheric samples from those originating from the planetary boundary layer (i.e., with recent land contact).*”

Line 288: I think “convolve” might be the more correct mathematical term instead of “convolute”. Footprints are convolved with flux to produce mole fraction enhancement at receptor site.

Line 452-453: “*The offset in 2019 was probably due to a small fossil contamination in our early $\Delta^{14}\text{CO}_2$ measurements that was remedied during the year.*” Not that I don’t believe this explanation, but as it stands by reading this manuscript alone it is not very clear that the “true” European background $\Delta^{14}\text{CO}_2$ in 2019 is closer to the ICOS/Heidelberg values than the LARA values. Maybe show/at least mention Mace Head (MHD) measurements as

independent dataset to support the idea that LARA sampling stream was slightly contaminated in 2019?

Line 526: Very minor thing, I might have missed it but I think “ $\Delta\Delta^{14}\text{CO}_2$ ” is not defined. Just say “*difference between ICOS sodium hydroxide-based integrated sampling vs. our JASS system*” to reduce jargon.

References

Basu, S., Lan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P., Apadula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini, J., and Manca, G.: Estimating emissions of methane consistent with atmospheric measurements of methane and $\delta^{13}\text{C}$ of methane, *Atmospheric Chemistry and Physics*, 22, 15351–15377, <https://doi.org/10.5194/acp-22-15351-2022>, 2022.

Hmiel, B., Petrenko, V. V., Dyonisius, M., Buizert, C., Smith, A. M., Place, P. F., Harth, C. M., Beaudette, R., Hua, Q., Yang, B., Vimont, I., Michel, S. E., Severinghaus, J. P., Etheridge, D. M., Bromley, T. M., Schmitt, J., Fain, X., Weiss, R. F., and Dlugokencky, E. J.: Preindustrial $^{14}\text{CH}_4$ indicates that anthropogenic fossil CH_4 emissions are underestimated, *Nature*, 2020.

Kutschera, W.: Applications of accelerator mass spectrometry, *International Journal of Mass Spectrometry*, 349–350, 203–218, <https://doi.org/10.1016/j.ijms.2013.05.023>, 2013.

Lassey, K. R., Lowe, D. C., and Smith, A. M.: The atmospheric cycling of radiomethane and the “fossil fraction” of the methane source, *Atmospheric Chemistry and Physics*, 7, 2141–2149, <https://doi.org/10.5194/acp-7-2141-2007>, 2007.

Lowe, D. C., Brenninkmeijer, C. A. M., Manning, M. R., Sparks, R., and Wallace, G.: Radiocarbon determination of atmospheric methane at Baring Head, New Zealand, *Nature*, 332, 522–525, <https://doi.org/10.1038/332522a0>, 1988.

MacFarling Meure, C., Etheridge, D., Trudinger, C., Steele, P., Langenfelds, R., van Ommen, T., Smith, A., and Elkins, J.: Law Dome CO_2 , CH_4 and N_2O ice core records extended to 2000 years BP, *Geophys. Res. Lett.*, 33, L14810, <https://doi.org/10.1029/2006GL026152>, 2006.

Oeschger, H., Siegenthaler, U., Schotterer, U., and Gugelmann, A.: A box diffusion model to study the carbon dioxide exchange in nature, *Tellus*, 27, 168–192, <https://doi.org/10.1111/j.2153-3490.1975.tb01671.x>, 1975.

Rafter, T. A. and Fergusson, G. J.: “Atom Bomb Effect”—Recent Increase of Carbon-14 Content of the Atmosphere and Biosphere, *Science*, 126, 557–558, <https://doi.org/10.1126/science.126.3273.557>, 1957.

Roberts, M. L. and Southon, J. R.: A Preliminary Determination of the Absolute $^{14}\text{C}/^{12}\text{C}$ Ratio of OX-I, *Radiocarbon*, 49, 441–445, <https://doi.org/10.1017/S0033822200042363>, 2007.

Sparrow, K. J., Kessler, J. D., Southon, J. R., Garcia-Tigreros, F., Schreiner, K. M., Ruppel, C. D., Miller, J. B., Lehman, S. J., and Xu, X.: Limited contribution of ancient methane to surface waters of the U.S. Beaufort Sea shelf, *Science Advances*, 4, eaao4842, <https://doi.org/10.1126/sciadv.aao4842>, 2018.

Stuiver, M. and Polach, H. A.: Discussion Reporting of ^{14}C Data, *Radiocarbon*, 19, 355–363, <https://doi.org/10.1017/S0033822200003672>, 1977.

Wahlen, M., Tanaka, N., Henry, R., Deck, B., Zeglen, J., Vogel, J. S., Southon, J., Shemesh, A., Fairbanks, R., and Broecker, W.: Carbon-14 in Methane Sources and in Atmospheric Methane: The Contribution from Fossil Carbon, *Science*, 245, 286–290, <https://doi.org/10.1126/science.245.4915.286>, 1989.