

**Macro constraints and local eutrophication shape sediment nitrogen
removal in the Eastern China Plain lakes**

Jiahao Wei^{1, 2}, Ke Yang^{1 3}, Yaru Liu², Yi Yin⁴, Lachun Wang², Enlou Zhang^{1, 3}, Zhijun
Gong^{1, 3}, Yongjiu Cai^{1, 3*}

1 State Key Laboratory of Lake and Watershed Science for Water Security, Nanjing Institute of Geography and
Limnology, Chinese Academy of Sciences, Nanjing 211135, China

2 School of Geography and Ocean Science, Nanjing University, Nanjing 210023, China

3 University of Chinese Academy of Sciences, Beijing 100049, China

4 School of Life Sciences, Fudan University, Shanghai 200433, China

Correspondence to: Yongjiu Cai (caiyj@niglas.ac.cn)


Abstract:

Eutrophication poses a profound threat to the provisioning of ecosystem services in lakes worldwide. Understanding how eutrophication, alongside other biotic and abiotic factors, drives sediment nitrogen removal is essential for maintaining the intrinsic purification capacity of ecosystems and safeguarding water security. To address this, we conducted a systematic investigation of 17 representative lakes across the Eastern China Plain using ¹⁵N stable isotope tracing, 16S rRNA amplicon sequencing, and metagenomics profiling. This study aims to quantify the contribution of the anammox to nitrogen removal, elucidate the impacts of eutrophication on overall nitrogen removal, and identify the critical biotic and abiotic factors driving these processes. Results revealed that while denitrification was the dominant process across all lakes, anammox made substantial contributions of up to 34.3%, exhibiting a distinct environmental dependency. Eutrophication significantly amplified both the overall nitrogen removal capacity and the relative contribution of anammox, with local physicochemical properties and microbial communities demonstrating a pronounced joint driving effect on these processes. Notably, the observed decoupling among microbial community structure, functional gene abundance, and nitrogen removal rates highlights a significant discrepancy between genetic potential and active expression. This discrepancy is further manifested by the superior environmental adaptability of functionally redundant denitrifying bacteria compared to narrow ecological niche anammox bacteria. Furthermore, we demonstrate that macro scale factors, including spatial, climatic, and anthropogenic, indirectly drive nitrogen removal by reshaping local water-sediment physicochemical properties, rather than acting as direct determinants of process rates. Collectively, this study enriches our fundamental understanding of nitrogen removal, particularly the anammox process, and its multidimensional drivers in the Eastern China Plain lakes. We advocate that future lake management strategies must rigorously account for the overarching constraints jointly imposed by trophic status, alongside spatial and climatic gradients, to ensure the sustainability of nitrogen purification ecosystem services.

Key Words: Denitrification; Anammox; Functional redundancy; Eutrophication; Shallow lakes

设置格式[JIAHAO WEI]: 两端对齐, 行距: 1.5 倍行距, 无孤行控制

Graphical Abstract



设置格式[JIAHAO WEI]: 字体: (中文) 黑体
设置格式[JIAHAO WEI]: 标题 1, 段落间距段前: 7.8 磅
设置格式[JIAHAO WEI]: 居中

1 Introduction

Lakes (including reservoirs) constitute critical components of the Earth's surface system, playing irreplaceable roles in regulating global biogeochemical cycles and energy fluxes, as well as maintaining regional ecological security (Mendonça et al., 2017; Woolway et al., 2025). However, since the Industrial Revolution, the dual stressors of climate change and intense anthropogenic activities have severely disrupted the homeostasis of lake ecosystems, precipitating a series of global lake environmental crises (Woolway et al., 2020; IPCC, 2022). Statistics indicate that algal blooms occur in over 20,000 lakes worldwide, covering approximately 2.57% of the total global lake surface area, with a continuing upward trend in both frequency and intensity (Ho et al., 2019; Hou et al., 2022). The synergistic between climate warming and anthropogenic nitrogen loading is recognized as the core mechanism driving the exacerbation of algal blooms (Huisman et al., 2018; Wang et al., 2025). This not only undermines ecosystem services but also poses severe challenges to global public health security (Dai et al., 2023).

Nitrogen, an essential element for the synthesis of proteins and nucleic acids, undergoes biogeochemical cycling that directly governs the regulation of primary productivity and the maintenance of homeostasis in aquatic ecosystems (Stüeken et al., 2024). When exogenous nitrogen inputs exceed the lake's carrying capacity, disrupted nitrogen cycling can induce ecological disasters such as algal blooms (Ho et al., 2019; Wang et al., 2025). As a primary limiting factor for algal blooms, the spatiotemporal distribution of nitrogen directly determines phytoplankton growth potential (Hutchins and Tagliabue, 2024). The removal of excess reactive nitrogen via microbially mediated pathways is a critical mechanism for maintaining lake health and self-purification capacity (Wang et al., 2024). Sediment nitrogen removal processes—primarily denitrification and anaerobic ammonium oxidation (anammox)—can eliminate 30% to 50% of nitrogen from the water column (Kuypers et al., 2018). Therefore, elucidating the spatiotemporal evolution and driving mechanisms of lake nitrogen removal is scientifically critical for controlling eutrophication and mitigating algal blooms (Helmond et al., 2020; Wu et al., 2025).

In China, driven by rapid socioeconomic development, approximately 69.5% of lakes face eutrophication, a phenomenon particularly pronounced in the densely populated Eastern Plain


Region (Zhou et al., 2022). Ecological degradation resulting from eutrophication causes annual economic losses exceeding \$10 billion, severely threatening the health of China's lake ecosystems and the realization of national water security strategies (Wang et al., 2025). However, previous studies in this region have predominantly focused on denitrification or dissimilatory nitrate reduction to ammonium (DNRA) (Jiang et al., 2023), often overlooking the potential contribution of anammox. Recent evidence suggests that under specific environmental, the contribution of anammox to nitrogen removal is non-negligible and may even become dominant (Deng et al., 2024). Consequently, the lack of systematic assessment of anammox limits our comprehensive understanding of the nitrogen budget and self-purification mechanisms in the Eastern China Plain lakes.


Therefore, we hypothesized that the lakes across the Eastern China Plain could serve as hotspots for the anammox process, and that eutrophication might induce significant shifts in anammox rates and their relative contribution to nitrogen removal. To test this hypothesis, we investigated 17 representative lakes across the Eastern China Plain, covering both mesotrophic and eutrophic lakes. By integrating ^{15}N stable isotope tracing, 16S rRNA amplicon sequencing, and metagenomics, this study aims to address the following questions: (1) What is the relative contribution of the anammox process to overall nitrogen removal in these lakes? (2) Does eutrophication significantly alter the dynamics of denitrification and anammox processes? (3) What are the relative roles of biotic and abiotic factors in driving these lacustrine nitrogen removal processes?

89 **2 Materials and Methods**

90 **2.1 Study area and sample collection**

91 The Eastern China Plain lakes region (ECPLR), one of China's five primary lake regions,
92 encompasses the middle and lower reaches of the Yangtze River, the lower Yellow River, the Huai
93 River, and the Eastern Route of the South-to-North Water Diversion Project. With a total surface
94 area exceeding 20,000 km², the ECPLR accounts for approximately 25.9% of China's total lake
95 area (Ma et al., 2010). Seventeen representative lakes within this region were selected for this
96 study (Fig. 1, Tab. S1). These lakes provide critical ecosystem services and possess significant
97 economic value, including potable water supply, navigation, flood control, fisheries, aquaculture,
98 and biodiversity conservation, among others. Considering the morphological characteristics,
99 hydrological connectivity, and historical monitoring data of each lake, a total of 81 sampling sites
100 were established. Further details regarding the study area are provided in Supplementary Method
101 S1.





设置格式[JIAHAO WEI]: 正文, 居中, 行距: 单倍行距

Figure 1 Overview of the Eastern China Plain lakes region and distribution of sampling sites.

设置格式[JIAHAO WEI]: 两端对齐

Note: SB, Lake Shaobo; GY, Lake Gaoyou; HZ, Lake Hongze; LM, Lake Luoma; NS, Lake Nansi; DP, Lake Dongping; HS, Lake Hengshui; TH, Lake Taihu; FB, Fangbian reservoir; YJ, Yaojia reservoir; ZS, Zhongshan reservoir; NY, Lake Nanyi; GC, Lake Gucheng; CH, Lake Chaohu; PY, Lake Poyang; DH, Lake Donghu; DT, Lake Dongting.

Field surveys were conducted from August to October 2024. At each site, 2 L of composite water samples were collected for physicochemical analysis and slurry incubation experiments. Concurrently, surface sediment (0–5 cm) was collected for physicochemical analysis and slurry incubation (~500 g), and microbial sequencing (~5 g). All samples were immediately stored at 4°C in darkness and transported to the laboratory.

2.2 Physicochemical analysis, climate data acquisition, and human activity intensity quantification

Water temperature (WT), electrical conductivity (EC), dissolved oxygen (DO), and pH were

measured on-site (YSI Professional Plus, USA). Water depth (WD) and secchi depth (SD) were determined using a Speedtech SM-5 depth sounder and Secchi disk, respectively. To minimize random errors, triplicate measurements were performed, and their average was taken to ensure data reliability. Detailed analytical parameters and methods for water and sediment samples are provided in Tab. S2. Trophic status of the lake water was evaluated using the Comprehensive Trophic Level Index (TLI) (Supplementary Method S2). Climate data were obtained from the CRU TS dataset (Harris et al., 2020). Using the "ncdf4" package in R (v4.5.2), the monthly mean precipitation (MP) and mean temperature (MT) were extracted for the sampling months. Additionally, annual-scale climate indices (MAP, MAT, etc., Tab. S3) were calculated. Human Activity Intensity (HAI) was quantified using construction land equivalent conversion coefficients (Tab. S4) (Xu et al., 2015). Detailed procedures regarding land use data sources, preprocessing, and calculations are provided in Supplementary Method S3.

2.4 Nitrogen removal rate determination

Sediment nitrogen removal processes were quantified using ^{15}N isotope tracing slurry incubations (Supplementary Method S4). Dissolved $^{28}\text{N}_2$, $^{29}\text{N}_2$, and $^{30}\text{N}_2$ concentrations were measured using membrane inlet mass spectrometer (MIMS-200, Bay Instruments, USA). Denitrification (V_{den}) and anammox rates (V_{ana}), and the relative contribution of anammox to nitrogen removal (Ra), were calculated following Thamdrup and Dalsgaard (2002).

2.5 DNA extraction, sequencing, and bioinformatic analyses

Total DNA was extracted using the FastDNA® Spin Kit for Soil (MP Biomedicals, USA). The integrity of genomic DNA was assessed by agarose gel electrophoresis, while the concentration and purity were measured using a Nanodrop 2000 and Qubit 3.0 Spectrophotometer. The 16S rRNA gene V3–V4 region was amplified (primers 338F/806R) (Dai et al., 2022), and sequenced on the Illumina MiSeq platform (Guangdong Magigene Biotechnology Co., Ltd., Guangzhou, China). Bioinformatic processing included quality trimming, paired-end read merging, and chimera removal. Amplicon Sequence Variants (ASVs) were resolved based on 100% sequence identity and assigned taxonomy using the SILVA database (v138.2).

Metagenomic libraries were constructed using the ALFA-SEQ DNA Library Prep Kit and sequenced on the Illumina NovaSeq 6000 platform (PE150 mode; Guangdong Magigene

删除[JIAHAO WEI]: Metagenomic libraries were constructed using the ALFA-SEQ DNA Library Prep Kit and sequenced on the Illumina NovaSeq 6000 platform (PE150 mode; Guangdong Magigene Biotechnology Co., Ltd., Guangzhou, China). Raw reads were quality-filtered and adapter-trimmed using fastp and Trimmomatic, yielding approximately 12 Gbp of high-quality clean data per sample. Subsequent bioinformatic analyses encompassed de novo assembly, Open Reading Frame (ORF) prediction, construction of a non-redundant gene catalog (unigenes), and gene abundance quantification (Steinegger and Söding, 2018). Nitrogen-cycling genes and their taxonomic affiliations were annotated using Diamond (v0.9.32) by aligning sequences against the NCycDB (Tu et al., 2019) and the NCBI non-redundant (NR) protein database, respectively. Taxonomic classification was determined using the Lowest Common Ancestor (LCA) algorithm to generate gene abundance profiles. Functional annotation was filtered based on an alignment coverage threshold of > 40%.

Biotechnology Co., Ltd., Guangzhou, China). Raw reads were quality-filtered and adapter-trimmed using fastp and Trimmomatic, yielding approximately 12 Gbp of high-quality clean data per sample. Subsequent bioinformatic analyses encompassed de novo assembly, Open Reading Frame (ORF) prediction, the construction of a non-redundant gene catalog (unigenes), and gene abundance quantification (Steinegger and Söding, 2018). Nitrogen-cycling genes and their taxonomic affiliations were annotated using Diamond (v0.9.32) by aligning sequences against the NCycDB (Tu et al., 2019) and the NCBI non-redundant (NR) protein database, respectively. Taxonomic classification was subsequently determined using the Lowest Common Ancestor (LCA) algorithm to generate comprehensive gene abundance profiles. Finally, functional annotation was filtered based on an alignment coverage threshold of > 40%.

2.6 Statistical analyses

All statistical analyses and data visualization were primarily performed using the R (v4.5.2) and ArcMap 10.4. Microbial diversity was calculated using the 'vegan' package based on a normalized sequencing depth of 34,235 reads per sample. Linear Discriminant Analysis Effect Size (LEfSe) was conducted using the 'microeco' package to identify discriminative taxa between groups (Liu et al., 2022). To quantify spatial structures, Principal Coordinates of Neighbor Matrices (PCNM) analysis was performed using the 'SoDA' package to extract spatial eigenvectors of the sampling sites (Borcard and Legendre, 2002). To disentangle the complex relationships driving nitrogen removal, a multi-tiered statistical approach was employed: (1) Spearman's correlations and Mantel tests were performed using the 'linkET' package to evaluate the interconnections among different factors. (2) Linear Mixed Models (LMMs) were constructed using the 'lme4' package to quantify the effects of geographic location and human activity intensity on nitrogen removal rates. (3) Random Forest (RF) regression models were built using the 'randomForest' package to identify key predictors. (4) Partial Least Squares Path Modeling (PLS-PM) was established using the 'plspm' package to quantitatively disentangle the direct and indirect causal pathways linking environmental factors, microbial communities, and functional genes to nitrogen removal rates (Breiman, 2001; Yuan et al., 2021; Deng et al., 2024).

3 Results

3.1 Spatial heterogeneity of nitrogen removal rates

Measurements based on the ^{15}N isotope pairing technique revealed substantial spatial heterogeneity in sediment nitrogen removal rates across the Eastern China Plain lakes ($p < 0.05$; Fig. 2a, Tab. S5). Mean denitrification rates (V_{den}) ranged from 26.58 to 117.54 $\text{nmol N}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$. Notably, lakes situated along the Eastern Route of the South-to-North Water Diversion Project (ER-SNWD)P—specifically Lakes Hongze, Chaohu, Nansi, and Luoma—exhibited elevated denitrification activities, with mean V_{den} values consistently exceeding 100 nmol N g . In contrast, Lake Dongting ($26.58 \pm 4.84 \text{ nmol N}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$) and certain reservoir-type lakes displayed significantly lower rates. The spatial distribution pattern of anammox rates (V_{ana}) was highly congruent with that of denitrification, with mean values ranging from 2.50 to 34.54 $\text{nmol N}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$. Although V_{den} consistently exceeded V_{ana} , indicating the predominance of denitrification across all studied lakes, the contribution of anammox was substantial in specific lakes such as Lake Dongping (Tab. S5), where the maximum relative contribution (Ra) reached 34.34%. Sampling sites were categorized into eutrophic and mesotrophic groups based on the TLI assessment (Fig. S1o). Comparative analysis demonstrated that V_{den} , V_{ana} , and Ra were all significantly higher in eutrophic sites compared to mesotrophic ones ($p < 0.001$; Fig. 2).




Figure 2 Spatial patterns of nitrogen removal rates in the Eastern China Plain lakes (a) and their variations across sites with different trophic states (b–d). Note: (1) The size of the pie charts is proportional to the total nitrogen removal potential (the sum of mean denitrification and anammox rates). (2) Statistical significance was determined using the Mann-Whitney U test with Benjamini-Hochberg correction, indicated by asterisks (***: $p < 0.001$).

3.2 Spatial patterns of nitrogen removal functional gene abundance and diversity

Metagenomic analysis revealed that the key functional genes mediating denitrification and anammox were ubiquitous across all studied lakes, yet their abundances varied substantially (16–910,554 reads; Fig. S2). Denitrification genes overwhelmingly dominated the functional gene

删除[JIAHAO WEI]: , Tab. S6

pool, with a mean abundance of 1.98×10^6 reads, approximately three orders of magnitude higher than anammox genes (1.33×10^3 reads). Spatially, the major denitrification genes maintained a consistent abundance hierarchy (*nirK* > *nosZ* > *nirS*) across all lakes (Fig. 3a). For major anammox genes, *hzsB* exhibited the highest abundance (Fig. 3b). No significant inter-lake differences were found for *hzo*, *hzsB*, and *hzsC* ($p > 0.05$). Conversely, *hzsA* displayed distinct spatial heterogeneity, with significantly higher abundance in Lake Gaoyou compared to Lakes Poyang, Dongting, and the reservoir-type lakes ($p < 0.05$).




Figure 3 Spatial distribution of functional genes associated with sediment denitrification (a) and anammox (b) in the Eastern China Plain lakes. (c) Beta-diversity of functional gene profiles visualized via Principal Coordinates Analysis (PCoA). (d) Linear Discriminant Analysis Effect Size (LEfSe) identifying discriminative functional genes between groups.

Beta-diversity elucidated the spatial patterns of the nitrogen removal functional gene (Fig. 3c). Sampling sites overlapped extensively in the two-dimensional ordination space, with the first principal coordinate axis (PCoA1) explaining the majority of the variance (73.41%). However,

PERMANOVA tests ($R^2 = 0.798$, $p = 0.001$) indicated that significant differences in functional gene profiles did exist among lakes. Combined with similarity matrices and hierarchical clustering (Fig. S3), these differences were primarily driven by specific sites within Lakes Shaobo, Dongting, and Hengshui. The generally low clustering coefficient (< 0.03) suggest that, with the exception of a few specific habitats, the functional gene composition of sediments was highly similar across most lakes. LEfSe analysis further identified denitrification genes as the decisive discriminative features driving the variation in functional gene profiles among lakes (Fig. 3d). These genes consistently showed LDA scores ($\log_{10}$) exceeding 3 and were significantly enriched in Lakes Hengshui, Shaobo, and Dongting ($p < 0.01$). In comparison, although anammox genes were enriched in Lakes Shaobo, Gaoyou, and Taihu, their LDA scores were generally below 2, indicating a minor contribution to inter-group differences compared to denitrification genes.

3.3 Characteristics of microbial community structure

16S rRNA sequencing revealed that sediment microbial communities in the study area were primarily dominated by the phyla Chloroflexi and Proteobacteria, each exceeding 20% in relative abundance (Fig. 4a). Although taxa such as Desulfobacterota and Firmicutes exhibited lower overall relative abundances, they showed significant enrichment in specific lakes, suggesting that regional environmental strongly influence the proliferation of specific taxa. Cluster analysis indicated that lakes along the ER-SNWD (e.g., Lakes Hongze and Gaoyou) grouped together, reflecting high community similarity (Fig. S4). In contrast, Lakes Poyang and Dongting formed distinct, independent branches, highlighting the profound influence of regional environmental.

Alpha-diversity (Fig. 4b) indicated that Lakes Gaoyou, Gucheng, and the reservoir-type lakes possessed high species richness (Observed species > 1000) and Shannon diversity (> 6). The Simpson evenness was generally above 0.95, indicating a high overall community evenness. Conversely, Lake Dongting exhibited relatively lower indices, suggesting a stronger dominance effect by a few specific taxa. This implies that strong environmental filtering within these lakes has selectively enriched taxa equipped with specific adaptive traits. Beta-diversity revealed highly significant differences in community structure among lakes ($R^2 = 0.642$, $p = 0.001$). Sampling sites appeared highly dispersed in the two-dimensional ordination space (Fig. 4c). Most lake sites formed independent branches with clustering coefficient generally exceeding 0.4 (Fig. S5),

indicative of substantial inter-lake variation. LEfSe analysis further identified specific microbial taxa significantly enriched in different lakes, with marked variations in the number of discriminative taxa and LDA scores (Fig. 4d). Collectively, these findings underscore the critical role of regional environmental in driving microbial community.




Figure 4 Sediment microbial community characteristics in the Eastern China Plain lakes. (a) Taxonomic composition of the top 20 phyla. (b) Alpha-diversity indices, where "Observed species" denotes the number of observed amplicon sequence variants (ASVs). (c) Microbial beta-diversity based on PCoA. (d) LEfSe analysis identifying discriminative taxa at the phylum level between groups.

3.4 Relationships between biotic/abiotic factors and nitrogen removal rates

Spearman's correlations and Mantel tests revealed complex interplay between biotic/abiotic factors and nitrogen removal rates (Fig. S6). Both V_{den} and V_{ana} rates exhibited significant correlations with multiple physicochemical parameters, spatial structural variables (PCNM), and functional genes (e.g., *nirS*, *napC*). However, V_{ana} showed a stronger correlation with microbial community structure (Shannon diversity), whereas V_{den} was more strongly associated with hydro-chemical parameters. This divergence suggests that the regulatory mechanisms governing these two nitrogen removal processes are not identical.

First, linear regression analysis indicated no significant correlation between Human Activity Intensity (HAI) and nitrogen removal rates across different spatial scales ($p > 0.05$; Fig. 5A, B). In contrast, latitude showed significant positive correlations with both V_{den} ($R^2 = 0.215$) and V_{ana} ($R^2 = 0.315$, $p < 0.001$), and anammox may be subject to stricter latitudinal constraints. Random Forest (RF) models further identified the spatial structural variable PCNM2 as the shared major predictor for both V_{den} and V_{ana} (Fig. 5c). Organic matter indicators (COD_{Mn} and DOC) were identified as key drivers for V_{den} (Fig. 5i). In contrast, V_{ana} was more significantly influenced by ionic conditions (Cl^-) and substrate ($\text{NO}_2\text{-N}$) (Fig. 5j). Regarding functional genes, major denitrification genes (*nirK* and *nosZ*) were jointly regulated by climatic (MAT and MAP) and ionic conditions (Cl^- and SO_4^{2-}) (Fig. 5k, l). The key anammox gene (*hzsB*) was primarily driven by PCNM1, DOC, and redox conditions (DO) (Fig. 5m). Additionally, variations in Shannon diversity were dominated by PCNM2, with sediment trophic status (STN, Fig. S8) serving as another crucial predictor (Fig. 5n). Collectively, these results underscore the pivotal role of environmental differences resulting from spatial heterogeneity in shaping nitrogen removal processes.





Figure 5 Drivers of nitrogen removal processes in the Eastern China Plain lakes. Linear regressions of denitrification (A) and anammox (B) rates against Human Activity Intensity (HAI)




删除[JIAHAO WEI]:

281 and geographic location. The solid lines represent ordinary least squares (OLS) linear regressions,
282 with statistical significance evaluated using two-tailed Student's t-tests. (C) Key predictors of
283 nitrogen removal rates, functional gene abundance, and microbial Shannon diversity identified via
284 Random Forest regression models. Note: PCNM, Principal Coordinates of Neighbor Matrices;
285 DOC, dissolved organic carbon; SOC, sediment organic carbon; DO, dissolved oxygen; MAT,
286 mean annual temperature; MAP, mean annual precipitation; SW, sediment water content; STN,
287 sediment total nitrogen.

288 To disentangle the causal pathways of these multi-factor interactions, a Partial Least Squares Path
289 Model (PLS-PM) was constructed (Fig. 6). The models exhibited good predictive power, with
290 Goodness of Fit (GoF) values of 0.609 for denitrification and 0.598 for anammox, explaining
291 54.4% and 42.9% of the variance in V_{den} and V_{ana} , respectively. The results indicated that
292 water-sediment physicochemical properties exerted significant direct driving effects on both V_{den}
293 ($r = 0.784, p < 0.001$) and V_{ana} ($r = 0.420, p < 0.01$). V_{den} was also significantly directly driven by
294 microbial community structure ($r = 0.206, p < 0.05$). Crucially, effect decomposition highlighted
295 that while spatial structure had no significant direct effect, it exerted substantial influence via
296 indirect pathways mediated through climate, environmental conditions, and microbial
297 communities (Fig. 6b, d). Overall, the two nitrogen removal processes exhibited distinct
298 sensitivities to different drivers, highlights the complex and hierarchical regulatory patterns
299 underlying nitrogen removal in the Eastern China Plain lakes.

300



删除[JIAHAO WEI]:




Figure 6 Causal mechanisms driving nitrogen removal in the Eastern China Plain lakes. Partial Least Squares Path Modeling (PLS-PM) disentangling the direct and indirect effects of multiple drivers on denitrification (a) and anammox (c) rates, alongside their standardized total effects (b, d). Note: (1) Arrows represent significant unidirectional causal relationships. Red and blue arrows denote positive and negative effects, respectively. Numbers adjacent to arrows indicate standardized path coefficients (*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$). (2) R^2 values represent the proportion of variance explained for each dependent variable. (3) Goodness of Fit (GoF) is displayed at the top of each model. (4) To ensure model robustness, indicator variables for each latent construct were selected based on ecological relevance and preliminary collinearity screening, with measurement validity confirmed through outer model loadings and cross-loadings (retaining indicators with loadings > 0.70 where applicable). (5) PCNM, Principal Coordinates of Neighbor Matrices; DOC, dissolved organic carbon; DO, dissolved oxygen; MAT, mean annual temperature; MAP, mean annual precipitation; HAI, Human Activity Intensity.

4 Discussion

4.1 Eutrophication enhances the contribution of anammox to nitrogen removal in the Eastern China Plain lakes

This study systematically characterized the significant spatial heterogeneity of V_{den} and V_{ana} in the Eastern China Plain lakes (Fig. 2a, Tab. S5). Although both processes were ubiquitous across all surveyed lakes, distinct variations in rates were evident among individual lakes and lake zones, indicating that lacustrine nitrogen removal capacity is non-uniformly distributed at the regional scale (Jiang et al., 2023). Furthermore, previous studies have highlighted substantial variability in the relative contributions of denitrification and anammox (Deng et al., 2023; Stüeken et al., 2024), our results show that denitrification overwhelmingly dominated in all surveyed lakes (Tab. S5). However, the contribution of anammox was non-negligible in specific systems (Fig. 2), underscoring its distinct environmental dependency (Zhu et al., 2013). Furthermore, variations in latitudinal gradients among lakes introduce climatic differences such as regional temperature and precipitation. These differences, coupled with variations in geological backgrounds and watershed runoff generation and concentration processes, collectively highlight the critical regulatory role of spatial constraints—exerted by local environments—in shaping nitrogen removal processes.

删除[JIAHAO WEI]: These findings highlight the critical regulating role of the local environmental in shaping nitrogen removal processes.

Nitrogen and dissolved organic carbon (DOC) serve as critical metabolic substrates for nitrogen-removing microorganisms and act as key environmental factors of these processes (Lambert et al., 2024; Pascal et al., 2025). Our results demonstrated that both V_{den} and V_{ana} were significantly elevated in eutrophic sites compared to mesotrophic ones (Fig. 2b, c). Subjected to prolonged exogenous nutrient loading, the Eastern China Plain lakes generally sustain high substrate availability (Fig. S1) (Zhou et al., 2022). This enrichment likely enhances microbial substrate utilization efficiency, thereby stimulating and accelerating the corresponding nitrogen cycling processes (Zhou et al., 2021). Notably, previous studies have demonstrated potential substrate competition between DNRA and both denitrification and anammox processes. However, the reaction flux of DNRA is typically one to two orders of magnitude lower than that of denitrification. In typical eutrophic lakes, where $\text{NO}_3\text{-N}$ is generally abundant, the impact of this substrate competition on overall nitrogen removal rates remains relatively limited (Jiang et al., 2023). We recommend that future studies incorporate multi-isotope tracing techniques to achieve comprehensive, multi-pathway monitoring of the nitrogen cycle. Furthermore, organic carbon and

删除[JIAHAO WEI]: Nitrogen and dissolved organic carbon (DOC) serve as critical metabolic substrates for nitrogen-removing microorganisms and act as key environmental factors of these processes (Liu et al., 2018; Pascal et al., 2025). Our results demonstrated that both V_{den} and V_{ana} were significantly elevated in eutrophic sites compared to mesotrophic ones (Fig. 2b, c). Subjected to prolonged exogenous nutrient loading, the Eastern China Plain lakes generally sustain high substrate availability (Fig. S1) (Zhou et al., 2022). This enrichment likely enhances microbial substrate utilization efficiency, thereby stimulating and accelerating the corresponding nitrogen cycling processes (Zhou et al., 2021; Jiang et al., 2023). Organic carbon and ionic conditions act as critical electron donors for denitrification (Yoon et al., 2015). The effective supply of electrons (Fig. S1c–f) directly fueled elevated V_{den} in lakes such as Nansi and Luoma (Fig. 2a, Tab. S5). However, high concentrations of SO_4^{2-} may exert an inhibitory effect on denitrification through competitive inhibition for electron donors or sulfide toxicity (Jiang et al., 2020), which likely accounts for the relatively lower V_{den} observed in Lake Hengshui (Fig. S1f).

347 ionic conditions act as critical electron donors for denitrification (Yoon et al., 2015). The effective
348 supply of electrons (Fig. S1c–f) directly fueled elevated V_{den} in lakes such as Nansi and Luoma
349 (Fig. 2a, Tab. S5). However, high concentrations of SO_4^{2-} may exert an inhibitory effect on
350 denitrification through competitive inhibition for electron donors or sulfide toxicity (Jiang et al.,
351 2020), which likely accounts for the relatively lower V_{den} observed in Lake Hengshui (Fig. S1f).

352 Intriguingly, although denitrification remained dominant under eutrophic conditions, the relative
353 contribution of anammox significantly increased (Fig. 2d). This shift may be attributed to the
354 susceptibility of denitrification to factors such as electron donor (organic carbon) allocation and
355 redox conditions (Carlson et al., 2013). Additionally, given that denitrifying communities possess
356 a broader tolerance range for oxygen compared to anammox communities (Deng et al., 2020),
357 processes such as DOC mineralization (Yang et al., 2025) and variations in sediment water content
358 (Tab. S6) and oxygen permeability (Sun et al., 2023) can alter local oxygen concentrations. This
359 result creates more intensely anoxic or anaerobic niches, which favor the survival of anammox
360 bacteria (Yang et al., 2026). This process also stimulates heterotrophic denitrification, which in
361 turn supplies crucial intermediate substrates $\text{NO}_2\text{-N}$ for anammox, thereby boosting the
362 contribution of anammox to lacustrine nitrogen removal (Tsementzi et al., 2016).

删除[JIAHAO WEI]: 7

363 Collectively, these findings suggest that under specific substrate concentrations and redox
364 conditions, anammox can form a synergistic or complementary relationship with denitrification.
365 This complementarity ultimately enhances both the magnitude and stability of the nitrogen
366 removal function in lake ecosystems (Kuypers et al., 2018), bearing profound ecological
367 significance for facilitating nitrogen cycling and buffering against environmental changes. As
368 "sinks" within watershed ecosystems, lakes integrate substantial material and energy inputs
369 (Bergström and Karlsson, 2019). This results in a complex scenario where nitrogen cycling is
370 simultaneously regulated by multiple environmental variables, leading to the pronounced spatial
371 heterogeneity observed in lacustrine nitrogen removal capacity.

删除[JIAHAO WEI]:

372 **4.2 Pronounced decoupling among microbial community, functional gene abundance, and** 373 **nitrogen removal rates**

374 As quintessential microbe-mediated biogeochemical processes, the intensities of denitrification
375 and anammox are fundamentally governed by microbial community structure and the distribution

of functional genes (Jiang et al., 2020). Our study revealed a certain degree of coupling between metagenome-derived functional gene abundance/diversity and nitrogen removal rates. Specifically, key denitrification genes (e.g., *nirK*, *nosZ*) exhibited significant correlations with nitrogen removal rates (Fig. S6), aligning with previous assertions that gene abundance can serve as a proxy for nitrogen removal potential (Xiong et al., 2017).

However, it is crucial to note that functional gene abundance does not always scale linearly with nitrogen removal rates (Dandie et al., 2011; Ma et al., 2020). We observed a highly consistent functional gene composition across the studied lakes, where diversity variations were dictated by only a few outlier samples (Fig. 3c, S3). This indicates a high similarity in the potential nitrogen removal capacity among lakes, which stands in sharp contrast to the significant spatial heterogeneity observed in the measured rates. Furthermore, the substantial discrepancy—approximately three orders of magnitude—between denitrification and anammox gene abundances (Tab. S5) was not proportionally reflected in their respective rate differences. This non-linear association likely stems from the fact that the presence of genes at the DNA level represents only the genetic potential and does not guarantee their transcription and translation into active proteins (Ma et al., 2020). Path modeling (PLS-PM), which showed no significant direct effect of functional genes on rates, further corroborated this phenomenon (Fig. 6a, c). Consequently, assessing regional nitrogen removal capacity based solely on functional gene abundance may lead to misjudgments of actual rates.

Beyond the genetic level, microbial community structure is also a critical driver of spatial differentiation in nitrogen removal (Fig. 6). Variations in the dominance of functional taxa fundamentally determine the system's potential to execute complete nitrogen removal processes (Sveen et al., 2025). Our results showed that core nitrogen-cycling taxa, such as Proteobacteria and Nitrospirota, exceeded 20% relative abundance in all lakes (Fig. 4a), with specific dominant taxa significantly enriched in different lakes (Fig. 4d). This reflects a robust potential for nitrogen cycling and organic matter degradation across the Eastern China Plain lakes (Han et al., 2024). However, shifts in the proportions of these dominant taxa may alter community response thresholds to substrates and environmental conditions, thereby driving the spatial divergence of nitrogen removal rates (Cline and Zak, 2015; Egidi et al., 2023).

Notably, V_{den} responded more significantly to microbial community structure than V_{ana} (Fig. 6a, b). This distinction may be attributed to the modular nature of denitrification, a stepwise process performed by a diverse array of microorganisms (Kuypers et al., 2018; Patchett et al., 2025). The generally high microbial diversity observed in the study area (Fig. 4b, c) implies a high level of functional redundancy (Louca et al., 2018). This taxonomic diversity allows distinct taxa to perform identical or similar metabolic functions, enabling denitrifying communities to occupy broader ecological niches. In contrast, anammox is a specialized, narrow ecological niche process restricted to a limited number of bacteria within the phylum Planctomycetes (Wang et al., 2019), making its rate strictly constrained by the population size and gene expression of key species. Consequently, the differential stability of these functional communities in the face of environmental fluctuations (Hernandez et al., 2023) results in the observed divergence in process rates.

4.3 Multiple driving effects of macro-environmental heterogeneity on nitrogen removal processes

Macro-ecological studies consistently demonstrate that nitrogen cycling is significantly correlated with geographic factors such as latitude, longitude, and altitude (Fuhrman et al., 2008; Tu et al., 2019). Our findings reveal that latitudinal gradients and spatial structural variables (e.g., PCNM2) constitute the primary macro-scale background governing nitrogen removal processes across the Eastern China Plain lakes (Fig. 5, S6). Specifically, latitude acts as a robust proxy for climatic drivers—including temperature, precipitation, and seasonal irradiance (Ding et al., 2025)—consistent with the subtropical to temperate monsoon climate characteristics of this region (Method S1, Tab. S3). Such climatic variations reshape microbial habitat quality by modulating hydraulic residence times, organic matter mineralization pathways, and the thermal, dissolved oxygen, and nutrient stratification regimes (Manzoni et al., 2014; Hu et al., 2020). These shifts subsequently exert profound effects on functional gene abundance, highlighting the critical roles of dispersal limitation and spatial heterogeneity in shaping microbial biogeography (Du et al., 2023). Notably, the response of V_{ana} to latitude was significantly stronger than that of V_{den} (Fig. 5c, g). This discrepancy is likely attributable to the broader metabolic versatility and phylogenetic

删除[JIAHAO WEI]: Macro-ecological studies consistently demonstrate that nitrogen cycling is significantly correlated with geographic factors such as latitude, longitude, and altitude (Fuhrman et al., 2008; Tu et al., 2019). Our study reveals that the latitudinal gradient and spatial structural variables (PCNM2) constitute the primary macro-scale background regulating nitrogen removal processes in the Eastern China Plain lakes (Fig. 5, S6). Essentially, latitude serves as a proxy for climatic effects—including temperature, precipitation, and seasonal irradiance (Ding et al., 2025)—which aligns with the subtropical and temperate monsoon climate characteristics of the region (Method S1, Tab. S3). Climate reshapes microbial habitat quality by modulating hydraulic residence time, organic matter mineralization pathways, and the stratification stability of thermal regimes, dissolved oxygen, and nutrients (Manzoni et al., 2014; Hu et al., 2020). And subsequently exert profound impacts on functional gene abundance, highlighting the critical roles of dispersal limitation and spatial heterogeneity in shaping microbial biogeography (Du et al., 2023). Notably, the response of V_{ana} to latitude was significantly stronger than that of V_{den} (Fig. 5c, g). This discrepancy is likely attributable to the broader metabolic versatility and phylogenetic distribution of denitrifying communities. In contrast, anammox communities possess a narrower ecological niche specificity and higher sensitivity to macro-environmental fluctuations, subjecting their spatial distribution to stricter geographic constraints (Gao et al., 2018).

设置格式[JIAHAO WEI]: 字体: (默认) Times New Roman, 五号

设置格式[JIAHAO WEI]: 字体: (默认) Times New Roman

设置格式[JIAHAO WEI]: 字体: (默认) Times New Roman

设置格式[JIAHAO WEI]: 字体: (默认) Times New Roman, 五号

distribution of denitrifying communities. Conversely, anammox communities exhibit narrower ecological niche specialization and heightened sensitivity to macro-environmental fluctuations, rendering their spatial distribution subject to stricter geographic constraints (Gao et al., 2018).

Despite the study area being situated in Eastern China—a region characterized by intense anthropogenic disturbance (Fig. S7)—the Human Activity Intensity (HAI) across various buffer zone scales showed no direct, significant effects on nitrogen removal rates (Fig. 5A, B). This decoupling suggests that anthropogenic influences undergo a sequential process of environmental filtering followed by micro-environmental redistribution (Keck et al., 2025). Specifically, while human activities drive external loadings of nitrogen, phosphorus, and organic matter, thereby altering lake trophic states and sediment characteristics (Wei et al., 2024), it is the local water-sediment physicochemical properties that constitute the immediate habitat for microbial metabolism. These local factors directly regulate functional genes and nitrogen removal rates through substrate availability and redox conditions, acting as the primary proximate drivers responsible for the spatial heterogeneity observed in this study (Fig. 6a, c).

It is noteworthy that anthropogenic activities and water-sediment physicochemical properties did not exert significant direct effects on microbial community structure, yet significantly shaped the distribution of functional genes (Fig. 6). This suggests that microbial communities in the Eastern China Plain lakes have achieved high structural stability following long-term environmental filtering. Given the generally high microbial diversity observed (Fig. 4b, c), the system likely possesses a high level of functional redundancy (Louca et al., 2018). When environmental physicochemical properties fluctuate due to anthropogenic or other factors, communities may prioritize regulating functional gene expression to maintain metabolic homeostasis rather than undergoing large-scale taxonomic restructuring. Consequently, the response of community structure to environmental changes exhibits a certain hysteresis or dampening effect (Martiny et al., 2023). Moreover, the robust buffering effects provided by microbial biofilm architectures and community dispersal processes (Yan et al., 2024) also partially explain why physicochemical fluctuations failed to translate directly and linearly into shifts in community structure (Kuypers et al., 2018).

设置格式[JIAHAO WEI]: 字体: 小五

删除[JIAHAO WEI]: Despite the study area being located in eastern China, a region characterized by intense anthropogenic disturbance (Fig. S7), the Human Activity Intensity (HAI) across different buffer zones scales exhibited no significant direct effects on nitrogen removal rates (Fig. 5A, B). This influence likely underwent a process of environmental filtering followed by redistribution (Keck et al., 2025). Specifically, while anthropogenic activities drive the input of nitrogen, phosphorus, and organic matter, thereby altering the trophic state and sediment characteristics (Wei et al., 2024), it is the local water-sediment physicochemical properties that constitute the immediate habitat for microbial metabolism. These local factors directly regulate functional genes and nitrogen removal rates through substrate and redox conditions, serving as the primary drivers explaining the spatial heterogeneity observed in this study (Fig. 6a, c).

设置格式[JIAHAO WEI]: 字体: (默认) Times New Roman, 五号

设置格式[JIAHAO WEI]: 字体: 小五

463 **5 Conclusion**

464 Elucidating the driving mechanisms of sediment nitrogen removal is crucial for understanding the
465 resilience of lacustrine ecosystems to eutrophication. This study reveals that while denitrification
466 dominates nitrogen removal in the Eastern China Plain lakes, anammox serves as a critical, highly
467 environmentally dependent complementary process. Notably, eutrophication significantly
468 amplified both the overall nitrogen removal capacity and the relative contribution of anammox,
469 with local physicochemical properties and microbial communities demonstrating a pronounced
470 joint driving effect on these processes. The observed decoupling among microbial community,
471 functional gene abundance, and nitrogen removal rates highlights the intricate complexity of
472 microbial responses to environmental perturbations. This complexity is further evidenced by the
473 stark contrast between the functional redundancy of denitrifying bacteria and the ecological niche
474 specificity of anammox bacteria. Furthermore, we demonstrate that macro-scale factors (spatial,
475 climatic, and anthropogenic) indirectly drive nitrogen removal by reshaping local water-sediment
476 physicochemical properties, rather than acting as direct determinants of process rates. These
477 findings enrich our fundamental understanding of nitrogen removal, particularly the anammox
478 process, and its multidimensional drivers in the Eastern China Plain lakes. Given the sensitivity of
479 nitrogen removal processes to habitat conditions, we suggest validating driving mechanisms using
480 key enzyme activity assays and targeted controlled experiments. Furthermore, we advocate that
481 future lake management strategies must rigorously account for the overarching constraints jointly
482 imposed by trophic status, alongside spatial and climatic gradients, to ensure the sustainability of
483 nitrogen purification ecosystem services.**Data availability**

484 All data analyzed in this study are available in figshare
485 (<https://doi.org/10.6084/m9.figshare.32005818>).

486 **Author contributions**

487 JHW: conceptualization, sampling, data analysis, manuscript preparation. KY: analyzed the
488 samples and original draft preparation. YRL: designed figures and tables, original draft
489 preparation. YY: original draft preparation. LCW: design and guidance of experiments, draft
490 revision. ELZ: draft revision, funding acquisition; ZJG: design and guidance of experiments, draft

删除[JIAHAO WEI]: Given the profound sensitivity of nitrogen removal processes to habitat conditions, we advocate that future lake management strategies must rigorously account for the overarching constraints jointly imposed by trophic status, alongside spatial and climatic gradients, to ensure the sustainability of nitrogen purification ecosystem services.

删除[JIAHAO WEI]:

revision, funding acquisition; YJC: design and guidance of experiments, draft revision, funding acquisition.

Competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Disclaimer

Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors. Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.

Acknowledgments

We thank the Institutional Center for Shared Technologies and Facilities of NIGALS, CAS for technical support with physicochemical analyses. We are also grateful to Jiashuai Wen, Weihang Wang, Chennan Wang, Xin Zhang, Yao Zhang, Ziyang Wang, and Zhen Shen for their assistance during field investigations, and to Prof. Hai Xu, Dr. Xiangming Tang, and Yu Qiu for their experimental guidance.

Financial support

This work was supported by the National Natural Science Foundation of China (U24A20639, 42171119), the State Key Laboratory of Lake and Watershed Science for Water Security (NKL2023-ZD02, NKL2023-KP01).

References

Bergström, A. K. and Karlsson, J.: Light and nutrient control phytoplankton biomass responses to global change in northern lakes, *Global Change Biology*, 25(6), 2021-2029, <https://doi.org/10.1111/gcb.14623>, 2019. 删除[JIAHAO WEI]: [1]

Borcard, D. and Legendre, P.: All-scale spatial analysis of ecological data by means of principal coordinates of neighbour matrices, *Ecological Modelling*, 153(1-2), 51-68, [https://doi.org/10.1016/S0304-3800\(01\)00501-4](https://doi.org/10.1016/S0304-3800(01)00501-4), 2002. 删除[JIAHAO WEI]: [2]

519 Breiman, L.: Random forests, Machine Learning, 45(1), 5-32, 删除[JIAHAO WEI]: [3]
520 <https://doi.org/10.1023/A:1010933404324>, 2001.

521 Carlson, H. K., Clark, I. C., Blazewicz, S. J., Iavarone, A. T., and Coates, J. D.: Fe (II)
522 oxidation is an innate capability of nitrate-reducing bacteria that involves abiotic and
523 biotic reactions, Journal of Bacteriology, 195(14), 3260-3268, 删除[JIAHAO WEI]: [4]
524 <https://doi.org/10.1128/Jb.00058-13>, 2013.

525 Cline, L. C. and Zak, D. R.: Soil microbial communities are shaped by plant-driven changes
526 in resource availability during secondary succession, Ecology, 96(12), 3374-3385, 删除[JIAHAO WEI]: [5]
527 <https://doi.org/10.1890/15-0184.1>, 2015.

528 Dai, T. J., Wen, D. H., Bates, C. T., Wu, L. W., Guo, X., Liu, S., Su, Y. F., Lei, J. S., Zhou, J.
529 Z., and Yang, Y. F.: Nutrient supply controls the linkage between species abundance and
530 ecological interactions in marine bacterial communities, Nature Communications, 13, 175, 删除[JIAHAO WEI]: [6]
531 <https://doi.org/10.1038/s41467-021-27857-6>, 2022.

532 Dai, Y. H., Yang, S. B., Zhao, D., Hu, C. M., Xu, W., Anderson, D. M., Li, Y., Song, X. P.,
533 Boyce, D. G., Gibson, L., Zheng, C. M., and Feng, L.: Coastal phytoplankton blooms
534 expand and intensify in the 21st century, Nature, 615(7951), 280-284, 删除[JIAHAO WEI]: [7]
535 <https://doi.org/10.1038/s41586-023-05760-y>, 2023.

536 Dandie, C. E., Wertz, S., Leclair, C. L., Goyer, C., Burton, D. L., Patten, C. L., Zebarth, B. J.,
537 and Trevors, J. T.: Abundance, diversity and functional gene expression of denitrifier
538 communities in adjacent riparian and agricultural zones, FEMS Microbiology Ecology,
539 77(1), 69-82, <https://doi.org/10.1111/j.1574-6941.2011.01084.x>, 2011. 删除[JIAHAO WEI]: [8]

540 Deng, D. L., Ding, B. J., He, G., Ji, M. F., Yang, Y. Y., Liu, G. H., Zhang, Q. F., and Liu, W. Z.:
541 The contribution of anammox to nitrogen removal is greater in bulk soils than in
542 rhizosphere soils in riparian wetlands along the Yangtze River, Global Biogeochemical
543 Cycles, 37(5), e2022GB007576, <https://doi.org/10.1029/2022GB007576>, 2023. 删除[JIAHAO WEI]: The Contribution of Anammox to
Nitrogen Removal Is Greater in Bulk Soils Than in
Rhizosphere Soils in Riparian Wetlands Along the Yangtze
River

544 Deng, D. L., Pan, Y. T., Liu, G. H., Liu, W. Z., and Ma, L.: Seeking the hotspots of nitrogen
545 removal: A comparison of sediment denitrification rate and denitrifier abundance among
546 wetland types with different hydrological conditions, Science of the Total Environment,
547 737, 140253, <https://doi.org/10.1016/j.scitotenv.2020.140253>, 2020. 删除[JIAHAO WEI]: [10]

548 Deng, D. L., Yang, Z. J., Yang, Y. Y., Wan, W. J., Liu, W. Z., and Xiong, X.: Metagenomic
549 insights into nitrogen-cycling microbial communities and their relationships with nitrogen
550 removal potential in the Yangtze River, Water Research, 265(1), 122229, 删除[JIAHAO WEI]: [11]
551 <https://doi.org/10.1016/j.watres.2024.122229>, 2024.

552 Ding, D. D., Shi, Z. J., Chen, J. S., Zhang, H., Meng, Q. Q., and Han, W. X.: Solar radiation
553 and water stress are major environmental drivers of the latitudinal pattern of leaf Si
554 concentration in woody plants, Global Ecology and Biogeography, 34, e70122, 删除[JIAHAO WEI]: [12]
555 <https://doi.org/10.1111/geb.70122>, 2025.

556 Du, X. F., Gu, S. S., Zhang, Z., Li, S. Z., Zhou, Y. Q., Zhang, Z. J., Zhang, Q., Wang, L. L., Ju,
557 Z. C., Yan, C. L., Li, T., Wang, D. R., Yang, X. S., Peng, X., and Deng, Y.: Spatial
558 distribution patterns across multiple microbial taxonomic groups, Environmental Research,
559 223, 115470, <https://doi.org/10.1016/j.envres.2023.115470>, 2023. 删除[JIAHAO WEI]: [13]

560 Egidi, E., Coleine, C., Delgado-Baquerizo, M., and Singh, B. K.: Assessing critical thresholds
561 in terrestrial microbiomes, Nature Microbiology, 8(12), 2230-2233, 删除[JIAHAO WEI]: [14]
562 <https://doi.org/10.1038/s41564-023-01536-2>, 2023.

- 563 Fuhrman, J. A., Steele, J. A., Hewson, I., Schwalbach, M. S., Brown, M. V., Green, J. L., and
564 Brown, J. H.: A latitudinal diversity gradient in planktonic marine bacteria, *PNAS*, 105(22),
565 7774-7778, <https://doi.org/10.1073/pnas.0803070105>, 2008. 删除[JIAHAO WEI]: [15]
- 566 Gao, D. W., Wang, X. L., Liang, H., Wei, Q. H., Dou, Y., and Li, L. W.: Anaerobic ammonia
567 oxidizing bacteria: ecological distribution, metabolism, and microbial interactions,
568 *Frontiers of Environmental Science and Engineering*, 12, 10,
569 <https://doi.org/10.1007/s11783-018-1035-x>, 2018. 删除[JIAHAO WEI]: [16]
- 570 Han, P., Tang, X. F., Koch, H., Dong, X. Y., Hou, L. J., Wang, D. H., Zhao, Q., Li, Z., Liu, M.,
571 Lückner, S., and Shi, G. T.: Unveiling unique microbial nitrogen cycling and nitrification
572 driver in coastal Antarctica, *Nature Communications*, 15, 3143,
573 <https://doi.org/10.1038/s41467-024-47392-4>, 2024. 删除[JIAHAO WEI]: [17]
- 574 Harris, I., Osborn, T. J., Jones, P., and Lister, D.: Version 4 of the CRU TS monthly
575 high-resolution gridded multivariate climate dataset, *Scientific Data*, 7, 109,
576 <https://doi.org/10.1038/s41597-020-0453-3>, 2020. 删除[JIAHAO WEI]: [18]
- 577 Hernandez, D. J., Kieseewetter, K. N., Almeida, B. K., Revillini, D., and Afkhami, M. E.:
578 Multidimensional specialization and generalization are pervasive in soil prokaryotes,
579 *Nature Ecology & Evolution*, 7, 1408-1418, <https://doi.org/10.1038/s41559-023-02149-y>,
580 2023. 删除[JIAHAO WEI]: [19]
- 581 Ho, J. C., Michalak, A. M., and Pahlevan, N.: Widespread global increase in intense lake
582 phytoplankton blooms since the 1980s, *Nature*, 574(7780), 667-670,
583 <https://doi.org/10.1038/s41586-019-1648-7>, 2019. 删除[JIAHAO WEI]: [20]
- 584 Hou, X. J., Feng, L., Dai, Y. H., Hu, C. A. M., Gibson, L., Tang, J., Lee, Z. P., Wang, Y., Cai,
585 X. B., Liu, J. G., Zheng, Y., and Zheng, C. M.: Global mapping reveals increase in
586 lacustrine algal blooms over the past decade, *Nature Geoscience*, 15(2), 130-134,
587 <https://doi.org/10.1038/s41561-021-00887-x>, 2022. 删除[JIAHAO WEI]: [21]
- 588 Hu, Y. L., Wang, S., Niu, B., Chen, Q. Y., Wang, J., Zhao, J. X., Luo, T. X., and Zhang, G. X.:
589 Effect of increasing precipitation and warming on microbial community in Tibetan alpine
590 steppe, *Environmental Research*, 189, 109917,
591 <https://doi.org/10.1016/j.envres.2020.109917>, 2020. 删除[JIAHAO WEI]: [22]
- 592 Huisman, J., Codd, G. A., Paerl, H. W., Ibelings, B. W., Verspagen, J. M. H., and Visser, P. M.:
593 Cyanobacterial blooms, *Nature Reviews Microbiology*, 16(8), 471-483,
594 <https://doi.org/10.1038/s41579-018-0040-1>, 2018. 删除[JIAHAO WEI]: [23]
- 595 Hutchins, D. A. and Tagliabue, A.: Feedbacks between phytoplankton and nutrient cycles in a
596 warming ocean, *Nature Geoscience*, 17(6), 495-502,
597 <https://doi.org/10.1038/s41561-024-01454-w>, 2024. 删除[JIAHAO WEI]: [24]
- 598 IPCC: Climate change 2022: Impacts, adaptation and vulnerability, p. 3056, Cambridge
599 University Press, Cambridge, UK and New York, USA, 2022. 删除[JIAHAO WEI]: [25]
- 600 Jiang, X. Y., Gao, G., Zhang, L., Tang, X. M., Shao, K. Q., and Hu, Y.: Denitrification and
601 dissimilatory nitrate reduction to ammonium in freshwater lakes of the Eastern Plain,
602 China: Influences of organic carbon and algal bloom, *Science of the Total Environment*,
603 710, 136303, <https://doi.org/10.1016/j.scitotenv.2019.136303>, 2020. 删除[JIAHAO WEI]: [26]
- 604 Jiang, X. Y., Liu, C. Q., Cai, J., Hu, Y., Shao, K. Q., Tang, X. M., Gong, Y., Yao, X. L., Xu, Q.
605 J., and Gao, G.: Relationships between environmental factors and N-cycling microbes
606 reveal the indirect effect of further eutrophication on denitrification and DNRA in shallow 删除[JIAHAO WEI]: [27]

607 lakes, *Water Research*, 245, 120572, <https://doi.org/10.1016/j.watres.2023.120572>, 2023.

608 Keck, F., Peller, T., Alther, R., Barouillet, C., Blackman, R., Capo, E., Chonova, T., Couton, 删除[JIAHAO WEI]: [28]
609 M., Fehlinger, L., Kirschner, D., Knüsel, M., Muneret, L., Oester, R., Tapolczai, K., Zhang,
610 H., and Altermatt, F.: The global human impact on biodiversity, *Nature*, 641, 395-400,
611 <https://doi.org/10.1038/s41586-025-08752-2>, 2025.

612 Kuypers, M. M. M., Marchant, H. K., and Kartal, B.: The microbial nitrogen-cycling network, 删除[JIAHAO WEI]: [29]
613 *Nature Reviews Microbiology*, 16(5), 263-276, <https://doi.org/10.1038/nrmicro.2018.9>,
614 2018.

615 Lambert, T., Dupas, R., and Durand, P.: The role of nitrogen and iron biogeochemical cycles 删除[JIAHAO WEI]: [30]
616 in the production and export of dissolved organic matter in agricultural headwater
617 catchments, *Biogeosciences*, 21(20), 4533–4547, <https://doi.org/10.5194/bg-21-4533-2024>,
618 2024.

619 Liu, C., Li, X. Z., Mansoldo, F. R. P., An, J. X., Kou, Y. P., Zhang, X., Zeng, J. X., Vermelho, 删除[JIAHAO WEI]: [31]
620 A. B., Wang, J. M., and Yao, M. J.: Microbial habitat specificity largely affects microbial
621 co-occurrence patterns and functional profiles in wetland soils, *Geoderma*, 418, 115866,
622 <https://doi.org/10.1016/j.geoderma.2022.115866>, 2022.

623 Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., O'Connor, M. I., 删除[JIAHAO WEI]: [32]
624 Ackermann, M., Hahn, A. S., Srivastava, D. S., Crowe, S. A., Doebeli, M., and Parfrey, L.
625 W.: Function and functional redundancy in microbial systems, *Nature Ecology &*
626 *Evolution*, 2(6), 936-943, <https://doi.org/10.1038/s41559-018-0519-1>, 2018.

627 Ma, L., Jiang, X. L., Liu, G. H., Yao, L. G., Liu, W. Z., Pan, Y. T., and Zuo, Y. X.: 删除[JIAHAO WEI]: [33]
628 Environmental factors and microbial diversity and abundance jointly regulate soil nitrogen
629 and carbon biogeochemical processes in Tibetan Wetlands, *Environmental Science &*
630 *Technology*, 54(6), 3267-3277, <https://doi.org/10.1021/acs.est.9b06716>, 2020. Ma, R. H., 删除[JIAHAO WEI]:
631 Duan, H. T., Hu, C. M., Feng, X. Z., Li, A. N., Ju, W. M., Jiang, J. H., and Yang, G. S.: A [34]
632 half-century of changes in China's lakes: Global warming or human influence?
633 *Geophysical Research Letters*, 37(24), L24106, <https://doi.org/10.1029/2010gl045514>,
634 2010.

635 Manzoni, S., Schaeffer, S. M., Katul, G., Porporato, A., and Schimel, J. P.: A theoretical 删除[JIAHAO WEI]: [35]
636 analysis of microbial eco-physiological and diffusion limitations to carbon cycling in
637 drying soils, *Soil Biology and Biochemistry*, 73, 69-83,
638 <https://doi.org/10.1016/j.soilbio.2014.02.008>, 2014.

639 Martiny, J. B. H., Martiny, A. C., Brodie, E., Chase, A. B., Rodriguez-Verdugo, A., Treseder, 删除[JIAHAO WEI]: [36]
640 K. K., and Allison, S. D.: Investigating the eco-evolutionary response of microbiomes to
641 environmental change, *Ecology Letters*, 26, S81-S90, <https://doi.org/10.1111/ele.14209>,
642 2023.

643 Mendonça, R., Müller, R. A., Clow, D., Verpoorter, C., Raymond, P., Tranvik, L. J., and Sobek, 删除[JIAHAO WEI]: [37]
644 S.: Organic carbon burial in global lakes and reservoirs, *Nature Communications*, 8, 1694,
645 <https://doi.org/10.1038/s41467-017-01789-6>, 2017.

646 Pascal, L., Cloutier-Artiwat, F., Zanon, A., Wallace, D. W. R., and Chaillou, G.: New 删除[JIAHAO WEI]: [38]
647 Deoxygenation Threshold for N₂, and N₂O Production in Coastal Waters and Sediments,
648 *Global Biogeochemical Cycles*, 39(8), e2024GB008218,
649 <https://doi.org/10.1029/2024GB008218>, 2025. Patchett, A., Rütting, L., Rütting, T., Bodé, 删除[JIAHAO WEI]:
650 S., Hallin, S., Juhanson, J., Stange, C. F., Björkman, M. P., Boeckx, P., Rosqvist, G., and [39]

651 Björk, R. G.: The role of mycorrhizal type and plant dominance in regulating nitrogen
652 cycling in Oroarctic soils, *Biogeosciences*, 22(22), 6841–6860,
653 <https://doi.org/10.5194/bg-22-6841-2025>, 2025.

654 Steinegger, M. and Söding, J.: Clustering huge protein sequence sets in linear time, *Nature*
655 *Communications*, 9, 2542, <https://doi.org/10.1038/s41467-018-04964-5>, 2018. 删除[JIAHAO WEI]: [40]

656 Stüeken, E. E., Pellerin, A., Thomazo, C., Johnson, B. W., Duncanson, S., and Schoepfer, S.
657 D.: Marine biogeochemical nitrogen cycling through Earth's history, *Nature Reviews Earth*
658 *& Environment*, 5(10), 732–747, <https://doi.org/10.1038/s43017-024-00591-5>, 2024. 删除[JIAHAO WEI]: [41]

659 Sun, F. S., Ma, C., Yu, G. H., Kuzyakov, Y., Lang, Y. C., Fu, P. Q., Guo, L. J., Teng, H. H.,
660 and Liu, C. Q.: Organic carbon preservation in wetlands: Iron oxide protection vs.
661 thermodynamic limitation, *Water Research*, 241, 120133,
662 <https://doi.org/10.1016/j.watres.2023.120133>, 2023. 删除[JIAHAO WEI]: [42]

663 Sveen, T. R., Viketoft, M., Bengtsson, J., Strengbom, J., Lejoly, J., Buegger, F., Pritsch, K.,
664 Fritscher, J., Hildebrand, F., Osburn, E., Hannula, S. E., and Bahram, M.: Functional
665 diversity of soil microbial communities increases with ecosystem development, *Nature*
666 *Communications*, 16, 10408, <https://doi.org/10.1038/s41467-025-66544-8>, 2025. 删除[JIAHAO WEI]: [43]

667 Thamdrup, B. and Dalsgaard, T.: Production of N through anaerobic ammonium oxidation
668 coupled to nitrate reduction in marine sediments, *Applied and Environmental*
669 *Microbiology*, 68(3), 1312–1318, <https://doi.org/10.1128/Aem.68.3.1312-1318.2002>,
670 2002. Tsementzi, D., Wu, J. Y., Deutsch, S., Nath, S., Rodriguez, L. M., Burns, A. S.,
671 Ranjan, P., Sarode, N., Malmstrom, R. R., Padilla, C. C., Stone, B. K., Bristow, L. A.,
672 Larsen, M., Glass, J. B., Thamdrup, B., Woyke, T., Konstantinidis, K. T., and Stewart, F. J.:
673 SAR11 bacteria linked to ocean anoxia and nitrogen loss, *Nature*, 536(7615), 179–183,
674 <https://doi.org/10.1038/nature19068>, 2016. 删除[JIAHAO WEI]: [44]

675 Tu, Q. C., Lin, L., Cheng, L., Deng, Y., and He, Z. L.: NCycDB: a curated integrative
676 database for fast and accurate metagenomic profiling of nitrogen cycling genes,
677 *Bioinformatics*, 35(6), 1040–1048, <https://doi.org/10.1093/bioinformatics/bty741>, 2019. 删除[JIAHAO WEI]: [45]

678 van Helmond, N. A. G. M., Robertson, E. K., Conley, D. J., Hermans, M., Humborg, C.,
679 Kubeneck, L. J., Lenstra, W. K., and Slomp, C. P.: Removal of phosphorus and nitrogen in
680 sediments of the eutrophic Stockholm archipelago, Baltic Sea, *Biogeosciences*, 17(10),
681 2745–2766, <https://doi.org/10.5194/bg-17-2745-2020>, 2020. 删除[JIAHAO WEI]: [47]

682 Wang, C. R., He, T. X., Zhang, M. M., Zheng, C. X., Yang, L., and Yang, L.: Review of the
683 mechanisms involved in dissimilatory nitrate reduction to ammonium and the efficacies of
684 these mechanisms in the environment, *Environmental Pollution*, 345, 123480,
685 <https://doi.org/10.1016/j.envpol.2024.123480>, 2024. 删除[JIAHAO WEI]: [48]

686 Wang, S. Y., Liu, W. Y., Zhao, S. Y., Wang, C., Zhuang, L. J., Liu, L., Wang, W. D., Lu, Y. L.,
687 Li, F. B., and Zhu, G. B.: Denitrification is the main microbial N loss pathway on the
688 Qinghai-Tibet Plateau above an elevation of 5000 m, *Science of the Total Environment*,
689 696, 133852, <https://doi.org/10.1016/j.scitotenv.2019.133852>, 2019. 删除[JIAHAO WEI]: [49]

690 Wang, Y., Zhao, D., Woolway, R. I., Yan, H. R., Paerl, H. W., Zheng, Y., Zheng, C. M., and
691 Feng, L.: Global elevation of algal bloom frequency in large lakes over the past two
692 decades, *National Science Review*, 12(3), nwaf011, <https://doi.org/10.1093/nsr/nwaf011>,
693 2025. 删除[JIAHAO WEI]: [50]

694 Wei, J. H., Zhang, Q. J., Yin, Y., Peng, K., Wang, L. C., Cai, Y. J., and Gong, Z. J.: Limited
删除[JIAHAO WEI]: [51]

695 impacts of water diversion on micro-eukaryotic community along the Eastern Route of
696 China's South-to-North Water Diversion Project, *Water Research*, 262, 122109,
697 <https://doi.org/10.1016/j.watres.2024.122109>, 2024.

698 ✓Woolway, R. I., Kraemer, B. M., Lenters, J. D., Merchant, C. J., O'Reilly, C. M., and Sharma,
699 S.: Global lake responses to climate change, *Nature Reviews Earth & Environment*, 1,
700 388-403, <https://doi.org/10.1038/s43017-020-0067-5>, 2020. 删除[JIAHAO WEI]: [52]

701 ✓Woolway, R. I., Zhang, Y. L., Jennings, E., Zohary, T., Jane, S. F., Jansen, J., Weyhenmeyer, G.
702 A., Long, D., Fleischmann, A., Feng, L., Qin, B. Q., Shi, K., Shi, H. R., Wang, W. J., Tong,
703 Y., Zhang, G. Q., Zscheischler, J., Ren, Z., and Jeppesen, E.: Extreme and compound
704 events in lakes, *Nature Reviews Earth & Environment*, 6, 593-611,
705 <https://doi.org/10.1038/s43017-025-00710-w>, 2025. 删除[JIAHAO WEI]: [53]

706 ✓Wu, Z., Guo, X., Shi, J., Ding, X., Nakakuni, M., and Tada, K.: Seasonal variations and
707 controlling factors of nitrogen fluxes at the sediment-water interface in a semi-enclosed
708 inland sea, *EGUsphere*, 1-37, <https://doi.org/10.5194/egusphere-2025-6187>, 2025. 删除[JIAHAO WEI]: [54]

709 ✓Xiong, Z. Q., Guo, L. D., Zhang, Q. F., Liu, G. H., and Liu, W. Z.: Edaphic conditions
710 regulate denitrification directly and indirectly by altering denitrifier abundance in wetlands
711 along the Han River, China, *Environmental Science & Technology*, 51(10), 5483-5491,
712 <https://doi.org/10.1021/acs.est.6b06521>, 2017. 删除[JIAHAO WEI]: [55]

713 ✓Xu, Y., Sun, X. Y., and Tang, Q.: Human activity intensity of land surface: Concept, method
714 and application in China, *Journal of Geographical Sciences*, 26, 1349-1361,
715 <https://doi.org/10.1007/s11442-016-1331-y>, 2016. 删除[JIAHAO WEI]: [56]

716 ✓Yan, P. H., Zhuang, S. Z., Li, M. J., Zhang, J., Wu, S. B., Xie, H. J., and Wu, H. M.:
717 Combined environmental pressure induces unique assembly patterns of micro-plastisphere
718 biofilm microbial communities in constructed wetlands, *Water Research*, 260, 121958,
719 <https://doi.org/10.1016/j.watres.2024.121958>, 2024. 删除[JIAHAO WEI]: [57]

720 ✓Yang, C. H., Zhang, M., Zhang, C. C., Wang, G. X., Yin, H. B., Zhang, L. Y., and Liu, J. A.:
721 Release pattern of internal nutrients after sediment dredging driven by particulate particles
722 deposition in eutrophic lakes, *Water Research*, 289, 124879,
723 <https://doi.org/10.1016/j.watres.2025.124879>, 2026. 删除[JIAHAO WEI]: [58]

724 ✓Yang, L., Zou, Y. C., Jia, Z. J., Luo, Y., Qin, L., Jiang, M., and Yu, Z. C.: Iron reduction
725 promotes carbon mineralization and nutrient release of iron-associated organic matter in
726 anoxic environments, *Water Research*, 284, 124032,
727 <https://doi.org/10.1016/j.watres.2025.124032>, 2025. 删除[JIAHAO WEI]: [59]

728 ✓Yoon, S., Cruz-García, C., Sanford, R., Ritalahti, K. M., and Löffler, F. E.: Denitrification
729 versus respiratory ammonification: environmental controls of two competing dissimilatory
730 $\text{NO}_3^-/\text{NO}_2^-$ reduction pathways in *Shewanella loihica* strain PV-4, *The ISME Journal*, 9(5),
731 1093-1104, <https://doi.org/10.1038/ismej.2014.201>, 2015. 删除[JIAHAO WEI]: [60]

732 ✓Yuan, M. M., Guo, X., Wu, L. W., Zhang, Y., Xiao, N. J., Ning, D. L., Shi, Z., Zhou, X. S.,
733 Wu, L. Y., Yang, Y. F., Tiedje, J. M., and Zhou, J. Z.: Climate warming enhances microbial
734 network complexity and stability, *Nature Climate Change*, 11, 343-348,
735 <https://doi.org/10.1038/s41558-021-00989-9>, 2021. 删除[JIAHAO WEI]: [61]

736 ✓Zhou, J., Leavitt, P. R., Zhang, Y. B., and Qin, B. Q.: Anthropogenic eutrophication of shallow
737 lakes: Is it occasional? *Water Research*, 221, 118728,
738 <https://doi.org/10.1016/j.watres.2022.118728>, 2022. 删除[JIAHAO WEI]: [62]

739 Zhou, Y. W., Xu, X. G., Song, K., Yeerken, S., Deng, M., Li, L., Riya, S. H., Wang, Q. L., and
740 Terada, A.: Nonlinear pattern and algal dual-impact in N₂O emission with increasing
741 trophic levels in shallow lakes, Water Research, 203, 117489,
742 <https://doi.org/10.1016/j.watres.2021.117489>, 2021.
743 Zhu, G. B., Wang, S. Y., Wang, W. D., Wang, Y., Zhou, L. L., Jiang, B., Op den Camp, H. J.
744 M., Risgaard-Petersen, N., Schwark, L., Peng, Y. Z., Hefting, M. M., Jetten, M. S. M., and
745 Yin, C. Q.: Hotspots of anaerobic ammonium oxidation at land-freshwater interfaces,
746 Nature Geoscience, 6(2), 103-107, <https://doi.org/10.1038/Ngeo1683>, 2013.

删除[JIAHAO WEI]: [63]

删除[JIAHAO WEI]: [64]