

Spectral Signatures of Zenith Sky Radiance from Surface-Based Sky Radiometers: Implications for Clear- and Cloudy-Sky Detection

Pradeep Khatri¹, Hitoshi Irie², Hiroshi Kobayashi³, Isamu Morino⁴, Yoshitaka Jin⁴, Tadahiro Hayasaka⁵,
5 Hironobu Iwabuchi⁶, Tamio Takamura², and Yoshiki Takayama⁷

¹Department of Green Technology, Faculty of Science and Engineering, Soka University, Hachioji, Tokyo, Japan

²Center for Environmental Remote Sensing, Chiba University, Chiba, Japan

³Department of Environmental Sciences, Faculty of Life and Environmental Sciences, University of Yamanashi, Yamanashi, Japan

⁴National Institute for Environmental Studies, Tsukuba, Japan

⁵Graduate School of Science, Tohoku University, Sendai, Japan

⁶Center for Atmospheric and Oceanic Studies, Tohoku University, Sendai, Japan

⁷ Graduate School of Environment and Information Sciences, Yokohama National University, Yokohama, Japan

15 Correspondence to: Pradeep Khatri (pradeep@soka.ac.jp)

Abstract. Accurate identification of clear- and cloudy-sky conditions is essential for reliable aerosol and cloud retrievals from ground-based remote sensing observations. This study investigates the spectral characteristics of zenith sky radiance, measured by SKYNET sky radiometers and evaluates their potential for distinguishing clear- and cloudy-sky conditions. For a given wavelength pair, the spectral slope derived from the measured signals (m_{obs}) was separated into atmospheric (m_{atm}) and wavelength-dependent instrumental (m_{ins}) contributions. Radiative transfer simulations demonstrated that m_{atm} is strongly negative under cloud-free conditions and systematically shifts toward less negative values under cloudy conditions, providing a physical basis for m_{atm} -based sky-state discrimination. Based on these theoretical analyses, together with observational analyses of spectral zenith radiances and the dependence of m_{atm} on solar geometry, aerosol properties, and cloud properties for observation data at multiple sites representing diverse atmospheric environments, including urban, maritime, tropical continental, and polar regions, practical criteria using m_{atm} were developed to classify observations into clear, cloudy, and undetermined categories. All four wavelength pairs examined (0.400 – 0.675, 0.400 – 0.870, 0.500 – 0.675, and 0.500 – 0.870 μm) provided robust discrimination, with the 0.500 – 0.675 μm pair showing the best and most consistent overall performance across the observation sites. Independent validation using collocated direct normal irradiance measurements confirmed the physical consistency of the m_{atm} -based classification. Further comparison with an existing cloud-screening procedure showed good agreement in clear-sky aerosol data extracted using the newly developed method across diverse atmospheric environments. Overall, these results demonstrate that spectral zenith sky radiance can provide a simple and physically interpretable basis for clear- and cloudy-sky detection, provided that the wavelength-dependent instrumental contribution is appropriately characterized. Incorporating such spectral diagnostics into ground-based radiometric networks can improve cloud screening and enhance the reliability of long-term aerosol and cloud climatological analyses.

1 Introduction

Aerosols and clouds play fundamental roles in the Earth's radiation budget and environmental systems by regulating the balance between incoming solar radiation and outgoing terrestrial radiation, while simultaneously influencing atmospheric thermodynamic structure and hydrological processes (IPCC 2023). Through scattering and absorption of solar and terrestrial radiation, aerosols modify atmospheric heating rates and surface energy availability, whereas clouds exert strong radiative forcing by reflecting shortwave radiation and trapping longwave radiation. These processes directly affect atmospheric circulation, precipitation formation, and climate feedback mechanisms. Consequently, accurate characterization of aerosols and clouds is essential for understanding the Earth system and reducing uncertainties in climate projections.

Deleted:	s
Deleted:	based on wavelength dependence
Formatted:	Font: Times New Roman, 10 pt
Formatted:	Font: Times New Roman, 10 pt, Subscript
Formatted:	Font: Times New Roman, 10 pt
Formatted:	Font: Times New Roman, 10 pt, Subscript
Formatted:	Font: Times New Roman, 10 pt
Formatted:	Font: Times New Roman, 10 pt, Subscript
Formatted:	Font: Times New Roman, 10 pt
Deleted:	Analyses of normalized
Formatted:	Subscript
Formatted:	Font: Times New Roman, 10 pt
Formatted:	Font: Times New Roman, 10 pt
Formatted:	Subscript
Deleted:	.
Formatted:	Font: Times New Roman, 10 pt
Formatted:	Font: Times New Roman, 10 pt
Formatted:	Font: Times New Roman, 10 pt, Subscript
Formatted:	Font: Times New Roman, 10 pt
Deleted:	revealed systematic differences between clear- and cloudy-sky conditions identified using quality-controlled MODIS cloud mask products. The spectral slope derived from the logarithmic contrast between zenith radiances at 0.400 and 0.675 μm showed strong sensitivity to cloudy-sky conditions but limited capability for detecting clear-sky conditions. In contrast, slopes based on the 0.500 and 0.675 μm and 0.400 and 0.675 μm wavelength pairs exhibited more balanced detection performance for both clear- and cloudy-sky conditions. Based on detailed analyses of the dependence of these slope values on solar geometry, aerosol properties, and cloud properties, together with classification scores evaluated against MODIS cloud mask data, this study proposes a simple, efficient ... [1]
Deleted:	slope
Deleted:	the
Deleted:	standard
Formatted:	Subscript
Deleted:	very
Deleted:	criterion
Deleted:	s
Formatted:	Font: Times New Roman, 10 pt
Deleted:	.
Formatted:	Space Before: 0 pt, After: 0 pt

Advances in remote sensing technologies over the past decades have substantially improved the capability to observe atmospheric constituents from both satellite platforms and ground-based systems. Satellite observations provide extensive spatial coverage and enable global monitoring of aerosol and cloud distributions, while ground-based measurements offer high temporal resolution that are often difficult to obtain from spaceborne sensors. Within such observational frameworks, ground-based remote sensing instruments such as the SKYNET sky radiometer network (Nakajima et al., 2020) have proven particularly valuable. SKYNET sky radiometers provide high-quality spectral direct solar irradiance and diffuse sky radiance measurements that enable retrievals of key aerosol properties including aerosol optical thickness, single scattering albedo, and particle size distributions (Hashimoto et al. 2012; Kudo et al., 2021; Nakajima et al. 1996, 2020). In addition to aerosol observations, recent methodological developments have demonstrated that these instruments can also be used to retrieve cloud optical properties such as cloud optical depth and cloud particle effective radius from spectral zenith radiance measurements (Khatri et al. 2018, 2019, 2022). Because the measurements are obtained continuously with high temporal resolution, sky radiometers provide valuable long-term datasets that complement satellite retrievals and serve as high-quality reference observations for validation and climatological analyses.

Such datasets have been widely used in studies aimed at improving the understanding of aerosol–cloud interactions, evaluating satellite products, and examining regional and long-term atmospheric variability (e.g., Bi et al., 2014; Damiani et al., 2019; Irie et al., 2019; Khatri et al., 2010, 2014a, 2018, 2022; Ningombam et al., 2024; Wang et al., 2014). In addition, under clear-sky conditions, i.e., sky conditions in which the observations are unaffected by clouds, the sky radiometers can provide information on atmospheric gases such as water vapor (Campanelli et al., 2014) and ozone (Khatri et al. 2014b) along with aerosol properties. These capabilities demonstrate that sky radiometer observations constitute a versatile dataset for investigating multiple atmospheric components within a unified observational framework.

Given these diverse applications, reliable discrimination between clear-sky and cloudy-sky conditions is a fundamental prerequisite for maximizing the utility of sky radiometer observations. Many aerosol retrieval algorithms assume cloud-free conditions because clouds strongly perturb the radiative field and can introduce substantial biases in aerosol optical property retrievals. Similarly, atmospheric absorbing gases retrieval methods relying on spectral absorption features require uncontaminated clear-sky measurements to ensure accuracy. The sky-condition classification is therefore essential for maintaining data quality, improving retrieval reliability, and strengthening the credibility of satellite validation and model evaluation studies that rely on ground-based reference observations.

Despite its importance, a discrimination between clear and cloudy skies remains challenging in practical observations. Thin cirrus clouds, subvisual cloud layers, and rapidly evolving cloud fields can produce radiative signatures that resemble clear-sky conditions when evaluated using traditional screening approaches. In addition, atmospheric conditions characterized by high aerosol loading—such as mineral dust outbreaks, biomass-burning plumes, or pollution events—can significantly increase atmospheric turbidity. Under such conditions, irradiance variability or other statistical indicators used for cloud screening may incorrectly classify aerosol-rich but cloud-free skies as cloudy. These factors introduce uncertainties in sky-condition classification and can degrade the quality of derived atmospheric products.

Several cloud detection approaches have been proposed to address this problem. For example, the method developed by Khatri and Takamura (2009), uses the temporal variability of global irradiance, wavelength dependency behaviour of aerosols, and statistical approach to identify cloud contamination, while Song et al. (2014) and Choi et al. (2020) use somewhat different approaches. Although these approaches have proven useful in many applications, they may exhibit limitations under conditions characterized by complex aerosol loading or rapidly evolving cloud fields. Consequently, there remains a need for a simple physically interpretable method for discriminating between clear and cloudy sky conditions using readily available observational parameters.

To address these issues, it is necessary to investigate observational quantities that exhibit direct sensitivity to cloud scattering processes. Among them, spectral zenith radiance, measured by sky radiometers provide particularly valuable information for sky-condition discrimination. Zenith sky radiances are directly influenced by scattering from cloud particles within the instrument’s instantaneous field of view, making them sensitive indicators of cloud presence even when cloud optical thickness is relatively small (Khatri et al., 2018, 2019, 2022). Despite this strong physical sensitivity to cloud scattering, zenith radiance observations have not yet been fully exploited in operational cloud-screening procedures for sky radiometer datasets. This gap suggests that additional investigation into the potential of zenith radiances could lead to improved methods for sky-condition classification.

Deleted:

Deleted: .

Deleted: SKYNET

Deleted:

Deleted:

Deleted: s

Deleted: statistical

Deleted: s

Deleted: s

Deleted:

Deleted: Because cloud particles typically produce stronger and spectrally distinct scattering signals compared with aerosols, zenith radiance measurements can reveal subtle cloud contamination that may not be captured by irradiance-based screening methods or purely statistical approaches.

Deleted: Most existing screening algorithms rely primarily on irradiance variability or derived statistical indicators rather than directly utilizing the spectral information contained in zenith radiance measurements.

150 The present study aims to systematically evaluate the capability of spectral zenith radiance measurements for discriminating
between clear and cloudy sky conditions. Rather than retrieving aerosol or cloud properties, the proposed method provides an
efficient preprocessing or postprocessing screening procedure to support subsequent aerosol and cloud retrievals. In this
155 framework, sky-state detection is performed based on the spectral dependence of zenith-sky radiance after removing the
wavelength-dependent instrumental contribution, thereby eliminating the need for preliminary aerosol retrievals, temporal
variability analyses, or ancillary observations. Based on theoretical analyses, together with observational analyses at multiple
sites representing diverse atmospheric environments, including urban, maritime, tropical continental, and polar regions, the
study establishes simple criteria for sky-state classification. The proposed method is intended to complement existing cloud-
160 screening techniques for aerosol studies while remaining independent of the subsequent retrieval methodology. Furthermore,
the method has direct relevance to cloud retrieval approaches based on zenith-sky radiance measurements, such as that
demonstrated by Khatri et al. (2019), because cloud presence can be identified from the same measurements used for cloud
retrieval. Thus, the method can provide an initial screening step prior to cloud retrieval. Because the proposed method relies
on the physical characteristics of spectral zenith radiance rather than network-specific processing procedures, it also has the
potential to be extended to other observational networks, such as AERONET (Holben et al., 1998), that acquire comparable
measurements.

Formatted: Font: Times New Roman, 10 pt

2 Observation Sites and Data

This study analyses observation data from eight SKYNET sites, including Chiba, Cape Hedo, Fukue, Miyako, Sendai, Burgos,
Syowa, and Ny-Ålesund, which collectively cover a wide range of atmospheric conditions across mid-latitude, tropical, and
polar environments (Table 1). As summarized in Table 1, Chiba represents an urban-coastal polluted environment influenced
170 by anthropogenic emissions from Tokyo metropolitan emissions and marine air from the Pacific, with episodic long-range
transport of continental aerosols from East Asia (Xue et al., 2021). Cape Hedo is located on Okinawa Island and is often
considered a marine background site in the western Pacific, although it is periodically affected by long-range transport of Asian
dust and anthropogenic aerosols (Khatri et al., 2010). Fukue Island, situated in the East China Sea region, represents
a continental outflow receptor site that receives transported aerosols from East Asia mixed with marine air masses (Khatri
175 et al., 2014a). Similar to Hedo, Miyako represents a mid-latitude maritime environment with generally clean oceanic air masses
but episodic influence from long-range transport of continental aerosols (Dim et al., 2013). Sendai is a mid-latitude coastal
site influenced by regional urban emissions and seasonal continental transport from East Asia (Shiobara et al., 1991). Burgos
in the northern Philippines represents a tropical maritime environment dominated by Pacific trade winds with occasional
influence from transported aerosols from East and Southeast Asia (Bagtasa et al., 2018). In addition, two polar sites are
180 included: Syowa Station in Antarctica, representing a polar pristine environment with extremely low aerosol loading and
occasional long-range transport from midlatitudes and low latitudes (Hara et al., 2025); and Ny-Ålesund in the high Arctic,
which represents an Arctic background environment influenced by seasonal long-range transport of pollution known as Arctic
haze emitted from midlatitudes (Pasquier et al., 2022). The data analyses periods in this study vary among the sites depending
on data availability, extending from January 2010 up to December 2024 (Table 1).

185 The study utilized observational data from multiple complementary sources, including a sky radiometer (Model: POM-02,
PREDE Co., Ltd.), the Moderate Resolution Imaging Spectroradiometer (MODIS), and ground-based direct solar irradiance
measurements obtained using a pyrheliometer.

The POM-02 is a multifunctional ground-based radiometer widely used in the SKYNET observation network. It performs
190 automated measurements of direct solar irradiance, spectral sky radiance distributions (e.g., almucantar and principal-plane
scans), and zenith sky radiance at multiple wavelengths, enabling the retrieval of aerosol, cloud, and atmospheric optical
properties (Choi et al., 2020; Nakajima et al., 2020; Khatri et al., 2010, 2019). In the present study, different categories of
measurements were employed. First, unprocessed spectral zenith radiances at wavelengths with negligible absorption by
atmospheric gases —0.380, 0.400, 0.500, 0.675, 0.870, and 1.020 μm —were used. The direct and almucantar radiances
observed at these wavelengths are commonly used in SKYNET for aerosol observations. The zenith radiances observed at
195 these wavelengths provide valuable information about sky brightness and atmospheric scattering characteristics under both
clear- and cloudy-sky conditions. Second, quality-controlled aerosol optical thickness (AOT) values at wavelengths of 0.400,
0.500, and 0.870 μm were used. These values were processed using skyrad.pack inversion software (Hashimoto et al., 2012)
and cloud-screening algorithm (Khatri and Takamura, 2009) both of which are incorporated in SR-CERES (Sky Radiometer-
Center for Environmental Remote Sensing) processing package. These data analyzed using SR-CERES data processing

Deleted: By examining the spectral characteristics of zenith radiances under different atmospheric states, the study seeks to establish physically based criterion for identifying cloud contamination in sky radiometer observations. The resulting methodology can be applied either independently or in combination with existing cloud-screening techniques to enhance the reliability and quality of atmospheric retrievals derived from SKYNET measurements.

Beyond improving SKYNET data processing, the findings of this study may also provide useful insights for other ground-based aerosol monitoring networks. In particular, networks such as AERONET (Holben et al., 1998), which also rely on ground-based radiometric observations for aerosol characterization, may benefit from improved sky-condition screening approaches informed by the radiative behaviour of spectral measurements. By strengthening the ability to distinguish between aerosol-dominated and cloud-contaminated observations, such improvements would contribute to more accurate aerosol climatology, more reliable satellite validation efforts, and a deeper understanding of aerosol-cloud-radiation interactions in the Earth's atmosphere.

Formatted: Space Before: 0 pt, After: 0 pt

Deleted:

Formatted: Font: Times New Roman, 10 pt

Deleted: From the sky radiometer

Deleted: two

Formatted: Font: Times New Roman, Not Italic

Formatted: Font: Times New Roman, 10 pt

Deleted: 0.340,

Deleted: ¶

Deleted: are

230 package are publicly available through the SKYNET data archive (<https://skynet.irie-lab.jp>). The study further uses calibration constants values determined using Improved Langely method (Campanelli et al., 2004; Nakajima et al., 1996, 2000) at those wavelengths.

Deleted: <http://atmos3.cr.chiba-u.jp/skynet/>

230 **Table 1.** Observation locations, representative air mass types, and study periods

Study Location	Representative Air Mass Type	Study Period
Chiba (Japan) (35.625°N, 140.104°E)	Urban-coastal air mass influenced by Tokyo emissions, marine air, and episodic East Asian aerosol transport.	January, 2010 ~ December, 2024
Cape Hedo (26.867°N, 128.248°E)	Marine background air mass influenced by long-range transport from East Asia	February, 2010 ~ February, 2021
Fukue (Japan) (32.752°N, 128.682°E)	Continental outflow with marine mixing	January, 2010 ~ July, 2019
Miayko (Japan) (24.737°N, 125.327°E)	Maritime background air mass with episodic Asian dust and pollution transport	August, 2017 ~ November, 2020
Sendai (Japan) (38.26°N, 140.84°E)	Mid-latitude coastal air mass influenced by continental and marine sources	July, 2018 ~ December, 2024
Burgos (Philippines) (18.533°N, 120.650°E)	Maritime tropical air mass with occasional long-range transport	March, 2018 ~ December, 2024
Syowa (Antarctica) (-69.006S, 39.590°E)	Polar background (pristine) air mass with extremely low aerosol loading and occasional long-range transport.	February, 2009~December, 2021
Ny-Ålesund (Norway) (78.929°N, 11.861°E)	Arctic background air mass with seasonal long-range transport influence.	March, 2011~ September, 2018

235 Satellite-based cloud information and cloud optical properties were obtained from MODIS Level-2 cloud mask and cloud property products. Specifically, the MOD35_L2 product from the Terra satellite and the MYD35_L2 product from the Aqua satellite were used for cloud mask information, while cloud optical properties were obtained from the MOD06_L2 and MYD06_L2 products corresponding to Terra and Aqua satellites, respectively. The MODIS cloud mask products provide pixel-level cloud detection using a series of multispectral threshold tests that analyze radiances measured in visible, near-infrared, and thermal infrared channels. The cloud mask algorithm classifies each pixel into categories such as clear, probably clear, probably cloudy, and cloudy based on spectral characteristics and spatial consistency tests. The fundamental framework of this detection approach is described by Ackerman et al., (1998). The MODIS cloud optical property products provide retrieved cloud optical thickness, cloud effective radius, and related cloud microphysical parameters using multispectral reflectance measurements, primarily from visible, near-infrared, and shortwave-infrared bands. These retrievals are based on the bispectral method, in which cloud optical thickness is mainly constrained by visible-band reflectance, while cloud particle effective radius is inferred from absorbing near-infrared or shortwave-infrared channels (Platnick et al., 2017; Nakajima and King, 1990). Both cloud mask and cloud property products have a nominal spatial resolution of approximately 1 km at nadir.

240 In addition to the sky radiometer and satellite observations, direct normal irradiance (DNI) measurements were also utilized where available. These measurements were obtained using a pyrheliometer manufactured by Kipp & Zonen, which measures the solar irradiance received directly from the sun within a narrow field of view aligned with the solar disk. Pyrheliometers provide highly accurate measurements of DNI and are widely used in atmospheric radiation studies as well as solar energy applications. DNI observations are particularly useful for identifying clear-sky conditions and for assessing the influence of clouds and aerosols on direct solar radiation reaching the Earth's surface.

255 3 Methodology

To investigate the spectral characteristics of zenith sky radiances under clear- and cloudy-sky conditions, sky radiometer-observed zenith radiance measurements were temporally collocated with MODIS overpass times by selecting observations

Deleted: ¶

- Deleted:
- Deleted:
- Deleted:
- Deleted:
- Deleted:
- Deleted:
- Deleted:
- Deleted:
- Deleted:
- Deleted:
- Deleted: cloud

with an absolute time difference of less than 5 minutes, noting that zenith sky radiances are measured at time interval of 10 minutes. Based on this temporal collocation, the corresponding spectral zenith radiances were categorized into two groups: confidently clear and confidently cloudy conditions as identified by the MODIS cloud mask product. The MODIS cloud mask data used in this study were quality-controlled using strict spatial and quality-assurance criteria. For each ground-based observation, the nine MODIS pixels closest to the observation site, corresponding to a 3×3 pixel area centered the observation site, were extracted. This criterion was adopted to ensure that the collocated satellite observations represented a uniformly cloudy scene. Only cases for which all nine closest pixels were available and the minimum distance between the observation site and the nearest MODIS pixel was less than 1 km were retained. A scene was classified as confidently clear only when all nine pixels were identified as confident clear with a MODIS quality-assurance level of 6 or higher. Similarly, a scene was classified as confidently cloudy only when all nine pixels were identified as cloudy with a MODIS quality-assurance level of 6 or higher. This strict screening criterion can substantially reduce uncertainties associated with cloud-edge contamination, geolocation mismatch, and mixed clear-cloudy scenes. However, some uncertainties may still remain, particularly under broken-cloud conditions, where the spatial distribution of clouds can change rapidly within the MODIS pixel area and during the short time interval between satellite overpass and ground-based observation. Therefore, although the applied criteria may provide a basis for separating clear and cloudy cases as rigorously as possible, possible residual mismatches between the satellite-derived cloud mask and the actual sky condition observed by the sky radiometer cannot be completely eliminated. For the zenith sky radiance signals $S(\lambda_1)$ and $S(\lambda_2)$ observed at any two wavelengths λ_1 and λ_2 , the slope value (m_{obs}) can be expressed using the following equation.

$$m_{\text{obs}}(\lambda_1, \lambda_2) = \frac{\ln[S(\lambda_1)/S(\lambda_2)]}{\ln(\lambda_1/\lambda_2)}. \quad (1)$$

The measured zenith sky radiance can be further represented as

$$S(\lambda) = C(\lambda)L(\lambda), \quad (2)$$

where $L(\lambda)$ is the band-averaged atmospheric zenith radiance and $C(\lambda)$ is the wavelength-dependent relative instrumental response. Substituting Eq. (2) into Eq. (1), we can write

$$m_{\text{obs}}(\lambda_1, \lambda_2) = \frac{\ln\left[\frac{C(\lambda_1)L(\lambda_1)}{C(\lambda_2)L(\lambda_2)}\right]}{\ln(\lambda_1/\lambda_2)} \quad (3)$$

Using $\ln(ab) = \ln(a) + \ln(b)$,

$$m_{\text{obs}}(\lambda_1, \lambda_2) = \frac{\ln\left[\frac{C(\lambda_1)}{C(\lambda_2)}\right]}{\ln(\lambda_1/\lambda_2)} + \frac{\ln\left[\frac{L(\lambda_1)}{L(\lambda_2)}\right]}{\ln(\lambda_1/\lambda_2)} \quad (4)$$

Thus,

$$m_{\text{obs}}(\lambda_1, \lambda_2) = m_{\text{atm}}(\lambda_1, \lambda_2) + m_{\text{inst}}(\lambda_1, \lambda_2), \quad (5)$$

where,

$$m_{\text{atm}}(\lambda_1, \lambda_2) = \frac{\ln\left[\frac{L(\lambda_1)}{L(\lambda_2)}\right]}{\ln(\lambda_1/\lambda_2)}, \quad (6)$$

and

$$m_{\text{inst}}(\lambda_1, \lambda_2) = \frac{\ln\left[\frac{C(\lambda_1)}{C(\lambda_2)}\right]}{\ln(\lambda_1/\lambda_2)}. \quad (7)$$

Deleted: cantered

Formatted: Font: 10 pt

Deleted: For each observation time, the radiance spectrum was normalized by its maximum value to reduce the influence of absolute radiance magnitude and to emphasize the spectral shape of the zenith radiance. The normalized radiance spectra were then analysed separately for the clear-sky and cloudy-sky groups. In particular, the median values of the normalized radiance spectra at each wavelength were calculated in order to characterize the typical spectral features associated with clear and cloudy atmospheric conditions.[¶] To further quantify the spectral dependence of zenith radiances

Deleted: , slope values were

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Subscript

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Subscript

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman, Subscript

Formatted: Font: (Default) Times New Roman

Deleted: calculated for all possible wavelength pairs for each observation time

Deleted: ¶

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Space Before: 0 pt, After: 0 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Space Before: 0 pt, After: 0 pt

Formatted: Font: 10 pt, Font color: Auto

Formatted: Justified, Space Before: 0 pt, After: 0 pt

330

Formatted: Justified

Formatted: Font: Times New Roman, 10 pt

Formatted: Subscript

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

Formatted: Font: Cambria Math

335

Formatted: Left

Formatted

Formatted: Subscript

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: 10 pt, Font color: Auto

Deleted: ¶

... [2]

Formatted: Subscript

Deleted: presence of

Deleted: ¶

Deleted: 2

Similarly, the extra-terrestrial instrument signal F_0 (i.e calibration constant) at wavelength λ can be expressed as

$$F_0(\lambda) = C(\lambda)E_0(\lambda). \quad (8)$$

where $E_0(\lambda)$ is band-averaged extra-terrestrial solar irradiance at wavelength λ . For the two wavelengths λ_1 and λ_2 ,

$$\frac{C(\lambda_1)}{C(\lambda_2)} = \frac{F_0(\lambda_1) E_0(\lambda_2)}{F_0(\lambda_2) E_0(\lambda_1)} \quad (9)$$

Consequently, from Eqs. (7) and (9),

$$m_{\text{inst}}(\lambda_1, \lambda_2) = \frac{\ln \left[\frac{F_0(\lambda_1) E_0(\lambda_2)}{F_0(\lambda_2) E_0(\lambda_1)} \right]}{\ln(\lambda_1 / \lambda_2)}. \quad (10)$$

Since $m_{\text{atm}}(\lambda_1, \lambda_2) = m_{\text{obs}}(\lambda_1, \lambda_2) - m_{\text{inst}}(\lambda_1, \lambda_2)$, from Eqs. (5), and (10) we can write,

$$m_{\text{atm}}(\lambda_1, \lambda_2) = \frac{\ln[S(\lambda_1)/S(\lambda_2)]}{\ln(\lambda_1/\lambda_2)} - \frac{\ln \left[\frac{F_0(\lambda_1) E_0(\lambda_2)}{F_0(\lambda_2) E_0(\lambda_1)} \right]}{\ln(\lambda_1/\lambda_2)} \quad (11)$$

Here, m_{atm} represents the atmospheric contribution arising from molecular, aerosol, and cloud scattering together with atmospheric absorption, and m_{inst} represents the wavelength-dependent instrumental contribution. To calculate m_{atm} , $F_0(\lambda)$ and $E_0(\lambda)$ are also required along with instrument-observed $S(\lambda)$ values. In this study, $F_0(\lambda)$ values determined from the Improved Langely method and $E_0(\lambda)$ values estimated by integrating the extraterrestrial solar spectrum of Gueymard (2004) over the spectral response function of each sky-radiometer channel were used.


The resulting correction should be regarded as a first-order estimate of the wavelength-dependent instrumental contribution, because a single set of median spectral calibration constants was applied to each instrument over the corresponding analysis period. Nevertheless, a key advantage of this approach is that, even with such a first-order approximation, the corrected atmospheric spectral slope, m_{atm} , retains sufficient sensitivity to distinguish between cloud-free and cloud-contaminated conditions, as demonstrated and discussed in the preceding Results and Discussion.

By analyzing the slope values derived from Eq. (11) for the grouped clear- and cloudy-sky datasets, threshold-based criterion was established to discriminate between clear and cloudy conditions. In addition, the collocated aerosol data and MODIS Level-2 cloud property data were statistically analyzed to examine the behavior of these m_{atm} values under varying atmospheric conditions while developing criteria for detecting sky condition.

The resulting classification of clear and cloudy sky conditions was evaluated using collocated direct normal irradiance (DNI) measurements, applying a temporal matching criterion of less than 2.5 minutes. DNI is highly sensitive to the clouds located along the direct solar beam, particularly those near the solar disk; under clear-sky conditions, DNI typically exhibits relatively high values because direct solar radiation reaches the surface without obstruction. In contrast, the presence of clouds significantly attenuates the direct solar beam, leading to substantially reduced DNI values. Therefore, DNI measurements provide an independent observational reference for assessing the reliability and robustness of the proposed classification approach. Additional validation was conducted by comparing aerosol optical thickness (AOT) at 0.5 μm and the Ångström exponent (AE) obtained for clear-sky cases identified by the proposed slope-based method with those derived using the cloud-screening algorithm of Khatri and Takamura (2009) implemented in SR-CERES data processing package. The Ångström exponent, which characterizes the spectral dependence of aerosol optical thickness and is commonly used as an indicator of aerosol particle size, was calculated as

$$AE = -\ln(AOT_{0.40} / AOT_{0.87}) / \ln(0.400 / 0.870), \quad (12)$$

385 where $AOT_{0.40}$ and $AOT_{0.87}$ represent aerosol optical thickness at wavelengths of 0.400 μm and 0.870 μm , respectively. This comparison provides an additional consistency check by examining whether the aerosol optical properties retrieved under the clear-sky conditions identified by the proposed slope-based method are physically reasonable and consistent with those obtained using an established cloud-screening algorithm.



390 **Figure 1.** Atmospheric spectral slope (m_{atm}) as a function of solar zenith angle (SZA) simulated for four selected wavelength pairs: (a) 0.400–0.500 μm , (b) 0.400–0.870 μm , (c) 0.500–0.675 μm , and (d) 0.500–0.870 μm . Simulations were performed with mid-latitude summer atmospheric model, precipitable water vapor of 1.0 cm, total column ozone of 0.30 cm-atm, Lambertian surface albedo of 0.10, and clouds represented as plane-parallel layers.

395 To provide the theoretical background for the applicability of atmospheric spectral slope (m_{atm}) for detecting sky conditions, radiative transfer calculations were performed using the Santa Barbara DISORT Atmospheric Radiative Transfer (SBDART) model (Ricchiazzi et al., 1998). The calculations used the atmospheric profile specified by mid-latitude summer atmospheric model, with precipitable water vapor and total column ozone set to 1.0 cm and 0.30 cm-atm, respectively. The surface was assumed to be Lambertian with an albedo of 0.10, and clouds were represented as plane-parallel layers. Simulations were performed for solar zenith angles (SZAs) ranging from 10° to 80°. Four atmospheric conditions were considered: a clean atmosphere with AOT at 0.55 μm of 0.1, a polluted atmosphere with AOT at 0.55 μm of 0.5, a thin-cloud atmosphere with

400

AOT at 0.55 μm of 0.1 and COT at 0.55 μm of 2, and a thick-cloud atmosphere with AOT at 0.55 μm of 0.1 and COT at 0.55 μm of 20. The simulated zenith-sky radiances were subsequently used to calculate m_{atm} for the selected wavelength pairs 0.400–0.675, 0.400–0.870, 0.500–0.675, and 0.500–0.870 μm . In the calculation, the spectral dependency of AOT was represented by assuming AEt value for 1.2. Similarly, the effective radius of cloud-droplet particles was set to 10 μm . Figure 1 shows the SZA dependence of m_{atm} under these atmospheric conditions. A distinct difference in m_{atm} is evident between the cloud-free and cloudy atmospheres for all four wavelength pairs. Under cloud-free conditions, m_{atm} is strongly negative, particularly for the clean atmosphere, reflecting the stronger wavelength dependence of molecular and aerosol scattering and the resulting decrease in zenith-sky radiance toward longer wavelengths. Increasing the aerosol loading from AOT at 0.55 μm to 0.50 shifts m_{atm} toward less negative values, indicating a weakening of the spectral dependence of zenith-sky radiance. Nevertheless, the polluted cloud-free atmosphere remains clearly separated from the cloudy cases over the investigated SZA range. In contrast, the presence of clouds produces substantially less negative m_{atm} values. This behavior is consistent with the weaker wavelength dependence of scattering by cloud droplets over the visible and near-infrared spectral region and the enhanced role of multiple scattering within clouds. In particular, the thick-cloud case exhibits relatively weak spectral dependence, whereas the thin-cloud case shows a stronger dependence on SZA; for the 0.400–0.675 μm pair, m_{atm} is positive at smaller SZAs and becomes negative as SZA increases. These results provide a physical basis for the applicability of m_{atm} to discriminate between clear- and cloudy-sky conditions. Although the magnitude and SZA dependence of m_{atm} vary with aerosol loading, cloud optical thickness, and wavelength pair, the transition from molecular- and aerosol-dominated scattering under cloud-free conditions to cloud-dominated multiple scattering systematically modifies m_{atm} toward less negative values. Importantly, increasing AOT alone does not reproduce the spectral behavior obtained under cloudy conditions, as demonstrated by the separation between the polluted atmosphere with AOT at 0.55 μm of 0.50 and the simulated cloudy cases. Thus, m_{atm} contains physically meaningful information on changes in the atmospheric scattering regime and provides a theoretical basis for discriminating cloud-contaminated from cloud-free zenith-sky observations.

4 Results and Discussion

4.1 Spectral Signatures of Zenith Radiance Under Clear and Cloudy Skies

4.1.1 General Features





Figure 2. Median normalized spectral zenith sky radiance under cloudy and confidently clear sky conditions at (a) Chiba, (b) Hedo, (c) Fukue, (d) Miyako, (e) Sendai, (f) Burgos, (g) Syowa, and (h) Ny-Ålesund. Zenith sky radiances were normalized

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Deleted: ¶



Deleted:

Deleted: 1

by the maximum value within each observation and averaged separately for MODIS-identified cloudy and confidently clear cases.

Figure 2 presents the median normalized spectral zenith radiances under confidently clear and cloudy sky conditions at eight selected SKYNET sites. The measured zenith sky radiances were first converted into physical zenith-sky radiances using $C(\lambda)$ values, which were determined from Eq. (8). For each observation, the radiance spectrum was normalized by its maximum value before calculating the median spectrum. Therefore, the figure emphasizes differences in spectral shape and relative spectral distribution rather than absolute radiance magnitude.

Across all sites, the spectral curves under cloudy and clear conditions exhibit systematically different patterns. Under cloudy conditions, the normalized spectra appear relatively weaker spectral dependence across the visible and near-infrared wavelength range in comparison to those for clear-sky spectra. This systematic shift is consistently observed across the diverse environments represented by the eight SKYNET sites, including maritime, continental, tropical, and polar locations. The consistency of this behavior suggests that the observed spectral differences are governed primarily by fundamental atmospheric scattering processes rather than by site-specific characteristics.

The physical basis for this behavior can be explained by the dominant scattering mechanisms in clear and cloudy atmospheres. Under clear-sky conditions, zenith radiance is mainly controlled by molecular (Rayleigh) scattering and aerosol scattering. Rayleigh scattering exhibits a strong inverse dependence on wavelength (approximately proportional to λ^{-4}), resulting in enhanced scattering at shorter wavelengths in the blue portion of the spectrum. Aerosol scattering also exhibits wavelength dependence, though typically weaker than Rayleigh scattering. In the near-ultraviolet to blue spectral region, aerosol scattering can be typically several times smaller than Rayleigh scattering under low-to-moderate aerosol loading, although its relative contribution increases under polluted or dust-influenced conditions. As a result, clear-sky zenith radiance spectra tend to show relatively higher normalized radiance at shorter visible wavelengths and a gradual decrease toward longer wavelengths. In contrast, under cloudy conditions the scattering of solar radiation is dominated by cloud droplets. The typical size of cloud droplets (on the order of several micrometers) is significantly larger than the wavelengths of visible light. Under these circumstances, scattering is primarily governed by Mie scattering, which exhibits much weaker wavelength dependence. Consequently, the scattered radiation becomes spectrally more uniform, producing relatively weaker normalized radiance spectra compared with clear-sky conditions. These characteristics are consistent with theoretical descriptions of cloud scattering behavior (Bohren and Huffman, 2007; Hansen and Travis, 1974).

Although the overall spectral behavior is consistent among the sites, minor variations in spectral shape are observed. These differences likely reflect variations in aerosol loading, aerosol type, solar geometry, and surface reflectance conditions. For example, sites influenced by maritime air masses or high aerosol concentrations may exhibit slightly modified spectral gradients due to changes in aerosol optical properties. Nevertheless, the fundamental contrast between clear- and cloudy-sky spectra remains evident across all environments.

4.1.2 Spectral Slopes

To better understand the spectral dependence of zenith sky radiances on wavelengths during clear- and cloudy-sky conditions, we analyzed the atmospheric spectral slope (m_{atm}) values for all possible wavelength pairs, as given by Eq. (11). Figure 3 presents box-and-whisker distributions of slope values calculated from zenith radiance pairs for cloudy and confidently clear sky conditions at the Chiba site. Although results from other sites are not shown here for brevity, they exhibit slope patterns similar to those observed for Chiba. For shorter first wavelength λ_1 of 0.38 μm , the slope values are predominantly positive and relatively large. This behavior reflects the strong wavelength dependence of molecular (Rayleigh) scattering and aerosol scattering in the shorter spectral region. Although cloudy conditions generally exhibit larger slopes than clear skies for this first wavelength λ_1 of 0.38 μm , most of the distributions partially overlap. As the first wavelength shifts toward the visible and near-infrared region, the slope magnitude decreases and, for several wavelength combinations, become negative. In particular, the wavelength pairs of 0.400 and 0.675 μm , 0.400 and 0.870 μm , 0.500 and 0.675 μm , and 0.500 and 0.870 μm show clear separation between the two sky conditions, with minimal overlaps. Other wavelength pairs, such as 0.675 and 0.870 μm and 0.870 and 1.020 μm , show smaller differences in slope values between clear- and cloudy-sky conditions. Overall, the results demonstrate that wavelength pairs connecting the visible and near-infrared spectral regions provide the greatest sensitivity to changes in atmospheric scattering associated with cloud presence. These wavelength combinations therefore represent the

- Deleted: 1
- Formatted: Line spacing: At least 0 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Deleted: flat
- Deleted: (approximately 0.400–0.870 μm), indicating weak spectral dependence within this region. In contrast,
- Deleted: show stronger wavelength-dependent variation, characterized by a more pronounced decrease toward longer wavelengths
- Deleted: More specifically, under clear-sky conditions the (... [3])
- Deleted: across the visible spectrum
- Deleted: flatter
- Deleted: Because shorter wavelengths are relatively less en (... [4])
- Deleted:
- Deleted: 2
- Deleted: slope
- Deleted: s
- Deleted: (
- Deleted: =
- Deleted: 0.34–
- Deleted:)
- Deleted: the ultraviolet and blue
- Deleted: s
- Formatted: Font: Times New Roman, 10 pt
- Deleted: some separation between cloudy and clear conditic (... [5])
- Deleted: the
- Deleted: at these shorter wavelength regions
- Deleted: The most distinct contrast between clear- and cloud (... [6])
- Deleted: 0.400 and 0.870 μm
- Deleted: predominantly negative slope values under clear sk (... [7])
- Deleted: 500
- Deleted: 675
- Deleted: 0.870
- Deleted: , also indicate
- Deleted: ; however, the boundary values separating the two (... [8])
- Formatted: Font: Times New Roman, 10 pt

540 [most suitable candidates for clear- and cloudy-sky discrimination based solely on the atmospheric spectral characteristics of zenith-sky radiance.](#)





Figure 3. Box-and-whisker distributions of spectral slope values of atmospheric slope (m_{atm}) for cloudy and confidently clear sky conditions for Chiba. Each row corresponds to a fixed first wavelength (λ_1), while the x-axis represents the second



Deleted:

Deleted: 2

Deleted: calculated from zenith radiance pairs

550


wavelength (λ_2). In the figure, the boxes represent the 25th and 75th percentiles, and the whiskers represent the 5th and 95th percentiles.

4.1.2.1 Dependence of Spectral Slope Values of Key Wavelength Pairs on Solar Geometry, Aerosol, and Cloud Conditions

555

As the four wavelength pairs—0.400 and 0.675 μm , 0.500 and 0.675 μm , 0.400 and 0.870 μm , and 0.500 and 0.870 μm —exhibited relatively stronger discrimination between clear- and cloudy-sky conditions, a more detailed analysis was performed to understand the dependencies of m_{atm} values corresponding to those wavelength pairs on solar position and aerosol and cloud optical properties. Specifically, we analyzed the dependence of m_{atm} on solar zenith angle (SZA), AOT, and AE under clear sky conditions identified using the reference MODIS cloud mask data. Similarly, for cloudy-sky conditions, we investigated the relationships between m_{atm} corresponding to those wavelength pairs and SZA, COT, and CER. These analyses are shown in Figures 4 and 5. In those figures, only the mean values are presented because the objective is to illustrate the dependence of m_{atm} values on the atmospheric parameters over different atmospheric conditions together.

560



565

Figure 4. Dependence of mean spectral atmospheric slope (m_{atm}) values on solar zenith angle (SZA), aerosol optical thickness (AOT), and Ångström exponent (AE) under clear-sky conditions identified using the reference MODIS cloud mask data. Mean values were calculated at each site using bin intervals of 2° for SZA, 0.1 for AOT, and 0.1 for the AE.

Deleted: ¶

Deleted: three

Deleted: and

Deleted: show predominant negative and positive slope values for

Deleted: , respectively

Deleted: their

Formatted: Subscript

Deleted:

Deleted: the spectral slope

Deleted: the spectral slope

Deleted: 3

Deleted: 4

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Deleted: ¶

... [9]

Deleted: 3

Deleted: slope

Deleted: Ångström exponent




Figure 5. Dependence of mean spectral atmospheric slope (m_{atm}) values on solar zenith angle (SA), cloud optical thickness (COT), and cloud-particle effective radius (CER) under cloudy-sky conditions identified using the reference MODIS cloud mask data. Mean values were calculated at each site using bin intervals of 2° for SA, 2 for COT, and $2 \mu\text{m}$ for CER.

Under clear-sky conditions, m_{atm} values generally decrease with increasing SA. This behavior indicates that atmospheric spectral contrast becomes stronger as the atmospheric optical path length increases. At larger SA, enhanced Rayleigh and aerosol scattering along the longer slant path modifies the shortwave spectral distribution, leading to more negative slope values at most sites. This tendency is especially evident at mid- and low-latitude sites, whereas polar sites show weaker or somewhat different behavior, likely due to differences in aerosol loading, surface reflectance, and illumination geometry. The dependence on AOT differs from the SA dependence. For all wavelength pairs, one can note the increase of m_{atm} value with the increase of AOT. Particularly, the 0.500 and 0.675 μm and 0.500 and 0.870 μm pairs remain predominantly negative even as AOT increases, although the value becomes less negative under enhanced aerosol loading. The relationship between m_{atm} and AE is weaker and more site-dependent than the relationship with AOT. In some cases, m_{atm} becomes slightly more negative with increasing AE, but the pattern is not consistently monotonic across sites. This suggests that aerosol amount exerts a stronger first-order control on m_{atm} magnitude than aerosol size information alone. The observed site-to-site variability may be related to differences in aerosol type, surface albedo, humidity-driven aerosol growth, and the limited sampling range of AE at some sites.

Under cloudy conditions, m_{atm} values increase substantially relative to clear-sky conditions for all examined wavelength combinations, clearly contrasting with the predominantly negative slopes observed under clear skies. Among the tested pairs, the 0.400 and 0.675 μm pair generally shows the largest m_{atm} values followed by 0.400 and 0.870 μm , 0.500 and 0.870 μm , and 0.500 and 0.675 μm . Although latter wavelength pairs often retain negative values under clouds, they consistently shift toward larger (less negative) values under cloudy conditions compared with clear-sky conditions. This contrast is physically meaningful as cloud scattering weakens the spectral distribution of zenith radiance. The dependence of cloudy-sky

Deleted: 1 ... [10]

Deleted: 4

Deleted: slope

Deleted:

Deleted: the mean slope

Deleted: s

Deleted: the

Deleted: of zenith radiance

Deleted: slope

Deleted: s

Deleted: slope

Deleted: This result indicates that the 0.500 and 0.675 μm ... [11]

Deleted: spectral slope

Deleted: Ångström exponent

Deleted: the slope

Deleted: Ångström exponent

Deleted: the

Deleted: slope

Deleted: c

Deleted: Ångström exponent

Deleted: the spectral slope

Deleted: s

Deleted: are predominantly positive

Formatted: Font: Times New Roman, 10 pt

Deleted: mostly

Deleted: positive

Deleted: slope

Deleted: , indicating strong sensitivity to cloudy-sky conditions

Deleted: The 0.500 and 0.675 μm pair

Deleted: also shows positive

Formatted: Font: Times New Roman, 10 pt

Deleted: although the magnitude is generally smaller and r ... [12]

Deleted: modifies

Deleted: in a way that reverses the sign of the slope relative ... [13]

slope on SZA is weaker than that observed under clear-sky conditions. Although some site-to-site variability is present, m_{atm} generally remains positive or relatively weak negative across most SZA ranges. This indicates that cloud scattering dominates the spectral behavior of zenith radiance, reducing the relative influence of solar geometry compared with clear-sky cases. The m_{atm} also varies with COT, but the relationship is generally nonlinear. At low COT, the m_{atm} may change rapidly, indicating strong sensitivity to the transition from optically thin to moderately thick clouds. As COT increases, m_{atm} tends to become more stable, suggesting that the zenith radiance spectrum approaches a saturated cloudy-sky regime. This is consistent with the expectation that, for optically thick clouds, additional increases in cloud optical thickness have a progressively smaller effect on radiance-based spectral signatures. The dependence on CER is weaker and more variable than the dependence on COT. For many sites, especially for water clouds, the mean slope remains relatively stable across the CER range. Ice cloud cases show larger variability, likely because ice cloud retrievals are more heterogeneous and are affected by particle habit, vertical structure, and possible mixed-phase contamination. Overall, the results demonstrate that m_{atm} is governed by different dominant physical factors under clear- and cloudy-sky conditions. Under clear skies, m_{atm} is mainly influenced by solar geometry and aerosol optical loading, while under cloudy skies, it is primarily controlled by cloud scattering and COT.

Deleted: the slope

4.1.2.2 Possibility of Clear- and Cloudy-Sky Discrimination

Accounting for the clear separation of m_{atm} values between clear- and cloudy-sky conditions discussed above, practical operational criteria were developed for the purpose of sky state detection using m_{atm} values. For this purpose, $m_{\text{atm}}(\lambda_1, \lambda_2)$ values corresponding to each wavelength pair from all observation sites were pooled using the reference MODIS cloud mask labels, and the discrimination performance was evaluated over the full range of observed $m_{\text{atm}}(\lambda_1, \lambda_2)$ values using receiver operating characteristic (ROC) analysis.

Deleted: the spectral slope

Deleted: ¶

For a given decision threshold, the sensitivity (a measure of true positive rate) and specificity (a measure of true negative rate) were calculated as

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: 10 pt

Formatted: Justified

$$\text{Sensitivity}(T) = \frac{TP(T)}{TP(T) + FN(T)} \quad (13)$$

$$\text{Specificity}(T) = \frac{TN(T)}{TN(T) + FP(T)} \quad (14)$$

where TP, TN, FP, and FN denote the numbers of true positives, true negatives, false positives, and false negatives, respectively. They are counts of classification outcomes, and are dimensionless. The classification performance at each decision threshold was evaluated using the Youden index,

Formatted: Justified

Formatted: Font: Times New Roman, 10 pt

Formatted: Justified

$$J(T) = \text{Sensitivity}(T) + \text{Specificity}(T) - 1. \quad (15)$$

The decision threshold corresponding to the maximum value of $J(T)$ was selected as the optimum binary decision boundary, T_y , between clear- and cloudy-sky conditions. The ROC curve describes the relationship between sensitivity and specificity over all possible decision thresholds, while the area under the ROC curve (ROC AUC) provides an overall measure of the discrimination capability of each wavelength pair, with a larger value closer to 1 indicating better separation between clear- and cloudy-sky conditions. Further, to avoid forcing observations located close to T_y into either class, an intermediate "undetermined" category was introduced by defining a symmetric uncertainly range ($\pm \varepsilon$) around T_y . Hence, the final sky state detection criterion can be defined as

Formatted: Font: 10 pt

Formatted: Justified, Space Before: 0 pt, After: 0 pt

Formatted: Font: 10 pt

Clear sky: $m_{\text{atm}}(\lambda_1, \lambda_2) < T_y(\lambda_1, \lambda_2) - \varepsilon$, (16)

Cloudy sky: $m_{\text{atm}}(\lambda_1, \lambda_2) > T_y(\lambda_1, \lambda_2) + \varepsilon$, (17)

Undetermined: $T_y(\lambda_1, \lambda_2) - \varepsilon \leq m_{\text{atm}}(\lambda_1, \lambda_2) \leq T_y(\lambda_1, \lambda_2) + \varepsilon$ (18)

Formatted: Subscript

Formatted: Font: 10 pt, English (UK)

Formatted: Justified, Space Before: 0 pt, After: 0 pt

Formatted: Subscript

Formatted: Font: Times New Roman, 10 pt

where λ_1 and λ_2 represent the two wavelengths of a wavelength pair. Table 2 summarizes T_y , ROC AUC, specificity and sensitivity values for four selected wavelength pairs. Here, the parameter ε controls the degree of conservativeness of the

classification. A smaller ϵ results in a narrower undetermined region and allows more observations to be classified as clear or cloudy, whereas a larger ϵ broadens the undetermined region and provides a more conservative classification. The value of ϵ may be adjusted according to the desired balance between classification coverage and confidence or to meet specific operational requirements.

Table 2. Optimum binary decision thresholds (T_c) and corresponding discrimination performance of the atmospheric spectral slope (α_{atm}) for the four selected wavelength pairs

Wavelength pair (μm)	T_c	ROC AUC	Sensitivity	Specificity
0.400 and 0.675	-0.56	0.981	0.946	0.967
0.500 and 0.675	-1.73	0.984	0.961	0.954
0.400 and 0.870	-1.03	0.981	0.951	0.956
0.500 and 0.870	-1.78	0.978	0.954	0.954

Table 2 shows that all four wavelength pairs provide strong discrimination between clear- and cloudy-sky conditions, with ROC AUC values ranging from 0.978 to 0.984. Sensitivity and specificity are also consistently high, ranging from 0.946 to 0.961 and from 0.954 to 0.967, respectively, indicating that the spectral slopes of those wavelength pairs can effectively identify both cloudy- and clear-sky conditions. Among the examined wavelength pairs, 0.500–0.675 μm shows the highest overall discrimination capability, with the largest ROC AUC (0.984) and sensitivity (0.961), while maintaining a high specificity (0.954). The 0.400–0.675 and 0.400–0.870 μm pairs show similarly strong performance. Although the 0.500–0.870 μm pair has the lowest ROC AUC (0.978), its sensitivity and specificity remain high (both 0.954), indicating that it also provides effective clear–cloud discrimination. Overall, the small differences in ROC AUC among the four wavelength pairs indicate that all four spectral combinations have substantial capability for sky-state discrimination, although the 0.500–0.675 μm pair provides the strongest individual criterion.

To further evaluate whether the criteria derived from the pooled observations described above are applicable across different atmospheric environments, T_c for each wavelength pair was applied independently to the eight observation sites to classify observation data into three classes (clear-sky, cloudy-sky, and undetermined) using ϵ value of 0.05. The resulting confusion matrices for classification for each wavelength pair are shown in Figures 6–9.

For the 0.400 and 0.675 μm wavelength pair (Figure 6), clear-sky agreement is consistently high across all sites, ranging from 92.2% at Hedo to 100% at Miyako and Burgos. Cloudy-sky agreement is also high at Chiba, Fukue, Hedo, Miyako, and Ny-Ålesund, where it ranges from 94.4% to 96.8%. Somewhat lower cloudy-sky agreement is observed at Sendai (81.6%), Syowa (84.0%), and Burgos (83.7%). For the 0.500 and 0.675 μm wavelength pair (Figure 7), clear-sky agreement ranges from 91.6% at Hedo to 100% at Miyako and Burgos, while cloudy-sky agreement ranges from 87.9% at Sendai to 98.2% at Ny-Ålesund. Cloudy-sky agreement exceeds 95% at Chiba, Fukue, Hedo, Miyako, and Ny-Ålesund and remains relatively high at Syowa (90.7%) and Burgos (93.9%). For 0.400 and 0.870 μm wavelength pair (Figure 8), clear-sky agreement is also consistently high, ranging from 92.9% at Hedo to 100% at Miyako and Burgos. Cloudy-sky agreement exceeds 94% at Chiba, Fukue, Hedo, Miyako, and Ny-Ålesund but decreases to 78.7% at Sendai, 85.5% at Syowa, and 85.7% at Burgos. The 0.500–0.870 μm wavelength pair (Figure 9) also shows that clear-sky agreement ranges from 89.0% at Hedo to 100% at Miyako and Burgos. Cloudy-sky agreement exceeds 94% at Chiba, Fukue, Hedo, Miyako, and Ny-Ålesund, while lower values are found at Sendai (79.9%) and Syowa (85.5%). Burgos shows a cloudy-sky agreement of 91.8%. The fraction of observations classified as undetermined is generally small for ϵ value of 0.05. This indicates that the selected tolerance margin removes only a small fraction of observations from definite clear/cloudy classification while providing an explicit category for observations located close to the optimum decision boundary. Further, to quantify the relative performance of the four wavelength-pair slope indices, we calculated three classification metrics: clear recall, cloudy recall, and balanced accuracy. These are defined as follows:

$$\text{Clear Recall} = \frac{N_{\text{clear-clear}}}{N_{\text{MODIS,clear}}} \quad (19)$$

$$\text{Cloudy Recall} = \frac{N_{\text{cloudy-cloudy}}}{N_{\text{MODIS,cloudy}}} \quad (20)$$

- Formatted ... [14]
- Formatted ... [15]
- Formatted ... [16]
- Formatted ... [17]
- Formatted ... [18]
- Formatted ... [21]
- Formatted ... [19]
- Formatted ... [22]
- Formatted Table ... [20]
- Formatted ... [23]
- Formatted ... [25]
- Formatted ... [26]
- Formatted ... [24]
- Formatted ... [27]
- Formatted ... [29]
- Formatted ... [28]
- Formatted ... [30]
- Formatted ... [32]
- Formatted ... [33]
- Formatted ... [31]
- Formatted ... [34]
- Formatted ... [36]
- Formatted ... [35]
- Formatted ... [37]
- Formatted ... [38]
- Formatted ... [39]
- Deleted: To identify wavelength-pair slope indices that are ... [41]
- Formatted ... [40]
- Formatted ... [42]
- Deleted: 5
- Formatted ... [43]
- Deleted: positive slope values successfully identify cloudy ... [44]
- Formatted ... [45]
- Deleted: clear-sky detection improves substantially. Clear-sky d ... [46]
- Formatted ... [47]
- Deleted: shows a more balanced performance. Clear-sky d ... [48]
- Formatted ... [49]
- Formatted ... [50]
- Deleted: $\uparrow$... [51]
- Deleted: TN
- Deleted: TP

$$\text{Balanced Accuracy} = \frac{1}{2} (\text{Clear Recall} + \text{Cloudy Recall}), \quad (21)$$

where, $N_{\text{clear} \rightarrow \text{clear}}$ and $N_{\text{cloudy} \rightarrow \text{cloudy}}$ denote the numbers of observations correctly classified as clear and cloudy, respectively, while $N_{\text{MODIS_clear}}$ and $N_{\text{MODIS_cloudy}}$ denote the total numbers of reference clear- and cloudy-sky observations. The site-averaged values of these metrics are summarized in Table 3.

Table 3. Performance comparison of three wavelength-pair slope indices for clear/cloudy discrimination using MODIS cloud mask as the reference

Wavelength pair	Mean clear recall	Mean cloudy recall	Mean balanced accuracy	Site-to-site Standard Deviation
0.400 and 0.675 μm	96.9%	90.9%	93.9%	2.9%
0.500 and 0.675 μm	96.1%	94.5%	95.3%	2.4%
0.400 and 0.870 μm	97.2%	91.0%	94.1%	3.2%
0.500 and 0.870 μm	96.2%	92.0%	94.1%	3.6%

All four wavelength pairs show high site-averaged performance, with mean balanced accuracies ranging from 93.9% to 95.3%. The 0.500–0.675 μm pair provides the highest mean balanced accuracy (95.3%) and the smallest site-to-site variability (2.4%), with mean clear and cloudy recalls of 96.1% and 94.5%, respectively. This result is consistent with the pooled ROC analysis, in which this pair also shows the highest ROC AUC. The 0.400–0.870 μm pair provides the highest mean clear recall (97.2%), although its lower mean cloudy recall (91.0%) results in a balanced accuracy of 94.1%. The 0.500–0.870 μm pair shows the same mean balanced accuracy (94.1%), with clear and cloudy recalls of 96.2% and 92.0%, respectively. The 0.400–0.675 μm pair has the lowest, although still high, mean balanced accuracy (93.9%), mainly because of its relatively lower cloudy recall (90.9%).

Overall, the pooled ROC analysis and independent site-level evaluation provide consistent evidence that the atmospheric m_{lim} corresponding to all four wavelength pairs are useful for clear- and cloudy-sky discrimination. Among them, the 0.500 – 0.675 μm pair provides the strongest overall performance. The other three wavelength pairs also provide high discrimination performance and may therefore serve as complementary criteria.




Figure 6. Confusion matrices for clear, undetermined, and cloudy sky discrimination using atmospheric slope (m_{lim}) values determined using 0.400 and 0.675 μm wavelength pair. MODIS cloud-mask classifications are used as the reference, while POM-02 slope-based classifications are treated as predictions. Each matrix is normalized by the MODIS reference class, so that each row sums to 100%.




Figure 7. Same as Fig. 6, but for the spectral slope between 0.500 and 0.675 μm .






Figure 8. Same as Fig. 6, but for the spectral slope between 0.400 and 0.870 μm .



Deleted:

Formatted: Centered

Deleted: 6



Deleted:

Formatted: Centered

Deleted: 7




Figure 9. Same as Fig. 6, but for the spectral slope between 0.500 and 0.870 μm .

4.2 Validation of the Spectral Slope Criterion Using Direct Normal Irradiance





Figure 10. Relationship between median values of observed direct normal irradiance (DNI) within a 2° solar zenith angle (SZA) bin and SZA for clear, cloudy, and undetermined conditions predicted using wavelength pair of 0.500 and 0.675 μm , with $\epsilon = 0.05$ at (a) Chiba, (b) Hedo, (c) Fukue, (d) Miyako, and (e) Sendai.



Formatted: Centered

Deleted: ¶

Formatted

Formatted: Indent: First line: 0 ch

Deleted:

Formatted: Centered

Deleted: 8

Deleted: s

Deleted: μm and 0.400 and 0.875 μm

To evaluate the applicability of the m_{sun} criteria for discriminating clear, cloudy, and ambiguous sky conditions, direct normal irradiance (DNI) measurements obtained from pyrheliometers were analyzed together with sky radiometer observations. DNI provides an independent physical indicator of sky condition because it represents the direct solar beam reaching the surface and is therefore highly sensitive to cloud attenuation. Under clear-sky conditions, DNI is expected to remain relatively high and to decrease smoothly with increasing solar zenith angle (SZA) mainly because of increasing atmospheric air mass. Under cloudy conditions, DNI is strongly reduced because clouds attenuate or block the direct solar beam.

Figure 10 shows the relationship between median observed DNI and SZA for five observation sites: Chiba, Fukue, Hedo, Miyako, and Sendai. The median DNI values were calculated within 2° SZA bins. Sky conditions were classified using 0.500 and 0.675 μm wavelength pair with conservative threshold of $\varepsilon = 0.05$. These wavelengths also exhibit relatively stable long-term calibration characteristics in POM-02 instruments compared with shorter-wavelength channels (Uchiyama et al., 2018).

Across all five sites, the clear-sky class exhibits high median DNI values over low to moderate SZAs, followed by a gradual decrease at larger SZAs. This behavior is physically consistent with direct-beam transmission under cloud-free conditions, where the dominant SZA dependence is controlled by geometric air-mass effects. The clear-sky curves are especially coherent among the sites, although site-to-site differences are visible because of differences in aerosol loading, water vapor, surface environment, and sampling conditions.

In contrast, the cloudy class is characterized by substantially lower median DNI values over most SZA ranges. For several sites, cloudy-sky DNI values rapidly decrease to near-zero levels, indicating strong attenuation of the direct solar beam by clouds. This confirms that the combined positive-slope criterion successfully identifies conditions in which direct solar radiation is strongly suppressed.

The undetermined class occupies an intermediate regime between the clear and cloudy classes. Its median DNI values are generally lower than those of the clear class but higher and more variable than those of the cloudy class. This behavior is physically reasonable because the undetermined category includes cases where the two slope criteria do not provide a consistent classification, such as thin clouds, broken clouds, cloud-edge conditions, aerosol-influenced cases, or transitional sky conditions. Therefore, the undetermined group should not be interpreted as classification failure, but rather as a conservative buffer zone that prevents uncertain cases from being forced into either the clear or cloudy category.

Overall, the clear, cloudy, and undetermined DNI regimes are well separated. The high and smoothly varying DNI values for the clear class, the strongly reduced DNI values for the cloudy class, and the intermediate behavior of the undetermined class demonstrate that m_{sun} criteria provides a physically consistent and conservative method for sky-condition discrimination. The results may also indicate that introducing an undetermined category is useful for improving the robustness of the method, especially under mixed or transitional atmospheric conditions.

Deleted: combined spectral-slope
Formatted: Subscript

Deleted: 8
Deleted: two spectral-slope features derived from the
Deleted: and 0.400 and 0.875 μm
Deleted: s. A
Formatted: Font: Times New Roman, 10 pt
Deleted: was applied: clear conditions were assigned when both slopes were smaller than $-\varepsilon$, cloudy conditions when both slopes were larger than $+\varepsilon$, and all remaining cases were classified as undetermined.

Deleted: the combined use of the 0.500 and 0.675 μm and 0.400 and 0.875 μm
Deleted: spectral-slope
Formatted: Subscript
Deleted: ¶
Deleted: ¶

4.3 Comparison with Results from Existing Clear- and Cloudy-Sky Detection Procedure




Figure 11. Scatter plots of Ångström exponent (AE) versus aerosol optical thickness (AOT) at 0.5 μm for eight SKYNET sites after cloud screening using two systems: the SR-CEReS package (left panels) and the slope-based method derived from the atmospheric slope of zenith radiance between 0.500 and 0.675 μm with ε value of 0.05 (right panels). Numbers indicate the total observations retained by each method.

To evaluate the performance of the $m_{\min}$ -based cloud screening approach over a wider range of atmospheric conditions, the relationship between AOT (0.5 μm) and AE derived by detecting clear-sky conditions using 0.500 and 0.675 nm wavelength pair was compared with results obtained using the SR-CEReS package. The SR-CEReS processing package employs a multi-step cloud-screening procedure to identify cloud-contaminated observations before aerosol retrieval. The algorithm follows the method of Khatri and Takamura (2009), but excludes the pyranometer-based global irradiance test to allow consistent processing across SKYNET sites where pyranometer observations are unavailable. The cloud-screening procedure combines (1) a spectral variability test (Kaufman et al., 2006) and (2) statistical quality-control tests based on Smirnov et al. (2000), including checks of the number of observations, diurnal stability, smoothness of aerosol optical thickness, and the three-standard-deviation criterion. Observations failing one or more of these tests are classified as cloud contaminated and excluded from the aerosol retrieval process.

Figure 11 compares the AOT – AE relationships obtained using the two cloud-screening approaches for eight SKYNET sites covering markedly different atmospheric environments. Overall, the distributions obtained using the $m_{\min}$ -based method closely resemble those derived using SR-CEReS. This agreement is evident in the dominant AOT and AE ranges as well as in the overall shape and spread of the joint distributions. Figures 12 and 13 provide a more direct comparison of the corresponding probability-density distributions. The probability density was calculated as,

$$f_j = \frac{n_j}{N \Delta x}, \quad (22)$$

where n_j is the number of observations within bin j , N is the total number of observations retained by the corresponding cloud-screening method, and Δx is the bin width. Fixed bin widths of 0.05 for AOT and 0.1 for AE were used for all sites and for both screening methods.




Figure 12. Probability density distributions (PDFs) of AOT (0.5 μm) for eight SKYNET sites after cloud screening using two systems: the SR-CEReS package (left panels) and the slope-based method derived from the atmospheric slope of zenith radiance between 0.500 and 0.675 μm with ε value of 0.05 (right panels). Numbers indicate the total observations retained by each method.

Formatted: Subscript
Formatted: Line spacing: At least 0 pt
Formatted: Font: Times New Roman
Formatted: Font: Times New Roman, Not Bold
Deleted:
Formatted: ... [56]
Formatted: ... [57]
Formatted: ... [58]
Formatted: ... [59]
Formatted: ... [60]
Formatted: Font: 10 pt
Formatted: Subscript
Formatted: Font: Times New Roman, 10 pt
Formatted: Font: Times New Roman, 10 pt
Formatted: Font: 10 pt

As shown in Fig. 12, the AOT probability density distributions obtained using the proposed method generally closely follow those obtained using SR-CEReS. The locations of the major peaks and the overall shapes of the distributions are similar at most sites, despite the large differences in aerosol loading among the sites. Particularly close agreement is observed at Chiba, Cape Hedo, Miyako, Sendai, Burgos, Syowa, and Ny-Ålesund. A somewhat more noticeable difference occurs at Fukue, where the SR-CEReS distribution exhibits a broader contribution toward relatively high AOT than the proposed method. Nevertheless, the dominant AOT range remains similar between the two methods. This overall agreement indicates that the proposed screening does not substantially modify the AOT distribution of the observations identified as clear sky. A similar result is obtained for AE (Fig. 13). The two methods reproduce not only the locations of the major distribution peaks but also the characteristic site-to-site differences in AE. For example, the distributions at Syowa and Ny-Ålesund are shifted toward considerably larger AE than those at the midlatitude and tropical sites, and this distinctive behavior is reproduced by both screening methods. At several sites, the distribution obtained using the proposed method shows a small shift toward lower AE relative to SR-CEReS, but the overall shapes and ranges remain similar. Thus, the proposed method preserves the major characteristics of both the AOT and AE distributions across atmospheric environments ranging from very low aerosol loading at the polar sites to substantially higher aerosol loading at the midlatitude and tropical sites.




Figure 12. Probability density distributions of aerosol optical thickness (AOT) at $0.5 \mu\text{m}$ obtained after clear-sky screening using the SR-CEReS procedure and the proposed atmospheric-slope-based method at the eight SKYNET sites. A fixed AOT bin width of 0.05 was used for the probability density calculation. For clarity, the displayed AOT range is limited to 0 – 2.0, although the probability density analysis was performed using all valid observations over the full analysis range.

Formatted: Justified, Space Before: 0 pt, After: 0 pt

Formatted: Font: 11 pt

Deleted: Figure 5 presents AE–AOT scatter plots for eight SKYNET sites representing diverse atmospheric environments, as summarized in Table 1. Here, we used the criteria of $S_{0.5-0.675} < -\epsilon$ and $S_{0.4-0.87} < -\epsilon$ with ϵ value of 0.05. Overall, the AE–AOT distributions obtained using the slope-based method closely resemble those derived from the SR-CEReS outputs across all sites. In particular, the slope-based screening method reproduces the distinct aerosol regimes characteristic of each observation site. For example, relatively large AE values and broader AOT variability observed at the sites within Japan suggest the influence of both anthropogenic fine-mode aerosols and particles transported from surrounding oceanic and continental regions, including desert areas, via long-range transport. In contrast, the tropical continental site, Burgos, shows relatively high AE values associated with biomass burning and anthropogenic aerosols commonly observed in Southeast Asia. The polar sites—Syowa and Ny-Ålesund—exhibit very low AOT values and relatively narrow AE distributions, reflecting the extremely clean atmospheric conditions typical of polar regions. The ability of the slope-based method to reproduce these site-dependent aerosol characteristics indicates that the proposed screening approach does not introduce significant biases in the cloud-screened aerosol optical properties. Because the clear-sky detection criterion given in Eq. (6) is flexible, the screening can be made either stricter or more relaxed through an appropriate choice of logical operator and/or ϵ , depending on the application requirements. Nonetheless, the good agreement with the SR-CEReS results demonstrates that the proposed slope-based approach provides a practical and physically consistent method for cloud screening in SKYNET aerosol retrievals.

Formatted: Font: Bold

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt




Figure 13. Same as Fig. 12, but for the Ångström exponent. A fixed Ångström exponent bin width of 0.10 was used for the probability density calculation. For clarity, the displayed Ångström exponent range is limited to -0.5 - 3.5, although the probability density analysis was performed using all valid observations over the full analysis range.

Table 4. Seasonal comparison of the number of retained observations, aerosol optical thickness (AOT) at 0.5 μm , and Ångström exponent obtained after clear-sky screening using SR-CEReS and the proposed atmospheric-slope-based method

Site	Period	Count		AOT(0.5 μm)		Ångström exponent	
		SR-CEReS	This study	SR-CEReS	This study	SR-CEReS	This study
Chiba	DJF	38,757	41,076	0.11 ± 0.09	0.12 ± 0.10	1.29 ± 0.31	1.25 ± 0.34
	MAM	43,750	44,301	0.28 ± 0.18	0.27 ± 0.16	1.15 ± 0.36	1.14 ± 0.36
	JJA	32,280	32,357	0.28 ± 0.21	0.27 ± 0.19	1.28 ± 0.34	1.25 ± 0.36
	SON	31,532	33,011	0.15 ± 0.11	0.15 ± 0.11	1.29 ± 0.29	1.25 ± 0.32
Hedo	DJF	3,805	4,261	0.26 ± 0.13	0.28 ± 0.16	1.11 ± 0.27	1.05 ± 0.30
	MAM	11,771	12,826	0.37 ± 0.21	0.37 ± 0.21	1.10 ± 0.30	1.05 ± 0.32
	JJA	9,684	10,985	0.21 ± 0.19	0.20 ± 0.17	1.01 ± 0.39	0.95 ± 0.40
	SON	8,355	9,292	0.21 ± 0.14	0.22 ± 0.15	1.10 ± 0.34	1.05 ± 0.36
Fukue	DJF	3,796	4,076	0.25 ± 0.20	0.24 ± 0.19	0.96 ± 0.32	0.93 ± 0.32
	MAM	13,754	13,385	0.38 ± 0.25	0.32 ± 0.17	0.96 ± 0.35	0.97 ± 0.33
	JJA	5,429	5,307	0.32 ± 0.26	0.27 ± 0.18	1.12 ± 0.36	1.08 ± 0.37
	SON	9,283	9,454	0.23 ± 0.18	0.22 ± 0.16	1.14 ± 0.35	1.10 ± 0.36
Miyako	DJF	1,195	1,324	0.23 ± 0.12	0.23 ± 0.12	1.11 ± 0.27	1.06 ± 0.30
	MAM	3,158	3,374	0.33 ± 0.19	0.32 ± 0.17	1.16 ± 0.25	1.11 ± 0.28
	JJA	2,601	2,865	0.17 ± 0.17	0.15 ± 0.13	0.83 ± 0.41	0.79 ± 0.40
	SON	3,038	3,231	0.16 ± 0.11	0.16 ± 0.11	1.05 ± 0.35	0.99 ± 0.37
Sendai	DJF	2,330	2,564	0.12 ± 0.12	0.12 ± 0.10	1.24 ± 0.32	1.19 ± 0.36
	MAM	11,105	11,474	0.25 ± 0.19	0.23 ± 0.14	1.12 ± 0.39	1.11 ± 0.37
	JJA	8,661	8,861	0.27 ± 0.17	0.26 ± 0.15	1.27 ± 0.29	1.24 ± 0.31

Formatted: Font: Bold

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Line spacing: At least 0 pt

	SON	5,048	5,432	0.12 ± 0.08	0.13 ± 0.08	1.20 ± 0.28	1.17 ± 0.31
Burgos	Nov–Apr	19,125	20,433	0.19 ± 0.16	0.19 ± 0.16	1.14 ± 0.43	1.10 ± 0.44
	May–Oct	7,046	7,557	0.15 ± 0.12	0.16 ± 0.13	1.05 ± 0.37	1.02 ± 0.38
Syowa	Mar–Apr	441	468	0.07 ± 0.03	0.07 ± 0.05	2.19 ± 0.43	2.12 ± 0.53
	SON	2,827	2,942	0.10 ± 0.04	0.11 ± 0.05	2.21 ± 0.51	2.17 ± 0.58
	DJF	6,959	7,251	0.10 ± 0.07	0.11 ± 0.06	2.33 ± 0.47	2.28 ± 0.54
Ny-Ålesund	MAM	8,427	9,013	0.11 ± 0.10	0.11 ± 0.08	1.68 ± 0.37	1.62 ± 0.43
	JJA	14,441	15,324	0.08 ± 0.07	0.08 ± 0.07	1.71 ± 0.32	1.66 ± 0.40
	Sep–Oct	0	0	==	==	==	==

300 The seasonal statistics summarized in Table 4 provide a quantitative evaluation of this agreement. For the five Japanese sites, the comparison was performed separately for DJF, MAM, JJA, and SON; Burgos was divided into November – April and May – October; and seasonally appropriate periods were used for the two polar sites. For each period, Table 4 compares the number of retained observations, mean AOT ± standard deviation, and mean AE ± standard deviation obtained using SR-CEReS and the proposed method. In most cases, the seasonal mean values from the two screening approaches are very similar and remain well within the corresponding standard deviations.

305 At Chiba, for example, the mean AOT values obtained using SR-CEReS and the proposed method are 0.11 ± 0.09 and 0.12 ± 0.10 in DJF, 0.28 ± 0.18 and 0.27 ± 0.16 in MAM, 0.28 ± 0.21 and 0.27 ± 0.19 in JJA, and 0.15 ± 0.11 and 0.15 ± 0.11 in SON, respectively. The corresponding AE values also show close agreement, differing by only approximately 0.01 – 0.04 in their seasonal means. Similar behavior is observed at Cape Hedo, Miyako, and Sendai, indicating that the proposed method reproduces the seasonal aerosol characteristics identified by SR-CEReS.

310 Fukue shows somewhat larger differences during periods of enhanced aerosol loading. In MAM, the mean AOT decreases from 0.38 ± 0.25 using SR-CEReS to 0.32 ± 0.17 using the proposed method, while in JJA it decreases from 0.32 ± 0.26 to 0.27 ± 0.18 . Despite these differences, the corresponding AE values remain similar, and the overall AOT and AE distributions retain the similar characteristic between the two methods. This suggests that the two methods differ somewhat in the treatment of observations under higher aerosol-loading conditions, while the resulting aerosol populations remain broadly consistent.

315 Burgos provides a contrasting tropical environment. During November – April, the mean AOT is 0.19 ± 0.16 for both methods, while the mean AE values are 1.14 ± 0.43 and 1.10 ± 0.44 for SR-CEReS and the proposed method, respectively. During May – October, the corresponding AOT values are 0.15 ± 0.12 and 0.16 ± 0.13 , and the AE values are 1.05 ± 0.37 and 1.02 ± 0.38 . These close seasonal statistics support the similarity of the aerosol populations retained by the two cloud-screening approaches under tropical conditions.

320 The polar sites provide an additional test under very low-AOT conditions. At Syowa, during DJF, the mean AOT is 0.10 ± 0.07 for SR-CEReS and 0.11 ± 0.06 for the proposed method, while the corresponding AE values are 2.33 ± 0.47 and 2.28 ± 0.54 . During SON, the AOT values are 0.10 ± 0.04 and 0.11 ± 0.05 , and the AE values are 2.21 ± 0.51 and 2.17 ± 0.58 . At Ny-Ålesund, the MAM AOT values are 0.11 ± 0.10 and 0.11 ± 0.08 , while the corresponding AE values are 1.68 ± 0.37 and 1.62 ± 0.43 . The similarly close agreement during JJA further indicates that the proposed method remains effective under extremely clean atmospheric conditions.

325 Table 4 also shows that the number of retained observations is generally comparable between the two methods and is often slightly larger for the proposed m_{lim} -based screening. For example, at Chiba the proposed method retains 41,076 observations in DJF compared with 38,757 using SR-CEReS, while at Cape Hedo the corresponding values are 4,261 and 3,805. Similar increases occur at several other sites and periods. Importantly, this increase in the number of retained observations is not accompanied by a substantial shift in the mean AOT or AE values. This suggests that the proposed method can retain additional usable observations without introducing an obvious systematic bias into the retrieved aerosol optical properties.

330 Taken together, Figs. 11 – 13 and Table 4 demonstrate that the proposed m_{lim} -based screening reproduces both the distributional characteristics and the seasonal statistics of aerosol properties obtained using SR-CEReS across diverse atmospheric environments. The evaluation includes midlatitude Japanese sites, a tropical site, and two polar sites, and therefore spans a wide range of aerosol loadings, aerosol types, and seasonal conditions. These results indicate that the proposed method

- Formatted: Line spacing: At least 0 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Line spacing: At least 0 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Subscript
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Font: Times New Roman, 10 pt
- Formatted: Subscript
- Formatted: Font: Times New Roman, 10 pt

is robust across substantially different atmospheric regimes and does not introduce a systematic distortion of the cloud-screened aerosol climatology.

Nevertheless, the present evaluation is statistical and climatological rather than event specific. Individual desert-dust outbreaks, biomass-burning smoke episodes, and thin-cirrus cases were not independently identified using external aerosol- or cloud-type observations. Therefore, the present results should not be interpreted as a direct event-by-event validation for those specific conditions. Such evaluation would require independent identification using coincident lidar, satellite cloud products, or other ancillary observations. However, the multi-site, seasonal, and distributional comparisons presented here substantially broaden the range of atmospheric conditions under which the method is evaluated and provide quantitative evidence that its performance remains consistent across those conditions.

5 Conclusions

This study investigated the spectral characteristics of zenith sky radiance measurements from SKYNET sky radiometers under clear- and cloudy-sky conditions identified using quality-controlled MODIS cloud mask products. The study also examined the potential for discriminating between these sky conditions using spectral information.

For a given wavelength pair, the spectral slope derived from the measured signal, m_{obs} , contains contributions from both the atmospheric spectral dependence and the wavelength-dependent instrumental response and can be expressed as $m_{\text{obs}} = m_{\text{atm}} + m_{\text{inst}}$, where m_{atm} represents the atmospheric contribution and m_{inst} represents the wavelength-dependent instrumental contribution. Provided that the relative spectral response of an individual instrument remains sufficiently stable, m_{inst} , primarily introduces an instrument-dependent offset, while temporal variations associated with changes in atmospheric scattering conditions are retained in m_{atm} . The radiative transfer simulations demonstrated that m_{atm} is strongly negative under cloud-free conditions, particularly under relatively low aerosol loading, whereas cloud scattering systematically shifts m_{atm} toward less negative values because of the weaker wavelength dependence of cloud-droplet scattering and the enhanced contribution of multiple scattering. These results established a physical basis for using m_{atm} to distinguish between cloud-free and cloud-contaminated atmospheric conditions.

The analysis of spectral zenith-sky radiances estimated from observation data revealed that cloudy skies exhibit relatively flatter spectral behavior in the visible to near-infrared region, whereas clear skies show relatively stronger wavelength dependence consistent with Rayleigh and aerosol scattering processes. These results are consistent with spectral behaviours of aerosol- and cloud-laden atmospheres revealed from radiative transfer simulations. Among the wavelength combinations available from the instrument, m_{atm} values derived from four wavelength pairs (0.400 – 0.675, 0.400 – 0.870, 0.500 – 0.675, and 0.500 – 0.870 μm) showed consistently high capability for discriminating between clear- and cloudy-sky conditions. Based on these results, practical sky-state detection criteria were developed to discriminate observations into clear-sky, cloudy-sky, and undetermined categories. Although all these four wavelength pairs showed the capability of robust discrimination, the 0.500 – 0.675 μm pair provided the best performance across the different observation sites, with well-balanced capability for detecting both clear- and cloudy-sky conditions.

The robustness of the proposed criteria was further validated using collocated DNI measurements across different atmospheric regimes. The cases identified as clear-skies corresponded to high DNI values that varied smoothly with solar zenith angle, characteristic of clear-sky radiative transfer. In contrast, cases identified as cloudy skies were associated with substantially reduced DNI values, reflecting strong attenuation of the direct solar beam by clouds. Further, DNI values corresponding to undetermined cases existed between those for clear and cloudy sky conditions. This clear separation in DNI distributions across geographically and climatologically distinct sites confirms the physical consistency of the proposed slope-based classification approach for detecting clear- and cloudy-sky conditions easily and smoothly.

To further evaluate the performance of the m_{atm} -based screening approach, cloud-screened aerosol optical properties using our method were compared with those derived from the SR-CEReS package. The joint AE–AOT relationships and their probability distributions obtained from both approaches showed substantial agreement across multiple observation sites representing urban, maritime, tropical continental, and polar environments. In particular, the m_{atm} -based method successfully reproduced the characteristic aerosol regimes observed at each site without introducing noticeable biases in the retrieved aerosol optical properties. Consistent with these distributional comparisons, the seasonal mean AOT and AE values derived using the two approaches were generally similar and mostly within their corresponding variability ranges, demonstrating that the proposed method also preserves the seasonal characteristics of aerosol loading and particle-size-related behavior. Furthermore, the m_{atm}

Formatted: Font: Times New Roman, 10 pt

Deleted: ¶ ... [61]

Formatted: Font color: Auto

Formatted: Space Before: 0 pt, After: 0 pt, Line spacing: At least 0 pt

Formatted: Line spacing: At least 0 pt

Deleted: o

Deleted: By analyzing normalized spectral radiance behavior across wavelengths with minimal absorption by atmospheric gases at multiple SKYNET sites representing diverse atmospheric environments, systematic and physically consistent differences between clear- and cloudy-sky conditions were identified.

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Deleted: examined

Formatted: Subscript

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Deleted: 0.400 and 0.675 μm wavelength pair showed high sensitivity to cloudy-sky conditions and enabled more accurate identification of cloudy cases; however, it performed poorly in detecting clear-sky conditions. In contrast, the 0.500 and 0.675 μm and 0.400 and 0.870 μm pairs showed relatively higher sensitivity to clear-sky conditions and provided more balanced discrimination between clear and cloudy skies

Deleted: Specifically, the spectral slope, defined as the ratio of the natural logarithm of the zenith radiance ratio at the shorter and longer wavelengths to the natural logarithm of the corresponding wavelength ratio, showed predominantly negative values under clear skies and positive under cloudy skies. Furthermore, the spectral slope was controlled mainly by solar geometry and aerosol optical ... [62]

Deleted: These observation-based results were systematic ... [63]

Deleted: criterion

Deleted: slope

Formatted: Subscript

Deleted: developed criterion

Deleted: standard

Formatted: Font: Times New Roman, 10 pt

Deleted: strong

Deleted: slope

Formatted: Subscript

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt, Subscript

Formatted: Font: Times New Roman, 10 pt

based method generally retained a comparable or slightly larger number of observations without producing a substantial systematic shift in AOT or AE.

Overall, the results demonstrate that the spectral sky zenith radiances observed by sky radiometer provide a very powerful information to detect sky states reasonably without substantial computational burden or auxiliary data from other sources, provided that wavelength-dependent instrumental contribution, m_{inst} , is removed from the observed zenith sky radiance signals.

Author Contributions: Conceptualization, P. K. and H. I.; methodology, P. K.; software, P. K.; formal analysis, P. K.; investigation, P. K.; resources, H. I., H. K., I. M., T. H., H. I. and T. T. ; writing—original draft preparation, P. K.; writing—review and editing, P. K., T. H., I. M., Y. J., and Y. T.; funding acquisition, P. K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by JSPS KAKENHI Grant Number 24K07129 and collaborative research fund of Center for Environmental Remote Sensing (CEReS), Chiba University of Fiscal year 2025.

Data Availability Statement: MODIS data used in this study were obtained from the NASA Earthdata Search portal (<https://search.earthdata.nasa.gov/>). Aerosol optical thickness (AOT) data were downloaded from the SKYNET website (<http://atmos3.cr.chiba-u.jp/skyenet/>). The spectral zenith sky radiance data used in this study are available from the respective site owners upon reasonable request. The Burgos SKYNET operation is supported in part by the GOSAT series project. Local support for Burgos is provided by the Energy Development Corporation (EDC, the Philippines). Similarly, the observation data from Showa Station in Antarctica and Ny-Ålesund in the Arctic were collected as part of the National Institute of Polar Research's monitoring observations.

Conflicts of Interest: Authors have no conflict of interest.

Acknowledgements: Authors would like to thank the two anonymous reviewers and the editor for their useful and insightful comments and suggestions, which greatly helped improve the manuscript.

References

- Ackerman, S. A., Strabala, K. I., Menzel, W. P., Frey, R. A., Moeller, C. C., and Gumley, L. E.: Discriminating clear sky from clouds with MODIS, *Journal of Geophysical Research Atmospheres*, 103, 32141–32157, <https://doi.org/10.1029/1998JD200032>, 1998.
- Bagtasa, G., Cayetano, M. G., and Yuan, C. S.: Seasonal variation and chemical characterization of PM2.5 in northwestern Philippines, *Atmos. Chem. Phys.*, 18, 4965–4980, <https://doi.org/10.5194/acp-18-4965-2018>, 2018.
- Bi, J., Shi, J., Xie, Y., Liu, Y., Takamura, T., and Khatri, P.: Dust aerosol characteristics and shortwave radiative impact at a gobi desert of northwest China during the spring of 2012, *Journal of the Meteorological Society of Japan*, 92A, 33–56, <https://doi.org/10.2151/jmsj.2014-A03>, 2014.
- Bohren, C. F. and Huffman, D. R.: *Absorption and Scattering of Light by Small Particles*, wiley, 1–530 pp., <https://doi.org/10.1002/9783527618156>, 2007.
- Campanelli, M., Nakajima, T., Khatri, P., Takamura, T., Uchiyama, A., Estelles, V., Liberti, G. L., and Malvestuto, V.: Retrieval of characteristic parameters for water vapour transmittance in the development of ground-based sun-sky radiometric measurements of columnar water vapour, *Atmos. Meas. Tech.*, 7, 1075–1087, <https://doi.org/10.5194/amt-7-1075-2014>, 2014.
- Campanelli, M., Nakajima, T., and Olivieri, B.: Determination of the solar calibration constant for a sun-sky radiometer: proposal of an in-situ procedure, *Appl. Opt.*, 43, 651–659, <https://doi.org/10.1364/AO.43.000651>, 2004.

Deleted: ai

Deleted: iton

Formatted: Subscript

Formatted: Line spacing: At least 0 pt

Deleted: . Because the method relies on spectral shape rather than absolute radiance magnitude, it is relatively insensitive to calibration uncertainties and can be applied across diverse atmospheric environments. Therefore, spectral slope diagnostics of zenith radiances can serve as a reliable and computationally efficient complementary tool for cloud detection in SKYNET observations and other ground-based radiometric networks measuring zenith radiance.

Formatted: Line spacing: At least 0 pt

Deleted: ¶

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Choi, Y., Ghim, Y. S., Kim, S.-W., and Yeo, H.: Development of a cloud-screening algorithm for direct and diffuse AODs from the Skyradiometer Network, *Atmos. Res.*, **243**, 104997, <https://doi.org/10.1016/j.atmosres.2020.104997>, 2020.

Damiani, A., Irie, H., Takamura, T., Kudo, R., Khatri, P., Iwabuchi, H., Masuda, R., and Nagao, T.: An Intensive Campaign-Based Intercomparison of Cloud Optical Depth from Ground and Satellite Instruments under Overcast Conditions, *Sola*, **15**, 198–204, <https://doi.org/10.2151/sola.2019-036>, 2019.

Dim, J. R., Takamura, T., Higurashi, A., Kathri, P., Kikuchi, N., and Nakajima, T. Y.: Validation of Two MODIS aerosols algorithms with SKYNET and prospects for future climate satellites such as the GCOM-C/SGLI, *Advances in Meteorology*, 2013, <https://doi.org/10.1155/2013/508064>, 2013.

Gueymard, C. A.: The Sun's total and spectral irradiance for solar energy applications and solar radiation models, *Sol. Energy*, **76**, 423–453, <https://doi.org/10.1016/j.solener.2003.08.039>, 2004.

Hansen, J. E. and Travis, L. D.: Light scattering in planetary atmospheres, *Space Sci. Rev.*, **16**, 527–610, 1974.

Hara, K., Osato, K., Yabuki, M., Osada, K., Hirasawa, N., Shiobara, M., and Yamanouchi, T.: Antarctic haze phenomena at Syowa Station, Antarctica: seasonal features and impacts on atmospheric chemistry, *NPJ Clim. Atmos. Sci.*, **8**, <https://doi.org/10.1038/s41612-025-01176-9>, 2025.

Hashimoto, M., Nakajima, T., Dubovik, O., Campanelli, M., Che, H., Khatri, P., Takamura, T., and Pandithurai, G.: Development of a new data-processing method for SKYNET sky radiometer observations, *Atmos. Meas. Tech.*, **5**, 2723–2737, <https://doi.org/10.5194/amt-5-2723-2012>, 2012.

Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET—A Federated Instrument Network and Data Archive for Aerosol Characterization, *Remote Sens. Environ.*, **66**, 1–16, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.

IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2391 pp., <https://doi.org/10.1017/9781009157896>, 2023.

Irie, H., Mohammed Syedul Hoque, H., Damiani, A., Okamoto, H., Mashroor Fatmi, A., Khatri, P., Takamura, T., and Jarupongsakul, T.: Simultaneous observations by sky radiometer and MAX-DOAS for characterization of biomass burning plumes in central Thailand in January–April 2016, *Atmos. Meas. Tech.*, **12**, 599–606, <https://doi.org/10.5194/amt-12-599-2019>, 2019.

Kaufman, Y. J., Gobbi, G. P., and Koren, I.: Aerosol climatology using a tunable spectral variability cloud screening of AERONET data, *Geophys. Res. Lett.*, **33**, L07817, <https://doi.org/10.1029/2005GL025478>, 2006.

Khatri, P. and Takamura, T.: An algorithm to screen cloud-affected data for sky radiometer data analysis, *Journal of the Meteorological Society of Japan*, **87**, 189–204, <https://doi.org/10.2151/jmsj.87.189>, 2009.

Khatri, P., Takamura, T., Shimizu, A., and Sugimoto, N.: Spectral dependency of aerosol light-absorption over the East China sea region, *Scientific Online Letters on the Atmosphere*, **6**, 1–4, <https://doi.org/10.2151/sola.2010-001>, 2010.

Khatri, P., Takamura, T., Shimizu, A., and Sugimoto, N.: Observation of low single scattering albedo of aerosols in the downwind of the East Asian desert and urban areas during the inflow of dust aerosols, *J. Geophys. Res.*, **119**, 787–802, <https://doi.org/10.1002/2013JD019961>, 2014.

Khatri, P., Takamura, T., Yamazaki, A., and Uchiyama, A.: Use of 315 nm channel data of the sky radiometer to estimate the columnar ozone concentration: A preliminary study, *Journal of the Meteorological Society of Japan*, **92A**, 185–194, <https://doi.org/10.2151/jmsj.2014-A12>, 2014b.

Khatri, P., Hayasaka, T., Iwabuchi, H., Takamura, T., Irie, H., and Nakajima, T. Y.: Validation of MODIS and AHI observed water cloud properties using surface radiation data, *Journal of the Meteorological Society of Japan*, **96B**, 151, <https://doi.org/10.2151/jmsj.2018-036>, 2018.

Khatri, P., Iwabuchi, H., Hayasaka, T., Irie, H., Takamura, T., Yamazaki, A., Damiani, A., Letu, H., and Kai, Q.: Retrieval of cloud properties from spectral zenith radiances observed by sky radiometers, *Atmos. Meas. Tech.*, **12**, 6037–6047, <https://doi.org/10.5194/amt-12-6037-2019>, 2019.

Khatri, P., Hayasaka, T., Irie, H., Letu, H., Nakajima, T. Y., Ishimoto, H., and Takamura, T.: Quality assessment of Second-generation Global Imager (SGLI)-observed cloud properties using SKYNET surface observation data, *Atmos. Meas. Tech.*, **15**, 1967–1982, <https://doi.org/10.5194/amt-15-1967-2022>, 2022.

Formatted: Font: Not Bold

Formatted: Font: Times New Roman, 10 pt, Bold

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Kudo, R., Diémoz, H., Estellés, V., Campanelli, M., Momoi, M., Marengo, F., Ryder, C. L., Ijima, O., Uchiyama, A., Nakashima, K., Yamazaki, A., Nagasawa, R., Ohkawara, N., and Ishida, H.: Optimal use of the Prede POM sky radiometer for aerosol, water vapor, and ozone retrievals, *Atmos. Meas. Tech.*, 14, 3395–3426, <https://doi.org/10.5194/amt-14-3395-2021>, 2021.

540 Nakajima, T., Campanelli, M., Che, H., Estellés, V., Irie, H., Kim, S. W., Kim, J., Liu, D., Nishizawa, T., Pandithurai, G., Kumar Soni, V., Thana, B., Tugisurn, N. U., Aoki, K., Go, S., Hashimoto, M., Higurashi, A., Kazadzis, S., Khatri, P., Kouremeti, N., Kudo, R., Marengo, F., Momoi, M., Ningombam, S. S., Ryder, C. L., Uchiyama, A., and Yamazaki, A.: An overview of and issues with sky radiometer technology and SKYNET, *Atmos. Meas. Tech.*, 13, 4195–4218, <https://doi.org/10.5194/amt-13-4195-2020>, 2020.

545 Nakajima, T. and King, M. D.: Determination of the Optical Thickness and Effective Particle Radius of Clouds from Reflected Solar Radiation Measurements. Part I: Theory, *J. Atmos. Sci.*, 47, 1878–1893, [https://doi.org/https://doi.org/10.1175/1520-0469\(1990\)047<1878:DOTOTA>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1990)047<1878:DOTOTA>2.0.CO;2), 1990.

Nakajima, T., Tonna, G., Rao, R., Boi, P., Kaufman, Y., and Holben, B.: Use of sky brightness measurements from ground for remote sensing of particulate polydispersions, *Appl. Opt.*, 35, 2672–2686, <https://doi.org/10.1364/AO.35.002672>, 1996.

550 Ningombam, S. S., Larson, E. J. L., Indira, G., Madhavan, B. L., and Khatri, P.: Aerosol classification by application of machine learning spectral clustering algorithm, *Atmos. Pollut. Res.*, 15, <https://doi.org/10.1016/j.apr.2023.102026>, 2024.

Pasquier, J. T., David, R. O., Freitas, G., Gierens, R., Gramlich, Y., Haslett, S., Li, G., Schäfer, B., Siegel, K., Wiedner, J., Adachi, K., Belosi, F., Carlsen, T., Decesari, S., Ebell, K., Gilardoni, S., Gysel-Beer, M., Henneberger, J., Inoue, J., Kanji, Z. A., Koike, M., Kondo, Y., Krejci, R., Lohmann, U., Maturilli, M., Mazzolla, M., Modini, R., Mohr, C., Motos, G., Nenes, A., Nicosia, A., Ohata, S., Paglione, M., Park, S., Pileci, R. E., Ramelli, F., Rinaldi, M., Ritter, C., Sato, K., Storelvmo, T., Tobo, Y., Traversi, R., Viola, A., and Zieger, P.: The Ny-Ålesund Aerosol Cloud Experiment (NASCENT) Overview and First Results, *Bull. Am. Meteorol. Soc.*, 103, E2533–E2558, <https://doi.org/10.1175/BAMS-D-21-0034.1>, 2022.

Platnick, S., Meyer, K., King, M. D., Wind, G., Amarasinge, N., Marchant, B., Arnold, G. T., Zhibo, Z., Hubanks, P. A., Holz, R. E., Yang, P., Ridgway, W. L., and Riedi, J.: The MODIS cloud optical and microphysical products: Collection 6 updates and examples from terra and aqua, *IEEE Transactions on Geoscience and Remote Sensing*, 55, 502–525, <https://doi.org/10.1109/TGRS.2016.2610522>, 2017.

560 Ricchiazzi, P., Yang, S., Gautier, C., and Sowle, D.: SBDART: A Research and Teaching Software Tool for Plane-Parallel Radiative Transfer in the Earth's Atmosphere, *Bulletin of the American Meteorological Society*, 79(10), 2101–2114, 1998. DOI: 10.1175/1520-0477(1998)079<2101:SARATS>2.0.CO;2.

565 Shiobara, M., Hayasaka, T., Nakajima, T., and Tanaka, M.: Aerosol Monitoring Using a Scanning Spectral Radiometer in Sendai, Japan, 1991.

Smirnov, A., Holben, B. N., Eck, T. F., Dubovik, O., and Slutsker, I.: Cloud-screening and quality control algorithms for the AERONET database, *Remote Sens. Environ.*, 73, 337–349, [https://doi.org/10.1016/S0034-4257\(00\)00109-7](https://doi.org/10.1016/S0034-4257(00)00109-7), 2000.

570 Song, H. J., Sohn, B. J., Chun, H. W., Chun, Y., and Lee, S. S.: Improved cloud screening method for the analysis of sky radiometer measurements and application to asian dust detection, *Journal of the Meteorological Society of Japan*, 92A, 167–183, <https://doi.org/10.2151/jmsj.2014-A11>, 2014.

Uchiyama, A., Matsunaga, T., and Yamazaki, A.: The instrument constant of sky radiometers (POM-02) – Part I: Calibration constant, *Atmos. Meas. Tech.*, 11, 5363–5388, <https://doi.org/10.5194/amt-11-5363-2018>, 2018.

575 Wang, Z., Liu, D., Wang, Z., Wang, Y., Khatri, P., Zhou, J., Takamura, T., and Shi, G.: Seasonal characteristics of aerosol optical properties at the SKYNET Hefei site (31.90°N, 117.17°E) from 2007 to 2013, *Journal of Geophysical Research: Atmospheres*, 119, 6128–6139, <https://doi.org/10.1002/2014jd021500>, 2014.

Xue, Z., Kuze, H., and Irie, H.: Retrieval of aerosol optical thickness with custom aerosol model using SKYNET data over the Chiba area, *Atmosphere (Basel)*, 12, <https://doi.org/10.3390/atmos12091144>, 2021.

1580

Formatted: Font: (Default) Times New Roman, 10 pt

Deleted: Nakajima, T. and King, M. D.: Determination of the Optical Thickness and Effective Particle Radius of Clouds from Reflected Solar Radiation Measurements. Part I: Theory, *J. Atmos. Sci.*, 47, 1878–1893, [https://doi.org/https://doi.org/10.1175/1520-0469\(1990\)047<1878:DOTOTA>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1990)047<1878:DOTOTA>2.0.CO;2), 1990.

Deleted: ¶

Formatted: Font: 10 pt

Formatted: Indent: Left: 0 cm, Hanging: 0.25 cm, Line spacing: At least 0 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Formatted: Font: Times New Roman, 10 pt

Moved up [1]: Nakajima, T. and King, M. D.: Determination of the Optical Thickness and Effective Particle Radius of Clouds from Reflected Solar Radiation Measurements. Part I: Theory, *J. Atmos. Sci.*, 47, 1878–1893, [https://doi.org/https://doi.org/10.1175/1520-0469\(1990\)047<1878:DOTOTA>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1990)047<1878:DOTOTA>2.0.CO;2), 1990.

Formatted: Font: (Default) Times New Roman, 10 pt