

Polythermal conditions in small glaciers in the Swiss Alps

Janosch Beer^{1,2}, Mylène Jacquemart^{1,2}, Matthias Huss^{1,2,6}, Ilaria Santin^{1,2}, Gabriela Clara Racz³, Christophe Ogier^{1,2}, Saskia Gindraux⁴, Leo Hösli^{1,2}, Raphael Moser^{1,2}, James Irving³, Mauro Fischer^{5,7}, and Daniel Farinotti^{1,2}

¹Laboratory of Hydraulics, Hydrology and Glaciology (VAW), ETH Zurich, Zurich, Switzerland

²Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), bâtiment ALPOLE, Sion, Switzerland

³University of Lausanne, Institute of Earth Sciences, Lausanne, Switzerland

⁴Centre de recherche sur l'environnement alpin (CREALP), Sion, Switzerland

⁵Institute of Geography, University of Bern, Bern, Switzerland

⁶Department of Geosciences, University of Fribourg, Fribourg, Switzerland

⁷Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

Correspondence: Janosch Beer (beer@vaw.baug.ethz.ch)

Abstract. Englacial temperature measurements in the Alps remain sparse and biased toward high-elevation accumulation areas, leaving the thermal state of ablation zones poorly constrained. Here, we present borehole thermometry and ground-penetrating radar surveys from six small Swiss glaciers (< 0.5 km², 2700–3800 m a.s.l.). Polythermal conditions are confirmed. We confirm polythermal conditions in three glaciers, with – Alphubel South, Chessjengletscher, and Hohsaasgletscher – where the cold-temperate transition surface at depths of 17–38 m lies 17–38 m below the surface and ice temperatures ranging range from temperate to 2.1°C. A fourth site, Glacier du Sex Rouge, likely retains its historically documented polythermal structure, but borehole measurements were limited to the near-surface cold layer and do not reach the temperate ice below. A consistent spatial pattern emerges, with temperate ice at higher elevations transitioning into fully or partially cold-based glacier termini. Ground-penetrating radar retrievals are generally consistent with borehole-derived thermal conditions, with low scattering in cold ice and enhanced scattering in temperate zones. The observed thermal structures are closely linked to the history of firn cover loss, reconstructed from long-term mass balance records, with sites that lost their firn cover earliest showing the most advanced cooling. Our findings suggest that polythermal conditions among small Alpine glaciers may be more widespread than previously recognised, with important implications for glacial hazard assessments and highlighting the need for systematic regional-scale thermal observations and modelling.

1 Introduction

The glacier thermal structure influences key glacier processes, including ice deformation, ice viscosity, and basal sliding, as well as the storage and drainage of meltwater (Jansson, 1996; Flowers and Clarke, 2002; Boon and Sharp, 2003; Cuffey and Paterson, 2010). Glacier thermal conditions are generally classified into three categories: (1) fully temperate, where all ice is at the pressure melting point (PMP), (2) cold, where ice temperatures remain below the PMP, or (3) polythermal, where both cold and temperate ice coexist within a glacier. Englacial temperature is influenced by multiple factors, including the surface

energy balance, ice advection, internal heat production due to strain heating, geothermal heat flux, and latent heat release from refreezing water (Cuffey and Paterson, 2010).

Despite their importance, englacial temperature measurements remain sparse. In the Alps, existing observations are biased toward high-altitude accumulation areas (Haeberli and Alean, 1985; Haeberli and Funk, 1991; Suter and Hoelzle, 2002; Hoelzle et al., 2011) accumulation areas above 3800 m a.s.l., including sites in the Monte Rosa, Mont Blanc and Bernese Alps massifs (Haeberli and Alean, 1985; Haeberli and Funk, 1991; Suter and Hoelzle, 2002; Hoelzle et al., 2011; Pavlova et al., 2014) (Supplementary Fig. S1), where limited energy availability has historically resulted in predominantly cold thermal structures. More recent studies using both modelling and direct measurements have reported increasing englacial temperatures at these elevations (Vincent et al., 2020; Mattea, 2020; Gastaldello et al., 2025)(Vincent et al., 2020; Mattea et al., 2021; Gastaldello et al., 2025; Huss et al., 2026), a trend linked to extended melt seasons and latent heat release from refreezing meltwater in firn. In contrast, observations from ablation areas at lower elevations are scarce. A small number of measurements exist from large valley glaciers below 3000 m a.s.l., primarily dating back to the 1970s (Haeberli, 1976), as well as early pioneering measurements from the nineteenth century (Hugi, 1842; Agassiz, 1847). More recent ablation-area measurements are limited to a few notable exceptions: Glacier de Tête Rousse (Vincent et al., 2012), Vadret dal Corvatsch (Haeberli et al., 2004), Vadret Muragl, Vadret da Rims (Engadin, Switzerland) and Glacier du Sex Rouge (Fischer, 2018).

At these lower elevations, the thermal response of glaciers (< 3800 m a.s.l.), glaciers may respond to atmospheric warming may differ fundamentally from their high-elevation counterparts, as firn loss reduces meltwater retention and the associated latent heat release from refreezing differently from higher-elevation glaciers that still retain firn-covered accumulation areas, because firn loss alters how latent heat from meltwater is retained (Hock, 2005). Where firn is absent, the loss of latent heat from refreezing present, meltwater percolates to depth and refreezes in ice that is less thermally coupled to the surface, retaining the released latent heat; where firn is absent, meltwater instead runs off over the impermeable ice or refreezes at the surface, where the heat is quickly lost to the atmosphere. This loss of retained latent heat can outweigh other heat inputs, allowing the ice to cool despite rising air temperatures (Wilson and Flowers, 2013; van Pelt et al., 2016). While this effect has been studied in Arctic glaciers (Irvine-Fynn et al., 2006, 2011; Delcourt et al., 2013; ?), it has received comparatively little attention elsewhere in alpine and high-mountain environments (Gilbert et al., 2012; Huss and Fischer, 2016). Such cooling may promote the development of polythermal structures and is expected to be particularly relevant may be particularly important for small glaciers, where limited ice flow and internal deformation (Huss and Fischer, 2016) result in reduced strain heating reduce strain heating and sparse crevassing limits deep meltwater infiltration. Despite this expected sensitivity, the englacial thermal regime of small Alpine glaciers undergoing rapid firn loss remains almost entirely unobserved.

Understanding where and when such polythermal structures develop has direct relevance for glacier hazard assessment. While thermal conditions alone do not directly cause instability (Faillettaz et al., 2015), they can promote preconditions for ice break-offs, avalanches (Kavanaugh and Clarke, 2001; Faillettaz et al., 2015; Bondesan and Francese, 2023), glacier lake and water pocket outburst floods (?Huggel et al., 2004; Vincent et al., 2012)(?Huggel et al., 2004; Vincent et al., 2010, 2012), and even large-scale glacier detachments (Gilbert et al., 2018; ?; Jacquemart et al., 2020). In particular, polythermal glaciers are thought to be prone to the development of hazardous conditions, as the presence of cold ice can inhibit the development of

efficient drainage systems, increasing a glacier's susceptibility to accumulate meltwater beyond its drainage capacity (Wadham et al., 2001; Vincent et al., 2012; ?; Jacquemart et al., 2024). A recent example is the 2022 collapse of the Marmolada Glacier (Dolomites, Italy), which caused 11 fatalities and has been linked to a polythermal configuration in which cold basal ice likely prevented efficient meltwater drainage, leading to a frontal ice break-off (Bondesan and Francese, 2023; Olivieri and Bettanini, 2023). Assessing the hazard potential of alpine glaciers therefore requires a better understanding of the distribution of polythermal glaciers. Unfortunately, the lack of observations currently precludes any predictions about which glaciers might fall into this category, especially in the **aforementioned mid-elevation range . poorly sampled 2700–3800 m a.s.l. range** (Supplementary Fig. S1).

To address this knowledge gap, we instrumented six small ($< 0.5 \text{ km}^2$) glaciers in the Swiss Alps, spanning an elevation range of 2700–3800 m a.s.l.. In addition to several new glaciers, we included Glacier du Sex Rouge and Vadret dal Corvatsch in our study, where previous englacial temperature measurements (Haeberli et al., 2004; Fischer, 2018) enable a direct assessment of thermal change over decadal timescales. At all sites we measured englacial temperatures in boreholes during the summers of 2024 and 2025 and performed ground-penetrating radar (GPR) surveys to provide spatially continuous proxies for the thermal conditions, exploiting the sensitivity of radar scattering to liquid water content. To contextualise the observed thermal structures, we reconstructed the evolution of the firn cover from the 1970s to present, drawing on measured and modelled mass balance data (GLAMOS - Glacier Monitoring Switzerland, 2025).

2 Study sites

To address the observational gap identified above, site selection was guided by the following criteria:

1. Elevation; targeting glaciers below the current firn line where ongoing firn loss enhances susceptibility to conductive cooling and englacial temperature observations remain scarce.
2. Size; with a preference for very small glaciers ($< 0.5 \text{ km}^2$) where strain heating can be expected to be minimal (Huss and Fischer, 2016)
3. Limited crevassing, which reduces surface permeability and meltwater infiltration (Huss and Fischer, 2016; Gilbert et al., 2019)
4. An availability of previous englacial temperature measurements, enabling direct comparison with past observations.
5. Accessibility.

Following these criteria, we selected six small glaciers ($< 0.5 \text{ km}^2$) in the Swiss Alps: Alphubel South, Chessjengletscher, Hohsaasgletscher, Glacier du Sex Rouge, Glacier de Tortin, and Vadret dal Corvatsch (hereafter Alphubel, Chessjen, Hohsaas, Sex Rouge, Tortin, and Corvatsch, respectively; Table 1; Fig. 1 and 2). The sites cover an elevation range from ca. **2700–3800 m 2700 m to 3800 m** a.s.l. and mean surface slopes between **159° and 3329° within the study areas**. The sites span all

Table 1. Key characteristics of the six study sites. Area and elevation range were derived from the SwissALTI3D Digital Elevation Model (DEM) (Swisstopo, 2023a) clipped to the 2023 Swiss Glacier Inventory (SGI2023) outlines (Huss et al., 2025). Centre coordinates are given in the Swiss coordinate system (LV95; EPSG:2056). Mean slope and was averaged over the study area (the GPR survey / borehole footprint); aspect was averaged over all glacier pixels. For Chessjengletscher, this study focused on the lower ice body (area and slope given in parenthesis). The year when in-situ measurements were started in the frame of GLAMOS measurements is given. Previous TT: references to englacial temperature measurements prior to this study.

Glacier	Centre (E, N) [m]	Area [km ²]	Elev. range [m a.s.l.]	Aspect	Slope [°]	in-situ (since)	Previous T
Alphubel South (AH)	2633189, 1100627	0.07	3684–3890	SW	29	2023	–
Chessjengletscher (CJ)	2637008, 1101741	0.21 (0.12)	2881–3252	N	21	2023	–
Hohsaasgletscher (HS)	2643538, 1109805	0.15	3229–3504	W	21	2023	–
Glacier du Sex Rouge (SR)	2582698, 1130658	0.22	2714–2862	NW	9	2011	Fischer (2018)
Glacier de Tortin (GT)	2589973, 1103477	0.49	2780–3251	NW	12	2014	–
Vadret dal Corvatsch (CV)	2783238, 1143337	0.14	3073–3365	E	22	2012	Haeberli et al. (2004)

aspects, although the majority are located on north- and west-facing slopes. Three sites (Sex Rouge, Tortin, and Corvatsch) are part of the GLAMOS mass balance monitoring network, providing multi-year surface mass balance records as context for the thermal observations (Huss et al., 2026). For Sex Rouge and Corvatsch past borehole obseravtions observations are available as indicated in Table 1.

90 **3 Data and Methods**

We conducted two field campaigns, in the summers of 2024 and 2025. Before detailing the individual methods (Sects. 3.2 to 3.4), we first give a chronological overview of this fieldwork. During the initial campaign in August 2024, we drilled shallow boreholes on all six glaciers to depths of approximately 15 m and equipped them with two thermistors(cf. Sec. 3.2.2). . In parallel, we conducted ground-based GPR surveys on all glaciers expect Alphubel(cf. Sec. 3.1). except Alphubel. Follow-up
95 visits in September and October 2024 were used to retrieve initial thermistor data after sensor equilibration. A second field campaign between 5 and 13 August 2025 included the retrieval of the complete shallow temperature records and the drilling of additional deep boreholes to the glacier bed (here defined as bed-reaching boreholes of 20 to 58.3 m depth) at Alphubel, Chessjen, and Hohsaas. We retrieved temperature data from these deeper boreholes after one month (16-17.09.2025), with the exception of Chessjen, where a second readout was conducted in mid-December 2025. Additional radar measurements included
100 a GPR survey at Alphubel in May 2025 and an uncrewed aerial vehicle (UAV)-based GPR survey at Hohsaas in September 2025. Furthermore, mass balance measurements were conducted at all six study sites. End-of-melt-season stake readings were performed between August and October 2024 and 2025, with stake networks ranging from two to seven mass balance stakes




Figure 1. Overview and detailed maps of the six glaciers investigated in this study. (a) Location of study sites with cantonal boundaries and glacierized area according to the Swiss Glacier Inventory 2023 (SGI2023) (Huss et al., 2025). (b-g) Alphubel (AH), Chessjen (CJ), Hohsaas (HS), Sex Rouge (SR), Tortin (GT), and Corvatsch (CV), showing field installations on orthophotos (Swisstopo, 2024) with contour lines (Swisstopo, 2023a) and SGI2023 glacier outlines. Red arrows indicate approximate flow direction. Boreholes are distinguished by the type of thermistor string (see legend) and labeled profiles indicate transects used for temperature interpolation (L4, L47, L1, L2). Annotations on SR and CV mark sites of prior temperature measurements (Haeberli et al., 2004; Fischer, 2018). For Hohsaas (d), the survey extent and the extracted centerlines of a UAV-based GPR survey are indicated.




Figure 2. Overview images of the six study sites taken in July–August 2025: (a) Alphubel, (b) Chessjen, (c) Hohsaas, (d) Sex Rouge, (e) Tortin, and (f) Corvatsch. Approximate borehole locations are marked on each image and color-coded by sensor type: Geoprecision (G, red) and Tinytag (TT, white). The red arrow indicates the approximate flow direction of each glacier. CV1TT lies in a small depression and is therefore not directly visible; the green arrow indicates its approximate location.

per glacier. Snow depth measurements were collected in April and May 2025. A timeline of all borehole instrumentation and GPR surveys across the six study sites is given in FigSupplementary Fig. S2. 3.

105 **Timeline of borehole instrumentation, GPR and mass balance surveys across all study sites.** Bars indicate thermistor time-series color coded by manufacturer. All loggers remain active; bar ends denote the date of last readout. Black dashed lines indicate GPR survey dates.

3.1 Ground-penetrating radar (GPR) surveys

Two GPR systems were used in this study: a ground-based system deployed at all sites, and a UAV-based system used at Hohsaas. An overview of the setups for both systems is provided in Fig. S7 (Supplementary).

All ground-based GPR surveys were conducted using a Sensor & Software pulseEKKO Pro system. The choice of antenna frequency balanced penetration and robustness against scattering on the one hand, and ease of handling on the other. We used 100 MHz antennas at most sites as a conservative choice, ensuring good-quality bedrock reflections given the expected ice thickness and the scattering from englacial water and inhomogeneities. At Alphubel, the steep terrain favored the more compact 250 MHz antennas which still imaged the bedrock reliably. Antenna positions were tracked with a Leica differential GNSS

115

receiver operating in real-time kinematic (RTK) mode, providing centimeter-level positioning accuracy. Antenna elevations were corrected for the offset between the GNSS and radar antennas and for the antenna height above the ground surface. Depending on surface conditions, the antennas were either carried approximately 50 cm above the surface or, in areas with sufficient snow cover (Sex Rouge, Corvatsch, Alphubel), mounted on a sledge and pulled at approximately 10 cm above the surface. In both configurations, a constant separation of 2 m between transmitter and receiver was maintained. At most sites, the survey included a profile approximately along the glacier centerline connecting the borehole locations.

At Hohsaas, we employed a recently developed UAV-based GPR system (Ruols et al., 2023), using an M300 RTK UAV (DJI) carrying a GPR unit with a single transmit receive antenna at a center frequency of 80 MHz. Survey line positions were determined using a D-RTK 2 high-precision GNSS mobile station (DJI), providing centimeter-level RTK positioning (nominal accuracy 1 cm + 1 ppm horizontal, 2 cm + 1 ppm vertical). By flying a regular grid of parallel lines with 4 m spacing oriented approximately along the glacier contours, we obtained a 3D data cube covering nearly the entire glacier area. For the thermal interpolation, we extracted two down-slope profiles from this grid by sampling across the survey lines, each profile intersecting three boreholes.

All ground-based GPR data were processed using a standard workflow in Geolix (Geolix Technologies Inc., 2020); details are provided in Sect. 2.7.1 of the Supplement. The UAV-based GPR data from Hohsaas were processed using a separate workflow described in Sect. 2.7.2 of the Supplement. In both cases, the basal reflection was manually picked and ice thickness computed as the difference between surface and bed elevations, establishing the geometric framework needed for interpolation of the temperature data. We estimate the resulting bedrock-depth uncertainty at approximately ± 5 m, arising from the manual picking of the basal reflection and the vertical resolution of the radar. We primarily used the radar data to characterise the englacial thermal structure: In GPR data, temperate ice containing free water at grain boundaries produces diffuse scattering and signal attenuation, whereas cold ice is largely transparent to radar (Pettersson et al., 2004; Vincent et al., 2012; Wilson et al., 2013; Sevestre et al., 2015; Gilbert et al., 2019; Forte et al., 2025). The cold-temperate transition surface (CTS) is therefore identifiable as the boundary between diffuse scattering and low-reflectivity zones, used here to independently validate borehole-derived CTS positions and to extend the thermal interpretation beyond the borehole network.

3.2 Borehole temperature measurements

3.2.1 Drilling and borehole locations

During the initial campaign in August 2024, we drilled two shallow boreholes on each of the six glaciers using a Heucke steam drill (Heucke, 1999). We selected this system for its portability, even though it limited maximum borehole depth to 15 m. Attained borehole depths ranged from 7 to 15.6 m, reaching bedrock at the shallowest sites. We generally selected one borehole at higher and one at lower elevation per glacier to capture contrasts in surface settings. At Corvatsch and Sex Rouge, locations were chosen to allow comparison with previous measurements. Shallow boreholes for ablation stakes were drilled using a Kovacs ice auger powered by an electrical hand drill, which was also used to pre-drill the upper few meters of

deeper boreholes. Ablation stake readings were used to monitor local melt and adjust the effective measurement depth of the thermistors.

150 Based on the preliminary insights from the 2024 data, in 2025 we drilled boreholes to the glacier bed at Alphubel, Chessjen, and Hohsaas to acquire complete vertical temperature profiles. Drilling was conducted using a hot-water drill system consisting of a Kärcher HD 9/23G pressure pump connected to a HG43 flow heater (Supplementary Fig. S2S3). In this manner we drilled eleven additional boreholes (four each at Alphubel and Chessjen, three at Hohsaas), with depths ranging from 20 to 58.3 m and with seven of them reaching bedrock. Boreholes were positioned near the glacier centerline and, where possible, in a way to
155 fill gaps from the 2024 campaign. We surveyed all borehole positions using a GNSS antenna, which also enabled calculation of annual glacier surface displacement rates for the 2024 boreholes (Supplementary Table S14S17). Across both campaigns, we drilled a total of 23 boreholes. Table A1 in the Appendix provides an overview of locations, elevations, depths, and sensor types, while Figs. 1b–g and 2 illustrate borehole positions and local topography.

3.2.2 Measurements and calibration

160 Depending on the borehole depth, we used two thermistor setups: Tinytag Plus 2 (TGP-4520) loggers from Gemini Data Loggers for shallow boreholes, and multi-sensor thermistor chains from Geoprecision for deeper boreholes.

During the 2024 campaign, we installed Tinytag loggers in weatherproof enclosures mounted on aluminum plates and secured to ablation stakes at the borehole sites (Supplementary Fig. S3aS4a). Each logger records two Tinytag PB-5010 negative temperature coefficient (NTC) thermistors with a nominal accuracy of $\pm\pm 0.2^{\circ}\text{C}$ (manufacturer specification) and a reading
165 resolution of 0.02°C . We positioned the sensors so that one reached the maximum borehole depth and the second was installed five meters above. Since our shallow boreholes were approximately 15 m deep, this configuration typically provided measurements at depths of about 10 m and 15 m. For the Tinytag sensors, which operated over a full year, we were able to compute the actual sensor depth at the time of readout by accounting for surface melt measured at the stake locations between installation and retrieval. Prior to deployment, we calibrated these sensors in an ice-bath setup after 1-2 hours of thermal equilibration to
170 quantify and correct sensor-specific offsets relative to 0°C . The mean offset was 0.3°C , larger than the nominal accuracy and thus a meaningful correction for these sensors.

The Geoprecision multi-sensor thermistor chains, deployed in the deep boreholes, were secured by taping the logger at the top of each chain to a 1 m wooden stick over the borehole (Supplementary Fig. S3bS4b). Depending on borehole depth, the chains contained 5–10 5-10 TNodeHD thermistors that were spaced between 2.5 and 5 m apart. Like the Tinytag sensors, the
175 TNodeHD thermistors use NTC sensors but offer higher measurement accuracy ($\pm\pm 0.05^{\circ}\text{C}$, manufacturer specification) and a temperature resolution of 0.0001°C .

Before field deployment, we calibrated all thermistors to quantify sensor-specific offsets relative to 0°C . We calibrated Tinytag thermistors using an ice-bath setup after 1–2 hours of thermal equilibration, and we calibrated the Geoprecision chains. These chains were calibrated in a temperature-controlled alcohol bath (Huber CC245 WL) with against a high-precision reference thermometer (ISOTECH F250) (; Supplementary Fig. S4). We then applied the derived offsets S5). The mean offset was
180 only -0.004°C .

The derived offsets were applied to all sensors before analysis. A detailed description of the calibration procedures, including stability criteria and a statistical overview of the offsets, is provided in Sect. 2.3 of the Supplement. The temperature contrasts underlying our interpretations are generally substantially larger than these sensor accuracies; where individual values fall within sensor accuracy of the melting point, we interpret them accordingly.

3.3 Ground-penetrating radar (GPR) surveys

Two GPR systems were used in this study: a ground-based system deployed at all sites, and a UAV-based system used at Hohsaas. An overview of the setups for both systems is provided in Fig. S6 (Supplementary).

All ground-based GPR surveys were conducted using a Sensor & Software pulseEKKO Pro system. Antennas operated at a central frequency of 100 MHz, except for the survey on Alphubel where we used 250 MHz. Antenna positions were tracked in real-time with a Leica differential GNSS receiver. Depending on surface conditions, the antennas were either carried approximately 50 cm above the surface or, in areas with sufficient snow cover (Sex Rouge, Corvatsch, Alphubel), mounted on a sledge and pulled at approximately 10 cm above the surface. In both configurations, a constant separation of 2 m between transmitter and receiver was maintained. At most sites, the survey included a profile approximately along the glacier centerline connecting the borehole locations.

At Hohsaas, we employed a recently developed UAV-based GPR system (Ruols et al., 2023), using an M300 RTK UAV (DJI) carrying a GPR unit with a single transmit receive antenna at a center frequency of 80 MHz. Survey line positions were determined using a D-RTK 2 high-precision GNSS mobile station (DJI). By flying a regular grid of parallel lines with 4 m spacing oriented approximately along the glacier contours, we obtained a 3D data cube covering nearly the entire glacier area. For the thermal interpolation, we extracted two down-slope profiles from this grid by sampling across the survey lines, each profile intersecting three boreholes.

All ground-based GPR data were processed using a standard workflow in Geolix (Geolix Technologies Inc., 2020); details are provided in Sect. 2.7.1 of the Supplement. The UAV-based GPR data from Hohsaas were processed using a separate workflow described in Sect. 2.7.2 of the Supplement. In both cases, the basal reflection was manually picked and ice thickness computed as the difference between surface and bed elevations, establishing the geometric framework needed for interpolation of the temperature data. We primarily used the radar data to characterise the englacial thermal structure: In GPR data, temperate ice containing free water at grain boundaries produces diffuse scattering and signal attenuation, whereas cold ice is largely transparent to radar (Pettersson et al., 2004; Vincent et al., 2012; Wilson et al., 2013; Sevestre et al., 2015; Gilbert et al., 2019; Forte et al., 2025). The cold-temperate transition surface (CTS) is therefore identifiable as the boundary between diffuse scattering and low-reflectivity zones, used here to independently validate borehole-derived CTS positions and to extend the thermal interpretation beyond the borehole network.

3.3 Borehole temperature analysis

3.3.1 Temperature data analysis

The temperature data were analysed with two objectives: characterising the seasonal thermal signal at each site, and constructing time-averaged vertical temperature profiles for spatial interpolation. The Tinytag time series (boreholes 1TT and 2TT), which cover nearly a full annual cycle (Supplementary Fig. 3S2), are used to characterise seasonal variations, estimate the depth of the zero annual amplitude (ZAA; See Sect. 2.6 Supplementary), and identify transient events such as short-term warming following water intrusion. This analysis is presented in Sect. 4.1. At Sex Rouge and Corvatsch, where historical temperature measurements are available, the continuous Tinytag record also allows a day-to-day comparison with the historical data, which consist of single transient measurements.

To construct vertical temperature profiles, we averaged temperatures over the measurement period for each sensor, excluding an initial equilibration phase of 14 days following installation to allow the borehole to refreeze and temperatures to stabilise. The thermal disturbance introduced by hot-water drilling is spatially confined to the immediate borehole surroundings and decays over timescales of days to weeks, depending on borehole diameter and ice temperature (Humphrey and Echelmeyer, 1990; Cuffey and Paterson, 2010). Our 14-day exclusion window is therefore sufficient for the deep boreholes to converge on stable, undisturbed ice temperatures, as confirmed directly by the equilibrated sensor records (Supplementary Fig. S14-16). Sensor depths were corrected for cumulative surface melt between installation and readout, using the melt-corrected final depth z_{final} as the effective measurement depth for interpolation. The interpolation was restricted to Alphubel, Chessjen, and Hohsaas, where the installation of additional thermistor chains provided a higher density of observations, including measurements at or near the glacier bed. The interpolated temperature fields were subsequently compared against ground-penetrating radar reflectivity patterns to assess their consistency and to avoid over-interpretation of areas without direct temperature measurements.

Finally, boreholes were assigned to one of three thermal regime categories based on the fraction of temperature measurements at or above 0°C within sensor precision accuracy (0.05°C for Geoprecision, 0.20°C for Tinytag): to-bed boreholes with less than 2% such measurements were classified as cold-based. Those cold, those with 50% or more as mostly temperate, temperate, and all others as mostly coldpolythermal. These categories are used in Fig. 8 to contextualise borehole locations thermal structure within the firn cover reconstruction.

3.3.2 Interpolation and CTS detection

Borehole temperature data were interpolated along downslope profiles, connecting as many borehole locations as possible. This resulted in one profile for Alphubel and Chessjen and two profiles for Hohsaas, extracted by sampling across the contour-parallel UAV-GPR survey lines. The profiles (L4, L47, L1, L2) are marked in Fig. 1a-c. Interpolation was performed using Radial Basis Function (RBF) interpolation (Hardy, 1971), implemented via the RBFInterpolator routine of the SciPy library (Virtanen et al., 2020). RBF interpolation is well suited for this application as it makes no assumptions about the spatial pattern of the temperature field and handles sparse, unevenly distributed observations well. To account for the strong vertical temperature gradients (compared to horizontal) (Cuffey and Paterson, 2010), the vertical coordinate was scaled prior

245 to interpolation. The scaling factor was determined empirically from the ratio of observed vertical to horizontal temperature gradients in the borehole data. A multiquadric kernel was used and the interpolated field was evaluated on a regular grid masked to the glacier geometry. A detailed description of the interpolation scheme is provided in Sect. 2.4 of the Supplement.

The position of the cold-temperate transition surface (CTS) CTS was estimated by identifying grid cells where the interpolated temperature approached the pressure-dependent melting point, following the Clausius-Clapeyron relation with
250 $\beta = 8.7 \times 10^{-4} \text{ }^{\circ}\text{C m}^{-1}$ (Cuffey and Paterson, 2010) and a tolerance of $\delta_{\text{CTS}} = 0.05 \text{ }^{\circ}\text{C}$. Details are given in Sect. 2.5 of the Supplement.

3.4 Mass balance and long-term firn cover development

Annual point mass balance was measured at all Tinytag borehole locations (1TT and 2TT). Snow depth measurements were additionally collected across each glacier in April 2025 using manual probing at 15-60 points per site (Supplementary Fig.
255 S8S9). Snow density was measured on site using a coring device, or adopted from measurements on nearby glaciers. At Corvatsch, Sex Rouge and Tortin, these measurements are part of monitoring programs started around 2010. Point measurements of ablation and accumulation were spatially extrapolated to the entire glacier using a distributed mass balance model (Huss et al., 2021)(Huss et al., 2021), which accounts for spatial variability and the inhomogeneous distribution of field observations. The model combines a distributed accumulation model with a temperature-index melt model that includes a potential-solar-radiation term (Hock, 1999), rather than a full surface energy balance. Melt is computed from air temperature and potential
260 clear-sky solar radiation, while accumulation is derived from precipitation, scaled by a correction factor and distributed using observed snow-accumulation patterns to capture the spatial variability of snow deposition across the glacier. Model parameters are calibrated against the seasonal stake and snow-probing measurements at each site. A full description of the model, including the melt and accumulation formulations and the calibration procedure, is given in Huss et al. (2021). This approach allows
265 inferring surface mass balance distribution on a 10 m spatial grid constrained by the seasonal measurements.

The calibrated mass balance model can also be driven by meteorological forcing outside of the period covered by in-situ measurements. In this case, the model is constrained by observed long-term changes in ice volume (Huss et al., 2015). Supported by the detailed measurements of the recent years, informing about the spatial variability of mass balance components, the model thus provides a gridded annual mass balance distribution for each glacier from 1970 to 2025. These grids were
270 used to derive spatially distributed firn cover maps by cumulatively tracking mass balance at each grid cell through time. Firn thickness was updated annually by adding positive mass balance (accumulation) and removing negative mass balance (ablation) from the existing firn column, with ablation removing the most recently deposited layers first. Grid cells were classified as firn-covered if they accumulated snow in two consecutive years. This criterion excluded single years with a larger end-of-season snow extent from the firn extent. Firn layers persisting for more than 20 years were considered ice. This threshold was
275 arbitrarily chosen and represents , a simple assumption consistent with typical densification timescales in the Alps (Ambach and Eisner, 1966). This approach yields annual maps of firn thickness and extent from 1970 to 2025 for each glacier. Because the reconstruction depends primarily on the accumulation component, its reliability rests on how well snow accumulation and its spatial distribution are captured. For Alphubel, Chessjen, and Hohsaas the accumulation model is constrained by only a

single year of snow-probing data, fewer than at the long-monitored sites (Corvatsch, Sex Rouge, Tortin). We therefore assessed
280 the reconstructed firn extent independently against firn/perennial-snow boundaries manually digitized from high-resolution
SWISSIMAGE orthophotos (Swisstopo, 2023b); the comparison shows no systematic bias and comparable agreement across
all six glaciers, including the less-constrained polythermal sites (Supplementary Sect. 3.1). From the annual firn cover maps
we computed the time since firn loss for each cell.

4 Results and discussion

285 4.1 Seasonal temperature variations and meltwater signals

Tinytag sensors installed in 2024 recorded nearly a full annual temperature cycle at all six sites (Fig. 3). Over the measurement
period, surface melt thinned the ice above each sensor, so that effective sensor depths changed between installation (Z_{init}) and
retrieval (Z_{final}). For simplicity, all depths stated in the following refer to the installation depth. The reduction in effective
depth ranged from approximately 0.4 m to more than 2 m, meaning that sensors progressively sampled shallower ice over
290 the course of the measurement period. We expect this to result in slightly larger observed seasonal amplitudes. Unless stated
otherwise, reported temperatures carry an uncertainty corresponding to the calibrated sensor accuracy, $\pm 0.2^\circ\text{C}$ for the Tinytag
and $\pm 0.05^\circ\text{C}$ for the Geoprecision sensors. Seasonal temperature amplitudes varied by nearly two orders of magnitude across
sensors and sites, with the largest recorded amplitude of 3.64°C at Tortin (GT1TT, 4.2 m), the smallest of 0.03°C at Hohsaas
(HS1TT, 15.6 m). As expected, amplitudes decreased systematically with depth and the temperatures at depths of 10 m to 15
295 m lagged roughly six months behind the surface (Fig. 3 a-d), meaning that the lowest temperatures occurred in the summer,
the highest in winter.

At Alphubel, Chessjen, Hohsaas, and Sex Rouge, temperatures showed no sign of approaching 0°C at the instrumented
depths (6.8-15.4 m), suggesting that the winter cold wave could not be compensated during summer. Two sensors are exceptions
to this: SR2TT (10 m) and HS1TT (15.6 m), whose temperature maxima (0.18°C and 0.1°C , respectively) fall within the
300 $\pm 0.2^\circ\text{C}$ sensor accuracy, meaning temperate conditions at these depths are likely. Shallow sensors at Tortin and Corvatsch
were trending towards 0°C by the end of the observation period and are most likely approaching temperate conditions later
in the year. The final readout was conducted before the 2025 summer warming signal had fully propagated through the ice
column. GT1TT (4.2 m) and CV1TT (2.0 m) are therefore expected to reach their annual maximum during or shortly after the
melt season, while sensors at 10 to 15 m depth likely do so between December and March.

305 At Chessjen, Hohsaas, and Sex Rouge, mean temperatures increase with depth. Deeper temperatures are on average 1.2-
1.5 $^\circ\text{C}$ warmer than shallower ones at Chessjen, 0.6-0.9 $^\circ\text{C}$ warmer at Hohsaas, and approximately 0.1 $^\circ\text{C}$ warmer at Sex Rouge.
This distinction is less obvious at Alphubel, where mean temperatures appear to vary more with borehole position than with
depth, ranging from -2.0 to -2.2 $^\circ\text{C}$ at AH2TT (the upper borehole) and from -2.3 to -2.7 $^\circ\text{C}$ at AH1TT further downslope. At
Alphubel, Chessjen, Hohsaas, and Sex Rouge, where the deepest sensors exhibit amplitudes below 0.3°C , we estimated the
310 depth of zero annual amplitude (ZAA; See Sect. 2.6 Supplementary) to range from 17 m at Alphubel to 23 m at Sex Rouge
(see annotation in panels a-d).




Figure 3. Seasonal evolution of englacial temperature at all six study sites from August 2024 to September 2025. Line colour indicates initial sensor installation depth, with solid lines denoting Tinytag borehole 1 (1TT) and dashed lines borehole 2 (2TT). For Glacier de Tortin, only borehole 1 is shown due to calibration issues affecting borehole 2. Each panel lists sensor installation depth (Z_{init}) and depth at last readout (Z_{final}). At CV2TT the lower sensor failed during the measurement period and is thus not shown. Where the seasonal amplitude at the deepest sensor was sufficiently small, the ZAA is indicated.

Short-term temperature jumps are visible in both the Tinytag and Geoprecision time series data at Alphubel and Chessjen. At CJ2TT, the shallow sensor recorded an abrupt warming of 1.64°C on 18-19 July (Fig.3b), and at Alphubel, the upper borehole (AH1G) showed a temperature jump of approximately 0.15°C on 17-18 August (Supplementary Fig. S11a-S14a). Since these events occurred during the melt season and have transient character, we interpret them as the effect of englacial meltwater, likely routed via crevasses and englacial channels. The resulting temperature jumps reflect heat input to the surrounding ice, through direct exchange with liquid water, latent heat release upon refreezing, or both - processes central to the thermal structures discussed in Sect. 4.3.

4.2 Decadal changes of **of** englacial temperatures at Sex Rouge and Corvatsch

320 The availability of previously published borehole data at Sex Rouge and Corvatsch provides a rare opportunity to investigate how englacial temperatures at these sites have evolved over decadal timescales (Fig. 4). The earlier measurements represent multiple, single-day snapshots, while our data comprise a full annual time series. To compare, we extracted the matching days of they year from our time series for each prior measurement date. **Matching the day of year controls for the seasonal phase of the signal but not for inter-annual weather variability, which still influences temperatures within the seasonally affected zone**
325 **sampled here. We therefore compare the historical single-day measurements against the full modern annual range at each depth (min-max envelope in 4), and interpret differences within the seasonal zone in terms of the broader thermal structure and its evolution rather than as precise climatic-trend magnitudes.**

At Sex Rouge, Fischer (2018, data from 2013-2015) documented a polythermal structure with a cold surface layer of approximately 20 m overlying a temperate core. We also found a persistent cold surface layer with a strong temperature gradient
330 trending to 0 °C with depth (Fig. 4a,b), suggesting that this polythermal structure likely still persists today. Relative to the 2013-2015 measurements, temperatures at 10 m rose by 0.65°C at SR2TT and by 0.23°C at SR1TT, while at 15.2 m temperatures fell by 0.11°C at SR1TT, remaining within sensor accuracy ($\pm 0.2^\circ\text{C}$). However, with approximately 20 m of surface ice lost through ablation since 2013 (GLAMOS - Glacier Monitoring Switzerland, 2025), our sensors sample ice that was located considerably deeper during the earlier measurement period. The hollow red diamonds in Fig. 4 show our current temperatures
335 projected to their estimated depths at the time of the prior measurements, illustrating this reference-frame shift. This vertical shift complicates our interpretations. The observed signal reflects the effects of downward cold wave penetration and surface removal of cold ice through ablation, rather than atmospheric forcing alone. These competing drivers are discussed further in Sect. 5.1.

At Corvatsch, Haeberli et al. (2004) documented cold-based conditions with temperatures of approximately -4°C at 13 m
340 depth in 1999/2000. The glacier has since thinned to a point where the entire ice column, now only 9.3 m thick, is subject to seasonal temperature fluctuations (Fig. 3f). Temperatures at 4.3 m depth exhibit a clear seasonal cycle but remain sub-zero throughout the year, indicating that summer warming is insufficient to bring the ice to the melting point. Our measured temperatures at 4.3 m depth were on average $-0.36^\circ - 0.36^\circ\text{C}$ colder than the measurements at comparable depths from Haeberli et al. (2004), **suggesting a slight cooling of the near-surface ice since 1999/2000.** . The lower sensor reached the bed at 9.3 m but
345 failed during equilibration, recording -1.3°C with temperatures still declining at the time of failure, suggesting the bed remains cold today. Among the six study sites, Corvatsch occupies a unique topographic position on a narrow ice ridge. The ridge geometry limits snow accumulation and firn formation while exposing both flanks to lateral conductive heat loss, providing an additional cooling pathway beyond surface forcing alone (Haeberli et al., 2004). The persistently cold conditions observed here are therefore at least partly a consequence of this inherently cold topographic setting, alongside the long firn-free history
350 of the site, discussed further in Sect. 5.1.

4.3 Polythermal sites (Alphubel, Chessjen, Hohsaas)




Figure 4. Englacial temperature profiles at Sex Rouge and Corvatsch compared with historical borehole data. (a) SR1TT versus four repeated profiles from BH1 by Fischer (2018). (b) SR2TT versus two single-point measurements from BH2 by Fischer (2018). (c) CV2TT versus five profiles by Haeberli et al. (2004). Red diamonds indicate the time-averaged temperature at each sensor depth over the Aug 2024–Sep 2025 measurement period; horizontal bars show the observed min–max range. (d–f) Temperature difference relative to historical measurements, computed by matching each historical snapshot to the same calendar day in our 2024/25 record; vertical lines indicate the mean difference $\overline{\Delta T}$ at each sensor depth.

Englacial temperature profiles Because deep boreholes reaching the glacier bed were drilled only at Alphubel, Chessjen, and Hohsaas reveal polythermal conditions at all three borehole sites (Fig. (Sec. 3.3.1)), the following analysis of the full vertical thermal structure is limited to these three sites. The shallow (15 m) boreholes at Corvatsch, Sex Rouge, and Tortin sample only the seasonally influenced near-surface layer. At all three deep-drilled sites, englacial temperature profiles reveal polythermal conditions (5, 6, 7), with a recurring spatial pattern: a temperate core in the upper reaches of the glacier transitioning to partially and fully cold-based conditions near the glacier terminus.

In all three glaciers, we determined the CTS to be at relatively shallow depths of 17–20 m in the upper reaches, while ice near the termini was found to be predominantly cold (Figs. 5a, 6a, 7a,b). At Alphubel, temperate ice was confirmed between 25 m and 33 m depth at AH1G and between 20 m and 25 m at AH3G, with the transition to cold conditions occurring approximately

110 m upstream of the terminus (Fig. 5b). Notably, in both AH1G and AH3G, temperatures approach 0°C at intermediate depths but drop below 0°C again towards the bed, suggesting the temperate zone forms a sort of core zone. As neither AH1G nor AH3G have sensors shallower than 17 m, the cold ice overlying the temperate zone is not directly sampled but is assumed to correspond to the seasonally influenced surface layer (Sect. 4.1). At Chessjen, the CTS was identified at the two higher-
365 elevation boreholes (CJ1G and CJ3TT) at depths of 17 m to 38 m, with the CTS descending towards the bed approximately 180 m upstream of the terminus, beyond which cold conditions dominate throughout (Fig. 6b). At Hohsaas, the CTS was identified at the higher-elevation borehole HS1G at a depth of 20 m and remains shallow (0 - 5 m) until approximately 150 m from the terminus, where it begins to descend towards the bed. At the northern profile L1, the ice becomes progressively thinner and colder, suggesting a more laterally extensive cold zone in this part of the glacier (Fig. 7a).

370 Despite a broadly similar thermal structure across the three sites, basal conditions differ. At Alphubel, bed temperatures remained cold throughout, reaching as low as -2.1°C at the terminus. At Chessjen, CJ2G reached the bed at 17 m depth confirming temperate conditions in early August 2025, but cooled to -0.43°C by mid-December (Supplementary Fig. S12bS15b). We interpret this as a seasonally evolving basal thermal regime: meltwater sustains temperate conditions through summer, while basal refreezing occurs once melt ceases in autumn. At Hohsaas, HS2G confirmed temperate basal conditions close to
375 the terminus despite cold conditions in the overlying ice column, though basal coverage between HS2G and HS1G remains limited.

To complement the point measurements from the boreholes with a spatially continuous view of the thermal structure, we next consider the radar profiles. The GPR profiles at all three sites are broadly consistent with the borehole-derived thermal conditions, though the degree of agreement varies. At Alphubel (Fig. 5c), the profile shows a substantially cleaner signal than
380 at the other two sites, with little scattering even in the area where boreholes confirmed temperate ice. This difference is at least partly attributable to survey timing: unlike the late-summer surveys at Chessjen and Hohsaas (August-September), the Alphubel GPR survey was conducted at the end of winter in mid-May (Supplementary Fig. S3S2). At this time of year and at 3800 m a.s.l., liquid water is expected to be at its seasonal minimum regardless of thermal state, potentially suppressing scattering that would otherwise indicate temperate ice. Still, the clean signal below the seasonally influenced layer suggests limited englacial water
385 content at depth, supporting the cold bed interpretation derived from the borehole temperatures. At Chessjen (Fig. 6c), the scattering pattern corresponds well with the interpolated temperature field. Radar scatter is high throughout the temperate zone up-glacier and diminishes towards the terminus below CJ1G. Notably, scattering extends to the glacier surface in the upper part of the profile, suggesting the ice column is temperate throughout and that the CTS detected here reflects the seasonal penetration depth of the winter cold wave rather than a stable year-round thermal boundary. At Hohsaas (Fig. 7c), a qualitatively similar
390 pattern is observed, with increased scattering around the upper borehole HS1G and a largely clean signal in the cold terminus. At both Chessjen and Hohsaas, enhanced near-bed scattering suggests that temperate basal conditions are likely more extensive than the borehole network resolves.

While polythermal glacier structures can arise through a variety of mechanisms (Irvine-Fynn et al., 2011), the consistent pattern observed across all three sites, with temperate ice confined to higher elevations and cold conditions prevailing towards

395 the terminus, points to a shared underlying control on englacial heat distribution, a point that we further examine in the following.

5 Discussion

5.1 Firn cover loss as primary control on the thermal structure of small glaciers

Temperate ice within glaciers can be generated or sustained by several heat sources, including geothermal heat flux, strain
400 heating associated with ice flow, latent heat release from refreezing of percolating meltwater, and advection of temperate ice
from upstream (Cuffey and Paterson, 2010; Wilson and Flowers, 2013; Oerlemans, 2013; Gilbert et al., 2019; Law et al.,
2021; Karlsson et al., 2021; Raspoet and Pattyn, 2025). The glaciers investigated here are all small ($<0.5 \text{ km}^2$), meaning that
internal heat production from strain heating is expected to be limited (Huss and Fischer, 2016), consistent with the low surface
displacement rates observed at all sites (Supplementary Table S14). **Advection of temperate ice S17). Since the temperate ice**
405 **identified here lies in the uppermost reaches of each glacier, where no upstream ice input exists, advection** from upstream can
likewise also be ruled out **, as temperate conditions are predominantly observed in the upper glacier where no upstream source**
exists as a means of supplying temperate ice. Geothermal heat flux in the region is modest ($< 80 \text{ mW m}^{-2}$, Medici and Rybach
1995) and, as a basal heat source, cannot account for the temperate conditions observed throughout substantial portions of the
ice column at these sites.

410 Instead, we believe that the observed thermal structure is primarily controlled by firn cover distribution and its temporal
evolution. Firn acts as a meltwater reservoir, retaining water that percolates to depth, refreezes, and releases latent heat, warming
the underlying ice and sustaining temperate conditions (Gilbert et al., 2012; Huss and Fischer, 2016; Gastaldello et al., 2025).
The densification of temperate firn into glacier ice provides an additional, if secondary, supply of temperate ice to the system.
Where firn is absent and meltwater runs off, both of these heat sources are lost and the ice can cool. This process chain
415 is supported by modelling studies showing that meltwater refreezing and entrapment can dominate the internal heat budget
of small polythermal glaciers, making the thermal structure highly sensitive to changes in surface melt and firn availability
(Gilbert et al., 2012; Wilson and Flowers, 2013; Gilbert et al., 2014).

To examine the role of firn cover as a control on englacial thermal structure, we reconstructed its evolution across the six
studied glaciers from the 1970s until today using modelled annual mass balance distribution constrained with GLAMOS mass
420 balance data (Huss et al., 2015; GLAMOS - Glacier Monitoring Switzerland, 2025). Across the six sites, firn-free durations at
the borehole locations span from less than a decade at Alphubel to more than three decades at Sex Rouge, Tortin and Corvatsch,
providing a range of exposure histories against which to assess the thermal response to firn loss (Fig. 8).

Firn-free exposure broadly reflects the thermal state across the six sites, with longer firn-free histories associated with more
advanced cooling. The clearest examples are Sex Rouge and Corvatsch, both firn-free for more than three decades, where cold
425 surface conditions prevail throughout the measured depth range. At Sex Rouge, the apparent near-surface warming relative to
previous measurements (Sect. 4.2) is most plausibly explained by surface ablation removing cold ice from above rather than a




Figure 5. Englacial temperature structure and radar reflectivity along Alphubel profile L4 (for profile location, see Fig. 1a). (a) Interpolated englacial temperature field derived from borehole measurements. Temperatures at AH1G–AH3G and AH3TT are averaged over the period 10.08.–16.09.2025; AH1TT–AH2TT over September 2024–September 2025. Modelled firm cover extent is shown for 2010 (orange) and 2025 (blue). Whiskers along the bed denote ± 5 m bedrock-depth uncertainty. **Black ticks along each borehole mark the depth of zero annual temperature amplitude (ZAA), below which seasonal temperature variations become negligible.** (b) Vertical temperature profiles for AH1G, AH2G, AH3G, and AH3TT all boreholes; the **dashed line grey-shaded area marks the temperatures above 0°C isotherm, and the solid black line marks the ZAA depth.** (c) Depth radargram along profile L4 with borehole positions and the CTS derived from the interpolated temperature field.




Figure 6. Englacial temperature structure and radar reflectivity along Chessjen profile L47 (for profile location, see Fig. 1b). (a) Interpolated englacial temperature field derived from borehole measurements. Temperatures at CJ1G, CJ2G, CJ3TT, and CJ4TT are averaged over the period 08.08.–03.09.2025; CJ1TT and CJ2TT over September 2024–September 2025. Modelled firn cover extent is shown for 2025. In the year 2010 no firn was present at Chessjen. **Black ticks along each borehole mark the ZAA depth.** (b) Vertical temperature profiles for all boreholes; the **dashed line grey-shaded area marks the temperatures above 0°C isotherm**, and the **solid black line marks the ZAA depth**. (c) Depth radargram along profile L47 with borehole positions and the CTS derived from the interpolated temperature field.




Figure 7. Englacial temperature structure and radar reflectivity at Hohsaas. (a) Interpolated temperature fields along profiles L1 (left) and L2 (right) (for profile location, see Fig. 1c). Temperatures at HS1G, HS2G, and HS3G are averaged over the period 13.08.–27.09.2025; HS1TT and HS2TT over September 2024–September 2025. Whiskers along the bed denote ± 5 m bedrock-depth uncertainty. Black ticks along each borehole mark the ZAA depth. (b) Vertical temperature profiles for all boreholes (HS1G, HS2G, HS3G, HS1TT, HS2TT); the dashed line grey-shaded area marks the temperatures above 0°C isotherm, and the solid black line marks the ZAA depth. (c) Depth radargram along profile L2 with borehole positions and the CTS derived from the interpolated temperature field.




Figure 8. Number of years a given area was firm free, mapped across all six study glaciers and draped over a shaded-relief DEM (Swisstopo, 2023a). Darker colors indicate longer firm-free periods (prior to 2025). The 2025 firm cover extent is indicated in light blue. Grid cells which have been part of the ablation area over the entire time series (1970-2025) are masked. Borehole colors indicate thermal regime: dark blue = cold-based cold (entire ice column frozen to the bed); light blue = mostly coldpolythermal; red = mostly temperate. The blue line marks the SGI2023 outline.

genuine change in thermal forcing. At Tortin, where borehole locations have similarly been firm-free for more than 30 years, shallow measurements alone are insufficient to characterise the full thermal structure.

At Chessjen and Hohsaas, the spatial pattern of time since firm loss broadly reflects the glaciers glacier's thermal structure, with the CTS deepest in areas that lost firm earliest, suggesting the up-glacier retreat of the firm line has left a spatial imprint on the thermal regime. At both sites, firm-free duration increases down-glacier, ranging from approximately 5-10 15-20 years at the uppermost borehole positions to 15-20 25-30 years towards the terminus. The persistence of temperate conditions despite firm-free periods of up to a decade at the uppermost borehole positions can be attributed to several factors. First, the thermal

diffusion timescale for That temperate ice nonetheless persists beneath the upper reaches, despite one to two decades without
 435 firn, reflects the slow pace of conductive cooling combined with continued heat input. Conductive cooling of a temperate layer
 of 20-30 m thickness may well be on the order of 10-15 years tens of meters thick proceeds on a timescale of decades (Cuffey
 and Paterson, 2010; Gilbert et al., 2012, 2014), meaning the cooling front may only recently have begun to penetrate to the
 observed CTS depths . ; over the firn-free period, the cold front driven by firn loss would be expected to penetrate roughly 20–
 25 m, broadly consistent with, though slightly deeper than, the 17-20 m CTS depths observed. That temperate ice is preserved
 440 to somewhat shallower depths than conduction alone would predict points to additional heat input. Two mechanisms contribute.
 First, the downward migration of the cold front is slowed by the latent heat released as residual pore water in the temperate ice
 refreezes ahead of it, so that temperate conditions persist longer than pure conduction would imply. Second, meltwater delivery
 to the firn-free temperate zone has likely not ceasedentirely. While the mechanism remains incompletely resolved, : at both
 sites, patches of firn persist in the highest reaches of the glacier, up-glacier of the borehole network (Fig. 8), and bergschrunds
 445 and crevasses at both sites provide potential provide englacial entry points for meltwater draining from thesestill firn-covered
 areas aboveinto the temperate zone below, sustaining it through latent-heat release. At Chessjen, the transient temperature jump
 observed in mid-July (Sect. 4.1) provides indirect support for the notion that such pathways are remain active.

The situation at Alphubel is more nuanced. While the firn-free terminus is cold and the firn-covered upper glacier is tem-
 perate, consistent with the pattern seen at Chessjen and Hohsaas, the underlying thermal history likely differs. The largely
 450 cold-based terminus is difficult to reconcile with a firn-free period of only 5-7 5-10 years, and the cold-temperate-cold vertical
 structure observed, where a temperate layer is sandwiched between a cold surface layer and a cold bed, suggests a fundamen-
 tally different thermal history at this site. Historical imagery from 1930 shows Alphubel as a much thicker hanging glacier
 with a prominent calving front (Supplementary Fig. S15S18), a morphology more characteristic of cold or polythermal high-
 elevation hanging glaciers with limited basal sliding (Gilbert et al., 2015). This is in contrast to the much larger valley glacier
 455 systems represented by Chessjen and Hohsaas, as documented in the historical imagery. We therefore interpret the cold termi-
 nus as largely predating the current period of firn loss, with the temperate core in the upper glacier representing a relatively
 recent feature sustained by meltwater percolation from the remaining firn cover above. A similar transition from a cold to a
 polythermal regime has been documented at Whymper hanging glacier on the Grande Jorasses south face (Troilo et al., 2022).

5.2 Implications for our understanding of Alpine glacier thermal regimes

460 The conceptual diagram in Fig. 9 illustrates our interpretation of how the thermal structure of small Alpine glaciers may have
 evolved in response to climate change and the associated firn cover retreat over the last decades. In the past (e.g., 1980s), the
 glaciers studied here were likely entirely temperate. Extensive firn cover, sustained by a cooler climate, provided an effective
 heat source through meltwater percolation and latent heat release upon refreezing (Gilbert et al., 2012; Huss and Fischer,
 2016; Gastaldello et al., 2025). Active glacier dynamics promoted crevassing, creating englacial entry points for meltwater and
 465 further supporting temperate conditions throughout the ice column via cryo-hydrologic warming (Phillips et al., 2010; Gilbert
 et al., 2019). Greater ice flow velocities also enhanced strain heating, providing an additional internal heat source (Cuffey and
 Paterson, 2010).




Figure 9. Conceptual illustration of the thermal consequences of firn cover loss under glacier recession, comparing the 1980s with present-day (2026) conditions.

Sustained atmospheric warming has driven firn retreat, progressively reducing latent heat input from meltwater percolation and refreezing. Where firn is lost, meltwater runs off at the surface rather than percolating into the pore space. The associated latent heat release upon refreezing is therefore lost (Wilson and Flowers, 2013; van Pelt et al., 2016). Simultaneously, ongoing glacier recession reduces ice flow velocities and with it strain heating, while also limiting crevassing and the availability of englacial meltwater pathways. In the absence of firn, both the loss of latent heat from refreezing and the reduced insulation allow the winter cold wave to penetrate the ice surface more effectively, initiating the development of a cold surface layer and, over time, cold ice. The thermal response to firn loss is neither instantaneous nor spatially uniform. This is illustrated by the contrast between profiles L1 and L2 at Hohsaas, which are only 20-30 m apart but show markedly different thermal structures (Fig. 7a). At L1, ice thickness (15-17 m) is comparable to the seasonally-influenced layer, leaving insufficient ice thickness for a temperate core to persist even where firn-loss history is similar to L2; at L2, the greater ice thickness (>30 m) provides enough insulated depth for temperate ice to remain despite ongoing cooling. This contrast highlights how ice thickness modulates the pace and completeness of thermal response to firn loss, independent of the surface forcing itself. Elevation, topography, residual meltwater supply, and the thermal inertia of the ice all modulate how quickly and how completely cold conditions develop following firn loss. In some cases, atmospheric warming may already offset the cooling initiated by firn loss, for instance where accelerated surface ablation removes cold ice faster than the winter cold wave can produce it. Alphubel may represent an exception to this thermal history, reminding us of the complexity of thermal responses to climate change. As a

high-elevation hanging glacier, it likely harboured cold or polythermal conditions at its terminus even historically, in contrast to
485 Chessjen and Hohnsaa which, as much larger valley glacier systems, were likely fully temperate (Supplementary Fig. S15S18).

The results presented here suggest that polythermal structures may be more widespread among small Alpine glaciers than previously recognised. The existing observational ice temperature record is strongly biased toward larger glaciers (Haeberli, 1976) and high-altitude sites (Haeberli and Alean, 1985; Haeberli and Funk, 1991; Hoelzle et al., 2011), largely overlooking smaller glaciers at lower elevations where firn is progressively being lost and polythermal structures may be actively emerging.
490 Regional-scale thermal modelling, informed by the growing body of englacial temperature observations such as those compiled in the glenlat database (Jacquemart and Welty, 2024) and the records presented here, offers a path toward assessing the current distribution of polythermal structures and projecting their evolution under continued atmospheric warming. Such projections are particularly relevant given the hazard implications of polythermal configurations.

5.3 Possible hazard implications

Polythermal glaciers are not dangerous per se. The thermal structures identified in this study are however consistent with a configuration known to promote meltwater accumulation beyond drainage capacity: a frozen terminus combined with a temperate upper glacier (?Wadham et al., 2001; Vincent et al., 2012; Jacquemart et al., 2024). During drilling at Alphubel, the only site with direct evidence for a fully frozen terminus, pressurised water escaped from borehole AH1TT at approximately 15 m depth, producing a water fountain 3 to 5 m high. This indicates that water pressure locally exceeded the ice overburden,
500 implying that meltwater had been trapped sub- or englacially and couldn't drain. If such conditions occur at the glacier bed, basal friction may be substantially reduced locally (Iken and Bindshadler, 1986; Cuffey and Paterson, 2010). A broadly similar configuration, combining a cold glacier tongue with pressurised englacial water, has been discussed in the context of the 2022 Marmolada collapse (Bondesan and Francese, 2023; Olivieri and Bettanini, 2023). Whether similar configurations exist at other small Alpine glaciers remains unknown, but the number of potentially affected sites may be substantial.

To give a rough sense of how many glaciers share the size and elevation characteristics of our study sites, we filtered the SGI2023 (Huss et al., 2025) for glaciers smaller than 0.5 km² within an elevation range of 2900 m a.s.l. to 3800 m a.s.l. The choice of 2900 m a.s.l. as a lower bound roughly excludes elevations where surface ablation dominates over cold ice formation, as suggested by our observations at Sex Rouge. Approximately 515 glaciers in Switzerland meet these criteria, a number that is likely an underestimate, as small hanging glaciers at high elevations are frequently mapped as part of glacier
510 accumulation areas. This underscores the need for a systematic assessment of the thermal structure of small Alpine glaciers and its implications for hazard potential.

6 Conclusions

In this study, we presented englacial temperature data for six small glaciers (< 0.5 km²) in the Swiss Alps. The observations span an elevation range of 2700–3800 m a.s.l., directly addressing the lack of data in mid- to lower-elevation ablation areas.

515 [key gap in the existing ablation-zone record](#). Our dataset combines borehole thermometry from a total of 23 boreholes with ground-penetrating radar surveys to assess glacier thermal structures.

Polythermal conditions were identified in three glaciers, Alphubel, Chessjen, and Hohsaas, with the CTS occurring at depths of 17 to 38 m. Beneath the cold seasonal layer, ice temperatures ranged from temperate conditions to 2.1 °C. A consistent pattern emerged across these three glaciers, with a generally temperate zone at higher elevations transitioning into fully or
520 partially cold-based glacier termini. However, evidence for a consistently frozen glacier terminus was only found at Alphubel, the highest-elevation glacier in our dataset. At the three additional shallow-drilled sites, Corvatsch appears predominantly cold-based, while Sex Rouge likely retains a polythermal structure with a cold surface layer overlying temperate basal ice (Fischer, 2018), though this remains poorly constrained at depth. At Tortin, sensors were limited to the seasonally influenced surface layer, precluding any assessment of the deeper thermal structure. Our GPR surveys confirmed the spatial thermal
525 patterns inferred from boreholes, with low scattering in cold ice and enhanced scattering in temperate zones. Enhanced near-bed scattering at Chessjen and Hohsaas further suggests that temperate basal conditions may extend beyond what the borehole network resolves.

The fact that three out of six glaciers were confirmed polythermal, with a fourth, likely retaining its historically documented polythermal structure (Fischer, 2018), suggests that this thermal regime may be more widespread among small Alpine glaciers
530 than previously recognised. A systematic regional-scale assessment of their distribution and hazard implications represents an important next step, and our dataset provides essential constraints for the thermal modelling efforts needed to address it.

Code and data availability. The source code used to process our data is publicly available on GitHub under https://github.com/sdrocer/polythermal_swiss_glaciers (Beer, 2026). The borehole thermometry data has been added to the Global Englacial Temperature database (glenglat) available on github under <https://github.com/mjacqu/glenglat> (Jacquemart and Welty, 2024). The ground based ground-penetrating
535 radar data presented in this study are available in the ETH Research Collection at <https://doi.org/10.3929/ethz-c-000799146> (Beer et al., 2026).

Appendix A: Borehole Inventory

Table A1. Overview of the boreholes drilled in this study, showing drilling date, borehole coordinates (X, Y; EPSG:2056), elevation (Z), and sensor depth at the time of installation. The borehole ID suffix indicates the sensor type (TT = Tinytag loggers; G = Geoprecision loggers). Boreholes marked with * reached the bedrock.

#	Borehole ID	Date	X [m]	Y [m]	Z [m a.s.l.]	Depth [m]
Alphubel South						
1	AH1G*	06/08/2025	2633226.81	1100604.12	3802	50.6*
2	AH2G*	06/08/2025	2633183.50	1100551.82	3759	20.3*
3	AH3G*	06/08/2025	2633245.17	1100634.30	3820	58.3*
4	AH1TT	21/08/2024	2633249.72	1100537.76	3775	15.0
5	AH2TT	21/08/2024	2633251.63	1100564.25	3789	14.0
6	AH3TT	06/08/2025	2633203.23	1100579.67	3781	17.0
Chessjengletscher						
7	CJ1G*	08/08/2025	2636970.56	1101731.07	3031	38.3*
8	CJ2G*	08/08/2025	2636963.22	1101802.26	3005	17.0*
9	CJ1TT	09/08/2024	2636956.76	1101858.36	2983	11.8
10	CJ2TT	09/08/2024	2636976.64	1101745.91	3027	13.5
11	CJ3TT	08/08/2025	2636969.90	1101687.21	3046	20.0
12	CJ4TT	08/08/2025	2637011.63	1101794.98	3004	20.0
Hohsaasgletscher						
13	HS1G*	13/08/2025	2643544.74	1109819.94	3349	29.0*
14	HS2G*	13/08/2025	2643372.47	1109737.94	3278	21.5*
15	HS3G	13/08/2025	2643420.68	1109732.65	3292	21.5
16	HS1TT	08/08/2024	2643397.51	1109785.68	3284	15.6
17	HS2TT	08/08/2024	2643519.71	1109839.68	3337	15.4
Glacier du Sex Rouge						
18	SR1TT	06/08/2024	2582609.56	1130598.61	2787	15.3
19	SR2TT	06/08/2024	2582586.65	1130751.59	2760	14.5
Glacier de Tortin						
20	GT1TT	07/08/2024	2589854.86	1103667.77	2858	9.2
21	GT2TT	07/08/2024	2589986.76	1103495.13	2891	15.25
Vadret dal Corvatsch						
22	CV1TT*	28/08/2024	2783252.23	1143066.66	3271	7.0*
23	CV2TT*	28/08/2024	2783141.08	1143188.93	3304	9.3*

Author contributions. Janosch Beer: Conceptualization, Data curation, Formal analysis, Investigation, Project administration, Visualization, Writing – original draft, Writing – review and editing. Mylène Jacquemart: Conceptualization, Supervision, Writing – review and editing.
540 Matthias Huss: Conceptualization, Software (firm cover reconstruction), Supervision, Writing – review and editing. Ilaria Santin: Investigation, Data curation (ground-based GPR post-processing), Writing – review and editing. Gabriela Clara Racz: Investigation, Methodology, Data curation (UAV-based GPR survey and post-processing, Hohsaas), Writing – review and editing. Christophe Ogier: Investigation, Data curation (ground-based GPR post-processing, Alphubel), Writing – review and editing. Saskia Gindraux: Investigation, Financing, Writing – review and editing. Leo Hösli: Investigation, Writing – review and editing. Raphael Moser: Investigation. James Irving: Conceptualization,
545 Methodology, Investigation (UAV-based GPR), Writing – review and editing. Mauro Fischer: Resources (historical englacial temperature data, Sex Rouge), Investigation, Writing – review and editing. Daniel Farinotti: Conceptualization, Investigation, Financing, Supervision, Writing – review and editing.

Competing interests. Daniel Farinotti is a member of the editorial board of The Cryosphere. The remaining authors declare that they have no conflict of interest.

550 *Acknowledgements.* This study draws on an extensive series of field campaigns that would not have been possible without the dedication, support, and hard work of all those who contributed. The first author made use of Claude (Anthropic) as an AI-assisted tool to support language editing and code development. All scientific interpretations, analyses, and conclusions are the sole responsibility of the authors. The first author would like to thank Dominika Teluchová for her personal support throughout this work.

Financial support. This study was primarily supported by internal research funds of ETH Zurich and WSL. Fieldwork was additionally
555 supported by the Section of Natural Hazards of the Canton of Valais, Switzerland. Previous borehole temperature measurements at Sex Rouge first published by Fischer (2018) received financial support from the Swiss National Science Foundation (SNSF), grant no. 200021 137586.

References

- Agassiz, L.: Nouvelles études et expériences sur les glaciers actuels, V. Masson, 1847.
- 560 Ambach, W. and Eisner, H.: Analysis of a 20 m. Firn Pit on the Kesselwandferner (Ötztal Alps), *Journal of Glaciology*, 6, 223–231, <https://doi.org/10.3189/S0022143000019237>, 1966.
- Beer, J.: Code for: Polythermal Conditions in Small Glaciers in the Swiss Alps, 2026.
- Beer, J., Santin, I., and Ogier, C.: GPR Data for Beer et al. 2026: Radar Dataset of Glacier Du Sex Rouge, Glacier de Tortin, Vadret Dal Corvatsch, Alphubel South, Chessjengletscher, Hohsaasgletscher, <https://doi.org/10.3929/ethz-c-000799146>, 2026.
- 565 Bondesan, A. and Francese, R. G.: The Climate-Driven Disaster of the Marmolada Glacier (Italy), *Geomorphology*, 431, 108 687, <https://doi.org/10.1016/j.geomorph.2023.108687>, 2023.
- Boon, S. and Sharp, M.: The Role of Hydrologically-Driven Ice Fracture in Drainage System Evolution on an Arctic Glacier, *Geophysical Research Letters*, 30, <https://doi.org/10.1029/2003GL018034>, 2003.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Academic Press, ISBN 978-0-08-091912-6, 2010.
- 570 Delcourt, C., Van Lieffring, B., Nolan, M., and Pattyn, F.: The Climate Memory of an Arctic Polythermal Glacier, *Journal of Glaciology*, 59, 1084–1092, <https://doi.org/10.3189/2013JoJ12J109>, 2013.
- Faillietaz, J., Funk, M., and Vincent, C.: Avalanching Glacier Instabilities: Review on Processes and Early Warning Perspectives, *Reviews of Geophysics*, 53, 203–224, <https://doi.org/10.1002/2014RG000466>, 2015.
- Fischer, M.: Understanding the Response of Very Small Glaciers in the Swiss Alps to Climate Change, Ph.D. thesis, 2018.
- 575 Flowers, G. E. and Clarke, G. K. C.: A Multicomponent Coupled Model of Glacier Hydrology 1. Theory and Synthetic Examples, *Journal of Geophysical Research: Solid Earth*, 107, ECV 9–1–ECV 9–17, <https://doi.org/10.1029/2001JB001122>, 2002.
- Forte, E., Gutgesell, P., Securo, A., Marcer, M., Citterio, M., Machguth, H., and Colucci, R. R.: Comparing GPR with Ice Thickness and Thermal Models: Insights from Two Polythermal Glaciers in West Greenland, *Journal of Glaciology*, 71, e97, <https://doi.org/10.1017/jog.2025.10067>, 2025.
- 580 Gastaldello, M., Mattea, E., Hoelzle, M., and Machguth, H.: Modelling Cold Firn Evolution at Colle Gnifetti, Swiss/Italian Alps, *The Cryosphere*, 19, 2983–3008, <https://doi.org/10.5194/tc-19-2983-2025>, 2025.
- Geolitix Technologies Inc.: Geolitix: Cloud-hosted Platform for GPR Data Visualization and Interpretation, 2020.
- Gilbert, A., Vincent, C., Wagnon, P., Thibert, E., and Rabatel, A.: The Influence of Snow Cover Thickness on the Thermal Regime of Tte Rouse Glacier (Mont Blanc Range, 3200 m a.s.l.): Consequences for Outburst Flood Hazards and Glacier Response to Climate Change, *Journal of Geophysical Research*, 117, <https://doi.org/10.1029/2011JF002258>, 2012.
- 585 Gilbert, A., Gagliardini, O., Vincent, C., and Wagnon, P.: A 3-D Thermal Regime Model Suitable for Cold Accumulation Zones of Polythermal Mountain Glaciers, *Journal of Geophysical Research: Earth Surface*, 119, 1876–1893, <https://doi.org/10.1002/2014JF003199>, 2014.
- Gilbert, A., Vincent, C., Gagliardini, O., Krug, J., and Berthier, E.: Assessment of Thermal Change in Cold Avalanching Glaciers in Relation to Climate Warming, *Geophysical Research Letters*, 42, 6382–6390, <https://doi.org/10.1002/2015GL064838>, 2015.
- 590 Gilbert, A., Leinss, S., Kargel, J., Kääb, A., Gascoin, S., Leonard, G., Berthier, E., Karki, A., and Yao, T.: Mechanisms Leading to the 2016 Giant Twin Glacier Collapses, Aru Range, Tibet, *The Cryosphere*, 12, 2883–2900, <https://doi.org/10.5194/tc-12-2883-2018>, 2018.
- Gilbert, A., Sinisalo, A., Gurung, T. R., Fujita, K., Maharjan, S. B., Sherpa, T. C., and Fukuda, T.: The Influence of Water Percolation through Crevasses on the Thermal Regime of Himalayan Mountain Glaciers, <https://doi.org/10.5194/tc-2019-172>, 2019.

- 595 GLAMOS - Glacier Monitoring Switzerland: Swiss Glacier Mass Balance (Release 2025), <https://doi.org/10.18750/MASSBALANCE.2025.R2025>, 2025.
- Haerberli, W.: Eistemperaturen in Den Alpen, Verlag nicht ermittelbar, 1976.
- Haerberli, W. and Alean, J.: Temperature and Accumulation of High Altitude Firn in the Alps, *Annals of Glaciology*, 6, 161–163, <https://doi.org/10.3189/1985AoG6-1-161-163>, 1985.
- 600 Haerberli, W. and Funk, M.: Borehole Temperatures at the Colle Gnifetti Core-Drilling Site (Monte Rosa, Swiss Alps), *Journal of Glaciology*, 37, 37–46, <https://doi.org/10.3189/S0022143000042775>, 1991.
- Haerberli, W., Frauenfelder, R., Käab, A., and Wagner, S.: Characteristics and Potential Climatic Significance of “Miniature Ice Caps” (Crest- and Cornice-Type Low-Altitude Ice Archives), *Journal of Glaciology*, 50, 129–136, <https://doi.org/10.3189/172756504781830330>, 2004.
- Hardy, R. L.: Multiquadric Equations of Topography and Other Irregular Surfaces, <https://doi.org/10.1029/JB076i008p01905>, 1971.
- 605 Heucke, E.: A Light Portable Steam-Driven Ice Drill Suitable for Drilling Holes in Ice and Firn, *Geografiska Annaler. Series A, Physical Geography*, 81, 603–609, 1999.
- Hock, R.: A Distributed Temperature-Index Ice- and Snowmelt Model Including Potential Direct Solar Radiation, *Journal of Glaciology*, 45, 101–111, <https://doi.org/10.3189/S0022143000003087>, 1999.
- Hock, R.: Glacier Melt: A Review of Processes and Their Modelling, *Progress in Physical Geography: Earth and Environment*, 29, 362–391, <https://doi.org/10.1191/0309133305pp453ra>, 2005.
- 610 Hoelzle, M., Darms, G., Lüthi, M. P., and Suter, S.: Evidence of Accelerated Englacial Warming in the Monte Rosa Area, Switzerland/Italy, *The Cryosphere*, 5, 231–243, <https://doi.org/10.5194/tc-5-231-2011>, 2011.
- Huggel, C., Haerberli, W., Käab, A., Bieri, D., and Richardson, S.: An Assessment Procedure for Glacial Hazards in the Swiss Alps, *Canadian Geotechnical Journal*, 41, 1068–1083, <https://doi.org/10.1139/t04-053>, 2004.
- 615 Hugli, F. J.: Ueber das Wesen der Gletscher und Winterreise in das Eismeer, Gotta’scher Verlag, 1842.
- Humphrey, N. and Echelmeyer, K.: Hot-Water Drilling And Bore-hole Closure In Cold Ice, *Journal of Glaciology*, 36, 287–298, <https://doi.org/10.3189/002214390793701354>, 1990.
- Huss, M. and Fischer, M.: Sensitivity of Very Small Glaciers in the Swiss Alps to Future Climate Change, *Frontiers in Earth Science*, 4, 2016.
- 620 Huss, M., Dhulst, L., and Bauder, A.: New Long-Term Mass-Balance Series for the Swiss Alps, *Journal of Glaciology*, 61, 551–562, <https://doi.org/10.3189/2015JoG15J015>, 2015.
- Huss, M., Bauder, A., Linsbauer, A., Gabbi, J., Kappenberger, G., Steinegger, U., and Farinotti, D.: More than a Century of Direct Glacier Mass-Balance Observations on Claridenfirn, Switzerland, *Journal of Glaciology*, 67, 697–713, <https://doi.org/10.1017/jog.2021.22>, 2021.
- Huss, M., Bauder, A., Hodel, E., and Linsbauer, A.: Swiss Glacier Bulletin 2024: Annual Report #145 on Glacier Observation in Switzerland, Tech. rep., GLAMOS - Glacier Monitoring Switzerland, https://doi.org/10.18752/GLBULLETIN_2024, 2025.
- 625 Huss, M., Bauder, A., Barandun, M., Hodel, E., and Linsbauer, A.: Swiss Glacier Bulletin 2025, Tech. rep., GLAMOS - Glacier Monitoring Switzerland, https://doi.org/10.18752/GLBULLETIN_2025, 2026.
- Iken, A. and Bindshadler, R. A.: Combined Measurements of Subglacial Water Pressure and Surface Velocity of Finde-
lengletscher, Switzerland: Conclusions about Drainage System and Sliding Mechanism, *Journal of Glaciology*, 32, 101–119, <https://doi.org/10.3189/S0022143000006936>, 1986.
- 630

- Irvine-Fynn, T. D. L., Moorman, B. J., Williams, J. L. M., and Walter, F. S. A.: Seasonal Changes in Ground-Penetrating Radar Signature Observed at a Polythermal Glacier, Bylot Island, Canada, *Earth Surface Processes and Landforms*, 31, 892–909, <https://doi.org/10.1002/esp.1299>, 2006.
- Irvine-Fynn, T. D. L., Hodson, A. J., Moorman, B. J., Vatne, G., and Hubbard, A. L.: Polythermal Glacier Hydrology: A Review, *Reviews of Geophysics*, 49, <https://doi.org/10.1029/2010RG000350>, 2011.
- Jacquemart, M. and Welty, E.: Glenglat: A Database of Global Englacial Temperatures, *Earth System Science Data Discussions*, pp. 1–47, <https://doi.org/10.5194/essd-2024-249>, 2024.
- Jacquemart, M., Loso, M., Leopold, M., Welty, E., Berthier, E., Hansen, J. S., Sykes, J., and Tiampo, K.: What Drives Large-Scale Glacier Detachments? Insights from Flat Creek Glacier, St. Elias Mountains, Alaska, *Geology*, 48, 703–707, <https://doi.org/10.1130/G47211.1>, 2020.
- Jacquemart, M., Weber, S., Chiarle, M., Chmiel, M., Cicoira, A., Corona, C., Eckert, N., Gaume, J., Giacona, F., Hirschberg, J., Kaitna, R., Magnin, F., Mayer, S., Moos, C., van Herwijnen, A., and Stoffel, M.: Detecting the Impact of Climate Change on Alpine Mass Movements in Observational Records from the European Alps, *Earth-Science Reviews*, 258, 104 886, <https://doi.org/10.1016/j.earscirev.2024.104886>, 2024.
- Jansson, P.: Dynamics and Hydrology of a Small Polythermal Valley Glacier, *Geografiska Annaler: Series A, Physical Geography*, 78, 171–180, <https://doi.org/10.1080/04353676.1996.11880463>, 1996.
- Karlsson, N. B., Solgaard, A. M., Mankoff, K. D., Gillet-Chaulet, F., MacGregor, J. A., Box, J. E., Citterio, M., Colgan, W. T., Larsen, S. H., Kjeldsen, K. K., Korsgaard, N. J., Benn, D. I., Hewitt, I. J., and Fausto, R. S.: A First Constraint on Basal Melt-Water Production of the Greenland Ice Sheet, *Nature Communications*, 12, 3461, <https://doi.org/10.1038/s41467-021-23739-z>, 2021.
- Kavanaugh, J. L. and Clarke, G. K. C.: Abrupt Glacier Motion and Reorganization of Basal Shear Stress Following the Establishment of a Connected Drainage System, *Journal of Glaciology*, 47, 472–480, <https://doi.org/10.3189/172756501781831972>, 2001.
- Law, R., Christoffersen, P., Hubbard, B., Doyle, S. H., Chudley, T. R., Schoonman, C. M., Bougamont, M., des Tombe, B., Schilperoort, B., Kechavarzi, C., Booth, A., and Young, T. J.: Thermodynamics of a Fast-Moving Greenlandic Outlet Glacier Revealed by Fiber-Optic Distributed Temperature Sensing, *Science Advances*, 7, eabe7136, <https://doi.org/10.1126/sciadv.abe7136>, 2021.
- Mattea, E.: Measuring and Modelling Changes in the Firn at Colle Gnifetti, 4400 m a.s.l., Swiss Alps, Ph.D. thesis, University of Fribourg, Fribourg, 2020.
- Mattea, E., Machguth, H., Kronenberg, M., van Pelt, W., Bassi, M., and Hoelzle, M.: Firn Changes at Colle Gnifetti Revealed with a High-Resolution Process-Based Physical Model Approach, *The Cryosphere*, 15, 3181–3205, <https://doi.org/10.5194/tc-15-3181-2021>, 2021.
- Medici, F. and Rybach, L.: Geothermal Map of Switzerland 1995 (Heat Flow Density), Tech. Rep. 30, Commission Suisse de Géophysique, Zurich, Switzerland, 1995.
- Oerlemans, J.: A Note on the Water Budget of Temperate Glaciers, *The Cryosphere*, 7, 1557–1564, <https://doi.org/10.5194/tc-7-1557-2013>, 2013.
- Olivieri, L. and Bettanini, C.: Preliminary Observation of Marmolada Glacier Collapse of July 2022 with Space-Based Cameras, *Remote Sensing Letters*, 14, 21–29, <https://doi.org/10.1080/2150704X.2022.2152754>, 2023.
- Pavlova, P. A., Schmid, P., Bogdal, C., Steinlin, C., Jenk, T. M., and Schwikowski, M.: Polychlorinated Biphenyls in Glaciers. 1. Deposition History from an Alpine Ice Core, *Environmental Science & Technology*, 48, 7842–7848, <https://doi.org/10.1021/es5017922>, 2014.
- Pettersson, R., Jansson, P., and Blatter, H.: Spatial Variability in Water Content at the Cold-Temperate Transition Surface of the Polythermal Storglaciären, Sweden, *Journal of Geophysical Research: Earth Surface*, 109, <https://doi.org/10.1029/2003JF000110>, 2004.

- Phillips, T., Rajaram, H., and Steffen, K.: Cryo-Hydrologic Warming: A Potential Mechanism for Rapid Thermal Response of Ice Sheets, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL044397>, 2010.
- Raspoet, O. and Pattyn, F.: Estimates of Basal and Englacial Thermal Conditions of the Antarctic Ice Sheet, *Journal of Glaciology*, 71, e104, <https://doi.org/10.1017/jog.2025.10087>, 2025.
- Ruols, B., Baron, L., and Irving, J.: Development of a Drone-Based Ground-Penetrating Radar System for Efficient and Safe 3D and 4D Surveying of Alpine Glaciers, *Journal of Glaciology*, 69, 2087–2098, <https://doi.org/10.1017/jog.2023.83>, 2023.
- Sevestre, H., Benn, D. I., Hulton, N. R. J., and Bælum, K.: Thermal Structure of Svalbard Glaciers and Implications for Thermal Switch Models of Glacier Surging, *Journal of Geophysical Research: Earth Surface*, 120, 2220–2236, <https://doi.org/10.1002/2015JF003517>, 2015.
- Suter, S. and Hoelzle, M.: Cold Firn in the Mont Blanc and Monte Rosa Areas, European Alps: Spatial Distribution and Statistical Models, *Annals of Glaciology*, 35, 9–18, <https://doi.org/10.3189/172756402781817059>, 2002.
- Swisstopo: SWISSALTI3D – The Digital Elevation Model of Switzerland, <https://www.swisstopo.admin.ch>, 2023a.
- Swisstopo: SWISSIMAGE 10 Cm, <https://map.geo.admin.ch>, 2023b.
- Swisstopo: SWISSIMAGE: Orthophoto Mosaic of Switzerland, 2024.
- Troilo, F., Paolo, P., Gottardelli, S., Mondardini, L., Giordan, D., Dematteis, N., Piard, L., Gagliardini, O., Gilbert, A., and Vincent, C.: Basal Thermal Regime Investigations at Whymper Hanging Glacier (Aosta Valley – Italy), Tech. Rep. EGU22-7208, Copernicus Meetings, <https://doi.org/10.5194/egusphere-egu22-7208>, 2022.
- van Pelt, W. J. J., Pohjola, V. A., and Reijmer, C. H.: The Changing Impact of Snow Conditions and Refreezing on the Mass Balance of an Idealized Svalbard Glacier, *Frontiers in Earth Science*, 4, <https://doi.org/10.3389/feart.2016.00102>, 2016.
- Vincent, C., Garambois, S., Thibert, E., Lefèvre, E., Meur, E. L., and Six, D.: Origin of the Outburst Flood from Glacier de Tête Rousse in 1892 (Mont Blanc Area, France), *Journal of Glaciology*, 56, 688–698, <https://doi.org/10.3189/002214310793146188>, 2010.
- Vincent, C., Descloîtres, M., Garambois, S., Legchenko, A., Guyard, H., and Gilbert, A.: Detection of a Subglacial Lake in Glacier de Tête Rousse (Mont Blanc Area, France), *Journal of Glaciology*, 58, 866–878, <https://doi.org/10.3189/2012JoG11J179>, 2012.
- Vincent, C., Gilbert, A., Jourdain, B., Piard, L., Ginot, P., Mikhalenko, V., Possenti, P., Le Meur, E., Laarman, O., and Six, D.: Strong Changes in Englacial Temperatures despite Insignificant Changes in Ice Thickness at Dôme Du Goûter Glacier (Mont Blanc Area), *The Cryosphere*, 14, 925–934, <https://doi.org/10.5194/tc-14-925-2020>, 2020.
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M., Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E., Carey, C. J., Polat, İ., Feng, Y., Moore, E. W., VanderPlas, J., Laxalde, D., Perktold, J., Cimrman, R., Henriksen, I., Quintero, E. A., Harris, C. R., Archibald, A. M., Ribeiro, A. H., Pedregosa, F., and van Mulbregt, P.: SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python, *Nature Methods*, 17, 261–272, <https://doi.org/10.1038/s41592-019-0686-2>, 2020.
- Wadham, J. L., Hodgkins, R., Cooper, R. J., and Tranter, M.: Evidence for Seasonal Subglacial Outburst Events at a Polythermal Glacier, Finsterwalderbreen, Svalbard, *Hydrological Processes*, 15, 2259–2280, <https://doi.org/10.1002/hyp.178>, 2001.
- Wilson, N. J. and Flowers, G. E.: Environmental Controls on the Thermal Structure of Alpine Glaciers, *The Cryosphere*, 7, 167–182, <https://doi.org/10.5194/tc-7-167-2013>, 2013.
- Wilson, N. J., Flowers, G. E., and Mingo, L.: Comparison of Thermal Structure and Evolution between Neighboring Subarctic Glaciers, *Journal of Geophysical Research: Earth Surface*, 118, 1443–1459, <https://doi.org/10.1002/jgrf.20096>, 2013.