

Chlorine enhances nocturnal heterogeneous uptake of NO₂ in coastal atmosphere under sea-land breeze circulation

Ziyi Lin^{1,2,3}, Xiuwen Yong^{1,2,4}, Lingjun Li^{1,2,3}, Yuping Chen^{1,2,3}, Lingling Xu^{1,2,3*}, Xiaoting Ji^{1,2,3}, Chen Yang^{1,2,3}, Keran Zhang^{1,2,3}, Feng Zhang^{1,2}, Ziyang Chen^{1,2,3}, Gaojie Chen^{1,2,3}, Xiaolong Fan^{1,2}, Mengren Li^{1,2}, Jinsheng Chen^{1,2,3*}

Affiliations:

¹State Key Laboratory of Advanced Environmental Technology, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

²Fujian Key Laboratory of Atmospheric Ozone Pollution Prevention, Institute of Urban Environment, Chinese Academy of Sciences, Xiamen 361021, China

³University of Chinese Academy of Sciences, Beijing 100049, China

⁴Fujian Agriculture and Forestry University, Fuzhou, 350002, China

*Correspondence to: jschen@iue.ac.cn (Jinsheng Chen); Linglingxu@iue.ac.cn (Lingling Xu)

Abstract: Heterogeneous uptake of NO₂ serves as a significant source for reactive nitrogen species, playing an important role in atmospheric chemistry. Laboratory studies have demonstrated that chlorine can promote the heterogeneous NO₂ uptake, yet this effect under real ambient conditions remains poorly elucidated. Based on comprehensive field observations, a machine learning technique, and a multiphase chemical box model in the coastal city of Xiamen, China, this study reveals the enhancement effect of chlorine (Cl) on NO₂ uptake and quantifies the impact of this enhanced uptake on reactive nitrogen species during nocturnal sea-land breeze (SLB) periods. Compared with non-SLB days, nocturnal concentrations of nitrous acid (HONO) and particulate nitrate (NO₃⁻) increased significantly during SLB days, with high mean value of NO₂ uptake rate constant (kNO₂) reaching $9.70 \times 10^{-6} \text{ s}^{-1}$. Machine learning revealed that chlorine was the most important influencing factor for the enhanced kNO₂. Incorporating this kNO₂ into the chemical box model substantially resolved the underestimation of HONO concentration and NO₃⁻ production under SLB conditions. Notably, nocturnal NO₂ uptake dominated HONO formation (83.9%), while making a substantial contribution (46.4%) to nitrate formation. This study highlights the critical role of chlorine-enhanced NO₂ uptake in atmospheric reactive nitrogen cycling and provides valuable insights for nocturnal chemistry in complex coastal environments.

1. Introduction

Nitrous acid (HONO) and nitrate (NO₃⁻) are key reactive nitrogen species in the atmosphere. HONO, a major source of OH radicals, significantly contributes to the formation of O₃ and secondary aerosols by influencing radical budget (Acker et al., 2006; Alicke et al., 2003). NO₃⁻ is an important chemical component of particulate matter (PM), contributing to PM pollution and acid rain (Cao et al., 2022; Qu and Han, 2021). As a crucial source of both HONO and NO₃⁻, the heterogeneous uptake of NO₂ has attracted considerable attention (Xuan et al., 2025; Carter et al., 1982).

Traditionally, the uptake of NO₂ in bulk solution was considered extremely slow, and thus this process was often negligible (Lee and Schwartz, 1981). However, numerous field observations have indicated that NO₂ uptake plays a significant role in the formation of HONO and NO₃⁻. Previous studies have revealed a strong correlation between NO₂ and HONO after excluding homogeneous formation and primary emissions, suggesting efficient heterogeneous conversion of NO₂ to HONO

(Su et al., 2008; Zhang et al., 2024). Studies have also found that incorporating NO₂ uptake mechanisms into models can substantially improve the HONO simulation, especially at night (Zhang et al., 2023; Xuan et al., 2024; Zhang et al., 2024; Shi et al., 2020). Using an aerosol-fog sampling system and a box model, Xu et al (Xu et al., 2024), found that NO₂ uptake was the primary driver of the rapid increase in NO₃⁻ during fog events. Currently, the reported reaction rate constant for NO₂ uptake vary widely across different types of surfaces, spanning several orders of magnitude from 10⁻⁷ s⁻¹ to 10⁻⁵ s⁻¹ (Zhang et al., 2025b). Multiple factors have been demonstrated to affect the NO₂ uptake rate constant (kNO₂) in laboratories, including NO₂ concentrations, aerosol particle size, ionic strength, pH, NH₃, and liquid water content (Liu and Abbatt, 2021; Gen et al., 2024; Zhang et al., 2025a). In the real atmosphere, research during haze events showed that parameterization schemes of kNO₂ considering relative humidity (RH) and NH₃ influences significantly improved the simulation results of heterogeneous HONO formation (Zhang et al., 2023). Long-term observations revealed that changes in aerosol chemical composition could alter aerosol pH, consequently affecting the uptake kinetics of NO₂ to HONO (Zhang et al., 2025b).

Investigations in coastal areas reported higher NO₂ uptake rates in marine air masses compared to continental ones (Zha et al., 2014; Yang et al., 2021), overall surpassing those measured at continental urban, suburban, and rural sites (Wentzell et al., 2010; Kleffmann et al., 2003; Zhang et al., 2020; Hao et al., 2020; Cui et al., 2018). These findings suggest that heterogeneous NO₂ uptake plays a particularly critical role in the coastal troposphere. Within the marine boundary layer, chloride ions are abundant due to sea spray. Chloride ions exhibit special surface propensity, making them tend to distribute at the air-water interface (Knipping et al., 2000; Piatkowski et al., 2014; Jungwirth and Tobias, 2006). Laboratory studies demonstrated that chloride ions can establish a surface-directed electric field, which attracts atmospheric NO₂ molecules and promotes their further uptake at air-water interface (Wang et al., 2025a; Zhang et al., 2025a). Nevertheless, the influence of chlorine on NO₂ uptake in realistic coastal atmosphere and the consequent impacts on reactive nitrogen cycling remain to be fully elucidated.

Sea-land breeze (SLB) events occur frequently in coastal urban areas (Chen et al., 2023; Liu et al., 2025; Xie et al., 2023). During SLB periods, the coastal boundary layer experiences intensive interactions between marine and continental air masses, providing a natural laboratory to investigate the effect of sea-salt chloride on NO₂ uptake. In this study, a multi-parameter field observation was conducted in Xiamen, a typical coastal city in China, during the autumn and winter seasons characterized by frequent SLB events (Chen et al., 2023). Based on observations, the characteristics of SLB days were revealed and kNO₂ under both SLB and non-SLB conditions were quantified. Using machine learning method, the significant promoting effect of chlorine on NO₂ uptake during nocturnal SLB periods was demonstrated. Furthermore, the obtained kNO₂ was incorporated into an advanced multiphase box model to evaluate the impact of chlorine-enhanced NO₂ uptake on atmospheric reactive nitrogen cycling. This study highlights the role of previously overlooked chlorine on NO₂ uptake for nocturnal chemistry in coastal atmosphere.

2. Methods

2.1 Field measurements and sea-land breeze identification

Our observations were conducted at the Atmospheric Observation Supersite (24.61°N, 118.06°E) of the Institute of Urban Environment (IUE), Chinese Academy of Sciences in Xiamen, from November 18, 2023 to January 6, 2024. Xiamen is a coastal city in southeastern China. The

observation site is located approximately 5 km from the shore and is surrounded by residential, commercial areas and major traffic arteries. Instruments were deployed on the roof of the IUE building at approximately 70 m above ground level. Virtually no high buildings obstructed the path between the site and the coast, making it a typical coastal urban site.

A suite of atmospheric parameters was observed simultaneously. Ambient HONO was measured online by a water-based long-path absorption photometer (Zhichen Beijing, China) (Xuan et al., 2024). Sulfuric acid (H₂SO₄) and HCl were measured using a long time-of-flight chemical ionization mass spectrometer, equipped with a nitrate reagent ion source (Aerodyne Research Inc., USA) (Yang et al., 2023). The chemical composition of PM_{2.5} was analyzed using both a MARGA ADI 2080 (Metrohm Applikon, Switzerland) and an ACSM (Aerodyne Research Inc., USA) (Liu et al., 2022; Chen et al., 2022). Particle number size distribution (PNSD) in the range of 2–300 nm was measured by two scanning mobility particle sizers (TSI inc., USA) equipped with a long-DMA (2.5–62 nm) and a nano-DMA (7–300 nm), respectively (Li et al., 2025). Additionally, a range of commercial instruments were employed to monitor trace gases (NO, NO₂, SO₂, O₃, and VOCs), meteorological parameters (ambient temperature, RH, wind-speed, wind-direction and ultraviolet radiation), and photolysis rates (*J*O¹D, *J*NO₂, *J*HONO, *J*NO₃, *J*HCHO and *J*H₂O₂) (Liu et al., 2022). Detailed descriptions of instrument operation, species identification, and calibration are provided in **Text S1**. Hourly BLH data were obtained from the ERA5 reanalysis dataset (Hersbach et al., 2020).

Sea-land breeze days were identified based on observed wind-speed and wind-direction according to the criteria established in our previous work (Chen et al., 2023). Briefly, the coastline in Xiamen is characterized by a northeast-southwest trend, the sea breeze is blowing from the east-south, and the land breeze is blowing from the west-north-northeast. According to observed wind direction, a natural day could be divided into four periods. Specifically, 01:00-08:00 LT (land breeze period), 13:00-20:00 LT (sea breeze period), and 09:00-12:00 LT and 21:00-24:00 LT (mixed sea-land breeze period). A day was classified as an SLB day if the following three conditions were met: (1) the 24-hour mean wind speed was < 10 m/s; (2) during the land breeze period, the land breeze persisted for ≥ 4 hours and the sea breeze lasted ≤ 2 hours; and (3) during the sea breeze period, sea breeze persisted for ≥ 4 hours and the land breeze lasted ≤ 2 hours. As shown in **Fig. S1** (light blue areas), a total of 23 SLB days were identified during the observation period. The remaining days were classified as non-SLB days. **Fig. S2** illustrates that the wind patterns on SLB days differed from those on non-SLB days, characterized by a higher prevalence of southeasterly sea breezes.

2.2 Calculation of HONO from heterogeneous formation processes and NO₂ uptake rate

Due to the absence of agricultural fields near the sampling site and its proximity to major traffic arteries, the primary emission of HONO was attributed to vehicles. Only nighttime data were used in our study, the uncertainties associated with photochemical process, such as NO₃⁻ photolysis were excluded. Therefore, the nighttime corrected HONO (HONO_{corr}) is calculated as follows:

$$\text{HONO}_{\text{corr},t} = \text{HONO}_t - E_{\text{vehicle},t} - P_{\text{NO}+\text{OH},t} \quad (1)$$

where HONO_t is the observed HONO concentration, *E*_{vehicle,t} is the HONO emission from vehicles, and *P*_{NO+OH,t} is the HONO production from homogeneous reactions. *E*_{vehicle,t} is calculated as the NO_x concentration multiplied by the emission factor 0.005 (Zhang et al., 2025b). *P*_{NO+OH,t} is calculated as follows:

$$P_{\text{NO}+\text{OH},t} = k_{\text{NO}+\text{OH}} \times [\text{NO}] \times [\text{OH}] \times 3600 \quad (2)$$

where *k*_{NO+OH} (cm³ molecules⁻¹ s⁻¹) is calculated in the Master Chemical Mechanism 3.3.1 (MCM

v3.3.1 (<https://mcm.york.ac.uk/MCM/>), 3600 s represents the 1 hour integration interval, and the concentration of OH radicals is estimated based on the steady-state balance of H₂SO₄ (Nie et al., 2022). Because H₂SO₄ is primarily produced by the reaction of SO₂ and OH and is mainly lost via condensation sink (CS) onto particle surfaces, the OH concentration can be estimated as below:

$$[\text{OH}] = \frac{[\text{H}_2\text{SO}_4] \times \text{CS}}{k_{\text{OH}+\text{SO}_2} \times [\text{SO}_2]} \quad (3)$$

where $k_{\text{OH}+\text{SO}_2}$ is 5×10^{-13} cm³/s (Seinfeld et al., 1998), and CS (/s) is calculated as follows (Kulmala et al., 2012):

$$\text{CS} = 2\pi D_v \int_0^{d_{p,\text{max}}} d_p \beta n(d_p) dd_p \quad (4)$$

where D_v is the diffusion coefficient of H₂SO₄, d_p is the particle diameter derived from the measured PNSD, β is the transitional correction factor, and $n(d_p)$ is the particle number concentration of diameter d_p .

In line with previous studies (Zhang et al., 2025b; Zhang et al., 2020; Zhang et al., 2024), the heterogeneous conversion of NO₂ to form HONO_{corr} at night is considered a pseudo-first-order reaction. The reaction rate constant (k_{NO_2} , s⁻¹) is calculated based on the conversion of NO₂ during a given period (from t_1 to t_2) as follows:

$$k_{\text{NO}_2} = \frac{\text{HONO}_{\text{corr},t_2} - \text{HONO}_{\text{corr},t_1}}{\overline{\text{NO}_2}(t_2 - t_1)} \quad (5)$$

where $\overline{\text{NO}_2}$ is the mean concentration of NO₂ between t_1 and t_2 .

To evaluate the sensitivity of the results to the assumed emission factor, the calculation of HONO_{corr} and k_{NO_2} (Fig. S3) were calculated using emission factors of 0.005, 0.008 and 0.012 (Xu et al., 2015), while the observational inputs, analysis periods and calculation procedures held constant.

2.3 Machine learning technique

Random Forest (RF) algorithm, a machine learning technique extensively applied in atmospheric (Yang et al., 2024; Lin et al., 2026a; Zhang et al., 2025b), was employed to elucidate the primary drivers of k_{NO_2} . The model was built using the “scikit-learn” library (<https://github.com/scikit-learn/scikit-learn/blob/fe2edb3cd/sklearn/ensemble/>, last access: June 10, 2026) in a python environment. Drawing upon previously identified factors influencing k_{NO_2} (e.g., pH, RH, NH₃) (Liu and Abbatt, 2021; Gen et al., 2024; Zhang et al., 2025a; Zhang et al., 2023) alongside the specific characteristics of the coastal urban environment, we incorporated meteorological parameters (RH and T), atmospheric reactive species (proxy of effective particle-phase chloride (Cl_{total}), NH₃, and NO₂), and heterogeneous-related parameters (pH, SA, and ground surfaces areas) as input variables for the RF model. Cl_{total} was defined as the sum of particulate Cl⁻ and gaseous HCl. Particle-surface Cl⁻ can facilitate heterogeneous NO₂ uptake, however, H⁺ produced during this process can protonate particulate Cl⁻ to form HCl, which subsequently partitions into the gas phase. Consequently, the measured Cl⁻ represents the residual concentration after chloride depletion and may underestimate the amount of particle-phase chloride present before and during NO₂ uptake. Adding gaseous HCl to the residual Cl⁻ partly accounts for this depleted fraction, therefore, Cl_{total} was used as a proxy for the particle-phase chloride involved in the uptake process, rather than assuming that gaseous HCl directly promotes NO₂ uptake. To evaluate whether the inferred chlorine dependence was sensitive to this proxy, two additional RF models were constructed by replacing

Cl_{total} with either Cl^- or gaseous HCl . Only nighttime data were used to train the model to eliminate the effects of photochemical processes on kNO_2 quantification, thereby enabling a more precise identification of key drivers. Detailed RF hyperparameter configurations and model evaluation are provided in Text S2. In this study, the dominant factors were identified based on RF factor importance, which quantifies the percentage contribution of each variable to kNO_2 (Breiman, 2001; Gregorutti et al., 2017). Furthermore, one-dimensional partial dependence plots (PDPs) were employed to delineate the main effects of individual variables, while bivariate PDPs were utilized to visualize and access the interaction effects between pairs of factors on kNO_2 (Goldstein et al., 2015).

2.4 Multiphase chemical box model. The multiphase chemical box model was established based on the Framework for 0-D Atmospheric Modeling (F0AM) (Wolfe et al., 2016). As shown in Table S1 and Table S2, comprehensive HONO and NO_3^- budgets were incorporated into the F0AM model to simulate the variations of HONO and NO_3^- . The ISORROPIA II model (Seinfeld et al., 1998) was integrated into the model to determine aerosol pH, aerosol liquid water content and the partitioning ratio of HNO_3 and NO_3^- . Six simulation scenarios were conducted: simulations without heterogeneous NO_2 uptake for (1) SLB and (2) non-SLB days; simulations with NO_2 uptake for (3) SLB and (4) non-SLB days, using the mean kNO_2 derived from SLB days for the SLB simulation and that derived from non-SLB days for the non-SLB simulation, respectively; and two additional sensitivity simulations for SLB days using the (5) 25th and (6) 75th percentiles of the kNO_2 distribution derived from SLB days. The ability of the model to reproduce real atmospheric processes was evaluated by comparing the simulated HONO and NO_3^- with observations. Due to the underestimation of total particulate nitrate concentrations measured by instruments, the performance of NO_3^- simulations was evaluated by comparing the trends of the simulated NO_3^- production rate with the observed NO_3^- concentration. In contrast, as HONO predominantly exists in the gas phase, the simulated HONO was evaluated directly against the observed HONO concentrations.

As for the model simulation, each simulation was preceded by a 3-day spin-up to allow intermediate species concentrations to stabilize. The mean diurnal pattern of trace gas concentrations, meteorological parameters, and the results from the ISORROPIA II model during SLB days and non-SLB days were used as input data to constrain the box model. Notably, the concentrations of HONO and NO_3^- were not constrained by observed values. More details about the model incorporating mechanisms, modules, and simulation settings are provided in Text S3 and our previous study (Lin et al., 2026b).

3 Results and discussion

3.1 Characteristics of atmospheric species during SLB days

Fig. S1 shows the time series of the observed parameters. Based on the identification criteria detailed in the Methods section, 23 days were classified as SLB days, accounting for 48% of entire observation period. The diurnal patterns of major observed parameters are summarized in Fig. S4, revealing that the primary differences between the two types of days predominantly emerged at night. Compared to non-SLB days, SLB days exhibited rapid nighttime accumulation of HONO and NO_3^- , accompanied by elevated chlorine levels, NO_x concentrations, and aerosol surface area (SA) concentrations. Additionally, consistent with our previous study, SLB days were characterized by

higher temperature and RH at night (Chen et al., 2022). **Fig. 1** and **Table S3** further summarizes the nighttime observational results on SLB and non-SLB days. By definition, SLB days are influenced by persistent sea breezes, which likely enhances the impact of sea-salt aerosols. As shown in **Fig. 1a**, the Cl^- concentrations were comparatively higher on SLB days ($0.154 \mu\text{g}/\text{m}^3$) than on non-SLB days ($0.139 \mu\text{g}/\text{m}^3$). Specifically, this overall increase (**Fig. S5**) was primarily driven by elevated levels during the sea breeze (18:00-20:00 LT) and mixed sea-land breeze (21:00-00:00 LT) periods, while concentrations declined during the land breeze (01:00-06:00 LT) period. The elevation pattern of Cl^- indicates that the more abundant Cl^- transported by stronger sea breezes on SLB days can persist into the subsequent mixed sea-land breeze periods. Additionally, **Fig. S6** shows a moderate positive correlation between Na^+ and Cl^- on SLB days ($r = 0.67$), compared with a weaker relationship on non-SLB days ($r = 0.28$). Because sodium chloride (NaCl) is a major component of sea salt, this contrast is consistent with a greater sea-salt influence on Cl^- on SLB days. As for gaseous chlorine, **Fig. 1b** shows that the concentration of HCl was significantly higher on SLB days (96.2 ppt) than on non-SLB days (67.5 ppt). This further confirms that, under the influence of sea-salt aerosols on SLB days, chlorine levels are distinctly elevated in the nighttime coastal troposphere.

Fig. 1c indicates that the concentration of NO_2 on SLB days (mean: 28.3 ppb) was significantly higher than that on non-SLB days (23.2 ppb). Within our study area, ship emissions, cross-sea bridge traffic, and road traffic contribute substantial NO_x , and the enhanced local circulation on SLB days can promote NO_2 accumulation (Chen et al., 2023). The mean concentrations of HONO and NO_3^- on SLB day were 0.836 ppb and $3.61 \mu\text{g}/\text{m}^3$, respectively, which were significantly higher than the corresponding values of 0.497 ppb and $3.13 \mu\text{g}/\text{m}^3$ on non-SLB days (**Figs. 1d-1e**). To better understand the elevated HONO levels, $\text{HONO}_{\text{corr}}$ concentrations, defined as the observed HONO concentration subtracts contributions from primary emissions and gas-phase production, were calculated to represent the nocturnal HONO produced from heterogeneous conversion. The ratio of $\text{HONO}_{\text{corr}}/\text{NO}_2$ was notably higher on SLB days (0.030) than on non-SLB days (0.021), indicating more active heterogeneous conversion of NO_2 to HONO on SLB days. Multiple studies have confirmed that NO_2 uptake is a primary source of nocturnal heterogeneous HONO formation (Zhang et al., 2023; Zhang et al., 2022). Here, the higher SA concentrations on SLB days were also conducive to heterogeneous reactions (**Fig. 1g**). Furthermore, **Figs. 2a-2b** indicate a moderate positive correlation between $\text{HONO}_{\text{corr}}$ and NO_3^- , with the correlation coefficient (r) increasing from 0.56 on non-SLB days to 0.76 on SLB days. The $\text{HONO}_{\text{corr}}$ concentrations under the two conditions overlapped mainly below 1.5 ppb. Within this overlapping range, the correlation on SLB days remained significant ($r = 0.63$, $p < 0.001$, **Fig. S7a**) but was close to that on non-SLB days ($r = 0.56$), indicating that the wider concentration range partly contributed to the higher full-range correlation on SLB days, while the positive relationship between $\text{HONO}_{\text{corr}}$ and NO_3^- persisted within the overlapping range. As shown in **R1**, NO_2 uptake via disproportionation simultaneously produces HONO and HNO_3 , the latter of which is further converted to NO_3^- via gas-particle partitioning. Thus, nocturnal NO_2 uptake is also likely an important contributor to the formations of NO_3^- on SLB days. **Figs. 2c-2d** show that the full-range correlation between $\text{HONO}_{\text{corr}}$ and gaseous HCl was stronger on SLB days than on non-SLB days ($r = 0.58$ versus 0.16). Within the overlapping $\text{HONO}_{\text{corr}}$ range below 1.5 ppb, the correlation on SLB days decreased to ($r = 0.47$) but remained significant ($p < 0.001$, **Fig. S7b**) and appreciably higher than that on non-SLB days. This result indicates that the stronger HCl - $\text{HONO}_{\text{corr}}$ relationship during SLB periods cannot be explained solely by the wider $\text{HONO}_{\text{corr}}$ range. The H^+ generated by NO_2 uptake can combine with Cl^- (**R2-R3**). Simultaneously,

the aerosol pH in the coastal region of southeastern China was relatively low (approximately 1.92 ± 0.34 during nocturnal SLB periods), chloride depletion could easily occur, leading to the observed positive correlation between $\text{HONO}_{\text{corr}}$ and HCl . The enhanced chloride depletion can also explain our observations of a significant increase in gaseous HCl but not in Cl^- concentrations on SLB days. Similarly, the rapid decline in Cl^- concentration during land breeze periods (**Fig. S4**) could also be explained by chloride depletion.





Figure 1. Box plots of key observed parameters during nocturnal SLB and non-SLB periods. (a) Cl^- , (b) HCl , (c) NO_2 , (d) HONO , (e) NO_3^- , (f) RH , (g) SA , and (h) ambient temperature (T). The box shows the 25th–75th percentiles with whiskers representing ± 1 standard deviation. The black dot and black line within the box represent the mean and the median. Differences between SLB and non-SLB days were tested using a two-sample t-test. Statistical significances are denoted as * ($p < 0.05$), ** ($p < 0.01$) and *** ($p < 0.001$). For all panels except (h), the number adjacent to the significance symbol denotes the ratio of the mean value on SLB days to that on non-SLB days; in panel (h), it denotes the absolute difference between the means.




Figure 2. Relationship between HONO_{corr} and the important products of heterogeneous chemical processes during nocturnal SLB and non-SLB periods. HONO_{corr} denotes the corrected HONO concentration from secondary conversion. The products include (a, b) NO₃⁻ and (c, d) HCl. Blue and red dots represent observations during SLB days and non-SLB days, respectively. Each panel displays the linear regression fits and Pearson correlation coefficients (r) for the corresponding periods.

3.2 Enhanced rate of NO₂ uptake and its influencing factors

The values of k_{NO_2} on SLB days and non-SLB days were derived as Fig. S8. Overall, the average k_{NO_2} values on SLB days ($9.70 \times 10^{-6} \text{ s}^{-1}$) were significantly higher than those on non-SLB days ($6.72 \times 10^{-6} \text{ s}^{-1}$), by a factor of approximately 1.44. Across the tested emission factors, the mean HONO_{corr} decreased from 0.58 to 0.51 ppb on SLB days, whereas k_{NO_2} varied by no more than 6.2%, indicating limited sensitivity of the derived rate constant to the assumed emission factor (Fig. S3). As presented in Table S4, a prolonged increase in the HONO_{corr}/NO₂ ratio was also observed during 17 of the 23 SLB days (see an example in Fig. S9). In contrast, this phenomenon was virtually absent on non-SLB days, indicating a sustained nocturnal enhancement of k_{NO_2} during SLB periods. Compared with existing field observations, the k_{NO_2} on SLB days was broadly consistent with the levels reported under the sea-case scenario in coastal cities of China such as Qingdao ($1.25 \times 10^{-5} \text{ s}^{-1}$) and Hong Kong ($8.81 \times 10^{-6} \text{ s}^{-1}$) (Yang et al., 2021; Zha et al., 2014), while it was markedly higher than those values observed in major non-coastal cities including Beijing ($7.78 \times 10^{-7} - 5.75 \times 10^{-6} \text{ s}^{-1}$) (Wang et al., 2017b; Zhang et al., 2020; Xuan et al., 2023; Jia et al., 2020; Zhang et al., 2022), Nanjing ($1.19 \times 10^{-6} - 2.22 \times 10^{-6} \text{ s}^{-1}$) (Zheng et al., 2020; Liu et al., 2019), and Guangzhou ($4.44 \times 10^{-6} - 6.67 \times 10^{-6} \text{ s}^{-1}$) (Su et al., 2008; Li et al., 2012; Qin et al., 2009). The consistently higher k_{NO_2} in coastal areas implies that coastal atmospheric environments are likely more conducive to NO₂ uptake.

A machine learning technique was used to further identify key drivers of the enhanced kNO_2 on SLB days. The RF models using Cl_{total} reproduced the calculated kNO_2 well during both SLB and non-SLB periods, with R^2 values of 0.91 and 0.87, respectively (Figs. S10c-10d). To evaluate the robustness of the Cl_{total} proxy, RF models were also constructed using residual particulate Cl^- or gaseous HCl separately. Both models retained good predictive performance on SLB days, with R^2 values of 0.90 and 0.89, respectively, compared with 0.91 for the Cl_{total} -based model (Fig. S10). When considered separately, Cl^- and HCl both ranked third after RH and SA (Fig. S11). In comparison, Cl_{total} ranked first and produced a modest improvement in model performance (Fig. 3a). These results indicate that the chlorine-related signal is captured partly by residual particulate Cl^- and partly by gaseous HCl formed through chloride depletion. Combining the two species therefore provides a more representative process-based proxy for the particulate chloride involved in NO_2 uptake. Notably, the role of Cl_{total} and RH diminished substantially on non-SLB days (Fig. 3b), underscoring the profound modulation of these two influencing factors by the SLB circulation. Fig. 3c-3e further elucidate the specific impacts of these primary drivers on nocturnal kNO_2 under SLB conditions. Specifically, Cl_{total} exhibits a positive correlation with kNO_2 (Fig. 3c). This phenomenon is likely attributable to the preferential partitioning of chloride ions at the gas-aqueous interface of aerosols (Zhang et al., 2025a; Shen et al., 2025; Wang et al., 2025a). The surface-bound Cl^- can attract gaseous NO_2 molecules (R4) to form interfacial intermediates $[\text{Cl-NO}_2]^-$, thereby facilitating the overall NO_2 uptake process. Subsequent interaction analysis was consistent with the mechanisms proposed in laboratory studies (R4-R5) and suggest that similar processes may operate in the coastal boundary layer. Fig. 3f shows that at a fixed NO_2 concentration, elevated chloride levels amplify the positive effect of NO_2 molecules on kNO_2 . This indicates that chloride ions actively promote the heterogeneous uptake of NO_2 , consequently accelerating the kNO_2 . Similar halogen-promoted effect has also been revealed in NO_2 -initiated aqueous-phase oxidation of dissolved SO_2 at the air-aqueous interface and of nitrate photolysis at the aqueous-phase surface (Wang et al., 2025a; Shen et al., 2025). Fig. 3d shows a significant positive relationship exists between RH and kNO_2 , suggesting that moist sea breezes effectively supply the water molecules essential for the heterogeneous NO_2 uptake. This finding aligns with previously reported field observations highlighting RH-driven NO_2 uptake (Xuan et al., 2024). Analogous to Cl_{total} , the interaction between RH and NO_2 reveals that at a constant NO_2 level, the positive impact of NO_2 on kNO_2 is enhanced under elevated RH. It implies that RH could also facilitate NO_2 uptake by promoting the interaction between NO_2 molecules and heterogeneous interface. Furthermore, Fig. 3h shows a threshold-dependent joint enhancement: kNO_2 increased when RH exceeded approximately 80% and Cl_{total} exceeded approximately $0.5 \mu\text{g}/\text{m}^3$, whereas the response was limited when either variable remained below these levels. This pattern suggests a synergistic effect under combined high-RH and high-chlorine conditions and may partly explain their higher relative importance on SLB days. As a primary factor on both SLB and non-SLB days, SA positively correlates with kNO_2 (Fig. 3e). This indicates that an increase in aerosol surface area concentrations provides effective reaction interfaces for NO_2 uptake process, consistent with most atmospheric heterogeneous processes (Wang et al., 2017a; Ma et al., 2026). Moreover, Fig. S12 shows that on SLB days, concentrations of the major NO_2 uptake products (HONO and NO_3^-) increased with those key factors including Cl_{total} , RH, and SA, providing additional support for the above driving factor analysis. In general, the reasons underlying the enhanced kNO_2 on SLB days help to explain the multi-fold higher kNO_2 observed in coastal areas compared to non-coastal urban areas, highlighting ambient chlorine levels

and RH as non-negligible contributors.





Figure 3. Main and interaction effects of influencing factors for k_{NO_2} . Factor importance determined by the random forest (a) SLB days and (b) non-SLB days. Partial dependence plots for the primary drivers on SLB days: (c) Cl_{total} , (d) RH, and (e) SA. Interactions between key factors on SLB days: (f) Cl_{total} and NO_2 , (g) RH and NO_2 , and (h) RH and Cl_{total} . In panels (a-b), S_{ground} denotes ground surface areas calculated by $1/\text{BLH}$ (boundary layer height). In panels (c-h), partial dependence reflects the extent to which the influencing factors affect k_{NO_2} .

3.3 Enhanced NO_2 uptake promotes the atmospheric reactive nitrogen cycle

The above correlations analysis suggests that NO_2 uptake could play an important role in the production of HONO and NO_3^- on SLB days. Since the HONO and NO_3^- formation were influenced by multiple meteorological factors and many atmospheric reaction pathways, a multiphase chemical box model was employed to evaluate the role of enhanced NO_2 uptake. As described in Methods, six simulation scenarios were set up. Incorporating NO_2 uptake mechanism substantially improved the simulated diurnal HONO profile during SLB periods: r increased from 0.57 to 0.94, while mean absolute error (MAE) and root mean square error (RMSE) decreased from 0.58 to 0.11 ppb and from 0.64 to 0.13 ppb, respectively (Fig. 4a). For NO_3^- , incorporating NO_2 uptake increased the mean simulated net NO_3^- formation rate by $2.14 \mu\text{g}/\text{m}^3/\text{h}$ and strengthened the temporal consistency between the simulated net formation rate and observed concentration, increasing r from 0.04 ($p > 0.05$) to 0.64 ($p < 0.001$, Fig. 4b). This improvement indicates that the added mechanism better

represented the observed temporal variability in NO_3^- during SLB periods. Additionally, the NO_2 uptake could robustly explain the enhancements in both atmospheric reactive nitrogen species on SLB days across various kNO_2 quantiles (Fig. S13). Conversely, applying the current calculated kNO_2 to non-SLB days overpredicted HONO and NO_3^- levels (Fig. S14). Given uncertainties in BLH, deposition, dilution and other loss processes, this bias cannot be attributed solely to kNO_2 . Nevertheless, it suggests that the effective difference in NO_2 uptake impacts between SLB and non-SLB conditions may exceed that inferred from the observation-derived mean kNO_2 values, which differed by a factor of 1.44.

Figs. 4c-4d illustrate the simulated HONO and NO_3^- budgets for the scenarios incorporating NO_2 uptake on SLB days. Despite potential overestimations of non-SLB kNO_2 , the mean NO_2 uptake rate remained significantly higher on SLB days than on non-SLB days, with an average rate of 1.03 ppb/h (Fig. S15). During nighttime, NO_2 uptake accounted for 83.8% of HONO formation (Fig. 4e). The predominance of NO_2 uptake in nocturnal HONO formation has also been observed during PM pollution episodes in urban areas (Xuan et al., 2024; Jia et al., 2020). However, unlike the chlorine-driven reason identified here, those prior cases were primarily attributed to favorable heterogeneous reaction conditions, such as larger SA concentrations, lower BLH, and higher RH. In addition to NO_2 uptake, primary emissions from vehicle exhaust, coal combustion, and biomass burning are also recognized as major contributors to nighttime HONO in urban areas (Wang et al., 2025b; Zhang et al., 2022). In this study, primary emissions contributed only 15.5% to nocturnal HONO formation, substantially less than the contribution from NO_2 uptake. As for nighttime NO_3^- formation, many studies had indicated that N_2O_5 uptake was the dominant pathway in urban areas (Chen et al., 2020; Wang et al., 2018). However, the situation changed under the impact of the enhanced NO_2 uptake. Fig. 4f shows that NO_2 uptake contributed 46.4% to nocturnal NO_3^- formation, which was almost comparable to that of N_2O_5 uptake (51.6%). These values were obtained using the BT09 parameterization of $\gamma\text{N}_2\text{O}_5$ (Bertram and Thornton, 2009b). Additional simulations using the EJ05 and GRI09 schemes yielded contributions of 47.0–47.9% for NO_2 uptake and 50.0–50.9% for N_2O_5 uptake (Fig. S16 and Table S5), indicating that their comparable importance was insensitive to the choice of $\gamma\text{N}_2\text{O}_5$ parameterization (Evans and Jacob, 2005; Hallquist et al., 2003; Griffiths et al., 2009). While chlorine can also promote the uptake of N_2O_5 (Bertram and Thornton, 2009a), the relative enhancement in the contribution of NO_2 uptake to nocturnal NO_3^- formation was more pronounced. This might be related to the suppression of N_2O_5 uptake caused by increased NO_3^- generating directly from NO_2 uptake. The increase in NO_3^- component of aerosol can shift the equilibrium between N_2O_5 and NO_3^- toward N_2O_5 , thereby reducing the contribution of N_2O_5 to NO_3^- formation (Bertram and Thornton, 2009a; Wagner et al., 2013). Additionally, the enhanced NO_2 uptake process also competed for NO_2 , an important precursor of N_2O_5 formation, especially in the NO_2 -limited regime for N_2O_5 formation (Lin et al., 2025), ultimately reducing the overall enhancement effect on N_2O_5 generation. Moreover, this comparable contribution of NO_2 and N_2O_5 uptake to NO_3^- formation was also found during fog events (Xu et al., 2024). Although the observed phenomenon was similar, the dominant drivers differed between fog conditions and coastal SLB periods. The former was primarily attributed to favorable uptake conditions such as abundant aerosol water content and high surface area concentration, whereas the latter was more due to the enhancement effect of chlorine levels and RH on the NO_2 uptake process. Overall, these results reveal the high contribution of NO_2 uptake to HONO and NO_3^- formation during SLB circulation periods, demonstrating that enhanced NO_2 uptake can

significantly influence the atmospheric reactive nitrogen cycle in coastal urban areas.




Figure 4. Quantified results from the multiphase chemical box model during SLB periods. (a) Diurnal variations in observed and simulated HONO concentrations with and without NO₂ uptake. (b) Relationship between the simulated NO₃⁻ production rate and observed concentration with and without NO₂ uptake. Diurnal variations of HONO (c) and NO₃⁻ budgets (d) during SLB days. Diurnal variations of source contributions to (e) HONO and (f) NO₃⁻. The grey shaded areas represent the nighttime period.

4. Conclusions

Overall, our observations show that nighttime sea-land breeze period in this coastal urban site were characterized by stronger sea-salt aerosol influence, higher relative humidity, and elevated HONO and nitrate concentrations. Nighttime conditions led to a significant enhancement in the NO₂ uptake rate constant, mainly driven by increased chlorine levels and relative humidity. Multiphase chemical box model simulations further confirmed that this enhanced NO₂ uptake promoted HONO and nitrate formation. Fig.5 illustrates a schematic diagram of chlorine-enhanced heterogeneous uptake

of NO_2 during sea-land breeze days. Under the influence of strong sea-land breeze circulation, the continuous influx of sea-salt aerosols transported by moist sea breezes leads to high chlorine levels during the night. Meanwhile, emission from vehicles and ships contribute substantial NO_2 , which accumulates in the coastal atmospheric boundary layer (Wen et al., 2023; Sun et al., 2025). Under these conditions, abundant chloride can effectively **increase** kNO_2 and promote the NO_2 uptake process, thereby influencing the formation of key atmospheric reactive nitrogen species, HONO and NO_3^- , as well as broader reactive nitrogen cycling. Notably, other sea-salt halides (Br^- , I^-) exhibit similar chemical properties to chlorine and may also contribute to the elevated kNO_2 (Zhang et al., 2025a).

Although the observations were limited to one coastal urban site in autumn and winter, **chlorine-enhanced NO_2 uptake may also occur in other coastal environments where chloride-rich sea-salt aerosols, high RH, and sufficient NO_2 coexist. Its magnitude is expected to vary seasonally and regionally with sea-salt loading, aerosol water content, aerosol composition, and NO_2 abundance. Therefore, the quantitative contribution derived here should be considered site- and period-specific, and multi-season observations at other coastal sites are needed to assess its broader relevance.** Additionally, our study focuses specifically on nocturnal NO_2 uptake to form HONO, given that daytime HONO formation involves more complex factors such as photo-enhanced heterogeneous reactions of NO_2 and nitrate photolysis (Stemmler et al., 2006; Han et al., 2016; Ye et al., 2017; Zhou et al., 2011). Future investigations are warranted to explicitly evaluate the impact of chlorine on daytime NO_2 uptake. Moreover, current air quality models often exhibit considerable biases in simulating secondary pollutants such as O_3 and $\text{PM}_{2.5}$ over coastal areas (Huang et al., 2025; Ma et al., 2025; Mao et al., 2022). One possible reason is that the influence of chlorine on chemical processes like NO_2 uptake has not yet been adequately considered. Given the dense traffic and substantial NO_2 emission in coastal cities of China, greater attention should be paid to the chlorine-enhanced NO_2 uptake process and its subsequent impact on nocturnal chemistry in coastal urban atmosphere.




Figure 5. Schematic of the chlorine-enhanced heterogeneous uptake of NO_2 during sea-land breeze days in coastal areas. Red arrows, black solid circles, and blue solid circles represent the

enhancement of the uptake process, halide ions and water molecular, respectively.

Code availability. Data analysis methods are available on request from Jinsheng Chen (jschen@iue.ac.cn).

Data availability. The dataset can be accessed at Zenodo website (Lin et al., 2026).

Author contributions:

Z.L. contributed to the conceptualization, investigation, methodology, data curation, software, analysis and writing of the original draft. L.X. and J.C. contributed to the data curation, reviewing and editing the text, supervision, and funding acquisition. X.Y., L.L., C.Y., X.J., Y.C., K.Z., F.Z., Z.C., G.C., X.F., and M.L. provided useful advice and revised the manuscript.

Competing interests:

The authors declare no competing interests.

Acknowledgements. The authors acknowledge all members of the atmospheric chemistry team at the Institute of Urban Environment.

Financial support. This work was funded by the National Natural Science Foundation of China (U22A20578), the guiding project of seizing the commanding heights of “self-purifying city” (IUE-CERAE-202402), the National Key Research and Development Program (2022YFC3700304), STS Plan Supporting Project of the Chinese Academy of Sciences in Fujian Province (2023T3013), and Xiamen Atmospheric Environment Observation and Research Station of Fujian Province.

References:

- Acker, K., Möller, D., Wieprecht, W., Meixner, F. X., Bohn, B., Gilge, S., Plass-Dülmer, C., and Berresheim, H.: Strong daytime production of OH from HNO₂ at a rural mountain site, *Geophysical Research Letters*. 33, <https://doi.org/10.1029/2005GL024643>, 2006.
- Alicke, B., Geyer, A., Hofzumahaus, A., Holland, F., Konrad, S., Pätz, H. W., Schäfer, J., Stutz, J., Volz-Thomas, A., and Platt, U.: OH formation by HONO photolysis during the BERLIOZ experiment, *Journal of Geophysical Research: Atmospheres*. 108, PHO 3-1-PHO 3-17, <https://doi.org/10.1029/2001JD000579>, 2003.
- Bertram, T. H. and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride, *Atmos. Chem. Phys.*, 9, 8351-8363, <https://doi.org/10.5194/acp-9-8351-2009>, 2009a.
- Bertram, T. H. and Thornton, J. A.: Toward a general parameterization of N₂O₅ reactivity on aqueous particles: the competing effects of particle liquid water, nitrate and chloride, *Atmos. Chem. Phys.*, 9, 8351-8363, <https://doi.org/10.5194/acp-9-8351-2009>, 2009b.
- Breiman, L.: Random Forests, *Machine Learning*. 45, 5-32, <https://doi.org/10.1023/A:1010933404324>, 2001.
- Cao, Y., Ma, Q., Chu, B., and He, H.: Homogeneous and heterogeneous photolysis of nitrate in the atmosphere: state of the science, current research needs, and future prospects, *Frontiers of Environmental Science & Engineering*. 17, 48, <https://doi.org/10.1007/s11783-023-1648-6>, 2022.
- Carter, W. P. L., Atkinson, R., Winer, A. M., and Pitts Jr, J. N.: Experimental investigation of

507 chamber-dependent radical sources, *International Journal of Chemical Kinetics*. 14, 1071-1103,
 508 <https://doi.org/10.1002/kin.550141003>, 1982.
 509 Chen, X., Wang, H., Lu, K., Li, C., Zhai, T., Tan, Z., Ma, X., Yang, X., Liu, Y., Chen, S., Dong, H.,
 510 Li, X., Wu, Z., Hu, M., Zeng, L., and Zhang, Y.: Field Determination of Nitrate Formation Pathway
 511 in Winter Beijing, *Environmental Science & Technology*. 54, 9243-9253,
 512 <https://doi.org/10.1021/acs.est.0c00972>, 2020.
 513 Chen, Y., Yang, C., Xu, L., Chen, J., Zhang, Y., Shi, J., Fan, X., Zheng, R., Hong, Y., and Li, M.:
 514 Chemical composition of NR-PM₁ in a coastal city of Southeast China: Temporal variations and
 515 formation pathways, *Atmospheric Environment*. 285, 119243, [https://doi.org/](https://doi.org/10.1016/j.atmosenv.2022.119243)
 516 [10.1016/j.atmosenv.2022.119243](https://doi.org/10.1016/j.atmosenv.2022.119243), 2022.
 517 Chen, Y., Yang, C., Xu, L., Fan, X., Shi, J., Zheng, R., Hong, Y., Li, M., Liu, T., Chen, G., Yin, L.,
 518 and Chen, J.: Variations of chemical composition of NR-PM₁ under the influence of sea land breeze
 519 in a coastal city of Southeast China, *Atmospheric Research*. 285, 106626, [https://doi.org/](https://doi.org/10.1016/j.atmosres.2023.106626)
 520 [10.1016/j.atmosres.2023.106626](https://doi.org/10.1016/j.atmosres.2023.106626), 2023.
 521 Cui, L., Li, R., Zhang, Y., Meng, Y., Fu, H., and Chen, J.: An observational study of nitrous acid
 522 (HONO) in Shanghai, China: The aerosol impact on HONO formation during the haze episodes,
 523 *Science of The Total Environment*. 630, 1057-1070, <https://doi.org/10.1016/j.scitotenv.2018.02.063>,
 524 2018.
 525 Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model
 526 budgets of tropospheric nitrogen oxides, ozone, and OH, *Geophysical Research Letters*. 32,
 527 <https://doi.org/10.1029/2005GL022469>, 2005.
 528 Gen, M., Zheng, H., Sun, Y., Xu, W., Ma, N., Su, H., Cheng, Y., Wang, S., Xing, J., Zhang, S., Xue,
 529 L., Xue, C., Mu, Y., Tian, X., Matsuki, A., and Song, S.: Rapid hydrolysis of NO₂ at High Ionic
 530 Strengths of Deliquesced Aerosol Particles, *Environmental Science & Technology*. 58, 7904-7915,
 531 <https://doi.org/10.1021/acs.est.3c08810>, 2024.
 532 Goldstein, A., Kapelner, A., Bleich, J., and Pitkin, E.: Peeking Inside the Black Box: Visualizing
 533 Statistical Learning With Plots of Individual Conditional Expectation, *Journal of Computational and*
 534 *Graphical Statistics*. 24, 44-65, <https://doi.org/10.1080/10618600.2014.907095>, 2015.
 535 Gregorutti, B., Michel, B., and Saint-Pierre, P.: Correlation and variable importance in random
 536 forests, *Statistics and Computing*. 27, 659-678, <https://doi.org/10.1007/s11222-016-9646-1>, 2017.
 537 Griffiths, P. T., Badger, C. L., Cox, R. A., Folkers, M., Henk, H. H., and Mentel, T. F.: Reactive
 538 Uptake of N₂O₅ by Aerosols Containing Dicarboxylic Acids. Effect of Particle Phase, Composition,
 539 and Nitrate Content, *The Journal of Physical Chemistry A*. 113, 5082-5090,
 540 <https://doi.org/10.1021/jp8096814>, 2009.
 541 Hallquist, M., Stewart, D. J., Stephenson, S. K., and Anthony Cox, R.: Hydrolysis of N₂O₅ on sub-
 542 micron sulfate aerosols, *Physical Chemistry Chemical Physics*. 5, 3453-3463,
 543 <https://doi.org/10.1039/B301827J>, 2003.
 544 Han, C., Yang, W., Wu, Q., Yang, H., and Xue, X.: Heterogeneous Photochemical Conversion of
 545 NO₂ to HONO on the Humic Acid Surface under Simulated Sunlight, *Environmental Science &*
 546 *Technology*. 50, 5017-5023, <https://doi.org/10.1021/acs.est.5b05101>, 2016.
 547 Hao, Q., Jiang, N., Zhang, R., Yang, L., and Li, S.: Characteristics, sources, and reactions of nitrous
 548 acid during winter at an urban site in the Central Plains Economic Region in China, *Atmos. Chem.*
 549 *Phys.*, 20, 7087-7102, <https://doi.org/10.5194/acp-20-7087-2020>, 2020.
 550 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J.,

551 Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G.,
 552 Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis,
 553 M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan,
 554 R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay,
 555 P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J.-N.: The ERA5 global reanalysis, Quarterly
 556 Journal of the Royal Meteorological Society. 146, 1999-2049, <https://doi.org/10.1002/qj.3803>,
 557 2020.
 558 Huang, L., Zhang, X., Emery, C., Mu, Q., Yarwood, G., Zhai, H., Sun, Z., Xue, S., Wang, Y., Fu, J.
 559 S., and Li, L.: Recommendations on benchmarks for numerical air quality model applications in
 560 China – Part 2: Ozone and uncertainty analysis, Atmos. Chem. Phys., 25, 4233-4249,
 561 <https://doi.org/10.5194/acp-25-4233-2025>, 2025.
 562 Jia, C., Tong, S., Zhang, W., Zhang, X., Li, W., Wang, Z., Wang, L., Liu, Z., Hu, B., Zhao, P., and
 563 Ge, M.: Pollution characteristics and potential sources of nitrous acid (HONO) in early autumn 2018
 564 of Beijing, Science of The Total Environment. 735, 139317, [https://doi.org/](https://doi.org/10.1016/j.scitotenv.2020.139317)
 565 [10.1016/j.scitotenv.2020.139317](https://doi.org/10.1016/j.scitotenv.2020.139317), 2020.
 566 Jungwirth, P. and Tobias, D. J.: Specific Ion Effects at the Air/Water Interface, Chemical Reviews.
 567 106, 1259-1281, <https://doi.org/10.1021/cr0403741>, 2006.
 568 Kleffmann, J., Kurtenbach, R., Lörzer, J., Wiesen, P., Kalthoff, N., Vogel, B., and Vogel, H.:
 569 Measured and simulated vertical profiles of nitrous acid—Part I: Field measurements, Atmospheric
 570 Environment. 37, 2949-2955, [https://doi.org/10.1016/S1352-2310\(03\)00242-5](https://doi.org/10.1016/S1352-2310(03)00242-5), 2003.
 571 Knipping, E. M., Lakin, M. J., Foster, K. L., Jungwirth, P., Tobias, D. J., Gerber, R. B., Dabdub, D.,
 572 and Finlayson-Pitts, B. J.: Experiments and Simulations of Ion-Enhanced Interfacial Chemistry on
 573 Aqueous NaCl Aerosols, Science. 288, 301-306, <https://doi.org/10.1126/science.288.5464.301>,
 574 2000.
 575 Kulmala, M., Petäjä, T., Nieminen, T., Sipilä, M., Manninen, H. E., Lehtipalo, K., Dal Maso, M.,
 576 Aalto, P. P., Junninen, H., Paasonen, P., Riipinen, I., Lehtinen, K. E. J., Laaksonen, A., and Kerminen,
 577 V.-M.: Measurement of the nucleation of atmospheric aerosol particles, Nature Protocols. 7, 1651-
 578 1667, <https://doi.org/10.1038/nprot.2012.091>, 2012.
 579 Lee, Y. N. and Schwartz, S. E.: Reaction kinetics of nitrogen dioxide with liquid water at low partial
 580 pressure, The Journal of Physical Chemistry. 85, 840-848, <https://doi.org/10.1021/j150607a022>,
 581 1981.
 582 Li, L., Lin, Z., Li, M., Fan, X., Yang, C., Hou, A., Chen, Y., Xu, L., Hong, Y., and Chen, J.:
 583 Importance of Methanesulfonic Acid in New Particle Formation in a Coastal City of Southeast China,
 584 Geophysical Research Letters. 52, e2025GL117238, <https://doi.org/10.1029/2025GL117238>, 2025.
 585 Li, X., Brauers, T., Häsel, R., Bohn, B., Fuchs, H., Hofzumahaus, A., Holland, F., Lou, S., Lu, K.
 586 D., Rohrer, F., Hu, M., Zeng, L. M., Zhang, Y. H., Garland, R. M., Su, H., Nowak, A., Wiedensohler,
 587 A., Takegawa, N., Shao, M., and Wahner, A.: Exploring the atmospheric chemistry of nitrous acid
 588 (HONO) at a rural site in Southern China, Atmos. Chem. Phys., 12, 1497-1513,
 589 <https://doi.org/10.5194/acp-12-1497-2012>, 2012.
 590 Lin, Z., Li, L., Xu, L., Ji, X., Zhang, K., Yang, C., Chen, G., Fan, X., and Chen, J.: Unexpected
 591 Daytime N₂O₅ in Urban Areas: Formation Mechanisms and Environmental Impacts, Geophysical
 592 Research Letters. 53, e2025GL121461, <https://doi.org/10.1029/2025GL121461>, 2026a.
 593 Lin, Z., Ji, X., Xu, L., Chen, G., Yang, C., Zhang, K., Zhang, F., Li, L., Chen, Y., and Chen, J.:
 594 Enhanced NO₂-driven multiphase formation of particulate nitrate and sulfate under high-humidity

conditions, *npj Climate and Atmospheric Science*. 9, 76, <https://doi.org/10.1038/s41612-026-01352-5>, 2026b.

Lin, Z., Ying, C., Xu, L., Ji, X., Zhang, K., Zhang, F., Chen, G., Li, L., Yang, C., Chen, Y., Chen, Z., and Chen, J.: Measurement report: High contribution of N_2O_5 uptake to particulate nitrate formation in NO_2 -limited urban areas, *Atmos. Chem. Phys.*, 25, 17747-17759, <https://doi.org/10.5194/acp-25-17747-2025>, 2025.

Liu, C., Wang, H., Li, L., Chen, X., Lu, X., and Fan, S.: Impacts of sea-land breeze on the coastal ozone in the Pearl River Delta, China, *Journal of Environmental Sciences*. <https://doi.org/10.1016/j.jes.2025.08.037>, 2025.

Liu, T. and Abbatt, J. P. D.: Oxidation of sulfur dioxide by nitrogen dioxide accelerated at the interface of deliquesced aerosol particles, *Nature Chemistry*. 13, 1173-1177, <https://doi.org/10.1038/s41557-021-00777-0>, 2021.

Liu, T., Hong, Y., Li, M., Xu, L., Chen, J., Bian, Y., Yang, C., Dan, Y., Zhang, Y., Xue, L., Zhao, M., Huang, Z., and Wang, H.: Atmospheric oxidation capacity and ozone pollution mechanism in a coastal city of southeastern China: analysis of a typical photochemical episode by an observation-based model, *Atmos. Chem. Phys.*, 22, 2173-2190, <https://doi.org/10.5194/acp-22-2173-2022>, 2022.

Liu, Y., Nie, W., Xu, Z., Wang, T., Wang, R., Li, Y., Wang, L., Chi, X., and Ding, A.: Semi-quantitative understanding of source contribution to nitrous acid (HONO) based on 1 year of continuous observation at the SORPES station in eastern China, *Atmos. Chem. Phys.*, 19, 13289-13308, <https://doi.org/10.5194/acp-19-13289-2019>, 2019.

Ma, W., Chen, X., Wang, Y., Zhang, Y., Zheng, F., Li, J., Hua, C., Zhao, Z., Yuan, Q., Wu, J., Jiang, J., Ye, P., Kulmala, M., Worsnop, D. R., Yang, H., and Liu, Y.: High Aerosol Ionic Strength Possibly Affecting Nocturnal Chlorine Formation via Heterogeneous Reaction in Urban Beijing, *Environmental Science & Technology*. 60, 902-912, <https://doi.org/10.1021/acs.est.5c09187>, 2026.

Ma, X., Liu, H., and Peng, Z.: Improving $\text{PM}_{2.5}$ and PM_{10} predictions in China from WRF Chem through a deep learning method: Multiscale depth-separable UNet, *Environmental Pollution*. 364, 125344, <https://doi.org/10.1016/j.envpol.2024.125344>, 2025.

Mao, J., Li, L., Li, J., Sulaymon, I. D., Xiong, K., Wang, K., Zhu, J., Chen, G., Ye, F., Zhang, N., Qin, Y., Qin, M., and Hu, J.: Evaluation of Long-Term Modeling Fine Particulate Matter and Ozone in China During 2013–2019, *Frontiers in Environmental Science*. Volume 10 - 2022, <https://doi.org/10.3389/fenvs.2022.872249>, 2022.

Nie, W., Yan, C., Huang, D. D., Wang, Z., Liu, Y., Qiao, X., Guo, Y., Tian, L., Zheng, P., Xu, Z., Li, Y., Xu, Z., Qi, X., Sun, P., Wang, J., Zheng, F., Li, X., Yin, R., Dallenbach, K. R., Bianchi, F., Petäjä, T., Zhang, Y., Wang, M., Schervish, M., Wang, S., Qiao, L., Wang, Q., Zhou, M., Wang, H., Yu, C., Yao, D., Guo, H., Ye, P., Lee, S., Li, Y. J., Liu, Y., Chi, X., Kerminen, V.-M., Ehn, M., Donahue, N. M., Wang, T., Huang, C., Kulmala, M., Worsnop, D., Jiang, J., and Ding, A.: Secondary organic aerosol formed by condensing anthropogenic vapours over China's megacities, *Nature Geoscience*. 15, 255-261, <https://doi.org/10.1038/s41561-022-00922-5>, 2022.

Piatkowski, L., Zhang, Z., Backus, E. H. G., Bakker, H. J., and Bonn, M.: Extreme surface propensity of halide ions in water, *Nature Communications*. 5, 4083, <https://doi.org/10.1038/ncomms5083>, 2014.

Qin, M., Xie, P., Su, H., Gu, J., Peng, F., Li, S., Zeng, L., Liu, J., Liu, W., and Zhang, Y.: An observational study of the HONO– NO_2 coupling at an urban site in Guangzhou City, South China, *Atmospheric Environment*. 43, 5731-5742, <https://doi.org/10.1016/j.atmosenv.2009.08.017>, 2009.

639 Qu, R. and Han, G.: A critical review of the variation in rainwater acidity in 24 Chinese cities during
 640 1982–2018, *Elementa: Science of the Anthropocene*. 9, 00142,
 641 <https://doi.org/10.1525/elementa.2021.00142>, 2021.
 642 Seinfeld, J. H., Pandis, S. N., and Noone, K. J.: Atmospheric Chemistry and Physics: From Air
 643 Pollution to Climate Change, *Physics Today*. 51, 88-90, 1998.
 644 Shen, H., Li, Q., Xu, F., Xue, L., Hu, Y., Saiz-Lopez, A., Wang, W., and Wang, T.: Aerosol iodide
 645 accelerates reactive nitrogen cycling in the marine atmosphere, *Nature Communications*. 16, 8148,
 646 <https://doi.org/10.1038/s41467-025-63420-3>, 2025.
 647 Shi, X., Ge, Y., Zheng, J., Ma, Y., Ren, X., and Zhang, Y.: Budget of nitrous acid and its impacts on
 648 atmospheric oxidative capacity at an urban site in the central Yangtze River Delta region of China,
 649 *Atmospheric Environment*. 238, 117725, <https://doi.org/10.1016/j.atmosenv.2020.117725>, 2020.
 650 Stemmler, K., Ammann, M., Donders, C., Kleffmann, J., and George, C.: Photosensitized reduction
 651 of nitrogen dioxide on humic acid as a source of nitrous acid, *Nature*. 440, 195-198,
 652 <https://doi.org/10.1038/nature04603>, 2006.
 653 Su, H., Cheng, Y. F., Cheng, P., Zhang, Y. H., Dong, S., Zeng, L. M., Wang, X., Slanina, J., Shao,
 654 M., and Wiedensohler, A.: Observation of nighttime nitrous acid (HONO) formation at a non-urban
 655 site during PRIDE-PRD2004 in China, *Atmospheric Environment*. 42, 6219-6232,
 656 <https://doi.org/10.1016/j.atmosenv.2008.04.006>, 2008.
 657 Sun, L., Zhang, J., Ducruet, C., Itoh, H., and Liu, X.: The impact of shipping activities on air quality
 658 and residents' health in China's port cities, *Journal of Transport Geography*. 123, 104099,
 659 <https://doi.org/10.1016/j.jtrangeo.2024.104099>, 2025.
 660 Wagner, N. L., Riedel, T. P., Young, C. J., Bahreini, R., Brock, C. A., Dubé, W. P., Kim, S.,
 661 Middlebrook, A. M., Öztürk, F., Roberts, J. M., Russo, R., Sive, B., Swarthout, R., Thornton, J. A.,
 662 VandenBoer, T. C., Zhou, Y., and Brown, S. S.: N₂O₅ uptake coefficients and nocturnal NO₂ removal
 663 rates determined from ambient wintertime measurements, *Journal of Geophysical Research:*
 664 *Atmospheres*. 118, 9331-9350, <https://doi.org/10.1002/jgrd.50653>, 2013.
 665 Wang, G., Zhang, S., Wu, C., Zhu, T., Xu, X., Ge, S., Sun, H., Sun, Z., Wang, J., Ji, Y., Gao, J., Ren,
 666 Y., Li, H., Zhang, F., Wang, Y., and Seinfeld, J. H.: Atmospheric sulfate aerosol formation enhanced
 667 by interfacial anions, *PNAS Nexus*. 4, pgaf058, <https://doi.org/10.1093/pnasnexus/pgaf058>, 2025a.
 668 Wang, H., Lu, K., Guo, S., Wu, Z., Shang, D., Tan, Z., Wang, Y., Le Breton, M., Lou, S., Tang, M.,
 669 Wu, Y., Zhu, W., Zheng, J., Zeng, L., Hallquist, M., Hu, M., and Zhang, Y.: Efficient N₂O₅ uptake
 670 and NO₃ oxidation in the outflow of urban Beijing, *Atmos. Chem. Phys.*, 18, 9705-9721,
 671 <https://doi.org/10.5194/acp-18-9705-2018>, 2018.
 672 Wang, H., Lu, K., Chen, X., Zhu, Q., Chen, Q., Guo, S., Jiang, M., Li, X., Shang, D., Tan, Z., Wu,
 673 Y., Wu, Z., Zou, Q., Zheng, Y., Zeng, L., Zhu, T., Hu, M., and Zhang, Y.: High N₂O₅ Concentrations
 674 Observed in Urban Beijing: Implications of a Large Nitrate Formation Pathway, *Environmental*
 675 *Science & Technology Letters*. 4, 416-420, <https://doi.org/10.1021/acs.estlett.7b00341>, 2017a.
 676 Wang, J., Zhang, X., Guo, J., Wang, Z., and Zhang, M.: Observation of nitrous acid (HONO) in
 677 Beijing, China: Seasonal variation, nocturnal formation and daytime budget, *Science of The Total*
 678 *Environment*. 587-588, 350-359, <https://doi.org/10.1016/j.scitotenv.2017.02.159>, 2017b.
 679 Wang, Y., Liu, P., Zhang, C., Xue, C., Zhao, X., Fu, S., Wu, Z., Song, Y., Wang, Y., Liu, C., and Mu,
 680 Y.: Evaluating the contribution of residential coal combustion to atmospheric nitrous acid (HONO)
 681 in winter, *Environmental Pollution*. 378, 126488, <https://doi.org/10.1016/j.envpol.2025.126488>,
 682 2025b.

Wen, Y., Liu, M., Zhang, S., Wu, X., Wu, Y., and Hao, J.: Updating On-Road Vehicle Emissions for China: Spatial Patterns, Temporal Trends, and Mitigation Drivers, *Environmental Science & Technology*. 57, 14299-14309, <https://doi.org/10.1021/acs.est.3c04909>, 2023.

Wentzell, J. J. B., Schiller, C. L., and Harris, G. W.: Measurements of HONO during BAQS-Met, *Atmos. Chem. Phys.*, 10, 12285-12293, <https://doi.org/10.5194/acp-10-12285-2010>, 2010.

Wolfe, G. M., Marvin, M. R., Roberts, S. J., Travis, K. R., and Liao, J.: The Framework for 0-D Atmospheric Modeling (F0AM) v3.1, *Geosci. Model Dev.*, 9, 3309-3319, <https://doi.org/10.5194/gmd-9-3309-2016>, 2016.

Xie, J., Sun, J., Li, Y., Chan, P. W., Li, L., Huang, C., Tang, L., Fan, S., Zheng, Y., and Fan, Z.: Sea-land breezes in the Guangdong- Hong Kong- Macau Greater Bay Area coastal zone from 2013 to 2022, *Atmospheric Research*. 296, 107044, <https://doi.org/10.1016/j.atmosres.2023.107044>, 2023.

Xu, W., Kuang, Y., Xu, W., Liu, L., Xu, H., Wang, X., Liu, Y., Cheng, H., Zhang, X., Zhai, M., Liu, C., Liang, L., Zhang, G., Luo, B., Tao, J., Liu, J., Zhao, H., Ren, S., Zhou, G., Liu, P., Xu, X., and Sun, Y.: Efficient Nitrate Formation in Fog Events Implicates Fog Interstitial Aerosols as Significant Drivers of Atmospheric Chemistry, *Environmental Science & Technology*. 58, 22298-22311, <https://doi.org/10.1021/acs.est.4c09078>, 2024.

Xu, Z., Wang, T., Wu, J., Xue, L., Chan, J., Zha, Q., Zhou, S., Louie, P. K. K., and Luk, C. W. Y.: Nitrous acid (HONO) in a polluted subtropical atmosphere: Seasonal variability, direct vehicle emissions and heterogeneous production at ground surface, *Atmospheric Environment*. 106, 100-109, <https://doi.org/10.1016/j.atmosenv.2015.01.061>, 2015.

Xuan, H., Liu, C., Zhang, P., Chu, B., Liang, L., Ma, Q., and He, H.: A Review of Laboratory Studies on the Heterogeneous Chemistry of NO₂: Mechanisms and Uptake Kinetics, *The Journal of Physical Chemistry A*. 129, 815-835, <https://doi.org/10.1021/acs.jpca.4c07943>, 2025.

Xuan, H., Zhao, Y., Ma, Q., Chen, T., Liu, J., Wang, Y., Liu, C., Wang, Y., Liu, Y., Mu, Y., and He, H.: Formation mechanisms and atmospheric implications of summertime nitrous acid (HONO) during clean, ozone pollution and double high-level PM_{2.5} and O₃ pollution periods in Beijing, *Science of The Total Environment*. 857, 159538, <https://doi.org/10.1016/j.scitotenv.2022.159538>, 2023.

Xuan, H., Liu, J., Zhao, Y., Cao, Q., Chen, T., Wang, Y., Liu, Z., Sun, X., Li, H., Zhang, P., Chu, B., Ma, Q., and He, H.: Relative humidity driven nocturnal HONO formation mechanism in autumn haze events of Beijing, *npj Climate and Atmospheric Science*. 7, 193, <https://doi.org/10.1038/s41612-024-00745-8>, 2024.

Yang, C., Dong, H., Chen, Y., Wang, Y., Fan, X., Tham, Y. J., Chen, G., Xu, L., Lin, Z., Li, M., Hong, Y., and Chen, J.: Machine Learning Reveals the Parameters Affecting the Gaseous Sulfuric Acid Distribution in a Coastal City: Model Construction and Interpretation, *Environmental Science & Technology Letters*. 10, 1045-1051, <https://doi.org/10.1021/acs.estlett.3c00170>, 2023.

Yang, C., Dong, H., Chen, Y., Xu, L., Chen, G., Fan, X., Wang, Y., Tham, Y. J., Lin, Z., Li, M., Hong, Y., and Chen, J.: New Insights on the Formation of Nucleation Mode Particles in a Coastal City Based on a Machine Learning Approach, *Environmental Science & Technology*. 58, 1187-1198, <https://doi.org/10.1021/acs.est.3c07042>, 2024.

Yang, J., Shen, H., Guo, M.-Z., Zhao, M., Jiang, Y., Chen, T., Liu, Y., Li, H., Zhu, Y., Meng, H., Wang, W., and Xue, L.: Strong marine-derived nitrous acid (HONO) production observed in the coastal atmosphere of northern China, *Atmospheric Environment*. 244, 117948, <https://doi.org/10.1016/j.atmosenv.2020.117948>, 2021.

Ye, C., Zhang, N., Gao, H., and Zhou, X.: Photolysis of Particulate Nitrate as a Source of HONO and NO_x, *Environmental Science & Technology*. 51, 6849-6856, <https://doi.org/10.1021/acs.est.7b00387>, 2017.

Zha, Q., Xue, L., Wang, T., Xu, Z., Yeung, C., Louie, P. K. K., and Luk, C. W. Y.: Large conversion rates of NO₂ to HNO₂ observed in air masses from the South China Sea: Evidence of strong production at sea surface?, *Geophysical Research Letters*. 41, 7710-7715, <https://doi.org/10.1002/2014GL061429>, 2014.

Zhang, R., Minamikawa, R., Gen, M., Singh, N., Daniel, D., Li, Y. J., Wang, X., and Chan, C. K.: Evidence on Interfacial Reaction Governing NO₂ Hydrolysis in Deliquesced Aerosol Particles, *Environmental Science & Technology*. 59, 11708-11719, <https://doi.org/10.1021/acs.est.5c05223>, 2025a.

Zhang, W., Tong, S., Jia, C., Wang, L., Liu, B., Tang, G., Ji, D., Hu, B., Liu, Z., Li, W., Wang, Z., Liu, Y., Wang, Y., and Ge, M.: Different HONO Sources for Three Layers at the Urban Area of Beijing, *Environmental Science & Technology*. 54, 12870-12880, <https://doi.org/10.1021/acs.est.0c02146>, 2020.

Zhang, X., Tong, S., Jia, C., Zhang, W., Wang, Z., Tang, G., Hu, B., Liu, Z., Wang, L., Zhao, P., Pan, Y., and Ge, M.: Elucidating HONO formation mechanism and its essential contribution to OH during haze events, *npj Climate and Atmospheric Science*. 6, 55, <https://doi.org/10.1038/s41612-023-00371-w>, 2023.

Zhang, X., Tong, S., Jia, C., Zhang, W., Li, J., Wang, W., Sun, Y., Wang, X., Wang, L., Ji, D., Wang, L., Zhao, P., Tang, G., Xin, J., Li, A., and Ge, M.: The Levels and Sources of Nitrous Acid (HONO) in Winter of Beijing and Sanmenxia, *Journal of Geophysical Research: Atmospheres*. 127, e2021JD036278, <https://doi.org/10.1029/2021JD036278>, 2022.

Zhang, Y., Zheng, F., Feng, Z., Lian, C., Wang, W., Fan, X., Ma, W., Lin, Z., Li, C., Zhang, G., Yan, C., Zhang, Y., Kerminen, V. M., Bianchi, F., Petäjä, T., Kangasluoma, J., Kulmala, M., and Liu, Y.: Concentration and source changes of nitrous acid (HONO) during the COVID-19 lockdown in Beijing, *Atmos. Chem. Phys.*, 24, 8569-8587, <https://doi.org/10.5194/acp-24-8569-2024>, 2024.

Zhang, Y., Liu, Y., Ma, W., Hua, C., Zheng, F., Lian, C., Wang, W., Xia, M., Zhao, Z., Li, J., Xie, J., Wang, Z., Wang, Y., Chen, X., Zhang, Y., Feng, Z., Yan, C., Chu, B., Du, W., Kerminen, V.-M., Bianchi, F., Petäjä, T., Worsnop, D., and Kulmala, M.: Changing aerosol chemistry is redefining HONO sources, *Nature Communications*. 16, 5238, <https://doi.org/10.1038/s41467-025-60614-7>, 2025b.

Zheng, J., Shi, X., Ma, Y., Ren, X., Jabbour, H., Diao, Y., Wang, W., Ge, Y., Zhang, Y., and Zhu, W.: Contribution of nitrous acid to the atmospheric oxidation capacity in an industrial zone in the Yangtze River Delta region of China, *Atmos. Chem. Phys.*, 20, 5457-5475, <https://doi.org/10.5194/acp-20-5457-2020>, 2020.

Zhou, X., Zhang, N., TerAvest, M., Tang, D., Hou, J., Bertman, S., Alaghmand, M., Shepson, P. B., Carroll, M. A., Griffith, S., Dusanter, S., and Stevens, P. S.: Nitric acid photolysis on forest canopy surface as a source for tropospheric nitrous acid, *Nature Geoscience*. 4, 440-443, <https://doi.org/10.1038/ngeo1164>, 2011.