

Supplementary Material

For

Linking the effects of environmental factors and heterotrophic bacteria to the variability of carbon isotope ($\delta^{13}\text{C}$) across crustacean zooplankton in shallow freshwater ecosystems

authored by

Wojciech Krztoń¹, Edward Walusiak¹, Iwona Gołaś², Elżbieta Wilk-Woźniak¹

¹Institute of Nature Conservation Polish Academy of Sciences, al. Adama Mickiewicza 33, 31-120 Kraków, Poland

²Department of Water Protection Engineering and Environmental Microbiology, Institute of Engineering and Environmental Protection, Faculty of Geoengineering, University of Warmia and Mazury in Olsztyn, ul. Prawocheńskiego 1, 10-720 Olsztyn, Poland

Corresponding author:

Dr. Wojciech Krztoń

Email: krzton@iop.krakow.pl

ORCID: 0000 – 0003 – 2224 – 2599

Table S1. One-way Analysis of Variance (ANOVA) of $\delta^{13}\text{C}$ values in studied groups of planktonic crustaceans

	Df	Sum of Squares	Mean Square	F value	Pr(>F)
group	2	39	19.50	1.081	0.342
Residuals	121	2182	18.03		

Table S2. Tukey HSD test for $\delta^{13}\text{C}$ vs. groups ANOVA

Group	Difference	Confidence Interval		Adjusted p-value
		Lower	Upper	
cyclopoida-calanoida	1.2673441	-0.7790798	3.313768	0.3091662
daphnia-calanoida	0.5897779	-1.8343555	3.013911	0.8324815
daphnia-cyclopoida	-0.6775662	-3.0927782	1.737646	0.7838003