



U-Pb dating of sub-ng/g U garnet by LA-MC-ICPMS

Aratz Beranoaguirre^{1,2}, Leo J. Millonig^{1,2}, Richard Albert^{1,2}, Horst R. Marschall^{1,2}, and Axel Gerdes^{1,2}

¹Frankfurt Isotope and Element Research Center (FIERCE), Goethe-University Frankfurt, 60438 Frankfurt am Main, Germany

²Goethe-University Frankfurt, Department of Geosciences, 60438 Frankfurt am Main, Germany

Correspondence to: Aratz Beranoaguirre (beranoaguirre@fierce.uni-frankfurt.de)

Abstract. Advances in laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) have largely focused on improving spatial resolution through progressively smaller laser spot sizes. Here, we explore the opposite end of the analytical limits by investigating the lower concentration limits of in-situ U–Pb geochronology in garnet. Using a Neptune Plus multi-collector ICP-MS equipped with seven ion counters, we developed an analytical workflow specifically designed for metamorphic garnet with ultra-low U concentrations. The method was used to date garnet from a wide range of ages and geological settings, including granulites, eclogites and hydrothermal demantoids. Uranium concentrations were exceptionally low, even below 1 ng/g in some cases. At these concentrations, the total amount of uranium ablated during a single analysis is at femtogram levels, more than three orders of magnitude lower than that of a typical in-situ analysis of zircon. Despite these extremely low signal intensities, geologically meaningful ages were obtained for the analysed samples. Analytical precision depends on the U and radiogenic Pb concentrations, but we have obtained precisions of ca. 5–6 % for garnet containing less than 1 ng/g U. In garnet with higher U concentrations or ages as old as the Archean, internal precision of ca. 1 % may be achieved. These results expand the applicability of in-situ garnet U–Pb geochronology to the vast majority of metamorphic garnet, providing a powerful new tool for constraining garnet growth, prograde metamorphism, and deep crustal evolution.

1 Introduction

The combination of laser ablation (LA) systems with inductively coupled plasma mass spectrometers (ICP-MS) during the 1980s revolutionised in-situ chemical analysis (Gray, 1985; Arrowsmith, 1987). Since the pioneering works, LA-ICP-MS U–Pb geochronology of high-U accessory minerals such as zircon, monazite, or titanite has become a routine analytical tool (e.g., Fryer et al., 1993; Machado and Gauthier, 1996; Willigers et al., 2002; Kosler and Sylvester, 2003; Horstwood et al., 2016). Besides, over the last decade, these analytical capabilities have also been expanded to minerals with lower uranium concentrations (<10 µg/g), including carbonate (Roberts and Walker, 2016; Ring and Gerdes, 2016), sulfate (Beranoaguirre et al., 2022), fluorite (Piccione et al., 2019), and garnet (Seman et al., 2017; Millonig et al., 2020; O’Sullivan et al., 2023), among others. Continuous advances in instrumentation have also been accompanied by progressively smaller laser spot sizes (review in Sylvester and Jackson, 2016), improving the spatial resolution and enabling the distinction of multiple domains within a single crystal (e.g., Chew et al., 2017). In turn, analysing low-U minerals requires a different strategy, in which the main challenge is no longer spatial resolution, but concentration sensitivity. This issue is particularly challenging for metamorphic garnet, since its U and Pb amounts are extremely low (Deng et al., 2022). Therefore, being



capable of developing a robust analytical strategy for analysing metamorphic garnet would represent a major advance in metamorphic petrology.

Unlike many geochronological systems that either record cooling and exhumation histories (e.g., apatite fission-track, (U–Th)/He, and Ar-based systems) or preferentially date high-temperature metamorphic events (e.g., zircon, monazite, and titanite U–Pb geochronology) (Dodson, 1973; McDougall and Harrison, 1999; Rubatto, 2002; Reiners et al., 2005; Flowers et al., 2009; Kohn et al., 2017), garnet can preserve prograde metamorphic growth histories over a wide range of pressure–temperature conditions (Spear and Parrish, 1996; Caddick and Kohn, 2013), in part due to its exceptionally high closure temperature for the U–Pb system (potentially exceeding 1000 °C; Dahl, 1997; Shu et al., 2024). Traditionally, garnet has been dated by Sm–Nd and Lu–Hf (e.g. Duchêne et al., 1997; Baxter et al., 2017). Although a very high precision can be achieved through these systems, they require laborious mineral separation and dissolution procedures. More recently, the use of tandem mass spectrometry (ICP-MS/MS) has also enabled the in-situ Lu–Hf garnet dating (Simpson et al., 2021). However, the application of in-situ Lu–Hf geochronology remains largely restricted to relatively old metamorphic systems (Caledonian or older; Tamblyn et al., 2022; Kirkland et al., 2025). This limitation may arise from the low radiogenic growth of ^{176}Hf in younger garnet, which hampers analytical precision and age resolution in younger orogens. In contrast, U–Pb garnet geochronology has been used to resolve younger metamorphic events (Peillod et al., 2024; Manzotti et al., 2025). Consequently, U–Pb garnet geochronology (eventually complemented with in-situ Lu–Hf dating) offers a unique opportunity to directly constrain the timing of prograde metamorphism and deep crustal processes.

In this study, we investigate the lower limits of the uranium and lead concentrations in garnet required to obtain meaningful garnet U–Pb dates by LA-MC-ICPMS. For that purpose, we have optimised the LA-MC-ICPMS method on a series of natural, low-U metamorphic garnet specimens from granulites, eclogites, and one gemstone-quality grossular-andradite (i.e. demantoid). The possibility to date garnet crystals with extremely low U concentrations (<1 ng/g) by LA-MC-ICPMS is expected to substantially increase the field of application of garnet U–Pb dating, as most of the metamorphic garnet have <50 ng/g U (Millonig et al., 2020; O’Sullivan et al., 2023; Bartoli et al., 2024; Shu et al., 2024).

2 Methodology

U–Pb data were acquired *in-situ* from polished garnet grain mounts and thin sections using a RESOLution 193 nm ArF excimer laser (CompexPro 102) equipped with a two-volume ablation cell (Laurin Technic S155) coupled to a multi-collector (MC)-ICPMS (Neptune Plus, ThermoScientific) at the Frankfurt Isotope and Element Research Center (FIERCE), Goethe University Frankfurt. The Neptune Plus at FIERCE is equipped with ten Faraday cups and seven ion counters, of which two are classical discrete dynode secondary electron multipliers (SEM) and five compact discrete dynode (CDD) electron multipliers. This results in two possibilities for the simultaneous detection of the U, Th, and Pb isotopes: I) for moderate U contents of ~ 0.2 to 30 $\mu\text{g/g}$, ^{238}U and ^{232}Th are detected on the Faraday cups with 10^{13} Ω amplifiers, ^{207}Pb , ^{206}Pb on the SEM and ^{208}Pb on a CDD (method described in Beranoaguirre et al., 2022); while II) at low U of < 0.2 $\mu\text{g/g}$, ^{238}U and, if necessary



64 ^{232}Th , can be measured on the CDD attached to the Faraday cups (H4 and H3, respectively). A summary report of the U-Pb
65 dating procedure is presented in the supplementary material (Table S1), and the analytical results are presented in
66 supplementary Tables S2 to S6.

67 Ablation was performed in a He atmosphere (0.3 l/min) and mixed in the ablation funnel with daily tuned 0.95–1 l/min
68 Ar and 5–10 ml/min N_2 . Signal strength at the MC-ICPMS was tuned for maximum sensitivity while keeping oxide formation
69 below 0.5 % (UO/U) and element fractionation low (e.g. Th/U $\sim$ 0.9). This was done by line ablating the SRMNIST 614 glass
70 (Jochum et al., 2011) using a 50 μm spot size, 6 Hz, *ca.* 3.5 J/cm² fluence and a 3 $\mu\text{m/s}$ line speed. The average sensitivity
71 obtained with this setup was *ca.* 250,000 cps per $\mu\text{g/g}$ for ^{238}U in the best of the cases. The garnet samples were ablated using
72 a round spot size of 193 μm and a fluence of *ca.* 2 J/cm² at 15 Hz. This yielded a depth penetration of *ca.* 0.8 $\mu\text{m/s}$. Each
73 analysis consisted of 16 s of background acquisition followed by 18 s of sample ablation and 20 s of washout. The analyses
74 were done in the static mode, measuring ^{206}Pb and ^{207}Pb with SEMs, and ^{202}Hg , ^{204}Pb , (^{208}Pb) and ^{238}U with the CDD and an
75 integration time of 0.131 s (due to electronic issues with the ion counter, ^{208}Pb was not measured in all the sessions).

76 In each analytical session, soda-lime glass SRMNIST 614 was used as the primary reference material to correct for
77 mass bias ($^{207}\text{Pb}/^{206}\text{Pb}$), inter-element fractionation and instrumental drift ($^{206}\text{Pb}/^{238}\text{U}$) throughout the analytical sessions. Mali
78 garnet (Seman et al., 2017) was used as a matrix-matched reference material to determine the difference in the Pb/U
79 fractionation between garnet and the synthetic glass matrix. As the CDDs saturate at around 1,000,000 cps, we have used a
80 low-uranium (*ca.* 20 ng/g) yellow-coloured Mali garnet from our collection (see below). Secondary reference garnet materials,
81 Lake Jaco garnet (34.0 ± 1.4 Ma, Seman et al., 2017), and an in-house garnet from Balochistan (TIMS age of *ca.* 45.5 Ma,
82 Maria Stiffee, pers. comm.) were measured for quality control. Due to the possible saturation of the ion counters, the
83 secondary garnet crystals were pre-screened to identify domains with sufficiently low-U concentrations. Those domains also
84 show lower Pb content, and therefore, the $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{206}\text{Pb}/^{238}\text{U}$ ratios are not affected.

85 Raw data were corrected offline using an in-house VBA spreadsheet program (Gerdes and Zeh, 2006, 2009).
86 Following background and interference corrections, outliers were rejected based on the time-resolved $^{207}\text{Pb}/^{206}\text{Pb}$ and
87 $^{238}\text{U}/^{206}\text{Pb}$ ratios, while the Pb and U signals were used to identify and reject data derived from the co-ablation of mineral
88 inclusions or zones with clearly distinct common-Pb and U concentrations compared to the pure garnet. Likewise, outliers in
89 U and/or Pb content were rejected for age calculations, assuming that those analyses reflect inclusions (Fig. 1). Data are
90 displayed in Tera-Wasserburg plots (Tera and Wasserburg, 1972), and U-Pb dates were calculated as lower Concordia-curve
91 intercepts using the same algorithms as Isoplot 4.15 (Ludwig, 2012). Age uncertainties shown in the Tera-Wasserburg plots
92 are the within-session uncertainty, considering the within-run precision, counting statistic uncertainties of each isotope, and
93 the excess of scatter and of variance (Horstwood et al., 2016), calculated from the SRMNIST 614 and the Mali garnet, and the
94 expanded uncertainty, which includes the long-term variance of our in-house quality control garnet crystals (2%). All
95 uncertainties are reported at the 2σ level.

96




Figure 1. Plots of U vs Pb concentration (ng/g), showing the analyses rejected and considered for the age calculations. A) Cabo Ortegal demantoid (session 2) and B) garnet G99-2 from Orlica-Śnieżnik felsic granulite.

Matrix-matched reference materials (RMs) are an important component of any LA-ICPMS method (e.g. Schaltegger et al., 2015). With respect to garnet, only a few well-characterised RMs are available to the LA-ICPMS community in sufficient quantities, although many research groups have developed new RMs over the last few years (e.g. Stifreeva et al., 2019, Salnikova et al., 2019, 2026; Li et al., 2022; Aysal et al., 2023; Beno et al., 2024, Yang et al., 2025). The most popular garnet RMs for U–Pb dating are still the ones published by Seman et al. (2017); the red and yellow varieties of Mali Grandite (LA-ICPMS U–Pb: 202 ± 2 Ma; ID-TIMS U–Pb: 202.0 ± 1.2 Ma) and the Lake Jaco Grossular (LA-ICPMS U–Pb: 35 ± 2 Ma; (U–Th)/He: 35 ± 5 Ma). However, U concentrations of $>2 \mu\text{g/g}$ of the aforementioned garnet RMs would saturate the ion counters, and thus, they are not usable for the method applied in this study. To overcome this issue, we investigated several low-U ($<0.5 \mu\text{g/g}$) garnet from the same localities investigated by Seman et al. (2017), and we found a yellow Mali Grandite with an average U content of 15 ng/g, which was subsequently used as the matrix-matched RM. The secondary reference materials, used for quality control, were compositionally and optically diverse grossular garnet crystals from Lake Jaco, and a dark brown grandite from Balochistan. When assuming the published age of 202 Ma for the yellow Mali, we obtained internally consistent results across multiple sequences for the secondary garnet RMs and different samples. Although no independent age estimates (i.e. TIMS ages) are available for our matrix-match and quality control RMs we consider the obtained U–Pb ages as accurate within their analytical uncertainties, based on the facts that (1) the U–Pb ages for the various Lake Jaco garnet crystals are within the uncertainty of the published ages from that locality (Seman et al., 2017), and (2) the U–Pb age of 384 ± 12 Ma for garnet from sample G99-2 is similar to a Lu–Hf age of 387 ± 5 Ma from that sample (Anczkiewicz et al., 2007; section 3.2). However, we will (re-)calibrate these low-U garnet crystals against well-characterised low-U garnet RMs, once they become available. Furthermore, this potential age ambiguity does not affect the purpose of this contribution.



3 Samples and results

3.1 Cabo Ortegal demantoid (Cabo Ortegal Complex, NW Iberian Massif)

The Cabo Ortegal Complex represents an allochthonous assemblage of continental- and oceanic-derived lithologies that record subduction-related metamorphism prior to their emplacement onto the Gondwanan margin during the Variscan orogeny (e.g. Martínez Catalán et al., 2019). Previous geochronological studies of the complex suggest that the magmatic protoliths (520–460 Ma) were metamorphosed during the Variscan at *ca.* 400–390 Ma followed by a rapid decompression to shallow crustal levels (Beranoaguirre et al., 2020). The demantoid andraditic garnet analysed here was described in detail by Madon et al. (1991). It occurs associated with chlorite and serpentine in fissures cutting across the foliation of the pyroxenite.





Figure 2. Tera-Wasserburg diagrams of garnet U-Pb analyses for Cabo Ortegal Demantoid, analysed with the ElementXR single-collector ICPMS (a) and Neptune multi-collector ICPMS (b and c), and sample G99-2 from Bohemian massif (d).



132 The first attempt to date the demantoid garnet (crystals was made using a single-collector ICPMS (the method is
133 described in Millonig et al., 2020 and Beranoaguirre et al., 2022). The resulting data defined a regression line with a lower
134 intercept at 284 ± 65 Ma (MSWD = 2, Fig. 2a), and only 43 out of 72 analytical spots had U or Pb concentrations above the
135 detection limits. Subsequently, two sessions were performed with the MC-ICPMS. The resulting lower intercept dates for the
136 sessions were $254.7 \pm 7.9/8.7$ Ma (MSWD = 1.19, $n = 47/64$, Fig. 2b) and $255.3 \pm 9.8/10.5$ Ma (MSWD = 0.37, $n = 31/40$,
137 Fig. 2c). The regression lines were anchored to a $^{207}\text{Pb}/^{206}\text{Pb}$ ratio of 0.870 ± 0.005 , derived from the mean of the unanchored
138 upper intercept of both sessions. Most of the analyses have extremely low U concentrations between 0.5 and 1 ng/g, with rare
139 exceptions up to 4 ng/g (Fig. 1a and 3; Tables S2 and S3).



140

141 **Figure 3.** Plot of U vs Pb concentration (ng/g) for all the analysed garnet. The rejected analyses (Fig. 1) are not considered, to avoid
142 scaling issues. Note the difference on the x-axis scale for the samples L212 and L02100. In the Kaapvaal samples, ^{208}Pb was not
143 measured and therefore, the Pb concentration is the sum of $^{204}\text{Pb}+^{206}\text{Pb}+^{207}\text{Pb}$.

144 3.2 Orlica-Śnieżnik felsic granulite (NE Bohemian Massif)

145 The Orlica-Śnieżnik Dome is located on the northeastern margin of the Bohemian Massif in SW Poland. It comprises
146 predominantly amphibolite-facies, partly migmatized, orthogneisses, hosting numerous inclusions of (ultra) high-pressure



rocks (e.g. Walczak et al., 2017). Their granitic precursors intruded supracrustal sedimentary successions of Neoproterozoic to Ordovician age and were subsequently metamorphosed during the Variscan orogeny (Szczepański and Ilnicki, 2014).

The garnet analysed here corresponds to the felsic granulite G99-2 studied in detail by Anczkiewicz et al. (2007), who obtained garnet Lu-Hf and Sm-Nd ages of 387 ± 5 Ma and 320 ± 3 Ma, respectively. From the U-Pb analyses performed in our laboratory, an unanchored lower intercept age of $384.0 \pm 10.0/11.5$ Ma (MSDW = 0.52, $n = 46$, Fig. 2d) was calculated. The U and Pb contents are homogeneous, varying between 2 and 4 ng/g, and between 0.3 and 3 ng/g, respectively (Fig. 1b and 3; Table S4).

3.3 Granulite xenoliths from the Star kimberlite (Kaalpvaal craton, South Africa)

Ultrahigh temperature (UHT) metamorphic crustal granulite xenoliths were found in four kimberlite diatremes from along the central axis of the Witwatersrand basin in the central Kaapvaal craton (Schmitz and Bowring, 2003). The xenoliths are K-free, Mg-Al-rich UHT granulites with abundant garnet, sillimanite, rutile, graphite and sulphides, and local occurrence of sapphirine, orthopyroxene, plagioclase and quartz. Metamorphic conditions have been estimated at temperatures above 1050°C and pressures between 0.9–1.2 GPa (Dawson et al., 1997). Zircon and monazite gave U–Pb ages of approximately 2.7 Ga (Schmitz and Bowring, 2003). The xenoliths were brought to the surface by Cretaceous kimberlite magmatism at approximately 120 Ma.


The results of these samples have already been published and discussed in detail by Shu et al. (2024). The analyses of the garnet of that work followed the method described here, and considering that the aim of this work is purely methodological, two examples of those rocks are shown: the one with the lowest U concentration (sample ST66) and the one with the highest precision (ST 70). The samples ST66 and ST70 define regression lines with lower intercepts at $3096 \pm 63/70$ Ma (MSWD = 2.18, $n = 53/53$, Fig. 4a) and $3096 \pm 37/48$ Ma (MSWD = 3.05, $n = 53/58$, Fig. 4b). As stated above, sample ST66 shows the lowest U content among the samples presented in this work, varying from 0.3 to 2.2 ng/g, whereas sample ST70 has a more variable U content, between 1 and 17 ng/g. Pb and U contents are positively correlated. Unfortunately, it was not possible to measure the ^{208}Pb isotope, so the calculated Pb content is the sum of $^{204}\text{Pb}+^{206}\text{Pb}+^{207}\text{Pb}$. The resulting Pb concentrations do not exceed 1 ng/g for sample ST66 and 6 ng/g for sample ST70, except in two spots (Fig. 3; Table S5).

3.4 Eclogite xenoliths from the Namaqua-Natal Belt (South Africa)

Multiple Cretaceous kimberlite pipes occurring on the Proterozoic Namaqua-Natal-Natal Belt, which surrounds the Archean Kaapvaal craton, have brought to the surface a significant amount of high-pressure eclogite, granulite and pyroxenite xenoliths (Le Roex et al., 2020). The eclogites are mainly biminerallitic (garnet-clinopyroxene) with accessory rutile and coesite/quartz (and even diamond), although kyanite-bearing eclogites are also described. As for the Kaapvaal craton granulites, we have analysed several samples from different kimberlites of the area, but only two of them are discussed in the present manuscript. Data points from sample L02100 define a regression line with a lower intercept at $102.7 \pm 3.6/3.9$ Ma (MSWD = 1.05; $n = 37/41$; Fig. 4c), whereas the analyses of sample L212 define a regression line with a lower intercept at



179 $105.5 \pm 2.5/2.9$ Ma (MSWD = 0.92; $n = 34/48$; Fig. 4d). Both samples have relatively homogeneous U contents of 14–16 ng/g
180 on average. Sample L212 also shows a homogeneous Pb content between 0.5 and 4 ng/g, whereas sample L02100 display a
181 higher variability with up to 14 ng/g (Fig. 3, Table S6).



182
183 **Figure 4. Tera-Wasserburg diagrams of garnet U-Pb analyses for the samples from Kaapvaal craton ST66 and ST70**
184 **(a and b) and Namaqua-Natal Belt, L02100 and L212 (c and d, respectively).**

185 4 Discussion

186 4.1 Bulk Uranium content: zircon vs. garnet

187 One of the main strengths of the LA-ICPMS technique is its good spatial resolution. In contrast with the bulk dating
188 method (TIMS), the laser ablation method only analyses a tiny volume of the targeted mineral, and therefore, the bulk amount
189 of U analysed is limited. Besides, not all the ablated material reaches the detector, and the useful yield for U (=ions



190 detected/total number of atoms in sample volume for a species of interest, Schaltegger et al., 2015) is variable depending on
191 the utilised instrument.

192 In the case of this study, the ablation of garnet with a 193 μm spot size, 15 Hz and $\sim 2 \text{ J/cm}^2$ for 18 seconds results in
193 a *ca.* 15 μm deep ablation pit. This corresponds to a total volume of *ca.* 600,000 μm^3 or 2.3 μg of ablated material, assuming
194 a mean garnet density of 3.9 g/cm^3 (Table 1). Li et al. (2022) recently reported skarn garnet with up to 75 $\mu\text{g/g}$ U. Applying
195 the above calculations, this corresponds to *ca.* 100 pg U per ablated spot, and *ca.* 8 pg U and 0.4 pg Pb for a Mali garnet (6
196 $\mu\text{g/g}$ U and 0.3 $\mu\text{g/g}$ Pb, Seman et al., 2017) analysis. However, the metamorphic garnet rarely exceeds U concentrations of
197 $\sim 100 \text{ ng/g}$ (Deng et al., 2022). For example, a regional metamorphic garnet with 90 ng/g U (sample SS-1 of Millonig et al.,
198 2020) would yield *ca.* 0.1 pg U per analytical spot. In the present study, sample ST66 has the lowest U concentrations with an
199 average concentration of 1.1 ng/g . The total Uranium ablated during such a spot analysis is *ca.* 1 fg (femtogram; 10^{-15} g). This
200 is *ca.* 3,500 times less U than for a renowned GJ-1 zircon analysis (287 $\mu\text{g/g}$ U_{avg} ; 26 $\mu\text{g/g}$ Pb_{avg} , Jackson et al., 2004), which
201 yields 3.5 pg of U and 0.30 pg of Pb per spot, calculated for a 20 μm spot size, 15 μm pit depth and *ca.* 10 ng of ablated
202 material (considering a zircon density of 4.65 g/cm^3) (Table 1).

Material / Sample	U concentration (ng/g)	Total U per spot (pg)
GJ-1 zircon (Jackson et al., 2004)	287,000	3.5
Mali garnet (Seman et al., 2017)	6,000	~ 8
Skarn garnet (Li et al., 2022)	75,000	~ 100
Metamorphic garnet SS-1 (Millonig et al., 2020)	90	~ 0.10
Namaqua–Natal eclogite L212	14	~ 0.032
Orlica–Śnieżnik granulite (G99-2)	2–4	$\sim 0.003\text{--}0.009$
Cabo Ortegal demantoid	1.3	$\sim 0.001\text{--}0.002$
Kaapvaal granulite ST66	1.1	~ 0.001

203

204 **Table 1. Comparison of Uranium content analysed in a single LA-ICP-MS spot analysis for different minerals.**
205 **Calculations are done for the regular ablation parameters (see text for further details). The figure illustrates the**
206 **extreme reduction in available parent isotope mass when dating low-U metamorphic garnet, with the ST66 granulite**
207 **and Cabo Ortegal demantoid analyses containing only $\sim 1\text{--}3 \text{ fg}$ U per spot, approximately 3,500 times less Uranium**
208 **than a typical GJ-1 zircon.**



209

210 Samples with even lower amounts of U could be analysed, if they are older and have higher ^{238}U than ^{206}Pb for the
211 majority of the spots (as well as $^{206}\text{Pb} > ^{207}\text{Pb}$), with the exception of the high common-Pb analyses. As an example, a 200 Ma
212 old concordant analysis will have *ca.* 30 times more ^{238}U than ^{206}Pb and *ca.* 20 times more ^{206}Pb than ^{207}Pb (or 600 times more
213 ^{238}U than ^{207}Pb). In turn, a 3 Ga sample intercepts the Concordia at *ca.* 1.5 ($^{238}\text{U}/^{206}\text{Pb}$) and 0.25 ($^{207}\text{Pb}/^{206}\text{Pb}$), the ^{238}U being
214 only *ca.* 6 times more abundant than ^{207}Pb . Hence, for such old samples, it would be possible to date a garnet with even lower
215 U concentrations.

216 4.2 Background level, limit of detection, limit of quantification and analytical precision

217 Such low U and Pb concentrations in garnet correspond to a few hundred counts per second (cps), and therefore, the
218 background level plays a vital role, especially for ^{207}Pb , the least abundant of the measured isotopes. Background Pb may come
219 from different sources, such as a memory effect in the tubing, contamination/dirt in gas-line connections, the cones or the laser
220 cell. Although careful cleaning of those parts is effective in reducing their Pb contribution, Pb derived from the Ar and He gas
221 supplies is unavoidable. This contribution varies from day to day (Fig. 5 shows the background level at the FIERCE laboratory
222 observed over one and a half years), and it is a critical factor for the analytical performance. Despite the variations in
223 background intensity between analytical sessions, the background is generally stable within a single session, which allows low
224 limits of detection. Following the approach proposed by Pettke et al. (2012), the limits of detection for average background
225 signals of 10 cps and 400 cps are approximately 6 pg/g and 30 pg/g, or 2 to 8 cps, respectively (considering the aforementioned
226 sensitivity of 250,000 cps per $\mu\text{g/g}$). However, the limit of quantification (Currie, 1995), defined as 10 times the standard
227 deviation of the measured background signal, drastically increases with increasing background intensity. For example, during
228 low-background sessions, the standard deviation of the ^{207}Pb background can be as good as 5-6 cps, corresponding to a limit
229 of quantification of approximately 50–60 cps (Fig.6). In contrast, during one of the highest-background sessions, the standard
230 deviation is *ca.* 200 cps, increasing the limit of quantification to 2000 cps. Thus, a high Pb background prevents reliable
231 analysis of low ^{207}Pb samples, as their signal falls below the quantification limit. In practice, samples with ultra-low-U and -
232 Pb garnet are only analysed during sessions in which the ^{207}Pb background remains below 50 cps.


233 However, even though the Pb can be reliably detected, the precision of each analysis is also crucial. Figure 6 shows
234 the relationship between the ^{207}Pb signal intensity and the within-run precision of the $^{207}\text{Pb}/^{206}\text{Pb}$ ratio, over several sequences
235 covering a large range of ^{207}Pb concentrations. As could be expected from counting statistics, a lower ^{207}Pb signal results in a
236 lower precision. Analyses yielding ^{207}Pb signals above 2,500 cps commonly achieve uncertainties of less than 1% (2 SE),
237 whereas signals below 2,500 cps typically result in uncertainties above 1% (2 SE). Furthermore, as the signal gets closer to the
238 limit of quantification (≤ 100 cps), there is a strong increase in the uncertainty, which highlights the limit of the technique.

239 Another potential source of uncertainty, although it is not considered in our calculations, is the variance of the
240 Uranium decay, or in other words, the statistical uncertainty coming from the decay of a small number of radioactive atoms.
241 In the case of dating high-U minerals, as well as old samples that have been decaying for a long time, such uncertainty can be



242 considered negligible and that is likely why it is not considered in Geochronology. However, physicists studying slow decay
243 processes have discussed this phenomenon (e.g. Opendak and Wildenhain, 1994). In the case of low-U and young minerals
244 (<1 ng/g U or <10 Ma old), this incertitude may be of importance and needs to be considered. Nonetheless, young samples
245 with such a low content of uranium are currently not measurable with the analytical setup discussed in this study, as the amount
246 of radiogenic ^{207}Pb is below the limit of detection.

247 In addition to the analytical precision of each individual spot measurement, the precision of U-Pb dating for non-
248 concordant minerals like garnet when using a Tera-Wasserburg diagram is also controlled by how accurately the regression
249 line can be defined. This strongly depends on the variability of the $^{238}\text{U}/^{206}\text{Pb}$ and $^{207}\text{Pb}/^{206}\text{Pb}$ ratios; the larger the spread in
250 these ratios, the smaller will be the uncertainty on the lower intercept of the regression line with the Concordia curve and, thus,
251 the more precise the obtained age will be. The internal precision for the analyses of the Cabo Ortegal demantoid are $\pm 4.7\%$
252 and $\pm 6.3\%$ for the absolute garnet ages, even with only 40 analytical spots and an extremely low U concentration (mainly
253 below 1 ng/g, Fig. 3). In turn, U concentrations of 2–4 ng/g, as for the Orlica-Śnieżnik felsic granulite garnet, significantly
254 improve the precision to ca. $\pm 3\%$ absolute age uncertainty. Furthermore, the analyses of sample ST70 resulted in an internal
255 age precision of ca. $\pm 1\%$. In this case, two factors concurred to result in such precise data: I) higher U and Pb contents and
256 II) the Archean age of the sample, as older samples benefit the most from the technique due to their more abundant daughter
257 isotopes ^{206}Pb and ^{207}Pb . However, in samples as young as the Cretaceous Namaqua-Natal belt eclogites, we have obtained
258 precisions of ca. $\pm 3\%$ at U concentrations of ca. 14–16 ng/g. The results obtained for the demantoid in two independent
259 sessions are reproducible within uncertainty. Likewise, the U-Pb age of the garnet from Orlica-Śnieżnik felsic granulite, 384
260 ± 12 Ma, is in agreement with the Lu-Hf age reported by Anczkiewicz et al. (2007, 387 ± 5 Ma). Thus, the procedure described
261 in this study produces accurate results within the internal precision of the technique.



262
263 **Figure 5. ^{207}Pb background (in cps) at the FIERCE laboratory during a period of ca. one and a half years. The spots**
264 **show the average ^{207}Pb background for each of the days.**




Figure 6. ^{207}Pb signal intensity (blue symbols) and calculated limit of quantification (orange symbols) plotted against the within-run precision of the $^{207}\text{Pb}/^{206}\text{Pb}$ ratio (2SE, %). The limit of quantification was calculated as ten times the standard error of the measured ^{207}Pb background signal, following Currie (1995). The correlation between the ^{207}Pb intensity and the within-run precision is clearly observable.

4.3 Applications of the technique

U-Pb dating of garnet is becoming a relatively common tool, above all for skarn garnet, as they typically contain several $\mu\text{g/g}$ of U (e.g., Reinhardt et al., 2022). However, the applicability of the technique is rather limited in metamorphic, typically almandine-pyrope, garnet due to their low U content (Millonig et al., 2020; Cerva-Alves et al., 2021; Schannor et al., 2021; O’Sullivan et al., 2023; Mark et al., 2023; Peillod et al., 2024; Manzotti et al., 2025), and the common presence of high-U mineral inclusions. The method described here allows for the analysis of very low-U garnet, or a reduction in laser spot sizes to avoid inclusions (cf. Walters et al., 2025) and thus has enormous potential to become a routine tool for petro(chrono)logical investigations.



279 Moreover, due to the high closure temperature of the U-Pb system in garnet (Shu et al., 2024), it appears to be the
280 most reliable method for dating its formation and thus the associated mineral reactions. In many cases, this cannot be
281 determined by traditionally datable minerals such as zircon, rutile, or monazite because of I) their lower closure temperatures
282 (in some contexts/minerals) or II) their tendency to recrystallise (e.g., Rubatto, 2017), in addition to the fact that they rarely
283 occur in mafic rocks. The granulite and eclogite xenoliths from the Kaapvaal craton and Namaqua-Natal belt analysed here
284 are good examples, similar to the peridotitic garnet dated by O’Sullivan et al. (2023). Even though scarce rutile has been found
285 in a few samples, the most common mineral assemblage is formed only by garnet and clinopyroxene. In addition, the low U
286 content and their small grain sizes do not necessarily make metamorphic rutile a better geochronometer. In this work, we have
287 successfully dated garnet from those geological contexts, despite their very low U contents. Larger sample sets of both the
288 Kaapvaal craton granulites and the Namaqua-Natal belt eclogites have been analysed, and the results are reported and discussed
289 in Shu et al. (2024).

290 Likewise, at present, the only possible way to date the hydrothermal garnet-chlorite-serpentine veins crosscutting the
291 pyroxenites at Cabo Ortegal is by U-Pb dating of the Cabo Ortegal demantoid. The rock is near-monomineralic garnet, and the
292 associated chlorite lacks K and Rb to date it by means of the K-Ar or Rb-Sr techniques. As the garnet occurs in fissures cutting
293 across the foliation of the pyroxenite, and likely the whole complex in depth, ages younger than the plutons of the area were
294 expected (youngest plutons are ca. 280 Ma, Rodriguez et al., 2007). The results of ca. 255 Ma obtained here are similar to the
295 Aia Pluton age in the westernmost Pyrenees (dated at 267 Ma, Denèle et al., 2012). Those authors interpreted the age as
296 reflecting the earliest evidence of an extensive regime that led later to the formation of the Bay of Biscay rift. Thus, the garnet-
297 bearing veins at the Cabo Ortegal Complex may have formed during the same event.

298 The speed of data acquisition provided by laser-ablation techniques also allows for obtaining a large dataset in a short
299 time, in contrast to the more precise acid-digestion-based garnet dating methods (Lu-Hf and Sm-Nd). This gives the
300 opportunity to investigate a larger number of samples, either with a higher sampling density in a given area of investigation,
301 or by extending the investigation to a much larger area. In some cases, a larger dataset can reveal small-scale metamorphic
302 processes that have been hidden due to sampling bias (including samples not analysed due to the appearance of alteration). For
303 instance, the U-Pb age of garnet from the NE Bohemian Massif obtained here is within the uncertainty of the Lu-Hf age of this
304 sample (Anczkiewicz et al., 2007). Walczak et al. (2017) dated other mesocratic and mafic granulites from the same outcrop
305 sampled by Anczkiewicz et al. (2007) using the same analytical protocols. Their results pointed to a younger event of garnet
306 formation at ca. 345–340 Ma and they attributed this inconsistency to a contribution of inherited Hf in the whole-rock analysis
307 of Anczkiewicz et al. (2007). The *in-situ* U-Pb dating in garnet circumvents the inherited Hf issue as potential inclusions can
308 be avoided by direct petrographic observations (i.e. setting spots in clean domains) or rejecting analysis with distinct U and/or
309 Pb content (Fig. 1). In any case, the fact that the sample analysed by Walczak et al. (2017) is not exactly the same, may indicate
310 that the geological evolution of the area is more complex (polymetamorphic?) than previously thought.



311 **5 Conclusions**

312 This study demonstrates that meaningful U–Pb ages can be obtained by LA-MC-ICP-MS from garnet containing U
313 concentrations lower than 1 ng/g. The analysis of all Pb and U isotopes using ion counters, combined with high-sensitivity
314 analytical conditions, permits reliable regression lines to be defined in Tera–Wasserburg space, even for minerals that contain
315 up to 3,500 times less U than commonly analysed accessory minerals such as zircon.

316 The method yields reproducible and geologically meaningful ages across a wide range of sample ages and
317 compositions. Independent analytical sessions on the Cabo Ortegal demantoid produced indistinguishable ages within
318 uncertainty, while the age obtained for the Orlica–Śnieżnik granulite garnet agrees with previously published Lu–Hf data. The
319 precision achieved depends on the U and Pb concentrations, with internal uncertainties of ca. ± 5 –6 % for garnet containing <1
320 ng/g U and ca. ± 3 % for samples with higher U contents (2–4 ng/g). Moreover, in Archean samples, uncertainties as low as ± 1
321 % were achieved, as older samples benefit from their higher radiogenic Pb contents.

322 These results significantly expand the applicability of in-situ garnet U–Pb geochronology. As most of the
323 metamorphic garnet contain only a few ng/g U or less, this analytical approach opens the possibility of dating a much broader
324 range of metamorphic rocks. The methodology described here provides a valuable new tool for petrochronology and for
325 directly constraining the timing of garnet growth, prograde metamorphism, and deep crustal processes. Garnet U–Pb
326 geochronology can be used together with other conventional accessory mineral geochronometers but may be particularly
327 powerful in geological settings where such chronometers are absent, scarce, or unsuitable.

328 **Data availability**

329 The data presented in this manuscript are available in the supplementary material.

330 **Author contributions**

331 AB conceptualised the study, obtained funding and carried out the formal analyses. Data curation, validation, and
332 writing were done by all the authors through a continuous discussion process.

333 **Competing interests**

334 The authors declare that they have no conflict of interest.

335 **Acknowledgements**

336 We would like to thank Drs. Sonja Aulbach, Jose Ignacio Gil Ibarguchi, Qiao Shu and Robert Anezkievich for sharing
337 the samples analysed in this study. The help of Linda Marko and Alex Schmidt with the day-to-day operation of the instruments
338 makes everything easier.



339 Financial support

340 This work has been supported by the Deutsche Forschungsgemeinschaft (DFG) project number 521366037, granted
341 to AB. FIERCE is financially supported by the Deutsche Forschungsgemeinschaft (DFG: INST 161/921-1 FUGG, INST
342 161/923-1 FUGG and INST 161/1073-1 FUGG), and received financial support from the Wilhelm and Else Heraeus
343 Foundation, which is gratefully acknowledged. This is FIERCE contribution No. xxx.

344 References

- 345 Anczkiewicz, R., Szczepański, J., Mazur, S., Storey, C., Crowley, Q., Villa, I. M., Thirlwall, M. F., and Jeffries, T. E.: Lu-Hf
346 geochronology and trace element distribution in garnet: implications for uplift and exhumation of ultra-high pressure granulites
347 in the Sudetes, SW Poland, *Lithos*, 95, 363–380, <https://doi.org/10.1016/j.lithos.2006.09.001>, 2007.
- 348 Arrowsmith, P.: Laser ablation of solids for elemental analysis by inductively coupled plasma mass spectrometry, *Anal. Chem.*,
349 59, 1437–1444, <https://doi.org/10.1021/ac00137a014>, 1987.
- 350 Aysal, N., Guillong, M., Bayanova, T., Fukuyama, M., Leonard, N., Yılmaz, İ., Varol, E., Tükel, F. Ş., Kadioğlu, Y. K.,
351 Haniç, N., Uzun, F., and Kaygısız, E.: A New Natural Secondary Reference Material for Garnet U-Pb Dating by TIMS and
352 LA-ICP-MS, *Geostand. Geoanal. Res.*, 47, 297–310, <https://doi.org/10.1111/ggr.12493>, 2023.
- 353 Bartoli, O., Millonig, L. J., Carvalho, B. B., Marschall, H. R., and Gerdes, A.: The age of granulite-facies metamorphism in
354 the Ivrea–Verbano Zone (NW Italy) determined through in situ U–Pb dating of garnet, *J. Petrol.*, 65, egae083,
355 <https://doi.org/10.1093/petrology/egae083>, 2024.
- 356 Baxter, E. F., Caddick, M. J., and Dragovic, B.: Garnet: A rock-forming mineral petrochronometer, *Rev. Mineral. Geochem.*,
357 83, 469–533, <https://doi.org/10.2138/rmg.2017.83.15>, 2017.
- 358 Beno, C. J., Lackey, J. S., Schmitz, M. D., Bowman, J. R., Stearns, M. A., Bartley, J. M., and Fernandez, D. P.: Assessment
359 of Natural Reference Materials for U–Pb Geochronology of Grossular-Andradite Garnet, *Geostand. Geoanal. Res.*, 48, 909–
360 925, <https://doi.org/10.1111/ggr.12561>, 2024.
- 361 Beranoaguirre, A., García de Madinabeitia, S., Sanchez-Lorda, M. E., Puellas, P., Ábalos, B., and Gil Ibarguchi, J. I.: U-Pb,
362 Hf isotope and REE constraints on high-pressure acid migmatites from the Cabo Ortegal Complex (NW Spain): New evidence
363 of short-duration metamorphism in a Variscan subduction channel, *Lithos*, 372–373, 105660,
364 <https://doi.org/10.1016/j.lithos.2020.105660>, 2020.
- 365 Beranoaguirre, A., Vasiliev, I., and Gerdes, A.: In-situ LA-ICPMS U-Pb dating of Sulfates: Applicability of carbonate
366 reference materials as matrix-matched standards, *Geochronology*, 4, 601–616, <https://doi.org/10.5194/gchron-4-601-2022>,
367 2022.
- 368 Caddick, M. J. and Kohn, M. J.: Garnet: Witness to the Evolution of Destructive Plate Boundaries Elements, 9, 427–432,
369 <https://doi.org/10.2113/gselements.9.6.427>, 2013.



- 370 Cerva-Alves, T., Hartmann, L. A., Queiroga, G. N., Lana, C., Castro, M. P., Maciel, L. A. C., and Remus, M. V. D.:
371 Metamorphic evolution of the juvenile Serrinha forearc basin in the southern Brasiliano Orogen, Precambrian Res., 365,
372 106394, <https://doi.org/10.1016/j.precamres.2021.106394>, 2021.
- 373 Chew, D. M., Petrus, J. A., Kenny, G. G., and McEvoy, N.: Rapid high-resolution U–Pb LA-Q-ICPMS age mapping of zircon,
374 J. Anal. At. Spectrom., 32, 262–276, <https://doi.org/10.1039/C6JA00404K>, 2017.
- 375 Currie, L. A.: Nomenclature in evaluation of analytical methods including detection and quantification capabilities (IUPAC
376 Recommendations 1995), Pure Appl. Chem., 67, 1699–1723, <https://doi.org/10.1351/pac199567101699>, 1995.
- 377 Dahl, P. S.: A crystal-chemical basis for Pb retention and fission-track annealing systematics in U-bearing minerals, with
378 implications for geochronology, Earth Planet. Sci. Lett., 150, 277–290, [https://doi.org/10.1016/S0012-821X\(97\)00108-8](https://doi.org/10.1016/S0012-821X(97)00108-8),
379 1997.
- 380 Dawson, J. B., Harley, S. L., Rudnick, R. L., and Ireland, T. R.: Equilibration and reaction in Archaean quartz-sapphirine
381 granulite xenoliths from the Lace kimberlite pipe, South Africa, J. Metamorph. Geol., 15, 253–266,
382 <https://doi.org/10.1111/j.1525-1314.1997.00017.x>, 1997.
- 383 Denèle, Y., Paquette, J. L., Olivier, P., and Barbey, P.: Permian granites in the Pyrenees: the Aya pluton (Basque Country),
384 Terra Nova, 24, 105–113, <https://doi.org/10.1111/j.1365-3121.2011.01043.x>, 2012.
- 385 Deng, Y., Zhong, R., Li, D., Li, Y., and Cui, H.: Hunting the Datable Garnet using the LA-ICP-MS U-Pb Method: Predicting
386 Garnet U Concentration, based on Major and Minor Elements, Acta Geol. Sin., 96, 2148–2157, <https://doi.org/10.1111/1755-6724.14921>, 2022.
- 387
388 Dodson, M. H.: Closure temperature in cooling geochronological and petrological systems, Contrib. Mineral. Petrol., 40, 259–
389 274, <https://doi.org/10.1007/BF00373790>, 1973.
- 390 Duchêne, S., Blichert-Toft, J., Luais, B., Télouk, P., Lardeaux, J. M., and Albarède, F.: The Lu–Hf dating of garnets and the
391 ages of the Alpine high-pressure metamorphism, Nature, 387, 586–589, <https://doi.org/10.1038/42446>, 1997.
- 392 Flowers, R. M., Ketcham, R. A., Shuster, D. L., and Farley, K. A.: Apatite (U–Th)/He thermochronometry using a radiation
393 damage accumulation and annealing model, Geochim. Cosmochim. Acta, 73, 2347–2365,
394 <https://doi.org/10.1016/j.gca.2009.01.015>, 2009.
- 395 Fryer, B. J., Jackson, S. E., and Longerich, H. P.: The application of laser ablation microprobe-inductively coupled plasma-
396 mass spectrometry (LAM-ICP-MS) to in situ U–Pb geochronology, Chem. Geol., 109, 1–8, [https://doi.org/10.1016/0009-2541\(93\)90058-Q](https://doi.org/10.1016/0009-2541(93)90058-Q), 1993.
- 397
398 Gerdes, A. and Zeh, A.: Combined U–Pb and Hf isotope LA-(MC)-ICP-MS analyses of detrital zircons: comparison with
399 SHRIMP and new constraints for the provenance and age of an Armorican metasediment in Central Germany, Earth Planet.
400 Sci. Lett., 249, 47–61, <https://doi.org/10.1016/j.epsl.2006.06.039>, 2006.
- 401 Gerdes, A. and Zeh, A.: Zircon formation versus zircon alteration – new insights from combined U–Pb and Lu–Hf in-situ LA-
402 ICPMS analyses, and consequences for the interpretation of Archean zircon from the Central Zone of the Limpopo Belt, Chem.
403 Geol., 261, 230–243, <https://doi.org/10.1016/j.chemgeo.2008.03.005>, 2009.



- 404 Gray, A. L.: Solid sample introduction by laser ablation for inductively coupled plasma source mass spectrometry, *Analyst*,
405 110, 551–556, <https://doi.org/10.1039/AN9851000551>, 1985.
- 406 Horstwood, M. S. A., Košler, J., Gehrels, G., Jackson, S. E., McLean, N. M., Paton, C., Pearson, N. J., Sircombe, K., Sylvester,
407 P., Vermeesch, P., and Bowring, J. F.: Community-derived standards for LA-ICP-MS U-(Th-)Pb geochronology-Uncertainty
408 propagation, age interpretation and data reporting, *Geostand. Geoanal. Res.*, 40, 311–332, [https://doi.org/10.1111/j.1751-](https://doi.org/10.1111/j.1751-908X.2016.00379.x)
409 [908X.2016.00379.x](https://doi.org/10.1111/j.1751-908X.2016.00379.x), 2016.
- 410 Jackson, S. E., Pearson, N. J., Griffin, W. L., and Belousova, E. A.: The application of laser ablation-inductively coupled
411 plasma-mass spectrometry to in situ U–Pb zircon geochronology, *Chem. Geol.*, 211, 47–69,
412 <https://doi.org/10.1016/j.chemgeo.2004.06.017>, 2004.
- 413 Jochum, K. P., Weis, U., Stoll, B., Kuzmin, D., Yang, Q., Raczek, I., Jacob, D. E., Stracke, A., Birbaum, K., Frick, D. A.,
414 Günther, D., and Enzweiler, J.: Determination of reference values for NIST SRM 610–617 glasses following ISO guidelines,
415 *Geostand. Geoanal. Res.*, 35, 97–429, <https://doi.org/10.1111/j.1751-908X.2011.00120.x>, 2011.
- 416 Kirkland, C. L., Ribeiro, B. V., and Slagstad, T.: In situ garnet Lu–Hf geochronology and constraints on metamorphism in the
417 northernmost Norwegian Caledonides, *J. Geol. Soc. Lond.*, <https://doi.org/10.1144/jgs2024-193>, 2025.
- 418 Kohn, M. J., Engi, M., and Lanari, P.: Petrochronology: Methods and applications, *Rev. Mineral. Geochem.*, 83,
419 <https://doi.org/10.2138/rmg.2017.83.0>, 2017.
- 420 Košler, J. and Sylvester, P. J.: Present trends and the future of zircon in geochronology: laser ablation ICP-MS, *Rev. Mineral.*
421 *Geochem.*, 53, 243–275, <https://doi.org/10.2113/0530243>, 2003.
- 422 Le Roex, A., Tinguely, C., and Gregoire, M.: Eclogite and Garnet Pyroxenite Xenoliths from Kimberlites Emplaced Along the
423 Southern Margin of the Kaapvaal Craton, Southern Africa: Mantle or Lower Crustal Fragments?, *J. Petrol.*, 61, ega040,
424 <https://doi.org/10.1093/petrology/egaa040>, 2020.
- 425 Li, D., Fu, Y., Hollings, P., Mitchell, R. H., Zurevinski, S., Kamo, S., Zhang, R., Zhang, Y., Liu, Q., Liao, J., Liang, Y., and
426 Sun, X.: PL57 garnet as a new natural reference material for in situ U–Pb isotope analysis and its perspective for geological
427 applications, *Contrib. Mineral. Petrol.*, 177, 19, <https://doi.org/10.1007/s00410-021-01884-4>, 2022.
- 428 Ludwig, K. R.: User’s Manual for Isoplot Version 3.75–4.15: a Geochronological Toolkit for Microsoft Excel, Berkeley
429 Geochronological Center Special Publication, 5, 2012.
- 430 Machado, N. and Gauthier, G.: Determination of 207Pb/206Pb ages on zircon and monazite by laser-ablation ICP-MS and
431 application to a study of sedimentary provenance and metamorphism in southeastern Brazil, *Geochim. Cosmochim. Acta*, 60,
432 5063–5073, [https://doi.org/10.1016/S0016-7037\(96\)00287-6](https://doi.org/10.1016/S0016-7037(96)00287-6), 1996.
- 433 Madon, M., Gil-Ibarguchi, J. I., Via, J., and Girardeau, J.: Characterization and thermodynamic properties of andradite,
434 *Ca₃Fe₂Si₃O₁₂*, *Am. Mineral.*, 76, 1249–1260, 1991.
- 435 Manzotti, P., Millonig, L. J., Gerdes, A., Whitehouse, M. J., Jeon, H., Poujol, M., and Ballèvre, M.: Protolith age, and timing
436 of burial and exhumation of the UHP Chasteiran Unit (Dora-Maira Massif, Western Alps), constrained by zircon, garnet and
437 rutile petrochronology, *Lithos*, 496–497, 107951, <https://doi.org/10.1016/j.lithos.2025.107951>, 2025.



- 438 Mark, C., O’Sullivan, G., Glorie, A., Simpson, A., Andò, S., Barbarano, M., Stutenbecker, L., Daly, J. S., and Gilbert, S.:
439 Detrital Garnet Geochronology by In Situ U-Pb and Lu-Hf Analysis: A Case Study From the European Alps, *J. Geophys. Res.*
440 *Earth Surf.*, 128, e2023JF007244, <https://doi.org/10.1029/2023JF007244>, 2023.
- 441 Martínez Catalán, J. R., Gómez Barreiro, J., Dias da Silva, I., Chichorro, M., López-Carmona, A., Castiñeiras, P., Abati, J.,
442 Andonaegui, P., Fernández-Suárez, J., González Cuadra, P., and Benítez-Pérez, J. M.: Variscan Suture Zone and Suspect
443 Terranes in the NW Iberian Massif: allochthonous Complexes of the Galicia-Trás os Montes Zone (NW Iberia), in: *The*
444 *Geology of Iberia: A Geodynamic Approach, Volume 2: The Variscan Cycle*, Springer, 99–130, [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-3-030-10519-8_4)
445 [3-030-10519-8_4](https://doi.org/10.1007/978-3-030-10519-8_4), 2019.
- 446 McDougall, I. and Harrison, T. M.: *Geochronology and Thermochronology by the ⁴⁰Ar/³⁹Ar Method*, 2nd Edn., Oxford
447 University Press, New York, 269 pp., ISBN 978-0195109207, 1999.
- 448 Millonig, L. J., Albert, R., Gerdes, A., Avigad, D., and Dietsch, C.: Exploring laser ablation U-Pb dating of regional
449 metamorphic garnet - The Straits Schist, Connecticut, USA, *Earth Planet. Sci. Lett.*, 552, 116589,
450 <https://doi.org/10.1016/j.epsl.2020.116589>, 2020.
- 451 Opendak, M. and Wildenhain, P.: On the analysis of the decay of a small number of radioactive atoms in radiochemical solar
452 neutrino experiments, *Nucl. Instrum. Methods Phys. Res. A*, 345, 570–575, [https://doi.org/10.1016/0168-9002\(94\)90517-7](https://doi.org/10.1016/0168-9002(94)90517-7),
453 1994.
- 454 O’Sullivan, G. J., Hoare, B. C., Mark, C., Drakou, F., and Tomlinson, E. L.: Uranium–lead geochronology applied to pyrope
455 garnet with very low concentrations of uranium, *Geol. Mag.*, 160, 1010–1019, <https://doi.org/10.1017/S0016756823000122>,
456 2023.
- 457 Peillod, A., Patten, C. G. C., Drüppel, K., Beranoaguirre, A., Zeh, A., Gudelius, D., Hector, S., Majka, J., Kleine, B. I., Karlson,
458 A., Gerdes, A., and Kolb, J.: Disruption of a high-pressure unit during exhumation: example of the Cycladic Blueschist Unit
459 (Thera, Ios and Naxos islands, Greece), *J. Metamorph. Geol.*, 42, 225–255, <https://doi.org/10.1111/jmg.12753>, 2024.
- 460 Pettke, T., Oberli, F., Audétat, A., Guillong, M., Simon, A. C., Hanley, J. J., and Klemm, L. M.: Recent developments in
461 element concentration and isotope ratio analysis of individual fluid inclusions by laser ablation single and multiple collector
462 ICP-MS, *Ore Geol. Rev.*, 44, 10–38, <https://doi.org/10.1016/j.oregeorev.2011.11.001>, 2012.
- 463 Piccione, G., Rasbury, E. T., Elliott, B. A., Kyle, J. R., Jaret, S. J., Acerbo, A. S., Lanzirotti, A., Northrup, P., Wootton, K., and
464 Parrish, R. R.: Vein fluorite U-Pb dating demonstrates post-6.2 Ma rare-earth element mobilization associated with Rio Grande
465 rifting, *Geosphere*, 15, 1958–1972, <https://doi.org/10.1130/GES02139.1>, 2019.
- 466 Reinhardt, N., Gerdes, A., Beranoaguirre, A., Frenzel, M., Meinert, L. D., Gutzmer, J., and Burisch, M.: Timing of magmatic-
467 hydrothermal activity in the Variscan belt - LA-ICP-MS U-Pb geochronology of skarn-related garnet from the Schwarzenberg
468 district, Erzgebirge, *Mineral. Deposita*, 57, 1071–1087, <https://doi.org/10.1007/s00126-021-01084-x>, 2022.
- 469 Reiners, P. W., Ehlers, T. A., and Zeitler, P. K.: Past, present, and future of thermochronology, *Rev. Mineral. Geochem.*, 58,
470 1–18, <https://doi.org/10.2138/rmg.2005.58.1>, 2005.



- 471 Ring, U. and Gerdes, A.: Kinematics of the Alpenrhein-Bodensee graben system in the Central Alps: Oligocene/Miocene
472 transtension due to formation of the Western Alps arc, *Tectonics*, 35, 1367–1391, <https://doi.org/10.1002/2015TC004085>,
473 2016.
- 474 Roberts, N. M. W. and Walker, R. J.: U–Pb geochronology of calcite-mineralized faults: absolute timing of rift-related fault
475 events on the northeast Atlantic margin, *Geology*, 44, 531–534, <https://doi.org/10.1130/G37868.1>, 2016.
- 476 Rodriguez, J., Gil Ibarguchi, J. I., and Paquette, J. L.: Variscan synchronous magmatism across the Iberian Massif: new U–Pb
477 ages in granitoids from the Fisteria area (A Coruña, Spain), XV Semana – VI Congresso Ibérico de Geoquímica, 2007.
- 478 Rubatto, D.: Zircon trace element geochemistry: Partitioning with garnet and the link between U–Pb ages and metamorphism,
479 *Chem. Geol.*, 184, 123–138, [https://doi.org/10.1016/S0009-2541\(01\)00355-2](https://doi.org/10.1016/S0009-2541(01)00355-2), 2002.
- 480 Rubatto, D.: Zircon: the metamorphic mineral, *Rev. Mineral. Geochem.*, 83, 261–295, <https://doi.org/10.2138/rmg.2017.83.9>,
481 2017.
- 482 Salnikova, E. B., Chakhmouradian, A. R., Stifeeva, M. V., Reguir, E. P., Kotov, A. B., Gritsenko, Y. D., and Nikiforov, A.
483 V.: Calcic garnets as a geochronological and petrogenetic tool applicable to a wide variety of rocks, *Lithos*, 338–339, 141–
484 154, <https://doi.org/10.1016/j.lithos.2019.03.032>, 2019.
- 485 Schaltegger, U., Schmitt, A. K., and Horstwood, M. S. A.: U–Th–Pb zircon geochronology by ID-TIMS, SIMS, and laser
486 ablation ICP-MS: Recipes, interpretations, and opportunities, *Chem. Geol.*, 402, 89–110,
487 <https://doi.org/10.1016/j.chemgeo.2015.02.028>, 2015.
- 488 Schannor, M., Lana, C., Nicoli, G., Cutts, K., Buick, I., Gerdes, A., and Hecht, L.: Reconstructing the metamorphic evolution
489 of the Araçuaí orogeny (SE Brazil) using in situ U–Pb garnet dating and P–T modelling, *J. Metamorph. Geol.*, 39, 1145–1171,
490 <https://doi.org/10.1111/jmg.12605>, 2021.
- 491 Schmitz, M. D. and Bowring, S. A.: Constraints on the thermal evolution of continental lithosphere from U–Pb accessory
492 mineral thermochronology of lower crustal xenoliths, southern Africa, *Contrib. Mineral. Petrol.*, 144, 592–618,
493 <https://doi.org/10.1007/s00410-002-0419-9>, 2003.
- 494 Seman, S., Stockli, D. F., and McLean, N. M.: U–Pb geochronology of grossular-andradite garnet, *Chem. Geol.*, 460, 106–
495 116, <https://doi.org/10.1016/j.chemgeo.2017.04.020>, 2017.
- 496 Shu, Q., Beranoaguirre, A., Albert, R., Millonig, L. J., Walters, J. B., Marschall, H. R., Gerdes, A., Hoefer, H. E., Hezel, D.,
497 and Brey, G. P.: Multi-stage ultrahigh temperature metamorphism in the lower crust of the Kaapvaal craton recorded by U–Pb
498 ages of garnet, *Contrib. Mineral. Petrol.*, 179, 49, <https://doi.org/10.1007/s00410-024-02121-4>, 2024.
- 499 Simpson, A., Gilbert, S., Tamblyn, R., Hand, M., Spandler, C., Gillespie, J., Nixon, A., and Glorie, S.: In situ Lu–Hf
500 geochronology of garnet, apatite and xenotime by LA-ICP-MS/MS, *Chem. Geol.*, 577, 120299,
501 <https://doi.org/10.1016/j.chemgeo.2021.120299>, 2021.
- 502 Spear, F. S. and Parrish, R. R.: Petrology and cooling rates of the Valhalla Complex, British Columbia, Canada, *J. Petrol.*, 37,
503 733–765, <https://doi.org/10.1093/petrology/37.4.733>, 1996.



504 Stifeeva, M. V., Salnikova, E. B., Samsonov, A. V., Kotov, A. B., and Gritsenko, Y. D.: U-Pb geochronology of garnet from
505 Daschkasan skarn deposit (Lesser Caucasus), Dokl. Earth Sci., 487, 953–956, <https://doi.org/10.1134/S1028334X19080178>,
506 2019.

507 Stifeeva, M., Salnikova, E., Adamskaya, E., Millonig, L. J., Ivanov, A., Mottram, C., Kovach, V., Kotov, A., Bryanskiy, N.,
508 Karimov, A., Gladkochub, E., Peytcheva, I., Gerdes, A., Bowie, S., and Gritsenko, J.: Kovdor-GRT – A New Natural Reference
509 Material for Garnet U-Pb Dating, Geostand. Geoanal. Res., <https://doi.org/10.1111/ggr.70040>, 2026.

510 Sylvester, P. J. and Jackson, S. E.: A Brief History of Laser Ablation Inductively Coupled Plasma Mass Spectrometry (LA–
511 ICP–MS), Elements, 12, 307–310, <https://doi.org/10.2113/gselements.12.5.307>, 2016.

512 Szczepański, J. and Ilnicki, S.: From Cadomian arc to Ordovician passive margin: geochemical records preserved in
513 metasedimentary successions of the Orlica-Snieżnik Dome in SW Poland, Int. J. Earth Sci., 103, 627–647,
514 <https://doi.org/10.1007/s00531-013-0993-2>, 2014.

515 Tamblyn, R., Hand, M., Simpson, A., Gilbert, S., Wade, B., and Glorie, S.: In situ laser ablation Lu–Hf geochronology of
516 garnet across the Western Gneiss Region: campaign-style dating of metamorphism, J. Geol. Soc. Lond., 179, jgs2021-094,
517 <https://doi.org/10.1144/jgs2021-094>, 2022.

518 Tera, F. and Wasserburg, G. J.: U-Th-Pb systematics in three Apollo 14 basalts and the problem of initial Pb in lunar rocks,
519 Earth Planet. Sci. Lett., 14, 281–304, [https://doi.org/10.1016/0012-821X\(72\)90128-8](https://doi.org/10.1016/0012-821X(72)90128-8), 1972.

520 Walczak, K., Anczkiewicz, R., Szczepański, J., Rubatto, D., and Košler, J.: Combined garnet and zircon geochronology of the
521 ultra-high temperature metamorphism: Constraints on the rise of the Orlica-Śnieżnik Dome, NE Bohemian Massif, SW Poland,
522 Lithos, 292–293, 388–400, <https://doi.org/10.1016/j.lithos.2017.09.013>, 2017.

523 Walters, J. B., Garber, J. M., Beranoaguirre, A., Millonig, L. J., Gerdes, A., Grützner, T., and Marschall, H. R.: Zircon micro-
524 inclusions as an obstacle for in situ garnet U-Pb geochronology: An example from the As Sifah eclogite locality, Oman,
525 Geochronology, 7, 309–333, <https://doi.org/10.5194/gchron-7-309-2025>, 2025.

526 Willigers, B. J. A., Baker, J. A., Krogstad, E. J., and Peate, D. W.: Precise and accurate in situ Pb-Pb dating of apatite, monazite,
527 and sphene by laser ablation multiple-collector ICP-MS, Geochim. Cosmochim. Acta, 6, 1051–1066,
528 [https://doi.org/10.1016/S0016-7037\(01\)00838-9](https://doi.org/10.1016/S0016-7037(01)00838-9), 2002.

529 Yang, Y., Wu, S., Wang, H., Kamo, S. L., Ma, Q., Liang, T., Xu, L., Xie, L., Huang, C., Wan, B., Yang, J., and Wu, F.: Three
530 new natural secondary reference materials for in situ andradite U–Pb geochronology, J. Anal. At. Spectrom., 40, 326–337,
531 <https://doi.org/10.1039/D4JA00290C>, 2025.

532

533