

GPU-accelerated three-dimensional variational data assimilation with automatic differentiation for WRF-Chem

Hancheng Ye¹, Zengliang Zang¹, Wei You¹, Yiwen Hu¹, Ning Liu¹, Yi Li¹

¹College of Meteorology and Oceanography, National University of Defense Technology, Changsha, China

Correspondence: Wei You (ywlx_1987@163.com) and Zengliang Zang (zzlqxy@163.com)

Abstract: This study developed a PyTorch-based three-dimensional variational (3DVar) data assimilation system (PY3DVar) for the Weather Research and Forecasting model coupled to Chemistry (WRF-Chem), which integrates automatic differentiation (AD) to replace traditional manual gradient derivation and adopts Graphics Processing Unit (GPU) acceleration for high computational efficiency. PY3DVar enables the simultaneous assimilation of gaseous pollutants (SO₂, NO₂, CO, O₃) and particulate matter (PM_{2.5}, PM₁₀) and supports flexible deployment on both Central Processing Unit (CPU) and GPU computing platforms. Idealized tests were first conducted to verify the physical consistency of the assimilation system. Twenty consecutive real-case assimilation experiments, each followed by a 24-hour forecast, were then conducted to assess computational performance and efficiency, benchmarked against the classic Fortran-based 3DVar system (FT3DVar) using CPU parallelization. The results of the idealized tests show PY3DVar effectively propagates observation information, generating increment fields consistent with FT3DVar. Long-term real-case experiments further verify that the two assimilation systems achieve nearly identical analysis accuracy for chemical initial fields. On the GPU platform, PY3DVar achieves an average assimilation time of 1.2 s, representing a 26.2-fold speedup compared to FT3DVar on 64 CPU cores (32.7 s). Parallel scalability tests across 1, 2, 4, 8, 16, 32 and 64 CPU cores reveal that FT3DVar runs faster per iteration across all core counts. Consequently, FT3DVar achieves shorter total assimilation time on 1, 2, 4 and 16 cores, whereas PY3DVar outperforms it at 8, 32 and 64 cores, owing to its intrinsically fewer required iterations for convergence. The average 24-hour forecast results show that the improved initial fields have sustained positive effects on short-term forecasts: the improvements persist for over 24 hours for SO₂, CO, PM_{2.5}, and PM₁₀, and for over 6 hours for NO₂ and O₃. These results demonstrate that PY3DVar delivers substantially faster assimilation than FT3DVar while maintaining equivalent analysis accuracy, offering an efficient and practical solution for operational atmospheric chemical data assimilation.

1. Introduction

Air quality models (AQMs) are crucial tools for atmospheric pollution simulation and forecasting (Li et al., 2013; Wang et al., 2022; Gao et al., 2024). Errors in initial conditions can degrade model performance, particularly within the early hours of simulation, leading to significant biases in short-term forecasts. Data assimilation (DA) combines multi-source observations, such as those from surface stations and satellites, with model background fields through statistically derived weighting to produce an optimal analysis field. This technique effectively improves the initial conditions of numerical models and has become a core method for enhancing simulation accuracy in both research and operational forecasting (Engelen et al., 2009; Ma et al., 2025).

Compared to the long-standing development of meteorological data assimilation, atmospheric chemical data assimilation started relatively late, and its theoretical framework and operational applications are still under continuous improvement. Early explorations in atmospheric chemical data assimilation were largely constrained by limitations in observational capabilities and the difficulty of simulating complex aerosol compositions. Consequently, research initially focused on gaseous precursors like SO_2 , NO_2 , O_3 and typical key chemical components like HNO_3 with clear chemical reaction mechanisms and mature simulation schemes. In contrast, the assimilation of particulate matter (PM) components was less common due to the high complexity of their sources and physicochemical mechanisms. In earlier work, Austin et al. (1992) assimilated LIMS satellite measurements of HNO_3 and O_3 into a 2D chemical model, effectively identifying and correcting systematic biases in the original chemical model, significantly improving the analysis accuracy of stratospheric chemical composition, and demonstrating the key role of four-dimensional assimilation in refining chemical models. Elbern et al. (1997) used the RADM2 (Second-Generation Regional Acid Deposition Model) gas-phase chemical mechanism and a variational method to assimilate ozone observations, successfully optimizing the initial concentrations of its precursors NO_x and volatile organic compounds (VOCs), and enabling quantitative analysis of ozone chemical sensitivity. Frydendall et al. (2009) developed a data assimilation algorithm based on statistical interpolation and coupled it with the Danish Eulerian Operational Model (DEOM), conducting assimilation studies on surface ozone and effectively improving forecast performance by optimizing assimilation parameters.

With advancements in observational technology and the continuous improvement of atmospheric chemistry modeling systems, the focus of atmospheric chemical data assimilation has gradually expanded to include particulate pollutants with more complex compositions and sources. In the early stages, owing to the limited simulation capability for aerosol physicochemical processes, assimilation studies mainly focused on adjusting total mass concentrations, such as $\text{PM}_{2.5}$ and PM_{10} , to improve model performance. For example, Denby et al. (2008) applied both Statistical Interpolation (SI) and Ensemble Kalman Filtering (EnKF) data assimilation methods to an atmospheric chemistry model. They found that both methods significantly corrected the model's systematic underestimation of PM_{10} concentration, with the SI method reducing the root mean square error (RMSE) of daily average

concentration from $16.7 \mu\text{g m}^{-3}$ to $9.2 \mu\text{g m}^{-3}$, and noted that the improvement from SI was significantly better than that from EnKF. Tombette et al. (2009) used the optimal interpolation (OI) method to assimilate surface PM_{10} observations from France into a regional chemical transport model, effectively enhancing the short-term forecast accuracy of total aerosol mass ($\text{PM}_{2.5}$ and PM_{10}) across Europe, demonstrating the potential application of this method in operational forecasting.

As a core method in data assimilation, the three-dimensional variational (3DVar) approach statistically combines observations with model background fields to derive improved initial conditions by minimizing a prescribed cost function (Lorenc, 1986; Parrish and Derber, 1992). Although more advanced methods such as four-dimensional variational (4DVar) and hybrid ensemble-variational assimilation have been developed for global operational systems, 3DVar remains widely used in regional and chemical data assimilation applications owing to its computational efficiency, implementation simplicity, and robust performance (Jiang et al., 2013; Seo et al., 2023; Chen et al., 2024; Zhou et al., 2025). Pagowski et al. (2010) introduced surface ozone and fine particulate matter ($\text{PM}_{2.5}$) observations into the WRF-Chem model using the GSI (Grid-point Statistical Interpolation) 3DVar system. Results showed that while assimilation significantly reduced forecast errors, these errors grew rapidly in the early forecast hours. With the advancement of observational techniques and the refinement of model mechanisms, simply correcting total aerosol mass often fails to fundamentally optimize the simulation of its internal chemical composition. The target variables for aerosol assimilation are shifting from total mass concentration towards a multivariate system encompassing key chemical components such as sulfate, nitrate, and organic carbon. Li et al. (2013) integrated a 3DVar algorithm with the WRF-Chem model employing the MOSAIC aerosol scheme. By assimilating $\text{PM}_{2.5}$ and its chemical component observations, they achieved a comprehensive analysis of the concentrations and size distributions of eight key species, including black carbon, organic carbon, sulfate, and nitrate, significantly improving the simulation and forecast accuracy of these components. Jiang et al. (2013) implemented the assimilation of PM_{10} observations using the 3DVar method, providing an improved initial field for the WRF-Chem model by optimizing various aerosol components. However, the assimilation effect diminished rapidly due to vertical mixing and advective processes, highlighting the importance of optimizing the aerosol vertical initial state. By designing various assimilation scenarios, Zang et al. (2016) confirmed that introducing aircraft-measured aerosol data into traditional surface-based aerosol data assimilation significantly optimized aerosol mass concentration, vertical distribution, and chemical composition. They also found that the forecast skill improvement from vertical observations lasted longer than that from surface data, underscoring the important influence of vertical resolution on forecast performance. Apart from experiments focusing on different observation types and pollutants, studies have optimized the intrinsic mathematical formulation of 3DVar algorithms. From a methodological perspective, Key et al. (2025) revealed the mathematical connection between 3DVar and Gaussian process regression, offering an alternative pathway for scalable assimilation computation.

Despite continuous improvements in assimilation algorithms and observation sources, traditional Fortran-based 3DVar systems still face prominent computational bottlenecks in modern high-resolution chemical assimilation scenarios. As observational data grow in volume and diversity, model resolutions become finer, and the number of simulated chemical species continues to expand, the state vector and matrix dimensions in atmospheric chemistry assimilation have increased substantially (Liang et al., 2020; Wang et al., 2022). The incorporation of satellite-derived aerosol optical depth and extinction profiles has proven effective for constraining the three-dimensional distribution of pollutants (Zhang et al., 2011; Chen et al., 2017; Ye et al., 2022; Wang et al., 2022). However, these observations require complex observation operators and their adjoints, which further compounds the computational challenge (Liang et al., 2020; Ye et al., 2021). These factors impose a heavy computational burden on traditional Fortran-based assimilation systems, prolonging assimilation time and challenging their ability to meet the timeliness requirements of operational air quality forecasting. In recent years, emerging high-performance computing techniques and PyTorch-based tensor computing frameworks have provided promising solutions to relieve the computational bottlenecks and tedious manual coding burden of traditional Fortran 3DVar systems (Boudier et al., 2023; De Luca et al., 2021; He et al., 2025). In parallel, modern deep learning (DL) frameworks such as PyTorch provide tensor-based computation and AD capabilities that are well suited to the matrix-dominated operations in variational data assimilation. Closely related to this study, Cheng et al. (2025) developed TorchDA, a Python package that implements multiple assimilation algorithms including 3DVar within PyTorch, demonstrating the feasibility of building variational assimilation systems in DL frameworks. However, TorchDA was demonstrated primarily on low-dimensional Lorenz models, and its applicability to full-scale atmospheric chemistry assimilation with three-dimensional grids and multiple species has not been established. Beyond updating classical algorithms, numerous full data-driven methods have been proposed to augment or replace components of traditional assimilation, such as using neural networks as ensemble surrogates (Chattopadhyay et al., 2023), adopting learned priors to replace background error covariance (Beauchamp et al., 2025), and constructing end-to-end frameworks that eliminate explicit observation operators (Xu et al., 2025). Although satisfactory computational efficiency, most existing fully data-driven assimilation methods are validated only under idealized low-dimensional experimental conditions. Meanwhile, their inherent black-box characteristic leads to poor physical interpretability and limited generalization ability. Such drawbacks not only hinder diagnostic analysis of pollution formation and evolution mechanisms, but also bring potential risks for reliable operational forecasting applications.

Overall, current atmospheric chemical data assimilation still faces a clear trade-off: traditional Fortran-based 3DVar owns rigorous physical basis and full mathematical traceability but suffers from low computational efficiency and tedious manual coding; pure DL-based assimilation achieves fast computation but sacrifices physical interpretability and practical robustness. To fill this research gap and balance reliable physical constraints and high computational performance, this study develops a novel PyTorch-based 3DVar assimilation system. The proposed framework inherits the complete

classical 3DVar variational optimization theory without modifying underlying physical equations, guaranteeing transparent physical meaning and traceable gradient calculation throughout the assimilation workflow. Benefiting from native PyTorch AD and hardware-accelerated tensor parallel computing, the new system avoids manual adjoint programming and supports flexible deployment on both CPU and GPU hardware. Meanwhile, compared with fully black-box DL assimilation methods, it retains all physical constraints of traditional variational assimilation. This study finally conducts comprehensive contrast experiments between the newly built PY3DVar system and the traditional manually coded Fortran 3DVar system (FT3DVar), focusing on computational efficiency, parallel scalability and assimilation analysis performance for multiple conventional air pollutants including SO₂, NO₂, CO, O₃, PM_{2.5}, and PM₁₀.

2. Model, data, and methodology


2.1 Air quality model

The Weather Research and Forecasting model with Chemistry (WRF-Chem), was jointly developed by the National Center for Atmospheric Research (NCAR), the National Oceanic and Atmospheric Administration (NOAA), and other institutions (Grell et al., 2005; Fast et al., 2006). Its core strength lies in the fully online, bidirectional coupling between meteorological and chemical fields, enabling a precise description of pollutant transport, transformation, and removal under complex meteorological conditions. The meteorological fields drive the spatiotemporal distribution and transformation of chemical species through physical processes such as dynamic advection, turbulent diffusion, and dry/wet deposition. In turn, the chemical fields can feedback on meteorological elements through mechanisms like aerosol radiative forcing and cloud condensation nuclei effects. This online coupling ensures spatiotemporal consistency and physical compatibility between atmospheric dynamic processes and chemical transformations.

The WRF-Chem (version 3.5) used in this study integrates the Aerosol Module 5 (AERO5) from the Community Multiscale Air Quality Modeling System (CMAQ) and employs the SAPRC-99 gas-phase chemical mechanism (Carter, 2000; Li et al., 2010, 2011a, 2011b, 2012). Through over two decades of development, this model incorporates advanced chemical reactions and processes crucial for atmospheric particulate formation. For example, it includes irreversible aqueous-phase processes of glyoxal and methylglyoxal on particle surfaces, which improves the simulation of secondary organic aerosol (SOA) over China (Xing et al., 2018). A parameterization for heterogeneous nitrous acid (HONO) processes has been implemented, enhancing the model's capability to simulate atmospheric oxidizing capacity and SOA (Feng et al., 2005, 2016). The suppression effect of organic coatings on the heterogeneous hydrolysis uptake of dinitrogen pentoxide (N₂O₅) is also parameterized, leading to better simulation of nitrate aerosols. Furthermore, the model incorporates sulfate formation via transition-metal catalysis and Criegee intermediate reactions, improving sulfate simulation. Additionally, a module accounting for aerosol–radiation–photolysis interactions has been developed,

182 enabling assessment of aerosol–cloud interactions, radiative effects of brown carbon aerosols, and
 183 direct and indirect effects of various aerosol components.

184 In this study, a single-domain configuration with 210 (west–east) $\times$ 168 (south–north) grid points
 185 at 27 km horizontal resolution and 34 vertical layers was employed. The vertical layers follow a terrain-
 186 following coordinate with higher resolution near the surface that gradually decreases with altitude,
 187 extending from the surface to approximately 20 km. The domain-averaged height corresponding to
 188 each vertical level is illustrated in Fig. 3, which mainly presents the vertical distribution of background
 189 error standard deviation. The simulation domain covers most of China and surrounding regions, as
 190 shown in Fig. 1.



191

192 **Figure 1.** Distribution of the WRF-Chem simulation domain.

192

193 2.2 Data

194 The meteorological fields for the WRF-Chem model were driven by the NCEP Final Operational
 195 Global Analysis (FNL, <https://gdex.ucar.edu/datasets/d083002/>) data, which has a horizontal
 196 resolution of $1^\circ \times 1^\circ$ and a temporal resolution of 6 hours. Provided by NCAR and NCEP, the FNL
 197 dataset assimilates multi-source global observations from surface stations, radiosondes, satellites, etc.
 198 It supplies physically consistent meteorological fields and serves as crucial input for generating reliable
 199 initial and boundary conditions for the model. This dataset has been widely used in various mesoscale
 200 numerical simulations.

201 Anthropogenic emissions were derived from the 2013 Multi-resolution Emission Inventory for
 202 China (MEIC, <http://www.meicmodel.org/index.html>), provided by Tsinghua University (Tong et al.,
 203 2020; Zheng et al., 2021). This inventory has a spatial resolution of 6 km, covers most regions of China,
 204 and includes emission data for the gaseous pollutants and aerosol precursors required in this study.

205 Chemical boundary conditions for the WRF-Chem model were obtained from the Whole
 206 Atmosphere Community Climate Model (WACCM, <https://www.acom.ucar.edu/waccm/download.shtml>) data released by NCAR. This dataset provides
 207 the global spatiotemporal distribution of various chemical components, including ozone and aerosol
 208 precursors, ensuring reasonable atmospheric chemical boundary conditions for the regional air quality
 209

simulations.

Observation data were sourced from the China National Environmental Monitoring Centre (CNEMC, <https://air.cnemc.cn:18007/>), primarily comprising hourly surface concentrations of major air pollutants: SO₂, NO₂, CO, O₃, PM_{2.5}, and PM₁₀. The locations of observation sites follow the distribution described in He et al. (2025). To evaluate the assimilation performance, observations are divided into assimilation and independent validation datasets following Hu et al. (2022). Considering distinct missing-value characteristics among different pollutants, the station split was performed independently for each of the six pollutants and executed on a daily basis, using the observations at the assimilation start time (00:00 UTC) as the basis. The partitioning procedure for an individual pollutant is described as follows. First, invalid missing records of the target pollutant at 00:00 UTC are removed, and the remaining valid stations are mapped onto model grids. If only one valid station is present within a grid cell, it is assigned to the assimilation set. When two or more valid stations exist in one grid cell, one station is randomly selected for the independent validation set, and the rest are assigned to the assimilation set. After the station assignment for each pollutant at 00:00 UTC is completed, all 25 time steps of the forecast period (from 00:00 UTC of the current day to 00:00 UTC of the next day) share the same station assignment. Assimilation is performed only at the initial forecast time, and the subsequent time steps of the forecast period are used solely for forecast verification. For later hours, valid measurements from stations belonging to the validation set are treated as independent validation samples. Observations from stations outside the corresponding validation sets are marked as missing and are not used for validation.

2.3. Control variable design

The WRF-Chem model employed in this study incorporates aerosol variables that encompass a variety of fine and coarse particle components. The fine particle components include: elemental carbon (ECJ), ammonium (NH₄AJ), sulfate (SO₄AJ), nitrate (NO₃AJ_A/B), organic aerosols (ORGAJ, ORGBAJ, SOA_RXT_G/M), nine categories of secondary organic aerosols generated from the semi-volatile gases of primary organic aerosol (CPO1AER–CPO9AER), chloride (CLAJ), and accumulation-mode fine particles (P25AJ), among others. The coarse particle components include: dust (DUST_1–DUST_5), anthropogenic coarse aerosol (ANTHA), and coarse-mode soil-derived particles (SOILA). Furthermore, the model also includes gaseous pollutants such as SO₂, NO₂, CO, and O₃. The fine particle components listed above collectively constitute PM_{2.5}, while the addition of the coarse particle components (DUST_1–DUST_5, ANTHA, and SOILA) forms the complete PM₁₀ composition.

Using all aerosol species directly as control variables would provide detailed compositional resolution but at the cost of substantially increased computational scale, storage demands, and reduced efficiency. To balance assimilation accuracy with computational cost, this study merges specific components based on their physicochemical properties and particle size characteristics: the dual nitrate

components are merged into a single variable; the four organic aerosol components are combined into one group; the nine secondary organic aerosols originating from semi-volatile primary organic gases are uniformly merged; the dust components are optimally merged according to particle size differences—DUST_1 is retained as an independent control variable, while DUST_2 to DUST_5 are merged into a single variable. Through this merging process, the number of aerosol control variables is reduced to 12. Gaseous pollutants (SO₂, NO₂, CO, O₃) are directly used as control variables for assimilation. Consequently, the assimilation system constructed in this study utilizes a total of 16 control variables. Table 1 provides the detailed mapping between the WRF-Chem model variables and the 3DVar control variables.

Model variables	3DVar control variables
ECJ	EC
NH4AJ	NH4
SO4AJ	SO4
NO3AJ_A NO3AJ_B	NO3
ORGAJ ORGBAJ SOA_RXT_G SOA_RXT_M	ORG
CPO1AER CPO2AER ... CPO9AER	CPO
CLAJ	CL
P25AJ	P25
DUST_1	DUST1
DUST_2 DUST_3 DUST_4 DUST_5	DUST2
ANTHA	ANTHA
SOILA	SOILA
SO2	SO2
NO2	NO2
CO	CO
O3	O3

Table 1. Mapping between WRF-Chem model variables and 3DVar control variables.

2.4 3DVar assimilation theory

The 3DVar method is based on probability and statistical theory, treating both the simulated background field and observations as useful information containing errors. This method constructs a cost function that incorporates the weighted errors of the background field and the observations, and seeks the state that minimizes this function, known as the analysis field. Statistically, this analysis field is considered the optimal estimate of the true atmospheric state. The 3DVar cost function is:

$$J(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \mathbf{x}_b)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + \frac{1}{2}(\mathbf{H}\mathbf{x} - \mathbf{y})^T \mathbf{R}^{-1}(\mathbf{H}\mathbf{x} - \mathbf{y}) \quad (1)$$

where $\mathbf{x}$ is the analysis field, representing the set of control variables at all grid points in the model. Its dimensionality is jointly determined by the number of control variables, vertical layers, and horizontal grid points. $\mathbf{x}_b$ is the background field, provided by the WRF-Chem model simulation, and

has the same dimensions as x . B is the background error covariance matrix, which describes the magnitude and spatial structure of the uncertainties in the background field. H is the observation operator that maps from model space to observation space. Since only surface in-situ observations are assimilated, H consists solely of horizontal bilinear interpolation from the model surface level to the observation locations, without any vertical interpolation. y is the observation vector, containing the assimilated observations, with dimensions of $\text{Obs_Station} \times 6$, where Obs_Station represents the number of valid observation sites, and 6 corresponds to the six types of pollutants. R is the observation error covariance matrix, representing the uncertainty of the observational data, and has dimensions consistent with y . Following Li et al. (2013), the observational error standard deviation in this study is set to half of the background error standard deviation.

The background error covariance matrix B has a very large dimension. Storing and performing iterative computations with it directly would pose serious challenges to memory and computational efficiency. To optimize the solution of the cost function, this study adopts a classic preconditioning framework for variational data assimilation, factorizing and structurally simplifying the B matrix (El Akkraoui et al., 2013; Li et al., 2013). The specific processing is as follows: First, the B matrix is decomposed into the form $B = DCD^T$, where D is a diagonal matrix composed of the background error standard deviations, and C is the background error correlation matrix. Furthermore, the background error correlation matrix C is decomposed into the tensor product of correlation matrices along the longitude, latitude, and vertical directions, i.e., $C = C_x \otimes C_y \otimes C_z$ (Derber and Rosati, 1989; Li et al., 2008). In practical assimilation, the analysis field x is typically not solved for directly. Instead, the incremental field $\delta x = x - x_b$ is introduced to solve for the analysis field indirectly, and the control variable is transformed to δz , where the control variables satisfy $\delta x = DC^{1/2}\delta z$. Defining the transformation operator $F = DC^{1/2}$, the cost function can be rewritten in terms of δz as:

$$J(\delta z) = \frac{1}{2}\delta z^T \delta z + \frac{1}{2}(HF\delta z - d)^T R^{-1}(HF\delta z - d) \quad (2)$$

where $d = y - Hx_b$ is the innovation vector (observation increment). After minimizing $J(\delta z)$ to obtain the optimal δz , the final analysis field x_a is obtained through the following transformation:

$$x_a = x_b + F\delta z \quad (3)$$

2.5 Implementation of the two assimilation systems

The two assimilation systems compared in this study, PY3DVar and FT3DVar, are built upon the same variational theory (Section 2.4) and share identical background error covariance modeling scheme (Section 2.6). Both systems employ the Limited-memory Broyden-Fletcher-Goldfarb-Shanno (L-BFGS) algorithm to achieve minimization with identical parameter settings: the memory size (history size) is set to 6, which determines the number of historical gradient-update pairs stored for the Hessian approximation (Liu and Nocedal, 1989); the maximum number of inner iterations per optimization step (max iteration) is set to 4; and convergence is declared when the scaled gradient

norm satisfies $\|\nabla J\| / \max(1, \|\delta z\|) \leq 10^{-2}$, a standard criterion in variational data assimilation. The inherent algorithmic difference between the two implementations lies in the internal line search strategy: FT3DVar uses the classical Moré-Thuente line search (Moré and Thuente, 1994), while PY3DVar relies on the Strong Wolfe line search built into PyTorch's L-BFGS optimizer.

FT3DVar is a Fortran-based 3DVar system independently developed by our research group and validated in several previous studies over China (Zang et al., 2016; Liang et al., 2020; Wang et al., 2020). It should be noted that FT3DVar is entirely independent of the WRFDA-Chem system distributed with the official WRF release. The full assimilation algorithm is written in Fortran with OpenMP loop parallelism for CPU execution. The L-BFGS outer driver follows the standard L-BFGS algorithm (Liu and Nocedal, 1989; Nocedal, 1980), adapted from publicly-shared Fortran implementations, with customized data interfaces for compatibility with the 3DVar framework and parallelized via multi-CPU threading. The gradient of the cost function is derived analytically:

$$\nabla J(\delta z) = \delta z + F^T H^T R^{-1} (H F \delta z - d) \quad (4)$$

This analytical gradient is then manually implemented in the code. Although AD tools are available for Fortran, the adjoint of the 3DVar cost function involves a limited set of modular operations whose analytical derivation is straightforward, allowing manual implementation with full control over memory and parallelization.

PY3DVar is developed within the PyTorch framework and can be deployed on both CPU and GPU platforms without code modification. In the PY3DVar assimilation framework, the forward assimilation process is the computation from the control variable δz to the scalar cost J . The only differentiable quantity in this process is δz (created with `requires_grad = True`); all other quantities — including the background-error transform operator F , the observation operator H , the innovation vector d , and the observation error covariance R — are treated as constants during the computation. The complete forward process that starts from δz and computes J is encapsulated in the closure function `closure()`. During this process, PyTorch's autograd engine records all tensor operations that depend on δz , building a complete computational graph from δz to J . Once the forward computation is complete, calling `cost.backward()` makes the autograd engine backpropagate along this graph, yielding the gradient of the cost with respect to the control variable, $\nabla J(\delta z)$, thereby achieving automatic gradient computation without any manually coded adjoint. During the minimization, the L-BFGS optimizer calls the closure function `closure()` at each iteration, completing four steps: zeroing the gradients, forward-computing J , backpropagating to obtain the gradient, and updating δz . Since L-BFGS calls the closure multiple times during line search, the computational graph is rebuilt at each call, ensuring that the gradient always corresponds to the δz of the current iteration step. This process is repeated until the convergence criterion is satisfied, yielding the final analysis increment δz . The simplified pseudo-code is shown in Fig. 2.

```

# Forward cost function
def compute_cost(δz):
    r = H @ F @ δz - d
    J = 0.5 * δz.T @ δz + 0.5 * r.T @ R_inv @ r
    return J

# L-BFGS closure
def closure():
    optimizer.zero_grad() # clear gradients
    J = compute_cost(δz) # forward pass
    J.backward() # autograd: ∂J/∂δz
    return J

# Minimization
optimizer = torch.optim.LBFGS([δz], line_search_fn='strong_wolfe', ...)
for _ in range(max_iter):
    optimizer.step(closure)

```

Figure 2. Simplified pseudo-code of the AD-based 3DVar minimization in PY3DVar. The control variable δz is the only differentiable quantity; F , H , d , and R are constants. R_inv denotes the inverse of the observation error covariance R .

2.6 Background error covariance modeling

The key components of the background error covariance matrix B , namely, the background error standard deviation matrix D , the vertical background error correlation matrix C_z , and the horizontal correlation scale parameter L , are statistically estimated using the National Meteorological Center (NMC) method (Parrish and Derber, 1992). In this procedure, 24-hour and 48-hour forecast experiments are initiated daily at 00:00 UTC for 30 consecutive days. The forecast difference field ε is then constructed from forecasts at the same time, calculated as $\varepsilon = X_{48h} - X_{24h}$. These forecasts were run with the same model configuration as the assimilation experiments, ensuring consistency between the estimated background error statistics and the assimilation cycles.

The background error standard deviation matrix D is a diagonal matrix. The values for each control variable in D are computed from the standard deviation of the 30-day forecast difference field:

$$D = \sqrt{\frac{1}{N-1} \sum_{i=1}^N (\varepsilon_i - \bar{\varepsilon})^2} \quad (5)$$

where N is the number of days ($N=30$ in this study). The vertical profiles of the background error standard deviation for each control variable are shown in Fig. 3.

The vertical correlation matrix C_z is obtained by calculating the correlation coefficients of the same variable between different model layers. Its matrix elements can be expressed as:

$$[C_z]_{mn} = \frac{\text{cov}(m,n)}{\sigma_{\varepsilon_m} \cdot \sigma_{\varepsilon_n}} \quad (6)$$

Here, ε_m and ε_n represent the forecast differences at the m -th and n -th layers, respectively, cov denotes the covariance, and σ represents the standard deviation. The distribution of the vertical autocorrelation matrix for the background error of each control variable is shown in Fig. 4.

360 For the horizontal correlation matrices C_x and C_y , horizontal isotropy is assumed. Consequently,
 361 they can be represented by the same decay function, differing only in their dimensions. Their physical
 362 meaning is to describe the horizontal correlation between different grid points in the longitudinal or
 363 latitudinal direction, specifically fitted using a Gaussian kernel function:

$$364 \quad C(x_1, x_2) = \exp[-(x_1 - x_2)^2 / (2L^2)] \quad (7)$$


365 Here, L is the horizontal correlation scale parameter for the background error, x_1 and x_2 represent
 366 horizontal grid points.

367 Horizontal correlation scale parameter L is estimated by analyzing the decay of correlation for
 368 the same variable across different horizontal grid distances. The correlation coefficient $\rho(d)$ between
 369 all pairs of horizontal grid points is computed as a function of their distance d . L is defined as the
 370 maximum grid distance d at which the horizontal correlation coefficient $\rho(d)$ decays to $e^{-1/2}$:

$$371 \quad L = \max \{ d \mid \rho(d) \geq e^{-1/2} \} \quad (8)$$

372 The horizontal correlation scale parameter L quantitatively describes the decay characteristics of
 373 background error correlation between different grid points with increasing distance. Its physical
 374 meaning is that when the distance between two horizontal grid points exceeds L , the correlation
 375 coefficient between their background errors falls below $e^{-1/2}$ (approximately 0.6065), indicating that
 376 their statistical correlation has significantly weakened. This parameter quantifies the decay rate of
 377 horizontal correlation and is a key parameter for describing the horizontal structure of the background
 378 error covariance. The computed values of L are used to construct the horizontal correlation matrices C_x
 379 and C_y , which characterize the spatial horizontal correlation structure and directly determine the shape
 380 of the Gaussian correlation kernel function. The decay of the horizontal background error correlation
 381 coefficient with distance and the corresponding L value for each control variable are shown in Fig. 5.


382 To achieve efficient computation, the three correlation matrices C_x , C_y and C_z are further
 383 decomposed via Cholesky decomposition. Each is expressed as the product of a lower triangular matrix
 384 and an upper triangular matrix (e.g., $C_x = C_{x_low} \cdot C_{x_up}$), where these two triangular matrices are
 385 transposes of each other (i.e., $C_{x_up} = C_{x_low}^T$, where T denotes transpose). During the actual
 386 assimilation process, only the upper triangular matrices for these three directions are retained and used,
 387 thereby avoiding direct manipulation of the full B matrix.



388

389 **Figure 3.** Vertical profiles of background error standard deviation for each control variable: (a) EC,
 390 NH₄, SO₄, NO₃; (b) ORG, CPO, CL, P25; (c) DUST1, DUST2; (d) ANTHA, SOILA; (e) SO₂, NO₂,
 391 CO; (f) O₃.


392 Fig. 3 shows the vertical profiles of the background error standard deviation, which quantifies the
 393 uncertainty of the model background field across different vertical layers and for different variables.
 394 In the 3DVar assimilation process, the D matrix assigns the relative weights of the analysis increments
 395 among the control variables in response to observational information. A larger background error
 396 standard deviation for a variable indicates higher uncertainty in the background field simulation,
 397 allowing for greater adjustment by observational information during assimilation. Conversely, a
 398 smaller standard deviation implies higher confidence in the background field, resulting in more
 399 moderate increments. Fig. 3 reveals significant differences in the near-surface values among the
 400 variables. The background error standard deviations for DUST1 and DUST2 are the largest,
 401 substantially higher than those of other variables. With increasing height, the background error
 402 standard deviations for most variables generally exhibit a decaying trend, and the differences between
 403 them gradually diminish. In contrast, O₃ exhibits an increasing trend with height, reflecting the
 404 enhanced background uncertainty associated with its photochemical production and longer lifetime in
 405 the upper troposphere. In terms of profile shape and decay rate, DUST1 and DUST2 decay most rapidly,
 406 while the decay of the other variables is relatively gradual. Additionally, some species show non-
 407 monotonic vertical variations, such as a near-surface peak for SO₂ and NO₂, indicating distinct vertical
 408 uncertainty characteristics for these species.



409

410 **Figure 4.** Distribution of the vertical autocorrelation matrix of background error for each control
411 variable.

412 Fig. 4 presents the vertical autocorrelation matrices of the background error for each control
413 variable. These matrices govern how surface observational information propagates across vertical
414 levels during assimilation. Although the detailed structures differ among variables, all matrices exhibit
415 strong correlations in the near-surface layers that weaken with increasing height. This indicates that
416 surface observations primarily constrain the lower levels of the model, while their influence on the
417 upper levels is more limited.



418

419 **Figure 5.** Decay of horizontal background error correlation coefficient with distance for each control
420 variable.

421 Fig. 5 presents the decay curves of the horizontal background error correlation coefficient with
422 grid distance for each control variable, along with their corresponding horizontal correlation scale
423 parameter L. These curves are derived directly from the empirical correlation statistics computed over
424 the 30-day forecast difference fields, rather than from a mathematical fitting process. This correlation
425 coefficient determines the horizontal spread and weight allocation of observational information. The

horizontal axis represents physical distance, calculated from a model grid spacing of approximately 27 km. To focus on the region of physical significance, the figure only displays distances up to 500 km, as correlation coefficients beyond this range are already negligible. These empirical curves exhibit an initially rapid then gradually slowing decay with distance, consistent with the Gaussian-type correlation model. The decay rates differ considerably among the variables, yielding L values that range from 1 to 4 grid units. This spread reflects the heterogeneity in the horizontal spatial correlation of background errors for different pollutants.

3. Experimental design and evaluation metrics

3.1 Experimental design

To systematically evaluate the performance and effectiveness of the PY3DVar assimilation system, this study designed three categories of numerical experiments: idealized experiment, 20-day real-case assimilation and forecast experiments, and multi-core CPU parallel performance benchmark tests.

Computing platforms. The CPU platform is a single node equipped with two AMD EPYC 7542 processors (32 physical cores per processor, 64 cores in total @ 2.9 GHz) and 256 GB of system memory. The GPU platform employs an NVIDIA RTX 3090 with 24 GB of device memory. The FT3DVar system runs exclusively on CPU, whereas PY3DVar is deployed on both CPU and GPU platforms. All calculations adopted 32-bit floating-point precision.

Idealized experiment. An idealized twin-observation experiment was designed based on the 27 km grid. Its purpose is to test the system's capability to generate physically reasonable increment fields with observations of varying magnitudes. Two observation sites were positioned far apart to avoid spatial correlation interference, ensuring the independence of constraints from different observation magnitudes. For the low-concentration observation site (37.2104°N, 116.0108°E), the prescribed concentrations were 0.01 $\mu\text{g m}^{-3}$ (SO_2 , NO_2 , O_3), 0.01 mg m^{-3} (CO), 1.00 $\mu\text{g m}^{-3}$ ($\text{PM}_{2.5}$) and 1.50 $\mu\text{g m}^{-3}$ (PM_{10}). For the high-concentration site (37.8666°N, 92.3785°E), the corresponding values were 200 $\mu\text{g m}^{-3}$ (SO_2 , NO_2 , O_3), 10 mg m^{-3} (CO), 500 $\mu\text{g m}^{-3}$ ($\text{PM}_{2.5}$) and 700 $\mu\text{g m}^{-3}$ (PM_{10}). By comparing the analysis field with the background field, the ability of the assimilation system to produce physically sound increment fields under different concentration magnitudes can be verified.

Real-case assimilation and forecast experiments. Twenty consecutive assimilation cycles were conducted at 00:00 UTC each day from October 10 to October 29, 2023. For each cycle, the assimilation was performed using two systems: (1) FT3DVar on the CPU platform with 64 cores, and (2) PY3DVar on the GPU platform. Following each assimilation, a 24-hour forecast was launched from the respective analysis field. A control forecast was also initiated from the original, unassimilated background field (labeled Control). All three forecasts for each cycle shared identical model configurations except for the initial conditions.

Multi-core CPU parallel benchmark tests. To evaluate the parallel scalability of both FT3DVar and PY3DVar on CPU platforms, the 20 assimilation cases were repeated using 1, 2, 4, 8, 16, 32, and 64 CPU cores. Only the assimilation step was performed in these tests, without the subsequent 24-hour forecast. This experiment enables a direct comparison of how the computational efficiency of each system scales with increasing parallel resources.

3.2 Evaluation metrics

To quantitatively evaluate the assimilation and forecasting performance of both PY3DVar and FT3DVar, this study employs three statistical metrics: the correlation coefficient (CORR), root mean square error (RMSE), and mean absolute error (MAE). CORR measures the linear correlation between simulated and observed values; RMSE is more sensitive to extreme errors and reflects the overall deviation of the simulation results; MAE measures the average absolute error between simulated and observed values and is less sensitive to outliers. The formulas for each metric are as follows:

$$\text{CORR} = \frac{\sum_{i=1}^N (M_i - \bar{M})(O_i - \bar{O})}{\sqrt{\sum_{i=1}^N (M_i - \bar{M})^2 \sum_{i=1}^N (O_i - \bar{O})^2}} \quad (9)$$

$$\text{RMSE} = \sqrt{\frac{1}{N} \sum_{i=1}^N (M_i - O_i)^2} \quad (10)$$

$$\text{MAE} = \frac{1}{N} \sum_{i=1}^N |M_i - O_i| \quad (11)$$

Where:

M_i and O_i represent the simulated value and observed value for the i -th sample, respectively.

$\bar{M}$ and $\bar{O}$ are the average values of the simulated and observed values, respectively.

N is the total number of samples.

4. Results and evaluation

4.1 Evaluation of the idealized experiment

Figs. 6 and 7 present the surface increment fields generated by PY3DVar and FT3DVar in the idealized experiment, while Figs. 8 and 9 show the corresponding vertical cross-sections. The $\text{PM}_{2.5}$ and PM_{10} increments are diagnosed as the sum of the relevant aerosol control variable increments, while the gaseous pollutant increments are those of the corresponding control variables.

For both systems, the increments exhibit Gaussian-type horizontal distributions centered on the observation points, decaying with distance, and columnar vertical structures that weaken with height. The vertical extent of the increments is controlled by the vertical correlation matrix C_z , while their

horizontal spread is governed by the horizontal correlation scale parameter L , consistent with the background error characteristics described in Section 2.6. Differences in L and C_z among pollutants account for the observed variations in horizontal spread and vertical column thickness.




Figure 6. Surface increment fields for the six pollutants ((a) SO_2 , (b) NO_2 , (c) CO , (d) O_3 , (e) $\text{PM}_{2.5}$, (f) PM_{10}) from the PY3DVar assimilation system in the idealized experiment. Blue and red colors represent negative and positive increments, respectively.




Figure 7. Surface increment fields for the six pollutants ((a) SO_2 , (b) NO_2 , (c) CO , (d) O_3 , (e) $\text{PM}_{2.5}$, (f) PM_{10}) from the FT3DVar assimilation system in the idealized experiment. Blue and red colors represent negative and positive increments, respectively.




Figure 8. Vertical cross-sections of the increment fields for the six pollutants ((a) SO₂, (b) NO₂, (c) CO, (d) O₃, (e) PM_{2.5}, (f) PM₁₀) from the PY3DVar assimilation system in the idealized experiment. Blue and red colors represent negative and positive increments, respectively.





Figure 9. Vertical cross-sections of the increment fields for the six pollutants ((a) SO₂, (b) NO₂, (c) CO, (d) O₃, (e) PM_{2.5}, (f) PM₁₀) from the FT3DVar assimilation system in the idealized experiment. Blue and red colors represent negative and positive increments, respectively.

Fig. 10 further compares the analysis fields under high and low concentration conditions, showing that both systems effectively correct the background field toward the observations, with negligible differences between their analysis values. Together, Figs. 6–10 demonstrate that PY3DVar and FT3DVar produce highly consistent increment fields and analysis values under idealized conditions, confirming that PY3DVar achieves numerical accuracy and physical consistency comparable to



514


515 **Figure 10.** Comparison of background field (gray), observation (black), FT3DVar analysis field (red), and
516 PY3DVar analysis field (blue) under extremely high (a) and low (b) concentration conditions.

517 **4.2 Evaluation of the analysis fields**


518 Figs. 11 and 12 present scatter plots comparing simulated and observed concentrations across the
519 20-day period for the Control run, the PY3DVar assimilation experiment, and the FT3DVar
520 assimilation experiment, respectively. In these figures, blue dots represent the control experiment,
521 while red dots represent the assimilation experiments. PY3DVar was run on the GPU platform, while
522 FT3DVar was run on the CPU platform with 64 cores. Regarding the distribution pattern, the blue dots
523 from the Control experiment are relatively scattered and deviate considerably from the 1:1 line. In
524 contrast, the red dots from the assimilation experiments show a clear convergence towards the 1:1 line,
525 indicating that both assimilation systems effectively improve the quality of the initial field, with a very
526 pronounced improvement effect.

527 Examining the specific evaluation metrics, the degree of improvement in the initial fields varies
528 among different pollutants. After assimilation, the CORR values for all pollutants increased
529 significantly. The CORR reached 0.58 for SO₂ and 0.48 for CO, those for NO₂, O₃ and PM_{2.5} were
530 above 0.8, and the CORR for PM₁₀ was 0.79. Concurrently, the RMSE and MAE for all pollutants

531 decreased substantially compared to the Control experiment, further confirming the corrective effect
 532 of assimilation on initial field biases. The scatter distributions from PY3DVar and FT3DVar are nearly
 533 identical across all pollutants, and the corresponding CORR, RMSE, and MAE values are essentially
 534 identical. This strong quantitative agreement, consistent with the findings of the idealized experiment,
 535 confirms that the two systems achieve equivalent assimilation performance in real-case applications.



536
 537 **Figure 11.** Scatter plots of observations against Control (blue) and PY3DVar analysis (red)
 538 concentrations. The black line is the 1:1 line.



539
 540 **Figure 12.** Scatter plots of observations against Control (blue) and FT3DVar analysis (red)
 541 concentrations. The black line is the 1:1 line.

542 4.3 Iteration behavior and computational efficiency

543 This section comprehensively evaluates the iteration behavior and computational efficiency of
544 the conventional FT3DVar and the developed PY3DVar (GPU-based implementation) using 20
545 consecutive days of real atmospheric assimilation experiments. The analysis encompasses iteration
546 counts, total and per-iteration time consumption, and cost function convergence characteristics, as
547 presented in Fig. 13.

548 Fig. 13a illustrates the number of iterations required per assimilation cycle throughout the 20-day
549 experiment. FT3DVar requires an average of 34.6 iterations to converge, whereas PY3DVar converges
550 in only 9.9 iterations on average. As discussed in Section 2.5, both systems adopt identical L-BFGS
551 parameter settings and the same convergence criterion; the primary difference lies in their internal line
552 search algorithms — the Moré-Thuente scheme for FT3DVar versus the Strong Wolfe scheme for
553 PY3DVar. The Strong Wolfe line search generally yields a larger per-iteration reduction in the cost
554 function, which partly explains the substantially fewer iterations required by PY3DVar to achieve
555 convergence.

556 Fig. 13b presents the total iteration time per cycle. On average, FT3DVar requires 32.7 s per cycle,
557 while PY3DVar requires only 1.2 s, yielding a $26.2\times$ speedup over the traditional system. The per-
558 iteration time shown in Fig. 13c further elucidates the source of this performance discrepancy. The
559 mean per-iteration time is 0.94 s for FT3DVar and 0.13 s for PY3DVar, corresponding to a $7.5\times$
560 speedup. Combined with the significantly reduced iteration count (Fig. 13a), these two factors
561 collectively account for the high overall efficiency of PY3DVar. The GPU-accelerated execution of
562 vectorized tensor operations, which is central to the PyTorch framework, is the primary contributor to
563 the drastic per-iteration time reduction.

564 Fig. 13d displays the mean descent curves of the cost function within the common iteration range.
565 All curves exhibit a typical nonlinear convergence pattern, characterized by a rapid initial drop
566 followed by a gradually slowing decline, eventually leveling off as the systems approach the optimal
567 solution. Consistent with the fewer iterations depicted in Fig. 13a, PY3DVar reduces the cost function
568 much more aggressively in the early iterations than FT3DVar. Specifically, the descent curve of
569 FT3DVar becomes notably flat and decreases only very slowly after the 10th iteration. The close
570 agreement between the final cost values of PY3DVar and FT3DVar further confirms that both systems
571 achieve highly consistent assimilation accuracy.




Figure 13. Statistical characteristics of iteration performance for different assimilation systems. (a) Number of iterations; (b) Total iteration time; (c) Mean time per iteration; (d) Mean descent curves of cost function within common iteration steps. Blue and red bars/lines represent FT3DVar and PY3DVar, respectively.

4.4 Evaluation of multi-core CPU scalability

Fig. 14 illustrates the parallel scalability and computational efficiency of FT3DVar and PY3DVar across different CPU core configurations (1, 2, 4, 8, 16, 32, and 64 cores), based on averaged results from a 20-day continuous assimilation experiment.

In terms of per-iteration time, FT3DVar outperforms PY3DVar across all tested core counts, decreasing from 5.47 s to 0.99 s, which reflects the inherent speed of the compiled Fortran implementation. PY3DVar drops more steeply, from 42.81 s to 2.04 s, yet remains consistently slower than FT3DVar at every core configuration. The modest absolute reduction in FT3DVar's per-iteration time is due to its already fast single-core baseline, which leaves limited room for further improvement.

Regarding total iteration time, both systems benefit substantially from increasing parallelism.

FT3DVar declines steadily from 224.3 s at a single core to 32.7 s at 64 cores. PY3DVar starts from a much higher baseline of 423.1 s but exhibits a steeper reduction, reaching 20.2 s at 64 cores. Despite its per-iteration disadvantage, PY3DVar achieves a shorter total assimilation time than FT3DVar at 8 cores (73.9 s vs. 85.7 s), 32 cores (42.4 s vs. 51.1 s) and 64 cores (20.2 s vs. 32.7 s). This reversal results from the combined effect of PY3DVar's substantially fewer iterations (9.9 vs. 34.6; Fig. 13a) and its stronger parallel scaling at higher core counts. At 1, 2, 4, and 16 cores, FT3DVar retains the total-time advantage, as its faster per-iteration speed outweighs the difference in iteration count. The speedup of PY3DVar relative to FT3DVar (Fig. 14c) exceeds unity at 8, 32, and 64 cores, and reaches 1.61 at 64 cores.


In summary, FT3DVar benefits from inherently faster single-core performance, while PY3DVar compensates through stronger parallel scalability and far fewer iterations. These two factors together enable PY3DVar to surpass FT3DVar in total assimilation time at 8, 32, and 64 cores.

599

Figure 14. Parallel scalability of FT3DVar (blue) and PY3DVar (red) across CPU core counts. (a) Mean per-iteration time. (b) Mean total iteration time. (c) Speedup of PY3DVar relative to FT3DVar. Dashed line in (c): speedup = 1.

4.5 Evaluation of 24-hour forecast performance

Figs. 15 and 16 display the 20-day averaged temporal evolution of the CORR, RMSE, and MAE




605 for various pollutants during the 24-hour forecast period, initialized from the analysis fields of
606 PY3DVar and FT3DVar, respectively. Comparing the forecast performance of the two systems reveals
607 that the curves of the forecast metrics for all pollutants are fundamentally consistent in terms of
608 morphological characteristics, temporal trends, and numerical values. This finding corroborates the
609 previous results showing minor differences in idealized experiment increment fields, similar analysis
610 field effectiveness in the real case, and comparable cost function convergence characteristics during
611 iteration. It fully demonstrates that the PY3DVar system achieves performance consistent with the
612 mature FT3DVar system not only in initial field assimilation but also in subsequent forecast
613 applications, thoroughly verifying the reliability and practical utility of its assimilation results.

614 Regarding the evolution of forecast performance, although the initial field is significantly
615 optimized by assimilation, this positive effect decays rapidly as the forecast lead time increases
616 (Pagowski et al., 2010; Jiang et al., 2013; Li et al., 2013). This decay primarily stems from inherent
617 model uncertainties, spatiotemporal errors in emission inventories, and inaccuracies in the simulated
618 meteorological fields affecting pollutant dispersion and transport. These errors accumulate during the
619 forecast process and gradually become dominant, thereby progressively diminishing the positive effect
620 of the assimilation improvements.


621 It is noteworthy that the duration of the positive assimilation effect varies significantly among
622 different pollutants. The improvement for SO₂, CO, PM_{2.5}, and PM₁₀ can be maintained for over 24
623 hours, whereas the positive effects for NO₂ and O₃ are shorter, lasting approximately 6 hours and 8
624 hours, respectively. This difference is mainly determined by the respective physicochemical properties
625 and dominant life-cycle processes of the pollutants. SO₂, CO, and particulate matter typically have
626 longer atmospheric lifetimes, and their concentration distributions are more influenced by large-scale
627 transport processes. Consequently, improvements in the initial field yield more lasting positive effects.
628 In contrast, NO₂ and O₃ have strong photochemical reactivity. Their concentrations during daytime are
629 more significantly governed by local photochemical processes and boundary layer variations, causing
630 the initial field improvements from assimilation to dissipate quickly.

631 In summary, PY3DVar achieves assimilation accuracy comparable to FT3DVar and delivers
632 sustained forecast improvements. Although these improvements decay with forecast time, the benefits
633 for short-term forecasts are evident, particularly for longer-lived species such as SO₂, CO, and
634 particulate matter.



635

636 **Figure 15.** Temporal evolution of forecast performance for PY3DVar (red) and Control (blue) over the
 637 24-hour period. From left to right: CORR, RMSE, MAE. From top to bottom: SO₂, NO₂, CO, O₃, PM_{2.5},
 638 PM₁₀.



639

640 **Figure 16.** Temporal evolution of forecast performance for FT3DVar (red) and Control (blue) over the
 641 24-hour period. From left to right: CORR, RMSE, MAE. From top to bottom: SO₂, NO₂, CO, O₃, PM_{2.5},
 642 PM₁₀.

643 5. Summary

644 Based on the three-dimensional variational (3DVar) assimilation principle and integrated with the
645 PyTorch DL framework, this study successfully developed a GPU-accelerated atmospheric chemical
646 3DVar assimilation system (PY3DVar). The system is built upon the CMAQ-AERO5 aerosol scheme
647 coupled with the WRF-Chem model. It consolidates 27 aerosol species into 12 control variables and
648 directly designates four gaseous pollutants as additional control variables. The background error
649 covariance matrix is constructed using the NMC method, enabling the simultaneous assimilation of
650 surface concentrations of SO₂, NO₂, CO, O₃, PM_{2.5}, and PM₁₀. At the algorithm level, the core
651 computations replace traditional Fortran loops with PyTorch tensor operations. Leveraging the AD
652 mechanism greatly simplifies the gradient derivation required for the minimization, and the PyTorch
653 built-in L-BFGS optimizer, along with the tensorized computation, facilitates efficient minimization
654 on both CPU and GPU hardware.

655 To comprehensively evaluate the system, this study conducted a series of experiments: an
656 idealized test to verify physical correctness, 20-day real-case assimilation cycles using both FT3DVar
657 and PY3DVar (GPU-accelerated), and parallel scalability assessments across 1, 2, 4, 8, 16, 32, and 64
658 CPU cores. The system's performance was benchmarked against the traditional FT3DVar in terms of
659 assimilation accuracy, computational efficiency, and forecast skill. The idealized experiments
660 demonstrate that PY3DVar propagates observation information through the background error
661 covariance to generate Gaussian-distributed increment fields. These fields exhibit high consistency
662 with those produced by FT3DVar, confirming the physical soundness and numerical correctness of the
663 new framework. In the 20-day real-case assimilation experiments, PY3DVar substantially improved
664 the model background field across all six target pollutants. Compared to the control run, the
665 assimilation significantly enhanced the CORR between the analysis and observations, while notably
666 reducing the RMSE and MAE. Furthermore, the final analysis fields produced by PY3DVar were
667 nearly identical to those of FT3DVar. This agreement verifies that the GPU-based implementation
668 achieves the same high assimilation accuracy as the traditional Fortran system. In terms of
669 computational efficiency, PY3DVar demonstrates a clear advantage, which is most pronounced on
670 GPU hardware. With FT3DVar on 64 CPU cores requiring an average of 32.7 s per assimilation cycle,
671 the GPU-accelerated PY3DVar completes the same task in 1.2 s, a 26.2-fold speedup. The mean
672 iteration count is correspondingly reduced from 34.6 (FT3DVar) to 9.9 (PY3DVar). This speedup
673 results from two factors: the massive parallelism of tensor operations on the GPU, and the faster
674 convergence of the PyTorch L-BFGS optimizer, whose Strong Wolfe line search yields a larger per-
675 iteration reduction in the cost function than the Moré-Thuente scheme used in FT3DVar. On CPU
676 platforms, although PY3DVar remains slower per iteration than FT3DVar across all tested core counts,
677 its substantially fewer iterations and stronger parallel scaling enable it to surpass FT3DVar in total
678 iteration time at 8, 32, and 64 cores. Overall, PY3DVar's computational efficiency arises from the
679 combination of hardware acceleration and an algorithmic advantage in convergence speed, ensuring

680 competitive performance across diverse computing environments. Finally, short-term forecast
681 experiments demonstrate that the positive impacts of assimilation persist over time. The improvements
682 for SO₂, CO, PM_{2.5}, and PM₁₀ last for more than 24 hours, while those for NO₂ and O₃ maintain
683 significance for over 6 hours. Importantly, the decay trends of these forecast improvements are
684 consistent between the PY3DVar and FT3DVar, further validating the robustness and stability of the
685 proposed system across different hardware architectures.

686 In summary, the PY3DVar system developed in this study achieves substantial improvements in
687 computational efficiency compared to the traditional Fortran-based FT3DVar, while maintaining
688 equivalent assimilation accuracy. This is accomplished through GPU-accelerated tensor computation
689 and the faster convergence of the PyTorch L-BFGS optimizer. Additionally, the use of AD significantly
690 simplifies code development by eliminating the need for manual adjoint coding. The results
691 demonstrate that PY3DVar offers an efficient and practical solution for operational atmospheric
692 chemical data assimilation.

693

694 *Code and data availability.* The WRF-Chem model source code can be downloaded at the WRF
695 model download page (<https://www.mmm.ucar.edu/models/wrf>). The FT3DVar and PY3DVar source
696 codes, together with the observation data, the observation-splitting code, and the background error
697 covariance modeling code, are available at <https://doi.org/10.5281/zenodo.23077072> (Ye, 2026).

698

699 *Author contributions.* HY conceived the study, developed the PY3DVar assimilation system,
700 performed all numerical simulations and assimilation experiments, conducted the formal analysis, and
701 wrote the original manuscript. ZZ and WY contributed to the methodology design and assisted in the
702 validation and evaluation of the assimilation system. YH, NL, and YL discussed the assimilation
703 effects of PY3DVar, provided supervision for the research, and reviewed and edited the manuscript.
704 All authors have read and agreed to the published version of the manuscript.

705

706 *Competing interests.* The contact author has declared that neither they nor their co-authors have any
707 competing interests.

708

709 *Acknowledgements.* We would like to express our sincere gratitude to the National Centers for
710 Environmental Prediction (NCEP) for providing the FNL Operational Global Analysis dataset, and to
711 the China National Environmental Monitoring Center (CNEMC) for making available the
712 observational data of atmospheric pollutants.

713

714 *Financial support.* This research is supported by the National Natural Science Foundation of China
715 (Grant Nos. 42430612, 42475166 and 42405149).

716

717 **References**

718 Austin, J.: Toward the four dimensional assimilation of stratospheric chemical constituents, *J. Geophys.*
719 *Res.-Atmos.*, 97, 2569-2588, <https://doi.org/10.1029/91JD02603>, 1992.

720 Beauchamp, M., Karagali, I., Gacitúa, G., Høyer, J. L., Ballarotta, M., and Fablet, R.: Multiscale neural
721 assimilation scheme for high-resolution sea surface temperature reconstruction from satellite
722 observations, *Sci Rep*, 15, 39928, <https://doi.org/10.1038/s41598-025-23682-9>, 2025.

723 Boudier, P., Fillion, A., Gratton, S., Gürol, S., and Zhang, S.: Data Assimilation Networks, *J Adv Model*
724 *Earth Syst*, 15, e2022MS003353, <https://doi.org/10.1029/2022MS003353>, 2023.

725 Carlton, A. G., Bhawe, P. V., Napelenok, S. L., Edney, E. O., Sarwar, G., Pinder, R. W., Pouliot, G. A.,
726 and Houyoux, M.: Model Representation of Secondary Organic Aerosol in CMAQv4.7, *Environ. Sci.*
727 *Technol.*, 44, 8553-8560, <https://doi.org/10.1021/es100636q>, 2010.

728 Carter, W. P. L.: Documentation of the SAPRC-99 chemical mechanism for VOC reactivity assessment,
729 Final Report to California Air Resources Board Contract No. 92-329, and South Coast Air Quality
730 Management District Contract No. 95-308, University of California, Riverside,
731 2000, https://ww2.arb.ca.gov/sites/default/files/classic/research/apr/past/92-392_1.pdf

732 Chattopadhyay, A., Nabizadeh, E., Bach, E., and Hassanzadeh, P.: Deep learning-enhanced ensemble-
733 based data assimilation for high-dimensional nonlinear dynamical systems, *J. Comput. Phys.*, 477,
734 111918, <https://doi.org/10.1016/j.jcp.2023.111918>, 2023.

735 Chen, D., Liu, Z., Davis, C., and Gu, Y.: Dust radiative effects on atmospheric thermodynamics and
736 tropical cyclogenesis over the Atlantic Ocean using WRF-Chem coupled with an AOD data
737 assimilation system, *Atmos. Chem. Phys.*, 17, 7917-7939, <https://doi.org/10.5194/acp-17-7917-2017>,
738 2017.

739 Chen, H., Gao, J., Sun, T., Chen, Y., Wang, Y., and Carlin, J. T.: Assimilation of Water Vapor Retrievals
740 from ZDR Columns Using the 3DVar Method for Improving the Short-Term Prediction of Convective
741 Storms, *Mon. Wea. Rev.*, 152, 1077-1095, <https://doi.org/10.1175/MWR-D-23-0196.1>, 2024.

742 Cheng, S., Min, J., Liu, C., and Arcucci, R.: TorchDA: A Python package for performing data
743 assimilation with deep learning forward and transformation functions, *Comput. Phys. Commun.*, 306,
744 109359, <https://doi.org/10.1016/j.cpc.2024.109359>, 2025.

745 De Luca, P., Galletti, A., Giunta, G., and Marcellino, L.: Recursive filter based GPU algorithms in a
746 Data Assimilation scenario, *J. Comput. Sci.*, 53, 101339, <https://doi.org/10.1016/j.jocs.2021.101339>,

2021.

Denby, B., Schaap, M., Segers, A., Builtjes, P., and Horálek, J.: Comparison of two data assimilation methods for assessing PM₁₀ exceedances on the European scale, *Atmos. Environ.*, 42, 7122 – 7134, <https://doi.org/10.1016/j.atmosenv.2008.05.058>, 2008.

Derber, J. and Rosati, A.: A Global Oceanic Data Assimilation System, *J. Phys. Oceanogr.*, 19, 1333-1347, [https://doi.org/10.1175/1520-0485\(1989\)019%3C1333:AGODAS%3E2.0.CO;2](https://doi.org/10.1175/1520-0485(1989)019%3C1333:AGODAS%3E2.0.CO;2), 1989.

El Akkraoui, A., Trémolet, Y., and Todling, R.: Preconditioning of variational data assimilation and the use of a bi-conjugate gradient method, *Q. J. R. Meteorol. Soc.*, 139, 731-741, <https://doi.org/10.1002/qj.1997>, 2013.

Elbern, H., Schmidt, H., and Ebel, A.: Variational data assimilation for tropospheric chemistry modeling, *J. Geophys. Res.-Atmos.*, 102, 15967-15985, <https://doi.org/10.1029/97JD01213>, 1997.

Engelen, R. J., Serrar, S., and Chevallier, F.: Four-dimensional data assimilation of atmospheric CO₂ using AIRS observations, *J. Geophys. Res.-Atmos.*, 114, 2008JD010739, <https://doi.org/10.1029/2008JD010739>, 2009.

Fast J D , Gustafson W I , Easter R C , et al. Evolution of Ozone, particulates, and aerosol direct radiative forcing in the vicinity of Houston using a fully coupled meteorology-chemistry-aerosol model. *J. Geophys. Res.-Atmos.*, 2006, 111(D21).DOI:10.1029/2005jd006721.

Feng, T., G. Li, J. Cao, N. Bei, Z. Shen, W. Zhou, S. Liu, T. Zhang, Y. Li, G., Zhang, R., Fan, J., and Tie, X.: Impacts of black carbon aerosol on photolysis and ozone, *J. Geophys. Res.*, 110, 10.1029/2005jd005898, 2005.

Feng, T., N. Bei, R. Huang, J. Cao, Q. Zhang, W. Zhou, X. Tie, S. Liu, T. Zhang, X. Su, W. Lei, L. T. Molina, and G. Li.: Summertime ozone formation in Xi'an and surrounding areas, China, *Atmos. Chem. Phys.*, 16, 4323-4342, 2016.

Frydendall, J., Brandt, J., and Christensen, J. H.: Implementation and testing of a simple data assimilation algorithm in the regional air pollution forecast model, DEOM, *Atmos. Chem. Phys.*, 9, 5475-5488, <https://doi.org/10.5194/acp-9-5475-2009>, 2009.

Gao, C., Zhang, X., Xiu, A., Tong, Q., Zhao, H., Zhang, S., Yang, G., Zhang, M., and Xie, S.: Intercomparison of multiple two-way coupled meteorology and air quality models (WRF v4.1.1-CMAQ v5.3.1, WRF-Chem v4.1.1, and WRF v3.7.1-CHIMERE v2020r1) in eastern China, *Geosci. Model Dev.*, 17, 2471-2492, <https://doi.org/10.5194/gmd-17-2471-2024>, 2024.

Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmos. Environ.*, 39, 6957 – 6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.

780 He, X., Zang, Z., Li, Y., Liu, N., Hu, Y., Liu, L., and You, W.: Inversion of a Near-Real-Time China
781 Gridded Hourly SO₂ Emission Inventory Using Deep Learning Combined With 3D-Variational
782 Assimilation, *Geophys. Res. Lett.*, 52, e2024GL114375, <https://doi.org/10.1029/2024GL114375>, 2025.

783 Hu, Y., Zang, Z., Ma, X., Li, Y., Liang, Y., You, W., Pan, X., and Li, Z.: Four-dimensional variational
784 assimilation for SO₂ emission and its application around the COVID-19 lockdown in the spring 2020
785 over China, *Atmospheric Chemistry and Physics*, 22, 13183 – 13200, [https://doi.org/10.5194/acp-22-](https://doi.org/10.5194/acp-22-13183-2022)
786 13183-2022, 2022.

787 Jiang, Z., Liu, Z., Wang, T., Schwartz, C. S., Lin, H., and Jiang, F.: Probing into the impact of 3DVar
788 assimilation of surface PM₁₀ observations over China using process analysis, *JGR Atmospheres*, 118,
789 6738-6749, <https://doi.org/10.1002/jgrd.50495>, 2013.

790 Key, O., Takao, S., Giles, D., and Deisenroth, M. P.: Scalable data assimilation with message passing,
791 *Environ. Data Sci.*, 4, e1, <https://doi.org/10.1017/eds.2024.47>, 2025.

792 Li, Z., Chao, Y., McWilliams, J. C., and Ide, K.: A three - dimensional variational data assimilation
793 scheme for the Regional Ocean Modeling System: Implementation and basic experiments, *J. Geophys.*
794 *Res.*, 113, 2006JC004042, <https://doi.org/10.1029/2006JC004042>, 2008.

795 Li, G., Lei, W., Zavala, M., Volkamer, R., Dusanter, S., Stevens, P., and Molina, L. T.: Impacts of
796 HONO sources on the photochemistry in Mexico City during the MCMA-2006/MILAGO Campaign,
797 *Atmos. Chem. Phys.*, 10, 6551-6567, 10.5194/acp-10-6551-2010, 2010.

798 Li, G., Bei, N., Tie, X., and Molina, L. T.: Aerosol effects on the photochemistry in Mexico City during
799 MCMA-2006/MILAGRO campaign, *Atmos. Chem. Phys.*, 11, 5169-5182, 10.5194/acp-11-5169-2011,
800 2011a.

801 Li, G., Zavala, M., Lei, W., Tsimpidi, A. P., Karydis, V. A., Pandis, S. N., agaratna, M. R., and Molina,
802 L. T.: Simulations of organic aerosol concentrations in Mexico City using the WRF-CHEM model
803 during the MCMA-2006/MILAGRO campaign, *Atmos. Chem. Phys.*, 11, 3789-3809, 10.5194/acp-11-
804 3789-2011, 2011b.

805 Li, G., Lei, W., Bei, N., and Molina, L. T.: Contribution of garbage burning to chloride and PM_{2.5} in
806 Mexico City, *Atmos. Chem. Phys.*, 12, 8751-8761, 10.5194/acp-12-8751-2012, 2012.

807 Li, Z., Zang, Z., Li, Q. B., Chao, Y., Chen, D., Ye, Z., Liu, Y., and Liou, K. N.: A three-dimensional
808 variational data assimilation system for multiple aerosol species with WRF/Chem and an application
809 to PM_{2.5} prediction, *Atmos. Chem. Phys.*, 13, 4265-4278, <https://doi.org/10.5194/acp-13-4265-2013>,
810 2013.

811 Liang, Y., Zang, Z., Liu, D., Yan, P., Hu, Y., Zhou, Y., and You, W.: Development of a three-dimensional
812 variational assimilation system for lidar profile data based on a size-resolved aerosol model in WRF-
813 Chem model v3.9.1 and its application in PM_{2.5} forecasts across China, *Geosci. Model Dev.*, 13, 6285-

814 6301, <https://doi.org/10.5194/gmd-13-6285-2020>, 2020.

815 Liu, D. C. and Nocedal, J.: On the limited memory BFGS method for large scale optimization, *Math.*
816 *Program.*, 45, 503-528, <https://doi.org/10.1007/BF01589116>, 1989.

817 Lorenc, A. C.: Analysis methods for numerical weather prediction, *Q. J. R. Meteor. Soc.*, 112, 1177-
818 1194, <https://doi.org/10.1002/qj.49711247414>, 1986.

819 Ma, X., Liu, H., and Peng, Z.: Improving WRF-Chem PM_{2.5} predictions by combining data
820 assimilation and deep-learning-based bias correction, *Environ. Int.*, 195, 109199,
821 <https://doi.org/10.1016/j.envint.2024.109199>, 2025.

822 Moré, J. J., and Thuente, D. J.: Line search algorithms with guaranteed sufficient decrease, *ACM Trans.*
823 *Math. Softw.*, 20(3), 286–307, <https://doi.org/10.1145/192115.192132>, 1994.

824 Nocedal, J.: Updating quasi-Newton matrices with limited storage, *Math. Comp.*, 35, 773 – 782,
825 <https://doi.org/10.1090/S0025-5718-1980-0572855-7>, 1980.

826 Pagowski, M., Grell, G. A., McKeen, S. A., Peckham, S. E., and Devenyi, D.: Three-dimensional
827 variational data assimilation of ozone and fine particulate matter observations: some results using the
828 Weather Research and Forecasting-Chemistry model and Grid-point Statistical Interpolation, *Quart. J.*
829 *Royal. Meteor. Soc.*, 136, 2013-2024, <https://doi.org/10.1002/qj.700>, 2010.

830 Parrish, D. F. and Derber, J. C.: The National Meteorological Center ' s Spectral Statistical-
831 Interpolation Analysis System, *Mon. Wea. Rev.*, 120, 1747-1763, [https://doi.org/10.1175/1520-0493\(1992\)120%3C1747:TNMCSS%3E2.0.CO;2](https://doi.org/10.1175/1520-0493(1992)120%3C1747:TNMCSS%3E2.0.CO;2), 1992.

833 Seo, M.-G. and Kim, H. M.: Effect of meteorological data assimilation using 3DVar on high-resolution
834 simulations of atmospheric CO₂ concentrations in East Asia, *Atmos. Pollut. Res.*, 14, 101759,
835 <https://doi.org/10.1016/j.apr.2023.101759>, 2023.

836 Tombette, M., Mallet, V., and Sportisse, B.: PM₁₀ data assimilation over Europe with the optimal
837 interpolation method, *Atmos. Chem. Phys.*, 9, 57-70, <https://doi.org/10.5194/acp-9-57-2009>, 2009.

838 Tong, D., Cheng, J., Liu, Y., Yu, S., Yan, L., Hong, C., Qin, Y., Zhao, H., Zheng, Y., Geng, G., Li, M.,
839 Liu, F., Zhang, Y., Zheng, B., Clarke, L., and Zhang, Q.: Dynamic projection of anthropogenic
840 emissions in China: methodology and 2015 – 2050 emission pathways under a range of socio-
841 economic, climate policy, and pollution control scenarios, *Atmos. Chem. Phys.*, 20, 5729 – 5757,
842 <https://doi.org/10.5194/acp-20-5729-2020>, 2020.

843 Wang, D., You, W., Zang, Z., Pan, X., He, H., and Liang, Y.: A three-dimensional variational data
844 assimilation system for a size-resolved aerosol model: Implementation and application for particulate
845 matter and gaseous pollutant forecasts across China, *Sci. China Earth Sci.*, 63, 1366-1380,
846 <https://doi.org/10.1007/s11430-019-9601-4>, 2020.

847 Wang, D., You, W., Zang, Z., Pan, X., Hu, Y., and Liang, Y.: A three-dimensional variational data
848 assimilation system for aerosol optical properties based on WRF-Chem v4.0: design, development,
849 and application of assimilating Himawari-8 aerosol observations, *Geosci. Model Dev.*, 15, 1821-1840,
850 <https://doi.org/10.5194/gmd-15-1821-2022>, 2022.

851 Ye, H.: GPU-accelerated three-dimensional variational data assimilation with automatic differentiation
852 for WRF-Chem, Zenodo [data set], <https://doi.org/10.5281/zenodo.23077072>, 2026.

853 Xing, L., Wu, J., Elser, M., Tong, S., Liu, S., Li, X., Liu, L., Cao, J., Zhou, J., El-Haddad, I., Huang,
854 R., Ge, M., Tie, X., Prévôt, A. S. H., and Li, G.: Wintertime secondary organic aerosol formation in
855 Beijing-Tianjin-Hebei (BTH): Contributions of HONO sources and heterogeneous reactions, *Atmos.*
856 *Chem. Phys. Discuss.*, 2018, 1-25, 10.5194/acp-2018-770, 2018.

857 Xu, X., Sun, X., Han, W., Zhong, X., Chen, L., Gao, Z., and Li, H.: FuXi-DA: a generalized deep
858 learning data assimilation framework for assimilating satellite observations, *npj Clim Atmos Sci*, 8,
859 156, <https://doi.org/10.1038/s41612-025-01039-3>, 2025.

860 Ye, H., Pan, X., You, W., Zhu, X., Zang, Z., Wang, D., Zhang, X., Hu, Y., and Jin, S.: Impact of
861 CALIPSO profile data assimilation on 3-D aerosol improvement in a size-resolved aerosol model,
862 *Atmos. Res.*, 264, 105877, <https://doi.org/10.1016/j.atmosres.2021.105877>, 2021.

863 Ye, H., You, W., Zang, Z., Pan, X., Wang, D., Zhou, N., Hu, Y., Liang, Y., and Yan, P.: Observing
864 system simulation experiment (OSSE)-quantitative evaluation of lidar observation networks to
865 improve 3D aerosol forecasting in China, *Atmos. Res.*, 270, 106069,
866 <https://doi.org/10.1016/j.atmosres.2022.106069>, 2022.

867 Zang, Z., Li, Z., Pan, X., Hao, Z., and You, W.: Aerosol data assimilation and forecasting experiments
868 using aircraft and surface observations during CalNex, *Tellus. B.: Chem. Phys. Meteo.*, 68,
869 <https://doi.org/10.3402/tellusb.v68.29812>, 2016.

870 Zang, Z., You, W., Ye, H., Liang, Y., Li, Y., Wang, D., Hu, Y., and Yan, P.: 3DVar Aerosol Data
871 Assimilation and Evaluation Using Surface PM_{2.5}, Himawari-8 AOD and CALIPSO Profile
872 Observations in the North China, *Remote. Sens.*, 14, 4009, <https://doi.org/10.3390/rs14164009>, 2022.

873 Zhang, J., Campbell, J. R., Reid, J. S., Westphal, D. L., Baker, N. L., Campbell, W. F., and Hyer, E. J.:
874 Evaluating the impact of assimilating CALIOP-derived aerosol extinction profiles on a global mass
875 transport model: CALIOP DATA ASSIMILATION, *Geophys. Res. Lett.*, 38, n/a-n/a,
876 <https://doi.org/10.1029/2011GL047737>, 2011.

877 Zheng, B., Cheng, J., Geng, G., Wang, X., Li, M., Shi, Q., Qi, J., Lei, Y., Zhang, Q., and He, K.:
878 Mapping anthropogenic emissions in China at 1 km spatial resolution and its application in air quality
879 modeling, *Sci. Bull.*, 66, 612-620, <https://doi.org/10.1016/j.scib.2020.12.008>, 2021.

880 Zhou, Y., Sun, W., Liu, Z., Gao, L., Chen, D., Feng, J., Zhang, T., and Zhou, Z.: Further development

881 and application of the WRFDA-Chem three-dimensional variational (3DVar) system: Joint
882 assimilation of satellite AOD retrievals and surface observations, *Atmos. Res.*, 316, 107942,
883 <https://doi.org/10.1016/j.atmosres.2025.107942>, 2025.