



Characterisation of hygroscopic aerosol processes using remote sensing and in situ techniques at the MeteoSwiss Payerne station

Arlett Díaz-Zurita^{1,2}, Simone Brunamonti³, Alexander Haeferle⁴, Martine Collaud Coen⁴, Giovanni Martucci⁴, Benjamin Tobias Brem⁵, Martin Gysel-Beer⁵, Hassan Lyamani², Gloria Titos^{1,2}, Daniel Pérez-Ramírez^{1,2}, Christoph Hüglin³, Lucas Alados-Arboledas^{1,2}, Alejandro Keller⁶, Frank Gunther Wienhold⁴, Maxime Hervé⁴, and Francisco Navas-Guzmán^{1,2}

¹Andalusian Institute for Earth System Research (IISTA), University of Granada, Granada, 18006, Spain

²Department of Applied Physics, University of Granada, Granada, 18071, Spain

³Empa, Laboratory for Air Pollution/Environmental Technology, Dübendorf, Switzerland

⁴Federal Office of Meteorology and Climatology MeteoSwiss, Payerne, Switzerland

⁵PSI Center for Energy and Environmental Sciences, 5232 Villigen PSI, Switzerland

⁶University of Applied Sciences Northwestern Switzerland, Windisch, 5210, Switzerland

Correspondence: Francisco Navas-Guzmán (fguzman@ugr.es)

Abstract.

This study presents a comprehensive assessment of aerosol hygroscopicity during the HYGRO19 field campaign conducted at the MeteoSwiss station in Payerne (Switzerland) from June to December 2019. The campaign provided a unique experimental framework by combining continuous Raman lidar observations with vertically resolved in situ measurements from balloon-borne Compact Backscatter Aerosol Detector coupled with RS41 Vaisala radiosondes (COBALD-RS41), ground-based tandem nephelometers, and complementary aerosol chemical analyses, enabling aerosol hygroscopicity to be evaluated using three independent approaches. Two contrasting aerosol regimes were observed: a summer period characterised by relatively high aerosol loading and frequent influence of long-range transported particles, and a second period dominated by low-pressure systems, persistent cloudiness, and reduced aerosol concentrations. Throughout these different atmospheric conditions, the Raman Lidar for Meteorological Observations (RALMO), which continuously provides independent profiles of aerosol backscatter, water vapour mixing ratio, and temperature during both daytime and nighttime, identified more than 200 well-mixed hygroscopic aerosol layers. Median hygroscopicity parameters were $f_{\beta}(355 \text{ nm}) = 2.64 \pm 0.64$ and $\gamma_{\beta}(355 \text{ nm}) = 0.70 \pm 0.18$. Despite differences in measurement principle, wavelength, and sampling technique, Raman lidar retrievals showed good agreement with COBALD-RS41 and tandem nephelometer observations. Statistical analyses revealed no systematic dependence of aerosol hygroscopicity on altitude, with comparable values in the atmospheric boundary layer and free troposphere. Enhanced hygroscopic growth was generally associated with a larger contribution of inorganic aerosol species, whereas no clear dependence on particle size was identified. These results demonstrate that combining remote sensing and in situ observations provides a robust framework for characterising aerosol hygroscopicity under ambient atmospheric conditions.



1 Introduction

20 Atmospheric aerosols have a significant impact on Earth's radiative balance both directly, through the scattering and absorption of solar radiation, and indirectly, by modifying the microphysical properties of clouds due to their ability to act as cloud condensation nuclei (CCN) or ice nucleating particles (INPs) (Seinfeld et al., 2016). According to the last Intergovernmental Panel on Climate Change (IPCC), the largest uncertainty in the net radiative forcing is primarily dominated by uncertainties associated with the aerosol component (Douville et al., 2021). These uncertainties are mainly associated with the high temporal and spatial variability of aerosol distribution and composition, as well as their complex interactions with clouds (aerosol-cloud interactions, ACI) and precipitation processes (Bindoff et al., 2013; Chen et al., 2020). One of the key aspects contributing to these uncertainties is aerosol hygroscopicity, which refers to the ability of particles to uptake water vapour, thereby altering particle size, chemical composition and optical properties (Chen et al., 2020). Aerosol hygroscopic growth can influence the direct radiative effect by modifying particle size and consequently the light scattering pattern (i.e. the phase function).
30 Moreover, the hygroscopic growth represents a fundamental aerosol parameter leading to indirect radiative effect, as the ability of aerosols to act as CCN is directly related to their affinity for water vapour uptake (Hänel, 1976; Seinfeld et al., 2016).

Several studies have addressed the effects of hygroscopic growth on aerosol optical and microphysical properties using ground-based in situ measurements. Techniques such as the Humidified Tandem Differential Mobility Analyser (HT-DMA) (e.g., Liu et al., 2018; Swietlicki et al., 2008; Pariyothon et al., 2023; Deshmukh et al., 2025; Zhang et al., 2025) and Humidified Tandem Nephelometers (e.g., Titos et al., 2014b; Alonso-Blanco et al., 2019; Burgos et al., 2019; Mouton et al., 2023; Yuan and Zhao, 2023) have been extensively used to quantify changes in particle diameter or aerosol optical properties due to water vapour uptake. However, these techniques also have certain limitations. Most in situ techniques alter the ambient conditions and are subject to particle losses in the sampling lines, which can modify the aerosol properties in ambient conditions (Bedoya-Velázquez et al., 2018). These techniques also present larger uncertainties under conditions close to saturation (high relative humidity (RH)), particularly when the aerosol is more hygroscopic (Titos et al., 2016). However, the most relevant issue is that these measurements are representative of only a few metres above the ground and do not allow for a complete understanding of aerosol hygroscopicity in the vertical.

Remote sensing techniques can complement observations of aerosol hygroscopicity by providing independent measurements of aerosol properties and relative humidity. Furthermore, these measurements can be obtained at high RH without altering the aerosol sample (Feingold and Morley, 2003). Measurements with ground-based Aerosol Robotic Network (AERONET; Holben et al., 1998) can estimate aerosol hygroscopicity through studying changes in aerosol size distributions in relation to column precipitable water vapour (PWV) (Schafer et al., 2008), and also in connection with the dissipation of fog or low-altitude clouds (Eck et al., 2012). However, AERONET measurements are representative of the entire atmospheric column. In contrast, lidar measurements provide vertically resolved information of aerosols, which make such measurements suitable for investigating aerosol hygroscopicity within certain atmospheric layers (Ferrare et al., 1998). To achieve this, well-mixed atmospheric conditions are required to isolate the effect of hygroscopic growth from other aerosol processes (Wulfmeyer and Feingold, 2000; Feingold and Morley, 2003). Nevertheless, as these measurements are performed under real atmospheric



conditions, neither the sample nor the ambient conditions can be controlled, limiting their applicability (Granados-Muñoz et al., 2015), and obtaining RH profiles with high vertical and temporal resolution remains challenging. Furthermore, lidar system quantifies the impact of RH on the backscatter coefficient at a polar angle of 180° , which requires assumptions and calculations to translate it to, for example, the RH effect on the hemispheric backscatter coefficient. Some studies relied on thermodynamic profiles from collocated radiosondes (RS) (Veselovskii et al., 2009; Sicard et al., 2022; Jing et al., 2026; Veselovskii et al., 2025), which limited the application of lidar measurements for aerosol hygroscopicity investigations. To extend the applicability of the technique to other studies, RH profiles have been derived by combining temperature (T) profiles from microwave radiometers (MWR) with water vapour mixing ratio profiles from Raman lidar observations (Navas-Guzmán et al., 2014; Bedoya-Velásquez et al., 2018; Miri et al., 2024). However, uncertainties in the MWR temperature profiles, due to their lower spatial resolution, can affect the accuracy in the derived RH (Navas-Guzmán et al., 2016) and in the retrieved aerosol hygroscopicity parameters (Pérez-Ramírez et al., 2021). Although these approaches have served as useful benchmarks, long-term analyses have been constrained (Sicard et al., 2022; Jing et al., 2026), and the need to also have correlative measurements of aerosol chemical composition and particle size has limited comprehensive aerosol hygroscopicity studies to just a few specific cases during special field campaigns (Pérez-Ramírez et al., 2021; Laly et al., 2025).

The Raman Lidar for Meteorological Observations (RALMO) system, operating at the aerological station of MeteoSwiss in Payerne, Switzerland, can overcome the previously mentioned limitations in aerosol hygroscopicity studies due to its capability of providing high-vertical-resolution profiles of aerosol, temperature and water vapour mixing ratio (Dinoev et al., 2013). The capabilities of the system allow the instrument to operate continuously, during both daytime and nighttime, and have already been proven to be ideal for aerosol hygroscopicity studies (Navas-Guzmán et al., 2019). In this context, the HYGRO19 field campaign, which was conducted from June to December 2019 in Payerne, represents a unique opportunity to continuously investigate aerosol hygroscopicity due to the capabilities of RALMO. Furthermore, vertical aerosol measurements performed with the Compact Backscatter Aerosol Detector (COBALD, Brunamonti et al., 2018), alongside RH profiles from operational radiosondes (RS41, Vaisala), permitted the direct validation of aerosol hygroscopicity parameters retrieved from RALMO against those derived from vertically resolved in situ observations. Additionally, complementary in situ measurements, including tandem nephelometers, were performed at ground level, enabling the evaluation of lidar retrievals under well-mixed atmospheric conditions and providing a proxy for aerosol hygroscopicity based on aerosol chemical properties. Nevertheless, some limitations arise from the different approaches and wavelengths used to obtain aerosol hygroscopicity parameters, since lidars rely solely on backscatter measurements, whereas tandem nephelometers use the scattering coefficient.

This work focuses on the analysis of hygroscopic aerosol growth during the HYGRO19 field campaign. The unique capabilities of RALMO enable the assessment of a large database of aerosol hygroscopicity parameters under real atmospheric conditions. Comprehensive closure studies are presented by combining RALMO observations with ground-based in situ measurements. Specific case studies are presented to investigate the validation of aerosol hygroscopicity against COBALD-RS41 sondes. To address these objectives, Sect. 2 describes the experimental site and instrumentation, Sect. 3 reviews the different methodologies used, Sect. 4 presents the main results and their discussion, and finally, Sect. 5 summarises the conclusions and provides perspectives for future work.



2 Experimental site and instrumentation

2.1 The HYGRO19 field campaign

90 The intercomparison campaign of hygroscopic aerosol properties using remote sensing and in situ techniques (HYGRO19 campaign) was carried out from June to December 2019 at the MeteoSwiss aerological station in Payerne, Switzerland (46.82° N, 6.95° E, 491 m above sea level (asl)), which forms part of ACTRIS (Aerosol, Clouds and Trace Gases Research Infrastructure), EARLINET (European Aerosol Research Lidar Network), E-PROFILE (European Profiling Network), and GRUAN (GCOS Reference Upper-Air Network). The main objective of the campaign was to assess the ability of the different techniques to characterise hygroscopic aerosol processes. To meet this goal, three distinct approaches were employed: the first, based on remote sensing measurements from the RALMO Raman lidar, the second, involved in situ measurements using a tandem nephelometers at ground level; and the third, consisted in situ vertical profiling with COBALD sondes operating alongside RS41 radiosondes (COBALD-RS41). The experimental station is located in a rural area of the Swiss Plateau, between the Jura Mountains (25 km to the Northwest) and the Alpine foothills (20 km to the Southeast). The site is characterised by a mid latitude continental climate with some influence of long-range transported pollution (Navas-Guzmán et al., 2019).

2.2 The RALMO Raman lidar system

The Raman Lidar for Meteorological Observations (RALMO) system is a custom-designed instrument developed by the Swiss Federal Institute of Technology (EPFL) in collaboration with MeteoSwiss. It has been operating in the Payerne station since August 2008 and joined EARLINET the same year. The RALMO system uses a Nd:YAG laser source that emits pulses at 105 355 nm (400 mJ), with a repetition rate of 30 Hz and a pulse duration of 8 ns. Before the laser emits to the atmosphere it is expanded to a diameter of 14 cm to reduce beam divergence (to 0.1 mrad). The receiving system consists of four telescopes with 30 cm parabolic mirrors symmetrically arranged with the vertically mounted beam expander to collect the backscattered photons, giving a field of views (FOV) of 0.2 mrad. These characteristics enable RALMO to operate during both daytime and nighttime. Optical fibres connect the telescope mirrors to two grating polychromators, which isolate the rotational-vibrational (VR) and pure-rotational (RR) signals. A detailed description of RALMO can be found in Dinov et al. (2013).

The system optimises its operability in the incomplete overlap region because the Raman backscatter is excited by a single monochromatic beam and the designed reflective-type lidar telescopes minimise chromatic aberrations. Moreover, the fibre-coupled diffraction grating-based polychromator further reduces uncertainties induced by range-dependent optical imperfections of the lidar receiver to a negligible level (Dinov et al., 2013). The water vapour mixing ratio is computed from 115 VR measurements of nitrogen and water vapour at wavelengths of 386.7 and 407.5 nm. The accuracy of RALMO water vapour measurements has been validated by Brocard et al. (2013), showing good agreement with radiosonde observations and integrated water vapour retrieved from the Global Positioning System (GPS) and MWR.

Temperature measurements with RALMO are obtained using four sections of the RR Raman spectrum around 355 nm, which relies on the rotational Raman scattering of oxygen and nitrogen molecules. The RR signal is separated from the elastic (Rayleigh and Mie) scattering using a two-stage temperature polychromator, which isolates specific rotational Raman lines 120



while guiding the elastic signal through optical fibres to the detector. High-efficiency parabolic mirrors collect the signals, which are then filtered, fibre-coupled, and diffracted by a reflective holographic grating to resolve Stokes and anti-Stokes lines from different rotational quantum numbers. This configuration enables accurate and stable retrieval of atmospheric temperature profiles. Further details and a complete validation of the RALMO temperature channel are provided by Martucci et al. (2021).

125 The system also registers the backscattered signal at 355 nm backscattered signal, which combined with one RR measurement allows the retrieval of independent aerosol backscatter and extinction measurements (Veselovskii et al., 2015).

For water vapour, nitrogen, and oxygen, the interference filters are centred at 407.5, 386.7 and 375.4 nm, respectively, each with a full width at half maximum (FWHM) of 0.33 nm. This configuration ensures minimal temperature sensitivity of the Raman scattering signal, with a relative change in transmitted intensity of 0.4 % per 100 K (Dinoev et al., 2013), in agreement
130 with values reported in the literature (Whiteman, 2003). It also minimises spectral differences between the laser emission and Raman channels (Veselovskii et al., 2013, 2015).

2.3 The COBALD-RS41 sondes

COBALD is a lightweight (500 g) aerosol backscatter detector for balloon-borne measurements developed at the Swiss Federal Institute of Technology (ETH Zürich) based on the original prototype by Rosen and Kjome (1991). Using two light-emitting
135 diodes (LEDs) as light sources and a photodiode detector with FOV of 6°, COBALD provides aerosol backscatter at 455 and 940 nm with an accuracy of 5 % and a precision of 1 % for both channels, and temporal and spatial resolutions of 1 s and 5 m, respectively (Vernier et al., 2015). During the HYGRO19 field campaign, COBALD was combined with an additional meteorological radiosonde (RS41, Vaisala) to obtain measurements of temperature, pressure, and relative humidity. The RS41 sonde specifications for nominal measurement uncertainties, as provided by the manufacturer, are 0.3 °C for temperature below
140 16 km and 0.4 °C above that altitude, 0.01 hPa (decreasing with altitude) for pressure, and 4 % RH (Vaisala, 2017).

The COBALD measurements are typically expressed as the backscatter ratio (BSR, Eq. 1), defined as the ratio of the total measured signal (normalised to the altitude-dependent LED power) to its molecular contribution (Brunamonti et al., 2021). The molecular contribution is computed following Bucholtz (1995) using the atmospheric number density of molecules derived from the collocated RS (Cirisan et al., 2014). A molecular extinction-to-backscatter ratio of $8\pi/3$ sr is assumed.

$$145 \quad BSR = \frac{\beta^{aer}(\lambda) + \beta^{mol}(\lambda)}{\beta^{mol}(\lambda)} \quad (1)$$

During the HYGRO19 field campaign, the COBALD–RS41 sonde was used to obtain vertically resolved in situ observations of aerosol hygroscopicity parameters, providing the reference measurements for direct comparison with those retrieved from RALMO. However, the COBALD and RALMO systems differ in receiver FOV (0.01° for RALMO), which results in systematic differences in backscatter measurements that are corrected using Mie scattering simulations (Brunamonti et al., 2021).

150 2.4 The tandem nephelometers

The tandem nephelometers consisted of two integrating nephelometers operating in parallel and measuring the same air sample. One instrument operated at RH lower than 40 % (dry conditions), whereas the second was equipped with a humidification



system allowing measurements at different RH conditions from 40 to 90 % (wet conditions) (Titos et al., 2014a). The dry scattering coefficients ($\sigma_{sp, dry}$) were measured using an Ecotech Aurora 3000 nephelometer at three wavelengths (425, 525, and 635 nm) with a 1 min sampling interval. The scattering coefficients under wet conditions were measured using a TSI 3563 integrating nephelometer at 450, 550, and 700 nm, recording data every 1 min. The nephelometer data were corrected to account for angular truncation errors and instrument non-idealities, following the method of Anderson and Ogren (1998) for the TSI instrument and Müller et al. (2011) for the Ecotech nephelometer. The setup consisted of a 30 min cycle beginning with a baseline (zero) measurement, followed by gradual increases of RH up to approximately 85–90 %. For the computations of the hygroscopicity parameters, the dry scattering coefficient measured at 525 nm by the Ecotech instrument was corrected to 550 nm using the Ångström exponent (Ångström, 1929).

2.5 Complementary instrumentation

During the HYGRO19 campaign, complementary instruments were also employed to provide a comprehensive characterisation of aerosols. Continuous measurements of aerosol optical depth (AOD) and column-integrated extinction Ångström exponent (AOD-AE) were obtained from an AERONET sun photometer (SP) operating at Payerne since 2000 (Level 2 Version 3, Giles et al., 2019). The station is also part of the European E-PROFILE network (Haefele et al., 2016) since 2008 and operates a CHM15k ceilometer (Lufft) emitting at 1064 nm, which continuously records aerosol attenuated backscatter. Furthermore, within the framework of E-PROFILE, a Doppler wind lidar profiler operates to measure horizontal and vertical wind components every 10 min, typically covering altitudes from 150 m to about 5 km above ground level (agl) (Nowak, 2008). Ground-based in situ measurements included an Aethalometer (AE31), operated by the Centre for Energy and Environmental Sciences at the Paul Scherrer Institute (PSI), capable of providing aerosol absorption coefficients (σ_{ab}) at seven wavelengths (from 370 to 950 nm). The Aethalometer can also estimate equivalent black carbon (EBC) concentrations using wavelength-dependent mass absorption cross sections (Petzold et al., 2013). Data processing followed Global Atmosphere Watch (GAW)/ACTRIS recommendations, including corrections for filter-loading and multiple-scattering effects and was harmonised as described in Zanatta et al. (2016).

Additional in situ measurements of aerosol properties were conducted by the Swiss Federal Laboratories for Materials Science and Technology (EMPA) as part of the NABEL (National Air Pollution Monitoring Network) programme (EMPA, 2013). Particle mass concentrations of $PM_{2.5}$ and PM_{10} were determined in real time with a Fidas 200 optical aerosol spectrometer and corrected versus gravimetric reference data (Hüglin and Schwarzenbach, 2016). Elemental carbon (EC) in $PM_{2.5}$ was obtained in the laboratory from $PM_{2.5}$ filter samples using a thermal optical EC analyser (Sunset Laboratory Inc.) following the EUSAAR2 protocol (Cavalli et al., 2010). The concentrations of sulphate (SO_4^{2-}) and nitrate (NO_3^-) in PM_{10} were determined from PM_{10} filter samples using ion chromatography (Hüglin et al., 2024). Ammonium (NH_4^+) was estimated assuming complete neutralisation of sulphate and nitrate (Jones et al., 2021; Liu et al., 2021), and organic matter (OM) was derived from PM mass closure ($PM = \text{inorganic species} + \text{black carbon (BC)} + \text{OM}$). Furthermore, total carbon (TC) concentrations were obtained using the novel FATCAT (Fast Thermal Carbon Totalizer) system, developed by the University of Applied Sciences and Arts Northwestern Switzerland (FHNW), with a detection limit of 0.16 $\mu\text{g-C}$ (micrograms of carbon) for a sampling interval



of 2 h (Keller et al., 2023). Additionally, aerosol size distributions in the range 10 to 487 nm, were measured using a scanning mobility particle sizer (SMPS, TSI 3034) (PSI, Durand et al., 2023).

3 Methodology

190 3.1 Aerosol optical properties from Raman lidar measurements

The combined use of elastic (Klett, 1981, 1985) and Raman (Ansmann et al., 1990) backscattered signals allows the independent calculation of the aerosol backscatter coefficient (β^{aer}) as described by Ansmann et al. (1990) and Whiteman (2003):

$$\beta^{aer}(z, \lambda_L) = \beta^{mol}(z, \lambda_L) \cdot \left[k \cdot \frac{P(z, \lambda_L)}{P(z, \lambda_i)} - 1 \right] \quad (2)$$

195 where the sub-index i indicates the Raman molecular species, $\beta^{mol}(z, \lambda_L)$ is the molecular backscatter coefficient, k is the calibration constant, and $P(z, \lambda)$ is the backscattered signal from range z at either the elastic (λ_L) or the Raman shifted λ_i wavelengths. Stokes and anti-Stokes portions of the pure-rotational spectrum with opposite temperature dependence are summed, giving in good approximation a temperature-independent inelastic signal (Dinoyev et al., 2013). The RALMO system provide accurate aerosol backscattered and extinction coefficients at 355 nm with a bias of $-2 \% \pm 37 \% (-0.018 \pm 0.237$
200 $\text{Mm}^{-1}\text{sr}^{-1})$ compared with COBALD at 455 nm (Brunamonti et al., 2021).

3.2 Water vapour mixing ratio, temperature and relative humidity profiles from Raman lidar measurements

Water vapour mixing ratio (r) is defined as the ratio of the mass of water vapour to the mass of dry air in a sample of the atmosphere (Goldsmith, 1998). Consequently, the ratio $N_{H_2O}(z)/N_{N_2}(z)$ is proportional to the water vapour mixing ratio (Whiteman et al., 1992, 2006). Assuming identical overlap factors and the range-independent Raman backscatter cross sections
205 for the two signals, this ratio can be expressed as (Whiteman et al., 2006):

$$\frac{N_{H_2O}(z)}{N_{N_2}(z)} = \frac{P(z, \lambda_{H_2O})}{P(z, \lambda_{N_2})} \cdot \frac{F_{N_2}[T(z)]}{F_{H_2O}[T(z)]} \cdot \frac{K_{N_2}\sigma_{N_2}}{K_{H_2O}\sigma_{H_2O}} \cdot \exp \left\{ \int_0^z [\alpha(z', \lambda_{H_2O}) - \alpha(z', \lambda_{N_2})] dz' \right\} \quad (3)$$

and thus,

$$r(z) = \frac{P(z, \lambda_{H_2O})}{P(z, \lambda_{N_2})} \cdot \frac{F_{N_2}[T(z)]}{F_{H_2O}[T(z)]} \cdot K \cdot \exp \left\{ \int_0^z [\alpha(z', \lambda_{H_2O}) - \alpha(z', \lambda_{N_2})] dz' \right\} \quad (4)$$

210 where the term $P(z, \lambda_{H_2O})/P(z, \lambda_{N_2})$ represents the backscattered signal ratio, $F_{N_2}[T(z)]/F_{H_2O}[T(z)]$ is the ratio of the temperature-dependent functions for the Raman nitrogen and water vapour channels, and K is the instrument calibration constant. For RALMO the calibration constant was determined by combining absolute calibration using collocated radiosondes with relative calibration based on the solar background (Martucci et al., 2018). The passband filter employed in RALMO



allows the ratio $F_{MR}[T(z)]/F_{H_2O}[T(z)]$ to be reasonably approximated as 1 (Sect. 3.1), resulting in essentially temperature-independent measurements (Whiteman, 2003).

215 The exponential term (Eq. 4) accounts for the difference in atmospheric transmission between the nitrogen and water vapour Raman wavelengths, mainly due to the λ^{-4} dependence of Rayleigh scattering by molecules and an additional contribution from wavelength-dependent Mie scattering by aerosols. Molecular extinction can be estimated assuming US Standard Atmosphere, while the differential aerosol extinction is generally negligible under typical Payerne conditions (Dinoev et al., 2013), though it can reach up to 10 % in hazy conditions (Whiteman, 2003; Díaz-Zurita et al., 2025).

220 Temperature measurements with RALMO are obtained using RR Raman lidar signals (Martucci et al., 2021), and combining these with water vapour mixing ratio measurements, it is possible to obtain RH profiles following Yau and Rogers (1996):

$$RH = \frac{e(z)}{e_w(z)} \cdot 100 \quad (5)$$

where $e(z)$ is the water vapour pressure, and $e_w(z)$ denotes the saturation vapour pressure. The water vapour pressure is related to the water vapour mixing ratio through:

$$225 \quad e(z) = \frac{p(z) \cdot r(z)}{0.622 + r(z)} \quad (6)$$

where $p(z)$ represents the air pressure profile, estimated from the closest RS41 operational at Payerne. The saturation vapour pressure $e_w(z)$ is computed using the empirical formulation:

$$e_w(z) = 6.107 \cdot \exp \left[\frac{M_A [T(z) - 273]}{M_B + [T(z) - 273]} \right] \quad (7)$$

where the constants M_A and M_B are 17.08 and 234.2, for temperature above 273 K (List, 1951). RALMO relative humidity
230 profiles have been validated against collocated RS41 radiosondes (Navas-Guzmán et al., 2019; Martucci et al., 2021), demonstrating strong performance. These capabilities make RALMO one of the few instruments worldwide capable of providing continuous daytime and nighttime profiles of water vapour mixing ratio, temperature, and relative humidity.

3.3 Aerosol hygroscopicity with tandem nephelometers

Aerosol hygroscopicity refers to the equilibrium water content of water-soluble particulate matter at RH below 100 % and,
235 thereby altering particle size, shape and optical properties. Particularly, the impact of hygroscopic growth on aerosol optical properties is commonly studied using the enhancement factor, $f_\zeta(RH)$, defined by Hänel (1976):

$$f_\zeta(RH) = \frac{\zeta(RH)}{\zeta(RH_{ref})} \quad (8)$$

where $\zeta(\lambda, RH)$ denotes an optical or microphysical aerosol property at RH, while the reference value, RH_{ref} , corresponds to the lowest value within the analysed layer or time interval.

240 In the case of laboratory measurements, when RH can be controlled, the scattering enhancement factor can be determined using two instruments that measure the same aerosol optical property, such as the aerosol scattering coefficient. One instrument operates at RH below 40 % (dry conditions), whereas the second is equipped with a humidification system that enables



measurements over an RH range from 40 to 90 % (wet conditions). This configuration, known as a tandem nephelometer (Titos et al., 2014a), allows the scattering enhancement factor to be determined using the following two-parameter equation (Clarke et al., 2002; Carrico et al., 2003):

$$f_{\sigma}(RH) = a(1 - RH)^{-\gamma_{\sigma}} \quad (9)$$

where a is the value at $RH = 0$ % and γ_{σ} is the Hänel parameter characterising the magnitude of hygroscopic growth for the scattering enhancement. In the case of laboratory measurements, different RH ranges can be controlled, allowing measurements over a wide RH range from 40 to 90 %. This enables a more robust determination of a through linear fitting. Moreover, the use of the same measurement approach at different sites allows a direct intercomparison of the scattering enhancement factor $f_{\sigma}(\lambda, 85/ < 40$ %). Hereafter, $f_{\sigma}(\lambda)$ for simplicity.

3.4 Aerosol hygroscopicity with lidar measurements

For lidar measurements, the aerosol optical parameter used to study aerosol hygroscopicity is the aerosol backscatter coefficient (β^{aer}) and the backscatter enhancement factor is described as Veselovskii et al. (2009):

$$f_{\beta}(RH) = \frac{\beta^{aer}(RH)}{\beta^{aer}(RH_{ref})} = \left(\frac{1 - RH}{1 - RH_{ref}} \right)^{-\gamma_{\beta}} \quad (10)$$

In the particular case of the RALMO system, the Hänel parameter (γ_{β}) can only be obtained at 355 nm. Since ambient RH cannot be controlled, it must be ensured that changes in aerosol optical properties are solely related to changes in RH. This is achieved by assuming a well-mixed atmospheric layer that typically occurs when water vapour mixing ratio and potential temperature profiles remain nearly constant with altitude (Feingold and Morley, 2003). When these conditions are fulfilled, RH_{ref} is selected as the point with the lowest RH, typically corresponding to the lowest altitude, as RH and β^{aer} increase simultaneously with altitude (Veselovskii et al., 2009). Consequently, humidograms ($f_{\zeta}(RH)$ versus RH) can be computed, and the Hänel parameter is obtained as the slope of the linear fit between the natural logarithms of β^{aer} and RH. Note that, for comparison with tandem nephelometers and previous studies, the backscatter enhancement factor is extrapolated or interpolated to the 40-85 % range ($f_{\beta}(\lambda, 85/40$ %)), depending on the measured RH range. In the following, it is denoted as $f_{\beta}(\lambda)$.

It should be highlighted that the Hänel parameter may depend on the aerosol optical property for its computation. Consequently, γ values derived from aerosol backscatter measurements are not directly comparable to those obtained from total scattering measurements from tandem nephelometers. Moreover, the spectral dependence of hygroscopic growth can vary with particle size, as shown by Mie simulations in previous studies (Zieger et al., 2013; Navas-Guzmán et al., 2019). Therefore, the enhancement factor for the same RH may differ between wavelengths and particle sizes, which further complicates comparisons between the different techniques. In addition, uncertainties associated with the retrieval of aerosol hygroscopicity parameters should also be considered. In the case of lidar-derived parameters, these uncertainties can reach up to 20 % for RH uncertainties below 10 % (Pérez-Ramírez et al., 2021).



4 Results and discussion

4.1 Overview of the HYGRO19 field campaign

Continuous measurements of AOD and extinction AE (AOD-AE) obtained from the AERONET sun photometer during the HYGRO19 campaign in Payerne were analysed to characterise the columnar aerosol properties (Fig. 1). Two clearly different periods were identified: the first, from June to August, characterised by relatively high aerosol loading, and the second, from September to December, dominated by low-pressure systems, persistent cloudiness, and lower aerosol concentrations (MeteoSwiss, 2026b). During the first period, a large number of measurements were available, with a median AOD (500 nm) of 0.24 ± 0.09 , where the uncertainty is expressed as half of the interquartile range ($IQR/2 = (Q3 - Q1)/2$). Fine-mode particles predominated, as indicated by the median AOD-AE (440–870 nm) value of 1.50 ± 0.19 . Backward trajectory analyses using the HYSPLIT model (Stein et al., 2015) (not shown) revealed that most air masses originated over mainland Europe, a major source of anthropogenic aerosols. However, on 26 and 27 June, an event influenced by North African air masses was observed, with high aerosol load (AOD (500 nm) up to 0.3) and low AOD-AE (reaching 0.4), typical of long-range transport of Saharan dust. In contrast, fewer measurements were available during the second period. The median AOD (500 nm) and AOD-AE (440–870 nm) values were 0.07 ± 0.02 and 1.15 ± 0.13 , respectively, characteristic of continental background aerosol.




Figure 1. Hourly time series of AERONET observations during the HYGRO19 campaign from June–December 2019: (a) aerosol optical depth (AOD) at 380, 500, and 1020 nm, and (b) extinction Ångström exponent (AOD-AE (440–870 nm)).

Figure 2 shows the temporal evolution of in situ ground-based measurements of aerosol scattering (a) and absorption (c) coefficients at different wavelengths, scattering Ångström exponent (SAE, b), absorption Ångström exponent (AAE, d), single scattering albedo (SSA, e), particulate matter (f), and chemical composition (g-h). Data are hourly values, except for TC concentrations and chemical compositions that correspond to 2-hourly and daily measurements, respectively (panels g and h). Unfortunately, during the HYGRO19 campaign, data from the Aurora dry nephelometer and the Aethalometer AE31 were available only since 6–8 August (a–e, g), whereas FATCAT data date since 27 July 2019 (g). The main statistical parameter for the entire campaign, as well as for the first and second periods separately, are summarised in Table 1.




Figure 2. Time series of in situ ground-based measurements of (a) aerosol scattering coefficient at 450, 525, and 635 nm and (b) scattering Ångström exponent (SAE (450–635 nm)) from the Aurora 3000 dry nephelometer; (c) absorption coefficient at 370, 520, and 880 nm and (d) absorption Ångström exponent (AAE (370–950 nm)) from the aethalometer AE31; (e) single scattering albedo (SSA (520 nm)) derived from the dry nephelometer and aethalometer data, (f) PM concentrations, (g) equivalent black carbon at 880 nm from the aethalometer and total carbon concentrations from FATCAT; and (h) chemical composition. The data represent hourly mean values, except for TC concentrations and chemical compositions that are 2-hourly and daily measurements.



During the campaign, the temporal evolution of the scattering coefficient (Fig. 2a) shows remarkable day-to-day variability, with values ranging from 0.97 to 206 Mm^{-1} , and median values of $28 \pm 21 \text{ Mm}^{-1}$ and $19 \pm 14 \text{ Mm}^{-1}$ at 450 and 550 nm, respectively. Several intense events with higher scattering coefficients values were observed between 21 August and 21 September, 17 October and 1 November, 9 and 26 November 2019, and 30 November and 7 December, coinciding with increases in aerosol loading (Fig. 1) and particulate matter (Fig. 2f). Concurrently, the analysis of the SAE (450–635 nm, Fig. 2b) suggests a predominance of fine particles (SAE = 1.64 ± 0.26), although some outliers are observed, in agreement with AOD-AE AERONET measurements (Fig. 1b). This predominance of fine particles is supported by the small differences observed between PM_{10} and $\text{PM}_{2.5}$ ($\text{PM}_{10-2.5} = 3.0 \pm 1.6$, Fig. 2f). The low PM concentrations ($\text{PM}_{10} = 13.2 \mu\text{g}/\text{m}^3$ and $\text{PM}_{2.5} = 9.2 \mu\text{g}/\text{m}^3$, (Hüglin and Grange, 2021)), characterised by a predominance of fine-mode ($\text{PM}_{2.5}$) over coarse-mode ($\text{PM}_{10-2.5}$) mass, are typical of a rural site such as Payerne, which is usually influenced by the transport of local and Swiss Plateau emissions, as well as anthropogenic pollution from mainland Europe (Minguillón et al., 2012; Dada et al., 2025).

Table 1. Statistical summary of the aerosol scattering and absorption coefficients at 450 and 520 nm, the scattering Ångström exponent (SAE (450–635 nm)) and absorption Ångström exponent coefficient (AAE (370–950 nm)), single scattering albedo (SSA (520 nm)), PM, equivalent and total black carbon concentrations, as well as chemical composition. Note that ammonium was derived assuming complete neutralisation and OM was estimated from PM mass closure.

	median $\pm$ IQR/2				median $\pm$ IQR/2		
	Entire	Jun-Aug	Sep-Dec		Entire	Jun-Aug	Sep-Dec
σ_{sp} (450 nm) (Mm^{-1})	28 ± 21	28 ± 17	28 ± 22	EBC (880 nm) ($\mu\text{g}/\text{m}^3$)	0.27 ± 0.15	0.23 ± 0.12	0.28 ± 0.16
SAE (450–635 nm)	1.64 ± 0.26	1.73 ± 0.14	1.61 ± 0.29	TC ($\mu\text{g}/\text{m}^3$)	2.8 ± 1.1	2.69 ± 0.67	2.8 ± 1.2
σ_{ab} (520 nm) (Mm^{-1})	3.4 ± 1.9	2.7 ± 1.3	3.5 ± 2.1	EC ($\mu\text{g}/\text{m}^3$)	0.28 ± 0.13 (4.6 %)	0.26 ± 0.10 (3.8 %)	0.31 ± 0.15 (5.2 %)
AAE (370–950 nm)	1.06 ± 0.10	0.92 ± 0.11	1.09 ± 0.10	SO_4^{2-} ($\mu\text{g}/\text{m}^3$)	0.30 ± 0.13 (4.4 %)	0.38 ± 0.12 (5.3 %)	0.22 ± 0.11 (3.7 %)
SSA (520 nm)	0.87 ± 0.05	0.90 ± 0.02	0.85 ± 0.05	NO_3^- ($\mu\text{g}/\text{m}^3$)	0.11 ± 0.08 (4.1 %)	0.08 ± 0.02 (1.2 %)	0.17 ± 0.22 (6.4 %)
PM_{10} ($\mu\text{g}/\text{m}^3$)	9.3 ± 3.9	10.1 ± 3.5	7.9 ± 4.2	NH_4^+ ($\mu\text{g}/\text{m}^3$)	0.17 ± 0.07 (2.8 %)	0.17 ± 0.04 (2.3 %)	0.16 ± 0.10 (3.2 %)
$\text{PM}_{2.5}$ ($\mu\text{g}/\text{m}^3$)	5.7 ± 3.2	6.2 ± 2.7	5.1 ± 3.5	OM ($\mu\text{g}/\text{m}^3$)	4.9 ± 2.5 (83 %)	5.5 ± 2.1 (87 %)	4.3 ± 2.7 (81 %)

The analysis of the aerosol absorption coefficient (Fig. 2c) exhibits values ranging from 0.06 to 26 Mm^{-1} , with a median of $3.4 \pm 1.9 \text{ Mm}^{-1}$ at 520 nm, suggesting relatively low contribution of absorbing aerosols during the campaign. This is consistent with the single scattering albedo values (SSA (520 nm) = 0.87 ± 0.05 , Fig. 2e). However, differences were observed between the two periods, with larger presence of absorbing particles during the second ($\sigma_{ab} = 3.5 \pm 2.1$, SSA = 0.85 ± 0.05 and AAE = 1.09 ± 0.10) than in the first period ($\sigma_{ab} = 2.7 \pm 1.3$, SSA = 0.90 ± 0.02 and AAE = 0.92 ± 0.11). Correlative analyses of carbonaceous measurements suggest that these changes can be associated with a higher frequency of local wood burning (Fig. 2g–h). Regarding chemical composition (Fig. 2h), inorganic species (sulphate, ammonium, and nitrate) show an important contribution to particulate matter, accounting for 11 %. However, the TC values compared with measured inorganic species and $\text{PM}_{2.5}$ support that organic matter is the main fraction (83 %), which is typically the largest fraction in summer (OM = 87 %), compared with the second part of the campaign (OM = 81 %), as has also been reported in the literature for the Payerne station (Minguillón et al., 2012; Hüglin and Grange, 2021; Nowak et al., 2023–2025; Dada et al., 2025).

Figure 3 shows the temporal evolution of vertical profiles of the aerosol backscatter coefficient at 355 nm, temperature, water vapour mixing ratio, relative humidity, and the backscatter enhancement factor ($f_{\beta}(355 \text{ nm})$) obtained with the RALMO system during the HYGRO19 campaign. For simplicity, the data are shown only in the lower troposphere (0–6 km agl). The temporal and spatial resolutions were 30 min and 30 m, respectively. Note that the white regions in the figure correspond to areas where retrievals were not possible due to cloudy conditions.




Figure 3. Temporal and vertical evolution of RALMO measurements during the HYGRO19 field campaign at the MeteoSwiss station in Payerne from June to December 2019. (a) Aerosol backscatter coefficient at 355 nm, (b) temperature, (c) water vapour mixing ratio, (d) relative humidity, and (e) backscatter enhancement factor at 355 nm and 85 % RH, normalised to 40 % RH.

The key result from Fig. 3 is the capability of RALMO to provide continuous daytime and nighttime observations with high temporal and vertical resolution, enabling continuous monitoring of atmospheric vertical profiles and aerosol hygroscopic



parameters (panel e). Temperature and water vapour mixing ratio (Fig. 3b-c) show a similar temporal evolution over the campaign period, with higher values in summer and lower values towards winter, and generally decreasing with altitude. Both parameters exhibited higher values during late June to early August, reaching maxima of 305.6 K and 15.2 g/kg below 1.5 km agl, respectively. Later, both temperature and water vapour mixing ratio decrease, reaching their minimum values in December. These behaviours correspond to the typical seasonal evolution of atmospheric conditions in Payerne (Bader et al., 2020; Hicks-Jalali et al., 2020). The median and half of the interquartile range of these parameters for the entire period, as well as for the aforementioned sub-periods, are summarised in Table 2. The values are separated below 1.5 km agl and above that, because aerosol, temperature, and water vapour profiles (Fig. 3a-c) reveal the highest values within the first 1.5 km agl corresponding approximately to the mean atmospheric boundary layer (ABL) height.

The analysis of $\beta^{aer}(355 \text{ nm})$ (Fig. 3a) reveals again two very distinct periods. The first corresponds to June–August, when the highest values are observed, while the second period is from September–December and exhibits considerably lower values. This is consistent with the analysis of AERONET measurements (Fig. 1a). During the first period, aerosol is frequently present throughout the entire ABL and even within the free troposphere (FT). Some decoupled aerosol layers are also observed in this period (e.g., July 2–5 and July 17–19). In the second period, however, almost no decoupled aerosol layers are detected, very low $\beta^{aer}(355 \text{ nm})$ values are measured (Table 2), and aerosol are typically confined below 1.0 km agl, consistent with a lower ABL height (Collaud Coen et al., 2014).

Table 2. Statistical analysis of the aerosol backscatter coefficient, temperature, water vapour mixing ratio, relative humidity retrieved with RALMO during the HYGRO19 campaign at different height ranges.

	0.0–1.5 km agl			1.5–3.5 km agl		
	median $\pm$ IQR/2			median $\pm$ IQR/2		
	Entire	Jun-Aug	Sep-Dec	Entire	Jun-Aug	Sep-Dec
$\beta^{aer}(355 \text{ nm}) (\text{Mm}^{-1}\text{sr}^{-1})$	1.50 ± 0.48	1.63 ± 0.43	1.09 ± 0.44	0.57 ± 0.34	1.35 ± 0.39	0.51 ± 0.07
Temperature (K)	285.1 ± 4.9	288.4 ± 3.3	279.9 ± 4.9	277.6 ± 4.1	279.6 ± 3.1	273.7 ± 4.9
Water vapour mixing ratio (g/kg)	6.6 ± 1.7	7.6 ± 1.2	4.7 ± 1.5	4.5 ± 1.3	5.4 ± 1.0	3.04 ± 0.75
RH (%)	61 ± 10	59 ± 10	63 ± 9.0	62 ± 12	66 ± 11	53 ± 12

4.2 Aerosol hygroscopicity analyses using remote sensing techniques

The backscatter enhancement factor retrieved with RALMO ($f_{\beta}(355 \text{ nm})$) during the campaign was computed within the layers fulfilling the criteria described in Sect. 3.4, considering only cases with no significant changes in air-mass origin with altitude (Navas-Guzmán et al., 2019). Note that for comparison with parameters derived with others instruments, $f_{\beta}(355 \text{ nm})$ was interpolated or extrapolated to the 40–85 % RH range (Fig. 3e). A total of 227 hygroscopic aerosol growth cases were recorded, with $f_{\beta}(355 \text{ nm})$ values ranging from 1.2 to 7.2 and a median of 2.64 ± 0.64 , corresponding to $\gamma_{\beta}(355 \text{ nm})$ of 0.70 ± 0.18 . The highest frequency of occurrence was observed in summer (177 hygroscopic layers, 78 %), whereas only a few cases were detected during October–December, not permitting a robust statistical comparison between the two periods.



Frequency histograms of the $f_{\beta}(355 \text{ nm})$ and γ_{β} are shown in Fig. 4. To study differences with altitude, two layers were distinguished: one below 1.5 km agl (orange line in Fig. 4), typically below the ABL, where most aerosol particles are found, and the other one above 1.5 km agl (red line), generally associated with the FT at the Payerne station. To evaluate whether the differences with altitude are statistically significant, the non-parametric Kruskal-Wallis and Mann-Whitney U tests were applied. The Kruskal-Wallis test determines whether three or more independent groups differ significantly based on ranked data, while the Mann-Whitney U test assesses differences between two independent groups based on ranked observations using the sum of ranks (Chicco et al., 2025).

For the backscatter enhancement factor (Fig. 4a), no significant differences were observed between all data and the different layer base heights (Kruskal-Wallis p-value = 0.99 and Mann-Whitney U p-values ≈ 1.0 ; at the 95 % confidence level), with the frequency distribution skewed towards values lower than 4.0. The Hänel parameter (Fig. 4b), however, follows an approximately normal frequency distribution, and the differences between layers are small and not statistically significant (Kruskal-Wallis p-value = 0.98). In general, the retrieved values indicated significant aerosol hygroscopicity, with 64 % of the cases exhibiting $f_{\beta}(355 \text{ nm})$ values between 2.0 and 4.0, and $\gamma_{\beta}(355 \text{ nm})$ values in the range 0.50-0.99. The most frequent hygroscopic aerosol layers were observed below 1.5 km agl, with 162 layers (71.4 %) and retrieved hygroscopicity parameters of $f_{\beta}(355 \text{ nm}) = 2.64 \pm 0.59$ and $\gamma_{\beta}(355 \text{ nm}) = 0.70 \pm 0.16$. Layers above 1.5 km agl were less frequent but exhibited similar hygroscopicity ($f_{\beta}(355 \text{ nm}) = 2.72 \pm 0.71$ and $\gamma_{\beta}(355 \text{ nm}) = 0.72 \pm 0.20$).




Figure 4. Frequency of occurrence of (a) $f_{\beta}(355 \text{ nm}, 85/40 \%)$ and (b) $\gamma_{\beta}(355 \text{ nm})$ obtained from RALMO measurements during the HYGRO19 field campaign from June to December 2019. The analysis includes all detected aerosol hygroscopicity layers, as well as cases recorded below and above 1.5 km agl. N represents the number of cases.

The large database of aerosol hygroscopic cases during the HYGRO19 campaign allows for an investigation of $f_{\beta}(355 \text{ nm})$ and $\gamma_{\beta}(355 \text{ nm})$ with height. In this context, Fig. 5 shows box-whisker plots of these parameters for different height ranges, which were determined by optimising the number of bins in the vertical histograms using established statistical rules for

histogram bin selection (Doane's rule (Doane, 1976), Rice's rule (Rubia, 2024), and Scott's normal reference rule (Scott, 2010)). The analyses are also presented separately for the convective boundary layer (CBL), the residual layer (RL), and the FT, using the ABL operational products at Payenne station developed by Collaud Coen et al. (2014). Additionally, the statistical parameters and the frequency of occurrence (rel. freq.) derived are summarised in Table 3.




Figure 5. Box-whisker plots of (a) $f_{\beta}(355 \text{ nm}, 85/40 \%)$ and (c) $\gamma_{\beta}(355 \text{ nm})$ by different height ranges. The analyses are also presented for the CBL, RL, and FT (b, d). In each box plot, the central mark represents the median, the red point indicates the mean, and the box extends vertically from the 25th to the 75th percentiles. N represents the number of cases.

Results from Fig. 5a,c and Table 3 show the vertical variability of aerosol hygroscopicity parameters. In general, some vertical variability is observed, with the highest hygroscopicity parameters found between 606-912 m agl ($f_{\beta}(355 \text{ nm}) = 3.22 \pm 0.75$ and $\gamma_{\beta}(355 \text{ nm}) = 0.84 \pm 0.18$), and between 1830 and 2136 m agl ($f_{\beta}(355 \text{ nm}) = 3.17 \pm 0.77$ and $\gamma_{\beta}(355 \text{ nm}) = 0.83 \pm 0.19$). Some height ranges show appreciable differences (e.g., between 606-912 m agl, and between 1218-1524 m agl), for which the Mann-Whitney U test yielded a p-value of 0.01. However, the low number of data, particularly for the highest layers, and the range of variability of median values are below 20 %, which is within the uncertainties of the methodology.



The most important point to remark is the large variability of the hygroscopicity parameters, as reflected by the large standard deviations and percentile ranges. Grouping $f_{\beta}(355 \text{ nm})$ and $\gamma_{\beta}(355 \text{ nm})$ into the CBL, RL and FT (Fig. 5b,d) increases the number of data and makes the statistics more robust. Nevertheless, the analysis did not reveal any significant differences, as the mean, median and percentile values are very similar. Particularly, slightly higher hygroscopicity was observed in the FT ($f_{\beta}(355 \text{ nm}) = 2.75 \pm 0.59$ and $\gamma_{\beta}(355 \text{ nm}) = 0.73 \pm 0.16$), lower values were obtained in the CBL ($f_{\beta}(355 \text{ nm}) = 2.56 \pm 0.60$ and $\gamma_{\beta}(355 \text{ nm}) = 0.68 \pm 0.17$) and the highest frequency of occurrence was found in the RL. Overall, these results suggest no clear vertical dependence of aerosol hygroscopicity, and the observed height-dependent patterns should be interpreted with caution, since apparent differences may be amplified by the limited number of observations per layer or by inherent methodological limitations. For instance, the different RH ranges used to compute $f_{\beta}(355 \text{ nm})$ and $\gamma_{\beta}(355 \text{ nm})$ in the humidograms (not all experimental cases cover the same minimum-to-maximum RH range), along with uncertainties in RH measurements, can affect the retrieval (Pérez-Ramírez et al., 2021).

Table 3. Statistical analysis of hygroscopic aerosol layers retrieved with RALMO during the HYGRO19 campaign at different height ranges.

Height range (m agl)	$f_{\beta}(355 \text{ nm}, 85/40 \text{ \%})$	$\gamma_{\beta}(355 \text{ nm})$	N. (rel. freq. (%))
	median $\pm$ IQR/2	median $\pm$ IQR/2	
300-606	2.62 ± 0.58	0.70 ± 0.15	37 (16.3)
606-912	3.22 ± 0.75	0.84 ± 0.18	37 (16.3)
912-1218	2.77 ± 0.40	0.74 ± 0.10	35 (15.4)
1218-1524	2.30 ± 0.44	0.60 ± 0.14	53 (23.4)
1524-1830	3.12 ± 0.52	0.82 ± 0.17	16 (7.1)
1830-2136	3.17 ± 0.77	0.83 ± 0.19	17 (7.5)
2136-2442	2.34 ± 0.37	0.61 ± 0.12	13 (5.7)
2442-2748	1.72 ± 0.02	0.39 ± 0.01	4 (1.8)
2748-3054	2.80 ± 0.65	0.74 ± 0.16	8 (3.5)
3054-3360	2.48 ± 0.42	0.65 ± 0.12	7 (3.1)

To further investigate temporal changes in aerosol hygroscopicity, Fig. 6 shows the monthly evolutions of $f_{\beta}(355 \text{ nm})$ and $\gamma_{\beta}(355 \text{ nm})$ during the HYGRO19 campaign. Results are presented as box-whisker plots, with the most important statistical parameters summarised in Table 4, which also includes surface chemical composition (Sect. 2.5). Note that chemical composition data are available at daily resolution and correspond to surface measurements, whereas hygroscopicity parameters are derived from RALMO measurements at higher temporal resolution and at different altitudes. Therefore, a direct relationship between chemistry and individual hygroscopicity retrievals may not be representative, and the chemistry data are only used to provide a general overview of aerosol composition during the campaign. The largest number of cases for studying aerosol hygroscopicity was in July (104 layers, 45.8 %), followed by August (55 layers, 24.3 %) and September (40 layers, 17.6 %), whereas fewer cases were recorded from October to December (Table 4). The most hygroscopic layers occurred in June ($f_{\beta}(355 \text{ nm}) = 3.65 \pm 0.70$ and $\gamma_{\beta}(355 \text{ nm}) = 0.94 \pm 0.15$), while lower hygroscopicity was observed in October and December ($f_{\beta}(355 \text{ nm}) = 2.36 \pm 0.26$ and $\gamma_{\beta}(355 \text{ nm}) = 0.60 \pm 0.08$), as well as in July ($f_{\beta}(355 \text{ nm}) = 2.47 \pm 0.49$ and $\gamma_{\beta}(355 \text{ nm}) = 0.65 \pm 0.15$). Consistent with these differences, the Mann-Whitney U test revealed statistically significant

differences between June and July (p-value = 0.001). High values were also recorded in September, which likewise showed statistically significant differences compared with July. High hygroscopicity was also observed in November, although only three cases were identified (Fig. 6 and Table 4).




Figure 6. Box-whisker plots of (a) $f_{\beta}(355 \text{ nm}, 85/40 \%)$ and (b) $\gamma_{\beta}(355 \text{ nm})$ from RALMO measurements for every month during the HYGRO19 field campaign. In each box plot, the central mark represents the median, the red point indicates the mean, and the box extends vertically from the 25th to the 75th percentiles. N represents the number of cases.

Table 4. Statistical analysis of aerosol hygroscopicity parameters retrieved from RALMO measurements and surface chemical composition obtained during the HYGRO19 campaign for hygroscopic aerosol cases detected during each month.

Month	Aerosol hygroscopicity parameters			Chemical composition				
	$f_{\beta}(355 \text{ nm})$	$\gamma_{\beta}(355 \text{ nm})$	N (rel. freq. (%))	median $\pm$ IQR/2				
	median $\pm$ IQR/2	median $\pm$ IQR/2		EC ($\mu\text{g}/\text{m}^3$) (%)	SO_4^{2-} ($\mu\text{g}/\text{m}^3$) (%)	NO_3^- ($\mu\text{g}/\text{m}^3$) (%)	NH_4^+ ($\mu\text{g}/\text{m}^3$) (%)	OM ($\mu\text{g}/\text{m}^3$) (%)
June	3.65 ± 0.70	0.94 ± 0.15	18 (7.9)	0.26 ± 0.04 (4.6)	0.28 ± 0.06 (5.0)	0.06 ± 0.01 (1.1)	0.12 ± 0.02 (2.2)	5.24 ± 0.60 (87)
July	2.47 ± 0.49	0.65 ± 0.15	104 (45.8)	0.26 ± 0.06 (3.5)	0.47 ± 0.07 (6.6)	0.09 ± 0.03 (1.2)	0.19 ± 0.03 (2.8)	6.7 ± 1.2 (86)
August	2.64 ± 0.66	0.70 ± 0.18	55 (24.3)	0.26 ± 0.11 (4.2)	0.43 ± 0.10 (6.3)	0.07 ± 0.02 (1.1)	0.18 ± 0.04 (2.7)	4.9 ± 2.5 (83)
September	3.15 ± 0.99	0.83 ± 0.22	40 (17.6)	0.26 ± 0.10 (5.0)	0.31 ± 0.08 (5.5)	0.15 ± 0.04 (3.1)	0.17 ± 0.03 (2.9)	4.6 ± 1.4 (83)
October	2.36 ± 0.26	0.60 ± 0.08	2 (0.90)	0.36 ± 0.02 (6.0)	0.30 ± 0.02 (4.9)	0.16 ± 0.02 (2.7)	0.16 ± 0.02 (2.6)	5.00 ± 0.78 (84)
November	2.94 ± 0.66	0.78 ± 0.16	3 (1.3)	0.40 ± 0.05 (5.7)	0.15 ± 0.01 (1.6)	0.40 ± 0.11 (4.3)	0.17 ± 0.04 (1.8)	4.7 ± 1.0 (87)
December	2.30 ± 0.18	0.60 ± 0.06	5 (2.2)	0.19 ± 0.13 (4.7)	0.15 ± 0.06 (3.1)	0.21 ± 0.21 (7.1)	0.12 ± 0.08 (3.3)	2.8 ± 1.9 (82)

The high aerosol hygroscopicity observed during the campaign might be associated with aerosols mainly composed by inorganic species (11 %), particularly sulphate ($0.30 \pm 0.13 \mu\text{g}/\text{m}^3$ (4.4 %)), nitrate ($0.11 \pm 0.08 \mu\text{g}/\text{m}^3$ (4.1 %)) and ammonium ($0.17 \pm 0.07 \mu\text{g}/\text{m}^3$ (2.8 %)), along with low concentrations of elemental carbon ($0.28 \pm 0.13 \mu\text{g}/\text{m}^3$ (4.6 %)), with organic matter as the major fraction ($4.9 \pm 2.5 \mu\text{g}/\text{m}^3$ (83 %); Table 1). This chemical composition is consistent with previous observations in Payerne (Minguillón et al., 2012; Hüglin and Grange, 2021; Dada et al., 2025), and was more pronounced during



summer, particularly in June, when higher and lower concentrations of inorganic components and carbonaceous species, respectively, (Tables 1 and 4), and can be attributed to enhanced photochemical activity during summer (Kammermann et al., 2010). In contrast, during autumn and winter, black carbon concentrations increase due to greater residential wood burning and a shallower boundary layer in the region (Minguillón et al., 2012; Dada et al., 2025), and these particles are typically less hygroscopic (Deshmukh et al., 2025; Lorenzo et al., 2025). The predominance of scattering particles, fine particles, and lower concentrations of carbonaceous species during summer can also explain the higher hygroscopicity observed during this period compared with the second part of the campaign (Figure 2 and Table 1).

Changes in the light-scattering and backscatter due to hygroscopic water uptake at a given RH also depend on particle size. This effect was assessed using the extinction AE from sun photometer measurements. The relationship between particle size and the enhancement factor shows no clear dependence on particle size, with differences not being statistically significant. Fine particles yielded $f_{\beta}(355 \text{ nm}) = 2.67 \pm 0.22$ and $\gamma_{\beta}(355 \text{ nm}) = 0.70 \pm 0.13$, compared to $f_{\beta}(355 \text{ nm}) = 2.30 \pm 0.45$ and $\gamma_{\beta}(355 \text{ nm}) = 0.60 \pm 0.22$ for larger particles. Note that the corresponding figure is not included here. A similarly weak size dependence has been reported in previous studies (Kammermann et al., 2010; Zieger et al., 2013; Titos et al., 2014b; Jefferson et al., 2017; Sicard et al., 2022).

Wind speed and wind direction allow investigating the relationship between air-mass origin and aerosol hygroscopic behaviour. The profile data were obtained using a Doppler wind lidar. Figure 7 illustrates the temporal evolution of wind direction for the detected hygroscopic aerosol layers (a), monthly wind-rose diagrams as a function of wind speed (b) and the corresponding $f_{\beta}(355 \text{ nm})$ (c). Data from Fig. 7 show that during summer more wind profiles could be monitored due to lower cloud cover and bird migration, allowing the observation of distinct vertical wind structures. Near the surface, Northeast (NE) winds were frequently observed, while above that winds predominantly originated from the Southwest (SW) and West (W). Wind speeds were around 5 to 8 m/s (inset in Fig. 7a), with a median value of $4.2 \pm 1.9 \text{ m/s}$, corresponding to a moderate breeze (WMO, 2017). These conditions are typical of high-pressure systems, which occur more frequently in summer over central Europe (MeteoSwiss, 2026a). In contrast, during autumn and winter months, layers were mostly monitored below 1.5 km agl, predominantly due to instrumental limitations under cloudy conditions associated with more frequent presence of low-pressure systems (MeteoSwiss, 2026b). Southwestern and Western air masses, originating in the Atlantic, travel through France before reaching Payerne, suggesting a possible mixture of particles (backward trajectories computed with HYSPLIT are not shown for clarity). Similar results were found by Kammermann et al. (2010) at the Jungfraujoch alpine site (Switzerland), where higher hygroscopicity was observed for winds from the SW or W and lower for other wind directions. Layers with $f_{\beta}(355 \text{ nm}) > 3$ in July–September also showed SW–W winds above 5 m/s. Lower hygroscopicity in late June and July can be associated to the presence of dust particles, due to the influence of air masses from Southern Europe and North Africa, consistent with other studies in Switzerland (Fierz-Schmidhauser et al., 2010; Navas-Guzmán et al., 2019; Eirund et al., 2022; Collaud Coen et al., 2026). Northeast winds are observed mostly at surface level, with wind speeds of 3–6 m/s, suggesting the influence of Central European urban and industrial emissions, containing less hygroscopic black carbon and organic matter.




Figure 7. Temporal and vertical evolution of wind direction (a) for the hygroscopic aerosol layers from wind profiler measurements during the HYGRO19 field campaign (June–December 2019). The inset in panel (a) shows the wind rose. Monthly wind rose plots depicting wind speed (b) and the $f_{\beta}(355 \text{ nm}, 85/40 \%)$ from RALMO measurements (c).

Table 5 summarises some of the most relevant studies on aerosol hygroscopicity based on aerosol optical properties. When reported, aerosol chemical composition is provided; otherwise, site classification (e.g., urban, rural, marine) is used as a proxy for aerosol type. These studies are also classified according to whether they are based on remote sensing (lidar and ceilometer) or in situ (tandem nephelometers) techniques. Note that RALMO hygroscopicity parameters are provided at 355 nm for the 40–85 % RH range, whereas for some other studies, values are provided at different wavelengths and RH ranges due to differences in system configurations and experimental conditions. The main advance provided by the HYGRO19 campaign is the investigation of the temporal and vertical variability of the hygroscopicity parameters. Long-term analyses using remote sensing techniques have been reported in only a few studies (e.g., Bedoya-Velázquez et al., 2019; Sicard et al., 2022; Jing et al., 2026), whereas most studies are limited to a small number of case studies or were conducted during specific field campaigns (e.g., Lv et al., 2017; Fernández et al., 2018; Chen et al., 2019; Navas-Guzmán et al., 2019; Pérez-Ramírez et al., 2021; Laly et al., 2025). On the other hand, for tandem nephelometers (f_{σ} and γ_{σ}), the results of this work are also comparable to those reported in previous studies based on in situ techniques (e.g., Zieger et al., 2014; Titos et al., 2021; Li et al., 2025), with the differences being within the uncertainties associated with each technique.



Table 5. Aerosol hygroscopicity parameters derived from aerosol light-scattering enhancement factors (in situ measurements) and aerosol backscatter enhancement factors (remote sensing observations) from this study and previous studies. Values are expressed as median $\pm$ half of the interquartile range for this study and Zieger et al. (2014), and as mean $\pm$ standard deviation for Jing et al. (2026).

Research type	Location	Instrument	Chemical composition or site type	Hygroscopicity	Reference
Statistics 227 cases	Payerne, Switzerland 46.82° N, 6.95° E	Raman lidar	Low fractions (EC: 4.6 %) mainly (SO_4^{2-} , NH_4^+ , NO_3^- : 11 %)	$f_\beta(355 \text{ nm}, 85/40 \%)$ 2.64 ± 0.64 $\gamma_\beta(355 \text{ nm}): 0.70 \pm 0.18$	This study
Campaign Feb-March	Melpitz, Germany 52.53° N, 12.93° E	Tandem nephelometers	rural	$f_\sigma(550 \text{ nm}, 85/40 \%)$ 2.78 ± 0.23	Zieger et al. (2014)
Case study	Xinzhou, Shanxi 38.39° N, 112.7° E	Raman Lidar and RS	mixed aerosols of anthropogenic origin	$\gamma_\beta(355 \text{ nm}): 0.75$	Lv et al. (2017)
Case study	Madrid, Spain 40.46° N, 3.73° W	Raman Lidar and RS	marine mixed with local aerosol	$\gamma_\beta(532 \text{ nm}): 0.62$	Fernández et al. (2018)
Long-term 8 cases	Paris, France 48.71° N, 2.21° E	Ceilometer Tower	Low fractions (BC: 8 %) higher (SO_4^{2-} , NO_3^- , NH_4^+)	$f_\beta(910 \text{ nm}, 85/ < 40 \%)$ $2.6\text{-}3.3$	Bedoya-Velásquez et al. (2019)
Case study	Baltimore, Washington 38.75-39.75° N 77.75-75.25° W	Remote sensing and In situ	mainly (SO_4^{2-} , 52 %)	$\gamma_\beta(355 \text{ nm})$ 0.65 ± 0.10	Pérez-Ramírez et al. (2021)
Long-term and campaign observations	23 Sites	Tandem nephelometers	urban and rural marine	$f_\sigma(550 \text{ nm}, 85/ < 40 \%)$ 1-3 2-4	Titos et al. (2021)
Campaign Feb-April	Xiamen, China	Tandem nephelometers	urban and rural	$f_\sigma(550 \text{ nm}, 85/ < 40 \%)$ 1.5-2	Li et al. (2025)
Case study	Saclay, France 50.6° N, 3.1° E	Raman Lidar	regional pollution	$\gamma_\beta(355 \text{ nm})$ 0.75 ± 0.05	Laly et al. (2025)
Statistics 192 cases	Wuhan, China 30.5° N, 114.4° E	Polarization lidar	local and regional pollution	$\gamma_\beta(532 \text{ nm})$ 0.62 ± 0.24	Jing et al. (2026)

4.3 Assessment of different techniques for studying aerosol hygroscopic growth

Hygroscopic growth retrievals depend on both the measurement wavelength and the aerosol optical property, while chemical composition may also play an important role (Gysel et al., 2007). Consequently, f_ζ and γ_ζ values derived from aerosol backscatter measurements (e.g. lidar or COBALD) are not directly comparable to those obtained from total scattering measurements using tandem nephelometers. In addition, previous studies based on Mie simulations have shown that the spectral dependence of hygroscopic growth can depend on particle size (Zieger et al., 2013; Navas-Guzmán et al., 2019). Therefore, enhancement factors for the same RH may differ between wavelengths and particle sizes, and values obtained from different techniques are expected to correlate in their variability, whereas their absolute values are not expected to be directly comparable.

4.3.1 RALMO retrievals versus ground-based tandem nephelometers

During the HYGRO19 campaign, tandem nephelometers operated from October to December, providing the enhancement factor at 550 nm ($f_{\sigma}(550 \text{ nm})$) derived from scattering coefficient measurements over a relative humidity range from below 40 % to 85% (Sect. 3.3). Note that the tandem nephelometers measurements are only representative only of near-surface
 465 conditions (approximately 3 m agl). Figure 8 shows the temporal evolution of $f_{\sigma}(550 \text{ nm})$ retrieved with the tandem (cyan curve) and those values obtained with the RALMO system ($f_{\beta}(355 \text{ nm})$), differentiating between cases detected within RL or FT (purple points and diamonds, respectively). Unfortunately no layers were detected in the CBL. Only seven hygroscopic aerosol layers could be identified using the RALMO data during this period, primarily due to the frequent presence of cloud, fog, and precipitation. The aerosol hygroscopicity parameters for coincident measurements between both methodologies are
 470 summarised in Table 6. During the period when tandem nephelometer measurements were available, fine ($\text{SAE} = 1.71 \pm 0.22$) and large particles ($\text{SAE} = 0.88 \pm 0.20$) exhibited very similar hygroscopic enhancement factors, with median $f_{\sigma}(550 \text{ nm})$ values of 1.34 in both cases and IQR/2 values of 0.07 and 0.06, respectively. Similarly, γ_{σ} values showed negligible differences (0.25 ± 0.03 and 0.26 ± 0.03 , respectively).




Figure 8. Temporal evolution of $f_{\sigma}(550 \text{ nm}, 85 / < 40 \%)$ obtained using a tandem nephelometers during the HYGRO19 field campaign. The purple points and diamonds indicate $f_{\beta}(355 \text{ nm}, 85 / 40 \%)$ from RALMO measurements in the RL and FT, respectively. Note that the values for RALMO were normalised to 40 % RH, whereas the tandem measurements correspond to RH values below 40%.

Figure 8 shows an approximately constant value of $f_{\sigma}(550 \text{ nm})$ below 1.5, with a median of 1.34 ± 0.07 . A similar be-
 475 haviour is observed for $\gamma_{\sigma}(550 \text{ nm})$, with a median of 0.26 ± 0.03 (graph not shown for clarity). However, clear outliers were observed on 9 December, coinciding with a marked increase in the aerosol scattering coefficient, $\text{PM}_{2.5}$ concentrations, and scattering AE (values greater than 2.0), along with increases in EBC and TC concentrations (Fig. 2). For this day, HYSPLIT backward trajectories showed that the air masses originated over North America, where many fire hotspots were detected by the NASA/NOAA Suomi National Polar-orbiting Partnership satellite (<https://firms.modaps.eosdis.nasa.gov/>). For the cases
 480 within the RL (12 November and 10 December 2019), good agreement was obtained between both techniques (Table 6). In contrast, the agreement for the FT cases was poorer, which is expected since aerosol layers in the free troposphere are largely decoupled from the boundary layer and typically exhibit different microphysical and optical properties than those near the surface. Nevertheless, the cases detected on 24 October and 13 December showed good agreement with the tandem nephelometers



values, which can be explained by the presence of a well-mixed layer extending from the boundary layer into the lower free troposphere, allowing the hygroscopic layers to be representative of ground-based measurements.

Overall, the comparisons of lidar-derived aerosol hygroscopicity parameters with those from the tandem nephelometers show good agreement for layers below the ABL, as expected because lidar techniques require a well-mixed layer. The slightly lower values obtained from the tandem nephelometers compared with RALMO can be attributed to differences in the sampled aerosol populations, the specific optical quantities measured (scattering coefficient vs. backscatter coefficient) and the different wavelengths used by each instrument. This comparison should be interpreted with caution, since the retrievals are based on different wavelengths and optical properties. A more quantitative comparison would require dedicated Mie simulations to ensure consistent wavelength and backscatter geometry conditions. Particularly, particle-size effects on aerosol hygroscopicity for near-surface measurements were also investigated using the scattering Ångström exponent (SAE (450–635 nm)) from the Aurora 3000 dry nephelometer. The results (not shown for clarity) indicate a weak dependence of the hygroscopic enhancement on particle size, with no statistically significant differences between small and large particles.

Table 6. Values of the enhancement factor and Hänel parameter retrieved using the tandem nephelometers and RALMO instruments. Note that the values for RALMO were normalised to 40 % RH, whereas the tandem measurements correspond to RH values below 40%.

Date	Tandem Nephelometers at 550 nm			RALMO at 355 nm			Layer
	Hour (UTC)	$f_{\sigma}(550 \text{ nm}, 85 / < 40 \%)$	$\gamma_{\sigma}(550 \text{ nm})$	Hour (UTC)	$f_{\beta}(355 \text{ nm}, 85 / 40 \%)$	$\gamma_{\beta}(355 \text{ nm})$	
24 October 2019	17:00	1.52 ± 0.09	0.25 ± 0.03	16:00	1.83 ± 0.07	0.26 ± 0.03	FT
9 November 2019	16:00	1.32 ± 0.07	0.24 ± 0.03	17:00	2.94 ± 0.08	0.88 ± 0.02	FT
12 November 2019	11:00	1.71 ± 0.14	0.40 ± 0.04	23:30	1.92 ± 0.05	0.47 ± 0.02	RL
6 December 2019	15:00	1.30 ± 0.03	0.22 ± 0.01	18:30	2.30 ± 0.06	0.60 ± 0.02	FT
6 December 2019	15:00	1.30 ± 0.03	0.22 ± 0.01	19:00	2.11 ± 0.06	0.54 ± 0.02	FT
10 December 2019	09:00	1.86 ± 0.17	0.48 ± 0.05	08:00	1.89 ± 0.24	0.46 ± 0.09	RL
13 December 2019	18:00	2.60 ± 0.27	0.59 ± 0.06	22:30	2.48 ± 0.10	0.65 ± 0.02	FT

4.3.2 Intercomparisons with COBALD-RS41 sondes I: 17 September 2019

The COBALD-RS41 sondes, which provide a unique opportunity to validate lidar-derived hygroscopicity parameters, were launched sporadically during the HYGRO19 campaign between 23:00 and 23:30 UTC. Comparisons with RALMO are performed using 30 m vertical intervals, resulting in a 6 s temporal resolution for the COBALD sondes. A key difference between remote sensing and balloon-borne techniques is that lidars probe the vertical air column directly above the laser, whereas balloon sondes can suffer of horizontal drifts with altitude due to winds. Consequently, in the presence of wind shear, the two instruments may sample different air masses, with the horizontal separation generally increasing with altitude. The COBALD and lidar systems also differ in receiver FOV (0.01° for RALMO and 6° for COBALD), resulting in systematic differences in backscatter measurements, which are corrected using Mie theory scattering simulations (Brunamonti et al., 2021).



505 The first case study focuses on 17 September 2019, when the region was affected by air masses originating from Northwest-
ern Europe, passing over the Swiss Plateau and arriving at Payerne (backward-trajectories not shown for clarity). These air
masses can transport anthropogenic and biogenic aerosols, especially considering that they travel at very low altitudes within
the ABL. In situ measurements (graphs shown in Fig. A1) indicate that scattering processes were dominated by fine particles
(SAE (450-635 nm) = 1.75 ± 0.17), while absorbing particles also contributed notably (SSA (520 nm) = 0.71 ± 0.08 and
510 AAE (370-950 nm) = 1.22 ± 0.05 , Fig. A1a-d), suggesting the presence of a mixture of anthropogenic and biogenic particles.
These observations are consistent with HYSPLIT backward trajectories, wind profiler measurements (wind speeds between 3
to 6 m/s from the NE) and PM concentrations (Fig. A1e). Consistently, the daily aerosol chemical composition was dominated
by OM ($5.6 \mu\text{g}/\text{m}^3$ (85 %)), followed by EC ($0.39 \mu\text{g}/\text{m}^3$ (5.9 %)), SO_4^{2-} ($0.31 \mu\text{g}/\text{m}^3$ (4.7 %)), NH_4^+ ($0.15 \mu\text{g}/\text{m}^3$ (2.3
%)), and NO_3^- ($0.12 \mu\text{g}/\text{m}^3$ (1.8 %)), suggesting minor contributions from local traffic emissions (Dada et al., 2025). Surface
515 size distribution from SMPS (Fig. A1, right panels) confirmed the dominance of fine-mode particles (diameter lower than
 $2.5 \mu\text{m}$), with a pronounced increase in Aitken-mode particles during the afternoon of 17 September (red curve, Fig. A1h),
along with a slight enhancement in coarse mode particles ($D > 2.5 \mu\text{m}$). In contrast, particle number concentrations on 18
September decreased from the first half of the day (00:00–12:00 UTC) to the second half (12:00–24:00 UTC), particularly
for particles smaller than 100 nm (green curve). The two-hourly mean particle size distributions on 17 September (Fig. A1i),
520 revealed nearly constant particle number concentrations between 12:00 and 14:00 UTC, followed by a progressive increase in
fine particles during the afternoon.

The temporal evolutions of aerosol backscatter coefficient, water vapour mixing ratio, and relative humidity on 17 September
2019 are illustrated in Fig. 9a–c. Ceilometer data were also available, which permits the retrieval of β^{aer} at 1064 nm but with
 β^{aer} derived with the Klett inversion algorithm (Klett, 1981, 1985) at constant lidar ratio of 50 sr. Aerosol particles are observed
525 up to approximately 3.5 km agl until 03:00 UTC, with peak of concentration below 2.5 km agl. Low clouds are also observed
around 3.0–4.0 km agl between 07:00 and 11:00 UTC (Fig. 9a). During nighttime, there is a considerable increase in the
aerosol backscatter coefficient after 21:00 UTC, which became even more pronounced after 23:30 UTC within the 0.48–0.87
km agl layer. For this time interval, the RH values (Fig. 9c) show a strong correlation with β^{aer} . At the same time, the water
vapour mixing ratio (Fig. 9b) exhibited greater homogeneity, suggesting a well-mixed layer. Therefore, these measurements
530 (dashed lines in Fig. 9a–c) fulfil the criterion for aerosol hygroscopicity studies and coincides with COBALD-RS41 sonde
launch. The vertical profiles of aerosol backscatter, water vapour mixing ratio, and RH retrieved from RALMO, the ceilometer,
and the COBALD-RS41 sonde are shown in Fig. 9d–f.

The vertical profiles of β^{aer} between 23:00 and 23:59 on 17 September show very similar patterns for the different in-
struments (RALMO, COBALD and ceilometer). RALMO recording the highest β^{aer} values, while the lowest values were
535 registered by ceilometer. COBALD, however, revealed intermediate values. This pattern is consistent with the spectral depen-
dence of light scattering using Mie theory for spherical particles, which indicates stronger backscattering in the ultraviolet than
in the infrared region. Particularly, within the 0.48–0.87 km agl layer, a marked increase in β^{aer} ($\Delta\beta = 2.3 \text{ Mm}^{-1}\text{sr}^{-1}$, Fig.
9d) was observed in coincidence with an increase in RH from 69 % to 86 % (Fig. 9e). The potential temperature and water

vapour mixing ratio profiles (Fig. 9f) exhibited relatively constant values within this layer, with $\theta = 296.29 \pm 0.05$ K and $r = 7.79 \pm 0.25$ g/kg, suggesting a well-mixed atmosphere.




Figure 9. Temporal evolution of (a) β^{aer} (355 nm), (b) water vapour mixing ratio, and (c) RH profiles from RALMO on 17 September 2019. (d) β^{aer} profiles from RALMO at 355 nm (purple), COBALD at 455 and 940 nm (blue and red), and ceilometer at 1064 nm (black). (e) RH profiles from RALMO (purple) and RS41 (red). (f) Water vapour mixing ratio (purple) and potential temperature (blue) profiles from RALMO. The vertical profiles correspond to 23:30–23:59 UTC for RALMO and ceilometer, and 23:00–23:30 UTC for COBALD-RS41.

Figure 10 shows the humidograms (dependency of $f_{\zeta}(\lambda)$ with RH) for RALMO (355 nm), COBALD-RS41 (450 and 940 nm), and the ceilometer (1064 nm) within the selected 0.48–0.87 km agl layer. Note that for homogeneity all the derived hygroscopicity parameters were normalised to the 40–85% RH range. These parameters obtained using the different techniques for the three case studies are summarised in Table 7. It should be noted that different wavelengths were used in the retrievals.

Overall, there is good agreement between remote sensing (RALMO and ceilometer) and in situ vertical measurements (COBALD-RS41), with the differences being within the uncertainties of each method. In particular, the slight differences can be attributed to balloon drift with altitude, differences in integration time, and the high spatial and temporal variability of aerosol optical properties, which can lead to discrepancies between both techniques (Brunamonti et al., 2021). More specifically, RALMO observed an increase in β^{aer} at 355 nm by a factor of 3.5, while RH increased from 68 % (reference RH) to 86 %, with the retrieved hygroscopicity parameters of $\gamma_{\beta}(355 \text{ nm}) = 1.40 \pm 0.04$ and $f_{\beta}(355 \text{ nm}, 85/40 \%) = 6.98 \pm 0.37$, indicating high aerosol hygroscopicity. Ceilometer retrieval, however, suggested lower wavelength sensitivity with $\gamma_{\beta}(1064 \text{ nm}) = 1.26 \pm 0.05$ and $f_{\beta}(1064 \text{ nm}) = 5.77 \pm 0.37$. Using the in situ vertical technique with COBALD-RS41 sondes, the values obtained were $\gamma_{\beta}(455 \text{ nm}) = 1.27 \pm 0.11$ and $f_{\beta}(455 \text{ nm}) = 5.86 \pm 0.88$, and $\gamma_{\beta}(940 \text{ nm}) = 1.40 \pm 0.11$ and $f_{\beta}(940 \text{ nm}) = 6.93 \pm 1.06$, showing no clear spectral dependence. The apparent differences in the spectral behaviour of γ_{β} are between instruments uncertainties and do not clearly indicate a consistent wavelength dependence.




Figure 10. Enhancement factor ($f_{\beta}(\lambda, RH)$) retrieved from RALMO, COBALD-RS41 and ceilometer within the 0.48–0.87 km agl layer. The time interval corresponds to 23:30–23:59 UTC for RALMO and ceilometer, and 23:00–23:30 UTC for COBALD-RS41. The retrieved parameters are summarised in Table 7.

4.3.3 Intercomparisons with COBALD-RS41 sondes II and aerosol hygroscopicity near clouds: 12 November 2019

The second case study of aerosol hygroscopicity using RALMO and COBALD-RS41 sondes was conducted on 12 November 2019, when air masses originated over the North Atlantic Ocean, passed through France, and reached Payerne. Wind profiler data at the height of this layer indicated relatively low wind speeds of 2.6 m/s from the South (265°), suggesting local and regional aerosol influence. In situ measurements (graphs shown in Fig. A2) during the period 11–13 November 2019 exhibited characteristics that could be attributed to local anthropogenic particles. On 11 November, both the scattering coefficient (a) and SSA (d) reached relatively high values ($\sigma_{sp}(450 \text{ nm}) = 54.0 \pm 5.1 \text{ Mm}^{-1}$ and $\text{SSA}(520 \text{ nm}) = 0.78 \pm 0.06$, respectively), indicating a clear predominance of scattering processes over absorption (Fig. A2b, $\sigma_{ab}(520 \text{ nm}) = 9.7 \pm 1.6 \text{ Mm}^{-1}$). From 12 November onwards, scattering and absorbing coefficients decreased markedly, coinciding with an increase in SAE (>1.5 , Fig. A2c), and a decrease in the albedo ($\text{SSA}(520 \text{ nm}) = 0.59 \pm 0.35$) suggesting a greater contribution of absorbing aerosol and increased anthropogenic influence. Daily aerosol chemical composition exhibited higher EC concentrations ($0.23 \mu\text{g}/\text{m}^3$ (7.0 %)) than NO_3^- , SO_4^{2-} , and NH_4 (5.2, 3.7 and 2.9 %, respectively) and slightly higher values than in the previous case study (EC = 5.9 %). Nevertheless, OM remained the dominant aerosol component ($2.7 \mu\text{g}/\text{m}^3$ (81 %)).

Analysis of the temporal and vertical evolution of RALMO measurements in the lower troposphere (Fig. 11, left panel) shows that the conditions required to investigate aerosol hygroscopicity were fulfilled between 23:30 and 23:59 UTC within the 0.47–0.99 km agl layer. Vertical profiles of aerosol and thermodynamic parameters for this layer (Fig. 11, right panel) show an increase in β^{aer} from 0.15 to $0.30 \text{ Mm}^{-1}\text{sr}^{-1}$ (Fig. 11d) and in RH from 65 % to 90 % (Fig. 11e). Meanwhile, the potential temperature and water vapour mixing ratio profiles (Fig. 11f) remained relatively constant ($\theta = 281.00 \pm 0.14 \text{ K}$ and $r = 2.90 \pm 0.03 \text{ g}/\text{kg}$). High values in backscatter above the hygroscopic aerosol layer were associated with the presence of clouds, suggesting that hygroscopic conditions near the cloud base could facilitate particle activation as cloud condensation nuclei (Pérez-Ramírez et al., 2021).




Figure 11. Temporal evolution of (a) β^{aer} (355 nm), (b) water vapour mixing ratio, and (c) RH profiles from RALMO on 12 November 2019. (d) β^{aer} profiles from RALMO at 355 nm (purple), COBALD at 455 and 940 nm (blue and red), and ceilometer at 1064 nm (black). (e) RH profiles from RALMO (purple) and RS41 (red). (f) Water vapour mixing ratio (purple) and potential temperature (blue) profiles from RALMO. The vertical profiles correspond to 23:30–23:59 UTC for RALMO and ceilometer, and 23:00–23:30 UTC for COBALD-RS41.

The uniqueness of the observations on this day is that a comprehensive assessment of aerosol hygroscopicity can be performed using all available techniques (RALMO, ceilometer, COBALD-RS41 sondes, and tandem nephelometers). RALMO humidogram (purple curve, panel a) reveals that β^{aer} at 355 nm increased by a factor of 2.0 ($f_{\beta}(355 \text{ nm}, 90/65 \%) = 2.0$), corresponding to a Hanel parameter of $\gamma_{\beta}(355 \text{ nm}) = 0.47 \pm 0.02$ and $f_{\beta}(355 \text{ nm}, 85/40 \%) = 1.92 \pm 0.05$, indicating lower aerosol hygroscopicity compared with the case on 17 September ($f_{\beta}(355 \text{ nm}, 85/40 \%) = 6.98 \pm 0.37$). Ceilometer humidogram (black curve, panel a) showed slightly lower hygroscopicity, with $\gamma_{\beta}(1064 \text{ nm}) = 0.43 \pm 0.04$ and $f_{\beta}(1064 \text{ nm}) = 1.83 \pm 0.05$. By contrast, COBALD-RS41 sondes (blue and red curves, panel a) yielded $\gamma_{\beta}(455 \text{ nm}) = 0.46 \pm 0.04$ and $f_{\beta}(455 \text{ nm}) = 1.88 \pm 0.10$ and $\gamma_{\beta}(940 \text{ nm}) = 0.62 \pm 0.05$ and $f_{\beta}(940 \text{ nm}) = 2.35 \pm 0.17$ (Fig. 12a). Despite these slight differences, the results indicate good agreement between the different techniques. On the other hand, hygroscopic values obtained with the tandem nephelometers (Fig. 12b) closest in time to the RALMO retrievals on the same day (12 November 2019 at 11:00 UTC) show consistent values, with $\gamma_{\sigma}(550 \text{ nm}) = 0.40 \pm 0.04$ and $f_{\sigma}(550 \text{ nm}) = 1.71 \pm 0.14$, regardless of differences in sampling time, sampled aerosol, and specific optical properties (Table 7). The analyses of different humidograms with time for this day (Fig. 12b) indicate an increase in aerosol hygroscopicity over time, reaching maximum values at 11:00 UTC (cyan curve), whereas the subsequent available humidogram at 03:00 UTC (pink curve) shows that these processes were decreasing. From these results, it can be concluded that the different techniques are in good agreement within the uncertainties of each method, and may be attributable to differences in wavelength and sampling methods. Overall, no consistent spectral dependence of the retrieved hygroscopic parameters was identified across the different instruments.




Figure 12. Enhancement factor ($f_{\lambda}(\lambda, RH)$) retrieved from (a) RALMO, COBALD-RS41 and ceilometer within the 0.47–0.99 km agl layer, and (b) from tandem nephelometers at different times on 12 November 2019. The parameters are summarised in Table 7. Note that the values in the left panel were normalised to dry conditions at 40 % RH, whereas the tandem measurements correspond to RH values below 40 %.

4.3.4 Aerosol hygroscopicity and clouds: 10 December 2019

595 The last case study corresponds to 10 December 2019. HYSPLIT backward trajectories calculated at 500, 1000, and 1500 m agl (not shown for clarity) were nearly identical within the first 2 km agl, suggesting that the origin and transport pathways of the air masses were independent of altitude. The air masses originated over the North Atlantic Ocean and travelled across France before reaching Payerne; therefore, the influence of a mixture of aerosol types is expected.

In situ measurements from 9 to 11 December 2019 (shown in Fig. A3) revealed a marked increase in EBC concentrations
 600 (f) from 10 December onwards, suggesting large influence of heavily polluted anthropogenic aerosols from local emissions. The concurrent increase in scattering and absorption further indicates enhanced aerosol loading, and the low values of single scattering albedo ($SSA = 0.63 \pm 0.08$, Fig. A3d) suggest large influence of absorbing particles. Consistently, the daily aerosol chemical composition revealed similar concentrations of EC and NO_3^- ($0.26 \mu g/m^3$ (6.3 %) and $0.27 \mu g/m^3$ (6.4 %), respectively), both higher than SO_4^{2-} ($0.17 \mu g/m^3$ (4.1 %)) and NH_4 ($0.14 \mu g/m^3$ (3.4 %)). This indicates a higher contribution of
 605 EC compared with 17 September (6.3 vs. 5.9 %), although lower than on 12 November (7.0 %).

Figure 13a–c shows the temporal evolution of $\beta^{aer}(355 \text{ nm})$, water vapour mixing ratio and RH profiles. The most relevant feature is the presence of clouds before 15:00 UTC as can be seen by the strong backscattering signal (Fig. 13a). Consequently, there are regions where retrievals were not possible. Later, low clouds dissipate and retrievals could be performed. From 08:00 to 08:30 UTC within the layer 0.70–0.97 km agl the requirements to study aerosol hygroscopicity are met (dashed line in
 610 Fig. 13). Aerosol backscatter coefficients (Fig. 13d) showed an increase from 1.8 to $2.1 \text{ Mm}^{-1}\text{sr}^{-1}$ at 355 nm, simultaneous with an increase in RH from 53 % to 66 % (Fig. 13e), while the potential temperature ($\theta = 279.18 \pm 0.29 \text{ K}$) and water vapour mixing ratio ($r = 1.90 \pm 0.01 \text{ g/kg}$) profiles remained relatively constant (Fig. 13f).




Figure 13. Temporal and vertical evolution of (a) β^{aer} (355 nm), (b) water vapour mixing ratio, and (c) RH from RALMO on 10 December 2019. (d) β^{aer} profiles from RALMO at 355 nm (purple), and ceilometer at 1064 nm (black). (e) RH, (f) Water vapour mixing ratio (purple) and potential temperature (blue) profiles from RALMO during the 08:00–08:30 UTC interval.

Figure 14a shows the humidograms obtained with RALMO (purple) and ceilometer (black) measurements. Humidograms retrieved with the tandem nephelometers at different times are also illustrated in Fig. 14b. Results from RALMO showed that β^{aer} at 355 nm increased by a factor of 1.2, corresponding to a Hänel parameter of $\gamma_{\beta}(355 \text{ nm}) = 0.46 \pm 0.09$, and $f_{\beta}(355 \text{ nm}, 85/40 \text{ \%}) = 1.89 \pm 0.24$, indicating lower aerosol hygroscopicity compared to those obtained on 17 September ($f_{\beta}(355 \text{ nm}) = 6.98 \pm 0.37$) and similar to that retrieved on 12 November ($f_{\beta}(355 \text{ nm}) = 1.92 \pm 0.05$). Values calculated using ceilometer measurements showed less agreement, which may be related to the lower signal-to-noise ratio (SNR) due to the cloud presence, yielding $\gamma_{\beta}(1064 \text{ nm}) = 0.19 \pm 0.01$ and $f_{\beta}(1064 \text{ nm}) = 1.30 \pm 0.03$. Data from the tandem nephelometers at 09:00 UTC (cyan curve in Fig. 14b) were of $\gamma_{\sigma}(550 \text{ nm}) = 0.48 \pm 0.05$ and $f_{\sigma}(550 \text{ nm}) = 1.86 \pm 0.17$, in strong agreement with the RALMO values (Table 7).

The good agreement between remote sensing and the tandem nephelometers provides a comprehensive picture of aerosol chemistry in the atmosphere, as under well-mixed conditions, aerosol chemical properties measured at the surface can be extrapolated to higher altitudes. The enhanced EC content combined with lower SO_4^{2-} concentration (4.1 vs 4.7 %), may explain why this case exhibited lower hygroscopicity ($f_{\beta}(355 \text{ nm}) = 1.89 \pm 0.24$) than those on 17 September ($f_{\beta}(355 \text{ nm}) = 6.98 \pm 0.37$). Actually, previous studies have also reported that particles with high EC concentrations tend to exhibit lower hygroscopicity, while those primarily composed of sulphuric acid and ammonia show higher hygroscopicity (Zieger et al., 2014; Jefferson et al., 2017; Bedoya-Velásquez et al., 2019; Chen et al., 2019; Pérez-Ramírez et al., 2021; Cedeño et al., 2023; Deshmukh et al., 2025; Li et al., 2025; Lorenzo et al., 2025). But the key aspect here is the close proximity to clouds, suggesting how aerosol hygroscopicity can facilitate particle activation as cloud condensation nuclei under favourable thermodynamic and dynamical conditions (Pérez-Ramírez et al., 2021).




Figure 14. Enhancement factor ($f_c(\lambda, RH)$) retrieved from (a) RALMO and ceilometer within the 0.70–0.97 km agl layer, and from (b) tandem nephelometers at different times on 10 December 2019. The parameters are summarised in Table 7. Note that the humidograms in the left panel were normalised to dry conditions at 40 % RH, whereas the tandem measurements correspond to RH values below 40 %.

Table 7. Values of the enhancement factor and Hänel parameter retrieved using remote sensing and in situ techniques. Note that the values for RALMO, COBALD-RS41 and ceilometer were normalised to 40 % RH, whereas the tandem measurements correspond to RH values below 40%. Values from different techniques are expected to correlate, whereas their absolute values are not directly comparable.

Case I: 17 September 2019				
Hour (UTC)	Technique	Layer (km agl)	$f_{\zeta}(\lambda, RH)$	$\gamma_{\zeta}(\lambda)$
23:30	RALMO (355 nm)	0.48-0.87	$(f_{\beta}(355 \text{ nm}, 85/40 \text{ \%}) = 6.98 \pm 0.37$	$\gamma_{\beta}(355 \text{ nm}) = 1.40 \pm 0.04$
23:00	COBALD-RS41 (455 nm)		$(f_{\beta}(455 \text{ nm}, 85/40 \text{ \%}) = 5.86 \pm 0.88$	$\gamma_{\beta}(455 \text{ nm}) = 1.27 \pm 0.11$
	COBALD-RS41 (940 nm)		$(f_{\beta}(940 \text{ nm}, 85/40 \text{ \%}) = 6.93 \pm 1.06$	$\gamma_{\beta}(940 \text{ nm}) = 1.40 \pm 0.11$
23:30	Ceilometer (1064 nm)		$(f_{\beta}(1064 \text{ nm}, 85/40 \text{ \%}) = 5.77 \pm 0.37$	$\gamma_{\beta}(1064 \text{ nm}) = 1.26 \pm 0.05$
Case II: 12 November 2019				
Hour (UTC)	Technique	Layer (km agl)	$f_{\zeta}(\lambda, RH)$	$\gamma_{\zeta}(\lambda)$
23:30	RALMO (355 nm)	0.47-0.99	$(f_{\beta}(355 \text{ nm}, 85/40 \text{ \%}) = 1.92 \pm 0.05$	$\gamma_{\beta}(355 \text{ nm}) = 0.47 \pm 0.02$
23:00	COBALD-RS41 (455 nm)		$(f_{\beta}(455 \text{ nm}, 85/40 \text{ \%}) = 1.88 \pm 0.10$	$\gamma_{\beta}(455 \text{ nm}) = 0.46 \pm 0.04$
11:00	Tandem (550 nm)	ground	$(f_{\sigma}(550 \text{ nm}, 85/ < 40 \text{ \%}) = 1.71 \pm 0.14$	$\gamma_{\sigma}(550 \text{ nm}) = 0.40 \pm 0.04$
23:00	COBALD-RS41 (940 nm)	0.47-0.99	$(f_{\beta}(940 \text{ nm}, 85/40 \text{ \%}) = 2.35 \pm 0.17$	$\gamma_{\beta}(940 \text{ nm}) = 0.62 \pm 0.05$
23:30	Ceilometer (1064 nm)		$(f_{\beta}(1064 \text{ nm}, 85/40 \text{ \%}) = 1.83 \pm 0.05$	$\gamma_{\beta}(1064 \text{ nm}) = 0.43 \pm 0.04$
Case III: 10 December 2019				
Hour (UTC)	Technique	Layer (km agl)	$f_{\zeta}(\lambda, RH)$	$\gamma_{\zeta}(\lambda)$
08:00	RALMO (355 nm)	0.70-0.97	$(f_{\beta}(355 \text{ nm}, 85/40 \text{ \%}) = 1.89 \pm 0.24$	$\gamma_{\beta}(355 \text{ nm}) = 0.46 \pm 0.09$
09:00	Tandem (550 nm)	ground	$(f_{\sigma}(550 \text{ nm}, 85/ < 40 \text{ \%}) = 1.86 \pm 0.17$	$\gamma_{\sigma}(550 \text{ nm}) = 0.48 \pm 0.05$
08:00	Ceilometer (1064 nm)	0.70-0.97	$(f_{\beta}(1064 \text{ nm}, 85/40 \text{ \%}) = 1.30 \pm 0.03$	$\gamma_{\beta}(1064 \text{ nm}) = 0.19 \pm 0.01$



5 Conclusions

This study has focused on aerosol hygroscopicity during the intercomparison campaign of hygroscopic aerosol properties using remote sensing and in situ techniques (HYGRO19 campaign), which was carried out at the MeteoSwiss aerological station in Payerne, Switzerland (46.82° N, 6.95° E, 491 m asl) from June to December 2019. The main objective of the campaign was to assess the ability of the different techniques to characterise hygroscopic aerosol processes. Several instruments were deployed during the campaign, including the Raman Lidar for Meteorological Observations (RALMO), which provides continuous measurements of temperature, water vapour mixing ratio, and aerosol backscattering and extinction, both during daytime and nighttime. RALMO measurements alone enable the retrieval of aerosol hygroscopicity parameters, such as the enhancement factor (f_{ζ}) and the Hänel parameter (γ_{ζ}). The methodology is based on a simultaneous increase in aerosol backscatter coefficient and relative humidity (RH) with height under well-mixed layer conditions, characterised by an approximately constant water vapour mixing ratio and potential temperature. The advantage of this methodology over other approaches based on tandem nephelometers is that it provides aerosol hygroscopicity parameters under ambient conditions, without requiring particle drying. RALMO measurements provided more than 200 cases for aerosol hygroscopicity analysis, constituting one of the largest databases for Raman lidar-based hygroscopicity studies. The use of launches of Compact Backscatter Aerosol Detectors coupled to RS41 Vaisala radiosondes (COBALD-RS41) enabled a unique evaluation of aerosol hygroscopicity parameters derived from RALMO. In situ measurements also included a tandem nephelometers, allowing intercomparison under well-mixed conditions in the lower layers above the ground. Additional in situ instrumentation provided measurements of the aerosol absorption coefficient and particle size distributions. Real-time measurements of equivalent black carbon (EBC), PM_{2.5}, and PM₁₀ concentrations were also available. Aerosol chemical composition was obtained through laboratory analyses of filter samples. Doppler lidar measurements were used to provide wind profiles.

The Payerne station is also an operational site of the AERONET network, and during the campaign, analyses based on Aerosol Optical Depth (AOD) and extinction Ångström exponent (AOD-AE) revealed two distinct periods. The first, from June to August 2019, featured numerous measurements due to the frequent presence of clear skies. The median values of AOD (500nm) and AOD-AE (440–870 nm) were 0.24 ± 0.09 and 1.50 ± 0.19 , respectively, representing a considerable aerosol load for a rural background site such as Payerne. Backward-trajectory analyses indicated the influence of air masses from mainland Europe, which typically act as a source of anthropogenic pollution. However, some cases of air masses originating from Southwestern Europe and North Africa were also detected, showing higher aerosol loads (AOD (500nm) approximately 0.3) and low AOD-AE (approximately 0.4), suggesting the presence of dust particles. The second period September–December was characterised by the predominance of low-pressure systems and cloudy skies, with only a few AERONET measurements available. Median values during this period were AOD (500nm) = 0.07 ± 0.02 and AOD-AE (440–870 nm) = 1.15 ± 0.13 , more typical of a rural background site. These meteorological conditions largely limited the number of cases available for aerosol hygroscopicity studies using RALMO.

Aerosol hygroscopicity was studied using the Hänel equation, which allows the retrieval of the enhancement factor and the Hänel parameter. These parameters depend on the selected aerosol optical property. For instance, RALMO uses the aerosol



backscatter coefficient (β^{aer}) at 355 nm, whereas the tandem nephelometer system uses the total scattering coefficient (σ_{sp}) at 500 nm. Other factors, such as the measurement wavelength and aerosol optical and microphysical properties, directly affect the retrieved hygroscopicity parameters. Despite these differences, comparisons between techniques are performed qualitatively because of the different measurement approaches. However, to facilitate intercomparisons, all parameters are derived for the
670 RH range from $\leq 40\%$ to 85% .

The uniqueness of the HYGRO19 campaign was the continuous operation of the RALMO system, which allowed the retrieval of aerosol hygroscopicity parameters for more than 200 cases. The campaign was characterised by high hygroscopic aerosol growth, with $f_{\beta}(355\text{ nm}, 85/40\%)$ values ranging from 1.2 to 7.2 and a median of 2.64 ± 0.64 , corresponding to median $\gamma_{\beta}(355\text{ nm})$ of 0.70 ± 0.18 . The highest occurrence was in summer (177 layers, 78 % of all cases), with aerosols
675 being more hygroscopic in June ($f_{\beta}(355\text{ nm}) = 3.65 \pm 0.70$ and $\gamma_{\beta}(355\text{ nm}) = 0.94 \pm 0.15$), which can be associated with an aerosol composition dominated by secondary inorganic species (sulphate, nitrate, and ammonium). Fewer cases were detected in the period October–December, and lower hygroscopicity was observed ($f_{\beta}(355\text{ nm}) = 2.36 \pm 0.26$ and $\gamma_{\beta}(355\text{ nm}) = 0.60 \pm 0.08$) probably explained by the high fractions of EC and organic matter (OM) and the low concentrations of sulphate. On the other hand, the evaluation of the hygroscopicity parameters with altitude was explored, and no statistically signifi-
680 cant differences were observed when differentiating between values above and below 1.5 km agl, which was assumed as the threshold separating the Atmospheric Boundary layer (ABL) from the Free Troposphere (FT). A more complete approach additionally distinguished between the Convective Boundary Layer (CBL) and the Residual Layer (RL), and again no statistically significant differences were found.

Comparisons between RALMO and in situ techniques (tandem nephelometers) during the second part of the campaign
685 showed good agreement for aerosol layers within the residual layer, when well-mixed conditions extended up to the RL (e.g., on 10 December 2019, $\gamma_{\beta}(355\text{ nm}) = 0.46 \pm 0.09$ for RALMO and $\gamma_{\beta}(550\text{ nm}) = 0.48 \pm 0.05$ for the tandem nephelometers). However, some discrepancies were found for some layers identified within the free troposphere, which may be due to the presence of decoupled layers, which can exhibit different microphysical and optical properties compared to those near the surface. Representative case studies were analysed to provide a comprehensive assessment of aerosol hygroscopicity using remote
690 sensing (RALMO and ceilometer), vertically resolved in situ observations (COBALD-RS41 sondes), and surface in situ measurements from tandem nephelometers. The results confirmed the capability of these techniques to robustly characterise aerosol hygroscopicity, as indicated by the close agreement between the retrieved hygroscopicity parameters and their associated uncertainties. A weak wavelength-dependence was also observed, with larger hygroscopicity parameters at shorter wavelengths. Aerosol chemistry from ground observations was used as a proxy to interpret aerosol hygroscopicity with height; however, fur-
695 ther analyses were not possible, and future work requires the combination of remote sensing with real-time chemistry profiles. Furthermore, Raman lidar measurements revealed high aerosol hygroscopicity in the vicinity of clouds, suggesting that aerosol hygroscopicity may facilitate particle activation as cloud condensation nuclei under favourable thermodynamic and dynamical conditions. Additional studies using numerical weather prediction models capable of accurately reproducing both clouds and the evolution of aerosol hygroscopicity are needed.



700 *Data availability.* RALMO data can be accessed through the EARLINET network (<https://data.earlinet.org/earlinet/>), while ceilometer and
Doppler wind lidar data can be obtained via the E-PROFILE network (<https://data.ceda.ac.uk/badc/eprofile/data/switzerland/payerne>). Co-
author Alexander Haeferle can be contacted for access to raw RALMO and radiosonde measurements (alexander.haeferle@meteoswiss.ch).
Sun photometer data can be retrieved from the AERONET network (<https://aeronet.gsfc.nasa.gov/>). In situ aerosol data can be requested
from co-author Benjamin T. Brem (benjamin.brem@psi.ch), while chemical composition data from the NABEL station can be accessed at
705 <https://www.bafu.admin.ch/en/data-query-nabel>.



Appendix A: In situ measurements for the case studies analysed




Figure A1. Time hourly series (left panels) over 16 to 18 September 2019 of: (a) aerosol scattering coefficient at 450, 525, and 635 nm from the Aurora 3000 dry nephelometer, (b) absorption coefficient at 370, 520 and 880 nm from the aethalometer AE31, (c) scattering (SAE (450–635 nm)) and absorption (AAE (370–950 nm)) Ångström exponents and (d) single scattering albedo (SSA) at 520 nm derived from the dry nephelometer and aethalometer, (e) PM concentrations, (f) equivalent black carbon at 880 nm from the aethalometer. Aerosol size distribution from the SMPS (right panels) for mean between (g) 00:00–12:00 UTC and (h) 12:00–24:00 UTC from 16 to 18 September 2019, and (i) 2 hourly mean on 17 September 2019.




Figure A2. Time hourly series (left panels) over 11 to 13 November 2019 of: (a) aerosol scattering coefficient at 450, 525, and 635 nm from the Aurora 3000 dry nephelometer, (b) absorption coefficient at 370, 520 and 880 nm from the aethalometer AE31, (c) scattering (SAE (450–635 nm)) and absorption (AAE (370–950 nm)) Ångström exponents and (d) single scattering albedo (SSA) at 520 nm derived from the dry nephelometer and aethalometer, (e) PM concentrations, (f) equivalent black carbon at 880 nm from the aethalometer. Aerosol size distribution from the SMPS (right panels) for mean between (g) 00:00–12:00 UTC and (h) 12:00–24:00 UTC from 11 to 13 November 2019, and (i) 2 hourly mean on 12 November 2019.




Figure A3. Time hourly series over 9 to 11 December 2019 of: (a) aerosol scattering coefficient at 450, 525, and 635 nm from the Aurora 3000 dry nephelometer, (b) absorption coefficient at 370, 520 and 880 nm from the aethalometer AE31, (c) scattering (SAE (450–635 nm)) and absorption (AAE (370–950 nm)) Ångström exponents and (d) single scattering albedo (SSA) at 520 nm derived from the dry nephelometer and aethalometer, (e) PM concentrations, (f) equivalent black carbon at 880 nm from the aethalometer.

Author contributions. F.N.G., S.B. and A.H. conceptualised the study. A.D.Z., F.N.G. and all co-authors contributed to the analysis of the results and the discussion. A.D.Z. prepared the original draft of the manuscript, which was reviewed and edited by all authors. F.N.G. supervised the study and acquired funding.

710 Competing interests. The authors declare no conflicts of interest.

Acknowledgements. This research was funded by Grant PID2021-128008OB-I00 funded by MICIU/AEI/ 10.13039/501100011033 by ERD-F/EU European Union. This work was also supported by the Swiss National Science Foundation through project PZ00P2 168114 and the Spanish national projects Grant PID2024-162154OB-I00, PID2023-151817OA-I00, CNS2023-145435, and by the Horizon Europe program under the Marie Skłodowska-Curie Staff Exchange Actions with the project GRASP-SYNERGY (grant agreement no. 101131631). This article is based upon work from COST Action EARLICOST, supported by COST (European Cooperation in Science and Technology). The work made use of the strategic networks RED2022-134824-E and RED2024-153821-E and infrastructure grants EQC2019-006192-P and EQC2019-006423-P founded by MCIN/AEI /10.13039/501100011033, ATMO-ACCESS grant agreement No 101008004, ACTRIS-IMP grant agreement No 871115, and Scientific Unit of Excellence: Earth System (UCE-PP2017-02).



References

- 720 Alonso-Blanco, E., Gómez-Moreno, F. J., and Artíñano, B.: Size-resolved hygroscopicity of ambient submicron particles in a suburban Atmos, Atmos Environ, 213, 349–358, <https://doi.org/10.1016/j.atmosenv.2019.05.065>, 2019.
- Anderson, T. L. and Ogren, J. A.: Determining Aerosol Radiative Properties Using the TSI 3563 Integrating Nephelometer, Aerosol Sci. Technol., 29, 57–69, <https://doi.org/10.1080/02786829808965551>, 1998.
- Ansmann, A., Riebesell, M., and Weitkamp, C.: Measurement of atmospheric aerosol extinction profiles with a Raman lidar, Opt Lett, 15, 725 <https://doi.org/10.1364/ol.15.000746>, 1990.
- Bader, S., Collaud Coen, M., Duguay-Tetzlaff, A., Frei, C., Fukutome, S., Gehrig, R., Maillard Barras, E., Martucci, G., Romanens, G., Scherrer, S., Schlegel, T., Spirig, C., Stübi, R., Vuilleumier, L., and Zubler, E.: Rapport climatologique 2019, Rapports et Bulletins 313.008.f, MeteoSwiss and OFCL, Vente des publications fédérales, CH-3303 Berne, Switzerland, ISSN 2296-1496, <https://www.meteosuisse.admin.ch/services-et-publications/publications/rapports-et-bulletins/2016/2020/rapport-climatologique-2019.html>, 2020.
- 730 Bedoya-Velásquez, A. E., Navas-Guzmán, F., Granados-Muñoz, M. J., Titos, G., Román, R., Casquero-Vera, J. A., Ortiz-Amezcu, P., Benavent-Oltra, J. A., Moreira, G. D. A., Montilla-Rosero, E., Hoyos, C. D., Artíñano, B., Coz, E., Olmo-Reyes, F. J., Alados-Arboledas, L., and Guerrero-Rascado, J. L.: Hygroscopic growth study in the framework of EARLINET during the SLOPE i campaign: Synergy of Remote Sens and in situ instrumentation, Atmos Chem Phys, 18, 7001–7017, <https://doi.org/10.5194/acp-18-7001-2018>, 2018.
- Bedoya-Velásquez, A. E., Titos, G., Bravo-Aranda, J. A., Haefelin, M., Favez, O., Petit, J. E., Casquero-Vera, J. A., Olmo-Reyes, 735 F. J., Montilla-Rosero, E., Hoyos, C. D., Alados-Arboledas, L., and Guerrero-Rascado, J. L.: Long-term aerosol optical hygroscopicity study at the ACTRIS SIRTa observatory: Synergy between ceilometer and in situ measurements, Atmos Chem Phys, 19, <https://doi.org/10.5194/acp-19-7883-2019>, 2019.
- Bindoff, N. L., Stott, P., AchutaRao, K., Allen, M., Gillett, N., Gutzler, D., Hansingo, K., Hegerl, G., Hu, Y., Jain, S., Mokhov, I., Overland, J., Perlwitz, J., Sebbari, R., and Zhang, X.: IPCC 2013 AR5 - Chapter 10: Detection and Attribution of Climate Change: From Global 740 to Regional, Climate Change 2013: The Physical Science Basis - Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change 2013, Change, edited by: Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, Cambridge, UK, New York, NY, USA, 9781107057999, 867–952, <https://doi.org/10.1017/CBO9781107415324.022>, 2013.
- Brocard, E., Philipona, R., Haefele, A., Romanens, G., Mueller, A., Ruffieux, D., Simeonov, V., and Calpini, B.: Raman lidar 745 for meteorological observations, RALMO - Part 2: Validation of water vapor measurements, Atmos Meas Tech, 6, 1347–1358, <https://doi.org/10.5194/amt-6-1347-2013>, 2013.
- Brunamonti, S., Jorge, T., Oelsner, P., Hanumanthu, S., Singh, B. B., Kumar, K. R., Sonbawne, S., Meier, S., Singh, D., Wienhold, F. G., Luo, B. P., Boettcher, M., Poltera, Y., Jauhiainen, H., Kayastha, R., Karmacharya, J., Dirksen, R., Naja, M., Rex, M., Fadnavis, S., and Peter, T.: Balloon-borne measurements of temperature, water vapor, ozone and aerosol backscatter on the southern slopes of the Himalayas during 750 StratoClim 2016-2017, Atmos Chem Phys, 18, 15 937–15 957, <https://doi.org/10.5194/acp-18-15937-2018>, 2018.
- Brunamonti, S., Martucci, G., Romanens, G., Poltera, Y., Wienhold, F. G., Hervo, M., Haefele, A., and Navas-Guzmán, F.: Validation of aerosol backscatter profiles from Raman lidar and ceilometer using balloon-borne measurements, Atmos Chem Phys, 21, 2267–2285, <https://doi.org/10.5194/acp-21-2267-2021>, 2021.
- Bucholtz, A.: Rayleigh-scattering calculations for the terrestrial Atmos, Appl Opt, 34, 2765–2773, <https://doi.org/10.1364/ao.34.002765>, 755 1995.



- Burgos, M. A., Andrews, E., Titos, G., Alados-Arboledas, L., Baltensperger, U., Day, D., Jefferson, A., Kalivitis, N., Mihalopoulos, N., Sherman, J., Sun, J., Weingartner, E., and Zieger, P.: A global view on the effect of water uptake on aerosol particle light scattering, *Sci Data*, 6, <https://doi.org/10.1038/s41597-019-0158-7>, 2019.
- Carrico, C. M., Kus, P., Rood, M. J., Quinn, P. K., and Bates, T. S.: Mixtures of pollution, dust, sea salt, and volcanic aerosol during ACE-Asia: Radiative properties as a function of relative humidity, *J Geophys Res Atmos*, 108, <https://doi.org/10.1029/2003jd003405>, 2003.
- Cavalli, F., Viana, M., Yttri, K. E., Genberg, J., and Putaud, J. P.: Toward a standardised thermal-optical protocol for measuring atmospheric organic and elemental carbon: The EUSAAR protocol, *Atmos Meas Tech*, 3, <https://doi.org/10.5194/amt-3-79-2010>, 2010.
- Cedeño, E., Sun, Z., Hong, Z., and Du, J.: Raman spectroscopy for profiling physical and chemical properties of atmospheric aerosol particles: A review, *Ecotoxicol. Environ. Saf.*, 249, <https://doi.org/10.1016/j.ecoenv.2022.114405>, 2023.
- 765 Chen, J., Li, Z., Lv, M., Wang, Y., Wang, W., Zhang, Y., Wang, H., Yan, X., Sun, Y., and Cribb, M.: Aerosol hygroscopic growth, contributing factors, and impact on haze events in a severely polluted region in northern China, *Atmos Chem Phys*, 19, <https://doi.org/10.5194/acp-19-1327-2019>, 2019.
- Chen, X., Chu, Y., Chu, Y., Lee, A. K., Gen, M., Kasthuriarachchi, N. Y., Chan, C. K., and Li, Y. J.: Relative Humidity History Affects Hygroscopicity of Mixed Particles of Glyoxal and Reduced Nitrogenous Species, *Environ Sci Technol*, 54, <https://doi.org/10.1021/acs.est.0c00680>, 2020.
- 770 Chicco, D., Sichenze, A., and Jurman, G.: A simple guide to the use of Student's t-test, Mann-Whitney U test, Chi-squared test, and Kruskal-Wallis test in biostatistics, *BioData Mining*, 18, <https://doi.org/10.1186/s13040-025-00465-6>, 2025.
- Cirisan, A., Luo, B. P., Engel, I., Wienhold, F. G., Sprenger, M., Krieger, U. K., Weers, U., Romanens, G., Levrat, G., Jeannet, P., Ruffieux, D., Philipona, R., Calpini, B., Spichtinger, P., and Peter, T.: Balloon-borne match measurements of midlatitude cirrus clouds, *Atmos Chem Phys*, 14, 7341–7365, <https://doi.org/10.5194/acp-14-7341-2014>, 2014.
- 775 Clarke, A. D., Howell, S., Quinn, P. K., Bates, T. S., Ogren, J. A., Andrews, E., Jefferson, A., Massling, A., Mayol-Bracero, O., Maring, H., Savoie, D., and Cass, G.: INDOEX aerosol: A comparison and summary of chemical, microphysical, and optical properties observed from land, ship, and aircraft, *J Geophys Res Atmos*, 107, <https://doi.org/10.1029/2001JD000572>, 2002.
- Collaud Coen, M., Praz, C., Haeefe, A., Ruffieux, D., Kaufmann, P., and Calpini, B.: Determination and climatology of the planetary boundary layer height above the Swiss plateau by in situ and Remote Sens measurements as well as by the COSMO-2 model, *Atmos Chem Phys*, 14, <https://doi.org/10.5194/acp-14-13205-2014>, 2014.
- 780 Collaud Coen, M., Brem, B. T., Gysel-Beer, M., Modini, R., Henne, S., Steinbacher, M., Putero, D., Gini, M. I., and Eleftheriadis, K.: Detection and climatology of Saharan dust frequency and mass at the Jungfrauoch (3580 m asl, Switzerland), *Atmos Chem Phys*, 26, 1623–1645, <https://doi.org/10.5194/acp-26-1623-2026>, 2026.
- 785 Dada, L., Brem, B. T., Amarandi-Netedu, L.-M., Collaud Coen, M., Evangelidou, N., Hüglin, C., Nowak, N., Modini, R., Steinbacher, M., and Gysel-Beer, M.: Sources of ultrafine particles at a rural midland site in Switzerland, *Aerosol Res*, 3, 315–336, <https://doi.org/10.5194/ar-3-315-2025>, 2025.
- Deshmukh, S., Poulain, L., Wehner, B., Henning, S., Petit, J.-E., Fombelle, P., Favez, O., Herrmann, H., and Pöhlker, M.: External particle mixing influences hygroscopicity in a sub-urban area, *Atmos Chem Phys*, 25, 741–758, <https://doi.org/10.5194/acp-25-741-2025>, 2025.
- 790 Díaz-Zurita, A., Naval-Hernández, V. M., Whiteman, D. N., Rodríguez-Navarro, O., Muñoz-Rosado, J., Pérez-Ramírez, D., Alados-Arboledas, L., and Navas-Guzmán, F.: Sensitivity Analysis of the Differential Atmospheric Transmission in Water Vapour Mixing Ratio Retrieval from Raman Lidar Measurements, *Remote Sens*, 17, 3444, <https://doi.org/10.3390/rs17203444>, 2025.



- Dinoyev, T., Simeonov, V. B., Arshinov, Y., Bobrovnikov, S., Ristori, P., Calpini, B., Parlange, M., and Bergh, H. V. D.: Raman Lidar for Meteorological Observations, RALMO - Part 1: Instrument description, *Atmos Meas Tech*, 6, 1329–1346, <https://doi.org/10.5194/amt-6-1329-2013>, 2013.
- Doane, D. P.: Aesthetic frequency classifications, *American Statistician*, 30, <https://doi.org/10.1080/00031305.1976.10479172>, 1976.
- Douville, H., Raghavan, K., Renwick, J., Allan, R., Arias, P., Barlow, M., Cerezo-Mota, R., Cherchi, A., Gan, T., Gergis, J., Jiang, D., Khan, A., Pokam Mba, W., Rosenfeld, D., Tierney, J., and Zolina, O.: Water Cycle Changes, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., chap. 8, pp. 1055–1210, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.010>, 2021.
- Durand, E., Durdina, L., Smallwood, G., Johnson, M., Spirig, C., Edebeli, J., Roth, M., Brem, B., Sevcenco, Y., and Crayford, A.: Correction for particle loss in a regulatory aviation nvPM emissions system using measured particle size, *J Aerosol Sci*, 169, <https://doi.org/10.1016/j.jaerosci.2023.106140>, 2023.
- Eck, T. F., Holben, B. N., Reid, J. S., Giles, D. M., Rivas, M. A., Singh, R. P., Tripathi, S. N., Bruegge, C. J., Platnick, S., Arnold, G. T., Krotkov, N. A., Carn, S. A., Sinyuk, A., Dubovik, O., Arola, A., Schafer, J. S., Artaxo, P., Smirnov, A., Chen, H., and Goloub, P.: Fog- and cloud-induced aerosol modification observed by the aerosol robotic network (AERONET), *J Geophys Res Atmos*, 117, <https://doi.org/10.1029/2011JD016839>, 2012.
- Eirund, G. K., Drossaert van Dusseldorp, S., Brem, B. T., Dedekind, Z., Karrer, Y., Stoll, M., and Lohmann, U.: Aerosol–cloud–precipitation interactions during a Saharan dust event—A summertime case-study from the Alps, *QJR Meteorol Soc*, 148, 943–961, <https://doi.org/10.1002/qj.4240>, 2022.
- EMPA: Technischer Bericht zum Nationalen Beobachtungsnetz für Luftfremdstoffe (NABEL), Tech. rep., EMPA, Swiss Federal Laboratories for Materials Science and Technology, Dübendorf, Switzerland, Technical report, 2013.
- Feingold, G. and Morley, B.: Aerosol hygroscopic properties as measured by lidar and comparison with in situ measurements, *J. Geophys. Res. Atmos*, 108, <https://doi.org/10.1029/2002jd002842>, 2003.
- Fernández, A. J., Molero, F., Becerril-Valle, M., Coz, E., Salvador, P., Artñano, B., and Pujadas, M.: Application of Remote Sens techniques to study aerosol water vapour uptake in a real Atmos, *Atmos. Res.*, 202, 112–127, <https://doi.org/10.1016/j.atmosres.2017.11.020>, 2018.
- Ferrare, R. A., Melfi, S. H., Whiteman, D. N., Evans, K. D., Poellot, M., and Kaufman, Y. J.: Raman lidar measurements of aerosol extinction and backscattering 2. Derivation of aerosol real refractive index, single-scattering albedo, and humidification factor using Raman lidar and aircraft size distribution measurements, *J Geophys Res Atmos*, 103, <https://doi.org/10.1029/98JD01647>, 1998.
- Fierz-Schmidhauser, R., Zieger, P., Gysel, M., Kammermann, L., DeCarlo, P. F., Baltensperger, U., and Weingartner, E.: Measured and predicted aerosol light scattering enhancement factors at the high alpine site Jungfraujoch, *Atmos Chem Phys*, 10, <https://doi.org/10.5194/acp-10-2319-2010>, 2010.
- Giles, D. M., Sinyuk, A., Sorokin, M. G., Schafer, J. S., Smirnov, A., Slutsker, I., Eck, T. F., Holben, B. N., Lewis, J. R., Campbell, J. R., Welton, E. J., Korkin, S. V., and Lyapustin, A. I.: Advancements in the Aerosol Robotic Network (AERONET) Version 3 database - Automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements, *Atmos Meas Tech*, 12, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- Goldsmith, J. E.: Turn-key Raman lidar for profiling atmospheric water vapor, clouds, and aerosols, *Appl Opt*, 37, <https://doi.org/10.1364/ao.37.004979>, 1998.



- Granados-Muñoz, M. J., Navas-Guzmán, F., Bravo-Aranda, J. A., Guerrero-Rascado, J. L., Lyamani, H., Valenzuela, A., Titos, G., Fernández-Gálvez, J., and Alados-Arboledas, L.: Hygroscopic growth of atmospheric aerosol particles based on active Remote Sens and radiosounding measurements: Selected cases in southeastern Spain, *Atmos Meas Tech*, 8, 705–718, <https://doi.org/10.5194/amt-8-705-2015>, 2015.
- Gysel, M., Crosier, J., Topping, D. O., Whitehead, J. D., Bower, K. N., Cubison, M. J., Williams, P. I., Flynn, M. J., McFiggans, G. B., and
835 Coe, H.: Closure study between chemical composition and hygroscopic growth of aerosol particles during TORCH2, *Atmos Chem Phys*, 7, <https://doi.org/10.5194/acp-7-6131-2007>, 2007.
- Haefele, A., Hervo, M., Turp, M., Lampin, J., Haeffelin, M., Lehmann, V., et al.: The E-PROFILE network for the operational measurement of wind and aerosol profiles over Europe, *Proceeding of TECO*, Madrid, Spain, available at: https://www.eumetnet.eu/wp-content/uploads/2016/10/E-PROFILE_TECO_Madrid_2016.pdf, 2016.
- 840 Hicks-Jalali, S., Sica, R. J., Martucci, G., Barras, E. M., Voirin, J., and Haefele, A.: A Raman lidar tropospheric water vapour climatology and height-resolved trend analysis over Payerne, Switzerland, *Atmos Chem Phys*, 20, <https://doi.org/10.5194/acp-20-9619-2020>, 2020.
- Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET - A federated instrument network and data archive for aerosol characterization, *Remote Sens Environ*, 66, [https://doi.org/10.1016/S0034-4257\(98\)00031-5](https://doi.org/10.1016/S0034-4257(98)00031-5), 1998.
- 845 Hänel, G.: The properties of atmospheric aerosol particles as functions of the relative humidity at thermodynamic equilibrium with the surrounding moist air, *Adv Geophys*, 19, [https://doi.org/10.1016/S0065-2687\(08\)60142-9](https://doi.org/10.1016/S0065-2687(08)60142-9), 1976.
- Hüglin, C. and Grange, S. K.: Chemical Characterisation and Source Identification of PM₁₀ and PM_{2.5} in Switzerland, Project report, Empa, Swiss Federal Laboratories for Materials Science and Technology, Dübendorf, Switzerland, <https://www.bafu.admin.ch/dam/en/sd-web/oyMXRQizcq24/chemical-characterisation-and-source-identification-of-pm-in-switzerland.pdf>, commissioned by the Federal Office for
850 the Environment (FOEN), 2021.
- Hüglin, C. and Schwarzenbach, B.: Feinstaubmessungen (PM₁₀ und PM_{2.5}) mit dem Fidas 200 der Firma Palas im NABEL, Tech. rep., Empa, Abteilung für Luftfremdstoffe und Umwelttechnik, https://www.empa.ch/documents/56101/246436/Fidas_200_NABEL.pdf/93790947-135a-4575-a718-370316f506c5, Technical report, 2016.
- Hüglin, C., Buchmann, B., Steinbacher, M., and Emmenegger, L.: The Swiss National Air Pollution Monitoring Network (NABEL) –
855 Bridging Science and Environmental Policy, *Chimia*, 78, 722–727, <https://doi.org/10.2533/chimia.2024.722>, 2024.
- Jefferson, A., Hageman, D., Morrow, H., Mei, F., and Watson, T.: Seven years of aerosol scattering hygroscopic growth measurements from SGP: Factors influencing water uptake, *J. Geophys. Res. Atmos.*, 122, <https://doi.org/10.1002/2017JD026804>, 2017.
- Jing, D., He, Y., Yin, Z., Müller, D., Huang, K., and Yi, F.: Hygroscopic growth characteristics of anthropogenic aerosols over central China revealed by lidar observations, *Atmos Meas Tech*, 19, 389–403, <https://doi.org/10.5194/amt-19-389-2026>, 2026.
- 860 Jones, A. C., Hill, A., Remy, S., Abraham, N. L., Dalvi, M., Hardacre, C., Hewitt, A. J., Johnson, B., Mulcahy, J. P., and Turnock, S. T.: Exploring the sensitivity of atmospheric nitrate concentrations to nitric acid uptake rate using the Met Office’s Unified Model, *Atmos Chem Phys*, 21, 15 901–15 927, <https://doi.org/10.5194/acp-21-15901-2021>, 2021.
- Kammermann, L., Gysel, M., Weingartner, E., and Baltensperger, U.: 13-month climatology of the aerosol hygroscopicity at the free tropospheric site Jungfraujoch (3580 m a.s.l.), *Atmos Chem Phys*, 10, <https://doi.org/10.5194/acp-10-10717-2010>, 2010.
- 865 Keller, A., Specht, P., Steigmeier, P., and Weingartner, E.: A novel measurement system for unattended, in situ characterization of carbonaceous aerosols, *Aerosol Res*, 1, <https://doi.org/10.5194/ar-1-65-2023>, 2023.
- Klett, J. D.: Stable analytical inversion solution for processing lidar returns, *Appl Opt*, 20, 211–220, <https://doi.org/10.1364/ao.20.000211>, 1981.



- 870 Klett, J. D.: Lidar inversion with variable backscatter/extinction ratios, *Appl Opt*, 24, 1638–1643, <https://doi.org/10.1364/ao.24.001638>, 1985.
- Laly, F., Chazette, P., Totems, J., Crenn, V., Ledur, D., and Marpillat, A.: How does humidity affect lidar-derived aerosol optical properties, and how do they compare with CAMS?, *Atmos Meas Tech*, 18, 7629–7649, <https://doi.org/10.5194/amt-18-7629-2025>, 2025.
- Li, L., Li, M., Fan, X., Chen, Y., Lin, Z., Hou, A., Zhang, S., Zheng, R., and Chen, J.: Measurement report: The variation properties of aerosol hygroscopic growth related to chemical composition during new particle formation days in a coastal city of Southeast China, *Atmos Chem Phys*, 25, 3669–3685, <https://doi.org/10.5194/acp-25-3669-2025>, 2025.
- 875 List, R. J.: *Smithsonian Meteorological Tables*, 6th rev. edn., Washington, DC, 1951.
- Liu, L., Tan, H., Fan, S., Cai, M., Xu, H., Li, F., and Chan, P.: Influence of aerosol hygroscopicity and mixing state on aerosol optical properties in the Pearl River Delta region, China, *Sci Total Environ*, 627, <https://doi.org/10.1016/j.scitotenv.2018.01.199>, 2018.
- Liu, Y., Feng, Z., Zheng, F., Bao, X., Liu, P., Ge, Y., Zhao, Y., Jiang, T., Liao, Y., Zhang, Y., Fan, X., Yan, C., Chu, B., Wang, Y., Du, W., Cai, J., Bianchi, F., Petäjä, T., Mu, Y., He, H., and Kulmala, M.: Ammonium nitrate promotes sulfate formation through uptake kinetic regime, *Atmos Chem Phys*, 21, 13 269–13 286, <https://doi.org/10.5194/acp-21-13269-2021>, 2021.
- 880 Lorenzo, G. R., Ziemba, L. D., Arellano, A. F., Barth, M. C., Crosbie, E. C., DiGangi, J. P., Diskin, G. S., Ferrare, R., Hilario, M. R. A., Shook, M. A., Tilmes, S., Wang, J., Xiao, Q., Zhang, J., and Sorooshian, A.: Measurement report: Characterization of aerosol hygroscopicity over Southeast Asia during the NASA CAMP²Ex campaign, *Atmos Chem Phys*, 25, 5469–5495, <https://doi.org/10.5194/acp-25-5469-2025>, 2025.
- 885 Lv, M., Liu, D., Li, Z., Mao, J., Sun, Y., Wang, Z., Wang, Y., and Xie, C.: Hygroscopic growth of atmospheric aerosol particles based on lidar, radiosonde, and in situ measurements: Case studies from the Xinzhou field campaign, *J Quant Spectrosc Radiat Transf.*, 188, <https://doi.org/10.1016/j.jqsrt.2015.12.029>, 2017.
- Martucci, G., Voirin, J., Simeonov, V., Renaud, L., and Haeefe, A.: A novel automatic calibration system for water vapor Raman LIDAR, in: *EPJ Web of Conferences*, vol. 176, p. 05008, EDP Sciences, ISSN 2100014X, <https://doi.org/10.1051/epjconf/201817605008>, 2018.
- 890 Martucci, G., Navas-Guzmán, F., Renaud, L., Romanens, G., Gamage, S. M., Hervé, M., Jeannet, P., and Haeefe, A.: Validation of pure rotational Raman temperature data from the Raman Lidar for Meteorological Observations (RALMO) at Payerne, *Atmos Meas Tech*, 14, 1333–1353, <https://doi.org/10.5194/amt-14-1333-2021>, 2021.
- MeteoSwiss: Air Pressure and Anticyclone Overview, <https://www.meteoswiss.admin.ch/weather/weather-and-climate-from-a-to-z/anticyclone-high-pressure.html>, accessed: 12 Jan 2026, 2026a.
- 895 MeteoSwiss: Air Pressure and Anticyclone Overview, <https://www.meteoswiss.admin.ch/weather/weather-and-climate-from-a-to-z/low-pressure-situation.html>, accessed: 12 Jan 2026, 2026b.
- Minguillón, M. C., Querol, X., Baltensperger, U., and Prévôt, A. S.: Fine and coarse PM composition and sources in rural and urban sites in Switzerland: Local or regional pollution?, *Sci Total Environ*, 427–428, <https://doi.org/10.1016/j.scitotenv.2012.04.030>, 2012.
- 900 Miri, R., Pujol, O., Hu, Q., Goloub, P., Veselovskii, I., Podvin, T., and Ducos, F.: Innovative aerosol hygroscopic growth study from Mie–Raman–fluorescence lidar and microwave radiometer synergy, *Atmos Meas Tech*, 17, 3367–3375, <https://doi.org/10.5194/amt-17-3367-2024>, 2024.
- Mouton, M., Malek, K. A., James, M. H., Pokhrel, R. P., Fiddler, M. N., Asa-Awuku, A. A., and Bililign, S.: The hygroscopic properties of biomass burning aerosol from Eucalyptus and cow dung under different combustion conditions, *Aerosol Sci. Technol.*, 57, 665–677, <https://doi.org/10.1080/02786826.2023.2198587>, 2023.
- 905



- Müller, T., Laborde, M., Kassell, G., and Wiedensohler, A.: Design and performance of a three-wavelength LED-based total scatter and backscatter integrating nephelometer, *Atmos Meas Tech*, 4, 1291–1303, <https://doi.org/10.5194/amt-4-1291-2011>, 2011.
- Navas-Guzmán, F., Fernández-Gálvez, J., Granados-Muñoz, M. J., Guerrero-Rascado, J. L., Bravo-Aranda, J. A., and Alados-Arboledas, L.: Tropospheric water vapour and relative humidity profiles from lidar and microwave radiometry, *Atmos Meas Tech*, 7, 1201–1211, <https://doi.org/10.5194/amt-7-1201-2014>, 2014.
- Navas-Guzmán, F., Kämpfer, N., and Haeefe, A.: Validation of brightness and physical temperature from two scanning microwave radiometers in the 60 GHz O₂ band using radiosonde measurements, *Atmos Meas Tech*, 9, 4587–4600, <https://doi.org/10.5194/amt-9-4587-2016>, 2016.
- Navas-Guzmán, F., Martucci, G., Coen, M. C., Granados-Muñoz, M. J., Hervó, M., Sicard, M., and Haeefe, A.: Characterization of aerosol hygroscopicity using Raman lidar measurements at the EARLINET station of Payerne, *Atmos Chem Phys*, 19, 11 651–11 668, <https://doi.org/10.5194/acp-19-11651-2019>, 2019.
- Nowak, D.: Radiation and clouds: observations and model calculations for Payerne BSRN site, Ph.D. thesis, ETH Zurich, 2008.
- Nowak, N., Brem, B., Gysel-Beer, M., and Simon, L.: Inorganics in air and particle phase at Payerne, EBAS at NILU, <https://doi.org/10.48597/HB8T-9C9B>, 2023–2025.
- Pariyothon, J., Bualert, S., Choomanee, P., Rungratanaubon, T., Thongyen, T., Fakkaew, N., Phuetfoo, C., Phupijit, J., and Szymanski, W. W.: Hygroscopic Growth Factors of Sub-micrometer Atmospheric Aerosols at Four Selected Sites in Thailand, *Aerosol Air Qual Res*, 23, <https://doi.org/10.4209/aaqr.220374>, 2023.
- Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S. M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, X. Y.: Recommendations for reporting black carbon measurements, *Atmos Chem Phys*, 13, <https://doi.org/10.5194/acp-13-8365-2013>, 2013.
- Pérez-Ramírez, D., Whiteman, D. N., Veselovskii, I., Ferrare, R., Titos, G., Granados-Muñoz, M. J., Sanchez-Hernandez, G., and Navas-Guzman, F.: Spatiotemporal changes in aerosol properties by hygroscopic growth and impacts on radiative forcing and heating rates during DISCOVER-AQ 2011, *Atmos Chem Phys*, 21, <https://doi.org/10.5194/acp-21-12021-2021>, 2021.
- Rosen, J. M. and Kjöme, N. T.: Backscattersonde: a new instrument for atmospheric Aerosol Res, *Appl Opt*, 30, 1552–1561, <https://doi.org/10.1364/ao.30.001552>, 1991.
- Rubia, J. M. D. L.: Rice University Rule to Determine the Number of Bins, *Open J Stat*, 14, <https://doi.org/10.4236/ojs.2024.141006>, 2024.
- Schafer, J. S., Eck, T. F., Holben, B. N., Artaxo, P., and Duarte, A. F.: Characterization of the optical properties of atmospheric aerosols in Amazônia from long-term AERONET monitoring (1993-1995 and 1999-2006), *J Geophys Res Atmos*, 113, <https://doi.org/10.1029/2007JD009319>, 2008.
- Scott, D. W.: Scott's rule, *Wiley Interdisciplinary Reviews: Computational Statistics*, 2, <https://doi.org/10.1002/wics.103>, 2010.
- Seinfeld, J. H., Bretherton, C., Carslaw, K. S., Coe, H., DeMott, P. J., Dunlea, E. J., Feingold, G., Ghan, S., Guenther, A. B., Kahn, R., Kraucunas, I., Kreidenweis, S. M., Molina, M. J., Nenes, A., Penner, J. E., Prather, K. A., Ramanathan, V., Ramaswamy, V., Rasch, P. J., Ravishankara, A. R., Rosenfeld, D., Stephens, G., and Wood, R.: Improving our fundamental understanding of the role of aerosol-cloud interactions in the climate system, *Proc Natl Acad Sci USA*, 113, <https://doi.org/10.1073/pnas.1514043113>, 2016.
- Sicard, M., Oliveira, D. C. F. D. S., Muñoz-Porcar, C., Gil-Díaz, C., Comerón, A., Rodríguez-Gómez, A., and Otín, F. D.: Measurement report: Spectral and statistical analysis of aerosol hygroscopic growth from multi-wavelength lidar measurements in Barcelona, Spain, *Atmos Chem Phys*, 22, <https://doi.org/10.5194/acp-22-7681-2022>, 2022.



- Stein, A. F., Draxler, R. R., Rolph, G. D., Stunder, B. J., Cohen, M. D., and Ngan, F.: NOAA's hysplit atmospheric transport and dispersion modeling system, *Bull Am Meteorol Soc*, 96, <https://doi.org/10.1175/BAMS-D-14-00110.1>, 2015.
- 945 Swietlicki, E., Hansson, H. C., Hämeri, K., Svenningsson, B., Massling, A., Mcfiggans, G., McMurry, P. H., Petäjä, T., Tunved, P., Gysel, M., Topping, D., Weingartner, E., Baltensperger, U., Rissler, J., Wiedensohler, A., and Kulmala, M.: Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments - A review, *Tellus, Series B: Chemical and Physical Meteorology*, 60 B, <https://doi.org/10.1111/j.1600-0889.2008.00350.x>, 2008.
- 950 Titos, G., Jefferson, A., Sheridan, P. J., Andrews, E., Lyamani, H., Alados-Arboledas, L., and Ogren, J. A.: Aerosol light-scattering enhancement due to water uptake during the TCAP campaign, *Atmos Chem Phys*, 14, 7031–7043, <https://doi.org/10.5194/acp-14-7031-2014>, 2014a.
- Titos, G., Lyamani, H., Cazorla, A., Sorribas, M., Foyo-Moreno, I., Wiedensohler, A., and Alados-Arboledas, L.: Study of the relative humidity dependence of aerosol light-scattering in southern Spain, *Tellus, B: Chem Phys Meteorol.*, 66, <https://doi.org/10.3402/tellusb.v66.24536>, 2014b.
- 955 Titos, G., Cazorla, A., Zieger, P., Andrews, E., Lyamani, H., Granados-Muñoz, M. J., Olmo, F. J., and Alados-Arboledas, L.: Effect of hygroscopic growth on the aerosol light-scattering coefficient: A review of measurements, techniques and error sources, <https://doi.org/10.1016/j.atmosenv.2016.07.021>, 2016.
- Titos, G., Burgos, M. A., Zieger, P., Alados-Arboledas, L., Baltensperger, U., Jefferson, A., Sherman, J., Weingartner, E., Henzing, B., Luoma, K., O'Dowd, C., Wiedensohler, A., and Andrews, E.: A global study of hygroscopicity-driven light-scattering enhancement in the context of other in situ aerosol optical properties, *Atmos Chem Phys*, 21, <https://doi.org/10.5194/acp-21-13031-2021>, 2021.
- 960 Vaisala: Vaisala Radiosonde RS41 Measurement Performance, Tech. rep., Vaisala, Helsinki, Finland, <https://www.vaisala.com/sites/default/files/documents/WEA-MET-RS41-Performance-White-paper-B211356EN-B-LOW-v3.pdf>, 2017.
- Vernier, J. P., Fairlie, T. D., Natarajan, M., Wienhold, F. G., Bian, J., Martinsson, B. G., Crumeyrolle, S., Thomason, L. W., and Bedka, K. M.: Increase in upper tropospheric and lower stratospheric aerosol levels and its potential connection with Asian pollution, *J. Geophys. Res. Atmos.*, 120, 1609–1619, <https://doi.org/10.1002/2014JD022372>, 2015.
- 965 Veselovskii, I., Whiteman, D. N., Kolgotin, A., Andrews, E., and Korenskii, M.: Demonstration of aerosol property profiling by multiwavelength lidar under varying relative humidity conditions, *J Atmos Ocean Technol*, 26, 1543–1557, <https://doi.org/10.1175/2009JTECHA1254.1>, 2009.
- Veselovskii, I., Whiteman, D. N., Korenskiy, M., Kolgotin, A., Dubovik, O., Perez-Ramirez, D., and Suvorina, A.: Retrieval of spatio-temporal distributions of particle parameters from multiwavelength lidar measurements using the linear estimation technique and comparison with AERONET, *Atmos Meas Tech*, 6, <https://doi.org/10.5194/amt-6-2671-2013>, 2013.
- 970 Veselovskii, I., Whiteman, D. N., Korenskiy, M., Suvorina, A., and Perez-Ramirez, D.: Use of rotational Raman measurements in multiwavelength aerosol lidar for evaluation of particle backscattering and extinction, *Atmos Meas Tech*, 8, <https://doi.org/10.5194/amt-8-4111-2015>, 2015.
- 975 Veselovskii, I., Hu, Q., Goloub, P., Podvin, T., Dubois, G., Kolgotin, A., and Korenskii, M.: Impact of water uptake on fluorescence of atmospheric aerosols: insights from Mie–Raman–fluorescence lidar measurements, *Atmos Meas Tech*, 18, 6039–6051, <https://doi.org/10.5194/amt-18-6039-2025>, 2025.
- Whiteman, D. N.: Examination of the traditional Raman lidar technique I Evaluating the temperature-dependent lidar equations, *Appl Opt*, 42, <https://doi.org/10.1364/ao.42.002571>, 2003.



- 980 Whiteman, D. N., Melfi, S. H., and Ferrare, R. A.: Raman lidar system for the measurement of water vapor and aerosols in the Earth's Atmos,
Appl Opt, 31, <https://doi.org/10.1364/ao.31.003068>, 1992.
- Whiteman, D. N., Demoz, B., Girolamo, P. D., Comer, J., Veselovskii, I., Evans, K., Wang, Z., Cadirola, M., Rush, K., Schwemmer, G.,
Gentry, B., Melfi, S. H., Mielke, B., Venable, D., and van Hove, T.: Raman lidar measurements during the International H2O Project. Part
I: Instrumentation and analysis techniques, J Atmos Ocean Technol, 23, <https://doi.org/10.1175/JTECH1838.1>, 2006.
- 985 WMO: Multilingual List of Terms used in Weather and Sea Bulletins, WMO-No. 471, World Meteorological Organization, Geneva, Switzer-
land, 6th edition edn., https://library.wmo.int/doc_num.php?explnum_id=3416, 2017.
- Wulfmeyer, V. and Feingold, G.: On the relationship between relative humidity and particle backscattering coefficient in the marine boundary
layer determined with differential absorption lidar, J Geophys Res Atmos, 105, <https://doi.org/10.1029/1999jd901030>, 2000.
- Yau, M. K. and Rogers, R. R.: A short course in cloud physics, Elsevier, 1996.
- 990 Yuan, L. and Zhao, C.: Quantifying particle-to-particle heterogeneity in aerosol hygroscopicity, Atmos Chem Phys, 23, 3195–3205,
<https://doi.org/10.5194/acp-23-3195-2023>, 2023.
- Zanatta, M., Gysel, M., Bukowiecki, N., Müller, T., Weingartner, E., Areskoug, H., Fiebig, M., Yttri, K. E., Mihalopoulos, N., Kouvarakis, G.,
Beddows, D., Harrison, R. M., Cavalli, F., Putaud, J. P., Spindler, G., Wiedensohler, A., Alastuey, A., Pandolfi, M., Sellegri, K., Swietlicki,
E., Jaffrezo, J. L., Baltensperger, U., and Laj, P.: A European aerosol phenomenology-5: Climatology of black carbon optical properties
995 at 9 regional background sites across Europe, Atmospheric Environment, 145, <https://doi.org/10.1016/j.atmosenv.2016.09.035>, 2016.
- Zhang, J., Wang, Y., Xu, J., Zuo, X., Lu, C., Zhu, B., Yang, Y., Yan, X., and Sun, Y.: Measurement report: Size-resolved and seasonal
variations in aerosol hygroscopicity dominated by organic formation and aging: insights from a year-long observation in Nanjing, Atmos
Chem Phys, 25, 17 413–17 428, <https://doi.org/10.5194/acp-25-17413-2025>, 2025.
- Zieger, P., Fierz-Schmidhauser, R., Weingartner, E., and Baltensperger, U.: Effects of relative humidity on aerosol light scattering: Results
1000 from different European sites, Atmos Chem Phys, 13, <https://doi.org/10.5194/acp-13-10609-2013>, 2013.
- Zieger, P., Fierz-Schmidhauser, R., Poulain, L., Müller, T., Birmili, W., Spindler, G., Wiedensohler, A., Baltensperger, U., and Weingartner,
E.: Influence of water uptake on the aerosol particle light scattering coefficients of the Central European aerosol, Tellus B Chem Phys
Meteorol., 66, <https://doi.org/10.3402/tellusb.v66.22716>, 2014.
- Ångström, A.: On the Atmospheric Transmission of Sun Radiation and on Dust in the Air, Geogr Ann Ser B, 11,
1005 <https://doi.org/10.2307/519399>, 1929.