

Riukojietna, a small low-altitude ice cap that may have persisted through the Holocene: Evidence from combining cosmogenic multi-nuclide dating and lacustrine sediment records

Arjen P. Stroeven^{1,2}, Gunhild C. Rosqvist^{1,2}, Alexandria J. Koester³, Jane L. Andersen^{1,4}, Carl-Anton Wahlström⁵, Nathaniel A. Lifton^{3,6}

¹ Department of Physical Geography, Stockholm University, 106 91 Stockholm, Sweden

² Bolin Centre for Climate Research, Stockholm University, 106 91 Stockholm, Sweden

³ Department of Earth, Atmospheric, and Planetary Sciences, Purdue University, West Lafayette IN 47907, USA

⁴ Research Centre for the Built Environment, Energy, Climate, and Water Technology, VIA University College, 8700 Horsens, Denmark

⁵ IKNI AS, PO Box 6646, 0129 Oslo, Norway

⁶ Purdue Rare Isotope Measurement Laboratory (PRIME Lab), Department of Physics and Astronomy, Purdue University, West Lafayette IN 47907, USA

Correspondence to: Arjen P. Stroeven (arjen.stroeven@natgeo.su.se)

Abstract. Riukojietna, a small, low-altitude, low-gradient plateau ice cap in northern Sweden, has been retreating rapidly over at least the last century. Its low surface gradient implies that it should be quite sensitive to, and therefore a potentially valuable indicator of, climate change since regional deglaciation at 9.8 ka. Here, we assess its former extent and activity by combining cosmogenic nuclide measurements in bedrock (*in situ* ¹⁴C, ¹⁰Be, and ²⁶Al) that constrain ice-free and ice-buried conditions with indirect evidence of glacial activity from proglacial lake sediment records, complemented by historical ice thickness reconstructions. These data are the basis for subsequent forward modeling of measured cosmogenic nuclide concentrations to constrain the Holocene history of Riukojietna.

The ice cap has an outlet glacier tongue that drains to the northeast, with a bouldery moraine deposit further down valley constraining its extent at the end of the Little Ice Age (LIA, ca. 1910 CE). Five cosmogenic nuclide samples were collected: two from bedrock on the plateau adjacent to the ice cap, two from a bedrock knob protruding from the outlet glacier tongue (exposed in 2011), and one from an outcrop embedded within the LIA moraine at the outlet of the most proximal of a series of four proglacial lakes. The latter sample yielded concentrations of ¹⁴C, ¹⁰Be, and ²⁶Al consistent with continuous exposure since 8.1 ± 0.1 ka (weighted mean). Nuclide measurements in the other four samples indicate complex exposure/burial histories. Lake cores from Pajep Luoktejaure, the third of the four down-valley proglacial lakes, indicate up to three periods of glacial sediment deposition >8.1 ka, from 5.4–5.0 ka, and after 1.8 ka, with intervening gyttja that indicates minimal or no glacial influence, with radiocarbon age constraints from bulk sediment and plant macrofossils.

We perform a forward modeling exercise to determine whether the cosmogenic-nuclide concentrations in the recently exposed bedrock samples are consistent with the glacial history inferred from the lake sediment record and the deglaciation age of 9.8 ka. Riukojietna persisted during the Holocene Thermal Maximum (ca. 8-5 ka), in contrast to earlier suggestions that Scandinavian glaciers vanished during the Holocene, as a result of an inferred increase in precipitation due to atmospheric circulation changes. The glacier has been in a retracted state similar or smaller than today during the late Holocene, as climate grew colder and drier. This approach combining short- and long-lived cosmogenic nuclides with lake sediments can thus provide new constraints on high-latitude Holocene glacial and paleoclimate history.

1 Introduction and Background

Mountain glaciers and ice caps are reliable indicators of regional climate change on decadal timescales because their mass balances are sensitive to changes in winter precipitation and summer ablation (Oerlemans and Fortuin, 1992; Oerlemans, 2005; Andreassen et al., 2020; Hugonnet et al., 2021; Rounce et al., 2023). Indeed, worldwide, glaciers have retreated at accelerating rates during the 20th century (Dussaillant et al., 2025; WGMS, 2025) and are projected for continued decline throughout the 21st century due to increased global temperatures (Oerlemans et al., 1998; Hock et al., 2019). Within this observed contemporary framework, this study explores the response of a small low-altitude Swedish ice cap, Riukojietna, to Holocene climate change.

A popular method to reconstruct the former extent of a glacier or ice sheet relies on exposure dating with cosmogenic nuclides in bedrock or erratics (e.g., Dunai, 2010). In glacial landscapes, the inventory of cosmogenic nuclides such as ¹⁰Be ($t_{1/2} = 1.39$ Myr; Chmeleff et al., 2010; Korschinek et al., 2010) and ²⁶Al ($t_{1/2} = 705$ kyr; Nishiizumi, 2004) in rock surfaces is governed by complex exposure, erosion, and ice burial histories (Gosse and Phillips, 2001; Fabel et al., 2002; Stroeve et al., 2002). When a surface is buried by >30 m of ice, production is reduced to <1% of that at the surface, and radionuclides, if present, will decay at known rates to concentrations supported by production at those depths. It is not possible to resolve complex glacial histories of ice burial and exposure arising from late Pleistocene and Holocene glacier fluctuations by pairing long-lived nuclides such as ¹⁰Be and ²⁶Al since neither will decay significantly in that time frame. However, pairing a long-lived nuclide with short-lived *in situ* cosmogenic ¹⁴C (*in situ* ¹⁴C, $t_{1/2} = 5.7$ kyr) enables such complex histories to be constrained. This has been demonstrated in the European Alps (Goehring et al., 2011; Wirsig et al., 2016; Schimmelpfennig et al., 2022), on Greenland and Baffin Island (Briner et al., 2014; Young et al., 2021), in Antarctica (Johnson et al., 2019; Nichols et al., 2019; Balco et al., 2023), and in Norway (Rand and Goehring, 2019).

Another common method to reconstruct past glacier extent and activity is to study proglacial lake sediment records (Karlén, 1988; Nesje et al., 2000; Rosqvist et al., 2004; Nielsen et al., 2016; Aa and Sønstegaard, 2019; Røthe et al., 2019). Typically, glacial sediments in a lake record (glacial flour) indicate an active glacier within the lake catchment while a prevalence of organic-rich sediments (gyttja) indicates deposition during a period of glacial inactivity or absence (Jansson et al., 2005). This is because basal sliding by a warm-based glacier is required for a glacier to erode its substrate and produce glacial flour. However, several other factors influence the sedimentation rate in proglacial lakes, such as hydrological regime, intermediate sediment storage capacity, and the activity of other geomorphological processes within the catchment (Leonard, 1986; Rubensdotter and Rosqvist, 2003; Jansson et al., 2005). In a setting with a sequence of lakes acting as sediment traps, glacial lake sedimentation may constrain the ice extent (Jansson et al., 2005; Nesje, 2009).

Climate reconstructions based on vegetation development and lake sediment proxy records indicate that the climate in northern Sweden warmed rapidly and that conditions were relatively wet (Shemesh et al., 2001; Sjögren and Damm, 2019) when the Fennoscandian Ice Sheet (FIS) melted (Kullman and Öberg, 2015; Sjögren, 2021). Even warmer but also drier summer conditions characterize the so-called Holocene Thermal Maximum (HTM; ~8 – 5 ka; Barnekow, 2000; Seppä et al., 2005; Sjögren, 2021; Wastegård, 2022). Based on sediment stratigraphies in proglacial lakes, Karlén (1976, 1981, 1988) proposed that some Swedish glaciers advanced during this time while others may have melted away, in line with records from some Norwegian glaciers (Bjune et al., 2005; Nesje et al., 2008; Nesje, 2009). Proglacial lake records show that glacier activity generally increased in response to an overall cooling trend after 5 ka, a progression that is documented across the Arctic (Karlén et al., 1995; Matthews and Dresser, 2008; Kaufman et al., 2009; Larocca and Axford, 2022). The latest period of glacier reactivation was during the Little Ice Age (1500-1900 CE), and evidence for this is rich compared to earlier periods (Grove, 2004; Matthews and Briffa, 2005).

Based on lichenometric dating of end moraine ridges, Karlén (1988) inferred that most Swedish glaciers reached their maximum Holocene position sometime during the 18th century and, supported by the oldest photographic evidence (Svenonius, 1910), again as late as ca. 1910 CE. Geomorphic evidence of earlier Holocene advances was potentially eradicated during this time (Jansson et al., 2005). Sediments from proglacial lakes therefore constitute the best continuous archive of Holocene glacier activity.

Here, we aim to constrain the former size and activity of Riukojietna (Sámi: Rivgojehkki), a rapidly retreating ice cap located in northern Sweden (Fig. 1). We do this by combining evidence for ice-free conditions and ice-burial durations from cosmogenic nuclide chronometry (*in situ* ^{14}C , ^{10}Be , ^{26}Al) and evidence of glacial extent and activity derived from proglacial lacustrine records, complemented with ice-

cap reconstructions based on historical records. This allows us to present the first full Holocene glacial history of a Swedish glacier based on multi-proxy records, and evaluate its sensitivity to Holocene climate change.




Figure 1: (a) Map showing deglacial isochrons of the Fennoscandian Ice Sheet (Stroeven et al., 2016) in 100-year intervals around the location of Riukojietna (box with white dot: panel b). (b) European Union Digital Elevation Model (EU-DEM) of northern Sweden/Norway depicting the location of glaciers, including Riukojietna (box: Fig. 2a). The numbers on the contours are deglacial isochrons in cal ka BP (Stroeven et al., 2016). Modified from Koester (2023).

2 Study site

Riukojietna is a small, thin (thickness <100 m), polythermal ice cap covering a plateau on the Swedish-Norwegian border (Fig. 2). This ice cap is ideal for studying Late Pleistocene to Holocene glacier fluctuations using cosmogenic nuclides and lake sediments for two reasons. First, the ice cap rests on quartz-rich bedrock of the Köli Nappe which allows for the application of multiple nuclides, including ^{10}Be , ^{26}Al , and *in situ* ^{14}C . In contrast, most glaciers in northern Sweden are situated in the higher alpine areas where amphibolite and metadolerite bedrock of the Seve Nappe dominates (Andréasson and Gee, 1989; Bergman et al., 2012). Second, a thin and blocky till cover, gentle slopes, and an absence of sediment-rich landforms in the catchment implies that glacial flour is the dominant source of minerogenic material in downstream lakes. As such, we expect lake records from the catchment to primarily reflect fluctuations in glacier activity and extent.




Figure 2: (a) Google Earth satellite image (2018; Map data: Google, © 2025 CNES / Airbus) of the Riukojietna ice cap depicting the location of bedrock samples (red circles). Simple exposure ages with internal uncertainties (in ka) are shown in white boxes. Thick black lines depict the location of Late Holocene moraines. White stars depict lakes that have been cored downstream of the ice cap. (b) Two bedrock samples collected directly adjacent to the ice cap (1290 m a.s.l.). (c) Two bedrock samples collected from a bedrock knob that was exposed in 2011 (1240 m a.s.l.). (d) One bedrock sample collected adjacent to Lake 1063 outlet (1064 m a.s.l.). Modified from Koester (2023).

Riukojietna had an area of 5.5 km² and 4.6 km² when surveyed from 1960 and 1978 aerial photographs, respectively (Rosqvist and Østrem, 1989; Table 1). However, the ice cap has continued to shrink, covering a mere 2.8 km² and thinned to span only 283 m vertically between the ice divide and the snout in 2015 (Table 1). The ice cap flows radially outwards from its highest point and is impounded by bedrock ridges to the south and northeast. Currently, Riukojietna has one outlet glacier tongue that terminates on a steep rise to the northeast (Fig. 2). The position of this tongue has been mapped since 1963 and the glacier mass balance has been monitored annually since 1986 revealing high sensitivity to changes in summer temperature and winter precipitation (WGMS, 2025). Two topographic maps were constructed based on aerial photography from 1960 and 1978 showing ice cap surface topography and extent (Rosqvist and Østrem, 1989). The eastern terminus calved into Lake 1063 (Fig. 2) before 1960 but retreated up-valley after 1975. Ice volume has been decreasing significantly during the 20th century (Table 1) due to a

dominantly negative annual mass balance (WGMS, 2025). Because it is relatively low-elevation and spans an unusually narrow elevation interval compared to most valley glaciers in the region, Riukojietna is regarded to be highly sensitive to climate change (Rosqvist and Østrem, 1989).

Four lakes on the eastern side of the ice cap act as a series of traps for sediment derived from glacial erosion. Ordered by decreasing elevation (in m a.s.l.), they are lakes 1063, 1009, Pajep Luoktejaure (913; where “Pajep” means upper and “jaure” means lake— so Upper Luokte lake), and Vuolep Luoktejaure (903) (or Lower Luokte lake, which is only partially visible in Fig. 2). Lakes at this latitude and altitude are generally ice-covered from October to June (Rosqvist et al., 2004). From the variations in relative densities and organic content in sediment cores from Lake 1009 and Pajep Luoktejaure, Karlén (1981, 1988) proposed that Riukojietna was inactive and small, or disappeared, between 10.9 ± 0.4 (Si-2860) and 2.7 ± 0.1 (Si-2859) cal ka BP (re-calibrated bulk sediment samples using the online CALIB Radiocarbon Calibration v 8.2; Stuiver and Reimer, 1993; Reimer et al., 2020) and that the ice cap subsequently reactivated.

The relatively prominent lateral/end moraine sequence consisting of large boulders located to the northeast and east of Lake 1063 (Fig. 2) indicates that the ice cap advanced to this position at least once during the Holocene. A depositional age of c. 1910 CE was inferred for this ridge (Pohjola et al., 2005) based on lichenometry of *Rhizocarpon geographicum* and *Rhizocarpon alpicola* (Karlén, 1973), which is similar to other lichenometry-dated Little Ice Age (LIA) moraines in the region (Karlén, 1976; Karlén and Denton, 1976). A less prominent moraine ridge indicates that the glacier once extended down the southwestern slope of the plateau (Fig. 2). A glacially molded plateau surface extends from the current ice cap margin towards the southeast, indicating a possible earlier extension direction of the ice cap towards Lake 1009 (Koester, 2023).

Table 1: Riukojietna area, thickness, and volume reconstructions since the LIA (Wahlström, 2016).

Year (CE)	1910	1960	1978	2015
Area (km ²)	6.7	5.5	4.6	2.8
Volume (km ³)	0.36	0.26	0.22	0.13
Min altitude (m a.s.l.)	1058	1058	1122	1146
Max altitude (m a.s.l.)	1488	1468	1459	1429
Elevation span (m)	430	410	337	283
Average thickness (m)	54	47	47	46
Volume decrease in % (from 1910 CE)	--	28	39	64

169

170 **3 Methods**

171 We combine cosmogenic nuclide chronometry, lacustrine sediment records, historic ice cap
172 reconstructions, and modeling of complex exposure histories to evaluate the Holocene history of
173 Riukojietna (Koester, 2023). Each method is described in detail below.

174 **3.1 Cosmogenic nuclide sampling and analysis**

175 We collected five granitic bedrock samples with angle grinder, hammer, and chisel from three locations
176 near Riukojietna in August 2016 and measured concentrations of *in situ* ^{10}Be , ^{26}Al , and ^{14}C (Fig. 2, Table
177 2). Two samples were collected from the base of a small bedrock ridge immediately adjacent to the ice
178 cap (<2 m from the modern ice; Riuko-16-001 and 16-002). Two samples were taken from a bedrock
179 knob protruding through the outlet glacier tongue that became exposed in 2011 (Riuko-16-003 and 16-
180 004). In the following, we collectively refer to these two sites as ‘ice-marginal bedrock’. The last sample
181 was collected from a bedrock outcrop adjacent to the outlet of Lake 1063 (Riuko-16-005), in the
182 following referred to as the ‘lake outlet sample’. The bedrock outcrops in line with the LIA moraine and
183 constitutes the subglacial meltwater outlet of Riukojietna at that time.

184 We separated quartz at the Purdue Rare Isotope Measurement Laboratory (PRIME Lab) using standard
185 mineral separation procedures ([https://www.physics.purdue.edu/primelab/labs/mineral-separation-](https://www.physics.purdue.edu/primelab/labs/mineral-separation-lab/procedure.php)
186 [lab/procedure.php](https://www.physics.purdue.edu/primelab/labs/mineral-separation-lab/procedure.php)). Samples were crushed, sieved to 250-500 μm , and magnetic minerals removed using
187 a Carpco magnetic separator. Micas and feldspars were removed with froth flotation, and the remaining
188 quartz separate was subsequently leached in weak hydrofluoric (HF) and nitric acids (HNO_3) to remove
189 meteoric ^{10}Be (Kohl and Nishiizumi, 1992). Quartz purity was assessed with inductively coupled plasma–
190 optical emission spectrometry (ICP-OES) at PRIME Lab (Al target: <200 ppm). Riuko-16-002 and
191 Riuko-16-005 had high Al and Na contents, indicating the presence of feldspars, and were therefore
192 subjected to an additional separation step using heavy liquids and etching in weak acids. We have never
193 observed any demonstrable effects from froth flotation on PRIME Lab *in situ* ^{14}C measurements using
194 these quartz purification procedures, thus no additional steps were taken (e.g., Nichols and Goehring,
195 2019).

196 Beryllium and aluminum were extracted at PRIME Lab following the procedures outlined in Andersen et
197 al. (2020). Beryllium samples were spiked with ~0.26 mg of Be carrier and aluminum samples were
198 spiked with 0.9-1.3 mg of Al carrier if they didn’t contain enough native Al before digestion in HF (target
199 ~1.5 mg Al). The samples were prepared in one batch with a procedural blank and an aliquot of the

intercomparison material CoQtz-N (Binnie et al., 2019). Be and Al were separated through anion and cation column chromatography, precipitated, oxidized, mixed with niobium powder, and pressed into cathodes for measurement. Isotopic ratios ($^{10}\text{Be}/^9\text{Be}$ and $^{26}\text{Al}/^{27}\text{Al}$) were measured by accelerator mass spectrometry (AMS) at PRIME Lab.

Carbon-14 was extracted at PRIME Lab using the automated Carbon Extraction and Graphitization System described by Lifton et al. (2023) and following procedures modified from that publication. Twenty grams of lithium metaborate (LiBO_2) flux is first degassed for one hour at 1100 °C in a reusable Pt-Rh boat (90% Pt, 10% Rh) in ~ 6.67 kPa of Research Purity O_2 , allowed to cool overnight under vacuum, and then approximately 5 g of sample quartz is evenly distributed over the solidified LiBO_2 . The sample is then combusted at 500 °C for one hour in ~ 6.67 kPa of Research Purity O_2 to remove atmospheric/organic contaminants, and the system is evacuated. Approximately 6.67 kPa of Research Purity O_2 is then added to the system, and the sample is heated to 1100 °C for three hours to release any trapped carbon species. All evolved C species are oxidized to CO_2 by passing the process gas over 2 mm quartz beads at ~ 950 °C. After extraction, the evolved gas is collected in a coil trap cooled with liquid nitrogen, purified using a variable temperature trap at -145 °C and passed over Cu mesh and Ag wool held at 600 °C to remove contaminants, before volume measurement and dilution with ^{14}C -free CO_2 to the equivalent of ~ 300 μg C. Finally, a ca. 9 μg C split is collected for stable C isotopic measurement, and the remaining sample is reduced to graphite in Research Purity H_2 over an Fe catalyst before packing into a cathode for AMS measurement (e.g., Santos et al., 2004, 2007).

Sample $^{14}\text{C}/^{13}\text{C}$ ratios were measured by AMS relative to Oxalic Acid II (NIST-4990C) at Purdue University. Stable carbon isotopic ratios were measured at the University of California Davis Stable Isotope Facility using isotope ratio mass spectrometry (e.g., Lifton et al., 2023). The ^{14}C concentration is calculated from the measured $^{14}\text{C}/\text{C}_{\text{total}}$ after subtracting out representative procedural background ^{14}C (Hippe and Lifton, 2014).

252

253

Table 2: Sample information and measured cosmogenic nuclide concentrations.

Sample ID	Latitude (°N)	Longitude (°E)	Elevation (m a.s.l.)	Sample thickness (cm)	Density (g cm ⁻³)	Shielding	[¹⁰ Be] ($\times 10^5$ at g ⁻¹)	[²⁶ Al] ($\times 10^6$ at g ⁻¹)	[¹⁴ C] ($\times 10^5$ at g ⁻¹)
16-001	68.08005	18.07339	1291	1.5	2.65	0.99	1.8107 ± 0.0452	1.2158 ± 0.0549	1.9242 ± 0.0820
16-002	68.08013	18.07315	1288	1.2	2.65	0.99	2.2140 ± 0.0534	1.5105 ± 0.0538	2.0705 ± 0.0856
16-003	68.08749	18.06473	1238	1.5	2.65	0.99	0.8825 ± 0.0442	0.6391 ± 0.0278	1.4580 ± 0.0815
16-004	68.08751	18.06409	1242	1.3	2.65	0.99	0.9251 ± 0.0590	0.6834 ± 0.0431	1.3336 ± 0.0800
16-005	68.08550	18.09376	1064	1.4	2.65	1.00	0.9918 ± 0.0531	0.7170 ± 0.0288	2.2698 ± 0.0830

We use the primary CRONUS-Earth global calibration spallation production rates for ^{10}Be (3.9 ± 0.3 at $\text{g}^{-1} \text{yr}^{-1}$), ^{26}Al (28.5 ± 3.1 at $\text{g}^{-1} \text{yr}^{-1}$) and *in situ* ^{14}C (12.8 ± 0.9 at $\text{g}^{-1} \text{yr}^{-1}$) to calculate exposure ages and model ice burial and erosion histories (Lifton et al., 2014, 2016; Borchers et al., 2016; Phillips et al., 2016). The *in situ* ^{14}C production rate also includes the Young et al. (2014) *in situ* ^{14}C calibration dataset (Koester and Lifton, 2023, 2024). All calculations utilize nuclide-specific LSDn scaling (Lifton et al., 2014, 2016) and the muon production formulation from Balco (2017). Simple exposure ages (i.e., assuming continuous exposure with no erosion or burial) and 1σ uncertainties were calculated using the online University of Washington cosmogenic calculator, v.3 (wrapper script 3.0.2, constants: 2024-08-26, muons: 1A; Balco et al., 2008; <https://hess.ess.washington.edu/math/>), with *in situ* ^{14}C production rates calibrated using the above datasets.

3.2 Lacustrine sediment cores

Pajep Luoktejaure has a relatively even bathymetry with one deeper section (12 m) in its northern part. We retrieved a deep core (1995) and five surface and deep cores (PL1 – PL5) from the deepest part of the lake in April 1995 and April 1998, respectively. We used a gravity corer to retrieve an undisturbed surface sample (top 25 cm; PL3) and a modified Livingstone piston corer (90 mm diameter) for the others (1995; PL1, 2, 4, and 5). Visual inspection and results from measurements of loss-on-ignition (LOI), determined at 550 °C with 1 cm resolution, for PL1 – PL4 show the same general stratigraphy for common sections (PL5 remains unopened, for reference). PL1-169 is the longest core (down to 169 cm below the lake floor) and the only one representing the early phase of lake development. The composite stratigraphic description of Pajep Luoktejaure is based on this core and surface core PL3-25.

Sedimentary structures and grayscale densities of cores PL1-169 and PL3-25 were determined by X-ray radiography (Rubensdotter and Rosqvist, 2003; Rosqvist et al., 2004). Line transects (1 mm width) of grayscale density values were taken at a resolution of 0.3 mm. In this study, grayscale density is merely used to identify areas of lamination. Grain size distribution was analyzed in a SediGraph 5100 on 1 cm slices of specific sedimentary structures (i.e., finely laminated minerogenic sections at 115-117 cm and 8-10 cm depths in core PL1-169, and replicated by results from finely laminated minerogenic sections at 128-131 cm and 6-9 cm depths in core PL2-133; Fig. S1, Table S1).

Age control is provided by four samples from Pajep Luoktejaure; three AMS ^{14}C -dates from core PL1-169 and one from near-surface sediments in the core of 1995 (Table 3). We also used a basal date derived from a terrestrial plant macrofossil from nearby lake Vuolep Allakasjaure to help constrain the timing of deglaciation (Table 3). The samples were radiocarbon dated at the Ångström Laboratory, Uppsala University, and converted to calendar ages using the IntCal20 calibration curve (Reimer et al., 2020).

Three of the radiocarbon constraints were used to derive an age–depth model using the Bayesian accumulation age-modelling software of Blaauw and Christen (2011; Fig. S2).

Table 3: Radiocarbon chronology for Pajep Luoktejaure. Three radiocarbon dates were derived from core PL1-169 at 116 cm depth (1 cm bulk sample), 90 cm depth (terrestrial macrofossil), and 9 cm depth (1 cm bulk sample). The latter replicated the date from a 12 cm depth terrestrial macrofossil sample of the same stratigraphic unit in the 1995 core from Pajep Luoktejaure. The basal bulk sediment age was replicated with a terrestrial macrofossil date from the same stratigraphic position (first radiocarbon accumulation after deglaciation) from nearby lake Vuolep Allakasjaure (12 km distance; Rosqvist et al., 2004; Fig. 1b). Calibrated ages derived using CALIB Radiocarbon Calibration v 8.2 (<http://calib.org/calib/>; Stuiver and Reimer, 1993; Reimer et al., 2020).

Sample Name	Depth (cm)	Material	¹⁴ C Age (yr BP)	δ ¹³ C (‰ VPDB*)	Calibrated Age (cal ka BP)
<i>Pajep Luoktejaure</i>					
Ua-14024	9	bulk sediment	1920 ± 70	-27.1	1.8 ± 0.1
Ua-14025	90	macrofossil	4024 ± 85	-27.5	4.5 ± 0.1
Ua-14026	116	bulk sediment	9650 ± 160	-26.9	11.0 ± 0.1
Ua-4751	12	macrofossil	1896 ± 70	-26.2	1.8 ± 0.1
<i>Vuolep Allakasjaure (Rosqvist et al., 2004)</i>					
Ua-16628	155	macrofossil	8740 ± 100	-23.0	9.8 ± 0.2

* Measured relative to the Vienna Pee Dee Belemnite standard

3.3 Modern ice thickness reconstructions, bed topography, and volume calculations

Detailed ice surface topographic maps were constructed from aerial photographs taken in 1960 and 1978 to determine the average mass balance of Riukojietna (Rosqvist and Østrem, 1989). These two maps, reproduced at 1:10 000 in the back sleeve of *Geogr. Ann.* 71A (1-2), were scanned and georeferenced in QGIS software to derive the ice-cap surface elevation during these years. The ice-cap topography was additionally mapped during the summer of 2014 and spring of 2015 using two differential GPS (dGPS) rover units. The subglacial bedrock topography was constrained with ground penetrating radar (GPR) towed behind a snowmobile in the springs of 2011, 2012, and 2015 (Fig. S3). We use these data sources to track ice cap area and volume changes relative to a 2-m-resolution LiDAR digital elevation model (DEM) from 2015 (Fig. S3).

The elevations of fixed bedrock points on the 1960 and 1978 maps were compared to the recent LiDAR DEM to evaluate their accuracies, and differences were calculated. These differences were interpolated using kriging in Surfer 11 to create misfit values across the glacier surface. The resulting spatial pattern of

misfit is used to correct the altitudes derived from the 1960 and 1978 maps to better match ice surface elevations to the 2015 LiDAR DEM (Fig. S4).

The perimeter of the presumed LIA glacial extent (1910 CE) was mapped from the 1960 and 1978 orthophotos and integrates the location of the eastern terminal moraine. As there are no moraines limiting previous extents on the western side, we assumed a 50 – 100 m more expanded glacier relative to its extent in 1960. The 1910 CE ice surface was calculated for the central flow line (Fig. S5) using the reconstructed bed topography, a target maximum elevation of 1490 m a.s.l. based on a summit ice thickness of c. 93 m estimated by Pohjola et al. (2005), and an adjustable yield stress following Benn and Hulton (2010). The surface elevations were interpolated in Surfer 11 using kriging to create an LIA ice surface reconstruction. For each of the four time slices, Surfer 11 was used to calculate the volume and average ice thickness of Riukojietna using the difference between the ice surface DEM and the bed topography DEM (Table 1).

3.4 Modeling complex exposure histories

We explore potential effects of temporal variations in ice cap thickness on our ice-marginal bedrock cosmogenic nuclide inventories (Riuko-16-001 to 16-004) using a Lagrangian forward modeling framework (e.g., Knudsen et al., 2019) implemented in MATLAB (code reference – to be added before publication). This model predicts the post-LGM time evolution of ^{10}Be , ^{26}Al , and ^{14}C concentrations in bedrock during subaerial exposure and when either completely (ice > 30 m thick) or incompletely shielded beneath thin ice (< 30 m thick) from both spallogenic and muogenic production, while allowing for subglacial erosion. Modeling details are provided in the Supplement text S1-S3.

4 Results

4.1 Cosmogenic nuclide concentrations and simple exposure ages

The ^{10}Be concentrations in bedrock samples Riuko-16-001 to 16-005 range from $8.8 \pm 0.4 \times 10^4$ to $2.2 \pm 0.1 \times 10^5$ atoms g^{-1} , while the ^{26}Al concentrations range from $6.4 \pm 0.3 \times 10^5$ to $1.5 \pm 0.1 \times 10^6$ atoms g^{-1} (Tables 2, S2; Koester, 2023). Measured *in situ* ^{14}C concentrations range from $1.3 \pm 0.1 \times 10^5$ to $2.3 \pm 0.1 \times 10^5$ atoms g^{-1} (Tables 2, S3; Koester, 2023). The corresponding simple exposure ages range between 3.2 ± 0.2 ka and 8.2 ± 0.5 ka for *in situ* ^{14}C , 6.3 ± 0.3 ka and 14.3 ± 0.5 ka for ^{26}Al , and 6.1 ± 0.3 ka and 14.7 ± 0.4 ka for ^{10}Be (Fig. 2; Tables S2-S4). When plotted on a two-isotope diagram, the ^{10}Be - ^{26}Al results of all five bedrock samples overlap with the simple exposure line at 1σ (Fig. 3a). In contrast, the ^{14}C - ^{10}Be two-isotope diagram shows that only sample Riuko-16-005 overlaps the simple exposure line, indicating continuous exposure since deglaciation for that sample (Fig. 3b). All simple exposure ages for this sample

agree within 1σ external (and internal) uncertainty (Fig. 2). Using external uncertainties, the weighted mean and the larger of either the standard error in inverse-error-weighted mean, or the inverse-error-weighted average variance (Bevington and Robinson, 2003) yields 8.0 ± 0.1 ka, while the straight mean and standard deviation yields 8.1 ± 0.1 ka. As these values are not significantly different, we take the straight mean ^{14}C - ^{10}Be - ^{26}Al exposure age of 8.1 ± 0.1 ka (1σ) as the deglacial age for that location (Koester, 2023).

The remaining samples (Riuko-16-001 to 16-004) plot in the complex exposure field of the ^{14}C - ^{10}Be two-isotope diagram (Fig. 3b). Because simple ^{10}Be and ^{26}Al exposure ages for the same samples overlap within 1σ , we focus dominantly on the ^{10}Be and ^{14}C values in the remainder of the paper. The highest elevation site, adjacent to the current ice cap margin (~ 1290 m a.s.l.; Riuko-16-001 and 16-002), has the oldest simple ^{10}Be (^{14}C) exposure ages of 12.0 ± 0.3 (4.9 ± 0.3) and 14.7 ± 0.4 (5.4 ± 0.3) ka, respectively (Fig. 2). The two samples collected on the recently emerged bedrock knob at ~ 1240 m a.s.l. (Riuko-16-003 and 16-004) have younger ^{10}Be (^{14}C) simple exposure ages of 6.1 ± 0.3 (3.5 ± 0.2) and 6.4 ± 0.4 (3.2 ± 0.2) ka, respectively – internally consistent for each nuclide but disagreeing between nuclides (Koester, 2023).




Figure 3: (A) Two-isotope diagram of normalized $^{26}\text{Al}/^{10}\text{Be}$ concentration ratio vs. normalized ^{10}Be concentration. Ellipses show 2σ uncertainty. Normalization (indicated by *) is to site production rates, hence the unit of $[^{10}\text{Be}^*]$ is yr. (B) Two-isotope diagram of normalized in situ $^{14}\text{C}/^{10}\text{Be}$ concentration ratio vs. normalized ^{10}Be concentration. Burial isochrons (dashed lines) are calculated relative to the simple exposure curve. Burial isochron spacing is 500 kyr in A), and 2 kyr in B). Near-vertical dot-dashed lines are decay trajectories followed during burial, plotted in A) for every fifth age in a logarithmic array of 100 ages generated from 100 to 10^7 yr that lie within the x-axis limits, and in B) for every other age in a logarithmic array of 100 ages generated from 100 to 2×10^5 yr that lie within the x-axis limits.

368

369 4.2 Lacustrine records

370 The bottom of core PL1-169 from 169 to 116 cm depth is characterized by homogenous glacial silt and
371 clay (Fig. 4b) with LOI less than 2%. Between 116 and 107 cm depth, sediments are finely laminated silt
372 and clay with LOI less than 10%. Laminations become more diffuse between 107 and 95 cm depth and
373 LOI values increase, reaching as high as 16% at 98 cm depth. A drop in LOI to 7% at 93 cm is caused by
374 an influx of minerogenic matter. Homogenous gyttja with up to 26% LOI characterizes the sediments
375 between 90 and 10 cm depth. The uppermost 10 cm of PL1-169 consist of finely laminated silt and clay
376 with high density and low LOI (between 5% and 8%). Results from grain size distribution analyses show
377 indeed that the sampled sections consist of silt and minor amounts of clay (>85% silt/clay; i.e. <63 μm ;
378 Table S1).

379 Surface core PL3-25 records the intact uppermost 25 cm of the stratigraphy in Pajep Luoktejaure (Fig.
380 4a). Its sedimentary structure is similar to PL1-169 in the overlapping interval. Well-preserved
381 laminations are observed between 19 and 3 cm. The topmost 3 cm of the core displays high LOI (up to
382 17% at the surface). No laminations are detectable in these non-compacted organic surface sediments.
383




Figure 4: Stratigraphy and loss on ignition (LOI %) from (a) surface gravity core PL3-25 and (b) Livingstone piston core PL1-169 in Pajep Luoktejaure (Fig. 2a). High LOI indicates high organic content, interpreted as relatively low glacial activity, while low LOI indicates low organic content associated with a relatively active glacier and minerogenic influx. Core PL1-169 has 3 radiocarbon tie points (in cal ka BP; Table 3). (c) Composite core of LOI record versus age, where 1 – 13 cm are from PL3-25 and 13 – 130 cm are from PL1-169. Age vs. depth for PL1-169 below 130 cm is poorly constrained. (Fig. S2). In this conversion we replaced the basal bulk sediment age of 11.0 ± 0.1 cal ka BP with a terrestrial macrofossil date of 9.8 ± 0.2 cal ka BP from the same stratigraphic position (first radiocarbon accumulation after deglaciation) from nearby lake Vuolep Allakasjaure (Table 3). Blue fields indicate our inferred periods of glacier activity. The lowest field is informed by the cosmogenic nuclide retreat age of Riukojietna from the outlet of Lake 1063 at 8.1 ± 0.1 ka. The youngest period of inferred glacier activity is informed by the oldest ages of glacier reactivation of 1.8 ± 0.1 cal ka BP (on macrofossil and bulk sediment; Table 3) and final retreat from the LIA moraine at 1910 CE. During the mid-Holocene short period of glacier activity between 5.4 and 5.0 cal ka BP, LOI values are on-par with the previous and subsequent periods of inferred glacier activity.

Three radiocarbon dates were obtained from PL1-169, with the oldest (116 cm depth; 11.0 ± 0.1 cal ka BP; 2% LOI) and youngest (9 cm depth; 1.8 ± 0.1 cal ka BP; 14% LOI) ages derived from bulk sediment and the intermediate age (90 cm depth; 4.5 ± 0.1 cal ka BP) derived from a terrestrial plant macrofossil (Table 3). An additional radiocarbon date of a terrestrial macrofossil at 12 cm depth in the 1995 core, also representing the first deposition of late Holocene laminations in the core, replicates the bulk sediment age at 9 cm depth of 1.8 ± 0.1 cal ka BP (Table 3). The basal radiocarbon age, from bulk sediment with only 2% organic content, is much older than the expected regional ~ 9.8 cal ka BP deglaciation age (Fig. 1b) of

Stroeven et al. (2016). Therefore, we adopt the basal sediment age from nearby lake Vuolep Allakasjaure (Fig. 1b; ~12 km NE of Pajep Luoktejaure) of 9.8 ± 0.2 cal ka BP derived from a terrestrial macrofossil in the same stratigraphic position, that is, representing the first radiocarbon accumulation after deglaciation (Table 3; Rosqvist et al., 2004). Together with the two youngest radiocarbon ages (at 90 and 9 cm depth) and a sediment surface age of -48 BP (i.e., 1998 CE) they create the four tie points constraining the age-depth model (Fig. S2). To provide a composite record we spliced the core PL3-25 LOI record representing the top 13 cm of the sediment record into the PL1-169 LOI record (Fig. 4c). The average sedimentation rate of the upper laminated section is ~ 0.1 mm yr⁻¹ if the uppermost 3 cm corresponds to the time since the glacier retreated from the moraine at Lake 1063 at the end of the LIA. The sedimentation rate of the organic section (90 – 9 cm) is ~ 0.3 mm yr⁻¹.

4.3 Ice thickness and volume reconstructions

Ice thickness reconstructions for the past century are shown in Fig. 5. At the end of the LIA, Riukojietna covered an area of 6.7 km² and had an estimated volume of 0.36 km³ (Table 1). Ice extent and volume have decreased by >58% and >64%, respectively, between 1910 CE and 2015 (Table 1; Fig. 5). Based on these reconstructions, we estimate that the ice thickness above our two ice-marginal cosmogenic sample locations (Riuko-16-001 to 16-004) started at 33-35 m one century prior to exposure (Table S5; Koester, 2023).

5 Discussion

5.1 Constraints on the Holocene history of Riukojietna from the lake outlet sample

The cosmogenic nuclide results offer constraints on the dimensions, including the minimum thickness of Riukojietna, through the Holocene (Koester, 2023). The agreement between ¹⁴C-¹⁰Be-²⁶Al simple exposure ages at 1σ for the lake outlet sample (Riuko-16-005; 1064 m a.s.l.) indicates continuous exposure at that site since deglaciation following erosional resetting of the surface. Thus, this result establishes that Riukojietna did not cover or expand beyond the Lake 1063 outlet sampling site after 8.1 ± 0.1 ka (mean of all three nuclide ages; Fig. 2). The result also reveals that the FIS, or Riukojietna, eroded sufficient bedrock at that site to remove any cosmogenic nuclide inventory produced prior to the Last Glacial Maximum (LGM).




Figure 5: Ice thickness reconstructions for Riukojietna from 1910 CE, 1960, 1978, and 2015 with respect to the maximum extent of 1910 CE (LIA extent). The 40 m ice thickness contour is shown in dashed grey. Sample sites are shown as red dots. Figure modified from Wahlström (2016).

5.2 Constraints on the Holocene history of Riukojietna from lacustrine sediments

Pajep Luoktejaure receives meltwater coming from the northeastern active outlet of Riukojietna (Fig. 2). Lakes 1063 and 1009 in its drainage system act as sediment traps. This will have a damping effect on sedimentary responses in Pajep Luoktejaure to changes in the activity and extent of Riukojietna.

Meltwater from the southeastern tongue of Riukojietna also enters Lake 1009 before it reaches Pajep Luoktejaure (Fig. 2). However, based on thermistor measurements in 1986 (Pohjola et al., 2005), the ice cap dome, and by inference its thin low-gradient southeastern tongue, were shown to be cold-based, which means that sub-glacial erosion is an unimportant component to downstream sediment delivery from that location during such restricted ice configurations. Although the Pajep Luoktejaure record may reflect some sediment contribution from this tongue during extended ice cap configurations (e.g., LIA), we estimate that its contribution is minor based on the topographic configuration (faint valley development and steep slope towards Lake 1009) implying that the ice would remain thin even in a more advanced position.

The basal section of homogenous glacial silt and clay in Pajep Luoktejaure (169-116 cm depth; Fig. 4b) represents an ice-dammed lake sequence. We suggest that this lake was formed by the retreating FIS margin obstructing water flow out of the valley, similar to the many ephemeral ice-dammed lakes that it dammed immediately to the north and east of Riukojietna (Ploeg and Stroeve, 2025). The subsequent laminated sediments that began accumulating from ca. 9.8 ± 0.2 cal ka BP (age constraint from nearby lake Vuolep Allakasjaure), with increasing LOI, indicate that organic production and seasonal input of meltwater from Riukojietna dominated the depositional signal after the northern FIS margin retreated from the site (Fig. 1b; Stroeve et al., 2016). The ice cap continued to deliver sediments to Pajep Luoktejaure after 9.8 ka, forming the 9 cm-thick laminated section, indicating that the glacier remained in a relatively advanced position during the early Holocene. The following section with higher organic

content and diffuse laminations between 107 and 91 cm depth (Fig. 4b), represents a time interval when the ice cap probably retreated sufficiently to also expose the Lake 1063 sediment trap. Our cosmogenic nuclide results indicate that Riukojietna covered Lake 1063 until 8.1 ± 0.1 ka, which we adopt as the constraining age for the start of this section. The increase in glacial sediment input and lowering of LOI between 94 and 92 cm depth indicates that Riukojietna delivered more sediment to downstream lakes, signaling a higher activity between 5.4 and 5.0 cal ka BP (Fig. 4b, c).

Subsequently, laminations disappear altogether and organic productivity increases to maximum Holocene levels. Possible paleoglaciological interpretations of this sedimentary section include that Riukojietna retreated and thinned to a configuration where it was as small or smaller than today with cold-based conditions inhibiting sub-glacial erosion, or that the ice cap melted away completely. Finely laminated glacial sediments in the top sections of PL1-169 and PL3-25 indicate a reactivation and expansion of Riukojietna starting shortly after 1.8 ± 0.1 cal ka BP and culminating in 1910 CE. Finally, the decrease in glacial sediment input and deposition of organic sediments in the topmost 3 cm reflects the time since Riukojietna retreated from its LIA and post-glacial maximum position marked by the moraine ridge fringing Lake 1063.

In summary, we infer that Riukojietna existed as an ice cap, and was more active and extensive than today, between 9.8 cal ka BP and 8.1 ka, during a short interval just before 5.0 cal ka BP, and between 1.8 cal ka BP and 1910 CE. While these periods will determine the durations of sample burial in cosmogenic nuclide forward modeling in the next section, we acknowledge that the duration of burial during the mid Holocene, which is solely constrained by the lacustrine sediment record and radiocarbon dating, has the largest uncertainty due to a poor control on sediment accumulation rates.

5.3 Constraints on Holocene history of Riukojietna from cosmogenic nuclide modeling

While the lake sediments represent an indirect measure of the extent and state of Riukojietna, cosmogenic nuclide inventories in bedrock surfaces more directly reflect the extent and state of Riukojietna through ice burial durations and erosion histories at the sampled sites. Here we test the ice history derived from the proglacial lake record by calculating the resulting cosmogenic nuclide concentration in ice-marginal bedrock sites arising from exposure and burial after FIS retreat.

The two samples from the highest altitude site (Riuko-16-001 and 16-002) from the bedrock ridge adjacent to the ice cap margin plot significantly below the continuous exposure field in the ^{10}Be - ^{14}C two-isotope plot (Fig. 3b), indicating a complex exposure history. If one assumes glacial erosion to have completely removed any inherited component from exposure prior to the LGM, their position on this plot is consistent with a cumulative burial duration of 4 – 5 kyr. However, this simplest interpretation is

untenable because their simple ^{10}Be exposure ages (of 12-15 ka) are older than the regional deglaciation age of ~ 9.8 ka (Stroeven et al., 2016; Fig. 2, Table 3), requiring additional contributions to ^{10}Be inventory from pre-LGM exposure. Therefore, their position on the ^{10}Be - ^{14}C two-isotope plot is skewed by inherited ^{10}Be and inferred burial durations are, therefore, difficult to quantify but likely shorter.

Like the highest-altitude samples, the samples from the bedrock knob (Riuko-16-003 and 16-004) also plot in the complex exposure field of the ^{10}Be - ^{14}C two-isotope plot (Fig. 3b), consistent with a mean burial duration of 4 kyr in the simplest interpretation of continuous exposure followed by burial and exposure by recent ice retreat. However, it is important to note that their position is also consistent with continuous postglacial exposure and an inherited ^{10}Be component (smaller than that of Riuko-16-001 and 16-002) that would displace the position below the simple exposure curve.

Our reconstructed ice history for the ice-marginal bedrock sites (Riuko-16-001 to 16-004) relies on a set of assumptions based on derived cosmogenic nuclide concentrations, the composite lacustrine record, and a LIA ice-thickness reconstruction. First, we prescribe zero initial cosmogenic nuclide concentrations consistent with a complete removal of pre-LGM inventories and a thick (>50 m) ice cover until regional deglaciation at 9.8 cal ka BP (Stroeven et al., 2016). Second, we infer that Riukojietna extended over Lake 1063 until 8.1 ± 0.1 ka as evidenced by the ^{10}Be - ^{26}Al - ^{14}C weighted mean exposure age from sample Riuko-16-005. We therefore assume an ice thickness corresponding to the reconstructed LIA ice thickness over our samples (33-35 m; Fig. S5) between 9.8 cal ka BP and 8.1 ka. Third, the lake record shows evidence for a short mid-Holocene glacial advance between ~ 5.4 and 5.0 cal ka BP, based on our age-depth model (Figs. 4c, S2). We therefore allow for complete sample site exposure from 8.1 ka to 5.4 cal ka BP followed by ice burial to 5.0 cal ka BP. Finally, our lake record indicates a re-advance after 1.8 cal ka BP. We therefore model full exposure from 5.0-1.8 cal ka BP, followed by ice burial until 1910 CE and ice thinning since, per our ice surface reconstructions (Section 4.3; Tables 1, S5; Figs. 5, 6, S5). However, uncertainties in the age-depth model are large (Fig. S2), and the median value we use might not capture shorter-term fluctuations in sedimentation rate. We thus also consider an alternate scenario without that mid-Holocene advance over our ice-marginal sites (Fig. 7).

The modeled *in situ* ^{14}C inventories for Riuko-16-001 and 16-002 (without erosion) slightly underestimate the corresponding measured concentrations (slight overlap at 2σ with Riuko-16-001, no overlap with Riuko-16-002; Fig. 6a), indicating slightly longer Holocene or late deglacial exposure duration than was modeled (~ 400 years). In contrast, the modeled ^{10}Be and ^{26}Al concentrations for those samples significantly underestimate the measured ^{10}Be and ^{26}Al concentrations, indicating a significant component of ^{10}Be and ^{26}Al derived from exposure prior to the LGM (Fig. 6c, 6e). Given its relatively rapid decay, we consider it highly unlikely that discrepancies between measured and modeled *in situ* ^{14}C

can be explained through inheritance. Including erosion in the model for these samples would result in a worse fit between measured and modeled concentrations. The position of these samples, at a distance from faster flowing outlets, likely implies that they experienced cold-based, non-erosive ice conditions during the Holocene.

The modeled *in situ* ^{14}C concentrations of Riuko-16-003 and 16-004 without erosion slightly overestimate the measured values but agree within 2σ (Fig. 6b). As such, the *in situ* ^{14}C inventories measured at this bedrock knob site are largely compatible with the history of Riukojietna inferred from downstream Pajep Luoktejaure and our Holocene ice shielding assumptions for that location. However, the modeled ^{10}Be and ^{26}Al results for Riuko-16-003 and 16-004 without erosion slightly underestimate the corresponding measured concentrations (although agree within 2σ), indicating slightly longer exposure durations than was modeled (Fig. 6d, 6f). The position of Riuko-16-003 and 16-004 on an abundantly-striated bedrock knob emerging from an outlet glacier (Fig. S6), renders it worthwhile to model the potential effects of Holocene erosion on resulting concentrations. Including a subglacial erosion component of 0.05 mm yr^{-1} during Holocene ice burial periods yields improved agreement with measured *in situ* ^{14}C values over the no-erosion scenario (Fig. 6b; black stippled line). However, including erosion worsens the fit with measured values for ^{10}Be and ^{26}Al (Fig. 6d, 6f). A 400 year-longer Holocene exposure duration than suggested by this reconstruction leads to an even better fit with the *in situ* ^{14}C results at both sites when simultaneously allowing for 0.05 mm yr^{-1} of subglacial erosion of the bedrock knob in the northeastern outlet glacier (Fig. 7a and b). A component of inheritance is required to explain observed ^{10}Be and ^{26}Al nuclide concentrations in all ice-marginal samples (Figs. 6c-f, 7c-f). In summary, the ice-cap history reconstructed from the lacustrine record, the lake-outlet cosmogenic sample, and the LIA ice thickness reconstruction for Riukojietna, largely fits the measured *in situ* ^{14}C concentrations in the two ice-marginal bedrock sites.




Figure 6: Modeled cosmogenic nuclide accumulation (red) as a function of ice thickness (blue) based on the model described in the text. The width of the red shading surrounding the modeled accumulation trace reflects the cumulative effect of the production rate uncertainty in the sample. Panels a, c, and e show the results for samples Riuko-16-001 and Riuko-16-002, while b, d, and f show results for samples Riuko-16-003 and Riuko-16-004 for cases without subglacial erosion and with 0.05 mm yr^{-1} of subglacial erosion (dotted lines). Measured values of each nuclide ($\pm 2\sigma$ uncertainty) are indicated by gray shaded bands. Note that we track the cosmogenic nuclide concentrations in the sample as it is exhumed to the surface (a Lagrangian approach) as opposed to tracking the concentrations in the bedrock surface as rock is advected toward the surface by erosion (a Eulerian approach).

A rate of 0.05 mm yr^{-1} appears highly modest in a Swedish context where valley glaciers may have eroded their bedrock at ten times that rate (e.g., Schneider and Bronge, 1996). Having a difference in erosion between the two sites studied (Figs. 6, 7) is reasonable considering that basal sliding, and thereby glacial erosion, is likely higher at the bedrock knob site in the middle of the northeastern outlet glacier (where striations were abundant) than at the higher-elevation site flanking the southeastern and rather inactive tongue (where striations were absent). Encouragingly, a lower degree of inheritance is evident in the outlet glacier bedrock knob samples, aligning with inferred low rates of subglacial erosion for that site.




Figure 7: Modeled cosmogenic nuclide accumulation (red) as a function of ice thickness (blue) based on the model described in the text – in this case with no mid-Holocene readvance modeled. The width of the red shading surrounding the modeled accumulation trace reflects the cumulative effect of the production rate uncertainty in the sample. Panels a, c, and e show the results for samples Riuko-16-001 and Riuko-16-002, while b, d, and f show results for samples Riuko-16-003 and Riuko-16-004 for cases without subglacial erosion and with 0.05 mm yr^{-1} of subglacial erosion (dotted lines). Measured values of each nuclide are indicated by gray shaded bands ($\pm 2\sigma$ uncertainty).

5.4 Paleoclimate constraints

From our cosmogenic nuclide results and lake sediment record, we infer that Riukojietna covered the uppermost lake until 8.1 ka after which it retreated and thinned. However, although the ice cap perhaps remained smaller than today until shortly before 5.4 ka, diffuse fine silt and clay laminations indicate that the glacier persisted and produced sediment. This in itself is quite remarkable, and contrary to some previously widely held belief of Riukojietna demise (Karlén, 1988) and more broadly Scandinavian glacier demise (e.g., Nesje, 2009) during the Holocene, that a low-elevation ice cap persisted and produced sediments throughout the HTM. How did the warming climate of the HTM adjust to sustain this ice cap and allowed it to produce glacial flour?

Temperature reconstructions based on vegetation proxies such as tree-line shifts and pollen composition indicate that summer temperatures were already relatively high in the early Holocene (9.5–8.5 cal ka BP;

Karlén, 1976; Seppä and Birks, 2001; Kullman and Öberg, 2015) and during the HTM (Larocca and Axford, 2022). Given the paleo-record evidence for higher summer temperatures, we infer that Riukojietna remained in an advanced position until 8.1 ka and then persisted throughout the HTM because accumulation season precipitation was high enough to sustain the glacier. Indeed, a switch to a wetter post 8.1 ka Arctic (Thomas et al., 2018) and a wet early Holocene climate for Scandinavia, inferred from pollen and lake isotope records, may have resulted from a shifting dominance of moist North Atlantic air masses over Arctic air masses (Seppä and Hammarlund, 2000; Seppä and Birks, 2001; Shemesh et al., 2001) and a diminished influence of the deglaciating FIS. The reason for this switch may reside in warmer Arctic seas delivering more moisture and hemispheric warming driving increased poleward moisture transport (Thomas et al., 2018). Lake records showing surviving and even advancing glaciers during the early Holocene come from Austre Okstindbreen in northern Norway (Bakke et al., 2010). With summer temperatures higher-than-today and winters wetter-than-today characterizing the HTM, Riukojietna would have had a larger mass throughput which means, with a configuration as small as or even smaller than today, higher ice flow velocities and more favorable conditions for wet-based ice and bedrock erosion or sediment evacuation.

A reduction in ablation season temperature (Seppä and Birks, 2001), may have promoted a brief advance or ice cap reactivation recorded in Pajep Luoktejaure sediments before 5.0 ka. A nearby glacier also advanced around this time (Rosqvist et al., 2004) and so did several glaciers in northern Norway (Bakke et al., 2010; Wittmeier et al., 2015; Jansen et al., 2016).

Thinning and retreat of Riukojietna is implied by the cessation of glacial sediments into Pajep Luoktejaure after 5.0 cal ka BP. Subglacial erosion must then have been inhibited under a thin and cold-based ice cap, until, perhaps, disappearance. Because low ablation season temperatures have been inferred from vegetation proxies (Seppä and Birks, 2001) relative to HTM, we suggest that the ice cap primarily responded to a reduction in winter snow accumulation. We cannot currently preclude the complete demise of Riukojietna without further studies of upstream lakes 1009 and 1063. We currently favor a retracted ice cap configuration over full deglaciation because glacier model studies (e.g., Stroeve, 1996) clearly show that it requires more climate deterioration to grow an ice cap anew than to expand a remnant ice cap (hysteresis).

The ice cap reactivated and advanced again after 1.8 cal ka BP, as evidenced by laminated glacial sediments in Pajep Luoktejaure and historical evidence. The advance was likely triggered by a simultaneous lowering of ablation season temperatures and an increase in accumulation season precipitation. The increase in winter precipitation likely also prompted glaciers on the Lyngen Peninsula in northern Norway to advance at 1.8 cal ka BP (Bakke et al., 2005). A pronounced cooling starting at ca.

2 cal ka BP has been inferred from carbonate and diatom oxygen isotope records from several lakes in the area (Shemesh et al., 2001; Rosqvist et al., 2004, 2007). Evidence of a cooling at this time also comes from other parts of the Arctic (Kaufmann et al., 2009; Larocca and Axford, 2022) and the Northern Hemisphere (Solomina et al., 2016). Riukojietna remained in a relatively advanced position until the end of the LIA (Pohjola et al., 2005) after which it retreated to its current state.

6 Conclusions

In this paper we reconstruct glacier extent during the Holocene by means of cosmogenic nuclide modeling, thus integrating uncertainties from two traditional glacier reconstruction techniques. We apply this approach to Riukojietna, a small, low-altitude, low-gradient ice cap in northern Sweden, by combining evidence for ice-free conditions and ice-burial durations from cosmogenic nuclide chronometry (*in situ* ^{14}C , ^{10}Be , ^{26}Al) and evidence of glacial extent and activity derived from proglacial lacustrine records. This allows us to present the strongest case yet for a full Holocene glacial history of a Swedish glacier, and evaluate its sensitivity to climate change. An increase in winter precipitation during the early Holocene allowed the ice cap to be relatively large until 8.1 ka and to persist through peak Holocene warming before a possible brief reactivation prior to 5.0 cal ka BP. Drier conditions in the late Holocene forced its retreat to a configuration smaller than today until colder conditions during the last two millennia allowed the glacier to advance to its 8.1 ka position at the end of the LIA.

Modeling shows that the glacier history derived from lake sediment studies, given appropriate assumptions, yields cosmogenic nuclide inventories showing that Riukojietna was as small or smaller than today during earlier parts of the Holocene, that including subglacial erosion can improve model results, and that even for such a low-elevation ice cap the likelihood that it persisted throughout the Holocene is larger than that it disappeared (but this remains to be conclusively demonstrated). This last inference has ramifications for other higher-seated glaciers in Sweden and northern Scandinavia, where some paleoenvironmental studies have previously inferred the Holocene demise and regrowth of these glaciers, but where considerations of cold-based conditions and application of cosmogenic nuclide studies could potentially reveal complete Holocene ice-occupational histories.

Code Availability

All MATLAB code is available on <https://zenodo.org/records/20616442>.

Author contributions

This study was conceived by APS, GCR, and NAL. Sample material was collected by APS, GCR, and NAL. Cosmogenic sample preparation and analysis was done by AJK. Ice reconstructions were done by C-AW. JLA, AJK, and NAL wrote the model code. This manuscript is based on a portion of the Ph.D. dissertation of AJK. All authors contributed to the manuscript.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

Funding for the research was provided by “Stiftelsen Carl Mannerfelts fond”, “Göran Gustafssons Stiftelse, Lappland”, and the “Albert and Maria Bergströms Stiftelse”. Funding for this research was also provided by US National Science Foundation grants EAR-1560658 and EAR-2300559. We thank Mikael Amlert for help during field collection, Adrian Singleton (Purdue University) for initial sample processing, and Simon Larsson (Stockholm University) for assistance with Rbacon (Figs 4c and S2). JLA was supported by a Wenner-Gren Foundations Stipend. We thank Irene Schimmelpfennig and an anonymous reviewer for helpful comments that improved the manuscript.

References

- Aa, A. R., and Sønstegaard, E.: Early-Holocene glacier fluctuations of northern Grovabreen, western Norway, *Holocene*, 29, 187–196, <https://doi.org/10.1177/0959683618810392>, 2019.
- Andersen, J. L., Newall, J. C., Blomdin, R., Sams, S. E., Fabel, D., Koester, A. J., Lifton, N. A., Fredin, O., Caffee, M. W., Glasser, N. F., Rogozhina, I., Suganuma, Y., Harbor, J. M., and Stroeven, A. P.: Ice surface changes during recent glacial cycles along the Jutulstraumen and Penck Trough ice streams in western Dronning Maud Land, East Antarctica, *Quaternary Sci. Rev.*, 249, 106636, <https://doi.org/10.1016/j.quascirev.2020.106636>, 2020.
- Andreassen, L. M., Elvehøy, H., Kjølmoen, B., and Belart, J. M. C.: Glacier change in Norway since the 1960s—an overview of mass balance, area, length and surface elevation changes, *J. Glaciol.*, 66, 313–328, <https://doi.org/10.1017/jog.2020.10>, 2020.
- Andréasson, P. G., and Gee, D. G.: Bedrock geology and morphology of the Tarfala area, Kebnekaise Mts., Swedish Caledonides, *Geogr. Ann.*, 71A, 235–239, <https://doi.org/10.1080/04353676.1989.11880290>, 1989.
- Bakke, J., Dahl, S. O., Paasche, Ø., Løvlie, R., and Nesje, A.: Glacier fluctuations, equilibrium-line altitudes and palaeoclimate in Lyngen, northern Norway, during the Lateglacial and Holocene, *Holocene*, 15, 518–540, <https://doi.org/10.1191/0959683605hl815rp>, 2005.

680 Bakke, J., Dahl, S. O., Paasche, Ø., Simonsen, J. R., Kvisvik, B., Bakke, K., and Nesje, A.: A complete record of
681 Holocene glacier variability at Austre Okstindbreen, northern Norway: an integrated approach, *Quaternary*
682 *Sci. Rev.*, 29, 1246–1262, <https://doi.org/10.1016/j.quascirev.2010.02.012>, 2010.

683 Balco, G.: Production rate calculations for cosmic-ray-muon-produced ^{10}Be and ^{26}Al benchmarked against
684 geological calibration data, *Quaternary Geochronol.*, 39, 150–173,
685 <https://doi.org/10.1016/j.quageo.2017.02.001>, 2017.

686 Balco, G., Stone, J. O., Lifton, N. A., and Dunai, T. J.: A complete and easily accessible means of calculating
687 surface exposure ages or erosion rates from ^{10}Be and ^{26}Al measurements, *Quaternary Geochronol.*, 3, 174–
688 195, <https://doi.org/10.1016/j.quageo.2007.12.001>, 2008.

689 Balco, G., Brown, N., Nichols, K., Venturelli, R. A., Adams, J., Braddock, S., Campbell, S., Goehring, B., Johnson,
690 J. S., Rood, D. H., Wilcken, K., Hall, B., and Woodward, J.: Reversible ice sheet thinning in the Amundsen
691 Sea Embayment during the Late Holocene, *The Cryosphere*, 17, 1787–1801, [https://doi.org/10.5194/tc-17-](https://doi.org/10.5194/tc-17-1787-2023)
692 [1787-2023](https://doi.org/10.5194/tc-17-1787-2023), 2023.

693 Barnekow, L. Holocene regional and local vegetation history and lake-level changes in the Torneträsk area, northern
694 Sweden, *J. Paleolimnol.*, 23, 399–420, <https://doi.org/10.1023/A:1008171418429>, 2000.

695 Benn, D. I., and Hulton, N. R. J.: An ExcelTM spreadsheet program for reconstructing the surface profile of former
696 mountain glaciers and ice caps, *Comput. Geosci.*, 36, 605–610, <https://doi.org/10.1016/j.cageo.2009.09.016>,
697 2010.

698 Bergman, S., Stephens, M. B., Andersson, J., Kathol, B., and Bergman, T.: Bedrock map of Sweden, Scale 1:1
699 million, Sveriges geologiska undersökning, K 423, 2012.

700 Bevington, P., and Robinson, D.: *Data Reduction and Error Analysis for the Physical Sciences*, McGraw-Hill, New
701 York, 320 p., 2003.

702 Binnie, S. A., Dewald, A., Heinze, S., Voronina, E., Hein, A., Wittmann, H., von Blanckenburg, F., Hetzel, R.,
703 Christl, M., Schaller, M., Léanni, L., ASTER Team, Hippe, K., Vockenhuber, C., Ivy-Ochs, S., Maden, C.,
704 Fülöp, R. H., Fink, D., Wilcken, K. M., Fujioka, T., Fabel, D., Freeman, S. P. H. T., Xu, S., Fifield, L. K.,
705 Akçar, N., Spiegel, C., and Dunai, T. J.: Preliminary results of CoQtz-N: A quartz reference material for
706 terrestrial in-situ cosmogenic ^{10}Be and ^{26}Al measurements, *Nucl. Instrum. Meth. B*, 456, 203–212,
707 <https://doi.org/10.1016/j.nimb.2019.04.073>, 2019.

708 Bjune, A. E., Bakke, J., Nesje, A., and Birks, H. J. B.: Holocene mean July temperature and winter precipitation in
709 western Norway inferred from palynological and glaciological lake-sediment proxies, *Holocene*, 15, 177–189,
710 <https://doi.org/10.1191/0959683605hl798rp>, 2005.

711 Blaauw, M., and Christen, J. A.: Flexible paleoclimate age-depth models using an autoregressive gamma process,
712 *Bayesian Anal.*, 6, 457–474, <https://doi.org/10.1214/11-BA618>, 2011.

713 Borchers, B., Marrero, S., Balco, G., Caffee, M., Goehring, B., Lifton, N., Nishiizumi, K., Phillips, F., Schaefer, J.,
714 and Stone, J.: Geological calibration of spallation production rates in the CRONUS-Earth project, *Quaternary*
715 *Geochronol.*, 31, 188–198, <https://doi.org/10.1016/j.quageo.2015.01.009>, 2016.

716 Briner, J. P., Lifton, N. A., Miller, G. H., Refsnider, K., Anderson, R., and Finkel, R.: Using in situ cosmogenic
717 ^{10}Be , ^{14}C , and ^{26}Al to decipher the history of polythermal ice sheets on Baffin Island, Arctic Canada,
718 *Quaternary Geochronol.*, 19, 4–13, <https://doi.org/10.1016/j.quageo.2012.11.005>, 2014.

- 719 Chmeleff, J., von Blanckenburg, F., Kossert, K., and Jakob, D.: Determination of the ^{10}Be half-life by multicollector
720 ICP-MS and liquid scintillation counting, *Nucl. Instrum. Meth. B*, 268, 192–199,
721 <https://doi.org/10.1016/j.nimb.2009.09.012>, 2010.
- 722 Dunai, T. J.: *Cosmogenic Nuclides: Principles, Concepts and Applications in the Earth Surface Sciences*, Cambridge
723 University Press, 187 p., 2010.
- 724 Dussaillant, I., Hugonnet, R., Huss, M., Berthier, E., Bannwart, J., Paul, F., and Zemp, M.: Annual mass change of
725 the world's glaciers from 1976 to 2024 by temporal downscaling of satellite data with in situ observations,
726 *Earth Syst. Sci. Data*, 17, 1977–2006, <https://doi.org/10.5194/essd-17-1977-2025>, 2025.
- 727 Fabel, D., Stroeven, A. P., Harbor, J., Kleman, J., Elmore, D., and Fink, D.: Landscape preservation under
728 Fennoscandian ice sheets determined from in situ produced ^{10}Be and ^{26}Al , *Earth Planet. Sc. Lett.*, 201, 397–
729 406, [https://doi.org/10.1016/S0012-821X\(02\)00714-8](https://doi.org/10.1016/S0012-821X(02)00714-8), 2002.
- 730 Goehring, B. M., Schaefer, J. M., Schluechter, C., Lifton, N. A., Finkel, R. C., Jull, A. J. T., Akçar, N., and Alley, R.
731 B.: The Rhone Glacier was smaller than today for most of the Holocene, *Geology*, 39, 679–682,
732 <https://doi.org/10.1130/G32145.1>, 2011.
- 733 Gosse, J. C., and Phillips, F. M.: Terrestrial in situ cosmogenic nuclides: theory and application, *Quaternary Sci.*
734 *Rev.*, 20, 1475–1560, [https://doi.org/10.1016/S0277-3791\(00\)00171-2](https://doi.org/10.1016/S0277-3791(00)00171-2), 2001.
- 735 Grove, J. M.: *The Little Ice Age*, 2nd Edition, Routledge, London, <https://doi.org/10.4324/9780203505205>, 2004.
- 736 Hippe, K., Lifton, N. A.: Calculating isotope ratios and nuclide concentrations for in situ cosmogenic ^{14}C analyses,
737 *Radiocarbon*, 56, 1167–1174, <https://doi.org/10.2458/56.17917>, 2014.
- 738 Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., Jackson, M., Kääb, A., Kang, S., Kutuzov,
739 S., Milner, A., Molau, U., Morin, S., Orlove, B., and Steltzer, H.: High Mountain Areas, In Pörtner, H.-O.,
740 Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Alegría, A.,
741 Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N. M. (Eds.): *IPCC Special Report on the Ocean*
742 *and Cryosphere in a Changing Climate*, 131–202, 2019.
- 743 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, I.,
744 Brun, F., and Kääb, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592,
745 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- 746 Jansen, H. L., Simonsen, J. R., Dahl, S. O., Bakke, J., and Nielsen, P. R.: Holocene glacier and climate fluctuations
747 of the maritime ice cap Høgtuvbreen, northern Norway, *Holocene*, 26, 736–755,
748 <https://doi.org/10.1177/0959683615618265>, 2016.
- 749 Jansson, P., Rosqvist, G., and Schneider, T.: Glacier fluctuations, suspended sediment flux and glacio-lacustrine
750 sediments, *Geogr. Ann.*, 87A, 37–50, <https://doi.org/10.1111/j.0435-3676.2005.00243.x>, 2005.
- 751 Johnson, J. S., Nichols, K. A., Goehring, B. M., Balco, G., and Schaefer, J. M.: Abrupt mid-Holocene ice loss in the
752 western Weddell Sea Embayment of Antarctica, *Earth Planet. Sc. Lett.*, 518, 127–135,
753 <https://doi.org/10.1016/j.epsl.2019.05.002>, 2019.
- 754 Karlén, W.: Holocene glacier and climatic variations, Kebnekaise Mountains, Swedish Lapland, *Geogr. Ann.*, 55A,
755 29–63, <https://doi.org/10.1080/04353676.1973.11879879>, 1973.

- 756 Karlén, W.: Lacustrine sediments and tree-limit variations as indicators of Holocene climatic fluctuations in
757 Lappland, northern Sweden, *Geogr. Ann.*, 58A, 1–34, <https://doi.org/10.1080/04353676.1976.11879921>,
758 1976.
- 759 Karlén, W.: Lacustrine sediment studies: A technique to obtain a continuous record of Holocene glacier variations,
760 *Geogr. Ann.*, 63A, 273–281, <https://doi.org/10.1080/04353676.1981.11880042>, 1981.
- 761 Karlén, W.: Scandinavian glacial and climate fluctuations during the Holocene, *Quaternary Sci. Rev.*, 7, 199–209,
762 [https://doi.org/10.1016/0277-3791\(88\)90006-6](https://doi.org/10.1016/0277-3791(88)90006-6), 1988.
- 763 Karlén, W., and Denton, G. H.: Holocene glacial variations in Sarek National Park, northern Sweden, *Boreas*, 5, 25–
764 56, <https://doi.org/10.1111/j.1502-3885.1976.tb00329.x>, 1976.
- 765 Karlén, W., Bodin, A., Kuylenstierna, J., and Naslund, J.-O.: Climate of northern Sweden during the Holocene, *J.*
766 *Coastal Res. Spec. Issue*, 17, 49–54, 1995.
- 767 Kaufman, D. S., Schneider, D. P., McKay, N. P., Ammann, C. M., Bradley, R. S., Briffa, K. R., Miller, G. H., Otto-
768 Bliessner, B. L., Overpeck, J. T., Vinther, B. M., Arctic Lakes 2K Project Members, Abbott, M., Axford, Y.,
769 Bird, B., Birks, H. J. B., Bjune, A. E., Briner, J., Cook, T., Chipman, M., Francus, P., Gajewski, K.,
770 Geirsdóttir, Á., Hu, F. S., Kutchko, B., Lamoureux, S., Loso, M., MacDonald, G., Peros, M., Porinchu, D.,
771 Schiff, C., Seppä, H., and Thomas, E.: Recent warming reverses long-term Arctic cooling, *Science*, 325,
772 1236–1239, <https://doi.org/10.1126/science.1173983>, 2009.
- 773 Knudsen, M. F., Egholm, D. L., and Jansen, J. D.: Time-integrating cosmogenic nuclide inventories under the
774 influence of variable erosion, exposure, and sediment mixing, *Quaternary Geochronol.*, 51, 110–119,
775 <https://doi.org/10.1016/j.quageo.2019.02.005>, 2019.
- 776 Koester, A. J.: Applications of in situ ^{14}C to glacial landscapes in Sweden and Antarctica [PhD Dissertation]: Purdue
777 University, 186 p., 2023.
- 778 Koester, A. J., and Lifton, N. A.: Technical note: A software framework for calculating compositionally dependent
779 in situ ^{14}C production rates, *Geochronol.*, 5, 21–33, <https://doi.org/10.5194/gchron-5-21-2023>, 2023.
- 780 Koester, A. J., and Lifton, N. A.: Corrigendum to “Technical note: A software framework for calculating
781 compositionally dependent in situ ^{14}C production rates” published in *Geochronology*, 5, 21–33, 2023,
782 *Geochronol.*, <https://doi.org/10.5194/gchron-5-21-2023-corrigendum>, 2024.
- 783 Kohl, C. P., and Nishiizumi, K.: Chemical isolation of quartz for measurement of in-situ-produced cosmogenic
784 nuclides, *Geochim. Cosmochim. Ac.*, 56, 3583–3587, [https://doi.org/10.1016/0016-7037\(92\)90401-4](https://doi.org/10.1016/0016-7037(92)90401-4), 1992.
- 785 Korschinek, G., Bergmaier, A., Faestermann, T., Gerstmann, U. C., Knie, K., Rugel, G., Wallner, A., Dillmann, I.,
786 Dollinger, G., Lierse von Gostomski, Ch., Kossert, K., Maiti, M., Poutivtsev, M., and Remmert, A.: A new
787 value for the half-life of ^{10}Be by Heavy-Ion Elastic Recoil Detection and liquid scintillation counting, *Nucl.*
788 *Instrum. Meth. B*, 268, 187–191, <https://doi.org/10.1016/j.nimb.2009.09.020>, 2010.
- 789 Kullman, L., and Öberg, L.: New aspects of high-mountain palaeobiogeography: A synthesis of data from forefields
790 of receding glaciers and ice patches in the Tärna and Kebnekaise mountains, Swedish Lapland, *ARCTIC*, 68,
791 141–152, <https://doi.org/10.14430/arctic4480>, 2015.
- 792 Larocca, L. J., and Axford, Y.: Arctic glaciers and ice caps through the Holocene: a circumpolar synthesis of lake-
793 based reconstructions, *Clim. Past*, 18, 579–606, <https://doi.org/10.5194/cp-18-579-2022>, 2022.

Leonard, E. M.: Use of lacustrine sedimentary sequences as indicators of Holocene glacial history, Banff National Park, Alberta, Canada, *Quaternary Res.*, 26, 218–231, [https://doi.org/10.1016/0033-5894\(86\)90106-7](https://doi.org/10.1016/0033-5894(86)90106-7), 1986.

Lifton, N., Sato, T., and Dunai, T. J.: Scaling in situ cosmogenic nuclide production rates using analytical approximations to atmospheric cosmic-ray fluxes, *Earth Planet. Sc. Lett.*, 386, 149–160, <https://doi.org/10.1016/j.epsl.2013.10.052>, 2014.

Lifton, N. A., Phillips, F. M., and Cerling, T. E.: Using Lake Bonneville features to calibrate in situ cosmogenic nuclide production rates, *Developments Earth Surf. Proc.*, 20, 165–183, <https://doi.org/10.1016/B978-0-444-63590-7.00009-3>, 2016.

Lifton, N., Wilson, J., and Koester, A.: Technical note: Studying lithium metaborate fluxes and extraction protocols with a new, fully automated in situ cosmogenic ^{14}C processing system at PRIME Lab, *Geochronol.*, 5, 361–375, <https://doi.org/10.5194/gchron-5-361-2023>, 2023.

Matthews, J. A., and Briffa, K. R.: The ‘Little Ice Age’: Re-evaluation of an evolving concept, *Geogr. Ann.*, 87A, 17–36, <https://doi.org/10.1111/j.0435-3676.2005.00242.x>, 2005.

Matthews, J. A., and Dresser, P. Q.: Holocene glacier variation chronology of the Smørstabbtindan massif, Jotunheimen, southern Norway, and the recognition of century- to millennial-scale European Neoglacial Events, *Holocene*, 18, 181–201, <https://doi.org/10.1177/0959683607085608>, 2008.

Nesje, A.: Latest Pleistocene and Holocene alpine glacier fluctuations in Scandinavia, *Quaternary Sci. Rev.*, 28, 2119–2136, <https://doi.org/10.1016/j.quascirev.2008.12.016>, 2009.

Nesje, A., Dahl, S. O., Andersson, C., and Matthews, J. A.: The lacustrine sedimentary sequence in Syngneskardvatnet, western Norway: A continuous, high-resolution record of the Jostedalbreen ice cap during the Holocene, *Quaternary Sci. Rev.*, 19, 1047–1065, [https://doi.org/10.1016/S0277-3791\(99\)00090-6](https://doi.org/10.1016/S0277-3791(99)00090-6), 2000.

Nesje, A., Bakke, J., Dahl, S. O., Lie, Ø., and Matthews, J. A.: Norwegian mountain glaciers in the past, present and future, *Glob. Planet. Change*, 60, 10–27, <https://doi.org/10.1016/j.gloplacha.2006.08.004>, 2008.

Nichols, K. A., and Goehring, B. M.: Isolation of quartz for cosmogenic in situ ^{14}C analysis, *Geochronol.*, 1, 43–52, <https://doi.org/10.5194/gchron-1-43-2019>, 2019.

Nichols, K. A., Goehring, B. M., Balco, G., Johnson, J. S., Hein, A. S., and Todd, C.: New Last Glacial Maximum ice thickness constraints for the Weddell Sea Embayment, Antarctica, *The Cryosphere*, 13, 2935–2951, <https://doi.org/10.5194/tc-13-2935-2019>, 2019.

Nielsen, P. R., Balascio, N. L., Dahl, S. O., Jansen, H. L., Støren, E. W. N., and Bradley, R. S.: A high-resolution 1200-year lacustrine record of glacier and climate fluctuations in Lofoten, northern Norway, *Holocene*, 26, 917–934, <https://doi.org/10.1177/0959683615622551>, 2016.

Nishiizumi, K.: Preparation of ^{26}Al AMS standards, *Nucl. Instrum. Meth. B*, 223–224, 388–392, <https://doi.org/10.1016/j.nimb.2004.04.075>, 2004.

Oerlemans, J.: Extracting a climate signal from 169 glacier records, *Science*, 308, 675–677, <https://doi.org/10.1126/science.1107046>, 2005.

Oerlemans, J., and Fortuin, J. P. F.: Sensitivity of glaciers and small ice caps to greenhouse warming, *Science*, 258, 115–118, <https://doi.org/10.1126/science.258.5079.115>, 1992.

- 831 Oerlemans, J., Anderson, B., Hubbard, A., Huybrechts, P., Jóhannesson, T., Knap, W. H., Schmeits, M., Stroeve,
832 A. P., van de Wal, R. S. W., Wallinga, J., and Zuo, Z.: Modelling the response of glaciers to climate warming,
833 *Clim. Dynam.*, 14, 267–274, <https://doi.org/10.1007/s003820050222>, 1998.
- 834 Phillips, F. M., Argento, D. C., Balco, G., Caffee, M. W., Clem, J., Dunai, T. J., Finkel, R., Goehring, B., Gosse, J.
835 C., Hudson, A. M., Jull, A. J. T., Kelly, M. A., Kurz, M., Lal, D., Lifton, N., Marrero, S. M., Nishiizumi, K.,
836 Reedy, R. C., Schaefer, J., Stone, J. O. H., Swanson, T., and Zreda, M. G.: The CRONUS-Earth Project: A
837 synthesis, *Quaternary Geochronol.*, 31, 119–154, <https://doi.org/10.1016/j.quageo.2015.09.006>, 2016.
- 838 Ploeg, K., and Stroeve, A. P.: History and dynamics of Fennoscandian Ice Sheet retreat, contemporary ice-dammed
839 lake evolution, and faulting in the Torneträsk area, northwestern Sweden, *The Cryosphere*, 19, 347–373,
840 <https://doi.org/10.5194/tc-19-347-2025>, 2025.
- 841 Pohjola, V. A., Cole-Dai, J., Rosqvist, G., Stroeve, A. P., and Thompson, L. G.: Potential to recover climatic
842 information from Scandinavian ice cores: An example from the small ice cap Riukojietna, *Geogr. Ann.*, 87A,
843 259–270, <https://doi.org/10.1111/j.0435-3676.2005.00257.x>, 2005.
- 844 Rand, C., and Goehring, B. M.: The distribution and magnitude of subglacial erosion on millennial timescales at
845 Engabreen, Norway, *Ann. Glaciol.*, 60, 73–81, <https://doi.org/10.1017/aog.2019.42>, 2019.
- 846 Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H.,
847 Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G.,
848 Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J.,
849 Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F.,
850 Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake,
851 F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A., and Talamo, S.: The IntCal20 Northern Hemisphere
852 radiocarbon age calibration curve (0–55 cal kBP), *Radiocarbon*, 62, 725–757,
853 <https://doi.org/10.1017/RDC.2020.41>, 2020.
- 854 Rosqvist, G., and Østrem, G.: The sensitivity of a small icecap to climatic fluctuations, *Geogr. Ann.*, 71A, 99–103,
855 <https://doi.org/10.1080/04353676.1989.11880277>, 1989.
- 856 Rosqvist, G., Jonsson, C., Yam, R., Karlén, W., and Shemesh, A.: Diatom oxygen isotopes in pro-glacial lake
857 sediments from northern Sweden: A 5000 year record of atmospheric circulation, *Quaternary Sci. Rev.*, 23,
858 851–859, <https://doi.org/10.1016/j.quascirev.2003.06.009>, 2004.
- 859 Rosqvist, G., Leng, M., and Jonsson, C.: North Atlantic region atmospheric circulation dynamics inferred from a
860 late-Holocene lacustrine carbonate isotope record, northern Swedish Lapland, *Holocene*, 17, 867–873,
861 <https://doi.org/10.1177/0959683607080508>, 2007.
- 862 Røthe, T. O., Bakke, J., and Støren, E. W. N.: Glacier outburst floods reconstructed from lake sediments and their
863 implications for Holocene variations of the plateau glacier Folgefonna in western Norway, *Boreas*, 48, 616–
864 634, <https://doi.org/10.1111/bor.12388>, 2019.
- 865 Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D.,
866 Compagno, L., Copland, L., Farinotti, D., Menounos, B., and McNabb, R. W.: Global glacier change in the
867 21st century: every increase in temperature matters, *Science*, 379, 78–83,
868 <https://doi.org/10.1126/science.abo1324>, 2023.
- 869 Rubensdotter, L., and Rosqvist, G.: The effect of geomorphological setting on Holocene lake sediment variability,
870 northern Swedish Lapland, *J. Quaternary Sci.*, 18, 757–767, <https://doi.org/10.1002/jqs.800>, 2003.

871 Santos, G. M., Southon, J. R., Druffel-Rodriguez, K. C., Griffin, S., and Mazon, M.: Magnesium Perchlorate as an
872 alternative water trap in AMS graphite sample preparation: A report on sample preparation at KCCAMS at the
873 University of California, Irvine, Radiocarbon, 46, 165–174, <https://doi.org/10.1017/S0033822200039485>,
874 2004.

875 Santos, G. M., Moore, R. B., Southon, J. R., Griffin, S., Hinger, E., and Zhang, D.: AMS ^{14}C sample preparation at
876 the KCCAMS/UCI Facility: Status report and performance of small samples, Radiocarbon, 49, 255–269,
877 <https://doi.org/10.1017/S0033822200042181>, 2007.

878 Schimmelpfennig, I., Schaefer, J. M., Lamp, J., Godard, V., Schwartz, R., Bard, E., Tuna, T., Akçar, N., Schlüchter,
879 C., Zimmerman, S., and ASTER Team: Glacier response to Holocene warmth inferred from in situ ^{10}Be and
880 ^{14}C bedrock analyses in Steingletscher's forefield (central Swiss Alps), Clim. Past, 18, 23–44,
881 <https://doi.org/10.5194/cp-2021-110>, 2022.

882 Schneider, T., and Bronge, C.: Suspended sediment transport in the Storglaciären drainage basin, Geogr. Ann., 78A,
883 155–161, <https://doi.org/10.1080/04353676.1996.11880461>, 1996.

884 Seppä, H., and Birks, H. J. B.: July mean temperature and annual precipitation trends during the Holocene in the
885 Fennoscandian tree-line area: pollen-based climate reconstructions, Holocene, 11, 527–539,
886 <https://doi.org/10.1191/095968301680223486>, 2001.

887 Seppä, H., and Hammarlund, D.: Pollen-stratigraphical evidence of Holocene hydrological change in northern
888 Fennoscandia supported by independent isotopic data, J. Paleolimnol., 24, 69–79,
889 <https://doi.org/10.1023/A:1008169800682>, 2000.

890 Seppä, H., Hammarlund, D., and Antonsson, K.: Low-frequency and high-frequency changes in temperature and
891 effective humidity during the Holocene in south-central Sweden: Implications for atmospheric and oceanic
892 forcings of climate, Clim. Dynam., 25, 285–297, <https://doi.org/10.1007/s00382-005-0024-5>, 2005.

893 Shemesh, A., Rosqvist, G., Rietti-Shati, M., Rubensdotter, L., Bigler, C., Yam, R., and Karlén, W.: Holocene
894 climatic change in Swedish Lapland inferred from an oxygen-isotope record of lacustrine biogenic silica,
895 Holocene, 11, 447–454, <https://doi.org/10.1191/095968301678302887>, 2001.

896 Sjögren, P., and Damm, C.: Holocene vegetation change in northernmost Fennoscandia and the impact on
897 prehistoric foragers 12 000–2000 cal. a BP – A review, Boreas, 48, 20–35, <https://doi.org/10.1111/bor.12344>,
898 2019.

899 Sjögren, P. J. E.: An overview of Holocene climate reconstructions in northernmost Fennoscandia, Septentrio
900 Reports 3, 31 p. <https://doi.org/10.7557/7.5747>, 2021.

901 Solomina, O. N., Bradley, R. S., Jomelli, V., Geirsdottir, A., Kaufman, D. S., Koch, J., McKay, N. P., Masiokas, M.,
902 Miller, G., Nesje, A., Nicolussi, K., Owen, L. A., Putnam, A. E., Wanner, H., Wiles, G., and Yang, B.: Glacier
903 fluctuations during the past 2000 years, Quaternary Sci. Rev., 149, 61–90,
904 <https://doi.org/10.1016/j.quascirev.2016.04.008>, 2016.

905 Stroeve, A. P.: The robustness of one-dimensional time dependent ice-flow models: A case study from
906 Storglaciären, northern Sweden, Geogr. Ann., 78A, 133–146,
907 <https://doi.org/10.1080/04353676.1996.11880459>, 1996.

908 Stroeve, A. P., Fabel, D., Harbor, J., Hättestrand, C., and Kleman, J.: Quantifying the erosional impact of the
909 Fennoscandian ice sheet in the Torneträsk–Narvik corridor, northern Sweden, based on cosmogenic
910 radionuclide data, Geogr. Ann., 84A, 275–287, <https://doi.org/10.1111/j.0435-3676.2002.00182.x>, 2002.

911 Stroeve, A. P., Hättestrand, C., Kleman, J., Heyman, J., Fabel, D., Fredin, O., Goodfellow, B. W., Harbor, J. M.,
 912 Jansen, J. D., Olsen, L., Caffee, M. W., Fink, D., Lundqvist, J., Rosqvist, G. C., Strömberg, B., and Jansson,
 913 K. N.: Deglaciation of Fennoscandia, *Quaternary Sci. Rev.*, 147, 91–121,
 914 <https://doi.org/10.1016/j.quascirev.2015.09.016>, 2016.

915 Stuiver, M., and Reimer, P. J.: Extended ^{14}C data base and revised CALIB 3.0 ^{14}C age calibration program,
 916 *Radiocarbon*, 35, 215–230, <https://doi.org/10.1017/S0033822200013904>, 1993.

917 Svenonius, F.: Studien über den Kårso- und die Kebnegletscher nebst Notizen über andere Gletscher im
 918 Jukkasjärvicebirge, Part 1 of Die Gletscher Schwedens im Jahre 1908, *Sveriges geologiska undersökning*,
 919 serie Ca, 5, 1910.

920 Thomas, E. K., Castañeda, I. S., McKay, N. P., Briner, J. P., Salacup, J. M., Nguyen, K. Q., and Schweinsberg, A.
 921 D.: A wetter Arctic coincident with hemispheric warming 8,000 years ago, *Geoph. Res. Lett.*, 45, 10637–
 922 10647, <https://doi.org/10.1029/2018GL079517>, 2018.

923 Wahlström, C.-A.: A century of volume reduction of a small ice cap; Riukojietna northern Sweden, Unpublished
 924 MSc Thesis, Stockholm University, 2016.

925 Wastegård, S.: The Holocene of Sweden – a review, *GFF*, 144, 126–149,
 926 <https://doi.org/10.1080/11035897.2022.2086290>, 2022.

927 WGMS: Fluctuations of Glacier (FoG) Database, World Glacier Monitoring Service (WGMS), Zurich, Switzerland,
 928 <https://doi.org/10.5904/wgms-fog-2025-02b>, 2025.

929 Wirsig, C., Ivy-Ochs, S., Akçar, N., Lupker, M., Hippe, K., Wacker, L., Vockenhuber, C., and Schlüchter, C.:
 930 Combined cosmogenic ^{10}Be , in situ ^{14}C and ^{36}Cl concentrations constrain Holocene history and erosion depth
 931 of Grueben glacier (CH), *Swiss J. Geosci.*, 109, 379–388, <https://doi.org/10.1007/s00015-016-0227-2>, 2016.

932 Wittmeier, H. E., Bakke, J., Vasskog, K., and Trachsel, M.: Reconstructing Holocene glacier activity at
 933 Langfjordjøkelen, Arctic Norway, using multi-proxy fingerprinting of distal glacier-fed lake sediments,
 934 *Quaternary Sci. Rev.*, 114, 78–99, <https://doi.org/10.1016/j.quascirev.2015.02.007>, 2015.

935 Young, N. E., Schaefer, J. M., Goehring, B., Lifton, N., Schimmelpfennig, I., and Briner, J. P.: West Greenland and
 936 global in situ ^{14}C production-rate calibrations. *J. Quaternary Sci.*, 29, 401–406,
 937 <https://doi.org/10.1002/jqs.2717>, 2014.

938 Young, N. E., Lesnek, A. J., Cuzzone, J. K., Briner, J. P., Badgley, J. A., Balter-Kennedy, A., Graham, B. L.,
 939 Cluett, A., Lamp, J. L., Schwartz, R., Tuna, T., Bard, E., Caffee, M. W., Zimmerman, S. R. H., and Schaefer,
 940 J. M.: In situ cosmogenic ^{10}Be – ^{14}C – ^{26}Al measurements from recently deglaciated bedrock as a new tool to
 941 decipher changes in Greenland Ice Sheet size, *Clim. Past*, 17, 419–450, [https://doi.org/10.5194/cp-17-419-](https://doi.org/10.5194/cp-17-419-2021)
 942 [2021](https://doi.org/10.5194/cp-17-419-2021), 2021.

943