Author Responses for NO_x Emissions Constraints from GEMS NO_2 Retrievals: Inversion Methodology and Air Quality Model Evaluation in Bangkok using ASIA-AQ Multi-Platform Observations

Christopoulos et al.

We thank the reviewers for their careful reading of the manuscript and for the constructive comments that have helped improve its quality. Below, we respond to each comment (italics). Revisions made in the manuscript are indicated in green text.

Response to the Editor

Comment: *Thank you for your detailed and thorough responses to the reviewers' questions and comments. I am pleased to accept the revised manuscript for publication subject to a minor change to Sections 7 and 8. In the current manuscript, both include "Discussion", at least indicated by their titles. I suggest to combine the two sections or, alternatively, modifying them so that Section 7 contains a discussion and Section 8 focuses on Conclusions. Please have a look at ACP's authors'guidelines for the concluding section (https://www.atmospheric-chemistry-and-physics.net/policies/guidelines_for_authors.html).*

Response: We thank the editor for this helpful suggestion. In the revised manuscript, we have updated Section titles accordingly to reflect this. Section 7 now serves as the discussion section, while Section 8 is dedicated to the conclusion and summary of the study which was the intended structure previously. Additionally, in this updated version we added a statement to the acknowledgements section and made minor typographical and grammatical corrections throughout the manuscript.

Anonymous Referee #1

General Comments:

Comment #1: *Section 3: The study relies heavily on the CSF method for deriving top-down NOx emissions. While this approach is computationally inexpensive, it has known limitations. For instance, how does the algorithm determine the optimal length for line density integration? Additionally, what is the sensitivity of the inferred emissions to transport wind speed errors and the fitting process for line density and chemical lifetime? I suggest the authors include a brief discussion on the structural uncertainties inherent to the CSF method.*

Response #1: Thank you for this helpful comment. We agree that the CSF method introduces several structural uncertainties that should be clearly described and so have revised Section 3 to explicitly address these points. Specifically, we now clarify that the integration bounds used to compute line densities are defined by the plume subregion identified using the *ddeq* framework, based on GEMS-observed

enhancements and ERA5 wind direction. We also added a discussion of the uncertainty of inferred emissions to transport wind speed, noting that errors in wind magnitude and direction directly propagate into emission estimates through the flux calculation (referring to Figure S5). In addition, we now describe uncertainties associated with the Gaussian fitting of line density and the estimation of the effective lifetime, both of which influence the derived emissions. These additions are included in a new paragraph (Section 3.3.5).

Revised Manuscript (Lines 304 – 307): The integration bounds y_1 and y_1 are defined by the plume subregion (polygon) identified using the ddeq framework, which delineates the area of enhanced NO_2 associated with the source based on plume detection and wind direction (Kuhlmann et al., 2024). As such, the effective integration length across the plume is determined by the spatial extend of this polygon and the fitted Gaussian representation.

Revised Manuscript (Lines 3336 – 348)

3.1.5. Structural uncertainties in the CSF framework

While the CSF framework provides a computationally efficient approach for deriving satellite-derived emissions, we acknowledge the method contains some structural uncertainties. First, the integration bounds used to compute the line densities (Eq. 4) are defined by the plume polygon identified from the satellite observations and wind field, such that the inferred top-down emissions depend on how the plume extent is delineated. As a result, inferred satellite-derived emissions can be sensitive to the selected plume extend, particularly under conditions of weak enhancements or overlapping sources. Second, the satellite-derived emission estimate scales directly with the effective wind speed (Eq. 1), such that errors in wind magnitude or direction propagate into the derived fluxes and satellite-derived emissions (see Fig. S4). Although ERA5 winds averaged over the lowest model layers are used to approximate plume transport, they may not fully represent local meteorological variability. Third, the inferred satellite-derived emission rate depends on the assumptions within the fitting framework, including the Gaussian representation of plume structure (Eq. 5), the parametrization of plume spreading (Eq. 6), and the effective lifetime τ (Eq. 7), which represent a simplified treatment of chemical decay and dispersion. Together, these factors represent inherent uncertainties in the CSF approach that should be considered when interpreting the satellite-derived emissions.

Comment #2: *The authors develop a novel approach for updating diurnal emission profiles; however, the inversion is essentially simplified to a 1-D problem (magnitude over time for a single urban box). Inaccuracy in the spatial distribution of NOx sources also contributes significantly to model error. Do the authors have thoughts on how to enhance this method to quantify emissions on a finer spatial scale?*

Response #2: Thank you for the comment. We would like to clarify that while the inversion is formulated as a temporal optimization (i.e., solving for hourly scaling factors), these scaling factors are applied to each grid cell within the Bangkok box of the domain. As a result, the emissions retain their spatial structure from the prior inventory but are adjusted uniformly in time across all grid cells.

We agree, however, that this approach does not explicitly optimize the spatial distribution of emissions, and therefore any spatial inaccuracies in the prior (e.g., EDGAR downscaled with ODIAC) spatial pattern

are not corrected by the inversion. Given the current framework, which treats emissions as area sources, improving spatial patterns is inherently limited. Addressing this would likely require a more detailed treatment, such as performing separate inversions for distinct source types (e.g., point and area sources, including features like the economic corridor southeast of Bangkok). In addition, the use of a higher-resolution local inventory could further improve the representation of regional variability. However, this study relied on the best available data at the time (EDGAR).

We have clarified this distinction and added a brief discussion in the revised manuscript.

Revised Manuscript (Lines 1051 – 1060):

7.2 Limitations and potential extensions of the inversion framework


The current model emission update framework optimizes the temporal evolution of emissions but does not explicitly resolve regional variability, as a single set of hourly scaling factors is applied uniformly across all grid cells within the BMR. As a result, the inversion preserves the spatial structure of the prior inventory, and any inaccuracies in the spatial distribution of emissions are not corrected.

Extending this approach to resolve emissions at finer scales would require allowing emissions to vary across grid cells or subregions, supported by additional constraints. These constraints could include regional higher-resolution regional prior inventories when available, as well as higher-resolution observations (e.g., GCAS) that can better resolve the urban variability in NO_2 as shown in this analysis.


Comment #3: *Biomass burning in Southeast Asia significantly influences regional air quality. Was the study period affected by any specific biomass burning events? I would appreciate a brief discussion on how such events might impact the NOx and O3 simulations, as well as the impact on the NOx emission inversion.*

Response #3: Thank you for your important comment. To address this, we include an additional plot of biomass burning emissions (aggregated NO emissions) below derived from the QFED (used in this study) and FINN v2.5 inventories. These inventories provide a reasonable range of fire activity, with QFED generally representing a lower-bound estimate (particularly with agricultural fires in SE Asia) and FINN representing an upper-bound estimate. We separate the plots to illustrate two different regimes during our study period (13 – 19 March and 20 – 27 March 2024). During the study period, biomass burning activity was present primarily outside of the Bangkok region with burning focused on regions in northern Thailand and Myanmar. Peak fire activity was seen from 10 – 15 March. During this period, weak synoptic flow likely reduced efficient long-range transport of biomass burning plumes, likely contributing to buildup in the source regions and minimal influence in the Bangkok area given chemical lifetime of NO_x. Following 20 March, precipitation in the region substantially reduced fire activity as depicted in the below plots (b), indicating substantially cleaner conditions. While burning can contribute to regional NO_x and O_3 enhancements, its impact on the inversion results presented here is expected to be somewhat limited given the burning locations, which substantial impacts likelier seen in cities in the north (e.g., Chiang Mai). We have added clarifying information on this in the independent evaluation to frame interpretation of the results.

QFED NO Emission:



FINN NO Emission:



Revised Manuscript (Lines 675 – 680): Biomass burning was active in the region during the study period, particularly over northern Thailand and Myanmar, with peak activity occurring in mid-March. However, transport to Bangkok was likely limited due to weak synoptic flow and short chemical lifetime of NO_x, reducing its impact on urban concentrations of NO_x in the BMR. Following March 20, precipitation suppressed fire activity, further minimizing its impact. While biomass burning may contribute to broader regional enhancements, its impact on the inversion results and coincident model evaluation over Bangkok is expected to be minor.

Comment #4: *Section 3&4: The current terminology used to describe emissions is inconsistent and may be confusing to the reader. Terms such as 'prior NOx emissions,' 'CSF-derived emissions,' and 'emissions after applying the forward operator' are used interchangeably or in ways that overlap. I suggest the author*

review these sections to consolidate the terminology and ensure that uniform terms are used consistently throughout the manuscript to distinguish between the different emissions datasets.

Response #4: Thank you for this helpful comment. We agree that the terminology describing the different emission datasets was not sufficiently consistent and may lead to confusion. As such, we have revised Section 3 and 4 to distinguish terms throughout the manuscript. Specifically, we now consistently refer to the input inventory as the *prior emissions*, the GEMS emissions derived from the CSF method as the *satellite-derived emissions*, the final daytime profile after applying the optimization as the *posterior emissions,* and the optimized input emissions as the *updated emissions*. We hope these revisions improve clarify and ensure a clear distinction between the different emissions referred to in this study.

Revised Manuscript (Lines 276 – 279): In the following subsections and in Section 4, we refer to the WRF-Chem input inventory based on EDGAR as the *prior emissions*, the CSF-based estimates from GEMS as the *satellite-derived emissions*, the CSF-based estimates from the model-simulated columns as the model-derived emissions, and the final emissions after applying the optimization framework, as the *posterior emissions*.

Comment #5: *The citations for figures in the Supplement appear disorganized. Please verify the numbering and ensure they are cited sequentially.*

Response #5: We have carefully reviewed all Supplementary figure citation and ensured they are numbered and referenced sequentially throughout the manuscript. The figures in the Supplement have been reordered to flow with order of reference in the manuscript to improve clarity and consistency.

Specific comments:

Comment #6: *Line 49-51: The statement that GEMS NO2 product bias "cannot be diagnosed using satellite data alone" seems somewhat redundant, as independent observations are inherently required to assess retrieval quality.*

Response #6: We have corrected the manuscript text as follows:

Revised Manuscript (Lines 53 – 54): Remaining negative biases are consistent with a systematic low bias in the GEMS v3 NO_2, highlighting the importance of multi-platform evaluation using independent observations.

Comment #7: *Line 125: The reanalysis dataset should be formally cited as NCEP FNL.*

Response #7: The manuscript has been updated to reflect this citation.

Revised Manuscript (Lines 128 – 130): The model was driven by reanalysis meteorology, NCEP FNL (Final), and Copernicus Modeling Service (CAMS) chemical boundary conditions (Commerce, 2000; Inness et al., 2019).

Comment #8: *Figure 1: Since NOx is primarily emitted as NO, it would be more appropriate to plot either NO or total NOx emissions rather than just NO2.*

Response #8: We have replaced Figure 1 with a plot of total NO_x emissions obtained from the EDGAR inventory.

Revised Manuscript (Lines 155 – 159):




Figure 1. Spatial illustration of the WRF-Chem model domain configuration, including D01 (20 km) and D02 (4 km), and average base-model NO_x input emissions from EDGAR v5. D02 is centered over Thailand, and the urban signal associated with the Bangkok Metropolitan Region (BMR) is highlighted by the star.

Comment #9: *Table 1 and Section 6.1.2: Since model NOx is sensitive to the accuracy of local wind field simulations, were any measures like grid or observational nudging applied to the model meteorology? Furthermore, why was 1-degree NCEP FNL data utilized instead of higher- resolution 0.25-degree data?*

Response #9: Yes, meteorology was constrained using four-dimensional data assimilation (FDDA) via grid nudging. Nudging was applied to d01 and d02. We also clarify that the simulations here used the 0.25° NCEP FNL data; the former reference was a typo. We have updated Table 1 and added clarifying information to the manuscript.

Revised Manuscript (Lines 130 – 131): Meteorology was constrained using four-dimension data assimilation (FDDA) via grid nudging applied to model domains using the 0.25° NCEP FNL reanalysis fields to improve large-scale wind representation.

Revised Manuscript (Lines 153 – 155): **Table 1** WRF-Chem base (WRF_{Base}) model configuration and input datasets.

Component	Configuration/Source
Model Version	WRF-Chem v4.2.2
Domain	D01: 20 km (regional); D02: 4 km (nested)
Meteorological IC/BC	NCEP FNL reanalysis (0.25v x 0.25°; 6-hourly)
Chemical IC/BC	CAMS global model output
Chemistry Mechanism	RACM-MADE-VBS (with aqueous reactions)
Anthropogenic Emissions	EDGAR v5; downscaled with ODIAC CO_2
Biogenic Emissions	MEGAN v2.1 (Guenther et al., 2006)
Biomass Burning Emissions	QFED v2.6 (Grell et al., 2011; Koster et al., 2015)
Dust Emissions	GOCART scheme (Zhao et al., 2010)
Sea Salt Emissions	Gong et al. (1997) parameterization (Gong et al., 1997)
Radiation Scheme	RRTMG (longwave); Goddard (shortwave)
Cumulus Parameterization	Grell-Freitas (D01 only)
Microphysics	Morrison (double-moment) (Ye et al., 2021)
PBL Scheme	Mellor-Yamada-Janjic TKE scheme
Simulation Period	14 – 27 March 2024

Comment #10: *Line 227: Correct the citation for Kuhlmann et al. to (2024)*

Response #10: The citation has been corrected in the manuscript text.

Revised Manuscript (Lines 254 – 255): A brief description of inversion methods and specifics is provided here. A complete description of the inversion methods and algorithm can be found in Kuhlmann et al. (2024).

Comment #11: *Line 298: Please provide more justification for the fixed conversion factor fq = 1.32, as the NO2-to-NOx ratio is highly dependent on environmental conditions and emissions sources. Could WRF-Chem provide a localized factor for Bangkok?*

Response #11: Thank you for the comment. We agree that the NO2-to-NOx ratio can vary with chemical regime, emissions source, and meteorological conditions. However, in our study case, we did not use a WRF-Chem-derived localized conversion factor because the prior simulation at the time showed substantial

biases in NO_x and related chemistry before inversion, including unrealistically long NO_x lifetimes. In that case a model-derived local NO2-to-NOx ratio would likely reflect model error rather than provide an independent or more accurate constraint. We therefore retained the fixed conversion factor, fq = 1.32, as suggested in the *ddeq* framework for consistency and to avoid introducing additional uncertainty from biases in the prior chemistry.

Revised Manuscript (Lines 331 – 334): Following the *ddeq* implementation, we use:

$$Q_{NO_x} = f_q \cdot Q_{NO_2} \quad (8)$$

to obtain NO_x estimates, where f_q= 1.32, as implemented in the *ddeq* algorithm; a localized factor was not derived from WRF-Chem due to known NO_2 biases in the prior simulation that would propagate into the conversion.

Comment #12: *Line 343-347: Do the author know why does the CSF method perform better than the GP or IME approaches in recovering the model's prior emissions?*

Response #12: We suspect the performance of the CSF method is likely related to how the framework separates and constrains physical processes. In the CSF method, emissions are derived from the downwind NO_2 flux, and multiple cross-sections along the plume are used to independently estimate fluxes at different downwind distances. This provides multiple constraints on plume evolution and might reduce the sensitivity to local errors in ERA5 wind fields, plume definition, or missing satellite data.

In contrast, the IME method relies on a single integrated plume mass, making it more sensitive to plume boundaries, wind errors, and assumed decay corrections, while the GP methods depend on fitting the full 2D plume structure, which introduces sensitivity to assigned plume shape and spatial variability. Overall, the CSF method might provide a more robust mapping between observed NO_2 and emissions by leveraging multiple cross-sectional constraints. We have clarified this point in the revised manuscript.

Revised Manuscript (Lines 404 – 407): Overlaying the prior emissions shows that the model-derived CSF method most accurately recovers them, likely due to its use of downwind fluxes and multiple cross-sections across the plume, which provide additional constraints on plume evolution and reduce sensitivity to local errors in wind fields, plume definition, and missing data compared to the IME and GP methods.

Comment #13: *Figure 3b: Is the black dashed line (WRF Prior) representing the three-hour backward average of EDGAR v5 emissions? It appears visually inconsistent with the prior curve shown in Figure 4c.*

Response #13: Thank you for the comment. Yes, the black dashed line in Figure 3b represents the three-hour backward average of EDGAR v5 emissions used as an input to the optimization. This is consistent with the prior emissions shown in Figure 4a.

The apparent discrepancy with Figure 4c arises from differences in the temporal window: Figures 3b and 4a include the 23 UTC hour from the previous day in the backward averaging, whereas Figure 4c does not. Figure 4c represents the full diurnal profile of the input emissions. Values from 00 – 07 UTC are identical across figures; differences at earlier times are solely due to inclusion of the 23 UTC hour in the backward averaging. We have edited the figure caption to clarify this distinction.

Revised Manuscript (Lines 550 – 555): **Figure 4.** Results of the emissions optimization, shown as averages over the BMR. (a) Posterior optimized (solid) and prior (dashed) daytime NO_x emissions in model space, where the prior represents the three-hour backward average of EDGAR v5 emissions used as input to the optimization. (b) Same emissions as in (a) but transformed into observational space using the observational operator. The GEMS profile is shown in blue. (c) and (d) are representative of optimization results using the GEMS bias-corrected CSF results as the constraint. (e) Full diurnal cycle of updated (solid), updated+BC (solid pink), and prior (dashed) NO_x emissions. Note that, unlike (a), the prior in (e) does not include the 23 UTC hour from the previous day in the backward averaging.

Comment #14: *Line 370-378: I have concerns regarding the GEMS NO2 column scaling factor in Equation 11. Early morning model errors in boundary layer height and NO2 profiles can introduce artifacts into this scaling factor. Would it be more robust to correct GEMS NO2 columns using independent observations (e.g., Pandora, MAX-DOAS) prior to the inversion?*

Response #14: Thank you for this comment. We agree that early morning model errors in boundary layer height and NO_2 vertical structure can influence the scaling factor in Eq. (11). Our intent in applying this correction is not to adjust absolute GEMS column biases, but to account for the reduced surface sensitivity inherent to the GEMS averaging kernel in a way that is internally consistent with the model vertical distribution used in the inversion framework.

At the recommendation of both reviewers, we conducted a sensitivity analysis in which GEMS NO_2 columns within the BMR are adjusted using an independent, Pandora-derived bias prior to inversion. Based on the Pandora Bangkok evaluation (new Fig. S1), we apply an approximate scaling factor of two to the GEMS columns within the BMR. To assess the impact of this retrieval bias, we perform the inversion and subsequent model optimization framework both with and without the bias corrected emission results. The results from both configurations are now included in the main manuscript figures to explicitly illustrate the sensitivity of inferred NO_x emissions to the assumed satellite bias. In addition to $WRF_{Updated}$, we introduce a bias-corrected simulation, $WRF_{Updated+BC}$. In most cases, $WRF_{Updated+BC}$ shows improved agreement relative to $WRF_{Updated}$; however, it can lead to overestimation at times as indicated by comparisons with DC-8 and GCAS observations. We therefore retain both simulations in the manuscript to highlight the influence of the bias correction and the associated trade-offs in model performance. The manuscript has been revised throughout the reflect these additions in addition to updated statistics in Table 2.

Revised Manuscript:

(Lines 48 – 49): GEMS-constrained NO_x emissions for March 2024 are estimated to range from 2.7 to 4.3 kT $month^{-1}$ after accounting for known low biases in the GEMS retrievals.

(Lines 149 – 151): Model cases with updated anthropogenic NO_x emissions will be referred to as $WRF_{Updated}$ and $WRF_{Updated+BC}$ (later introduced).

(Lines 213 – 232): **3.1 Correcting for GEMS bias prior to inversion**

Recent independent validation studies of the operational GEMS v3 product over Bangkok and South Korea report low biases in NO_2 columns relative to ground-based sun-photometer and DOAS measurements (Bae et al., 2025; Jung et al., 2025). Bae et al. (2025) shows that GEMS v3 increasingly underestimates NO_2 relative to Pandora under high-NO_2 conditions (>1 × 10^{16} molecules cm^{-2}) as is the case for Bangkok pollution levels. In Jung et al. (2025), validation results over Bangkok indicate a pronounced low bias in GEMS tropospheric NO_2 columns relative to Pandora, with regression slopes of ~0.35 for v2.0 and ~0.28 for v3.0, indicating increasing underestimation at higher NO_2 levels. While moderate correlations ($r \approx 0.6$–0.7) suggest that GEMS captures temporal variability, column magnitudes are substantially underestimated, particularly under polluted conditions. The persistence of this behavior in the v3.0 product indicates that the low bias is not fully corrected by recent algorithm updates and is consistent with retrieval sensitivity limitations in highly polluted urban environments (Jung et al., 2025).

To address the low bias in GEMS prior to the NO_x satellite emission inversion, we first quantify the GEMS bias relative to Pandora measurements over our period of interest (14 – 27 March 2024). We compare total column GEMS NO_2 to Pandora Level 2 direct-sun total column retrieval, filtering for high quality measurements (quality flag = 10) and averaging to hourly means. GEMS columns are sampled at the nearest grid cell to each Pandora site and temporally collocated. Results are shown in Fig. S1. GEMS NO_2 columns are generally about a factor of two lower than Pandora measurements (mean bias ≈ -9.2×10^{15} molecules cm^{-2}). This factor-of-two difference persists throughout March 2024 (Fig. S1).

To assess the sensitivity of the top-down NO_x inversion framework to this bias, we apply a simple correction factor to GEMS prior to inversion. Specifically, GEMS NO_2 columns over the BMR are scaled by a factor of 1.67 derived from the Pandora comparison. The inversion is then performed with and without the bias-corrected columns.

(Lines 428 – 429): Two versions of the optimization are performed: one using the constraint derived from the bias-corrected retrievals, and one without it.

(Lines 657 – 672):

Table 2. Summary of WRF-Chem D02 validation statistics for WRF_{Base} (B) $WRF_{Updated}$ (U), and $WRF_{Updated+BC}$ (U+BC) simulations evaluated against independent observational datasets. Metrics include mean bias (MB), mean error (ME), normalized mean bias (NMB), normalized mean error (NME), root-mean-square error (RMSE), and Pearson correlation coefficient (CORR).

Dataset	Species	Model Run	MB	ME	NMB	NME	RMSE	CORR
GCAS	Tropospheric NO_2 column (molecules cm^{-2})	**B**	4.0E+15	5.1E+15	93	1.2E+2	9.3E+15	0.64
		U	-2.4E+14	1.9E+15	-5.6	45	3.4E+15	0.54

		U+BC	6.8E+14	2.3E+15	16	53	4.1E+15	0.4
DC-8	NO_xO_3 NO_2 (ppbV)	**B**	1.3	1.4	82	87	2.8	0.97
		U	-0.81	0.8	-49	49	1.4	0.93
		U+BC	-0.38	0.49	-23	30.3	0.89	0.94
	CANOE NO_2 (ppbV)	**B**	1.4	1.5	87	94	2.9	0.97
		U	-0.76	0.76	-48	48	1.3	0.93
		U+BC	-0.34	0.50	-21	31	0.88	0.94
Ground Monitors	NO_2 (ppbV)	**B**	12	12	1.3E+2	1.3E+2	15	0.61
		U	-3.0	4.5	-32	47	5.6	0.53
		U+BC	0.1	4.0	0.6	43	5.5	0.55
	NO_x (ppbV)	**B**	14	15	1.4E+2	1.4E+2	18	0.59
		U	-3.6	4.9	-34	47	6.7	0.61
		U+BC	0.02	4.8	0.2	46	7.1	0.57
	O_3 (ppbV)	**B**	-5.6	8.2	-18	27	10	0.87
		U	4.9	8.5	16	27	11	0.81
		U+BC	3.1	6.6	10	22	9.2	0.86
Pandora	Tropospheric NO_2 column (molecules cm^{-2})	**B**	7.9E+15	8.6E+15	55	59	9.6E+15	0.71
		U	-6.6E+15	6.8E+15	-46	47	8.2E+15	0.67
		U+BC	-3.8E+15	4.9 E+15	-26	33	6.4 E+15	0.70

(Lines 991 – 1042): **7.1 Implications of GEMS NO_2 retrieval biases**

In our study, the comparisons with independent aircraft and ground-based observations indicate the satellite-constrained emissions without bias-corrected presented here may represent a low estimate. In particular, the systematic low bias in GEMS NO_2 is consistent with the negative biases often seen in the $WRF_{Updated}$ simulations. Importantly, this bias could not be diagnosed using the satellite data alone.

To further evaluate the relative behavior of GEMS and airborne GCAS NO_2 columns, we compare their tropospheric columns using $WRF_{Updated+BC}$ as a common transfer framework (Fig. S21). Figure S21 compares the observed GCAS/GEMS NO_2 column ratios with ratios calculated after both datasets are mapped through $WRF_{Updated+BC}$. In the observations, GCAS columns are consistently higher than the raw GEMS columns as illustrated by the orange line, with GCAS/GEMS ratios of ~2–3 on most days and values reaching ~6–7 on 21 March. In contrast, the corresponding $WRF\text{-}GCAS_{Updated+BC}$/$WRF\text{-}GEMS_{Updated+BC}$ ratios are closer to 1:1 (~1.1–1.7), even on 21 March. This behavior is consistent with a low bias in GEMS NO_2 columns relative to GCAS. After applying the bias correction, the $GCAS/GEMS_{BC}$ ratios approach 1:1 on most days, indicating improved consistency between the datasets, and supporting the effectiveness of the applied bias correction.

When incorporating bias-corrected retrievals into the top-down inversions and NO_x emission optimization process, the resulting $WRF_{Updated\ +BC}$ run reduces the negative biases in the independent validation, in some cases bringing the model results even closer to observations as seen in the surface air quality analysis. However, this improvement is not uniform. In certain comparisons (e.g., GCAS and DC-8), $WRF_{Updated\ +BC}$ introduces a tendency toward overprediction in morning hours, highlighting tradeoffs associated with the bias correction. These results suggest that while accounting for the retrieval bias can improve mean model performance, additional uncertainties such as overestimated wind speeds and associated transport errors continue to influence the representation of modeled NO_2.

Here, the integration of ground-based, airborne, satellite, and model data provides a powerful framework not only for improving emissions but also for identifying limitations within individual observing systems. While $WRF_{Updated}$ clearly outperforms the baseline model, and $WRF_{Updated\ +BC}$ offers targeted improvements in reducing systematic bias, the combined observational evidence highlights the necessity of a multi-platform validation to fully interpret the satellite-based emission estimates.

Despite this low bias in the GEMS retrieval, the high-frequency daytime sampling provided by geostationary observations offers critical constraints on daytime variability and plume evolution that are particularly valuable for emission inversion and air quality modelling. For example, the GEMS-constrained emission adjustments presented here were critical for improving the temporal evolution of NO_x in WRF-Chem, resulting in substantial and robust improvements in model performance across independent evaluations. Future work may further benefit from continued refinement of bias-corrected GEMS products and upcoming algorithm improvements (e.g., v4), alongside improved representation of spatial representation of emissions within the model.

Comment #15: *Section 4.1: Since the authors did not directly assimilate NO2 columns, the term "satellite observations" in this context is imprecise (e.g., Line 422-424, Line 434-435). I suggest using "GEMS-informed emissions" or a similar term.*

Response #15: Yes, following the revised wording in this section, we have changed the term to reflect "satellite-derived emissions."

Revised Manuscript (Lines 490 – 492): The result is a corrected model hourly emission vector, $\hat{x}$, that balances the satellite-derived emissions with the model's prior estimate while accounting for chemistry and transport behavior (Fig. 4). Figure 4b shows the posterior profile closely reproduces the GEMS-derived daytime pattern, indicating the optimization effectively aligns model emissions with satellite-derived emissions while preserving components of the prior model behavior.

Comment #16: *Figure 4: The caption for Figure 4a and 4b requires more detail to clearly distinguish between the optimized profiles and those viewed through the observational operator.*

Response #16: We have edited the caption for Figure 4 accordingly.

Revised Manuscript (Lines 550 – 555): **Figure 4.** Results of the emissions optimization, shown as averages over the BMR. (a) Posterior optimized (solid) and prior (dashed) daytime NO_x emissions in model space, where the prior represents the three-hour backward average of EDGAR v5 emissions used as input to the optimization. (b) Same emissions as in (a) but transformed into observational (GEMS) space using the observational operator. The GEMS profile is shown in blue. (c) Full diurnal cycle of updated (solid purple), updated+BC (solid pink), and prior (dashed) NO_x emissions. Note that, unlike (a), the prior in (c) does not include the 23 UTC hour from the previous day in the backward averaging.

Comment #17: *Line 527-528: Beyond meteorology, how do inaccuracies in the spatial pattern of emissions affect the inversion framework? Do the authors have any thoughts on addressing this issue under this study's emissions inversion framework?*

Response #17: Thank you for the comment. We agree that this approach does not explicitly optimize the spatial distribution of emissions, and therefore any spatial inaccuracies in the prior are not inherently corrected by the inversion. Such errors can propagate into the forward model through biases in plume structure and transport, potentially affecting the agreement with observations as shown in this section.

As noted above, extending this framework to resolve emissions at finer scales would require additional steps. This could be achieves by performing separate inversions for different source types. A first step could be to incorporate high-resolution local priors when available, together with higher-resolution observations (e.g., GCAS) that can better resolve variability in NO_2.

Revised Manuscript (Lines 621 – 623): Additionally, while the inversion framework adjusts emission magnitudes temporally, it does not correct spatial errors in plume transport or source distribution (see Sect. 7.2 for further discussion).

Comment #18: *Line 643-644: Note that the highest observed O3 peak actually occurred on March 22, rather than March 20–21.*

Response #18: We have updated the phrasing here to address this comment.

Revised Manuscript (Lines 747 – 750): The 20 – 21 March is shaded in grey to indicate the period where the model mis-represented synoptic conditions resulting in enhancements in the observed concentrations due to stagnation. During this time, NO_2 and NO_x reached as high as 49, and 59 ppbV, respectively. O_3 peaked at 103 ppbV the following day, on 22 March.

Comment #19: *Line 655-661: The shift in O3 peak time to the early morning in the WRFupdated rum (Figure S12a) does not align with observations. Does this discrepancy suggest the emission inversion timing might be incorrect?*


Response #19: Thank you for the comment. We agree that the earlier O_3 peak in $WRF_{Updated}$ D02 indicates the model does not fully reproduce the observed diurnal evolution of ozone. However, we do not interpret this as evidence that the inferred NO_x emission timing is incorrect.

O_3 production in Bangkok occurs under a mixed NO_x-VOC sensitivity regime, as supported by recent ASIA-AQ analyses (Cho et al., 2026). In such regimes, the timing and magnitude of O_3 are not only controlled by NO_x emissions alone, but also depend strongly on VOC availability, boundary layer evolution, and transport processes. Importantly, the same emission inversion is applied in both D01 and D02, yet the early O_3 peak is only present in D02 (Figure S12a-b). This suggests that the discrepancy is more likely driven by domain-dependent differences in meteorology and chemical processing, rather than an error in emission timing. We therefore interpret the early O_3 peak in D02 as reflecting remaining model chemistry (as VOC emission was not altered in this study) rather than a direct bias in the inferred NO_x emission timing.

We have updated the supplemental figure (now Figure S12) to illustrate the NO_x diurnal cycles as well to support this comparison.

Revised Manuscript (Lines 762 – 768): Average diurnal cycles of NO_x for D02, D01 are further illustrated in Fig. S12a, c. Observed NO_x shows a clear morning peak and lower midday concentrations driven by boundary layer evolution. WRF_{Base} overestimates NO_x and exaggerates the morning peak while $WRF_{Updated}$ (dark purple) substantially improves both magnitude and timing but underestimates the evening peak values. $WRF_{Updated+BC}$ shows mixed performance with an overprediction of the morning peak while the evening peak aligns more closely with observations compared to $WRF_{Updated}$, indicating a trade-off in how the bias correction might redistribute NO_x across the diurnal cycle in the updated models.

Revised Manuscript (Lines 776 – 780): While $WRF_{Updated}$ exhibits an earlier O_3 peak in D02 compared to observations, this shift is not present in D01 despite the same emission inversion being applied. This likely suggests the discrepancy is driven by domain-dependent differences in meteorological or chemical processes, rather than an error in the inferred NO_x emission timing.



Revised Supplement (Lines245 - 250): **Figure S12.** Average diurnal cycles of NO_x and O_3 for D02 (a,b) and D01 (c,d). Comparison of WRF_{Base} (dotted), $WRF_{Updated}$, and $WRF_{Updated+BC}$ simulations against Thailand Pollution Control Department (PCD) ground monitor network observations during the ASIA-AQ deployment period (14-27 March 2024).

Comment #20: *Line 677-679: Please clarify which Pandora data (e.g., direct sun or sky scan), data screening, and temporal collocation methods were used for the evaluation*

Response #20: Thank you for the comment. The Pandora evaluation was conducted using the Bangkok Level 2 Pandora NO_2 direct-sun total column product. Only high-quality measurements (quality flag = 10) were retained. The observations were aggregated to hourly means prior to comparison. To approximate the tropospheric columns, the Pandora total column was corrected by subtracting the closest GEMS stratospheric NO_2 column at the nearest satellite pixel. WRF-Chem NO_2 columns were computed at the nearest model grid cell to the Pandora site and were temporally collocated with the hourly Pandora data. We have revised the manuscript to provide this clarifying information.

Revised Manuscript (Lines 796 – 800): Pandora NO_2 evaluation used Level 2 direct-sun total column retrievals, filtered for high-quality measurements (quality flag = 10), and averaged to hourly means. Tropospheric columns from Pandora were estimated by subtracting coincident or closest GEMS stratospheric NO_2 at the nearest satellite pixel. Model columns were sampled at the nearest grid cell to the Pandora site and temporally matched to observations.

Comment #21: *Equation 17: Please define SW (Scattering Weights).*

Response #21: We have revised the manuscript to include this definition.

Revised Manuscript (Lines 879 – 880): Here, SW_i represents the scattering weight for layer i, representing the sensitivity of the measured radiance to NO_2 in that layer.

Comment #22: *Figures 8–9: I suggest converting the time from UTC to Local Standard Time (LST) for better interpretability.*

Response #22: In Figure 8 and corresponding Figure S16 we have changed the times to reflect local standard time in the figure. We have edited the GEMS image to reflect local time in Figure 9c.

Comment #23: *S19 is provided in supplement materials but not used in the manuscript.*

Response #23: As suggested in the earlier comment, we have revised the manuscript and Supplement figure references to ensure each figure is referenced accordingly.

Comment #24: *Line 804: This reference should likely be Figure S17.*

Response #24: We have adjusted the figure reference accordingly to match the updated Supplement.

Comment #25: *Lines 925–929: Any model output (e.g., WRFbase) can be used as a transfer standard for accessing GEMS vs. GCAS, though high-resolution models are preferred.*

Response #25:
Thanks for the comment. We agree that the model output can serve as a transfer standard to facilitate comparisons between different observing systems, including GEMS and GCAS and have revised the text to clarify this point.

Revised Manuscript (Lines 1003 – 1005): While model output (e.g., WRF_{Base}) can serve as a transfer standard for comparing different observing systems, these results highlight the importance of integrating ground-based, airborne, satellite, and model information to robustly identify biases and improve emission estimates.

Anonymous Referee #2

General Comments:

This manuscript investigates daytime hourly Nitrogen Oxides (NOx) emissions over the Bangkok Metropolitan Region (BMR) using hourly Geostationary Environment Monitoring Spectrometer (GEMS) Nitrogen Dioxide (NO2) observations. The authors apply the Cross-Sectional Flux (CSF) method via the ddeq Python library to derive satellite-based emission estimates, which are then used to update the diurnal NOx profile in nested WRF-Chem simulations at 20 km and 4 km resolutions. The updated model is evaluated against independent multi-platform measurements from the 2024 ASIA-AQ campaign, including

the Thailand Pollution Control Department (PCD) surface monitors, Pandora, the GEOstationary Coastal and Air Pollution Events (GEO-CAPE) Airborne Simulator (GCAS), DC-8, and the High Spectral Resolution Lidar (HSRL)-2 lidar observations.

This work is well-motivated given the growing availability of hourly Geostationary Earth Orbit (GEO)-based composition retrievals and the persistent uncertainty in urban NOx inventories across Southeast Asia. The effort to constrain the daytime structure of emissions, rather than applying a single scaling factor, is a meaningful methodological advance, and the multi-platform evaluation substantially strengthens confidence in the results.

However, several methodological assumptions, particularly the inconsistency in meteorological fields used across different stages of the analysis, require further clarification. Additionally, while the authors correctly acknowledge a systematic low bias in the GEMS v3 NO2 product, the manuscript would benefit from a clearer discussion of how this bias propagates into the derived emission estimates.

Response: Thank you for the helpful comments. We address these assumptions in detail in the responses below. In the revised manuscript, we now explicitly account for the systematic low bias in the GEMS product and introduce an additional simulation that incorporates this correction ($WRF_{Updated+BC}$). Details of these updates are provided in the responses that follow.

Specific Comments:

Comment #1: *The use of different meteorological fields at different stages of the analysis needs clarification. ERA5 winds are used for the CSF inversion because WRF-Chem winds are biased high, yet the NOx lifetimes fed into the optimization operator H appear to come from WRF-Chem. The remaining negative biases in WRF-Updated are subsequently attributed to WRF-Chem wind overprediction. The authors should add a short paragraph that clearly separates (a) how the wind bias affects the initial inversion step and (b) how it affects the final model evaluation. It should also be clarified whether the lifetimes used in H are physically consistent with ERA5- or WRF-Chem-based transport, and whether using ERA5-derived lifetimes would materially change the optimized emission profile.*

Response #1: We thank the reviewer for this comment and have clarified this point in the manuscript. The NO_x lifetimes used in the forward operator (H) are not derived-from WRF-Chem. Instead, they are obtained from the GEMS-based CSF analysis, where lifetime is fitted from the observed along-plume decay using ERA5 wind fields. As such, the lifetimes are internally consistent with ERA5-based transport. This makes the satellite emission estimates completely independent of the WRF-Chem simulations.

WRF-Chem wind biases affect primarily the model evaluation stage. When the updated emissions are simulated within WRF-Chem, the model's tendency to overestimate wind speed contributes to the negative remaining biases in $WRF_{Updated}$. Because the lifetime is derived from the GEMS + ERA5 framework, we are already including it within our optimization approach.

Revised Manuscript (Lines 429 – 434): The three-hour window provided the best agreement with the model-derived CSF emissions as shown in Fig 3b, 3c. Within this window, the contribution from each prior

hour depends on the atmospheric lifetime of NO_2 for that hour. We use hourly lifetimes, τ, derived from the GEMS-based CSF inversion in Eq. (7), where τ is fitted from observed along-plume decay using ERA5 winds, and is therefore independent of WRF-Chem meteorology. This ensures the satellite-derived emission estimates are completely independent of the WRF-Chem simulations. For reference, WRF-Chem-derived decay times are generally longer than those inferred from GEMS, reflecting differences in modeled vs. observed loss and dilution processes.

Comment #2: *As the authors acknowledge, the remaining negative biases in WRF-Updated are consistent with the known low bias in GEMS v3 NO2. However, this bias is compounded under high-NO2 conditions, precisely the conditions characteristic of Bangkok, meaning that emissions derived directly from the uncorrected satellite data are likely biased low. Consequently, the conclusion that EDGAR v5 overestimates NOx by ~75% should be stated with appropriate caveats, as the true magnitude of overestimation may be smaller once the satellite bias is accounted for.*

Response #2: Thank you for this helpful suggestion. At the recommendation of both reviewers, we conducted a sensitivity analysis in which GEMS NO_2 columns within the BMR are adjusted using an independent, Pandora-derived bias prior to inversion. Based on the Pandora Bangkok evaluation (new Fig. S1), we apply an approximate scaling factor of two to the GEMS columns within the BMR. To assess the impact of this retrieval bias, we perform the inversion and subsequent model optimization framework both with and without the bias corrected emission results.

The results from both configurations are now included in the main manuscript figures to explicitly illustrate the sensitivity of inferred NO_x emissions to the assumed satellite bias. In addition to $WRF_{Updated}$, we introduce a bias-corrected simulation, $WRF_{Updated+BC}$. In most cases, $WRF_{Updated+BC}$ shows improved agreement relative to $WRF_{Updated}$; however, it can lead to overestimation at times as indicated by comparisons with DC-8 and GCAS observations. We therefore retain both simulations in the manuscript to highlight the influence of the bias correction and the associated trade-offs in model performance. The manuscript has been revised throughout the reflect these additions in addition to updated statistics in Table 2.

Revised Manuscript:

(Lines 48 – 49): GEMS-constrained NO_x emissions for March 2024 are estimated to range from 2.7 to 4.3 kT $month^{-1}$ after accounting for known low biases in the GEMS retrievals.

(Lines 149 – 151): Model cases with updated anthropogenic NO_x emissions will be referred to as $WRF_{Updated}$ and $WRF_{Updated+BC}$ (later introduced).

(Lines 213 – 232): **3.1 Correcting for GEMS bias prior to inversion**

Recent independent validation studies of the operational GEMS v3 product over Bangkok and South Korea report low biases in NO_2 columns relative to ground-based sun-photometer and DOAS measurements (Bae et al., 2025; Jung et al., 2025). Bae et al. (2025) shows that GEMS v3 increasingly underestimates NO_2 relative to Pandora under high-NO_2 conditions (>1 × 10^{16} molecules cm^{-2}) as is the case for Bangkok

pollution levels. In Jung et al. (2025), validation results over Bangkok indicate a pronounced low bias in GEMS tropospheric NO_2 columns relative to Pandora, with regression slopes of ~0.35 for v2.0 and ~0.28 for v3.0, indicating increasing underestimation at higher NO_2 levels. While moderate correlations ($r \approx 0.6–0.7$) suggest that GEMS captures temporal variability, column magnitudes are substantially underestimated, particularly under polluted conditions. The persistence of this behavior in the v3.0 product indicates that the low bias is not fully corrected by recent algorithm updates and is consistent with retrieval sensitivity limitations in highly polluted urban environments (Jung et al., 2025).

To address the low bias in GEMS prior to the NO_x satellite emission inversion, we first quantify the GEMS bias relative to Pandora measurements over our period of interest (14 – 27 March 2024). We compare total column GEMS NO_2 to Pandora Level 2 direct-sun total column retrieval, filtering for high quality measurements (quality flag = 10) and averaging to hourly means. GEMS columns are sampled at the nearest grid cell to each Pandora site and temporally collocated. Results are shown in Fig. S1. GEMS NO_2 columns are generally about a factor of two lower than Pandora measurements (mean bias $\approx -9.2 \times 10^{15}$ molecules cm^{-2}). This factor-of-two difference persists throughout March 2024 (Fig. S1).

To assess the sensitivity of the top-down NO_x inversion framework to this bias, we apply a simple correction factor to GEMS prior to inversion. Specifically, GEMS NO_2 columns over the BMR are scaled by a factor of 1.67 derived from the Pandora comparison. The inversion is then performed with and without the bias-corrected columns.

(Lines 428 – 429): Two versions of the optimization are performed: one using the constraint derived from the bias-corrected retrievals, and one without it.

(Lines 657 – 672):

Table 2. Summary of WRF-Chem D02 validation statistics for WRF_{Base} (B) $WRF_{Updated}$ (U), and $WRF_{Updated+BC}$ (U+BC) simulations evaluated against independent observational datasets. Metrics include mean bias (MB), mean error (ME), normalized mean bias (NMB), normalized mean error (NME), root-mean-square error (RMSE), and Pearson correlation coefficient (CORR).

Dataset	Species	Model Run	MB	ME	NMB	NME	RMSE	CORR
GCAS	Tropospheric NO_2 column (molecules cm^{-2})	**B**	4.0E+15	5.1E+15	93	1.2E+2	9.3E+15	0.64
		U	-2.4E+14	1.9E+15	-5.6	45	3.4E+15	0.54
		U+BC	6.8E+14	2.3E+15	16	53	4.1E+15	0.4
DC-8	NO_xO_3 NO_2 (ppbV)	**B**	1.3	1.4	82	87	2.8	0.97
		U	-0.81	0.8	-49	49	1.4	0.93

		U+BC	-0.38	0.49	-23	30.3	0.89	0.94
	CANOE NO_2 (ppbV)	**B**	1.4	1.5	87	94	2.9	0.97
		U	-0.76	0.76	-48	48	1.3	0.93
		U+BC	-0.34	0.50	-21	31	0.88	0.94
Ground Monitors	NO_2 (ppbV)	**B**	12	12	1.3E+2	1.3E+2	15	0.61
		U	-3.0	4.5	-32	47	5.6	0.53
		U+BC	0.1	4.0	0.6	43	5.5	0.55
	NO_x (ppbV)	**B**	14	15	1.4E+2	1.4E+2	18	0.59
		U	-3.6	4.9	-34	47	6.7	0.61
		U+BC	0.02	4.8	0.2	46	7.1	0.57
	O_3 (ppbV)	**B**	-5.6	8.2	-18	27	10	0.87
		U	4.9	8.5	16	27	11	0.81
		U+BC	3.1	6.6	10	22	9.2	0.86
Pandora	Tropospheric NO_2 column (molecules cm^{-2})	**B**	7.9E+15	8.6E+15	55	59	9.6E+15	0.71
		U	-6.6E+15	6.8E+15	-46	47	8.2E+15	0.67
		U+BC	-3.8E+15	4.9 E+15	-26	33	6.4 E+15	0.70

(Lines 991 – 1042): **7.1 Implications of GEMS NO_2 retrieval biases**

In our study, the comparisons with independent aircraft and ground-based observations indicate the satellite-constrained emissions without bias-corrected presented here may represent a low estimate. In particular, the systematic low bias in GEMS NO_2 is consistent with the negative biases often seen in the $WRF_{Updated}$ simulations. Importantly, this bias could not be diagnosed using the satellite data alone.

To further evaluate the relative behavior of GEMS and airborne GCAS NO_2 columns, we compare their tropospheric columns using $WRF_{Updated+BC}$ as a common transfer framework (Fig. S21). Figure S21 compares the observed GCAS/GEMS NO_2 column ratios with ratios calculated after both datasets are mapped through $WRF_{Updated+BC}$. In the observations, GCAS columns are consistently higher than the raw GEMS columns as illustrated by the orange line, with GCAS/GEMS ratios of ~2–3 on most days and values reaching ~6–7 on 21 March. In contrast, the corresponding $WRF\text{-}GCAS_{Updated+BC}/WRF\text{-}GEMS_{Updated+BC}$ ratios are closer to 1:1 (~1.1–1.7), even on 21 March. This behavior is consistent with a low bias in GEMS NO_2 columns relative to GCAS. After applying the bias correction, the $GCAS/GEMS_{BC}$ ratios approach 1:1 on most days, indicating improved consistency between the datasets, and supporting the effectiveness of the applied bias correction.

When incorporating bias-corrected retrievals into the top-down inversions and NO_x emission optimization process, the resulting $WRF_{Updated\ +BC}$ run reduces the negative biases in the independent validation, in some cases bringing the model results even closer to observations as seen in the surface air quality analysis. However, this improvement is not uniform. In certain comparisons (e.g., GCAS and DC-8), $WRF_{Updated\ +BC}$ introduces a tendency toward overprediction in morning hours, highlighting tradeoffs associated with the bias correction. These results suggest that while accounting for the retrieval bias can improve mean model performance, additional uncertainties such as overestimated wind speeds and associated transport errors continue to influence the representation of modeled NO_2.

Here, the integration of ground-based, airborne, satellite, and model data provides a powerful framework not only for improving emissions but also for identifying limitations within individual observing systems. While $WRF_{Updated}$ clearly outperforms the baseline model, and $WRF_{Updated\ +BC}$ offers targeted improvements in reducing systematic bias, the combined observational evidence highlights the necessity of a multi-platform validation to fully interpret the satellite-based emission estimates.

Despite this low bias in the GEMS retrieval, the high-frequency daytime sampling provided by geostationary observations offers critical constraints on daytime variability and plume evolution that are particularly valuable for emission inversion and air quality modelling. For example, the GEMS-constrained emission adjustments presented here were critical for improving the temporal evolution of NO_x in WRF-Chem, resulting in substantial and robust improvements in model performance across independent evaluations. Future work may further benefit from continued refinement of bias-corrected GEMS products and upcoming algorithm improvements (e.g., v4), alongside improved representation of spatial representation of emissions within the model.

Comment #3: *The 24-hour emission scaling approach warrants further justification. Since GEMS constrains emissions only between 08:00 and 14:00 LT, applying a single average scaling factor to unconstrained evening and nighttime hours is a strong assumption, particularly given Bangkok's well-documented bimodal traffic patterns. It is unclear whether this extrapolation is physically reasonable, especially during the evening rush hour. The authors should provide a brief justification or, where possible, compare the implied evening scaling against an independent diurnal reference such as the THAI-KMUTT local inventory profile.*


Response #3: Thank you for the helpful comment. We agree that applying a single mean scaling factor derived from the GEMS-constrained window (08:00-14:00 LT) to the full 24-hour profile is an assumption and warrants justification. In our framework, the satellite observations primarily constrain the emissions during the daytime window. Applying the scaling uniformly during the nighttime is necessary to maintain physical consistency. If the scaling was applied only during the GEMS-constrained hours, it would introduce an artificial discontinuity between daytime and nighttime emissions as the GEMS constrain substantially reduces daytime NO_x (leaving nighttime emissions unscaled would result in an unrealistic step change in the diurnal profile).

To evaluate the physical plausibility of the resulting 24-hour profile, we compared it with the THAI-KMUTT local inventory (see below). The prior inventory (EDGAR) already includes a bimodal structure, although the second (evening peak) is partially obscured by an overall high emission bias. The updated profile preserves this temporal structure during the evening hours while reducing the overall magnitude. The amplitude/timing of the peak is weaker than in THAI-KMUTT, suggesting greater uncertainty outside the GEMS-constrained window. It is also important to note that NO_x emissions in the model represent a combination of sources (e.g., traffic, area sources, and other anthropogenic activities), not traffic alone. As such, perfect agreement with a traffic-based or locally derived diurnal profile is not expected.

We also emphasize that the prior inventory provides a regionally representative temporal profile (developed for broader Asian emissions rather than specifically for Bangkok) which was available at the time and thus may not fully capture the local diurnal variability. Future work should incorporate the best available temporal profiles where available.

Lastly, this approach is consistent with prior satellite-constrained emission studies. For example, a similar methodology was applied using a single daytime satellite constraint to scale emissions without resolving the full diurnal profile in a KORUS-AQ field study over Seoul (Goldberg et al. 2019). Our use of GEMS extends this framework by leveraging multiple daytime observations but relies on the same assumption for extrapolation beyond the observational window.

We have added some sentences to the main manuscript to further clarify this.



Revised Manuscript (Lines 482 – 489): Because GEMS provides constraints only during the daytime overpass window, the derived scaling below primarily reflects daytime emission adjustments. We therefore apply the mean scaling factor uniformly during the nighttime hours to preserve the prior temporal structure and avoid introducing artificial discontinuities between the constrained and unconstrained hours, particularly given the large daytime emission reductions. This approach is consistent with previous satellite-constrained emission studies (e.g., Goldberg et al. (2019)) which constrain emission magnitude based on a single daytime value derived from OMI. Where available, locally derived temporal profiles should be used to better represent regional emission patterns; in this study, our prior profile reflects the best available regional representation for Asia at the time of study.

Technical Corrections:

Page 1, Line 24: *"College Park, postal code, USA" → "College Park, USA"*
Page 4, Line 103*: "e.g..," → "e.g.,"*
Page 4, Line 106*: "(Kim et al., 2020; Park et al., 2025).Recent" → "(Kim et al., 2020; Park et al., 2025). Recent"*
Page 4, Line 122*: "1; Fig 1)(Agarwal et.." → "1; Fig 1) (Agarwal et …"*
Page 5, Line 134: *Does "1 km2 resolution" mean the 1 km × 1 km resolution?*
Page 8, Line 220: *I don't know if it is better to write from "Sentinel5P-TROPOMI" to "Sentinel-5P TROPOMI".*
Page 8, Line 225: *"LEO observations For our" → "LEO observations. For our".*
Page 8, Line 227: *"Kuhlmann et al. (2014)" → "Kuhlmann et al. (2024)".*
Page 14, Line 376: *"where larger correction (e.g., 1.33) where present" → "where larger correction (e.g., 1.33) were present".*
Page 25, Line 610: *"from that fact that" → "from the fact that".*
Page 32, Line 803: *"(St. Clair et al., 2019) .We" → "(St. Clair et al., 2019). We".*
There is no definition of the HSRL-2.
This is not significant, but authors can use the consistent word:
1. modelled or modeled
2. re-grided or re-gridded

Response: Thank you for the clarifications. In the revised manuscript we have addressed the above technical corrections through tracked changes.

NO_x Emissions Constraints from GEMS NO_2 Retrievals: Inversion Methodology and Air Quality Model Evaluation in Bangkok using ASIA-AQ Multi-Platform Observations

Julianna A. Christopoulos[1], Pablo E. Saide[1,2], Manas R. Mohanty[1], Nattamon Maneenoi[1], Jhoon Kim[3], Laura Judd[4], Katherine R. Travis[4], Savitri Garivait[5,6], Agapol Junpen[5,6], Kazuyuki Miyazaki[7,8], Jinkyul Choi[7,8], Takashi Sekiya[98], David Peterson[109], Theodore M. McHardy[109], Nicholas Gapp[1110], Jason M. St. Clair[12,1311,12], Erin Delaria[14,1513,14], Glenn M. Wolfe[1211], Abby Sebol[1514], Alessandro Franchin[1615], Changmin Cho[1615], Morgan L. Silverman[4,174,16], James H. Crawford[4]

[1]Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, Los Angeles, CA, USA
[2]Institute of the Environment and Sustainability, University of California, Los Angeles, Los Angeles, CA, USA
[3]Department of Atmospheric Sciences, Yonsei University, Seoul, South Korea
[4]NASA Langley Research Center, Hampton, VA, USA
[5]The Joint Graduate School of Energy and Environment, King Mongkut's University of Technology, Thonburi, Thailand
[6]Center of Excellence on Energy Technology and Environment, PERDO, Ministry of Higher Education, Science, Research and Innovation, Bangkok, Thailand
[7]Jet Propulsion Laboratory (JPL), California Institute of Technology, Pasadena, 91109 CA, USA
[8]Joint Institute for Regional System Science and Engineering (JIFRESSE), University of California, Los Angeles, Los Angeles, CA, USA
[98]Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan
[109]U.S. Naval Research Laboratory, Monterey, CA, USA
[1110]Science Applications International Corporation, Monterey, CA
[1211]Atmospheric Chemistry and Dynamics Laboratory, NASA Goddard Space Flight Center, Greenbelt, USA
[1312]GESTAR II, University of Maryland Baltimore Country, Baltimore, USA
[1413]ESSIC, University of Maryland, College Park, ~~postal code,~~ USA
[1514]Department of Atmospheric and Oceanic Science, University of Maryland, College Park, USA
[1615]Atmospheric Chemistry Observations and Modeling Laboratory, NSF National Center for Atmospheric Research, Boulder, CO, USA
[1716]Science Systems and Applications, Inc. Hampton, VA, USA

Correspondence to: Julianna A. Christopoulos (juliechristo@g.ucla.edu)

Abstract. Nitrogen Dioxide (NO_2) is a key component of tropospheric chemistry and air quality, yet large uncertainties persist in regional NO_x emissions across rapidly developing megacities in Southeast Asia. Observations from the Geostationary Emissions Monitoring Spectrometer (GEMS) provide new constraints on anthropogenic NO_2 variability, while the 2024 NASA Airborne and Satellite Investigation of Asian Air Quality (ASIA-AQ) campaign, offers an extensive, independent dataset for model evaluation. Here, we examine air quality in Bangkok using coarse (20 km) and high-resolution (4 km) WRF-Chem simulations during ASIA-AQ. We develop a top-down framework that uses hourly GEMS NO_2 columns to derive constraints on the daytime cycle of NO_x emissions. Emissions are first estimated from GEMS using a Cross-Sectional Flux (CSF) inversion and then incorporated into WRF-Chem through a novel optimization that reshapes the magnitude and daytime structure of NO_x while accounting for lifetime and satellite vertical sensitivity. GEMS-constrained NO_x emissions for March 2024 are estimated to range from 2.7 to 4.3 kT $month^{-1}$ after accounting for known low biases in the GEMS retrievals. ~~GEMS-constrained NO_x emissions for March 2024 are estimated at 2.7 kT $month^{-1}$ over Bangkok, approximately 75% lower than EDGAR v5.~~ ≈Re-running WRF-Chem with the updated emissions leads to substantial improvements in modeled NO_2 magnitude and temporal variability when evaluated against independent ground-based, Pandora, and airborne measurements. Remaining negative biases are consistent with a systematic low bias in the GEMS v3 NO_2 product ~~product that cannot be diagnosed using satellite data alone~~, highlighting the importance of multi-platform evaluation using independent observations. Together, these results demonstrate the value of hourly geostationary observations combined with high-resolution modeling as a scalable pathway for improving urban NO_x emissions estimates and air quality simulations in Southeast Asia.

1 Introduction

The troposphere contains a variety of pollutants and aerosols that degrade air quality and affect human health (Chen and Chen, 2021; Fuller et al., 2022; Shetty et al., 2023). Nitrogen oxides (NO_x = NO + NO_2) are primary pollutants with significant variability in space and time (Seinfeld and Pandis, 2016). Information on NO_x sources is crucial in defining its concentration and distribution (Miyazaki et al., 2019). Nitrogen dioxide (NO_2) is fundamental to air quality and atmospheric chemistry, as it is the primary precursor to surface ozone (O_3) and nitrate aerosols (Pörtner et al., 2022) and is independently linked to the development of pediatric asthma (Anenberg et al., 2022). It is thus inherent that NO_2, and its sources, are well quantified and studied to gain insights into its environmental impact.

Over the past two decades, satellite observations of tropospheric NO_2 have revealed substantial regional variability in emissions, particularly over industrial and urban areas (Goldberg et al., 2024; Park et al., 2025; Rey-Pommier et al., 2025). While strict air quality policies in North America and Europe have resulted in significant NO_2 reductions, many developing

regions, especially the megacities of Southeast Asia, have seen increases (Elguindi et al., 2020; Georgoulias et al., 2019; Miyazaki et al., 2017; Park et al., 2025; Sicard et al., 2023). Thailand, for example, has undergone rapid industrialization, urbanization, and economic growth over the past 30 years, with most of this development occurring in the Bangkok Metropolitan Region (BMR) (Thailand Office of the National Economic and Social Development Board, World Bank, 2017) (Uttamang et al., 2018). This has led to increased emissions from vehicular traffic and industrial activity, resulting in a sustained degradation of air quality (Uttamang et al., 2018). Since the mid 1990s, the BMR has frequently exceeded Thailand's National Ambient Air Quality Standards (NAAQS) for particulate matter (PM) (25 $\mu g\ m^{-3}$) and O_3 (100 ppb), particularly during the dry season (February-May) (Kumar et al., 2012; Uttamang et al., 2018, 2020, 2023). Within the BMR, there have been initiatives to reduce health impacts and exposure related to $PM_{2.5}$. For example, the National Agenda Action Plan on "Solving the Pollution Problems of Particulate Matter" motivated implementation measures in transport, industry, and waste sectors (Aung et al., 2025). However, achieving the air quality standard has remained an issue. Modeling studies have shown that O_3 levels typically peak between January and March, coinciding with increased solar radiation, higher temperatures, elevated humidity, and prevailing northeasterly winds during the Northeast monsoon season. These meteorological conditions, combined with rising emissions, often contribute to O_3 pollution episodes in the region (Uttamang et al., 2020). A key limitation of these air quality modeling studies is the uncertainty in bottom-up anthropogenic emissions inventories, which remain a significant source of error. In the BMR, uncertainties in regional emissions can be as large as a factor of 2 or higher (Bond et al., 2004, 2007; Smith et al., 2011; Uttamang et al., 2020). These emissions uncertainties limit our ability to accurately simulate pollutant concentrations and assess the effectiveness of emission control strategies. Thus, to address this issue, new observational capabilities that can directly capture emission variability at fine spatial (< 10 km) and temporal (hourly) scales are needed.

We are currently entering a new era of satellite atmospheric composition monitoring with the launch of three geostationary (GEO) imaging spectrometers covering a large component of the Northern Hemisphere. The Geostationary Environmental Monitoring Spectrometer, GEMS (Kim et al., 2020), Tropospheric Emissions: Monitoring of Pollution, TEMPO (Zoogman et al., 2017), and Sentinel-4 (Gulde et al., 2017), now provide unprecedented spatial and temporal resolution of atmospheric constituents. These instruments offer hourly observations that resolve daytime variability of key pollutants, including NO_2. Until now, most studies of top-down NO_x emissions have relied on once-daily measurements from low Earth orbit (LEO) satellites (i.e., OMI, TROPOMI) requiring assumptions about diurnal emission and chemistry patterns that introduce uncertainties when coupled with chemical transport models, particularly those arising from coarse spatial resolution (> 20 km), emissions inventories, and chemical mechanisms (Park et al., 2025). Although these measurements have been invaluable for global and long-term NO_x assessments, they cannot fully capture the pronounced sub-daily variability in NO_2 driven by emissions, chemistry, and transport. GEO satellites now provide the capability to directly observe this hourly variability (Park et al., 2025). For example, using 12 km WRF-Chem simulations Hsu et al. (2026) showed that TEMPO-derived top-down NO_x

emissions are broadly consistent with TROPOMI and bottom-up inventories, but noted coarse model resolution (e.g., 12 km) can limit the representation of coastal meteorology and chemical nonlinearity.

GEMS was the first UV-visible hyper-spectrometer in geostationary orbit and was launched on GEO-KOMPSAT-2B on February 18, 2020. GEMS measurements allow the observation of air quality constituents (e.g.~~,~~ NO_2, SO_2, O_3, HCHO, CHOCHO, and aerosols) at a spatial resolution of 3.5 x 7.7 km^2 at the center of its field of regard and was the first space-based instrument to provide hourly observations of these species. The GEMS field of regard covers 20 countries in Asia, E-W from Japan to India, and N-S from Mongolia to Indonesia (Kim et al., 2020; Park et al., 2025). Recent work has demonstrated the potential of GEMS and air quality models to estimate top-down NO_x emissions over major Asian cities during the summertime (de Foy and Schauer, 2022; Park et al., 2024, 2025). These studies highlight the need for comprehensive validation involving independent observations. The Airborne and Satellite Investigation of Asian Air Quality, ASIA-AQ, has provided an excellent opportunity to conduct extensive validation of air pollutants across multiple Asian megacities, including Bangkok (ASIA-AQ White Paper | ASIA-AQ, 2025).

In this manuscript, we aim to address these gaps specifically near Bangkok, Thailand by deriving emissions and examining fine- (4 km) and coarse- (20 km) resolution Weather Research and Forecasting model coupled with Chemistry (WRF-Chem) simulations driven by them during mid-March 2024. First, we derive daytime hourly NO_x emissions from GEMS over the BMR in March 2024. Next, we use the GEMS emissions to constrain hourly model emissions through a novel optimization technique. To conclude, we assess model performance using GEMS and independently with ground-monitor information, Pandora site measurements, and airborne measurements collected during the ASIA-AQ campaign.

2 Model configuration and experimental design

To simulate air quality over the BMR, we use a regional simulation of WRF-Chem v4.2.2 in a research configuration (Table 1; Fig. 1) (Agarwal et al., 2024; Anav et al., 2024; Gao and Zhou, 2024; Skamarock et al., 2019). The model setup follows previous WRF-Chem implementations in the Korea-United States Air Quality field study (KORUS-AQ) for air quality studies over the Seoul Metropolitan Area (Choi et al., 2020; Goldberg et al., 2019; Lennartson et al., 2018; Park et al., 2021; Saide et al., 2020). The model was driven by reanalysis meteorology, NCEP Final Reanalysis (NCEP FNL) ~~Final Operational Global Analysis data (FNL)~~, and Copernicus Modeling Service (CAMS) chemical boundary conditions (Commerce, 2000; Inness et al., 2019). Meteorology was constrained using four-dimension data assimilation (FDDA) via grid nudging applied to model domains using the 0.25° NCEP FNL reanalysis fields to improve large-scale wind representation. A two-domain approach with one-way nesting was incorporated to gauge strengths between coarse- and high- resolution simulations. D01 (20 km)

spans a large portion of the GEMS field of regard, covering primary transboundary pollution sources (e.g., deserts in China/India, anthropogenic emissions from China/India). A nested 4 km domain (D02) is centered over Bangkok (Fig. 1).

Original anthropogenic emissions of trace gases and primary aerosols were based on the Emissions Database for Global Atmospheric Research (EDGAR v5) (Crippa et al., 2020) inventory at 0.1° x 0.1° horizontal resolution. For D02, EDGAR emissions for the transport and industrial sectors were downscaled using the Open-source Data Inventory for Anthropogenic Carbon Dioxide emission inventory (ODIAC) at 1 km x 1km^2 resolution (Oda et al., 2018). This is done by distributing EDGAR emissions into the 1 km^2 grid using ODIAC CO_2 as a spatial proxy in a mass conserving way. Biogenic, dust, sea salt, and fire emissions were computed within WRF-Chem using the model's full-chemistry emission modules (Table 1). In this configuration, dust and sea-salt emissions are calculated every chemistry timestep, while biogenic emissions (MEGAN) are updated every 30 minutes. For fire emissions, plume rise is enabled, in which the injection heights are diagnosed using the plume-rise parameterization and emissions are vertically distributed (typically ~80% at the surface with the remainder aloft). The model was configured using the RACM-MADE-VBS (Ahmadov et al., 2012; Tuccella et al., 2015) chemical mechanism and physics parameterizations selected based on prior campaign experience (e.g., KORUS-AQ), with updates to better represent secondary organic aerosol formation, heterogenous chemistry, and aerosol properties. The modal aerosol scheme in MADE-VBS tracks both particle mass and number, allowing mode diameters to evolve dynamically with aerosol aging and growth processes. Hereafter, we will refer to this model configuration as WRF_{Base}. Model cases with updated anthropogenic NO_x emissions will be referred to as $WRF_{Updated}$ and $WRF_{Updated+BC}$ (later introduced). ~~The model with updated anthropogenic NO_x emissions (later introduced) will be referred to as $WRF_{Updated}$.~~

Table 1 WRF-Chem base (WRF_{Base}) model configuration and input datasets.

Component	Configuration/Source
Model Version	WRF-Chem v4.2.2
Domain	D01: 20 km (regional); D02: 4 km (nested)
Meteorological IC/BC	NCEP FNL reanalysis (0.25~~1°~~ v x 0.25~~1~~°; 6-hourly)
Chemical IC/BC	CAMS global model output
Chemistry Mechanism	RACM-MADE-VBS (with aqueous reactions)
Anthropogenic Emissions	EDGAR v5; downscaled with ODIAC CO_2
Biogenic Emissions	MEGAN v2.1 (Guenther et al., 2006)
Biomass Burning Emissions	QFED v2.6 (Grell et al., 2011; Koster et al., 2015)
Dust Emissions	GOCART scheme (Zhao et al., 2010)
Sea Salt Emissions	Gong et al. (1997) parameterization (Gong et al., 1997)

Radiation Scheme	RRTMG (longwave); Goddard (shortwave)
Cumulus Parameterization	Grell-Freitas (D01 only)
Microphysics	Morrison (double-moment) (Ye et al., 2021)
PBL Scheme	Mellor-Yamada-Janjic TKE scheme
Simulation Period	14 – 27 March 2024




Figure 1. Spatial illustration of the WRF-Chem model domain configuration, including D01 (20 km) and D02 (4 km), and average base-model NO_x input emissions from EDGAR v5. D02 is centered over Thailand, and the urban signal associated with the Bangkok Metropolitan Region (BMR) is highlighted by the star.



1 Estimate Daytime NO_x Emissions from GEMS
Detect Bangkok urban plume from GEMS v3 NO_2 VCDs with *ddeq* v1
Estimate hourly NO_x with *ddeq* v1 and ERA5 winds through top-down methods
Inversion Methods
CSF, GP, IME
Obtain GEMS average daytime NO_x profile
2 Correct WRF-Chem's NO_x Profile Based on GEMS
Define prior: model daytime NO_x profile (EDGAR v5)
Define constraint: GEMS daytime NO_x profile (CSF)
Construct lifetime/decay informed emission mapping (H)
Optimize model daytime NO_x emission profile based on cost function, J(x)
Scale 24 hr WRF-Chem NO_x input emissions based on optimized daytime profile
Obtain WRF-Chem updated 24 hr input NO_x emissions
3 Test WRF-Chem NO_x Sensitivity
Run model with original NO_x emissions
Run model with updated NO_x emissions
WRF_{Base}
$WRF_{Updated}$
4 Evaluate WRF_{Base} and $WRF_{Updated}$
Datasets
Evaluate wind speed, direction, and PBLH
Evaluate surface and column chemistry
Thailad PCD
HSRL-2 Lidar
GEMS
GCAS
DC-8
Thailad PCD
Pandora

Figure 2. Schematic overview of the workflow used to derive satellite-constrained daytime NO_x emission profiles and assess their impact in WRF-Chem over the BMR. (1) Urban NO_2 plumes are detected using GEMS tropospheric NO_2 columns, and hourly NO_x emissions are estimated using top-down methods implemented in the *ddeq* v1 Python library. (2) a GEMS-derived daytime NO_x profile is used to constrain the prior emission profile through an optimization framework, yielding updated input 24 h NO_x emissions for WRF-Chem. (3) Model simulations are performed using both the original and updated emissions. (4) Model performance is evaluated against surface, column, airborne, and observations.

3 Satellite-derived ~~top-down~~ NO_x emission estimates

3.1 Correcting for GEMS bias prior to inversion

Recent independent validation studies of the operational GEMS v3 product over Bangkok and South Korea report low biases in NO_2 columns relative to ground-based sun-photometer and DOAS measurements (Bae et al., 2025; Jung et al., 2025). Bae et al. (2025) shows that GEMS v3 increasingly underestimates NO_2 relative to Pandora under high-NO_2 conditions (>1 × 10^{16} molecules cm^{-2}) as is the case for Bangkok pollution levels. In Jung et al. (2025), validation results over Bangkok indicate a pronounced low bias in GEMS tropospheric NO_2 columns relative to Pandora, with regression slopes of ~0.35 for v2.0 and ~0.28 for v3.0, indicating increasing underestimation at higher NO_2 levels. While moderate correlations ($r \approx 0.6–0.7$) suggest that GEMS captures temporal variability, column magnitudes are substantially underestimated, particularly under polluted conditions. The persistence of this behavior in the v3.0 product indicates that the low bias is not fully corrected by recent algorithm updates and is consistent with retrieval sensitivity limitations in highly polluted urban environments (Jung et al., 2025).

To address the low bias in GEMS prior to the NO_x satellite emission inversion, we first quantify the GEMS bias relative to Pandora measurements over our period of interest (14 – 27 March 2024). We compare total column GEMS NO_2 to Pandora Level 2 direct-sun total column retrieval, filtering for high quality measurements (quality flag = 10) and averaging to hourly means. GEMS columns are sampled at the nearest grid cell to each Pandora site and temporally collocated. Results are shown in Fig. S1. GEMS NO_2 columns are generally about a factor of two lower than Pandora measurements (mean bias ≈ −9.2 × 10^{15} molecules cm^{-2}). This factor-of-two difference persists throughout March 2024 (Fig. S1).

To assess the sensitivity of the top-down NO_x inversion framework to this bias, we apply a simple correction factor to GEMS prior to inversion. Specifically, GEMS NO_2 columns over the BMR are scaled by a factor of 1.67 derived from the Pandora comparison. The inversion is then performed with and without the bias-corrected columns.

3.2 GEMS inversion framework

Previous studies have used satellite data (OMI, TROPOMI, SCIAMACHY, GOME (-2), OMPS) to estimate top-down NO_x emissions over urban areas, but were limited to once-a-day, mid-afternoon measurements, leaving temporal variability in emissions largely unaddressed (Beirle et al., 2011; Goldberg et al., 2017, 2019). Ground-based and aircraft measurements of emissions can be challenging to constrain given boundary layer dynamics, such as changes in boundary layer height, stability, and vertical mixing, which can strongly influence observed concentrations of trace gases (Goldberg et al., 2017, 2019). Additionally, temporal allocation in bottom-up emission inventories remain a significant uncertainty, particularly at hourly and daily timescales because default temporal profiles (diurnal, weekly, seasonal) often fail to capture real activity patterns, meteorological influences, and sector-specific variability (Goldberg et al., 2017, 2019; Mues et al., 2014). GEMS observations, however, offer a unique opportunity to provide hourly emissions rates over Bangkok, complementing existing inventories that are usually provided at monthly scales. To estimate NO_x emissions over Bangkok in March 2024, we apply the methodology of Kuhlmann et al. (2024), implemented through the openly available data-driven emission quantification (*ddeq v1*) Python library (see step 1 Fig. 2). The standard *ddeq* library implements computationally inexpensive methods (e.g., Gaussian Plume Inversion (GP), Cross-Sectional Flux (CSF) method, Integrated Mass Enhancement (IME) method) to estimate emissions from Sentinel5P -TROPOMI images (Graziosi and Manca, 2025; Meier et al., 2024) . One study found limited temporal sampling from polar-orbiting satellites (i.e., TROPOMI) can lead to systematic underestimation of NO_x emissions, particularly because wintertime conditions with higher emissions are often poorly observed due to cloud cover (Meier et al., 2024) . Geostationary observations, including GEMS, address this limitation by providing hourly measurements that increase the likelihood of usable data on partially cloudy days and enable direct resolution of daytime emission variability, which is not accessible from once-daily LEO observations. For our application, we estimate emissions for daylight hours from GEMS tropospheric column NO_2 data over Bangkok during the ASIA-AQ campaign. A brief description of inversion methods and specifics is provided here. A complete description of the inversion methods and algorithm can be found in Kuhlmann et al. (20241~~4~~).

First, within the *ddeq* framework, the source location (Bangkok; 13.7563°N, 100.5018°E), GEMS v3 NO_2 column data, and ERA5 wind fields are read. We chose to incorporate ERA5 wind fields as opposed to those provided by our high resolution WRF-Chem simulation in this analysis given they presented significantly lower bias when compared to surface observations (Fig. S2; Sect. 6.1). The ERA5 wind fields were downloaded and prepared by the *ddeq* library. Cloudy pixels (CF > 0.3) were removed. Next, a plume detection algorithm is implemented to identify the Bangkok urban plume within the GEMS image.

The Bangkok plume subregion is estimated based on the source location and ERA5 wind field. Generally, a wind vector is taken at the source location, and the plume is assumed to be located downwind, and a rectangular polygon can be drawn with the along- and across- wind direction (Kuhlmann et al., 2024) . An example of these plume detections and associated rectangular polygons is shown in Fig. S3~~1~~. To aid the algorithm's plume detection process, we identify two additional sources north of Bangkok in the Saraburi province to avoid overlapping plumes from other NO_x sources. These correspond to the Khao Wong (14.692856°N, 100.817204°E), and Thap Kwang (14.645313°N, 101.077650°E) regions, which we identified through NO_2 patterns visible in the GEMS data in Fig. S3~~1~~. These areas are not representative of Bangkok urban emissions as they are a source of industrial activity related to limestone quarries and mining operations outside the inversion domain (Makkwao and Prueksasit, 2021). Next, a center curve is fitted to the ~~data~~data, and natural coordinates are calculated for the detected plume associated to Bangkok to prep for the inversion algorithm. The coordinates are computed as the distance along and perpendicular to the wind vector and for curved plumes this distance is computed as the arc length. Lastly, the data is prepared for the satellite-derived emissions estimation by calculating and removing the background field and converting the NO_2 columns to kg m^{-2}. We estimate satellite-derived NO_x emissions using three inversion techniques included in *ddeq*: CSF, GP, and IME methods. Below is a summary of the CSF method defined in Kuhlmann et al. (2024). GP and IME method descriptions are available in the Supplement (e.g., S1) and Kuhlmann et al. (2024). In the following subsections and in Section 4, we refer to the WRF-Chem input inventory based on EDGAR as the *prior emissions*, the CSF-based estimates from GEMS as the *satellite-derived emissions*, the CSF-based estimates from the model-simulated columns as the model-derived emissions, and the final emissions after applying the optimization framework, as the *posterior emissions*.

3.3~~3.1~~ Cross-Sectional Flux (CSF) framework

The CSF approach applies mass conservation to quantify the NO_x flux transported downwind of Bangkok using (i) the wind speed perpendicular to the plume and (ii) the NO_2 enhancement integrated along the plume (line density). The NO_x flux, F (kg s^{-1}), is defined as:

$$F = u \cdot q \quad (1)$$

where u (m s^{-1}) is the effective transport wind speed and q (kg m^{-1}) is the NO_2 line density. The emission rate, Q (kg s^{-1}), is then inferred by correcting for chemical decay along the plume using:

$$Q = \frac{F(x)}{D(x,\tau)} \quad (2)$$

Here, $D(x,\tau)$ represents the along-plume decay for lifetime τ. Here, τ represents an effective plume lifetime that accounts for both chemical decay and plume dispersion (Kuhlmann et al., 2024).

3.31.1. Effective wind speed estimation

The *ddeq* algorithm computes an effective wind speed, i.e., the plume-transport wind, that supports an unbiased emission estimate. Ideally, this is a NO_2-enhancement-weighted average of the along-plume wind:

$$u(x,y) = \frac{\int_0^{z_T} \rho_e(x,y,z)u(x,y,z)dz}{\int_0^{z_T} \rho_e(x,y,z)dz} \quad (3)$$

Here, $\rho_e(x,y,z)$ is the NO_2 enhancement, $u(x,y,z)$ is the along-plume wind speed and z_T is the plume-top height. Because ρ_e is not directly observed, *ddeq* follows the approximation of Fioletov et al. (2015), in which the effective wind speed is estimated using the mean of the lowest three ERA5 layers, representing the typical transport level for near-surface urban plumes.

3.31.2. NO_2 line density calculation

The line density $q(x)$, is the integral of the NO_2 enhancement across the plume cross-section:

$$q(x) = \int_{y_1}^{y_2} \big(V(x,y) - V_{bg}(x,y)\big)\, dy \quad (4)$$

where $\big(V(x,y) - V_{bg}(x,y)\big)$ represents the NO_2 enhancement above the background in kg m^{-2}. The integration bounds y_1 and y_1 are defined by the plume subregion (polygon) identified using the *ddeq* framework, which delineates the area of enhanced NO_2 associated with the source based on plume detection and wind direction (Kuhlmann et al., 2024). As such, the effective integration length across the plume is determined by the spatial extend of this polygon and the fitted Gaussian representation. To further compute this and account for missing satellite pixels, *ddeq* fits a Gaussian function to all GEMS pixels within a plume polygon:

$$g(y) = \frac{q}{2\sqrt{\pi}\sigma} exp\ \exp\left(-\frac{(y-\mu)^2}{2\sigma^2}\right) + my + b \quad (5)$$

Here, μ and $\sigma(x)$ are the plume center and width in meters, and $my + b$ approximates a linear background. The plume width is further defined as:

$$\sigma(x) = \sqrt{\frac{2Kx^{\kappa}}{u}} \quad (6)$$

Where K is the eddy diffusivity coefficient ($m^2\ s^{-1}$), κ accounts for nonlinear plume spreading under varying meteorology, and u is the effective transport wind speed defined above. The plume width, $\sigma(x)$ is derived from the Gaussian fit, and the parameters, K and κ are then determined by fitting, $\sigma(x)$ within the CSF framework. After fitting the Gaussian function, line densities are converted to fluxes using wind speeds at each corresponding downwind cross-section. Fluxes can be estimated for several cross-sections or polygons located downwind of the urban source (see Fig. S31,S42) (Kuhlmann et al., 2024). Using multiple points along the plume ensures different portions of the plume are sampled robustly.

3.31.3. Emission rate and lifetime estimation

Because NO_2 decays as the plume travels, the flux decreases with distance. To estimate the true emission rate, Q , at the source, *ddeq* fits a lifetime τ by matching modeled and observed flux decay:

$$F_a(x, \tau, \mu_a, \sigma_a) = Q \int_{-\infty}^{\infty} D(x', \tau) g(x - x', \mu_a, \sigma_a) dx' \quad (7)$$

Where μ_a and σ_a describe the city-scale plume location and extent, $D(x', \tau)$ describes the exponential decay along the plume, and g is the Gaussian parameter described above. Fitting Eq. (7) to the derived fluxes yields both the emission rate Q and plume lifetime τ (Kuhlmann et al., 2024).

3.31.4. Conversion from NO_2 to NO_x emissions

GEMS provides NO_2 column densities; therefore, a NO_2-to-NO_x conversion factor is needed to obtain NO_x emissions. Following the *ddeq* implementation, we use:

$$Q_{NO_x} = f_q \cdot Q_{NO_2} \quad (8)$$

to obtain NO_x estimates, where f_q= 1.32, as implemented in the *ddeq* algorithm~~.~~ a localized factor was not derived from WRF-Chem due to known NO_2 biases in the prior simulation that would propagate into the conversion.

3.3.5 Structural uncertainties in the CSF framework

While the CSF framework provides a computationally efficient approach for deriving satellite-derived emissions, we acknowledge the method contains some structural uncertainties. First, the integration bounds used to compute the line densities (Eq. 4) are defined by the plume polygon identified from the satellite observations and wind field, such that the inferred top-down emissions depend on how the plume extent is delineated. As a result, inferred satellite-derived emissions can be sensitive to the selected plume extent, particularly under conditions of weak enhancements or overlapping sources. Second, the satellite-derived emission estimate scales directly with the effective wind speed (Eq. 1), such that errors in wind magnitude or direction propagate into the derived fluxes and satellite-derived emissions (see Fig. S5). Although ERA5 winds averaged over the lowest model layers are used to approximate plume transport, they may not fully represent local meteorological variability. Third, the inferred satellite-derived emission rate depends on the assumptions within the fitting framework, including the Gaussian representation of plume structure (Eq. 5), the parametrization of plume spreading (Eq. 6), and the effective lifetime τ (Eq. 7), which represent a simplified treatment of chemical decay and dispersion. Together, these factors represent inherent uncertainties in the CSF approach that should be considered when interpreting the satellite-derived emissions.

3.42 Results for satelliteGEMS-derived NO_x emission estimates

GEMS NO_x emissions for Bangkok were estimated for daytime hours (00:45:00 – 06:45:00 UTC) during the ASIA-AQ deployment, 14 – 27 March 2024. The bias-corrected satellite inversion results are denoted with the subscript "BC". Days with significant cloud contamination were removed from the analyses on a case-by-case basis, leaving satellite-derived emissions estimates for 14,15,17,18,22,24,25 March 2024. Satellite-derived eEmissions were computed for both weekends and weekdays, however, due to fewer weekends in the study period, weekday estimates are estimated to be more robust. We aggregate satellite-derived emissions hourly across weekdays to produce a summary daytime profile as illustrated in Fig. 3a and Fig. 3b. Satellite-derived eEmissions generally range between 0.5 – 4 kg s^{-1} dependent on the inversion method. Bias-corrected results (Fig. 3b) illustrate a larger NO_x range, between 0.4 – 5 kg s^{-1}. Differences between CSF, GP, and IME satellite-derived emission estimates are expected, as each method relies on distinct assumptions regarding transport, plume geometry, and chemical loss. Similar spreads between methods have been reported in previous satellite-based emission studies (Hakkarainen et al., 2023, 2024; Santaren et al., 2025), particularly in urban environments, and can be interpreted as a measure of structural uncertainty rather than inconsistency between methodologies. Nevertheless, in this application, the CSF method illustrates a distinct daytime pattern with satellite-derived emissions peaking between 08:00:00 – 09:00:00 LT (bias-corrected)at ~~01:00:00 UTC (or 08:00:00 LT)~~ coinciding with morning rush hour traffic, decreasing by approximately 65% by 14:00:00 LT. The GP and IME results illustrate a relatively stable daytime pattern and overall lower (< 50%) morning satellite-derived emission compared to the CSF method.

These results reflect both methodological differences and uncertainties in the underlying NO_2 observations. While the satellite-derived emissions might provide strong constraints on daytime variability, uncertainties in the GEMS NO_2 retrieval, particularly under high NO_2 conditions may influence the magnitude and timing of the inferred emissions as depicted in Fig. 3a,3b. The bias-corrected emissions are approximately 70% higher on average, indicating a substantial sensitivity of the inversion to the assumed retrieval bias.

3.53 Selection of the inversion method

To identify the optimal inversion method to be used for model correction, we test which method can successfully recover WRF-Chem's prior daytime NO_x emission profile. To do so, we apply the *ddeq* inversion algorithms to WRF_{Base}, to produce "top-down" estimates from the model, which we will refer to as model-derived emissions. To replicate the satellite output, we re-gridded the model output from its native resolution to the satellite swath grid using the Universal Regridder for Geospatial Data (xESMF) Python package (Jiawei Zhuang et al., 2025). A nearest-neighbor interpolation scheme was used to preserve spatial gradients and align the model resolution to that of the satellite observations. This spatial re-gridding was performed on an hourly basis, corresponding to the nearest observation time (00:45:00 – 06:45:00 UTC). Following re-gridding, model NO_2

mixing ratios were vertically interpolated to the satellite pressure levels using a one-dimensional linear interpolation to ensure consistency between the model and observed data for averaging kernel application. After interpolating model output to GEMS's spatial and pressure grid, we computed the model's tropospheric column NO_2 (molecules cm^{-2}). This is done by converting the model's standard output of NO_2 volume mixing ratio (ppmV) to a number density with the ideal gas law:


$$NO_{2_{Number\ Density}} = \frac{P \cdot \chi_{NO_2} \cdot N_A}{R \cdot T} \quad (9)$$

Where χ_{NO_2} is the NO_2 mixing ratio in ppmV, P is pressure in Pa, T is temperature in K, R is the gas constant (8.314 J/mol/K), and N_A is Avogadro's number (6.022 x 10^{23} mol^{-1}). The number density is then multiplied by the thickness of each vertical model layer Δz to obtain the NO_2 amount per unit area. Summing across all layers yields the total column NO_2:

$$NO_{2_{WRF-GEMS}} = \frac{1}{10^4} \times \sum_k (A_k \cdot NO_{2_{Number\ Density,k}} \cdot \Delta z_k) \quad (10)$$

Where A_k is the averaging kernel value at layer k, Δz_k is the thickness of the model layer k, in meters, and $\frac{1}{10^4}$ converts the units to molecules cm^{-2}. As WRF-Chem has no stratosphere, we regard this as the tropospheric column NO_2. The satellite averaging kernel weights each model layer based on how sensitive GEMS is to that part of the atmosphere. This allows for a direct comparison with GEMS retrievals.

We compute model-derived NO_x emissions from re-gridded WRF_{Base} (which we denote as WRF-GEMS) with *ddeq* and the resulting average daytime profile for Bangkok is illustrated in Fig. ~~3b,~~ 3c. The ~~model's~~ shape and magnitude of model-derived NO_x emissions differ substantially from the satellite-derived~~GEMS~~ emission data. Model-derived emissions are initially > 50% larger than GEMS-derived emissions. Additionally, there are minimal changes in the daytime pattern present in any of the inversion methods. In fact, WRF-GEMS D02 sees an increase in emission between 08:00:00-10:00:00 LT, a pattern that disagrees with the observed satellite-derived emissions. The results here further illustrate the need for an hourly correction to the model's emission profile, rather than applying a single scaling factor across all hours. The WRF_{Base} prior emission profile is shown by the black dashed line. For this comparison, we smoothed the prior emission profile using a 3 h backward-looking average to account for the accumulation and chemical evolution represented implicitly in the inversion results. Overlaying the prior emissions shows that the model-derived CSF method most accurately recovers them, likely due to its use of downwind fluxes and multiple cross-sections across the plume, which provide additional constraints on plume evolution and reduce sensitivity to local errors in wind fields, plume definition, and missing data compared to the IME and GP methods. As a result, we select the CSF inversion method moving forward to correct and scale WRF-Chem's daytime NO_x emission profile.


NOx Emissions
BMR Average (Weekday)
(a)
GEMS
CSF
GP
IME
NOx Emissions [kg s−1]
(b)
GEMSBC
CSF
GP
IME
NOx Emissions [kg s−1]
(c)
D02
CSF
GP
IME
WRF Prior
NOx Emissions [kg s−1]
Hour [Local Time]




Figure 3. Comparison of daytime NO_x emission estimates derived from (a) GEMS, (b) GEMS with bias correction applied ($GEMS_{BC}$) ~~WRF-GEMS D02~~, and (c) WRF-GEMS D02~~1~~ using the Cross-Sectional Flux (blue), Gaussian Plume inversion (red), and Integrated Mass Enhancement (green) methods. Emissions (kg s^{-1}) represent averages across daytime hours and weekdays during the ASIA-AQ deployment period (14 – 27 March 2024). Model-derived ~~inversion~~ emissions in ~~(b) and~~ (c) were obtained by re-gridding WRF_{Base} output to the GEMS spatial resolution and applying the same inversion methods. The

WRF_{Base} prior emission profile is shown as a black dashed line and represents a backward-looking three-hour average to account for emission accumulation embedded in the inversion estimates.

4 Model emission adjustment framework

To better represent NO_x emissions over Bangkok, we developed an optimization framework to re- scale and shape WRF-Chem's prior~~input~~ daytime emissions across the BMR (see step 2 Fig. 2). The prior emissions come from monthly EDGAR v5 values. The optimization uses a cost function $J(x)$ to adjust daytime prior~~model~~ emissions based on GEMS NO_x retrievals, incorporating the lifetime and uncertainties from both the model and observations.

4.1 Optimization of daytime anthropogenic NO_x emissions

4.1.1. Observational constraint

The GEMS average daytime NO_x profile (blue lines in Fig. 3a,3b) from the CSF method provides the observational constraint. Two versions of the optimization are performed: one using the constraint derived from the bias-corrected retrievals, and one without it. To improve consistency between the observed and modeled columns, we applied a correction factor to the constraint to account for the portion of the NO_2 column that GEMS does not capture because of reduced sensitivity near the surface. Although the air mass factor (AMF) calculation includes vertical weighting, the retrieved GEMS column can still underestimate tropospheric NO_2 in conditions where a substantial fraction resides in the lowest few hundred meters and is weakly sensed by the instrument. To address this sensitivity, satellite~~GEMS~~-derived NO_x emission estimates were scaled using the ratio of model-derived~~WRF-Chem~~ emissions obtained from the full column (i.e., Eq. (10) excluding the A_k) and from the same column after application of the GEMS averaging kernel:

$$y = e_{GEMS} \times \left(\frac{{e_{WRF}}^{TC}}{{e_{WRF}}^{AK}}\right) \quad (11)$$

Where e_{GEMS} , ${e_{WRF}}^{TC}$, and ${e_{WRF}}^{AK}$ denote the satellite-derived ~~CSF~~-derived daytime NO_x emission profiles derived from GEMS, model-derived emissions from WRF-Chem (D02) total columns, and model-derived emissions from the same columns after applying the GEMS averaging kernel, respectively. Hourly correction factors ranged between 1.05 – 1.33, where larger corrections (e.g., 1.33) were~~where~~ present in the morning hours, 01:00:00 – 02:00:00 UTC. This vertically adjusted satellite-derived emission vector was used as the observational constraint, y, in the optimization.

4.1.2. Prior assumptions, lifetimes, and temporal weighting

The prior for the optimization, x_b, refers to the bottom-up daytime NO_x emission profile shown in the black dashed line in Fig. 3c~~b~~. For the optimization, it is necessary to map the prior~~model~~ emissions to the satellite-derived emissions through the

observational operator H. Satellite emissions of NO_x represent not only emissions from the current hour, but also contributions from preceding hours. To address this, we introduce an operator H that links the prior~~model~~ daytime hourly emissions to the satellite-derived~~observed GEMS~~ NO_x amounts. Each row of H defines how emissions from a set of prior model hours contribute to an observation at time, t_i.

For each GEMS observation hour t_i, we assume that the column enhancement is influenced by emissions from a backward-looking window of three model hours:

$$t_i - 2,\ t_i - 1, t_i$$

The window choice was determined by comparing inversion results to model priors using averaging windows from 1 – 6 hours. The three-hour window provided the best agreement with the model-derived CSF emissions~~CSF inversion~~ as shown in Fig ~~3b,~~ 3c. Within this window, the contribution from each prior hour depends on the atmospheric lifetime of NO_2 for that hour. We use hourly lifetimes, τ, derived from the GEMS-based ~~which were fitted within the~~ CSF inversion in Eq. (7), where τ is fitted from observed along-plume decay using ERA5 winds, and is therefore independent of WRF-Chem ~~.~~meteorology. This ensures the satellite-derived emission estimates are completely independent of the WRF-Chem simulations. For reference, WRF-Chem-derived decay times are generally longer than those inferred from ~~GEMS,~~ GEMS, reflecting differences in modeled vs. observed loss and dilution processes.~~with partial overlap at earlier hours but increasing divergence later in the day, indicating differences in the effective loss and dilution processes represented by the model~~. Hours with no satellite information, are assigned a default lifetime of 3 hours. We then apply a simple exponential decay law:

$$w(\Delta t) = exp\ (-\tfrac{\Delta t}{\tau})\ (12)$$

where:

$$\Delta t\ \epsilon\ \{0,1,2\}$$

is the time lag between when NO_x was emitted and when its NO_2 enhancement is observed. These raw weights are normalized within each row of H to ensure mass conservation:

$$H_{i,j} = \frac{w(\Delta t_j)}{\Sigma_{j'} w(\Delta t_{j'})}\ (13)$$

Thus, $H_{i,j}$, represents the fraction of the observed column at hour i, that is attributable to emissions from model hour j. This framework of H ensures that shorter lifetimes contribute to a stronger emphasis on the most recent emissions, appropriately reflecting rapid chemical loss, and longer lifetimes spread the influence of emissions over multiple hours, consistent with slower decay. By embedding this chemical persistence into the operator, H, we obtain a physically representative mapping between model emissions and the satellite observations that are used to constrain them.

4.1.3. Solution for posterior~~optimized~~ daytime emissions

A model corrected daytime NO_x emission profile (posterior), $\hat{x}$ was obtained by minimizing the following cost function:

$$J(x) = \frac{1}{2}(y - Hx)^T R^{-1}(y - Hx) + \frac{1}{2}(x - x_b)^T B^{-1}(x - x_b) \quad (14)$$

Where y is the satellite-derived~~GEMS~~, column-corrected NO_x emission vector, x is the unknown corrected model daytime emission profile, x_b is the model prior profile, $R = {\sigma_I}^2 I, B = {\sigma_B}^2 I$ are the observation and background error covariance matrices, and H is the observational operator matrix described above. In this study, we use R = 1 and B = 10. These values were based on sensitivity tests in which we vary R and B over several orders of magnitude. The chosen values represent a compromise that allowed the posterior~~optimized~~ emission profile to closely follow the satellite constrains while preventing excessive deviations from the prior emission profile. To account for temporal correlation in prior emission errors, the elements of B are defined as:

$$B_{i,j} = {\sigma^2}_B exp\left(-\frac{|t_i - t_j|}{L}\right) \quad (15)$$

Where $|t_i - t_j|$ is the temporal separation between emission hours i and j, and L is the temporal correlation length, set to $L = 2\ hr$ in this study. This setup ensures that prior errors vary smoothly in time rather than independently hour-to-hour. Finally, the cost function was minimized using a Sequential Least Squares Quadratic Programming (SLSQP) algorithm (minimize(method='SLSQP') — SciPy v1.16.2 Manual, 2025; Nocedal and Wright, 2006). The result is a corrected model hourly emission vector, $\hat{x}$, that balances the satellite-derived emissions ~~observations~~ with the model's prior estimate while accounting for chemistry and transport behavior (Fig. 4).

Figure 4a and 4c illustrate~~s~~ the raw hourly daytime NO_x for both the prior model (dashed purple) and the optimized version (solid purple), based on constraints applied without and with the GEMS bias correction, respectively. In both bases, t~~T~~he posterior~~optimized~~ profile departs notably from the prior in both magnitude and ~~shape,~~ shape. In the case without the bias-corrected constraint (Fig. 4a), ~~with a pronounced~~ a morning peak is visible at 07:00:00 LT, followed by a steady decline until 13:00:00 LT. This pattern is consistent with temporal patterns in weekday traffic intensity in Bangkok found in Ly et al. (2015), which found the morning traffic to peak at 07:30:00 LT ending at 09:30:00 LT, and the evening peak from 17:00:00 to 20:00:00 LT (outside of GEMS observing window). Activity begins to increase at 13:00:00 LT which corresponds to the NO_x increase we begin to see in the posterior~~optimized~~ result after 06:00:00 UTC. The case with the bias-corrected constraint (Fig. 4c) places peak morning emission closer to the model prior at 07:00:00 – 08:00:00 LT, with NO_x declining slightly later, at 09:00:00 LT. NO_x emission begins to increase after 13:00:00 LT similar to what is depicted in Fig. 4a. When the prior and posterior~~optimized~~ hourly emissions are passed through H, we obtain a temporally smoothed emission profile consistent with what GEMS would observe (Fig. 4b,4d). Figure~~s~~ 4b and 4d show~~s~~ the posterior~~optimized~~ profile closely reproduces the GEMS-derived daytime pattern, indicating the optimization effectively aligns model emissions with satellite-derived emissions ~~observations~~ while preserving components of the prior model behavior.

4.2 Adjustment of WRF-Chem diurnal emission profiles

Using satelliteGEMS-derived NO_x emissions and the posterioroptimization results above, we correct the WRF-Chem 24 h input emissions over the BMR. Since GEMS provides estimates over 01:00:00-07:00:00 UTC, we only have posterioroptimized model emissions over this window. To apply these updates to the full gridded 24 h WRF-Chem anthropogenic emission input (00:00:00-23:00:00 UTC), while preserving the model's spatial distribution, we compute hourly scaling factors which we use to adjust the gridded priorinput NO_x emissions. Because GEMS provides constraints only during the daytime overpass window, the derived scaling below primarily reflects daytime emission adjustments. We therefore apply the mean scaling factor uniformly during the nighttime hours to preserve the prior temporal structure and avoid introducing artificial discontinuities between the constrained and unconstrained hours, particularly given the large daytime emission reductions. This approach is consistent with previous satellite-constrained emission studies (e.g., Goldberg et al., 2019) which constrain emission magnitude based on a single daytime value derived from OMI. Where available, locally derived temporal profiles should be used to better represent regional emission patterns; in this study, our prior profile reflects the best available regional representation for Asia at the time of study.

To compute the factors, hourly WRF-Chem NO and NO_2 prior emission fields were summed across the vertical levels and masked to the Bangkok region bounded by N-S:13.5°-15.0°, W-E:100.2°-100.9°, using the WRF-Chem grid. In WRF-Chem, NO_x emissions are stored as NO_2-equivalent mass ('NO_x-as-NO_2'), so all conversions use the molecular weight of NO_2. The total emissions were converted to mol hr^{-1} using a molecular weight of 46.01 g mol^{-1} and an area of 16 km^2 per grid cell (following the resolution of D02). Each posterioroptimized emission, $\hat{x}_h$ was converted from kg s^{-1} to mol hr^{-1} and divided by the corresponding original WRF-Chem total to compute a scale factor, f for each hour h:


$$f = \frac{\hat{x}_h \times \frac{1000}{46.01} \times 3600}{E_H{}^{WRF}} \quad (16)$$

Where $E_H{}^{WRF}$ is the original domain-integrated prior NO_x emissions for hour h. Since the optimization covered hours with GEMS measurements, the scaling factors for the remaining model hours were computed using the ratio of average daytime posterior emissions where averages are taken over the constrained hours. Finally, we apply the scaling factors spatially over the BMR. The scaling factor was applied to the hourly WRF-Chem anthropogenic emissions priorinput files and only to NO and NO_2 grid cells within the BMR mask. This preserves the original spatial distribution but adjusts the total prior emissions to match the posterioroptimized values. The updated emission files were then used to initialize an updated model simulation, ran for the ASIA-AQ deployment period, which we will refer to as $WRF_{Updated}$ (see step 3 Fig. 2).


Figure 4ce compares the average diurnal prior profile over BMR with the updated profile derived from the optimization and scaling procedure. The optimized diurnal profile based on the bias-corrected GEMS constraint is illustrated by the light pink line. The updated profiles shows a notable increase in NO_x between 07:00:00-10:00:00 UTC (14:00:00 to 17:00:00 LT), which may reflect the beginning of a rush-hour signal over the city. Nighttime emissions retain the overall shape of the prior profile

but are scaled downward based on the daytime average. Figure S6~~S4~~ maps the spatial distribution of these updates across BMR, showing the largest differences in the late afternoon and early evening, with localized reductions of up to 400 mol km^{-2} hr^{-1} in the updated input emissions.


(a)




(c)




(b)



(d)



(e)






Figure 4. Results of the emissions optimization, shown as averages over the BMR. (a) Posterior optimized (solid) and prior (dashed) daytime NO_x emissions in model space, where the prior represents the three-hour backward average of EDGAR v5 emissions used as input to the optimization. (b) Same emissions as in (a) but transformed into observational space using the observational operator. The GEMS profile is shown in blue. (c) and (d) are representative of optimization results using the GEMS bias-corrected CSF results as the constraint. (e) Full diurnal cycle of updated (solid purple), updated+BC (solid pink), and prior (dashed) NO_x emissions. Note that, unlike (a), the prior in (e) does not include the 23 UTC hour from the previous day in the backward averaging. ~~Results of the emissions optimization, shown as averages over the BMR. (a) Comparison of optimized (solid) and prior (dashed) daytime NO_x emissions. (b) Same as (a) but viewed through the GEMS observational operator. The GEMS profile is shown in blue. (c) Comparison of full optimized (solid) and prior (dashed) diurnal NO_x emission profiles.~~


4.3 Comparison with bottom-up emission inventories

To place the updated NO_x emissions in context of existing emission inventories, we compared total monthly emissions over the BMR against several widely used bottom-up inventories, our top-down emissions (including bias-corrected results – This Study$_{BC}$), and emissions from a chemical reanalysis product (Fig. 5). These include HTAP v3.2 (base 2019), a local Thailand Inventory (THAI-KMUTT) (base 2019), MIX v2.3 (base 2017), EDGAR v5 (base 2015), CEDS (base 2024), our study (2024),

and the Tropospheric Chemistry Reanalysis (TCR-3) top-down emissions TCR-3 (2024) (Miyazaki et al., 2020). Across inventories, monthly totals for March can vary nearly by an order of magnitude, highlighting large uncertainty in regional NO_x sources and emissions processing methodology. For example, although both HTAP v3.2 and MIX v2.3 rely on the REAS framework (Kurokawa and Ohara, 2020) for Asian anthropogenic emissions, they are derived from different REAS versions, temporal coverage, and processing assumptions**,** which can lead to substantial differences in absolute emission magnitudes over Bangkok. HTAP v3.2 is based on REAS v3.2.1, which provides monthly emissions for 2000–2015 at a spatial resolution of 0.25°. In the HTAP product, these emissions are re-gridded to 0.1° by assuming a uniform spatial distribution within each grid cell (Guizzardi et al., 2025). In contrast, the MIX inventory is based on REAS v2, which covers 2000–2008 and reflects earlier emission estimates and activity data (Li et al., 2024). REAS v2 was subsequently extended to 2010 following the scaling approach described by Kurokawa & Ohara (2020).

Overall, HTAP v3.2 depicts the highest NO_x totals at 31kTons $month^{-1}$. THAI-KMUTT follows with emissions at ~20 kTons $month^{-1}$, reflecting its incorporation of detailed regional activity from local emissions data. The coarser-resolution MIX v2.3, and EDGAR v5, and CEDS inventories reflect lower totals, at ~13,11,10 kTons $month^{-1}$ respectively, consistent with the use of different emission estimation methodology and older base years for MIX and EDGAR which correspond to lower macro-economic (e.g., GDP) indicators in Thailand compared to 2019. Lastly, GEMS top-down NO_x estimates (including bias-corrected results) and TCR-3 estimates based on the chemical reanalysis using TROPOMI NO_2 indicate substantially lower emissions (2-~~43~~ kTons $month^{-1}$). These reduced magnitudes may result from both recent emission declines and structural differences in how emissions are computed (i.e., top-down perspective).

It is important to note uncertainties are present and differ amongst emission methods. For example, bottom-up inventories depend on activity data, emission factors, spatial approximations, and assumptions that may not capture rapid socio-economic changes or region-specific behavior. Top-down estimates incorporate uncertainties from their respective satellite retrievals, reanalysis data (e.g., ERA5), averaging kernels, chemical lifetime assumptions, and forecast-model transport and chemistry (e.g., TCR-3). Together, these differences highlight the value of combining observational constraints with updated bottom-up information to define NO_x emission estimates.






Figure 5. Comparison of March NO_x emissions over the BMR derived from bottom-up, top-down, and assimilated approaches. Bottom-up inventories (HTAP v3.2 (Guizzardi et al., 2025), THAI-KMUTT, MIX v2.3 (Li et al., 2024), EDGAR v5 (Crippa et al., 2020), and CEDS (Hoesly et al., 2018)) exhibit substantial variability. In contrast, top-down estimates from GEMS and TROPOMI (TCR-3) for 2024 indicate considerably lower emissions (2.6–4.3~~2.7~~ kt month^{-1}), highlighting uncertainties in bottom-up inventory methodologies, satellite sensitivity, and potential changes in anthropogenic NO_x activity.


5 Evaluation against satellite observations


To evaluate the performance of WRF-Chem (Base + Updated) NO_2 simulations during the ASIA-AQ period (14 – 27 March 2024) in Bangkok, we first compared model output with satellite tropospheric column NO_2 retrievals from GEMS v3 data. For GEMS observations, pixels were filtered based on cloud fraction (CF < 0.3). The GEMS averaging kernel and air mass factor data were used to isolate tropospheric contributions and account for satellite instrument sensitivity. To enable a direct comparison with our model output, we re-gridded model output from its native resolution to the satellite swath grid as explained in Sect. 3.3. As before, refer to the re-gridded model simulations with the GEMS averaging kernel applied as WRF-GEMS. To evaluate regional NO_2 variability, we aggregated satellite and model-derived tropospheric column NO_2 values over the BMR, by defining a N-S (13.5-14°N), W-E (100.3-100.9°E) bounding box for the area. For pixels within this boxed region, we calculate the mean NO_2 concentration, minimum NO_2, maximum NO_2, standard deviation, and the 10^{th} and 90^{th} percentiles to characterize regional variability. This analysis was repeated on an hourly basis for each co-located WRF-Chem and GEMS dataset, providing a regionally aggregated view of column NO_2 for intercomparison.


The results shown here are based on satellite comparisons using the non-bias corrected model results and are intended to illustrate that the optimization brings the model closer to GEMS observations. The maps shown in Fig. 6 illustrate a spatial comparison of (a) GEMS, (b) updated model results, WRF-GEMS$_{Updated}$ D02, and (c) base model, WRF-GEMS$_{Base}$ D02 tropospheric columns NO_2 over the BMR for several snapshots on 18 March 2024. Results for D01 can be found in Fig. S6~~5~~, S7~~6~~. GEMS generally places BMR city enhancements between 5 x 10^{15} – 2 x 10^{16} molecules cm^{-2} in a N-S direction. Both WRF$_{Updated}$ D01, D02 generally reflect the spatial distribution and magnitude of tropospheric column NO_2 observed during daylight hours. WRF$_{Updated}$ D02 (Fig. 6b) tends to place the city plume more northward at 04:00:00 and 07:00:00 UTC compared to what is observed in GEMS data (Fig 6a). This spatial discrepancy is likely related to overpredicted model winds (further discussed in Sect. 6.1) which displaces the plume in a northward direction. Additionally, while the inversion framework adjusts emission magnitudes temporally, it does not correct spatial errors in plume transport or source distribution (See Sect. 7.2 for further discussion). Nevertheless, the daytime pattern seen in GEMS is accurately reflected in WRF$_{Updated}$ with column NO_2 values at their peak in the morning hours (08:00:00 LT), decreasing substantially throughout the afternoon (14:00:00 LT) due to photochemical interactions. This is a substantial improvement from WRF$_{Base}$, which sees a large overestimation of NO_2 in the morning hours followed by persistent and growing NO_2 columns throughout the afternoon as seen in Fig. 6c.

The time series in Fig. 6d illustrates a full comparison of daytime tropospheric column NO_2 values across model types in D02 (see Fig. S7~~8~~ for D01) for the duration of ASIA-AQ. WRF-GEMS$_{Updated}$ indicates a clear reduction in error and bias from WRF$_{Base}$ as illustrated by the reduced NO_2 columns and model spread. WRF$_{Updated}$ also indicates corrected daytime patterns

633 compared to WRF_{Base}, indicating better consistency with GEMS observations for the deployment period with few exceptions
634 such as 22 March where GEMS has peaks in the afternoon (further discussed in Sect. 6.1).


(a)
(b)
(c)
Mar 18th, 2024
GEMS
WRF-GEMS$_{Updated}$
WRF-GEMS$_{Base}$
D02
08:45 LT
10:45 LT
12:45 LT
09:00 LT
11:00 LT
13:00 LT
3 × 10^{15}
4 × 10^{15}
5 × 10^{15}
6 × 10^{15}
7 × 10^{15}
8 × 10^{15}
1 × 10^{16}
1.5 × 10^{16}
2 × 10^{16}
Tropospheric Column NO$_2$ [molecules/cm^2]


(d)
GEMS vs. WRF-Chem
D02
1e16
GEMS
WRF-GEMS$_{Base}$
WRF-GEMS$_{Updated}$
Tropospheric Column NO$_2$ [molec/cm^2]
Mar 15
Mar 16
Mar 17
Mar 18
Mar 19
Mar 21
Mar 22
Mar 23
Mar 24
Mar 25
Mar 26


Mar 18th,2024
(a) GEMS
(b) WRF-GEMS$_{Updated}$
(c) WRF-GEMS$_{Base}$
D02
01:45 UTC
03:45 UTC
06:45 UTC
02:00:00 UTC
04:00:00 UTC
06:00:00 UTC
3 x 10^{15}
4 x 10^{15}
5 x 10^{15}
6 x 10^{15}
7 x 10^{15}
8 x 10^{15}
1 x 10^{16}
1.5 x 10^{16}
2 x 10^{16}
Tropospheric Column NO$_2$
[molecules/cm^2]



(d)
GEMS vs. WRF-Chem
D02
1e16
Tropospheric Column NO$_2$
[molec/cm^2]
GEMS
WRF-GEMS$_{Base}$
WRF-GEMS$_{Updated}$
Mar 15
Mar 16
Mar 17
Mar 18
Mar 19
Mar 21
Mar 22
Mar 23
Mar 24
Mar 25
Mar 26

Figure 6. Spatial comparison over the BMR on 18 March 2024 of tropospheric NO_2 columns from (a) GEMS, (b) WRF-$GEMS_{Updated}$ D02, and (c) WRF-$GEMS_{Base}$ D02 for snapshots at 02:00:00, 04:00:00, and 07:00:00 UTC, corresponding to approximately 09:00, 11:00:00, and 14:00:00 LT. (d) Tropospheric NO_2 column for daytime hours between GEMS (blue), WRF-$GEMS_{Base}$ D02 (dotted purple), and WRF-$GEMS_{Updated}$ D02 (solid purple) during the ASIA-AQ deployment period. Shaded regions indicate variability, represented by the 10th and 90th percentiles.

6 Evaluation using independent observations

To independently assess the performance of the optimized WRF-Chem simulations, we compared three model configurations: WRF_{Base}, $WRF_{Updated}$, and $WRF_{Updated+BC}$ ~~results~~ for D01 and D02 against a suite of independent airborne and ground-based observations distinct from the satellite measurements used in the emission optimization process (see step 4 Fig. 2). Comparisons between $WRF_{Updated}$ and $WRF_{Updated+BC}$ reflect the sensitivity of the model results to biases in the GEMS retrieval used to constrain emissions in the optimization process. This evaluation primarily emphasizes the results of D02 (4 km), which more directly represents urban-scale processes and exhibits improved performance compared to D01 across multiple observational datasets. These datasets provide an unbiased test of how well the model reproduces meteorological conditions and trace gas amounts beyond those directly constrained from GEMS observations. By evaluating WRF-Chem against surface monitoring networks (Thailand PCD), Pandora spectrometer measurements, and airborne ASIA-AQ data (e.g., GCAS NO_2 columns, DC-8 vertical profiles), we examine the model's ability to capture daytime variability, vertical structure, and surface concentrations of NO_2 across different spatial and temporal scales. Evaluation amongst these diverse datasets, therefore, helps to support the robustness of the emission correction framework described in this paper.

Table 2. Summary of WRF-Chem D02 validation statistics for WRF_{Base} (B) ~~and~~ $WRF_{Updated}$ (U), and $WRF_{Updated+BC}$ (U+BC) simulations evaluated against independent observational datasets. Metrics include mean bias (MB), mean error (ME), normalized mean bias (NMB), normalized mean error (NME), root-mean-square error (RMSE), and Pearson correlation coefficient (CORR).

670

671

Dataset	Species	Model Run	MB	ME	NMB	NME	RMSE	CORR
GCAS	Tropospheric NO_2 column (molecules cm^{-2})	**B**	4.0E+15	5.1E+15	93	1.2E+2	9.3E+15	0.64
		U	-2.4E+14	1.9E+15	-5.6	45	3.4E+15	0.54
		U+BC	6.8E+14	2.3E+15	16	53	4.1E+15	0.4
DC-8	NO_xO_3 NO_2 (ppbV)	**B**	1.3	1.4	82	87	2.8	0.97
		U	-0.81	0.8	-49	49	1.4	0.93
		U+BC	-0.38	0.49	-23	30.3	0.89	0.94
	CANOE NO_2 (ppbV)	**B**	1.4	1.5	87	94	2.9	0.97
		U	-0.76	0.76	-48	48	1.3	0.93
		U+BC	-0.34	0.50	-21	31	0.88	0.94
Ground Monitors	NO_2 (ppbV)	**B**	12	12	1.3E+2	1.3E+2	15	0.61
		U	-3.0	4.5	-32	47	5.6	0.53
		U+BC	0.1	4.0	0.6	43	5.5	0.55
	NO_x (ppbV)	**B**	14	15	1.4E+2	1.4E+2	18	0.59
		U	-3.6	4.9	-34	47	6.7	0.61
		U+BC	0.02	4.8	0.2	46	7.1	0.57
		B	-5.6	8.2	-18	27	10	0.87

	O_3 (ppbV)	**U**	4.9	8.5	16	27	11	0.81
		U+BC	3.1	6.6	10	22	9.2	0.86
Pandora	Tropospheric NO_2 column (molecules cm^{-2})	**B**	7.9E+15	8.6E+15	55	59	9.6E+15	0.71
		U	-6.6E+15	6.8E+15	-46	47	8.2E+15	0.67
		U+BC	-3.8E+15	4.9 E+15	-26	33	6.4 E+15	0.70

672

673 **6.1 Meteorological evaluation: winds and PBL height**

674 **6.1.1. Evaluation methodology**

675 To assess the performance of WRF-Chem meteorological fields, model~~l~~ed wind speed and direction were evaluated against
676 surface observations across the BMR during the ASIA-AQ campaign, 14 – 27 March 2024. Biomass burning was active in the
677 region during the study period, particularly over northern Thailand and Myanmar, with peak activity occurring in mid-March.
678 However, transport to Bangkok was likely limited due to weak synoptic flow and short chemical lifetime of NO_x, reducing its
679 impact on urban concentrations of NO_x in the BMR. Following March 20, precipitation suppressed fire activity, further
680 minimizing its impact. While biomass burning may contribute to broader regional enhancements, its impact on the inversion
681 results and coincident model evaluation over Bangkok is expected to be minor.
682
683
684 Surface observations of wind speed (m s^{-1}) and direction are obtained from Thailand Pollution Control Department (PCD)
685 ground monitor network data. Hourly surface wind observations from PCD network data were co-located with corresponding
686 model outputs for D01 and D02. For each site, model data were extracted from the nearest grid cell and interpolated to
687 observation timestamps. Daily wind roses were constructed to visualize and compare the frequency distribution of wind speed
688 and direction between the model and observations. For each station, wind speeds were binned into intervals of 0-1, 1-3, 3-5,
689 and 5-8 m s^{-1} and wind directions were grouped into 16 compass sectors. Normalized frequency counts were computed to
690 highlight dominant flow patterns. Data was averaged across stations within a larger bounding box around BMR, N-S (12.5-
691 15°N), W-E (99.5-101.5°E). For meteorology, we use a large box that spans the regional flow influencing Bangkok so that we
692 capture plume transport into and out of the city. Figure S8~~7~~a illustrates the station locations and averaging domains in Bangkok.
693

Additionally, we performed an evaluation of planetary boundary layer height (PBLH) to gauge the model's ability to capture daily PBLH magnitude and development during the deployment period. Observed PBLHs were derived from the NASA Langley airborne High Spectral Resolution Lidar Generation -2- (HSRL-2) data following a machine learning approach described in Christopoulos et al., (2025) for ASIA-AQ flight days, 18 March, 19 March, 23 March, and 25 March 2024. We evaluate PBLHs for aircraft raster periods (i.e., Raster 1: morning, Raster 2: afternoon, Raster 3: late afternoon/evening hours) and co-locate the observed PBLHs to modelled PBLHs based on a nearest grid-cell approximation.

6.1.2. Model-observation wind discrepancies

Within the observational data, a pronounced synoptic transition occurred over Thailand on 20 – 21March 2024, during which winds reversed from the typical southwesterlies that were present during the deployment period to northeasterlies, weakened substantially, coinciding with a drop in PBLH, and increased atmospheric stability (see PCD observations; Fig. S98-10). This transition played a major role in $PM_{2.5}$ and ozone exceedances by promoting stagnation over the BMR. The WRF-Chem simulation failed to capture the timing or strength of this abrupt wind shift and associated stagnation (Fig. S2a, S2b, S9b, S9c)S8a, S8b). The model instead maintained stronger southwesterly flow and higher PBLH, which led to unrealistic vertical mixing patterns compared to the observations (Fig. S98-S100). Since this mismatch drives discrepancies between model and observations that are unrelated to emissions adjustments/inversions, we exclude 20 – 21 March from the inversion/air quality evaluation. Although the simulation is driven by FNL, it is possible for WRF-Chem to mis-time a rapid synoptic transition. This event evolved on a relatively short spatial and temporal scale, and the coarse FNL boundary conditions may not have fully resolved the sharp changes in low-level flow. In addition, local processes (i.e., convection, PBL mixing, land-sea interactions) around Bangkok can cause the model to drift from the observed timing once it is integrated forward. We suspect these factors likely contributed to the later wind shift in WRF-Chem. We additionally found WRF-Chem (D01, D02) systematically overpredicts wind speed throughout the deployment (Fig. S2a, S2bS8a, S8b; Table S2) with better agreement from the high-resolution 4 km domain. This bias likely arises from thcat fact that many PCD monitors are embedded within the complex urban environment of Bangkok, which are not fully resolved at WRF-Chem's spatial scale.

The consequences of this mis-timed synoptic transition are most pronounced on 21 March as demonstrated in the next section, when weak large-scale flow made the simulation highly sensitive to local processes. Under these conditions, WRF-Chem misses the stagnation and dilutes pollutants too quickly. This helps explain why the model performs worst during the highest observed $PM_{2.5}$ and O_3 episode. These results indicate that forecast performance during similar weak-flow pollution events would benefit from improved representation of urban boundary-layer processes (e.g., surface roughness, urban canopy effects, and land–sea breeze structure).

6.1.3. Model-observation PBLH discrepancies

As shown in Fig. S10 and summarized in Table S3, WRF-Chem generally captures the temporal evolution of the PBLH observed by the aircraft. During the daytime (afternoon hours represented by Raster 2-3), the simulated PBL is broadly comparable to the observations, with median values that are often similar. However, the magnitude and sign of the model bias vary by day, with WRF-Chem demonstrating both over- and under- estimation likely related to the meteorological conditions. This day-to-day variability suggests PBLH biases are not systematic during the afternoon and may influence surface pollution dilation differently across individual cases, potentially contributing to variability in simulated daytime NO_2 concentrations rather than a consistent model bias.

6.2 Surface air quality evaluation

To evaluate model performance of surface trace gases, model~~l~~ed surface mixing ratios of NO_2, NO_x, and O_3 were evaluated against observations across BMR during 14 – 27 March 2024. Surface observations of these constituents (in ppbV) were obtained from Thailand PCD ground monitor network data. Like the evaluation of wind speeds and direction, data was co-located by extracting model information from the nearest grid cell and matched to the observation timesteps. Station data and model data were averaged across stations located within the Bangkok urban plume, bounded by N-S (13.5-14.6°N), W-E (100.2-101°E). Figure S8~~7~~b illustrates the station locations and averaging domain in Bangkok. For the air quality evaluation, we restrict the station averaging domain to a smaller area over the Bangkok urban plume. This smaller domain allows us to assess how the inversion and optimization specifically correct the local plume structure as opposed to a broader regional background.

6.2.1. NO_2 and NO_x

Figure 7a,7b,7c shows a comparison of average surface NO_2, NO_x, and O_3 (ppbV) for stations located within the Bangkok urban plume during the ASIA-AQ deployment for D02. D01 results are available in Fig. S11~~11~~. The 20 – 21 March is shaded in grey to indicate the period where the model mis-represented synoptic conditions resulting in enhancements in the observed concentrations due to stagnation. During this time, NO_2~~,~~ and NO_x ~~, and O_3~~ reached as high as 49, and 59 ppbV, ~~and 103 ppbV,~~ respectively. O_3 peaked at 103 ppbV the following day, on 22 March. Excluding this event, observed NO_2 and NO_x for this period generally ranged between 3 – 17 ppbV, and 3.5 – 22 ppbV respectively. Overall, there is a clear improvement in the representation of NO_2 and NO_x during this period in $WRF_{Updated}$ and $WRF_{Updated+BC}$ as shown in Fig 7a, 7b. Table 2 depicts summary statistics for this analysis. The baseline simulation, WRF_{Base} overestimates both NO_2 and NO_x with mean biases of +12 ppbV and +14 ppbV, respectively. With updated emissions, these large positive biases in $WRF_{Updated}$ were eliminated and

reversed, yielding mean biases of -3.0 ppbV for NO_2 and -3.6 ppbV for NO_x. The remaining negative bias is likely linked to the overprediction of winds, associated advection, and inversion related biases, which leads to locally diluted concentrations near urban sites as shown in the previous analyses. $WRF_{Updated+BC}$ further reduces the residual bias related to $WRF_{Updated}$, bringing mean biases closer to zero (e.g., +0.1 ppbV for NO_2 and +0.02 ppbV for NO_x), while maintaining comparable error and correlation, indicating accounting for the GEMS retrieval bias prior to inversion provides a modest additional improvement in model performance. Correlations with observed NO_2 and NO_x remain moderate (r = 0.5–0.6) reflecting potential spatial discrepancies in $WRF_{Updated}$ as seen in the previous analyses. Average diurnal cycles of NO_x for D02, D01 are further illustrated in Fig. S12a, c. Observed NO_x shows a clear morning peak and lower midday concentrations driven by boundary layer evolution. WRF_{Base} overestimates NO_x and exaggerates the morning peak while $WRF_{Updated}$ (dark purple) substantially improves both magnitude and timing but underestimates the evening peak values. $WRF_{Updated+BC}$ shows mixed performance with an overprediction of the morning peak while the evening peak aligns more closely with observations compared to $WRF_{Updated}$, indicating a trade-off in how the bias correction might redistribute NO_x across the diurnal cycle in the updated models.

6.2.2. O_3

Observed O_3 generally ranged between 14–57 ppbV, excluding the stagnation event on 20 – 21 March, and the baseline model captures this range reasonably well (Fig. 7c). After updating the emissions, O_3 increases in $WRF_{Updated}$ and $WRF_{Updated+BC}$ primarily at night, consistent with reduced NO titration following the decrease in NO_x emissions. Average diurnal cycles in D02 (Fig. S12~~a~~b,d) show that daytime O_3 production changes only modestly between base and updated simulations, indicating the emission update mainly affects nighttime chemistry, as opposed to shifting photochemical O_3 formation in the daytime. This pattern is reflected in the change in overall mean bias in Table 2, which shifts from an underprediction (-5.6 ppbV) to an overprediction (+4.9 ppbV) in $WRF_{Updated}$. The overprediction is not as substantial in $WRF_{Updated+BC}$ (+3.1 ppbV). While $WRF_{Updated}$ exhibits an earlier O_3 peak in D02 compared to observations, this shift is not present in D01 despite the same emission inversion being applied. This likely suggests the discrepancy is driven by domain-dependent differences in meteorological or chemical processes, rather than an error in the inferred NO_x emission timing. The O_3 peak is better represented in $WRF_{Updated+BC}$.

The relatively weak daytime O_3 response to decreased NO_x emissions is consistent with recent analyses of O_3 formation sensitivity during ASIA-AQ with in situ measurements, which indicate that the BMR exhibits mixed sensitivity to NO_x and VOCs, in contrast to the predominantly NO_x-limited regimes observed at other ASIA-AQ locations (e.g., Manila) (Cho et al., 2026). In this mixed-sensitivity regime, changes in NO_x emissions alone are not expected to strongly perturb daytime O_3 production, further providing validation for the minimal daytime O_3 response observed here. The magnitude of the daytime O_3

response varies by domain/model as shown by the D01 results (Fig. S12d~~12b~~). Although $WRF_{Updated}$ D01 better matches observed daytime O_3, this apparent improvement likely reflects NO_x titration associated with spatial dilution.


Overall, these results suggest that while the updated emissions improve the model's NO_x representation, further improvements in VOC representation and local mixing processes may be needed to fully capture daytime O_3 levels in the BMR. Nevertheless, the updated emissions substantially improve the model's simulation of surface NO_2 and NO_x, resulting in a more realistic overall representation of air quality in Bangkok.

6.3 Evaluation with Pandora column observations

We additionally evaluate model~~l~~ed NO_2 columns against Pandora measurements for the Bangkok site. Pandora NO_2 evaluation used Level 2 direct-sun total column retrievals, filtered for high-quality measurements (quality flag = 10), and averaged to hourly means. Tropospheric columns from Pandora were estimated by subtracting coincident or closest GEMS stratospheric NO_2 at the nearest satellite pixel. Model columns were sampled at the nearest grid cell to the Pandora site and temporally matched to observations.

Figure 7d depicts a comparison of WRF_{Base} (light purple triangles), $WRF_{Updated}$ (dark purple squares), $WRF_{Updated+BC}$ (magenta dots) and Pandora (pink stars), tropospheric NO_2 column measurements for days with high quality data. Throughout this period, Pandora measurements generally ranged between approximately $7 \times 10^{15} – 3.5 \times 10^{16}$ molecules cm^{-2} and fit between WRF_{Base} which places columns higher (e.g., up to 5×10^{16} molecules cm^{-2}), $WRF_{Updated+BC}$, and ~~and~~ $WRF_{Updated}$ which places columns lower (e.g. 2×10^{15} molecules cm^{-2}). Average statistics between model cases for this analysis are shown in Table 2. Biases are generally similar between model cases. WRF_{Base} overestimates column NO_2 as illustrated by the mean bias ($+7.9 \times 10^{15}$ molecules cm^{-2}) whereas emissions updates contribute to an underestimation (-6.6×10^{15} molecules cm^{-2}). However, there are some improvements in absolute error metrics with ME and RMSE decreasing by roughly 20% and 12% respectively, and NME dropping from 61% to 48%. The greatest bias improvements overall are seen in $WRF_{Updated+BC}$ (-3.8×10^{15} molecules cm^{-2}). Figure S13~~14~~ illustrates the spatial distribution of mean tropospheric model NO_2 bias for GEMS and Pandora during the time reflected in Fig. 7d. In the WRF_{Base} simulation, a strong positive bias is seen within and north of the Pandora site, indicating the overestimation of NO_2 columns (up to 2×10^{16} molecules cm^{-2}) in the urban plume relative to GEMS. This pattern reiterates the point that the prior anthropogenic emissions (based on EDGAR v5) were too large with pollutant accumulation occurring downwind of Bangkok as a result. After applying the emission updates, the $WRF_{Updated,}$ simulation essentially eliminates this bias. The overall bias near the Pandora site becomes close to neutral or slightly negative, demonstrating the optimization effectively corrected the spatial overprediction. $WRF_{Updated+BC}$ further reduces the residual negative bias relative to $WRF_{Updated}$ with NMB improving from -46% to -26% and RMSE decreasing from 8.2×10^{15} to 6.4×10^{15} molecules cm^{-2}. This indicates that accounting for the GEMS retrieval bias in the initial optimization process partially corrects the remaining underestimation,

although some spatial discrepancies persist dude to transport and plume placement error. ~~The remaining negative bias close to the city center/Pandora site is likely related to discrepancies in the WRF$_{Updated}$ urban plume placement, overpredicted wind speeds, and inversion biases related to the GEMS retrievals.~~ Examples of ~~D~~daily biases for WRF$_{Base}$ D01, D02, and WRF$_{Updated}$ D01, D02 can be found in Fig. S14~~15~~ and Fig. S15~~16~~.



825
826
827
828

829
830
831
832
833




Figure 7. Comparison of WRF_{Base} D02 (dotted purple), ~~and~~ $WRF_{Updated}$ D02 (solid purple), and $WRF_{Updated+BC}$ D02 (solid magenta) simulations against Thailand Pollution Control Department (PCD) ground-monitor network observations for (a) NO_2 mixing ratio (ppbV~~v~~), (b) NO_x mixing ratio (ppbV~~v~~), and (c) O_3 mixing ratio (ppbV~~v~~) during the ASIA-AQ deployment period

(14 – 27 March 2024). represent averages across stations within the Bangkok urban plume. (d) Comparison of tropospheric NO_2 columns from WRF_{Base} D02 (light purple triangles), $WRF_{Updated}$ D02 (dark purple squares), $WRF_{Updated+BC}$ D01 (magenta dots) and Pandora (pink stars) for high-quality observations during the ASIA-AQ deployment period (18 – 27 March 2024). The shaded region represents the stagnation period, 20 – 21 March excluded from the evaluation.

6.4 Evaluation with ASIA-AQ aircraft measurements

The Airborne and Satellite Investigation of Asian Air Quality (ASIA-AQ) was a NASA field campaign conducted in February – March 2024 to advance the understanding of urban and regional air quality across East and Southeast Asia. Targeting several megacities (e.g., Manila, Seoul, Bangkok, Chiang Mai), the campaign combined satellite observations, aircraft measurements, ground-based monitoring, and modeling approaches to characterize pollution sources and validate satellite retrievals. A main objective of ASIA-AQ was to evaluate the data from GEMS. Airborne observations were collected using the NASA DC-8 in situ and LaRC G-III remote sensing aircraft, with coordinated support from ground-based networks such as Pandora and AERONET. Additionally, chemical transport models (e.g., GEOS-Chem, GEOS-FP, MUSICA, WRF-Chem, WRF-CMAQ) played a key role in real-time flight planning and post-campaign interpretation.

6.4.1. GCAS

The GEOstationary Coastal and Air Pollution Events (GEO-CAPE) Airborne Simulator (GCAS) is an airborne UV-Vis spectrometer that was flown on the G-III aircraft during the ASIA-AQ campaign. GCAS was designed to simulate the spectral capabilities of TEMPO and GEMS, but with a much finer pixel resolution of approximately 250 x 560 m at flight altitude (Janz et al., 2019; Lee et al., 2024). GCAS uses a push-broom remote sensing technique and consists of two spectrometer channels: a UV-Vis channel (300-490 nm) optimized for air quality measurements, and a Vis-NIR channel (480-900 nm) for ocean color observations (Kowalewski and Janz, 2014; Lee et al., 2024). This work focuses on the NO_2 retrieval from spectra in the UV-Vis channel. Retrieval details and validation results can be found in Judd et al., (2020) but were previously found to be unbiased with uncertainties within +/-25%. As in previous field campaigns, the aircraft executed a "lawnmower" flight pattern with parallel flight lines spaced 6.3 km apart, providing about 10% overlap between flight lines assuming a flight altitude of 28,000 feet. This flight strategy, combined with the instrument's 45° field of view, allowed for the generation of gap-free NO_2 column maps up to three times per day, period referred to as a "raster". Due to the short duration of each raster period (~3 hours), local meteorological conditions often influence the fine-scale structures observed in the GCAS NO_2 data (Goldberg et al., 2024). Here, we evaluate the WRF-Chem runs against GCAS for several flight days: 18 March, 19 March, 23 March, and 25 March 2024.

We perform the evaluation for each raster separately to isolate specific flight patterns and accurately evaluate spatial gradients in NO_2 between the model and observations. For each analysis, GCAS pixels corresponding to the flagged raster were retained for comparison. Additional filters were applied to remove poor-quality retrievals. We masked pixels with cloud or sun glint contamination based on a provided flag variable (cloud_glint_flag = 1) and discarded retrievals with missing data or undefined AMFs. GCAS provides separate NO_2 vertical columns above and below the aircraft, as well as model-derived scattering weights, and AMFs for both portions. The above and below aircraft contributions can be approximated as the stratospheric and tropospheric contributions, respectively. For this evaluation, we focus exclusively on the NO_2 column below the aircraft or the tropospheric column NO_2, which is the portion most relevant to surface-level air quality and most comparable to our WRF-Chem results. We compute the below-aircraft averaging kernel, A_i^{below}, as:

$$A_i^{below} = \frac{SW_i}{AMF_{below}} \quad (17)$$

Here, SW_i represents the scattering weight for layer i, representing the sensitivity of the measured radiance to NO_2 in that layer. This averaging kernel represents the satellite-equivalent vertical sensitivity to NO_2 below the aircraft and was used to weigh the WRF-Chem vertical profile.


WRF-Chem output including, NO_2 mixing ratio, pressure, temperature, and height were used to compute air density and convert volume mixing ratios to number densities as previously done in the GEMS evaluation. Each GCAS pixel was temporally matched to the nearest model output time (rounded to nearest hour) and spatially co-located by finding the nearest WRF-Chem grid cell. To isolate the portion of the model column below the aircraft, we filtered the model levels based on the aircraft altitude reported at each pixel. We compute the tropospheric column as shown in Eq. (10). To generate a model column that reflects the vertical sensitivity of the GCAS retrieval, we interpolate the WRF-Chem profile to the number of GCAS vertical layers (49), converted the mixing ratios to number density, and applied the GCAS averaging kernel to yield WRF-GCAS.

The maps in Fig. 8 illustrate a spatial comparison between (a) GCAS, (b) WRF-GCAS$_{Updated+BC}$ D02, (c) WRF-GCAS$_{Updated}$ D02, and (~~d~~e) WRF-GCAS$_{Base}$ D02 over BMR for a flight day on 18 March 2024. D01 spatial comparisons are available in Fig. S16. 18 March represents a typical example of local pollution dominating BMR with minimal influences from long-range pollution transport and biomass burning. The GCAS instrument generally places tropospheric column NO_2 values in BMR between 5 x 10^{15} – 2 x 10^{16} molecules cm^{-2}, with the largest enhancements observed in the city center, as also seen in the GEMS data (Fig. 7). A clear N-S plume is visible in the data, reflecting persistent southerly onshore flow from the Gulf of Thailand. This pattern coincides with the seasonal shift from the northeast to southwest monsoon. Overall, WRF-GCAS$_{Updated}$ and WRF-GCAS$_{Updated+BC}$ ~~can~~ generally better capture the spatial differences and magnitudes of tropospheric column NO_2 in BMR for different raster periods compared to WRF-GCAS$_{Base}$. WRF$_{Updated+BC}$ outperforms WRF$_{Updated}$ in capturing NO_2 column enhancements, particularly during raster 2 (late morning – afternoon LT). GCAS also illustrates enhanced NO_2 columns

southeast of Bangkok in a region known as the Eastern Economic Corridor (EEC), a major hub for industrial activity (e.g., automotive manufacturing, petrochemicals, electronics). $WRF_{Updated}$, $WRF_{Updated+BC}$, and WRF_{Base} tend to underestimate pollution levels in the EEC. This is likely related to wind speed overprediction and a lack of updated regional source data in the EDGAR v5 inventory (since this is a region outside of the performed inversion). Information from a local emissions inventory, or additional inversions performed on this region could aid the model in better representing the air quality in this region, which has similar magnitudes of column NO_2 (2×10^{16} molecules cm^{-2}) to the Bangkok city center.

Statistics for this analysis are shown in Table 2. Overall, statistics are representative of substantial bias and error improvements in the updated runs from WRF_{Base}, which significantly overpredicts column NO_2 amounts in the region for all raster periods. For example, ME and RMSE are improved by ~62% from WRF_{Base} and $WRF_{Updated}$. Mean biases are negative which indicate $WRF_{Updated}$ is underpredicting NO_2 columns. This is likely the result of the low inversion biases and windspeed overprediction. However, this underprediction is improved throughout the day as seen in Fig. S1720a, S1720b. This is in opposition to WRF_{Base}, which increases in biases and error for each raster as also shown in Fig. S1720a, S1720b. These daytime patterns also highlight an important distinction between the benefits of geostationary (GEMS) versus LEO observations. Morning improvements are particularly strong because the inversion directly constrains the rapid rise in emissions during the morning, something that LEO sensors generally under sample due to their limited overpass frequency. Midday and afternoon performance remains improved as well, but the morning hours show the clearest advantage of daytime GEO sampling for capturing short-timescale emission variability. Similar results are demonstrated in $WRF_{Updated}$ D01 (Fig. S18). $WRF_{Updated+BC}$ is indicative of a partial correction to the negative bias in $WRF_{Updated}$, shifting the NMB from -5.6% to +16%. While this brings the simulated NO_2 columns closer to observations in a mean sense, is it followed by increases in error, suggesting a trade-off between bias reduction and overall model performance.

Correlations remain moderate ~~high~~ (0.4~~0.6~~ – 0.6~~0.8~~) for ~~both~~ model cases, however, $WRF_{Updated}$ and $WRF_{Updated+BC}$ illustrate ~~has~~ weaker correlations likely due to (i) subtle shifts in the urban plume placement (i.e., shift in a more northerly direction), and (ii) a reduction in dynamic range after emissions corrections. For example, when plume magnitudes are lowered, the variability shrinks allowing the correlations to become less sensitive to spatial-temporal agreement and more sensitive to small plume-placement differences as is the case here.


Mar 18th,2024
(a)
GCAS
(b)
WRF-GCAS Updated + BC
(c)
WRF-GCAS Updated
(d)
WRF-GCAS Base
Raster 1
07:44 - 10:30 LT
Raster 2
10:30 - 12:59 LT
Raster 3
14:12 - 16:20 LT
D02
15′
14°N
45′
30′
13°N
100°E 15′ 30′ 45′ 101°E 15′
3 x 10^15
4 x 10^15
5 x 10^15
6 x 10^15
7 x 10^15
8 x 10^15
1 x 10^16
1.5 x 10^16
2 x 10^16
Tropospheric Column NO2
[molecules/cm2]




Figure 8. Spatial comparison over the BMR on 18 March 2024 of tropospheric NO_2 columns from (a) GCAS, (b) WRF-GCAS$_{Updated+BC}$ D02, (c) WRF-GCAS$_{Updated}$ D02 and (d) WRF-GCAS$_{Base}$ D02 for raster periods corresponding to morning, afternoon, and early evening local time.


6.4.2. NO_xO_3 and CANOE (DC-8)

We additionally compare the model simulations to in situ NO_2 data gathered by the National Center for Atmospheric Research's (NCAR) NO_xO_3 and NASA's GSFC's Compact Airborne NO_2 Instrument (CANOE) aboard the DC-8 aircraft. The NO_xO_3 instrument is a 3-channel chemiluminescence instrument designed for the measurement of NO, NO_2, and O_3 (Ridley et al., 1992; Ridley and Grahek, 1990). CANOE measures NO_2 using non-resonant laser induced fluorescence (LIF) (St. Clair et al., 2019). We use the 1-sec DC-8 data for the flights conducted in Thailand during March 2024 (18 March, 25 March). A visual depiction of the DC-8 flight path is shown in Fig. S19. The typical flight path included several low-altitude descents/ascents over several airports (see Fig. S19b) along with long-distance transects across Central Thailand.

To evaluate model performance, we match hourly WRF-Chem outputs from both the 20 km (D01) and 4 km (D02) domains to the aircraft location and time. For each observation, we extract co-located model NO_2 values along the aircraft track. Profiles are filtered by location (e.g., Don Mueang International Airport in Bangkok) and direction (e.g., ascent or descent) using flags in the provided observational datasets. Individual profiles are grouped by their unique number, and we bin the observations and model output by altitude (50 m vertical bins). For each bin, we compute mean NO_2 from the aircraft and model datasets. These vertically resolved comparisons allow us to assess model skill in capturing the observed structure and magnitude of NO_2 within the boundary layer and lower troposphere.

Figure 9 depicts a comparison of aggregated in situ NO_2 vertical profiles over the Don Mueang International airport for (a) 18 March and (b) 25 March. Model scenarios are shown in shades of purple (WRF_{Base}: dotted dark purple; $WRF_{Updated}$: solid dark purple; $WRF_{Updated+BC}$ solid magenta), and in situ information from two instruments aboard the DC-8, NO_xO_3 (solid) and CANOE (dashed) are displayed in black. The location of the Don Mueang International Airport with respect to the Bangkok urban plume is depicted in (c). We separate and group the data by time of day to gauge how the model runs perform with respect to time of day. Corresponding plots for D01 are available in Fig. S20. As shown in Fig. 9, the NO_xO_3 and CANOE NO_2 measurements exhibit excellent agreement throughout the analysis period. This consistency between two independent in situ instruments strengthens confidence in the observational data used for model evaluation. Comparison with DC-8 in situ NO_2 profiles show that WRF_{Base} consistently overestimated observed concentrations, with a mean bias of +1.3 ppb and normalized mean bias near 80%. Figure 9a and Figure 9b indicate the overestimation is most pronounced near the surface during the morning, where WRF_{Base} exceeds observations by nearly 25 ppb. $WRF_{Updated+BC}$ also suggests an overestimation compared to aircraft observations in the morning. In contrast, $WRF_{Updated}$ captures the observed morning surface enhancements accurately. As time progresses, however, $WRF_{Updated}$ and $WRF_{Updated+BC}$ begin~~s~~ to underestimate the column (average bias ~ -0.3 to -0.8 ppb), though both mean error and RMSE remain improved in the updated cases relative to WRF_{Base} (Table 2). This daytime behavior likely reflects the nature of the GEMS-based emission constraints, which are directly applied only until 14:00:00 LT, after which a single daily scaling factor is applied on the input emissions. As afternoon PBL growth and photochemistry evolve, the fixed scaling likely results in an underestimation in the afternoon profiles. Additionally, overestimated model winds speeds may further dilute surface concentrations, likely contributing to the midday and afternoon biases shown in $WRF_{Updated}$, and $WRF_{Updated+BC}$. The afternoon underestimation is slightly reduced in $WRF_{Updated+BC}$, likely reflecting the impact of the bias correction applied prior to inversion/NO_x emission optimization.

Correlations for $WRF_{Updated}$ ($r = 0.93 – 0.97$) and $WRF_{Updated+BC}$ ($r = 0.94$) remain high ~~($r = 0.93 – 0.97$)~~ indicating the simulation preserved the observed structure while improving the overprediction. Overall, these results highlight the effectiveness of the emission updates in accurately capturing morning surface air quality. This suggests that while the updated emissions substantially improve the magnitude of model~~l~~ed NO_2, remaining discrepancies are driven primarily by meteorological transport rather than emission magnitude.


(a)
(b)
(c)
March 18th
D02 – Morning
March 18th
D02 – Midday
March 18th
D02 – Afternoon
March 25th
D02 – Morning
March 25th
D02 – Midday
March 25th
D02 – Afternoon
DC-8 (NO_xO_3)
DC-8 (CANOE)
WRF_{Base}
$WRF_{Updated}$
$WRF_{Updated+BC}$
Altitude [m]
NO_2 [ppb]
GEMS
Mar 18th
01:45 UTC
Tropospheric Column NO_2 [molecules/cm^2]




Figure 9. Comparison of WRF-Chem D02 simulations (WRF_{Base}: dotted dark purple; $WRF_{Updated}$: solid dark purple; $WRF_{Updated+BC}$: solid magenta) with airborne in situ measurements from the CANOE (dashed black) and NO_xO_3 (solid black) instruments aboard the DC-8 for (a) 18 March and (b) 25 March 2024. Profiles are grouped by morning (06:00:00–11:00:00 LT), midday (11:00:00–13:00:00 LT), and afternoon (13:00:00–17:00:00 LT) approaches at Don Mueang International Airport in northern Bangkok, whose location relative to the Bangkok urban plume is shown within a GEMS snapshot in (c).

7 Discussion

7.1 Implications of GEMS NO_2 retrieval biases

~~Comparisons between GEMS tropospheric NO_2 columns and independent column references indicate a systematic low bias in the GEMS product, particularly over the Bangkok region. In our analysis during ASIA-AQ, GEMS total NO_2 columns are generally lower than coincident Pandora total column NO_2 observations (Fig. S13) (mean bias ≈ -9.2×10^{15} molecules cm^{-2}), consistent with recent independent validation studies of the operational GEMS v3 product over Bangkok and South Korea that report low biases relative to ground-based sun-photometer and DOAS measurements (Bae et al., 2025; Jung et al., 2025). Bae et al. (2025) shows that GEMS v3 increasingly underestimates NO_2 relative to Pandora under high NO_2 conditions (>1×10^{16} molecules cm^{-2}) as is the case for Bangkok pollution levels. In Jung et al. (2025), validation results over Bangkok indicate a~~

~~pronounced low bias in GEMS tropospheric NO_2 columns relative to Pandora, with regression slopes of ~0.35 for v2.0 and ~0.28 for v3.0, indicating increasing underestimation at higher NO_2 levels. While moderate correlations (r ≈ 0.6–0.7) suggest that GEMS captures temporal variability, column magnitudes are substantially underestimated, particularly under polluted conditions. The persistence of this behavior in the v3.0 product indicates that the low bias is not fully corrected by recent algorithm updates and is consistent with retrieval sensitivity limitations in highly polluted urban environments (Jung et al., 2025).~~

~~To further evaluate the relative behavior of GEMS and airborne GCAS NO_2 columns, we compare their tropospheric columns using $WRF_{Updated}$ as a common transfer framework (Fig. 10). Figure 10 compares the observed GCAS/GEMS NO_2 column ratios with ratios calculated after both datasets are mapped through $WRF_{Updated}$. In the observations, GCAS columns are consistently higher than GEMS, with GCAS/GEMS ratios of ~2–3 on most days and values reaching ~6–7 on 21 March. In contrast, the corresponding WRF-$GCAS_{Updated}$/WRF-$GEMS_{Updated}$ ratios are substantially smaller (~1.1–1.7), even on 21 March. This is consistent with a low bias in GEMS NO_2 columns as found in previous comparisons.~~

In our study, the comparisons with independent aircraft and ground-based observations indicate the satellite-constrained emissions without bias-corrections presented here may represent a low estimate. In particular, the systematic low bias in GEMS NO_2 is consistent with the negative biases often seen in the $WRF_{Updated}$ ~~updated~~ simulations. Importantly, this bias could not be diagnosed using the satellite data alone.


To further evaluate the relative behavior of GEMS and airborne GCAS NO_2 columns, we compare their tropospheric columns using $WRF_{Updated+BC}$ as a common transfer framework (Fig. S21~~0~~). Figure S21~~0~~ compares the observed GCAS/GEMS NO_2 column ratios with ratios calculated after both datasets are mapped through $WRF_{Updated+BC}$. In the observations, GCAS columns are consistently higher than the raw GEMS columns as illustrated by the orange line, with GCAS/GEMS ratios of ~2–3 on most days and values reaching ~6–7 on 21 March. In contrast, the corresponding WRF-$GCAS_{Updated+BC}$/WRF-$GEMS_{Updated+BC}$~~red~~ ratios are closer to 1:1 ~~substantially smaller~~ (~1.1–1.7), even on 21 March. ~~This is consistent with a low bias in GEMS NO_2 columns as found in previous comparisons.~~ This behavior is consistent with a low bias in GEMS NO_2 columns relative to GCAS. After applying the bias correction, the $GCAS/GEMS_{BC}$ ratios approach 1:1 on most days, indicating improved consistency between the datasets, and supporting the effectiveness of the applied GEMS bias correction.

When incorporating bias-corrected retrievals into the top-down inversions and NO_x emission optimization process, the resulting $WRF_{Updated+BC}$ run reduces the negative biases in the independent validation, in some cases bringing the model results even closer to observations as seen in the surface air quality analysis. However, this improvement is not uniform. In certain comparisons (e.g., GCAS and DC-8), $WRF_{Updated+BC}$ introduces a tendency toward overprediction in morning hours, highlighting tradeoffs associated with the bias correction. These results suggest that while accounting for the retrieval bias can

improve mean model performance, additional uncertainties such as overestimated wind speeds and associated transport errors continue to influence the representation of modeled NO_2.

Here, the integration of ground-based, airborne, satellite, and model data provides a powerful framework not only for improving emissions but also for identifying limitations within individual observing systems. While $WRF_{Updated}$ clearly outperforms the baseline model, and $WRF_{Updated\ +BC}$ offers targeted improvements in reducing systematic bias, the combined observational evidence highlights the necessity of a multi-platform validation to fully interpret the satellite-based emission estimates.

Despite this low bias in the GEMS retrieval, the high-frequency daytime sampling provided by geostationary observations offers critical constraints on daytime variability and plume evolution that are particularly valuable for emission inversion and air quality modelling. For example, the GEMS-constrained emission adjustments presented here were critical for improving the temporal evolution of NO_x in WRF-Chem, resulting in substantial and robust improvements in model performance across independent evaluations. Future work may further benefit from continued refinement of bias-corrected GEMS products and upcoming algorithm improvements (e.g., v4), alongside improved representation of spatial representation of emissions within the model. ~~the application of bias-corrected GEMS products or future GEMS algorithm updates and corrections (e.g., v4 products).~~



~~**Figure 10.** Daily raster (e.g., R1, R2, R3) comparison of tropospheric NO_2 columns from $WRF_{Updated}$ D02 (dashed purple), $WRF\text{-}GCAS_{Updated}$ D02 (dashed green), $WRF\text{-}GEMS_{Updated}$ D02 (dashed blue), GCAS (green), and GEMS (blue) for ASIA-AQ flight days. Dashed model lines indicate simulations with instrument-specific averaging kernels applied. The bottom panel illustrates the ratio GCAS/GEMS (orange) and WRF-GCAS/WRF-GEMS (black).~~

7.2 Limitations and potential extensions of the inversion framework

The current model emission update framework optimizes the temporal evolution of emissions but does not explicitly resolve regional variability, as a single set of hourly scaling factors is applied uniformly across all grid cells within the BMR. As a result, the inversion preserves the spatial structure of the prior inventory, and any inaccuracies in the spatial distribution of emissions are not corrected.

Extending this approach to resolve emissions at finer scales would require allowing emissions to vary across grid cells or subregions, supported by additional constraints. These constraints could include regional higher-resolution regional prior

inventories when available, as well as higher-resolution observations (e.g., GCAS) that can better resolve the urban variability in NO_2 as shown in this analysis.

8 Conclusion ~~and Discussion~~

8.1 A GEO-constrained framework for anthropogenic NO_x emissions

Accurate urban NO_x emissions remain a major challenge for air quality modeling efforts, but geostationary satellites now offer a path forward. This work takes a novel approach in improving urban NO_x emissions using daytime GEO satellite observations. We quantify and reduce biases in modeled NO_2 over the Bangkok Metropolitan Region (BMR), by integrating GEMS constraints into a high-resolution model's daytime prior emission profile. We first used top-down inversions of hourly GEMS NO_2 columns with the Cross-Sectional Flux (CSF) method to develop an average daytime NO_x profile for the BMR in March 2024. Using this information, we developed and applied an optimization framework that incorporates physical constraints (i.e., with regards to emissions accumulation and lifetime) to reshape the WRF-Chem's daytime emission pattern and magnitude to better reflect observed emissions variability in the BMR.

This represents one of the first applications of the *ddeq* framework to estimate hourly urban NO_x emissions with geostationary observations in Southeast Asia. In contrast to a full chemical reanalysis, the *ddeq*-based framework provides an efficient and scalable alternative that does not require repeated model reinitialization of extensive chemical state optimization. This makes the approach particularly suitable for regional applications and for broader application across different modeling systems and urban environments within the domain of GEO sensors.

8.2 Model improvements across independent observational platforms

Re-running WRF-Chem (D01 – 20 km; D02 – 4 km) with an updated NO_x emission profile ($WRF_{Updated}$; $WRF_{Updated+BC}$) led to substantial improvements across multiple independent datasets. Model evaluation with ground monitors, Pandora, GCAS, and DC-8 observations consistently showed reduced biases and errors relative to the baseline simulation based on EDHAR v5 emissions. For example, surface-level comparisons with Thailand PCD ground network data confirmed some of these trends in $WRF_{Updated}$, with mean biases in NO_2 and NO_x decreasing from +12 - +14 ppbV in WRF_{Base} to roughly -3 ppbV in $WRF_{Updated}$, and 0.02 to 0.1 ppbV in $WRF_{Updated+BC}$. At the Bangkok Pandora site, mean bias shifted from strongly positive ($+7.9 \times 10^{15}$ molecules cm^{-2}) to negative, and normalized mean errors decreased by ~20-~~425~~% in $WRF_{Updated}$ and $WRF_{Updated+BC}$. Evaluation against GCAS airborne column retrievals further showed that the emissions updates improved spatial variability and magnitude of NO_2 across the Bangkok urban plume, reducing mean and root mean square errors by ~55-6~~0~~% in $WRF_{Updated}$ and $WRF_{Updated+BC}$ compared to WRF_{Base}. The DC-8 in situ vertical profiles further illustrated that $WRF_{Updated}$ could substantially reduce near-surface overestimation in morning hours and preserve the observed vertical structure of NO_2 mixing ratios.

8.3 Assessing GEMS bias through multi-platform integration

Overall, these results demonstrate that incorporating geostationary satellite constraints into regional, high-resolution (4 km) chemical transport models can substantially improve the representation of urban air quality over Bangkok. Large biases in the baseline simulation are likely driven by reliance on outdated global emissions inventories (e.g., EDGAR v5), which do not reflect recent changes in regional anthropogenic activity, as well as uncertainties in bottom-up methodologies where updated local estimates appear systematically high. The satellite-constrained emission estimates derived here are consistent with independent top-down approaches (e.g., TCR-3) based on different satellite platforms and methodologies, increasing confidence in the inferred reductions.

Comparisons with independent aircraft and ground-based observations further indicate that the GEMS-constrained emissions presented here may represent an underestimate. In particular, the systematic low bias reported for the GEMS v3 NO_2 product is consistent with the remaining negative biases observed in $WRF_{Updated}$. By accounting for this bias through the application of bias-corrected retrievals, $WRF_{Updated+BC}$ reduces these negative biases, although in some cases introduces a tendency toward overprediction, highlighting tradeoffs associated with the correction. While model output (e.g., WRF_{Base}) can serve as a transfer standard for comparing different observing systems, these results highlight the importance of integrating ground-based, airborne, satellite, and model information to robustly identify biases and improve emission estimates. ~~This bias could not be diagnosed using the satellite data alone. Instead, the integration of ground-based, airborne, satellite and model information together provides a powerful framework not only for improving emission estimations, but also for identifying limitations within individual observing systems.~~

8.4 Value of hourly GEO constraints, high-resolution modelling and future directions

A key strength of this framework is the use of hourly daytime constraints uniquely provided by geostationary observations, which enable direct characterization of daytime emission variability and plume evolution that cannot be captured by once-daily low-Earth-orbit measurements. These daytime constraints are particularly important for urban environments, where emissions, chemistry, and boundary-layer dynamics vary rapidly and strongly influence air quality impacts.

Coarse (~12 km) simulations have been shown to inadequately represent circulations in coastal environments, and nonlinear NO_x chemistry, leading to systematic biases in simulations of NO_2 (Hsu et al., 2026; Valin et al., 2011; Verreyken et al., 2025; Yu et al., 2023). These limitations in model resolution are expected in coastal megacities such as Bangkok with complex local topography and land-sea contrasts. Consistent with recent TEMPO-based emission studies indicating that model resolution can limit the robustness of GEO-based top-down NO_x constraints (Hsu et al., 2026), these considerations motivate the use of

4 km WRF-Chem simulations in this work as a necessary framework for accurately interpreting geostationary satellite observations and constraining urban NO_x emissions.

While remaining discrepancies, including the negative model biases, are likely influenced in part by overpredicted model wind speeds and associated transport errors, future work could benefit from finer-resolution simulations (< 4 km) and more advanced urban parameterizations (e.g., multi-layer urban canopy models; Liu et al., (2025)) to better represent urban flow, drag, and mixing. Continued validation and further development of bias-corrected for updated GEMS retrievals will also strengthen the use of geostationary NO_2 products. Nevertheless, the high-frequency daytime sampling provided by GEMS already offers critical information for emission inversion, and the integration of GEMS-derived constraints into WRF-Chem represents a scalable pathway toward near-real-time, satellite-informed emissions estimation and improved air quality forecasting for rapidly developing megacities.

Code Availability.

The code repository for *ddeq v1* is available on GitLab: https://gitlab.com/empa503/remote-sensing/ddeq, Kuhlmann, 2024a. The implementation of *ddeq* v1 for top-down NO_x estimates and model optimization scripts are publicly available via Zenodo (https://doi.org/10.5281/zenodo.18381169).

Data Availability.

All ASIA-AQ field campaign data used in this study (i.e., GEMS NO_2 v3, NOxyO3, CANOE, GCAS, Thailand PCD air quality monitoring data) are openly available and were acquired from their NASA Langley Research Center maintained archive: https://www-air.larc.nasa.gov/cgi-bin/ArcView/asiaaq. Pandora total column NO_2 observations are available through the Pandonia Global Network website: https://www.pandonia-global-network.org/home/documents/pgn-data/. ERA5 meteorological reanalysis data were obtained from the Copernicus Climate Data Store: https://cds.climate.copernicus.eu/. Global emissions used in this study are openly available as well, HTAP v3.2 (https://edgar.jrc.ec.europa.eu/dataset_htap_v32), MIXv3.1(https://csl.noaa.gov/groups/csl4/modeldata/data/Li2023/),EDGARv5(https://edgar.jrc.ec.europa.eu/dataset_ghg50), ODIAC (https://db.cger.nies.go.jp/dataset/ODIAC/). THAI-KMUTT, and TCR-3 emissions can be made available upon request.

Supplement.

The supplement related to this article is available online.

Author Contributions

JAC designed the study, performed the model simulations, conducted the data analysis, prepared all figures, and wrote the initial draft of the manuscript. PS supervised the project, contributed ideas, guidance, and discussions regarding the model framework and interpretation of results. MM provided additional guidance on the modeling framework and interpretation of results. All authors discussed the results and contributed to the final version of the manuscript.

Competing Interests

At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

Acknowledgements

We thank the entire ASIA-AQ team for their contributions to the field experiment. We extend our gratitude to the instrument teams for their efforts in conducting in situ measurements, to the airborne remote sensing teams (HSRL-2 and GCAS) for their data collection and support, and the GEMS team for providing the data utilized in this work. We also thank Dr. Daniel Goldberg for helpful discussions related to top-down emissions estimation methodologies, including recommendations on the *ddeq* library and guidance on inversion methodology development. The TCR-3 product was generated by calculations using the Earth Simulator with the support of the Japan Agency for Marine-Earth Science and Technology.

Financial Support

The funding for this research came from the National Aeronautics and Space Administration under awards 80NSSC22M0266 and 80NSSC23K0786. Additionally, part of this work was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under contract to NASA (80NM0018D0004~~The~~). The views expressed in this manuscript are those of the authors and do not reflect the views of NASA.

References

Agarwal, P., Stevenson, D. S., and Heal, M. R.: Evaluation of WRF-Chem-simulated meteorology and aerosols over northern India during the severe pollution episode of 2016, Atmospheric Chemistry and Physics, 24, 2239–2266, https://doi.org/10.5194/acp-24-2239-2024, 2024.

Ahmadov, R., McKeen, S. A., Robinson, A. L., Bahreini, R., Middlebrook, A. M., Gouw, J. A. de, Meagher, J., Hsie, E.-Y., Edgerton, E., Shaw, S., and Trainer, M.: A volatility basis set model for summertime secondary organic aerosols over the eastern United States in 2006, https://doi.org/10.1029/2011JD016831, 2012.

Anav, A., Sorrentino, B., Collalti, A., Paoletti, E., Sicard, P., Coulibaly, F., Manzini, J., Hoshika, Y., and De Marco, A.: Meteorological, chemical and biological evaluation of the coupled chemistry-climate WRF-Chem model from regional to urban scale. An impact-oriented application for human health, Environmental Research, 257, 119401, https://doi.org/10.1016/j.envres.2024.119401, 2024.

Anenberg, S. C., Mohegh, A., Goldberg, D. L., Kerr, G. H., Brauer, M., Burkart, K., Hystad, P., Larkin, A., Wozniak, S., and Lamsal, L.: Long-term trends in urban NO2 concentrations and associated paediatric asthma incidence: estimates from global datasets, The Lancet Planetary Health, 6, e49–e58, https://doi.org/10.1016/S2542-5196(21)00255-2, 2022.

ASIA-AQ White Paper | ASIA-AQ: https://espo.nasa.gov/asia-aq/document/ASIA-AQ_White_Paper, last access: 6 January 2025.

minimize(method='SLSQP') — SciPy v1.16.2 Manual: https://docs.scipy.org/doc/scipy/reference/optimize.minimize-slsqp.html, last access: 4 November 2025.

Aung, S. H., Gheewala, S. H., Winijkul, E., Panyametheekul, S., and Prapaspongsa, T.: Environmental impacts and costs of ozone formation in Bangkok Metropolitan Region, Atmospheric Pollution Research, 16, 102450, https://doi.org/10.1016/j.apr.2025.102450, 2025.

Bae, K., Richter, A., Lange, K., Friedrich, M. M., Pinardi, G., Roozendael, M. V., Merlaud, A., Fayt, C., Bösch, T., Zilker, B., Latsch, M., Behrens, L. K., Lee, H., Jung, Y., Hong, H., Chang, L.-S., and Song, C.-K.: Comparison of GEMS v3.0 tropospheric NO2 columns with ground-based DOAS instruments in Ulsan, GIScience & Remote Sensing, 2025.

Beirle, S., Boersma, K. F., Platt, U., Lawrence, M. G., and Wagner, T.: Megacity Emissions and Lifetimes of Nitrogen Oxides Probed from Space, Science, 333, 1737–1739, https://doi.org/10.1126/science.1207824, 2011.

Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, https://doi.org/10.1029/2003JD003697, 2004.

Bond, T. C., Bhardwaj, E., Dong, R., Jogani, R., Jung, S., Roden, C., Streets, D. G., and Trautmann, N. M.: Historical emissions of black and organic carbon aerosol from energy-related combustion, 1850–2000, Global Biogeochemical Cycles, 21, https://doi.org/10.1029/2006GB002840, 2007.

Chen, F. and Chen, Z.: Cost of economic growth: Air pollution and health expenditure, Science of The Total Environment, 755, 142543, https://doi.org/10.1016/j.scitotenv.2020.142543, 2021.

Cho, C., Franchin, A., Flocke, F., Lesko, K., Owen, C., Hall, S. R., Ullmann, K., Apel, E. C., Hills, A. J., Hornbrook, R. S., Roozitalab, B., Jeong, D., Diskin, G. S., Choi, Y., DiGangi, J. P., Miech, J., Wolfe, G. M., Hanisco, T. F., St. Clair, J. M., Liao, J., Delaria, E. R., Sebol, A., Hannun, R. A., Wennberg, P. O., Ball, K., Lee, Y. R., Huey, L. G., Tanner, D. J., Arterburn, L., Blake, D. R., Blake, N. J., Barletta, B., Meinardi, S., Min, K.-E., Kang, H., Nam, W., Wisthaler, A., Piel, F., Wojnowski, W., Dibb, J., and Crawford, J.: Insights on Ozone Formation Sensitivity in Southeast and East Asian Megacities during ASIA-AQ, EGUsphere, 1–29, https://doi.org/10.5194/egusphere-2025-6434, 2026.

Choi, S., Lamsal, L. N., Follette-Cook, M., Joiner, J., Krotkov, N. A., Swartz, W. H., Pickering, K. E., Loughner, C. P., Appel, W., Pfister, G., Saide, P. E., Cohen, R. C., Weinheimer, A. J., and Herman, J. R.: Assessment of NO_2 observations during DISCOVER-AQ and KORUS-AQ field campaigns, Atmospheric Measurement Techniques, 13, 2523–2546, https://doi.org/10.5194/amt-13-2523-2020, 2020.

Christopoulos, J. A., Saide, P. E., Ferrare, R., Collister, B., Barton-Grimley, R. A., Scarino, A. J., Collins, J., Hair, J. W., and Nehrir, A.: Improving Planetary Boundary Layer Height Estimation From Airborne Lidar Instruments, Journal of Geophysical Research: Atmospheres, 130, e2024JD042538, https://doi.org/10.1029/2024JD042538, 2025.

Christopoulos, J.: GEMS Top-Down NOx Emissions and Model Optimization, Zenodo, https://doi.org/10.5281/zenodo.18381169, 2026.

Commerce, N. C. for E. P. W. S. S. D. of: NCEP FNL Operational Model Global Tropospheric Analyses, continuing from July 1999, https://doi.org/10.5065/D6M043C6, 2000.

Crippa, M., Solazzo, E., Huang, G., Guizzardi, D., Koffi, E., Muntean, M., Schieberle, C., Friedrich, R., and Janssens-Maenhout, G.: High resolution temporal profiles in the Emissions Database for Global Atmospheric Research, Sci Data, 7, 121, https://doi.org/10.1038/s41597-020-0462-2, 2020.

Elguindi, N., Granier, C., Stavrakou, T., Darras, S., Bauwens, M., Cao, H., Chen, C., Denier van der Gon, H. a. C., Dubovik, O., Fu, T. M., Henze, D. K., Jiang, Z., Keita, S., Kuenen, J. J. P., Kurokawa, J., Liousse, C., Miyazaki, K., Müller, J.-F., Qu, Z., Solmon, F., and Zheng, B.: Intercomparison of Magnitudes and Trends in Anthropogenic Surface Emissions From Bottom-

Up Inventories, Top-Down Estimates, and Emission Scenarios, Earth's Future, 8, e2020EF001520, https://doi.org/10.1029/2020EF001520, 2020.

de Foy, B. and Schauer, J. J.: An improved understanding of NOx emissions in South Asian megacities using TROPOMI NO2 retrievals, Environ. Res. Lett., 17, 024006, https://doi.org/10.1088/1748-9326/ac48b4, 2022.

Fuller, R., Landrigan, P. J., Balakrishnan, K., Bathan, G., Bose-O'Reilly, S., Brauer, M., Caravanos, J., Chiles, T., Cohen, A., Corra, L., Cropper, M., Ferraro, G., Hanna, J., Hanrahan, D., Hu, H., Hunter, D., Janata, G., Kupka, R., Lanphear, B., Lichtveld, M., Martin, K., Mustapha, A., Sanchez-Triana, E., Sandilya, K., Schaefli, L., Shaw, J., Seddon, J., Suk, W., Téllez-Rojo, M. M., and Yan, C.: Pollution and health: a progress update, The Lancet Planetary Health, 6, e535–e547, https://doi.org/10.1016/S2542-5196(22)00090-0, 2022.

Gao, Z. and Zhou, X.: A review of the CAMx, CMAQ, WRF-Chem and NAQPMS models: Application, evaluation and uncertainty factors, Environmental Pollution, 343, 123183, https://doi.org/10.1016/j.envpol.2023.123183, 2024.

Georgoulias, A. K., van der A, R. J., Stammes, P., Boersma, K. F., and Eskes, H. J.: Trends and trend reversal detection in 2 decades of tropospheric NO_2 satellite observations, Atmospheric Chemistry and Physics, 19, 6269–6294, https://doi.org/10.5194/acp-19-6269-2019, 2019.

Goldberg, D. L., Lamsal, L. N., Loughner, C. P., Swartz, W. H., Lu, Z., and Streets, D. G.: A high-resolution and observationally constrained OMI NO_2 satellite retrieval, Atmospheric Chemistry and Physics, 17, 11403–11421, https://doi.org/10.5194/acp-17-11403-2017, 2017.

Goldberg, D. L., Saide, P. E., Lamsal, L. N., de Foy, B., Lu, Z., Woo, J.-H., Kim, Y., Kim, J., Gao, M., Carmichael, G., and Streets, D. G.: A top-down assessment using OMI NO_2 suggests an underestimate in the NO_x emissions inventory in Seoul, South Korea, during KORUS-AQ, Atmospheric Chemistry and Physics, 19, 1801–1818, https://doi.org/10.5194/acp-19-1801-2019, 2019.

Goldberg, D. L., Tao, M., Kerr, G. H., Ma, S., Tong, D. Q., Fiore, A. M., Dickens, A. F., Adelman, Z. E., and Anenberg, S. C.: Evaluating the spatial patterns of U.S. urban NOx emissions using TROPOMI NO2, Remote Sensing of Environment, 300, 113917, https://doi.org/10.1016/j.rse.2023.113917, 2024.

Gong, S. L., Barrie, L. A., and Blanchet, J.-P.: Modeling sea-salt aerosols in the atmosphere: 1. Model development, Journal of Geophysical Research: Atmospheres, 102, 3805–3818, https://doi.org/10.1029/96JD02953, 1997.

Graziosi, F. and Manca, G.: Quantification of Hotspot Methane Emissions Using Sentinel-5P TROPOMI Observations, JRC Publications Repository, https://doi.org/10.2760/4473858, 2025.

Grell, G., Freitas, S. R., Stuefer, M., and Fast, J.: Inclusion of biomass burning in WRF-Chem: impact of wildfires on weather forecasts, Atmospheric Chemistry and Physics, 11, 5289–5303, https://doi.org/10.5194/acp-11-5289-2011, 2011.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), Atmospheric Chemistry and Physics, 6, 3181–3210, https://doi.org/10.5194/acp-6-3181-2006, 2006.

Guizzardi, D., Crippa, M., Butler, T., Keating, T., Wu, R., Kaminski, J., Kuenen, J., Kurokawa, J., Chatani, S., Morikawa, T., Pouliot, G., Racine, J., Moran, M. D., Klimont, Z., Manseau, P. M., Mashayekhi, R., Henderson, B. H., Smith, S. J., Hoesly, R., Muntean, M., Banja, M., Schaaf, E., Pagani, F., Woo, J.-H., Kim, J., Pisoni, E., Zhang, J., Niemi, D., Sassi, M., Duhamel, A., Ansari, T., Foley, K., Geng, G., Chen, Y., and Zhang, Q.: The HTAP_v3.2 emission mosaic: merging regional and global monthly emissions (2000–2020) to support air quality modelling and policies, Earth System Science Data, 17, 5915–5950, https://doi.org/10.5194/essd-17-5915-2025, 2025.

Gulde, S. T., Kolm, M. G., Smith, D. J., Maurer, R., Courrèges-Lacoste, G. B., Sallusti, M., and Bagnasco, G.: Sentinel 4: a geostationary imaging UVN spectrometer for air quality monitoring: status of design, performance and development, in: International Conference on Space Optics — ICSO 2014, International Conference on Space Optics — ICSO 2014, 1158–1166, https://doi.org/10.1117/12.2304099, 2017.

Hakkarainen, J., Nurmela, J., and Lindqvist, H.: D4.4 Benchmarking of plume detection and quantification methods, 2023.

Hakkarainen, J., Kuhlmann, G., Koene, E., Santaren, D., Meier, S., Krol, M. C., van Stratum, B. J. H., Ialongo, I., Chevallier, F., Tamminen, J., Brunner, D., and Broquet, G.: Analyzing nitrogen dioxide to nitrogen oxide scaling factors for data-driven satellite-based emission estimation methods: A case study of Matimba/Medupi power stations in South Africa, Atmospheric Pollution Research, 15, 102171, https://doi.org/10.1016/j.apr.2024.102171, 2024.

Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O'Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), Geoscientific Model Development, 11, 369–408, https://doi.org/10.5194/gmd-11-369-2018, 2018.

Hsu, C.-H., Henze, D. K., Mizzi, A. P., Harkins, C., Lyu, C., Cooper, O. R., Schwantes, R. H., He, J., Li, M., Wang, S., Stockwell, C. E., Warneke, C., Rollins, A. W., Waxman, E. M., Zuraski, K., Peischl, J., Kondragunta, S., Li, F., Xu, C., Pierce, R. B., Abad, G. G., Nowlan, C. R., Liu, X., and McDonald, B. C.: Top-Down Estimates of U.S. NOx Emissions Using TEMPO and TROPOMI NO2 Remote Sensing Observations With WRF-Chem/Chem-DART, Journal of Geophysical Research: Atmospheres, 131, e2025JD044223, https://doi.org/10.1029/2025JD044223, 2026.

Inness, A., Ades, M., Agustí-Panareda, A., Barré, J., Benedictow, A., Blechschmidt, A.-M., Dominguez, J. J., Engelen, R., Eskes, H., Flemming, J., Huijnen, V., Jones, L., Kipling, Z., Massart, S., Parrington, M., Peuch, V.-H., Razinger, M., Remy, S., Schulz, M., and Suttie, M.: The CAMS reanalysis of atmospheric composition, Atmospheric Chemistry and Physics, 19, 3515–3556, https://doi.org/10.5194/acp-19-3515-2019, 2019.

Janz, S. J., Kowalewski, M., Lamsal, L., Nowlan, C., and Judd, L.: Airborne hyperspectral trace gas sensors as testbeds for geostationary air quality missions, in: Sensors, Systems, and Next-Generation Satellites XXIII, Sensors, Systems, and Next-Generation Satellites XXIII, 509–518, https://doi.org/10.1117/12.2533765, 2019.

Jiawei Zhuang, raphael dussin, David Huard, Pascal Bourgault, Anderson Banihirwe, Stephane Raynaud, Brewster Malevich, Martin Schupfner, Filipe, Charles Gauthier, Sam Levang, André Jüling, Mattia Almansi, RichardScottOZ, RondeauG, Stephan Rasp, Trevor James Smith, Ben Mares, Jemma Stachelek, Matthew Plough, Pierre, Ray Bell, Romain Caneill, and Xianxiang Li: pangeo-data/xESMF: v0.8.10, , https://doi.org/10.5281/ZENODO.4294774, 2025.

Judd, L. M., Al-Saadi, J. A., Szykman, J. J., Valin, L. C., Janz, S. J., Kowalewski, M. G., Eskes, H. J., Veefkind, J. P., Cede, A., Mueller, M., Gebetsberger, M., Swap, R., Pierce, R. B., Nowlan, C. R., Abad, G. G., Nehrir, A., and Williams, D.: Evaluating Sentinel-5P TROPOMI tropospheric NO_2 column densities with airborne and Pandora spectrometers near New York City and Long Island Sound, Atmospheric Measurement Techniques, 13, 6113–6140, https://doi.org/10.5194/amt-13-6113-2020, 2020.

Jung, Y., Park, J., Hong, H., Lee, H., and Jeong, U.: Evaluation of GEMS NO Retrieval Algorithm Version 2.0 and 3.0 Using TROPOMI and Pandora Observations, Korean Journal of Remote Sensing, 41, 803–811, https://doi.org/10.7780/kjrs.2025.41.5.8, 2025.

Kim, J., Jeong, U., Ahn, M.-H., Kim, J. H., Park, R. J., Lee, H., Song, C. H., Choi, Y.-S., Lee, K.-H., Yoo, J.-M., Jeong, M.-J., Park, S. K., Lee, K.-M., Song, C.-K., Kim, S.-W., Kim, Y. J., Kim, S.-W., Kim, M., Go, S., Liu, X., Chance, K., Miller, C. C., Al-Saadi, J., Veihelmann, B., Bhartia, P. K., Torres, O., Abad, G. G., Haffner, D. P., Ko, D. H., Lee, S. H., Woo, J.-H., Chong, H., Park, S. S., Nicks, D., Choi, W. J., Moon, K.-J., Cho, A., Yoon, J., Kim, S., Hong, H., Lee, K., Lee, H., Lee, S., Choi, M., Veefkind, P., Levelt, P. F., Edwards, D. P., Kang, M., Eo, M., Bak, J., Baek, K., Kwon, H.-A., Yang, J., Park, J., Han, K. M., Kim, B.-R., Shin, H.-W., Choi, H., Lee, E., Chong, J., Cha, Y., Koo, J.-H., Irie, H., Hayashida, S., Kasai, Y.,

Kanaya, Y., Liu, C., Lin, J., Crawford, J. H., Carmichael, G. R., Newchurch, M. J., Lefer, B. L., Herman, J. R., Swap, R. J., Lau, A. K. H., Kurosu, T. P., Jaross, G., Ahlers, B., Dobber, M., McElroy, C. T., and Choi, Y.: New Era of Air Quality Monitoring from Space: Geostationary Environment Monitoring Spectrometer (GEMS), https://doi.org/10.1175/BAMS-D-18-0013.1, 2020.

Koster, R. D., Darmenov, A. S., and da Silva, A. M.: The Quick Fire Emissions Dataset (QFED): Documentation of Versions 2.1, 2.2 and 2.4: Technical Report Series on Global Modeling and Data Assimilation - Volume 38, 2015.

Kowalewski, M. G. and Janz, S. J.: Remote sensing capabilities of the GeoCAPE Airborne Simulator, in: Earth Observing Systems XIX, Earth Observing Systems XIX, 496–507, https://doi.org/10.1117/12.2062058, 2014.

Kuhlmann, G., Koene, E., Meier, S., Santaren, D., Broquet, G., Chevallier, F., Hakkarainen, J., Nurmela, J., Amorós, L., Tamminen, J., and Brunner, D.: The *ddeq* Python library for point source quantification from remote sensing images (version 1.0), Geoscientific Model Development, 17, 4773–4789, https://doi.org/10.5194/gmd-17-4773-2024, 2024.

Kumar, R., Naja, M., Pfister, G. G., Barth, M. C., Wiedinmyer, C., and Brasseur, G. P.: Simulations over South Asia using the Weather Research and Forecasting model with Chemistry (WRF-Chem): chemistry evaluation and initial results, Geoscientific Model Development, 5, 619–648, https://doi.org/10.5194/gmd-5-619-2012, 2012.

Kurokawa, J. and Ohara, T.: Long-term historical trends in air pollutant emissions in Asia: Regional Emission inventory in ASia (REAS) version 3, Atmospheric Chemistry and Physics, 20, 12761–12793, https://doi.org/10.5194/acp-20-12761-2020, 2020.

Lee, S., Bae, K., Janz, S. J., Judd, L. M., Xiong, S., Boehmler, J., Jung, Y., Lee, H.-J., Hong, H., Chang, L.-S., Kang, M., Ahn, M.-H., Song, C.-K., and Park, S. S.: Sensitivity analysis of NO2 differential slant column density according to spatial resolution using GCAS data from the SIJAQ 2022 campaign, Atmospheric Environment, 335, 120723, https://doi.org/10.1016/j.atmosenv.2024.120723, 2024.

Lennartson, E. M., Wang, J., Gu, J., Castro Garcia, L., Ge, C., Gao, M., Choi, M., Saide, P. E., Carmichael, G. R., Kim, J., and Janz, S. J.: Diurnal variation of aerosol optical depth and $PM_{2.5}$ in South Korea: a synthesis from AERONET, satellite (GOCI), KORUS-AQ observation, and the WRF-Chem model, Atmospheric Chemistry and Physics, 18, 15125–15144, https://doi.org/10.5194/acp-18-15125-2018, 2018.

Li, M., Kurokawa, J., Zhang, Q., Woo, J.-H., Morikawa, T., Chatani, S., Lu, Z., Song, Y., Geng, G., Hu, H., Kim, J., Cooper, O. R., and McDonald, B. C.: MIXv2: a long-term mosaic emission inventory for Asia (2010–2017), Atmospheric Chemistry and Physics, 24, 3925–3952, https://doi.org/10.5194/acp-24-3925-2024, 2024.

Liu, J., Gao, H., Jia, R., Wang, R., Han, D., Liu, L., Xu, X., and Qiao, Z.: A downscaling framework with WRF-UCM and LES/RANS models for urban microclimate simulation strategy: Validation through both measurement and mechanism model, Building and Environment, 269, 112361, https://doi.org/10.1016/j.buildenv.2024.112361, 2025.

Makkwao, K. and Prueksasit, T.: PM10 Concentration Emitted from Blasting and Crushing Processes of Limestone Mines in Saraburi Province, Thailand, 2021.

Meier, S., Koene, E. F. M., Krol, M., Brunner, D., Damm, A., and Kuhlmann, G.: A lightweight NO_2-to-NO_x conversion model for quantifying NO_x emissions of point sources from NO_2 satellite observations, Atmospheric Chemistry and Physics, 24, 7667–7686, https://doi.org/10.5194/acp-24-7667-2024, 2024.

Miyazaki, K., Eskes, H., Sudo, K., Boersma, K. F., Bowman, K., and Kanaya, Y.: Decadal changes in global surface NO_x emissions from multi-constituent satellite data assimilation, Atmospheric Chemistry and Physics, 17, 807–837, https://doi.org/10.5194/acp-17-807-2017, 2017.

Miyazaki, K., Sekiya, T., Fu, D., Bowman, K. W., Kulawik, S. S., Sudo, K., Walker, T., Kanaya, Y., Takigawa, M., Ogochi, K., Eskes, H., Boersma, K. F., Thompson, A. M., Gaubert, B., Barre, J., and Emmons, L. K.: Balance of Emission and Dynamical Controls on Ozone During the Korea-United States Air Quality Campaign From Multiconstituent Satellite Data Assimilation, Journal of Geophysical Research: Atmospheres, 124, 387–413, https://doi.org/10.1029/2018JD028912, 2019.

Miyazaki, K., Bowman, K., Sekiya, T., Eskes, H., Boersma, F., Worden, H., Livesey, N., Payne, V. H., Sudo, K., Kanaya, Y., Takigawa, M., and Ogochi, K.: Updated tropospheric chemistry reanalysis and emission estimates, TCR-2, for 2005–2018, Earth System Science Data, 12, 2223–2259, https://doi.org/10.5194/essd-12-2223-2020, 2020.

Mues, A., Kuenen, J., Hendriks, C., Manders, A., Segers, A., Scholz, Y., Hueglin, C., Builtjes, P., and Schaap, M.: Sensitivity of air pollution simulations with LOTOS-EUROS to the temporal distribution of anthropogenic emissions, Atmospheric Chemistry and Physics, 14, 939–955, https://doi.org/10.5194/acp-14-939-2014, 2014.

Nocedal, J. and Wright, S. J.: Numerical optimization, 2nd ed., Springer, New York, 664 pp., 2006.

Oda, T., Maksyutov, S., and Andres, R. J.: The Open-source Data Inventory for Anthropogenic CO_2 , version 2016 (ODIAC2016): a global monthly fossil fuel CO_2 gridded emissions data product for tracer transport simulations and surface flux inversions, Earth Syst. Sci. Data, 10, 87–107, https://doi.org/10.5194/essd-10-87-2018, 2018.

Park, J., Choi, Y., Jung, J., Lee, K., and Yeganeh, A. K.: First top-down diurnal adjustment to NOx emissions inventory in Asia informed by the Geostationary Environment Monitoring Spectrometer (GEMS) tropospheric NO2 columns, Sci Rep, 14, 24338, https://doi.org/10.1038/s41598-024-76223-1, 2024.

Park, J., Hong, H., Lee, H., Kim, S.-W., Kim, J., Van Roozendael, M., Fayt, C., Ahn, M.-H., Jacob, D. J., Seo, S., Kim, K.-M., Kim, D., Choi, W., Lee, W.-J., Lee, D.-W., Wagner, T., Richter, A., Krotkov, N. A., Lamsal, L. N., Ko, D. H., Lee, S. H., and Woo, J.-H.: Tropospheric nitrogen dioxide levels vary diurnally in Asian cities, Commun Earth Environ, 6, 389, https://doi.org/10.1038/s43247-025-02272-7, 2025.

Park, R. J., Oak, Y. J., Emmons, L. K., Kim, C.-H., Pfister, G. G., Carmichael, G. R., Saide, P. E., Cho, S.-Y., Kim, S., Woo, J.-H., Crawford, J. H., Gaubert, B., Lee, H.-J., Park, S.-Y., Jo, Y.-J., Gao, M., Tang, B., Stanier, C. O., Shin, S. S., Park, H. Y., Bae, C., and Kim, E.: Multi-model intercomparisons of air quality simulations for the KORUS-AQ campaign, Elementa: Science of the Anthropocene, 9, 00139, https://doi.org/10.1525/elementa.2021.00139, 2021.

Pörtner, H.-O., Roberts, D. C., Tignor, M. M. B., Poloczanska, E. S., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A., and Rama, B. (Eds.): Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change., 2022.

Rey-Pommier, A., Héraud, A., Chevallier, F., Ciais, P., Christoudias, T., Kushta, J., and Sciare, J.: Global gridded NO_x emissions using TROPOMI observations, Earth System Science Data, 17, 3329–3351, https://doi.org/10.5194/essd-17-3329-2025, 2025.

Ridley, B. A. and Grahek, F. E.: A Small, Low Flow, High Sensitivity Reaction Vessel for NO Chemiluminescence Detectors, 1990.

Ridley, B. A., Grahek, F. E., and Walega, J. G.: A Small High-Sensitivity, Medium-Response Ozone Detector Suitable for Measurements from Light Aircraft, 1992.

Saide, P. E., Gao, M., Lu, Z., Goldberg, D. L., Streets, D. G., Woo, J.-H., Beyersdorf, A., Corr, C. A., Thornhill, K. L., Anderson, B., Hair, J. W., Nehrir, A. R., Diskin, G. S., Jimenez, J. L., Nault, B. A., Campuzano-Jost, P., Dibb, J., Heim, E., Lamb, K. D., Schwarz, J. P., Perring, A. E., Kim, J., Choi, M., Holben, B., Pfister, G., Hodzic, A., Carmichael, G. R., Emmons, L., and Crawford, J. H.: Understanding and improving model representation of aerosol optical properties for a Chinese haze event measured during KORUS-AQ, Atmospheric Chemistry and Physics, 20, 6455–6478, https://doi.org/10.5194/acp-20-6455-2020, 2020.

Santaren, D., Hakkarainen, J., Kuhlmann, G., Koene, E., Chevallier, F., Ialongo, I., Lindqvist, H., Nurmela, J., Tamminen, J., Amorós, L., Brunner, D., and Broquet, G.: Benchmarking data-driven inversion methods for the estimation of local CO_2 emissions from synthetic satellite images of XCO_2 and NO_2, Atmospheric Measurement Techniques, 18, 211–239, https://doi.org/10.5194/amt-18-211-2025, 2025.

Seinfeld and Pandis: Atmospheric Chemistry and Physics: From Air Pollution to Climate Change, 3rd Edition | Wiley, 2016.

Shetty, S. S., D, D., S, H., Sonkusare, S., Naik, P. B., N, S. K., and Madhyastha, H.: Environmental pollutants and their effects on human health, Heliyon, 9, https://doi.org/10.1016/j.heliyon.2023.e19496, 2023.

Sicard, P., Agathokleous, E., Anenberg, S. C., De Marco, A., Paoletti, E., and Calatayud, V.: Trends in urban air pollution over the last two decades: A global perspective, Science of The Total Environment, 858, 160064, https://doi.org/10.1016/j.scitotenv.2022.160064, 2023.

Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., and Huang, X.-Y.: A Description of the Advanced Research WRF Model Version 4, https://doi.org/10.5065/1DFH-6P97, 2019.

Smith, S. J., van Aardenne, J., Klimont, Z., Andres, R. J., Volke, A., and Delgado Arias, S.: Anthropogenic sulfur dioxide emissions: 1850–2005, Atmospheric Chemistry and Physics, 11, 1101–1116, https://doi.org/10.5194/acp-11-1101-2011, 2011.

St. Clair, J. M., Swanson, A. K., Bailey, S. A., and Hanisco, T. F.: CAFE: a new, improved nonresonant laser-induced fluorescence instrument for airborne in situ measurement of formaldehyde, Atmospheric Measurement Techniques, 12, 4581–4590, https://doi.org/10.5194/amt-12-4581-2019, 2019.

Thailand Office of the National Economic and Social Development Board, World Bank: Industrial Change in the Bangkok Urban Region, 2017.

Tuccella, P., Curci, G., Grell, G. A., Visconti, G., Crumeyrolle, S., Schwarzenboeck, A., and Mensah, A. A.: A new chemistry option in WRF-Chem v. 3.4 for the simulation of direct and indirect aerosol effects using VBS: evaluation against IMPACT-EUCAARI data, Geoscientific Model Development, 8, 2749–2776, https://doi.org/10.5194/gmd-8-2749-2015, 2015.

Uttamang, P., Aneja, V. P., and Hanna, A. F.: Assessment of gaseous criteria pollutants in the Bangkok Metropolitan Region, Thailand, Atmospheric Chemistry and Physics, 18, 12581–12593, https://doi.org/10.5194/acp-18-12581-2018, 2018.

Uttamang, P., Campbell, P. C., Aneja, V. P., and Hanna, A. F.: A multi-scale model analysis of ozone formation in the Bangkok Metropolitan Region, Thailand, Atmospheric Environment, 229, 117433, https://doi.org/10.1016/j.atmosenv.2020.117433, 2020.

Uttamang, P., Choomanee, P., Phupijit, J., Bualert, S., and Thongyen, T.: Investigation of Secondary Organic Aerosol Formation during O3 and PM2.5 Episodes in Bangkok, Thailand, Atmosphere, 14, 994, https://doi.org/10.3390/atmos14060994, 2023.

Valin, L. C., Russell, A. R., Hudman, R. C., and Cohen, R. C.: Effects of model resolution on the interpretation of satellite NO_2 observations, Atmospheric Chemistry and Physics, 11, 11647–11655, https://doi.org/10.5194/acp-11-11647-2011, 2011.

Verreyken, B. W. D., Harkins, C., Li, M., Angevine, W., Stockwell, C. E., Xu, L., Coggon, M., Gilman, J., Warneke, C., Strobach, E., Brown, S., McCarty, B., Marchbanks, R., Baidar, S., Brewer, A., Pfannerstill, E. Y., Arata, C., Goldstein, A. H., Brioude, J., and McDonald, B. C.: Top-Down Evaluation of Volatile Chemical Product Emissions Using a Lagrangian Framework, Environ. Sci. Technol., 59, 7211–7221, https://doi.org/10.1021/acs.est.4c10117, 2025.

Ye, J., Liu, L., Wang, Q., Hu, S., and Li, S.: A Novel Machine Learning Algorithm for Planetary Boundary Layer Height Estimation Using AERI Measurement Data, IEEE Geoscience and Remote Sensing Letters, 19, https://doi.org/10.1109/lgrs.2021.3073048, 2021.

Yu, K. A., Li, M., Harkins, C., He, J., Zhu, Q., Verreyken, B., Schwantes, R. H., Cohen, R. C., McDonald, B. C., and Harley, R. A.: Improved Spatial Resolution in Modeling of Nitrogen Oxide Concentrations in the Los Angeles Basin, Environ. Sci. Technol., 57, 20689–20698, https://doi.org/10.1021/acs.est.3c06158, 2023.

Zhao, C., Liu, X., Leung, L. R., Johnson, B., McFarlane, S. A., Gustafson, W. I. J., Fast, J. D., and Easter, R.: The spatial distribution of mineral dust and its shortwave radiative forcing over North Africa: modeling sensitivities to dust emissions and aerosol size treatments, Atmospheric Chemistry and Physics, 10, 8821–8838, https://doi.org/10.5194/acp-10-8821-2010, 2010.

Zoogman, P., Liu, X., Suleiman, R. M., Pennington, W. F., Flittner, D. E., Al-Saadi, J. A., Hilton, B. B., Nicks, D. K., Newchurch, M. J., Carr, J. L., Janz, S. J., Andraschko, M. R., Arola, A., Baker, B. D., Canova, B. P., Chan Miller, C., Cohen, R. C., Davis, J. E., Dussault, M. E., Edwards, D. P., Fishman, J., Ghulam, A., González Abad, G., Grutter, M., Herman, J. R., Houck, J., Jacob, D. J., Joiner, J., Kerridge, B. J., Kim, J., Krotkov, N. A., Lamsal, L., Li, C., Lindfors, A., Martin, R. V., McElroy, C. T., McLinden, C., Natraj, V., Neil, D. O., Nowlan, C. R., O'Sullivan, E. J., Palmer, P. I., Pierce, R. B., Pippin, M. R., Saiz-Lopez, A., Spurr, R. J. D., Szykman, J. J., Torres, O., Veefkind, J. P., Veihelmann, B., Wang, H., Wang, J., and Chance, K.: Tropospheric emissions: Monitoring of pollution (TEMPO), Journal of Quantitative Spectroscopy and Radiative Transfer, 186, 17–39, https://doi.org/10.1016/j.jqsrt.2016.05.008, 2017.

1440

1441

1442

1443

1444

Supplementary Information of "NO_x Emissions Constraints from GEMS NO_2 Retrievals: Inversion Methodology and Air Quality Model Evaluation in Bangkok using ASIA-AQ Multi-Platform Observations"

Christopoulos et al.

Correspondence to: Julianna Christopoulos (juliechristo@g.ucla.edu)

S1 Supplementary Methods

S1.1 Gaussian Plume (GP) Inversion Method

We apply a Gaussian plume model method to estimate NO_x emissions over Bangkok as outlined in Kuhlmann et al., 2024. This approach fits a vertically integrated Gaussian plume model to GEMS-observed NO_2 column densities, accounting for wind advection, plume dispersion, and chemical loss. The Gaussian plume model for a point source is expressed as:

$$G(x,y) = \frac{QH(x)}{2\sqrt{\pi}u\sigma(x)} exp\left(-\frac{y^2}{2\sigma(x)^2}\right) + V_{bg}(x,y) \quad (1)$$

$$\sigma(x) = \sqrt{\frac{2Kx^{\kappa}}{u}} \quad (2)$$

Where Q is the emission rate, u is the wind speed, $H(x)$ is the Heaviside function, $\sigma(x)$ is the plume width across wind, and V_{bg} is the background column. K is the eddy diffusivity coefficient ($m^2\ s^{-1}$) and κ accounts for nonlinear spreading depending on the meteorological conditions. To obtain values for Q, K, V_{bg}, and κ, a least-squares method is used to minimize the cost function:

$$J(Q,K,V_{bg},\kappa) = ||V_{i,j} - G(x_i,y_i)||_2^2 \quad (3)$$

Where $V_{i,j}$ is the observed NO_2 column density for a pixel with center (x_i,y_i). Given NO_2's short lifetime, it is necessary to multiply the Gaussian plume model by the decay term:

$$D(x,\tau) = H(x)exp\left(-\frac{x}{u\tau}\right) \quad (4)$$

Where the lifetime, τ, is another fitting parameter. Since we are estimating emissions for a city source (Bangkok), the Gaussian model used in this context, accounts for fluxes increasing slowly across the source area. The city emissions are thus described by an emission map, p(x, y) and a decay term:

$$G_a(x,y) = G(x,y)\int_{-\infty}^{\infty} D(x',\tau)p(x-x',y)dx' \quad (5)$$

The emission map can represent a two-dimensional Gaussian surface $p(x,y)$:

$$p(x,y) = \frac{1}{2\pi\sigma_x\sigma_y\sqrt{1-r^2}} \cdot exp\left(-\frac{(x-x_0)^2}{2\sigma^2{}_x(1-r^2)} - \frac{(y-y_0)^2}{2\sigma^2{}_y(1-r^2)} + \frac{r(x-x_0)(y-y_0)}{\sigma_x\sigma_y(1-r^2)}\right) \quad (6)$$

S1.2 Integrated Mass Enhancement (IME) Inversion Method

Lastly, we estimate NO_2 emissions with the Integrated Mass Enhancement (IME) method, which relates the total mass of NO_2 observed within the detected plume to the underlying emission rate (Kuhlmann et al., 2024). This method is derived by integrating the Gaussian Plume model, $G(x, y)$, once the background, $V_{bg}(x, y)$ is subtracted over a singular polygon:

$$M = \int_{y_1}^{y_2} \int_{x_1}^{x_2} G(x, y) - V_{bg}(x, y) dx\, dy \quad (7)$$

Where:

$$M = \int_{x_1}^{x_2} \frac{Q}{u} dx \quad (8)$$

$$Q = \frac{u}{L} M \quad (9)$$

Here, u is the effective wind speed, $L = x_2 - x_1$ is the plume length, Q is the emission rate, and M is the integrated total mass enhancement of the plume. M in practice is computed as:

$$M = \sum_{(i,j) \in \mathcal{P}_a} (V_{i,j} - V_{bg}) \cdot A_{i,j} \quad (10)$$

Where $A_{i,j}$ is the pixel area and $\mathcal{P}_a$ is the integration area. For NO_2, a decaying gas, the decay term, $D(x)$, is included as well:

$$M = \int_{y_1}^{y_2} \int_{x_1}^{x_2} G(x, y) \cdot D(x) dx\, dy \quad (11)$$

The final emission rate, Q, is determined as:

$$Q = \frac{1}{c} \frac{u}{L} M \quad (12)$$

where c is a correction factor that corrects for decay along the plume direction:

$$c = \frac{u\tau}{L} \left(\exp\left(-\frac{x_1}{u\tau}\right) - \exp\left(-\frac{x_2}{u\tau}\right) \right) \quad (13)$$

S2 Supplementary Figures

Table S1. Summary of WRF-Chem D01 validation statistics for Base (B), ~~and~~ Updated (U), and bias-corrected (U+BC) simulations against independent observational datasets. Metrics include mean bias (MB), mean error (ME), normalized mean bias (NMB), normalized mean error (NME), root-mean-square error (RMSE), and Pearson correlation coefficient (CORR). Details of the evaluation methodology are provided in Sect. 6.

Dataset	Species	Model Run	MB	ME	NMB	NME	RMSE	CORR
GCAS	Tropospheric NO_2 column (molecules cm^{-2})	**B**	4.0E+15	5.5E+15	91.8	1.3E+2	8.9E+15	0.47
		U	-2.2E+14	2.1E+15	-5.0	49	3.1E+15	0.46
		U+BC	7.1 E+14	2.6 E+15	16	60.0	3.8 E+15	0.49
DC-8	NO_xO_3 NO_2 (ppbV)	**B**	0.27	0.53	17	33	1.2	0.94
		U	-1.0	1.0	-63	63	1.7	0.94
		U+BC	-0.77	0.77	-47	47	1.3	0.94
	CANOE NO_2 (ppbV)	**B**	0.3	0.57	20.0	36	1.3	0.94
		U	-1.0	1.0	-62	62	1.6	0.9
		U+BC	-0.73	0.74	-46	46	1.2	0.94
PCD Ground Monitors	NO_2 (ppbV)	**B**	6.9	8.2	73	87	11	0.45
		U	-4.6	4.9	-49	53	6.3	0.50
		U+BC	-2.4	3.9	-25	41	5.2	0.50
	NO_x (ppbV)	**B**	7.1	9.0	68	86	12	0.41
		U	-5.4	5.8	-51	55	7.8	0.57
		U+BC	-3.0	4.6	-28	44	6.7	0.53
	O_3 (ppbV)	**B**	5.2	11.4	17	37	16	0.86
		U	9.0	10.4	29	34	13	0.86

		U+BC	9.1	10.1	30.0	34	13	0.87
Pandora	Tropospheric NO_2 column (molecules cm^{-2})	**B**	4.5E+15	9.0E+15	31	62	1.1+16	-0.23
		U	-8.6E+15	8.6E+15	-59	59	1.1E+16	0.23
		U+BC	-6.4 E+15	6.5 E+15	-44	45	8.9 E+15	0.27

Table S2. Summary of WRF_{Base} wind validation statistics against surface observations (Thailand PCD) for 14 – 27 March 2024. Metrics include mean bias (MB), mean error (ME), root-mean-square error (RMSE), and Pearson correlation coefficient (CORR) included for wind speeds. ERA5 wind speeds align best with ground monitor observations in the BMR.

Dataset	Species	Model Run	MB	ME	RMSE	CORR
PCD Ground Monitors	Wind Speed ($m\ s^{-1}$)	WRF_{Base} D02	1.7	1.7	1.9	0.8
		WRF_{Base} D01	2.2	2.2	2.6	0.6
		ERA5	1.3	1.4	1.4	0.8
	Wind Direction (°)	WRF_{Base} D02	-13	19	24	-
		WRF_{Base} D01	7.2	23	32	-
		ERA5	-14	18	21	-

Table S3. Summary of WRF_{Base} PBLH statistics against airborne lidar observations (HSRL-2) for ASIA-AQ flight days, averaged over raster periods. Metrics include mean bias (MB), mean error (ME), normalized mean bias (NMB), normalized mean error (NME), root-mean-square error (RMSE), and Pearson correlation coefficient (CORR). Details of the evaluation methodology are provided in Sect. 6.

Date	MB	ME	NMB	NME	RMSE	CORR
18 March	190	420	28	62	590	0.53
19 March	-190	290	-31	48	340	0.27
21 March	330	510	43	67	650	0.31
23 March	150	410	21	56	480	0.54

25 March	190	450	27	63	550	0.72




Figure S15. (a) Comparison of Pandora (pink) and GEMS (blue) total column NO_2 for high-quality observations during the March 2024. The ASIA-AQ deployment and model simulation period is representative of 14 – 27 March 2024~~ASIA-AQ deployment period~~. GEMS reproduces the temporal variability observed by Pandora (CORR = 0.7) but systematically underestimates column magnitudes (mean bias ≈ -9.2×10^{15} molecules cm^{-2}; mean ratio ≈ 0.6), with the largest discrepancies occurring during the stagnation event on 21 March 2024. GEMS is generally a factor of two lower than Pandora during this period. ~~(b) Sensitivity of CSF-derived NO_x emissions to GEMS NO_2 bias correction over the BMR. The solid line shows the nominal inversion, while the solid line represents emissions derived after scaling GEMS NO_2 columns by a factor of 1.67.~~




Figure S2~~1~~. Comparison of (a) WRF_{Base} D02 (purple), (b) WRF_{Base} D01 (red), (c) ERA5 (pink) to Thailand Pollution Control Department (PCD) ground-monitor network (grey) wind speed (m s^{-1}), wind direction (from), and wind direction error during the ASIA-AQ deployment period (14–27 March 2024). Lines represent averages across stations within the BMR and surrounding regions. Both model domains and ERA5 overestimate wind speeds relative to surface observations, potentially due to unresolved urban canopy effects. These results suggest that improved urban parameterizations (e.g., a multi-layer urban canopy model) could enhance model performance. Error diagnostics also highlight misrepresentation of a short-lived northeasterly wind event on 20–21 March, leading to temporary errors in simulated transport direction.




Figure S3~~2~~. Example of the cross-sectional flux approach applied on 15 March 2024 using GEMS daytime measurements (00:45:00–06:45:00 UTC; 07:45:00–13:45:00 LT), generated with the *ddeq* Python package (Kuhlmann et al., 2024). Detected satellite pixels associated with the Bangkok urban plume are marked with blue ticks. A vector originating from Bangkok indicates the plume direction based on ERA5 wind fields at the source location. Khao Wong and Thap Kwang districts in Saraburi Province represent point sources distinct from the Bangkok urban plume, influenced by limestone quarry operations and heavy-truck transport (Makkhao and Prueksasit, 2023).




Figure S43. Examples of cross-sectional flux fits on 17 March 2024 for (a) 00:45:00 UTC (07:45:00 LT), (b) 01:45:00 UTC (08:45:00 LT), (c) 02:45:00 UTC (09:45:00 LT), (d) 03:45:00 UTC (10:45:00 LT), (e) 04:45:00 UTC (11:45:00 LT), (f) 05:45:00 UTC (12:45:00 LT), and (g) 06:45:00 UTC (13:45:00 LT), generated with the *ddeq* Python package (Kuhlmann et al., 2024). Derived NO_x fluxes vary throughout the daytime, with peak values in the morning hours and decreasing toward the afternoon, consistent with the daytime pattern of vehicular traffic emissions in the Bangkok Metropolitan Region (BMR).

(a)

(b)






Figure S54. (a) ERA5 (solid pink) and Thailand Pollution Control Department (PCD) ground-monitor network (grey) wind speed (m s^{-1}) during the ASIA-AQ deployment period (14–27 March 2024). ERA5 wind speeds scaled by a factor of 0.6 are shown as dashed pink lines. (b) GEMS-derived daytime NO_x emissions (kg s^{-1}) estimated using the cross-sectional flux (blue), Gaussian plume inversion (red), and integrated mass enhancement (green) methods. Emissions represent averages across daytime hours and weekdays during the ASIA-AQ deployment. As a sensitivity test, ERA5 wind speeds were scaled by a factor of 0.6 prior to inversion to assess wind speed uncertainty. Reduced wind speeds lead to lower inferred emissions (≈40%), consistent with enhanced NO_2 accumulation under weaker transport and a reduced emission flux required to reproduce the observed columns. These adjusted emissions fall below those inferred from other observational constraints, suggesting that further wind speed reductions may not be appropriate for representing PBL-averaged transport in the BMR.



~~**Figure S6.** Differences in gridded WRF-Chem input emissions (updated − base) expressed in mol km^{-2} h^{-1} over the Bangkok Metropolitan Region (BMR) for D01 at (a) 00:00:00 UTC, (b) 06:00:00 UTC, (c) 12:00:00 UTC, and (d) 18:00:00 UTC, and for D02 at (e) 00:00:00 UTC, (f) 06:00:00 UTC, (g) 12:00:00 UTC, and (h) 18:00:00 UTC. The largest differences occur during the afternoon local time (e.g., 06:00:00 UTC), reaching up to ~ 400 mol km^{-2} h^{-1} in D02.~~


Mar 18th, 2024
(a)
GEMS
(b)
WRF-GEMS Updated
(c)
WRF-GEMS Base
08:45 LT
10:45 LT
12:45 LT
09:00 LT
11:00 LT
13:00 LT
D01
3 x 10^15
4 x 10^15
5 x 10^15
6 x 10^15
7 x 10^15
8 x 10^15
1 x 10^16
1.5 x 10^16
2 x 10^16
Tropospheric Column NO2
[molecules/cm²]



Mar 18th,2024
(a) GEMS
(b) WRF-GEMS$_{Updated}$
(c) WRF-GEMS$_{Base}$
01:45 UTC
03:45 UTC
06:45 UTC
D01
02:00:00 UTC
04:00:00 UTC
06:00:00 UTC
40′
20′
14°N
13°N
45′
15′
30′
3 x 10^{15}
4 x 10^{15}
5 x 10^{15}
6 x 10^{15}
7 x 10^{15}
8 x 10^{15}
1 x 10^{16}
1.5 x 10^{16}
2 x 10^{16}
Tropospheric Column NO$_2$
[molecules/cm^2]

Figure S6~~7~~. Spatial comparison over the BMR on 18 March 2024 of tropospheric NO_2 columns from (a) GEMS, (b) WRF-$GEMS_{Updated}$ D02, and (c) WRF-$GEMS_{Base}$ D02 for snapshots at 02:00:00, 04:00:00, and 07:00:00 UTC, corresponding to approximately 09:00:00, 11:00:00, and 14:00:00 LT.




Figure S7~~8~~. Daytime tropospheric NO_2 column comparison between GEMS (blue), WRF-$GEMS_{Base}$ D01 (dotted red), and WRF-$GEMS_{Updated}$ D02 (solid red) during the ASIA-AQ deployment period. Shaded regions indicate variability (10th–90th percentiles).




Figure S89. Locations of Thailand Pollution Control Department (PCD) ground-monitor stations used for evaluation of (a) meteorological variables (wind speed and direction) and (b) trace gases (NO_2, NO_x, O_3). Black boxes indicate stations included in the analysis. For trace gas evaluation, stations within the Bangkok urban plume were selected to assess the region where the inversion was applied, whereas meteorological evaluation includes a broader regional domain to better capture spatial variability in winds.




Figure S9~~10~~. Comparison of observed wind roses (a) with modelled wind roses from (b) WRF_{Base} D02 and (c) WRF_{Base} D01 over the BMR for 18–22 March 2024. Panels show the frequency distribution of wind speed and direction. While the model captures the dominant southerly flow, wind speeds are consistently overpredicted in both domains. The largest discrepancies occur on 20–21 March, when the model fails to capture the observed shift toward northeasterly winds, likely contributing to transport and concentration errors during this period.





Figure S1011. Evaluation of observed lidar-derived (blue) and modelled WRF_{Base} D02 (magenta) planetary boundary layer height (PBLH; m) for ASIA-AQ flight days on (a) 18 March, (b) 19 March, (c) 23 March, and (d) 25 March 2024. PBLH was derived from airborne HSRL-2 lidar observations following the methodology of Christopoulos et al. (2025). Overall, WRF-Chem reproduces the temporal evolution of the daytime boundary layer reasonably well on several days (CORR = 0.5–0.7), particularly during periods of strong convective mixing. However, the model exhibits biases in PBL depth. These results suggest that while WRF reasonably captures the timing of PBL evolution, it frequently misrepresents the magnitude of boundary layer development, particularly over-deepening the PBL during rapid daytime growth.


Thailand PCD Air Monitoring Data vs. WRF-Chem
NO2
D01
Obs
WRFBase
WRFUpdated
WRFUpdated + BC
(a)
NO2 [ppbV]
Thailand PCD Air Monitoring Data vs. WRF-Chem
NOx
D01
Obs
WRFBase
WRFUpdated
WRFUpdated + BC
(b)
NOx [ppbV]
Thailand PCD Air Monitoring Data vs. WRF-Chem
O3
D01
Obs
WRFBase
WRFUpdated
WRFUpdated + BC
(c)
O3 [ppbV]
Mar 14
Mar 15
Mar 16
Mar 17
Mar 18
Mar 19
Mar 20
Mar 21
Mar 22
Mar 23
Mar 24
Mar 25
Mar 26
Mar 27

231

232

233





Figure S11~~12~~. Comparison of WRF_{Base} D01 (dotted red), ~~and~~ $WRF_{Updated}$ D01 (solid red), and $WRF_{Updated+BC}$ D01 (solid coral) simulations against Thailand Pollution Control Department (PCD) ground-monitor network observations for (a) NO_2 mixing ratio (ppbv), (b) NO_x mixing ratio (ppbv), and (c) O_3 mixing ratio (ppbv) during the ASIA-AQ deployment period (14–27 March 2024). Lines represent averages across stations within the Bangkok urban plume.






Figure S12~~13~~. Average diurnal cycles of NO_x and O_3 for D02 (a,b) and D01 (c,d). Comparison of WRF_{Base} (dotted), $WRF_{Updated}$, and $WRF_{Updated+BC}$ simulations against Thailand Pollution Control Department (PCD) ground monitor network observations during the ASIA-AQ deployment period (14-27 March 2024). ~~Average diurnal cycles of O_3 for (a) D01 and (b) D02 comparing~~

~~WRF_{Base} (dotted) and $WRF_{Updated}$ simulations against Thailand Pollution Control Department (PCD) ground-monitor network observations during the ASIA-AQ deployment period (14–27 March 2024).~~




Figure S13~~14~~. Spatial distribution of mean NO_2 column bias over the BMR averaged over 18–26 March 2024. Background shading represents WRF–GEMS bias, with the Pandora site location and corresponding model bias indicated. Results are shown for (a) $WRF_{Updated}$ and (b) WRF_{Base} simulations.




Figure S1415. Statistical performance metrics for WRF$_{Base}$ D01 (light red) and WRF$_{Updated}$ D01 (dark red) evaluated against Pandora observations during the ASIA-AQ deployment period (14–27 March 2024). Only days with high-quality Pandora data were included. Metrics include (a) mean bias (MB), (b) mean error (ME), (c) normalized mean bias (NMB), (d) normalized mean error (NME), (e) root-mean-square error (RMSE), and (f) correlation coefficient (CORR).





Figure S15~~16~~. Same as Figure S14, but for WRF_{Base} D02 (light purple) and $WRF_{Updated}$ D02 (dark purple) evaluated against Pandora observations during the ASIA-AQ deployment period (14–27 March 2024).


Mar 18th,2024
(a)
GCAS
(b)
WRF-GCAS Updated + BC
(c)
WRF-GCAS Updated
(d)
WRF-GCAS Base
Raster 1
07:44 - 10:30 LT
Raster 2
10:30 - 12:59 LT
Raster 3
14:12 - 16:20 LT
D01
15′
14°N
45′
30′
13°N
100°E 15′ 30′ 45′ 101°E 15′
3 x 10^15
4 x 10^15
5 x 10^15
6 x 10^15
7 x 10^15
8 x 10^15
1 x 10^16
1.5 x 10^16
2 x 10^16
Tropospheric Column NO₂
[molecules/cm²]




Figure S16~~17~~. Spatial comparison over the BMR on 18 March 2024 of tropospheric NO_2 columns from (a) GCAS, (b) WRF-GCAS$_{Updated}$+BC D01, ~~and~~ (c) WRF-GCAS$_{Updated}$~~Base~~ D01, and (d) WRF-GCAS$_{Base}$ D01 for raster periods corresponding to morning, midday, and afternoon local time.




Figure S17~~18~~. (a) Mean bias (MB) and (b) normalized mean bias (NMB) comparing WRF-GCAS$_{Base}$ D02 (light purple) and WRF-GCAS$_{Updated}$ D02 (dark purple) against GCAS observations for each raster period and ASIA-AQ flight day. Bias reductions in WRF-GCAS$_{Updated}$ D02 increase throughout the day, indicating improved representation of the daytime NO_2 evolution relative to the base simulation.




Figure S1819. (a) Mean bias (MB) and (b) normalized mean bias (NMB) comparing WRF-GCAS$_{Base}$ D01 (light red) and WRF-GCAS$_{Updated}$ D01 (red) against GCAS observations for each raster period and ASIA-AQ flight day. Bias reductions in WRF-GCAS$_{Updated}$ D01 increase throughout the day, indicating improved representation of the daytime NO_2 evolution relative to the base simulation.






Figure S19~~20~~. (a) Typical DC-8 flight track during the Thailand-led ASIA-AQ field campaign. The aircraft departed from U-Tapao Rayong Pattaya International Airport and flew northward across Bangkok and central Thailand toward Chiang Mai. (b) Locations of airports where the DC-8 conducted low approaches for profiling and sampling surface air quality (U-Tapao, Don Mueang, Phitsanulok, Sukhothai, Phrae, and Chiang Mai). NO_2 mixing ratios (ppbv) measured by NCAR's NO_xO_3 instrument are shown along the flight track, with time annotations indicated.



(a)
March 18th
D01 – Morning
March 18th
D01 – Midday
March 18th
D01 – Afternoon
(b)
March 25th
D01 – Morning
March 25th
D01 – Midday
March 25th
D01 – Afternoon
DC-8 (NO_xO_3)
DC-8 (CANOE)
WRF_{Base}
$WRF_{Updated}$
$WRF_{Updated+BC}$
Altitude [m]
NO_2 [ppb]
(c)
GEMS
Mar 18th
01:45 UTC
Tropospheric Column NO_2 [molecules/cm²]

369


(a)
March 18th
D01 – Morning
March 18th
D01 – Midday
March 18th
D01 – Afternoon
(b)
March 25th
D01 – Morning
March 25th
D01 – Midday
March 25th
D01 – Afternoon
DC-8 (NO_xO_3)
DC-8 (CANOE)
WRF_{Base}
$WRF_{Updated}$
Altitude [m]
NO_2 [ppb]
(c)
GEMS
Mar 18th
01:45 UTC
Tropospheric Column NO_2 [molecules/cm²]

370

Figure S2021. Comparison of WRF-Chem D01 simulations (WRF_{Base}: dotted red; $WRF_{Updated}$: solid red; $WRF_{Updated+BC}$: solid coral) with airborne in situ measurements from the CANOE (dashed black) and NO_xO_3 (solid black) instruments aboard the DC-8 for (a) 18 March and (b) 25 March 2024. Profiles are grouped by morning (06:00:00–11:00:00 LT), midday (11:00:00–13:00:00 LT), and afternoon (13:00:00–17:00:00 LT) approaches at Don Mueang International Airport, whose location relative to the Bangkok urban plume is shown in (c).


NO2 Column Raster Evaluation
BMR
D02
1e16
Tropospheric Column NO2 [molec/cm2]
Ratio
GCAS
GEMSBC
WRF-ChemUpdated + BC
WRF-GCASUpdated + BC
WRF-GEMSUpdated + BC
WRF-GCASUpdated + BC / WRF-GEMSUpdated + BC
GCAS / GEMSBC
GCAS / GEMS
Mar 18 R1
Mar 18 R2
Mar 18 R3
Mar 19 R1
Mar 19 R2
Mar 21 R1
Mar 21 R2
Mar 21 R3
Mar 23 R1
Mar 23 R2
Mar 23 R3
Mar 25 R1
Mar 25 R2
Mar 25 R3




Figure S2122. Daily raster comparison of tropospheric NO_2 columns from $WRF_{Updated+BC}$ D021 (dashed dark purplered), WRF-$GCAS_{Updated+BC}$ D021 (dashed green), WRF-$GEMS_{Updated+BC}$ D01 (dashed blue), GCAS (green), and GEMS (blue) for ASIA-AQ flight days. Dashed model lines indicate simulations with instrument-specific averaging kernels applied. The difference in sensitivity between GCAS and GEMS measurements is illustrated by the ratio of WRF-GCAS to WRF-GEMS, plotted in black.