

Seasonal prediction of springtime tornado activity in the United States using a hybrid model

Authors: Matthew Graber¹, Zhuo Wang¹, & Robert J. Trapp¹

¹Department of Climate, Meteorology, & Atmospheric Sciences, University of Illinois Urbana-Champaign, Urbana, 61820, United States

Correspondence to: Zhuo Wang (zhuowang@illinois.edu)

Abstract: Tornado activity in the contiguous United States (CONUS) causes fatalities and financial losses every spring, motivating attempts to skillfully predict springtime tornadoes. Such predictions would facilitate decision-making and resource management for both public and private stakeholders. Using ERA5 reanalysis, we analyze five April-May weather regimes (WRs) from 1981-2023, some of which strongly modulate tornado activity, ~~and the~~. The WR information is incorporated into a hybrid model to predict April-May CONUS tornado activity, including tornado outbreaks (days with >10 EF-1+ tornadoes). ECMWF seasonal forecasts initialized on April-1st are applied to predict WR frequency, including persistent and non-persistent WRs (lasting ≥ 5 and < 5 consecutive days, respectively). Prediction skill is evaluated using leave-one-year-out cross-validation. Predicted and observed tornado outbreak frequencies are significantly correlated ($cc=0.38$). Outbreak predictions are more skillful during the positive phase of the Arctic Oscillation (AO) and Pacific North American pattern (PNA), with a proportion correct of 0.75 and 0.71, respectively. In general, model skill is higher during climate mode phases that favor suppressed tornado activity. This implies that climate modes of low-frequency variability can be used to identify forecasts of opportunity ~~for low tornado activity.~~ SSTs over the North Pacific and North Atlantic may help explain the predictability of tornado activity, specifically a +PNA pattern for years of low tornado activity, but further ~~work needs to be done~~ research is needed to confirm those results. Our study demonstrates the potential for skillful prediction of spring tornado outbreaks using WR forecasts and should be prioritized in future work.

1. Introduction

Tornadoes in the contiguous United States (CONUS) result in significant losses of life and property (Ashley, 2007; NCEI, 2024; Strader et al., 2024). From 1981-2023 there were 2833 tornado-related fatalities in the CONUS, accounting for ~13% of weather-related fatalities ~~(National Weather Service, 2025).~~ (National Weather Service, 2025). Of these tornado-related fatalities, roughly 76% are associated with tornado outbreaks (TOs; hereinafter, days with > 10 EF/F-1 tornadoes; Graber et al. 2025), consistent with Schneider et al. (2004). Recent studies show a statistically significant increasing trend of +2.5 TOs per decade since 1960, the majority of which occur in the boreal spring (Brooks et al., 2014; Graber et al., 2024). Given these trends of tornado activity and their societal impacts, skillful seasonal predictions of springtime tornado activity would improve decision-making and resource management for both public and private stakeholders.

Subseasonal-to-seasonal (S2S) predictability of tornado activity has been previously investigated. Deterministic and probabilistic GEFS forecasts demonstrated skill in predicting tornado activity out to day 9 (Gensini and Tippet, 2019). The Extended-Range Tornado Activity Forecast (ERTAF) project (Gensini et al., 2020) produced tercile-forecasts 2-3 weeks in advance during boreal spring and found predictive skill at both lead times. ERTAF emphasized the role of large-

scale circulation patterns in conjunction with favorable severe thunderstorm environments, such as adequate convective available potential energy (CAPE) and vertical wind shear (VWS). Baggett et al. (2018) developed an empirical model using Madden Julian Oscillation phases and skillfully predicted weekly severe weather forecasts out 2-5 weeks in March-June. Similarly, Lepore et al. (2017) applied extended logistic regression based on the winter El Niño Southern Oscillation (ENSO) phase to predict tornadoes in March-May, and found that the tornado prediction skill is higher during La Niña years compared to El Niño years. This background motivates the work herein that seeks to further explore skillful seasonal prediction of tornado activity.

Modes of climate variability are important drivers of seasonal tornado activity and can serve as sources of their predictability. Since tornado activity depends on favorable thermodynamic and dynamic environments (Thompson et al., 2003), shifts in jet streams or moisture transport associated with low-frequency climate variability may enhance or suppress springtime severe weather potential across the CONUS (Cook and Schaefer, 2008). A major mode of interannual climate variability is ENSO (Chen and Van den Dool, 1997), which is characterized by the anomalous warming (El Niño) and cooling (La Niña) of sea-surface temperatures (SSTs) in the central and eastern equatorial Pacific (Bjerknes, 1969). ENSO modulates the strength and position of the Pacific jet stream (Bjerknes, 1969) and can affect the large-scale circulations that influence the frequency, intensity, and spatial locations of CONUS severe weather. Previous studies have linked the winter ENSO phase to the variability of tornado activity, [with increased springtime tornado activity during the La Niña phase, particularly in the Southeast](#) (Allen et al., 2015, 2018; Cook and Schaefer, 2008; Knowles and Pielke, 2005; Lee et al., 2016; [Malloy and Tippett, 2024](#)).

The Arctic Oscillation (AO) and North Atlantic Oscillation (NAO) are prominent modes of climate variability in the extratropics (Cohen et al., 2014). Specifically, the AO refers to the leading EOF of variability in ~~northern hemisphere~~[Northern Hemisphere](#) sea-level pressures at mid and polar latitudes, while the NAO refers to the sea-level pressure difference between the Subtropical High and Subpolar Low over the North Atlantic (Hurrell, 1995; Hurrell and Deser, 2010; Thompson and Wallace, 2000; Wyburn-Powell and Jahn, 2024). NAO and AO are strongly correlated and modulate the strength and position of the midlatitude jet stream. ~~Previous studies have linked the variability of NAO and AO to the variability of tornado activity, primarily over the central and eastern CONUS (Childs et al., 2018; Elsner et al., 2016; Niloufar et al., 2021; Tippett et al., 2022), suggesting that NAO and AO provide additional sources of predictability for springtime tornado activity.~~[Previous work has demonstrated that NAO and AO influence tornado activity across the central and eastern CONUS. The positive AO phase is associated with enhanced winter and early-spring tornado activity in the Southeast \(Childs et al., 2018; Hoogewind et al., 2025a; Niloufar et al., 2021; Tippett et al., 2022\), while the positive NAO phase corresponds to anomalously low springtime tornado activity in the Southeast and Central Plains \(Elsner et al., 2016; Niloufar et al., 2021\). These relationships indicate that the NAO and AO may provide additional sources of predictability for springtime tornado activity.](#)

Another important mode is the Pacific North American (PNA) pattern, which is a leading mode of low-frequency variability over the North Pacific and North America (Phillips et al., 2014). Although the PNA is influenced by ENSO (Wallace and Gutzler, 1981), it exhibits some

independent variability (Li et al., 2019) and can exist in the absence of interannual SST variability (Lau, 1981). PNA modulates the position and intensity of the East Asian jet stream which influences large-scale circulations relevant for severe weather over the CONUS (Leathers et al., 1991; Li et al., 2019; Ning and Bradley, 2015). ~~The negative phase of PNA has been associated with enhanced springtime tornado activity (Munoz and Enfield, 2011), and previous work has linked the variability of tornado activity to SST anomalies that are connected to PNA-like patterns over the Pacific (Chu et al., 2019).~~ The negative phase of PNA has been associated with enhanced springtime tornado activity and high-impact tornado outbreaks (Kim et al., 2024; Munoz and Enfield, 2011). These relationships suggest that PNA, along with ENSO, AO, and NAO, potentially serve as sources of predictability for springtime tornado activity.

~~Since climate modes evolve on the seasonal and longer timescales, they may serve as sources of predictability for seasonal tornado activity. However, they do not fully capture the synoptic-scale variability of atmospheric circulation. In addition to low-frequency climate modes, there is growing evidence that SST variability could provide an additional source of predictability for seasonal tornado activity. Increasing CONUS tornado activity has been previously linked to global SST increases, with anomalously warm North Pacific SSTs associated with enhanced tornado activity (Zhao et al., 2025). Additional work shows that North Pacific SST anomalies can influence the variability of tornado activity through PNA-like circulation anomalies, including an eastward shift of the Aleutian Low and enhanced moisture transport from the Gulf of Mexico (Chu et al., 2019). The North Atlantic tripole SST pattern, which reinforces the NAO through positive air-sea interactions (Czaja and Frankignoul, 2002), has been linked to CONUS tornado outbreaks (Lee et al., 2016). Gulf of Mexico SSTs also modulate spring tornado frequency with anomalously warm (cool) Gulf SSTs favoring more (fewer) CONUS tornadoes (Hoogewind et al., 2025b; Molina et al., 2016). These Gulf SSTs are also somewhat dependent on the phase of ENSO with La Niña (El Niño) favoring warmer (cooler) SSTs. These studies motivate a closer examination of how SSTs may serve as an additional source of predictability for springtime tornado activity.~~

~~A limitation of such SST variability as well as of low-frequency climate modes is that they do not fully capture the synoptic-scale variability of atmospheric circulation.~~ Weather regimes (WRs) can effectively bridge the gap. WRs represent a finite number of equilibrium, recurring states of the atmospheric circulation (Charney and DeVore, 1979; Hannachi et al., 2017; Michelangeli et al., 1995). Miller et al. (2020) is among the first to apply WRs to tornado prediction. They constructed a hybrid prediction model for weekly tornado activity in May using WRs and achieved skillful prediction out to week 3. Tippet et al. (2024) explored the modulation of tornado activity using year-longround WRs (Grams et al. 2017; Lee et al. 2023) and found statistically significant ~~relations~~relationships between tornado reports and WRs during all months except June-August, with ~~results mainly driven by years with more~~the frequency of the Pacific Ridge WR days ~~being the main driver of the relationship with tornado reports~~. Graber et al. (2025) identified two WRs that strongly affect the warm-season tornado activity, especially TOs, and developed an empirical model using WR frequency, persistence, and ~~TD probabilities~~probability of tornado days (TDs; ~~hereinafter, days with ≥ 1 EF/F-1+ tornadoes~~). The empirical model skillfully ~~captures~~captured the interannual variability of both TDs and TOs using WR information derived from the ERA5 reanalysis ~~and motivate its application to seasonal prediction~~. Such a demonstration of empirical model skill by Graber et al. (2025) suggests that WR-based models provide an avenue for improving seasonal tornado prediction beyond the capabilities of current operational model guidance.

The remainder of this paper is organized as follows. Section 2 describes the data and methodology, including the hybrid model. Section 3 presents the hybrid predictions and a discussion of the sources of predictability that help connect the WRs, tornado activity, and climate modes together. The paper ~~will culminate~~[culminates](#) with a summary and discussion in Section 4.

2. Methodology

2.1 Weather Regimes

Daily 500-hPa heights (500H) from the European Centre for Medium-Range Weather Forecasts (ECMWF) reanalysis, version 5 (ERA5) (Hersbach et al., 2020) were used to analyze weather regimes (WRs) during April-May, the season of peak tornado activity (Graber et al., 2024), from 1981-2023. The seasonal cycle, defined as the long-term mean 500H for each calendar day, was removed from daily 500H. Unlike Graber et al. (2025), we did not remove the long-term linear trend in 500H of ERA5 to be consistent with the ECMWF seasonal forecast data. Rather than re-deriving WRs for this period, we used the WRs derived by Graber et al. (2025) using the K-means clustering method for 1960-2022 April-July as the reference WR patterns. WRs were assigned by finding the reference WR pattern with the smallest Euclidean distance from the daily 500H anomalies. We took this approach because K-means clustering yields slightly different WR patterns in different time periods and employing WRs during a longer time period as the reference patterns improves the robustness of the results.

Seasonal forecasts of daily 500H anomalies from the ECMWF were used to predict April-May WR frequency from 1981-2023 (Copernicus Climate Change Service, 2018; Vitart et al., 2017). The 25-ensemble forecasts are initialized once per month, are available on a $1^\circ \times 1^\circ$ grid, are on 12-hour temporal resolution, and go out 7 months. The forecasts initialized on April 1st were selected to mitigate the effect of the spring-predictability barrier (~~Duan and Wei, 2012~~)([Duan and Wei, 2012](#)). WRs were assigned for each ensemble forecast by projecting the forecast 500H anomalies to the reference WR patterns, and the resultant long-term mean WR frequencies are similar to those derived from ERA5. Persistent and nonpersistent WRs are defined as WRs lasting for ≥ 5 consecutive days and for < 5 consecutive days, respectively.

2.2 Tornado observations

Tornado reports during April-May 1981-2023 were obtained from the NOAA Storm Prediction Center Severe Weather Database. Reports are georeferenced with time, date, and EF/F rating. To be consistent with previous work, tornado days (TDs) are defined as days with ≥ 1 tornadoes ranked EF/F-1 or greater, and tornado outbreaks (TOs) are defined as any day with > 10 tornadoes ranked EF/F-1 or greater (Graber et al., 2024, 2025). EF/F-0 reports were excluded due to reporting uncertainties (Brooks et al., 2014; Trapp, 2013). Known biases remain in this database, which a focus on TDs rather than raw tornado reports attempts to alleviate (Brooks et al., 2014; Graber et al., 2024; Trapp, 2014).

2.3 Hybrid model

Following Graber et al. (2025), an empirical model was used to evaluate the seasonal prediction of tornado activity:

$$TI(t) = \sum_{i=1}^5 f(i, t)_p \times P_{i,p} + \sum_{i=1}^5 f(i, t)_{np} \times P_{i,np} \quad (1)$$

where P_i denotes the TD probability for each WR, which is defined as the number of TDs with WR- i divided by the number of total WR- i days. The predictand, $TI(t)$, which is the tornado index for year t , $TI(t)$, is computed by using the seasonal count of WR- i days in year t , $f(i,t)$, multiplied by P_i . TD probabilities and counts were calculated for persistent and nonpersistent WRs separately, denoted by subscripts p and np , respectively. The same procedure was applied to TOs.

To evaluate tornado prediction skill, a leave-one-year-out cross-validation method was employed. Specifically, 1981 was first held for testing, and $P_{i,np}$ and $P_{i,p}$ were determined using the WR information derived from the ERA5 reanalysis and tornado reports during 1982-2023. The seasonal WR counts from the ECMWF ensemble forecasts for 1981 were used in Eq. 1 to predict the tornado index value in 1981. This process was repeated for each year, yielding a predicted time series of the tornado index. The tornado index time series for all ensemble members were averaged to form the ensemble mean of tornado index, which was standardized using z-score normalization.

The prediction skill was quantified using the Pearson correlation between the predicted and observed tornado index time series. Additionally, a three-tier categorical verification was used by classifying both observed and predicted tornado index values into lower, (0th-33rd percentile), middle, (33rd-67th percentile), and upper (67th-100th percentile) terciles. A correct prediction was recorded when the observed and predicted tornado index values fell within the same tercile for a given year. The proportion correct (PC) is defined as the number of correct predictions divided by the number of total predictions.

To assess the potential impacts of climate modes on tornado predictability, PC was calculated separately for positive, negative, and neutral phases of various low-frequency climate modes (see more information on climate modes in section 2.4). Significance ($p \leq 0.05$) between PC of each phase and random chance (33% for three-tier verification) was determined using a Monte Carlo test with 10,000 resamples.

2.4 Sources of Predictability

The impacts of some climate modes, ENSO (via Nino3.4), PNA, NAO, and AO, on tornado activity and WR counts were investigated to provide a physical basis for TD and TO predictability. We focus on their low-frequency variability on the seasonal and longer time scales and derived their April-May seasonal indices using the data from the NOAA Climate Prediction Center (2024b, a), (2024a, b). Positive and negative phases of a climate mode are defined as the years when the standardized springtime (April-May) index exceeded ± 0.9 . 0.9 was used instead of 1.0 to slightly increase the sample size.

TD and TO probability anomalies were calculated for each climate mode phase as:

$$P_a = \frac{P_r - P_c}{P_c} \times 100\% \quad (2)$$

where the climatological mean TD (TO) probability (P_c) is defined as the total number of TDs (TOs) divided by the total number of days; P_r is TD (TO) probability for the given phase of the climate mode; and P_a is the percentage anomaly.

3. Results

3.1 Model Prediction

For completeness, the five WRs (Fig. 1a-e) are briefly described here. WR-A features anomalous highs over the Pacific Northwest and off the U.S. east coast. WR-B is characterized by a prevailing anomalous low centered over central North America and an anomalous high over the Southeast. WR-C exhibits a three-cell wave pattern with anomalous lows over both coasts. WR-D and WR-E display west-east dipole patterns that nearly mirror each other. The WR spatial structures closely resemble WRs in Miller et al. (2020), Zhang et al. (2024), and Lee et al. (2023). Specifically, WR-AB and WR-E resemble the Pacific Ridge and Alaskan Ridge regimes in Lee et al. (2023), respectively, but noticeable differences exist due to differences in data processing procedures and the time periods analyzed. The impacts of the WRs on CONUS tornado activity during April-May 1981-2023 (Fig. 1f and 1h) are similar to those during April-July 1960-2022 from Graber et al. (2025; their Fig. 4). WR-A is the least favorable WR for springtime tornado activity, and WR-B is the most favorable. Tornado activity in WR-C and WR-E are slightly unfavorable, and in WR-D is slightly favorable. The estimated TDs and TOs based on the empirical model (Eq. 1) and the ERA5 during April-May 1981-2023 are significantly correlated with the observed time series (Figs. 1g and 1i).




Figure 1: 500H anomaly patterns of WRs A-E with long-term frequency indicated in panel title (a-e). ERA5 tornado day (f) and tornado outbreak (h) CONUS probability anomalies for persistent (≥ 5 days; light green, pink) and

nonpersistent (< 5 days; dark green, indigo) WRs. Time series of standardized tornado days (g) and tornado outbreak days (i) from the observation (blue) and estimation of the empirical modeling (red). Spearman rank correlation and p-value are indicated in panels g and i.

Before assessing the hybrid predictions of tornado activity, we first evaluate the prediction skill of WRs by the ECMWF springtime forecasts. The ECMWF ensemble mean prediction of springtime WR counts are significantly correlated with the ERA5 springtime WR counts (Fig. 2a-e). This indicates the seasonal predictability of WRs and provides the basis for tornado prediction using the hybrid framework (i.e., Eq. 1).





Figure 2: Z-score normalized ERA5 (blue) and ECMWF ensemble mean (red) springtime frequency of each WR with Spearman rank correlation and p-value indicated in each panel (a-e).

We assess the hybrid predictions of TDs and TOs using leave-one-year-out cross-validation method (Fig. 3). The hybrid prediction of TOs has a significant Pearson correlation of **0.38** with the observed TO time series, while TD prediction shows no skill, with a Pearson correlation close to zero. The skill contrast between the TO and TD predictions is probably because TOs usually occur under strong and persistent synoptic-scale patterns (Mercer et al. 2012; Cwik et al. 2022; Jiang et al. 2025; Graber et al. 2025), while transient (*i.e., non-persistent*) and weak patterns (*e.g., patterns with weaker 500H anomalies than those associated with TOs*), which are not well captured by the ECMWF springtime forecasts, may have strong impacts on TDs. In addition, it is worth noting that the percentage anomalies of TOs associated with various WRs are generally stronger than those of TDs (Fig. 1).

Cross-Validation



Cross-Validation





Figure 3: Standardized TDs (a) and TOs (b) time series from observation (red) and the hybrid prediction (blue). Thin gray lines represent the individual ensemble prediction (grey) time series. Pearson correlations (cc) and p-values (p) are shown at the upper left corner of each panel.

The tercile-based TO prediction indicates that the hybrid model correctly predicts **53.5%** of years. Additionally, the PC is strongly modulated by some climate modes (Fig. 4). In particular, the hybrid model performs significantly better during the positive phases of the AO (**75%** of correct prediction), NAO (**71.4%**), PNA (**71.4%**), and ENSO (**70.0%**). PC also improves in the negative phases of NAO (**62.5%**), AO (**55.6%**), and ENSO (**55.6%**), but the increases in PC are not significant ($p > 0.05$). In contrast, the PC in a neutral phase is below **53.5%**. Although the PC of the tercile-based TD prediction is close to random chance (**34.9%**), the hybrid model performs better in +PNA years (**57.1%**, $p=0.23$) (figure not shown). This suggests that climate modes may be used to identify forecasts of opportunity for springtime tornado prediction.

Additional cross-validation tests by leaving two-, three-, four-, and six-year out yield similar results (Fig. S1). TO predictions maintain skillful across all tests, while TD predictions remain unskillful. Given the low model skill (Fig. 3) and weaker connection to WRs for TDs (Fig. 1), the remainder of this paper focuses solely on TOs.

Proportion Correct






Figure 4: Proportion of correct TO predictions for negative (blue), neutral (grey), and positive (red) phases of AO (a) and NAO (b), PNA (c), and ENSO (d). Numbers above the bars represent p-values using a Monte Carlo test with 10,000 resamples, and bold values represent significant ($p \leq 0.05$) differences from random chance. The dashed, horizontal line represents the overall PC (i.e., 53.5%).

3.2 Low-frequency climate modes as sources of Predictability

To investigate predictability sources for tornado activity, the relationship between tornado activity and low-frequency climate modes is examined using springtime TD and TO probability anomalies associated with the AO, NAO, PNA, and ENSO phases (Figs. 5a–d). In addition to CONUS, we examined different regions, Midwest (MW), Southern Great Plains (SGP), Southeast (SE), and Southern Great Plains (SGP) (Fig. S2). The region definitions follow Moore (2018), with exception of the SGP, which herein contains New Mexico because it has storm events similar to those in the Texas Panhandle.

Significantly enhanced TO probability in the CONUS and Midwest is associated with the -AO phase. TO probability is also enhanced in the SE, NGP and SGP, but the anomalies are not statistically significant. In contrast, the probability anomalies of TDs during +AO and -AO phases

are much weaker and all are statistically insignificant, consistent with Tippet et al. (2022) showing a weaker tornado-AO relationship in April than in February. TO probability is also enhanced in the Southeast, NGP and SGP, but the anomalies are not statistically significant. +AO years are characterized by an anomalous high over the southeastern CONUS and negative anomalies over the north-central CONUS (Fig. S3a), implying enhanced westerly flow aloft and promoting moisture and warm-air advection from the Gulf of Mexico. These circulation anomalies lead to positive MUCAPE and TO probability anomalies over the Southeast and SGP (Fig. S3a). In contrast, -AO years feature negative 500H anomalies over the central CONUS, accompanied by anomalously positive VWS. Although reduced MUCAPE tends to suppress TD probability, the increased (Fig. S3b). Increased VWS supports positive TO probability anomalies when it occurs on days where MUCAPE is high (Diffenbaugh et al., 2013; Sherburn et al., 2016). These positive TO probability anomalies during -AO are only statistically significant over the Midwest, consistent with the region of anomalously positive VWS anomalies.

The TD and TO probability anomalies during the NAO phases are generally consistent in sign with those during AO phases when one phase is statistically significant, but quantitative differences exist. TD probability is reduced significantly over CONUS, SGP, and NGP, and TO probability is reduced significantly over the NGP during the +NAO phase. +NAO years feature an anomalous 500H high over the west-central CONUS, which hinders moisture and heat transport from the Gulf of Mexico. These circulation changes result in anomalously low TD probabilities over the SGP and NGP (Fig. S3e). Additionally, anomalously low VWS over the SGP further limits tornado potential during +NAO years. However, +NAO years also feature an anomalous 500H high over the western Atlantic and anomalous 500H low over the Southeast, which enhances southerly flow and moisture transport into the Southeast (Zhao et al., 2025), supporting positive MUCAPE anomalies (Fig. S3eS3c) and corresponding positive TO probability anomalies in that region (Fig. S4b).


The positive PNA phase (+PNA) is unfavorable for tornado activity TOs across the CONUS, MWMidwest, SGP, and SW, particularly for TOs Southeast (Fig. S4c). +PNA years feature an anomalous 500H high over the western CONUS and an anomalous 500H low over the eastern CONUS (Fig. S3eS3e), implying reduced moisture transport from the Gulf of Mexico and anomalously low MUCAPE and VWS. These circulation anomalies hinder tornado activity across the MWMidwest, SGP, and SW-Southeast. Positive TO probability anomalies are present across the CONUS in the negative phase of PNA (Fig. 5c), though there is no clear circulation pattern over the CONUS during these years (Fig. S3f). Springtime PNA has become more negative over time (Fig. S4), possibly explaining the increasing trend in TOs (Graber et al., 2024), though further work with other seasons needs to be done to confirm this, which goes beyond the scope of this study.

TO probability is enhanced in La Nina years in various regions and is accompanied by positive VWS anomalies, although the TO enhancement is significant only in the SESoutheast. Additionally, TDTO probability over the SGP and TO probability over NGP are enhanced in El Nino years. The La Nina-tornado-activity link is largely consistent with previous work (e.g., Cook and Schaefer 2008; Cook et al. 2017; Tippet et al. 2022; Allen et al. 2015; Moore 2019), but it is worth noting that the ENSO state in the spring timespringtime is examined here, while some previous studies focus on the winter ENSO state. Previous studies have found that ENSO modulates the phase of PNA during boreal winter (Li et al., 2019; Soulard et al., 2019); however, these spring results are not consistent with those findings. Accordingly, the correlation between


the ENSO and PNA indices are much stronger during the peak stage of ENSO in December than during the ENSO decay stage in April-May (Fig. S5).

The model appears to perform better (Fig. 4) in years when climate modes unfavorable for TOs, such as +NAO and +PNA, are present. To further evaluate skill, the Threat Score (Schaefer, 1990) was calculated for upper-tercile TO years. These years yield a Threat Score of **0.375** with 6 hits, 2 misses, and 8 false alarms, indicating a tendency for the model to overpredict TOs. For lower-tercile years, the Threat Score is slightly higher at **0.429**, with 9 hits, 7 misses, and 5 false alarms. While lower-tercile years have greater overall skill, they also include more misses. Overall, the model demonstrates higher skill for lower-tercile years, and modest skill for upper-tercile years.

Overall, the link between climate modes and tornado activity suggests that low-frequency climate modes provide a source of predictability for springtime [tornado TO](#) prediction, but the predictability seems slightly higher during inactive years for tornado activity. ~~Additionally, low-frequency climate modes modulate TO probabilities more strongly than TD probabilities, consistent with the stronger TO anomalies associated with WRs (Fig. 1h) and the higher predictive skill of TOs (Fig. 3).~~



Tornado Outbreak Probability Anomalies by Climate Mode and Region



347

Figure 5: ~~TD (left column) and TO (right column)~~ probability anomalies by region and climate mode: AO (a-b), NAO (c-d), PNA (e-f), and ENSO (g-h). Asterisks represent significant anomalies ($p \leq 0.05$) based on a Monte Carlo test with 10,000 resamples of each regions' data.


The connection between climate modes and WRs is examined next to investigate whether climate modes modulate tornado activity via WRs. Figure 6 shows the frequency/relative likelihood of WRs in each WR during different phases of low-frequency climate modes, ~~which is defined~~ expressed as the number/ratio of WR days during a given its phase of the climate mode divided by the total number of days in that phase. The specific frequency to its climatological frequency of WR-A is only ~15%~36% less likely to occur during the -AO phase, while ~~the frequency of~~ WR-B is ~33%~35% more likely to occur (Fig. 6a). The increased occurrence of WR-B and decreased occurrence of WR-A in -AO years compared to the long-term mean (Fig. 1) are consistent with the positive TO probability anomalies in the CONUS and Midwest during the -AO years (Fig. 5). WR-D is ~27% less likely to occur during -AO, so the above average tornado activity in -AO (Fig. 5a) is mainly due to above average frequency of WR-B. Additionally, WR-A is ~21% more likely to occur during +AO years.

During +NAO years, WR-A occurs on approximately 35% of days, making it the dominant WR during this phase. This predominance of WR-A is ~42% more likely to occur, which is consistent with the reduced TD probability in the CONUS, NGP, and SGP, and reduced TO probability in the NGP during NAO+ years and SGP (Fig. 6). Composite 500H anomalies for +NAO years resemble WR-A, while composite 500H anomalies for +AO show a mix of WR-A and WR-B (Fig. S3). WR-A's increased occurrence helps explain the high predictive skill of inactive TO years and indicates that +NAO offers a forecast of opportunity for inactive TO years (Fig. 6b). Additionally, WR-A is ~30% less likely to occur and WR-E is ~30% more likely to occur during -NAO years.

During +PNA years, WR-B occurs on ~15% of days is ~35% less likely to occur, whereas WR-E occurs on ~25% of days, making it the most frequent WR during this phase is ~62% more likely to occur (Fig. 6c). The reduced occurrence of WR-B and increased occurrence of WR-E are consistent with reduced TD and TO activity over CONUS, MWMidwest, and SGP during +PNA (Fig. 5). 5), and also help explain the high predictive skill of inactive TO years and indicate that +PNA offers a forecast of opportunity for inactive TO years (Fig. 6c). Composite 500H anomalies during +PNA years resembles WR-E, with an anomalous high over the western CONUS and an anomalous low over the eastern CONUS (Fig. S3c).

Neither positive nor negative phase of ENSO is not associated with significant changes in WR-A or WR-B frequency. Instead, the frequency of, except for WR-E (Fig. 6d). WR-E is reduced significantly in both phases of ENSO, and WR-E occurs ~32% more frequently in the positive phase of ENSO likely to occur during El Niño, which is consistent with the reduced tornado activity during El Niño years reported in previous studies (Cook et al., 2017). Relative to the neutral phase, WR-E also occurs more frequently during the negative phase of ENSO, which is statistically significant. However, WR-E's occurrence during the negative phase of ENSO is still less than any other WR. Additionally, WR-E is ~18% less likely to occur during the neutral phase.

To summarize, WRs associated with increased springtime tornado activity occur more frequently during the climate mode phases that favor increased tornado activity, while WRs associated with suppressed tornado activity preferentially occur during climate mode phases that are unfavorable for tornado activity. These consistent relationships provide additional confidence in the robustness of the WR-based prediction framework and help to understand the predictability of tornado activity.






Figure 6: ~~WR frequencies~~Relative likelihood (RL: %) of each WR in different phases of AO (a), NAO (b), PNA (c), and ENSO (d). An ~~asterisks~~asterisk indicates that the frequency is significantly different ($p \leq 0.05$) from ~~that during the corresponding neutral phase via~~climatology using a ~~student t-scipy binomial~~ test. Black horizontal line representing the climatological frequency is at RL=0.0.

3.3 Possible role of SST

In addition to low-frequency climate modes, slowly-evolving SSTs can also be an important source of predictability, sometimes through ~~their coupling with these modes. The role of SSTs is further examined in Fig. 7.~~air-sea coupling in those climate modes. Composite SST and 500H anomalies during active and inactive TO years are shown in Fig. 7. Composite years were selected as years in which the April-May TO count exceeds +1 standard deviation (active) or falls below -1 standard deviation (inactive) relative to the 1960-2023 mean. Extended Range SST (ERSST) (Huang et al., 2017) of $2^\circ \times 2^\circ$ resolution is used here, and we focus on ~~the month of~~SST anomalies in April because SST signals ~~for TOs~~ in April are more pronounced than in May based on a previous study (Chu et al., 2019) and our own analysis. In addition, the analysis is done for 1960-2023 to increase the sample size. ~~For each WR, active and inactive regime years are defined as the years when the springtime WR count exceeds ± 1 standard deviation from the mean, and composite anomalies of SST and 500H are constructed for these active and inactive years of the WR.~~

+WR-A years are characterized by an anomalous 500H high over the north-central CONUS, as part of a well-defined wavetrain pattern extending from the North Pacific to southern Greenland, in addition to an anomalous low over the Southeast and Gulf of Mexico. Although there are coherent cold and warm SST anomalies associated with the anomalous low and high over the North Pacific, respectively, the SST anomalies are not significant. Significant SST anomalies exist over the North Atlantic associated with anomalous 500H high. In contrast, the -WR-A years feature an anomalous low over the central CONUS, nearly opposite to the +WR years. However, the anomalous circulation patterns over the North Pacific and North Atlantic are not a simple mirror image to those during +WR-A years. The anomalous 500H field is characterized by an elongated anomalous low over the midlatitude North Pacific and North Atlantic, reminiscent of the -AO pattern. Significant SST anomalies are found over the North Pacific and North Atlantic, resembling the Atlantic tripole SST pattern, which helps to maintain the NAO pattern via positive air-sea interaction (Czaja and Frankignoul, 2002). Consistent with this circulation, -WR-A is suppressed during -NAO and -AO years (Fig. 6).

-WR-B is most favorable for tornado activity over the CONUS. Though positive SST anomalies exist over the central Pacific in +WR-B years, consistent with that shown in Chu et al. (2019) for April tornadoes, the anomalies are not significant. The 500H pattern over the North Atlantic is reminiscent of a -AO pattern, consistent with increased -WR-B occurrence during -AO (Fig. 6). Significant, negative SST anomalies exist over the North Pacific in -WR-B years. The associated North Pacific negative SSTs and the 500H high over Canada are reminiscent of a +PNA pattern, consistent with reduced -WR-B occurrence during +PNA (Fig. 6). Though statistically insignificant, the warm SST anomalies in the northern Pacific during +WR-B have been previously linked to enhanced CONUS tornado activity (Zhao et al., 2025) through a weakening and eastward shift of the Aleutian Low (Chu et al., 2019), which promotes enhanced southerly flow of moisture and heat from the Gulf of Mexico.

+WR-C years feature anomalously cold SSTs in the midlatitude North Pacific and North Atlantic as well as in the Gulf of Mexico. The cold SST anomalies over the Gulf of Mexico may limit moisture availability (Molina et al., 2016), partially offsetting the otherwise favorable southerly 500H anomalies over the central US. This compensating thermodynamical and dynamical effects help explain the near-climatological tornado activity during -WR-C (Fig. 1f, h). In contrast, -WR-C years are associated with prevailing warm SST anomalies in the North Pacific. Significant SST anomalies, however, are confined to the subtropical North Atlantic and portions of the subtropical eastern Pacific.

+WR-D and +WR-E years feature anomalously warm and cold SSTs in the Gulf of Mexico, respectively, though these anomalies are not statistically significant. Over the subtropical western and central North Pacific, respectively, +WR-D years exhibit significant warm SST anomalies, while +WR-E years show significant cold SST anomalies. Given that -WR-D and -WR-E are linked to enhanced and reduced TD probability, respectively (Fig. 1), this suggests that SST anomalies in the subtropical Pacific may be a source of predictability for tornado activity over the CONUS. However, the robustness of this relationship requires further examination given the incoherent pattern of the significant SST signals, and the underlying mechanisms also need to be investigated. In addition, significant cold SST anomalies exist over the North Atlantic for +WR-E years, suggesting cold SSTs in this region may be associated with reduced TD probabilities over the CONUS.

We also examined the composite SST and 500h anomalies for active and inactive TO years (Fig. S6). Interestingly, significant SST anomalies are found over equatorial eastern Pacific, North Pacific, and subtropical western North Atlantic during inactive TO years, while significant SST anomalies are nearly absent during active TO years. The lack of significant SST anomalies in the active tornado years are consistent with the relatively limited SST signals during the active WR-B and WR-D years and during the inactive WR-A and WR-E years, and it helps to explain the higher predictability of inactive tornado years discussed in the previous section.

Overall, Fig. 7 suggests that SST anomalies over the North Pacific and North Atlantic may help explain the predictability of some WRs and thus the predictability of tornado activity within the hybrid framework. However, WR-B and WR-D, which favor tornado activity over the CONUS (Fig. 1), are not associated with many coherent, statistically significant SST anomalies. Further investigation using longer observational records and focusing on different seasons is needed to better understand the predictability or lack thereof of tornado activity.

Composite SST Anomalies by WR Phases




Figure 7: April SST anomalies for active and inactive WR years, and hatching. During inactive TO years, significant positive SST anomalies are found over equatorial eastern Pacific and significant negative SST anomalies over the North Pacific and subtropical western Atlantic. Inactive TO years feature an anomalous 500H high over the west-central CONUS and anomalous 500H lows over the East Pacific and Southeast CONUS, resembling WR-A. A positive phase of the North Atlantic tripole SST pattern is present, consistent with (Lee et al., 2016). In contrast, significant SST anomalies are nearly absent during active TO years except a limited region of cold SST anomalies over the East Pacific just off California. The 500H anomalies are characterized by an anomalous trough over CONUS, resembling WR-B, but the anomalies are much weaker than those during the inactive TO years. The lack of significant SST anomalies and weak 500H anomalies in the active TO years are consistent with the reduced model skill during inactive TO years and during climate mode phases that are unfavorable for tornado activity and suggest that inactive TO years are more predictable than active TO years.

To quantify the link between the SST anomalies and climate modes, we calculated the pattern correlations between SST composites in Fig. 7 and those associated with various climate modes (Table S1). Stronger correlations tend to be found in inactive TO years. The highest correlation occurs for inactive TOs and +PNA ($cc=0.71$), thus demonstrating that +PNA can be used to identify a forecast of opportunity for inactive TO years.

We also examined composite SST and 500H anomalies for active and inactive WR years, defined as the years when the springtime WR count exceeds ± 1 standard deviation from the mean (Fig. S6). WR-A shows opposite circulation patterns over the CONUS between its active and inactive years, with the inactive years resembling a -AO/-NAO pattern consistent with reduced WR-A occurrence during those phases. Active WR-B years exhibit a North Atlantic 500H pattern resembling a -AO pattern. In contrast, inactive WR-B years feature anomalously low SSTs over the North Pacific and an anomalous 500H high over Canada, resembling a +PNA pattern. These patterns are consistent with how PNA and AO modulate WR-B frequency (Fig. 6). The SST composite for active WR-B years have the highest pattern correlation with active TO years ($cc=0.45$, Table S1). Active WR-C years feature a favorable 500H pattern for tornado activity with an anomalous low over the western Atlantic and enhanced southerly flow from the Gulf of Mexico. However, anomalously low Gulf of Mexico SSTs in active WR-C years demonstrates offsetting thermodynamical and dynamical components, which helps explain the near-climatology tornado activity during WR-C. WR-D and WR-E show opposite and statistically significant SST anomalies in the subtropical Pacific, suggesting some potential predictability. Since WR-D is favorable for tornado activity and WR-E suppresses it, these relationships are noteworthy, though the robustness of the relationship requires further examination given the overall weak SST patterns. Overall, the SSTs over the Pacific and Atlantic may help explain the predictability of WRs, though further investigation using longer observational records is needed to better understand the predictability of tornado activity.

Composite SST Anomalies by Active and Inactive Tornado Outbreak Phases




Figure 7: April composite SST anomalies for active and inactive tornado outbreak years. Hatching indicates significant SST anomalies ($p \leq 0.05$) using a one-sample, two-sided t-test. 500hPa anomalies are shown in gray contours. Sample sizes are included in each panel title.

4. Summary

A hybrid model for the seasonal prediction of CONUS springtime tornado activity is evaluated during 1981-2023. The model employs WR forecasts from the ECMWF ensemble springtime forecasts initialized on April-1st. Using the leave-one-year-out cross-validation, the WR-based model shows no predictive skill for TDs, but the predicted time series of TO days is significantly correlated with the observed time series. Predictive skill for TOs is modulated by low-frequency climate modes and is significantly enhanced during +AO, +NAO, +PNA, and El Niño years, with the proportion correct of 75%, 71.4%, 71.4%, and 70%, respectively. Given the difficulty of predicting TDs beyond subseasonal timescales (Gensini et al., 2020), springtime prediction efforts should prioritize TOs due to their higher predictive skill and societal impacts, and low-frequency climate modes can help to identify forecasts of opportunity.

To investigate the predictability sources of springtime tornado activity, the composite ~~TD and~~ TO probability anomalies were examined during different phases of climate modes. Significantly enhanced TO probability anomalies were associated with -AO and La Niña years, while significantly reduced TO probability anomalies were associated with +NAO and +PNA years. Corresponding circulation patterns during these years provide physical support, through CAPE and vertical shear anomalies. ~~TO probability anomalies are generally stronger than TD anomalies associated with climate modes, which helps explain the higher predictive skill of TOs. These results suggest that~~ In general, the model performs better during years that are unfavorable for tornado activity, possibly due to a lower signal-to-noise ratio, and use of the Threat Score ~~confirmsscore results confirm this result, although~~ the model is better in inactive TO years. However, this skill test still demonstrates shows modest skill ~~in~~during active TO years.

Additionally, WR occurrence is modulated by some climate modes. WR-A is the dominant WR during +NAO years and WR-E is the dominant WR during +PNA years, consistent with reduced tornado activity during +NAO and +PNA years. Given the increased model skill during inactive TO years, +NAO and +PNA may provide forecasts of opportunity for reduced springtime tornado activity. Furthermore, WR-B is the dominant WR during -AO years and is the least frequent WR during +PNA years, consistent with enhanced (reduced) tornado activity during the -AO (+PNA) phase. WRs associated with increased springtime tornado activity tend to occur more frequently during the climate mode phases that favor increased tornado activity, while WRs associated with

suppressed tornado activity preferentially occur during climate mode phases that are unfavorable for tornado activity. This physical consistency supports the use of WRs as a physically meaningful framework for predicting TOs.

The role of SSTs in the predictability of ~~WRs (with implication for the predictability of tornado activity)~~ TOs was examined, with the focus on April SST anomalies. ~~WR-A years feature a NAO SST and 500H pattern. Statistically significant SST anomalies are present over the eastern equatorial Pacific, North Pacific, and Western Atlantic, consistent with WR-A being suppressed during NAO patterns. WR-B years feature positive, though statistically insignificant, inactive TO years. In contrast, significant SSTs over the central Pacific consistent with patterns shown in Chu et al. (2019) for April tornadoes are mostly absent in active TO years. In addition, significantly negative SST anomalies exist over the northern Pacific the 500H pattern is weaker during WR-B active TO years, consistent with suppressed WR-B and TO. This suggests that TO anomalies are more predictable during +PNA inactive years. Overall than active years. In addition, SST anomalies can help explain WR predictability of some WRs, but SST signals for some other WRs are statistically insignificant or incoherent, making it challenging to identify specific SST-based predictors for WRs or tornado activity. Further investigation based on longer observational records would help better understand the role of SST in the predictability of WRs and tornado activity.~~

This study demonstrates that skillful prediction of springtime TOs is indeed possible with current resources via the connection between tornado activity and WRs. Questions remain on whether SSTs are a reliable driver of springtime TOs as well as the temporal extent to which TOs can be predicted.

Code Availability

Important python files for WR identification and modeling are included at the following link:

<https://github.com/Matt0604/Springtime-Prediction-Manuscript>

Data Availability

The ERA5 data are available through the NCAR research data archive (RDA) (d633000) and the Copernicus Climate Data Store (CDS):

<https://doi.org/10.24381/cds.bd0915c6> (Hersbach et al., 2023a)

<https://doi.org/10.24381/cds.adbb2d47> (Hersbach et al., 2023b)

The ECMWF data are available through the Copernicus Climate Data Store (CDS):

<https://doi.org/10.24381/cds.50ed0a73> (Copernicus Climate Change Service, 2018)

April SST data are available through the Extended-range sea surface temperatures data (Huang et al., 2017).

The tornado report data used in this study are available through the NOAA Storm Prediction Center severe weather database:

<https://www.spc.noaa.gov/wcm/#data>

Author Contributions

Conceptualization: MG, ZW, RJT

Methodology: MG, ZW, RJT

Project Administration: ZW, RJT

Supervision: ZW, RJT

Writing – Original Draft: MG

Writing – Review and Edits: MG, ZW, RJT

Competing Interests

Authors declare they have no competing interest.

Acknowledgements

[ZW is supported by the NSF grant 2518299.](#)

We acknowledge the NCAR Computation and Information Systems Laboratory (CISL) for providing computing resources through Derecho. All ERA5 data used in this study are available at the research data archive. All ECMWF seasonal forecast data are available via the Copernicus Climate Change Service. Tornado report data are available through the NOAA Storm Prediction Center severe weather database.

References

- Allen, J. T., Tippet, M. K., and Sobel, A. H.: Influence of the El Nino/Southern Oscillation on tornado and hail frequency in the United States, *Nat. Geosci.*, 8, 278–283, <https://doi.org/https://doi.org/10.1038/ngeo2385>, 2015.
- Allen, J. T., Molina, M. J., and Gensini, V. A.: Modulation of annual cycle of tornadoes by El-Nino-Southern Oscillation, *Geophys. Res. Lett.*, 45, 5708–5717, <https://doi.org/https://doi.org/10.1029/2018GL077482>, 2018.
- Ashley, W. S.: Spatial and temporal analysis of tornado fatalities in the United States: 1880-2005, *Weather Forecast.*, 22, 1214–1228, <https://doi.org/https://doi.org/10.1175/2007WAF2007004.1>, 2007.
- Bjerknes, J.: Atmospheric Teleconnections from the equatorial Pacific, *Mon. Weather Rev.*, 97, 163–172, [https://doi.org/https://doi.org/10.1175/1520-0493\(1969\)097%3C0163:ATFTEP%3E2.3.CO;2](https://doi.org/https://doi.org/10.1175/1520-0493(1969)097%3C0163:ATFTEP%3E2.3.CO;2), 1969.
- Brooks, H. E., Carbin, G. W., and Marsh, P. T.: Increased variability of tornado occurrence in the United States, *Science*, 346, 349–352, <https://doi.org/https://doi.org/10.1126/science.1257460>, 2014.
- Charney, J. G. and DeVore, J. G.: Multiple flow equilibria in the atmosphere and blocking, *J. Atmospheric Sci.*, 36, 1205–1216, [https://doi.org/https://doi.org/10.1175/1520-0469\(1979\)036%3C1205%3E1205:MFEITA%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1979)036%3C1205%3E1205:MFEITA%3E2.0.CO;2), 1979.

Chen, W. Y. and Van den Dool, H. M.: Atmospheric predictability of seasonal, annual, and decadal climate means and the role of the ENSO cycle: A model study, *J. Clim.*, 10, 1236–1254, [https://doi.org/https://doi.org/10.1175/1520-0442\(1997\)010%3C1236:APOSAA%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0442(1997)010%3C1236:APOSAA%3E2.0.CO;2), 1997.

Childs, S., Schumacher, R. S., and Allen, J. T.: Cold-season tornadoes: Climatological and meteorological insights, *Weather Forecast.*, 33, 671–691, <https://doi.org/https://doi.org/10.1175/WAF-D-17-0120.1>, 2018.

Chu, J.-E., Timmermann, A., and Lee, J.-Y.: North American April tornado occurrences linked to global sea surface temperature anomalies, *Sci. Adv.*, 5, <https://doi.org/https://doi.org/10.1126/sciadv.aaw9950>, 2019.

Cohen, J., Screen, J. A., Furtado, J. C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J., and Jones, J.: Recent Arctic amplification and extreme mid-latitude weather, *Nat. Geosci.*, 7, 627–637, <https://doi.org/https://doi.org/10.1038/ngeo2234>, 2014.

Cook, A. R. and Schaefer, J. T.: The relation of El Nino-Southern Oscillation (ENSO) to winter tornado outbreaks, *Mon. Weather Rev.*, 136, 3121–3137, <https://doi.org/https://doi.org/10.1175/2007MWR2171.1>, 2008.

Cook, A. R., Leslie, L., Parsons, D., and Schaefer, J.: The impact of El Nino-Southern Oscillation (ENSO) on Winter and early Spring U.S. tornado outbreaks, *J. Appl. Meteorol. Climatol.*, 56, 2455–2478, <https://doi.org/https://doi.org/10.1175/JAMC-D-16-0249.1>, 2017.

Copernicus Climate Change Service, C. D. S.: Seasonal forecast subdaily data on pressure levels, <https://doi.org/10.24381/cds.50ed0a73>, 2018.

Cwik, P., McPherson, R. A., Richman, M. B., and Mercer, A. E.: Climatology of 500-hPa geopotential height anomalies associated with May tornado outbreaks in the United States., *Int. J. Climatol.*, 43, 893–913, <https://doi.org/https://doi.org/10.1002/joc.7841>, 2022.

Czaja, A. and Frankignoul, C.: Observed Impact of Atlantic SST Anomalies on the North Atlantic Oscillation, *J. Clim.*, 15, 606–623, [https://doi.org/https://doi.org/10.1175/1520-0442\(2002\)015%3C0606%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0442(2002)015%3C0606%3E2.0.CO;2), 2002.

Diffenbaugh, N. S., Scherer, M., and Trapp, R. J.: Robust increases in severe thunderstorm environments in response to greenhouse forcing, *Proc. Natl. Acad. Sci. U. S. A.*, 110, 16361–16366, <https://doi.org/https://doi.org/10.1073/pnas.1307758110>, 2013.

Duan, W. and Wei, C.: The “spring predictability barrier” for ENSO predictions and its possible mechanism: Results from a fully coupled model, *Int. J. Climatol.*, 33, 1280–1292, <https://doi.org/https://doi.org/10.1002/joc.3513>, 2012.

Elsner, J. B., Jagger, T. H., and Fricker, T.: Statistical models for tornado climatology: Long and short-term views, *PLoS ONE*, 11, <https://doi.org/https://doi.org/10.1371/journal.pone.0166895>, 2016.

Gensini, V. A. and Tippett, M. K.: Global Ensemble Forecast System (GEFS) predictions of days 1–15 U.S. tornado and hail frequencies, *Geophys. Res. Lett.*, 46, 2922–2930, <https://doi.org/https://doi.org/10.1029/2018GL081724>, 2019.

654 Gensini, V. A., Barrett, B. S., Allen, J. T., Gold, D., and Sirvatka, P.: The Extended-Range Tornado Activity
655 Forecast (ERTAF) Project, *Bull. Am. Meteorol. Soc.*, 101, E700–E709,
656 <https://doi.org/https://doi.org/10.1175/BAMS-D-19-0188.1>, 2020.

657 Graber, M., Trapp, R. J., and Wang, Z.: The regionality and seasonality of tornado trends in the United
658 States, *Npj Clim. Atmospheric Sci.*, 7, <https://doi.org/https://doi.org/10.1038/s41612-024-00698-y>,
659 2024.

660 Graber, M., Wang, Z., and Trapp, R. J.: Linking weather regimes to the variability of warm-season
661 tornado activity over the United States, *Weather Clim. Dyn.*, 6, 807–816,
662 <https://doi.org/https://doi.org/10.5194/wcd-6-807-2025>, 2025.

663 Hannachi, A., Straus, D. M., Franzke, C. L. E., Corti, S., and Woollings, T.: Low-frequency nonlinearity and
664 regime behavior in the northern hemisphere extratropical atmosphere, *Rev. Geophys.*, 55, 199–234,
665 <https://doi.org/https://doi.org/10.1002/2015RG000509>, 2017.

666 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C.,
667 Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati,
668 G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming,
669 J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M.,
670 Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume,
671 S., and Thépaut, J.-N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–2049,
672 <https://doi.org/10.1002/qj.3803>, 2020.

673 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C.,
674 Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thepaut, J.-N.: ERA5 hourly data on
675 pressure levels from 1940 to present, <https://doi.org/10.24381/cds.bd0915c6>, 2023a.

676 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C.,
677 Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thepaut, J.-N.: ERA5 hourly data on
678 single levels from 1940 to present, <https://doi.org/10.24381/cds.adbb2d47>, 2023b.

679 [Hoogewind, K. A., Gensini, V. A., and Brooks, H. E.: On the relationship between monthly mean surface](https://doi.org/https://doi.org/10.1038/s41612-025-00993-2)
680 [temperature and tornado days in the United States, *Npj Clim. Atmospheric Sci.*, 8,](https://doi.org/https://doi.org/10.1038/s41612-025-00993-2)
681 <https://doi.org/https://doi.org/10.1038/s41612-025-00993-2>, 2025a.

682 [Hoogewind, K. A., Galarneau Jr., T. J., and Gensini, V. A.: Severe convective weather outbreaks on 10 and](https://doi.org/10.1175/MWR-D-24-0213.1)
683 [15 December 2021: Large-scale antecedent conditions, *Mon. Weather Rev.*, 153, 1171–1194,](https://doi.org/10.1175/MWR-D-24-0213.1)
684 <https://doi.org/10.1175/MWR-D-24-0213.1>, 2025b.

685 Huang, B., Thorne, P. W., Banzon, V. F., Boyer, T., Chepurin, G., Lawrimore, J. H., Menne, M. J., Smith, T.
686 H., Vose, R. S., and Zhang, H.-M.: Extended Reconstructed Sea Surface Temperature, version 5
687 (ERSSTv5): Upgrades, validations, and intercomparisons, 15 Oct 2017, 30, 8179–8205,
688 <https://doi.org/https://doi.org/10.1175/JCLI-D-16-0836.1>, 2017.

689 Hurrell, J. W.: Decadal trends in the North Atlantic Oscillation: Regional temperatures and precipitation,
690 *Science*, 269, 676–679, <https://doi.org/https://doi.org/10.1126/science.269.5224.676>, 1995.

691 Hurrell, J. W. and Deser, C.: North Atlantic climate variability: The role of the North Atlantic Oscillation, *J.*
692 *Mar. Syst.*, 79, 231–244, <https://doi.org/https://doi.org/10.1016/j.jmarsys.2009.11.002>, 2010.

693 Jiang, Q., Dawson II, D. T., Funing, L., and Chavas, D. R.: Classifying synoptic patterns driving tornadic
694 storms and associated spatial trends in the United States, *Npj Clim. Atmospheric Sci.*, 8,
695 <https://doi.org/https://doi.org/10.1038/s41612-025-00897-1>, 2025.

696 [Kim, D., Lee, S.-K., Lopez, H., Jeong, J.-H., and Hong, J.-S.: An unusually prolonged Pacific-North American](https://doi.org/https://doi.org/10.1038/s41612-024-00688-0)
697 [pattern promoted the 2021 winter quad-state tornado outbreaks, *Npj Clim. Atmospheric Sci.*, 7,](https://doi.org/https://doi.org/10.1038/s41612-024-00688-0)
698 <https://doi.org/https://doi.org/10.1038/s41612-024-00688-0>, 2024.

699 Knowles, J. B. and Pielke, R. A.: The Southern Oscillation and its effects on tornadic activity in the United
700 States, 15 pp, 2005.

701 Lau, N.-C.: A diagnostic study of recurrent meteorological anomalies appearing in a 15-year simulation
702 with a GFDL general circulation model, *Mon. Weather Rev.*, 109, 2287–2311,
703 [https://doi.org/https://doi.org/10.1175/1520-0493\(1981\)109%3C2287:ADSORM%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0493(1981)109%3C2287:ADSORM%3E2.0.CO;2), 1981.

704 Leathers, D. J., Yarnal, B., and Palecki, M. A.: The Pacific/North American teleconnection pattern and
705 United States climate. Part I: Regional temperature and precipitation associations, *J. Clim.*, 4, 517–528,
706 [https://doi.org/https://doi.org/10.1175/1520-%200442-0442\(1991\)004,0517:TPATPA.2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-%200442-0442(1991)004,0517:TPATPA.2.0.CO;2), 1991.

707 Lee, S. H., Tippett, M. K., and Polvani, L. M.: A new year-round weather regime classification for North
708 America, *J. Clim.*, 36, 7091–7108, <https://doi.org/https://doi.org/10.1175/JCLI-D-23-0214.1>, 2023.

709 Lee, S.-K., Wittenberg, A. T., Enfield, D. B., Weaver, S. J., Wang, C., and Atlas, R.: US regional tornado
710 outbreaks and their links to spring ENSO phases and North Atlantic SST variability, *Environ. Res. Lett.*, 11,
711 <https://doi.org/10.1088/1748-9326/11/4/044008>, 2016.

712 Lepore, C., Tippett, M. K., and Allen, J. T.: ENSO-based probabilistic forecasts of March-May U.S. tornado
713 and hail activity, *Geophys. Res. Lett.*, 44, 9093–9101,
714 <https://doi.org/https://doi.org/10.1002/2017GL074781>, 2017.

715 Li, X., Hu, Z.-Z., Liang, P., and Zhu, J.: Contrastive influence of ENSO and PNA on variability and
716 predictability of North American winter precipitation, *J. Clim.*, 32, 6271–6284,
717 <https://doi.org/https://doi.org/10.1175/JCLI-D-19-0033.1>, 2019.

718 [Malloy, K. and Tippett, M. K.: A stochastic statistical model for U.S. outbreak-level tornado occurrence](https://doi.org/https://doi.org/10.1175/MWR-D-23-0219.1)
719 [based on the large-scale environment, *Mon. Weather Rev.*, 152, 1141–1161,](https://doi.org/https://doi.org/10.1175/MWR-D-23-0219.1)
720 <https://doi.org/https://doi.org/10.1175/MWR-D-23-0219.1>, 2024.

721 Mercer, A. E., Shafer, C. M., Doswell III, C. A., Leslie, L. M., and Richman, M. B.: Synoptic composites of
722 tornadic and nontornadic outbreaks, *Mon. Weather Rev.*, 140, 2590–2608,
723 <https://doi.org/https://doi.org/10.1175/MWR-D-12-00029.1>, 2012.

724 Michelangeli, P.-A., Vautard, R., and Legras, B.: Weather regimes: Recurrence and quasi stationarity, *J.*
725 *Atmospheric Sci.*, 52, 1237–1256, [https://doi.org/https://doi.org/10.1175/1520-](https://doi.org/https://doi.org/10.1175/1520-0469(1995)052%3C1237:WRRSQS%3E2.0.CO;2)
726 [0469\(1995\)052%3C1237:WRRSQS%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0469(1995)052%3C1237:WRRSQS%3E2.0.CO;2), 1995.

727 Miller, D., Wang, Z., Trapp, R. J., and Harnos, D. S.: Hybrid prediction of weekly tornado activity out to
 728 week 3: Utilizing weather regimes, *Geophys. Res. Lett.*, 47,
 729 <https://doi.org/https://doi.org/10.1029/2020GL087253>, 2020.

730 Molina, M. J., Timmer, R. P., and Allen, J. T.: Importance of the Gulf of Mexico as a climate driver for U.S.
 731 severe thunderstorm activity, *Geophys. Res. Lett.*, 43, 12295–12304,
 732 <https://doi.org/https://doi.org/10.1002/2016GL071603>, 2016.

733 Moore, T. W.: Annual and seasonal tornado trends in the Contiguous United States and its regions, *Int. J.*
 734 *Climatol.*, 38, 1582–1594, <https://doi.org/https://doi.org/10.1002/joc.5285>, 2018.

735 Moore, T. W.: Seasonal frequency and spatial distribution of tornadoes in the United States and their
 736 relationship to the El Nino/Southern Oscillation, *Ann. Am. Assoc. Geogr.*, 109, 1033–1051,
 737 <https://doi.org/https://doi.org/10.1080/24694452.2018.1511412>, 2019.

738 Munoz, E. and Enfield, D.: The boreal spring variability of the Intra-Americas low-level jet and its relation
 739 with precipitation and tornadoes in the eastern United States, *Clim. Dyn.*, 36, 247–259,
 740 <https://doi.org/https://doi.org/10.1007/s00382-009-0688-3>, 2011.

741 National Weather Service: 80-year list of severe weather fatalities, available at:
 742 https://www.weather.gov/media/hazstat/80year_2024.pdf, 2025.

743 NCEI: U.S. billion-dollar weather and climate disasters, DOI: <https://doi.org/10.25921/stkw-w737w73>,
 744 2024.

745 Niloufar, N., Devineni, N., Were, V., and Khanbilvardi, R.: Explaining the trends and variability in the
 746 United States tornado records using climate teleconnections and shifts in observational practices, *Sci.*
 747 *Rep.*, 11, <https://doi.org/https://doi.org/10.1038/s41598-021-81143-5>, 2021.

748 Ning, L. and Bradley, R. S.: NAO and PNA influences on winter temperature and precipitation over the
 749 eastern United States in CMIP5 GCMs, *Clim. Dyn.*, 46, 1257–1276,
 750 <https://doi.org/https://doi.org/10.1007/s00382-015-2643-9>, 2015.

751 NOAA Climate Prediction Center: Daily climate mode indices, NOAA/NCEP/CPC [dataset], available at:
 752 <https://ftp.cpc.ncep.noaa.gov/cwlinks/>, 2024a.

753 NOAA Climate Prediction Center: Oceanic Niño Index (ONI) v5 - Monthly SST anomalies in the Nino 3.4
 754 region, NOAA/NCEP/CPC [dataset], available at:
 755 https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v5.php, 2024b.

756 Phillips, A. S., Deser, C., and Fasullo, J.: Evaluating modes of variability in climate models, *Eos Trans. Am.*
 757 *Geophys. Union*, 95, 453–471, <https://doi.org/https://doi.org/10.1002/2014EO490002>, 2014.

758 Schaefer, J. T.: The critical success index as an indicator of warning skill, *Weather Forecast.*, 5, 570–575,
 759 [https://doi.org/https://doi.org/10.1175/1520-0434\(1990\)005%3C0570:TCSIAA%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0434(1990)005%3C0570:TCSIAA%3E2.0.CO;2), 1990.

760 Schneider, R. S., Schaefer, J. T., and Brooks, H. E.: Tornado outbreak days: An updated and expanded
 761 climatology (1875–2003), 22nd Conference on Severe Local Storms, 2004.

762 Sherburn, K. D., Parker, M. D., King, J. R., and Lackmann, G.: Composite environments of severe and
763 nonsevere high-shear, low-CAPE convective events, *Weather Forecast.*, 31, 1899–1927,
764 <https://doi.org/https://doi.org/10.1175/WAF-D-16-0086.1>, 2016.

765 Soulard, N., Lin, H., and Yu, B.: The changing relationship between ENSO and its extratropical response
766 patterns, *Sci. Rep.*, 9, <https://doi.org/https://doi.org/10.1038/s41598-019-42922-3>, 2019.

767 Strader, S. M., Gensini, V. A., Ashley, W. S., and Wagner, A. N.: Changes in tornado risk and societal
768 vulnerability leading to greater tornado impact potential, *Npj Nat. Hazards*, 1,
769 <https://doi.org/https://doi.org/10.1038/s44304-024-00019-6>, 2024.

770 Thompson, D. W. J. and Wallace, J. M.: Annular Modes in the Extratropical Circulation. Part 1: Month-to-
771 month Variability, *J. Clim.*, 13, 1000–1016, [https://doi.org/https://doi.org/10.1175/1520-0442\(2000\)013%3C1000:AMITEC%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0442(2000)013%3C1000:AMITEC%3E2.0.CO;2), 2000.

773 Thompson, R. L., Edwards, R., and Hart, J. A.: Close proximity soundings within supercell environments
774 obtained from the rapid update cycle, *Weather Forecast.*, 18, 1243–1261,
775 [https://doi.org/10.1175/1520-0434\(2003\)018%3C1243:CPSWSE%3E2.0.CO;2](https://doi.org/10.1175/1520-0434(2003)018%3C1243:CPSWSE%3E2.0.CO;2), 2003.

776 Tippett, M. K., Lepore, C., and L’Heureux, M. L.: Predictability of a tornado environment index from El
777 Nino Southern Oscillation (ENSO) and the Arctic Oscillation, *Weather Clim. Dyn.*, 3, 1063–1075,
778 <https://doi.org/https://doi.org/10.5194/wcd-3-1063-2022>, 2022.

779 Tippett, M. K., Malloy, K., and Lee, S. H.: Modulation of U.S. tornado activity by year-round North
780 American weather regimes, *Mon. Weather Rev.*, 152, 2189–2202,
781 <https://doi.org/https://doi.org/10.1175/MWR-D-24-0016.1>, 2024.

782 Trapp, R. J.: Mesoscale-convective processes in the atmosphere, Cambridge University Press, 377 pages,
783 <https://doi.org/10.1017/CBO9781139047241>, 2013.

784 Trapp, R. J.: On the significance of multiple consecutive days of tornado activity, *Mon. Weather Rev.*,
785 142, 1452–1459, <https://doi.org/https://doi.org/10.1175/MWR-D-13-00347.1>, 2014.

786 Vitart, F., Ardilouze, C., Bonet, A., Brookshaw, A., Chen, M., Codorean, C., Deque, M., Ferranti, L., Fucile,
787 E., Fuentes, M., Hendon, H., Hodgson, J., Kang, H.-S., Kumar, A., Lin, H., Liu, G., Liu, X., Malguzzi, P.,
788 Mallas, I., Manoussakis, M., Mastrangelo, D., MacLachlan, C., McLean, P., Minami, A., Mladek, R.,
789 Nakazawa, T., Najm, S., Nie, Y., Rixen, M., Robertson, A. W., Ruti, P., Sun, C., Takaya, Y., Tolstykh, M.,
790 Venuti, F., Waliser, D., Woolnough, S., Wu, T., Won, D.-J., Xiao, H., Zaripov, R., and Zhang, L.: The
791 subseasonal to seasonal (S2S) prediction project database, *Bull. Am. Meteorol. Soc.*, 98, 163–173,
792 <https://doi.org/https://doi.org/10.1175/BAMS-D-16-0017.1>, 2017.

793 Wallace, J. M. and Gutzler, D. S.: Teleconnections in the geopotential height field during the Northern
794 Hemisphere winter, *Mon. Weather Rev.*, 109, 784–812, [https://doi.org/https://doi.org/10.1175/1520-0493\(1981\)109%3C0784:TITGHF%3E2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0493(1981)109%3C0784:TITGHF%3E2.0.CO;2), 1981.

796 Wyburn-Powell, C. and Jahn, A.: Large-scale climate modes drive low-frequency regional Arctic sea ice
797 variability, *J. Clim.*, 37, 4313–4333, <https://doi.org/https://doi.org/10.1175/JCLI-D-23-0326.1>, 2024.

798 Zhang, W., S-Y Wang, S., Chikamoto, Y., Gillies, R., LaPlante, M., and Hari, V.: A weather pattern
799 responsible for increasing wildfires in the western United States, *Environ. Res. Lett.*, 20,
800 <https://doi.org/10.1088/1748-9326/ad928f>, 2024.

801 Zhao, J., Bai, L., Zhou, B., and Zhang, L.: Contributions of Interdecadal Pacific Oscillation, Atlantic
802 Multidecadal Oscillation, and global ocean warming to the secular change in United States tornado
803 occurrence, *Atmospheric Res.*, 331, <https://doi.org/https://doi.org/10.1016/j.atmosres.2025.108689>.,
804 2025.

805

806