

Different paths, same destination: similar benthic nitrogen cycling sustained by distinct biogeochemical mechanisms in artificialized coastal systems

Ludovic Pascal¹, Éléonore Dansereau-Macias¹, Carole-Anne Guay¹, David Didier², Christian Nozais³,
5 Luc Sirois³, Faten Zouaghi², & Gwénaëlle Chaillou¹

¹ Institut des Sciences de la Mer de Rimouski (ISMER), Université du Québec à Rimouski, 310 Allée des Ursulines, Rimouski, QC, G5L 3A1, Canada

² Northern and Arctic Coastal Research Lab., Center for Northern Studies, Département de biologie, chimie et géographie, Université du Québec à Rimouski, 300 Allée des Ursulines, Rimouski, QC, G5L 3A1, Canada

10 ³ Département de biologie, chimie et géographie, Université du Québec à Rimouski, 300 Allée des Ursulines, Rimouski, QC, G5L 3A1, Canada

Correspondence to: Ludovic Pascal (Ludovic_pascal@uqar.ca)

Abstract. Human activities are increasingly affecting coastal ecosystems, with shoreline stabilization structures becoming a prevalent response to sea-level rise and extreme climatic events. While these structures aim to protect coastal communities and
15 infrastructures, their effects on ecosystem functioning, and particularly nitrogen cycling and its removal processes such as denitrification, remain poorly understood. To assess the ecological impacts of breakwater construction, we employed a space-for-time substitution approach to examine changes in sediment biogeochemistry, macrobenthic community structure, and nitrogen cycling in an intertidal salt marsh in the St. Lawrence Estuary, Canada. We measured benthic fluxes, denitrification rates, and macrofaunal assemblages at five locations: two landward of the breakwater (impacted sites), one situated
20 immediately seaward of the breakwater (intermediate site) and two reference sites (one vegetated and one unvegetated outside the breakwater system). Landward sites exhibited finer sediment with higher organic carbon content compared to other sites and supported distinct macrobenthic communities dominated by opportunistic Oligochaetes while reference sites were dominated by Molluscs and Crustaceans. The intermediate seaward site closely resembled the unvegetated reference site in both sediment characteristics and oxygen dynamics. Surprisingly, despite these substantial physical and biological changes,
25 dark total benthic oxygen uptake, nitrate and ammonium benthic fluxes and benthic denitrification rates showed no significant differences between impacted and reference sites. This functional similarity reflects an apparent functional convergence, where distinct combinations of abiotic and biotic factors can sustain similar ecosystem functions. However, this convergence might not apply to all ecosystem services offered by natural salt marshes and may depend on specific biological communities, emphasizing the importance of careful consideration in coastal management decisions.

30

1 Introduction

Coastal areas provide essential ecological, economic, and social services. They host various ecosystems and support fisheries, aquaculture, recreational and cultural activities. Around 10% of the global population lives in coastal areas less than 10 meters above sea-level, with coastal development rising in recent decades, a trend expected to continue (Neumann et al., 2015).

35 However, these areas face significant threats from sea-level rise and shoreline erosion (Mentaschi et al., 2018). For instance, a global temperature rise of 2°C above pre-industrial temperatures could lead to a global sea-level rise exceeding 0.2 m by 2040 (Jevrejeva et al., 2016). The most common approach to these threats is shoreline hardening (Gittman et al., 2015) which involves the construction of rigid, engineered structures to stabilize shorelines and protect infrastructures. The main types of shorelines hardening structure include bulkheads, seawalls and breakwaters. Bulkheads are vertical walls typically deployed
40 in environments with low to moderate hydrodynamic energy, built directly on the shoreline to retain soil and prevent land loss due to wave action. Seawalls are robust, vertical or sloped structures designed to protect the shoreline in high-energy environment by reflecting and/or dissipating wave energy and preventing coastal flooding and erosion. Breakwaters are offshore, barrier-like structures built nearly parallel to the coast to dissipate wave energy and reduce current velocity before the waves reach the shore. While effective at protecting human settlements, these structures artificialize coastal ecosystems by
45 fundamentally modifying natural habitats, leading to significant socio-ecological changes such as loss of biodiversity and reduction of the intertidal habitat (Bernatchez and Fraser, 2012; Gittman et al., 2016).

Salt marshes are important coastal ecosystems developing in low-energy shorelines, in temperate and cooler regions. They provide some of the most valuable ecosystem services (Barbier et al., 2011). These intertidal habitats play a crucial role in coastal protection by stabilizing sediments, attenuating wave energy, and enhancing sediment deposition, thereby helping to
50 mitigate coastal erosion (Duarte et al., 2013; Gedan et al., 2009). Due to their high productivity and the capacity to enhance sediment deposition, salt marshes contribute to long-term carbon sequestration (Duarte et al., 2013), provide habitat for numerous organisms (Barbier et al., 2011), and play a significant role in nutrient cycling, particularly by acting as nitrogen sinks (Sousa et al., 2008; Valiela and Cole, 2002). This function has gained increasing importance in recent decades with the rise of coastal eutrophication (Pascal et al., 2022; Rabalais et al., 2009). Eutrophication, the excessive supply of water bodies
55 with organic matter (Nixon, 1998), largely driven by nitrogen enrichment, can lead to significant ecosystem degradation (Rabalais et al., 2009).

In this context, understanding nitrogen removal processes in salt marshes is crucial when assessing the impacts of shoreline hardening on coastal ecosystem functioning (Sousa et al., 2008; Valiela and Cole, 2002). Shoreline hardening can alter sediment characteristics, hydrodynamic conditions, and benthic communities (Gittman et al., 2016), all of which influencing
60 nitrogen cycling processes (Albert et al., 2021; Stief, 2013). Two key microbial processes drive nitrogen removal from soft-sediment systems. (i) Denitrification which involves the oxidation of organic matter (OM) using nitrate (NO_3^-) as an electron acceptor under anaerobic conditions, resulting mainly in the production of dinitrogen gas (N_2), but can also release as nitrous oxide (N_2O), a potent greenhouse gas, together with carbon dioxide (CO_2). (ii) Anammox (anaerobic ammonium oxidation),

produces N_2 through the oxidation of ammonium (NH_4^+) coupled to the reduction of nitrite (NO_2^-) although in many organic-rich coastal sediments its contribution can be smaller than that of heterotrophic denitrification (Dalsgaard et al., 2005; Devol, 2015). In addition to denitrification and anammox, dissimilatory nitrate reduction to ammonium (DNRA) can also occur, competing for nitrate and retaining nitrogen within the system. These processes are further influenced by benthic macrofauna, which play key roles in OM mineralization and nitrogen cycling through bioturbation (*sensu* Kristensen et al., 2012), enhancing solute exchanges and nutrient regeneration at the sediment-water interface (Pascal et al., 2016; Snelgrove et al., 2018). Therefore, alterations to coastal environments through shoreline hardening could significantly impact these nitrogen removal pathways, with potential consequences for coastal water quality and eutrophication mitigation. Given the vulnerability of salt marshes to such modification, understanding how nitrogen cycling is impacted is critical. Salt marshes face increasing threats worldwide (Mcowen et al., 2017) including shoreline hardening (Gedan et al., 2009) that can exceed 50% of the total shoreline length in some regions (Bulleri and Chapman, 2010; Gittman et al., 2015). This extensive shoreline hardening contributes to a phenomenon known as coastal squeezing (Doody, 2013) where natural habitat are trapped between rising seas and hardened shoreline, resulting in habitat loss and degradation (Bernatchez and Quintin, 2016; Bozek and Burdick, 2005). Previous research on bulkhead implementation found no direct effects on nitrogen removal rates per unit area, but showed that these structures can reduce the overall nitrogen removal capacity of the ecosystem by decreasing the total area of salt marshes through coastal squeezing, rather than impacting nitrogen removal process rates (O'Meara et al., 2015). However, breakwaters differ from bulkheads as they are built offshore. When emergent and built with sufficient vertical distance between the crest of the breakwater and the high-tide level, breakwater create low-energy environments on their landward side (Wang et al., 2021). These conditions promote the sedimentation of fine, organic-rich particles (Gittman et al., 2016), which can substantially alter macrobenthic communities, and biogeochemical processes. Such changes may enhance favourable conditions for denitrification and anammox in the sediment, thereby increasing nitrogen removal rates (Albert et al., 2021; Stief, 2013). Yet, the extent and underlying mechanisms of these impacts remain poorly understood, limiting our ability to predict how different shoreline hardening strategies will affect nitrogen removal processes and associated ecosystem services. This study aims to compare sedimentary biogeochemistry and nitrogen cycling across contrasting areas within an intertidal salt marsh seascape, including areas dominated by salt marsh vegetation, unvegetated sediments, and areas located directly seaward and landward of a breakwater. We hypothesize that the presence of the breakwater causes organic-rich sediment accumulation on its landward side, which alters macrobenthic community structure, increases the sediment oxygen demand and the rates of nitrogen removal processes compared to areas farther from the breakwater.

2 Methods

2.1 Study site

The St. Lawrence Estuary area (Québec, Canada) has already seen an accelerated sea-level rise since the early 1990s (Gibeault et al., 2016), with anticipated relative sea-level rise to 0.67 m by 2100 (Savoie-Ferron et al., 2021), threatening infrastructure

and human populations with erosion and flooding. As in many regions worldwide, coastal hardening has been implemented in response to these threats; recent estimates indicate that 10% of the Estuary and Gulf of St. Lawrence coastline is now artificialized by rigid structure (Sauvé et al., 2020). Among these, the Rivière-du-Loup embayment (47°50'29.30" N; 69°33'11.84" W; WSG 84), located on the south shore of the Upper St. Lawrence Estuary (Fig. 1), provides a representative example of coastal hardening.




Figure 1: Study site. (a) Location of the study area (open square) in the St. Lawrence system, (b) location of the study site (black rectangle) in the bay of Rivière-du-Loup and (c) aerial view of the sampling location around the breakwater (from Zouaghi et al., 2024) with saltmarsh areas highlighted by yellow contours. Reference sites: Vegetated Sediment (VS), and Unvegetated Sediment (US); Impacted sites: Southern area landward of the breakwater (LBWS), Northern area landward of the breakwater (LBWN); and Intermediate site: Seaward of the breakwater (SBW). (a) © Stamen Design (CC BY 4.0) / © Stadia Maps; © OpenStreetMap contributors (ODbL), (b) © Gouvernement du Québec (CC BY 4.0; MNRF 2025).

The Rivière-du-Loup embayment receives freshwater inputs from a river located in its northern part, with a mean annual discharge of $14.7 \text{ m}^3 \text{ s}^{-1}$ (from 3.4 to $141.1 \text{ m}^3 \text{ s}^{-1}$) and mean annual concentrations of 2.7 , 13.7 and $0.5 \text{ } \mu\text{mol L}^{-1}$ for NH_4^+ , NO_3^- and PO_4^{3-} , respectively. This bay is covered by sea ice during winter and is characterized by a semi-diurnal tide with an amplitude between 0.5 and 3.2 m , and a salt marsh partially covers its intertidal area with cordgrass (*Sporobolus alterniflorus*, formerly known as *Spartina alterniflora*) in the high foreshore. The shoreline has experienced erosion rates ranging from 2.3

to 6.4 m y⁻¹ since 1984 (Dionne, 2007), endangering a major highway in addition to reducing the salt marsh area by approximately 62,750 m² (Zouaghi et al., 2024). In 2009, the Quebec Government implemented a 300 m-wide breakwater made of boulders to mitigate erosion. It was estimated that the construction of the breakwater *per se* induced a loss of 6,318 m² of salt marsh cover between 2009 and 2011, which has not been compensated for since (Zouaghi et al., 2024). A distinctive feature of this breakwater is that the access path for its construction (perpendicular and towards the center of the breakwater; Fig. 1C) was never removed and now acts as a jetty that is only submerged during high tide events (Zouaghi et al., 2024). In 2010, following the implementation of the breakwater, a storm event coupled with high tides transported and deposited a substantial amount of sediment, which became trapped landward of the breakwater.

2.2 Experimental design

To assess the impacts of the breakwater structure on ecosystem functioning and nitrogen cycling, we adopted a space-for-time approach comparing sites with and without breakwater influence within the same coastal system. This approach assumes that spatial differences in biogeochemical processes and ecosystem functioning reflect the long-term impacts of the breakwater. Five sites were selected to assess how the breakwater affects sediment characteristics and biogeochemical processes within the intertidal zone. (Fig. 1). Two unvegetated sites located landward of the breakwater, the southern (LBWS) and northern (LBWN) areas, were classified as “impacted”. Two other sites, a vegetated (VS) and an unvegetated (US) sediment areas served as “reference” conditions. A site located seaward of the breakwater (SBW) was designated as “intermediate” to help to distinguish the breakwater’s effects from natural environmental gradients between reference and impacted sites. We acknowledge however that the SBW site could also capture potential offshore effects of the breakwater.

2.3 Seawater, sediment and macrobenthic community sampling procedure

Sampling was performed in June 2021 and 2022, targeting the beginning of the salt marsh growing season following winter ice cover break-up. This period was chosen because salt marsh plants are entering their active growth phase, during which root oxygen release and associated rhizosphere processes are expected to influence sediment nitrogen cycling (Koop-Jakobsen and Giblin, 2010; Zheng et al., 2016). Additionally, the relatively short leaf canopy at this stage allows intact sediment cores with vegetation to be incubated without cutting shoots and associated artefacts. Sea water temperature and salinity during sampling were similar between years: 9 °C and 19.6 in 2021 and 10 °C and 18.7 in 2022. In this region, winter ice cover and associated ice scouring can drastically reduce plant cover and height as already reported for *Zostera marina* (Pascal et al., 2020). A 20 L volume of seawater was collected at high tide on the sampling day in the bay. This seawater was filtered on-site using a GF/F filter and aerated in an opaque container for later use. At each of the five sites, eight large sediment cores (10 cm diameter, ~ 15 cm of sediment and ~ 15 cm of air headspace) were manually collected at low tide using transparent acrylic liners (30 cm long), with compaction < 0.5 cm. The sediment surface was immediately capped with Styrofoam® to prevent disturbance during transport. Additionally, five small sediment cores (2.5 cm diameter, 1 cm long) were collected at each site for analysis of surficial sediment characteristics (see section 2.4), and stored at – 20 °C until analysis. Six of the eight large sediment cores

145 were kept intact and immediately transported to the laboratory to assess total benthic fluxes. Of the two remaining cores, one was used to measure the vertical distribution of the dissolved O_2 concentrations at the sediment-water interface and subsequently sliced for particle phase analysis, while the other was dedicated for porewater extraction. To assess macrobenthic community structure, sediment from three box corers ($40 \times 40 \times 15$ cm depth) was collected at each site. This sediment was sieved through a 0.5 mm mesh and preserved in a 4% buffered formaldehyde solution for further analysis. Due to sampling
150 failure at SBW, total benthic fluxes were assessed on only four sediment cores, and no isotope pairing incubation, porewater extraction, or macrofauna sampling were performed at this site. To address this, a June 2022 campaign was conducted to (i) perform O_2 microprofiles at SBW and (ii) increase replication of sediment characteristics at all site. Cross-validation of sediment characteristics and O_2 microprofiles between 2021 and 2022 confirmed the temporal stability of spatial gradients.

2.4 Characterization of the sediment

155 Sediment characteristics were analyzed using samples from the top layer (0 - 0.5 cm) of the five small sediment cores collected at each location. One core was dedicated to measuring sediment grain size (D_{50}), porosity (ϕ), and organic carbon (C_{org}) and total nitrogen (TN) contents, as well as C_{org} natural isotope composition ($\delta^{13}C_{org}$). From this core, D_{50} was assessed by laser diffraction (Malvern Instruments[®], 2 μ m detection limit) after OM digestion with hydrogen peroxide and porosity was determined by measuring water loss after freeze-drying, corrected for sea salt content, using a dry sediment density of 2.65 g
160 cm^{-3} . Prior the C_{org} and $\delta^{13}C_{org}$ analyses, inorganic carbon was removed by fuming of sediments in a closed container in the presence of an open beaker of concentrated HCl. Elemental characteristics of the OM, (i.e., C_{org} , TN) of the sediment were determined using an elemental analyzer (Carlo Erba[®] NC2500). The $\delta^{13}C_{org}$ was then measured using a Micromass Isoprime 100 isotope ratio mass spectrometer coupled to an Elementar Vario MicroCube elemental analyzer. Carbon isotope ratios are reported using the delta notation, relative to the Vienna Pee Dee Belemnite standard. The remaining sediment from this first
165 core, along with the other four cores, was used to determine the OM content via loss on ignition (LOI, n=5), as the weight loss after combustion at 550 °C during 4 h. Due to storage failure, LOI was not assessed at US.

2.5 Oxygen and nutrient porewater concentrations

A 500 mL volume of aerated seawater was gently added at the top of the two cores and the overlying water was then aerated. The vertical oxygen distribution within the sediment was measured within an hour of sampling using Clark-type
170 microelectrodes in one core. The core was maintained at in situ temperature (i.e., 9 °C) using ice packs and the overlying water was stirred and kept at 100% of air saturation by gently bubbling it during all the following operations. For each location, we conducted five depth profiles of O_2 concentration, randomly distributed within a single large sediment core. Microelectrodes (OX100, Unisense[®] with a repeatability better than 2% as specified by the manufacturer) were inserted into the sediment at 100 μ m intervals using a motor-driven micromanipulator (MU1, Pyroscience[®]). A two-point linear calibration of the electrodes
175 was achieved from measurements of the O_2 concentration of air bubbled overlying water (100% saturation) and the anoxic

zone of the sediment (0%). Oxygen diffusive fluxes (DOU in mmol m⁻² h⁻¹) were computed from the O₂ gradient between 0 and 500 µm depth below the sediment-water interface using Fick's first law of diffusion, as denoted in the following equation:

$$\text{DOU} = \phi \times \frac{D_0}{(1-\ln(\phi))^2} \times \frac{\delta C}{\delta z} \quad (1)$$

180 where ϕ is the sediment porosity, D_0 the diffusion coefficient (m² h⁻¹) and $\frac{\delta C}{\delta z}$ is the concentration gradient (mmol m⁻³ m⁻¹).

Oxygen penetration depth (OPD) was determined from the micro-profiles as the depth at which O₂ concentration was < 1% of the concentration at the sediment-water interface. Porewater extraction was performed on another large core (< 2 h after the core recovery) using CSS rhizons (Rhizosphere Research Products®). Rhizons allow the extraction of porewater under a closed vacuum, minimizing the contact with atmospheric oxygen. After removing the overlying water to minimize potential
185 contamination of surface porewater samples, Rhizons were horizontally inserted in the sediment core at 0.5, 1, 1.5, 2, 3, 4, 5, 7, and 10 cm depth. To avoid significant overlap between successive samples, porewater extraction was limited to 8 mL, we cannot exclude that nutrient profiles in the uppermost cm of the sediment column are slightly smoothed due to limited smearing from adjacent layers. After extraction, porewater samples were stored at – 20 °C until nutrient analyses.

2.6 Reactive Fe oxyhydroxide concentrations

190 The reactive fraction of iron-oxyhydroxides was extracted from homogenized, freeze-dried sediment samples using a buffered ascorbate solution. The term “reactive” refers to the easily reducible fraction of Fe oxides, typically amorphous Fe(III) oxides (Hyacinthe et al., 2001; Kostka and Luther, 1994). The ascorbate reagent consisted of 10 g of sodium citrate and 10 g of sodium bicarbonate dissolved in 200 ml of Nanopure® water, with 4 g of ascorbic acid added to reach pH 8 (Kostka and Luther, 1994). Approximately 250 mg of ground, freeze-dried sediment was digested in 15 ml of the ascorbate reagent. After 24 hours of
195 shaking at room temperature, the sediment slurries were centrifuged, and a fraction of the supernatant was diluted with a 1 M HNO₃ solution before Fe was analyzed using Microwave Plasma Atomic Emission Spectroscopy (4210 MP-AES, Agilent®). Fe_{asc} concentrations were corrected for sediment porosity and adjusted for salinity, with a measurement reproducibility better than 2%.

2.7 Intact sediment core incubations and isotope pairing experiments

200 For vegetated sediment cores containing *S. alterniflorus*, the relatively short leaf length at the beginning of the growing season allowed the vegetation to fit entirely within the core dimensions without modification. Six sediment cores were fitted with PVC lids equipped with a magnetic stirrer. The dissolved O₂ concentration was measured in real time using an optode (O₂ sensor SP; PreSens® with a repeatability better than 0.5% as specified by the manufacturer) connected to a multichannel O₂ meter (OXY-10 SMA; PreSens®) through optical fibre cables. Incubations were performed in the dark at the in situ water
205 temperature (i.e., 9 °C). Prior to the initial incubation, the overlying water in the sediment cores was carefully filled with previously collected and bubbled seawater, and the cores were left open and bubbled overnight for acclimation. The following

day, the overlying water was completely replaced with fresh, previously collected and bubbled seawater, and left open while being gently bubbled with air for two hours before starting the experiment. The incubation duration was adjusted so that O₂ saturation in the overlying water never dropped below 80% of its initial value (typically between 1.5 and 4 h). During the incubations, discrete water samples were collected each hour, filtered (0.2 µm) and stored at - 20 °C for subsequent nutrient analyses. Water removed by sampling was automatically replaced with previously collected and bubbled seawater. Overall, cumulative volume of replacement waters corresponded to <5% of the overlying water and its dilution effect was thus considered as negligible. After the initial incubation period, two thirds of the overlying water volume in each sediment core was replaced with previously collected and bubbled seawater. The sediment cores were then left open, and the overlying water was gently bubbled with air for two hours. Then, an initial water sample was extracted from the overlying water of all six sediment cores to determine the ambient NO₃⁻ concentration before the addition of ¹⁵N - NO₃⁻ stable isotope. A calculated volume of a 20 mM ¹⁵N - NO₃⁻ solution (99 % atom ¹⁵N, Sigma-Aldrich®) was then introduced into the overlying water of each core. The added volume was designed to achieve a nominal concentration equivalent to either 50% or 100% of the ambient concentration, with two sets of three cores. A 3 h pre-incubation period was implemented to ensure sufficient diffusion of ¹⁵N - NO₃⁻ to the sediment nitrate reduction zone (Nielsen, 1992), including shallow rhizosphere and bioturbated layers. During this period, an air bubbling system was used to keep the overlying bottom-water homogeneously saturated with air, maintaining the in situ conditions. Following the pre-incubation, water samples were collected to determine the exact concentration of ¹⁵N - NO₃⁻ and the initial ²⁹N₂ and ³⁰N₂ concentrations. Then, the incubations were performed as described above. Discrete water samples were collected using 12 mL Exetainers and directly analysed for ²⁹N and ³⁰N measurements. At the end of the incubations, the remaining water was mixed with the top 5 cm of the sediment column. From the slurry, 12 mL of sample was transferred to an Exetainer to which 200 µL of 7 M ZnCl₂ were added for further ¹⁵N - NH₄⁺ analysis to assess DNRA rates.

For porewater sample, NO₃⁻ (quantification limit 0.6 µM), NO₂⁻ (quantification limit 0.05 µM), NH₄⁺ (quantification limit 0.1 µM) and PO₄³⁻ (quantification limit 0.1 µM) concentrations were analyzed using an autoanalyzer AA500 (Seal analytical®) following Aminot et al. (2009). For water incubation samples, where sample volume was limited to avoid dilution due to replacement water, NH₄⁺ was determined manually according to Hansen and Koroleff (1999), while NO_x were analyzed by continuous flow analysis following Anderson (1979) both with measurements reproducibility better than 5%. Analytical accuracy was assessed using certified reference materials (MOOS-4, NRC Canada).

The isotope ²⁸N₂, ²⁹N₂ and ³⁰N₂ measurements were completed within one week of collection using a membrane inlet mass spectrometer (MIMS, Bay Instruments®) (Kana et al., 1994). The protocol followed Kana et al. (2006). Calibration factors for standard sample were computed based on solubility equations (Hamme and Emerson, 2004). Analytical precision for N₂/Ar ratios (coefficient of variation of five replicates standards) was consistently < 0.5%. To quantify the production of ¹⁵N - NH₄⁺, the water samples were first gently bubbled with He for 5 min to remove all ¹⁵N - N₂ produced during the incubation. The

water samples were then treated with 200 μL of 2 M KCl to desorb any particle-bound NH_4^+ followed by the addition of
 240 alkaline hypobromite iodine to oxidize NH_4^+ to N_2 prior to analysis using MIMS as described in Yin et al. (2014).
 The slope (in $\text{mmol m}^{-3} \text{ h}^{-1}$) from the linear regression of the changes in O_2 , nutrients and $^{29,30}\text{N}_2$ concentrations over
 incubation duration was used to compute benthic fluxes as follows:

$$J_{\text{benthic}} = \text{slope} \times \frac{V}{A} \quad (2)$$

245

where J_{benthic} is the flux across the sediment-water interface ($\text{mmol m}^{-2} \text{ h}^{-1}$), V is the volume of overlying water (m^3), and A is
 the surface of the sediment-water interface (m^2).

The validity of the original isotope pairing technique (Nielsen, 1992) was tested by checking the independence of the estimated
 $^{28}\text{N}_2$ production and the amount of added $^{15}\text{N} - \text{NO}_3^-$ (Risgaard-Petersen et al., 2003). No production of $^{15}\text{N} - \text{NH}_4^+$ was
 250 detected over the incubation periods, DNRA was thus considered to be negligible. Since no deviation from the original isotope
 pairing technique was detected, anammox was also considered negligible. Both processes are therefore not considered further
 in the manuscript and denitrification rates were computed on the six sediment cores following Nielsen 1992:

$$D_{15} = p29 + 2 \times p30 \quad (3)$$

255 where D_{15} is the denitrification rate of the $^{15}\text{N} - \text{NO}_3^-$ and $p29$ and $p30$ the production of $^{29}\text{N}_2$ and $^{30}\text{N}_2$, respectively.

$$D_{14} = D_{15} \times (p29/2p30) \quad (4)$$

where D_{14} is the denitrification rate of $^{14}\text{N} - \text{NO}_3^-$. From D_{14} and D_{15} , the denitrification of nitrate diffusing from the overlying
 260 water to the nitrate reduction zone (D_w) and the denitrification of nitrate produced within the sediment through nitrification
 (*i.e.*, coupled nitrification-denitrification; D_n) were calculated as followed:

$$D_w = (^{14}\text{NO}_3^- / ^{15}\text{NO}_3^-) \times D_{15} \quad (5)$$

$$D_n = D_{14} - D_w \quad (6)$$

265

where $^{14}\text{NO}_3^-$ is the ambient nitrate concentration and $^{15}\text{NO}_3^-$ is the concentration of labelled nitrate added to the overlying
 water.

2.8 Macrofauna identification

Macrofauna (epi- and infauna) was sorted and identified to the lowest possible taxonomic level (typically species level).
 270 Biomass (blotted wet mass; $\pm 0.0001 \text{ g}$) was determined as well. Using this macrofauna dataset, total density, total biomass

and species richness were calculated. In this study, we identified six biological traits characterized by multiple categories to describe the organisms' functional behaviors (Table 1). Taxa were assigned trait modalities using a standard fuzzy coding approach which allowed each taxon to represent multiple trait modalities and thus account for intraspecific variation. Scoring ranges from 0 to 3 were adopted with 0 corresponding to no affinity for a given trait modality, 1 and 2 reflecting a low and high affinity, respectively, and 3 an exclusive affinity. Information on trait expression for all taxa were extracted from Queiros et al. (2013) and Degen and Faulwetter (2019). When information was not available for a given taxon, information was obtained from that of closely related species at the nearest taxonomic level. The community trait expression in each sample was computed from the multiplication of the species density matrix by trait categories matrix.

Table 1: *Biological traits.* List of traits, trait categories and associated abbreviations.

Trait	Categories	Abbreviation
Size	Small (< 1cm)	S_SMA
	Small-medium (1-5 cm)	S_SMM
	Medium (5-10 cm)	S_MED
	Medium-large (10-30 cm)	S_MDL
Mobility	Low	M_LOW
	Medium	M_MD
	High	M_HI
Bioturbation	None	B_NO
	Surface mixing	B_SRM
	Diffusive mixing	B_DIF
Environmental position	Epifauna	P_EPI
	Infauna	P_IN
Feeding Habit	Filter feeder	F_FIL
	Surface deposit feeder	F_SRD
	Subsurface deposit feeder	F_SBD
	Predator	F_PRE
	Scavenger	F_SCA

2.9 Data analyses

Data analyses were performed using R software (R Core Team, 2024). Data values are given as means $\pm$ SD. First, linear models (lm function from *stats* package; R Core Team 2024) were fitted to test the effects of *Site* (fixed factor; five levels) on the total macrofauna density, biomass and species number, DOU, OPD, as well as O₂, nutrients and N₂ total benthic fluxes.

285 Test assumptions were assessed for each model (plotQQunif function from *DHARMA* package; Hartig 2022). Effect sizes were calculated as generalized eta squared (η^2) to quantify the magnitude of *Site* effects, with η^2 values of 0.01, 0.06, and 0.14 representing small, medium, and large effects, respectively. Both p-values and effect sizes are reported (Table S2) given the limited replication ($n = 3$ per site), which may affect statistical power. If a significant effect of the *Site* factor was observed ($p < 0.05$), pairwise tests (Tukey contrast) were performed to characterize the differences between its levels. Second, macrofauna

290 density data were transformed using Hellinger transformation. Sites were grouped using a hierarchical clustering analysis with a ward agglomeration method (hclust function from *stat* package; the number of clusters was determined using the HCPC function from *FactoMineR* package; Lê et al. 2008). Patterns of macrobenthic community assemblage were visualized using principal component analysis (PCA, RDA function from *vegan* package; Oksanen et al. 2022). In addition, the difference in macrobenthic community between clusters was tested using a multivariate analysis of variance (anova.cca function from *vegan*

295 package). Finally, functional diversity was assessed using the dbFD function from the *FD* package (Laliberté and Legendre, 2010), computing functional richness (FRic) and functional evenness (FEve). Community-weighted mean trait values (CWM) were calculated and visualized using PCA as described above. Site effects on functional indices were tested using linear models as described above.

2.10 Methodological considerations

300 Sediment characteristics were measured with variable replication levels ($n = 1$ to 5) due to analytical constraints. LOI was replicated five times, some parameters were duplicated, while others were measured once per station. The representativeness of these measurements is supported by two observations: (1) low variability among replicated measurements where available, and (2) strong consistency between highly replicated LOI values and corresponding Corg and TN measurements (duplicate or single measurements).

305 Oxygen microprofiles and porewater nutrients were each measured on a separate single sediment core per location. This approach was chosen because biogeochemical heterogeneity of the sediment is often associated with microenvironments (Stockdale et al., 2009). Such approach is considered qualitatively robust (Bull and Tallefert, 2001). Nevertheless, since this sampling design may involve pseudo-replication, statistical results on these measurements should be interpreted cautiously. Measurement representativeness is supported by: (1) consistency among sediment characteristics, oxygen microprofiling, and

310 porewater nutrient profiles measured in three independent cores per location, and (2) absence of large infauna in cores, limiting potential bias from bioturbation.

Our benthic flux incubations were conducted in the dark to standardize experimental conditions and ensure robust comparisons among sampling sites. This approach inherently excludes potential in situ effects of microphytobenthos or *S. alterniflorus* photosynthesis, both of which can significantly influence sediment oxygenation and nutrient uptake during daylight. However, in intertidal zones near breakwaters, high turbidity and suspended sediment loads, often limit light penetration and thereby reduce photosynthetic activity (Pratt et al., 2014). Moreover, belowground oxygen release by marsh roots correlates with photosynthesis but can persist in darkness (Koop-Jakobsen et al., 2018). As such, our dark incubations represent potential, near-maximum benthic fluxes, under low-light conditions typical of these habitats. While this standardized approach supports reliable site comparisons, we acknowledge that it does not capture the full spectrum of benthic fluxes. We therefore recommend future studies include incubations under natural light to better capture daytime flux dynamics, especially where benthic primary producers are exposed to substantial light during immersion.

3 Results

3.1 Sediment characteristics and biogeochemical properties

Among all sampling locations, surficial sediments consisted of silty clay (Table 2). Overall, impacted sites (LBWN and LBWS) showed finer median grain size compared to reference sites (VS and UV). Correspondingly, sediment organic carbon (OC) and total nitrogen (TN) contents tended to be higher in impacted sites, while VS exhibited the lowest OC and TN contents among sites. The $\delta^{13}\text{C}_{\text{org}}$ signatures remained relatively uniform among all sites, suggesting a common mixture of terrestrial and marine carbon sources through the system. The intermediate sites (SBW) showed similar sediment characteristics to the two reference sites.

Table 2: *Surficial Sediment characteristics.* Median sediment grain size (D_{50} , in μm), porosity (ϕ , volume ratio), loss on ignition (LOI in %), organic carbon content (C_{org} , in %), total nitrogen content (TN, in %), $\delta^{13}\text{C}_{\text{org}}$ signature (in ‰ vs. VPDB). Values are means $\pm$ SD with replication and sampling year (SY) indicated by superscripts (^an = 2, SY: 2021-2022; ^bn = 5, SY: 2021; none : n = 1, SY: 2021).

Site	D_{50}	ϕ	LOI	C_{org}	TN	$\delta^{13}\text{C}_{\text{org}}$
LBWN	12.9 ± 0.9^a	0.83 ± 0.03^a	11.94 ± 2.11^b	4.08 ± 0.14^a	0.38 ± 0.02^a	-24.3 ± 0.0^a
LBWS	15.3 ± 0.6^a	0.78	11.14 ± 2.13^b	3.52	0.28	-24.7
SBW	22.4 ± 0.2^a	0.75	6.51 ± 0.40^b	2.52	0.23	-24.7
VS	20.5 ± 2.6^a	0.65	6.13 ± 0.89^b	1.09	0.10	-24.6
US	26.8 ± 2.5^a	0.81	--	2.72	0.30	-24.5

Oxygen in the sediment column showed contrasting dynamics among sites (Table S1). Diffusive oxygen uptake (DOU) rates tended to be higher at impacted sites compared to intermediate and reference sites (Fig. 2a). Correspondingly, oxygen penetration depths (OPD) were systematically shallower (Fig. 2b). Similarly, porewater nutrient distributions varied among

sites (Fig. 2d). While NO_2^- concentration was near zero throughout the sediment column at all sites, NO_3^- concentration consistently reached its peak at 0.5 cm depth, and then sharply decreased below the detection limit around 4 cm deep in impacted sites and around 6 cm deep in reference sites (Fig. 2d). The peak concentration was particularly high at VS. NH_4^+ and PO_4^{3-} concentration increased with depth with a larger increase at impacted sites compared to reference ones (Fig. 2d). The increase in PO_4^{3-} concentration started from the sediment surface at impacted sites. In comparison, it started deeper (5 cm deep) at US and PO_4^{3-} concentration was near zero along the sampled sediment column at VS (Fig. 2d). Fe_{asc} showed a decreasing trend from the sediment surface to the bottom for LBWN and LBWS, no trend for US and an increase in the first 3.5 cm then a decrease for VS (Fig. 2e).




Figure 2: Porewater solute concentrations. (a) Diffusive O₂ uptake (DOU) by the sediment and (b) O₂ penetration depth (OPD). (c) O₂ concentration microprofile. (d) porewater NO_3^- , NO_2^- , NH_4^+ and PO_4^{3-} concentrations at each site (n = 1 with 5 pseudo replicates for DOU and OPD). (e) Solid-phase iron oxyhydroxide concentration at each site. Different letters indicate significant (p < 0.05, Tuckey contrast) differences among the different levels of Site (a and b)

3.2 Macrobenthic community structure

A total of 2,657 individuals from 12 samples (4 sites × 3 replicates) were sorted, and 14 taxa were identified (Table S2). Total density, biomass and species richness significantly differed among sites (Table S1). However, these univariate metrics do not allow a clear separation between impacted and reference sites; comparisons must instead be made at the site level. Total densities tended to be higher at the sites LBWS than at LBWN, VS and US (Fig. 3). Total biomass was significantly lower at LBWN, LBWS and VS than at US. Finally, species richness was lower at LBWN than at LBWS, VS and US.




Figure 3: *Univariate descriptors of macrofauna communities.* Total biomass, density and species richness at the sampled sites (n = 3 per site). Different letters indicate significant (p < 0.05, Tuckey contrast) differences among the different levels of *Site*.

Based on hierarchical clustering on species density, the sites could be discriminated into three groups (Fig. 4a). Cluster 1 included the impacted sites, cluster 2 corresponded to VS, and cluster 3 corresponded to the US. The statistical significance of these clusters was confirmed by MANOVA (Table S3). The three clusters exhibited distinct macrofaunal assemblages (Figure 4b and c). The impacted sites (cluster 1) were characterized by a high density of Oligochaetes and Polychaete *Hediste diversicolor*, while VS (cluster 2) were characterized by relatively high density of small body grazers (*Gammarus* spp. and Gastropods). US (cluster 3) were distinguished by a relatively high density of the bivalve *Mya* sp. Both clusters 2 and 3 showed notable densities of the bivalve *Macoma balthica* as compared to cluster 1. It is worth noting that while gathered in the same cluster, LBWS showed greater densities of *H. diversicolor* and Oligochaete than LBWN.




Figure 4: *Macrofauna community assemblages.* (a) Hierarchical clustering and (b–c) principal component analyses (PCA) of Hellinger-transformed species densities. Panels show (b) scaling 1, and (c) scaling 2. Only variables representing at least 20 % of the total variation are shown. Abbreviations: MYA = *Mya* sp.; MAC = *Macoma balthica*; GAM = *Gammarus* spp.; OLI = Oligochaeta; GAS = Gastropoda.

Functional richness (FRic) differed significantly among sites (Table S1) and was higher at VS and US, intermediate at LBWS and lower at LBWN (Fig. 5a). Functional evenness (FEve) did not differ significantly among sites, although a large effect size (Table S1) suggests the functional evenness differences may exist but require greater sampling effort to confirm. Overall, FEve tended to be higher at LBWN, VS and US than at LBWS (Fig. 5b).




Figure 5: *Macrofauna community trait expression.* (a) Functional richness (FRic) and (b) functional evenness (FEve); (c–d) PCA of community-weighted mean (CWM). Panels show (b) scaling 1, and (c) scaling 2. Only variables representing at least 20 % of the total variation are shown. See Table 1 for trait abbreviations.

Beyond diversity metrics, the community-weighted mean trait composition revealed contrasting spatial patterns (Fig. 5c). Although LBWN and US differed markedly in species composition, they exhibited partial convergence in trait expression (Fig. 5d). US was characterized by larger, more mobile organisms dominated by filter-feeding, predatory and scavenger trait expressions, while LBWN showed a tendency toward this trait profile. In contrast, LBWS and VS maintained clear segregation in trait expression. VS was characterized by epifauna with minimal or surface-restricted bioturbation, while LBWS trait expression was dominated by small-bodied infaunal organisms characterized by subsurface deposit feeding and biodiffusive bioturbation.

3.3 Benthic fluxes

Total O_2 , NO_3^- , and NH_4^+ fluxes differed significantly among sites (Table S1). Total O_2 and NO_3^- fluxes were always negative, indicating an uptake by the sediment while positive total NH_4^+ flux indicated a release by the sediment (Fig. 6a). Similarly to the macrobenthic community structure results, comparison should be made at the site level. Overall, all benthic fluxes followed the same pattern with higher fluxes at LBWS and VS than at LBWN, SBW and US (Fig. 5a). Total denitrification (D_{tot}) also differed significantly among sites (Table S1). Similarly to benthic fluxes, D_{tot} was highest at sites VS and LBWS and lowest at sites LBWN and US (Fig. 6b). The contribution of sedimentary nitrification to total denitrification (coupled nitrification-denitrification; D_n) was higher than 50% at sites LBWN, LBWS and VS and lower than 10% at site US.




Figure 6: *Benthic fluxes* (a) O_2 , NH_4^+ and NO_3^- fluxes (in $mmol\ m^{-2}\ h^{-1}$) and (b) N_2 flux (D_{tot} in $\mu mol\ m^{-2}\ h^{-1}$) across the sediment-water interface. The dashed bar plots depict the denitrification of nitrate produced within the sediment through nitrification (D_n) and the percentage represents the proportion of D_{tot} explained by D_n (B). Different letters indicate significant ($p < 0.05$, Tuckey contrast) differences among the different levels of *Site*. Note that positive values correspond to an uptake by the sediment, and negative values correspond to a release by the sediment. $n = 6$ per site except at SBW where $n = 4$.

4 Discussion

Shoreline stabilization structures have become increasingly prevalent features of coastal ecosystems worldwide (Gittman et al., 2016). This proliferation is driven by the urgent need to protect coastal communities from the impacts of sea-level rise and intensifying extreme climatic events. While previous studies have documented changes in sediment characteristics and macrobenthic communities associated with coastal hardening (see review by Gittman et al., 2016 and references therein), the influence such structure on biogeochemical processes and nitrogen cycling remains poorly documented (but see O'Meara et al., 2015). Our study addresses this knowledge gap by comparing ecosystem functioning among contrasting habitats created by a breakwater construction within a salt marsh system.

410 4.1 Impact of breakwater construction on benthic habitats

Our study highlights marked alterations in surficial sediment characteristics, with impacted sites (i.e., LBWS and LBWN) accumulating finer and more organic-rich sediment compared to reference and intermediate sites. These observations are consistent with expectations for low-energy depositional environments landward of high-crested breakwaters (Da Silva et al., 2022; van Rijn, 2011; Wang et al., 2021). In contrast, sediments of the intermediate site (SBW) resembled reference sites (i.e.,
415 US and VS), suggesting minimal breakwater influence in this area. The consistent sedimentary $\delta^{13}\text{C}_{\text{org}}$ signature among sites suggested a shared origin of organic carbon, likely a mix of terrestrial and marine inputs typical of intertidal zones. This homogeneity suggests that low-energy hydrodynamic conditions landward of the breakwater enhance deposition of fine, organic-rich particles without selective retention of organic matter from specific sources. Therefore, differences among sampled areas are driven by organic carbon quantity, not origin.

420 Geochemical measurements further support this pattern. Sediment cores from impacted sites showed up to 2.3 times higher diffusive O_2 uptake (DOU) and shallower O_2 penetration depth (OPD) than reference sites (Fig. 2a and b), consistent with enhanced organic carbon mineralization driven by higher organic carbon content (Glud, 2008). These results align well with our porewater nutrient profiles extracted from another sediment core. The decrease in porewater NO_3^- concentration with depth in all sites indicates a consumption by the sediment (Fig. 2d), consistent with our measured NO_3^- benthic fluxes (Fig. 6). This
425 consumption results from the denitrification process, which has been detected at all sites (Fig. 6). VS exhibited a subsurface NO_3^- peak, suggesting active nitrification consistent with rhizosphere oxygenation (Koop-Jakobsen et al., 2018), this pattern is consistent with the relatively high proportion of coupled nitrification-denitrification detected in incubation experiment. This interpretation is further corroborated by the solid-phase ascorbate extractable iron (Fe_{asc}) profiles (Fig. 2e). At VS, Fe_{asc} increase progressively from the surface to a peak at 3-4 cm depth, well below the NO_3^- maximum (Fig. 2d), suggesting
430 subsurface oxidation consistent with rhizosphere oxygenation. The absence of NO_3^- peak co-located with the Fe_{asc} peak suggests a tight coupling between nitrification in the rhizosphere area and denitrification in adjacent anoxic sediment (Koop-Jakobsen et al., 2018; Zheng et al., 2016). The quicker NO_3^- consumption in the porewater of impacted sites in the upper four centimeters compared to six centimeters in reference sites suggests a more reduced condition in impacted sites. Similarly, deeper sediment layers in impacted sites exhibit greater increases in NH_4^+ and PO_4^{3-} compared to reference sites, indicating
435 larger ammonification driven by their higher organic matter content, along with PO_4^{3-} release, typically associated with the reductive dissolution of iron-oxyhydroxides in anoxic sediment layers (Slomp, 2011). Consistently, Fe_{asc} concentrations at impacted sites are highest at the sediment surface and decrease rapidly with depth (Fig. 2e). This indicates that aerobic processes, including nitrification, are restricted to the very surface of the sediment at these sites. Overall, these O_2 and nutrient profiles suggest more reduced conditions and greater nutrient accumulations in the sediment column in impacted sites
440 compared to reference sites. Despite these changes in sedimentary conditions, denitrification rates were not substantially altered, suggesting that the capacity of these sediments to remove excess nitrogen remains resilient to these perturbations.

4.2 Impact of breakwater construction on macrobenthic community structure

Consistent with previous studies on the ecological impact of shoreline stabilization structure (Bulleri and Chapman, 2010; Dugan et al., 2018; Gittman et al., 2016), impacted sites exhibited distinct assemblages compared to reference sites. Impacted sites were notably dominated by high densities of small-bodied *H. diversicolor* and Oligochaetes, the former being tolerant to organic enrichment, and the latter considered opportunistic under such conditions and often associated with fine-grained, organic-rich sediments (Borja et al., 2000). Both *H. diversicolor* and Oligochaetes are significant bioturbators (Queirós et al., 2013) that influence sediment structure, biogeochemistry and microbial activity at the sediment-water interface (Kristensen et al., 2012). In contrast, reference sites supported less dense but more diverse communities in terms of species and function. Notably, US appeared to be colonized by larger organisms than VS, as suggested by species traits analysis (Fig. 5d) and the relatively low total density and high total biomass in US (Fig. 3). This difference may reflect the dense root and rhizome network of *S. alterniflorus* in VS, which could hamper the movement and burrowing of larger infauna (Pascal et al., 2020; van Wesenbeeck et al., 2007). The higher density of biodiffusive bioturbation at impacted sites, and particularly LBWS (Fig. 5c), suggests that some ecosystem processes, such as sediment oxygenation and nutrient regeneration, may persist despite habitat alteration. However, the lower biodiversity and functional richness and evenness, at these sites may limit functional redundancy thereby reducing the resilience of ecosystem functions to additional stressors.

4.3 Potential convergence in ecosystem functioning

Despite marked differences in sediment characteristics, macrobenthic community structure and trait expression between impacted and reference sites (i.e., LBWN vs. US and particularly LBWS vs. VS), total benthic fluxes and denitrification rates remained similar. A similar convergence in process rates despite contrasting biological drivers has been reported in other benthic systems (Woodin et al., 2016). Here, however, it results from distinct mechanisms, namely root-driven oxygenation and bioturbation. These findings, derived from dark incubations (see section 2.10), suggest that distinct benthic environments can yield similar functional outcomes regarding oxygen uptake, nutrient cycling and notably denitrification rates. Our measured denitrification rates ranging from 6 to 75 $\mu\text{mol m}^{-2} \text{h}^{-1}$ fall within the range of reported denitrification rates for temperate intertidal mudflats and salt marshes (Hinshaw et al., 2017; Koch et al., 1992; Poulin et al., 2007; Sousa et al., 2012). Coupled nitrification-denitrification (D_n) was the dominant N-removal pathway at impacted sites and VS, indicating that denitrification relies on locally produced NO_3^- . Conversely, at US, denitrification was primarily driven by water-column nitrate. The high density of subsurface bioturbators at LBWS (Fig. 5c) likely enhances oxygen and nutrient exchange at the sediment-water interface. The sediment irrigation enhances aerobic mineralisation of the OM, stimulates nitrification fuelling coupled nitrification-denitrification, a pattern supported by our isotope pairing results (Fig. 6). In VS, although macrofauna is present, bioturbation is absent or constrained to the sediment surface (Fig. 5c). However, as evidenced by the deep subsurface Fe_{asc} peak (Fig. 2e; Section 4.1), radial O_2 loss from *S. alterniflora* roots actively creates deep oxidized zones (Koop-Jakobsen et al., 2018). This rhizosphere oxygenation sustains subsurface nitrification, which is tightly coupled to denitrification, thereby

achieving high denitrification rates similar to those driven by bioturbation at LBWS. By contrast, LBWN showed a trait
475 expression tendency toward US that could yield similar benthic fluxes. The low denitrification rates at US likely reflect the
relatively low organic carbon availability compared to impacted sites, which limits local nitrification fuelling denitrification.
Moreover, macrofauna at US is dominated by a few large burrowers (e.g., *Mya* sp.) whose local influence may not be captured
at the core scale, leading to low benthic fluxes. Interestingly, despite being both impacted sites and having similar sediment
characteristics, LBWN and LBWS exhibited different ecosystem functions based on total benthic fluxes and denitrification
480 rates. This most likely results from the higher density of macrobenthic organisms at LBWS than at LBWN, highlighting the
critical role of macrobenthic community activity in shaping biogeochemical processes (Snelgrove et al., 2018). In the absence
of dense bioturbators, the reduced sediment at LBWN (Fig. 2d and e) prevents nitrification despite the high availability of
organic carbon, effectively limiting denitrification. However, the factors driving the variation in macrobenthic community
structure between these two areas remain elusive. The absence of a detectable breakwater effect on denitrification rates
485 therefore does not reflect an absence of biogeochemical change, but rather a functional compensation mediated by distinct
biological drivers at each site.

Placed in a broader context, although the impacts of coastal protection structures on benthic nitrogen cycling remain poorly
documented, our results suggest that these effects are structure specific. Indeed, vertical, hard-engineered structures like
seawalls induce coastal squeeze and habitat loss. This reduces the seascape-scale nitrogen-removal capacity, even if unit-area
490 denitrification rates remain unaffected (O'Meara et al., 2015). Conversely, in the present study, our offshore breakwater
accumulates organic-rich sediment. This accumulation, associated with dense populations of bioturbators (e.g., *H.
diversicolor*), sustains a bioirrigation pathway that functionally substitutes for the lost sediment oxygenation by roots. While
this functional equivalence aligns with observations from nature-based protections like living shorelines (Onorevole et al.,
2018; Schoell et al., 2023), our study shows that it can also occur in a completely different, fully artificial environment. Taken
495 together, these results show that even in artificialized settings, nitrogen-removal processes can remain functionally equivalent,
as long as biogeochemical mechanisms capable of coupling nitrification and denitrification are maintained. Whether such
compensatory mechanisms occur in other artificialized systems will depend on local environmental conditions and biological
assemblages.

From a coastal-management perspective, one of the main issues is whether areal flux rates measured in LBWS can offset, at
500 the seascape level, the functional loss caused by salt marsh vegetation removal. LBWS area (~18,800 m²) is about three times
larger than the vegetated area directly lost due to the breakwater implementation (~6,300 m²). Assuming that the measured
fluxes capture the whole area heterogeneity, LBWS could compensate for these specific functional capacity losses of that part
of the salt marsh. However, it is worth noting that this area is still more than three times smaller than the total vegetated area
lost due to combined erosion and breakwater implementation (~62,700 m²) so any functional compensation remains only
505 partial at the scale of the whole seascape.

5 Conclusion

Our study highlights the complex relationships between sediment characteristics, macrobenthic communities, and ecosystem functioning in coastal environments impacted by shoreline hardening structures. We observed that some ecosystem functions (i.e., benthic respiration, nutrient cycling, and denitrification rates) can be maintained through several combinations of abiotic and biotic factors. This finding reflects a convergence in ecosystem functioning that can arise from distinct underlying mechanisms. However, it is important to interpret these results carefully. While we observed functional similarities in specific ecosystem functions, natural coastal habitats provide an extensive collection of ecosystem functions and services (Friess et al., 2020; Gedan et al., 2009) that are far from being fully captured by our measurements. The observed functional convergence may contribute to the resilience of some ecosystem functions at the seascape level. However, these localized similarities do not mean that these functions are unaltered, as the underlying mechanisms are completely modified and potentially more vulnerable to additional environmental pressures. Furthermore, it does not imply that artificialized environments can sustain all ecosystem functions and services provided by natural habitats. These findings highlight the challenges in predicting and managing ecosystem responses to shoreline hardening, coastal squeeze and probably other coastal disturbances. It also emphasizes that physical, geochemical, and biological factors should be considered when assessing the functional impact of coastal development projects. Given the increasing pressures on coastal ecosystems (Halpern et al., 2019), future research should assess the ability of artificialized ecosystems to maintain their functions under multiple stressors and investigate their long-term stability under various climate change scenarios. Finally, broader ecological implications of apparent functional convergence in coastal ecosystems including potential cascading effects on higher trophic levels should be explored. This understanding will be crucial for developing more accurate predictive models and effective management strategies in the face of ongoing coastal development and climate change.

Code and data availability

The data and code supporting this study are archived at the Observatoire global du Saint-Laurent (OGSL) and are accessible at <https://doi.org/10.26071/a974587e-583e-4a2c>.

Supplement link

The link to the supplement will be included by Copernicus, if applicable.

Author contributions

LP: Conceptualization (lead), Methodology (lead), Investigation (lead), Formal analysis (lead), Visualization (lead), Writing - Original Draft Preparation (lead), Writing – Review and Editing (lead); **EDM:** Investigation (supporting), Formal analysis

(supporting), Writing – Review and Editing (supporting); **CAG:** Investigation (supporting), Formal analysis (supporting); **DD:**
535 Funding acquisition (lead) , Writing – Review and Editing (supporting); **CN:** Investigation (supporting), Funding acquisition
(supporting), Supervision (equal), Writing – Review and Editing (supporting); **LS:** Investigation (supporting), Funding
acquisition (supporting), Writing – Review and Editing (supporting); **FZ:** Investigation (supporting); **GC:** Conceptualization
(supporting), Investigation (supporting), Funding acquisition (supporting), Supervision (equal), Writing – Review and Editing
(supporting).

540 **Competing interests**

The authors declare that they have no conflict of interest.

Disclaimer

Copernicus Publications adds a standard disclaimer: “Copernicus Publications remains neutral with regard to jurisdictional
claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While
545 Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.
Views expressed in the text are those of the authors and do not necessarily reflect the views of the publisher.”
Please feel free to add disclaimer text at your choice, if applicable.

Acknowledgements

We acknowledge that this research was conducted on the traditional, ancestral, and unceded territory of the Wolastoqiyik
550 Wampanoag First Nation in the Rivière-du-Loup region (Québec, Canada). We thank Joannie Cool and Carole-Anne Guay for
their assistance and support during the sampling, Marie-Pomme Poissant and Eve Tremblay-Morel for their assistance in
macrofauna identification and Eve Beaudet and Pascal Rioux for their assistance in nutrients analyses. We are grateful to the
GEOTOP facilities (Light stable isotope geochemistry laboratory, UQAM, Montréal, QC, Canada) for the analyses of sediment
characteristics. This project is a contribution to the Québec-Océan and the CFREF Transforming Climate Action research
555 program.

Financial support

This work was financially supported through multiple postdoctoral fellowships from FRQNT (B3X grant numbers 281864
<https://doi.org/10.69777/281864> and 322485; <https://doi.org/10.69777/322485>) and MEOPAR (grant number PDF-25-2020)
to LP. Funding was also provided by the Ministry of Transportation and Sustainable Mobility of Quebec to DD and by NSERC
560 Discovery Grant to CN (DDG-2019-05993), DD (RGPIN-2021-03333) and GC (RGPIN-2018-556538).

Review statement

The review statement will be added by Copernicus Publications listing the handling editor as well as all contributing referees according to their status anonymous or identified.

References

- 565 Albert, S., Bonaglia, S., Stjärnkvist, N., Winder, M., Thamdrup, B., and Nascimento, F. J. A.: Influence of settling organic matter quantity and quality on benthic nitrogen cycling, *Limnology and Oceanography*, 66, 1882–1895, <https://doi.org/10.1002/lno.11730>, 2021.
- Aminot, A., K  rouel, R., and Coverly, S. C.: Nutrients in Seawater Using Segmented Flow Analysis, in: *Practical Guidelines for the Analysis of Seawater*, CRC Press, 408, 2009.
- 570 Anderson, L.: Simultaneous spectrophotometric determination of nitrite and nitrate by flow injection analysis, *Analytica Chimica Acta*, 110, 123–128, [https://doi.org/10.1016/S0003-2670\(01\)83537-6](https://doi.org/10.1016/S0003-2670(01)83537-6), 1979.
- Barbier, E. B., Hacker, S. D., Kennedy, C., Koch, E. W., Stier, A. C., and Silliman, B. R.: The value of estuarine and coastal ecosystem services, *Ecological Monographs*, 81, 169–193, <https://doi.org/10.1890/10-1510.1>, 2011.
- Bernatchez, P. and Fraser, C.: Evolution of Coastal Defence Structures and Consequences for Beach Width Trends, Qu  bec, Canada, *Journal of Coastal Research*, 285, 1550–1566, <https://doi.org/10.2112/JCOASTRES-D-10-00189.1>, 2012.
- 575 Bernatchez, P. and Quintin, C.: Potentiel de migration des   cosyst  mes c  tiers meubles qu  b  cois de l’estuaire et du golfe du Saint-Laurent dans le contexte de la hausse appr  hend  e du niveau de la mer, *natcan*, 140, 91–104, <https://doi.org/10.7202/1036507ar>, 2016.
- Borja, A., Franco, J., and P  rez, V.: A Marine Biotic Index to Establish the Ecological Quality of Soft-Bottom Benthos Within European Estuarine and Coastal Environments, *Marine Pollution Bulletin*, 40, 1100–1114, [https://doi.org/10.1016/S0025-326X\(00\)00061-8](https://doi.org/10.1016/S0025-326X(00)00061-8), 2000.
- 580 Bozek, C. M. and Burdick, D. M.: Impacts of Seawalls on Saltmarsh Plant Communities in the Great Bay Estuary, New Hampshire USA, *Wetlands Ecol Manage*, 13, 553–568, <https://doi.org/10.1007/s11273-004-5543-z>, 2005.
- Bull, D. C. and Taillefert, M.: Seasonal and topographic variations in porewaters of a southeastern USA salt marsh as revealed by voltammetric profiling†, *Geochem Trans*, 2, 104, <https://doi.org/10.1186/1467-4866-2-104>, 2001.
- 585 Bulleri, F. and Chapman, M. G.: The introduction of coastal infrastructure as a driver of change in marine environments, *Journal of Applied Ecology*, 47, 26–35, <https://doi.org/10.1111/j.1365-2664.2009.01751.x>, 2010.
- Da Silva, R. F., Hansen, J. E., Rijnsdorp, D. P., Lowe, R. J., and Buckley, M. L.: The influence of submerged coastal structures on nearshore flows and wave runup, *Coastal Engineering*, 177, 104194, <https://doi.org/10.1016/j.coastaleng.2022.104194>, 2022.
- 590 Dalsgaard, T., Thamdrup, B., and Canfield, D. E.: Anaerobic ammonium oxidation (anammox) in the marine environment, *Research in Microbiology*, 156, 457–464, 2005.

- Degen, R. and Faulwetter, S.: The arctic traits database - a repository of arctic benthic invertebrate traits, *Earth System Science Data*, 11, 301–322, <https://doi.org/10.5194/essd-11-301-2019>, 2019.
- 595 Devol, A. H.: Denitrification, Anammox, and N₂ Production in Marine Sediments, *Annu. Rev. Mar. Sci.*, 7, 403–423, <https://doi.org/10.1146/annurev-marine-010213-135040>, 2015.
- Dionne, J.-C.: Érosion récente des marais intertidaux de l’estuaire du Saint-Laurent, Québec, *gpq*, 40, 307–323, <https://doi.org/10.7202/032651ar>, 2007.
- 600 Doody, J. P.: Coastal squeeze and managed realignment in southeast England, does it tell us anything about the future?, *Ocean & Coastal Management*, 79, 34–41, <https://doi.org/10.1016/j.ocecoaman.2012.05.008>, 2013.
- Duarte, C. M., Losada, I. J., Hendriks, I. E., Mazarrasa, I., and Marbà, N.: The role of coastal plant communities for climate change mitigation and adaptation, *Nature Clim Change*, 3, 961–968, <https://doi.org/10.1038/nclimate1970>, 2013.
- Dugan, J. E., Emery, K. A., Alber, M., Alexander, C. R., Byers, J. E., Gehman, A. M., McLenaghan, N., and Sojka, S. E.: Generalizing Ecological Effects of Shoreline Armoring Across Soft Sediment Environments, *Estuaries and Coasts*, 41, 180–196, <https://doi.org/10.1007/s12237-017-0254-x>, 2018.
- 605 Friess, D. A., Yando, E. S., Alemu I, J. B., Wong, L.-W., Soto, S. D., and Bhatia, N.: Ecosystem services and disservices of mangrove forests and salt marshes, in: *Oceanography and Marine Biology*, edited by: Hawkins, S. J., Allcock, A. L., Bates, A. E., Evans, A. J., Firth, L. B., McQuaid, C. D., Russell, B. D., Smith, I. P., Swearer, S. E., and Todd, P. A., CRC Press, 107–141, <https://doi.org/10.1201/9780429351495-3>, 2020.
- 610 Gedan, K. B., Silliman, B. R., and Bertness, M. D.: Centuries of Human-Driven Change in Salt Marsh Ecosystems, *Annu. Rev. Mar. Sci.*, 1, 117–141, <https://doi.org/10.1146/annurev.marine.010908.163930>, 2009.
- Gibeault, C., Neumeier, U., and Bernatchez, P.: Spatial and Temporal Sediment Dynamics in a Subarctic Salt Marsh (Gulf of St. Lawrence, Canada), *coas*, 32, 1344–1361, <https://doi.org/10.2112/JCOASTRES-D-15-00160.1>, 2016.
- 615 Gittman, R. K., Fodrie, F. J., Popowich, A. M., Keller, D. A., Bruno, J. F., Currin, C. A., Peterson, C. H., and Piehler, M. F.: Engineering away our natural defenses: an analysis of shoreline hardening in the US, *Frontiers in Ecol & Environ*, 13, 301–307, <https://doi.org/10.1890/150065>, 2015.
- Gittman, R. K., Scyphers, S. B., Smith, C. S., Neylan, I. P., and Grabowski, J. H.: Ecological Consequences of Shoreline Hardening: A Meta-Analysis, *BioScience*, 66, 763–773, <https://doi.org/10.1093/biosci/biw091>, 2016.
- 620 Glud, R. N.: Oxygen dynamics of marine sediments, *Marine Biology Research*, 4, 243–289, <https://doi.org/10.1080/17451000801888726>, 2008.
- Halpern, B. S., Frazier, M., Afflerbach, J., Lowndes, J. S., Micheli, F., O’Hara, C., Scarborough, C., and Selkoe, K. A.: Recent pace of change in human impact on the world’s ocean, *Sci Rep*, 9, 11609, <https://doi.org/10.1038/s41598-019-47201-9>, 2019.
- Hamme, R. C. and Emerson, S. R.: The solubility of neon, nitrogen and argon in distilled water and seawater, *Deep Sea Research Part I: Oceanographic Research Papers*, 51, 1517–1528, <https://doi.org/10.1016/j.dsr.2004.06.009>, 2004.
- 625 Hansen, H. P. and Koroleff, F.: Determination of nutrients, in: *Methods of Seawater Analysis*, edited by: Grasshoff, K., Kremling, K., and Ehrhardt, M., Wiley, 159–228, <https://doi.org/10.1002/9783527613984.ch10>, 1999.
- Hartig, F.: DHARMa: Residual diagnostics for hierarchical (multi-level/mixed) regression models, 2022.

- 630 Hinshaw, S. E., Tatariw, C., Flournoy, N., Kleinhuizen, A., Taylor, C., Sobecky, P. A., and Mortazavi, B.: Vegetation Loss Decreases Salt Marsh Denitrification Capacity: Implications for Marsh Erosion, *Environ. Sci. Technol.*, 51, 8245–8253, <https://doi.org/10.1021/acs.est.7b00618>, 2017.
- Hyacinthe, C., Anschütz, P., Carbonel, P., Jouanneau, J.-M., and Jorissen, F. J.: Early diagenetic processes in the muddy sediments of the Bay of Biscay, *Marine Geology*, 177, 111–128, [https://doi.org/10.1016/S0025-3227\(01\)00127-X](https://doi.org/10.1016/S0025-3227(01)00127-X), 2001.
- Jevrejeva, S., Jackson, L. P., Riva, R. E. M., Grinsted, A., and Moore, J. C.: Coastal sea level rise with warming above 2 °C, *Proc. Natl. Acad. Sci. U.S.A.*, 113, 13342–13347, <https://doi.org/10.1073/pnas.1605312113>, 2016.
- 635 Kana, T. M., Darkangelo, Christina., Hunt, M. Duane., Oldham, J. B., Bennett, G. E., and Cornwell, J. C.: Membrane Inlet Mass Spectrometer for Rapid High-Precision Determination of N₂, O₂, and Ar in Environmental Water Samples, *Anal. Chem.*, 66, 4166–4170, <https://doi.org/10.1021/ac00095a009>, 1994.
- Kana, T. M., Cornwell, J. C., and Zhong, L.: Determination of denitrification in the Chesapeake Bay from measurements of N₂ accumulation in bottom water, *Estuaries and Coasts: J ERF*, 29, 222–231, <https://doi.org/10.1007/BF02781991>, 2006.
- 640 Koch, M. S., Maltby, E., Oliver, G. A., and Bakker, S. A.: Factors controlling denitrification rates of tidal mudflats and fringing salt marshes in south-west England, *Estuarine, Coastal and Shelf Science*, 34, 471–485, [https://doi.org/10.1016/S0272-7714\(05\)80118-0](https://doi.org/10.1016/S0272-7714(05)80118-0), 1992.
- Koop-Jakobsen, K. and Giblin, A. E.: The effect of increased nitrate loading on nitrate reduction via denitrification and DNRA in salt marsh sediments, *Limnology & Oceanography*, 55, 789–802, <https://doi.org/10.4319/lo.2010.55.2.0789>, 2010.
- 645 Koop-Jakobsen, K., Mueller, P., Meier, R. J., Liebsch, G., and Jensen, K.: Plant-Sediment Interactions in Salt Marshes – An Optode Imaging Study of O₂, pH, and CO₂ Gradients in the Rhizosphere, *Front. Plant Sci.*, 9, <https://doi.org/10.3389/fpls.2018.00541>, 2018.
- Kostka, J. E. and Luther, G. W.: Partitioning and speciation of solid phase iron in saltmarsh sediments, *Geochimica et Cosmochimica Acta*, 58, 1701–1710, [https://doi.org/10.1016/0016-7037\(94\)90531-2](https://doi.org/10.1016/0016-7037(94)90531-2), 1994.
- 650 Kristensen, E., Penha-Lopes, G., Delefosse, M., Valdemarsen, T., Quintana, C., and Banta, G.: What is bioturbation? The need for a precise definition for fauna in aquatic sciences, *Marine Ecology Progress Series*, 446, 285–302, <https://doi.org/10.3354/meps09506>, 2012.
- Laliberté, E. and Legendre, P.: A distance-based framework for measuring functional diversity from multiple traits, *Ecology*, 91, 299–305, <https://doi.org/10.1890/08-2244.1>, 2010.
- 655 Lê, S., Josse, J., and Husson, F.: FactoMineR: An R Package for Multivariate Analysis, *Journal of Statistical Software*, 25, 1–18, <https://doi.org/10.18637/jss.v025.i01>, 2008.
- Mcowen, C., Weatherdon, L., Bochove, J.-W., Sullivan, E., Blyth, S., Zockler, C., Stanwell-Smith, D., Kingston, N., Martin, C., Spalding, M., and Fletcher, S.: A global map of saltmarshes, *BDJ*, 5, e11764, <https://doi.org/10.3897/BDJ.5.e11764>, 2017.
- 660 Mentaschi, L., Voutsdoukas, M. I., Pekel, J.-F., Voukouvalas, E., and Feyen, L.: Global long-term observations of coastal erosion and accretion, *Sci Rep*, 8, 12876, <https://doi.org/10.1038/s41598-018-30904-w>, 2018.
- Ministère des ressources naturelles et des forêts: Imagerie orthorectifiée du Québec, <https://doi.org/https://www.donneesquebec.ca/recherche/dataset/imagerie-orthorectifiee-du-quebec>, 2025.

- Neumann, B., Vafeidis, A. T., Zimmermann, J., and Nicholls, R. J.: Future Coastal Population Growth and Exposure to Sea-Level Rise and Coastal Flooding - A Global Assessment, PLoS ONE, 10, e0118571, <https://doi.org/10.1371/journal.pone.0118571>, 2015.
- Nielsen, L. P.: Denitrification in sediment determined from nitrogen isotope pairing, FEMS Microbiology Letters, 86, 357–362, <https://doi.org/10.1111/j.1574-6968.1992.tb04828.x>, 1992.
- Nixon, S. W.: Enriching the Sea to Death, Scientific American, 1998.
- Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., Evangelista, H. B. A., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill, M. O., Lahti, L., McGlinn, D., Ouellette, M.-H., Ribeiro Cunha, E., Smith, T., Stier, A., Ter Braak, C. J. F., and Weedon, J.: vegan: Community ecology package, 2022.
- O'Meara, T., Thompson, S. P., and Piehler, M. F.: Effects of shoreline hardening on nitrogen processing in estuarine marshes of the U.S. mid-Atlantic coast, Wetlands Ecol Manage, 23, 385–394, <https://doi.org/10.1007/s11273-014-9388-9>, 2015.
- Onorevole, K. M., Thompson, S. P., and Piehler, M. F.: Living shorelines enhance nitrogen removal capacity over time, Ecological Engineering, 120, 238–248, <https://doi.org/10.1016/j.ecoleng.2018.05.017>, 2018.
- Pascal, L., Maire, O., Volkenborn, N., Lecroart, P., Bichon, S., de Montaudouin, X., Grémare, A., and Deflandre, B.: Influence of the mud shrimp *Upogebia pusilla* (Decapoda: Gebiidea) on solute and porewater exchanges in an intertidal seagrass (*Zostera noltei*) meadow of Arcachon Bay: An experimental assessment, Journal of Experimental Marine Biology and Ecology, 477, 69–79, <https://doi.org/10.1016/j.jembe.2016.01.008>, 2016.
- Pascal, L., Bernatchez, P., Chaillou, G., Nozais, C., Lapointe Saint-Pierre, M., and Archambault, P.: Sea ice increases benthic community heterogeneity in a seagrass landscape, Estuarine, Coastal and Shelf Science, 243, 106898, <https://doi.org/10.1016/j.ecss.2020.106898>, 2020.
- Pascal, L., Chaillou, G., Nozais, C., Cool, J., Bernatchez, P., Letourneux, K., and Archambault, P.: Benthos response to nutrient enrichment and functional consequences in coastal ecosystems, Marine Environmental Research, 175, 105584, <https://doi.org/10.1016/j.marenvres.2022.105584>, 2022.
- Poulin, P., Pelletier, E., and Saint-Louis, R.: Seasonal variability of denitrification efficiency in northern salt marshes: An example from the St. Lawrence Estuary, Marine Environmental Research, 63, 490–505, <https://doi.org/10.1016/j.marenvres.2006.12.003>, 2007.
- Pratt, D. R., Pilditch, C. A., Lohrer, A. M., and Thrush, S. F.: The effects of short-term increases in turbidity on sandflat microphytobenthic productivity and nutrient fluxes, Journal of Sea Research, 92, 170–177, <https://doi.org/10.1016/j.seares.2013.07.009>, 2014.
- Queirós, A. M., Birchenough, S. N. R., Bremner, J., Godbold, J. A., Parker, R. E., Romero-Ramirez, A., Reiss, H., Solan, M., Somerfield, P. J., Colen, C. V., Hoey, G. V., and Widdicombe, S.: A bioturbation classification of European marine infaunal invertebrates, Ecology and Evolution, 3, 3958–3985, <https://doi.org/10.1002/ece3.769>, 2013.
- R Core Team: R: A language and environment for statistical computing, 2024.
- Rabalais, N. N., Turner, R. E., Díaz, R. J., and Justić, D.: Global change and eutrophication of coastal waters, ICES Journal of Marine Science, 66, 1528–1537, <https://doi.org/10.1093/icesjms/fsp047>, 2009.

- 700 van Rijn, L. C.: Coastal erosion and control, *Ocean & Coastal Management*, 54, 867–887, <https://doi.org/10.1016/j.ocecoaman.2011.05.004>, 2011.
- Risgaard-Petersen, N., Nielsen, L. P., Rysgaard, S., Dalsgaard, T., and Meyer, R. L.: Application of the isotope pairing technique in sediments where anammox and denitrification coexist, *Limnology and Oceanography: Methods*, 1, 63–73, <https://doi.org/10.4319/lom.2003.1.63>, 2003.
- 705 Sauvé, P., Bernatchez, P., and Glaus, M.: The role of the decision-making process on shoreline armoring: A case study in Quebec, Canada, *Ocean & Coastal Management*, 198, 105358, <https://doi.org/10.1016/j.ocecoaman.2020.105358>, 2020.
- Savoie-Ferron, F., Drouet, M., Drejza, S., Fraser, C., Marie, G., and Bernatchez, P.: Modélisation de l'effet de la hausse du niveau de la mer sur les marais maritimes, Chaire de recherche en géoscience côtière, Laboratoire de dynamique et de gestion intégrée des zones côtières, Université du Québec à Rimouski. Rapport remis au ministère de l'Environnement et de la Lutte
710 contre les changements climatiques, 2021.
- Schoell, M., Ayvazian, S., Cobb, D., Grunden, D., Chintala, M., Gerber-Williams, A., Pimenta, A., Strobel, C., and Rocha, K.: Exploring the Use of Living Shorelines for Stabilization and Nutrient Mitigation in New England, *Ecological Rest.*, 41, 84–98, <https://doi.org/10.3368/er.41.2-3.84>, 2023.
- Slomp, C. P.: Phosphorus cycling in the estuarine and coastal zones, in: *Treatise on estuarine and coastal science*, Elsevier, 715 201–229, <https://doi.org/10.1016/b978-0-12-374711-2.00506-4>, 2011.
- Snelgrove, P. V. R., Soetaert, K., Solan, M., Thrush, S., Wei, C.-L., Danovaro, R., Fulweiler, R. W., Kitazato, H., Ingole, B., Norkko, A., Parkes, R. J., and Volkenborn, N.: Global Carbon Cycling on a Heterogeneous Seafloor, *Trends in Ecology & Evolution*, 33, 96–105, <https://doi.org/10.1016/j.tree.2017.11.004>, 2018.
- Sousa, A., Lillebø, A., Risgaard-Petersen, N., Pardal, M., and Caçador, I.: Denitrification: an ecosystem service provided by
720 salt marshes, *Mar. Ecol. Prog. Ser.*, 448, 79–92, <https://doi.org/10.3354/meps09526>, 2012.
- Sousa, A. I., Lillebø, A. I., Caçador, I., and Pardal, M. A.: Contribution of *Spartina maritima* to the reduction of eutrophication in estuarine systems, *Environmental Pollution*, 156, 628–635, <https://doi.org/10.1016/j.envpol.2008.06.022>, 2008.
- Stief, P.: Stimulation of microbial nitrogen cycling in aquatic ecosystems by benthic macrofauna: mechanisms and environmental implications, *Biogeosciences*, 10, 7829–7846, <https://doi.org/10.5194/bg-10-7829-2013>, 2013.
- 725 Stockdale, A., Davison, W., and Zhang, H.: Micro-scale biogeochemical heterogeneity in sediments: A review of available technology and observed evidence, *Earth-Science Reviews*, 92, 81–97, <https://doi.org/10.1016/j.earscirev.2008.11.003>, 2009.
- Valiela, I. and Cole, M. L.: Comparative Evidence that Salt Marshes and Mangroves May Protect Seagrass Meadows from Land-derived Nitrogen Loads, *Ecosystems*, 5, 92–102, <https://doi.org/10.1007/s10021-001-0058-4>, 2002.
- Wang, H., Capurso, W., Chen, Q., Zhu, L., Niemoczynski, L., and Snedden, G.: Assessment of Wave Attenuation, Current
730 Patterns, and Sediment Deposition and Erosion During Winter Storms by Living Shoreline Structures in Gandys Beach, New Jersey., <https://doi.org/https://doi.org/10.3133/ofr20211040>, 2021.
- van Wesenbeeck, B. K., van de Koppel, J., Herman, P. M. J., Bakker, J. P., and Bouma, T. J.: Biomechanical warfare in ecology; negative interactions between species by habitat modification, *Oikos*, 116, 742–750, <https://doi.org/10.1111/j.0030-1299.2007.15485.x>, 2007.

- 735 Woodin, S. A., Volkenborn, N., Pilditch, C. A., Lohrer, A. M., Wetthey, D. S., Hewitt, J. E., and Thrush, S. F.: Same pattern, different mechanism: Locking onto the role of key species in seafloor ecosystem process, *Sci Rep*, 6, 26678, <https://doi.org/10.1038/srep26678>, 2016.
- Yin, G., Hou, L., Liu, M., Liu, Z., and Gardner, W. S.: A Novel Membrane Inlet Mass Spectrometer Method to Measure $^{15}\text{NH}_4^+$ for Isotope-Enrichment Experiments in Aquatic Ecosystems, *Environ. Sci. Technol.*, 48, 9555–9562, <https://doi.org/10.1021/es501261s>, 2014.
- 740 Zheng, Y., Hou, L., Liu, M., Yin, G., Gao, J., Jiang, X., Lin, X., Li, X., Yu, C., and Wang, R.: Community composition and activity of anaerobic ammonium oxidation bacteria in the rhizosphere of salt-marsh grass *Spartina alterniflora*, *Appl Microbiol Biotechnol*, 100, 8203–8212, <https://doi.org/10.1007/s00253-016-7625-2>, 2016.
- Zouaghi, F., Dansereau-Macias, E., Caulet, C., Dugas, S., Pascal, L., Chaillou, G., Sirois, L., Bernatchez, P., Nozais, C., Roberge, B., Noël, B., Bujold, É., and Didier, D.: Analyse, modélisation et suivi des interactions entre le briselames et la dynamique sédimentaire et végétale du marais le long de l’autoroute 20 à Rivière-du-Loup, Présenté au ministère des transports du Québec, 2024.
- 745