

Planktonic foraminifera Iodine/Calcium ratio: is it a proxy for dissolved oxygen in the ocean?

Vincent Guarinos^{1*}, Kazuyo Tachikawa¹, Marta Garcia¹, Giuseppe Siani², Marie Revel³, Hartmut Schulz⁴, Francisco Sierro⁵, Thomas Chalk¹

¹Aix Marseille Univ, CNRS, IRD, INRAE, Coll France, CEREGE, Aix-en-Provence, France.

²Laboratoire de Géosciences Paris-Saclay (GEOPS), CNRS, Paris Saclay, France.

³Université de la Cote d'Azur, CNRS, OCA, IRD, Geoazur, Valbonne, France.

⁴Department of Geosciences, University of Tübingen, Tübingen, Germany.

⁵Dept. de Geología, Univ. de Salamanca, Plaza de los Caídos s/n, 37008, Salamanca, Spain

Correspondence to: Guarinos Vincent (guarinos@cerege.fr)

Abstract

Direct observations indicate a declining trend in ocean oxygen concentrations, which is not quantitatively captured by models. The complexity of oxygenation variability, linked to both physical and biochemical parameters, must be investigated across different modern climate contexts. Foraminiferal iodine-to-calcium (I/Ca) ratios have emerged as a proxy for subsurface oxygen concentrations although ~~their~~ capacity for quantitative reconstructions remains to be elucidated. We provide a new database of modern foraminiferal I/Ca, including the first results from the Mediterranean Sea and new samples in the Arabian Sea together with the parameters of biochemistry (nutrient concentration, pH, chlorophyll, oxygen, net primary productivity), physical and geographical (temperature, salinity, latitude, distance from the coast, water depths) and diagenesis potential (depth in core, sediment age) to better understand the proxy behaviour. Considering the strong spatiotemporal variability in dissolved oxygen in subsurface ocean, we propose to use statistically robust 25th percentile of oxygen concentration (p25 [O₂]) ~~in the~~ upper 500 m in the water column ~~instead of minimum concentration~~. Our results affirm that oxygen concentration is the primary driver of foraminiferal I/Ca and we propose a new calibration equation of foraminiferal I/Ca against p25 [O₂]. Based on a new database, we identify a complex relationship between p25 [O₂] and iodine speciation, which is one of the main sources of scatter. A comparison with synthetic calcite reveals that planktonic foraminiferal tests can incorporate either more or less iodate at high p25 [O₂] than abiotic calcite, probably due to “vital effects”. The Mediterranean Sea samples present a wide range from low to high I/Ca (0.6 to 6.9 μmol/mol) in well-oxygenated water, which cannot be explained solely by authigenic calcite precipitation. Our results highlight the complex behaviour of the I/Ca proxy, while ~~its~~ semi-quantitative reconstruction in palaeoceanography ~~remains promising~~.

Introduction

Deleted: its

Deleted: instead of minimum concentration

Deleted: reinforcing

Oxygen concentration [O₂] in the global ocean is decreasing, and the decline is highly influenced by warming patterns induced by global climate change (Oschlies et al., 2018). Future oxygenation change is difficult to predict as it depends on various parameters. Dissolved oxygen depletion can be related to changes in solubility with seawater temperature, generally decreasing ventilation of water masses (Oschlies et al., 2017), and through biological productivity triggered oxygen consumption in the ocean, especially in the subsurface, where the productivity and remineralisation are at a maximum. Models can currently only explain about half the observed oxygen depletion (Bopp et al., 2013; Frölicher et al., 2016) and thus a proxy for subsurface oxygen content is highly sought to study longer variability under different climate conditions.

Deleted: (Oschlies et al., 2017)

Deleted: /or

The iodine to calcium ratios (I/Ca) from planktonic foraminifera tests have been investigated and used as a proxy for subsurface/intermediate depth minimum oxygen concentration (Lu et al., 2010, 2016, 2020; Winkelbauer et al., 2021; Hess et al., 2025; Zhou et al., 2014, 2022; Hoogakker et al., 2025) as inorganic iodine speciation depends on redox potential, which is mainly controlled by oxygen content (Luther, 2023). Severe oxygen depletion, less than 1 nM of oxygen, will chemo-dynamically trigger iodate reduction to iodide (I⁻) (Luther, 2023). In oxygenated water, iodine is mainly in its oxidized form, iodate (IO₃⁻) (Chance et al., 2014, 2019). Iodide, the reduced form of inorganic iodine can also be found in small concentrations near surface because of biological iodate uptake and iodide release during phytoplankton growth (Hepach et al., 2020). In hypoxic and anoxic waters, iodide is the major iodine species, thought to increase as iodate is reduced. The reduction reaction takes place under short time scales, from days to weeks (Chance et al., 2010) whereas the full (re)oxidation of iodide back to iodate has been reported to take place from several weeks to several months (Wadley et al., 2020). Total inorganic iodine in oxygenated water typically ranges between 400 and 500 nM, corresponding to a variability of about 20% (Chance et al., 2010). Iodine is also present as organically bound iodine, primarily in the form of dissolved organic iodine. In the open ocean, it is typically about 5% of total dissolved iodine, but this proportion can increase to up to ~20% in coastal areas due to interaction with higher concentrations of dissolved organic matter (Jones et al., 2024). Iodate is assumed to be the only iodine species to be incorporated to foraminiferal calcite tests as a substitute to carbonate ion (Feng and Redfern, 2018; Lu et al., 2010; Podder et al., 2017). The processes and kinetics of the chemical reactions in the water column, as well as its incorporation mechanisms into biologically mediated calcite are still poorly understood (Luther, 2023).

Deleted: (Lu et al., 2010, 2016, 2020; Winkelbauer et al., 2021; Hess et al., 2025; Zhou et al., 2014, 2022; Hoogakker et al., 2024)

Calibration of the I/Ca–O₂ proxy was first focused on the quantitative reconstruction of subsurface oxygen concentration as synthetic calcite showed linear incorporation of iodate depending on its concentration (Lu et al., 2010). Later, foraminiferal core-top calibrations started to relate the I/Ca ratio in respect with minimum oxygen from 0–500 m in the water column, considering a tight relationship between the minimum oxygen concentration and surface water iodate concentration (Lu et al., 2016). The proxy was thus proposed as a semi-quantitative tool with thresholds I/Ca value of 0.85, 2.3 and 5 μmol/mol that corresponds to [O₂] 50, 90 and 140 μmol/kg (Lu et al., 2016; Wang et al., 2026). Data from the Southeast Atlantic including sites close to the Benguela upwelling system highlighted the possibility of low I/Ca in apparently oxygenated water (Lu et al., 2020). The minimum oxygen values of the study region were highly spatially variable and the use of oxygen data at higher resolution (0.25°) reduced the number data points with low I/Ca under high oxygenation condition (Lu et al., 2020). However, substantial scatter remains in the relationship between foraminiferal I/Ca and minimum oxygen concentration that is characterised by two broad domains: oxygen depleted waters (I/Ca <2.5 μmol/mol) and oxygenated water (I/Ca >4 μmol/mol). Nilsson-Kerr et al. (2025) expanded the I/Ca–O₂ core top calibration dataset with measurements

Deleted: The proxy was thus proposed as a semi-quantitative tool with a threshold I/Ca value of 2.5 μmol/mol that corresponds to [O₂] <70 μmol/kg (Lu et al., 2016).

from regions with low subsurface O₂, confirming the presence of significant scatter. Hess et al. (2025) proposed that I/Ca dispersion is linked with the different iodate concentrations at different foraminiferal calcification depths. Scattering could then be reduced by using Mg/Ca ratios in the foraminifera, a proxy for temperature that estimates their calcification depth. Although they showed that I/Ca varies predictably based on iodate concentration and water depth, substantial dispersion still persists and the effect of calcification depth cannot explain all the results.

Moreover, low I/Ca from plankton tow in highly oxygenated waters have been measured (Winkelbauer et al., 2023). It was suggested that iodine is incorporated during particle sinking and could explain those low values, thus challenging the use of the proxy for subsurface oxygenation, an interpretation that has not yet been further interrogated. Besides, given the lack of knowledge on the underlying mechanism, application of this proxy needs caveating when carried out for samples from coastal shelves and restricted basins where the mixing with open ocean water is not efficient (Hess et al., 2025; Lu et al., 2020).

The previous studies have mostly focused on trying to reduce scattering by investigating the Y-axis (I/Ca), but the definition of X-axis (oxygen concentration) has not been treated systematically, despite its importance (Lu et al., 2020). Some studies suggest that the complexity of iodine speciation and its coupling with dissolved oxygen is responsible for the observed data scatter (Hess et al., 2025; Nilsson-Kerr et al., 2025). However, the systematic study with a large number of data focusing on both the definition of minimum dissolved oxygen concentration and iodine speciation has not yet been realised.

We aim to determine whether the definition of “oxygen concentration” can help to better calibrate the proxy, and further investigate the possible reasons for foraminiferal I/Ca data scattering. We also examine if the proxy can be used in a semi-enclosed basin, the Mediterranean Sea as important oceanographical variations over small distances may limit the use of the proxy (Hess et al., 2025). In this paper, our approach is to (1) propose a statistically robust definition for minimum O₂, (2) investigate the coupling between iodine speciation and dissolved oxygen as a potential control for foraminiferal I/Ca scattering (3) examine the differences in iodine incorporation between foraminiferal and abiotic calcite, and (4) evaluate the influence of authigenic carbonate precipitation on foraminiferal I/Ca under high alkalinity conditions.

1 Material and Methods

1.1 Material

In this study we present core top planktonic foraminiferal I/Ca from 7 locations including 5 from the Mediterranean region (Fig. 1b). Core MD99-2341 (577 mbsl) from Gulf of Cadiz together with ODP Site 977 (1984 mbsl) from Alboran Sea are selected to characterise the Atlantic inflow signature. MD04-2797 (771 mbsl) is located next to the Strait of Sicily, and is used to indicate mixing ratio at the connection between the Western and Eastern Mediterranean basins. SL95 (1390 mbsl), in the Gulf of Sirte, is thought to represent the southern part of the Ionian Sea. Core MD04-2724 (1659 mbsl) has been recovered in the deep Nile delta and it is the easternmost core of this study. Core MD04-2722 (1780 mbsl) was recovered in the Levantine basin. MD90-917 (1010 mbsl), was recovered in the Adriatic Sea, where the Eastern Mediterranean deep water is formed and corresponds to a well oxygenated area. The two other cores MD04-2873 (2196 mbsl) and MD04-2876 (828 mbsl) were recovered in the Pakistan margin, one below the Oxygen Minimum Zone (OMZ), and the other inside it, respectively (Fig. 1c). They are selected in order to identify any impact of the bottom water oxygenation state on I/Ca ratios.

Deleted: (Hess et al., 2025)

Deleted: We also examine if the proxy can be used in a semi-enclosed basin, the Mediterranean Sea.

The core top materials of the two new cores, SL95 and MD04-2724, were dated to $1,835 \pm 30$ (1sd, at 0–1 cm) and $5,515 \pm 30$ radiocarbon years BP (1sd, at 0–1cm) respectively, using mono-species planktonic foraminifera performed at the ARTEMIS facility in Gif-sur-Yvette, France.

Deleted: We present

Deleted: first measurements from 2

Deleted: . The core tops




Figure 1: Map of 25th percentile of oxygen concentration (p25 [O₂]) in μmol/kg in ±0.25° (latitude and longitude) at 0–500 mbsl and symbols corresponding to modern foraminiferal I/Ca (μmol/mol) less than 8000 years (8 ka) before present (BP) at global scale (a) and in the Mediterranean Sea (b). The locations of new core top data from this study are represented with triangles and published values with circles. Plankton tow data from Winkelbauer

et al. (2023) are represented with diamonds. Position of dissolved inorganic iodine species concentrations in the Mediterranean Sea are represented in black (Tian and Nicolas, 1995; Ullman et al., 1990). Oxygen distributions were obtained from World Ocean Database 2023 (WOD23) (Mishonov et al., 2024). Arabian Sea mean oxygen section (c) shows the position of the two core-tops studied. MD04-2876 is in the core of the OMZ, and MD04-2873 is below. Oxygen profile is made with Ocean Data View (Schlitzer and Reiner, 2025).

Deleted: (Mishonov et al., 2024)

Deleted: top

142

1.2 Analysis of Iodine/Calcium ratio from planktonic foraminifera

For foraminiferal I/Ca analysis, 30 monospecific planktonic foraminiferal tests were picked from 250–355 µm fraction and cleaned following Barker et al. (2003) short-cleaning protocol. Only the oxidative cleaning was realised. Depending on the core, different species are used. For Gulf of Cadiz, *Globigerinoides ruber albus* (hereafter referred to as *G. ruber*), *Globigerina bulloides*, *Trilobatus sacculifer* (without sack-like chamber), *Globigerinella siphonifera* and *Pulleniatina obliquiloculata* were picked. Given the influence of calcification depth on I/Ca ratios (Hess et al., 2025), species-specific calcification depths are reported here. *Globigerinoides ruber* and *T. sacculifer* are mixed layer calcifiers. *G. bulloides*, *G. siphonifera* and *P. obliquiloculata* are thermocline-dwelling species. For the Alboran Sea, *G. ruber* and *G. bulloides* ratios were analysed. For MD04-2797, SL95, MD90-917 and the two cores from the Arabian Sea, only *G. ruber* was used. After cleaning, samples were dissolved with diluted nitric acid at 0.1 M (0.66%). The obtained solution was then centrifuged to separate any residual non-dissolved particles. After a calcium (Ca) concentration test to determine the necessary dilution factor for each sample, an aliquot for iodine measurement is stabilised with tetramethylammonium hydroxide (TMAH, SIGMA-ALDRICH) to obtain a 0.33% HNO₃/0.48% TMAH matrix. The solution is then diluted to a Ca concentration of 50 µg/mL (50 ppm). The remainder is diluted to obtain Ca concentration of 100 µg/mL to determine Ca, U, Mn, Fe and Al concentrations (Guarinos et al., in prep). In this study only I/Ca ratio is discussed. Elemental analysis is carried out using ICP-MS (Agilent 7500ce) at the CEREGE, France.

Deleted: protocol. Depending on the core, different species are used. For Gulf of Cadiz, *Globigerinoides ruber albus* (hereafter referred to as *G. ruber*), *Globigerina bulloides*, *Trilobatus sacculifer* (without sack-like chamber), *Globigerinella siphonifera* and *Pulleniatina obliquiloculata* were picked....

Table 1: Description of cores and planktonic foraminiferal species used for I/Ca analysis

Core name	Location	Latitude	Longitude	Depth (mbsl)	Depth in core (cm)	Core top age (ka cal BP)	Species picked	Comment	Age model reference
MD99-2341	Gulf of Cadiz	36.383	-7.533	577	2–4	1.02	<i>G. ruber</i> , <i>G. bulloides</i> , <i>T. sacculifer</i> , <i>G. siphonifera</i> , <i>P. obliquiloculata</i>	Atlantic water inflow	Eynaud et al., 2009
ODP Site 977	Alboran Sea	36.032	-1.955	1984	23–25	1.37	<i>G. ruber</i> and <i>G. bulloides</i>	Atlantic water inflow	Martrat et al., 2004
MD04-2797	Sicily Strait	36.959	11.666	771	0–1	0.70	<i>G. ruber</i>	Connection between basins	Cornuault et al., 2018
SL95	Gulf of Sirte	32.774	19.191	1390	0–1	1.20	<i>G. ruber</i>	South Ionian Sea	This study
MD04-2724	Levantine Sea	33.033	29.150	1659	5–6	6.00	<i>G. ruber</i> , <i>T. sacculifer</i>	Deep Nile delta	This study

MD04-2722	Levantine Sea	33.100	33.500	1780	0–1	3.60	G. ruber	Eastern basin edge	Cornuault et al., 2018
MD90-917	Adriatic Sea	41.280	17.620	1010	2–3	0.65	<i>G. ruber</i>	Well oxygenated	Siani et al., 2010
MD04-2873	Pakistan Margin	23.532	63.827	2196	1–2	NA	<i>G. ruber</i>	Below the OMZ	Pichevin et al., 2007
MD04-2876	Pakistan Margin	24.849	64.014	828	2–3	NA	<i>G. ruber</i>	Within the OMZ	Bard et al., 2013

169

170 Each sample is analysed and standard-sample bracketing is used to characterise instrumental drift during
171 measurement. There is no currently certified geostandard for I/Ca but JCp-1, a coral powder, is commonly used in
172 this role. In our analyses JCp-1 is measured every five samples. Our measure of I/Ca of JCp-1 is 4.34 ± 0.11
173 $\mu\text{mol/mol}$ ($n=27$, 1sd). This is in good agreement with published values of $4.27 \pm 0.10 \mu\text{mol/mol}$ ($n=13$, 1sd,
174 Winkelbauer et al., 2023), $4.33 \pm 0.06 \mu\text{mol/mol}$ ($n=6$, 1sd, Chai and Muramatsu, 2007), $4.27 \pm 0.06 \mu\text{mol/mol}$
175 (1sd, Lu et al., 2010), and $4.20 \pm 0.43 \mu\text{mol/mol}$ (1sd, Hess et al., 2025). Calculated I/Ca precision is better than
176 3% and blank levels were low, often below detection limits, so no correction is applied.

177 1.3 Inorganic iodine speciation depending on dissolved oxygen concentration

178 The concentration of inorganic iodine species in the ocean is primarily controlled by redox conditions, with iodate
179 prevailing in oxic conditions and iodide dominating in reducing environments (Chance et al., 2014; Ullman et al.,
180 1990). To establish the relationship between iodate and oxygen concentration, iodine data are expressed as iodate
181 proportion in total inorganic iodine in order to avoid potential impacts from changing iodine concentration. [We](#)
182 [have selected data shallower than 500 mbsl and within \$\pm 1\sigma\$ of total inorganic iodine concentration \(Fig. 3c\) to](#)
183 [avoid heavily iodine enriched/depleted areas. Only data with both iodate and iodide concentrations are selected to](#)
184 [better constrain the speciation and avoid assumption on total inorganic iodine concentration.](#) Iodine speciation data
185 from Bermuda inshore water are excluded (Jickells et al., 1988), as they originate from a highly restricted
186 environment, and not representative of the conditions considered in this study. Data analysis is performed using
187 least-squares fitting analysis in Python 3.12. The 95% prediction intervals are computed by propagating the
188 uncertainties of the fitted model parameters contained in the covariance matrix. This combines both the variance
189 of the fitted parameters and the residual variance of the data within the selected area.

190 To explore the influence of redox conditions on inorganic iodine speciation, oxygen data were combined with
191 published iodine speciation measures. Inorganic iodine speciation in the upper 500 m is investigated using
192 previously published data (Winkelbauer et al., 2023). Oxygen concentrations were obtained from World Ocean
193 Database 2023 (WOD23) from [Mishonov et al. 2024](#) from within a 0.25° area around each sampling site, where
194 possible. From every obtained oxygen distribution, we determined the percentiles of oxygen concentration. Each
195 measurement from 0–500 m at a given station is attributed to the same percentile value.

Deleted: Mishonov et al. 2024

196

197 1.4 Planktonic foraminiferal I/Ca database

199 Planktonic foraminiferal I/Ca database is obtained with 15 new core top data (this study), and previously published
200 core top and plankton tow measurements (Hess et al., 2025; Lu et al., 2020, 2016; Nilsson-Kerr et al., 2025;
201 Winkelbauer et al., 2021, 2023; Zhou et al., 2022).

202 The database contains 567 core top data points, 438 of which are younger than 2,000 years BP. Oxygen distribution
203 in the first 500 m are obtained from WOD23 accepted values (class 0) from i) Bottle (Winkler titration) and ii)
204 Conductivity–Temperature–Depth (CTD) with oxygen sensor data in a 0.25° area around each sampling site
205 (Mishonov et al., 2024). For sites where no oxygen profile is available, the area of search is extended to 0.5° and
206 the data flagged, no data from >0.5° is used. The obtained oxygen concentration data are used to calculate the
207 minimum, maximum, mean, and percentile values. CTD measurements are only used for 3 I/Ca values as they are
208 considered to be less accurate because of excursion artifacts. For this reason, we prioritise bottle data which are
209 consider to be more accurate.

Deleted: (Mishonov et al., 2024)

210 Additional environmental variables are included to assess their potential contribution to I/Ca variability. Other
211 parameters, seawater temperature, salinity, nitrate, phosphate, chlorophyll, pH and alkalinity are similarly obtained
212 from class 0 bottle measurements in 0.25° around sampling site at the first 500 m. Investigating the 0–500 m is
213 crucial because this is where the planktonic I/Ca signal is presumed to originate (Lu et al., 2016). Oxycline depth
214 and mean oxygen concentration at oxycline were also tested as they may highly influence foraminiferal I/Ca (Wang
215 et al., 2026). They were recovered from class 0 bottle measurements in 0.25° area around sampling site. No value
216 is attributed to sites without any available bottle measurement. Other parameters, net primary productivity (NPP)
217 (Behrenfeld and Falkowski, 1997; NASA Earth Data Base, 2025), current speed, and mixed layer depth, are
218 obtained using their corresponding raster resolution. NPP is derived from annual mean satellite data integrated
219 from 1998 to 2022 at a resolution of 25 km and using Eppley-Vertically Generalized Production Model (Eppley-
220 VGPM). Parameters related to ocean dynamics are derived from Global Ocean Physics Reanalysis annual mean
221 and maximum values (Lellouche et al., 2021). We use absolute Northward and Eastward velocities from surface
222 to 500 m and mixed layer depth. Data sources are listed in Table 2.

Deleted: No value is attributed to sites without any available bottle measurement.

Deleted: (Behrenfeld and Falkowski, 1997; Oregon State University ORCA, n.d.).

Deleted: (“Global Ocean Physics Reanalysis,” 2025).

223 The database construction we propose here is a table format with two entries. Each line is a specific measurement
224 of foraminiferal I/Ca. Columns correspond to different characteristics of the measurement, sampling site, and all
225 the extracted site-specific parameters previously mentioned. Information of symbionts presence and habitat depth
226 is also shown. We also specify the cleaning protocol of the samples as reductive cleaning has been shown to reduce
227 the I/Ca value by 30% (Zhou et al., 2022). Samples processed by reductive cleaning are corrected accordingly,
228 with an increase of 30% of I/Ca. This impacts 197 measures (35% of the full dataset). All the figures presented
229 are using the corrected I/Ca. The full database is available in supplementary material.

Deleted: The database construction we propose here is a table format with two entries. Each line is a specific measure of foraminiferal I/Ca. Columns correspond to different characteristics of the measure, sampling site, and all the extracted site-specific parameters previously mentioned. Information of symbionts presence and habitat depth is also shown....

230 To assess the relationships between foraminiferal I/Ca and environmental variables, we use both Spearman’s ρ
231 and Kendall’s τ correlation coefficients. Even if some individual uncertainties of the selected variables may follow
232 normal distributions, the dataset as a whole is heterogeneous and non-normally distributed, making non-parametric
233 tests more appropriate. Moreover, as the resulting relationships are likely non-linear, these rank-based methods
234 provide a robust evaluation of correlation. The use of both correlation coefficients captures two complementary
235 aspects of monotonic relationships: the strength of the association and how well the observations agree with each
236 other. These analyses are intended to identify statistical covariation.

250 **2 Results**

251 **2.1 Definition of p25 [O₂] and Iodine speciation**

252 Oxygen concentration ranges from 0 to 300 $\mu\text{mol/kg}$. Obtained oxygen distribution in the upper 500 m layer at
253 sites where foraminiferal I/Ca is available are mostly unimodal or multimodal (Fig. 2). Previous studies use the
254 minimum [O₂] value measured in 0–500 m to calibrate foraminiferal I/Ca (Hess et al., 2025; Lu et al., 2020). The
255 use of the minimum value can provide questionable representativeness of actual conditions because the minimum
256 value could reflect exceptional events or data reporting issues. As the distribution of oxygen concentration at 0 to
257 500 m water depth is not normally distributed, we use the 25th percentile (p25) to represent average oxygen
258 depletion instead of the minimum value. Its use is thought to partially eliminate outliers or errors in the oxygen
259 dataset. The use of percentiles takes into account the full distribution in 0–500 m in space and time as much as the
260 oxygen dataset allows. However, this indicator is not perfect as important errors or sampling bias may still
261 influence the values. Furthermore, the goal of the use of p25 is not to propose a quantitative interpretation of the
262 proxy. Iodine concentration and speciation dynamics may not be fully captured by oxygen variability and result in
263 subsequent dispersion. We have tested percentiles from 5 to 30 at each 5% (Fig. S1). All tested definition of X-
264 axis shows the same slope, with only the intercept being different. We choose the higher R² (0.43) together with
265 the lower Root Mean Square Error (RMSE) (1.370), corresponding to p25 (Fig. S1). We note here that p25 value
266 is close to mean value minus 1 standard deviation when the distribution was normal or quasi-normal. With p25
267 being the oxygen parameter with the best fit, it is possible to remove part of the oxygen influence on the proxy.

Deleted: 39

Deleted: 356




Figure 2: Histograms of common types of oxygen distributions encountered in this study, with OMZ-type from Mauritanian upwelling (a), intermediate from southwestern Pacific Ocean (b) and well oxygenated from North Atlantic (c) distributions. Oxygen depletion representative value p25 [O₂] is highlighted in full red line. Other percentiles (p5, p50, p75, p95) are in dotted grey lines. Dotted red line is the minimum oxygen value previously used. Dotted yellow and purple lines are 1 sigma (68%) and 2 sigma (95%) limits, respectively. The difference between [O₂] min and p25 [O₂] is 0–142 μmol/kg. Note that the shown oxygen concentration range is different between (a), (b) and (c).

In strongly oxygen-depleted areas, such as OMZ, the oxygen distributions are mainly multimodal with high values near surface and lower values in the heart of OMZ (Fig. 2a). Those areas are represented with p25 [O₂] values ranging from 0 to 100 μmol/kg.

In areas with p25 [O₂] value between 100 and 180 μmol/kg (tropical Atlantic Ocean, Ontong Java Plateau in the tropical Pacific), the distribution of oxygen concentration is mostly multimodal, reflecting vertical change with water depth but with moderate oxygen depletion (Fig. 2b).

284 Oxygenated waters are marked by p25 [O₂] values from 180 to 300 μmol/kg. The corresponding distributions are
285 mostly unimodal (Fig. 2c). Some sites were not attributed to a distribution type due to their relative rarity.

286

287 **2.2 Iodine speciation and oxygen concentration**

288 Total inorganic iodine is in the range of 400 to 500 nM and is more or less constant in the ocean (Chance et al.,
289 2010) (Fig. 3c). Inorganic iodine depletion near coastal regions have been attributed to biological uptake
290 (Elderfield and Truesdale, 1980; Wong and Zhang, 1992). But inorganic iodine speciation primarily varies with
291 redox potential, largely governed by oxygen availability (Luther, 2023). In order to compare inorganic iodine
292 speciation to foraminiferal I/Ca, we represent it from the first 500 m as a function of p25 [O₂] in 0–500 m (Fig.
293 3a) and [O₂]min in 0–500 m. Speciation is shown as iodate/(iodate + iodide) in Fig. 3a, that corresponds to the
294 iodate proportion of the inorganic iodine concentration (Tian and Nicolas, 1995). This representation allows us to
295 minimise the effect of inorganic iodine concentration (Elderfield and Truesdale, 1980; Tian and Nicolas, 1995).
296 As this approach is based on real data, it may partly account for other factors affecting inorganic iodine speciation
297 such as mixing of water masses and primary productivity (Cheng et al., 2024; Evans et al., 2020; Hardisty et al.,
298 2021; Moriyasu et al., 2020). The aim of Fig. 3 is not to compare the two x-axes, but rather to present a different
299 concept for inorganic iodine speciation (Fig. 3a) than the common iodate concentration (Fig. 3b). This is especially
300 visible with samples from the Indian Ocean with lower iodate concentration than the other oceans (Fig. 3b) being
301 in the cloud of points with inorganic iodine speciation (Fig. 3a) (Chance et al., 2020; Farrenkopf and Luther, 2002).

302 Main result shows high dispersion of iodine speciation, and non-linear relationship (Fig. 3a and 3b). It also shows
303 that iodate might exist in oxygen depleted waters, especially in Pacific Ocean. Iodine speciation is highly variable
304 at 0–500 m. Iodate absence is mainly found at p25 [O₂] <10 μmol/kg although it is not systematically absent in
305 those areas. In well oxygenated water (>200 μmol/kg), inorganic iodine is principally under the oxidised form,
306 with a proportion above 60%. However, measurements from one site in Pacific and three in Atlantic are low in
307 iodate proportions (<0.5) in oxygenated areas (>150 μmol/kg), and fall outside the prediction interval.

308 We choose an exponential fit function to represent the rapid iodate decrease when p25 [O₂] is below 20 μmol/kg
309 and the plateau in oxygenated waters (p25 [O₂] >180 μmol/kg) (Fig. 3a). Prediction interval for exponential fit is
310 represented in shaded red area. The objective of this representation is not to find the best fit, but rather to determine
311 whether the relationship between iodate proportion and dissolved oxygen content could explain the variation in
312 foraminiferal I/Ca-based oxygen reconstruction. Compared to min[O₂] and iodate concentration (Fig. 3b), the use
313 of p25 [O₂] with proportion of iodate in inorganic iodine (Fig. 3a) limits the influence of total inorganic iodine
314 concentration change. The main difference in the x-axis definition is the change in range, from 0–280 μmol/kg for
315 [O₂] min to 0–330 μmol/kg for p25 [O₂] (Fig. 3a, b). This results in a visually stretched dataset in the x-direction.
316 Moreover, the use of a different y-axis leads to fewer values outside the 95% prediction interval in the high-iodate-
317 proportion / low-p25 range compared to iodate concentration. A few points fall below the 95% prediction interval
318 at intermediate oxygen levels when using p25 [O₂].

Deleted:),

Deleted: min

Deleted: slightly improves the consistency of the relationship by limiting


Deleted: but does not affect the dispersion along the y-axis.

Deleted: fall

Deleted: [O₂] min

326 In the Mediterranean Sea, inorganic iodine speciation profiles are scarce, with only one time series and four
327 different locations reported (Tian and Nicolas, 1995; Ullman et al., 1990). Nevertheless, consistently high iodate
328 concentrations are observed in the upper 0–500 m (370–515 μM) corresponding to well oxygenated water ($[\text{O}_2]_{\text{min}}$
329 100–180 $\mu\text{mol/kg}$) (Fig. 3b). Under these conditions, iodate represents more than 80% of total inorganic iodine,
330 with $p25 [\text{O}_2]$ between 190 and 200 $\mu\text{mol/kg}$.

Deleted: nmol/L



331 **Figure 3:** Iodate proportion in total inorganic iodine, iodate/(iodate+iodide) in 0–500 m as a function of $p25$ (a)
332 (Spearman $\rho = 0.625$, Kendall $\tau = 0.447$, p-values < 0.01) and iodate concentration as a function of minimum (b)
333 (Spearman $\rho = 0.650$, Kendall $\tau = 0.470$, p-values < 0.01) $[\text{O}_2]$ ($\mu\text{mol/kg}$) at 0–500 m. The proportion of iodate in
334 inorganic iodine was obtained from Winkelbauer et al. (2023) and are classified by ocean. **Mediterranean sites are**
335 **highlighted with white contour.** Red dotted line is the exponential fit of this relation ($y = -0.6975 \times \exp(-0.0181 \times x)$
336 $+ 0.8945$ for $p25$), with the 95% prediction interval filled in red. Total inorganic iodine distribution with mean and
337 $\pm 1\sigma$ values (c).

Deleted: 6175

Deleted: 0138

Deleted: 9202

338 2.3 New core top planktonic foraminiferal I/Ca

339 Low iodine concentration in foraminifera, ranging from 0 to 2 $\mu\text{mol/mol}$, are mostly found with $p25 [\text{O}_2]$ between
340 0 and 180 $\mu\text{mol/kg}$ (Fig. 4). In the 180–300 $\mu\text{mol/kg}$ of $p25 [\text{O}_2]$, most foraminiferal I/Ca ratios show higher values
341 ranging from 1 to 8 $\mu\text{mol/kg}$.

Deleted:)




Figure 4: Core top foraminiferal I/Ca ($\mu\text{mol/mol}$) as a function of p25 [O₂] concentration ($\mu\text{mol/kg}$) for samples more recent than 8 ka BP (8,000 years Before Present). New measures (n=15) from this study are coloured as water depth (mbsl) and identified with colour code. Previously published values are represented in grey. All foraminiferal species are differentiated by symbols. Red linear regression was obtained using all data < 2 ka BP.

New core top data from the Pakistan margin present I/Ca values lower than 2.5 $\mu\text{mol/mol}$ (0.7 $\mu\text{mol/mol}$ for MD04-2876 and 1.8 $\mu\text{mol/mol}$ for MD04-2873). Difference in p25 [O₂] could be related to oxygen data availability

Deleted: .

357 with wider area of oxygen distribution of 0.5° for MD04-2873 instead of 0.25° for MD04-2876 and different
358 number of oxygen profiles (n=2 and n=7 respectively).

359 In the Mediterranean Sea, foraminiferal I/Ca values are very low in highly oxygenated areas (1.0 µmol/mol in the
360 Gulf of Cadiz and 1.6 to 3.1 µmol/mol in Alboran Sea) ~~that corresponds to p25 [O₂] between 180 and 220 µmol/kg~~
361 ~~(Fig. 4), or 168 and 190 µmol/kg for min[O₂]~~. The observed trend cannot be explained by vertical oxygen structure
362 or by seasonal oxygen-depletion events, as dissolved oxygen profiles in the upper 500 m show very limited
363 seasonal or inter-annual variability, even when more than 20 profiles per site are considered. Regarding I/Ca
364 difference between planktonic foraminiferal species, three cores provide information (Table 1). The dispersion of
365 different species of foraminifera is around 40% for the Gulf of Cadiz (MD99-2341) and the Alboran Sea (ODP
366 Site 977) whereas in the Eastern Mediterranean Sea the high I/Ca ratios (from 2 to 6 µmol/mol) are associated
367 with a dispersion of 50% for the deep Nile Delta core (MD04-2724).

368 Iodine incorporation during particle sinking has been proposed to explain the low I/Ca values from plankton tows
369 (Winkelbauer et al., 2023). However, we find no systematic relation with water depth and I/Ca with our new data
370 (Fig. 4). Deep sites in oxygenated water like ODP Site 977 at 1984 mbsl also show very low I/Ca. Other cores at
371 comparable depths and oxygenation levels, like MD04-2724 at 1659 mbsl show I/Ca above 4 µmol/mol.

372 2.4 Planktonic I/Ca values and the environmental parameters on global scale

373 The relationship between environmental parameters such as biogeochemical characterisation of 0–500 m at each
374 site, primary productivity and physical parameters (current velocity) and foraminiferal I/Ca is investigated using
375 statistical test Kendall's τ and Spearman's ρ correlation coefficients (Table 2). We only include samples more
376 recent than 2 ka BP in the statistical analysis because older samples are less likely to be representative of modern
377 conditions. The results are reported in Table 2.

378 Table 2: Results of statistical correlation test for environmental variables and planktonic foraminiferal I/Ca for
379 samples more recent than 2 ka BP ~~(core-tops only)~~. Tested parameters are listed from highest to lowest absolute
380 Spearman's ρ .

Deleted: 4).

Deleted: .

Variable	n	Spearman ρ	Spearman p-value	Kendall τ	Kendall p-value	Data source
p25_Oxygen*	406	0.78	<0.01	0.58	<0.01	WOD23
min_Oxygen*	406	0.78	<0.01	0.56	<0.01	WOD23
all_depth_mean_Oxygen*	404	0.77	<0.01	0.58	<0.01	WOD23
mean_Oxygen*	406	0.76	<0.01	0.57	<0.01	WOD23
oxycline_oxygen_mean*	224	0.72	<0.01	0.53	<0.01	WOD23
mean_deep_Oxygen*	427	0.65	<0.01	0.45	<0.01	WOD23
max_500m_depth MLD*	431	0.64	<0.01	0.46	<0.01	CMEMS GLORYS12V1
mean_500m_depth MLD*	431	0.63	<0.01	0.44	<0.01	CMEMS GLORYS12V1
mean_Phosphate*	266	-0.62	<0.01	-0.46	<0.01	WOD23
mean_Alkalinity*	71	-0.56	<0.01	-0.40	<0.01	WOD23

NPP_VGPM*	426	-0.44	<0.01	-0.30	<0.01	Orca Ocean Productivity
min_pH*	109	0.42	<0.01	0.26	<0.01	WOD23
Age (ka BP)*	61	-0.39	<0.01	-0.29	<0.01	See database
mean_pH*	109	0.33	<0.01	0.31	<0.01	WOD23
mean_Nitrate*	236	-0.28	<0.01	-0.18	<0.01	WOD23
mean_500m_depth velocity*	431	0.27	<0.01	0.18	<0.01	See database
distance to coast (km)*	431	0.27	<0.01	0.18	<0.01	Calculated
mean_Salinity*	389	0.24	<0.01	0.17	<0.01	See database
Latitude*	431	-0.14	<0.01	-0.10	<0.01	See database
max_Chlorophyll_depth	170	-0.14	0.08	-0.08	0.13	WOD23
Depth in core (cm)*	408	-0.13	<0.01	-0.09	<0.01	See database
oxycline_depth_mean	376	-0.10	0.06	-0.05	0.17	WOD23
max_Chlorophyll	170	-0.08	0.31	-0.08	0.16	WOD23
Water depth (m)	431	0.08	0.11	0.04	0.20	WOD23
max_500m_depth velocity	431	-0.06	0.22	-0.04	0.24	CMEMS GLORYS12V1
mean_Temperature	389	0.00	0.94	0.02	0.50	WOD23
max_Oxygen	406	0.00	0.98	-0.02	0.63	WOD23

383 *: statistically significant correlations

384 All oxygen related variables (minimum, maximum, mean, p25, oxycline mean oxygen concentration, mean (all
385 depths) and bottom mean oxygen) are significantly linked to foraminiferal I/Ca with coefficients above 0.6 and
386 0.4 for Spearman's ρ and Kendall's τ , respectively, and p-values less than 0.01 for both cases (Table 2). Mixed
387 layer depth and mean phosphate concentration show high correlation coefficients, but these parameters are also
388 tightly linked to oxygen (Fig. S2). This observation supports an oxygen control on foraminiferal I/Ca, although
389 the quantitative relationship remains uncertain.

390 Minimum pH, alkalinity, mean nitrate concentration and NPP present correlation with I/Ca (Table 2) but they also
391 covary with p25 [O₂] and are thus not independent (Fig. S2). Distance to coast, mean salinity, latitude and current
392 velocity also show significant correlations with foraminiferal I/Ca. However, their correlation coefficients are less
393 than half those of the strongest correlations identified. These variables are therefore not further discussed as
394 primary drivers of foraminiferal I/Ca variability. He et al. (2026) documented a latitudinal trend in I/Ca that
395 mirrored the iodate/latitude relationship. Similar pattern is found with I/Ca in Fig. S5. However, p25 [O₂] also
396 shows very similar trend with latitude with the two variables being correlated.

397 Depth in core and sediment age are examined to evaluate a potential early diagenesis effect. Most of the core top
398 selected for this study are sampled close to the water-sediment interface and in various environments. Those two
399 variables do show statistically significant correlation with foraminiferal I/Ca on global scale. However, the number
400 of data used for age is low (n=61) and the correlation coefficients for depth in core is low. Water depth at sampling
401 site is not correlated to foraminiferal I/Ca, as it was shown with Fig. 4.

402 2.5 Multi-species foraminiferal I/Ca and calcification depths on global scale

Deleted: Maximum mixed

Deleted: shows the highest

Deleted: this parameter is

Deleted: phosphate

Deleted: maximum chlorophyll depth

Deleted: temperature

Deleted: , maximum

Deleted: , and mean salinity

Deleted: not

Deleted: any

Deleted: This is equally true with water depth above

Deleted: core sites as already examined with the new

Deleted: (Fig. 4), no clear

Deleted: visible.

417 Symbiont presence in foraminifera could also alter the incorporation of specific elements like anions because of
418 so-called “vital effects” (Hönisch et al., 2003). We have classified the symbiosis behaviour of the species and
419 represent their I/Ca as a function of p25 [O₂] (Fig. S3), we note the absence of specific patterns. No group is
420 enriched or depleted in iodine compared to the others. The slopes for linear regression are very close (0.016 or
421 0.017). Symbiont barren species are found in all defined domains, but the low I/Ca plankton tow with p25 [O₂]
422 concentration >250 µmol/kg are exclusively symbiont barren species.

423 In the database, we have also classified species in three zones of calcification depths as the local concentration of
424 iodate changes with depth and influences foraminiferal I/Ca (Hess et al., 2025). Species are classified as surface,
425 shallow subsurface, and deep subsurface dwellers following Hess et al. (2025). Figure S4 presents the linear fit
426 of the I/Ca ratios and p25 [O₂] ~~for Atlantic, Pacific and Indian oceans, which are obtained by classifying~~
427 ~~foraminifera according to their calcification depth. The main result is a similar slope for the three groups in Atlantic~~
428 ~~and Pacific oceans (0.017 to 0.025). Linear regressions for the Indian ocean were not plotted as 48 points are~~
429 ~~represented and the majority are located below 60 µmol/kg. Differences in intercepts are not interpreted due to the~~
430 ~~important impact of scattering. The pattern of lower I/Ca for deep subsurface dwellers in Pacific OMZs (Hess et~~
431 ~~al., 2025) is still visible, but is now located between 100 and 180 µmol/kg due to the use of p25. The Pacific Ocean~~
432 ~~is the only one where this specificity is observed.~~

Deleted:],

Deleted: is

Deleted: (0.015 to 0.019). Differences in intercepts are not interpreted due to the important impact of scattering

Formatted: Strikethrough

433 3 Discussion

434 A gain in iodine in foraminiferal test during their post-mortem settling to the seafloor was proposed to explain low
435 plankton tow values (Winkelbauer et al., 2023). This implies an increase in I/Ca with water depth. However, water
436 depth does not show any significant covariation with foraminiferal I/Ca (Table 2 and Fig. S5). We thus consider
437 that the influence of iodine accumulation during the foraminifera settling would be secondary and could be hidden
438 under a stronger oxygen control. No specific pattern for foraminiferal I/Ca as a function of oxygen can be
439 highlighted by using foraminiferal species, calcification depth groups, or symbiont presence/absence.

440 In the following, we will at first discuss the possible reasons for a larger scatter observed between iodate proportion
441 and p25 [O₂] (Fig. 3). Then, a new calibration of foraminiferal I/Ca ratios against p25 [O₂] will be proposed (Fig.
442 5). The influence of the scattered relationship between iodate proportion and p25 [O₂] on the calibration will be
443 evaluated (Fig. 6). Then, we will quantify residuals that are not supported by oxygenation changes to identify
444 parameters affecting foraminiferal I/Ca (Table S1). A comparison between foraminiferal I/Ca and the value of
445 synthetic calcite will provide us possible impact of vital effects (Fig. 6). Finally, we will examine the contribution
446 of authigenic calcite to the Mediterranean samples (Fig. 7) that are characterised by low I/Ca ratios under oxic
447 condition (Fig. 4).

448

449 3.1 Principal role of oxygen and possible reasons for a larger scatter observed between iodate proportion 450 and p25 [O₂]

451 First order results suggest clear covariation of foraminiferal I/Ca with oxygen and oxygen related variables (Table
452 2 and Fig. S5). Oxygen variables show a positive trend with foraminiferal I/Ca. This covariation ranges from linear

to exponential, and displays substantial scatter. Other parameters exhibit no consistent relationship, except oxygen-linked variables such as nutrients, NPP, and mixed-layer depth (Table 2 and Fig. S2).

Inorganic iodine speciation is theoretically controlled by redox potential, largely governed by oxygen availability (Luther, 2023). In practice, however, field observations frequently deviate from this thermodynamic framework (Fig. 3). The relationship between iodate and p25 [O₂] is strongly non-linear and exhibits important scatter. The dispersion likely reflects spatial and temporal variability in iodine redox cycling across regions and seasons, and water mass mixing. This mismatch between oxygen and iodate concentration may at least partly explain the apparent presence of iodide in waters here defined as well-oxygenated.

As iodine speciation is sensitive to short-term oxygen fluctuations and depth-specific redox structure, this statistical approach better represents the oxygen conditions than relying on the minimum value. The observed scatter likely reflects a combination of true environmental heterogeneity and the limitations of matching discrete iodine measurements with oxygen data. The iodate–oxygen relationship in Fig. 3 shows that iodate proportion does not decline uniformly with decreasing oxygen. Consequently, at low p25 [O₂], the variability in the proportion of iodate can generate substantial scatter in foraminiferal I/Ca, potentially producing a factor of ~5 spread (1.0–5.0 μmol/mol) near 50 μmol/kg. Therefore, uncertainty in inorganic iodine speciation, both estimation errors or natural variability, remains a major source of scatter in the I/Ca–oxygen relationship.

3.2 A new calibration of planktonic foraminiferal I/Ca using p25 [O₂]

In our calibration, we use foraminiferal I/Ca as a function of p25 [O₂] concentration in 0–500 m, a representative value for oxygen concentration at each sampling site. This calibration should be used with caution for quantitative oxygen reconstruction due to substantial residual scatter. Its purpose is rather to explore the extent to which oxygen relationships can explain the observed variance.

Figure 5 shows core top samples more recent than 2 ka BP considering possible temporal changes in dissolved oxygen concentration in the ocean. The linear fit shown in red in Fig. 5 corresponds to a Least Absolute Deviations (LAD) regression. This method was selected because it is more robust to outliers and large deviations than ordinary least squares. Its reliance on medians rather than means makes it particularly suitable when the error structure is non-Gaussian or affected by outliers.

The obtained new calibration equation is as follows:

$$\text{Planktonic foraminiferal I/Ca } (\mu\text{mol/mol}) = 0.021 * p25 [\text{O}_2] (\mu\text{mol/kg}) - 0.058 \quad (1)$$

Equation 1 provides a slightly better fit regarding precedent calibration from Hess et al. (2025) as shown by RMSE (Fig. 5a, b). This comparison integrates the recent results published by Nilsson-Kerr et al. (2025). The main difference between the two calibrations is that the clustered point at near-zero oxygen has been expanded in our case, covering a wider range of oxygen concentrations. Also, the intercept is lower, approaching I/Ca = 0 μmol/mol in strongly oxygen depleted waters (p25 [O₂] < 10 μmol/kg). The points with low I/Ca (< 2 μmol/mol) in oxygenated water correspond to samples from the Mediterranean Sea (this study) and from Southern Benguela (Lu et al., 2020).

Deleted: .

Deleted: iodate

Deleted: oxygen-depleted

Deleted:

Deleted: 0174

Deleted: +

Deleted: 04

Deleted: one

In order to identify secondary parameters that may influence foraminiferal I/Ca, the residuals are calculated using equation (1). The residuals are then plotted against multiple ocean variables (Fig. S6). Table S1 summarises the correlation coefficients and significance of the parameters tested. No clear nor visible trend can be highlighted within the global dataset including water depth, distance to coast, or oxycline depth.




Figure 5: Foraminiferal I/Ca <2 ka BP as a function of minimum (a) and p25 (b) [O₂] in 0–500m. The same dataset is used for the two figures. Only the X-axis definition is different. Black line is the linear model calibration curve from (Hess et al., 2025). The red line is the linear model proposed in this study. Samples with oxygen distribution from 0.25° area are represented as black circles, and from 0.5° as grey squares. Horizontal lines in (a) illustrate the effect of X-axis change on dataset, highlighting that many low [O₂]min sites are shifted to higher representative oxygen depletion p25[O₂]. This is especially true for [O₂]min <50 μmol/kg and I/Ca >2 μmol/mol. No specific pattern emerges within the wider area.

3.3 A comparison of the I/Ca ratio of foraminifera with the value of synthetic calcite.

We further compare the new calibration with the relationship proposed for synthetic calcite (Zhou et al., 2014). The experience of synthetic calcite precipitation in iodate-rich medium indicated a linear relationship between calcite I/Ca and iodate concentration (Lu et al., 2010; Zhou et al., 2014). For the comparison, the iodate concentration in the medium is converted to oxygen concentration using the relationship between iodate proportion and p25 [O₂] concentration (Fig. 3a). Total inorganic iodine concentration is considered to be 471 nM (Fig. 3c) and calcite is assumed to be precipitated at 19°C. The modelled abiotic I/Ca is based on inorganic iodine speciation (Fig. 3a) giving the proportion of iodate at each p25 [O₂] value. Using the mean total inorganic iodine concentration (Fig. 3c) we obtain a mean iodate concentration at different p25 [O₂]. The next step of the model is the incorporation of the iodine in the calcite lattice. Zhou et al. (2014) have synthesised calcite at different iodate concentrations at 6, 19 and 33°C. They showed linear incorporation of iodine in calcite with iodate concentration.

Deleted: or

Deleted: ¶

Deleted: Total inorganic iodine concentration is considered to be 471 nM (Fig. 3c) and calcite is assumed to be precipitated at 19°C.

We choose to use 19 °C and examine the influence of temperature considering the case of 33 and 6 °C. The aim of this model is to investigate the possible I/Ca values considering only abiotic incorporation in calcite to evaluate possible vital effects.

In Figure 6, the synthetic calcite I/Ca relationship is shown in blue, with the 95% prediction interval (blue shaded area). The 95% prediction interval is large, implying a large potential impact on our calibration. This indicates that iodate concentration relative to oxygen plays a key role in generating the observed dispersion in foraminiferal I/Ca.

The modelled abiotic I/Ca only accounts for inorganic iodine speciation and temperature between 6 and 33°C, not for total inorganic iodine concentration.

The abiotic incorporation of iodine is of the same order of magnitude as that found in foraminifera. Iodine incorporation into carbonates likely reflects predominantly abiotic processes.

However, some foraminiferal I/Ca ratio falls out of the prediction interval, both with p25 or minimum oxygen.

Below 200 µmol/kg p25[O₂], most I/Ca values are below the values that can be explained by abiotic calcite precipitation, until severe oxygen depletion (< 20 µmol/kg) where foraminifera I/Ca return to the abiotic level.

The highest foraminiferal I/Ca values (> 6 µmol/mol) are in the highly oxygenated areas (> 180 µmol/kg) and are above the abiotic interval (Fig. 6a). Those observations suggest the presence of unknown parameters that are specific for foraminiferal calcite (e.g. microenvironment, changes to growth rate...). Zhou et al. (2014) showed the temperature dependency of partition coefficient (Kd) of iodine for abiotic calcite. But no systemic behaviour could be highlighted explaining higher or lower foraminiferal I/Ca anomalies with temperature differences (Fig. 6).

The use of min[O₂] leads to similar conclusions with I/Ca values below the 95 % prediction interval of abiotic iodine incorporation (Fig. S7b). The main difference is the high density of data points next to origin, resulting from the definition of the x-axis.

High I/Ca ratios may reflect the presence of other calcium carbonate phases, as aragonite incorporates more iodine than calcite (Hardisty et al., 2017; Lu et al., 2022). The highest I/Ca value in the Mediterranean Sea is found in the Gulf of Sirte in core SL95. This area shows high aragonite concentrations due to the presence of *Halimeda* (Reitz and De Lange, 2006). Remaining aragonite needles after cleaning may result in higher I/Ca. We investigate the influence of the presence of aragonite with binary mixing between aragonite and foraminifera. We consider an I/Ca of 10 µmol/mol in aragonite, similar as the ratio measured in scleractinian corals (Sun et al. 2023). For foraminiferal I/Ca, we consider an expected value of 4 µmol/mol using the new calibration and an oxygen concentration at SL95 site (Fig. 6). Aragonite contribution should be around 40% to account for an I/Ca between 6 and 7 µmol/mol, a similar range of the observed I/Ca of 6.9 µmol/mol at site SL95. Sr/Ca of these mixtures would be 4.8 and 3.9 mmol/mol, respectively, considering a Sr/Ca of 10 mmol/mol for aragonite and 1.3 mmol/mol for foraminifera (Hathorne et al., 2013; Lea et al., 1999). As the Sr/Ca from SL95 core-top is 1.345 mmol/mol, the mixture does not correspond to the observed value. The contribution of aragonite alone cannot explain the measured I/Ca value.

To summarise, inorganic iodine speciation has a non-linear relationship with dissolved oxygen concentration although the observed pattern for foraminiferal I/Ca does not follow the exponential trend that is expected from the relationship between iodate proportion and p25 [O₂] (Fig. 6a). Low I/Ca values found in oxygenated areas might be caused by reduced iodine incorporation, related to unknown mechanisms/conditions. Since no mechanism has yet been identified to justify a specific fitting curve for I/Ca ratio, we consider that the linear

Deleted: foraminifera

Deleted: .

Deleted: ,

Deleted: area (>

Deleted: 6

Deleted: 6

Deleted: area

576 regression is a reasonable approximation. The dispersion of I/Ca values might be caused by the relationship
577 between iodine speciation and oxygen concentration that could account for ~70% of the variance compatible with
578 the model's 95% prediction interval (assuming constant total inorganic iodine), and up to ~87% by considering
579 $\pm 1\sigma$ uncertainty in total inorganic iodine from Fig. 4b (Fig. S8a). This dispersion likely represents the second key
580 explanation for I/Ca values in foraminifera. Given its systematic offset from the predicted values, low I/Ca in
581 highly oxygenated water seems to be related to an incorporation mechanism which remains unknown. No
582 geographical trend could be highlighted for I/Ca below the 95% prediction interval using p25 (Fig. 6) or using the
583 minimum (Fig. 8c and d) oxygen. Temperature may also affect the I/Ca (Zhou et al., 2014). However, Gulf of
584 Cadix I/Ca are 0.75 and 0.69 $\mu\text{mol/mol}$ for *G. ruber* and *G. bulloides*, respectively. The two species are known to
585 calcify in different seasons, where minimum temperature measured is 11.26°C, and maximum is 24.21°C in 0–
586 500 m. At ODP Site 977, I/Ca from *G. bulloides* is 0.69 and from *G. ruber* is 1.64 $\mu\text{mol/mol}$. This is the inverse
587 of what the temperature effect predicts, with here the warm season calcifier having a higher I/Ca. So, the
588 temperature effect may not explain all the observed values below 95% prediction interval of abiotic I/Ca. This
589 offset could potentially be related to the vital effects of discriminating iodine incorporation, or to local iodine
590 speciation itself. Alternatively, it could be due to the mixing of iodate-depleted water in oxygenated water masses.
591 However, iodate/oxygen-depleted water cannot be identified close to the Gulf of Cadix (Fig. S8b and d).

Deleted: an appropriate

Deleted: 75

Deleted: 93

Deleted: S7

Deleted: explanation

Deleted: However, it remains that

Deleted: that

Deleted: , given its systematic offset from the predicted values....

Deleted: is

Deleted: .




Figure 6: Foraminiferal I/Ca < 2 ka BP as a function of p25 [O₂] in 0–500 m. The red line is the linear model proposed in this study. Blue curve represents the abiotic incorporation of iodine following iodate proportion in total inorganic iodine determined in Figure 3 and assuming a mean concentration of total inorganic iodine of 471 nM (Fig. 3c). The abiotic incorporation was determined following Zhou et al. (2014). Blue area corresponds to 95% prediction interval also determined in Figure 4 for a temperature of 19 °C. 61% of the data fall in this prediction interval. Temperature from 6 to 33 °C are considered for the maximum and minimum 95% prediction interval (grey dashed lines).

Deleted: The abiotic incorporation was determined following (Zhou et al., 2014)

Deleted: 57

3.4 Authigenic contribution via diagenetic calcite precipitation

Deleted: secondary

Depth in core and age were not significantly correlated with foraminiferal I/Ca based on first-order statistical tests (Table 2). However, these parameters do not fully capture the potential influence of diagenesis or diagenetic calcite precipitation, which may be particularly important within the sediments from within OMZs and the Mediterranean

Deleted: secondary

Deleted: OMZ

622 Sea which are characterised by high alkalinity pore/bottom waters. This section therefore examines whether early
623 diagenetic carbonate precipitation could contribute to the observed variability in I/Ca.

624 Because authigenic carbonates typically have very low I/Ca (Lu et al., 2010), the lower I/Ca value of MD04-2876
625 (Arabian Sea, OMZ core) relative to MD04-2873 (below the OMZ) could partly reflect the presence of authigenic
626 carbonates formed under reducing conditions. In the marine sediments of the OMZ, the increase in alkalinity of
627 the pore water due to anaerobic respiration leads to the precipitation of authigenic carbonates (Suess, 1979; Meister
628 et al., 2022). In the Mediterranean Sea, high water-column alkalinity driven by excess evaporation (Schneider et
629 al., 2007) promotes authigenic carbonate precipitation, as shown by high Mg/Ca ratios in foraminifera (Boussetta
630 et al., 2011).

631 Figure 7 presents p25 [O₂] alongside the expected foraminiferal I/Ca predicted from Equation (1) with a red line.
632 Sensitivity tests were performed using a binary mixing model between the expected foraminiferal I/Ca and
633 authigenic carbonates assuming an I/Ca of 0.06 µmol/mol for authigenic carbonate (Lu et al., 2010). For the
634 Arabian Sea OMZ core, measured I/Ca values fall below the expected ratio and could correspond to a 30–40%
635 mass contribution of low I/Ca diagenetic carbonate. In contrast, MD04-2873 (below the OMZ) displays higher
636 I/Ca than expected, likely due to poorly constrained oxygen variability resulting from the limited number of
637 available oxygen profiles (section 3.2).

638 In the Mediterranean Sea, the two eastern basin cores (SL95 and MD04-2724) show I/Ca values above expectation,
639 while others fall below. Cores MD04-2722, MD04-2797 and MD90-917 are slightly depleted in I/Ca relative to
640 the value estimated from equation (1) and could reflect a 20–30% contribution of authigenic carbonates. However,
641 the samples from the western basin (ODP Site 977) and from the Gulf of Cadiz (MD99-2341) would require an
642 implausibly high 60–90% contribution of authigenic calcite to explain their (much-)lower-than-expected I/Ca.
643 Such a contribution is unrealistic, as authigenic calcite precipitation is far less common in the western basin, and
644 extensive overgrowths would have been excluded (Crudeli et al., 2004).

645 Overall, diagenetic calcite precipitation cannot fully explain the lower I/Ca deviations. Low I/Ca values occur in
646 regions where early diagenetic calcite formation is low, while no mechanism has been identified to account for
647 I/Ca values higher than expected, nor for the scatter observed among species dwelling at the same depth within
648 the same core top samples, except for as yet undefined vital effects.

649 We conclude that foraminiferal I/Ca primarily reflects local oxygen variability through its control on iodine
650 speciation. The Mediterranean Sea samples exhibit a similar behaviour and degree of scatter between I/Ca and p25
651 [O₂] as those from other ocean basins, indicating that the proxy remains applicable in this semi-enclosed
652 environment. While foraminiferal I/Ca provides a semi-quantitative estimation of local oxygenation conditions,
653 the core-top values observed in the western Mediterranean Sea is lower than the values expected from the
654 calibration. The use of the proxy must focus on relative variability combined with another independent approach.
655 The relationship between foraminiferal I/Ca and oxygen concentration may be site-specific, because it could be
656 influenced by local environmental characteristics or processes that are not yet fully understood. This limitation is
657 particularly relevant for coastal sites, including those within the Mediterranean Sea, where additional factors may
658 modulate iodine speciation or iodine incorporation.

Deleted: 6

Deleted: black

Deleted: secondary

Deleted: 10–

Deleted: secondary

Deleted: However, the




Figure 7: Foraminiferal I/Ca from the Mediterranean and Arabian Sea as a function of p25 [O₂] 0-500 m in whole p25 [O₂] range (a) and for p25 [O₂] >180 μmol/kg (b). Expected foraminiferal I/Ca from LAD from figure 5 is represented as red line. Blue dotted lines indicates deviation from expected value by increasing authigenic carbonate contribution from 10 to 90 %. Authigenic carbonate I/Ca was fixed at 0.06 μmol/mol following Lu et al. (2010).

3.5 Other possible factors

Oceanographic variables investigated other than oxygen did not show strong relationship with foraminiferal I/Ca (Table 2). Other parameters may also influence iodine speciation but were not highlighted through the statistical analysis conducted on foraminiferal I/Ca in this study. Previous works have shown that iodine speciation varies with distance to the coast, with iodate depletion toward productive coastal water (Wong and Zhang, 1992). The effect of distance to coast is absent on global scale, but it could be important at regional scale, such as in the Mediterranean Sea. This depletion is thought to be driven by enhanced biological iodine uptake through primary productivity. Although the Mediterranean Sea is highly oligotrophic, coastal and shelf regions can sustain episodes of high primary productivity (Packard et al., 1988; Powley et al., 2017), which could potentially influence iodate concentration. Only limited inorganic iodine vertical profiles (iodate and iodide) exists in the Mediterranean Sea (Fig. 1b), with one time series in the north western basin (Tian and Nicolas, 1995). The study showed limited variability across the year with iodate concentrations of 410–480 nM, but iodine speciation was measured in different oceanographic settings as the investigated locations in this study. Even though different inorganic iodine behaviours cannot be ruled out, the variability of iodate concentration observed in the north western Mediterranean Sea, about 10%, does not explain the scattering in I/Ca observed.

Part of the variability may also not be fully constrained with the statistical analysis conducted (Tables 2 and S1, Figures S2 and S5). In fact, we conducted the analysis on the global dataset, with every site being a unique context

Deleted: black

Deleted: Grey

Deleted: (Packard et al., 1988; Powley et al., 2017)

Deleted: So,

Deleted: may occur. Moreover

694 and oceanographic settings. If we highlighted the first and most important role of oxygen as an I/Ca control,
695 secondary parameters may not be fully captured. Inorganic iodine speciation remains poorly understood. Also, the
696 influence of growth rate (calcification kinetics) could not be ruled out, and was never tested for I/Ca ratio (e.g.
697 Allen et al., 2016). This has not been demonstrated but competition with other anions remains plausible as coupled
698 substitution is required for the incorporation of IO_3^- into CaCO_3 (Ram and Erez, 2023).

699 In light of these observations, we emphasise the use of foraminiferal I/Ca as a semi-quantitative proxy for changes
700 in ocean oxygenation. The proxy primarily responds to oxygen and oxygen-related parameters (Table 2, Fig. S5,
701 Table S1 and Fig. S6). The $\text{p25}[\text{O}_2]$ gave the best fit with less data dispersion (Fig. S1). The dispersion of the
702 relationship between inorganic iodine speciation and $\text{p25}[\text{O}_2]$ partly accounts for mixing and differences in
703 inorganic iodine redox kinetics. For these reasons together with potential non-investigated parameters,
704 foraminiferal I/Ca alone cannot be interpreted as a quantitative proxy. Combining it with other proxies and/or a
705 modelling approach is highly recommended. However, it may still be highly informative about changes in oxygen
706 at a specific site. The calibration is useful, not for absolute oxygen reconstruction, but to understand site-specific
707 I/Ca variation relative to the observed oxygen range and identify sites where high I/Ca in oxygenated waters
708 provides the best sensitivity to past oxygenation change. However, most studies focused oxygen depleted areas,
709 leading to less representation of oxygenated and high latitudes sites in the dataset. Wang et al. (2026) proposed
710 oxygen thresholds to interpret foraminiferal I/Ca. However, the current global data set would be biased towards
711 oxygen-depleted areas. Our Mediterranean core-tops data show a wide I/Ca range in oxygenated waters, which
712 cannot be explained by authigenic calcite precipitation or aragonitic carbonate presence, suggesting additional
713 scatter introduced in oxygenated settings (e.g., vital effects, water mass mixing, local inorganic iodine speciation).
714 Further core-top data spanning the whole oxygenation range is needed to validate the threshold behaviour and
715 determine the reason for low foraminiferal I/Ca in oxygenated water.

716 For future work, foraminifera culture with isolation of single parameters/monitoring of multiple parameters could
717 help better constrain the proxy behaviour, alongside field studies. In particular, plankton tow observations showing
718 systematically low foraminiferal I/Ca values suggest that limited incorporation at the surface may happen, leaving
719 open the question of when and where iodine is incorporated. Inorganic iodine speciation is still poorly constrained
720 and further research is needed to understand its dynamics and incorporation in calcite lattice. **Iodine incorporation**
721 **sites and mechanisms, are hardly needed to distinguish the origin of the signal and better understand the observed**
722 **trends.**

723

724 Conclusions

725 In this study, we investigate planktonic foraminiferal I/Ca as a proxy for subsurface oxygen changes by integrating
726 multiple environmental parameters. We first proposed a new definition of a representative oxygen depletion value
727 in subsurface waters, accounting for local oxygen variability. Local oxygen distributions were unimodal or
728 multimodal, reflecting both seasonal and depth-related oxygen variation. Planktonic foraminiferal I/Ca from core
729 top samples showed a positive correlation with this representative minimum oxygen value, $\text{p25}[\text{O}_2]$, supporting
730 the dominant control of oxygen on iodine presence and reinforcing the utility of the proxy to reconstruct past
731 oxygenation changes.

Deleted: , inside the structure of the shell or outside, as secondary precipitation, is

Deleted: iodine provenance in

Deleted: foraminiferal test

736 Despite this relationship, substantial amount of data dispersion remains. We explored several potential sources of
737 scatter and found that the largest contribution likely arises from variability in inorganic iodine speciation. The
738 global relationship between iodate and p25 [O₂] alone could account for more than 85% of the observed scatter.

739 No first-order relationship was found between water depth and foraminiferal I/Ca. This observation implies that
740 iodine gain during particle sinking (Winkelbauer et al., 2023) could not be confirmed.

741 We propose a new calibration for subsurface oxygen depletion reconstruction based on planktonic foraminiferal
742 I/Ca. By using p25 [O₂] values, we reduce the dispersion observed in previous calibrations. To maintain simplicity
743 for paleo-applications, core top data younger than 2 ka BP were included regardless of species. Remaining
744 dispersion may result from vital (integration of iodate into the foraminifera shell) or species-specific effects,
745 (difference in iodate integration between species), or other uninvestigated parameters, such as inorganic iodine
746 speciation, modulated by water mass mixing, NPP, or differences in oxidoreduction kinetics.

747 Finally, diagenetic calcite contribution was considered as a potential contributor to the low foraminiferal I/Ca
748 values, given that diagenetic calcite formed in high alkalinity environment could dilute the signal. However, such
749 an effect would require more than 60% of diagenetic calcite to explain western Mediterranean Sea values, which
750 is implausible.

751 We therefore support the use of planktonic foraminiferal I/Ca as a semi-quantitative proxy for oxygenation state,
752 consistent with Hess et al. (2025). This proxy can also be applied to semi-enclosed basins to reconstruct relative
753 variability when complemented with another independent approach. However, foraminiferal I/Ca cannot be used
754 as a quantitative proxy for oxygen reconstruction. The tool must be used alongside another approach to support
755 past environmental change. These recommendations are still prone to further evolution as we better constrain the
756 proxy and the inorganic iodine cycle in the ocean.

757

758 **Appendices**

759 **Data availability**

760 Data supporting this paper will be available in the following repository: <https://doi.org/10.5281/zenodo.18390413>

761 The data is under review, the DOI is not activated yet.

762 **Author contribution**

763 KT and VG designed the study; KT, MG and VG performed the laboratory procedures; VG have conducted the
764 statistical analysis. VG wrote the manuscript with contributions from all authors.

765 **Competing interests**

766 The authors declare that they have no known competing financial interests or personal relationships that could
767 have appeared to influence the work reported in this paper.

768 **Acknowledgements**

Deleted: ,

Deleted: , which is difficult to constrain.

Deleted: secondary

Deleted: secondary

Deleted: secondary

774 This work is supported by the French national project ANR MedSens (ANR-19-CE01-0019) granted to KT.
 775 Radiocarbon dating was performed at the LMC14 laboratory as a part of the French ARTEMIS program. Sample
 776 preparation and analytical measurements were supported by the LPM platform (Sandrine Conrod), the LGO2i
 777 platform, and the ENVITOP platform. We also thanks [Rosie Chance and Helge A. Winkelbauer](#) for providing
 778 global iodine speciation dataset, and Rosalind E.M. Rickaby for the inspiring discussions on the topic. [We thank](#)
 779 [Dalton Hardisty and the anonymous referee for their comments which greatly helped improve the manuscript.](#)

780 Financial support

781 References

- 782 Allen, K.A., Hönisch, B., Eggins, S.M., Haynes, L.L., Rosenthal, Y., Yu, J., 2016. Trace element proxies
 783 for surface ocean conditions: A synthesis of culture calibrations with planktic foraminifera.
 784 *Geochim. Cosmochim. Acta* 193, 197–221. <https://doi.org/10.1016/j.gca.2016.08.015>
- 785 Bard, E., Ménot, G., Rostek, F., Licari, L., Böning, P., Edwards, R.L., Cheng, H., Wang, Y., Heaton, T.J.,
 786 2013. Radiocarbon Calibration/Comparison Records Based on Marine Sediments from the
 787 Pakistan and Iberian Margins. *Radiocarbon* 55, 1999–2019.
 788 https://doi.org/10.2458/azu_js_rc.55.17114
- 789 Barker, S., Greaves, M., Elderfield, H., 2003. A study of cleaning procedures used for foraminiferal
 790 Mg/Ca paleothermometry: MG/CA PALEOTHERMOMETRY. *Geochem. Geophys. Geosystems*
 791 4, n/a-n/a. <https://doi.org/10.1029/2003GC000559>
- 792 Behrenfeld, M.J., Falkowski, P.G., 1997. Photosynthetic rates derived from satellite - based
 793 chlorophyll concentration. *Limnol. Oceanogr.* 42, 1–20.
 794 <https://doi.org/10.4319/lo.1997.42.1.0001>
- 795 Bopp, L., Resplandy, L., Orr, J.C., Doney, S.C., Dunne, J.P., Gehlen, M., Halloran, P., Heinze, C., Ilyina,
 796 T., Séférian, R., Tjiputra, J., Vichi, M., 2013. Multiple stressors of ocean ecosystems in the
 797 21st century: projections with CMIP5 models. *Biogeosciences* 10, 6225–6245.
 798 <https://doi.org/10.5194/bg-10-6225-2013>
- 799 Boussetta, S., Bassinot, F., Sabbatini, A., Caillon, N., Nouet, J., Kallel, N., Rebaubier, H., Klinkhammer,
 800 G., Labeyrie, L., 2011. Diagenetic Mg-rich calcite in Mediterranean sediments: Quantification
 801 and impact on foraminiferal Mg/Ca thermometry. *Mar. Geol.* 280, 195–204.
 802 <https://doi.org/10.1016/j.margeo.2010.12.011>
- 803 Chai, J.Y., Muramatsu, Y., 2007. Determination of Bromine and Iodine in Twenty-three Geochemical
 804 Reference Materials by ICP-MS. *Geostand. Geoanalytical Res.* 31, 143–150.
 805 <https://doi.org/10.1111/j.1751-908X.2007.00856.x>
- 806 Chance, R., Baker, A.R., Carpenter, L., Jickells, T.D., 2014. The distribution of iodide at the sea surface.
 807 *Env. Sci Process. Impacts* 16, 1841–1859. <https://doi.org/10.1039/C4EM00139G>
- 808 Chance, R., [Tinel, L., Sarkar, A., Sinha, A.K., Mahajan, A.S., Chacko, R., Sabu, P., Roy, R., Jickells, T.D.,](#)
 809 [Stevens, D.P., Wadley, M., Carpenter, L.J., 2020. Surface Inorganic Iodine Speciation in the](#)
 810 [Indian and Southern Oceans From 12°N to 70°S. Front. Mar. Sci. 7, 621.](#)
 811 <https://doi.org/10.3389/fmars.2020.00621>
- 812 [Chance, R.,](#) Weston, K., Baker, A.R., Hughes, C., Malin, G., Carpenter, L., Meredith, M.P., Clarke, A.,
 813 Jickells, T.D., Mann, P., Rossetti, H., 2010. Seasonal and interannual variation of dissolved
 814 iodine speciation at a coastal Antarctic site. *Mar. Chem.* 118, 171–181.
 815 <https://doi.org/10.1016/j.marchem.2009.11.009>
- 816 Chance, R.J., Tinel, L., Sherwen, T., Baker, A.R., Bell, T., Brindle, J., Campos, M.L.A.M., Croot, P.,
 817 Ducklow, H., Peng, H., Hopkins, F., Hoogakker, B., Hughes, C., Jickells, T.D., Loades, D.,
 818 Macaya, D.A.R., Mahajan, A.S., Malin, G., Phillips, D., Roberts, I., Roy, R., Sarkar, A., Sinha,
 819 A.K., Song, X., Winkelbauer, H., Wuttig, K., Yang, M., Peng, Z., Carpenter, L.J., 2019. Global
 820 sea-surface iodide observations, 1967–2018. *Sci. Data* 6, 286.
 821 <https://doi.org/10.1038/s41597-019-0288-y>

Cheng, K., Ridgwell, A., Hardisty, D.S., 2024. Characterizing the marine iodine cycle and its relationship to ocean deoxygenation in an Earth system model. *Biogeosciences* 21, 4927–4949. <https://doi.org/10.5194/bg-21-4927-2024>

Cornuault, M., Tachikawa, K., Vidal, L., Guihou, A., Siani, G., Deschamps, P., Bassinot, F., Revel, M., 2018. Circulation Changes in the Eastern Mediterranean Sea Over the Past 23,000 Years Inferred From Authigenic Nd Isotopic Ratios. *Paleoceanogr. Paleoclimatology* 33, 264–280. <https://doi.org/10.1002/2017PA003227>

Crudeli, D., Young, J.R., Erba, E., De Lange, G.J., Henriksen, K., Kinkel, H., Slomp, C.P., Ziveri, P., 2004. Abnormal carbonate diagenesis in Holocene–late Pleistocene sapropel-associated sediments from the Eastern Mediterranean; evidence from *Emiliana huxleyi* coccolith morphology. *Mar. Micropaleontol.* 52, 217–240. <https://doi.org/10.1016/j.marmicro.2004.04.010>

Elderfield, H., Truesdale, V.W., 1980. On the biophilic nature of iodine in seawater. *Earth Planet. Sci. Lett.* 50, 105–114. [https://doi.org/10.1016/0012-821X\(80\)90122-3](https://doi.org/10.1016/0012-821X(80)90122-3)

Evans, N., Boles, E., Kwiecinski, J.V., Mullen, S., Wolf, M., Devol, A.H., Moriyasu, R., Nam, S., Babbín, A.R., Moffett, J.W., 2020. The role of water masses in shaping the distribution of redox active compounds in the Eastern Tropical North Pacific oxygen deficient zone and influencing low oxygen concentrations in the eastern Pacific Ocean. *Limnol. Oceanogr.* 65, 1688–1705. <https://doi.org/10.1002/lno.11412>

Eynaud, F., De Abreu, L., Voelker, A., Schönfeld, J., Salgueiro, E., Turon, J., Penaud, A., Toucanne, S., Naughton, F., Sánchez Goñi, M.F., Malaizé, B., Cacho, I., 2009. Position of the Polar Front along the western Iberian margin during key cold episodes of the last 45 ka. *Geochim. Geophys. Geosystems* 10, 2009GC002398. <https://doi.org/10.1029/2009GC002398>

Farrenkopf, A.M., Luther III, G.W., 2002. Iodine chemistry reflects productivity and denitrification in the Arabian Sea: evidence for flux of dissolved species from sediments of western India into the OMZ. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 49, 2303–2318. [https://doi.org/10.1016/S0967-0645\(02\)00038-3](https://doi.org/10.1016/S0967-0645(02)00038-3)

Feng, X., Redfern, S.A.T., 2018. Iodate in calcite, aragonite and vaterite CaCO₃: Insights from first-principles calculations and implications for the I/Ca geochemical proxy. *Geochim. Cosmochim. Acta* 236, 351–360. <https://doi.org/10.1016/j.gca.2018.02.017>

Frölicher, T.L., Rodgers, K.B., Stock, C.A., Cheung, W.W.L., 2016. Sources of uncertainties in 21st century projections of potential ocean ecosystem stressors. *Glob. Biogeochem. Cycles* 30, 1224–1243. <https://doi.org/10.1002/2015GB005338>

Hardisty, D.S., Horner, T.J., Evans, N., Moriyasu, R., Babbín, A.R., Wankel, S.D., Moffett, J.W., Nielsen, S.G., 2021. Limited iodate reduction in shipboard seawater incubations from the Eastern Tropical North Pacific oxygen deficient zone. *Earth Planet. Sci. Lett.* 554, 116676. <https://doi.org/10.1016/j.epsl.2020.116676>

Hardisty, D.S., Lu, Z., Bekker, A., Diamond, C.W., Gill, B.C., Jiang, G., Kah, L.C., Knoll, A.H., Loyd, S.J., Osburn, M.R., Planavsky, N.J., Wang, C., Zhou, X., Lyons, T.W., 2017. Perspectives on Proterozoic surface ocean redox from iodine contents in ancient and recent carbonate. *Earth Planet. Sci. Lett.* 463, 159–170. <https://doi.org/10.1016/j.epsl.2017.01.032>

Hathorne, E.C., Gagnon, A., Felis, T., Adkins, J., Asami, R., Boer, W., Caillon, N., Case, D., Cobb, K.M., Douville, E., deMenocal, P., Eisenhauer, A., Garbe-Schönberg, D., Geibert, W., Goldstein, S., Hughen, K., Inoue, M., Kawahata, H., Kölling, M., Cornec, F.L., Linsley, B.K., McGregor, H.V., Montagna, P., Nurhati, I.S., Quinn, T.M., Raddatz, J., Rebaubier, H., Robinson, L., Sadekov, A., Sherrell, R., Sinclair, D., Tudhope, A.W., Wei, G., Wong, H., Wu, H.C., You, C., 2013. Interlaboratory study for coral Sr/Ca and other element/Ca ratio measurements. *Geochim. Geophys. Geosystems* 14, 3730–3750. <https://doi.org/10.1002/ggge.20230>

He, R., Pohl, A., Zhang, X., Chang, C., Prow-Fleischer, A., Payne, J.L., Xiao, S., Ridgwell, A., Lu, Z., 2026. A reversed latitudinal ocean oxygen gradient in the Proterozoic Eon. *Nat. Geosci.* 19, 325–330. <https://doi.org/10.1038/s41561-025-01896-w>

- Hepach, H., Hughes, C., Hogg, K., Collings, S., Chance, R., 2020. Senescence as the main driver of iodide release from a diverse range of marine phytoplankton. *Biogeosciences* 17, 2453–2471. <https://doi.org/10.5194/bg-17-2453-2020>
- Hess, A.V., Rosenthal, Y., Zhou, X., Bu, K., 2025. The I/Ca paleo-oxygenation proxy in planktonic foraminifera: A multispecies core-top calibration. *Geochim. Cosmochim. Acta* S0016703725000298. <https://doi.org/10.1016/j.gca.2025.01.018>
- Hönisch, B., Bijma, J., Russell, A.D., Spero, H.J., Palmer, M.R., Zeebe, R.E., Eisenhauer, A., 2003. The influence of symbiont photosynthesis on the boron isotopic composition of foraminifera shells. *Mar. Micropaleontol.* 49, 87–96. [https://doi.org/10.1016/S0377-8398\(03\)00030-6](https://doi.org/10.1016/S0377-8398(03)00030-6)
- Hoogakker, B.A.A., Davis, C., Wang, Y., Kusch, S., Nilsson-Kerr, K., Hardisty, D.S., Jacobel, A., Reyes Macaya, D., Glock, N., Ni, S., Sepúlveda, J., Ren, A., Auderset, A., Hess, A.V., Meissner, K.J., Cardich, J., Anderson, R., Barras, C., Basak, C., Bradbury, H.J., Brinkmann, I., Castillo, A., Cook, M., Costa, K., Choquel, C., Diz, P., Donnenfeld, J., Elling, F.J., Erdem, Z., Filipsson, H.L., Garrido, S., Gottschalk, J., Govindankutty Menon, A., Groeneveld, J., Hallmann, C., Hendy, I., Hennekam, R., Lu, W., Lynch-Stieglitz, J., Matos, L., Martínez-García, A., Molina, G., Muñoz, P., Moretti, S., Morford, J., Nuber, S., Radionovskaya, S., Raven, M.R., Somes, C.J., Studer, A.S., Tachikawa, K., Tapia, R., Tetard, M., Vollmer, T., Wang, X., Wu, S., Zhang, Y., Zheng, X.-Y., Zhou, Y., 2025. Reviews and syntheses: Review of proxies for low-oxygen paleoceanographic reconstructions. *Biogeosciences* 22, 863–957. <https://doi.org/10.5194/bg-22-863-2025>
- Jickells, T.D., Boyd, S.S., Knap, A.H., 1988. Iodine cycling in the Sargasso Sea and the Bermuda inshore waters. *Mar. Chem.* 24, 61–82. [https://doi.org/10.1016/0304-4203\(88\)90006-0](https://doi.org/10.1016/0304-4203(88)90006-0)
- Jones, M.R., Chance, R., Bell, T., Jones, O., Loades, D.C., May, R., Tinel, L., Weddell, K., Widdicombe, C., Carpenter, L.J., 2024. Iodide, iodate & dissolved organic iodine in the temperate coastal ocean. *Front. Mar. Sci.* 11, 1277595. <https://doi.org/10.3389/fmars.2024.1277595>
- Lea, D.W., Mashiotta, T.A., Spero, H.J., 1999. Controls on magnesium and strontium uptake in planktonic foraminifera determined by live culturing. *Geochim. Cosmochim. Acta* 63, 2369–2379. [https://doi.org/10.1016/S0016-7037\(99\)00197-0](https://doi.org/10.1016/S0016-7037(99)00197-0)
- Lellouche, Eric, G., Romain, B.-B., Gilles, G., Angélique, M., Marie, D., Clément, B., Mathieu, H., Olivier, L.G., Charly, R., Tony, C., Charles-Emmanuel, T., Florent, G., Giovanni, R., Mounir, B., Yann, D., Pierre-Yves, L.T., 2021. The Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis. *Front. Earth Sci.* 9, 698876. <https://doi.org/10.3389/feart.2021.698876>
- Lu, W., Dickson, A.J., Thomas, E., Rickaby, R.E.M., Chapman, P., Lu, Z., 2020. Refining the planktic foraminiferal I/Ca proxy: Results from the Southeast Atlantic Ocean. *Geochim. Cosmochim. Acta* 287, 318–327. <https://doi.org/10.1016/j.gca.2019.10.025>
- Lu, W., Wang, Y., Oppo, D.W., Nielsen, S.G., Costa, K.M., 2022. Comparing paleo-oxygenation proxies (benthic foraminiferal surface porosity, I/Ca, authigenic uranium) on modern sediments and the glacial Arabian Sea. *Geochim. Cosmochim. Acta* 331, 69–85. <https://doi.org/10.1016/j.gca.2022.06.001>
- Lu, Z., Hoogakker, B.A.A., Hillenbrand, C.-D., Zhou, X., Thomas, E., Gutchess, K.M., Lu, W., Jones, L., Rickaby, R.E.M., 2016. Oxygen depletion recorded in upper waters of the glacial Southern Ocean. *Nat. Commun.* 9.
- Lu, Z., Jenkyns, H.C., Rickaby, R.E.M., 2010. Iodine to calcium ratios in marine carbonate as a paleo-redox proxy during oceanic anoxic events. *Geology* 38, 1107–1110. <https://doi.org/10.1130/G31145.1>
- Luther, G.W., 2023. Review on the physical chemistry of iodine transformations in the oceans. *Front. Mar. Sci.* 10, 1085618. <https://doi.org/10.3389/fmars.2023.1085618>
- Martrat, B., Grimalt, J.O., Lopez-Martinez, C., Cacho, I., Sierro, F.J., Flores, J.A., Zahn, R., Canals, M., Curtis, J.H., Hodell, D.A., 2004. Abrupt Temperature Changes in the Western Mediterranean over the Past 250,000 Years. *Science* 306, 1762–1765. <https://doi.org/10.1126/science.1101706>
- Meister, P., Herda, G., Petrishcheva, E., Gier, S., Dickens, G.R., Bauer, C., Liu, B., 2022. Microbial Alkalinity Production and Silicate Alteration in Methane Charged Marine Sediments:

924 Implications for Porewater Chemistry and Diagenetic Carbonate Formation. *Front. Earth Sci.*
925 9, 756591. <https://doi.org/10.3389/feart.2021.756591>

926 Mishonov, A.V., Boyer, T.P., Baranova, O.K., Bouchard, C.N., Cross, S.L., Garcia, H.E., Locarnini, R.A.,
927 Paver, C.R., Reagan, J.R., Wang, Z., Seidov, D., Grodsky, A.I., Beauchamp, J.G., 2024. World
928 Ocean Database 2023 [WWW Document]. <https://doi.org/10.25923/z885-h264>

929 [Moriyasu, R., Evans, N., Bolster, K.M., Hardisty, D.S., Moffett, J.W., 2020. The Distribution and Redox
930 Speciation of Iodine in the Eastern Tropical North Pacific Ocean. *Glob. Biogeochem. Cycles*
931 34, e2019GB006302. <https://doi.org/10.1029/2019GB006302>](#)

932 [NASA Earth Data Base, 2025. ORCA Science. Ocean Productivity Data Server.](#) [WWW Document]. Or.
933 State Univ. ORCA. URL <http://orca.science.oregonstate.edu/index.php> (accessed 6.17.25).

934 Nilsson-Kerr, K., Hoogakker, B.A.A., Macaya, D.A.R., Winkelbauer, H.A., Hamilton, E., Chenery, S.,
935 Davis, C.V., Leng, M.J., 2025. Late Cenozoic intensification of deoxygenation in the Pacific
936 Ocean. *Earth Planet. Sci. Lett.* 655, 119253. <https://doi.org/10.1016/j.epsl.2025.119253>

937 Oschlies, A., Brandt, P., Stramma, L., Schmidtke, S., 2018. Drivers and mechanisms of ocean
938 deoxygenation. *Nat. Geosci.* 11, 467–473. <https://doi.org/10.1038/s41561-018-0152-2>

939 Oschlies, A., Duteil, O., Getzlaff, J., Koeve, W., Landolfi, A., Schmidtke, S., 2017. Patterns of
940 deoxygenation: sensitivity to natural and anthropogenic drivers. *Philos. Trans. R. Soc. Math.*
941 *Phys. Eng. Sci.* 375, 20160325. <https://doi.org/10.1098/rsta.2016.0325>

942 Packard, T.T., Minas, H.J., Coste, B., Martinez, R., Bonin, M.C., Gostan, J., Garfield, P., Christensen, J.,
943 Dortch, Q., Minas, M., Copin-Montegut, G., Copin-Montegut, C., 1988. Formation of the
944 Alboran oxygen minimum zone. *Deep Sea Res. Part Oceanogr. Res. Pap.* 35, 1111–1118.
945 [https://doi.org/10.1016/0198-0149\(88\)90003-9](https://doi.org/10.1016/0198-0149(88)90003-9)

946 Pichevin, L., Bard, E., Martinez, P., Billy, I., 2007. Evidence of ventilation changes in the Arabian Sea
947 during the late Quaternary: Implication for denitrification and nitrous oxide emission. *Glob.*
948 *Biogeochem. Cycles* 21, 2006GB002852. <https://doi.org/10.1029/2006GB002852>

949 Podder, J., Lin, J., Sun, W., Botis, S.M., Tse, J., Chen, N., Hu, Y., Li, D., Seaman, J., Pan, Y., 2017. Iodate
950 in calcite and vaterite: Insights from synchrotron X-ray absorption spectroscopy and first-
951 principles calculations. *Geochim. Cosmochim. Acta* 198, 218–228.
952 <https://doi.org/10.1016/j.gca.2016.11.032>

953 Powley, H.R., Cappellen, P.V., Krom, M.D., 2017. Nutrient Cycling in the Mediterranean Sea: The Key
954 to Understanding How the Unique Marine Ecosystem Functions and Responds to
955 Anthropogenic Pressures, in: Fuerst-Bjelis, B. (Ed.), *Mediterranean Identities - Environment,*
956 *Society, Culture.* InTech. <https://doi.org/10.5772/intechopen.70878>

957 Ram, S., Erez, J., 2023. Anion elements incorporation into corals skeletons: Experimental approach
958 for biomineralization and paleo-proxies. *Proc. Natl. Acad. Sci.* 120, e2306627120.
959 <https://doi.org/10.1073/pnas.2306627120>

960 [Reitz, A., De Lange, G.J., 2006. Abundant Sr-rich aragonite in eastern Mediterranean sapropel S1:
961 Diagenetic vs. detrital/biogenic origin. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 235, 135–
962 148. <https://doi.org/10.1016/j.palaeo.2005.10.024>](#)

963 Schlitzer, Reiner, 2025. Ocean Data View [WWW Document]. Ocean Data View. URL
964 <https://odv.awi.de/> (accessed 12.10.25).

965 Schneider, A., Wallace, D.W.R., Körtzinger, A., 2007. Alkalinity of the Mediterranean Sea. *Geophys.*
966 *Res. Lett.* 34, 2006GL028842. <https://doi.org/10.1029/2006GL028842>

967 Siani, G., Paterne, M., Colin, C., 2010. Late glacial to Holocene planktic foraminifera bioevents and
968 climatic record in the South Adriatic Sea. *J. Quat. Sci.* 25, 808–821.
969 <https://doi.org/10.1002/jqs.1360>

970 Suess, E., 1979. Mineral phases formed in anoxic sediments by microbial decomposition of organic
971 matter. *Geochim. Cosmochim. Acta* 43, 339–352. [https://doi.org/10.1016/0016-7037\(79\)90199-6](https://doi.org/10.1016/0016-7037(79)90199-6)

972

973 Tian, R.C., Nicolas, E., 1995. Iodine speciation in the northwestern Mediterranean Sea, method and
974 vertical profile. *Mar. Chem.* 48, 151–156. [https://doi.org/10.1016/0304-4203\(94\)00048-I](https://doi.org/10.1016/0304-4203(94)00048-I)

975 Ullman, W.J., Luther, G.W., de Lange, G.J., Woititz, J.R.W, 1990. Iodine chemistry in deep anoxic
 976 basins and overlying waters of the Mediterranean Sea.
 977 Wadley, M.R., Stevens, D.P., Jickells, T.D., Hughes, C., Chance, R., Hepach, H., Tinel, L., Carpenter, L.J.,
 978 2020. A Global Model for Iodine Speciation in the Upper Ocean. *Glob. Biogeochem. Cycles*
 979 34, e2019GB006467. <https://doi.org/10.1029/2019GB006467>
 980 [Wang, X., Rickaby, R.E.M., Lu, Z., 2026. A deep dive into the planktic foraminiferal I/Ca in global core-](#)
 981 [tops. *Glob. Planet. Change* 105412. <https://doi.org/10.1016/j.gloplacha.2026.105412>](#)
 982 Winkelbauer, H., Cordova - Rodriguez, K., Reyes - Macaya, D., Scott, J., Glock, N., Lu, Z., Hamilton, E.,
 983 Chenery, S., Holdship, P., Dormon, C., Hoogakker, B., 2021. Foraminifera Iodine to Calcium
 984 Ratios: Approach and Cleaning. *Geochem. Geophys. Geosystems* 22.
 985 <https://doi.org/10.1029/2021GC009811>
 986 Winkelbauer, H.A., Hoogakker, B.A.A., Chance, R.J., Davis, C.V., Anthony, C.J., Bischoff, J., Carpenter,
 987 L.J., Chenery, S.R.N., Hamilton, E.M., Holdship, P., Peck, V.L., Poulton, A.J., Stinchcombe,
 988 M.C., Wishner, K.F., 2023. Planktic foraminifera iodine/calcium ratios from plankton tows.
 989 *Front. Mar. Sci.* 10, 1095570. <https://doi.org/10.3389/fmars.2023.1095570>
 990 Wong, G.T.F., Zhang, L., 1992. Changes in iodine speciation across coastal hydrographic fronts in
 991 southeastern United States continental shelf waters. *Cont. Shelf Res.* 12, 717–733.
 992 [https://doi.org/10.1016/0278-4343\(92\)90027-H](https://doi.org/10.1016/0278-4343(92)90027-H)
 993 Zhou, X., Hess, A.V., Bu, K., Sagawa, T., Rosenthal, Y., 2022. Simultaneous Determination of I/Ca and
 994 Other Elemental Ratios in Foraminifera: Comparing Results From Acidic and Basic Solutions.
 995 *Geochem. Geophys. Geosystems* 23, e2022GC010660.
 996 <https://doi.org/10.1029/2022GC010660>
 997 Zhou, X., Thomas, E., Rickaby, R.E.M., Winguth, A.M.E., Lu, Z., 2014. I/Ca evidence for upper ocean
 998 deoxygenation during the PETM. *Paleoceanography* 29, 964–975.
 999 <https://doi.org/10.1002/2014PA002702>
 1000