

Characteristics of tropical clouds with strong updrafts revealed by Doppler-velocity measurements from EarthCARE/CPR

Haruka Hotta^{1,2}, Kentaroh Suzuki¹, Maki Kikuchi², Shunsuke Aoki², Takuji Kubota²
¹Atmosphere and Ocean Research Institute, University of Tokyo, Kashiwa, 277-8564, Japan
²Earth Observation Research Center, Japan Aerospace Exploration Agency, Tsukuba, 305-8505, Japan
Correspondence to: Haruka Hotta (hotta@aori.u-tokyo.ac.jp)

Abstract. Updrafts within clouds are important for the climate system, yet global assessments have relied on indirect proxies. EarthCARE's 94 GHz Cloud Profiling Radar (CPR) provides the first global, spaceborne Doppler velocity (V_d) profiles of clouds, enabling observational constraints on convective vertical motions, while intense convective cores can be affected by velocity folding. To evaluate how EarthCARE/CPR Doppler-velocity measurements offer more direct insight into convective intensity relative to conventional convective-intensity proxies based on reflectivity fields, we analyze tropical CPR cloud-property products and extract convectively driven columns. We define $MaxV_d$ as the maximum upward V_d within the subfreezing portion of each non-folded column. Non-folded columns with $MaxV_d > 2.0$ m s⁻¹ are classified as measurable strong-updraft (SU) columns, whereas folded columns are treated separately as extreme-updraft (EU) candidates. Both SU and EU exhibit systematically higher echo-top heights at 0 and 10 dBZ than other columns. Their occurrence depends strongly on the separation between the cloud top and the 0 dBZ echo top, with a small separation robustly identifying both categories, even in relatively low-topped systems. Spatiotemporally, SU occurrence varies moderately across regions and local time, whereas EU occurrence is especially high over land at the 14:00 local-time overpass. These results show that $MaxV_d$ -based SU statistics and folding-based EU statistics capture different parts of the convective spectrum: measurable upward hydrometeor motion and likely more extreme convection affected by velocity folding, respectively. The combined constraints from Doppler-derived updraft intensity and radar-echo structure offer a process-oriented benchmark for evaluating convective dynamics-microphysics coupling in numerical models.

1 Introduction

Updrafts within tropical deep clouds promote the activation and condensational growth of cloud droplets and the formation of precipitation particles, thereby exerting strong control over the timing, amount, and intensity of rainfall. The strength of these updrafts also contributes to the maximum height reached by convective clouds and the horizontal extent of their anvils. Within the Earth's climate system, convective updrafts therefore constitute a key dynamical element. Through the rapid ascent of moist air and the associated release of latent heat, they drive large-scale overturning circulations such as the Hadley and Walker cells and contribute to meridional energy transport from the tropics to the midlatitudes (Manabe et al., 1965; Tiedtke, 1989; Bechtold et al., 2001). Vigorous updrafts shape the spatial distribution of rainfall and the occurrence of extreme precipitation events. In addition, the heights and lifetimes of deep clouds, which are affected by updrafts, exert a strong influence on the planetary energy budget by reflecting incoming shortwave radiation and trapping outgoing longwave radiation (Ramanathan et al., 1989; Hartmann et al., 2001). Updrafts are also crucial in the context of global warming projections: the representation of convective mixing and mass flux in global models affects simulated climate sensitivity and helps explain the spread in equilibrium climate sensitivity across models (Zhao, 2014; Sherwood et al., 2014). Despite this central role, direct observations of updraft vertical velocity (w) in tropical deep clouds remain sparse, limiting our ability to evaluate and improve its representation in atmospheric models. Surface-based and airborne radars have provided

削除:

書式変更: 幅 : 210 mm, 高さ : 297 mm

削除: traditionally

削除: The EarthCARE satellite's

削除: ,

削除: V_d)

削除: direct

削除: updraft

削除: beyond proxy-

削除: diagnostics. We

削除: over the tropics

削除: $MaxV_d$

削除: V_d

書式を変更: フォントの色 : 自動, 英語 (英国)

書式を変更: フォントの色 : 自動, 英語 (英国)

削除: Columns

削除: $MaxV_d$

削除: 5

削除: . They

削除: weaker-updraft columns, linking microphysics to dynamics. The probability of SU

削除: depends

削除: defined as ΔH . A

削除: ΔH

削除: identifies SU, including

削除: enhanced

削除: , with notably higher values

削除: than at 02:00. In contrast, oceanic regions have a smaller SU fraction and exhibit a weaker difference between the two overpass times.

削除: SU enhancements primarily reflect a shift toward horizontally compact, small- ΔH systems rather than higher cloud tops alone. Doppler folding preferentially occurs in small- ΔH structures, with maxima during the continental afternoon, providing a qualitative tracer of extreme updrafts.

削除: the

削除: between convective dynamics and microphysics

削除: determines

削除: controlled

74 invaluable snapshots of w (Heymsfield et al., 2010; Collis et al., 2013; Giangrande et al., 2016; Schiro and Neelin 2018;
 75 Schumacher et al., 2015), but these measurements are episodic, geographically restricted, and heavily concentrated over land.
 76 Consequently, global model evaluation still relies primarily on bulk fields and rainfall statistics rather than on the convective
 77 processes themselves (Maloney et al., 2019). General circulation models that employ cumulus parameterizations exhibit
 78 several persistent, systematic biases, including the double-ITCZ bias and excessively early diurnal peaks in precipitation
 79 amount and frequency. These deficiencies point to structural shortcomings in current parameterizations and motivate a
 80 fundamental reassessment (Christopoulos and Schneider, 2021; Tian and Dong, 2020). Even at cloud-resolving scales, state-
 81 of-the-art numerical models struggle to reproduce the observed statistics and vertical structure of w and its interaction with
 82 cloud microphysics. Recent intercomparisons of global km-scale models reveal substantial spread in the magnitude of updrafts:
 83 the fraction of columns in which the maximum vertical velocity exceeds $10\text{--}20\text{ m s}^{-1}$ can differ by more than an order of
 84 magnitude across models, and simulated regional patterns of convective strength vary widely (Abbott et al., 2025). Takahashi
 85 et al. (2025) similarly demonstrated that global km-scale simulations fail to adequately represent the regional variability of
 86 convective intensity and exhibit an unrealistic relationship between updrafts and precipitation formation processes. Together,
 87 these discrepancies underscore a critical need for global, observation-based constraints on w to calibrate both parameterized
 88 convection in general circulation models (GCMs) and explicitly simulated convection in global km-scale models.
 89 Spaceborne measurements have helped address the lack of comprehensive observations of w for model evaluation by providing
 90 systematic, globally distributed samples of deep cloud systems. However, because most satellite instruments cannot directly
 91 observe vertical motion, satellite-based studies have relied on proxy diagnostics for convective intensity. For instance, infrared
 92 imagers use $11\text{--}12\text{ }\mu\text{m}$ brightness temperatures to track overshooting tops and to document the diurnal cycle of cloud-top
 93 extent (Bedka et al., 2010; Yang and Slingo, 2001). Attempts have been made to infer cloud-top vertical velocities from the
 94 temporal evolution of cloud-top temperature; however, methodological uncertainties and a focus on developing clouds have
 95 so far prevented these approaches from yielding a comprehensive, global-scale picture across cloud regimes (Adler and Fenn,
 96 1979; Luo et al., 2014; Hamada and Takayabu, 2016). Passive microwave radiometers exploit scattering and absorption at
 97 approximately 89 and $166\text{--}183\text{ GHz}$ to infer convective intensity (Mohr and Zipser, 1996; Skofronick-Jackson et al., 2017).
 98 Other studies use lightning flash rates—which are closely linked to the collision rates of ice particles in strong updrafts—as
 99 indicators of the most vigorous convection (Christian et al., 2003; Williams and Stanfill, 2002; Deierling and Petersen, 2008;
 100 Cecil et al., 2005). The Tropical Rainfall Measuring Mission (TRMM; 1997–2015) and its successor, the Global Precipitation
 101 Measurement (GPM; 2014–present) mission, have provided unprecedented global views of precipitation using spaceborne
 102 precipitation radars (Kummerow et al., 1998; Iguchi et al., 2000; Hou et al., 2014; Skofronick-Jackson et al., 2017). From
 103 these observations, researchers have developed widely used proxies for convective intensity. One common proxy is the echo-
 104 top height (ETH), defined as the highest altitude at which a specified reflectivity threshold (e.g., 20, 30, or 40 dBZ) is detected.
 105 ETH is interpreted as the altitude reached by large, precipitating hydrometeors (Zipser et al., 2006; Liu and Zipser, 2005;
 106 Romatschke et al., 2010). Collectively, these proxy-based studies have revealed robust climatological features, including
 107 enhanced convective intensity over land compared with the ocean (Williams et al., 2005; Liu and Zipser, 2008; Zipser et al.,
 108 2006).
 109 The A-Train constellation further advanced vertical profiling through CloudSat’s 94 GHz Cloud Profiling Radar (CPR), which
 110 is sensitive to clouds and light-to-moderate precipitation and resolves the vertical structure of clouds with horizontal and
 111 vertical resolutions of approximately 1 km and 240 m, respectively (Stephens et al., 2002; Tanelli et al., 2008; Stephens et al.,
 112 2008). Compared with the TRMM (13.8 GHz) and GPM (13.6 and 35.55 GHz) precipitation radars, CPR has higher sensitivity,
 113 detecting clouds with reflectivity as low as about -28 dBZ (Stephens et al., 2008), and its smaller horizontal footprint provides
 114 more detailed information on convective cloud structure. Several proxy metrics for convective intensity based on CPR profiles
 115 have been proposed. One class of metrics uses cloud-top height (CTH) and echo-top height (ETH) at thresholds of 0 or 10
 116 dBZ. Stronger updrafts are expected to loft larger hydrometeors to higher levels and therefore result in greater ETH (Luo et

删除: et al.,

al., 2011; Takahashi and Luo, 2014). Another class uses the separation between CTH and ETH at 0 or 10 dBZ: when both small and large particles are carried to similarly high levels, CTH and ETH converge (Luo et al., 2008; Luo et al., 2011; Takahashi and Luo, 2014). Regional composites of these proxies reaffirm geographic variations in convective intensity and relate them to environmental controls. For example, Takahashi et al. (2023) demonstrated a strong land–ocean contrast, with hotspots over equatorial Africa and the Amazon, whereas convection over the Maritime Continent is comparatively weaker. Another diagnostic, the cloud center of gravity (COG), defined as the reflectivity-weighted mean height of the cloud within a column, has also been used to characterize convective structure (Koren et al., 2009). COG is especially high over the tropical West African Basin and the Congo Basin (Pilewskie and L’Ecuyer, 2022).

Although many previous studies have sought to characterize convective updraft intensity at the global scale, two major limitations remain. First, a fundamental shortcoming of these approaches is that they do not directly measure vertical motion. Proxy indicators do not have a simple or universal correspondence with w , and their relationships vary across different dynamical regimes (Takahashi and Luo, 2014; Liu et al., 2007). Consequently, even with extensive satellite archives, direct, observation-based constraints on w at the global scale have remained out of reach. Second, most proxy-based climatologies are designed to preferentially sample specific types of clouds. The precipitation radars aboard TRMM and GPM primarily detect fully developed, tall convective systems with strong radar echoes because, owing to their relatively low sensitivity, they must rely on high reflectivity thresholds (20–30 dBZ). These thresholds are intrinsically insensitive to weaker hydrometeor populations. Lightning-based metrics emphasize only the most strongly electrified convective cores, while infrared-based ascent-rate approaches focus exclusively on rapidly developing cloud tops. Even studies using CloudSat, which is capable of detecting weak echoes, have often concentrated on tall, well-developed clouds, partly because only a limited set of observable parameters—essentially radar reflectivity—can be robustly analyzed. Consequently, the resulting “intensity” maps that have guided our understanding of convection primarily reflect the upper tail of convective vigor and under-represent relatively low-topped or weak-echo clouds, as well as life-cycle stages outside peak development (Takahashi and Luo, 2014; Hamada et al., 2015; Xu et al., 2022).

A decisive advance has become possible with the Earth Cloud, Aerosol and Radiation Explorer (EarthCARE), a joint mission of ESA and JAXA launched in May 2024. EarthCARE carries a 94-GHz Doppler Cloud Profiling Radar (CPR), together with a high-spectral-resolution lidar, a multispectral imager, and broadband radiometers (Illingworth et al., 2015; Wehr et al., 2023). Crucially, the EarthCARE CPR provides nadir profiles of both radar reflectivity and Doppler velocity (V_d), enabling the first direct measurements of vertical motions within clouds at the global scale. The high sensitivity of EarthCARE’s 94-GHz radar allows the detection of relatively thin clouds, thereby systematically sampling shallow, weak-echo convective clouds as well as deep convective systems. EarthCARE is therefore well suited for evaluating vertical motions across a broad range of convective clouds (Illingworth et al., 2015). This capability opens a new route for constraining convective vertical motions, but it also requires careful interpretation. The CPR Nyquist velocity is only about $\pm 5\text{--}6\text{ m s}^{-1}$, so the most intense convective cores can be affected by velocity folding, which limits reliable quantitative interpretation of V_d .

Building on this novel capability with explicit consideration of its limitations, this study presents the first global-scale characterization of convective vertical-motion signatures in tropical deep convective clouds based on Doppler-velocity measurements from the EarthCARE CPR. The primary objectives are threefold: (1) to develop a method for identifying intense convective columns using quality-controlled V_d measurements, distinguishing non-folded measurable strong-updraft (SU) columns from folded columns treated as extreme-updraft (EU) candidates; (2) to investigate how the occurrence of SU and EU columns relates to the vertical structure of radar reflectivity, thereby probing links between cloud dynamics and microphysical processes and comparing these relationships with those inferred from legacy satellite proxies; and (3) to map the global distributions of the SU and EU-candidate columns and examine their dependence on variations in cloud morphological properties. Sect. 2 describes the EarthCARE CPR data and analysis methodology used in this study, Sect. 3 presents the results, and Sect. 4 summarizes the findings and discusses their implications.

删除: ,

删除: weak reflectivity

删除: updrafts

删除:

删除: updraft strength in tropical deep

删除: clouds

删除: Doppler velocities

删除: updraft strength

删除: distribution

删除: updraft intensity

删除: its

172 **2 Data and analysis method**

173 **2.1 Datasets**

174 We use measurements from the Cloud Profiling Radar (CPR) aboard the EarthCARE satellite. EarthCARE flies in a sun-
175 synchronous, near-polar orbit (inclination 97.05°) with nominal local equator-crossing times near 02:00 local time (ascending)
176 and 14:00 local time (descending) (Wehr et al., 2023). The 94-GHz CPR provides nadir profiles of radar reflectivity and
177 Doppler radial velocity. Its intrinsic vertical resolution is 500 m, while the profiles are sampled every 100 m in the vertical.
178 The instantaneous, beam-limited horizontal footprint is approximately 750 m at nadir, with profiles sampled at about 500 m
179 intervals along track.
180 The variables used in this analysis are taken from the standard CPR one-sensor cloud property product (CPR_CLP, version
181 Bb; JAXA, 2025a), specifically the cloud mask, radar reflectivity factor (Ref), Doppler velocity (V_d), and air temperature.
182 These data have a horizontal resolution of 1 km and a vertical sampling of 100 m. In CPR_CLP, the Level-1B received echo
183 power and V_d values are remapped onto the same grid as the cloud mask. Second-trip echoes (mirror images) are screened
184 following Battaglia (2021), as implemented by Aoki et al. (2026). The cloud mask is derived using noise screening, echo-
185 continuity tests, and surface-echo diagnostics (Sato et al., 2025). Air temperature is provided by a JAXA auxiliary product that
186 interpolates temperatures from ECMWF forecasts (Eisinger et al., 2024) to the EarthCARE Level-2 grid (JAXA, 2025b). In
187 this study, positive (negative) V_d denotes upward (downward) motion.

188 **2.2 Column selection criteria**

189 We exploit EarthCARE’s Doppler capability to analyze V_d in clouds for radar echoes exceeding the reflectivity threshold of
190 -19 dBZ, which includes clouds that would not be detected by precipitation-radar echo-top proxies. We focus on the tropical
191 region (30° S–30° N) for the period January–September 2025. Data collected prior to early December 2024 are discarded to
192 avoid known noise in V_d within weaker echoes associated with operation of a redundant side of the signal processing unit in
193 the CPR instrument (JAXA, 2025c), thereby ensuring data quality. Because Doppler-based diagnostics are restricted to range
194 gates with temperatures below 273.15 K (as discussed in Sect. 2.3), we retain only columns that contain a substantial cold
195 portion. To focus on convectively driven clouds, we extract columns that satisfy all of the following conditions:
196 (1) Single-layer cloud. Using the cloud mask, a gate is regarded as “cloudy” when the cloud-mask value is ≥ 20 (weak echo).
197 All gates between the highest and lowest cloudy gates must be classified as cloudy.
198 (2) Thermodynamic bounds. The cloud-top temperature (CTT) is < 258.15 K, and the cloud-base temperature is > 273.15 K.
199 (3) Geometrical thickness. The difference between cloud-top height and cloud-base height exceeds 5 km.
200 (4) Reflectivity strength. The column-maximum Ref exceeds 0 dBZ.

201 **2.3 Doppler-velocity quality control**

202 CPR Doppler-velocity measurements are affected by several sources of error: random noise arising from reduced correlation
203 between successive pulses caused by the rapid motion of the satellite platform (ϵ_{random} ; Doviak and Zrnic, 2014; Hagihara
204 et al., 2023); pointing uncertainty due to satellite attitude perturbations and antenna thermal distortion ($\epsilon_{\text{pointing}}$; Tanelli et
205 al., 2005); multiple scattering (ϵ_{MS} ; Battaglia et al., 2011); non-uniform beam-filling effects (ϵ_{NUBF}); and velocity aliasing
206 beyond the Nyquist limit ($\epsilon_{\text{Nyquist}}$; Sy et al., 2014).
207 To correct for $\epsilon_{\text{pointing}}$, a bias correction is applied based on the assumption that the V_d averaged horizontally over 100 km
208 at the surface is zero (Aoki et al., 2026; JAXA, 2026). We note that surface V_d values over heterogeneous land can be affected
209 by surface type and orography, and this approach may not fully remove the bias over land (Puigdomènech Treserras et al.,

删除: (V_d),

删除: Doppler velocities

删除: V_d

删除: vertical velocity w even

删除: with weak

删除: Doppler velocity

删除: velocities

删除: we apply

删除: Doppler velocity

删除:).

2025). To limit ϵ_{NUBF} and ϵ_{random} , which are large near cloud boundaries and in weak-reflectivity regions, we retain only gates that simultaneously satisfy a cloud-mask value ≥ 20 and $\text{Ref} \geq -19$ dBZ. This threshold is used as a quality-control criterion to exclude low-SNR Doppler estimates with large random uncertainty, following the reflectivity-dependent Doppler-error assessment of Hagihara et al. (2023), and should not be interpreted as a physical boundary between cloud regimes. To mitigate ϵ_{MS} , we follow Battaglia et al. (2011) and exclude range bins for which the cumulative reflectivity from the top of the atmosphere down to that gate exceeds a prescribed threshold. The CPR Nyquist (folding) velocity is approximately $\pm 5\text{--}6$ m s⁻¹; therefore, large downward (negative) velocities associated with the sedimentation of large particles can be aliased, especially at temperatures above 273.15 K. For this reason, all Doppler-based diagnostics are restricted to subfreezing gates (i.e., temperatures below 273.15 K).

To further reduce ϵ_{random} before identifying updraft signatures, we average V_d in pulse-pair covariance space. After applying the Doppler-velocity quality-control filters described above, we construct two averaged V_d fields for different purposes. First, to detect velocity folding, V_d is averaged over a 3 km along-track window while retaining the native 100 m vertical sampling. This 3 km $\times$ 100 m field is used because fold-like signatures appear as sharp vertical discontinuities and can be weakened if additional vertical smoothing is applied. Second, to define the column-maximum V_d , $\text{Max}V_d$ is averaged over the same 3 km along-track window and over two vertical gates, corresponding to 200 m in the vertical. This 3 km $\times$ 200 m field is used to reduce random errors and small-scale turbulent variability before extracting $\text{Max}V_d$.

For both averaged fields, the averaging is performed in pulse-pair covariance space rather than directly in velocity space. For each valid gate, the observed Doppler velocity V_d is converted to a phase using the local Nyquist velocity V_{nyq} . We then compute reflectivity-weighted real and imaginary components as,

$$\begin{aligned} \text{Re} &= Z \cos\left(\frac{\pi V_d}{V_{\text{nyq}}}\right), \\ \text{Im} &= Z \sin\left(\pi V_d / V_{\text{nyq}}\right), \end{aligned}$$

where Z is linear radar reflectivity. The real and imaginary components are summed over the averaging window, and the averaged Doppler velocity $\bar{V}_d$ is recovered as

$$\bar{V}_d = \frac{V_{\text{nyq}}}{\pi} \text{atan2}(\sum \text{Im}, \sum \text{Re}).$$

This procedure avoids artificial cancellation and wrap-around artifacts that can occur when velocities close to the Nyquist limit are averaged directly in velocity space. We require all gates in an averaging window to satisfy the cloud-mask, reflectivity, temperature, and multiple-scattering filters; if even one gate in the window fails any of these filters, the averaged value for that window is discarded and is not used in subsequent statistics.

We tested horizontal averaging windows of 1, 3, 5, and 10 km and vertical averaging windows of 100–500 m. For horizontal windows of 1, 3, 5, and 10 km, the folded-column fractions were 16.5, 7.2, 4.8, and 2.5%, respectively. The $\text{Max}V_d$ distribution for the 1 km case retained a noisy tail near the Nyquist limit, whereas the 10 km case excessively suppressed the $\text{Max}V_d$ tail (Fig. S1). Sensitivity tests for the vertical averaging scale showed that increasing the vertical window from 100 to 500 m progressively reduced the occurrence of large $\text{Max}V_d$ values, while the 200 m case retained the main positive tail more effectively than cases with stronger vertical smoothing (Fig. S2). Based on these sensitivity analyses, we use a 3 km horizontal and 200 m vertical averaging window as a compromise between reducing Doppler-velocity random error and retaining coherent convective structures on the scale of deep-convective updraft cores (Kollias et al., 2022; Galfione et al., 2025).

2.4 Updraft-signature column classification

Using the filtered and pulse-pair-averaged Doppler-velocity fields described in Sect. 2.3, we classify selected convective columns into three categories: non-strong-updraft (NonSU) columns, $\text{Max}V_d$ -based strong-updraft (SU) columns, and columns

削除: is

削除: , despite the first-pass dealiasing applied following Hagihara et al. (2023).

書式変更: ACP平文

移動 (挿入) [1]

削除: 2.4 Metric for updraft strength⁴¹

We identify columns affected by residual folding (aliasing) at temperatures below 273.15 K as follows. First, for gates colder than 273.15 K, we compute along-track means of V_d over 3 km windows to reduce random error and small-scale turbulent variability. If any pixel within a 3 km window fails the quality filters (cloud-mask value < 20 or $\text{Ref} < -19$ dBZ), that window is excluded from the analysis. We then examine vertical differences in the 3 km-averaged V_d profiles and flag as folded those columns in which the jump in V_d between adjacent vertical gates exceeds the Nyquist velocity. The folded columns are used to characterize the strength of convective updrafts, rather than being completely excluded from the analysis.⁴¹

For columns that are not flagged as folded, we define a V_d -based updraft metric to quantitatively characterize the convective intensity of each column. For all remaining gates colder than 273.15 K, V_d is averaged over 3-km along-track windows and then over two adjacent vertical gates (200 m) to reduce random errors and small-scale variability. If any pixel within an averaging window fails the validity filters (cloud-mask value < 20 or reflectivity < -19 dBZ), that window is discarded. This procedure effectively removes cloud-edge regions. We then define $\text{Max}V_d$, the column-maximum Doppler velocity, as the largest V_d within the portion of the 3-km- and 200-m-averaged profile that is colder than 273.15 K for columns that are not flagged as folded. As a result, columns associated with extremely strong updrafts, for which folding is more likely to occur (Galfione et al., 2025), may be excluded from the $\text{Max}V_d$ -based analysis.⁴¹

293 affected by Doppler-velocity folding, which are treated as extreme-updraft (EU) candidates. For brevity, these columns
294 affected by Doppler-velocity folding are referred to as EU columns hereafter. However, they should be interpreted as EU
295 candidates rather than quantitatively retrieved extreme updrafts.

296 **2.4.1 Detection of columns affected by Doppler-velocity folding as extreme-updraft candidates**

297 Velocity folding is detected from the 3 km-averaged V_d profiles with 100 m vertical sampling in the subfreezing portion of
298 each selected convective column. A vertical discontinuity between adjacent 100 m gates is counted as fold-like when three
299 conditions are simultaneously satisfied: (i) the velocity jump exceeds $1.5V_{nyq}$, (ii) the two adjacent velocities have opposite
300 signs, and (iii) at least one of the two velocities exceeds $0.6V_{nyq}$ in magnitude. Columns with at least one such discontinuity
301 are classified as columns affected by Doppler-velocity folding. Because the aliased velocities in these columns cannot be
302 interpreted quantitatively as upward hydrometeor Doppler motion, we exclude them from the $MaxV_d$ calculation and analyze
303 them separately as extreme-updraft (EU) candidates.

304 **2.4.2 $MaxV_d$ -based strong-updraft definition and occurrence ratios**

305 For selected convective columns that are not flagged as folded, we define $MaxV_d$ using the 3 km $\times$ 200 m averaged Doppler-
306 velocity field. Specifically, $MaxV_d$ is the maximum value of the quality-controlled, pulse-pair-averaged V_d over all valid
307 subfreezing gates in each non-folded column. Thus, the maximum is searched over the full valid subfreezing portion of each
308 column after 3 km horizontal and 200 m vertical averaging. Non-folded columns with $MaxV_d > 2.0 \text{ m s}^{-1}$ are classified as
309 $MaxV_d$ -based strong-updraft (SU) columns, and the remaining non-folded columns are classified as non-SU (NonSU) columns.
310 Because EU columns are excluded from the $MaxV_d$ calculation, the $MaxV_d$ -based SU category should be interpreted as a subset
311 of columns with measurable upward hydrometeor Doppler motion, not as the absolute strongest convective cores observed by
312 EarthCARE/CPR.

313 Using an extreme-value metric such as the column-maximum Doppler velocity ($MaxV_d$) preferentially captures embedded
314 updraft cores that dominate momentum and mass transport. In contrast, the mean or median V_d is more sensitive to
315 measurement noise. Similar “maximum” or high-percentile Doppler metrics have been used as proxies for convective intensity
316 and for evaluating model clouds (Heymsfield et al., 2010; Abbott et al., 2025). We also tested an alternative column metric,
317 namely the volume fraction of gates for which V_d exceeds a fixed threshold. The qualitative conclusions are unchanged when
318 this alternative metric is used instead of $MaxV_d$, indicating that our findings are robust to the specific choice of updraft indicator.
319 All occurrence ratios in this paper are defined using the same denominator, namely, the total number of selected convective
320 columns in the relevant bin or region, including both non-folded columns and EU columns. The SU ratio is therefore defined
321 as the number of non-folded columns with $MaxV_d > 2.0 \text{ m s}^{-1}$ divided by the number of all selected convective columns,
322 whereas the EU ratio is defined as the number of columns affected by Doppler-velocity folding divided by the same
323 denominator. This common basis allows direct comparison between the occurrence of measurable SU columns and that of
324 potentially more extreme, Nyquist-limited EU candidates.

325 **2.5 Case examination of V_d quality control and the updraft metric**

326

削除: $MaxV_d$

削除: V_d

削除: The 3-km along-track and two-gate (200 m) vertical averaging scales are tunable choices; however, the main results are insensitive to reasonable variations in these parameters. ...

削除: V_d

削除: $MaxV_d$,

削除: V_d





Figure 1: Example illustrating Doppler-velocity quality control, folding detection, and the definition of the updraft metric MaxV_d for an EarthCARE/CPR scene. (a) Radar reflectivity factor (Ref; dBZ). (b) Input CPR CLP Doppler velocity (V_d ; m s^{-1}) after applying only the cloud-mask threshold (cloud mask ≥ 20); positive values indicate upward motion. The black line marks the 273.15 K isotherm. (c) V_d after applying the column-selection criteria and quality control. Gaps indicate columns or gates for which no valid V_d remains after column selection and quality control. Grey shading near column indices approximately 70, 500, and 510 denotes gates flagged as being affected by multiple scattering. (d) Column-maximum V_d as a function of column index, computed from the input field in (b) for visual comparison ("Original"; green) and from the filtered and averaged field in (c) for the quantitative metric ("MaxV_d"; blue). Light-orange shading denotes columns flagged as folded and classified as extreme-updraft (EU) candidate columns. Light-blue shading denotes columns with $\text{MaxV}_d > 2.0 \text{ m s}^{-1}$, classified as strong-updraft (SU) columns. To illustrate how the column-selection criteria, range restrictions, folding detection, and spatial averaging described in Sects. 2.2–2.4 affect the Doppler-velocity diagnostics, we examine the example case shown in Fig. 1. In the radar-reflectivity field (Fig. 1a), the cloud structure is readily identifiable. Several active convective cores with spreading anvils (column indices around 80 and 400), together with an extensive stratiform region exhibiting a bright band (column indices 180–380), are evident. Figure 1b shows the input CPR CLP Doppler-velocity field at $1 \text{ km} \times 100 \text{ m}$ resolution, before the filtering and averaging used in this study, with only the cloud-mask threshold (≥ 20) applied. Above the 273.15 K isotherm, indicated by the black line, the field is speckled with incoherent patches, particularly near cloud edges where reflectivity is weak. These



- 削除: MaxV_d
- 削除: Native
- 削除: V_d
- 削除: Doppler velocity
- 削除: Doppler velocity
- 削除: raw
- 削除: blue
- 削除: ("Quality-controlled";
- 削除:). Grey
- 削除: being affected by residual velocity folding
- 削除: native $1 \text{ km} \times 200 \text{ m}$

365 features likely arise from random noise or small-scale turbulence rather than from coherent convective updrafts. Below the
366 273.15 K isotherm, although negative values are more prevalent, sporadic large positive values also appear, suggesting that
367 strongly negative velocities may have been aliased to positive values.

368 Figure 1c shows the gates that satisfy the convective-column selection and Doppler-velocity quality-control criteria. Columns
369 that fail the convective-column selection, and gates lacking valid Doppler estimates after quality control, are not shown. The
370 selection criteria in Sect. 2.2 remove multilayer cloud regions (column indices 100–120) and anvil regions near column indices
371 400 and 480. The quality-control and averaging procedure removes weak-signal outliers that are prone to error, damps small-
372 scale turbulent fluctuations, excludes the impact of folding at temperatures warmer than 273.15 K, and yields smoother and
373 more coherent structures. Multiple-scattering-contaminated gates are explicitly marked as grey shading and are excluded from
374 the V_d statistics.

375 Figure 1d compares the column-maximum V_d computed directly from the raw V_d field in Fig. 1b (“Original”; green) with the
376 $MaxV_d$, defined as the column-maximum V_d computed from the filtered and averaged field in Fig. 1c for non-folded columns.
377 The $MaxV_d$ varies smoothly across adjacent columns and shows distinct peaks near column indices 300 and 500, albeit with
378 reduced amplitudes. This example motivates our use of $MaxV_d$, computed from the quality-controlled V_d field, as the updraft
379 metric. The processing suppresses major sources of error and retains coherent 3 km-scale convective structures, but the 3 km
380 along-track and 200 m vertical averaging reduces the peak amplitude of V_d in narrow updraft cores. The resulting $MaxV_d$
381 should therefore be interpreted as a scale-dependent indicator of upward hydrometeor Doppler motion rather than a point-scale
382 convective-core velocity. The shaded columns summarize the two updraft-related categories. Light-blue shading denotes
383 $MaxV_d$ -based strong-updraft (SU) columns, defined as non-folded columns with $MaxV_d > 2.0 \text{ m s}^{-1}$, whereas light-orange
384 shading denotes columns affected by Doppler-velocity folding, which are treated separately as extreme-updraft (EU) columns.

365 3 Characteristics of Doppler-based updraft regimes

386 We identified approximately 1.20×10^6 columns after applying the selection criteria described in Sect. 2.2. Among them, about
387 1.10×10^6 columns are non-folded and are suitable for quantitative $MaxV_d$ analysis, whereas about 9.95×10^4 columns are
388 classified as Doppler-velocity folded. We report $MaxV_d$ -based SU and folding-based EU statistics together where relevant.
389 The only exception is Fig. 2 because it characterizes the $MaxV_d$ distribution, while folded columns cannot be assigned reliable
390 $MaxV_d$.

391 3.1 Doppler-classified convective columns and vertical reflectivity structure

392 To examine the relationship between updraft strength, cloud development, and precipitation formation, and to evaluate the
393 proxies used in CloudSat-based studies, we relate the updraft-intensity metric $MaxV_d$ to several diagnostics derived from the
394 vertical structure of radar reflectivity. We first examine the frequency distribution of $MaxV_d$ for all non-folded columns to
395 characterize its behavior and compare it with representative updraft velocities reported in previous studies (Fig. 2a). Owing to
396 the Nyquist velocity constraint, $MaxV_d$ is limited to approximately $+5 \text{ m s}^{-1}$. The distribution peaks between 0 and 1 m s^{-1} ,
397 which is smaller than typical values reported from aircraft-borne Doppler radar observations, where column-maximum
398 updrafts often exceed 10 m s^{-1} (Heymsfield et al., 2010), as well as from cloud-top ascent rates inferred from temporal changes
399 in infrared brightness temperature, which are on the order of $1\text{--}3 \text{ m s}^{-1}$ (Hamada and Takayabu, 2016; Li et al., 2021). A likely
400 reason for this apparent underestimation is the difference in the cloud life-cycle stages targeted. Whereas those previous studies
401 primarily sampled fully developed or rapidly developing convective clouds, our statistics also include clouds in their decaying
402 stages, thereby shifting the distribution toward weaker updrafts. It is also important to note that CPR V_d is a reflectivity-
403 weighted hydrometeor Doppler velocity, not the air vertical velocity. Large precipitating ice particles can dominate the

削除: field

削除: applying the column-selection and averaging procedures described in Sects. 2.2–2.3.

削除: 80–180

削除: We then apply the validity filters (cloud-mask value ≥ 20 , Ref $> -19 \text{ dBZ}$, and $T < 273.15 \text{ K}$) to V_d and average the remaining data over 3 km along track and two vertical gates (200 m), as described in Sect. 2.3. This

削除: Doppler velocity

削除: V_d

削除: blue

書式変更: 標準

削除: Doppler velocity

削除: 1c (“Quality-controlled”; orange). Columns whose 3 km-averaged V_d profiles exhibit discrete vertical jumps with magnitudes exceeding the Nyquist velocity are flagged as folded (as described in Sect. 2.3) and are shaded in ... [1]

削除: 80,

削除: ,

削除: and 520,

削除: $\leftarrow$

削除: $MaxV_d$,

削除: V_d

削除: primary

削除: without erasing

削除: signals, and

削除: modest

削除: (

削除:) $\times$

削除: (

削除:)

削除: increases precision while preserving the loc... [2]

削除: embedded

削除: $\leftarrow$

書式を変更: フォント: 10 pt

削除: updrafts

書式を変更: フォント: 10 pt

削除: ,171,643

削除: 2.2. We then flagged columns as folded usin... [4]

削除: -3.2 that use $MaxV_d$

削除: restricted to non-folded columns,

削除: in folded columns, large negative V_d values... [5]

削除: are assessed separately in Sect. 3.4.

削除: 3.1 Relationship between updraft strength

書式を変更: タブ位置: 13.5 字, 左揃え

削除: $MaxV_d$

削除: $\leftarrow$

削除: $MaxV_d$

書式を変更: フォント: 斜体

削除: $MaxV_d$

469 reflectivity-weighted signal, and their terminal fall speed can offset the upward air motion. Consequently, V_d can be
 470 substantially smaller than the air updraft velocity, or even indicate downward motion, within ascending cloudy air (Heymsfield
 471 et al., 2010). In addition, the exclusion of aliased columns and the spatial averaging applied to V_d both tend to shift $\downarrow\text{Max}V_d$
 472 toward smaller values. In terms of its shape, the $\downarrow\text{Max}V_d$ distribution exhibits clear positive skewness, with a long positive
 473 (upward) tail, consistent with previous findings (LeMone and Zipser, 1980; Lucas et al., 1994; Hamada and Takayabu, 2016).
 474 This indicates that columns with strong updrafts are relatively rare compared with those with weak or moderate updrafts.
 475 To relate updraft strength to cloud vertical development, we construct a two-dimensional histogram of CTH versus $\downarrow\text{Max}V_d$
 476 (Fig. 2b). CTH is defined as the highest altitude in a column where the cloud-mask value is ≥ 20 . The histogram clearly shows
 477 that, as CTH increases, the mode of the $\downarrow\text{Max}V_d$ distribution systematically shifts toward higher values. For example, clouds
 478 with CTHs of 8–10 km exhibit a modal $\downarrow\text{Max}V_d$ near 0 m s⁻¹, whereas those with CTHs of 14–16 km show a mode close to 1
 479 m s⁻¹. This pattern indicates that taller convective systems are generally more likely to contain strong updraft cores. This height
 480 dependence is qualitatively consistent with previous studies, which have reported that higher cloud tops tend to be associated
 481 with stronger convective intensity (Price and Rind, 1992; Song et al., 2020).
 482 The relationship between $\downarrow\text{Max}V_d$ and the column-maximum radar reflectivity (MaxRef), which is closely linked to
 483 hydrometeor size and precipitation intensity, is even more apparent (Fig. 2c). The occurrence frequency of large $\downarrow\text{Max}V_d$ values
 484 increases nonlinearly with increasing MaxRef, indicating that large particles are more readily generated and/or maintained in
 485 environments with relatively strong updrafts. This statistical covariation between reflectivity and updraft strength is consistent
 486 with previous observations from aircraft radars, although the absolute magnitudes depend on observational conditions, such
 487 as radar frequency (Heymsfield et al., 2010). However, attenuation, non-Rayleigh scattering, and multiple scattering at 94 GHz
 488 complicate the interpretation of very high reflectivity values (Matrosov, 2007; Kollias et al., 2022; Chase et al., 2025).

删除: V_d

删除: $\text{Max}V_d$

删除: $\text{Max}V_d$

删除: $\text{Max}V_d$

删除: $\text{Max}V_d$

删除: $\text{Max}V_d$

删除: and Sohn,

删除: $\text{Max}V_d$

删除: $\text{Max}V_d$

删除: Because of strong

删除: , however, MaxRef is likely underestimated in regions with ...




Figure 2: Statistical distribution of the Doppler-derived updraft metric (MaxV_d) and its relationship to cloud vertical development and precipitation intensity for selected convective cloud columns that are not flagged as affected by Doppler-velocity folding. (a) Histogram of MaxV_d (m s^{-1}). (b) Two-dimensional histogram of MaxV_d versus cloud-top height. (c) Two-dimensional histogram of MaxV_d versus column-maximum radar reflectivity (MaxRef ; dBZ). Shading in (b) and (c) indicates sample counts (logarithmic color scale); only bins with ≥ 100 samples are shown.

To further investigate the vertical structure of radar reflectivity, we classified the non-folded columns into two categories based on their MaxV_d . Non-folded measurable strong-updraft (SU) columns are defined as those with $\text{MaxV}_d > 2.0 \text{ m s}^{-1}$, totaling 6.36×10^4 columns (5.29% of all convective columns). The remaining non-folded columns are classified as non-SU (NonSU). This threshold isolates a relatively rare but dynamically distinct subset of strong-updraft cases.

We then use normalized Contoured Frequency-by-Altitude Diagrams (normalized CFADs; Luo et al., 2009; Yuter and Houze, 1995) to compare the statistical distributions of reflectivity profiles for the NonSU, SU, and EU categories (Fig. 3). The normalized CFADs reveal clear structural contrasts between SU and NonSU columns. SU columns exhibit a significantly larger fraction of high-reflectivity bins ($\geq 10 \text{ dBZ}$) at high altitudes ($> 8 \text{ km}$), indicating that strong updrafts loft large hydrometeors deep into the upper troposphere and promote ice production and subsequent growth by deposition, aggregation,

削除: ... [7]

- 削除: MaxV_d
- 削除: all analyzed non-folded tropical
- 削除: .
- 削除: MaxV_d
- 削除: MaxV_d
- 削除: MaxV_d
- 削除: MaxRef
- 書式を変更: フォント : 12 pt
- 削除: 3.1.2 Vertical structure of reflectivity depending on updraft strength
- 削除: MaxV_d . Strong
- 削除: MaxV_d
- 削除: 5
- 削除: 75,449
- 削除: 6.66
- 削除: non-folded
- 削除: two
- 削除: . The observed Doppler velocity (V_d)

and riming at higher altitudes. The observed V_d is the sum of the particle fall speed and the vertical motion of the ambient air. Because greater radar reflectivity generally corresponds to faster particle fall speeds (Seiki et al., 2025), which act to reduce V_d , the occurrence of high reflectivity at high altitudes in SU columns implies particularly strong updrafts. EU columns occupy an even more extreme structural regime: high reflectivities extend to high altitudes, indicating that these columns are closely related to the most vigorous convective structures even though their V_d values cannot be used quantitatively. Below the melting layer (<5 km), by contrast, reflectivity in EU columns is smaller than in NonSU and SU columns, likely due to stronger radar attenuation, a feature consistent with previous studies (Luo et al., 2014).





Figure 3: Normalized frequency-by-altitude diagrams (CFADs) of EarthCARE/CPR radar reflectivity (Ref; dBZ) for (a) non-strong-updraft (NonSU) columns, (b) strong-updraft (SU) columns, and (c) extreme-updraft (EU) candidate columns. Reflectivity is binned in 2 dBZ intervals, and the vertical coordinate is binned in 500 m intervals. Each panel is normalized by its total number of height-reflectivity samples such that the color indicates probability (i.e., the sum over all bins in each panel equals 1).

This structural difference in reflectivity among NonSU, SU, and EU columns is further quantified using box-and-whisker plots of cloud-top height and echo-top heights (Fig. 4). The median CTH increases from 12.8 km for NonSU columns to 13.5 km for SU columns and 14.0 km for EU columns, corresponding to increases of 0.7 and 1.2 km relative to NonSU columns, respectively. The contrast is more pronounced for echo-top heights of 0 dBZ and 10 dBZ. The median ETH(0) increases from 9.5 km for NonSU to 11.3 km for SU and 12.6 km for EU columns, while the median ETH(10) increases from 7.4 to 9.6 and 10.9 km, respectively. Thus, relative to NonSU columns, the median ETH(0) is higher by about 1.8 km for SU and 3.1 km for EU, and the median ETH(10) is higher by about 2.2 km for SU and 3.5 km for EU. These differences are substantially larger than those in CTH, indicating that echo-top heights of 0 dBZ and 10 dBZ are more sensitive than cloud-top height to the dynamical and microphysical structure associated with intense convection. Although EU columns cannot be assigned quantitative $MaxV_d$ values because of Doppler-velocity folding, their systematically higher ETH values indicate that folding preferentially occurs in columns where high-reflectivity hydrometeors are lofted to significantly high altitudes.

删除: , submitted

删除: V_d ,

删除: SU



删除:

删除: and

删除: where SU is defined by a column-maximum Doppler velocity of $MaxV_d > 2.5 \text{ m s}^{-1}$.

删除: sample count

删除: 3.1.3 Relationship between CTH, ETH, and Updraft Strength

删除: between the SU and

删除: various

删除: metrics

删除: of SU columns is approximately 900 m higher than that of ...

删除: .

删除: even

删除: ETH at higher reflectivity thresholds.

删除: at 0 dBZ, denoted as ETH(0), is approximately 2

删除: higher in SU

删除: at 10 dBZ, denoted as ETH

删除:), is approximately

删除: 0 km higher. The

删除: difference

删除: ETH compared with



删除: indicates

删除: ETH is a

删除: direct and

删除: proxy for updraft strength

删除: CTH.



删除:) and

删除: interplay
删除:)
删除: SU
删除: 5c

删除: ← ... [8]

Figure 4: Box-and-whisker summaries of cloud vertical extent for non-strong-updraft (NonSU), strong-updraft (SU), and extreme-updraft (EU) columns, shown for cloud-top height (CTH) and echo-top heights at reflectivity thresholds of 0 and 10 dBZ [ETH(0) and ETH(10)]. For each metric, the box spans the 10th–90th percentiles and the central line indicates the median (50th percentile). Numeric labels denote the 10th, 50th, and 90th percentile heights (km).

To further explore the relation between these height-based metrics, we examine the joint distributions of CTH and ETH(0) separately for NonSU (Fig. 5b), SU (Fig. 5e), and EU columns (Fig. 5h). Because ETH(0) is, by definition, equal to or lower than CTH, the populated region in the CTH–ETH(0) plane is confined to the triangular area above the 1:1 line. For both SU and NonSU columns, ETH(0) and CTH are strongly and positively correlated, consistent with previous satellite- and radar-based studies that have related differences between cloud-top and precipitation-top heights to convective intensity and life-cycle stage (e.g., Masunaga et al., 2005; Masunaga and Kummerow, 2006; Kikuchi and Suzuki, 2019). However, their detailed characteristics differ markedly. NonSU columns are, on average, shallower and exhibit a broad distribution with a large separation between ETH(0) and CTH. In contrast, SU columns are systematically taller, and their joint distribution lies much closer to the 1:1 line, indicating a smaller gap between ETH(0) and CTH.

EU columns show a further shift of population toward the upper-right part of the CTH–ETH(0) plane. Their distribution is concentrated at high CTH and high ETH(0), and it lies close to the 1:1 line over a wide height range. This indicates that the 0 dBZ echo top frequently approaches the cloud top in folded columns, suggesting that relatively large hydrometeors are lofted to near-cloud-top levels. Although the Doppler velocities in these columns cannot be used to quantify $\text{Max}V_d$ because of velocity folding, their radar-reflectivity structure is consistent with a more extreme convective regime. Together, the three distributions indicate that the separation between CTH and ETH(0), hereafter denoted as ΔH , is a useful structural measure for identifying both SU and EU columns.




Figure 5: Relationship between cloud-top height (CTH) and the 0 dBZ echo-top height [ETH(0)] for non-strong-updraft (NonSU), strong-updraft (SU) columns, and extreme-updraft (EU) columns. Two-dimensional histograms of CTH versus ETH(0) are shown for (b) NonSU, (e) SU, and (h) EU; shading indicates sample counts per bin, and only bins with ≥ 100 samples are colored. The corresponding marginal distributions are shown for ETH(0) in (a) NonSU, (d) SU, and (g) EU, and for CTH in (c) NonSU, (f) SU, and (i) EU.

Building on Fig. 5, Fig. 6 further quantifies how the occurrence ratios of SU and EU columns vary with CTH and ETH(0), thereby more clearly characterizing the distinct nature of the two Doppler-based categories. Using the common denominator defined in Sect. 2.4.2, Fig. 6 compares the occurrence ratios of SU and EU columns as functions of CTH and ETH(0). In the one-dimensional projections (Figs. 6a, 6c, 6d, 6f), both ratios increase with each of CTH and ETH(0), but the increase is more pronounced for EU. The SU ratio rises gradually as CTH increases and as ETH(0) reaches the upper troposphere, whereas the EU ratio increases more sharply for the highest echo tops. This contrast indicates that high echo tops are more discriminating indicators of both SU and EU occurrence than CTH alone.

The two-dimensional maps (Figs. 6b and 6e) further show that the separation between CTH and ETH(0) is a key structural control on updraft velocity. For both SU and EU, the occurrence ratios are highest near the 1:1 line, where ETH(0) approaches CTH. For a fixed CTH, bins with larger ETH(0) exhibit higher occurrence ratios, indicating a clear dependence on the separation between CTH and ETH(0). Conversely, when the separation between ETH(0) and CTH is large, i.e., far from the 1:1 line, the occurrence of both SU and EU is strongly suppressed. In the most extreme part of this phase space, where CTH exceeds 11 km and $\Delta H = \text{CTH} - \text{ETH}(0)$ is less than 0.5 km, the EU ratio exceeds 0.5, indicating that more than half of the selected convective columns are affected by Doppler-velocity folding. Notably, even at moderate CTH, columns with small CTH-ETH(0) separation still show enhanced SU occurrence, suggesting that this metric can identify relatively low-topped vigorous convection that may be missed by conventional high-altitude echo-top proxies.

These results highlight the importance of the separation between CTH and ETH(0) for characterizing vertical structure of convection. We therefore define $\Delta H \equiv \text{CTH} - \text{ETH}(0)$ as a compact diagnostic that links radar-echo structure to convective dynamics and microphysics. Small ΔH corresponds to the presence of relatively large hydrometeors near the cloud top in developing convective cores, whereas large ΔH characterizes decaying or weakly forced columns in which larger hydrometeors are confined to lower levels. This interpretation is consistent with conceptual models of convective evolution and with CloudSat-based intensity proxies that emphasize small CTH-ETH(0) separation (e.g., Takahashi and Luo, 2014); however, it is here evaluated using EarthCARE CPR Doppler-velocity measurements for non-folded columns and extended to EU-candidate columns affected by Doppler-velocity folding. In this framework, small ΔH identifies not only measurable SU columns but also Nyquist-limited convective columns in which reliable MaxV_d quantification is difficult. Thus, ΔH provides a common structural diagnostic for both measurable SU and EU-candidate regimes.

- 削除:) and
- 削除: .
- 削除: e) SU
- 削除: and
- 削除: f) SU
- 書式を変更: フォント: 9 pt, 太字
- 削除: probability of SU
- 削除: varies
- 削除: SU columns.
- 削除: (defined as the number of SU columns divided by the total number of columns, i.e., SU plus NonSU)
- 削除: monotonically with CTH (Fig. 6c), remaining below 0.05 for shallow clouds ($\text{CTH} \lesssim 8$ km) and approaching 0.10
- 削除: tallest clouds ($\text{CTH} \gtrsim 16$ km). A similar but even sharper increase is observed for ETH(0); the SU ratio rises monotonically with ETH(0) (Fig. 6a), from below 0.03 for $\text{ETH}(0) \lesssim 7$ km to more
- 削除: 0.30 for $\text{ETH}(0) \gtrsim 15$ km
- 削除: map of the SU ratio (Fig.
- 削除:) shows that the likelihood of SU occurrence increases jointly with CTH and ETH(0) and is maximized in the upper-right region of the ETH(0)-CTH phase space,
- 削除: is, for tall cloud systems in which
- 削除: SU fractions
- 削除: (
- 削除:), SU
- 削除: ($\text{CTH} < 10$ km),
- 削除: a substantial probability of being classified as SU. This suggests that a steep reflectivity gradient near the cloud top is a strong indicator of
- 削除: updrafts. This behavior directly addresses a historical blind spot in proxy climatologies of intense
- 削除: , which often impose reflectivity thresholds at high altitudes and therefore overlook low-topped convective columns (Zipser et al., 2006; Romatschke et al., 2010; Liu and Zipser, 2015).
- 削除:).
- 削除: particles
- 削除: validated
- 削除: direct
- 削除:
- 削除: low-topped
- 削除: populations





Figure 6: Occurrence ratios of SU columns and EU columns as functions of ETH(0) and CTH. (a–c) SU ratio and (d–f) EU ratio. The SU ratio is defined as the number of non-folded columns with $\text{Max}V_d > 2.0 \text{ m s}^{-1}$ divided by the total number of selected convective columns in each bin. The EU ratio is defined as the number of folded columns divided by the same denominator. Note that the SU and EU panels use different color scales. Panels (b) and (e) show two-dimensional ratios in the CTH–ETH(0) phase space; marginal ratios are shown as functions of ETH(0) and CTH.

3.2 Spatial and diurnal characteristics

3.2.1 Spatial and local-time patterns of SU and EU columns

To place the column-wise updraft statistics in a geographical and diurnal context, we analyze where and when $\text{Max}V_d$ -based SU columns and folding-based EU columns occur at two local times, 02:00 and 14:00 LT, and how their occurrence relates to regional cloud structure. We begin by examining the spatial distribution of all selected convective columns within the analysis domain (Fig. 7a). These columns are predominantly concentrated over well-known hotspots of deep convection (Zipser et al., 2006; Liu and Zipser, 2015), including the eastern Pacific Intertropical Convergence Zone (EPAC-ITCZ), the Amazon basin, central Africa, the Maritime Continent, and the South Pacific Convergence Zone (SPCZ), and are relatively sparse over the subtropical oceans. This geographical pattern confirms that our procedure effectively isolates columns embedded within convective cloud systems. We define five large-sample regions (Fig. 7a): the EPAC-ITCZ, central Africa, the Amazon basin, the Maritime Continent, and the SPCZ. These regions are used throughout the subsequent analysis to quantify regional differences in the occurrence and structure of SU and EU columns.

The $\text{Max}V_d$ -based SU ratio, defined as in Sect. 2.4.2 for each $2.5^\circ \times 2.5^\circ$ grid box, exhibits modest regional variability (Fig. 7b). In most grid boxes, the SU ratio remains below 0.10, and only a limited number of boxes exceed 0.15. Representative values for convectively active regions are summarized in Table 1. For all local times combined, the tropical-mean SU ratio is 5.29%. Regional values range from 4.55% over the Maritime Continent region and 4.76% over central Africa to 6.02% over the Amazon, 6.12% over the SPCZ, and 6.58% over the EPAC-ITCZ. Thus, the all-time SU ratio does not show a simple land–ocean contrast; instead, measurable SU occurrence is enhanced in several organized convective regimes, including both continental and oceanic convergence regions.



削除: ratio

削除: (SU + NonSU) within

削除: diurnal pattern

削除: strong-updraft

削除: (SU; $\text{Max}V_d > 2.5 \text{ m s}^{-1}$)

削除: fraction

When the statistics are separated by local overpass time—late night (ascending node, around 02:00 LT) versus early afternoon (descending node, around 14:00 LT)—the regional contrast becomes clearer (Figs. 7c–d, Table 1). Over oceanic convergence zones, the SU ratio is generally higher at night or shows only a weak afternoon enhancement. For example, the SU ratio decreases from 8.45% at 02:00 LT to 4.79% at 14:00 LT over the EPAC-ITCZ, and from 7.00% to 5.19% over the SPCZ. In contrast, continental regions show a moderate afternoon enhancement: the SU ratio increases from 4.27% to 5.60% over central Africa and from 5.54% to 6.76% over the Amazon. These values are higher than the 14:00 LT tropical mean of 5.04%, but the contrast is not as strong as in conventional proxy-based climatologies of the most extreme convection. This muted contrast is expected because the MaxV_d -based SU category includes only non-folded columns for which quantitative Doppler velocities can be retrieved. The most intense convective cores may instead enter the folded-column category.

The spatial distribution of EU ratio highlights this limitation and provides complementary information (Fig. 8). The EU ratio is defined as the number of columns affected by Doppler-velocity folding divided by the same total number of selected convective columns used for the SU ratio. For all local times combined, the tropical-mean EU ratio is 8.27% (Table 1). Regional values are largest over central Africa and the Amazon, where the all-time EU ratios reach 12.03% and 10.44%, respectively. In contrast, the Maritime Continent region has a lower EU ratio of 7.00%, while the EPAC-ITCZ and SPCZ have values of 8.11% and 8.74%, respectively.

The local-time dependence of the EU ratio is much stronger than that of the SU ratio. At 02:00 LT, the EU ratio is relatively low over central Africa and the Amazon, with values of 6.73% and 5.96%, respectively, whereas oceanic convergence zones such as the EPAC-ITCZ and SPCZ show higher nighttime values of 9.56% and 10.51%. At 14:00 LT, however, the EU ratio increases sharply over the continental regions, reaching 21.03% over central Africa and 17.34% over the Amazon. These values are far higher than the 14:00 LT tropical mean of 8.76% and much larger than the corresponding SU ratios of 5.60% and 6.76%. By contrast, the 14:00 LT EU ratios remain relatively modest over the Maritime Continent region, the EPAC-ITCZ, and the SPCZ, with values of 6.49%, 6.72%, and 6.88%, respectively.

Taken together, Figs. 7 and 8 and Table 1 indicate that MaxV_d -based SU and folding-based EU statistics capture complementary aspects of intense convection. The SU ratio quantifies the occurrence of columns with measurable upward hydrometeor Doppler motion and shows moderate regional and diurnal variability. The EU ratio, by contrast, identifies Nyquist-limited columns for which MaxV_d cannot be quantitatively retrieved, and it reveals a much stronger continental afternoon enhancement. The pronounced 14:00 LT increase in EU occurrence over central Africa and the Amazon suggests that a substantial fraction of the most intense continental convective cores is not represented in the non-folded MaxV_d -based SU population. Therefore, the weaker regional contrast in the SU ratio should not be interpreted as evidence that continental afternoon convection is only moderately intense; rather, it reflects the fact that the most extreme Doppler signatures are partly shifted into the Doppler-velocity-folded EU category.

This geographical and diurnal pattern is consistent with prior satellite-based characterizations using other convective-intensity metrics, such as echo-top height and lightning flash rate (Nesbitt and Zipser, 2003; Christian et al., 2003; Liu and Zipser, 2005; Zipser et al., 2006; Liu et al., 2008; Cecil et al., 2014; Pilewskie and L'Ecuyer, 2022). The regional contrast found here is also consistent with the satellite-inferred characteristics of vertical velocity by Jeyaratnam et al. (2021), which show enhanced convective vertical motions over continental regions. The EarthCARE CPR Doppler observations provide a complementary, solely observation-based constraint on this regional variability.

移動 (挿入) [2]

削除: , highlighting an enhancement over


削除: (Amazon: 5.3%,

削除: : 5.9%) relative to oceanic convective zones (EPAC-ITCZ: 4.6%,

削除: : 3.8%, SPCZ: 4.2%).

移動 (挿入) [3]

削除: ←






Figure 7: Spatial distribution of the analyzed columns and the occurrence of strong-updraft (SU) columns. (a) Number of selected convective columns per $2.5^\circ \times 2.5^\circ$ grid box over 30°S – 30°N . (b) Fraction of SU columns ($\text{Max}V_d > 2.0 \text{ m s}^{-1}$) relative to all convective columns in each grid box. (c–d) Same as (b), but separated by local overpass time: (c) late night ($\sim 02:00$ LT; ascending node) and (d) early afternoon ($\sim 14:00$ LT; descending node). Grid boxes with fewer than 100 total columns are masked in (b–d). Black rectangles labeled a–e indicate the regions used for the regional analyses (EPAC–ITCZ, Amazon, central Africa, Maritime Continent, and SPCZ).

削除: non-folded

削除: $\text{Max}V_d$

削除: 5

削除: non-folded




Figure 8: Spatial distribution of the occurrence of extreme-updraft (EU) columns. (a) Fraction of EU candidate columns (folded-column) relative to all convective columns in each grid box. (b, c) Same as (a), but separated by local overpass time: (b) late night (~02:00 LT; ascending node) and (c) early afternoon (~14:00 LT; descending node). Grid boxes with fewer than 100 total columns are masked in (a-c). Black rectangles labeled a-e indicate the regions used for the regional analyses (EPAC-ITCZ, Amazon, central Africa, Maritime Continent, and SPCZ).

3.2.2 Structural origin of regional and diurnal contrasts

What differences in cloud properties account for the moderate early-afternoon enhancement of SU columns and the much more pronounced enhancement of EU columns over central Africa and the Amazon (Sect. 3.2.1; Table 1)? Guided by the column-wise radar-reflectivity structural analysis in Sect. 3.1, we address this question by examining the median CTH, ETH(0), and their separation, $\Delta H \equiv \text{CTH} - \text{ETH}(0)$. In addition, we consider a simple morphology metric—the cloud length—defined as the along-track length of contiguous cloudy segments.

As summarized in Table 2, clouds over central Africa and the Amazon during the early afternoon do not stand out as being exceptionally tall. The tropical-mean median CTH at 14:00 LT is 13.0 km, compared with 12.4 km over central Africa and 13.1 km over the Amazon. Similarly, the median ETH(0) is 9.7 km in the tropical mean and 9.5 and 10.1 km over central Africa and the Amazon, respectively, during the early afternoon. These values are comparable to, or even slightly lower than, those in other convective regions, such as the Maritime Continent, the EPAC-ITCZ, and the SPCZ. Thus, the enhanced occurrence of intense convective columns in continental afternoon regimes cannot be explained simply by systematically higher cloud or echo tops.

Instead, the most distinctive feature of early-afternoon convection over central Africa and the Amazon is a markedly smaller ΔH . The tropical-mean median ΔH at 14:00 LT is 2.5 km, whereas it is only 1.8 km over central Africa and 1.9 km over the Amazon—smaller than in the other major convective regions listed in Table 2. By contrast, the Maritime Continent and the

移動 (挿入) [4]

書式を変更: フォント: 9 pt, 太字

上へ移動 [2]: When the statistics are separated by local overpass time—late night (ascending node, around 02:00 LT) versus early afternoon (descending node, around 14:00 LT)—the regional contrast becomes clearer (Figs. 7c–d, Table 1). Over the Maritime Continent, the EPAC-ITCZ, and

上へ移動 [3]: and lightning flash rate (Nesbitt and Zipser, 2003; Christian et al., 2003; Liu and Zipser, 2005; Zipser et al., 2006; Liu et al., 2008; Cecil et al., 2014; Pilewskie and L'Ecuyer, 2022).

削除: Over the Maritime Continent, the EPAC-ITCZ, and the SPCZ, SU fractions are slightly higher at night than in the early afternoon. In contrast, central Africa and the Amazon basin exhibit a pronounced daytime enhancement, with early-afternoon SU fractions of 9.3% and 8.1%, respectively, nearly twice the tropical-mean value of 4.3% computed across all regions and local times. This geographical and diurnal pattern is consistent with prior satellite-based characterizations using other convective-intensity metrics, such as echo-top height (ETH)

削除: ←

削除: particularly elevated

削除: fractions

削除: , while remaining modest in other convective regions

削除: across all columns and local times

削除: 12.7

削除: 1

削除: 12.8

削除: at 14:00 LT

削除: 5

削除: 1 km

削除: 9.8

削除: SU fractions

削除: across all local times

削除: 4

削除: 9

削除: during the afternoon

削除: any

削除: region–local-time combination

SPCZ exhibit substantially larger ΔH values of 3.0 km and 2.9 km, respectively, at 14:00 LT. A complementary perspective is obtained by comparing the median ΔH with the difference between the median CTH and median ETH(0). For afternoon central Africa and the Amazon, the differences between the median CTH and median ETH(0) are approximately 2.9 km and 3.0 km, respectively, whereas the median ΔH values are only 1.8 km and 1.9 km. These comparisons indicate that the ΔH distributions are skewed toward small values and that clouds with small CTH–ETH(0) separation occupy a disproportionately large fraction of the population, with this tendency being particularly pronounced in early-afternoon convection over central Africa and the Amazon.

Horizontal cloud length, summarized in Table 2, provides an additional clue to the nature of the underlying convective systems. The tropical-mean median cloud length at 14:00 LT is 482 km, but it is only 292 km and 294 km over central Africa and the Amazon, respectively, during the early afternoon. In contrast, the SPCZ and the Maritime Continent exhibit substantially larger median cloud lengths of 579 km and 517 km, respectively, at 14:00 LT. These statistics suggest that early-afternoon convection over central Africa and the Amazon is dominated by relatively compact systems, whereas oceanic regions—especially the SPCZ and the Maritime Continent—more frequently host large, organized systems.

To further characterize regional differences in the CTH–ETH(0) relationship, we examine deviations of the two-dimensional histograms of CTH versus ETH(0) from a tropics-wide reference distribution, constructed by combining all regions and local times, as shown in Fig. 9a–c. For early-afternoon convection over central Africa and the Amazon, the difference maps indicate a reduced occurrence of stratiform-like profiles characterized by large ΔH , together with an enhanced occurrence of profiles exhibiting small ΔH and moderate-to-high cloud tops. When considered alongside the relatively short cloud lengths, these features indicate a cloud population skewed toward developing or mature convective columns, rather than decaying systems with extensive stratiform regions.

We next examine in greater detail how these structural differences relate to the occurrence of MaxV_d -based SU columns. To this end, we analyze deviations in the SU ratio as a function of CTH and ETH(0) from the tropics-wide relationship shown in Fig. 6, as shown in Figs. 10a–c. Over early-afternoon convection in central Africa and the Amazon, the SU ratio deviations are generally weak and spatially patchy. Although localized positive anomalies appear in some bins near the 1:1 line, the continental regions do not show a robust enhancement of the conditional SU ratio across the CTH–ETH(0) phase space. In other words, columns with small CTH–ETH(0) separation are more common over central Africa and the Amazon, but they are not necessarily much more likely to be classified as MaxV_d -based SU columns than columns with the same CTH–ETH(0) structure in the tropics-wide reference distribution. The elevated regional SU occurrence therefore arises primarily from a shift in the underlying cloud population toward small- ΔH structures, rather than from a systematic increase in the probability that a given small- ΔH column contains measurable strong upward hydrometeor Doppler motion.

The folded-column ratio, interpreted here as the EU ratio, provides an additional perspective on the same structural regime (Figs. 10d–f). In contrast to the weak SU ratio deviations, the EU ratio shows clear positive anomalies over central Africa and the Amazon at 14:00 LT, particularly near the 1:1 line and in the high-CTH and high-ETH(0) regime. This indicates that Doppler-velocity folding occurs preferentially in columns where the 0 dBZ echo top approaches the cloud top, consistent with vigorous convective cores that loft relatively large hydrometeors to near-cloud-top levels. Because reliable MaxV_d quantification is difficult in these folded columns, they are excluded from the MaxV_d -based SU statistics. The positive EU anomalies therefore suggest that SU statistics alone do not capture the full population of intense continental afternoon convection; part of this population is instead identified through Doppler-velocity folding.

The Maritime Continent provides a useful contrast. Despite having relatively high median CTH and ETH(0), comparable to or exceeding those over central Africa and the Amazon (Table 2), SU occurrence over the Maritime Continent remains close to or slightly below the tropical mean and is therefore substantially lower than that over these continental regions (Table 1). Structurally, clouds over the Maritime Continent tend to exhibit larger ΔH and longer horizontal extents, indicating a greater prevalence of deep convective systems with broad stratiform regions. This pattern is also evident in Fig. 9c, which shows an

- 删除: (
- 删除:).
- 删除: tropics as a whole
- 删除: median ΔH is 2.4 km, whereas the difference
- 删除: is 3.2 km. The contrast is even more pronounced for afternoon central Africa and the Amazon, for which the corresponding ...
- 删除: 9
- 删除: 3.0
- 删除: 283
- 删除: (567
- 删除: 504
- 删除:).
- 删除: (
- 删除:),
- 删除: 8a
- 删除: (CTH $\approx$ 6–15 km).
- 删除: enhanced
- 删除: fraction
- 删除: tropical-mean distribution (cf.
- 删除:),
- 删除: 9a
- 删除: fraction is enhanced across much of the (CTH, ETH(0)) phase space, with particularly strong
- 删除: along and
- 删除: where ΔH is small.
- 删除: both
- 删除: host strong updrafts
- 删除: in the tropical mean.
- 删除: fractions
- 删除: arise
- 删除: combination of (i) a
- 删除: and (ii) an increased
- 删除: updrafts, rather than from an overall increase in extremely tall clouds
- 删除: fractions
- 删除: remain
- 删除: are
- 删除: those
- 删除: 8c

enhanced frequency of clouds with high CTH but relatively low ETH(0). Furthermore, the SU ratio at a given CTH-ETH(0) pair (Fig. 10c) tends to be weak or negative over parts of the small-ΔH developing-cloud regime. The EU anomalies are also weaker over the Maritime Continent than over central Africa and the Amazon (Fig. 10f). Thus, over the Maritime Continent, the more frequent occurrence of stratiform-like profiles, together with weaker SU and EU enhancements at fixed CTH and ETH(0), leads to an overall smaller occurrence of intense convective columns.

These regional patterns are consistent with previous satellite-based analyses of deep convective intensity across the three major tropical “chimney zones,” which show that convection over tropical Africa exhibits the strongest updrafts, Amazonia is intermediate, and the tropical Maritime Continent regime displays the weakest convective intensity when measured using radar echo-top heights and related proxies (e.g., Takahashi and Luo, 2014; Takahashi et al., 2017, 2023; Liu et al., 2007; Zipser et al., 2006). These studies argue that stronger land convection arises from a more efficient conversion of convective available potential energy into vertical kinetic energy, associated with deeper and drier boundary layers and higher lifting condensation levels, which produce broader and more buoyant convective cores that are less susceptible to dilution by entrainment. In contrast, convection over the Maritime Continent regime tends to exhibit narrower cores and larger entrainment rates, resulting in weaker updrafts (Lucas et al., 1994; Williams and Stanfill, 2002; Takahashi et al., 2017, 2023). Our results support these findings by showing a broadly consistent regional ordering using EarthCARE CPR Doppler-velocity measurements while adding two constraints that were not available from proxy-based studies alone. First, the continental-afternoon enhancement of MaxV_d-based SU occurrence is mainly linked to a higher frequency of compact small-ΔH cloud structures. Second, the stronger EU signal indicates that part of the most intense continental-afternoon convection lies in the Nyquist-limited regime, where MaxV_d cannot be quantified reliably. The combined SU and EU statistics therefore show that regional contrasts in convective intensity are controlled not simply by cloud-top height, but by the joint structure of CTH, ETH(0), ΔH, cloud morphology, and Doppler-velocity folding.

Table 1: Regional occurrence of MaxV_d-based strong-updraft (SU) columns and columns affected by Doppler-velocity folding. For each domain and local overpass time, the table lists the total number of selected convective columns, the number of non-folded columns, the number of non-folded SU columns, the number of folded columns, the SU ratio, and the folded-column ratio. SU columns are defined as non-folded columns with MaxV_d > 20 m s⁻¹, and folded columns are interpreted as extreme-updraft (EU) candidates. Both ratios use the same denominator: the total number of selected convective columns in the corresponding region and local-time category. “All” denotes the tropical aggregate over 30°S–30°N. Region definitions are shown in Fig. 7a (see Sect. 3.2.1).

time	region	Total count	Non-folded count	SU count	Folded count	SU ratio	Folded ratio
	All	1.20 × 10 ⁶	1.10 × 10 ⁶	6.36 × 10 ⁴	9.95 × 10 ⁴	5.29%	8.27%
	Africa	6.54 × 10 ⁴	5.76 × 10 ⁴	3.12 × 10 ³	7.87 × 10 ³	4.76%	12.03%
	Amazon	9.83 × 10 ⁴	8.81 × 10 ⁴	5.92 × 10 ³	1.03 × 10 ⁴	6.02%	10.44%
	Maritime Continent	2.96 × 10 ⁵	2.75 × 10 ⁵	1.35 × 10 ⁴	2.07 × 10 ⁴	4.55%	7.00%
	ITCZ	5.59 × 10 ⁴	5.13 × 10 ⁴	3.68 × 10 ³	4.53 × 10 ³	6.58%	8.11%
	SPCZ	8.03 × 10 ⁴	7.33 × 10 ⁴	4.91 × 10 ³	7.02 × 10 ³	6.12%	8.74%
	All	6.33 × 10 ⁵	5.83 × 10 ⁵	3.49 × 10 ⁴	4.95 × 10 ⁴	5.51%	7.82%

- 削除: fraction
- 削除: warm-pool/
- 削除: .
- 削除: 5
- 書式を変更: フォント: 9 pt, 太字
- 削除: Central Africa
- 書式を変更
- 書式付きの表
- 削除: Local Time
- 削除: ALL
- 削除: Amazon
- 削除: Maritime Continent
- 削除: EPAC-ITCZ
- 削除: SPCZ
- 書式変更: 左揃え
- 削除: 20E+06
- 削除: 6.27E+04
- 削除: 1.04E+05
- 削除: 2.93E+05
- 削除: 38E+04
- 削除: 30E+04
- 削除: 6.29E+04
- 削除: 1.52E+05
- 削除: 29E+05
- 削除: 3.78E+04
- 書式変更: 左揃え
- 削除: 02:00 LT
- 削除: 2.59E+04
- 削除: 4.20E+04
- 削除: Total Count
- 書式を変更: フォントの色: 自動
- 結合されたセル
- 書式付きの表
- 移動 (挿入) [5]
- 削除: 5.74E+05
- 書式変更: 左揃え
- 削除: 49E+04
- 削除: 4.12E+04
- 削除: 1.41E+05
- 書式を変更: フォントの色: 自動
- 削除: 14:00 LT
- 削除: 2.79E+04
- 削除: 4.09E+04
- 書式変更: 両端揃え
- 削除: 5.54E+03
- 削除: 1.12E+04
- 削除: 12E+04
- 削除: 3.68E+03
- 書式変更: 左揃え
- 削除: All
- 削除: 2.47E+03
- 削除: 3.51E+03
- 削除: SU count
- 書式変更: 両端揃え
- 書式を変更: フォントの色: 自動
- 削除: 2.19E+03
- 削除: 6.05E+03
- 削除: 2.55E+04
- 削除: 1.36E+03
- 書式変更: 左揃え
- 削除: 02:00 LT
- 削除: 1.47E+03
- 削除: 2.11E+03
- 分割されたセル

02:00 LT							
	Africa	4.12×10^4	3.84×10^4	1.76×10^3	2.77×10^3	4.27%	6.73%
	Amazon	5.97×10^4	5.61×10^4	3.30×10^3	3.56×10^3	5.54%	5.96%
	Maritime Continent	1.54×10^5	1.42×10^5	7.60×10^3	1.15×10^4	4.94%	7.48%
	ITCZ	2.73×10^4	2.47×10^4	2.31×10^3	2.61×10^3	8.45%	9.56%
	SPCZ	4.10×10^4	3.67×10^4	2.87×10^3	4.31×10^3	7.00%	10.51%
14:00 LT							
	All	5.70×10^5	5.20×10^5	2.88×10^4	5.00×10^4	5.04%	8.76%
	Africa	2.42×10^4	1.91×10^4	1.36×10^3	5.10×10^3	5.60%	21.03%
	Amazon	3.87×10^4	3.20×10^4	2.61×10^3	6.71×10^3	6.76%	17.34%
	Maritime Continent	1.42×10^5	1.33×10^5	5.87×10^3	9.22×10^3	4.13%	6.49%
	ITCZ	2.86×10^4	2.66×10^4	1.37×10^3	1.92×10^3	4.79%	6.72%
	SPCZ	3.93×10^4	3.65×10^4	2.04×10^3	2.70×10^3	5.19%	6.88%

Table 2: Regional median structural metrics. Regional medians of cloud-top height (CTH; km), 0 dBZ and 10 dBZ echo-top height [ETH(0) and ETH(10); km], their separation from CTH (km), and along-track cloud length (km) for the tropical aggregate and five convective regions (central Africa, Amazon, Maritime Continent, EPAC-ITCZ, and SPCZ), stratified by local overpass time (All, ≈02:00 LT, and ≈14:00 LT). Region definitions are shown in Fig. 7a (see Sect. 3.2.1).

time	region	cloud top height	echo top height (0 dBZ)	echo top height (10 dBZ)	cloud top height - echo top height (0 dBZ)	cloud top height - echo top height (10 dBZ)	cloud length
ALL	All	12.9	9.7	7.7	2.4	4.6	494
	Africa	12.4	9.3	7.7	2.2	4.3	458
	Amazon	12.8	9.6	7.9	2.2	4.3	417
	Maritime Continent	13.5	10.0	8.0	2.7	5.2	516
	ITCZ	13.1	10.1	8.3	2.3	4.4	482
	SPCZ	13.0	9.8	7.7	2.6	5.0	581
02:00 LT	All	12.8	9.7	7.8	2.3	4.5	510
	Africa	12.4	9.2	7.6	2.5	4.5	532
	Amazon	12.6	9.3	7.6	2.4	4.4	536
	Maritime Continent	13.4	10.1	8.1	2.5	5.0	514

	ITCZ	13.2	10.5	8.7	2.0	4.2	449
	SPCZ	13.0	10.0	8.1	2.4	4.7	593
	All	13.0	9.7	7.7	2.5	4.7	482
	Africa	12.4	9.5	7.8	1.8	3.9	292
	Amazon	13.1	10.1	8.4	1.9	4.0	294
	Maritime Continent	13.6	10.0	7.8	3.0	5.4	517
	ITCZ	13.0	10.0	7.9	2.5	4.6	500
14:00 LT	SPCZ	13.0	9.6	7.5	2.9	5.2	579




Figure 9: Deviations in the joint probability density function (PDF) of cloud-top height (CTH) and the 0 dBZ echo-top height [ETH(0)] relative to the tropics-wide reference distribution (all regions and local times combined; non-folded columns only). Panels show early afternoon (~14:00 LT) convection over (a) central Africa, (b) the Amazon, and (c) the Maritime Continent. Bin size is 0.5 km in both dimensions; only bins with $N \geq 100$ columns are shown.




Figure 10: Deviations in Doppler-based occurrence ratios conditioned on cloud-top height (CTH) and the 0 dBZ echo-top height [ETH(0)] for early afternoon (~14:00 LT). (a–c) show deviations in the MaxV_d-based SU ratio, and panels (d–f) show deviations in

the Doppler-velocity folded-column ratio, for (a, d) central Africa, (b, e) the Amazon, and (c, f) the Maritime Continent. Bin size is 0.5 km in both dimensions; only bins with $N \geq 100$ columns are shown.

4 Summary and Discussion

This study exploits the first global Doppler-velocity observations from space using the EarthCARE/CPR to diagnose updraft strength in tropical convective clouds and to relate it to the reflectivity-based vertical structure. While most previous satellite studies have inferred convective intensity using indirect proxies, we directly analyze vertical particle motions, with explicit consideration of Doppler-velocity folding. Importantly, we examine not only mature deep convective systems but also lower-topped convective clouds, which are often underrepresented in legacy intensity metrics.

In this study, we developed a method to extract measurable strong-updraft (SU) columns and columns affected by Doppler-velocity folding, treated as extreme-updraft (EU) candidates, from quality-controlled EarthCARE/CPR Doppler-velocity profiles. Here, the quality-control procedure consists of: (1) masking weak echoes, (2) restricting quantitative diagnostics to range gates colder than 273.15 K, (3) applying modest spatial averaging, and (4) flagging velocity folding. We define $\text{Max}V_d$ as the maximum upward (positive) Doppler velocity within the subfreezing portion of each non-folded profile and classify non-folded measurable strong-updraft columns (SU) as those with $\text{Max}V_d > 2.0 \text{ m s}^{-1}$, whereas columns affected by Doppler-velocity folding are analyzed separately as extreme-updraft (EU) candidates. Thus, SU and EU are treated as complementary diagnostics: the former quantifies measurable upward hydrometeor Doppler motion, while the latter identifies Nyquist-limited columns likely associated with more extreme updraft regimes. We analyzed EarthCARE/CPR observations from January to September 2025 over the tropical region (30°S – 30°N).

Using the common denominator of all the selected convective columns, SU columns account for 5.29% of the tropical sample, whereas EU columns account for 8.27%. These occurrence ratios indicate that both categories are relatively rare yet dynamically important, and that a substantial fraction of intense convective columns is represented by the folded EU population rather than by the non-folded $\text{Max}V_d$ -based SU population alone. Examining SU and EU occurrence as a function of cloud-top height (CTH), the 0 dBZ echo-top height [ETH(0)], and the 10 dBZ echo-top height [ETH(10)], we find that both SU and EU columns exhibit systematically higher CTH and, more notably, higher ETH(0) and ETH(10) than NonSU columns. Mapping their occurrence onto the two-dimensional space spanned by CTH and ETH(0) further shows that both SU and EU occurrence are concentrated where $\Delta H \equiv \text{CTH} - \text{ETH}(0)$ is small. This small ΔH suggests that large hydrometeors are lofted near the cloud top.

Figure 11 summarizes the relationship between updraft strength and the vertical structure of radar reflectivity. When the 0 and 10 dBZ echo-top heights rise close to the cloud top, the separations $\text{CTH}-\text{ETH}(0)$ and $\text{CTH}-\text{ETH}(10)$ become small, indicating that high-reflectivity hydrometeors are lofted to near-cloud-top levels. Columns with small separations can meet the $\text{Max}V_d$ -based SU criterion even when the cloud-top height itself is modest (Fig. 11a), whereas EU columns occur preferentially when the same small-separation structure is accompanied by much taller cloud and echo tops (Fig. 11b). In contrast, when reflectivity increases gradually downward from the cloud top and both the 0 and 10 dBZ levels remain well below the CTH, updrafts are typically weak (Fig. 11c). These results indicate that the reflectivity structure beneath the cloud top can serve as a robust proxy for both measurable strong-updraft columns and Nyquist-limited extreme-updraft candidates: smaller $\text{CTH}-\text{ETH}(0)$ separations correspond to stronger updraft regimes, whereas larger separations correspond to weaker updrafts.

This finding provides direct observational support for CloudSat-era intensity proxies based on small $\text{CTH}-\text{ETH}(0)$ separations (e.g., Takahashi and Luo, 2014; Takahashi et al., 2017, 2023; Liu et al., 2007). The novelty of this study lies in leveraging direct Doppler-velocity measurements to move beyond the proxy-based estimates of convective intensity used in previous satellite studies, and in extending the same structural interpretation to EU columns. Our results support the central premise of

上へ移動 [4]: Black rectangles labeled a–e indicate the regions used for the regional analyses (EPAC-ITCZ, Amazon, central Africa, Maritime Continent, and SPCZ).

削除: 3.3 Implication from Doppler-velocity folding

In Sects. 3.1–3.2, our analysis was restricted to non-folded columns to ensure robust quantitative statistics of Doppler-derived updraft strength. In this subsection, we shift focus to columns in which Doppler-velocity folding (aliasing) is detected. Previous Doppler-radar studies have shown that the most intense convective cores often generate radial velocities exceeding the instrument Nyquist limit, resulting in folded spectra and aliased velocity measurements (Battaglia et al., 2011; Kollias et al., 2014; Sy et al., 2014). Although these folded velocities cannot be used directly as quantitative estimates of updraft magnitude, the presence of ve...

削除: After applying the column-selection and D...

削除: Table 3: Regional folding occurrence. R...

書式を変更: 英語 (英国)

書式を変更: フォント: 9 pt, 太字

書式を変更: フォント: 9 pt, 太字

書式を変更: フォント: 10 pt, フォントの色: 黒

削除: , spaceborne

削除:

削除: of W-band (94 GHz) radar reflectivity. Whereas

削除: .

削除: We analyzed EarthCARE/CPR observation...

削除: that masks

削除: restricts

削除: applies

削除: flags residual

削除: $\text{Max}V_d$

削除: $\text{Max}V_d$

削除: 5

削除: $\sim$.

削除: These SU columns account for approximat...

削除: . Examining SU

削除:).

削除: the SU fraction

削除: is

削除: to

削除: Although large ETH and small ΔH have t...

削除: 12

削除: and columns tend

削除: host strong updrafts

削除: 11a,b

削除: strength

削除: updrafts

削除:

削除:

削除: .

proxy approaches but also refine it by ranking the tested indicators for the measurable SU population: the cloud-top–echo-top separation ΔH ($CTH - ETH(0)$) is the most informative metric for identifying strong updrafts, followed by $ETH(10)$, $ETH(0)$, and finally CTH . The same small- ΔH structure is also preferentially associated with EU, consistent with the physical picture described by Takahashi et al. (2023), in which strong updrafts in developing convective cores are accompanied by a rapid upward extension of the radar echo top.

Regionally, the combined SU and EU statistics reveal complementary aspects of tropical convective intensity. The all-time SU ratio shows moderate regional variability. By contrast, the all-time EU ratio shows a clearer continental enhancement, reaching 12.03% over central Africa and 10.44% over the Amazon, compared with 7.00% over the Maritime Continent, 8.11% over the EPAC-ITCZ, and 8.74% over the SPCZ. When comparing the 02:00 and 14:00 LT overpasses, oceanic convergence zones exhibit either a nighttime enhancement or a weak early-afternoon signal in both diagnostics, whereas continental regions show an afternoon enhancement that is modest for SU but pronounced for EU. At 14:00 LT, the SU ratios over central Africa and the Amazon are 5.60% and 6.76%, respectively, only moderately higher than the tropical mean of 5.04%; the corresponding EU ratios, in contrast, reach 21.03% and 17.34%, far above the tropical mean of 8.76%.

Structurally, during the afternoon over Africa and the Amazon, CTH and $ETH(0)$ are not exceptionally large in absolute terms; instead, the separation $\Delta H = CTH - ETH(0)$ remains consistently small. The median ΔH at 14:00 LT is 1.8 km over central Africa and 1.9 km over the Amazon, compared with the tropical mean of 2.5 km, and the median cloud lengths are only 292 and 294 km, respectively, compared with the tropical mean of 482 km. In contrast, the Maritime Continent and the SPCZ exhibit larger ΔH and longer horizontal extents, indicating a greater prevalence of deep convective systems with broad stratiform regions. For SU, the regional enhancement therefore primarily reflects a greater contribution from horizontally compact, developing-to-mature convective cores characterized by small ΔH , rather than from unusually tall clouds. For EU, the same structural regime produces a stronger signal: Doppler-velocity folding occurs preferentially where $ETH(0)$ approaches CTH and where high-reflectivity hydrometeors are lofted to near-cloud-top levels.

Taken together, these results reinforce the interpretation in Fig. 11 that ΔH is a key structural discriminator of both SU and EU occurrences. Regional and diurnal variations in SU and EU are governed not simply by how high clouds grow but by how closely the radar echo top approaches the cloud top. Continental afternoon environments, characterized by a high frequency of horizontally compact convective columns with small ΔH , are particularly conducive to strong Doppler-derived updraft signatures and to Nyquist-limited EU occurrence, whereas regimes dominated by horizontally extensive, stratiform-rich systems tend to exhibit lower SU and EU occurrence even when cloud tops are high. This establishes a physically consistent link between regional contrasts in Doppler-derived updraft signatures and the vertical reflectivity structure of tropical convective systems.

These measurements help close a long-standing observational gap in global constraints on convective updrafts and enable the development of a new dynamical climatology. This climatology can be used to calibrate convective parameterizations in global climate models (GCMs) and to benchmark explicitly resolved convection in global kilometer-scale simulations. A key question for numerical models is whether they reproduce (i) regional and diurnal contrasts in the frequency of measurable strong-updraft columns and EU-candidate columns and (ii) the joint statistics of updraft signatures and radar-echo structure documented here.

Takahashi et al. (2025) compared deep convective clouds simulated by global storm-resolving models with CloudSat observations over three tropical “chimney” regions: tropical Africa, tropical Amazonia, and the tropical warm pool. Within that framework, cloud-top height and radar-derived echo-top metrics could be evaluated against CloudSat observations; however, updraft intensity could not be validated because CloudSat lacks Doppler-velocity measurements. Consequently, comparisons of updraft strength were necessarily limited to inter-model differences and proxy-based diagnostics. These studies identified substantial biases in convective intensity and precipitation formation: updraft indicators in the tropical warm pool are often too strong, while precipitation forms at unrealistically high altitudes under weak vertical velocities, indicating inconsistencies between cloud dynamics and microphysics. Our EarthCARE-based diagnostics overcome this limitation by

删除: This interpretation is

删除: columns occur more frequently over

删除: areas such as

删除: Basin and less frequently over oceanic convergence zones (e.g., the eastern Pacific

删除: ITCZ and the SPCZ) and mixed land–sea regions such as

删除: .

删除: regions

删除: slight

删除: of SU

删除: little diurnal contrast,

删除: tend to

删除: significantly higher

删除: occurrence at

删除: .

删除: Moreover, across most (CTH , $ETH(0)$) combinations, the local SU fraction in these continental hotspots substantially exceeds the tropical-mean value, indicating that the environment favors stronger updrafts even for clouds with comparable vertical development. We interpret this enhancement as reflecting a greater contribution from

删除: , thereby explaining the elevated SU occurrence

删除: 12

删除: strong updrafts.

删除: occurrence

删除: updrafts

删除: fractions

删除: updrafts

删除: Even after quality control, velocity folding is detected in approximately 3.3% of columns. The folding frequency increases with increasing CTH and $ETH(0)$ and is especially high where ΔH is small. Folding occurs most frequently over continental early-afternoon hotspots of intense convection, such as central Africa and the Amazon Basin. In both their vertical radar-reflectivity structure and their spatiotemporal characteristics, columns affected by folding closely resemble those with high SU fractions. Although folding cannot be used as a quantitative metric of updraft strength, its occurrence serves as a useful qualitative tracer of extreme updrafts. Because the analyses in Sects. 3.1 and 3.2 are restricted to unfolded columns, it is likely that the most intense convective cases are not fully represented in these analyses.

删除: updrafts

删除: intensity

删除:

1656 directly linking MaxV_d-based SU occurrence and folding-based EU occurrence to the vertical structure of radar reflectivity.
 1657 These constraints enable rigorous tests of whether models not only reproduce observed cloud-top and precipitation-top heights
 1658 but also associate these structures with realistic updraft velocities and observationally comparable Doppler-velocity folding
 1659 occurrence, thereby providing a process-oriented pathway to improve the coupling between convective dynamics and
 1660 microphysics in both parameterized and explicitly resolved convection.
 1661 Despite these advances, this study has several limitations that highlight priorities for future work. First, our analysis is based
 1662 on a nine-month record, which is too short to characterize the full range of intraseasonal and interannual variability, including
 1663 the Madden–Julian Oscillation (MJO) and the El Niño–Southern Oscillation (ENSO). Extending the analysis to multi-year
 1664 data sets will be essential for quantifying how long-period variability modulates convective updrafts. Second, to ensure data
 1665 quality, MaxV_d-based SU statistics exclude columns affected by Doppler-velocity folding. Because folding preferentially
 1666 occurs in the strongest updraft regimes, SU statistics alone likely underestimate both the frequency and magnitude of the most
 1667 extreme convective events. Treating folded columns as EU candidates partly addresses this under-sampling by retaining them
 1668 as a qualitative diagnostic, but it does not provide quantitative updraft magnitudes. Developing a physically based unfolding
 1669 algorithm for the CPR is therefore a high priority to mitigate this bias, and to convert EU-candidate occurrence into quantitative
 1670 velocity information. Third, we restricted the diagnosis to subfreezing regions (temperature < 273.15 K) to avoid Doppler-
 1671 velocity aliasing from large-particle sedimentation. This restriction excludes cloud systems with cloud-top temperatures above
 1672 273.15 K from the analysis and limits the assessment of updrafts to the upper levels of the targeted clouds. Fourth, because the
 1673 CPR operates at W band, it is susceptible to strong attenuation in heavy precipitation, non-Rayleigh scattering, and multiple
 1674 scattering, which can obscure reflectivity at lower altitudes, (Chase et al., 2025), as suggested by the lower-level reflectivity
 1675 patterns in SU and EU columns. As a result, we cannot yet directly evaluate echo-top-height climatologies at 20–40 dBZ
 1676 derived from the TRMM and GPM precipitation radars.
 1677 Future work should integrate this unique Doppler-velocity dataset with complementary observing systems. Combining
 1678 EarthCARE Doppler-velocity measurements with GPM precipitation estimates (Aoki et al., 2026), lightning observations, and
 1679 cloud life-cycle metrics from geostationary satellites would enable a more holistic characterization of deep convection. Such
 1680 an integrated framework would allow for the systematic evaluation and recalibration of legacy intensity proxies and support
 1681 the development of a comprehensive intensity index that integrates ETH, ΔH, electrification, life-cycle phase, and Doppler-
 1682 velocity folding.

删除: MaxV_d

删除: and by quantifying regional and diurnal variations in the occurrence of strong updrafts (SU).

删除: in the quantitative analyses in Sects. 3.1 and 3.2, we excluded...

删除:

删除: updrafts, our

删除: This

删除: may also reduce the apparent land–ocean contrast relative to climatologies based on precipitation-radar echo-top heights from GPM (Houze et al., 2015; Liu and Zipser, 2015) and lightning-based proxies (Albrecht et al., 2016).

删除: .

删除:

删除: ,

删除: (Fig. 5).

删除: and




Figure 11: Conceptual schematic of typical radar-reflectivity structures associated with strong and weak updraft columns. Grey, dotted, and filled-dot outlines indicate the regions exceeding -30 , 0 , and 10 dBZ, respectively. Orange arrows illustrate updraft strength qualitatively. **Panel (a)** shows strong-updraft cases in which the 0 and 10 dBZ echo tops rise close to the cloud top (small CTH-ETH(0)) and moderately deep cloud. **Panel (b)** shows extreme-updraft cases in which the 0 and 10 dBZ echo tops rise close to the cloud top (small CTH-ETH(0)) and exceedingly deep cloud. **Panel (c)** shows a weak-updraft case characterized by a top-light structure in which reflectivity increases gradually downward and both ETH(0) and ETH(10) remain well below the cloud top.

Data availability

The EarthCARE/CPR L2A CPR one-sensor Cloud Products (<https://doi.org/10.57746/EO.01jdvd2gqq34e6yz9p8kfe68x5>, JAXA, 2025a) used in this study can be downloaded from the JAXA G-Portal (<https://gportal.jaxa.jp/gpr/>).

Author contributions

HH conceptualized the study, performed the formal analysis, developed the methodology, and prepared the visualizations. KS contributed to the conceptualization and acquired the funding. MK and SA conducted the investigation and provided feedback on the results. TK conducted the investigation and supervised the study. HH wrote the original draft of the manuscript. KS, MK, SA, and TK contributed to the review and editing of the manuscript.

Competing interests

The contact author has declared that none of the authors has any competing interests.

削除: ←

... [17]

削除: 12

削除: Panels

削除: -b) show

削除:)), including a moderately deep cloud (a).

1726 **Special issue statement**

1727 This article is part of the special issue “Early results from EarthCARE (AMT/ACP/GMD inter-journal SI)”. It is not associated
1728 with a conference.

1729 **Acknowledgements**

1730 ChatGPT was used to assist the editing the language of parts of the manuscript. All content was subsequently reviewed and
1731 edited by the authors, who take full responsibility for the final version of the manuscript. We would also like to thank Editage
1732 (www.editage.jp) for English language editing. [We are grateful to the anonymous reviewers for their careful reading of the](#)
1733 [manuscript and for their constructive comments, which greatly helped us improve both the analysis and the presentation of the](#)
1734 [results.](#)

1735 **Financial support**

1736 This study was supported by JAXA/EarthCARE project, MEXT program for the Advanced Studies of Climate Change
1737 Projection (SENTAN) (Grant JPMXD0722680395), and JST Moonshot R&D (Grant JPMJMS2281).

1738 **References**

- 1739 Abbott, T. H., Jeevanjee, N., Cheng, K.-Y., Zhou, L., and Harris, L.: The land-ocean contrast in deep convective intensity in
1740 a global storm-resolving model, *J. Adv. Model. Earth Syst.*, 17, 5, <https://doi.org/10.1029/2024MS004467>, 2025.
- 1741 Adler, R. F. and Fenn, D. D.: Thunderstorm vertical velocities estimated from satellite data, *Mon. Weather Rev.*, 107, 1747–
1742 1754, [https://doi.org/10.1175/1520-0469\(1979\)036<1747:TVVEFS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1979)036<1747:TVVEFS>2.0.CO;2), 1979.
- 1743 Albrecht, R. I., Goodman, S. J., Buechler, D. E., Blakeslee, R. J., and Christian, H. J.: Where Are the Lightning Hotspots on
1744 Earth?, *B. Am. Meteorol. Soc.*, 97, 2051–2068, <https://doi.org/10.1175/BAMS-D-14-00193.1>, 2016.
- 1745 Aoki, S., Kubota, T., and Turk, F. J.: Exploring vertical motions in convective and stratiform precipitation using spaceborne
1746 radar observations: Insights from EarthCARE and GPM coincidence dataset, *Atmos. Meas. Tech.*, 19, 79–100,
1747 <https://doi.org/10.5194/amt-19-79-2026>, 2026.
- 1748 Battaglia, A., Augustynek, T., Tanelli, S., and Kollias, P.: Multiple scattering identification in spaceborne W-band radar
1749 measurements of deep convective cores, *J. Geophys. Res.-Atmos.*, 116, D19, <https://doi.org/10.1029/2011JD016142>, 2011.
- 1750 Battaglia, A.: Impact of second-trip echoes for space-borne high-pulse-repetition-frequency nadir-looking W-band cloud
1751 radars, *Atmos. Meas. Tech.*, 14, 7809–7820, <https://doi.org/10.5194/amt-14-7809-2021>, 2021.
- 1752 Bechtold, P., Bazile, E., Guichard, F., Mascart, P., and Richard, E.: A mass-flux convection scheme for regional and global
1753 models, *Q. J. R. Meteorol. Soc.*, 127, 869–886, <https://doi.org/10.1002/qj.49712757309>, 2001.
- 1754 Bedka, K., Brunner, J., Dworak, R., Feltz, W., Otkin, J., and Greenwald, T.: Objective satellite-based detection of overshooting
1755 tops using infrared window channel brightness temperature gradients, *J. Appl. Meteor. Climatol.*, 49, 181–202,
1756 <https://doi.org/10.1175/2009JAMC2286.1>, 2010.
- 1757 Cecil, D. J., Goodman, S. J., Boccippio, D. J., Zipser, E. J., and Nesbitt, S. W.: Three years of TRMM precipitation features.
1758 Part I: Radar, radiometric, and lightning characteristics, *Mon. Weather Rev.*, 133, 543–566, [https://doi.org/10.1175/MWR-](https://doi.org/10.1175/MWR-2876.1)
1759 [2876.1](https://doi.org/10.1175/MWR-2876.1), 2005.
- 1760 [Cecil, D. J., Buechler, D. E., and Blakeslee, R. J.: Gridded lightning climatology from TRMM-LIS and OTD: Dataset](#)
1761 [description, Atmos. Res., 135–136, 404–414, https://doi.org/10.1016/j.atmosres.2012.06.028, 2014.](#)

Chase, R. J., Dolan, B., Rasmussen, K. L., Schulte, R. M., Stephens, G. L., Turk, F. J., and van den Heever, S. C.: A multifrequency spaceborne radar perspective of deep convection, *J. Appl. Meteor. Climatol.*, 64, 133–145, <https://doi.org/10.1175/JAMC-D-24-0109.1>, 2025.

Christian, H. J., Blakeslee, R. J., Boccippio, D. J., Boeck, W. L., Buechler, D. E., Driscoll, K. T., Goodman, S. J., Hall, J. M., Koshak, J. M., Mach, D. M., and Stewart, M. F.: Global frequency and distribution of lightning as observed from space by the Optical Transient Detector, *J. Geophys. Res. Atmos.*, 108, ACL 4-1-ACL 4-15, <https://doi.org/10.1029/2002JD002347>, 2003.

Christopoulos, C. and Schneider, T.: Assessing biases and climate implications of the diurnal precipitation cycle in climate models, *Geophys. Res. Lett.*, 48, e2021GL093017, <https://doi.org/10.1029/2021GL093017>, 2021.

Collis, S., Protat, A., May, P. T., and Williams, C.: Statistics of storm updraft velocities from TWP-ICE including verification with profiling measurements, *J. Appl. Meteor. Climatol.*, 52, 1909–1922, <https://doi.org/10.1175/JAMC-D-12-0230.1>, 2013.

Deierling, W. and Petersen, W. A.: Total lightning activity as an indicator of updraft characteristics, *J. Geophys. Res. Atmos.*, 113, D16, <https://doi.org/10.1029/2007JD009598>, 2008.

Doviak, R. J. and Znić, D. S.: *Doppler Radar & Weather Observations: Second Edition*, Academic Press, ISBN 148329482X, 9781483294827, 2014.

Eisinger, M., Marnas, F., Wallace, K., Kubota, T., Tomiyama, N., Ohno, Y., Tanaka, T., Tomita, E., Wehr, T., and Bernaerts, D.: The EarthCARE mission: science data processing chain overview, *Atmos. Meas. Tech.*, 17, 839–862, <https://doi.org/10.5194/amt-17-839-2024>, 2024.

Galfione, A., Battaglia, A., Puigdomènech Treserras, B., and Kollias, P.: First insights into deep convection by the Doppler velocity measurements of the EarthCARE Cloud Profiling Radar, *Atmos. Meas. Tech.*, 18, 6747–6763, <https://doi.org/10.5194/amt-18-6747-2025>, 2025.

Giangrande, S. E., Toto, T., Jensen, M. P., Bartholomew, M. J., Feng, Z., Protat, A., Williams, C. R., Schumacher, C., and Machado, L.: Convective cloud vertical velocity and mass-flux characteristics from radar wind profiler observations during GoAmazon2014/5, *J. Geophys. Res.-Atmos.*, 121, 12891–12913, <https://doi.org/10.1002/2016JD025303>, 2016.

Houze, R. A., Jr., Rasmussen, K. L., Zuluaga, M. D., and Brodzik, S. R.: The variable nature of convection in the tropics and subtropics: A legacy of 16 years of the Tropical Rainfall Measuring Mission (TRMM) satellite, *Rev. Geophys.*, 53, 994–1029, <https://doi.org/10.1002/2015RG000488>, 2015.

Hagihara, Y., Ohno, Y., Horie, H., Roh, W., Satoh, M., and Kubota, T.: Global evaluation of Doppler velocity errors of EarthCARE cloud-profiling radar using a global storm-resolving simulation., *Atmos. Meas. Tech.*, 16, 3211–3219, <https://doi.org/10.5194/amt-16-3211-2023>, 2023.

Hamada, A., Takayabu, Y. N., C. Liu, and E. J. Zipser: Weak linkage between the heaviest rainfall and tallest storms, *Nat. Commun.*, 6, 6213, <https://doi.org/10.1038/ncomms7213>, 2015.

Hamada, A. and Takayabu, Y. N.: Convective cloud top vertical velocity estimated from geostationary satellite rapid-scan measurements, *Geophys. Res. Lett.*, 43, 5435–5441, <https://doi.org/10.1002/2016GL068962>, 2016.

Hartmann, D. L., Moy, L. A., and Fu, Q.: Tropical convection and the energy balance at the top of the atmosphere, *J. Climate*, 14, 4495–4511, [https://doi.org/10.1175/1520-0442\(2001\)014<4495:TCATEB>2.0.CO;2](https://doi.org/10.1175/1520-0442(2001)014<4495:TCATEB>2.0.CO;2), 2001.

Heymsfield, G. M., Tian, L., Heymsfield, A. J., Li, L., and Guimond, S.: Characteristics of deep tropical and subtropical convection from nadir-viewing high-altitude airborne Doppler radar, *J. Atmos. Sci.*, 67, 285–308, <https://doi.org/10.1175/2009JAS3132.1>, 2010.

Hou, A. Y., Kakar, R. K., Neeck, S., Azarbarzin, A. A., Kummerow, C. D., Kojima, M., Oki, R., Nakamura, K., and Iguchi, T.: The Global Precipitation Measurement mission, *Bull. Amer. Meteor. Soc.*, 95, 701–722, <https://doi.org/10.1175/BAMS-D-13-00164.1>, 2014.

Iguchi, T., Kozu, T., Meneghini, R., Awaka, J., and Okamoto, K.: Rain-profiling algorithm for the TRMM precipitation radar, *J. Appl. Meteor.*, 39, 2038–2052, [https://doi.org/10.1175/1520-0450\(2001\)040<2038:RPAFTT>2.0.CO;2](https://doi.org/10.1175/1520-0450(2001)040<2038:RPAFTT>2.0.CO;2), 2000.

Illingworth, A. J., Barker, H. W., Beljaars, A., Ceccaldi, M., Chepfer, H., Clerbaux, N., Cole, J., Delanoë, J., Domenech, C.,
 Donovan, D. P., Fukuda, S., Hirakata, M., Hogan, R. J., Huenerbein, A., Kollias, P., Kubota, T., Nakajima, T., Nakajima, T.,
 Y., Nishizawa, T., Ohno, Y., Okamoto, H., Oki, R., Sato, K., Satoh, M., Shephard, M., Velázquez-Blázquez, A., Wandinger,
 U., Wehr, T., and van Zadelhoff, G.-J.: The EarthCARE Satellite: The next step forward in global measurements of clouds,
 aerosols, precipitation and radiation, B. Am. Meteorol. Soc., 96, 1311–1332, <https://doi.org/10.1175/BAMS-D-12-00227.1>,
 2015.
 JAXA, EarthCARE/CPR L2A CPR one-sensor Cloud Products.
<https://doi.org/10.57746/EO.01jdv2gq34e6yz9p8kfe68x5>, last access: 5 January 2026, 2025a.
 JAXA, EarthCARE JAXA Level 2 Algorithm Theoretical Basis Document (L2 ATBD),
[https://www.eorc.jaxa.jp/EARTHCARE/document/JAXAL2ProductList/ATBD/NDX-](https://www.eorc.jaxa.jp/EARTHCARE/document/JAXAL2ProductList/ATBD/NDX-110018N_EarthCARE_L2_ATBD_draft_rev2.pdf)
[110018N_EarthCARE_L2_ATBD_draft_rev2.pdf](https://www.eorc.jaxa.jp/EARTHCARE/document/JAXAL2ProductList/ATBD/NDX-110018N_EarthCARE_L2_ATBD_draft_rev2.pdf), last access: 5 January 2026, 2025b.
 JAXA, Release notes for the EarthCARE/CPR Level-1b Product, available from
[https://www.eorc.jaxa.jp/EARTHCARE/document/L1ProductList/ReleaseNote/CPR/SEC-2024057-](https://www.eorc.jaxa.jp/EARTHCARE/document/L1ProductList/ReleaseNote/CPR/SEC-2024057-0D_Release_Notes_for_EarthCARE_CPR_L1b_Product.pdf)
[0D_Release_Notes_for_EarthCARE_CPR_L1b_Product.pdf](https://www.eorc.jaxa.jp/EARTHCARE/document/L1ProductList/ReleaseNote/CPR/SEC-2024057-0D_Release_Notes_for_EarthCARE_CPR_L1b_Product.pdf), last access: 5 January 2026, 2025c.
 JAXA: EarthCARE JAXA Level 2 Algorithm Theoretical Basis Document (L2 ATBD), ver. P, Japan Aerospace Exploration
 Agency, available at:
[https://www.eorc.jaxa.jp/EARTHCARE/document/JAXAL2ProductList/ATBD/EarthCARE_JAXA_L2_ATBD_verP_Apr2](https://www.eorc.jaxa.jp/EARTHCARE/document/JAXAL2ProductList/ATBD/EarthCARE_JAXA_L2_ATBD_verP_Apr2026.pdf)
[026.pdf](https://www.eorc.jaxa.jp/EARTHCARE/document/JAXAL2ProductList/ATBD/EarthCARE_JAXA_L2_ATBD_verP_Apr2026.pdf) (last access: 19 May 2026), 2026.
 Jeyaratnam, J., Luo, Z. J., Giangrande, S. E., Wang, D., and Masunaga, H.: A Satellite-Based Estimate of Convective Vertical
 Velocity and Convective Mass Flux: Global Survey and Comparison With Radar Wind Profiler Observations, Geophys. Res.
 Lett., 48, e2020GL090675, <https://doi.org/10.1029/2020GL090675>, 2021.
 Kikuchi, M. and Suzuki, K.: Characterizing vertical particle structure of precipitating cloud system from multiplatform
 measurements of A-Train constellation, Geophys. Res. Lett., 46, 1040–1048, <https://doi.org/10.1029/2018GL081244>, 2019.
 Kollias, P., Tanelli, S., Battaglia, A., and Tatarevic, A.: Evaluation of EarthCARE cloud profiling radar Doppler velocity
 measurements in particle sedimentation regimes, J. Atmos. Oceanic Technol., 31, 366–386, [https://doi.org/10.1175/JTECH-](https://doi.org/10.1175/JTECH-D-11-00202.1)
[D-11-00202.1](https://doi.org/10.1175/JTECH-D-11-00202.1), 2014.
 Kollias, P., Battaglia, A., Lamer, K., Treserras, B. P., and Braun, S. A.: Mind the Gap - Part 3: Doppler velocity measurements
 from space, Front. Remote Sens., 3, 860284, <https://doi.org/10.3389/frsen.2022.860284>, 2022.
 Koren, I., Altaratz, O., Feingold, G., Levin, Z., and Reisn, T.: Cloud's Center of Gravity – a compact approach to analyze
 convective cloud development, Atmos. Chem. Phys., 9, 155–161, <https://doi.org/10.5194/acp-9-155-2009>, 2009.
 Kummerow, C., Barnes, W., Kozu, T., Shiue, J., and Simpson, J.: The Tropical Rainfall Measuring Mission (TRMM) sensor
 package, J. Atmos. Oceanic Technol., 15, 809–817, [https://doi.org/10.1175/1520-0426\(1998\)015<0809:TTRMMT>2.0.CO;2](https://doi.org/10.1175/1520-0426(1998)015<0809:TTRMMT>2.0.CO;2),
 1998.
 LeMone, M. A., and E. J. Zipser: Cumulonimbus vertical velocity events in GATE. Part I: Diameter, intensity and mass flux, J.
 Atmos. Sci., 37, 2444–2457, [https://doi.org/10.1175/1520-0469\(1980\)037<2444:CVVEIG>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2444:CVVEIG>2.0.CO;2), 1980.
 Li, Y., Liu
 , Y., Chen, Y., Chen, B., Zhang, X., Wang, W., Shu, Z., and Huo, Z.: Characteristics of Deep Convective Systems and Initiation
 during Warm Seasons over China and Its Vicinity, Remote Sensing, 13, 4289, <https://doi.org/10.3390/rs13214289>, 2021.
 Liu, C. and Zipser, E. J.: Global distribution of convection penetrating the tropical tropopause, J. Geophys. Res. Atmos., 110,
 D23104, <https://doi.org/10.1029/2005JD006063>, 2005.
 Liu, C., Zipser, E. J., and Nesbitt, S. W.: Global distribution of tropical deep convection: different perspectives from TRMM
 infrared and radar data, J. Climate, 20, 489–503, <https://doi.org/10.1175/JCLI4023.1>, 2007.

削除: -

削除: .

削除: gap-part

削除: 860 84

1851 Liu, C. and Zipser, E. J.: Diurnal cycles of precipitation, clouds, and lightning in the tropics from 9 years of TRMM
1852 observations, *Geophys. Res. Lett.*, 35, L04819, <https://doi.org/10.1029/2007GL032437>, 2008.

1853 Liu, C., E. J. Zipser, D. J. Cecil, S. W. Nesbitt, and S. Sherwood: A cloud and precipitation feature database from nine years
1854 of TRMM observations, *J. Appl. Meteorol. Climatol.*, 47, 2712–2728, <https://doi.org/10.1175/2008JAMC1890.1>, 2008.

1855 Liu, C. and Zipser, E. J.: The global distribution of largest, deepest, and most intense precipitation systems, *Geophys. Res.*
1856 *Lett.*, 42, 3591–3596, <https://doi.org/10.1002/2015GL063776>, 2015.

1857 Lucas, C., Zipser, E. J., and LeMone, M. A.: Vertical velocity in oceanic convection off tropical Australia in early morning:
1858 Roles of buoyancy and dynamic pressure, *J. Atmos. Sci.*, 51, 3183–3193, [https://doi.org/10.1175/1520-0469\(1994\)051<3183:VVIOCO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<3183:VVIOCO>2.0.CO;2), 1994.

1860 Luo, Y., Zhang, R., and Wang, H.: Comparing occurrences and vertical structures of hydrometeors between eastern China and
1861 the Indian monsoon region using CloudSat/CALIPSO data. *J. Climate*, 22, 1052–1064,
1862 <https://doi.org/10.1175/2008JCLI2606.1>, 2009.

1863 Luo, Y., Zhang, R., Qian, W., Luo, Z., and Hu, J.: Intercomparison of deep convection over the Tibetan Plateau–Asian
1864 monsoon region and subtropical North America in boreal summer using CloudSat/CALIPSO data, *J. Climate*, 24, 2164–2177,
1865 <https://doi.org/10.1175/2010JCLI4032.1>, 2011.

1866 Luo, Z. J., Liu, C., and Stephens, G. L.: CloudSat adding new insight into tropical penetrating convection, *Geophys. Res. Lett.*,
1867 35, 19, <https://doi.org/10.1029/2008GL035330>, 2008.

1868 Luo, Z. J., Jeyaratnam, J., Iwasaki, S., Takahashi, H., and Anderson, R.: Convective vertical velocity and cloud internal vertical
1869 structure: An A-Train perspective, *Geophys. Res. Lett.*, 41, 723–729, <https://doi.org/10.1002/2013GL058922>, 2014.

1870 Masunaga, H. and Kummerow, C. D.: Observations of tropical precipitating clouds ranging from shallow to deep convective
1871 systems, *Geophys. Res. Lett.*, 33, L16805, <https://doi.org/10.1029/2006GL026547>, 2006.

1872 Maloney, E. D., Gettelman, A., Ming, Y., Neelin, J. D., Barrie, D., Mariotti, A., Chen, C.-C., Coleman, D. R. B., Kuo, Y.-H.,
1873 Singh, B., Annamalai, H., Berg, A., Booth, J. F., Camargo, S. J., Dai, A., Gonzalez, A., Hafner, J., Jiang, X., Jing, X., Kim,
1874 D., Kumar, A., Moon, Y., Naud, C. M., Sobel, A. H., Suzuki, K., Wang, F., Wang, J., Wing, A. A., Xu, X., and Zhao, M.:
1875 Process-Oriented Evaluation of Climate and Weather Forecasting Models, *B. Am. Meteorol. Soc.*, 100, 1665–
1876 1686, <https://doi.org/10.1175/bams-d-18-0042.1>, 2019.

1877 Manabe, S., Smagorinsky, J., and Strickler, R. F.: Simulated climatology of a general circulation model with a hydrologic
1878 cycle, *Mon. Weather Rev.*, 93, 769–798, [https://doi.org/10.1175/1520-0493\(1965\)093<0769:SCOAGC>2.3.CO;2](https://doi.org/10.1175/1520-0493(1965)093<0769:SCOAGC>2.3.CO;2), 1965.

1879 Masunaga, H., L'Ecuyer, T. S., and Kummerow, C. D.: Variability in the characteristics of precipitation systems in the tropical
1880 Pacific. Part I: Spatial structure. *J. Climate*, 18, 823–840, <https://doi.org/10.1175/JCLI-3304.1>, 2005.

1881 Matrosov, S. Y.: Potential for attenuation-based estimations of rainfall rate from CloudSat, *Geophys. Res. Lett.*, 34, 5,
1882 <https://doi.org/10.1029/2006GL029161>, 2007.

1883 Mohr, K. I. and Zipser, E. J.: Defining mesoscale convective systems by their 85-GHz ice scattering signatures, *B. Am.*
1884 *Meteorol. Soc.*, 77, 1179–1190, [https://doi.org/10.1175/1520-0477\(1996\)077<1179:DMCSBT>2.0.CO;2](https://doi.org/10.1175/1520-0477(1996)077<1179:DMCSBT>2.0.CO;2), 1996.

1885 Nesbitt, S. W., and Zipser, E. J.: The diurnal cycle of rainfall and convective intensity according to three years of TRMM
1886 measurements. *J. Climate*, 16, 1456–1475, [https://doi.org/10.1175/1520-0442\(2003\)016<1456:TDCORA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016<1456:TDCORA>2.0.CO;2), 2003.

1887 Pilewskie, J. A. and L'Ecuyer, T. S.: The Global Nature of Early-Afternoon and Late-Night Convection Through the Eyes of
1888 the A-Train, *J. Geophys. Res.-Atmos.*, 127, e2022JD036438, <https://doi.org/10.1029/2022JD036438>, 2022.

1889 Price, C. and Rind, D.: A simple lightning parameterization for calculating global lightning distributions, *J. Geophys. Res.-*
1890 *Atmos.*, 97, 9919–9933, <https://doi.org/10.1029/92JD00719>, 1992.

1891 Puigdomènech Treserras, B., Kollias, P., Battaglia, A., Tanelli, S., and Nakatsuka, H.: EarthCARE's cloud profiling radar
1892 antenna pointing correction using surface Doppler measurements, *Atmos. Meas. Tech.*, 18, 5607–5618,
1893 <https://doi.org/10.5194/amt-18-5607-2025>, 2025.

削除: ..

1895 Ramanathan, V. L., Cess, R. D., Harrison, E. F., Minnis, P., Barkstrom, B. R., Ahmad, E., and Hartmann, D.: Cloud-radiative
1896 forcing and climate: Results from the Earth Radiation Budget Experiment, *Science*, 243, 57-63,
1897 <https://doi.org/10.1126/science.243.4887.57>, 1989.

1898 Romatschke, U., Medina, S., and Houze, R. A., Jr.: Regional, seasonal, and diurnal variations of extreme convection in the
1899 South Asian region, *J. Climate*, 23, 419–439, <https://doi.org/10.1175/2009JCLI3140.1>, 2010.

1900 Sato, K., Okamoto, H., Nishizawa, T., Jin, Y., Nakajima, T. Y., Wang, M., Satoh, M., Roh, W., Ishimoto, H., and Kudo, R.:
1901 JAXA Level 2 cloud and precipitation microphysics retrievals based on EarthCARE radar, lidar, and imager: the CPR_CLP,
1902 AC_CLP, and ACM_CLP products, *Atmos. Meas. Tech.*, 18, 1325–1338, <https://doi.org/10.5194/amt-18-1325-2025>, 2025.

1903 Schiro, K. A. and Neelin, J. D.: Tropical continental downdraft characteristics: mesoscale systems versus unorganized
1904 convection, *Atmos. Chem. Phys.*, 18, 1997–2010, <https://doi.org/10.5194/acp-18-1997-2018>, 2018.

1905 Schumacher, C., Stevenson, S. N., and Williams, C. R.: Vertical motions of the tropical convective cloud spectrum over Darwin,
1906 Australia, *Q. J. Roy. Meteorol. Soc.*, 141, 2277–2288, <https://doi.org/10.1002/qj.2520>, 2015.

1907 Seiki, T., Horie, H., Hagihara, Y., Aoki, S., and Noda, A. T.: Global identification of dominant ice-particle growth in cirrus
1908 clouds using EarthCARE satellite observations, *EGUsphere [preprint]*, doi:10.5194/egusphere-2025-4819, 2025.

1909 Sherwood, S. C., Bony, S., and Dufresne, J.-L.: Spread in model climate sensitivity traced to atmospheric convective mixing,
1910 *Nature*, 505, 37–42, <https://doi.org/10.1038/nature12829>, 2014.

1911 Skofronick-Jackson, G., Petersen, W., Berg, W., Kidd, C., Stocker, E., Kirschbaum, D., Kakar, R., Braun, S., Huffman, G.,
1912 Iguchi, T., Kirstetter, P., Kummerow, C., Meneghini, R., Oki, R., Olson, W., Takayabu, Y., Furukawa, K., and Wilheit, T.:
1913 The Global Precipitation Measurement (GPM) mission for science and society, *B. Am. Meteorol. Soc.*, 98, 1679–1695,
1914 <https://doi.org/10.1175/BAMS-D-15-00306.1>, 2017.

1915 Song, H.J., Kim, S., Roh, S., and Lee, H.: Difference between cloud top height and storm height for heavy rainfall using TRMM
1916 measurements, *J. Meteorol. Soc. Jpn.*, 98, 901-914, <https://doi.org/10.2151/jmsj.2020-044>, 2020.

1917 Stephens, G. L., Vane, D. G., Boain, R. J., Mace, G. G., Sassen, K., Wang, Z., Illingworth, A. J., O'Connor, E. J., Rossow, W.
1918 B., Durden, S. L., Miller, S. D., Austin, R. T., Benedetti, A., Mitrescu, C., and the CloudSat Science Team: The cloudsat
1919 mission and the A-train, *Bull. Amer. Met. Soc.*, 83, 1771–1790, <https://doi.org/10.1175/BAMS-83-12-1771>, 2002.

1920 Stephens, G., Vane, D. G., Tanelli, S., Im, E., Durden, S., Rokeby, M., Reinke, D., Partain, P., Mace, G. G., Austin, R., L'Ecuyer,
1921 T., Haynes, J., Lebsock, M., Suzuki, K., Waliser, D., Wu, D., Kay, J., Gettelman, A., Wang, Z., and Marchand, R.: CloudSat
1922 mission: Performance and early science after the first year of operation, *J. Geophys. Res.-Atmos.*, 113, D8,
1923 <https://doi.org/10.1029/2008JD009982>, 2008.

1924 Sy, O. O., Tanelli, S., Takahashi, N., Ohno, Y., Horie, H., and Kollias, P.: Simulation of EarthCARE spaceborne Doppler radar
1925 products using ground-based and airborne data: Effects of aliasing and nonuniform beam-filling, *IEEE Trans. Geosci. Remote*
1926 *Sens.*, 52, 1463–1479, <https://doi.org/10.1109/TGRS.2013.2251639>, 2014.

1927 Takahashi, H. and Luo, Z. J.: Characterizing tropical overshooting deep convection from joint analysis of CloudSat and
1928 geostationary satellite observations, *J. Geophys. Res.-Atmos.*, 119, 112–121, <https://doi.org/10.1002/2013JD020972>, 2014.

1929 Takahashi, H., Luo, Z. J., and Stephens, G. L.: Level of neutral buoyancy, deep convective outflow, and convective core: New
1930 perspectives based on 5 years of CloudSat data, *J. Geophys. Res. Atmos.*, 122, 2958–2969,
1931 <https://doi.org/10.1002/2016JD025969>, 2017.

1932 Takahashi, H., Luo, Z. J., Stephens, G., and Mulholland, J. P.: Revisiting the Land-Ocean Contrasts in Deep Convective Cloud
1933 Intensity Using Global Satellite Observations, *Geophysical Research Letters*, 50, e2022GL102089,
1934 <https://doi.org/10.1029/2022GL102089>, 2023.

1935 Takahashi, H., Wu, L., Smalley, M. A., Stephens, G., Suzuki, K., and Posselt, D. I.: Land - Ocean Differences in Tropical
1936 Deep Convective Clouds: Intercomparison of DYAMOND Simulations and CloudSat Observations, *J. Geophys. Res.-Atmos.*,
1937 130, 23, <https://doi.org/10.1029/2025JD044688>, 2025.

削除: ..

削除: .

削除: Marc hand

削除: ..

Tanelli, S., Im, E., Mascelloni, S. R., and Facheris, L.: Spaceborne Doppler radar measurements of rainfall: correction of errors induced by pointing uncertainties, *J. Atmos. Oceanic Technol.*, 22, 1676–1690, <https://doi.org/10.1175/JTECH1797.1>, 2005.

Tanelli, S., Durden, S. L., Im, E., Pak, K. S., Reinke, D. G., Partain, P., Haynes, J. M., and Marchand, R. T.: CloudSat’s Cloud Profiling Radar After Two Years in Orbit: Performance, Calibration, and Processing, *IEEE T. Geosci. Remote*, 46, 3560–3573, doi:10.1109/TGRS.2008.2002030, 2008.

Tian, B. and Dong, X.: The Double-ITCZ Bias in CMIP3, CMIP5, and CMIP6 Models Based on Annual Mean Precipitation, *Geophys. Res. Lett.*, 47, e2020GL087232, <https://doi.org/10.1029/2020GL087232>, 2020.

Tiedtke, M.: A comprehensive mass flux scheme for cumulus parameterization in large-scale models, *Mon. Weather Rev.*, 117, 1779–1800, [https://doi.org/10.1175/1520-0493\(1989\)117<1779:ACMFSF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1779:ACMFSF>2.0.CO;2), 1989.

Wehr, T., Kubota, T., Tzeremes, G., Wallace, K., Nakatsuka, H., Ohno, Y., Koopman, R., Rusli, S., Kikuchi, M., Eisinger, M., Tanaka, T., Taga, M., Deghaye, P., Tomita, E., and Bernaerts, D.: The EarthCARE mission – science and system overview, *Atmos. Meas. Tech.*, 16, 3581–3608, <https://doi.org/10.5194/amt-16-3581-2023>, 2023.

Williams, E. R. and Stanfill, S.: The physical origin of the land-ocean contrast in lightning activity, *C. R. Physique*, 3, 1277–1292, [https://doi.org/10.1016/S0169-8095\(02\)00025-1](https://doi.org/10.1016/S0169-8095(02)00025-1), 2002.

Williams, E. R., Mushtak, V., Rosenfeld, D., Goodman, S., and Boccippio, D.: Thermodynamic conditions favorable to superlative thunderstorm updraft, mixed phase microphysics and lightning flash rate, *Atmos. Res.*, 76, 288–306, <https://doi.org/10.1016/j.atmosres.2004.11.010>, 2005.

Xu, W., Chen, H., Wei, H., Luo, Y., and Zhao, T.: Extreme Precipitation Produced by Relatively Weak Convective Systems in the Tropics and Subtropics, *Geophys. Res. Lett.*, 49, e2022GL098048, <https://doi.org/10.1029/2022GL098048>, 2022.

Yang, G.-Y. and Slingo, J.: The diurnal cycle in the tropics, *Mon. Weather Rev.*, 129, 784–801, [https://doi.org/10.1175/1520-0493\(2001\)129<0784:TDCITT>2.0.CO;2](https://doi.org/10.1175/1520-0493(2001)129<0784:TDCITT>2.0.CO;2), 2001.

Yuter, S. E. and Houze Jr., R. A.: Three-dimensional kinematic and microphysical evolution of Florida cumulonimbus. Part II: Frequency distributions of vertical velocity, reflectivity, and differential reflectivity, *Mon. Weather Rev.*, 123, 1941–1963, doi:10.1175/1520-0493(1995)123<1941:TDKAME>2.0.CO;2, 1995.

Zhao, M.: An investigation of the connections among convection, clouds, and climate sensitivity in a global climate model, *J. Climate*, 27, 1845–1862, <https://doi.org/10.1175/JCLI-D-13-00145.1>, 2014.

Zipser, E. J., Cecil, D. J., Liu, C., Nesbitt, S. W., and Yorty, D. P.: Where are the most intense thunderstorms on Earth?, *Bull. Amer. Meteor. Soc.*, 87, 1057–1071, <https://doi.org/10.1175/BAMS-87-8-1057>, 2006.

ページ 8: [1] 削除	Haruka Hotta	2026/06/03 11:46:00
---------------	--------------	---------------------



ページ 8: [2] 削除	Haruka Hotta	2026/06/03 11:46:00
---------------	--------------	---------------------



ページ 8: [3] 削除	Haruka Hotta	2026/06/03 11:46:00
---------------	--------------	---------------------



ページ 8: [4] 削除	Haruka Hotta	2026/06/03 11:46:00
---------------	--------------	---------------------



ページ 8: [5] 削除	Haruka Hotta	2026/06/03 11:46:00
---------------	--------------	---------------------



ページ 8: [6] 削除	Haruka Hotta	2026/06/03 11:46:00
---------------	--------------	---------------------



ページ 10: [7] 削除	Haruka Hotta	2026/06/03 11:46:00
----------------	--------------	---------------------



ページ 12: [8] 削除	Haruka Hotta	2026/06/03 11:46:00
----------------	--------------	---------------------



ページ 19: [9] 書式を変更	Haruka Hotta	2026/06/03 11:46:00
-------------------	--------------	---------------------

フォント : 10 pt, フォントの色 : 自動

ページ 20: [10] 削除

Haruka Hotta

2026/06/03 11:46:00

					cloud top height	cloud top height	
time	region	cloud top height	echo top height (0 dBZ)	echo top height (10 dBZ)	- echo top height (0 dBZ)	- echo top height (10 dBZ)	cloud length

ページ 22: [11] 削除

Haruka Hotta

2026/06/03 11:46:00

ページ 22: [12] 削除

Haruka Hotta

2026/06/03 11:46:00

ページ 22: [13] 削除

Haruka Hotta

2026/06/03 11:46:00

ページ 22: [14] 削除

Haruka Hotta

2026/06/03 11:46:00

ページ 22: [15] 削除

Haruka Hotta

2026/06/03 11:46:00

ページ 22: [16] 削除

Haruka Hotta

2026/06/03 11:46:00

